

# ***Interactive comment on “Water-use dynamics of an alien invaded riparian forest within the summer rainfall zone of South Africa” by Bruce C. Scott-Shaw and Colin S. Everson***

**Bruce C. Scott-Shaw and Colin S. Everson**

brucecharless@gmail.com

Received and published: 5 December 2018

RC1: 'Water-Use Dynamics of an Alien Invaded Riparian Forest Within the Summer Rainfall Zone of South Africa', Anonymous Referee #1, 2 July 2018

HESS-2018-227

Anonymous referee #1 (AR1) is thanked for their thorough review. The comments and suggestions provided were insightful and beneficial to the progress of this paper.

1. AR1 stated that the description of the sampling design and its rationale needs more detail in the text. Papers should be written to allow some level of replicability of the

method to other sites. The comments relating to up-scaling are itemised as follows:

â€¢ How and why were sites, species, and particular trees chosen? â€¢ This has been detailed in Chapter 2.2 which leads into the new Chapter 2.6.

â€¢ How did sites differ topographically, distance from river, soil properties, etc? â€¢ All sites were in close proximity to one another and the river. There were no variations in soils, climate and access to water. This has been mentioned in Chapter 2.2.

â€¢ Why were different numbers of trees of different species chosen? â€¢ This was due to availability of equipment and the associated budget constraints. Additionally, some trees which need to be in close proximity to the logger, do not provide good flow measurements. This limitation has been mentioned in Chapter 2.6.

â€¢ What is the likely or known age range or age structure in the indigenous and the invasive trees at the sites? â€¢ The growth stage of each tree has been included in Table 1.

â€¢ Are the indigenous trees necessarily older? â€¢ The growth stage of each tree has been included in Table 1. However, the authors were measuring the current state and therefore the water-use of the stand as it was during the measurement period. The selection of different size classes was more important than the age of each tree in this regard. Water-use does change with age but the size and LAI are more relevant to water-use. The state of the forest is addressed in the next comment.

â€¢ How did the species and the size classes of the trees compare to that of mature indigenous Eastern Mistbelt forest? â€¢ A description of the typical pristine composition and characteristics of Mistbelt forest has been provided in Chapter 2.1.

â€¢ Are any early successional? â€¢ The growth stage of each tree has been included in Table 1.

â€¢ What is the typical composition of Eastern Mistbelt forest in terms of the proportion of trees that are deciduous? â€¢ Provided in Chapter 2.1. The proportion of deciduous

species is variable in this forest type but the forest has been classified as Mistbelt forest.

â€¢ Deciduousness effects the water use of the trees. Was this proportion mimicked in the selection of trees to monitor? â€¢ Approximately 10% of species in the forest are deciduous, with the remainder being evergreen and semi-deciduous. The sampling design used this proportion in the selection of trees to monitor.

â€¢ Were the Acacia and Eucalyptus trees near maturity? â€¢ The growth stage of each tree has been included in Table 1.

2. Two of the three Acacia's and one of the two Eucalyptus trees measured had larger diameters than the indigenous trees, except for the *L. sericea*. Is statement on pg 7 In 11 that "the introduced species used 2.4 times more water than the indigenous species" made by comparing individual trees of similar sizes or of similar ages? â€¢ This was a general statement based on an average water-use of each species. The size differences are noted and the results represent the status quo of how much water the stand was using during the measurement period. The statement provides an indication of what the invasion is using in comparison to typical indigenous trees that it has replaced.

3. The description of how the scaling up from individual tree water use to stand scale water use also needs more detail. Was the water use from different indigenous trees measured and applied across all trees of all species and size classes across the stand? â€¢ This reiteration of the previous comments has been addressed by Chapter 2.6 and links to the selection of trees and sampling design.

4. Were water use figures of the individual measured trees applied to trees of the same species and/or functional group (e.g. deciduous or not, similar growth form or not, similar wood density) or similar size class? â€¢ Yes, as described in the new Chapter 2.6, water-use was extrapolated per representative size class and species class identified in the density measurements.

[Printer-friendly version](#)

[Discussion paper](#)



5. Wouldn't water use of mature trees be different to the young ones measured in this study? → Yes, however the objective was to capture the age and size distribution of the stand and measure this as accurately as possible with the available equipment.

6. What was assumed to be the size class distribution in the invaded and restored scenarios? The same as current or larger more mature trees assumed? → The hypothetical scenarios used the existing size class distribution for each species class and extrapolated this based on an assumed invasion level.

7. These aspects need to be described in the methods and the effects of the assumptions made, and alternatives, discussed in the discussions. As such the figures of species level and stand level water use should also have some estimates of likely uncertainty. → Comment no. 6 was described in the text as per this recommendation in Chapter 3.4. Due to the number of trees measured in each species class, the statistical level of uncertainty was not practical to include.

#### Specific Comments

1. Pg 2, Ln 21-23: unnecessary to cite the paper twice in the sentence. → Corrected.  
2. Pg3, Ln 35-37: The statement "invasive species use 189% more water than indigenous dominated stands" needs more clarification: this number is too specific to apply to all three of the cited studies. Was this the highest or lowest value from these three studies? Perhaps give the range of values across multiple studies. Does this only refer riparian forests compared to invaded stands? → This was made more clear in the text. This was a global literature review of published studies. It provided a baseline from which the findings of this paper are compared to. "At the ecosystem scale, a comprehensive review of numerous internationally published studies indicate that invasive species use up to 189 % more water than indigenous dominated stands, particularly in tropical moist forests (Nesetto et al., 2005; Yepez et al., 2005; Fritzsch et al., 2006). These findings, typically outside of South Africa are limited to mostly herbaceous species with very few recent studies focusing on measurement of introduced

[Printer-friendly version](#)

[Discussion paper](#)



trees."

3. Pg4, Section 2.2 requires more details in the text as to the numbers of trees of different species and why they were chosen as well as how the scaling up calculations were done. → A new chapter in the Methods (2.6) has been added to provide detail on the up-scaling approach.

4. Pg 7 Ln 43 There is no citation for Everson et al 2016 in the reference list. → This has been added to the reference list.

5. Pg 8 Ln 4 typo: "bcome". → Corrected.

6. Pg 15 Table 1 – typo: "Eucalyptus nitenss". → Corrected.

Please also note the supplement to this comment:

<https://www.hydrol-earth-syst-sci-discuss.net/hess-2018-227/hess-2018-227-AC3-supplement.pdf>

Interactive comment on Hydrol. Earth Syst. Sci. Discuss., <https://doi.org/10.5194/hess-2018-227>, 2018.

Printer-friendly version

Discussion paper

