Hydrol. Earth Syst. Sci. Discuss., https://doi.org/10.5194/hess-2018-264-AC3, 2018 © Author(s) 2018. This work is distributed under the Creative Commons Attribution 4.0 License.



HESSD

Interactive comment

# *Interactive comment on* "Vegetation vulnerability to drought on southeastern Europe" *by* Patrícia Páscoa et al.

### Patrícia Páscoa et al.

ppramos@fc.ul.pt

Received and published: 4 September 2018

1. The title of the manuscript is a mistake, since the content does not refer to other territory than Romania. Eastern Europe holds a very diverse geography, so that doing a study on Romania and pretending it is about Eastern Europe is huge mistake. Such a market strategy is used to attract public and citations, but it is not a fair approach.

Response: It was not our intention to mislead the readers when choosing this title. The study area encompasses several other countries including the whole Moldova, and parts of Ukraine, Hungary, Serbia, Bulgaria, and Slovakia. However, and as we akcnowledge the reviewer's concern, the title was changed to: "Vegetation vulnerability to drought in Romania and Moldova regions".

Printer-friendly version



2. Lines 185-190: "Negative correlations are more frequently found on the months of May and June" - can you explain why the correlations are more frequent in May and June?

Response: This result is discussed on the Discussion section as follows:

'Levanič et al. (2013), when analysing the relation between Pinus nigra tree-ring widths from southern Romania and monthly precipitation, as well as 3-month SPI, found July to be the month with the highest correlation with precipitation, whereas July, August, and September showed a high correlation with SPI. Moreover, SPEI and NDVI correlations respond to the water balance, and SPEI tends to increase when NDVI decreases (Vicente-Serrano et al., 2013). In the study area, monthly mean precipitation reaches its maximum in June (Koleva et al., 2008; Cheval et al., 2011; Spinoni et al., 2015), and the maximum temperature occurs in July (Spinoni et al., 2015; Dascălu et al., 2016), which points to increased soil water stress that could explain the increase in the correlations between SPEI and NDVI.'

3. The same amendments would be nice for the next paragraphs, as they are mostly descriptive and not enough explanations are provided.

Response: A short explanation will be added in order to clarify the referred issue. We initially decided to discuss the results only in the last section, but taking into account the comments from all the reviewers we will include in the Results section some interpretation, so it won't be as descriptive. Accordingly, we will also improve the Discussion section.

4. Why no correlation in June for "the area south of the Carpathian Mountains" (L194) and "The areas showing positive correlations from April to June are mostly agricultural land" (L197), so including the areas south of the Carpathians?

Response: The sentence 'The areas showing positive correlations from April to June are mostly agricultural land' refers to Figure 3, (as is stated). It is also stated that

# HESSD

Interactive comment

Printer-friendly version



this area 'exceeds 37%', which is the value obtained in June at 6 months SPEI. This implies that around 63% of the agricultural land does not present positive correlations in June, and therefore we do not consider that the sentences quoted by the Reviewer are contradictory.

5. Lines 214-219: as mentioned before, the explanations are almost missing, while the very simple description fills the lines. For example, why the standard deviation is lower until July in some forests?

Response: We agree that some details are needed in the results section and accordingly to the comments of the reviewers we will add additional information. In particular, a short explanation will be added in order to clarify the referred issue.

6. The section 3.3 is a very poor description of the drought event 2000/2001, with no explanation and no positioning in the general context of the droughts in Romania, not to say SE Europe as pretended in the title.

Response: The 2000/2001 drought event in SE Europe was chosen for analysis because it is well known and documented, being categorized as one of the worst in the region (http://www.geo.uio.no/edc/droughtdb/edr/DroughtEvents/\_2000\_Event.php) with strong impacts in several sectors, namely agriculture (Sepulcre-Canto et al., 2012). This drought affected several countries, namely Romania, Hungary and Bulgaria (Sepulcre-Canto et al., 2012), as well as Greece, Turkey and the Balkan countries (http://www.geo.uio.no/edc/droughtdb/edr/DroughtEvents/\_2000\_Event.php). The extent of this drought episode is depicted on Fig.1, showing the drought classification, as assessed by SPEI 12, across Europe, in February 2001 (http://spei.csic.es/map/maps.html). A short explanation will be added in Introduction and sections 2.4.3 and 3.3 in order to describe with more detail the severe character of the drought event of 2000/2001.

7. The authors claims that "In this section we assess the impact of the strong drought episode of 2000/2001". In reality, the impact has been mentioned very poorly men-

HESSD

Interactive comment

Printer-friendly version



### tioned in the section.

Response: In this topic we will clarify in our work that the aim is to analyse and assess the drought (characterized by means of SPEI) impacts on vegetation activity, as obtained using remote sensing data (NDVI). The type of analysis throughout the manuscript was already adopted in several works (Ji and Peters, 2003; Lotsch et al., 2003; 2005; Quiring et al., 2010; Brown et al., 2008; Wang et al., 2015; Vicente Serrano et., 2012; 2013; 2014; Gouveia et al., 2017). The scientific recognition of these methodologies to achieve the goals we proposed lead us to apply this well-known methodology to a very interesting region that we thought lacked studying. Accordingly with our goal and the methodology used in the manuscript, the results described on section 3.3 were obtained considering the months of April to October of the year 2000, corresponding to the months of high vegetation activity. We included two figures (Fig. 9 and 10) that fully support our results. The methodology used allowed to guantify the drought duration and area stricken and presenting vegetation under stress conditions (showing low NDVI anomalies). Fig.9 also allow the evaluation of drought occurrence. Additionally, we also assessed the stress shown on each land cover type in Fig.10. These impacts of drought on vegetation dynamics, namely their duration, freguency and extension, were discussed on the Discussion and Conclusions section on Line 345 Moreover, the analysis of the impacts of drought on different socio-economic sectors, such as agriculture or water storage is not the aim of this work, but will be the purpose of an additional work, now in preparation. However, and acknowledging that the main target of our analysis is misleading, we opted by clarify this issue in the above mentioned sentence, as follow: 'In this section we assess the impact of the strong drought episode of 2000/2001 on vegetation activity, as described using satellite data (NDVI)'

8. This is valid with section 4 too, where the authors claim again that "impacts of droughts on vegetation were analysed". The drought impact is actually indirectly tack-led in this paper in the form of the well-known relations between SPEI and NDVI.

# **HESSD**

Interactive comment

Printer-friendly version



Response: Please see answer to previous comment.

References:

Brown, J.F., Wardlow, B.D., Tadesse, T., Hayes, M., J., Reed, B.C., The Vegetation Drought Response Index (VegDRI): A New Integrated Approach for Monitoring Drought Stress in Vegetation, GIScience & Remote Sensing, 45(1), 16-46, 2008

Gouveia, C.M., Trigo, R.M., Beguería, S., Vicente-Serrano, S.M.: Drought impacts on vegetation activity in the Mediterranean region: An assessment using remote sensing data and multi-scale drought indicators, Global Planet. Chan., 151: 15-27. https://doi.org/10.1016/j.gloplacha.2016.06.011, 2017.

Ji, L., Peters, A.J., Assessing vegetation response to drought in the northern Great Plains using vegetation and drought indices, Remote Sensing of Environment, 87(1), 85-98, 2003, https://doi.org/10.1016/S0034-4257(03)00174-3.

Lotsch, A., Friedl, M.A., Anderson, B.T., Tucker, C.J., Coupled vegetationâĂŘprecipitation variability observed from satellite and climate records, Geophysical Research Letters, 30(14), 1774, 2003, doi:10.1029/2003GL017506

Lotsch, A., Friedl, M.A., Anderson, B.T., Tucker, C.J., Response of terrestrial ecosystems to recent Northern Hemispheric drought, Geophysical Research Letters, 32, L06705, 2005, doi:10.1029/2004GL022043

Quiring,S.M., Ganesh, S., Evaluating the utility of the Vegetation Condition Index (VCI) for monitoring meteorological drought in Texas, Agric. For. Meteorol., 150, pp. 330-339, 2010

Vicente-Serrano, S., Beguería, S., Lorenzo-Lacruz, J., Camarero, J.j., López-Moreno, J.I., Azorin-Molina, C., Revuelto, J., Morán-Tejeda, E., Sanchez-Lorenzo, A.: Performance of drought índices for ecological, agricultural, and hydrological applications. Earth Interact, 16, 1-27, https://doi.org/10.1175/2012EI000434.1, 2012

HESSD

Interactive comment

Printer-friendly version



Vicente-Serrano, S.M., Gouveia, C., Camarero, J.J., Beguería, S., Trigo, R., López-Moreno, J.I,m Azorín-Molina, C., Pasho, E., Lorenzo-Lacruz, J., Revuelto, J., Morán-Tejeda, E., Sanchez-Lorenzo, A.: Response of vegetation to drought timescales across global land biomes. P. Natl. Acad. Sci., 110(1), 52-57, https://doi.org/10.1073/pnas.1207068110, 2013.

Vicente-Serrano, S. M., Camarero, J. J. and Azorin-Molina, C.: Diverse responses of forest growth to drought time-scales in the Northern Hemisphere. Global Ecol. Bio-geogr., 23: 1019–1030, doi:10.1111/geb.12183, 2014.

Wang, H., Chen, A., Wang, Q., He, B., Drought dynamics and impacts on vegetation in China from 1982 to 2011, Ecological Engineering, 75, 303-307, 2015. https://doi.org/10.1016/j.ecoleng.2014.11.063


Interactive comment on Hydrol. Earth Syst. Sci. Discuss., https://doi.org/10.5194/hess-2018-264, 2018.

# **HESSD**

Interactive comment

Printer-friendly version







Interactive comment

Printer-friendly version

**Discussion paper** 



Fig. 1. SPEI 12 across Europe, in February 2001

1960

1970

-3 1950

1980

1990

2000

2010