

Dear Dr. Freer,

Please find our reviewer responses and descriptions of revisions to HESS-2018-266 below. In addition to the requested revisions, we have also improved the manuscript with these additional revisions: 1) revised all figures for improved clarity, 2) added one paragraph of new results that highlights additional nuances regarding stream temperature dynamics and associated basin properties, 3) fully revised and expanded the discussion section. We hope you find the revision and responses to reviewer comments satisfactory and look forward to future correspondence regarding this manuscript. Best regards, Dan Isaak

Reviewer #1 comments

General comments

This manuscript presents an elegant analysis of different components (magnitude, frequency, duration, timing) of thermal events for a large number of time series points in the United states. Two variations of Principal Components Analysis (T-mode and S-mode analyses) refine the analysis very nicely into spatial regions and temporal seasons of thermal homogeneity and seasonality. By disaggregating time series into metrics, and accounting for high levels of redundancy between metrics, together with the PC analyses, this research presents a novel approach to optimising site locations for water temperature gauging networks. In my opinion, this is a very useful addition to thermal research in lotic systems. The approach is generic and applicable to a global audience. The manuscript is clearly and well written, methodologically elegant and scientifically sound. I recommend publication given minor comments corrected below.

[Our response: We much appreciate the reviewer's kind words and attention to detail in their comments. We have made many of the suggested revisions as described below.](#)

Specific comments

Section 3.1 – Does one need to specify that the study assumed stationarity in the data, in order to generate temperatures for 365 days based on five-year time series?

[Our response: There has been some focus in the recent literature on the possibility of nonstationary responses in stream temperatures due to climate forcing. However, that type of nonstationarity is generally expected over multi-decadal timespans and the prediction is based largely on mechanistic models rather than documentation from empirical trends in monitoring datasets. During the short five year study period we considered, nonstationarity was unlikely to be important and the 12% of missing daily observations were reconstructed from nearby sites with strong covariance using the missMDA statistical package in R.](#)

Line 207 – Please provide a summary of the environmental gradients; it may be worth including a table on these.

[Our response: Table 1 was expanded to include summaries of these gradients, which were elevation, drainage area, annual flow, and reach slope.](#)

Line 212 – Please explain why the thermal year started on 1 December. In South Africa, we typically use 1 October – 30 September for the Hydrological year, but I am aware that this varies regionally, being based on the onset of the highest discharge season.

Our response: The water year in North America is also considered to start October 1 but in the climatological literature, seasons are slightly different and considered to be winter (Dec/Jan/Feb), spring (March/April/May), summer (June/July/Aug), and fall (Sept/Oct/Nov). Climatological considerations seemed more relevant in the present context, and the conventional three month seasonal periods also conveniently matched the water temperature patterns common to our study site. We describe our rationale for starting the thermal year on Dec 1 in the manuscript where we state “We started the thermal year on December 1 because temperatures usually reach their annual lows by this date and the 3-month period thereafter constituted a logical winter season (i.e., December, January, February).”

Line 234 – It would make more sense to me to represent the thermal gradient per 100m. This would be a useful figure in defining a water temperature lapse rate. For air temperatures, this is typically expressed as something like 0.7°C per 100m.

Our response: We agree and revised the values in Figure 2 accordingly.

Technical comments

Title and elsewhere in text: please check for correct spelling of “Principle [as in components]”, which needs to be corrected to PRINCIPAL and checked throughout text, as there are instances of both. Nothing serious – I get confused between these two spellings!

Our response: Embarrassing on our part that we missed this. Inconsistent usage was changed to “principal” throughout.

Lines 37-40: Sentence does not read well. Suggested revision “Knowledge of the local thermal regime, based on the annual sequence of temperatures characteristic to specific locations within a river network, is key to understanding natural conditions and diagnosing anthropogenic impairments.”

Our response: This revision was made.

Lines 62-64: Suggested revision “While that may bring..., most warm stream...correlated with each other and therefore redundant. If redundancy is also reflected across a broader...”

Our response: This revision was made.

Paragraph beginning line 146: Be explicit that these time series refer to water temperatures, as later on in the manuscript air temperatures are also used.

Our response: This revision was made.

Line 218 – “sites, an S-mode”

Our response: This change was made.

Line 253 – Figure 4a

Our response: This change was made.

Line 257 – insert Figure 4c

Our response: This change was made.

Table 1 – write US in full; standardise on number of decimal points down columns (also applies for Table 3).

Our response: United States was spelled out. In the tables, we generally standardized on having two or three significant digits rather than the number of decimal points. We are glad to adjust this either way depending on the convention in HESS but have left the values unchanged for now.

Figure 2 – I like this figure! Please include the range of R2 values, and I would recommend that the caption explicitly describes the month(s) with the highest thermal gradient.

Our response: We added the R2 values to the figure but did not modify the caption to highlight a subset of months with the highest thermal gradients because we don't think that information is inherently more useful than that for other months.

Figure 4 – caption revision to say "...show principal component scores for axes 1-2...". Please also check there are no other occurrences of "principle".

Our response: These revisions were made.

Figure 7 – "...and discharge (c-f)"

Our response: Caption was revised accordingly.

References: Carlisle et al. 2017; Fuhrman et al. 2018; Isaak et al. 2016b; Josse and Husson 2012; Steel et al. 2017 not cited in text.

Our response: These errors were corrected.

Inconsistencies in citations: Line 51 – Rieman et al 2015a; Line 80 Piechota 2001 or 1997?; Line 84 Gallacher 2016 or 2017?; line 90 Trumbo et al. Not referenced; line 175 – correct to R Development Core Team; Line 205 correct to SAS Institute Inc.; line 326 – spelling of Nusslé; line 352 – Jackson et al. 2017 or 2018?

Our response: These inconsistencies were rectified.

Table 3 not cited in text.

Our response: An appropriate citation was added.

Reviewer #2 comment:

Anonymous Referee #2

Received and published: 28 August 2018

This manuscript provides a nice analysis, characterizing the spatial and temporal characteristics and controls of thermal regimes of stream water. The work is based on a novel application of Principal Component Analysis, including the highly interesting differentiation of T-mode and S-mode PCA to illustrate both, temporal and spatial consistency of the stream temperature pattern. The paper is very well and concisely written, including a clear and complete description of the data and methods used. However and despite the flawless implementation of the analysis, the interpretation of the results and their implications remain somewhat superficial. After reading the manuscript, it seemed to me that the authors contented themselves with demonstrating how a well-known statistical tool can be applied with stream temperature data. The one finding that I found most interesting to demonstrate the value of PCA was that the authors could pin down the timing of the phase transitions. I may not see the forest for the trees but apart from that I am not sure what can be learned from the analysis. As far as I understand, the results essentially suggest that (1) stream temperature is mostly controlled by temperature magnitudes and lengths of winter periods (which again is related to temperature magnitude one would assume) and (2) stream temperature is more spatially homogeneous in winter than in summer. While the first does not really come as a surprise, it seems that the latter can also be inferred without PCA (or in other words: how is the information content of Figure 2 different to that of Figure 6?). I would thus be glad if the authors could invest a bit more effort in (1) highlighting the benefits of PCA with respect to other methods and (2) providing a somewhat stronger synthesis of their results – what are the novel aspects that can be learned from these results?

Our response: We agree with the overall critique that greater interpretation of results would be beneficial so have revised and expanded the discussion in a subsequent revision. As for the reviewer's first comment that "(1) stream temperature is mostly controlled by temperature magnitudes and lengths of winter periods (which again is related to temperature magnitude one would assume)", the statement in the parenthetical clause is incorrect in conflating temperature magnitude with the length of the winter period. Our analysis reveals that these are instead two distinct aspects of thermal regimes in the mountain streams we studied. Streams with similar mean or maximum summer temperatures appear to vary considerably with regards to their winter period lengths when temperatures are largely homothermous. Exploring why that variation occurs was a useful addition to a discussion revision. The reviewer's latter point that "(2) stream temperature is more spatially homogeneous in winter than in summer. [...], it seems that the latter can also be inferred without PCA (or in other words: how is the information content of Figure 2 different to that of Figure 6?)" is accurate but had previously been documented only at a few sites using time series plots like Figure 2. The T-mode PCA results put that site-level pattern into a broader context composed of hundreds of sites across large river basins. In this particular dataset, the thermal pattern across all the sites during the winter was largely consistent but that consistency was unknown prior to the analysis. Moreover, it is unlikely to be repeated in subsequent analyses we are planning with larger datasets that encompass greater climatic and hydrological diversity, so these PCA tools may help us identify subdomains regionally wherein stream thermal regimes behave differently. Here again, we think the revised discussion section has done much to bring out these points.

Technical comments:

p.7,1.204: what is a “Princomp procedure”?

Our response: This was the statistical script run in the SAS software to perform the analysis. The reference to SAS was moved forward in this sentence so it is adjacent to the “Princomp procedure” reference for clarity.

p.7,1.212: is there a specific reason to run the T-mode PCA on the 5-year mean values of the daily mean temperatures? In other words, why use 365 days (i.e. columns) and not the full data set of 1826 as in the S-mode analysis?

Our response: We judged it unlikely that appreciable inter-annual differences would be observed in the spatial phases revealed by the T-mode analysis given the large elevational gradient in the study area and because the dominant patterns in PC loadings were driven by cold and warm season cycles (Figure 5). Showing one annual cycle of tradeoffs between PC1 and PC2 was easier to present and read, so we elected to run the analysis on the 5-year mean daily values. We were less certain regarding the potential consistency of inter-annual variation in temporal patterns described using the S-mode analysis, so ran that analysis on the disaggregated water temperature records as well. In retrospect, the results based on the disaggregated records yielded similar insights as those based on the 5-year mean dailies, so little new information was gained except to re-enforce the fact that water temperatures respond strongly to variation in air temperature and discharge across a range of climate year conditions.

other than being repeated 4 more times in the plot of loadings. Displaying the pattern over the course the dominant annual scale variability should be more informative for readers and more easily grasped.

p.18,table 1: the values for reach slope seem excessively small. Should the unit perhaps be [m/m]? Please check.

Our response: Yes, the units were in m/m rather than % and the label was changed accordingly.

1 **Principal components of thermal regimes in mountain river networks**

2 Daniel J. Isaak, Charles H. Luce, Gwynne L. Chandler, Dona L. Horan, Sherry P. Wollrab
3 U.S. Forest Service, Rocky Mountain Research Station, Aquatic Sciences Lab, Boise, ID 83702
4 Correspondence to: Daniel Isaak (disaak@fs.fed.us)

5
6 **Abstract.** Description of thermal regimes in flowing waters is key to understanding physical
7 processes, enhancing predictive abilities, and improving bioassessments. Spatially and temporally
8 sparse datasets, especially in logistically challenging mountain environments, have limited studies
9 on thermal regimes but inexpensive sensors coupled with crowd-sourced data collection efforts
10 provide efficient means of developing large datasets for robust analyses. Here, thermal regimes are
11 assessed using annual monitoring records compiled from several natural resource agencies in the
12 northwestern United States that spanned a five-year period (2011–2015) at 226 sites across several
13 contiguous montane river networks. Regimes were summarized with 28 metrics and principal
14 components analysis (PCA) was used to determine those metrics which best explained thermal
15 variation on a reduced set of orthogonal axes. Four principal components (PC) accounted for 93.4%
16 of the variation in the temperature metrics, with the first PC (49% of variance) associated with
17 metrics that represented magnitude and variability and the second PC (29% of variance) associated
18 with metrics representing the length and intensity of the winter season. Another variant of PCA, T-
19 mode analysis, was applied to daily temperature values and revealed two distinct phases of spatial
20 variability—a homogeneous phase during winter when daily temperatures at all sites were < 3 °C
21 and a heterogeneous phase throughout the year’s remainder when variation among sites was more
22 pronounced. Phase transitions occurred in March and November, and coincided with the abatement
23 and onset of subzero air temperatures across the study area. S-mode PCA was conducted on the
24 same matrix of daily temperature values after transposition and indicated that two PCs accounted
25 for 98% of the temporal variation among sites. The first S-mode PC was responsible for 96.7% of
26 that variance and correlated with air temperature variation ($r = 0.92$) whereas the second PC
27 accounted for 1.3% of residual variance and was correlated with discharge ($r = 0.84$). Thermal
28 regimes in these mountain river networks were relatively simple and responded coherently to
29 external forcing factors, so sparse monitoring arrays and small sets of summary metrics may be
30 adequate for their description. PCA provided a computationally efficient means of extracting key
31 information elements from the temperature dataset used here, and could be applied broadly to
32 facilitate comparisons among more diverse stream types and develop classification schemes for
33 thermal regimes.

34
35 **1 Introduction**

36 Temperatures of flowing waters control many physicochemical processes (Likens and Likens, 1977;
37 Gordon et al., 1991; Ducharme, 2007) and affect the ecology of aquatic organisms and communities
38 (Isaak et al., 2017a; Neuheimer and Taggart, 2007; Woodward et al., 2010). Knowledge of thermal
39 regimes, characterized as the annual sequence of temperature conditions specific to locations within
40 river networks (Caissie, 2006), is key to understanding natural conditions and diagnosing
41 anthropogenic impairments. Seminal work by Poff and colleagues (Poff and Ward, 1989; Poff et al.,
42 1997) created a robust framework for describing flow regimes based on metric descriptions of
43 magnitude, frequency, timing, duration, and variability that are largely transferrable to thermal

Deleted: nce

Deleted: s

Deleted: large

Deleted: s

Moved (insertion) [4]

Moved up [4]: Knowledge of thermal regimes, characterized as the annual sequence of temperature conditions specific to locations within river networks (Caissie, 2006), is key to understanding natural conditions and diagnosing anthropogenic impairments.

52 regimes (Poole et al., 2004; Olden and Naiman, 2010). Recent studies have contributed useful
53 derivations of temperature metrics (Arismendi et al., 2013; Chu et al., 2010; Rivers-Moore et al.,
54 2013; Steel et al., 2016) or classification schemes based on a small number of pre-selected metrics
55 (Maheu et al., 2016) but the limited availability of annual temperature records (Orr et al., 2015;
56 Isaak et al., 2018a) has slowed broad development and adoption of thermal regime concepts. Data
57 inadequacies are often compounded for montane riverscapes that are difficult to sample (Brown and
58 Hannah, 2008; Isaak et al., 2013), a shortfall that needs to be overcome given the importance of
59 these areas as climate refugia for cold-water biodiversity (Brown et al., 2009; Isaak et al., 2016a;
60 Quaglietta et al. 2018) and as the focus of costly regional conservation strategies (Roni et al., 2002;
61 Rieman et al., 2015).

62
63 Despite existing limitations, the importance of temperature to stream biota is well recognized and
64 inculcated to regulatory standards based on metrics used within threshold-based approaches (Poole
65 et al., 2004; Todd et al., 2008). Most often, those metrics represent some aspect of conditions during
66 warm summer months when temperature sensitive species or life stages are thought to be most
67 vulnerable (Ice et al., 2004; McCullough, 2010), which contributes to the preponderance of short
68 monitoring records spanning only these months (Isaak et al., 2017b). However, thermally mediated
69 ecological processes occur throughout the year (Neuheimer and Taggart, 2007; Olden and Naiman,
70 2010), so adequate understanding requires broader characterization of thermal conditions from
71 annual datasets. While that may bring additional complexity, most warm season metrics are strongly
72 correlated and therefore redundant (Isaak and Hubert, 2001; Dunham et al., 2005; Steel et al., 2016).
73 If redundancy is also the norm among a broader array of annual temperature metrics, then
74 multivariate data reduction techniques might be useful for identifying a few key aspects of thermal
75 regimes.

76
77 Supporting that idea, Rivers-Moore et al. (2013) used Principal Components Analysis (PCA) to
78 describe covariation among 39 temperature metrics calculated for 82 South African stream sites and
79 found that two PCs accounted for 75% of the total variation among metrics. Similarly in the field of
80 hydrology, Olden and Poff (2003) examined 171 flow metrics calculated from 420 gage sites across
81 the United States (U.S.) and found that two to four PCs accounted for 76–97% of variation in the
82 dataset. In addition to metric-based PCA that is commonly used in the hydrological sciences,
83 several other PCA variants are standard analytical tools in the field of climatology and may be

Deleted: are

Deleted: .

86 relevant for characterizing the dynamics of thermal regimes (Richman, 1986; Demsar et al., 2013).
87 Most notably, PCA can be done on repeated measurements of a single variable to identify common
88 spatial or temporal behavior among monitoring stations. In the climatology literature, for example,
89 empirical orthogonal function analysis (S-mode PCA in the taxonomy of Richman (1986)) is used
90 to determine which sites covary temporally as a means of developing regionalization schemes for
91 precipitation, air temperatures, or wind speeds (Piechota et al., 1997; Jimenez et al., 2008; Martins
92 et al., 2012). If common temporal patterns are identified, it suggests potential redundancy in the
93 monitoring network and the information can be used to refine future sampling designs. The closely
94 allied T-mode PCA identifies dominant spatial patterns in datasets and the times when these phases
95 occur (Richman, 1986; Gallacher et al., 2017). A single dominant spatial pattern suggests the spatial
96 distribution of a variable is temporally consistent whereas more than one spatial phase suggests
97 change points and different states.

Deleted: alternate

98
99 The advent of inexpensive sensors, combined with regulatory requirements and concerns about
100 climate change, have led to the recent expansion in temperature monitoring networks for rivers and
101 streams (Isaak et al., 2010; Rivers-Moore et al., 2013; Hilderbrand et al., 2014; Luce et al., 2014a;
102 Trumbo et al., 2014; Hannah and Garner, 2015; Jackson et al., 2016; Molinero et al., 2015; Daigle
103 et al., 2016; Mauger et al., 2016; Steele et al., 2016). What was once a data dearth is becoming a
104 deluge and opportunities exist to study thermal regimes with robust datasets. Here, we use annual
105 temperature records compiled from several natural resource agencies for 226 monitoring sites in a
106 mountainous landscape to conduct an initial assessment of thermal regimes. We limit the
107 geographic scope of our effort to several adjacent river basins in the northwestern U.S. that are
108 geologically and topographically similar but which have particularly dense monitoring networks to
109 maximize analytical flexibility. Our objectives were to: 1) provide a basic description of the annual
110 thermal characteristics in mountain rivers and streams because these are rare within the literature, 2)
111 develop metrics to describe thermal regime characteristics based on magnitude, frequency, timing,
112 duration, and variability, and 3) explore spatiotemporal variation among those metrics and
113 temperature dynamics in relation to basin morphology and hydroclimatic conditions to better
114 discern the principal components of thermal regimes and their regulating factors.

Deleted: Luce et al., 2014

Deleted: and

Deleted: redundancy and patterns of

Deleted: co

Deleted: nce

Deleted: monitoring sites to

Deleted: determine the principal components of thermal regimes in montane river networks

116 2 Study area

126 The study area encompasses 79,500 km² of mountainous, topographically complex terrain that
127 spans a broad elevation range of 200–3,600 m [at a latitude of 45° N](#) (Figure 1). Climate is
128 characterized by cold, wet winters with moderate to heavy snow accumulations at high elevations
129 and hot, dry summers. Hydrographs are typical of snowmelt runoff systems, with high flows during
130 spring and early summer and low flows during late summer, fall, and winter (Figure 2). Vegetation
131 is dominated by conifer forests except at low elevations and south facing aspects where grasses and
132 shrubs predominate. Wildfires are common within the landscape and burned 8% of the area from
133 2011 to 2015 (Morgan et al., 2014). Parent geology consists mostly of resistant granites of the Idaho
134 Batholith and a smaller easterly portion of intrusive volcanics (Bond and Wood, 1978; Meyer et al.,
135 2001). Both geologies are heavily dissected and stream valleys are V-shaped except for some alpine
136 valleys at the highest elevations that were once glaciated. Human population densities are low
137 except along wider segments of river valleys where fertile floodplains and easy access to water
138 accommodate small amounts of agriculture and ranching. Most of the study area is publically
139 owned (81%) and federally administered by the United States National Forest Service and Bureau
140 of Land Management for a variety of land-use, recreational, and conservation purposes. Unpaved
141 road networks have been developed in some drainages for timber harvest but many drainages are
142 protected in large wilderness areas with minimal anthropogenic effects or roads (Swanson, 2015).

143

144 **2.1 River networks and temperature dataset**

145 Rivers and streams within the study area were delineated using the 1:100,000-scale National
146 Hydrography Dataset (NHD; <http://www.horizon-systems.com/NHDPlus/index.php>; McKay et al.,
147 2012), which was attributed with mean annual flow values from data at the Western U.S. Stream
148 Flow Metrics website
149 (http://www.fs.fed.us/rm/boise/AWAE/projects/modeled_stream_flow_metrics.shtml; Wenger et
150 al., 2010). To highlight the perennial subset of the network where temperature monitoring occurred,
151 reaches with annual flows less than 0.03 m³/s were removed from the network, as were reaches with
152 channel slopes >15%, and those coded as intermittent in the NHD (Fcode = 46003). Filtering
153 reduced the original network extent from 58,000 km to 29,600 km with streams flowing at
154 elevations of 221–3,105 m. To visualize thermal heterogeneity in the network, a scenario
155 representing mean August temperatures for a baseline climate period of 1993–2011 was
156 downloaded from the Northwestern Stream Temperature website (NorWeST:
157 <https://www.fs.fed.us/rm/boise/AWAE/projects/NorWeST.html>; Isaak et al., 2016b) and linked to

158 the NHD reaches (Figure 1). Several large rivers drain the area in a generally westerly direction, the
159 largest of which is the Salmon River with a mean annual discharge of 315 m³/s and a basin that
160 comprised 44% of the study area. Six large dams and reservoirs occur in downstream portions of the
161 network (three in the Boise River basin, two in the Payette River basin, and one in the Clearwater
162 River basin) but these affect thermal conditions in less than 300 km of river and no temperature data
163 were used from these sections. Spatial attributes and environmental characteristics of the study area
164 network are summarized in Table 1.

Deleted: s

166 To obtain a water temperature dataset for analysis, we intersected the filtered network with the
167 NorWeST database of daily temperature summaries (Chandler et al., 2016) and extracted data for
168 sites that had mean daily temperature values on at least 70% of the days from December 1, 2010 to
169 November 30, 2015. We started the thermal year on December 1 because temperatures usually
170 reach their annual lows by this date and the 3-month period thereafter constituted a logical winter
171 season (i.e., December, January, February). Subsequent three-month periods were considered to be
172 spring (March, April, May), summer (June, July, August), and fall seasons (September, October,
173 November). NorWeST temperature records were supplemented with additional data solicited from
174 hydrologists and fisheries biologists employed by the Idaho Department of Fish and Game and the
175 U.S. Forest Service, and we also downloaded data from online databases maintained by the
176 Columbia Habitat Monitoring Program (<https://www.champmonitoring.org/Home/Index>) and the
177 NOAA Northwest Fisheries Science Center (<https://www.webapps.nwfsc.noaa.gov/WaterQuality/>).
178 Geographic gaps in monitoring were identified using geospatial analysis (e.g., Jackson et al., 2016)
179 and additional sensors were strategically deployed where needed (Isaak et al., 2010; 2013). Data
180 from the different sources were recorded at a variety of sub-daily intervals, so records were
181 summarized to mean daily temperatures for standardization. Data were collected using different
182 sensor models (TidbiT, Stowaway, and Pendant models from Onset Computer Corporation,
183 Pocasset, Massachusetts, USA; Temp101a model from MadgeTech, Warner, New Hampshire,
184 USA), which had measurement accuracies of +/-0.2°C to +/-0.5°C and resolutions of 0.02°C to
185 0.14°C based on manufacturer specifications and calibration tests we performed. Sensors were
186 deployed using underwater epoxy or steel cables for connection to large boulders and other
187 immobile channel structures and were shielded from direct sunlight (Isaak et al., 2013; Stamp et al.,
188 2014). Temperature records were subject to standard quality assurance-quality control measures as
189 described elsewhere (Chandler et al., 2016).

191
192 The stream temperature dataset consisted of records from 226 sites across a range of elevations,
193 stream sizes, and reach slopes (Figure 1; Table 1). Although we set the minimum threshold for
194 record completeness at 70% during the five-year period, the average completeness of records was
195 higher at 88%. Missing daily values were imputed using the MissMDA package (Missing Values
196 with Multivariate Data Analysis; Josse and Husson, 2016) in R (R Development Core Team, 2014)
197 because temporal covariation among proximate stream temperature sites is usually strong. That was
198 confirmed in our dataset by the high correlations between observed daily temperatures and
199 predictions from the imputation technique, which ranged from $r = 0.98$ to 0.99 . All temperature
200 records at the 226 sites were complete after imputation and consisted of 1,826 mean daily
201 temperatures from December 1, 2010 to November 30, 2015. Climatological variation during the
202 same period was described using discharge data downloaded from the National Water Information
203 System database (<https://waterdata.usgs.gov/usa/nwis/nwis>) for a high-elevation gage site at 1,850
204 m and a low-elevation gage site at 294 m and air temperature data from monitoring stations in the
205 Cooperative Observer Network (<https://www.ncdc.noaa.gov/data-access>) that were near the gage
206 sites (Figure 1).

207

208 **3 Data analysis**

209 **3.1 PCA of thermal metrics**

210 Prior to calculating metrics for thermal characteristics, mean daily temperatures for 365 days were
211 calculated from the five years of data at each site to provide representative values. Twenty-eight
212 temperature metrics were then calculated to describe aspects of those annual records based on five
213 categories associated with magnitude, variability, frequency, timing, and duration (Tables 2 and 3).
214 Metrics were similar to those used in previous studies of thermal regimes (Arismendi et al., 2013;
215 Chu et al., 2010; Rivers-Moore et al., 2013; Steel et al., 2016) and in studies assessing the effects of
216 peak summer temperatures on the distribution and abundance of aquatic organisms (Dunham et al.,
217 2003; Huff et al., 2005; Isaak et al., 2017a). A wide range of variability occurred among sites where
218 mean annual temperatures ranged from 3.1 °C to 10.3 °C and annual standard deviations ranged
219 from 2.51 °C to 7.40 °C (Table 3). Relationships among the thermal metrics were described by
220 conducting PCA on a data matrix in which columns represented the 28 metrics and rows were the
221 226 monitoring sites. Linear combinations of the data were estimated with coefficients equal to the
222 eigenvectors of their correlation matrix, which were the principal components (PCs; Pearson, 1901;

223 Sergeant et al., 2016). The first principal component accounted for the largest possible variance in
224 the dataset and succeeding components accounted for the largest portions of the remaining variance
225 while being orthogonal (i.e., uncorrelated) to the preceding components. Correlations, or loadings,
226 between each metric and the PCs were also calculated to assist in subsequent interpretations. The
227 Princomp Procedure in SAS (SAS Institute Inc., 2015) was used to conduct the PCA. To describe
228 geographical relationships, PC scores were mapped to the 226 temperature sites and bivariate
229 correlations were calculated with descriptors of network conditions such as elevation, reach slope,
230 and discharge summarized in Table 1.

Deleted: (SAS Institute Inc., 2015)

231

232 3.2 PCA of daily water temperatures

233 To assess the consistency of spatial temperature patterns among monitoring sites, a T-mode PCA
234 (Richman, 1986) was done on a data matrix of mean daily temperatures in which the columns were
235 the 365 days starting on December 1 and the rows were the 226 monitoring sites. In this analysis,
236 the number of principal components explaining significant variation indicates the number of distinct
237 spatial phases that occur throughout the year (Gallacher et al., 2016). Eigenvector loadings on the
238 dominant PCs were plotted for each day of the year to describe when each phase occurred, and
239 mean daily temperatures were mapped during these periods for visualization.

240

241 To assess temporal covariance among sites, an S-mode PCA (Richman, 1986) was done by
242 transposing the T-mode data matrix so that monitoring sites were columns and the time ordered
243 daily mean temperatures were rows. Because hydroclimatic conditions among years could have
244 affected the results, the S-mode PCA was done not only for the five-year averages of daily water
245 temperatures but also on the disaggregated time series of 1,826 daily values at the 226 monitoring
246 sites. Concordance between the S-mode PC scores, air temperature, and discharge were examined
247 posthoc by plotting standardized time-series and calculating bivariate correlations.

248

249 4. Results

250 Water temperatures within the study area network exhibited spatial and temporal variation that
251 reflected the local topography and annual hydroclimatic cycle. The annual temperature cycle is
252 illustrated in Figure 2 by the slopes of linear regressions between mean monthly temperatures and
253 elevation at the 226 monitoring sites throughout the course of the year in 2013. No elevation trend
254 occurred during cold winter months when many sites had water temperatures at or near 0°C and

Deleted: ive of

Deleted: s

Deleted: one

259 were frequently exposed to subzero air temperatures. As temperatures warmed during the spring a
260 small elevation trend appeared, which became most pronounced (approximately $-0.37^{\circ}\text{C} / 100 \text{ m}$)
261 during peak temperatures in the months of July and August. Examples of inter-annual variation are
262 shown in Figure 3, which contrasts the extreme conditions observed in 2011 and 2015. The former
263 year was relatively cool with a large winter snow accumulation and spring runoff, whereas 2015
264 had below average snowfall, low runoff, and particularly warm early summer air temperatures. As a
265 result, the median discharge date occurred 1–2 months earlier in 2015 than in 2011 and peak water
266 temperatures were 4–5 °C warmer.

267
268 Four PCs accounted for 93.4% of the variation in the 28 temperature metrics (Table 4). The first PC
269 explained 49% of the variation and was strongly correlated with metrics that represented magnitude
270 and variability during most seasonal periods. Correlations between PC1 scores and elevation ($r = -$
271 0.59) and mean flow ($r = 0.58$) suggested gradients in these network characteristics were important
272 controls on this component of thermal regimes (Table 5). PC2 explained 29% of thermal variation
273 and represented the length and intensity of the winter period, with strong loadings for mean winter
274 temperature, minimum temperature, and timing metrics that determined growing season length. PC3
275 accounted for 9.8% of total variation and was associated with summer temperature variability and
276 two timing metrics, whereas PC4 accounted for 5.6% of thermal variance. An ordination plot of
277 scores from the two dominant PCs showed a symmetrical distribution except for several sites with
278 large positive scores on the first axis that were from large rivers at low elevations and had the
279 warmest temperatures (Figure 4a). A map of PC1 scores indicated that the spatial pattern in
280 magnitude and variability (Figure 4b) was congruent with the network scenario of mean August
281 temperatures as would be expected (Figure 1). In fact, the correlation between PC1 scores and the
282 NorWeST August scenario predictions at the 226 monitoring sites was strong at $r = 0.86$. The PC2
283 map showed several clusters of stream sites with high scores scattered throughout the study area
284 (Figure 4c), which tended to be associated with lower reach slopes (Table 5).

285
286 In the T-mode analysis, the first two PCs explained 88% of the total variation in mean daily
287 temperatures. A plot of the daily eigenvector loadings indicated that one distinct spatial phase
288 occurred in the winter and a second phase spanned the year's remainder (Figure 5). Phase
289 transitions occurred around days 100 and 350, which closely aligned with the abatement and onset
290 of subzero air temperatures in the study area (Figure 2). Figure 6 illustrates the spatial patterns

Deleted: where

Deleted: were low

293 characteristic of the two phases by mapping mean daily water temperatures at the monitoring sites
294 on days 50 and 250, which occurred in mid-January and late July, respectively. Temperatures
295 during the winter phase were spatially homogenous and exhibited a narrow range from 0 °C to 2.5
296 °C whereas the non-winter phase was heterogeneous and had a broader temperature range from 7.6
297 °C to 23.4 °C.

298
299 In the S-mode analysis, the first PC accounted for 98% of the variation when applied to the average
300 year of 365 daily temperatures at the 226 monitoring sites. Nearly an identical result was obtained
301 when the analysis was repeated on the disaggregated time-series of 1,826 daily temperatures, as
302 PC1 then explained 96.7% of total variation (Figure 7a). The correlation between PC1 scores and
303 mean daily air temperatures in the disaggregated series was strong ($r = 0.92$), suggesting that water
304 temperatures were responding coherently to variability in air temperatures across the study area. A
305 second PC accounted for 1.3% of water temperature variation in the disaggregated series and was
306 strongly correlated with variation in mean daily discharge ($r = 0.84$). A plot of PC1 versus PC2
307 indicated that variation along these axes corresponded to monthly and seasonal periods (Figure 7b).
308 As was expected, little variation occurred during the cold winter months but during spring and early
309 summer, variation was observed along both axes as air temperatures warmed and snowmelt runoff
310 created a large discharge pulse. Once discharge returned to baseflow conditions in late summer,
311 variability along PC1 was the primary signal until air temperatures cooled significantly in late fall
312 and the homothermous period began.

313
314 Although PC1 and PC2 are linearly uncorrelated, the loop structure of Figure 7b indicates there was
315 some mutual information and that one driver of temperature variation was out of phase with the
316 other. Examining this more closely by plotting the site loading values on each component from the
317 S-mode analysis, we see little variability among the loadings for PC1 relative to the much larger
318 range of loading values for PC2 (Figure 8). This confirms that PC1 represented the common
319 behavior among all stream sites and that deviations in timing of water temperature increases and
320 decreases were dictated by PC2. As a result, when annual temperature signals were reconstructed
321 for two sites from the PCs based on the mean loading value for PC1 and +/- 0.16 for PC2 to
322 represent strong negative and positive loadings, the expected timing shift was apparent (Figure 9).
323 Notably, the site with the -0.16 PC2 loading had a later, sharper rise in water temperature that
324 peaked in late summer approximately one month after the site with the positive loading. The

325 correspondence of PC2 with stream discharge in Figure 7a suggests the timing shift could be related
326 to runoff patterns. And indeed, the annual unit-area runoff for the basins associated with the 226
327 sites was a strong predictor of the PC2 loadings in a linear regression ($r^2 = 0.51$; Figure 10). Site
328 elevation provides some indication of rainfall-snowfall fraction that may help explain timing shifts
329 but this covariate added little predictive capacity beyond annual runoff when examined across all
330 sites ($r^2 = 0.54$). However, when sites with basin sizes less than 50 km² were examined (because
331 site elevation relates more strongly to mean basin elevation in smaller basins), elevation accounted
332 for a large increase in the explainable variance of PC2 loadings beyond that attributable to annual
333 runoff ($r^2 = 0.69$). Although orographic enhancement of precipitation is evident in the study area,
334 there is enough difference in circulation patterns across the north-south extent of the area that
335 elevation and annual runoff were only weakly correlated in the small basins ($r = -0.2$), so the
336 elevation effect was largely independent of annual precipitation. As a result, both factors appeared
337 to contribute to the PC2 loadings such that either wetter or colder locations had more negative
338 loadings and later rises in water temperatures.

Formatted: Font: Italic

Formatted: Superscript

Formatted: Font: Italic

Formatted: Superscript

Formatted: Superscript

Formatted: Font: Italic

Formatted: Superscript

Formatted: Font: Italic

Deleted: ¶

Deleted:Page Break.....

340 5 Discussion

341 5.1 Thermal regimes in mountain settings

342 Thermal regimes in the mountain river networks we studied were simple and responded relatively
343 coherently to climatic variability across a geomorphically consistent area with few reservoirs,
344 Strong seasonal patterns in water temperatures characteristic of temperate latitudes were apparent in
345 response to the primary signal set by the annual air temperature cycle and accompanying changes in
346 solar radiation. Not surprisingly given the pronounced elevational gradients in the study landscape,
347 the dominant regime aspect represented by PC1 in the metric-based PCA was associated with
348 magnitude. Less expected was that many of the variability metrics also loaded heavily on the first
349 PC because variation has been treated as a distinct element of thermal regimes (e.g., Steel et al.,
350 2012; Kovach et al., 2018). The concurrence of magnitude and variability metrics probably also
351 relates to elevation and changes in the importance of groundwater buffering, which both cools
352 streams and dampens diurnal and seasonal variations (Cassie and Luce, 2017). For example, the
353 coldest streams at the highest elevations are usually strongly buffered by groundwater inputs
354 derived from large annual snowpacks in mountain environments and often show limited thermal
355 variability (Luce et al., 2014a; Isaak et al., 2016). Downstream from the headwaters, the
356 proportional inputs of groundwater decrease and streams are more coupled to climatic variability

Deleted: .

Deleted: treams

Deleted: mountain

Deleted: were relatively

Deleted: conditions across a geomorphically consistent mountainous area with few reservoirs

Deleted: cycles

Deleted: , which the S-mode analysis indicated were correlated with variation in air temperatures and associated physical processes.

368 even as their average temperatures increase due to solar gains over longer flow distances (Caissie
369 2006). In contrast to the metrics associated with PC1, metrics that described the winter period and
370 the extent of the growing season largely defined PC2. This “winter” PC is probably common to
371 stream thermal regimes in mountain landscapes where subzero air temperatures are frequent and
372 result in prolonged periods with water temperatures near 0 °C. The orthogonal nature of PC1 and
373 PC2 suggests that streams with otherwise similar magnitude and variance structures will sometimes
374 differ substantially with regards to their winter and growing seasons—a distinction that could have
375 important implications for biological communities or stream physicochemical processes.

376
377 Our results also suggest that important local nuances in water temperature dynamics like the
378 differences in timing of spring warming and peak temperatures may emerge from the interactions
379 among annual climate cycles, basin geomorphology, and hydrology. Because precipitation, air
380 temperatures, snowpack, runoff volume, and runoff timing are all evolving in response to climate
381 change in mountain environments across the study region (Mote et al., 2005; Luce et al., 2013) and
382 globally (Stewart, 2009), better understanding of these connections is needed. In particular, more
383 insight to the relationship of water temperatures with annual unit-area runoff and whether the
384 underlying mechanisms relate to changes in snowpack accumulation (Luce et al., 2014b; Luce and
385 Luce, 2017), snowmelt timing and rate (Musselman et al., 2017), the volume of water stored in
386 groundwater (e.g. Tague et al., 2007), or the outcomes of extreme low flows (e.g. Kormos et al.,
387 2016; Luce and Holden, 2009) could lead to better predictions about water temperatures and the
388 evolution of thermal regimes in response to expected changes in air temperatures and precipitation.

390 **5.2 Implications for modeling and monitoring**

391 Water temperature models are often developed for use in ecological assessments and to understand
392 how habitat degradation or restoration efforts may affect thermal regimes (Benyahya et al., 2007;
393 Gallice et al., 2015; Dugdale et al., 2017). Our results, like several previous studies that have
394 compared multiple temperature metrics (Isaak and Hubert, 2001; Rivers-Moore et al., 2013; Steele
395 et al., 2016), confirm that numerous metrics are strongly correlated and provide redundant
396 information. The specific choice of a metric, therefore, may not be critical as long as it represents an
397 important aspect of a thermal regime and is suited to the goals of a study. Metrics associated with
398 temperature magnitude and variability, which have been the focus of most modeling efforts, are
399 good choices because they represent significant portions of the information about thermal regimes

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Moved down [5]: (Isaak and Hubert, 2001; Rivers-Moore et al., 2013; Steele et al., 2016)

Moved (insertion) [5]

Deleted: Discharge played a secondary role and mediated water temperature sensitivity to air temperature to greater or lesser degrees based on proximity to each year’s pulse of cold snowmelt runoff. Similar to previous studies that used multiple metrics to describe thermal regimes (Isaak and Hubert, 2001; Rivers-Moore et al., 2013; Steele et al., 2016), many metrics were redundant (Isaak and Hubert, 2001; Rivers-Moore et al., 2013; Steele et al., 2016), so a small number of PCs was adequate for summarizing most of the regime information. Not surprisingly given the large elevational gradients in the study landscape, the most important PC was associated with metric representations of magnitude. Less expected was that most of the variability metrics also loaded heavily on the first PC because variation is often considered a distinct element of thermal regimes (e.g., Steel et al., 2012; Kovach et al. 2018). That distinction appears instead to be associated with metrics describing the winter period and the extent of the growing season that define the second PC. This “winter” PC is probably common to thermal regimes in mountain landscapes where frequent subzero air temperatures occur and may result in prolonged periods of temperatures near 0 °C. The orthogonality of PC1 and PC2 is interesting in that it suggests streams with otherwise similar magnitude and variance structures will sometimes differ substantially with regards to their winter and growing seasons.

Moved (insertion) [3]

Moved (insertion) [1]

Moved up [1]: Similar to previous studies that have assessed metrics (Isaak and Hubert, 2001; Dunham et al., 2005; Steele et al., 2016), we observed strong correlations and redundancies among metrics.

Moved up [3]: Two distinct T-mode spatial phases as during 2-6 months of year, homothermous and no response but outside of this time, S-mode revealed strong response to air temperatures. Discharge effect was much smaller but showed noteworthy mediation and hysteresis wherein sensitivity to air temperature variation was muted on the ascending limb as others have noted (Mohseni 98) and enhance on descending limb

Moved up [2]: Would fit within Maheu et al. (2015) scheme as cold stable/cold unstable that are characterized by low magnitude and winter homothermous period.

Moved (insertion) [2]

Deleted: Would fit within Maheu et al. (2015) scheme as cold stable/cold unstable that are characterized by low magnitude and winter homothermous period. Two distinct T-mode spatial phases as during 2-6 months of year, homothermous and no response but outside of this time, S-mode revealed strong response to air temperatures. Discharge effect was much smaller but showed noteworthy mediation and hysteresis wherein sensitivity to air temperature variation was muted on the ascending limb as others have noted (Mohseni 98) and enhance on descending limb. Little evidence existed for subdomains or regionalization within the study area, probably because streams drained similar geologies and sample sites spanned a limited spatial extent relative to the scales at which heterogeneity in weather patterns occurred. Moreover, strong

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

536 and have been shown on many occasions to be important determinants of ecological attributes such
537 as species distributions and abundance or physical processes in streams and rivers (Isaak et al.,
538 2017a; Webb et al., 2008). Our preferred metrics in previous research have been mean August or
539 mean summer temperatures because the data records for their calculation are typically available at
540 the largest number of sites in mountain environments, which maximizes sample sizes and
541 minimizes the distances over which interpolations are made when developing and applying
542 network-scale temperature models (e.g., Detenbeck et al., 2016; Isaak et al. 2017b). Metrics based
543 on longer-term means rather than short-term daily or weekly maxima are also more stable and easier
544 to predict (Isaak et al. 2010; Turschwell et al., 2016), although a focus on the latter metrics is often
545 mandated within regulatory environments and may negate these considerations (Todd et al., 2008;
546 McCullough 2010). Comparatively little effort has gone towards modeling temperature metrics
547 associated with growing season length or the dates of spring and winter season onset, despite the
548 significant information these metrics provide about thermal regimes and their relevance to the
549 phenology and life histories of organisms that constitute aquatic communities (Huryn and Wallace,
550 2000; Neuheimer and Taggart, 2007). These aspects of thermal regimes, as well as magnitude and
551 variability characteristics, are also likely to be evolving in response to climate change, so new
552 models are needed to provide forecasting abilities about changes later this century. Rather than
553 focusing on individual metrics, researchers could also instead use PCA to efficiently summarize
554 multiple temperature metrics and then model the eigenvector loadings that define one or more of the
555 principal components. This approach would maximize the amount of thermal information
556 represented by a response metric but would yield results that were more ambiguous to interpret.

557
558 The growth of new stream and river temperature monitoring and data collection activities has been
559 remarkable in recent years. Although optimization of those efforts ultimately depends on local
560 considerations, some general guidelines emerge from this work that may be applicable to other
561 areas. For example, the coherent behavior we observed among temperatures at many sites suggests
562 that a limited number of monitoring stations will often be sufficient to represent the temporal
563 dynamics of thermal regimes. Those stations would need to be spread geographically and along
564 major environmental gradients and replicated to mitigate against sensor losses, but 20–30 stations
565 might prove sufficient at scales comparable to our study area. Given low sensor costs and the
566 availability of standardized data collection protocols (Isaak et al., 2013; Stamp et al., 2014),
567 monitoring arrays could also be crowd-sourced effectively if site locations were coordinated and

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Moved (insertion) [6]

Deleted: (Daigle et al., 2016; Isaak et al., 2017b)

Deleted: Within mountain landscapes characterized by uniform geohydroclimates, the

Deleted: can

Deleted: in

Deleted: ites

Deleted: 1

Deleted: ,

Moved up [6]: (Daigle et al., 2016; Isaak et al., 2017b)

Deleted:

Deleted:), and growing interest in temperature monitoring

579 chosen strategically using geospatial analyses to describe and stratify networks for sample
580 allocation (Jackson et al., 2016). Monitoring networks might also be supplemented by incorporating
581 data from sites established for other purposes, such as documenting thermal responses to habitat
582 restoration efforts (Nichols and Ketcheson, 2013) or disturbances associated with land management,
583 wildfires, or livestock grazing (Mahlum et al., 2011; Nusslé et al., 2015). In fact, those factors
584 motivated collection of many of the datasets compiled for this analysis, although supplementation
585 with additional sites was needed to ensure adequate coverage within the study area.

Deleted: factors

Deleted: representative

587 If one of the goals of temperature data collection efforts is to develop accurate prediction maps that
588 show spatial variation in one or more thermal metrics (e.g., Isaak et al., 2017b; Steel et al., 2016),
589 monitoring sites may need to be established more densely than the temporal considerations
590 discussed above otherwise suggest. Spatial autocorrelation in temperature metric values is minimal
591 in mountain river networks beyond distances of 10–100 km (Isaak et al., 2010; Zimmerman and Ver
592 Hoef 2017), so this level of sensor spacing would be required to generate the most accurate maps.
593 Given the extent of many river networks, that could translate into a large number of sites but most
594 of these could be monitored for short periods while temporal dynamics were represented by a subset
595 of long-term sites because temporal covariance among sites would be strong. Costs associated with
596 numerous sensor deployments could be prohibitive, so aggregation of existing data sets from
597 multiple natural resource agencies into a centralized database often becomes an attractive option.
598 Moreover, if those central databases are made publically accessible, professionals from the
599 contributing agencies may begin to coordinate data collection activities more consistently and
600 effectively across larger areas (e.g., Isaak et al. 2018b).

Deleted: one monitoring

Deleted: goal is

Deleted: ing

Deleted: also

Deleted: sampled

Deleted: above

Deleted: network

Formatted: Font: Not Bold

Deleted: similar

Deleted: is

Deleted: ost

Deleted: would

Deleted: often

Deleted: since

Formatted: Font: Not Bold

Formatted: Indent: First line: 0", Line spacing: 1.5 lines

602 As new data collection and database development efforts proceed, it is commonly the case that
603 temperature records have inconsistent period lengths or missing values. Usually it is desirable to
604 have complete records for analysis, so missing values are sometime imputed based on the
605 correlations between two monitoring site records that strongly covary (e.g., Rivers-Moore et al.,
606 2013). However, the process can be tedious if required at more than a few sites, so an efficient
607 improvement is offered by the imputation technique described by Josse and Husson (2012) that is
608 easily used in the MissMDA software package (Josse and Husson, 2016) for the R statistical
609 program (R Core Team, 2014). This technique examines and uses correlations among multiple site
610 records simultaneously to estimate missing values by first applying standard PCA to the incomplete

626 data set where missing values are replaced with the respective record mean. Data are then
627 reconstructed from the PCs, and the initial analysis step is repeated but with missing values replaced
628 using estimates from the reconstructed data. The process is repeated until convergence, and the
629 missing values in the original data records are ultimately replaced with estimates from the last PCA
630 reconstruction (Josse and Husson, 2012). Care should be taken against overreliance on the
631 technique to impute particularly sparse records but the MissMDA package provides a useful tool for
632 addressing gaps when working with large temperature datasets or time-series of other measurements
633 common to hydrology such as gage discharge records (e.g., Isaak et al., 2018a).

634

635 **5.3 Conclusions**

636 Our analysis of thermal regimes follows previous work that has proven fundamental to advancing
637 the understanding of hydrologic regimes (Poff et al., 1997; Olden and Poff, 2003) but also adds
638 novel applications of PCA variants from the field of climatology that hold utility for stream
639 temperature research and monitoring design. Insights from those analyses indicate that thermal
640 conditions in the mountain river networks studied here were strongly coherent through time,
641 exhibited two distinct spatial phases, could be adequately described by a few principal components
642 or allied metrics, and reflected landscape geomorphology and hydroclimatic conditions. A logical
643 next step involves application of these PCA techniques to larger stream and river temperature
644 datasets at regional, continental, or intercontinental scales to encompass greater heterogeneity and
645 discern the geographic domains over which distinct thermal regimes are operable. Across
646 sufficiently diverse landscapes, we might expect to find classes of thermal regimes that, at a
647 minimum, mimicked previously described classes of hydrologic regimes (e.g., rainfall, snowmelt,
648 spring-groundwater, and regulated) but possible divergences from, or additions to, those categories
649 would be useful to ascertain. In a national-scale assessment for the United States, Maheu et al.
650 (2015) classified stream thermal regimes into six categories but the 135 temperature stations that
651 supported the analysis were limited in comparison to a drainage network comprised of millions of
652 kilometers. Subsequent iterations on that effort could document additional, undescribed thermal
653 classes and might also prove beneficial by developing detailed maps of classification schemes to aid
654 in assessments of ecological conditions or anthropogenic effects on stream thermal regimes. As
655 research on the topic of thermal regimes matures, syntheses with flow regime concepts and
656 databases could also be sought to more fully describe the hydroclimatic conditions of flowing
657 waters.

Deleted: ¶
Formatted: No Spacing, Line spacing: 1.5 lines

659
660 *Data availability.* All water temperature data used in this study are available at the NorWeST website
661 (<https://www.fs.fed.us/rm/boise/AWAE/projects/NorWeST.html>) whereas the full data set that includes air temperature
662 and discharge data are available at the lead author's ResearchGate profile entry for this study
663 (https://www.researchgate.net/profile/Daniel_Isaak).
664
665 *Competing interests.* The authors declare that they have no conflict of interest.
666
667 *Acknowledgements.* We thank Dave Schoen, Bart Gamett, Dan Garcia, Scott Vuono, Caleb Zurstadt, and Clayton
668 Nalder with the U.S. Forest Service, Tim Copeland, Eric Stark, and Ron Roberts with the Idaho Department of Fish and
669 Game, Eric Archer and Jeff Ojala with the Pacfish-Infish Biological Opinion monitoring program, and Boyd Bowes and
670 Chris Jordan with the CHaMP monitoring program that contributed water temperature data to enable this research.
671 Comments from Nicholas Rivers-Moore and one anonymous reviewer improved the quality of the final manuscript. The
672 authors of this work were supported by the U.S. Forest Service Rocky Mountain Research Station.

Deleted:Page Break.....

674 **References**

675 Arismendi, I., Johnson, S.L., Dunham, J.B., and Haggerty, R.: Descriptors of natural thermal regimes in streams and their
676 responsiveness to change in the Pacific Northwest of North America, *Freshwater Biology*, 58, 880-894, 2013.
677 [Benyahya, L., Caissie, D., St-Hilaire, A., Ouarde, T., and Bobee, B.: A review of statistical water temperature models,
678 *Canadian Water Resources Journal*, 32, 179-192, 2007.](#)
679 Bond, J.G. and Wood, C.H.: Geologic map of Idaho, 1:500 000 scale. Idaho Department of Lands, Bureau of Mines and
680 Geology, Moscow, Idaho, 1978.
681 Brown, L.E., Cereghino, R., and Compin, A.: Endemic freshwater invertebrates from southern France: diversity,
682 distribution and conservation implications, *Biological Conservation*, 142, 2613-2619, 2009.
683 Brown, L.E. and Hannah, D.M.: Spatial heterogeneity of water temperature across an alpine river basin, *Hydrological
684 Processes*, 22, 954-967, 2008.
685 Caissie, D.: The thermal regime of rivers: a review, *Freshwater Biology*, 51, 1389-1406, 2006.
686 [Caissie, D., and Luce, C. H.: Quantifying streambed advection and conduction heat fluxes, *Water Resour. Res.*, 53,
687 1595-1624, 10.1002/2016WR019813, 2017.](#)
688 [Chandler, G.L., Wollrab, S., Horan, D., Nagel, D., Parkes, S., Isaak, D.J., Wenger, S.J., Peterson, E.E., Ver Hoef, J. M.,
689 Hostetler, S., Luce, C.H., Dunham, J.B., Kershner, J., and Roper, B.B.: NorWeST stream temperature data
690 summaries for the western United States. U.S. Forest Service, Rocky Mountain Research Station Research Data
691 Archive, Fort Collins, CO, <https://doi.org/10.2737/RDS-2016-0032>, 2016.](#)
692 Chu, C., Jones, N.E., and Allin, L.: Linking the thermal regimes of streams in the Great Lakes Basin, Ontario, to
693 landscape and climate variables, *River Research and Applications*, 26, 221-241, 2010.
694 Daigle, A., Caudron, A., Vigier, L., and Pella, H.: Optimization methodology for a river temperature monitoring
695 network for the characterization of fish thermal habitat, *Hydrological Sciences Journal*, 62, 483-497, 2016.

Deleted: ¶

698 Demsar, U., Harris, P., Brunson, C., Fotheringham, A.S., and McLoone, S.: Principal components analysis on spatial
699 data: An overview, *Annals of the Association of American Geographers*, 103, 106-128, 2013.

700 [Detenbeck, N.E., Morrison, A., Abele, R.W., and Kopp, D.: Spatial statistical network models for stream and river
701 temperature in New England, USA, *Water Resources Research*, 52, 6018-6040, 2016.](#)

702 [Ducharme, A.: Importance of stream temperature to climate change impact on water quality, *Hydrology and Earth
703 System Sciences Discussions*, 4, 2425-2460, 2007.](#)

704 [Dugdale, S.J., Hannah, D.M. and Malcolm, I.A.: River temperature modelling: A review of process-based approaches
705 and future directions, *Earth-Science Reviews*, 175, 97-113, 2017.](#)

706 [Dunham, J., Rieman, B., and Chandler, G.: Influences of temperature and environmental variables on the distribution of
707 bull trout within streams at the southern margin of its range, *North American Journal of Fisheries Management*, 23,
708 894-904, 2003.](#)

709 Gallacher, K., Miller, C., Scott, E.M., Willows, R., Pope, L., and Douglass, J.: Flow-directed PCA for monitoring
710 networks, *Environmetrics*, 28, e2434, 2017.

711 Gallice, A., Schaeffli, B., Lehning, M., Parlange, M.B. and Huwald, H.: Stream temperature prediction in ungauged
712 basins: review of recent approaches and description of a new physics-derived statistical model, *Hydrology and
713 Earth System Sciences*, 19, 3727-3753, 2015.

714 [Gordon, N.D., McMahon, T.A., and Finlayson, B.L.: Stream hydrology: an introduction for ecologists, *Stream
715 hydrology: an introduction for ecologists*, John Wiley and Sons, Chichester, UK, 1991.](#)

716 Hannah, D. and Garner, G.: River water temperature in the United Kingdom: Changes over the 20th century and
717 possible changes over the 21st century, *Progress in Physical Geography*, 39, 68–92, 2015.

718 Hilderbrand, R.H., Kashiwagi, M.T., and Prochaska, A.P.: Regional and local scale modeling of stream temperatures
719 and spatio-temporal variation in thermal sensitivities, *Environmental Management*, 54, 14-22, 2014.

720 Huff, D.D., Hubler, S.L., and Borisenko, A.N.: Using field data to estimate the realized thermal niche of aquatic
721 vertebrates, *North American Journal of Fisheries Management*, 25, 346–360, 2005.

722 [Huryn, A.D., and Wallace, J.B.: Life history and production of stream insects, *Annual Review of Entomology*, 45, 83–
723 110, 2000.](#)

724 Ice, G.G., Light, J., and Reiter, M.: Use of natural temperature patterns to identify achievable stream temperature
725 criteria for forest streams, *West. J. Appl. For.*, 19, 252–259, 2004.

726 Isaak, D.J. and Hubert, W.A.: A hypothesis about factors that affect maximum summer stream temperatures across
727 montane landscapes, *Journal of the American Water Resources Association*, 37, 351-366, 2001.

728 Isaak, D.J., Horan, D.L., and Wollrab, S.P.: A simple protocol using underwater epoxy to install annual temperature
729 monitoring sites in rivers and streams, U.S. Forest Service General Technical Report, Rocky Mountain Research
730 Station, 314, Fort Collins, Colorado, USA, 2013.

731 Isaak, D.J., Luce, C.H., Horan, D.L., Chandler, G.L., Wollrab, S.P., Nagel, D.: Global warming of salmon and trout
732 rivers in the northwestern United States: Road to ruin or path through purgatory?, *Transactions of the American
733 Fisheries Society*, 147:566-587, 2018a.

Deleted: DeWeber, J. and Wagner, T.: Probabilistic measures of climate change vulnerability, adaptation action benefits, and related uncertainty from maximum temperature metric selection, *Global Change Biology*, 2018. ¶
Dillon, M.E., Woods, H.A., Wang, G., Fey, S.B., Vasseur, D.A., Telemeco, R.S., Marshall, K., and Pincebourde, S.: Life in the frequency domain: the biological impacts of changes in climate variability at multiple time scales, *Integrative and Comparative Biology*, 56, 14-30, 2016. ¶

Deleted: Dunham, J.B., Chandler, G., Rieman, B.E., and Martin D.: Measuring stream temperature with digital dataloggers: A user's guide, General Technical Report RMRS-GTR-150WWW. U.S. Forest Service, Rocky Mountain Research Station, Fort Collins, Colorado, USA, 2005. ¶

Deleted: Garcia, R.A., Cabeza, M., Rahbek, C., and Araújo, M.B.: Multiple dimensions of climate change and their implications for biodiversity, *Science*, 344, 1247579, 2014. ¶

751 Isaak, D.J., Luce, C.H., Rieman, B.E., Nagel, D.E., Peterson, E.E., Horan, D.L., Parkes, S., and Chandler, G.L.: Effects
752 of climate change and wildfire on stream temperatures and salmonid thermal habitat in a mountain river network,
753 Ecol. Appl., 20, 1350–1371, 2010.

754 Isaak, D.J., Wenger, S.J., and Young, M.K.: Big biology meets microclimatology: Defining thermal niches of aquatic
755 ectotherms at landscape scales for conservation planning, Ecological Applications, 27, 977-990, 2017a.

756 Isaak, D.J., Wenger, S.J., Peterson, E.E., Ver Hoef, J.M., Nagel, D.E., Luce, C.H., Hostetler, S.W., Dunham, J.B.,
757 Roper, B.B., Wollrab, S., Chandler, G., Parkes, S., and Horan, D.: The NorWeST summer stream temperature
758 model and scenarios: A crowd-sourced database and new geospatial tools foster a user community and predict
759 broad climate warming of rivers and streams in the western United State, Water Resources Research, 53, 9181-
760 9205, 2017b.

761 Isaak, D.J., Wenger, S.J., Peterson, E.E., Ver Hoef, J.M., Nagel, D.E., Luce, C.H., Hostetler, S.W., Dunham, J.B.,
762 Roper, B.B., Wollrab, S., Chandler, G., Parkes, S., and Horan, D.: NorWeST modeled summer stream temperature
763 scenarios for the western United States. U.S. Forest Service, Rocky Mountain Research Station Research Data
764 Archive, Fort Collins, CO, <https://doi.org/10.2737/RDS-2016-0033>, 2016b.

765 ~~Isaak, D.J., Young, M.K. Luce, C.H. Hostetler, S. Wenger, S. Peterson, E.E. Ver Hoef, J.M. Groce, M. Horan, D.L. and~~
766 ~~Nagel D.: Slow climate velocities of mountain streams portend their role as refugia for cold-water biodiversity,~~
767 ~~Proceedings of the National Academy of Sciences, 113, 4374-4379, 2016a.~~

768 ~~[Isaak, D.J., Young, M.K., McConnell, C., Roper, B.B., Archer, E.K., Staab, B., Hirsch, C., Nagel, D.E., Schwartz,](#)~~
769 ~~[M.K., Chandler, G.L.: Crowd-sourced databases as essential elements for Forest Service partnerships and aquatic](#)~~
770 ~~[resource conservation, Fisheries, 43, 423-430, 2018b.](#)~~

771 Jackson, F.L., Malcolm, I.A., and Hannah, D.M.: A novel approach for designing large-scale river temperature
772 monitoring networks, Hydrology Research, 47, 569-590, 2016.

773 ~~Jiménez, P.A., García-Bustamante, E., González-Rouco, J.F., Valero, F., Montávez, J.P., and Navarro, J.: Surface wind~~
774 ~~regionalization in complex terrain, Journal of Applied Meteorology and Climatology, 47, 308–325, 2008.~~

775 Josse, J., and ~~Husson F.:~~ ~~Handling missing values in exploratory multivariate data analysis methods. Journal of the~~
776 ~~Société Française de Statistique, 153, 79–99, 2012.~~

777 Josse, J. and Husson, F.: ~~MjssMDA: a package for handling missing values in multivariate data analysis, Journal of~~
778 ~~Statistical Software, 70, 1-31, 2016.~~

779 ~~[Kormos, P., Luce, C., Wenger, S. J., and Berghuijs, W. R.: Trends and Sensitivities of Low Streamflow Extremes to](#)~~
780 ~~[discharge Timing and Magnitude in Pacific Northwest Mountain Streams, Water Resour. Res, 52, 4990-5007,](#)~~
781 ~~[10.1002/2015WR018125, 2016.](#)~~

782 ~~[Kovach, R.P., Muhlfeld, C.C., Al-Chokhachy, R., Ojala, J.V., and Archer, E.K.: Effects of land use on summer thermal](#)~~
783 ~~[regimes in critical salmonid habitats of the Pacific Northwest, Canadian Journal of Fisheries and Aquatic Sciences,](#)~~
784 ~~<https://doi.org/10.1139/cjfas-2018-0165>, 2018.~~

785 ~~[Likens, G.E. and Likens, G.E.: Biogeochemistry of a forested ecosystem, Springer-Verlag, 1977.](#)~~

786 ~~[Luce, C. H., Abatzoglou, J. T., and Holden, Z. A.: The Missing Mountain Water: Slower Westerlies Decrease](#)~~
787 ~~[Orographic Enhancement in the Pacific Northwest USA, Science, 342, 1360-1364, 2013.](#)~~

Deleted: ¶

Deleted: Jackson, F.L., Fryer, R.J., Hannah, D.M., Millar, C.P., and Malcolm, I.A.: A spatio-temporal statistical model of maximum daily river temperatures to inform the management of Scotland's Atlantic salmon rivers under climate change, Science of the Total Environment, 612, 1543-1558, 2018, ¶

Deleted: Husson.

Deleted: 2012.

Deleted: ¶

Deleted: ¶

Deleted: :

Deleted: .

Deleted: m

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

801 [Luce, C. H., and Holden, Z. A.: Declining annual streamflow distributions in the Pacific Northwest United States, 1948-](#)
802 [2006, *Geophysical Research Letters*, 36, L16401, 2009.](#)

803 [Luce, C. H., Lopez-Burgos, V., and Holden, Z.: Sensitivity of snowpack storage to precipitation and temperature using](#)
804 [spatial and temporal analog models, *Water Resources Research*, 50, 9447-9462, 2014b.](#)

805 [Luce, C.H., Staab, B., Kramer, M., Wenger, S., Isaak, D., and McConnell, C.: Sensitivity of summer stream](#)
806 [temperatures to climate variability in the Pacific Northwest, *Water Resoures Research*, 50, 3428-3443, 2014a.](#)

807 [Lute, A. C., and Luce, C. H.: Are model transferability and complexity antithetical? Insights from validation of a](#)
808 [variable-complexity snow model in space and time, *Water Resoures Research*, 53, 8825–8850, 2017.](#)

809 [Maheu, A., Poff, N.L., and St-Hilaire, A.: A classification of stream water temperature regimes in the conterminous](#)
810 [USA, *River Research and Applications*, 32, 896-906, 2016.](#)

811 [Mahlum, S.K., Eby, L.A., Young, M.K., Clancy, C.G., and Jakober, M.: Effects of wildfire on stream temperatures in](#)
812 [the Bitterroot River Basin, Montana, *International Journal of Wildland Fire*, 20, 240-247, 2011.](#)

813 [Martins, D.S., Razei, T., Paulo, A.A., and Pereira, L.S.: Spatial and temporal variability of precipitation and drought in](#)
814 [Portugal, *Nat. Hazards Earth Syst. Sci.*, 12, 1493–1501, 2012.](#)

815 [Mauger, S., Shaftel, R., Leppi, J.C., and Rinella, D.J.: Summer temperature regimes in southcentral Alaska streams:](#)
816 [watershed drivers of variation and potential implications for Pacific salmon, *Canadian Journal of Fisheries and*](#)
817 [Aquatic Sciences](#), 74, 702-715, 2016.

818 [McCullough, D.A.: Are coldwater fish populations of the United States actually being protected by temperature](#)
819 [standards?, *Freshwater Reviews*, 3, 147-199, 2010.](#)

820 [McKay, L., Bondelid, T., Dewald, T., Johnston, J., Moore, R., and Reah, A.: NHDPlus Version 2: User Guide.](#)
821 ftp://ftp.horizon-systems.com/NHDPlus/NHDPlusV21/Documentation/NHDPlusV2_User_Guide.pdf, 2012.

822 [Meyer, G.A., Pierce, J.L., Wood, S.H., and Jull, A.J.T.: Fire, storms, and erosional events in the Idaho batholith,](#)
823 [Hydrol. Process.](#), 15, 3025–3038, 2001.

824 [Molinero, J., Larrañaga, A., Pérez, J., Martínez, A., and Pozo, J.: Stream temperature in the Basque Mountains during](#)
825 [winter: thermal regimes and sensitivity to air warming, *Climatic Change*, 134, 593-604, 2016.](#)

826 [Morgan, P., Heyerdahl, E., Miller, C., Wilson, A., and Gibson, C.: Northern Rockies pyrogeography: an example of fire](#)
827 [atlas utility, *Fire Ecology*, 10, 14–30, 2014.](#)

828 [Mote, P.W., Hamlet, A. F., Clark, M. P., and Lettenmaier, D. P.: Declining mountain snowpack in western North](#)
829 [America, *Bulletin of the American Meteorological Society*, 86, 39–49, 2005.](#)

830 [Musselman, K. N., Clark, M. P., Liu, C., Ikeda, K., and Rasmussen, R.: Slower snowmelt in a warmer world, *Nature*](#)
831 [Climate Change](#), 7, 214-219, 2017.

832 [Neuheimer, A.B. and Taggart, C.T.: The growing degree-day and fish size-at-age: the overlooked metric, *Canadian*](#)
833 [Journal of Fisheries and Aquatic Sciences](#), 64, 375-385, 2007.

834 [Nichols, R.A., and Ketcheson, G.L.: A two-decade watershed approach to stream restoration log jam design and stream](#)
835 [recovery monitoring: Finney Creek, Washington, *Journal of the American Water Resources Association*, 49, 1367-](#)
836 [1384, 2013.](#)

837 [Nusslé, S., Matthews, K.R., and Carlson, S.M.: Mediating water temperature increases due to livestock and global](#)
838 [change in high elevation meadow streams of the Golden Trout Wilderness, *PLoS ONE*, 10, e0142426, 2015.](#)

Deleted: ¶

Likens, G.E. and Likens, G.E.: Biogeochemistry of a forested ecosystem, Springer-Verlag, 1977.¶
Luce, C.H., Staab, B., Kramer, M., Wenger, S., Isaak, D., and McConnell, C.: Sensitivity of summer stream temperatures to climate variability in the Pacific Northwest, *Water Resour Res.* 50, 3428-3443, 2014.¶

Deleted: ¶

847 Olden, J.D. and Poff, N.L.: Redundancy and the choice of hydrologic indices for characterizing streamflow regimes,
848 River Research and Applications, 19, 101-121, 2003.

849 Olden, J.D. and Naiman, R.J.: Incorporating thermal regimes into environmental flows assessments: modifying dam
850 operations to restore freshwater ecosystem integrity, Freshwater Biology, 55, 86-107, 2010.

851 Orr, H.G., Johnson, M.F., Wilby, R.L., Hatton-Ellis, T., and Broadmeadow, S.: What else do managers need to know
852 about warming rivers? A United Kingdom perspective, Wiley Interdisciplinary Reviews—Water, 2, 55-64, 2015.

853 Pearson, K.: On Lines and Planes of Closest Fit to Systems of Points in Space, Philosophical Magazine, 6, 559–572,
854 1901.

855 Piechota, T.C., Dracup, J.A., and Fovell, R.G.: Western US streamflow and atmospheric circulation patterns during El
856 Nino-Southern Oscillation, Journal of Hydrology, 201, 249-271, 1997.

857 Poff, N.L. and Ward, J.V.: Implications of streamflow variability and predictability for lotic community structure: a
858 regional analysis of streamflow patterns, Canadian Journal of Fisheries and Aquatic Sciences, 46, 1805–1818,
859 1989.

860 Poff, N.L., Allan, J.D., Bain, M.B., Karr, J.R., Prestegard, K.L., Richter, B.D., Sparks, R.E., and Stromberg, J.C.: The
861 natural flow regime, BioScience, 47, 769-784, 1997.

862 Poole, G.C., Dunham, J.B., Keenan, D.M., Sauter, S.T., McCullough, D.A., Mebane, C., Lockwood, J.C., Essig, D.A.,
863 Hicks, M.P., Sturdevant, D.J., and Materna, E.J.: The case for regime-based water quality standards, BioScience,
864 54, 155-161, 2004.

865 Quaglietta, L., Paupério, J., Martins, F.M.S., Alves, P.C., and Beja, P.: Recent range contractions in the globally
866 threatened Pyrenean desman highlight the importance of stream headwater refugia, Animal Conservation,
867 <https://doi.org/10.1111/acv.12422>.

868 R Development Core Team, R: A language and environment for statistical computing, R Foundation for Statistical
869 Computing, Vienna, Austria, 2014.

870 Richman, M.B.: Rotation of principal components, International Journal of Climatology, 6, 293-335, 1986.

871 Rieman, B.E., Smith, C.L., Naiman, R.J., Ruggeronee, G.T., Wood, C.C., Huntly, N., Merrill, E.N., Alldredge, J.R.,
872 Bisson, P.A., Congleton, J., Fausch, K.D., Levings, C., Percy, W., Scarnecchia, D., and Smouse, P.: A
873 comprehensive approach for habitat restoration in the Columbia Basin, Fisheries, 40, 124-135, 2015.

874 Rivers-Moore, N.A., Dallas, H.F., and Morris, C.: Towards setting environmental water temperature guidelines: A
875 South African example, Journal of Environmental Management, 128, 380-392, 2013.

876 Roni, P., Beechie, T.J., Bilby, R.E., Leonetti, F.E., Pollock, M.M., and Pess, G.R.: A review of stream restoration
877 techniques and a hierarchical strategy for prioritizing restoration in Pacific Northwest watersheds, North American
878 Journal of Fisheries Management, 22, 1-20, 2002.

879 SAS Institute Inc. SAS/STAT 14.1 User's Guide Cary, NC, USA, 2015.

880 Sergeant, C.J., Starkey, E.N., Bartz, K.K., Wilson, M.H., and Mueter, F.J.: A practitioner's guide for exploring water
881 quality patterns using principal components analysis and Procrustes, Environ Monit Assess, 188, 249, 2016.

882 Stamp, J., Hamilton, A., Craddock, M., Parker, L., Roy, A., Isaak, D., Holden, Z., Passmore, M., and Bierwagen, B.:
883 Best practices for continuous monitoring of temperature and flow in wadeable streams, Global Change Research
884 Program, National Center for Environmental Assessment, EPA/600/R-13/170F, 2014.

Deleted: Roberts, J.J., Fausch, K.D., Peterson, D.P., and Hooten, M.B.: Fragmentation and thermal risks from climate change interact to affect persistence of native trout in the Colorado River basin, Global Change Biology, 19, 1383-1398, 2013.¶

889 Steel, E.A., Tillotson, A., Larsen, D.A., Fullerton, A.H., Denton, K.P., and Beckman, B.R.: Beyond the mean: the role
890 of variability in predicting ecological effects of stream temperature on salmon, *Ecosphere*, 3, 1-11, 2012.

891 Steel, E.A., Sowder, C., and Peterson, E.E.: Spatial and temporal variation of water temperature regimes on the
892 Snoqualmie River network, *Journal of the American Water Resources Association*, 52, 769-787, 2016.

893 [Stewart, I.T.: Changes in snowpack and snowmelt runoff for key mountain regions, *Hydrological Processes*, 23, 78–94,](#)
894 [2009.](#)

895 Swanson, F.: Where roads will never reach: Wilderness and its visionaries in the Northern Rockies, University of Utah
896 Press. Salt Lake City, Utah, 2015.

897 [Tague, C., Farrell, M., Grant, G., Lewis, S., and Rey, S.: Hydrogeologic controls on summer stream temperatures in the](#)
898 [McKenzie River Basin, Oregon, *Hydrologic Processes*, 21, 3288–3300, 2007.](#)

899 ▼ [Todd, A.S., Coleman, M.A., Konowal, A.M., May, M.K., Johnson, S., Vieira, N.K.M., and Saunders, J.F.: Development](#)
900 [of new water temperature criteria to protect Colorado's fisheries, *Fisheries*, 33, 433-443, 2008.](#)

901 Trumbo, B.A., Nislow, K.H., Stallings, J., Hudy, M., Smith, E.P., Kim, D., Wiggins, B., and Dolloff, C.A.: Ranking site
902 vulnerability to increasing temperatures in southern appalachian brook trout streams in Virginia: an exposure-
903 sensitivity approach, *Transactions of the American Fisheries Society*, 143,173–187, 2014.

904 [Turschwell, M.P., Peterson, E.E., Balcombe, S.R. and Sheldon, F.: To aggregate or not? Capturing the spatio-temporal](#)
905 [complexity of the thermal regime, *Ecological indicators*, 67, 39-48, 2016.](#)

906 [Webb, B. W., Hannah, D. M., Moore, R. D., Brown, L. E., and Nobilis, F.: Recent advances in stream and river](#)
907 [temperature research, *Hydrological Processes*, 22, 902–918, 2008.](#)

908 Wenger, S.J., Luce, C.H., Hamlet, A.F., Isaak, D.J., and Neville, H.M.: Macroscale hydrologic modeling of ecologically
909 relevant flow metrics, *Water Resources Research*, 46:W09513, 2010.

910 Woodward, G., Perkins, D.M., and Brown, L.E.: Climate change and freshwater ecosystems: impacts across multiple
911 levels of organization, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365, 2093-2106,
912 2010.

913 [Zimmerman, D.L., and Ver Hoef, J.M.: The Torgegram for fluvial variography: characterizing spatial dependence on](#)
914 [stream networks, *Journal of Computational and Graphical Statistics*, 26, 253-264, 2017.](#)

915

Deleted: ¶

917 **Table 1.** Descriptive statistics for spatial attributes of the study network and 226 monitoring sites
 918 with annual temperature data in the northwestern United States.

Network reaches	Mean	Median	SD	Minimum	Maximum
<u>Elevation (m)</u>	<u>1,493</u>	<u>1,533</u>	<u>536</u>	<u>221</u>	<u>3,105</u>
<u>Drainage area (km²)</u>	<u>915</u>	<u>17.7</u>	<u>4,359</u>	<u>0.005</u>	<u>34,865</u>
<u>Mean annual flow (m³/s)</u>	<u>9.73</u>	<u>0.229</u>	<u>43.2</u>	<u>0.0253</u>	<u>379</u>
<u>Reach slope (m/m)</u>	<u>0.0584</u>	<u>0.0519</u>	<u>0.0429</u>	<u>0</u>	<u>0.150</u>
<u>Monitoring sites</u>					
Elevation (m)	1,392	1,407	464	280	2,369
Drainage area (km ²)	687	47.3	3,011	2.18	34,865
Mean annual flow (m ³ /s)	7.37	0.692	26.4	0.0253	281
Reach slope (<u>m/m</u>)	0.0389	0.0273	0.0403	0	0.150

919

920 **Table 2.** Temperature metrics used to describe thermal regimes of mountain rivers and streams.

Category	Thermal metric	Definition
Magnitude	M1. Mean annual temperature	Average of mean daily temperatures during a year
	M2. Mean winter temperature	Average of mean daily temperatures during December, January, and February
	M3. Mean spring temperature	Average of mean daily temperatures during March, April, and May
	M4. Mean summer temperature	Average of mean daily temperatures during June, July, and August
	M5. Mean August temperature	Average of mean daily temperatures during August
	M6. Mean fall temperature	Average of mean daily temperatures during September, October, and November
	M7. Minimum daily temperature	Lowest mean daily temperature during a year
	M8. Minimum weekly average temperature	Lowest seven-day running average of mean daily temperature during a year
	M9. Maximum daily temperature	Highest mean daily temperature during a year
	M10. Maximum weekly average temperature	Highest seven-day running average of mean daily temperature during a year
	M11. Annual degree days	Cumulative total of degree days during a year (1°C for 24 hours = 1 degree day)
Variability	V1. Annual standard deviation	Standard deviation of mean daily temperature during a year
	V2. Winter standard deviation	Standard deviation of mean daily temperature during winter months
	V3. Spring standard deviation	Standard deviation of mean daily temperature during spring months
	V4. Summer standard deviation	Standard deviation of mean daily temperature during summer months
	V5. Fall standard deviation	Standard deviation of mean daily temperature during fall months
	V6. Range in extreme daily temperatures	Difference between minimum and maximum mean daily temperatures during a year (M9 minus M7)
	V7. Range in extreme weekly temperatures	Difference between minimum and maximum weekly average temperatures during a year (M10 minus M8)
Frequency	F1. Frequency of hot days	Number of days with mean daily temperatures >20 °C
	F2. Frequency of cold days	Number of days with mean daily temperatures <2 °C
Timing	T1. Date of 5% of degree days	Number of days from December 1st until 5% of degree days are accumulated
	T2. Date of 25% of degree days	Number of days from December 1st until 25% of degree days are accumulated
	T3. Date of 50% of degree days	Number of days from December 1st until 50% of degree days are accumulated
	T4. Date of 75% of degree days	Number of days from December 1st until 75% of degree days are accumulated
	T5. Date of 95% of degree days	Number of days from December 1st until 95% of degree days are accumulated
Duration	D1. Growing season length	Number of days between the 95% and 5% of degree days (T5 minus T1)
	D2. Duration of hot days	Longest number of consecutive days with mean daily temperatures >20 °C
	D3. Duration of cold days	Longest number of consecutive days with mean daily temperatures <2 °C

921

922 **Table 3.** Descriptive statistics for temperature metrics used to describe thermal regimes at 226
 923 monitoring sites in mountain river networks. Statistics were calculated from the imputed time-series
 924 and mean daily values for the period 2011–2015.

	Mean (°C)	Median (°C)	SD (°C)	Minimum (°C)	Maximum (°C)
M1. Mean annual temperature	5.36	5.10	1.44	3.10	10.34
M2. Mean winter temperature	0.75	0.63	0.60	-0.10	4.03
M3. Mean spring temperature	3.67	3.47	1.61	1.14	9.38
M4. Mean summer temperature	11.2	10.9	2.68	6.55	19.1
M5. Mean August temperature	12.5	12.1	2.78	7.78	22.5
M6. Mean fall temperature	5.71	5.50	1.53	3.04	11.5
M7. Minimum daily temperature	0.21	0.14	0.35	-0.45	2.18
M8. Minimum weekly average temperature	0.31	0.23	0.40	-0.42	2.69
M9. Maximum daily temperature	13.5	13.0	3.00	8.26	23.5
M10. Maximum weekly average temperature	13.2	12.7	2.99	7.96	23.2
M11. Annual degree days	1956	1863	527	1132	3775
V1. Annual standard deviation	4.43	4.27	1.05	2.51	7.40
V2. Winter standard deviation	0.30	0.29	0.16	0.00	0.87
V3. Spring standard deviation	1.62	1.57	0.72	0.33	5.36
V4. Summer standard deviation	1.99	1.88	0.61	0.61	4.45
V5. Fall standard deviation	3.43	3.34	0.73	2.13	6.05
V6. Range in extreme daily temperatures	13.3	12.8	3.06	7.50	23.3
V7. Range in extreme weekly temperatures	12.9	12.3	3.06	6.99	22.9
F1. Frequency of hot days	0.81	0	5.82	0	61
F2. Frequency of cold days	131	132	35.6	0	212
T1. Date of 5% of degree days	109	113	25.5	44	168
T2. Date of 25% of degree days	193	194	10.9	148	217
T3. Date of 50% of degree days	237	238	5.01	215	251
T4. Date of 75% of degree days	276	276	2.99	264	288
T5. Date of 95% of degree days	323	323	4.78	309	340
D1. Growing season length	214	210	29.7	141	296
D2. Duration of hot days	0.691	0	5.61	0	61
D3. Duration of cold days	124	124	39.0	0	207

925

926 **Table 4.** Loadings of 28 temperature metrics on the first four principal components in a PCA of
 927 annual temperature records from mountain river networks in the northwestern United States.

Temperature metric	PC1	PC2	PC3	PC4
M1. Mean annual temperature	0.99	-0.07	-0.05	-0.03
M2. Mean winter temperature	0.26	-0.92	0.14	0.00
M3. Mean spring temperature	0.91	-0.19	-0.25	0.04
M4. Mean summer temperature	0.97	0.21	-0.06	-0.05
M5. Mean August temperature	0.95	0.22	0.16	-0.10
M6. Mean fall temperature	0.96	-0.18	0.14	-0.08
M7. Minimum daily temperature	-0.02	-0.86	0.08	-0.02
M8. Minimum weekly average temperature	-0.03	-0.90	0.08	0.00
M9. Maximum daily temperature	0.95	0.26	0.09	-0.08
M10. Maximum weekly average temperature	0.95	0.25	0.09	-0.07
M11. Annual degree days	0.99	-0.07	-0.05	-0.03
V1. Annual standard deviation	0.90	0.41	0.01	-0.07
V2. Winter standard deviation	0.69	-0.54	0.16	0.00
V3. Spring standard deviation	0.71	0.30	-0.55	0.04
V4. Summer standard deviation	0.42	0.32	0.78	-0.14
V5. Fall standard deviation	0.87	0.39	0.19	-0.12
V6. Range in extreme daily temperatures	0.93	0.33	0.08	-0.07
V7. Range in extreme weekly temperatures	0.93	0.33	0.08	-0.07
F1. Frequency of hot days	0.47	-0.01	0.30	0.82
F2. Frequency of cold days	-0.70	0.61	0.09	0.11
T1. Date of 5% of degree days	0.02	0.96	-0.10	0.01
T2. Date of 25% of degree days	-0.43	0.74	0.46	-0.08
T3. Date of 50% of degree days	-0.45	0.37	0.79	-0.16
T4. Date of 75% of degree days	-0.19	-0.51	0.72	-0.19
T5. Date of 95% of degree days	0.30	-0.88	0.12	-0.09
D1. Growing season length	0.03	-0.97	0.11	-0.03
D2. Duration of hot days	0.44	-0.03	0.32	0.84
D3. Duration of cold days	-0.64	0.66	0.07	0.11
Variance explained (%):	49.0%	29.0%	9.8%	5.6%
Cumulative variance (%):	49.0%	78.0%	87.8%	93.4%
Eigenvalue:	13.73	8.12	2.74	1.56


928

929 **Table 5.** Correlations among stream temperature principal components and spatial attributes of 226
 930 monitoring sites with annual data from river networks in the northwestern United States.

	Elevation	Mean flow	Reach slope	PC1	PC2	PC3	PC4
Elevation	1						
Mean flow	-0.34	1					
Reach slope	-0.10	-0.23	1				
PC1	-0.59	0.58	-0.34	1			
PC2	0.27	-0.06	-0.49	0.00	1		
PC3	-0.23	0.35	0.13	0.00	0.00	1	
PC4	0.12	0.54	-0.02	0.00	0.00	0.00	1

931
 932


933 **Fig. 1.** Locations of 226 monitoring sites overlaid on an August stream temperature scenario for the
934 29,600 km network in the study area. Stars denote where air temperature and stream discharge data
935 were obtained from a low-elevation site (294 m, northern station) and a high-elevation site (1850 m,
936 southern station).



937


938 **Fig. 2.** Linear regression trends between elevation and mean monthly temperatures at 226 river and
 939 stream sites during 2013 (data values are not shown for clarity). Values next to the trend lines are
 940 regression slopes and r^2 values from the regressions.

Formatted: Superscript




941

942 **Fig. 3.** Annual cycle of mean daily water temperatures (a), air temperatures (b), and discharge (c) at
943 a high-elevation site and a low-elevation site during two contrasting climate years. Discharge values
944 at the high elevation site are multiplied by ten for better visibility.




945

946 **Fig. 4.** Ordination plot that shows principal component scores of the first two axes derived from
947 water temperature data measured at 226 sites and summarized with 28 thermal metrics (a). Panels b
948 and c show principal component scores mapped to network locations.



949


950 **Fig. 5.** T-mode PCA results showing times when dominant spatial phases occurred in water
951 temperatures at 226 sites based on principal component eigenvector loadings during an average
952 year.



Deleted: in the northwestern United States

953


955 **Figure 6.** Thermal patterns during two periods with distinct spatial phases based on T-mode PCA
956 results (a). Day 50 occurs in mid-January and represents the homogenous winter period (b) whereas
957 day 250 occurs in late July and represents the heterogeneous period (c).



958

959 **Fig. 7.** S-mode PCA results showing principal component scores that describe temporal patterns in
 960 mean daily water temperatures for 226 stream sites during five years (a). Average daily air
 961 temperatures and discharge values from two monitoring stations are aligned with the principal
 962 component scores for comparative purposes. A plot of PC1 versus PC2 reveals that variation along
 963 the two axes differs by monthly and seasonal periods (b).

Deleted: among
 Deleted: we




964
 965

968
969
970
971

Fig. 8. Plot of S-mode eigenvector loadings from 226 stream sites on PC1 and PC2. Note that the range of variation in the PC1 loadings is small relative to the loadings along PC2, which indicates that most of the differences among sites were associated with the second principal component.

Formatted: Font: Not Bold

Formatted: Font: Not Bold




972
973

974
975
976
977

Fig. 9. Annual water temperature timing patterns reconstructed from S-mode PCs using the mean eigenvector loading value for PC1 and +/- 0.16 for PC2 to demonstrate the effects of strong negative loadings and positive loadings on PC2.

Formatted: Font: Not Bold

Formatted: Font: Not Bold




978
979

980
981
982

Fig. 10. Relationship between the S-mode eigenvector loadings from PC2 and the annual unit-area runoff in basins upstream of 226 water temperature sites.

Formatted: Left
Formatted: Font: Not Bold
Formatted: Font: Not Bold



983