

## Responses on the Referees 1 comments on the submitted manuscript "Geostatistical interpolation by Quantile Kriging" hess-2018-276

We are very thankful to the anonymous referees for their remarks on our submitted manuscript. We believe that they will significantly improve the quality of the manuscript. We use the nomination e.g. A1.13 (i.e. A<sub>(nswer)</sub>1 (no. of reviewer).13 (no. of comment)) and numbers (page, line, figures, tables) of the original manuscript submitted in order to address all queries raised:

### RC 1:

The proposed manuscript presents a new geo-statistical interpolation method (Quantile Kriging – QK) that is able to relax three of the main assumption/limitations of the most used Ordinary Kriging: 1) spatial stationarity of the process mean, 2) Gaussianity of the interpolated variable and 3) independence of the uncertainty on the estimation value. The work extends the formulations of other well-known kriging methods with logic and statistical rigour. Although the presented technique still has a major limitation in the ability to handle the presence of many zero values (as often happens when dealing with rainfall, especially at finer scales than the presented one), it can be considered an improvement on the state of the art and a contribution to the advancement of the field. Additionally, although the authors do not mention it in the manuscript (and should) there are many applications to a variety of environmental variables where the presence of zeros is not a problem and the presented technique could be better applied. The manuscript is very well written and easy to follow. I suggest the following improvements:

1. The introduction explains a lot about the evolution of kriging techniques. However a little bit more discussion about applications (especially to rainfall) their limitations in hydrology, the main challenges, etc... could help defining the framework.

A1.1: see A1.3

2. P.4 l. 18, many of the presented geostatistical techniques were developed in geological sciences, where the temporal evolution of the studied variables is often irrelevant. I would mention this to explain why the temporal variability is often ignored in kriging.

A1.2: see A1.3

3. I would mention spatio-temporal kriging and other similar techniques as attempts to incorporate the temporal variability. How is this method better/different (e.g. Gaussianity)? Examples:

- Snepvangers, J. J. J. C., Heuvelink, G. B. M., & Huisman, J. A. (2003). Soil water content interpolation using spatio-temporal kriging with external drift. *Geoderma*, 112, 253–271. [https://doi.org/10.1016/S016-7061\(02\)00310-5](https://doi.org/10.1016/S016-7061(02)00310-5)
- Sideris, I. V., Gabella, M., Erdin, R., & Germann, U. (2014). Real-time radar-rain-gauge merging using spatio-temporal co-kriging with external drift in the alpine terrain of Switzerland. *Quarterly Journal of the Royal Meteorological Society*, 140(April), 1097–1111. <https://doi.org/10.1002/qj.2188>


A1.3: We try to address all three comments (i.e. A1.1, A1.2 and A1.3) by rephrasing and extending the existing paragraph by the following, starting at p.4, l.18:

*[...]. The inclusion of a temporal behavior into the geostatistic models is mostly irrelevant for the original geological variables. However, the temporal variability of a variable becomes more prominent for other sciences, e.g. hydrology, where observations from raingauges over several time steps are implemented into the geostatistical models in order to generate spatial precipitation estimates. These estimates subsequently serve as input to the hydrological modelling (e.g. Syed et al. 2003; Basistha et al. 2008; Cole et al. 2008) over multiple time steps. Associated errors in the precipitation estimates may ultimately lead to greater errors in the subsequent discharge modelling (Kobold et al. 2005). These errors*

strongly depend on the spatial and temporal distribution of the input precipitation (Gabellani et al. 2007, Moulin et al. 2009) and may limit the accuracy of rainfall-runoff simulations.

There are geostatistical space-time models in order to incorporate the temporal variability of the variable, but they are primarily aiming on the extrapolation of the variable in time (Snepvangers et al. 2003). Therefore, they require a strong dependence of the variable over time, suited e.g. for groundwater modeling where temporal changes occur relative slowly. This temporal dependence might be absent for other variables, e.g. monthly precipitation. [...]

For demonstration purposes, we remove the seasonality from our observations by subtracting the mean precipitation of the specific calendar month from the observed precipitation. The graph below shows the autocorrelation (with lag 1) for all 226 raingauges (see graph below). There is hardly any (linear) dependence between the monthly precipitation of two successive months, i.e. if a specific month becomes “wet” or “dry” does hardly depend on if the preceding month was relative “wet” or “dry”.



#### References:

- Syed, K.H., Goodrich, D.C.. & Myers, D. E. (2003). Spatial characteristics of thunderstorm rainfall fields and their relation to runoff. *Journal of Hydrology*, 271, 1–21. [https://doi.org/10.1016/S0022-1694\(02\)00311-6](https://doi.org/10.1016/S0022-1694(02)00311-6)
- Basistha, A., Arya, D.S.. & Goel, N. K. (2008). Spatial Distribution of Rainfall in Indian Himalayas -- A Case Study of Uttarakhand Region. *Water resources Management*, 22, 1325–1346. <http://dx.doi.org/10.1007/s11269-007-9228-2>
- Cole, S.J., Moore, R.J. (2008). Hydrological modelling using raingauge- and radar-based estimators of areal rainfall. *Journal of Hydrology*, 358, 159 - 181. <https://doi.org/10.1016/j.jhydrol.2008.05.025>
- Kobold, M., Suselj, K. (2005). Precipitation forecasts and their uncertainty as input into hydrological models. *Hydrology and Earth System Sciences*, 9, 322-332. <https://www.hydrol-earth-syst-sci.net/9/322/2005/>
- Gabellani, S., Boni, G. Ferraris, L., von Hardenberg J., & Provenzale A. (2007). Propagation of uncertainty from rainfall to runoff: A case study with a stochastic rainfall generator. *Advances in Water Resources*, 30, 2061 - 2071. <https://doi.org/10.1016/j.advwatres.2006.11.015>
- Moulin, L., Gaume, E. & Obled C. (2009). Uncertainties on mean areal precipitation: assessment and impact on streamflow simulations. *Hydrology and Earth System Sciences*, 13, 99-114. <https://www.hydrol-earth-syst-sci.net/13/99/2009/>
- Snepvangers, J. J. J. C., Heuvelink, G. B. M., & Huisman, J. A. (2003). Soil water content interpolation using spatio-temporal kriging with external drift. *Geoderma*, 112, 253–271. [https://doi.org/10.1016/S0016-7061\(02\)00310-5](https://doi.org/10.1016/S0016-7061(02)00310-5)

4. Eq. 5: I am not sure why you fit a Beta distribution to the quantiles: isn't the Normal Score Transformation (NST) designed to work with empirical distributions?

A1.4: We used the Beta-distribution due to its definition on the interval  $[0, 1]$ , thus avoiding conditioning of the resulting distribution (from interpolation) at the extremes.

5. Eq.6 and Eq.7: You applied the NST, so isn't this  $E[F(U)] = m$  and same for Variance? Maybe I'm missing something

A1.5: Yes, you could rewrite Eq.6 and Eq.7 by using  $E[U(F_x(Z(x,t)))]$  instead. However, to our opinion, it does not make a difference.

6. Nowhere is explained how you calculate the variograms for all the interpolations you do. Maybe worth mentioning it somewhere.

A1.6: We add the following sentence at the end of the paragraph (p.8, l.4):

*[...]. The corresponding variograms are calculated using Kendall's tau for a robust interpolation (Lebrenz et al. 2017). [...]*

*Reference:*

- Lebrenz, H. & Bárdossy, A. (2017). Estimation of the variogram using Kendall's tau for a robust geostatistical interpolation. Journal of Hydrological Engineering, 22(9), 04017038. 10.1061/(ASCE)HE.1943-5584.0001568

7. Pg 7 top: you introduce the elevation dataset, but you don't explain why. Mention you use it for both EDK of the parameters and for the reference EDK of the rainfall process.

A1.7: We extend the adding the additional sentence at p.7, l.2:

*[...] The upscaled elevation ultimately serves as external drift for EDK of the parameters within QK and for the reference EDK with the original variable. [...]*

8. is the dry ratio the number of stations that recorded zero rainfall over the whole month over the total number of stations? Can you state this a bit more explicitly?

A1.8: We rephrase the sentence at p.7,l.9 in order to clarify the definition of the "dry ratio":

*[...] The observed average monthly precipitation over the twelve calendar months  $c$  is illustrated in Fig.3 along with the percentage of zero-value observations over all observations of the specific calendar month  $c$  (hereafter referred to as dry ratio), revealing a seasonal variation. [...]*

9. P.7, l. 18: if you fit a PDF for each month for each station, you have only 22 points to do it, it seems a very little number to be statistically robust. maybe one of the reasons why you need to fit mean and variance rather than the parameters?

A1.9: The 22 points might be a contributing factor but we rather believe that the resulting (very small) parameters  $\vartheta_{2,c}$  are an outcome from the extrapolation.

10. Eq. 8 and eq. 9: you here present both the distributions but don't explain why. Do you want to compare their performance? How did you select Gamma and Weibull distributions? Nowhere in the paper you comment on which one performs best overall.

A1.10: We used Gamma & Weibull – distributions as exemplary distribution, because:

1. they are both defined on the interval  $[0, \infty]$ ;
2. they are frequently used for the variable of monthly distribution;

3. they have only 2 parameters to be interpolated.

The intention of this paper is not to evaluate the distributions but rather to implement the general idea of Quantile Kriging. However, we agree on the inclusion of a statement on which one performs best (see A1.16)

11. P.8, l.27: You need to state that you do EDK with elevation as the drift. One of the problems I have in this comparison is that often EDK is performed with radar data, which probably would do better than elevation in defining the spatial pattern of rainfall. Can you comment on this?

A1.11: We didn't use radar data for two reasons: First, the availability of radar data is limited in South Africa: they are only available for a relatively short time, limited to urban centers and are not preprocessed/converted into rainfall sums. Secondly, radar images might be useful for real-time predictions but not for long-time (i.e. monthly or yearly) sums, where they show strong systematic errors (Pfaff, 2013).

*Reference:*

- Pfaff, T. (2013). Processing and analysis of weather radar data for use in hydrology. Ph.D. Thesis, Institute for Modelling Water and Environmental Systems, University of Stuttgart, <http://dx.doi.org/10.18419/opus-487>

12. P.9, l. 19: One of the drawbacks I observe is that QK does not estimate a higher uncertainty where there are less rain gauges, eg. top left corner of Figure 5f.

A1.12: Since the entire area (e.g. top left corner, Fig. 5f) shows the same standard deviation  $\sigma_K$ , the estimation uncertainty appears to be less dependent from the position of the rain gauges.

13. Explain what rho (eq. 12) represent, why you use it, what is its range, and what the optimal value)

A1.13: We use the Spearman rank correlation  $\rho_S$  as a non-parametric measure to describe the monotonic relation between estimator  $Z^*$  and estimation standard deviation  $\sigma_K$ , instead of the standard Pearson correlation coefficient, describing only the linear relation. We add the following explanation at p.9, l.26:

*[...]. The non-parametric Spearman rank correlation  $\rho_S$  describes the monotonic relation between the estimator  $Z^*$  and estimation standard deviation  $\sigma_K$ , ranging from -1 (negative) to +1 (positive) with 0 indicating its absence. [...].*

14. P 13: I find the explanation about chi squared a bit confusing. I could not understand what had to be uniform and why, until later on you introduce the histogram. Maybe worth introducing the histograms first? or at least explain more in details.

A1.14: Yes, we agree: the explanation could be more precise. We will explain in more detail by rewording the existing explanation on p.13, l.14 by:

*[...]. The test on uniformity verifies the estimated, conditional distribution  $F_{Z^*}$  by calculating its value  $F_{Z^*}(z(x_i, t))$  for every original observation  $z(x_i, t)$ . The resulting values (or quantiles) should be uniformly distributed on the interval  $[0, 1]$  (Bárdossy and Li, 2008). [...].*

15. Conclusions: You need to write more here, and remove one of the two paragraphs that are repeated (l. 20-26 or 27-3).

A1.15: Yes, they are actually repeating and we remove the first paragraph (p.14, l.20-26) and write more in an additional, subsequent paragraph (see A1.16)

16. I feel in general a little bit more discussion of the overall results could be introduced either in the Results and Discussion or the Conclusion section, including many of the comments I mentioned before.

A1.16: We include an additional paragraph at p.15,l.4, including comments from above:

*[...]. The variable of monthly precipitation, observed at 226 raingauges over 264 consecutive time steps, serves as input data. We selected the two parametric  $\Gamma$ -distribution and Weibull distribution, because they are defined on the interval  $[0, \infty]$  and are suitable to describe the variable of monthly precipitation. The selected distributions are fitted to the observations of a specific calendar month, implying an absence of temporal dependence between two sample members (e.g. between the monthly precipitation of December 2002 and December 2003). However, QK does accommodate temporal independence between consecutive observations, unlike existing spatio-temporal Kriging methods. In general, other types of distributions, with a higher number of parameters could be selected, especially in case of other variables of interest. Finally, we used elevation as external drift, both for the interpolation of the parameters within QK as well as for the reference EDK. [...].*

And add the following sentences into the last paragraph:

*at p.15,l.6: [...] In case of the estimator, QK- $\Gamma$  performs slightly better than QK-Wei for most of the selected objective functions. [...].*

*at p.15,l.8: [...] In general, QK-Wei shows a superior estimation of the associated uncertainty than QK- $\Gamma$ . [...].*

## Response to reviewer 2

We thank the reviewer for the valuable comments. We provide a detailed point by point answer to the reviewers remarks.

The authors propose a new kriging technique developed to regionalize non-normally distributed spatio-temporal variables. The approach, as I understand it, involves (i) fitting time series observations to a predetermined distribution type at each gauge independantly; (ii) using the fitted distribution to assign a probability of non-exceedance to each observation at each gauge; (iii) fit the subset of these probabilities corresponding to each observation time to (different) beta distributions; (iv) map the fitted probabilities to normal quantiles; (v) use the (now normally distributed and \*assumed\* second-order stationary) outcomes as a basis to apply OK or EDK, as described (rather cryptically) in section 2.1.2. While the approach is intriguing, its description in the paper lacks statistical rigor and minimal proofs and intuitions. Its application in cross validation suggests that it does a decent job at predicting some measure of rainfall, but so would most properly fitted interpolation methods. This does not mean that the estimator is BLUE (best or even efficient unbiased linear estimator) of the considered stochastic process. Actually, the apparent correlation between  $Z^*(x)$  and  $\sigma^2(x)$  (Figure 4) suggests that the process has some degree of heteroskedasticity. Under these conditions, linear models (like most kriging estimators) are not necessarily efficient (see, for instance, the Gauss Markov Theorem for ordinary least squares). Again, the approach is promising, but its exposition needs a major overhaul to be convincing.

The reviewer described the procedure we applied reasonably well. However note that the distributions fitted for each location are the same type (gamma for example). The parameters of the distribution have to be interpolated, this step is missing from the reviewers description.

An intuitive description of the procedure is based on the following properties of precipitation fields:

- The monthly (and daily) precipitation amounts for a given month often follow a skewed distribution.
- Monthly (even daily) precipitation amounts cannot be considered as stationary in space. Differences in expected precipitation amounts become clear if one considers long time accumulations.
- The precipitation generating meteorological processes are usually of large spatial extent. This means if there is heavy rainfall at one location it likely that other locations also have heavy rainfall.
- Correlations between time series of precipitation indicate a strong spatial dependence, while the spatial dependence of precipitation on a given time accumulations (day, month) usually show a much weaker spatial dependence.

A possible process model reflecting the above properties can be described as follows:

Let  $Y_0(x, t)$  be independent (for each different  $t$ ) normal stationary spatial fields with  $E[Y_0] = 0$  and  $D^2(Y_0) = 1$  for each  $t$ .

In order to reflect large scale meteorological processes the process  $M(t)$  is introduced. High  $M(t)$  values correspond to heavy rainfall covering the area - while low correspond to dry conditions. This  $M$  modifies the spatial process:

$$Y_1(x, t) = Y_0(x, t) + M(t) \quad (1)$$

Were  $M(t)$  is a process (in time) with zero mean. We may assume that the distribution of  $M(t)$  is normal. In this case for each  $x$   $Y_1(x, t)$  is normally distributed with  $N(0, d)$  with  $d = \sqrt{1 + \sigma_M^2}$ .

For each fixed  $t$  the distribution of  $Y_1(x, t)$  is  $N(M(t), 1)$ . Now  $Y_2$  is temporal non-exceedence probability at location  $x$  - formally:

$$Y_2(x, t) = \Phi_{0,d}(Y_1(x, t)) \quad (2)$$

where  $\Phi_{0,d}$  is the distribution function of  $N(0, d)$ . (By definition  $0 \leq Y_2(x, t) \leq 1$ .)

The rainfall is then *generated* as:

$$Z(x, t) = F_x^{-1}(Y_2(x, t)) \quad (3)$$

where  $F_x$  is the distribution function of rainfall at location  $x$ . The  $F_x$ s can be different for different  $x$  locations due to topography and other influencing factors. (These  $F_x$ -s can be interpolated - example see also in Mosthaf and Bardossy (2017)).

We use  $Y_2(x, t)$  for each  $t$  and assume that it follows a beta distribution. In fact its distribution depends on  $M(t)$ . If  $M(t) = 0$  for all  $t$ -s then monthly rainfall is fully characterised by independent realizations over space. In this case the distribution of  $Y_2$  is uniform for each  $t$ .

This however is not the case with observed data. The reason is that wet and dry conditions occur simultaneously over the whole area. This is controlled by  $M(t)$ . One can take  $M(t)$  for example as independent random variables or to follow an ARMA process. If  $M(t) \neq 0$  then the distribution of  $Y_2(x, t)$  for this  $t$  is not uniform. The reason for assuming it as beta was due to the fact that beta distributions are very flexible and can well describe distributions in  $[0, 1]$ . The exact form of the corresponding distribution would be something like:

$$G_t(v) = \Phi_{0,1} \left( \Phi_{M(t),1}^{-1}(v) \right)$$

However the use of this would require the estimation of  $M(t)$  for each  $t$ . We decided to use a simple beta distribution instead.

The introduction of  $M(t)$  is reasonable as it explains the difference between the correlation between stations and the spatial correlation calculated using a variogram type approach for a given time. The later correlations are usually lower (smaller ranges) which are increased by the common large scale weather described with  $M(t)$ . Note that the introduction of  $M(t)$  leads to a correlation of the precipitation time series even if the individual *snapshots*  $Y_0(x, t)$  are independent in space.

In our procedure we start with  $Z(x, t)$ , estimate and interpolate  $F_x$ . Then calculate  $Y_2$  for the observation locations. We interpolate  $Y_2$  and come back to  $Z(x, t)$ .

Spatial variograms are calculated for  $Y_2$  for each  $t$ , and  $Y_2$  is stationary in space. Non-stationarity and non-Gaussian distributions occur only for  $Z$ . That is the reason why we concentrate on  $Y_2$ .

**Non-Gaussianity:** In its canonical form, Ordinary Kriging is based on the method of moment (i.e. variance minimization subject to unbiasedness) and so is not technically restricted to normally distributed processes. Gaussianness is, however, required for maximum likelihood (ML) type estimations, which some studies have shown can be more efficient for kriging-based regionalization (e.g., Lark 2000). ML approaches become necessary to have unbiased prediction of both the mean and the variance when it comes to EDK or Universal Kriging, particularly when subject to more intricate error correlation structures (e.g., Muller 2015). Since non-gaussianity appears to be a key rationale for the proposed approach, it is important to be specific on that.

Non-Gaussianity is considered because of the usually skewed distribution of precipitation amounts for a given time step. The suggested model should enable a simulation of the precipitation amounts. Non-Gaussianity is only in the sense of the marginal distribution at a given time-step. The spatial dependence is considered to correspond to a multi-Gaussian copula. This kind of transformation is frequently used - for example for Lognormal Kriging.

**Beta-distribution:** The use of the beta distribution is intriguing, but more intuition is needed on the assumed underlying stochastic process. Please describe clearly the properties of the (space-time) stochastic process that you assume gives rise to the observed sample and use that to demonstrate Eqn 6 and 7 in a rigorous mathematical proof. I find the use of the beta distribution promising because a common interpretation is that it describes the distribution of the probabilities associated to a binomial process observed over a finite sample. Lets say that the binomial process in question is the exceedance of a given threshold (as eluded to in the manuscript). Then, if the underlying point process is identically distributed in space and if an identically sized sample is taken at each observation point, the proportion of observations lower than a given

threshold across all gauges will be beta-distributed. Perhaps thats a start?

The above description shows that the beta distribution is only a convenient tool, not a statistically rigorous approach. As the beta distribution is very flexible it provided an easy and quick approximation of the distribution of the non exceedence probabilities.

Stationarity: More fundamentally, a main issue that I have is that your approach involves fitting distributions independantly at different points in space (gamma or weibull) and time (beta), which implies that the underlaying random point process follows a different distribution at each point in space. Granted, EDK and Universal kriging allow the first moment of the underlaying distribution to vary through space, as allowed by the scaling properties of the expectedated value estimator. However, I am not aware of any existing geostatistical approach that allows for higher order moments to vary through space. This may be fine, but please demonstrate that your approach does not violate the second order stationarity assumption (i.e that the variogram is constant through space), which is critical (and arguably more important than gaussianity) for kriging.

The distributions fitted to the individual locations are supposed to have a spatial dependence. Further they are assumed to follow the same distribution. These distributions are then interpolated. Here we use the assumption that the distributions show a much clearer effect of the large scale rainfall generating meteorological processes than a sigle mothly (or daily) realization would show. Therefore the use of extrenal covariates , such as topography is more appropriate for this interpolation. The use of these distributions transforms the process to a staionary one. The stationary process is interpolated using the beta distribution of the non-exceedance probabilities. The reason for this is that we intended to avoid problems with interpolated probabilities being outside the  $[0, 1]$  interval.

We intend to to revise the paper and to include the above description and discussions.

Mosthaf, T. and A. Bárdossy, Regionalizing non-parametric precipitation amount models on different temporal scales, *Hydrology and Earth System Sciences*, **21** , 2463-2481, 2017

## Responses on the Referees 3 comments on the submitted manuscript "Geostatistical interpolation by Quantile Kriging" hess-2018-276

We are very thankful for your remarks and reply on them below:

### **RC 3:**

The article presents an interesting approach to kriging with skewed variables and to non-stationarity: i) For every single location the distribution over time is estimated and quantiles are estimated. ii) To the quantiles of a given time-step a Beta-distribution is fit-ted. iii) The quantiles of the Beta-distribution are transformed by a Normal-Score trans-formation into standard Gaussian variables. iv) Ordinary kriging of the transformed variables. v) Backtransformation of the kriging results to the original scale

One to my opinion main result now is that the variance of the prediction is dependent on the data values themselves, too, and not as in ordinary kriging only dependent on the kriging location. The methodology reminds me somewhat to trans-Gaussian kriging, where you have a similar effect, with the difference that you are still stationary. Maybe you could a little bit comment on this and also on the relationship to copulas.

A3.1: The interpolation of the beta distributed distribution function values can be seen as a Trans-Gaussian Kriging. Trans-Gaussian Kriging (Spöck et al., 2009) can also be interpreted as a Gaussian copula based linear interpolation.

#### *Reference:*

- Spöck, G., Kazianka, H. & Pilz, J. (2009). Modeling and Interpolation of Non-Gaussian Spatial Data: A Comparative Study. Dept. of Statistics, Alpen-Adria Universität, Klagenfurt.  
[https://www.stat.aau.at/Tagungen/statgis/2009/StatGIS2009\\_Spoeck\\_2.pdf](https://www.stat.aau.at/Tagungen/statgis/2009/StatGIS2009_Spoeck_2.pdf)

Non-stationarity comes into play because you estimate at each spatial location the quantiles separately. You calculate quantiles, and quantiles are always related to copulas,- is there also here a relationship to copulas? Please, elaborate on that. I am also not completely sure, why you need the Beta-distribution at all and not directly calculate the Normal-Score transformation.

A3.2: compare to reply on comment 4 from Referee 1: we used the Beta-distribution due to its definition on the interval  $[0,1]$ , thus avoiding conditioning of the resulting distribution at the extremes of 0 and 1.

## hess-2018-276 – List of changes

We undertook the following, major changes (compare reviewers comments):

(Pagennumber/ linenumber acc. "hess-2018-276\_QuantileKriging\_markedup.pdf")

p4, l.18 – l.29 :	added an paragraph	(acc. comment Reviewer 1)
p5,l.5 p7, l.5 – p8, l.31	rephrase sentence and added chapter	(acc. comment Reviewer 2,3)
p9, l.8 – l.9 :	added sentence	(acc. comment Reviewer 1)
p9, l.17 :	rephrased sentence	(acc. comment Reviewer 1)
p10, l.12 – l.13 :	added sentence	(acc. comment Reviewer 1)
p12, l.7 – l.9 :	rephrased sentence	(acc. comment Reviewer 1)
p15, l.18 – l.20 :	rephrased sentence	(acc. comment Reviewer 1)
p16, l.25 – p17, l.23:	rephrased/rewrote paragraph	(acc. comment Reviewer 1)

# Geostatistical interpolation by Quantile Kriging

Henning Lebrez<sup>1,2</sup> and András Bárdossy<sup>2</sup>

<sup>1</sup>University of Applied Sciences and Arts - Northwestern Switzerland, Institute of Civil Engineering, Switzerland

<sup>2</sup>University of Stuttgart, Institute for Modelling Hydraulic and Environmental Systems, Germany

**Correspondence:** Henning Lebrez (henning.lebrez@fhnw.ch)

**Abstract.** The widely applied geostatistical interpolation methods of Ordinary Kriging (OK) or External Drift Kriging (EDK) interpolate the variable of interest to the unknown location, providing a linear estimator and an estimation variance as measure of uncertainty. The methods implicitly pose the assumption of Gaussianity on the observations, which is not given for many variables. The resulting 'best linear and unbiased estimator' from the subsequent interpolation optimizes the mean error over many realizations for the entire spatial domain and, therefore, allows a systematic under- (over-) estimation of the variable in regions of relatively high (low) observations. In case of a variable with observed time-series, the spatial marginal distributions are estimated separately for one time step after the other, and the errors from the interpolations might accumulate over time in regions of relatively extreme observations.

Therefore, we propose the interpolation method of Quantile Kriging (QK) with a two step procedure prior to interpolation: we firstly estimate distributions of the variable over time at the observation locations and then estimate the marginal distributions over space for every given time step. For this purpose, a distribution function is selected and fitted to the observed time-series at every observation location, thus converting the variable into quantiles and defining parameters. At a given time step, the quantiles from all observation locations are then transformed into a Gaussian-distributed variable by a twofold quantile-quantile transformation with the Beta- and the Normal-distribution function. The spatio-temporal description of the proposed method accommodates skewed marginal distributions and resolves the spatial non-stationarity of the original variable. The Gaussian-distributed variable and the distribution parameters are now interpolated by OK and EDK. At the unknown location, the resulting outcomes are reconverted back into the estimator and the estimation variance of the original variable. As a summary, QK newly incorporates information from the temporal axis for its spatial marginal distribution and subsequent interpolation and, therefore, could be interpreted as a space-time version of Probability Kriging.

In this study, QK is applied for the variable of observed monthly precipitation from raingauges in South Africa. The estimators and estimation variances from the interpolation are compared to the respective outcomes from OK and EDK. The cross-validations shows that QK improves the estimator and the estimation variance for most of the selected objective functions. QK further enables the reduction of the temporal bias at locations of extreme observations. The performance of QK, however, declines when many zero-value observations are present in the input data. It is further revealed that QK relates the magnitude of its estimator with the magnitude of the respective estimation variance as opposed to the traditional methods of OK and EDK, whose estimation variances do only depend on the spatial configuration of the observation locations and the model settings.

*Copyright statement.* TEXT

## 1 Introduction

Many environmental variables (e.g. precipitation, ore grades) are only measured at some distinct observation locations, but possess a highly variable and unknown spatial distribution (Armstrong, 1998). The regionalization of the variable, i.e. the interpolation of the variable to the unknown locations, attempts the full description of its spatial distribution as a prerequisite for practical objectives (e.g. hydrological modeling, efficient exploitation of resources). However, a deterministic description of the spatial distribution is severely hampered for many variables since they incorporate a complex genesis which is neither fully known nor understood.

Therefore, the assessment of the distribution by geostatistical models arose, whose theoretical fundamentals were firstly laid out by Matheron (1962). The theory regards the (observed) values of a variable  $z$  as one realization  $z(\mathbf{x})$  of a Random Variable (RV)  $Z$  at the specific location  $\mathbf{x}$  ( $= \{x_1, x_2\}$  for  $\mathbb{R}^2$ ). Since the variable is often only partially known at a few distinct measurement locations  $\mathbf{x}_i$ , ergodicity is assumed and the intrinsic hypothesis for Ordinary Kriging (OK, Matheron (1965)) is given as:

$$E[Z(\mathbf{x})] = m(\mathbf{x}) = \text{const.} \quad (1)$$

$$\text{VAR}[Z(\mathbf{x} + \mathbf{h}) - Z(\mathbf{x})] = 2\gamma(\mathbf{h}) \quad (2)$$

in which  $\gamma(\mathbf{h})$  is the semi-variogram. The increment  $Z(\mathbf{x} + \mathbf{h}) - Z(\mathbf{x})$  is assumed as a stationary random function and its variance only depends on the translation vector  $\mathbf{h}$ . The set of outcomes by the interpolation to the unknown location  $\mathbf{x}$  is described by the linear estimator  $Z^*(\mathbf{x})$  as a measure of their centrality and by the estimation (or Kriging) variance  $\sigma_K^2(\mathbf{x})$  as a measure of the associated uncertainty.

The stated hypothesis entails three implications: the first condition of the intrinsic hypothesis (Eq.(1)) demands the variable to be spatially stationary and yields an unbiased estimation error in the entire domain (Chilès and Delfiner, 1999). Therefore, a systematic under- (over-) estimation is induced for regions of high (low) observations. Secondly, the marginal distribution of the observed data should ideally be Gaussian in order to be adequately described. Unfortunately, the distribution often departs from the ideal case (Journal and Alabert, 1989), necessitating an a-priori transformation of the marginal distribution. And last, the second condition (Eq.(2)) implies that the magnitude of the Kriging variance  $\sigma_K^2(\mathbf{x})$  only depends on the spatial configuration of the observation locations, the a-priori variance of all observations and on the selected variogram model, but not on the magnitude of the linear estimator  $Z^*(\mathbf{x})$  itself (Goovaerts, 2000).

The theoretical extension of External Drift Kriging (EDK, Ahmed and deMarsily (1987)) addresses the first implication. EDK can be attributed to the non-stationary geostatistical interpolation methods and has been frequently applied in various disciplines of practice and science (e.g.: Bourennane et al. (2000), van de Kasstele et al. (2009), Motaghian and Mohammadi (2011)). It incorporates additional information from external variables (or drifts)  $Y_j(\mathbf{x})$  for the estimation of the variable at the

unknown location. The mean  $m(\mathbf{x})$  is non-stationary but linearly depends on the external variables. Thus, the first condition (Eq.(1)) is reformulated to:

$$E[Z(\mathbf{x})] = a + \sum_{i=1}^k b_i \cdot Y_i(\mathbf{x}) \quad (3)$$

5 where  $k$  is the number of the incorporated external drifts  $Y_i(\mathbf{x})$ , while  $a, b_i$  are the unknown constants. The drifts are required to be known prior to interpolation at all relevant locations. Ideally, EDK still requires a marginal Gaussian distribution.

The non-parametric methods of Indicator Kriging and Probability Kriging ( Journel, 1983) are further derivatives and allow an a-priori transformation of the skewed marginal distribution. Indicator Kriging transforms the variable  $z$  into a binary variable by defining a threshold value  $z_{th}$  and restates the first condition (Eq.(1)) of the random function  $I(\mathbf{x}; z_{th}) = 1_{Z(\mathbf{x}) < z_{th}}$  to:

$$E[I(\mathbf{x}; z_{th})] = m = const. \quad (4)$$

10 This non-linear transformation is relatively robust and limits the effect of high values on the description of the variable at the unknown location. However, a loss of information comes along by the transformation into a binary variable. Therefore, Probability Kriging defines multiple thresholds and uses the order relation of the observed variable ( Carr and Mao, 1993), being implemented by using co-kriging in the derived multi-variate context. Both non-parametric methods have been subject to research, especially for detection limit problems of groundwater contamination (e.g. Goovaerts et al. (2005), Lee et al. (2007),  
15 Adhikary et al. (2011)).

In summary, geostatistical methods have been derived in the past in order to address the stated shortfalls of the intrinsic hypothesis. However, all present methods only regard the observations from the one respective time step for the estimation of their marginal spatial distribution, but do not incorporate observations from other time steps. The inclusion of a temporal behavior into the geostatistic models is mostly irrelevant for the original geological variables. However, the temporal variability of a variable becomes more prominent for other sciences, e.g. hydrology, where observations from raingauges over several time steps are implemented into the geostatistical models in order to generate spatial precipitation estimates. These estimates subsequently serve as input to the hydrological modelling (e.g. Syed et al. (2003), Basistha et al. (2008), Cole and Moore (2008)) over multiple time steps. Associated errors in the precipitation estimates may ultimately lead to greater errors in the subsequent discharge modelling (Kobold and Sušelj, 2005). These errors strongly depend on the spatial and temporal distribution of the input precipitation (Gabellani et al. (2007), Moulin et al. (2009)) and may limit the accuracy of rainfall-runoff simulations. There are geostatistical space-time models in order to incorporate the temporal variability of the variable, but they are primarily aiming on the extrapolation of the variable in time (Snepvangers et al. (2003)). Therefore, they require a strong dependence of the variable over time, suited e.g. for groundwater modeling where temporal changes occur relative slowly. This temporal dependence might be absent for other variables, e.g. monthly precipitation.

20  
25

In the following section, we introduce Quantile Kriging as a spatio-temporal description of the variable  $Z$ , addressing the three shortfalls: non-stationary variables, skewed marginal distributions over space and the independence of the error distribution from the magnitude of the observation.

## 2 Materials and Methods

- 5 The theory of Quantile Kriging (QK) is outlined along with the major theoretical implications, [followed by a general discussion of the underlying geostatistical model](#) and a case study for the variable of monthly precipitation is presented.

FIGURE 1

A preliminary analysis of the selected variable exemplary reveals (Fig. 1) the non-Gaussianity within the data and that the first assumption of the intrinsic hypothesis (see Eq.(1)) is not fulfilled since e.g.  $E_t[Z(\mathbf{x}_{57}, t)] \neq E_t[Z(\mathbf{x}_{29}, t)]$ .

### 10 2.1 Theory of Quantile Kriging

QK presumes the existence of observations of the variable  $z$  over consecutive time steps  $t$  ( $= 1, 2, \dots, J$ ) at every observation location  $\mathbf{x}_i = (x1_i, x2_i)$  [for the two-dimensional space  $\mathbb{R}^2$ ], providing an observed time series  $z(\mathbf{x}_i, t)$  at every observation location  $i$  ( $= 1, 2, \dots, n_i$ ). QK proceeds first with a two step procedure prior to interpolation (see Sect. 2.1.1) and second with the interpolation itself (see Sect. 2.1.2):

#### 15 2.1.1 Estimation of the temporal and the spatial marginal distribution

At first, the distribution over time is estimated at every observation location  $\mathbf{x}_i$ : an appropriate theoretical cumulative distribution function (cdf)  $F$  is selected and fitted to the corresponding time series of observations  $z(\mathbf{x}_i, t)$ , yielding  $n_i$  specific distributions  $F(z(\mathbf{x}_i, t))$ . The distributions are defined by their corresponding parameter sets  $\Theta(\mathbf{x}_i) (= \vartheta_k(\mathbf{x}_i)$  with  $k = 1, 2, \dots, K$ ) of the  $K$ -parametric distribution function  $F$ . The quantiles  $w(\mathbf{x}_i, t) (= F(z(\mathbf{x}_i, t); \Theta(\mathbf{x}_i)))$  are calculated from the observed values of the variable  $z(\mathbf{x}_i, t)$  and the defining parameter set  $\Theta(\mathbf{x}_i)$ . The quantiles  $w(\mathbf{x}_i, t)$  possess a uniform distribution over time on the interval  $[0, 1]$  for a given observation location  $\mathbf{x}_i$ . However, their empirical distribution in space is not uniform on  $[0, 1]$ . In order to profit from the optimality of Kriging, it requires a transformation into a Gaussian distribution as a prerequisite of the subsequent interpolation.

The marginal spatial distribution, corresponding to a time step  $t$  is, therefore, estimated by a twofold quantile-quantile conversion as the second step: the two-parametric Beta- distribution is fitted to the quantiles  $w(\mathbf{x}_i, t)$  of a given time step  $t$ , whose cdf  $G(w; \alpha, \beta)$  is defined as:

$$G(w; \alpha, \beta) = \sum_{n=\alpha}^{\alpha+\beta-1} \left[ \frac{(\alpha + \beta - 1)!}{n! \cdot (\alpha + \beta - 1 - n)!} \cdot w^n \cdot (1 - w)^{(\alpha+\beta-1-n)} \right] \quad (5)$$

on the interval  $[0, 1]$  by the two parameters  $\alpha > 0$  and  $\beta > 0$ . The quantiles  $G(w(\mathbf{x}_i, t); \alpha(t), \beta(t))$  from Eq.(5) are finally transformed by a Normal Score Transformation into the standard Gaussian variable  $u(\mathbf{x}_i, t)$  with  $N_u[0|1]$ , which ultimately serves as spatial marginal distribution to the subsequent geostatistical interpolation.

The transformation via the quantiles into the variable  $u$  accounts for spatially non-stationary distributions of the original variable  $z$  with  $E[Z(\mathbf{x}, t)] \neq m$  and exchanges the two conditions of Eq.(1) and (2) to:

$$E[F_x(Z(\mathbf{x}, t))] = m = \text{const.} \quad (6)$$

$$\text{VAR}[F_{x+h}(Z(\mathbf{x} + \mathbf{h}, t)) - F_x(Z(\mathbf{x}, t))] = 2\gamma(\mathbf{h}) \quad (7)$$

resolving the problem of spatial non-stationarity. QK can accommodate skewed marginal distributions of the original variable  $Z$ , which is similar to Probability Kriging, but it newly incorporates the temporal behavior of  $Z$  into its estimation of the spatial marginal distribution.

### 2.1.2 Interpolation to the unknown location

The outlined conversion of the variable  $z(\mathbf{x}_i, t)$  into the variable  $u(\mathbf{x}_i, t)$  and its corresponding parameter set  $\Theta(\mathbf{x}_i)$  entails separate interpolations to the unknown location  $\mathbf{x}$ .

The inherent assumption of second-order stationarity implies the existence of a constant spatial mean for the variable  $u$  within the domain for every time step  $t$ . The transformed quantiles are implicitly assumed to be more homogeneously distributed over space than the original variable  $z$ . The variable  $u$  is subject to a stationary geostatistical interpolation method (e.g. OK), providing a linear estimator  $U^*(\mathbf{x}, t)$  and the estimation variance  $\sigma_{K,U}^2(\mathbf{x}, t)$ . They jointly describe the Gaussian distribution of the random variable  $U(\mathbf{x}, t)$  with  $N[U^* | \sigma_{K,U}^2]$ .

The defining parameters  $\vartheta_k$  (for  $k = 1, 2, \dots, K$ ) of the K-parametric distribution function  $F$  are independent from the time step  $t$  and they are separately interpolated to the unknown location  $\mathbf{x}$ . The separate interpolation, however, requires the independence of the parameters  $\vartheta_k$  from each other. Therefore, a Principal Component Analysis examines the 'observed' parameters  $\vartheta_k(\mathbf{x}_i)$  in the Cartesian coordinate system  $(\vartheta_1 | \vartheta_2 | \dots | \vartheta_K)$  and determines the corresponding rotation angle  $\alpha$  and translation vector  $\mathbf{k}$ . The coordinate system is then rotated and translated accordingly prior to interpolation in order to ensure independence. A possible spatial non-stationarity of the parameters can be accounted for by the choice of an appropriate non-stationary interpolation method (e.g. EDK). The interpolation of the independent parameters yields their estimators at the unknown location  $\mathbf{x}$ , which are rotated and translated back to the original coordinate system. Thus, the estimators  $\vartheta_k^*(\mathbf{x})$  are defining the distribution function at the unknown location  $\mathbf{x}$ .

At last, the resulting Gaussian distribution of the random variable  $U(\mathbf{x}, t)$  is reconverted into a distribution of the original variable  $Z(\mathbf{x}, t)$  by the outlined steps of conversion (Section 2.1.1), but in reverse order: first the distribution of the quantiles  $G(W(\mathbf{x}, t); \alpha, \beta)$  of the Beta-distribution are calculated by the inverse of the Normal Score Transformation. The distribution of the quantiles  $W(\mathbf{x}, t)$  are calculated next using the inverse of Eq.(5) and last, the distribution of the original variable  $Z(\mathbf{x}, t)$  is estimated by using the inverse of the selected cdf, being defined by the estimators of its parameters  $\vartheta_k^*(\mathbf{x})$ . The reversion

of the distribution of  $U(\mathbf{x}, t)$  to the distribution of  $Z(\mathbf{x}, t)$  can be implemented by the simple numerical Rosenblueth point estimation method (Rosenblueth, 1975). The resulting distribution of the original variable  $Z$  is then described by the expectation value  $Z^*(\mathbf{x}, t)$  and the variance  $\sigma_K^2(\mathbf{x}, t)$ . Note that the resulting asymmetrical distribution of  $Z(\mathbf{x}, t)$  is non-Gaussian due to the conversion with the non-linear, but monotonic theoretical cdf  $F$ .

## 5 2.2 Discussion of the geostatistical model

Since the proposed method of QK is applied for the variable of monthly precipitation (see Chapter 2.3), the discussion of the underlying process model is based on the following properties of precipitation fields:

- The monthly (and even daily) precipitation amounts  $z(\mathbf{x}, t)$  for a given time step  $t$  often show a skewed distribution and cannot be considered as stationary over space. The differences in expected precipitation amounts become especially obvious for long time accumulations.
- The meteorological processes, which are generating precipitation, are usually of large spatial extent: if one location receives heavy precipitation, it is likely that other locations also receive heavy precipitation.
- Correlations between time series of precipitation indicate a strong spatial dependence, while the spatial dependence of precipitation at one given time step (e.g. day, month) usually show a much weaker spatial dependence.

15 A possible process model reflecting the above properties can be described as follows:

Let  $U(\mathbf{x}, t)$  be independent (for each different time step  $t$ ) normal stationary spatial fields with  $E[U] = 0$  and  $D^2(U) = 1$  for each time step  $t$ . Now, the process  $M(t)$  is introduced in order to reflect large scale meteorological processes. High  $M(t)$  values correspond to heavy precipitation covering the area, while low values correspond to dry conditions, as it is reflected by seasonal variations of precipitation amounts. The introduced  $M$  modifies the spatial process to:

$$20 \quad G(\mathbf{x}, t) = U(\mathbf{x}, t) + M(t) \quad (8)$$

where  $M(t)$  is a process (only in time) with a mean of zero. We may assume that the distribution of  $M(t)$  is normal and, therefore,  $G(\mathbf{x}, t)$  would be normally distributed with  $N(0, d)$  (with  $d = \sqrt{1 + \sigma_M^2}$ ) at every location  $\mathbf{x}$ .

For each individual time step  $t$ , the distribution of  $G(\mathbf{x}, t)$  is  $N(M(t), 1)$  and the resulting spatial field  $W$  is the temporal non-exceedance probability at location  $\mathbf{x}$  being confined to  $0 \leq W(\mathbf{x}, t) \leq 1$  and formally described as:

$$25 \quad W(\mathbf{x}, t) = \Phi_{0,d}(G(\mathbf{x}, t)) \quad (9)$$

where  $\Phi_{0,d}$  is the distribution function of  $N(0, d)$ . The precipitation is then generated as:

$$Z(\mathbf{x}, t) = F_x^{-1}(W(\mathbf{x}, t)) \quad (10)$$

where  $F_x$  is the distribution function of precipitation at the location  $\mathbf{x}$ . The distribution functions  $F_x$  may vary between different locations  $\mathbf{x}$  due to topography and other influencing factors, and they could be subject to interpolation (e.g. Mosthaf and Bárdossy (2010)).

We use  $W(\mathbf{x}, t)$  for each time step  $t$  and assume that it follows a Beta-distribution. In fact, its distribution depends on  $M(t)$ .

5 If  $M(t) = 0$  for all time steps  $t$ , then monthly precipitation can be fully characterized by independent realizations over space. In this case, the distribution of  $W$  is uniform for each  $t$ .

10 However, this is not the case with observed data because wet and dry conditions occur simultaneously over the entire domain. This is controlled by  $M(t)$ , which can be taken e.g. as an independent random variable or to follow an ARMA process. If  $M(t) \neq 0$  then the distribution of  $W(\mathbf{x}, t)$  is not uniform for this specific time step  $t$ . The exact form of the corresponding distribution would be something like:

$$G_t(v) = \Phi_{0,1} \left( \Phi_{M(t),1}^{-1}(v) \right) \quad (11)$$

However, the use of Eq.11 would require the estimation of  $M(t)$  for each time step  $t$ . We decided to use a simple Beta-distribution instead. The reason for assuming a Beta-distribution is due to their flexibility and their ability to describe distributions well within the interval  $[0, 1]$ .

15 The introduction of  $M(t)$  is reasonable as it explains the difference between the correlation between stations and the spatial correlation calculated using a variogram type approach for a given time step. The later correlations are usually lower (smaller ranges) which are increased by the common large scale weather described by  $M(t)$ . Note that the introduction of  $M(t)$  leads to a correlation of the precipitation time series even if the individual snapshots of  $U(\mathbf{x}, t)$  are independent in space.

20 We estimate and subsequently interpolate  $F_x$  within the proposed methodology by the preceding conversion of the variable  $Z(\mathbf{x}_i, t)$ . In addition, we calculate  $W(\mathbf{x}_i, t)$  for the observation locations  $\mathbf{x}_i$  and interpolate it to the unknown location  $\mathbf{x}$  in order to come back to  $Z(\mathbf{x}, t)$ . In here, spatial variograms are calculated for  $W$  for each time step  $t$ , assuming  $W$  to be spatially stationary. Non-Gaussian and non-stationary distributions only occur for the precipitation amounts (i.e. the variable  $Z$ ).

25 Non-Gaussianity should be considered due to the usually skewed distribution of precipitation amounts and it only applies to the marginal distribution at a given time step  $t$ . The suggested model should enable a simulation of the precipitation amounts. The spatial dependencies are considered to correspond to a multi-Gaussian copula, being a type of transformation frequently used (e.g. for Lognormal Kriging).

30 The distributions  $F_x$ , fitted to the individual locations, are supposed to have a spatial dependence. They are further assumed to follow the same distribution (e.g.  $\Gamma$ - or Weibull-distribution) and are subsequently interpolated. In here, we assume that the large scale meteorological processes, generating precipitation, are better reflected by the distributions than by a single monthly (or daily) realization. Therefore, the use of external covariates, e.g. elevation, is deemed more appropriate for their interpolation. The usage of these distributions transforms the process into a stationary one, which is then interpolated using the Beta-distribution of the non-exceedance probabilities.

## 2.3 Application of Quantile Kriging

The proposed method of QK is applied for the variable of monthly precipitation in South Africa and the outcomes are compared to those from OK and EDK.

### 2.3.1 Study Area and Data

- 5 The rectangular study area ( $3.5^\circ \times 3.5^\circ$ , Fig. 2) covers approx.  $132000 km^2$  and is located within the Republic of South Africa. The second release of the digital elevation model from the Shuttle Radar Topography Mission (USGS, 2003) serves as elevation input. The original resolution was upscaled from 3 arcseconds (approx.  $92m$ ) to 2 arcminutes (approx.  $3700m$ ) by spatial averaging, resulting in a mean of  $1442m$  and ranging from  $669m$  to  $2197m$  (a.m.s.l.). [The upscaled elevation ultimately serves as external drift for EDK of the parameters within QK and for the reference EDK with the original variable.](#)

10

FIGURE 2

- The observations of monthly precipitation were retrieved from raingauges of four different sources: the Department of Water Affairs (DWA, 2008), the Global Historical Climatology Network (Vose et al., 1992), the Climate Research Unit (Mitchell and Jones, 2005) and the internal database of the University of KwaZulu-Natal (Lynch, 2004). Accumulation of daily recordings yield monthly values for the 264 ( $= J$ ) consecutive months from January 1986 to December 2007. A total of 226 ( $= n_i$ ) raingauges (Fig. 2) provided 32226 ( $= N$ ) monthly precipitation values, which ultimately serve as input data.

- The observed average monthly precipitation over the twelve calendar months  $c$  is illustrated in Fig. 3 along with the percentage of zero-value observations [over all observations of the specific calendar month  \$c\$](#)  (hereafter referred to as the dry ratio), revealing a seasonal variation. High precipitation is typically encountered in the calendar months from October to March, being characterized by a low dry ratio  $< 3\%$ . The study area receives relatively low precipitation amounts during the calendar months from April (dry ratio =  $11\%$ ) to September (dry ratio =  $25\%$ ).

20

FIGURE 3

### 2.3.2 Adaptation to monthly precipitation

- At first, we subdivided the observations of monthly precipitation into the corresponding calendar month  $c$  ( $= 1, 2, \dots, 12$ ) prior to the fitting of the selected distribution function due to two reasons: the seasonal variation in monthly precipitation (Fig. 3) and to ensure independence of the individual sample members as a theoretical requirement for the fitting method. We used the maximum likelihood estimation method for fitting the selected distribution function to the respective measurements values  $z(\mathbf{x}_i, t_c)$  of every calendar month  $c$  and every measurement location  $\mathbf{x}_i$ , resulting in a total of 2712 ( $= 12 \times 226$ ) fittings. In this context, the two-parametric  $\Gamma$ - and Weibull- distribution were selected, whose cdf  $F(z; \Theta)$  are defined as:

$$\Gamma \text{ - distribution : } F(z; \Theta) = \frac{\gamma(\mu, \lambda \cdot z)}{\Gamma(\mu)} \quad (12)$$

$$\text{Weibull - distribution : } F(z; \Theta) = 1 - \exp\left[-\left(\frac{z}{\lambda}\right)^k\right] \quad (13)$$

where  $\Gamma(\mu)$  is the gamma function and  $\gamma(\mu, \lambda \cdot z)$  is the lower incomplete gamma function. The parameter set  $\Theta_c(\mathbf{x}_i)$  is composed for the  $\Gamma$ - distribution out of  $\mu_c(\mathbf{x}_i)$  ( $= \vartheta_{1,c}(\mathbf{x}_i)$ ) and  $\lambda_c(\mathbf{x}_i)$  ( $= \vartheta_{2,c}(\mathbf{x}_i)$ ) and for the Weibull- distribution out of  $k_c(\mathbf{x}_i)$  ( $= \vartheta_{1,c}(\mathbf{x}_i)$ ) and  $\lambda_c(\mathbf{x}_i)$  ( $= \vartheta_{2,c}(\mathbf{x}_i)$ ). All parameters are restrained to values greater than zero and both cdfs are defined for  $z(\mathbf{x}_i, t) \geq 0$ .

Thus, the original observations of monthly precipitation  $z(\mathbf{x}_i, t)$  are converted by Eq.(12) or Eq.(13) into the corresponding quantiles  $w(\mathbf{x}_i, t)$  ( $= F(z(\mathbf{x}_i, t); \Theta_c)$ ) and their defining parameter set  $\Theta_c(\mathbf{x}_i)$ . As outlined in Sect. 2.1, the quantiles  $w(\mathbf{x}_i, t)$  were further converted into the standard Gaussian variable  $u(\mathbf{x}_i, t)$ , ultimately subject to the subsequent OK as our chosen geostatistical interpolation method. Note that the stationary assumption of more homogeneously distributed quantiles in space appear more plausible in the case of monthly precipitation. In total, the variable  $u(\mathbf{x}_i, t)$  was interpolated 264 times by OK to the unknown location  $\mathbf{x}$ . [The corresponding variograms are calculated using Kendall's tau for a robust interpolation \(Lebrez and Bárdossy, 2017\).](#)

We further selected EDK as non-stationary interpolation method for the defining parameters  $\vartheta_{1,c}$  and  $\vartheta_{2,c}$ . Elevation data (Sect. 2.3.1) are taken as external drift since the distributions of monthly precipitation are assumed to depend on the altitude of the terrain. However, it was revealed that the direct use of the parameters may lead to negative or zero estimators at locations of extreme external drifts. Therefore, the sample mean  $\bar{z}_c(\mathbf{x}_i)$  and the sample variance  $\sigma_c^2(\mathbf{x}_i)$  are estimated instead, using the two statistical moments of the selected distribution functions, defined as:

$$\Gamma \text{ - distribution : } E[Z] = \frac{\mu}{\lambda}; \quad \text{VAR}[Z] = \frac{\mu}{\lambda^2} \quad (14)$$

$$\text{Weibull - distribution : } E[Z] = \lambda \cdot \Gamma(1 + 1/k); \quad \text{VAR}[Z] = \lambda^2 \cdot \Gamma(1 + 2/k) - E[Z]^2 \quad (15)$$

The dependence of the two derived parameters  $\bar{z}$  and  $\sigma^2$  on each other appears obvious in the case of monthly precipitation: a high mean is likely to be associated with a high variance and v.v.. Their dependence is exemplary illustrated for the calendar month 'May' in Fig. 4.

FIGURE 4

The principal component analysis allows for the transformation into the new Cartesian coordinate system with the new coordinates  $r_c(\mathbf{x}_i)$  and  $s_c(\mathbf{x}_i)$ . They are now independent and subject to a separate interpolation by EDK. A total of 24 ( $= 12 \times 2$ ) interpolations by EDK to the unknown location  $\mathbf{x}$  is performed for each selected type of distribution.

### 3 Results and Discussion

The proposed interpolation method of QK, using either a  $\Gamma$ -distribution (QK- $\Gamma$ ) or a Weibull-distribution (QK-Wei), is implemented and compared to the traditional geostatistical interpolation methods of OK and EDK. The respective performances are evaluated by cross-validation for the resulting estimators  $Z^*$  and the associated Kriging variances  $\sigma_K^2$ . In here, cross-validation eliminates all values  $z(\mathbf{x}_i, t)$  in turns from the input data, and subsequently calculates the estimator  $Z^*(\mathbf{x}_i)$  and the associated Kriging variance  $\sigma_K^2(\mathbf{x}_i)$  from the remaining data. Only the 32226 data points of the actually recorded values were considered for the cross-validation and the resulting outcomes are compared to the actual observations.

#### 3.1 Implementation of Quantile Kriging

The outcomes from the interpolation by OK, EDK and QK- $\Gamma$  are exemplary displayed and examined for a month with low precipitation and a high dry ratio (August 1993) and a month with high precipitation and a low dry ratio (January 1996). The respective spatial patterns of the estimator  $Z^*(\mathbf{x})$  and the associated standard deviation  $\sigma_K(\mathbf{x})$  are illustrated in Figs. 5 and 6.

FIGURE 5

FIGURE 6

The estimator  $Z^*$  displays similar spatial patterns and value ranges for all the interpolation methods. However, the local contours of the isohyets are more rugged for QK- $\Gamma$  (Figs. 5(e) and 6(e)) than for OK (Figs. 5(a) and 6(a)), but smoother than for EDK (Figs. 5(c) and 6(c)).

QK utilizes elevation for the interpolation of the two distribution parameters  $\vartheta_{1,c}$  and  $\vartheta_{2,c}$ . The two parameters incorporate information from all time steps  $t_c$  of the specific calendar month  $c$  and, thus, transfer information over time. They are further combined with the ordinary kriged quantiles  $W(\mathbf{x}, t)$ , leading to more smooth contours of the isohyets than EDK (compare Fig. 2). We regard the resulting spatial patterns of QK as more plausible, assuming that the accumulated monthly precipitation is hardly affected by local features in elevation.

The standard deviations  $\sigma_K$  of the associated estimation error show notable deviations in spatial pattern for the implemented interpolation methods. The range of error is notably higher for QK (Figs. 5(f) and 6(f)) and its spatial patterns deviates from the typical, bull-eye shaped patterns of OK (Figs. 5(b) and 6(b)) or EDK (Figs. 5(d) and 6(d)).

The estimation error from OK and EDK depends on the spatial configuration of the observation locations  $\mathbf{x}_i$ , their global variance and the selected variogram model. This typical spatial pattern of the error distribution from the ordinary kriged quantiles  $W(\mathbf{x})$  is converted within QK by the monotonic cdf (Eq. (12)). The resulting  $\sigma_K(\mathbf{x})$  of the original variable  $Z(\mathbf{x})$  is, therefore, increased by relatively flat slopes of the cdfs, which are encountered for relatively high values of  $W(\mathbf{x})$ . A relationship between the magnitude of the estimator  $Z^*$  and the magnitude of the associated standard deviation  $\sigma_K$  is suggested by Figs. 5 and 6.

### 3.1.1 Relationship between estimator and standard deviation

A relationship between the magnitude of the estimator  $Z^*$  and the associated standard deviation  $\sigma_K$  would be possibly an improvement to geostatistical interpolation and is, therefore, examined next by cross-validation. The Spearman rank correlation coefficient  $\rho_S$  is chosen for its description, being defined as:

$$\rho_S = \frac{\sum_{i=1}^n (rg(Z^*(\mathbf{x}_i, t)) - \overline{rg}_{Z^*}) \times (rg(\sigma_K(\mathbf{x}_i, t)) - \overline{rg}_\sigma)}{\sqrt{\sum_{i=1}^n (rg(Z^*(\mathbf{x}_i, t)) - \overline{rg}_{Z^*})^2 \times \sum_{i=1}^n (rg(\sigma_K(\mathbf{x}_i, t)) - \overline{rg}_\sigma)^2}} \quad (16)$$

where  $rg(Z^*(\mathbf{x}_i, t))$  and  $rg(\sigma_K(\mathbf{x}_i, t))$  are the ranks of the estimator  $Z^*$  and the associated standard deviation  $\sigma_K$  within a set of data, while  $\overline{rg}_{Z^*}$  and  $\overline{rg}_\sigma$  are the respective average ranks. [The non-parametric Spearman rank correlation  \$\rho\_S\$  describes the monotonic relation between the estimator  \$Z^\*\$  and estimation standard deviation  \$\sigma\_K\$ , ranging from  \$-1\$  \(negative\) to  \$+1\$  \(positive\) with  \$0\$  indicating its absence.](#) A set of data consists of all  $n$  values of the corresponding calendar month  $c$ . The evolution of the Spearman rank correlation coefficient  $\rho_S$  over all 12 calendar months is displayed in Fig. 7.

FIGURE 7

The rank correlation varies over the calendar months for all implemented interpolation methods and reach their seasonal maximums in June or July (Fig. 7), being characterized by a high dry ratio and low precipitation.

An improvement in the relationship between the estimator  $Z^*$  and the associated standard deviation  $\sigma_K$  can be observed for QK- $\Gamma$  and QK-Wei, exhibiting superior rank correlation coefficients for all calendar months with the exception of QK-Wei in December (Fig. 7). QK- $\Gamma$  deploys the strongest relation during the wetter months from October to March, while QK-Wei is superior from May to September. The resulting spread of the error distribution is increased by decreasing slopes of the theoretical cdfs (Eq. 12 and 13) and v.v.. The slope is effectively the probability density function (pdf). Both selected theoretical distributions imply a monotonic decrease in their respective pdfs for small parameters, being typically encountered during the dry season, and evoke a higher spread of the error distribution for higher monthly precipitation. Thus, the almost perfect rank correlation  $\rho_S(c)$  of QK during the months of low precipitation can be explained. The rank correlation between the estimator and the standard deviation is weakened for the months with higher precipitation due to the departure from the strict monotonic decrease of the pdfs, which is induced by increasing distribution parameters.

The inferior correlation coefficients of OK and EDK are nearly congruent due to their inherent geostatistical definition: although the Kriging weights are altered by the drift, they influence the linear estimator  $Z^*$  and the standard deviation  $\sigma_K$  by the same extent. Therefore, the non-parametric Spearman descriptor hardly differentiates between OK and EDK.

### 3.2 Cross-validation of the estimator

The estimator  $Z^*(\mathbf{x}_i, t_j)$  from the cross-validation is evaluated by six objective functions: the Pearson correlation coefficient  $\rho$ , the Nash-Sutcliffe-Efficiency coefficient NSE (Nash and Sutcliffe, 1970), the overall bias B1 and the Root Mean Square Error

(RMSE) are complemented by the temporal bias B2 and the spatial bias B3 (Bárdossy and Pegram, 2012), which are defined as:

$$\rho = \frac{\sum_{i=1}^{n_i} \sum_{j=1}^J (Z^*(\mathbf{x}_i, t_j) - \bar{Z}^*) \times (z(\mathbf{x}_i, t_j) - \bar{z})}{\sqrt{\sum_{i=1}^{n_i} \sum_{j=1}^J [Z^*(\mathbf{x}_i, t_j) - \bar{Z}^*]^2 \times \sum_{i=1}^{n_i} \sum_{j=1}^J [z(\mathbf{x}_i, t_j) - \bar{z}]^2}} \quad [-] \quad (17)$$

$$\text{NSE} = 1 - \sum_{i=1}^{n_i} \sum_{j=1}^J \frac{[z(\mathbf{x}_i, t_j) - Z^*(\mathbf{x}_i, t_j)]^2}{[z(\mathbf{x}_i, t_j) - \bar{z}]^2} \quad [-] \quad (18)$$

$$5 \quad \text{B1} = \frac{1}{n_{tot}} \sum_{i=1}^{n_i} \sum_{j=1}^J [z(\mathbf{x}_i, t_j) - Z^*(\mathbf{x}_i, t_j)] \quad [\text{mm}] \quad (19)$$

$$\text{B2} = \frac{1}{n_{tot}} \sum_{i=1}^{n_i} \left[ \sum_{j=1}^J (z(\mathbf{x}_i, t_j) - Z^*(\mathbf{x}_i, t_j)) \right]^2 \quad [\text{mm}^2] \quad (20)$$

$$\text{B3} = \frac{1}{n_{tot}} \sum_{j=1}^J \left[ \sum_{i=1}^{n_i} (z(\mathbf{x}_i, t_j) - Z^*(\mathbf{x}_i, t_j)) \right]^2 \quad [\text{mm}^2] \quad (21)$$

$$\text{RMSE} = \sqrt{\frac{1}{n_{tot}} \sum_{i=1}^{n_i} \sum_{j=1}^J [z(\mathbf{x}_i, t_j) - Z^*(\mathbf{x}_i, t_j)]^2} \quad [\text{mm}] \quad (22)$$

where  $J$  is the number of time steps,  $n_i$  is the number of observation locations and  $n_{tot}$  is the total number of cross-validated observations. Note that the cross-validation for only one time step ( $J = 1$ ) would yield the following relations:  $n_{tot} \times \text{B1}^2 = \text{B3}$  and  $\text{B2} = \text{RMSE}^2$ .

### 3.2.1 Summary results

The overall values of the six objective functions from all 32226 original observations, along with a separation into dry season (calender months: 4 - 9) and wet season (calender months: 1 - 3 and 10 - 12) are given in Table 1.

15

TABLE 1

The total values of the correlation coefficient  $\rho$ , the NSE coefficient, the temporal bias B2 and the RMSE are better for QK- $\Gamma$  and QK-Wei than for OK and EDK, evoking from a superior performance especially during the wet season when not many of many zero-values are present (see Table 1).

20 Complementary, OK and EDK have superior values for the biases B1 and B3 as a result of the implicit definition as best linear and unbiased estimator. OK (and to some extent EDK) optimize the spatial bias B3 for a given month by adapting their global mean to the observed mean, according to Eq. 1 (Eq.3). However, this evokes a systematic underestimation in regions of high precipitation and a systematic overestimation in regions of low precipitation. Therefore, a temporal bias B2 accumulates for a

location, which consistently experiences extreme precipitation over time. Especially during the wet season, QK outperforms OK and EDK with respect to the temporal bias. The following investigations on raingauge 'Wilgervier' exemplary serve as illustration for the evolution of a temporal bias.

### 3.2.2 Temporal bias at the raingauge 'Wilgervier'

5 The raingauge 'Wilgervier' ( $i = 125$ , see Fig. 2) records a relatively high monthly precipitation of  $70.1 \text{ mm}$  in average compared to the average monthly precipitation of  $54.7 \text{ mm}$  in the entire domain.

The evolution of the temporal bias  $B_2$  at the raingauge 'Wilgervier' is calculated from cross-validation according to Eq. 20 and illustrated in Fig. 8 (left). In addition, the relative estimation error  $\epsilon_r$  is estimated from the 218 (out of the 264 possible) original observations at 'Wilgervier', being defined as:

$$10 \quad \epsilon_r(\mathbf{x}_{125}, t_j) = \frac{Z^*(\mathbf{x}_{125}, t_j) - z(\mathbf{x}_{125}, t_j)}{z(\mathbf{x}_{125}, t_j)} \quad (23)$$

The 218 values of  $\epsilon_r(\mathbf{x}_{125})$  are smoothed by a Gaussian Kernel with a defined range  $d_G (= 0.35)$ . The distribution of the relative estimation errors should ideally be symmetrical around zero. However, the respective distributions are truncated due the confinement to  $\epsilon_r \geq -1$  for the variable of monthly precipitation. The smoothed distributions are factually a summary of the estimation errors and are illustrated in Fig. 8 (right).

15

FIGURE 8

OK displays the highest systematic underestimation over time (Fig. 8 (left)) and the relative estimation errors have a mode of  $-20\%$  (Fig. 8 (right)). EDK slightly improves the systematic bias of the interpolation, but the relative error distribution still possess a mode of  $-15\%$ . QK- $\Gamma$  and QK-Wei can further improve the systematic underestimation over time and exhibit error distributions with modes of  $-12\%$ , and  $-10\%$  respectively.

20

The raingauge 'Wilgervier' illustrates that OK and EDK might optimize the spatial bias (Table 1), but they are hampered to minimize the temporal bias in locations of extreme observations. QK, as a spatio-temporal interpolation method, is capable of reducing the temporal bias within regions of relatively high (or low) precipitation, which is potentially important for possible successive water balance considerations.

### 3.2.3 Cross-validation for different calendar months

25

The effects of the increased occurrence of zero-value observations on the Pearson correlation coefficient  $\rho$  (Eq. 17) and the RMSE (Eq. 22) is exemplary examined next. The respective values are calculated for each calendar month from the cross-validation and are illustrated in Fig. 9 along with the dry ratio (Fig. 3).

FIGURE 9

QK- $\Gamma$  and QK-Wei display improved values in comparison to OK and EDK for the two selected objective functions from October to March (Fig. 9). However, their performance deteriorates from May to September, when many zero-value observations are present, indicated by a dry ratio of at least 25% or above. The correlation coefficient  $\rho$  plunges in July for both versions of QK (Fig. 9 (left)) and the respective RMSE shows a similar qualitative behavior (Fig. 9 (right)).

5 The performance of QK is considerably influenced by the dry ratio. The presence of many zero-values in the data leads to very steep or nearly vertical theoretical cdfs, hampering the allocation of the quantiles to the respective precipitation values.

### 3.3 Cross-validation of the uncertainty

The estimated error distribution of the estimator  $Z^*(\mathbf{x})$  is described by the associated standard deviation  $\sigma_K(\mathbf{x})$  as a measure of associated uncertainty. The quality of the uncertainty from the cross-validation is assessed by two objective functions: the adapted Linear Error in the Probability Space LEPS (Ward and Folland, 1991) and a test on uniformity (Bárdossy and Li, 2008).

LEPS compares the values of the estimator  $Z^*(\mathbf{x}_i, t)$  and the observation  $z(\mathbf{x}_i, t)$  within the estimated cdf  $F_{Z^*}$  of the error distribution as:

$$\text{LEPS} = \frac{1}{n_{tot}} \cdot \sum_{i=1}^{n_{tot}} |F_{Z^*}(z(\mathbf{x}_i, t)) - F_{Z^*}(Z^*(\mathbf{x}_i, t))| \quad (24)$$

15 LEPS is defined on the interval  $[0, 1]$ : low values indicate a higher probability for the observation to originate from the estimated probability density distribution and v.v.. The average over the differences of all observations  $n_{tot}$  yields the overall LEPS value.

The test on uniformity ~~supposes that the values of the observations~~ verifies the estimated, conditional distribution  $F_{Z^*}$  by calculating its value  $F_{Z^*}(z(\mathbf{x}_i, t))$  for every original observation  $z(\mathbf{x}_i, t)$  from the estimated cdf  $F_{Z^*}(z(\mathbf{x}_i, t))$ . The resulting values (or quantiles) should be uniformly distributed on the interval  $[0, 1]$  (Bárdossy and Li, 2008). We defined ten classes of equal width, which should have the same resulting relative frequency. The deviation from uniformity is quantified by the  $\chi^2$ -test variable as the sum of the relative squared differences between uniformity and empirical distribution, ranging from zero (perfect) to nine (improper).

#### 3.3.1 Summary results

25 The values of the two objective functions from cross-validation of all 32226 original observations of the entire year, and divided into dry (calendar months: 4 - 9) and wet season (calendar months: 1 - 3 and 10 - 12) are displayed in Table 2.

TABLE 2

The best overall LEPS values are received from the traditional EDK and OK (Table 2). QK-Wei is superior to QK- $\Gamma$ , but both versions of QK are displaying higher LEPS values than OK or EDK, originating from the dry season when many zero-values are present in the data.

However, the  $\chi^2$ -test variables (Table 2) exhibit a reverse hierarchy among the implemented interpolation methods: QK is superior during the dry season and similar during the wet season. The  $\chi^2$ -test variables should be read in conjunction with the corresponding histograms of the  $F_{Z^*}$  values (Fig. 10). Note that the outer classes of the histograms host all the observations  $z(\mathbf{x}_i, t)$ , which are situated outside the estimated distribution. These classes exhibit the largest deviation from the ideal uniform distribution.

FIGURE 10

QK- $\Gamma$  and QK-Wei provide in general a more uniform distribution of  $F_{Z^*}$  than OK and EDK (Table 2), which possess the same value of the  $\chi^2$ -test variable and similar histograms (Table 2 and Fig. 10) due to their implicit affinity in definition.

### 3.3.2 Cross-validation for different calendar months

The effect of many zero-value observations on the error distribution is investigated by the differentiation into calendar months. The objective functions are recalculated accordingly and illustrated in Fig. 11.

FIGURE 11

The temporal evolution of the LEPS values for the two versions of QK is influenced by the presence of many zero-value observations. QK- $\Gamma$  and QK-Wei exhibit LEPS values superior to OK and EDK from September to April, characterized by a dry ratio of less than 26% (Fig. 11 (left)). However, the performance of QK deteriorates from May to August when many zero-value observations are present. This dependence explains the overall inferior LEPS values for QK in Table 2. The LEPS values for OK and EDK are hardly influenced by the dry ratio (Fig. 11 (left)) and show a congruent behavior.

The temporal evolution of the  $\chi^2$  test variable (Fig. 11 (right)) shows better values for QK- $\Gamma$  and QK-Wei than for OK or EDK during most calendar months. QK maintains a more uniform distribution of the  $F_{Z^*}$  values even for the months with a high dry ratio when OK and EDK deteriorate.

The cross-validation for the uncertainty suggests an improvement by QK under the prerequisite of a low dry ratio within the input data. This improvement is attributed to the wider range of the error distribution and the increased relation between the magnitude of the estimator and the spread of the distribution (see Section 3.1).

## 4 Conclusions

The geostatistical interpolation method of QK ~~firstly estimates the distribution of the variable over time and thereafter deducts the spatial distribution prior to interpolation, thus addressing the spatial~~ addresses the spatial non-stationarity of the a variable of interest ~~. The spatio-temporal by its conversion into quantiles and defining distribution parameters. The spatial-temporal~~ description of the variable by QK is a novelty in applied geostatistics and can be regarded as a temporal extension of Probability Kriging. The proposed method accommodates skewed marginal distributions ~~, converting and converts~~ them into an ideal Gaussian distribution prior to interpolation as a major theoretical advantage over the traditional OK or EDK. QK describes an

asymmetrical distribution of the random variable  $Z(\mathbf{x})$  by the non-linear estimator  $Z^*(\mathbf{x})$  and the estimation variance  $\sigma_K^2(\mathbf{x})$  of the error. QK additionally further establishes a relation between the magnitude of both descriptors.

The ~~geostatistical interpolation method of QK addresses the spatial non-stationarity of a variable of interest by its conversion into quantiles and defining distribution parameters. The spatial-temporal description of the variable by QK is a novelty in applied geostatistics and can be regarded as a temporal extension of Probability Kriging. The proposed method accommodates skewed marginal distributions and converts them into an ideal Gaussian distribution prior to interpolation as a major theoretical advantage over the traditional OK or EDK. QK describes an asymmetrical distribution of~~ variable of monthly precipitation, observed at 226 raingauges over 264 consecutive time steps, serves as input data. We selected the two parametric  $\Gamma$ -distribution and Weibull distribution, because they are defined on the interval  $[0, \infty]$  and are suitable to describe the variable of monthly precipitation. The selected distributions are fitted to the ~~random variable  $Z(\mathbf{x})$  by the non-linear estimator  $Z^*(\mathbf{x})$  and the estimation variance  $\sigma_K^2(\mathbf{x})$  of the error. QK further establishes a relation between the magnitude of both descriptors~~ observations of a specific calendar month, implying an absence of temporal dependence between two sample members (e.g. between the monthly precipitation of December 2002 and December 2003). However, QK does accommodate temporal independence between consecutive observations, unlike existing spatio-temporal Kriging methods. In general, other types of distributions, with a higher number of parameters could be selected, especially in case of other variables of interest. Finally, we used elevation as external drift, both for the interpolation of the parameters within QK as well as for the reference EDK.

The cross-validation of the estimator revealed an improvement for most of the selected objective functions. In particular, QK addresses the temporal bias, which remains unattended by the traditional geostatistical methods, which only optimize the mean spatial bias. In case of the estimator, QK- $\Gamma$  performs slightly better than QK-Wei for most of the selected objective functions.

20 The cross-validation of the associated uncertainty shows an improvement by QK in the description of the distribution of the estimation errors in comparison to the traditional geostatistical interpolation methods. However, its performance depends on the percentage of zero-values in the input data and declines when many zero-values are present. In general, QK-Wei shows a superior estimation of the associated uncertainty than QK- $\Gamma$ .

*Code availability.* Respective codes can be obtained from the corresponding author.

25 *Data availability.* Precipitation and elevation data can be obtained from the respective sources mentioned in Sect. 2.3.1

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* This research was executed at the Institute for Modeling Hydraulic and Environmental Systems of the University of Stuttgart.

## References


- Adhikary, P. P., Dash, C. J., Bej, R., and Chandrasekharan, H.: Indicator and probability kriging methods for delineating Cu, Fe, and Mn contamination in groundwater of Najafgarh Block, Delhi, India, *Environmental Monitoring and Assessment*, 176, 663–676, <https://doi.org/10.1007/s10661-010-1611-4>, <http://dx.doi.org/10.1007/s10661-010-1611-4>, 2011.
- 5 Ahmed, S. and deMarsily, G.: Comparison of geostatistical methods for estimating transmissivity using data on transmissivity and specific capacity, *Water Resources Research*, 23, 1717–1737, 1987.
- Armstrong, M.: *Basic Linear Geostatistics*, Springer, <http://books.google.de/books?id=-9vp11VuMCsC>, 1998.
- Basistha, A., Arya, D. S., and Goel, N. K.: Spatial Distribution of Rainfall in Indian Himalayas – A Case Study of Uttarakhand Region, *Water Resources Management*, 22, 1325–1346, <https://doi.org/10.1007/s11269-007-9228-2>, <https://doi.org/10.1007/s11269-007-9228-2>,  
10 2008.
- Bourennane, H., King, D., and Couturier, A.: Comparison of kriging with external drift and simple linear regression for predicting soil horizon thickness with different sample densities, *Geoderma*, 97, 255 – 271, [https://doi.org/http://dx.doi.org/10.1016/S0016-7061\(00\)00042-2](https://doi.org/http://dx.doi.org/10.1016/S0016-7061(00)00042-2), <http://www.sciencedirect.com/science/article/pii/S0016706100000422>, 2000.
- Bárdossy, A. and Li, J.: Geostatistical interpolation using copulas, *Water Resources Research*, 44, 1–15,  
15 <http://www.agu.org/pubs/crossref/2008/2007WR006115.shtml>, 2008.
- Bárdossy, A. and Pegram, G.: Interpolation of precipitation under topographic influence at different time scales, *Water Resources Research*, 49, 4545–4565, <https://doi.org/10.1002/wrcr.20307>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/wrcr.20307>, 2012.
- Carr, J. R. and Mao, N.-h.: A general form of probability kriging for estimation of the indicator and uniform transforms, *Mathematical Geology*, 25, 425–438, <https://doi.org/10.1007/BF00894777>, <http://dx.doi.org/10.1007/BF00894777>, 1993.
- 20 Chilès, J.-P. and Delfiner, P.: *Geostatistics: modeling spatial uncertainty*, Wiley, New York [u.a.], 1999.
- Cole, S. J. and Moore, R. J.: Hydrological modelling using raingauge- and radar-based estimators of areal rainfall, *Journal of Hydrology*, 358, 159 – 181, <https://doi.org/https://doi.org/10.1016/j.jhydrol.2008.05.025>, <http://www.sciencedirect.com/science/article/pii/S0022169408002394>, 2008.
- DWA: Hydrological Information System (HIS), <http://www.dwa.gov.za/Hydrology/>, 2008.
- 25 Gabellani, S., Boni, G., Ferraris, L., von Hardenberg, J., and Provenzale, A.: Propagation of uncertainty from rainfall to runoff: A case study with a stochastic rainfall generator, *Advances in Water Resources*, 30, 2061 – 2071, <https://doi.org/https://doi.org/10.1016/j.advwatres.2006.11.015>, <http://www.sciencedirect.com/science/article/pii/S030917080700053X>, recent *Developments in Hydrologic Analysis*, 2007.
- Goovaerts, P.: Geostatistical approaches for incorporating elevation into the spatial interpolation of rainfall, *Journal of Hydrology*, 228, 113  
30 – 129, [https://doi.org/10.1016/S0022-1694\(00\)00144-X](https://doi.org/10.1016/S0022-1694(00)00144-X), <http://www.sciencedirect.com/science/article/pii/S002216940000144X>, 2000.
- Goovaerts, P., Avruskin, G., Meliker, J., Slotnick, M., Jacquez, G., and Nriagu, J.: Geostatistical modeling of the spatial variability of arsenic in groundwater of southeast Michigan, *Water Resources Research*, 41, n/a–n/a, <https://doi.org/10.1029/2004WR003705>, <http://dx.doi.org/10.1029/2004WR003705>, w07013, 2005.
- Journal, A. and Alabert, F.: Non-Gaussian data expansion in the Earth Sciences, *Terra Nova*, 1, 123–134,  
35 <https://doi.org/10.1111/j.1365-3121.1989.tb00344.x>, <http://dx.doi.org/10.1111/j.1365-3121.1989.tb00344.x>, 1989.
- Journal, A. G.: Nonparametric estimation of spatial distributions, *Journal of the International Association for Mathematical Geology*, 15, 445–468, <https://doi.org/10.1007/BF01031292>, <http://dx.doi.org/10.1007/BF01031292>, 1983.

- Kobold, M. and Sušelj, K.: Precipitation forecasts and their uncertainty as input into hydrological models, *Hydrology and Earth System Sciences*, 9, 322–332, <https://doi.org/10.5194/hess-9-322-2005>, <https://www.hydrol-earth-syst-sci.net/9/322/2005/>, 2005.
- Lebrezn, H. and Bárdossy, A.: Estimation of the Variogram Using Kendall's Tau for a Robust Geostatistical Interpolation, *Journal of Hydrologic Engineering*, 22, 04017 038, [https://doi.org/10.1061/\(ASCE\)HE.1943-5584.0001568](https://doi.org/10.1061/(ASCE)HE.1943-5584.0001568),  
5 <https://ascelibrary.org/doi/abs/10.1061/%28ASCE%29HE.1943-5584.0001568>, 2017.
- Lee, J.-J., Jang, C.-S., Wang, S.-W., and Liu, C.-W.: Evaluation of potential health risk of arsenic-affected groundwater using indicator kriging and dose response model, *Science of The Total Environment*, 384, 151 – 162, <https://doi.org/http://dx.doi.org/10.1016/j.scitotenv.2007.06.021>, <http://www.sciencedirect.com/science/article/pii/S0048969707006833>, 2007.
- 10 Lynch, S.: Development of a Raster Database of Annual, Monthly and Daily Rainfall for Southern Africa, Report 1156/1/03, Water Research Commission, Pretoria, RSA, 2004.
- Matheron, G.: *Traité de Géostatistique Appliquée*, in: *Mémoires du Bureau de Recherches Géologiques et Minières*, vol. I, p. 333, Paris, 1962.
- Matheron, G.: *Les variables régionalisées et leur estimation: une application de la théorie des fonctions aléatoires aux sciences de la nature*,  
15 Ph.D. thesis, Université de Paris, Paris, 1965.
- Mitchell, T. D. and Jones, P. D.: An improved method of constructing a database of monthly climate observations and associated high-resolution grids, *International Journal of Climatology*, 25, 693–712, <https://doi.org/10.1002/joc.1181>, 2005.
- Mosthaf, T. and Bárdossy, A.: Regionalizing nonparametric models of precipitation amounts on different temporal scales, *Hydrology and Earth System Sciences*, 21, 2463–2481, <https://doi.org/10.5194/hess-21-2463-2017>, <https://www.hydrol-earth-syst-sci.net/21/2463/2017/>,  
20 2017.
- Motaghian, H. and Mohammadi, J.: Spatial Estimation of Saturated Hydraulic Conductivity from Terrain Attributes Using Regression, Kriging, and Artificial Neural Networks, *Pedosphere*, 21, 170 – 177, [https://doi.org/http://dx.doi.org/10.1016/S1002-0160\(11\)60115-X](https://doi.org/http://dx.doi.org/10.1016/S1002-0160(11)60115-X), <http://www.sciencedirect.com/science/article/pii/S100201601160115X>, 2011.
- Moulin, L., Gaume, E., and Obled, C.: Uncertainties on mean areal precipitation: assessment and impact on streamflow simulations, *Hydrology and Earth System Sciences*, 13, 99–114, <https://doi.org/10.5194/hess-13-99-2009>, <https://www.hydrol-earth-syst-sci.net/13/99/2009/>,  
25 2009.
- Nash, J. and Sutcliffe, J.: River flow forecasting through conceptual models part I — A discussion of principles, *Journal of Hydrology*, 10, 282 – 290, [https://doi.org/10.1016/0022-1694\(70\)90255-6](https://doi.org/10.1016/0022-1694(70)90255-6), <http://www.sciencedirect.com/science/article/pii/0022169470902556>, 1970.
- Rosenblueth, E.: Point estimates for probability moments, *Proceedings of the National Academy of Sciences of the United States of America*,  
30 72, 3812–3814, <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=433085tool=pmcentrezrendertype=abstract>, 1975.
- Snepvangers, J., Heuvelink, G., and Huisman, J.: Soil water content interpolation using spatio-temporal kriging with external drift, *Geoderma*, 112, 253 – 271, [https://doi.org/https://doi.org/10.1016/S0016-7061\(02\)00310-5](https://doi.org/https://doi.org/10.1016/S0016-7061(02)00310-5), <http://www.sciencedirect.com/science/article/pii/S0016706102003105>, *pedometrics* 2001, 2003.
- Syed, K. H., Goodrich, D. C., Myers, D. E., and Sorooshian, S.: Spatial characteristics of thunderstorm rainfall fields and their relation to runoff, *Journal of Hydrology*, 271, 1 – 21, [https://doi.org/https://doi.org/10.1016/S0022-1694\(02\)00311-6](https://doi.org/https://doi.org/10.1016/S0022-1694(02)00311-6), <http://www.sciencedirect.com/science/article/pii/S0022169402003116>, 2003.
- USGS: SRTM Topography, United States Geological Survey, 2003.


van de Kasstele, J., Stein, A., Dekkers, A. L. M., and Velders, G. J. M.: External drift kriging of NO<sub>x</sub> concentrations with dispersion model output in a reduced air quality monitoring network, *Environmental and Ecological Statistics*, 16, 321–339, <https://doi.org/10.1007/s10651-007-0052-x>, <http://dx.doi.org/10.1007/s10651-007-0052-x>, 2009.

5 Vose, R. S., Schmoyer, R. L., Steurer, P. M., Peterson, T. C., Heim, R., T.R., K., and Eischeid, J. K.: The Global Historical Climatology Network: Long-term monthly temperature, precipitation, sea-level pressure, and station pressure data, Rep. ornl/cdiac-53, Carbon Dioxide Inf. Anal. Cent., Oak Ridge Natl. Lab., Oak Ridge, TN, 25pp, 1992.


Ward, M. N. and Folland, C. K.: Prediction of seasonal rainfall in the north nordeste of Brazil using eigenvectors of sea-surface temperature, *International Journal of Climatology*, 11, 711–743, <https://doi.org/10.1002/joc.3370110703>, <http://dx.doi.org/10.1002/joc.3370110703>, 1991.




**Figure 1.** Histogram from the times series of observed monthly precipitation for two random stations: 'Laingsnek',  $\mathbf{x}_{57}$  with  $\bar{z} = 81.3 \text{ mm}$  (left) and 'Tambotieboom',  $\mathbf{x}_{29}$  with  $\bar{z} = 38.3 \text{ mm}$  (right).




**Figure 2.** Study area, elevation and location of raingauges.




**Figure 3.** Average monthly precipitation (in mm) and dry ratio (in %) from 226 raingauges. Note that the dry ratio (dashed brown line) is indicated on the right axis.




**Figure 4.** Scatterplot of sample mean  $\bar{z}_5$  and sample variance  $\sigma^2_{z_5}$  of calendar month 'May' along with the principle components  $s_5$  and  $r_5$ .




**Figure 5.** Spatial patterns of the estimator  $Z^*(\mathbf{x})$  and the standard deviation  $\sigma_K(\mathbf{x})$  from OK, EDK and QK- $\Gamma$  for August 1993. Note that crosses indicate positions of raingauges.




**Figure 6.** Spatial patterns of the estimator  $Z^*(\mathbf{x})$  and the standard deviation  $\sigma_K(\mathbf{x})$  from OK, EDK and QK- $\Gamma$  for January 1996. Note that crosses indicate positions of raingauges.




**Figure 7.** Evolution of the Spearman rank correlation coefficient  $\rho_S$  between the estimator  $Z^*$  and the standard deviation  $\sigma_K$ .




**Figure 8.** Errors of the estimator  $Z^*(x_{125})$  at raingauge 'Wilgervier': Evolution of temporal bias B2 over the study period (left) and Smoothed distribution of the relative estimation error  $\epsilon_r$  (right).



**Figure 9.** Evolution of two objective functions for the estimator over the twelve calendar months: Correlation coefficient  $\rho$  (left) and RMSE (right). Note that the dry ratio (dashed brown line) is indicated as percentage on the right axis.



**Figure 10.** Histograms for the  $F_{Z^*}$ -values of four different interpolation methods.



**Figure 11.** Evolution of the two objective functions for the error distribution over twelve calendar months: LEPS (left) and  $\chi^2$  (right). Note that the dry ratio (dashed brown line) is indicated as percentage on the right axis.

**Table 1.** Summary results from the cross-validation of the estimator  $Z^*$  for the twelve calendar months of the entire year, and spit into dry (calendar months: 4 - 9) and wet (calendar months: 1 - 3 and 10 - 12) season.

	$n_{tot}$	$\rho$	NSE	B1	B2	B3	RMSE
	[ - ]	[ - ]	[ - ]	[mm]	[mm <sup>2</sup> ]	[mm <sup>2</sup> ]	[mm]
<i>Entire year:</i>							
OK:	32226	0.902	0.813	0.07	3486.34	49.67	26.48
EDK:	32226	0.897	0.803	0.02	3539.73	76.41	27.16
QK- $\Gamma$ :	32226	0.905	0.819	-0.45	3480.90	1346.32	26.06
QK-Wei:	32226	0.903	0.814	0.75	3418.60	2105.09	26.40
<i>Dry season:</i>							
OK:	16256	0.908	0.824	0.00	2663.82	11.42	11.05
EDK:	16256	0.904	0.816	0.01	2731.59	15.82	11.30
QK- $\Gamma$ :	16256	0.897	0.803	0.46	2164.02	1147.99	11.69
QK-Wei:	16256	0.875	0.748	2.38	2153.02	3323.72	13.24
<i>Wet season:</i>							
OK:	15970	0.801	0.637	0.14	2939.87	88.60	35.93
EDK:	15970	0.790	0.618	0.02	3034.41	138.09	36.86
QK- $\Gamma$ :	15970	0.809	0.654	-1.39	2509.63	1548.21	35.10
QK-Wei:	15970	0.810	0.654	-0.91	2523.36	864.63	35.05

**Table 2.** Summary results from the cross-validation of the estimation error for the entire year, and split into dry (calender months: 4 - 9) and wet (calender months: 1 - 3 and 10 - 12) season.

	$n_{tot}$	LEPS	$\chi^2$	$n_{tot}$	LEPS	$\chi^2$	$n_{tot}$	LEPS	$\chi^2$
	[ - ]	[ - ]	[ - ]	[ - ]	[ - ]	[ - ]	[ - ]	[ - ]	[ - ]
	<i>Entire year:</i>			<i>Dry Season:</i>			<i>Wet Season:</i>		
OK:	32226	0.25	0.13	16256	0.19	0.32	15970	0.30	0.17
EDK:	32226	0.24	0.13	16256	0.19	0.33	15970	0.30	0.17
QK-Γ:	32226	0.32	0.11	16256	0.36	0.10	15970	0.28	0.18
QK-Wei:	32226	0.26	0.12	16256	0.27	0.10	15970	0.25	0.18