Point-by-point response to the Referee Reports

This document is formatted as follows:

- 5 comments from the referees are shown in blue
 - author's responses are shown in black
 - author's changes in the manuscript are shown in italics
 - 1. Response to Anonymous Referee #1:
- 10

General comments

The paper goal is to examine the importance of soil moisture to influence the hydrological response of a catchment located in the Mediterranean (Southern France). Despite

- 15 very interesting the topic is not new as many studies (also not cited by the authors) have tried to understand the role of soil moisture in flood modelling. In particular, many of them demonstrated that soil moisture is a good proxy of the catchment initial conditions and it behaves better than many API-based indexes (see the SC1 comment in the Interactive discussion). In this respect, this study has the advantage of relying upon
- 20 a really dense network of soil moisture monitoring stations that can help better the understanding of the underlying rainfall-runoff generation processes. Said that, I think that the paper is of interest for the journal readership and potentially very interesting. It is also well detailed and written. I have to major comments that the authors should take into consideration:
- 25

Thank you very much for your review of the manuscript. Your comments are highly appreciated and will helped us to improve the the manuscript. Thank you for the positive evaluation of the topic of the study and the appreciation of the data set.

Regarding the missing literature references, we want to stress that our study is not a modelling exercise.

- 30 Thus, in the literature review, we chose to focus on papers that rely on the analysis of soil moisture, discharge and precipitation measurements as does our study. But of course flood modelling studies are undisputedly very insightful to address the topic of the paper. We re-examined the literature on the subject and added the references listed below under "List of additional literature" (recommended by the reviewers and additional ones) and referred to them in the Introduction and Discussion section.
- 35 Concerning the finding of several studies that soil moisture is an asset for the estimation of catchment initial conditions we added the sentence "Numerous modelling studies have shown the high sensitivity of the modelled runoff response to dini and the importance of estimates of dini at the catchment scale as initial conditions in event-based models (e.g. Castillo et al., 2003; Huang et al., 2007; LeLay and Saulnier, 2007; Berthet et al., 2009; Brocca et al., 2009b; Tramblay et al., 2010, 2012; Li et al., 2012; Massari et al., 2010, 2012; Li et al., 2010; Massari et al., 2010; Massa
- 40 2014a,b, 2015; Grillakis et al., 2016)." (p.3, l.34).

1) The paper is really too long and the richness of details and number despite commendable sometimes distracts from the general objectives and make the reading of the paper not really easy. I suggest to shorten the manuscript and reduce the number of

45 figures trying to generalize a little bit the results and reducing the numbers in the text which should already evident from the figures. I think this will make the paper gaining

in readability. Please, define clearly in method why certain indexes are introduced and their scope. Consider the use of a table the role of each index in case.

This was also brought up by Referee #2 (H. Gao). We agree that the manuscript was too long and
shortened the paper substantially. The main text was shortened by three pages. We deleted figures 2,12 and 13 of the original manuscript and figure 4 was integrated into figure 3. To do so, two subfigures of figure 3 were deleted. Several numbers were deleted in the text in the results section (p.14, l. 10, l.15 and l. 16; p. 15 l. 8-10 and l. 16; p. 16 l.20). As you proposed we included a table (Table 1 in the new manuscript) that gives the role of each index and helps to describe the soil moisture data analysis more

10 concisely.

2) The study of the temporal/spatial variability of soil moisture and its connection with land use is surely important but at times seem disconnected with the main research question (RQ) which is the understanding of the impact of soil moisture on runoff generation.

15 The result is that the two RQs higleted in the paper seem two distinct chapter. If the authors want to maintain such a similar structure I am convinced that connections exist and thus they should emphasized. With connections I mean the role of the land use, spatial variability and temporal variability of certain plot/station soil moisture

values in the runoff generation mechanism. If this was done, it is not really immediate

20 to get. The intro section lacks of a significant part of the literature in this topic. The authors could refer to the SC1 comment in the interactive discussion to improve this part.

Thank you very much for this valuable comment. We regret that we could not convey the connections

- 25 between the two research questions well enough. In the revised version much of the analysis of the impact of land use on soil moisture was shortened. In this way we hope that the impression of two distinct chapters is diminished. We emphasize that the first objective (to study the variability of soil moisture) is necessary to obtain a relevant and representative value for soil moisture at the catchment scale. This is crucial to address the main research question of the role of initial soil moisture on the
- 30 hydrologic response of a catchment. Thus we rephrased the description of the objectives in the abstract: "Thus, the objectives are twofold: (1) obtaining meaningful estimates of soil moisture at catchment scale from a dense network of in-situ measurements and (2) using this estimate of \mathscr{T}_{ini} to analyze its relation with ϕev calculated for many runoff events."(p.1 l.19)

Further, in the introduction we no longer formulate our objectives as 2 research questions but rephrased

35 it: "Relying solely on in-situ data, we aim to (i) obtain a meaningful estimate of catchment scale soil moisture and its uncertainty and (ii) answer the research question how does soil moisture at the event onset affect the hydrological response?" (p.5 l.1)

This was done in the conclusion as well by rephrasing the introductory sentence to: "To this end, two issues were addressed and exemplarily examined in the nested Gazel (3.4 km²) and Claduègne (43 km²)

- 40 catchments: (1) Obtaining a meaningful estimate of soil moisture at the catchment scale and (2) Analyzing the relation between initial soil moisture $\tilde{\theta}_{ini}$ and the hydrological response quantified as the event-based runoff coefficient ϕ_{ev} ." (p.21 I.29)
 - Based on my comment above I recommend the paper accepted after major revisions.
- 45 I would be happy to revise the paper again and to provide a more detailed technical

revision once the above comments will have been addressed.

Thank you for this recommendation and for your offer to provide a second revision of a reworked manuscript.

5

2. Response to H. Gao (Referee 2)

This paper reports a case study to investigate the impact of initial soil moisture on hydrological response. Although many similar experiments have been carried out in

10 tremendous papers, I still believe that this kind of field experiments shall be encouraged for publication in hydrological journals, because experiments and observations are fundamental to test or reject our scientific hypotheses. But considering the quality of the writing, further revision is needed for consideration to publish in HESS.

We want to thank you for your review of the manuscript and for appreciating the usefulness of ourstudy. We revised the manuscript according to your comments and the ones of the other reviewers.

1. This paper is too long with too many details. The abstract shall be shorten. Introduction looks okay, but more literature shall be discussed as mentioned by previous two reviewers. There are too many details in the Results and Discussion. Particularly, the Conclusions have two pages, which is absolutely not necessary.

20 We shortened the manuscript as proposed esp. the sections you pointed out. Also, as recommended by reviewer 1, we deleted figures 2,12 and 13 of the original manuscript and figure 4 was integrated into figure 3.

Thank you for the further literature recommendations that you provided.

We re-examined the literature and added the references listed below under "List of additional literature"
 (recommended by the reviewers and additional ones) and referred to them in the Introduction and Discussion section.

2. For the science part, the authors missed an important factor while discussing the relationship between soil moisture and hydrological response – the topography. As we all know, as hydrologists, topography has great impact on initial soil moisture and runoff

- 30 generation. For example, hillslopes, riparian areas, and plateau have significantly different runoff generation mechanisms (cf. Seibert et al., 2003; Savenije, 2010; Gao et al., 2014). Figure 1 shows that many continuous soil moisture observations are located in the areas near the channel network, right? This means that the observations are mainly reflecting the soil moisture dynamic on riparian areas. This can explain
- 35 the immediate response of rainfall soil moisture runoff. But what is happening on hillslopes? What is the impact of topography on your conclusions?

We agree that topography plays an important role in the relationship of soil moisture and the hydrologic response. It is true that we only mention it among other factors that determine runoff generation. In our study we desided to factor an land use rather than on topography as our initial hyperbasis was that in

40 study we decided to focus on land use rather than on topography as our initial hypothesis was that –in

our catchments - land use and the associated different soil types determine this relationship more than other factors such as topography and geology etc.

Following your comment, we looked into the relationship between topography and initial soil moisture.
There seems to be no direct relationship between the distance to the river and the plot mean soil moisture. That is illustrated by the following figure where every point represents the plot mean of initial soil moisture (the different colors of the points represent the different events).



Distance to stream [m]

We agree that it is not ideal that the locations of the measurements are clustered in the south of the

- 10 Gazel catchment close to the stream and not evenly distributed in the catchment. This is due to practical reasons (e.g. protected sites for the installation of the probes available at Le Pradel, linked with the urgency of on-alert measurements that had to be completed before the rain started, which was always challenging). The distance of the continuous soil moisture measurement sites at le Pradel is > 40 m for all sites. As the stream is incised into the bedrock, all measurement sites are at a considerably higher
- 15 elevation than the stream bed. They do represent hillslopes that are not connected to the river via influent groundwater. Only two of the three sensors installed in the north of the catchments are located in grasslands that are potentially connected to the river during rain events.

We added a sentence in the Section 2.2 Data availability to give information on topography: "Concerning
 topography, most of the sensors are located on hillslopes which is the dominant topographic zone
 according to Savenije (2010) in the catchment. Only two plots are located in the riparian area and are
 potentially connected to the stream during rain events." (p.6 l. 1)

We agree, that the information on distance to the river is not evident from figure 1, which is misleading in this regard and we thank you for pointing this out. Your comment is in agreement with the comment of Reviewer 3 who also demands a more detailed map. We revised figure 1 and show a closer zoom into the area in the south of the Gazel catchment, a scale bar in the zoom window and by provide information on the land use of the sites of the measurement sites.

5

3. Response to Anonymous Referee #3:

The manuscript "How does initial soil moisture influence the hydrological response? A case study from southern France" by Uber et al. sets out to explore the relevance of

- 10 soil moisture for the generation of floods. Although the overall topic is highly appreciated, I am not sure what the novelty of this paper is, nor did the authors convince me about what they really want to do. There are several critical points I would the authors encourage to invest a bit more effort in to develop this study:
- 15 Thank you very much for the review of our paper. Your comments are highly appreciated and helped us to improve the manuscript. We regret that we could not convey the objectives and the novelty of the paper sufficiently. Thank you for pointing out this important fact. In the revised version of the paper we reformulated the objectives and stress the novelty of the paper in the last sentences of the introduction: *"Thus, this study's novelty is to analyze the relation between* ϕ_{ev} and ϑ_{ini} when both are obtained from a
- 20 comprehensive, high resolution data set allowing the assessment of the uncertainty of the two variables. Relying solely on in-situ data, we aim to (i) obtain a meaningful estimate of catchment scale soil moisture and its uncertainty and (ii) answer the research question how does soil moisture at the event onset affect the hydrological response?" (p. 4 I. 33)

(1) It is not entirely clear what the objective of this study is. Is it intended to give an

- 25 overview of the spatial pattern of soil moisture and how these evolve over time? In spite of a relatively rich data set, is such an analysis really warranted by the data? In other words, are the conclusions drawn generally applicable or do they merely describe the relatively local study sites? If the latter, what can be learned from that? Or is the study rather meant to improve our understanding of soil moisture to generate flows?
- 30 Then much of analysis and discussion of the spatial pattern can be condensed.

As suggested, we shortened large parts of the analysis of the spatial patterns and the influence on land use of soil moisture (Methods: section 2.4 was considerably shortened; Results: sections 3.1.1, 3.1.2 and 3.2 in the original manuscript are combined in the much shorter sect. 3.1 in the new manuscript;

35 Discussion: section 4.1 and 4.2 are combined in the shorter sect. 4.1 of the new manuscript).

Furthermore, we reformulated our objectives. Please see our response to comment 2) of referee 1 for the changes made in the manuscript to reformulate the objectives.

- 40 In this way we focus much more concisely on the objectives of (i) obtaining meaningful estimates of soil moisture at catchment scale as well as its uncertainty and (ii) using this information to analyze the relation between initial soil moisture and the hydrological response. The latter is quantified with the event based runoff coefficient while also considering its uncertainty.
- 45 What I also found surprising is that in the introduction much is made of the importance of flash

floods, but this is not been really picked up and discussed explicitly later with respect to the results. Why?

Thank you for pointing that out. The importance of flash floods is the motivation of our study but we did

5 not study flash floods in this manuscript. That aspect was shortened in the introduction (p.2 l.9 and l.12, l. 19-27).

In the discussion the potential of our results for threshold based flash flood warning systems as proposed by other authors is picked up (p.21 l.1 -14).

- 10 (2) Soil moisture and its spatio-temporal pattern have been subject to a vast body of studies in the past. In that context, neither the introduction nor the discussion of the results here do any justice to these earlier efforts. How is this study placed in the context of this earlier work? What is different? What is novel? What is the same?
- 15 We added the references listed below under "List of additional literature". In the revised version of the manuscript, we stress the novelty of the study (see above). Our studies is different to many studies that address this topic using indirect data or model output. In order to stress this point, we added the sentences "*These approaches [indirect information] offer many advantages, such as the global availability of remote sensing data and the easier acquisition of this data (e.g. Brocca et al., 2009c).*
- 20 Numerous studies found good agreement of soil moisture data obtained from in-situ measurements and remote sensing (e.g. Brocca et al., 2009c, 2013, Huza et al., 2014) Nonetheless, case studies are important to confirm the results obtained with indirect data as well as the results from modelling exercises and to thoroughly understand the hydrologic functioning of local sites." (p.4, l.26) Further, we added term "Relying solely on in-situ data, we [...]" (p.5, l.1) in the last sentence of the introduction that sums up our objectives.

Could your results and interpretation improve some of the earlier interpretations? If so, how? In which aspects are the conclusions you draw similar/not similar to other studies? Why? I would argue that there is quite a lot to discover on this topic and I

30 would strongly encourage the authors to do so to allow the reader to better appreciate the authors' efforts.

In the discussion, we added the following phrase to compare our results concerning the maximum number of samples needed to estimate mean soil moisture to other studies results: "This is consistent

- 35 with the results of Zucco et al. (2014) who found a maximum number of 11 or 20 required samples at the plot scale and catchment scale respectively and the ones of Molina et al. (2014) who concluded that plot mean soil moisture in a Mediterranean mountain area was well represented with 9 probes. The review by Vereecken et al. (2014) shows that there is a wide range of estimates for these numbers and that they are site specific." (p. 17, 1.11)
- 40

Concerning the threshold observed in the relationship between initial soil moisture and event based runoff coefficients we rephrased the original sentence and included the literature you recommended: "Threshold effects in the relation of ϕ_{ev} and $\tilde{\theta}_{ini}$ are also observed by other authors (e.g. McMillan et al., 2014; Hrachowitz et al., 2011). In the Mediterranean context, the thresholds obtained by Huza et al.

(2014) in the Gazel catchment and Braud et al. (2014) in the Valescure catchment (22 and 25 vol% respectively) are lower than the one obtained here. The threshold at 45 vol% observed by Penna et al. (2011) in a 1.9 km² headwater catchment in the Italian Dolomites on the other hand is higher than the one obtained here. McMillan et al. (2014) show that thresholds in different subcatchments of a 50 km²

catchment in New Zealand are highly variable: They range between 27 and 58 vol% and are more or less pronounced in different subcatchments." (p. 20, 1.7)

The conclusions that we draw from our work are similar to the ones of other authors. Our thorough
analysis of the uncertainties inherent in the estimation of mean catchment soil moisture and the event
based runoff coefficient further supports the conclusion that including soil moisture thresholds in
operational flood warning systems is not applicable yet (p.20 l. 1 - 18). We suggest further research to
advance on this problem by adding the phrase *"Further research and instrumentation could include the installation of piezometers in the catchment to understand subsurficial flow in the catchment, using*

10 tracers other than EC to differentiate subsurficial stormflow as a third flow component during hydrograph separation as well as the application of multivariate regression analysis methods that systematically examine different controls on ϕ_{ev} such as meteorological forcing as well as $\tilde{\theta}_{ini}$ and their interactions".

In addition to the other Reviewers' suggestions, I would also think

15 that (at least) the following references are highly relevant and provide necessary context and should thus be considered: McMillan et al. (2014), McMillan and Srinivasan (2015), Hrachowitz et al. (2011) and Li et al. (2011)

Thank you for the additional literature recommendations. They were included in the revised manuscript.

20

3) More information and context is needed for the soil moisture data. It would be nice to have a more detailed map of the locations of the sensors (maybe also crosssections), to get a better idea of what is observed where.

25 Information on topography of the soil moisture measurement sites was added: "Concerning topography, most of the sensors are located on hillslopes which is the dominant topographic zone according to Savenije (2010) in the catchment. Only two plots are located in the riparian area and are potentially connected to the stream during rain events." (p.5, l.1). And "All of these sites [on-alert measurement sites] are located on hillslopes." (p.5, l. 13).

30

Improvement of figure 1 was also demanded by Reviewer 2. We will provide a revised figure with a closer zoom into the area in the south of the Gazel catchment. We will provide information on the land use of the location of the measurement sites. Our experimental plan was not organized with a real upstream-downstream geometry which makes, in our opinion, cross-sections not relevant to this

- 35 particular study. However, Fig. 1 caption will be modified to give precisions about the sensors depths, additionally to the main text so that the vertical profile of the measurements is better conveyed. (Just as a reminder: continuous measurements probes were installed at different depths at one location per plot, the on-alert measurements were conducted at approx. 10 randomly chosen points per field).
- 40 We revised figure 1 and show a closer zoom into the area in the south of the Gazel catchment, a scale bar in the zoom window and by provide information on the land use of the sites of the measurement sites.
- Our experimental plan was not organized with a real upstream-downstream geometry which makes, in our opinion, cross-sections not relevant to this particular study. However, Fig. 1 caption was modified to give precisions about the sensors depths, additionally to the main text so that the vertical profile of the measurements is better conveyed. (Just as a reminder: continuous measurements probes were installed at different depths at one location per plot, the on-alert measurements were conducted at approx. 10 randomly chosen points per field).

Related to that, much is made of the, admittedly quite extensive soil moisture data set. However, in section 4.1 no consideration is given to the limitations of what is actually measured. In a simplified way, the role of unsaturated water storage lies in the temporal storage of water between

- 5 permanent wilting point and field capacity (i.e. water held against gravity). At any point in time, this storage is controlled by plant water use (transpiration) and soil evaporation, whereby plant water use extracts water more efficiently than soil evaporation. Thus, to make sure to meaningfully measure soil moisture, it therefore needs to be measured exactly where plant roots extract water from the soil. Is this the case here? it was
- 10 mentioned that in the vineyards the sensors were place between the vines. Is this were the most important parts of the root system of vines is to be found? I could imagine that the measurements obtained are thus largely biased towards high soil moisture, as plant water extraction may be underrepresented in these locations. How would that change the interpretation? It would be good if the authors not only acknowledged this
- 15 common problem but also discussed the limitations that come with it.

We agree that transpiration is crucial in determining initial soil moisture. The locations of the sensors in the vineyards where carefully chosen to represent a compromise between feasibility and representativeness. The root system of the vines can reach up to 4 m depth, so it was impossible to

- 20 measure soil moisture throughout the root system. By installing the permanent sensors in the vine rows in a depth of up to 50 cm the sensors are as close as possible to the root system. Considering a vineyard plot that consists of the vine rows and the (nearly) bare soil between the rows, we do not believe that we underestimate plot-scale soil moisture due to transpiration.
- The manual measurements in the topsoil were conducted between the rows, because this is where surface runoff started (Visual inspection during heavy rain). We also want to stress here that measuring soil moisture in vineyards with clayey and extremely stony soils is very challenging. To insert the 6 cm rods of the manual soil moisture probes in the stony soil without damaging the instrument, we often had to try more than 10 times before we managed to completely insert the rods.
- 30 We agree that this justification was not conveyed in the original version of the manuscript and rephrased the sentence on the location of the sensors in the vineyard as follows: "*The sensors in the vineyards were installed between two vine plants in a row which is a compromise between feasibility and representativeness of soil moisture in the vineyards which is heterogeneous due to transpiration.*" (p.5, I.5) To better justify the location of on-alert measurements in the vineyards we rephrased the sentence:
- 35 "In the vineyards the measurements were conducted in between the rows of vine plants because this is where surface runoff started (visual inspection)." (p.6, l.17)

(4) Throughout the manuscript, methods could be explained in a clearer and more consistent way

40

We regret that the methods were not explained clearly and consistently. As proposed by Reviewer 1 we inserted a table to explain the different variables calculated in section 2.4 (Table 1 of the new manuscript).

45 [...] and some of the results could be provided in a more quantitative manner. Some examples: it is stated that regression methods require the assumption of "normal distribution of dependent and independent variables". What does that mean? Why should x and y have to be normally distributed? That would only lead to clustering. Do you rather mean that the residuals need to be normally distributed? Please clarify.

Thank you for pointing that out. In order to shorten the manuscript we decided to delete the sentence you are referring to as this sentence is indeed not very clear and not necessary for the paper.

5 Another example section 3.1.1. What do you intend to say with this paragraph? Why is it important to have normally distributed soil moisture?

This paragraph is aimed to describe soil moisture at the plot scale, which is important for the first objective, to obtain a meaningful method for the catchment mean value. The distribution of the values helps interpreting the significance of the mean since an exhaustive sample related to all factors (land use, geology, slope, ...) could not be obtained. Thus, only a statistical approach can be used to assess its uncertainty. As you proposed in your comment (1) we will shortened this section and the following.

Similarly, in section 3.1.2 it

10

30

- 15 is stated that "[:::] pdf [:::] agree with either normal distribution or [:::]". Please use a more formal language here and provide quantification. Do you want to say that the hypothesis that the pdf is a sample from a normal distribution cannot be rejected on a 0.0x significance level? Then please say so.
- 20 Thank you for pointing that out. The respective sentence was also deleted to shorten the manuscript.
 - 4. List of additional literature
- 25 Baffaut, C., Ghidey, F., Sudduth, K. A., Lerch, R. N. and Sadler, E. J.: Long-term suspended sediment transport in the Goodwater Creek Experimental Watershed and Salt River Basin , Missouri , USA, Water Resour. Res., 49, 7827–7830, doi:10.1002/wrcr.20511, 2013.

Beck, H. E., de Jeu, R. A. M., Schellekens, J., van Dijk, A. I. J. M. and Bruijnzeel, L. A.: Improving Curve Number Based Storm Runoff Estimates Using Soil Moisture Proxies, IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens., 2(4), doi:10.1109/JSTARS.2009.2031227, 2009.

Berthet, L., Andréassian, V., Perrin, C. and Javelle, P.: How crucial is it to account for the antecedent moisture conditions in flood forecasting? Comparison of event-based and continuous approaches on 178 catchments, Hydrol. Earth Syst. Sci., 13, 819–831, doi:10.5194/hessd-6-1707-2009, 2009.

Brocca, L., Melone, F., Moramarco, T. and Morbidelli, R.: Antecedent wetness conditions based on ERS scatterometer data, J. Hydrol., 364, 73–87, doi:10.1016/j.jhydrol.2008.10.007, 2009a.

Brocca, L., Melone, F., Moramarco, T. and Singh, V. P.: Assimilation of Observed Soil Moisture Data in Storm Rainfall-Runoff Modeling, J. Hydrol. Eng., 14(2), 153–165, doi:10.1061/(ASCE)1084-0699(2009)14:2(153), 2009b.

Brocca, L., Melone, F., Moramarco, T. and Morbidelli, R.: Soil moisture temporal stability over
experimental areas in Central Italy, Geoderma, 148(3–4), 364–374, doi:10.1016/j.geoderma.2008.11.004, 2009c.

Brocca, L., Melone, F., Moramarco, T., Penna, D., Morga, M., Matgen, A., Gumuzzio, A., Martinez-Fernandez, J. and Wagner, W.: Detecting threshold hydrological response through satellite soil moisture data, Die Bodenkultur, 64(3–4), 7–12, 2013. Bronstert, A. and Bárdossy, A.: The role of spatial variability of soil moisture for modelling surface runoff generation at the small catchment scale, Hydrol. Earth Syst. Sci., 3(4), 505–516, 1999.

Castillo, V. M., Gómez-Plaza, a. and Martínez-Mena, M.: The role of antecedent soil water content in the runoff response of semiarid catchments: A simulation approach, J. Hydrol., 284(1–4), 114–130, doi:10.1016/S0022-1694(03)00264-6, 2003.

Crow, W. T., Bindlish, R. and Jackson, T. J.: The added value of spaceborne passive microwave soil moisture retrievals for forecasting rainfall-runoff partitioning, Geophys. Res. Lett., 32(18), 1–5, doi:10.1029/2005GL023543, 2005.

5

40

Grillakis, M. G., Koutroulis, A. G., Komma, J., Tsanis, I. K., Wagner, W. and Blöschl, G.: Initial soil moisture
effects on flash flood generation – A comparison between basins of contrasting hydro-climatic conditions, J. Hydrol., 541, 206–217, doi:10.1016/j.jhydrol.2016.03.007, 2016.

Hardie, M. A., Cotching, W. E., Doyle, R. B., Holz, G., Lisson, S. and Mattern, K.: Effect of antecedent soil moisture on preferential flow in a texture-contrast soil, J. Hydrol., 398(3–4), 191–201, doi:10.1016/j.jhydrol.2010.12.008, 2011.

15 Hrachowitz, M., Bohte, R., Mul, M. L., Bogaard, T. A., Savenije, H. H. G. and Uhlenbrook, S.: On the value of combined event runoff and tracer analysis to improve understanding of catchment functioning in a data-scarce semi-arid area, Hydrol. Earth Syst. Sci., 15(6), 2007–2024, doi:10.5194/hess-15-2007-2011, 2011.

Huang, M., Gallichand, J., Dong, C., Wang, Z. and Shao, M.: Use of soil moisture data and curve number
method for estimating runoff in the Loess Plateau of China, Hydrol. Process., 21(11), 1471–1481,
doi:10.1002/hyp.6312, 2007.

Javelle, P., Fouchier, C., Arnaud, P. and Lavabre, J.: Flash flood warning at ungauged locations using radar rainfall and antecedent soil moisture estimations, J. Hydrol., 394(1–2), 267–274, doi:10.1016/j.jhydrol.2010.03.032, 2010.

²⁵Jawson, S. D. and Niemann, J. D.: Spatial patterns from EOF analysis of soil moisture at a large scale and their dependence on soil, land-use, and topographic properties, Adv. Water Resour., 30(3), 366–381, doi:10.1016/j.advwatres.2006.05.006, 2007.

Korres, W., Reichenau, T. G., Fiener, P., Koyama, C. N., Bogena, H. R., Cornelissen, T., Baatz, R., Herbst, M., Diekkrüger, B., Vereecken, H. and Schneider, K.: Spatio-temporal soil moisture patterns – A meta-

analysis using plot to catchment scale data, J. Hydrol., 520, 326–341, doi:10.1016/j.jhydrol.2014.11.042, 2015.

Le Lay, M. and Saulnier, G. M.: Exploring the signature of climate and landscape spatial variabilities in flash flood events: Case of the 8-9 September 2002 C??vennes-Vivarais catastrophic event, Geophys. Res. Lett., 34(13), 1–5, doi:10.1029/2007GL029746, 2007.

35 Li, H., Sivapalan, M. and Tian, F.: Comparative diagnostic analysis of runoff generation processes in Oklahoma DMIP2 basins: The Blue River and the Illinois River, J. Hydrol., 418–419, 90–109, doi:10.1016/j.jhydrol.2010.08.005, 2012.

Massari, C., Brocca, L., Moramarco, T., Tramblay, Y. and Didon Lescot, J.-F.: Potential of soil moisture observations in flood modelling: Estimating initial conditions and correcting rainfall, Adv. Water Resour., 74, 44–53, doi:10.1016/j.advwatres.2014.08.004, 2014a.

Massari, C., Brocca, L., Barbetta, S., Papathanasiou, C., Mimikou, M. and Moramarco, T.: Using globally

available soil moisture indicators for flood modelling in Mediterranean catchments, Hydrol. Earth Syst. Sci., 18(2), 839–853, doi:10.5194/hess-18-839-2014, 2014b.

Massari, C., Brocca, L., Ciabatta, L., Moramarco, T., Gabellani, S., Albergel, C., De Rosnay, P., Puca, S. and Wagner, W.: The Use of H-SAF Soil Moisture Products for Operational Hydrology: Flood Modelling over Italy, Hydrology, 2(1), 2–22, doi:10.3390/hydrology2010002, 2015.

McMillan, H., Gueguen, M., Grimon, E., Woods, R., Clark, M. and Rupp, D. E.: Spatial variability of hydrological processes and model structure diagnostics in a 50 km2 catchment, Hydrol. Process., 28(18), 4896–4913, doi:10.1002/hyp.9988, 2014.

 McMillan, H. K. and Srinivasan, M. S.: Characteristics and controls of variability in soil moisture and
 groundwater in a headwater catchment, Hydrol. Earth Syst. Sci., 19(4), 1767–1786, doi:10.5194/hess-19-1767-2015, 2015.

Molina, A. J., Latron, J., Rubio, C. M., Gallart, F. and Llorens, P.: Spatio-temporal variability of soil water content on the local scale in a Mediterranean mountain area (Vallcebre, North Eastern Spain). How different spatio-temporal scales reflect mean soil water content, J. Hydrol., 516, 182–192, doi:10.1016/j.jbudgel.2014.01.040.2014

15 doi:10.1016/j.jhydrol.2014.01.040, 2014.

5

30

Moran, M. S., Holifield Collins, C. D., Goodrich, D. C., Qi, J., Shannon, D. T. and Olsson, A.: Long-term remote sensing database, Walnut Gulch Experimental Watershed, Arizona, United States, Water Resour. Res., 44(5), 1–8, doi:10.1029/2006WR005777, 2008.

Morbidelli, R., Saltalippi, C., Flammini, A., Corradini, C., Brocca, L. and Govindaraju, R. S.: An investigation
 of the effects of spatial heterogeneity of initial soil moisture content on surface runoff simulation at a
 small watershed scale, J. Hydrol., 539, 589–598, doi:10.1016/j.jhydrol.2016.05.067, 2016.

Penna, D., Tromp-van Meerveld, H. J., Gobbi, A., Borga, M. and Dalla Fontana, G.: The influence of soil moisture on threshold runoff generation processes in an alpine headwater catchment, Hydrol. Earth Syst. Sci., 15(3), 689–702, doi:10.5194/hess-15-689-2011, 2011.

25 Renard, K. G., Nichols, M. H., Woolhiser, D. A. and Osborn, H. B.: A brief background on the U.S. Department of Agriculture Agricultural Research Service Walnut Gulch Experimental Watershed, Water Resour. Res., 44(5), 1–11, doi:10.1029/2006WR005691, 2008.

Slaughter, C. W., Marks, D., Flerchinger, G. N., Van Vactor, S. S. and Burgess, M.: Thirty-five years of research data collection at the Reynolds Creek Experimental Watershed, Idaho, United States, Water Resour. Res., 37(11), 2819–2823, doi:10.1029/2001WR000413, 2001.

Tramblay, Y., Bouvier, C., Martin, C., Didon-Lescot, J. F., Todorovik, D. and Domergue, J. M.: Assessment of initial soil moisture conditions for event-based rainfall-runoff modelling, J. Hydrol., 387(3–4), 176–187, doi:10.1016/j.jhydrol.2010.04.006, 2010.

Tramblay, Y., Bouaicha, R., Brocca, L., Dorigo, W., Bouvier, C., Camici, S. and Servat, E.: Estimation of
 antecedent wetness conditions for flood modelling in northern Morocco, Hydrol. Earth Syst. Sci., 16(11),
 4375–4386, doi:10.5194/hess-16-4375-2012, 2012.

Zucco, G., Brocca, L., Moramarco, T. and Morbidelli, R.: Influence of land use on soil moisture spatial–temporal variability and monitoring, J. Hydrol., 516, 193–199, doi:10.1016/j.jhydrol.2014.01.043, 2014.

40 5. Marked-up manuscript version

How does initial soil moisture influence the hydrological response? A case study from southern France

Magdalena Uber^{1,3*}, Jean-Pierre Vandervaere¹, Isabella Zin¹, Isabelle Braud², Maik Heistermann³, Cédric Legoût¹, Gilles Molinié¹, Guillaume Nord¹

- ¹Univ. Grenoble Alpes, CNRS, IRD, Grenoble-INP, IGE Grenoble, 38000, France
 ²Irstea, UR RiverLy, Lyon-Villeurbanne Centre, Villeurbanne, 69625, France
 ³Institute of Earth and Environmental Science, University of Potsdam, Potsdam, 14476, Germany
 *Now at: Univ. Grenoble Alpes, CNRS, IRD, Grenoble-INP, IGE Grenoble, 38000, France
- 10

15

Correspondence to: Magdalena Uber (magdalena.uber@univ-grenoble-alpes.fr)

Abstract. The Cévennes-Vivarais region in southern France is prone to high intensity and long lastingheavy rainfalls that can lead to flash floods which are one of the most hazardous natural risks in Europe. The results of numerous studies show that besides rainfall-depth and intensity and physical catchment characteristics such as topography, geology, land use and hydraulie routing, the catchment's initial soil moisture also impacts the hydrological response to rain events. The aim of this paper is to analyze the relationship between catchment mean initial soil moisture $\tilde{\theta}_{ini}$ and the hydrological response that is quantified using the event-based runoff coefficient ϕ_{ev} in the two nested catchments of the Gazel (3.4 km²) and the Claduègne (43 km²). To this end, two research questions are addressed: (1) How heterogeneous are soil moisture patterns in space and time and do

they correlate with land use? (2) How does soil moisture at the event onset affect the hydrological response? Thus, the

20 <u>objectives are twofold: (1) obtaining meaningful estimates of soil moisture at catchment scale from a dense network of in-situ measurements and (2) using this estimate of $\tilde{\theta}_{ini}$ to analyze its relation with ϕ_{ev} calculated for many runoff events. The estimation of soil moisture at catchment scale is hindered by high spatial and temporal variability. A sampling setup</u>

The estimation of soil moisture at catchment scale is hindered by high spatial and temporal variability. A sampling setup including 45 permanently installed frequency domain reflectancy probes that continuously measure volumetric soil moisture at three depths is applied. Additionally, on-alert surface measurements of soil moisture in the topsoil at \approx 10 locations in each one of 11 plots are conducted. Thus, catchment mean soil moisture can be confidently assessed with a standard error of the

25 one of 11 plots are conducted. Thus, catchment mean soil moisture can be confidently assessed with a standard error of the mean of ≤1.7 vol% over a wide range of soil moisture conditions.

 ϕ_{ev} is calculated from high-resolution discharge and precipitation data for several rain events with a cumulative precipitation P_{cum} ranging from less than 5 mm to more than 80 mm. Because of the high uncertainty of ϕ_{ev} associated to the hydrograph separation method, ϕ_{ev} is calculated with several methods, including graphical methods, digital filters and a tracer based

30 method. The results indicate that the hydrological response depends on $\tilde{\theta}_{ini}$: the seasonal as well as the within event discharge dynamics follow that of soil moisture. Dduring dry conditions ϕ_{ev} is consistently below 0.1-close to zero, even for events with high and intense precipitation. Above a threshold of $\tilde{\theta}_{ini} = 34$ vol% ϕ_{ev} can reach values up to 0.99 but there is a high scatter.

Some variability can be explained with a weak correlation of ϕ_{ev} with P_{cum} and rain intensity, but a considerable part of the variability remains unexplained.

It is concluded that threshold-based methods can be helpful to prevent overestimation of the hydrological response during dry catchment conditions. The impact of soil moisture on the hydrological response during wet catchment conditions, however, is

5 still insufficiently understood and cannot be generalized based on the present results.

1 Introduction

The Cévennes-Vivarais region in southern France is prone to intense rainfall that can lead to the occurrence of flash floods in catchments of various scales ranging from small headwater catchments to ones of several thousand km² (Boudevillain et al., 2011; Braud et al., 2014). Flash floods are sudden floods with high peak discharges of $> 0.5 \text{ m}^3 \text{ s}^{-1} \text{ km}^{-2}$ (Gaume et al., 2009) and a short rise of the hydrograph, i.e. a time to peak of few hours for catchments with a size of up to 100 km² and less than 24 h for catchments of up to 1000 km² (Braud et al., 2014). They are one of the most destructive natural hazards in Europe, both in terms of number of fatalities and economic damage (Gaume et al., 2009). Unlike lowland floods they often result in losses of life, s Striking examples arebeing the October 2015 flash flood of the Brague river that hit the French Riviera and the 2002 flash flood of the Gard river with 23 deaths and an estimated direct tangible damage of 1.2 billion Euro (Huet et al., 2002 flash flood of the Gard river with 23 deaths and an estimated direct tangible damage of 1.2 billion Euro (Huet et al., 2002 flash flood of the Gard river with 23 deaths and an estimated direct tangible damage of 1.2 billion Euro (Huet et al., 2002 flash flood of the Gard river with 23 deaths and an estimated direct tangible damage of 1.2 billion Euro (Huet et al., 2002 flash flood of the Gard river with 23 deaths and an estimated direct tangible damage of 1.2 billion Euro (Huet et al., 2002 flash flood of the Gard river with 23 deaths and an estimated direct tangible damage of 1.2 billion Euro (Huet et al., 2002 flash flood of the Gard river with 23 deaths and an estimated direct tangible damage of 1.2 billion Euro (Huet et al., 2002 flash flood of the Gard river with 23 deaths and an estimated direct tangible damage of 1.2 billion Euro (Huet et al., 2002 flash flood of the Gard river with 23 deaths and an estimated direct tangible damage of 1.2 billion Euro (Huet et al., 2002 flash flood of t

15 2003).

10

Despite the recognition of their high damage potential, the hydrological processes leading to the generation of flash floods are still insufficiently understood at a scale that is important for prediction and management (Gaume et al., 2009; Braud et al., 2014). Three <u>One of the main problems that are recognized to hinder flash flood prediction is the ignorance of the (Creutin and Borga, 2003): the change of scale problem (Blöschl and Sivapalan, 1995) which is especially relevant for capturing the</u>

- 20 highly heterogenous rainfall fields causing flash floods (Borga, 2002; Creutin and Borga, 2003; Delrieu et al., 2014); the predictions in ungauged basins (PUB) problem (Sivapalan, 2003) and the problem of knowing the soil water retention capacity (Creutin and Borga, 2003). Soil moisture is known to govern overland flow generation (Zehe and Sivapalan, 2008). As it controls threshold behavior, it implies qualitative changes of hydrological processes and the hydrologic system's response to rain events (Zehe and Sivapalan, 2008). Because they are spatially and temporally distinct events, flash floods are difficult to
- 25 capture with the operational hydro meteorological measuring systems that are not dense enough to document discrete, rapidly occurring flood events at small scales. Thus, they remain a poorly documented phenomenon (Creutin and Borga, 2003; Borga et al., 2008; Gaume et al., 2009; Braud et al., 2014). Other issues concern Tthe lack of high-resolution data measured during flash flood events as well as the variety of catchment characteristics that influence their occurrence. of flash floods Tand the high degree of non-linearity in the hydrological response of catchments hinders the predictability of flash floods (Braud et al., 2014).
- 30 2014). This has motivated the installation of several measurement networks in first-order catchments especially in the USA and Australia (Slaughter et al., 2001; Renard et al., 2008; Moran et al., 2008; Baffaut et al., 2013) and at the mesoscale and in a Mediterranean context the FloodScale project in the Cévennes-Vivarais region (Braud et al., 2014, Nord et al., 2017).

Flash floods are usually associated with intense rainfall of > 100 mm in a few hours or long lasting rainfall (\approx 24 h) with moderate intensities (Braud et al., 2014) often generated by mesoscale convective systems and / or orographic precipitation (Marchi et al., 2010; Molinié et al., 2012; Panziera et al., 2015). However, the hydrological response to rain events varies greatly between catchments and between events. It can be quantified using the event-based runoff coefficient ϕ_{ev} , i.e. the ratio

- 5 of event runoff volume to total event rainfall volume. The major drawback of this quantity is the lack of standard procedures for obtaining event runoff volumes and for defining the beginning and end of an event, which impedes comparisons between studies (Blume et al., 2007). Yet, event-based runoff coefficients of flash-flood events have been found to differ substantially, spanning nearly the full range of values from zero to one, with a high positive skewness in their frequency distribution (Merz et al., 2006; Blume et al., 2007; Norbiato et al., 2008; Merz and Blöschl, 2009; Marchi et al., 2010). They were shown to differ
- 10 considerably between regions (Marchi et al., 2010), seasons (Li et al., 2012) and flood types (Merz et al., 2006), to increase with mean annual precipitation and event rainfall depth (Merz et al., 2006; Norbiato et al., 2009) and to depend on rain intensity, soil types and antecedent soil moisture conditions (Wood et al., 1990; Crow et al., 2005; Marchi et al., 2010; Hrachowitz et al., 2011; Penna et al., 2011; Li et al., 2012; Huza et al., 2014). Furthermore, a multitude of catchment characteristics also determine runoff generation and concentration, namely topography, geology, hydraulic routing and geomorphological controls
- 15 (Braud et al., 2014).

Soil moisture is known to govern overland flow generation (Zehe and Sivapalan, 2008). As it controls threshold behavior, it implies qualitative changes of hydrological processes and the hydrologic system's response to rain events (Zehe and Sivapalan, 2008; Hardie et al., 2011).

- 20 Initial soil moisture θ_{ini} , i.e. the soil water content at the onset of a rain event, is a crucial factor that influences the water storage capacity of the catchment as well as soil hydraulic properties and thus the hydrological response to rainfall events. It controls the soil moisture deficit, consequently, in the interplay with rainfall forcing, it determines whether soil saturation and saturation excess overland flow (Dunne and Black, 1970) occur during a rain event or not. Moreover, soil moisture controls the unsaturated hydraulic conductivity and thus the occurrence of infiltration excess overland flow (Horton, 1933). Given the high
- 25 spatial and temporal variability of soil moisture and the incoherence of scale of the measurements with the catchment size, it remains challenging to obtain meaningful estimates of θ_{ini} at the catchment scale (Brocca et al., 2009ae; Vereecken et al., 2014; Korres et al., 2015). There are multiple controls on soil moisture such as soil texture, topography and vegetation as well as small scale random variability (Jawson and Niemann, 2007; Garcia-Estringana et al., 2013; McMillan and Srinivasan, 2015). This problem is addressed by Vachaud et al. (1985) by introducing the concept of temporal stability, based on the finding that
- 30 deviations of point measurements from the catchment mean can be persistent in time. Thus, optimum sampling locations can be identified and the number of samples required can be reduced (e.g. Brocca et al., 2009ae, Huza et al., 2014). Several studies consider the impact of initial soil moisture on the hydrological response of catchments on heavy rain events. Seasonality in the occurrence of flash floods (Gaume et al., 2009) and discharge magnitudes (Borga et al., 2007, Li et al., 2012) have been attributed to initial soil moisture conditions. Numerous modelling studies have shown the high sensitivity of the

modelled runoff response to θ_{ini} and the importance of estimates of θ_{ini} at the catchment scale as initial conditions in eventbased models (e.g. Castillo et al., 2003; Huang et al., 2007; LeLay and Saulnier, 2007; Berthet et al., 2009; Brocca et al., 2009b; Tramblay et al., 2010, 2012; Li et al., 2012; Massari et al., 2014a,b, 2015; Grillakis et al., 2016). The dependence of catchment responses to initial soil moisture is also observed by Marchi et al. (2010) in a dataset comprising data for 25 flood

- 5 events in 60 basins across Europe and on this study's site by Huza et al. (2014). This relationship is characterized by high non-linearity and threshold effects (Zehe et al., 2005; Huza et al., 2014). There is no consent on the importance of initial soil moisture during extreme events. Wood et al. (1990) conclude that catchment characteristics are important only for flood events with a low return period (up to ca. 10 y) whereas rainfall characteristics dominate those with a higher return period. On the other hand this finding is rejected in analyses of historic flash floods (Gaume et al., 2004; Borga et al., 2007; Le Lay and Saulnier, 2007) -or flash flood data bases (Marchi et al., 2010) whose authors conclude that soil moisture plays an important
- 10 <u>Saulnier, 2007</u>) -or flash flood data bases (Marchi et al., 2010) w role in the hydrological response, also under extreme conditions.
 - The aim of this study is to assess how soil moisture controls the hydrological response in a flash-flood prone area in southern France. The study is conducted in the two nested catchments of the Claduègne (43 km²) and Gazel (3.4 km²), Ardèche, France. Thanks to an exceptionally good data baseis, it is possible to obtain reliable estimates of the two catchments' initial soil
- 15 moisture states for several rain events and to quantify the hydrological response with the event-based runoff coefficient. To this end, the spatio-temporal heterogeneity is assessed to obtain reliable estimates for mean initial soil moisture at the catchment scale as well as its uncertainty. It is examined whether soil moisture correlates with land use as this finding could improve the interpolation of point measurements to catchment means. Land use was chosen for this analysis as it controls soil moisture and its variability via interception and evapotranspiration (Reynolds, 1970; Grayson et al., 1997; Garcia Estringana et al., 2013)
- 20 and has a crucial influence on soil hydraulic properties, in particular soil hydraulic conductivity (Gonzales Soza et al., 2010; Jarvis et al., 2013). Moreover, it is easy to assess in the field or from remote sensing data. In the study site land use correlates well with soil texture in that the vineyards are found on finer textured soils than the other land use classes. Other studies results that suggest a dependence of ϕ_{ev} on initial soil moisture (e.g. Merz et al., 2006; Blume et al., 2007; Merz

and Blöschl, 2009; Norbiato et al., 2009). However, most of these studies use indirect information such as remote sensing data,

- 25 antecedent precipitation indices, initial baseflow, continuous soil moisture accounting models, or the ratio of actual evaporation to precipitation or remote sensing data. These approaches offer many advantages, such as the global availability of remote sensing data and the easier acquisition of this data (e.g. Brocca et al., 2009c). Numerous studies found good agreement of soil moisture data obtained from in-situ measurements and remote sensing (e.g. Brocca et al., 2009c, 2013, Huza et al., 2014) Nonetheless, case studies are important to confirm the results obtained with indirect data as well as the results from modelling
- 30 exercises and to thoroughly understand the hydrologic functioning of local sites. At this studies site the impact of θ_{ini} on ϕ_{ev} was already considered by Huza et al. (2014). However, these authors used soil moisture data obtained from ASCAT satellite data which is fitted to in-situ measurements of topsoil moisture that were conducted on grasslands only. They considered five rain events only, so this relation could not be quantified unambigously. Thus, this study's <u>novelty objective</u> is to analyze the relation between ϕ_{ev} and θ_{ini} when both are obtained from a comprehensive, high resolution data set allowing the assessment

of the uncertainty of the two variables. Relying solely on in-situ data, wWe aim to (i) obtain a meaningful estimate of catchment scale soil moisture and its uncertainty and (ii) answer the work research question_s: (1) How heterogeneous are soil moisture patterns in space and time and do they correlate with land use? And (2) how does soil moisture at the event onset affect the hydrological response?

5 2 Methods

2.1 Study site

For this study two nested sub-catchments of the Ardèche river in the Cévennes-Vivarais region of southern France are considered: the catchments of the intermittent Gazel stream and the perennial Claduègne river with an area of 3.4 and 43 km² respectively (Fig. 1).

- 10 Both catchments can be clearly divided into two distinct geologies: the northern part is constituted by the Coiron basaltic plateau that is bounded by a steep cliff of basaltic columns in the south, whereas the southern part of both catchments is a landscape of piedmont hills underlain by sedimentary limestone lithology (Nord et al., 2017). The basaltic plateau covers 51 % of the Claduègne catchment whereas its fraction of the Gazel catchment is only 23 %. Thus, the northern part is dominated by silty and stony soils on pebble deposit of basaltic component, while the soils in the southern part are predominantly
- 15 rendzinas or other clay-stony soils, cultivated soils of loam and clay-loam and in the south of the Claduègne catchment lithosols and regosols (Nord et al., 2017). The terrain is hilly, ranging from about 820 m above sea level (asl) to 230 m asl at the outlet (650–260 m asl for the Gazel catchment) havingand has a mean slope of about 20 %. The area is characterized by extensive agriculture and natural vegetation. Hence, the main land use / land cover types are grasslands, pastures, vineyards, forests and Mediterranean open woodlands. The vineyards are predominantly found on the finer textured soils in the southern part of the
- 20 Claduègne catchment while the other land use types are found throughout the catchments. The average annual precipitation at Le Pradel at the outlet of the Gazel catchment is 1030 mm (Huza et al. (2014), original data: daily rain gauge data for 1958– 2000 from Méteo-France). For further details see Nord et al. (2017).

2.2 Data availability

As part of the HyMex (Hydrological Cycle in the Mediterranean Experiment, Ducrocq et al. (2014)) and FloodScale (Braud et al., 2014) projects and the Cévennes-Vivarais Mediterranean Hydrometeorological Observatory (OHM-CV, Boudevillain et al. (2011)), the area is exceptionally well monitored, thus high-resolution spatio-temporal data on rainfall, discharge and soil moisture is available. The data used for this study was published in Nord et al. (2017) and the link to download the data can be found at the publishers website: https://www.earth-syst-sci-data.net/9/221/2017/essd-9-221-2017-assets.html.

Soil moisture θ: two different sets of soil moisture data are available: continuous measurements and on-alert measurements.
Soil moisture is continuously measured with 45 fixed soil moisture probes at nine plots (two vineyards, one fallow, six grasslands) within the Claduègne catchment since June 2013. Six of the plots are located in the piedmont hills and three on the

basaltic plateau (Fig. 1). Concerning topography, most of the sensors are located on hillslopes which is the dominant topographic zone according to Savenije (2010) in the catchment. Only two plots are located in the riparian area and are potentially connected to the stream during rain events. At each plot, five frequency domain reflectometry (FDR) probes (Decagon 10HS soil moisture sensors) are installed at different depths: 10 cm (n = 2), 20–25 cm (n = 2) and in the subsoil (33–

- 5 50 cm, n = 1). The temporal resolution is 15–20 min (Nord et al., 2017). The sensors in the vineyards were installed between two vine plants in a row which is a compromise between feasibility and representativeness of soil moisture in the vineyards which is heterogeneous due to transpiration. The accuracy and the range of the probe as provided by the manufacturer are ±3 vol% and 0–57 vol%. The data is available from June 20<u>13</u>06 November 2014 in the dataset doi: 10.17178/OHMCV.SMO.CLA.13-14.1 presented in Nord et al. (2017).
- In addition, following forecasts of heavy rain events on-alert measurements of soil moisture in 0–5 cm depth were conducted at 11 plots within the Gazel catchment with a hand-held FDR soil moisture sensor (Delta-T SM200). The accuracy and the range of the probe are ±3 vol% and 0–50 vol%. The plots comprised four vineyards, five grasslands, one fallow and one cultivated field. <u>All of these sites are located on hillslopes.</u> The sampling sites were selected for reasons of accessibility, congruence with other measurements conducted during the FloodScale project and representativeness of the catchments'
- 15 landscapes. All on-alert measurements were conducted in about 1 h at ≈ 10 randomly chosen measurement points within each plot. The distance between the measurement points was at least 1 m to ensure spatial independence (Huza et al., 2014). In the vineyards the measurements were conducted in between the rows of vine plants because this is where surface runoff started (visual inspection). On-alert measurements were done before and after 11 heavy rain events during the special observation periods of the HyMex Project in autumn (September–December) of the years-period 2012–2015. The dataset is found in the supplementary metarial of this article (\$1).
- 20 supplementary material of this article (S1).

Precipitation *P*: rainfall data was obtained from the HPiconet rain gauge network at a resolution of 1 min. The network consists of 19 tipping bucket rainfall gauges with a sampling surface of 1000 cm² and a resolution of 0.2 mm out of which 12 are located in the Claduègne catchment or its close vicinity (Nord et al. (2017), DOI: 10.17178/OHMCV.RTS.AUZ.10-14.1 Fig. 1).

- 25 **Discharge** *Q*: water level is continuously measured at the outlets of the two catchments with water level gauges at 2 min resolution (Gazel) <u>or-and_10</u> min resolution (Claduègne) respectively (Nord et al., 2017). The water level is converted to discharge with a stage-discharge relationship established using the BaRatin framework (Le-Coz et al., 2014) that also gives the uncertainty of the rating curve that is quantified as the 90 % confidence interval of discharge. The rating curve is based on numerous discharge measurements <u>performed</u> in 2012–2014 (Nord et al., 2017).
- 30 Additional data: spatial data used for this study include a digital elevation model with a resolution of 5 m (DOI: 10.6096/MISTRALS-HyMeX.1389) and the Ardèche soil data base by the French National Institute for Agricultural Research, Bureau de Recherches Géologiques et Minières and the French Department of Agriculture (DOI: 10.6096/MISTRALS-HyMeX.1385, Nord et al., 2017). Furthermore, a 0.5 m resolution land use map of the Claduègne catchment based on quickbird satellite images is available (DOI: 10.14768/mistrals-hymex.1381, Andrieu, 2015). Data on soil properties such as porosity,

texture and saturated hydraulic conductivity was obtained during a measurement campaign in 2012 by Braud and Vandervaere (2015; DOI: 10.6096/MISTRALS-HyMeX.1321). Electrical conductivity (*EC*) of stream flow is continuously measured at the outlets of both catchments and measurements of *EC* of overland flow from two runoff and erosion plots in a vineyard in the south of the Gazel catchment are available (Cea et al., 2015).

5 2.3 Precipitation data processing

The catchment mean hyetographs for both catchments are calculated from the HPiconet rain gauge data with the method of Thiessen polygons. Rain events are separated by using a threshold of 12 h without precipitation being recorded at any rain gauge. The onset of an event was defined as the first time rain occurred after a dry period of at least 12 h, the end as the last time with rain being recorded by at least one rain gauge before the next dry period. The threshold of 12 h provides a good compromise between having a high number of events and excluding two separate events that are not independent from each other. Averaged catchment rainfall is then summed over the whole period of the rain event to calculate cumulative event precipitation P_{cum} . Furthermore, mean rain intensity I_{μ} over the whole event as well as maximum rain intensity I_{max} at 2, 10,

20, 30 and 60 minutes are calculated using the averaged catchment rainfall.

2.4 Soil moisture analysis

15 From both data sets (continuous and on-alert measurements) plot and catchment mean values are calculated for the initial and final state of each rain event. From the continuous data, mean values are calculated for all three depths and the profile mean value is calculated.

 $\theta(x_{i,j}, t_{ev})$ refers to a spatially and temporally discrete on-alert soil moisture measurement, with the subscript i denoting the index of the n_i (usually ten) measurements within the plot, j denoting the plot and ev the event and the state (initial or final).

20 Mean soil moisture was calculated at the plot scale $(\overline{\theta}_j(t_{ev}))$, for each land use class $(\overline{\overline{\theta}}_{lu}(t_{ev}))$ and at the catchment scale

$\underline{\overline{\theta}}_{ev}$). See <u>T</u>table <u>XX1</u> for the formulas.

Plot mean soil moisture $\overline{\theta}_{j}(t_{ev})$ of the n_{i} local measurements were calculated for all plots as well as land use class means $\overline{\overline{\theta}}_{tu}(t_{ev})$ of $n_{j,tw} n_{jtw}$ plots belonging to the same of the four land use classes grassland, vineyard, fallow and cultivated field (c_{tw}) = {g,v,f,c}):

25

10

$$-\overline{\theta}_{j}(t_{\text{ev}}) = \frac{1}{n_{\text{f}}} \sum_{i=1}^{n_{\text{f}}} \theta\left(x_{i,j}, t_{\text{ev}}\right) \tag{1}$$

$$= \frac{1}{\overline{\theta}_{\mathrm{lu}}(t_{\mathrm{ev}})} = \frac{1}{n_{j_{\mathrm{tur}}}} \sum_{j_{\mathrm{lur}}=1}^{n_{j_{\mathrm{tur}}}} \overline{\theta}_{j}(t_{\mathrm{ev}}); \qquad j_{\mathrm{lur}} \in \mathrm{lu}; \mathrm{lu} \in c_{\mathrm{lur}}$$
(2)

with n_{thr} = number of plots in the respective land use class.

Catchment mean values $\overline{\overline{\theta}}_{ev}$ are computed as a mean of the different land use classes weighted with the number of measurements per land use class:

$$\overline{\overline{\theta}}_{ev} = \frac{1}{n_{\overline{p}}} \sum_{u}^{n_{\overline{e}_{\overline{h}\overline{u}}}} \overline{\overline{\theta}}_{u} \cdot n_{\overline{h}u}, \quad u \in c_{u}$$
(3)

with $n_{c_{tw}}$ = number of land use classes $(n_{c_{tw}} = 4)$ and n_p = total number of plots $(n_p = \sum n_{j_{tw}} = 11)$.

15

5 Plot means and catchment means obtained from for the continuously measured data are computed in the same way for all three

layers $I_{\underline{\cdot}}$ (therefore denoted $\overline{\theta}_{ev,1}$) with the exception, that <u>Here</u>, the plot mean is obtained by averaging not only the probes installed at the same depth and the same location, but also <u>several-all</u> measurements in a dry period of two hours before the onset or after the end of the rain event in order to diminish noise. <u>The catchment mean averaged over the three layers $\tilde{\theta}_{ev}$, i.e. considering the topmost 60 cm, is calculated from the continuously measured data (Eq. 4, Table XX1).</u>

10 The profile mean $\tilde{\theta}_{ev}$ over the three layers (n_{\perp} = number of layers) is calculated with the following equation. The thicknesses m_{\perp} of the layers are assumed to be 175, 150 and 275 mm respectively, hence, the topmost 60 cm are considered:

$$\tilde{\theta}_{ev} = \frac{\sum_{l=1}^{n_{t}} \frac{\Xi}{\theta_{ev,l}} \cdot m_{l}}{\sum_{l=1}^{n_{t}} m_{l}} \tag{4}$$

Finally, for all events the soil moisture storage change ΔS [mm/ev] in the upper 60 cm is computed from the difference between =initial $\frac{\Xi}{(\theta_{\text{ini},l} \text{[vol\%]})}$ and final soil moisture (Eq. 5, Table XX1). $\frac{\Xi}{(\theta_{\text{fin},l} \text{[vol\%]})}$ that is converted to [vol/vol] via division by 100:

$$\Delta S = \sum_{l=1}^{n_{\rm f}} \frac{1}{100} \left(\overline{\overline{\overline{\theta}}}_{\rm fin,l} - \overline{\overline{\overline{\theta}}}_{\rm ini,l} \right) m_{\rm f} \tag{5}$$

For all plots and all events, the frequency distributions of the on alert soil moisture measurements are tested for normality with a Shapiro Wilk test at a significance level of $\alpha = 0.05$. It is also assessed whether significant differences between the four land use classes exist by performing a visual inspection of boxplots or histograms and Student t-tests. Moreover, standard deviations σ as measures of spatial variability at different scales are calculated at the plot scale (σ_j^{inner} , Eq. XX6, Table XX1) and at the catchment scale ($\sigma_{cat.}^{inter}$, Eq. XX7, Table XX1). Furthermore, σ is calculated between plots of the same land use ($\sigma_{lu}^{inter}(t_{ev})$, Eq. XX8, Table XX1) and between land use classes ($\sigma^{betw}(t_{ev})$, Eq. XX9, Table 1): the inner plot standard deviation σ_j^{inner} and the inter plot standard deviation for the land use classes grassland and vineyard (σ_{lu}^{inter} ; $n_{j\in lu} > 1$) and the whole catchment ($\sigma_{eat.}^{inter}$) are calculated for each event:

$$\frac{1}{\sigma_{j}^{\text{inner}}(t_{ev}) - \sqrt{\frac{1}{n_{i} - 1} \sum_{i=1}^{n_{i}} (\theta(x_{i,j}, t_{ev}) - \overline{\theta}_{j}(t_{ev}))^{2}}} \qquad (6)$$

$$\frac{\sigma_{iu}^{\text{inter}}(t_{ev}) - \sqrt{\frac{1}{n_{ju} - 1} \sum_{ju=1}^{n_{ju}} (\overline{\theta}_{j}(t_{ev}) - \overline{\theta}_{iu}(t_{ev}))^{2}}} \qquad j_{iu} \in lu; lu \in c_{iu} \qquad (7)$$

$$\frac{\sigma_{eut}^{\text{inter}}(t_{ev}) - \sqrt{\frac{1}{n_{jv} - 1} \sum_{j=1}^{n_{jv}} (\overline{\theta}_{j}(t_{ev}) - \overline{\theta}(t_{ev}))^{2}}} \qquad (8)$$

Furthermore, the between-land use standard deviation σ^{betw} is computed:

10

5
$$-\frac{1}{\sqrt{n_{e_{tw}}}} = \sqrt{\frac{1}{\frac{1}{n_{e_{tw}}}} - 1} \sum_{lw}^{n_{e_{tw}}} (\overline{\overline{\theta}}_{lw}(t_{ev}) - \overline{\overline{\theta}}_{ev})^2}$$
(9)

As an estimate of the uncertainty of the calculated plot and catchment mean values, the standard error of the plot_-mean SEM_{j}^{inner} and the one of the catchment mean $SEM_{cat.}^{inter}$ are calculated with Eq. (10) and Eq. (11). It should be noted that $SEM_{cat.}^{inter}$ the latter is calculated from the on-alert measurements in the topsoil as well as from the continuous measurements over the soil profile, SEM_{j}^{inner} the former_-only from the on-alert measurements. The SEM is used as a measure of the confidence that the sample mean corresponds to the universal mean; it increases with the standard deviation and decreases with the number of sampling points.

 $\frac{SEM_{j}^{\text{inner}}(t_{\text{ev}})}{SEM_{j}^{\text{inter}}(t_{\text{ev}})} - \frac{\sigma_{j}^{\text{betw}}(t_{\text{ev}})}{\sqrt{n_{j}}}$ (10)

Moreover, it is assessed whether temporal stability, i.e. consistency of soil moisture patterns at the catchment scale at different times of measuring (Vachaud et al., 1985), as reported by Huza et al. (2014) for six grassland plots in the Gazel catchment, is also found in the present on-alert data set: the relative spatial difference $\delta_{j,ev}$ of each plot corresponds to the relative difference between the plot mean and the catchment mean (Eq. 12); its temporal mean $\overline{\delta_j}$ is calculated with Eq. (213):

$$\delta_{j,ev} = \frac{\overline{\theta}_j(t_{ev}) - \overline{\overline{\theta}}_{ev}}{\overline{\overline{\theta}}_{ev}}$$
(12)

$$\overline{\delta_{j}} = \frac{1}{n_{ev}} \sum_{ev=1}^{n_{ev}} \delta_{j,ev}$$
(213)

The plot with the smallest $\delta_{j,ev}$ is the one that agrees best with the catchment mean on a given time of measurement. The temporal variability of the spatial difference σ_{δ_i} serves as an auxillary variable to assess whether this behavior is stable in time:

$$\sigma_{\delta_j} = \sqrt{\frac{1}{n_{\rm ev} - 1} \sum_{\rm ev=1}^{n_{\rm ev}} (\delta_{j,\rm ev} - \overline{\delta_j})^2}$$
(314)

5 The temporal dynamics of soil moisture during the autumn season and during single events that are captured with the continuous soil moisture measurements are analyzed and compared with the hyetographs and hydrographs. Furthermore, crosscorrelations are calculated for all continuously measuring sensors and all depths with the R function ccf (Gilbert and Plummer, 2016). For each plot the maxima of the crosscorrelation functions *L*_{CCmax} between rainfall and the sensors in 10 cm depth, between the ones in 10 and 25 cm and the ones in 25 and 40 cm are calculated.

10 **2.5 Hydrological response**

2.5.1 Event based runoff coefficients

In order to quantify the hydrological response of the catchment to different rainfall events, the dimensionless event-based runoff coefficient ϕ_{ev} is calculated for all events:

$$\phi_{\rm ev} = \frac{Q_{\rm ev,cum}}{P_{\rm cum}} \tag{154}$$

- 15 P_{eum} is calculated for each event as described in Sect. 2.3. To obtain cumulative event discharge $Q_{ev,cum}$, the time series of stream discharge Q_{tot} has to be separated into baseflow Q_b and event flow Q_{ev} . Q_{ev} is defined here to be the fast responding part of discharge that occurs during or directly after the rain event. It usually encompasses surface runoff or overland flow and fast subsurface flow. Q_b on the other hand is the slow responding part of discharge that lasts long after the rain event and feeds the stream between rain events. To obtain $Q_{ev,cum}$, Q_{ev} is summed up over the whole period of the event. The onset of a discharge
- 20 event is defined as the first increase of discharge in response to a rain event. Defining the end of event discharge is more complicated and depends on the hydrograph separation method (Blume et al., 2007). Usually, the end of event flow is defined as the moment when Q_b equals Q_{tot} , but for some events the onset of a second event impedes this procedure which causes errors. Taking into consideration that there is no standard method for hydrograph separation and that results obtained with different methods can differ substantially (Blume et al., 2007), seven different hydrograph separation techniques that are
- 25 described in the following section are applied and compared (Sect. 2.5.2) (Fig. 2). The uncertainty of Q_{tot} associated with the

stage-discharge relation can be important especially for high-flow conditions. This was taken into account by calculating ϕ_{ev} with the upper and the lower limit of the 90 % confidence interval of discharge obtained with the BaRatin framework. Because several of the assumptions underlying standard regression analysis methods (normal distribution of dependent and

independent variables, only the dependent variable is subject to error, homoscedasticity) are not met, it is not attempted to set-

5 up linear or non linear regression models that predict ϕ_{ev} from possible explanatory variables such as initial soil moisture, initial baseflow, rainfall depth or rain intensity. The relation of these variables and ϕ_{ev} is considered nonetheless.

2.5.2 Hydrograph separation

Straight line method. The straight line method (SL) is a simple, graphical method where baseflow during the event is interpolated by connecting the point in the event hydrograph at which discharge first increases as a response to the rain event

10 with the first point on the falling limb of the hydrograph with the same discharge value. As this condition is often never met, the end of event flow is often determined by the onset of the next event or discharge below a threshold.

Constant-k method. The CK-method proposed by Blume et al. (2007) is based on the assumption that baseflow recession behaves similar to the outflow of a linear storage. Thus, baseflow at time step *t* can be described as exponential recession with the recession parameter *k* and initial flow Q_0 :

$$Q_{\rm b}(t) = Q_0 \,{\rm e}^{-k\,t} \tag{165}$$

k is calculated at each time step by differentiating eq. $\frac{16-5}{2}$ and division by $Q_{b}(t)$:

$$k = -\frac{\mathrm{d}Q}{\mathrm{d}t} \frac{1}{Q_{\mathrm{b}}(t)} \tag{176}$$

Event flow is assumed to terminate at time step t_e which is defined as the end of event runoff, once k becomes approximately constant. Baseflow is assumed to be equal to the discharge before the onset of event flow up to t_e when it equals Q_{tot} .

20 Electrical conductivity method. Hydrograph separation is also conducted based on electrical conductivity (*EC*) which serves as a natural tracer (Miller et al., 2014; Pellerin et al., 2008). The method relies on the assumption that stream flow Q_{tot} with electrical conductivity EC_{tot} is a mixture of subsurface flow Q_{sb} and surface flow Q_s , which have significantly different *EC* signals EC_{sb} and EC_s (Nakamura, 1971):

$$Q_{\rm tot} = Q_{\rm sb} + Q_{\rm s} \tag{187}$$

$$Q_{\rm tot} \cdot EC_{\rm tot} = Q_{\rm sb} \cdot EC_{\rm sb} + Q_{\rm s} \cdot EC_{\rm s} \tag{198}$$

25

30

Thus, with given values for
$$EC_{sb}$$
 (interpolated EC between values before the onset and after the end of event discharge) and

*EC*_s (measured in overland flow collected at the outlet of four erosion plots representative of the signature of the rainfall flowing at the surface of soils developed on sedimentary rocks), a time series of Q_{sb} can be calculated. Furthermore, the maximum signal of *EC* is calculated for each event as $\Delta EC_{max} = max(|EC_{sb} - EC_{tot}|)$. As no *EC*_s values were available for overland flow occurring on soils developed on basalts, covering half of the Claduègne catchment, the method could only be applied to the Gazel catchment, where the proportion of basaltic geology to total catchment surface is much smaller.

It should be noted that this method considers only the surficial part of event flow and is not able to separate the fast responding subsurficial flow occurring in the unsaturated zone. Thus, event flow is likely to be underestimated, especially in conditions when the latter plays an important role.

Recursive digital filter. The RDF method proposed by Lyne and Hollick (1979) uses a signal analysis low-pass filter to 5 separate high-frequency event flow signals from low frequency baseflow signals:

$$f(t) = a f(t-1) + \frac{1+a}{2} (Q_{\text{tot}}(t) - Q_{\text{tot}}(t-1))$$
(199)

where f(t) is filtered event flow at time t, a is a filter parameter that is usually in the range of 0.00 < a < 0.95 (Nathan and McMahon, 1990) and $Q_{tot}(t)$ is original stream flow at time t. The data is passed to the filter several times, forwards and backwards. Recommendations for the number of passes vary depending on the time increment resolution of the discharge series (Ladson et al., 2013). The method is implemented in the R function BaseflowSeparation of the package EcoHydRology (Archibald, 2014).

Hysep filters (HS1 – HS3). Three further filtering approaches are implemented in the Unites Stated Geological Survey's (USGS) HySep program (Sloto and Crouse, 1996). It is based on finding minima in the discharge time series. The minima are determined either within fixed (HS1) or sliding (HS2) intervals or with a local minima algorithm (HS3). The interval width is

15 adjusted according to Gonzales et al. (2009). It is applied using the R code of the USGS (2015).

3 Results

10

3.1 Spatial and temporal variability of soil moisture

3.1.1 Soil moisture at the plot scale

The probability density function (pdf) of the on alert measurements of soil moisture of the topsoil measured within one plot at 20 a discrete moment in time $\theta(x_{i,i},t_{vv})$ can usually be described with a normal distribution (230 out of 243 within plot measurements; Shapiro Wilk test with $\alpha = 0.05$, Fig. 3a). In the present data set, normal distribution is found during dry, medium and wet conditions. Out of the 23 measurements conducted during dry conditions ($\overline{\theta}_{t}(t_{ev}) < 19$ vol%, the 10 % quantile) and the 23 measurements during wet conditions ($\overline{\theta}_{i}(t_{ev}) > 37$ vol%, the 90 % quantile) only one and zero respectively data sets are not normally distributed.

- 25 The variability of soil moisture at the plot scale, determined from the on-alert measurements in the topsoil, is very high: the median range between the highest and the lowest measurement in one plot is 7.8 vol% but maximum values can get up to > 30 vol%. The mean of the inner-plot standard deviation $\sigma_i^{\text{inner}}(t_{\text{ev}})$ is 2.7 vol%. Values range from 1–8 vol% with no significant difference between the land-use classes. There is no significant correlation between plot means and standard deviations (Fig. 2a3e). The inner-plot standard deviation in the deeper layers, determined with the continuously measuring 30
- probes, cannot be confidently assessed because of the low number of probes installed in each plot at the same depth. However,

the difference of two sensors installed at the same depth indicate, that the variability is in the order of the one derived from the on-alert measurements (Table 24).

The mean $SEM_j^{\text{inner}}(t_{\text{ev}})$ is 0.8 vol% with only three out of 228 data sets exceeding 2.0 vol%. Thus, the confidence that the population plot mean lies within the sample mean $\overline{\theta}_j(t_{\text{ev}}) \pm 2.0$ vol% is very high. This accuracy was achieved with about ten measurements per plot.

3.1.2 Soil moisture at the catchment scale

5

The pdf of the on-alert measurements that are conducted at the same time in different plots of the same land use $\theta(x_{i,j}, t_{ev})$ agree either with the normal distribution (29 out of 50 data sets) or show a bi-modal or multi-modal distribution (remaining 21 data sets, Fig. 3b). This is an indication of inter-plot variability within the same land use class. In fact, the mean of the inter-plot variability within one land use class $\sigma_{1u}^{inter}(t_{ev})$ exceeds the mean inner-plot variability of the same land use class $\sigma_{j\in u}^{inner}(t_{ev})$ almost everywhere, except of in the vineyards at depth 10 cm (Table 1). There is a weak correlation between $\tilde{\theta}(t_{ev})$ and $\sigma_{cat.}^{inter}(t_{ev})$ (Fig. 3d). The variability at catchment scale is also high (Table 24), the catchment mean can be confidently assessed nonetheless. The mean of the standard error of the Claduègne catchment *SEM*_{cat.}^{inter} is 1.5 vol%, the maximum is 1.7 vol%. The mean *SEM*_{cat.}^{inter} of the Gazel catchment is 1.3 vol%, the maximum is 1.7 vol%.

- 15 Both the on-alert and the continuous measurements were analyzed for differences between the land use classes. The results of the student t tests that are conducted for all on alert measurements to detect differences between soil moisture in the different land use classes are given in Table 2. The null hypothesis (true difference in means is equal to zero) can be rejected under a significance level of $\alpha = 0.05$ only a limited number of times (upper panel). Furthermore, only the difference between the grasslands and the cultivated field is a systematic one, meaning that for all cases with significant differences the grasslands
- 20 have a higher soil moisture that the cultivated field (lower panel). In the present study site the plot means of grasslands, vineyards and a fallow are not significantly different from each other (Student t test, significance level $\alpha = 0.05$). Only in the cultivated field mean soil moisture is significantly and systematically different from the one in the grasslands.

A comparison of the variability between the four land use classes expressed as $\sigma^{\text{betw}}(t_{\text{ev}})$ to the one within land use classes $\sigma_{\text{lu}}^{\text{inter}}(t_{\text{ev}})$ or within plots $\sigma_{\text{l}}^{\text{inner}}(t_{\text{ev}})$ also reveals that it is smaller than both other standard deviations (Table 24). The initial

- and final soil moisture profile of the first major event in 2013 (ev. 27) shows, nonetheless, that there are differences in the profile shape and in the wetting behavior between grasslands and vineyard (Fig. 4<u>2c and 2d</u>). While the grasslands have a nearly homogenous profile before the rain event, the vineyards have a much more pronounced vertical soil moisture profile with higher values in the deeper layers. In response to the rain event, the profile of the grasslands shifts towards higher soil moisture, with similar differences in each depth. In the vineyards, mainly the moisture in the topmost layer increases, whereas
- 30 soil moisture in the subsoil hardly changes.

Figure <u>35</u> shows the relative spatial difference $\delta_{j,ev}$ of all plots for all on-alert measurements conducted from 2012 to 2015. It can be seen that temporal stability is found to some degree. The mean values of some plots are (nearly) consistently below the catchment mean (v4, v3, g5, f1, c1), others above (g4, g3, g1). This is also the case if deviations were related to the land use

mean instead of the catchment mean (see e.g. the noticeable difference between grasslands g2 and g5 on the one hand and g1, g3 and g4 on the other hand). However, there are also plots with above average soil moisture for a certain period of time and below average soil moisture during other periods, indicated by a change in signs of $\delta_{j,ev}$ between events (v1, v2, g2). The plots with the lowest mean spatial difference $\overline{\delta_j}$ are v1, v2 and c1 (3, 4 and 8 % respectively). The one with the lowest temporal variability of the relative spatial difference ($\sigma_{\delta_i} = 5$ %) is the cultivated field c1.

3.2 Temporal dynamics

5

25

Figure 6-4 shows the evolution of soil moisture in 10, 25 and 40 cm depth in autumn 2013. Due to several large rain events in July and August 2013 (not shown here), soil moisture at the beginning of the season is already relatively high-ca. 25 vol% at 10 cm depth and ca. 30 vol% in the deeper layers. In the topsoil, however, soil moisture is much lower at the beginning of the

10 season: the catchment mean value of the first measurement conducted on 28 September 2013 is 10.2 vol% (Fig. 75). After the first major rain events, it remains constantly above 30 vol% at 10 cm depth and above 36 vol% in the deeper layers, with maximum values of around 42 vol% reached after major rain events. This value is not exceeded, even after rain events that occur during wet initial conditions.

Temporal variability of soil moisture varies considerably between wet and dry conditions. Soil moisture in all continuously

- 15 sampled depths increases rapidly as a response to rain events by up to 12.6 vol% in less than 1 d (Fig. 64). Differences between initial and final state in the topsoil can be even larger (maximum difference: 16 vol% in November 2014, Fig. 75). The rapid response is evident from the small lag between the peak of rainfall and the peak of soil moisture (usually less than two hours for all soil layers). The maximum of the crosscorrelation function ($L_{CC_{max}}$) between hourly rainfall and hourly catchment mean soil moisture at 10 cm depth is at the lag time of 1 h. Mean $L_{CC_{max}}$ between the probes in 10 cm and the probes in 25 cm
- 20 depths is 0.3 h and the one between 25 and 45 cm is 1.3 h, which indicates a fast percolation to deeper layers. There are differences between grasslands (median $L_{CC_{max}} = 0.15$ h), vineyards (1.35 h) and the fallow (3.3 h). As these values are in the same order of magnitude as the resolution of the underlying data, their uncertainty is high. Thus, they represent only estimates for the order of magnitude of percolation time.

A few days after the rain events, soil moisture decrease slows down and soil moisture during dry conditions can persist for long time periods as in the first half of December 2013 (Fig. 6).

3.23 Event-based runoff coefficients

The event-based runoff coefficients ϕ_{ev} calculated for the Gazel catchment with seven hydrograph separation methods for 54 events range from 0 to 0.99 with large differences between the methods and a high positive skewness because of many low values (Fig. <u>8a6a</u>). In the Claduègne catchment, ϕ_{ev} was only calculated with the recursive digital filter method RDF and the

30 Hysep filter methods, but values still range from 0 to 0.97. The electric conductivity and constant-K methods result in the lowest values for ϕ_{ev} while the 3 Hysep filters yield considerably larger values than all other methods. Except of the Hysep filter methods, the other four methods correlate well with each other (Fig. 8b6b). The Hysep filters correlate very well with

each other ($R^2 \ge 0.96$ for all three pairs; not shown here), but to a lesser degree with the other methods (Fig. <u>8b6b</u>). For the following sections, values of ϕ_{ev} determined with the RDF method are used for reasons explained in the discussion (Sect. 4.4). The uncertainty of ϕ_{ev} associated with the stage-discharge relation is quantified with the BaRatin method and is shown in Fig. 9-7 as black vertical error bars. In both catchments this uncertainty is very small for events with low ϕ_{ev} while it can get up to

- 5 0.28 (difference between ϕ_{ev} calculated with the 5 % and the 95 % confidence interval of discharge) for event 40 which is the one with the highest ϕ_{ev} and highest discharge in both catchments (Table 3 and Table 4). The uncertainty due to different ϕ_{ev} obtained by different hydrograph separation methods is visualized as gray vertical error bars in Fig. 97. It can be very high for any event regardless of the ϕ_{ev} and is often due to the discordance of the HySep methods with the other methods. The mean range in ϕ_{ev} calculated with different hydrograph separation methods is 0.08, the one of ϕ_{ev} calculated with 5 % CI and 95 %
- 10 CI of discharge is 0.02. The mean standard deviation of ϕ_{ev} calculated with different hydrograph separation methods is 0.03, when the HySep methods are excluded it decreases to 0.02. However, these measures are biased by the important positive skewness of the distribution of ϕ_{ev} .

Factors that are suggested to influence ϕ_{ev} include rainfall depth and rain intensity. Figure 9–7 shows the correlation of the meteorological forcing quantified as cumulative <u>catchmentevent</u> rainfall depth P_{cum} , mean rain intensity I_{μ} , and maximum 20

- 15 min rain intensity $I_{\text{max},20}$ with ϕ_{ev} . In the present data set there is a weak correlation between ϕ_{ev} and P_{cum} , I_{μ} and $I_{\text{max},20}$ (Fig. 97) 97) ecoefficients of determination $R^2 = 0.24$, 0.17 and 0.27 respectively). The correlations of ϕ_{ev} with maximum rain intensity calculated at 2, 10, 30 and 60 min time steps were worse that the one at 20 min. None of these <u>variablesmeasures</u> can, therefore, explain more than 30 % of the variability of ϕ_{ev} . Figure 9-7 shows that events with similar rainfall characteristics (events 30 and 40, similar intensity) can have very different ϕ_{ev} . Additionally, similar ϕ_{ev} are obtained for events with very different 20 min fills the second data are the second data and the second data are the second data
- 20 rainfall characteristics (events 22 and 39). These striking differences in catchment behavior can partly be attributed to differences in initial soil moisture as shown in the following section.

3.34 Soil moisture's impact on runoff generation

The hydrological responses concerning the temporal dynamics of soil moisture and discharge in reaction to rain events of the two catchments vary greatly. In Table 3 and Table 4 the characteristics of all rain events in autumn 2013 that generate event
flow at the river gauges of the Gazel and / or Claduègne are given. The hyetographs, hydrographs and time series of catchment mean soil moisture of four of these events with very different behavior are exemplarily shown in Fig. 108: event 27 and event 30 occur at the beginning of the season when soil moisture is still relatively low. Rainfall leads to a considerable increase in soil moisture in all three layers and to a storage change ΔS in the 50 topmost cm of the soil profile that constitutes a notable share of cumulative precipitation. For event 30 on-alert surface soil moisture measurements are available as well and show a
sharp increase from 10.2 vol% before to 25.7 vol% after the event (Fig. 75). The runoff coefficients φ_{ev} of both events are very low. The within-event temporal dynamic of rainfall, soil moisture and runoff during event 27 is also noteworthy: the discharge peak does not follow the rainfall peak, which is closely followed by the steepest increase in soil moisture, but the second rainfall pulse that occurs when soil moisture is considerably higher than at the beginning of the event. As a response to this

much smaller rainfall impulse, soil moisture rises only slightly. This behavior is also observed during event 40 where the first rainfall impulse leads to a sharp increase in soil moisture and only a small discharge peak, while the second rainfall pulse generates a substantial discharge peak and only a slight increase in soil moisture. Event 40 and event 53 both occur during wet initial soil moisture conditions, but event 53 has a much smaller ϕ_{ev} than event 40. During these two events also an inversion

5 of the vertical soil moisture profile, i.e. temporally higher soil moisture at the topsoil than in deeper layers, can be observed approximately at the time of peak discharge. This inversion is an indicator of Hortonian overland flow. <u>Overland flow was</u> indeed observed during event 40 in vast areas in the north of the Claduègne catchment (Supplmentary material S2).

The large range of ϕ_{ev} of events with high $\tilde{\theta}_{ini}$ can also be seen in Fig. <u>449</u>. While the three events with low $\tilde{\theta}_{ini}$ consistently have very low ϕ_{ev} , <u>afterabove</u> a threshold of approximately 34 vol%, ϕ_{ev} can have a value anywhere between <u>zero</u> θ and <u>one</u>4.

- 10 An examination of Fig. <u>11–9</u> shows that both high $\tilde{\theta}_{ini}$ and high P_{cum} are necessary but not sufficient criteria for high ϕ_{ev} and that the relation between $\tilde{\theta}_{ini}$ and ϕ_{ev} is characterized by strong non-linearity and threshold effects. This is observed in both catchments and the threshold value is very similar for the Gazel and the Claduègne catchment. Further analysis of the relation between $\tilde{\theta}_{ini}$ and ϕ_{ev} for events with high P_{cum} and high $\tilde{\theta}_{ini}$ is limited because of the low number of events fulfilling these criteria and the uncertainty of both $\tilde{\theta}_{ini}$ and ϕ_{ev} .
- 15 Consideration of single events shows, that $\tilde{\theta}_{ini}$ can partly explain the high scatter in Fig. 97. The contrary behavior of events 30 and 40 can be explained by different initial soil moisture conditions. It can also be hypothesized that the high ϕ_{ev} of event 22 despite low P_{cum} , I_{μ} and $I_{max,20}$ is due to high initial soil moisture. For this event, that started on November 26th, 2012, only on-alert soil moisture is available and initial topsoil moisture is relatively high at 23.5 vol%. The event ocurredoccurred late in the season (26 November 2012, Fig. 75) two weeks after a heavy rain event with $P_{cum} = 72.5$ mm. Event 39 which has a 20 similar ϕ_{ev} but higher P_{cum} , I_{μ} and $I_{max,20}$, however also occurred during high initial moisture conditions (36 vol%)-which
- indicates that the relation between ϕ_{ev} , $\tilde{\theta}_{ini}$ and rainfall characteristics cannot easily be generalized. The results of hydrograph separation with the electric conductivity method give some insights into runoff generating processes in the Gazel catchment. During many events (e.g. Fig. 12a), there is a very large contribution of subsurface flow to total event flow. This is also reflected in rather low values for ϕ_{ev} obtained with the EC method compared to other methods. The subsurface
- 25 runoff component reacts very quickly during the rising limb of the hydrograph (e.g Fig 12a). Furthermore, the response of EC differs greatly between events (Fig. 12). A high signal in electric conductivity, ΔEC_{max} indicates that surface runoff contributes strongly to total stream discharge while a low value indicates a high impact of subsurface stormflow. Values for ΔEC_{max} for different events are given in Table 3 and Table 4 and shown in Fig. 13. The highest values for ΔEC_{max} are observed during events 30,39 and 40 suggesting a strong contribution of surface runoff for these events. In fact, for event 40, overland flow
- 30 was observed during a field visit on vast areas in the north of the Claduègne catchment (see pictures in the supplementary material S2). ΔEC_{max} -generally increases with P_{eum} and peak discharge Q_{p} , but there is no correlation with $\tilde{\theta}_{\text{imi}}$ ($R^2 = 0.00$) and only a moderate one with I_{tr} and $I_{\text{max},20}$ ($R^2 = 0.51$ and 0.64 respectively). Figure 13 shows that high values of ΔEC_{max} -are

observed for several events with high $\tilde{\theta}_{imi}$ (39, 40, 52). A high value of ΔEC_{max} is also measured during event 30, with low $\tilde{\theta}_{imi}$. This event has the highest $I_{max,20}$ and considerable cumulative precipitation.

4 Discussion

4.1 Validity of the soil moisture sampling strategy Soil moisture estimation at the catchment scale

- 5 The sampling design applied in this project proved to be efficient to assess spatial variability of soil moisture across scales, as well as to document temporal dynamics. The on-alert measurements of soil moisture allow a good estimate of the plot mean with a low mean SEM_j^{inner} of 0.8 vol% as well as an accurate estimate of the inner-plot variability, quantified as $\sigma_j^{\text{inner}}(t_{\text{ev}})$ during wet and dry conditions. On the other hand, the continuous soil moisture measurements cover a larger extent of the two studied catchments and different depths in the soil profile. Due to the higher variability at the catchment scale, the mean
- 10 $SEM_{cat.}^{inter}$ is somewhat higher (1.5 vol%). The values obtained for SEM_{j}^{inner} and $SEM_{cat.}^{inter}$ show that at an accepted uncertainty of the mean of ± 2 vol%, the number of 10 measurements per plot is sufficient. This is consistent with the results of Zucco et al. (2014) who found a maximum number of 11 or 20 required samples at the plot scale and catchment scale respectively and the ones of Molina et al. (2014) who concluded that plot mean soil moisture in a Mediterranean mountain area was well represented with 9 probes. The review by Vereecken et al. (2014) shows that there is a wide range of estimates for these
- 15 <u>numbers and that they are site specific.</u> Furthermore, tThe continuous measurements reveal the temporal evolution of soil moisture over the season and within events. The only drawback is the lack of continuous soil moisture estimates in the topsoil. The sampling at the plot scale and in nested catchments is considered to be a good approach to assess heterogeneity across scales and to cope with the change of scale problem (Braud et al., 2014).

This study's result of normally distributed soil moisture at the plot scale with a high spatial variability at scales smaller than

- 20 ten meters agrees well with other studies reviewed in Vereecken et al. (2014) and with results that Huza et al. (2014) obtained in the grasslands of the Gazel catchments. The latter authors also analyzed semi-variograms of six grasslands and report a very high nugget which indicates systematically high differences between measurements of less than 1 m distance. Unlike other studies that report positively or negatively skewed distributions for dry and wet conditions (Vereecken et al., 2014), the data of the present study is normally distributed during wet and dry conditions as well. The finding of a lower variability during
- 25 dry conditions (Huza et al., 2014; Vereecken et al., 2014) is not observed here ($R^2 = 0.04$), even though the range of plot mean soil moisture covers very diverse conditions. The differences of this study's results and the ones found by Huza et al. (2014) may be due to a different sampling locations as Huza et al. (2014) did not sample vineyards. If in this study's data set only grasslands are considered there is a slightly better but still poor correlation ($R^2 = 0.13$) between $\overline{\theta}_j(t_{ev})$ and $\sigma_j^{inner}(t_{ev})$.
- The observation that the probability density functions of soil moisture measurements on various plots in the catchment either agree with the normal distribution function or show a bi-or multi-modal behavior also agrees with other study's findings (Vereecken et al., 2014). In this study the inter-plot variability within one land use class $\sigma_{lu}^{inter}(t_{ev})$ usually exceeds the inner-

plot variability $\sigma_j^{\text{inner}}(t_{\text{ev}})$. This is not consistent with findings of Huza et al. (2014) which may-again be due to different sampling strategies: whereas Huza et al. (2014) conducted measurements along 50 m transects, for this study random locations were sampled within one field in an area of ≈ 20 by 20 m. At scales larger than 10 m, they found a spatial structure revealed by a higher semi-variance at distances of more than 10 m in at least one of their transects.

5 The temporal dynamics with a quick response to rain events and the strong accordance between the evolution of soil moisture and discharge of small catchments is also observed by Braud et al. (2014) in the Valescure catchment. This catchment is also located in the Cévennes Vivarais region and shows a similar dynamic of the interaction of soil moisture and discharge to the one observed here.

4.2 Comparison between land use classes and analysis of temporal stability

- 10 The results of this study indicate differences between grasslands and vineyards in the vertical soil profile <u>and</u>, in the response of the profile to rain events, in percolation behavior and in the persistence of soil moisture conditions. These differences are most likely due to differences in soil texture, as vineyards are usually found on soils with higher clay content than the ones of the other land use types. The different percolation behavior of grasslands may also be due to differences in root structure as the dense grass roots may accelerate downward water movement along preferential flow paths more efficiently than the ones
- 15 of the vine plants. However, there are no significant and systematic differences between the plot means of different land use classes. Thus, land use cannot be used as additional information to improve spatially distributed soil moisture estimation in the study site.

The cultivated field c1 shows a remarkable temporal stability of the difference of this plot's mean soil moisture and the catchment mean $\delta_{j,ev}$. This suggests that if the catchment mean has to be approximated by measurements in just one field, this

- 20 one is the best choice (Vachaud et al., 1985; Vanderlinden et al., 2012). Other fields show, however, that $\delta_{j,ev}$ is not consistent in time. The observation that several sites change the sign of $\delta_{j,ev}$ between measurements was also made <u>on-at</u> the plot scale on a grassland, a field cultivated with wheat and an olive grove by Vachaud et al. (1985) and <u>on-at</u> the catchment scale on grasslands by Huza et al. (2014). Here, notably <u>the vineyards</u> v1 and v2 are considerably wetter than the catchment mean throughout the autumn seasons of 2012 and 2013, dryer in 2014 and again wetter in 2015. Possible reasons include changes in
- 25 cultivation. Especially tillage practices play an important role in the vineyards (not shown here). Therefore, conclusions based on this finding should be considered carefully. Moreover, the choice of the plot which best represents the catchment mean should include the temporal variability of $\delta_{j,ev}$ and should not be solely based on the minimal mean difference $\overline{\delta_j}$ which is in this case the one of v1 and v2.

4.23 Quantification of the hydrological response

30 Besides the extensive soil moisture data set used for this study, t<u>T</u>he available precipitation and discharge data at a high spatiotemporal resolution is a major asset that is necessary to understand the hydrological processes at small scales and during short time spans that lead to flash flood generation (Nord et al., 2017). It allows to calculate the event-based runoff coefficient ϕ_{ev} and to estimate its uncertainty. The main sources of its uncertainty comprise that of the stage discharge relation which is especially important for events with high discharge and which was assessed with the BaRatin framework, the uncertainty associated to the choice of the method used for hydrograph separation and the uncertainty of the catchment mean precipitation. The latter source of uncertainty is not considered in this study. It stems from the rainfall measurements with tipping buckets

5 and the interpolation between the rain gauges. Tipping buckets are known to underestimate precipitation at high intensities (Marsalek, 1981; Molini et al., 2005), thus, including radar data could improve the estimation of catchment mean rainfall even in relatively well gauged catchments such as the ones of the Gazel and Claduègne (e.g. Creutin and Borga, 2003; Delrieu et al., 2014; Abon et al., 2015).

In this study, the uncertainty associated to the hydrograph separation method exceeds that of the stage-discharge relation. The

- 10 high range and positive skewness of event-based runoff coefficients is consistent with other study's results (Merz et al., 2006; Blume et al., 2007; Merz and Blöschl, 2009; Norbiato et al., 2009; Marchi et al., 2010). The dependence of ϕ_{ev} on rain characteristics suggested by other authors (Merz et al., 2006; Norbiato et al., 2009) was not entirely confirmed in this study as none of the rain characteristics examined here (P_{cum} , I_{μ} , $I_{max,20}$) could explain more than 30 % of the variability in ϕ_{ev} . 4.4 Validity of the hydrograph separation method
- 15 Each of the hydrograph separation methods used here has advantages and disadvantages. The method based on electric conductivity (*EC*) has a physically based foundation as it distinguishes components with different *EC* and represents subsurface flow dynamics. This method could not be applied to both catchments because values for <u>surface flow</u> EC_{sev} were only available from Le Pradel in the south of the Gazel catchment and it is assumed that EC_{sev} on the basaltic plateau differs considerably while this geology accounts for a large part of the Claduègne catchment. Furthermore, it is not possible to conduct a three
- 20 component hydrograph decomposition with the available data, so unlike with the other methods, the fast reacting subsurface flow is considered to be baseflow. Thus, event discharge is underestimated. Unlike the other methods, the Constant-k (CK) method offers a physical explanation for the end of event flow. The method builds on the assumption of baseflow behaving like the slow responding outflow of a linear reservoir. For the discharge data of the Gazel, this method could not always be applied because of the low discharge that results in "steps" in the data and high
- 25 noise so the threshold for defining that k is constant as proposed by Blume et al. (2007) was never reached. An adjusted threshold yielded reasonable results for some but not all events. The straight line method was rejected because it does not consider baseflow dynamics and the end of event flow has to be determined arbitrarily. The filter methods have the advantage of being easy to apply to all data sets without further data treatment or demand of additional data but these methods are very sensitive to parameters such as the interval width (HySep)
- 30 filters) or the number of passes (RDF). The HySep filters were discarded because of the disagreement with the other methods. Thus, the RDF method was used for all further analyses because it correlates well with the EC and CK methods and can easily be applied to all events and both catchments. The number of passes had to be calibrated as suggested by Ladson et al. (2013) in a manner that ϕ_{ev} is below 1 for all cases and that it is slightly higher than the value obtained with the EC method in order

to compensate for the underestimation of event discharge. Nonetheless, underestimation of event discharge is still a source of uncertainty.

4.35 The impact of initial soil moisture on the hydrological response

- The relation between ϕ_{ev} and $\tilde{\theta}_{ini}$ is not as clear as one might have expected from other studies results which suggest a dependence of ϕ_{ev} on $\tilde{\theta}_{ini}$. Moreover, both variables are still subject to large uncertainties. Catchment mean initial soil moisture $\tilde{\theta}_{ini}$ below a threshold of 34 vol% inhibits high ϕ_{ev} . However, only three of the events considered here occur during such dry conditions, so further measurements would be useful to corroborate this finding. Threshold effects in the relation of ϕ_{ev} and $\tilde{\theta}_{ini}$ are also observed by other authors (e.g. McMillan et al., 2014; Hrachowitz et al., 2011). It is consistent with the observations made by Huza et al. (2014) in the Gazel catchment and Braud et al. (2014) in the Valescure catchment. In the
- 10 Mediterranean context, Fishe thresholds obtained by these authorsHuza et al. (2014) in the Gazel catchment and Braud et al. (2014) in the Valescure catchment (22 and 25 vol% respectively) are lower than the one obtained here. The threshold at 45 vol% observed by Penna et al. (2011) in a 1.9 km² headwater catchment in the Italian Dolomites on the other hand is higher than the one obtained here. McMillan et al. (2014) show that thresholds in different subcatchments of a 50 km² catchment in New Zealand are highly variable: They range between 27 and 58 vol% and are more or less pronounced in different.
- 15 <u>subcatchments.</u> These differences might be due to different soil and land use features, <u>climate</u> and different sampling designs. The values for $\tilde{\theta}_{ini}$ that Huza et al. (2014) used are obtained from satellite data, while this study uses in situ data from several land use classes. Moreover, a profile mean is considered here, while Huza et al. (2014) used only values of topsoil moisture. Furthermore, different methods were applied for hydrograph separation. Huza et al. (2014) used a method similar to the HySep 3 filter, which yielded different results than the other methods applied for this study.
- 20 The high range of ϕ_{ev} obtained at high $\tilde{\theta}_{ini}$ also agrees with findings of Huza et al. (2014). It indicates, indicates that the hydrological response is influenced by other factors as well. The parameters describing the impact of meteorological forcing (P_{cum} , I_{μ} and $I_{max,20}$) neither explain that variability. When only events with high cumulative precipitation are considered, the range is still very high. Results obtained in virtual experiments (Merz and Plate, 1997; Bronstert and Bárdossy, 1999; Zehe and Blöschl, 2004) by Merz and Plate (1997) and Zehe and Blöschl (2004) showed that spatial patterns of soil moisture and
- 25 threshold effects strongly impact the runoff response. The latter authors show, that especially during initial moisture conditions close to the threshold, the runoff response depends strongly on the resampling of spatially distributed soil moisture. Therefore, actual, small-scale soil moisture patterns that control connectivity of pathways but are not reflected in the catchment mean value used for Fig. 11-are a possible explanation for the very diverse runoff behavior. On the other hand, Morbidelli et al. (2016) conclude that spatial heterogeneity of $\tilde{\theta}_{ini}$ does not affect the runoff response for events that are associated with floods,
- 30 so more research is needed on this topic. Additionally, subsurface flow along preferential flowpaths can contribute to high ϕ_{ev} and Hortonian overland flow is not directly related to $\tilde{\theta}_{ini}$ but produces a substantial proportion of event flow.

The results of this study partly confirm the suggestions of other authors (i.e. Brocca et al., 2009b; Javelle et al., 2010; Grillakis et al., 2016) to consider estimates of initial soil moisture in flash flood warning based on the dependence of ϕ_{ev} on $\tilde{\theta}_{ini}$. This offers high potential for predictions in poorly gauged or ungauged basins given the global availability of remote sensing soil moisture data (Crow et al., 2005; Beck et al., 2009; Brocca et al., 2009c; Massari et al., 2014a,b). Threshold-based warning

- 5 systems are advocated e.g. by Norbiato et al. (2008). Including a threshold value for initial soil moisture could prevent false positive flash flood warnings in cases when high precipitation is expected under dry initial catchment conditions, while above-threshold soil moisture in combination with high precipitation increases the likelihood of high runoff coefficients. This threshold seems not to be scale-dependent. However, the threshold values differ between catchments and depend to a high degree on the methodology to determine it as indicated by the different values in this study and the one by Huza et al. (2014).
- 10 Furthermore, there are high data requirements to determine such thresholds and it is not known whether they can be transferred from one catchment to another, so it is not applicable for operational flash-flood warning. Moreover, the high scatter of ϕ_{ev} under high initial soil moisture conditions suggests that the relation between ϕ_{ev} and $\tilde{\theta}_{ini}$ is very complex and depends on other factors and processes that are still insufficiently understood. Thus, the impact of soil moisture on the hydrological response during wet catchment conditions cannot be generalized based on the results obtained here. Further research and instrumentation
- 15 could include the installation of piezometers in the catchment to understand subsurficial flow in the catchment, using tracers other than *EC* to differentiate subsurficial stormflow as a third flow component during hydrograph separation as well as the application of multivariate regression analysis methods that systematically examine different controls on ϕ_{ev} such as meteorological forcing as well as $\tilde{\theta}_{ini}$ and their interactions.

The analysis of the baseflow dynamics as obtained by the hydrograph separation with *EC* shows that for almost all events there is a quick response of the subsurface stormflow which constitutes a large proportion of total event flow. This indicates that subsurface stormflow plays an important role in the Gazel catchment and should be considered thoroughly. The maximum *EC* signal during discharge events ΔEC_{max} varies considerably between events. Figure 13 shows that possible reasons include high $\tilde{\theta}_{imi}$ and high P_{eum} which might be indicators for saturation excess overland flow but also high $I_{max,20}$, a possible indicator of infiltration excess Hortonian overland flow. Even though $I_{max,20}$ is usually below 10 mm h⁻¹ the rather low values for saturated conductivity *Ks* measured by Braud and Vandervaere (2015) indicate that rain intensity can locally exceed infiltration

capacities.

5 Conclusions

This study aimed at assessing the influence of initial soil moisture on the hydrological response in a flash-flood prone area in southern France. To this end, two research questionsissues were addressed and exemplarily examined in the nested Gazel (3.4

30 km²) and Claduègne (43 km²) catchments: (1) How can soil moisture heterogeneity in time and space be described and how does soil moisture correlate with land use? Obtaining a meaningful estimate of soil moisture at the catchment scale and (2) What is the relationship between initial soil moisture $\tilde{\theta}_{mr}$ and the hydrological response quantified as the event-based runoff

coefficient ϕ_{ev} ? Analyzing the relation between initial soil moisture $\tilde{\theta}_{ini}$ and the hydrological response quantified as the eventbased runoff coefficient ϕ_{ev} .

The main findings of this study related to the first research questions objective are:

- (1.1) Spatial variability of soil moisture at the plot scale and at the catchment scale is very high, and the probability density
- 5 function (pdf) of plot-scale soil moisture measurements usually follows a normal distribution. Soil moisture at the catchment scale is also very variable, and the pdf of the measurements often resembles a bi-or multi-modal distribution, which is an indicator for inter plot variability. This is corroborated as inter plot standard deviation σ^{inter} exceeds inner plot standard deviation σ^{inner} .

(1.2) There are differences between land use classes in the vertical soil moisture profile, notably that the grasslands have a

- 10 relatively homogeneous initial soil moisture profile whereas the vineyards have a curved profile. There are further differences in and in wetting behavior and in percolation, but no significant and systematic differences in catchment mean soil moisture values between land use classes <u>exist</u>. Hence, land use cannot be used as an auxiliary variable to determine the spatial distribution of soil moisture. Between land use standard deviation σ_{lu}^{betw} exceeds neither σ^{inter} nor σ^{inner} .
- (1.23) There is one plot, c1, with remarkable temporal stability of the spatial difference between plot mean and catchment
 mean. Thus, this field should be opted for, if the catchment mean had to be assessed from measurements in just one plot.
 However, none of the other plots shows this temporal stability.

(1.4) The temporal dynamics of soil moisture show a seasonal trend, a quick reaction to rain events and fast percolation to lower layers. During dry periods moisture conditions can persist for a long time with just a slow decrease in soil moisture.

The sampling design applied for this study allowed a detailed characterization of soil moisture heterogeneity across scales as well as the assessment of temporal dynamics. The catchment mean soil moisture was derived with a mean standard error of the catchment mean of 1.3 vol% or 1.5 vol% for the Gazel and Claduègne catchments respectively.

Main findings concerning the impact of initial soil moisture on the hydrological response quantified with the event-based runoff coefficient ϕ_{ev} are:

 $(2.1) \phi_{ev}$ obtained with different hydrograph separation methods can differ considerably, but results obtained with EC, CK, SL

and RDF methods correlate well. The RDF method was preferred for this study because it is easy to apply and because of the good correlation with the more physically based methods EC and CK which could not be applied to all events and both catchments.

(2.2) There is a week correlation between ϕ_{ev} and cumulative event precipitation P_{cum} , mean rain intensity I_{μ} and maximum 20 min rain intensity $I_{max,20}$ ($R^2 = 0.24, 0.17$ or 0.27 respectively).

30 (2.3) The hydrological response depends on initial soil moisture $\tilde{\theta}_{ini}$: below a threshold of 34 vol%, ϕ_{ev} remains very low, even during high precipitation events. However, there is a large scatter in ϕ_{ev} above that threshold indicating that other factors and processes also have an important impact on ϕ_{ev} . The threshold is identical for both catchments which indicates that at this study's site it might be scale invariant.

(2.4) Regarding the seasonal and within event evolution of soil moisture and discharge shows that discharge peaks of two considered events did not follow the peaks in rainfall, but a second, smaller rain impulse, while the rainfall peaks lead to a considerable refilling of soil water storage.

(2.5) Some events reacted with a strong signal in electrical conductivity, indicating an important contribution of overland flow.

5 There are several possible explanations for high *EC* signals, such as high P_{cum} , $\tilde{\theta}_{\text{ini}}$ or $I_{\text{max},20}$. Many events are dominated by a considerable contribution of subsurface flow to total event flow and show a fast response in subsurface stormflow to rain events.

These results indicate, that $\tilde{\theta}_{ini}$ does impact the hydrological response. For single events ϕ_{ev} or *EC* signals can be attributed to $\tilde{\theta}_{ini}$, P_{cum} or $I_{max,20}$. However, these results cannot be generalized and no systematic and unequivocal relationship between $\tilde{\theta}_{ini}$

- 10 and ϕ_{ev} was found. Thus, the second research question could only partly be answered. Even though the present data set is exceptionally detailed, there still is substantial uncertainty in the values for $\tilde{\theta}_{ini}$, P_{cum} and cumulative event flow $Q_{ev,cum}$. The results of this study support suggestions by other authors to include estimates of initial soil moisture in flash flood warning based on the dependence of ϕ_{ev} on $\tilde{\theta}_{ini}$ by including a threshold that could prevent false positive flash flood warnings under dry initial conditions.
- 15 The results of this study partly confirm the suggestions of other authors to consider estimates of initial soil moisture in flash flood warning based on the dependence of ϕ_{ev} on $\tilde{\theta}_{Inf}$. Threshold-based warning systems are advocated e.g. by Norbiato et al. (2008). Including a threshold value for initial soil moisture could prevent false positive flash flood warnings in cases when high precipitation is expected under dry initial eatchment conditions, while above threshold soil moisture in combination with high precipitation increases the likelihood of high runoff coefficients. This threshold seems not to be scale dependent.
- 20 However, the threshold values differ between catchments and depend to a high degree on the methodology to determine it as indicated by the different values in this study and the one by Huza et al. (2014). Furthermore, there are high data requirements to determine such thresholds and it is not known whether they can be transferred from one catchment to another, so it is not applicable for operational flash flood warning. Moreover, the high scatter of ϕ_{ev} under high initial soil moisture conditions suggests that the relation between ϕ_{ev} and $\tilde{\theta}_{im}$ is very complex and depends on other factors and processes that are still
- 25 insufficiently understood. Thus, the impact of soil moisture on the hydrological response during wet catchment conditions cannot be generalized based on the results obtained here.

In the Gazel catchment subsurface flow seems to constitute an important part of event flow, so further information on preferential flow path that could be obtained from tracer experiments would be helpful to understand the hydrological behavior of the catchment. Further research could also focus on the role of subsurface flow andor on elaborating multivariate regression

30 analysis methods. that systematically examine different controls on ϕ_{ev} such as meteorological forcing as well as $\tilde{\theta}_{imi}$ and their interactions.

Author contributions

Jean-Pierre Vandervaere and Isabella Zin conceptualized the study and Magdalena Uber carried out data analysis and its interpretation. Jean-Pierre Vandervaere, Isabelle Braud, Cédric Legoût, Gilles Molinié and Guillaume Nord were principal investigators in the FloodScale project responsible for instrumentation and criticized and provided the data used for this study.

5 All authors contributed to data analysis and interpretation of results. Magdalena Uber prepared the manuscript with contributions from all co-authors. The authors declare that they have no conflict of interest.

Acknowledgements

This work was funded by the French National Research Agency (ANR) via the FloodScale project under contract no. ANR 2011 BS56 027 which contributes to the HyMeX program. OHM-CV is supported by the Institut National des Sciences de

l'Univers (INSU/CNRS), the French Ministry for Education and Research, the Environment Research Cluster of the Rhône-Alpes Region, the Observatoire des Sciences de l'Univers de Grenoble (OSUG) and the SOERE Réseau des Bassins Versants (Alliance Allenvi) and belongs to the OZCAR Reasearch Infrastructure. Magdalena Uber received financial support from German Academic Exchange Service (DAAD). The authors want to thank Brice Boudevillain, Sahar Hachani, Simon Gérard and Cindy Nicoud for their help during field work, and Luca Brocca, Hongkai Gao and two anonymous referees for their comments on earlier versions of this manuscript that were very valuable to improve this paper.

References

- Abon, C.-C., Kneis, D., Crisologo, I., Bronstert, A., David, C. P. C. and Heistermann, M.: Evaluating the potential of radarbased rainfall estimates for streamflow and flood simulations in the Philippines, Geomat. Nat. Haz. Risk, 7(4), 1–16, doi:10.1080/19475705.2015.1058862, 2015.
- Andrieu, J.: Landcover map Claduègne catchment, doi:10.14768/mistrals-hymex.1381, http://mistrals.sedoo.fr/?editDatsId=1381, 2015.
 Archibald, J.: BaseflowSeparation{EcoHydRology}, https://cran.r-project.org/web/packages/EcoHydRology/, 2014.
 Baffaut, C., Ghidey, F., Sudduth, K. A., Lerch, R. N. and Sadler, E. J.: Long-term suspended sediment transport in the Goodwater Creek Experimental Watershed and Salt River Basin, Missouri, USA, Water Resour. Res., 49, 7827–7830, doi:10.1002/wrcr.20511, 2013.

Beck, H. E., de Jeu, R. A. M., Schellekens, J., van Dijk, A. I. J. M. and Bruijnzeel, L. A.: Improving Curve Number Based Storm Runoff Estimates Using Soil Moisture Proxies, IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens., 2(4), doi:10.1109/JSTARS.2009.2031227, 2009. Berthet, L., Andréassian, V., Perrin, C. and Javelle, P.: How crucial is it to account for the antecedent moisture conditions in flood forecasting? Comparison of event-based and continuous approaches on 178 catchments, Hydrol. Earth Syst. Sci., 13, 819–831, doi:10.5194/hessd-6-1707-2009, 2009.

5 Blöschl, G. and Sivapalan, M.: Scale Issues in Hydrological Modeling, Hydrol. Process., 9, 251 – 290, doi:10.1029/96EO00131, 1995.

Blume, T., Zehe, E. and Bronstert, A.: Rainfall—runoff response, event-based runoff coefficients and hydrograph separation, Hydrol. Sci. J., 52(5), 843–862, doi:10.1623/hysj.52.5.843, 2007.

Borga, M.: Accuracy of radar rainfall estimates for streamflow simulation, J. Hydrol., 267(1-2), 26-39, doi:10.1016/S0022-1694(02)00137-3, 2002.

10

Borga, M., Boscolo, P., Zanon, F. and Sangati, M.: Hydrometeorological Analysis of the 29 August 2003 Flash Flood in the Eastern Italian Alps, J. Hydrometeorol., 8(5), 1049–1067, doi:10.1175/JHM593.1, 2007.

Borga, M., Gaume, E., Creutin, J. D. and Marchi, L.: Surveying flash floods: gauging the ungauged extremes, Hydrol. Process., 22(18), 3883–3885, doi:10.1002/hyp.7111, 2008.

15 Boudevillain, B., Delrieu, G., Galabertier, B., Bonnifait, L., Bouilloud, L., Kirstetter, P. E. and Mosini, M. L.: The Cévennes-Vivarais Mediterranean Hydrometeorological Observatory database, Water Resour. Res., 47(7), 1–6, doi:10.1029/2010WR010353, 2011.

Braud, I. and Vandervaere, J. P.: Analysis of infiltration tests and performed in the Claduègne catchment in May-June 2012, Report to the FloodScale project, 66 pp., 2015.

- 20 Braud, I., Ayral, P. A., Bouvier, C., Branger, F., Delrieu, G., Le Coz, J., Nord, G., Vandervaere, J. P., Anquetin, S., Adamovic, M., Andrieu, J., Batiot, C., Boudevillain, B., Brunet, P., Carreau, J., Confoland, A., Didon-Lescot, J. F., Domergue, J. M., Douvinet, J., Dramais, G., Freydier, R., Gérard, S., Huza, J., Leblois, E., Le Bourgeois, O., Le Boursicaud, R., Marchand, P., Martin, P., Nottale, L., Patris, N., Renard, B., Seidel, J. L., Taupin, J. D., Vannier, O., Vincendon, B. and Wijbrans, A.: Multi-scale hydrometeorological observation and modelling for flash flood understanding, Hydrol. Earth Syst. Sci., 18(9), 3733–
- 3761, doi:10.5194/hess-18-3733-2014, 2014.
 Brocca, L., Melone, F., Moramarco, T. and Morbidelli, R.: Antecedent wetness conditions based on ERS scatterometer data, J. Hydrol., 364, 73–87, doi:10.1016/j.jhydrol.2008.10.007, 2009a.
 Brocca, L., Melone, F., Moramarco, T. and Singh, V. P.: Assimilation of Observed Soil Moisture Data in Storm Rainfall-Runoff Modeling, J. Hydrol. Eng., 14(2), 153–165, doi:10.1061/(ASCE)1084-0699(2009)14:2(153), 2009b.
- Brocca, L., Melone, F., Moramarco, T. and Morbidelli, R.: Soil moisture temporal stability over experimental areas in Central Italy, Geoderma, 148(3–4), 364–374, doi:10.1016/j.geoderma.2008.11.004, 2009c.
 Brocca, L., Melone, F., Moramarco, T., Penna, D., Morga, M., Matgen, A., Gumuzzio, A., Martinez-Fernandez, J. and Wagner, W.: Detecting threshold hydrological response through satellite soil moisture data, Die Bodenkultur, 64(3–4), 7–12, 2013.

Bronstert, A. and Bárdossy, A.: The role of spatial variability of soil moisture for modelling surface runoff generation at the small catchment scale, Hydrol. Earth Syst. Sci., 3(4), 505–516, 1999.

Castillo, V. M., Gómez-Plaza, a. and Martínez-Mena, M.: The role of antecedent soil water content in the runoff response of semiarid catchments: A simulation approach, J. Hydrol., 284(1–4), 114–130, doi:10.1016/S0022-1694(03)00264-6, 2003.

5 Cea, L., Legout, C., Grangeon, T. and Nord, G.: Impact of model simplifications on soil erosion predictions: application of the GLUE methodology to a distributed event-based model at the hillslope scale, Hydrol. Process., 30(7), 1096–1113, doi:10.1002/hyp.10697, 2015.

Creutin, J. D. and Borga, M.: Radar hydrology modifies the monitoring of flash-flood hazard, Hydrol. Process., 17(7), 1453–1456, doi:10.1002/hyp.5122, 2003.

- 10 Crow, W. T., Bindlish, R. and Jackson, T. J.: The added value of spaceborne passive microwave soil moisture retrievals for forecasting rainfall-runoff partitioning, Geophys. Res. Lett., 32(18), 1–5, doi:10.1029/2005GL023543, 2005.
 - Delrieu, G., Wijbrans, A., Boudevillain, B., Faure, D., Bonnifait, L. and Kirstetter, P. E.: Geostatistical radar-raingauge merging: A novel method for the quantification of rain estimation accuracy, Adv. Water Resour., 71, 110–124, doi:10.1016/j.advwatres.2014.06.005, 2014.
- 15 Ducrocq, V., Braud, I., Davolio, S., Ferretti, R., Flamant, C., Jansa, A., Kalthoff, N., Richard, E., Taupier-Letage, I., Ayral, P. A., Belamari, S., Berne, A., Borga, M., Boudevillain, B., Bock, O., Boichard, J. L., Bouin, M. N., Bousquet, O., Bouvier, C., Chiggiato, J., Cimini, D., Corsmeier, U., Coppola, L., Cocquerez, P., Defer, E., Delanoe, J., Di Girolamo, P., Doerenbecher, A., Drobinski, P., Dufournet, Y., Fourrié, N., Gourley, J. J., Labatut, L., Lambert, D., Le Coz, J., Marzano, F. S., Molinié, G., Montani, A., Nord, G., Nuret, M., Ramage, K., Rison, W., Roussot, O., Said, F., Schwarzenboeck, A., Testor, P., Van Baelen,
- 20 J., Vincendon, B., Aran, M. and Tamayo, J.: HyMeX-SOP1: The field campaign dedicated to heavy precipitation and flash flooding in the northwestern mediterranean, B. Am. Meteorol. Soc., 95(7), 1083–1100, doi:10.1175/BAMS-D-12-00244.1, 2014.

Dunne, T. and Black, R. D.: Partial Area Contributions to Storm Runoff in a Small New England, Water Resour. Res., 6(5), 1296 – 1311, 1970.

- Garcia-Estringana, P., Latron, J., Llorens, P. and Gallart, F.: Spatial and temporal dynamics of soil moisture in a Mediterranean mountain area (Vallcebre, NE Spain), Ecohydrology, 6, 741–753, doi:10.1002/eco.1295, 2013. Gaume, E., Livet, M., Desbordes, M. and Villeneuve, J. P.: Hydrological analysis of the river Aude, France, flash flood on 12 and 13 November 1999, J. Hydrol., 286(1-4), 135–154, doi:10.1016/j.jhydrol.2003.09.015, 2004. Gaume, E., Bain, V., Bernardara, P., Newinger, O., Barbuc, M., Bateman, A., Blaškovičová, L., Blöschl, G., Borga, M.,
- 30 Dumitrescu, A., Daliakopoulos, I., Garcia, J., Irimescu, A., Kohnova, S., Koutroulis, A., Marchi, L., Matreata, S., Medina, V., Preciso, E., Sempere-Torres, D., Stancalie, G., Szolgay, J., Tsanis, I., Velasco, D. and Viglione, A.: A compilation of data on European flash floods, J. Hydrol., 367(1-2), 70–78, doi:10.1016/j.jhydrol.2008.12.028, 2009. Gilbert, P. D. and Plummer, M.: acf{stats} Auto and Cross Covariance and Correlation Function Estimation, http://stat.ethz.ch/R-manual/R-devel/library/stats/html/acf.html, 2016.

Gonzales, -A. L., Nonner, J., Heijkers, J. and Uhlenbrook, S.: Comparison of different base flow separation methods in a lowland catchment, Hydrol. Earth Syst. Sci., 13(11), 2055–2068, doi:10.5194/hess-13-2055-2009, 2009.
 Gonzales Soza, E., Braud, I., Dehotin, J., Lassabatère, L., Angulo Jaramillo, R., Lagouy, M., Branger, F., Jacqueminet, C., Kermadi, S. and Michel, K.: Impact of land use on the hydraulic properties of the topsoil in a small French catchment, Hydrol.

- 5 Process., 24, 2382–2399, doi:10.1002/hyp.7640, 2010.
 Grayson, R. B., Western, A. W., Chiew, F. H. S. and Blöschl, G.: Preferred states in spatial soil moisture patterns: Local and nonlocal controls, Water Resour. Res., 33(12), 2897–2908, 1997.
 Grillakis, M. G., Koutroulis, A. G., Komma, J., Tsanis, I. K., Wagner, W. and Blöschl, G.: Initial soil moisture effects on flash flood generation A comparison between basins of contrasting hydro-climatic conditions, J. Hydrol., 541, 206–217,
- 10 doi:10.1016/j.jhydrol.2016.03.007, 2016.
 Hardie, M. A., Cotching, W. E., Doyle, R. B., Holz, G., Lisson, S. and Mattern, K.: Effect of antecedent soil moisture on preferential flow in a texture-contrast soil, J. Hydrol., 398(3–4), 191–201, doi:10.1016/j.jhydrol.2010.12.008, 2011.
 Horton, R. E.: The rôle of infiltration in the hydrologic cycle, Eos, Trans. Am. Geophys. Union, 14(1), 446–460, 1933.
 Hrachowitz, M., Bohte, R., Mul, M. L., Bogaard, T. A., Savenije, H. H. G. and Uhlenbrook, S.: On the value of combined
- 15 event runoff and tracer analysis to improve understanding of catchment functioning in a data-scarce semi-arid area, Hydrol. Earth Syst. Sci., 15(6), 2007–2024, doi:10.5194/hess-15-2007-2011, 2011.
 Huang, M., Gallichand, J., Dong, C., Wang, Z. and Shao, M.: Use of soil moisture data and curve number method for estimating runoff in the Loess Plateau of China, Hydrol. Process., 21(11), 1471–1481, doi:10.1002/hyp.6312, 2007.
 Huet, P., Martin, X., Prime, J.-L., Foin, P., Laurain, C. and Cannard, P.: Retour D'Expérience des Crues de Septembre 2002
- 20 dans les Departements du Gard de l'Hérault, du Vaucluse, des Bouches-du-Rhône, de l'Ardèche et de la Drôme. Rapport Consolidé après Phase Contradictoire, <u>http://www.ladocumentationfrancaise.fr/var/storage/rapports_publics/034000547.pdf</u>, 2003.

Huza, J., Teuling, A. J., Braud, I., Grazioli, J., Melsen, L. A., Nord, G., Raupach, T. H. and Uijlenhoet, R.: Precipitation, soil moisture and runoff variability in a small river catchment (Ardèche, France) during HyMeX Special Observation Period 1, J.

- Hydrol., 516, 330–342, doi:10.1016/j.jhydrol.2014.01.041, 2014.
 Javelle, P., Fouchier, C., Arnaud, P. and Lavabre, J.: Flash flood warning at ungauged locations using radar rainfall and antecedent soil moisture estimations, J. Hydrol., 394(1–2), 267–274, doi:10.1016/j.jhydrol.2010.03.032, 2010.
 Jawson, S. D. and Niemann, J. D.: Spatial patterns from EOF analysis of soil moisture at a large scale and their dependence on soil, land-use, and topographic properties, Adv. Water Resour., 30(3), 366–381, doi:10.1016/j.advwatres.2006.05.006,
- 30 <u>2007.</u>

Jarvis, N., Koestel, J., Messing, I., Moeys, J. and Lindahl, A.: Influence of soil, land use and climatic factors on the hydraulic conductivity of soil, Hydrol. Earth Syst. Sci., 17, 5185–5195, doi:10.5194/hess-17-5185-2013, 2013.

Korres, W., Reichenau, T. G., Fiener, P., Koyama, C. N., Bogena, H. R., Cornelissen, T., Baatz, R., Herbst, M., Diekkrüger, B., Vereecken, H. and Schneider, K.: Spatio-temporal soil moisture patterns – A meta-analysis using plot to catchment scale data, J. Hydrol., 520, 326–341, doi:10.1016/j.jhydrol.2014.11.042, 2015.

Ladson, A., Bronw, R., Neal, B. and Nathan, R.: A standard approach to baseflow separation using the Lyne and Hollick filter,
Aust. J. Water Resour., 17(1), 25–34, doi:http://dx.doi.org/10.7158/W12-028.2013.17.1., 2013.

Le Coz, J., Renard, B., Bonnifait, L., Branger, F. and Le Boursicaud, R.: Combining hydraulic knowledge and uncertain gaugings in the estimation of hydrometric rating curves: A Bayesian approach, J. Hydrol., 509, 573–587, doi:10.1016/j.jhydrol.2013.11.016, 2014.

Le Lay, M. and Saulnier, G. M.: Exploring the signature of climate and landscape spatial variabilities in flash flood events:

10 <u>Case of the 8-9 September 2002 Cévennes-Vivarais catastrophic event, Geophys. Res. Lett., 34(13), 1–5, doi:10.1029/2007GL029746, 2007.</u>

Li, H., Sivapalan, M. and Tian, F.: Comparative diagnostic analysis of runoff generation processes in Oklahoma DMIP2 basins: The Blue River and the Illinois River, J. Hydrol., 418–419, 90–109, doi:10.1016/j.jhydrol.2010.08.005, 2012.

15 Lyne, V. D. and Hollick, M.: Stochastic time-variable rainfall-runoff modeling, Institute of Engineers Australia National Conference Publications, 10, 89–92, 1979.

Marchi, L., Borga, M., Preciso, E. and Gaume, E.: Characterisation of selected extreme flash floods in Europe and implications for flood risk management, J. Hydrol., 394(1-2), 118–133, doi:10.1016/j.jhydrol.2010.07.017, 2010.

Marsalek, J.: Calibration of the tipping-bucket raingage, J. Hydrol., 53(3-4), 343–354, doi:10.1016/0022-1694(81)90010-X, 1981.

Massari, C., Brocca, L., Moramarco, T., Tramblay, Y. and Didon Lescot, J.-F.: Potential of soil moisture observations in flood modelling: Estimating initial conditions and correcting rainfall, Adv. Water Resour., 74, 44–53, doi:10.1016/j.advwatres.2014.08.004, 2014a.

Massari, C., Brocca, L., Barbetta, S., Papathanasiou, C., Mimikou, M. and Moramarco, T.: Using globally available soil
 moisture indicators for flood modelling in Mediterranean catchments, Hydrol. Earth Syst. Sci., 18(2), 839–853, doi:10.5194/hess-18-839-2014, 2014b.

Massari, C., Brocca, L., Ciabatta, L., Moramarco, T., Gabellani, S., Albergel, C., De Rosnay, P., Puca, S. and Wagner, W.: The Use of H-SAF Soil Moisture Products for Operational Hydrology: Flood Modelling over Italy, Hydrology, 2(1), 2–22, doi:10.3390/hydrology2010002, 2015.

30 McMillan, H., Gueguen, M., Grimon, E., Woods, R., Clark, M. and Rupp, D. E.: Spatial variability of hydrological processes and model structure diagnostics in a 50 km2 catchment, Hydrol. Process., 28(18), 4896–4913, doi:10.1002/hyp.9988, 2014. McMillan, H. K. and Srinivasan, M. S.: Characteristics and controls of variability in soil moisture and groundwater in a headwater catchment, Hydrol. Earth Syst. Sci., 19(4), 1767–1786, doi:10.5194/hess-19-1767-2015, 2015. Merz, B. and Plate, E. J.: An analysis of the effects of spatial variability of soil and soil moisture on runoff, Water Resour. Res., 33(12), 2909–2922, doi:10.1029/97WR02204, 1997.

Merz, R. and Blöschl, G.: A regional analysis of event runoff coefficients with respect to climate and catchment characteristics in Austria, Water Resour. Res., 45(1), 1–19, doi:10.1029/2008WR007163, 2009.

5 Merz, R., Blöschl, G. and Parajka, J.: Spatio-temporal variability of event runoff coefficients, J. Hydrol., 331(3-4), 591–604, doi:10.1016/j.jhydrol.2006.06.008, 2006.

Miller, M. P., Susong, D. D., Shope, Ch. L., Heilweil, V. M. and Stolp, B. J.: Continuous estimation of baseflow in snowmeltdominated streams and rivers in the Upper Colorado River Basin: A chemical hydrograph separation approach, Water Resour. Res., 50(8), 6986–6999, doi:10.1002/2013WR014939, 2014.

10 Molina, A. J., Latron, J., Rubio, C. M., Gallart, F. and Llorens, P.: Spatio-temporal variability of soil water content on the local scale in a Mediterranean mountain area (Vallcebre, North Eastern Spain). How different spatio-temporal scales reflect mean soil water content, J. Hydrol., 516, 182–192, doi:10.1016/j.jhydrol.2014.01.040, 2014. Molini, A., Lanza, L. G. and La Barbera, P.: The impact of tipping-bucket raingauge measurement errors on design rainfall

for urban-scale applications, Hydrol. Process., 19(5), 1073–1088, doi:10.1002/hyp.5646, 2005.

- Molinié, G., Ceresetti, D., Anquetin, S., Creutin, J. D. and Boudevillain, B.: Rainfall regime of a mountainous mediterranean region: Statistical analysis at short time steps, J. Appl. Meteorol. Clim., 51(3), 429–448, doi:10.1175/2011JAMC2691.1, 2012. Moran, M. S., Holifield Collins, C. D., Goodrich, D. C., Qi, J., Shannon, D. T. and Olsson, A.: Long-term remote sensing database, Walnut Gulch Experimental Watershed, Arizona, United States, Water Resour. Res., 44(5), 1–8, doi:10.1029/2006WR005777, 2008.
- 20 Morbidelli, R., Saltalippi, C., Flammini, A., Corradini, C., Brocca, L. and Govindaraju, R. S.: An investigation of the effects of spatial heterogeneity of initial soil moisture content on surface runoff simulation at a small watershed scale, J. Hydrol., 539, 589–598, doi:10.1016/j.jhydrol.2016.05.067, 2016.

Nakamura, R.: Runoff analysis by electrical conductance of water, J. Hydrol., 14(3-4), 197–212, doi:10.1016/0022-1694(71)90035-7, 1971.

- Nathan, R. J. and McMahon, T.: Evaluation of Automated Techniques for Base Flow and Recession Analyses, Water Resour. Res., 26(7), 1465–1473, doi:10.1029/WR026i007p01465, 1990.
 Norbiato, D., Borga, M., Degli Esposti, S., Gaume, E. and Anquetin, S.: Flash flood warning based on rainfall thresholds and soil moisture conditions: An assessment for gauged and ungauged basins, J. Hydrol., 362(3-4), 274–290, doi:10.1016/j.jhydrol.2008.08.023, 2008.
- Norbiato, D., Borga, M., Merz, R., Blöschl, G. and Carton, A.: Controls on event runoff coefficients in the eastern Italian Alps, J. Hydrol., 375(3-4), 312–325, doi:10.1016/j.jhydrol.2009.06.044, 2009.
 Nord, G., Boudevillain, B., Berne, A., Branger, F., Braud, I., Dramais, G., Gérard, S., Le Coz, J., Legout, C., Molinié, G., Van Baelen, J., Vandervaere, J. ., Andrieu, J., Aubert, C., Calianno, M., Delrieu, G., Grazioli, J., Hachani, S., Horner, I., Huza, J., Le Boursicaud, R., Raupach, T., Teuling, A. J., Uber, M., Vincendon, B. and Wijbrans, A.: A high space–time resolution

dataset linking meteorological forcing and hydro-sedimentary response in a mesoscale Mediterranean catchment (Auzon) of the Ardèche region, France, Earth Syst. Sci. Data, 9, 221 – 249, 2017.

Panziera, L., James, C. N. and Germann, U.: Mesoscale organization and structure of orographic precipitation producing flash floods in the Lago Maggiore region, Q. J. Roy. Meteor. Soc., 141(686), 224–248, doi:10.1002/qj.2351, 2015.

- 5 Pellerin, B. A., Wollheim, W. M., Feng, X. and Vörösmarty, C. J.: The application of electrical conductivity as a tracer for hydrograph separation in urban catchments, Hydrol. Process., 22, 1810–1818, doi:10.1002/hyp.6786, 2008. Penna, D., Tromp-van Meerveld, H. J., Gobbi, A., Borga, M. and Dalla Fontana, G.: The influence of soil moisture on threshold runoff generation processes in an alpine headwater catchment, Hydrol. Earth Syst. Sci., 15(3), 689–702, doi:10.5194/hess-15-689-2011, 2011.
- 10 <u>Renard, K. G., Nichols, M. H., Woolhiser, D. A. and Osborn, H. B.: A brief background on the U.S. Department of Agriculture Agricultural Research Service Walnut Gulch Experimental Watershed, Water Resour. Res., 44(5), 1–11, doi:10.1029/2006WR005691, 2008.</u>

Reynolds, S. G.: The Gravimetric Method of Soil Moisture Determination Part III An Examination of Factors influencing Soil Moisture Variability, J. Hydrol., 11, 288 – 300, 1970.

Sivapalan, M.: Prediction in ungauged basins: a grand challenge for theoretical hydrology, Hydrol. Process., 17, 3163–3170, doi:10.1002/hyp.5155, 2003.

15

20

Slaughter, C. W., Marks, D., Flerchinger, G. N., Van Vactor, S. S. and Burgess, M.: Thirty-five years of research data collection at the Reynolds Creek Experimental Watershed, Idaho, United States, Water Resour. Res., 37(11), 2819–2823, doi:10.1029/2001WR000413, 2001.

Sloto, R. and Crouse, M. Y.: Hysep: a computer program for streamflow hydrograph separation and analysis, 1996.
 <u>Tramblay, Y., Bouvier, C., Martin, C., Didon-Lescot, J. F., Todorovik, D. and Domergue, J. M.: Assessment of initial soil moisture conditions for event-based rainfall-runoff modelling, J. Hydrol., 387(3–4), 176–187, doi:10.1016/j.jhydrol.2010.04.006, 2010.
</u>

25 Tramblay, Y., Bouaicha, R., Brocca, L., Dorigo, W., Bouvier, C., Camici, S. and Servat, E.: Estimation of antecedent wetness conditions for flood modelling in northern Morocco, Hydrol. Earth Syst. Sci., 16(11), 4375–4386, doi:10.5194/hess-16-4375-2012, 2012.

U.S. Geological Survey (USGS): hysep.R, GitHub Repository, https://github.com/USGS-R/DVstats/blob/master/R/hysep.R, 2015.

30 Vachaud, G., Passerat De Silans, A., Balabanis, P. and Vauclin, M.: Temporal Stability of Spatially Measured Soil Water Probability Density Function, Soil Sci. Soc. Am. J., 49(4), 822 – 828, doi:10.2136/sssaj1985.03615995004900040006x, 1985. Vanderlinden, K., Vereecken, H., Hardelauf, H., Herbst, M., Martínez, G., Cosh, M. H. and Pachepsky, Y. A: Temporal Stability of Soil Water Contents: A Review of Data and Analyses, Vadose Zone J., 11(4), doi:10.2136/vzj2011.0178, 2012. Vereecken, H., Huisman, J. A., Pachepsky, Y., Montzka, C., van der Kruk, J., Bogena, H., Weihermüller, L., Herbst, M., Martinez, G. and Vanderborght, J.: On the spatio-temporal dynamics of soil moisture at the field scale, J. Hydrol., 516, 76–96, doi:10.1016/j.jhydrol.2013.11.061, 2014.

Wood, E. F., Sivapalan, M. and Beven, K.: Similarity and Scale in Catchment Storm Response, Rev. Geophys., 28(1), 1–18, 1990.

Zehe, E. and Blöschl, G.: Predictability of hydrologic response at the plot and catchment scales: Role of initial conditions, Water Resour. Res., 40, 1–21, doi:10.1029/2003WR002869, 2004.

Zehe, E. and Sivapalan, M.: Threshold behavior in hydrological systems and geo-ecosystems: manifestations, controls and implications for predictability, Hydrol. Earth Syst. Sci., 13, 3247–3312, doi:10.5194/hessd-5-3247-2008, 2008.

Zehe, E., Becker, R., Bárdossy, A. and Plate, E.: Uncertainty of simulated catchment runoff response in the presence of 10 threshold Role of initial soil J. processes: moisture and precipitation, Hydrol., 315. 183 - 202,doi:10.1016/j.jhydrol.2005.03.038, 2005.

Zucco, G., Brocca, L., Moramarco, T. and Morbidelli, R.: Influence of land use on soil moisture spatial-temporal variability and monitoring, J. Hydrol., 516, 193–199, doi:10.1016/j.jhydrol.2014.01.043, 2014.

15





Figure 1: Location of the study site and measurement network. At every on-alert site measurements were taken at about 10 randomly chosen locations. At every continuous measurement sites two sensors were installed at 10 cm depth, two sensors at 20-25 cm depth and one sensor at a depth of 33-50 cm.



Figure 2: Hydrograph separation into baseflow and event flow of two different events with different methods: electric conductivity (EC), straight line (SL), constant K (CK), HySep filter 1 (HS1) and recursive digital filter (RDF). See the text



Figure 23: (a) Normalized histogram of soil moisture measurements within one grassland and one vineyard plot and (b) at the catchment scale within the land use classes grassland and vineyard during the first on alert measurement in 2012. Figure (c) shows the <u>rR</u>elationship between plot mean soil moisture $\overline{\theta}_j(t_{ev})$ and inner-plot standard deviation $\sigma_j^{\text{inner}}(t_{ev})$ as calculated with Eq. 1 and Eq. 6 in Table 1. Figure (db) shows the same at the catchment scale for catchment mean soil moisture $\overline{\overline{\theta}}_{ev}$ and inter-plot standard deviation $\sigma_{\text{cat.}}^{\text{inter}}(t_{ev})$ as calculated with Eq. 3 and Eq. 87 in Table 1. The right figures show initial (c) and final (d) soil moisture profile in plots of different land use during event 27 (06–09 September 2013).



Figure 4: Initial (a) and final (b) soil moisture profile in plots of different land use during event 27 (06 09 September 2013).



Figure 35: Temporal stability (Vachaud et al., 1985): the relative difference between the plot mean and the catchment mean $\delta_{j,ev}$ for each on-alert measurement and each plot where on-alert measurements were conducted (four vineyards v1 – v4, five grasslands g1 – g5, one fallow f1 and one cultivated field c1). Note that the time axis represents a sequence of events, no equidistant

time line. Blue squares show plots with plot means that exceed the catchment mean, red squares those with plot means below the catchment mean, white squares indicate plots that were not sampled during the respective measurement.



Figure <u>46</u>: Rainfall and soil moisture in autumn 2013 measured in 10, 25 and 40 cm depth. The line represents the catchment mean <u>calculated with Eq. 3 in Table 1</u> in the respective depth and the shaded area the mean $\pm SEM_{cat.}^{inter}$. The dashed lines represent the onsets of several rain events and the labels refer to the event numbers as in Table 3 and Table 4. Note that on 27 November 2013 one of the probes stopped working which is the reason for the step in the data and the higher $SEM_{cat.}^{inter}$.



Figure 57: Soil moisture in the topsoil (0–5 cm) measured on alert basis before and after major rain events in autumn seasons in 2012–2015. The points show catchment mean values, the lines the range of the mean $\pm SEM_{cat.}^{inter}$. The numbers above selected events give the event number as in Table 3 and Table 4 and as referred to in the text.



Figure <u>86</u>: (a) Differences of event-based runoff coefficients ϕ_{ev} calculated for 54 rain events in the Gazel catchment in autumn 2012 and autumn 2013 derived with different methods for hydrograph separation. (b) Correlation between different methods. The upper panel gives the coefficient of determination (R^2) and Spearman's rank correlation coefficient ρ , the dashed line in

5 the lower panel is the line of identity. The methods used for hydrograph separation are described in Sec. 2.5.2: constant-k (CK), electric conductivity (EC), Hysep filter with fixed or sliding interval (HS1–HS2), Hysep filter with local minima algorithm (HS3), recursive digital filter (RDF) and straight line (SL).



Figure 79: Correlation of three variables describing meteorological forcing with event-based runoff coefficients ϕ_{ev} of the Gazel catchment calculated with the recursive digital filter method. The lines represent the uncertainty associated to the

5 hydrograph separation method (gray vertical lines: range of ϕ_{ev} calculated with the seven different methods) and the stagedischarge relation (black vertical lines: range between ϕ_{ev} calculated with the 5 % and the 95 % confidence interval of discharge obtained

with the BaRatin framework). The point labels give the numbers of selected events as in Table 3 and Table 4 and described in the text.



Figure <u>8</u>+ Θ : Hyetographs, hydrographs of total discharge and evolution of soil moisture in the Gazel catchment during four different events in 2013. The event-based runoff coefficient ϕ_{ev} is also given for all events. The representation of soil moisture gives the mean $\pm SEM_{cat.}^{inter}$ in the respective depth and the one of discharge the best estimate \pm the uncertainty of the stage-discharge relation.



Figure 449: Relationship between initial soil moisture $\tilde{\theta}_{ini}$ and event-based runoff coefficients ϕ_{ev} in the Gazel (a) and Claduègne (b) catchments. On the x-axis the point represents the profile mean initial soil moisture and the horizontal line the range of the mean $\pm SEM_{cat.}^{inter}$. On the y-axis the point represents ϕ_{ev} calculated with the recursive digital filter method and the line the uncertainty as in Fig. 97. The color of the points indicates whether cumulative precipitation is low ($P_{cum} < 1.5 \text{ mm}$),

medium (1.5 mm $< P_{cum} < 13$ mm) or high ($P_{cum} > 13$ mm).



Figure 12: Streamflow signal of electric conductivity (*EC*) during two events in the Gazel catchment and hydrograph separation into surface and subsurface flow with the electric conductivity method. The maximum signal in streamflow *EC*, ΔEC_{max} that is calculated as the maximum of the difference between streamflow *EC*, EC_{tot} and interpolated baseflow *EC*, EC_{b} differs substantially between events.



Figure 13: Log-log scatter plot of the maximum electric conductivity signal ΔEC_{max} of rain events in the Gazel catchment in 2013 and 2012 (gray dots: no data for $\tilde{\theta}_{max}$ available) versus peak discharge Q_{p} , initial soil moisture $\tilde{\theta}_{max}$, and cumulative Precipitation P_{cum} .

Table 1: Calculated measures of averaged soil moisture and its variability at different scales. a) Thickness of layers 1-3 is assumed to be 175, 150 and 275 mm. b) only calculated for grasslands and vineyards, because the number of plots is 1 for land use classes fallow and cultivated field.

$\begin{array}{c} \begin{array}{c} \text{Plot mean} \\ \underline{\text{soil moisture}} \\ \underline{\text{soil moisture}} \\ \underline{\text{soil moisture}} \\ \hline \overline{\theta}_{1}(t_{\text{ev}}) = \frac{1}{n_{i}} \sum_{t=1}^{n_{i}} \theta\left(x_{1,j}, t_{\text{ev}}\right) \\ \underline{\text{Land use}} \\ \underline{\text{mean soil}} \\ \underline{\text{moisture}} \\ \hline \overline{\theta}_{1u}(t_{\text{ev}}) = \frac{1}{n_{j}} \sum_{i_{u}}^{n_{j}} \overline{\theta}_{1}(t_{\text{ev}}) \\ \hline \overline{\theta}_{1}(t_{\text{ev}}) \\ \hline \overline{\theta}_{1}(t_{\text{ev}}) = \frac{1}{n_{j}} \sum_{i_{u}}^{n_{j}} \overline{\theta}_{1}(t_{\text{ev}}) \\ \hline \overline{\theta}_{1}(t_{v}) \\ \hline \overline{\theta}_{1}$	Name	Formula		Purpose and Abbreviations
$\begin{array}{llllllllllllllllllllllllllllllllllll$	Plot mean		(1)	Best estimate at plot scale;
$\frac{ \operatorname{Land use} }{ \operatorname{mean soil} } = \overline{\theta}_{1u}(t_{ev}) = \frac{1}{n_{j_{1u}}} \sum_{j_{1u}=1}^{n_{j_{1u}}} \overline{\theta}_{j}(t_{ev})$ $\frac{ \operatorname{Land use} }{ \operatorname{moisture} } = \overline{\theta}_{1u}(t_{ev}) = \frac{1}{n_{j_{1u}}} \sum_{j_{1u}=1}^{n_{j_{1u}}} \overline{\theta}_{j}(t_{ev})$ $\frac{ \operatorname{Catchment} }{ \operatorname{mean soil} } = \overline{\theta}_{ev} = \frac{1}{n_{p}} \sum_{j_{1u}}^{n_{c_{1u}}} \overline{\theta}_{i_{1u}} \cdot n_{j_{1u}}}$ $\frac{ \operatorname{Catchment} }{ \operatorname{mean soil} } = \overline{\theta}_{ev} = \frac{1}{n_{p}} \sum_{j_{1u}}^{n_{c_{1u}}} \overline{\theta}_{i_{1u}} \cdot n_{j_{1u}}} $ $\frac{ \operatorname{Catchment} }{ \operatorname{mean soil} } = \overline{\theta}_{ev} = \frac{1}{n_{p}} \sum_{j_{1u}}^{n_{c_{1u}}} \overline{\theta}_{i_{1u}} \cdot n_{j_{1u}}}$ $\frac{ \operatorname{Catchment} }{ \operatorname{mean soil} } = \overline{\theta}_{ev} = \frac{1}{n_{p}} \sum_{j_{1u}}^{n_{c_{1u}}} \overline{\theta}_{i_{1u}} \cdot n_{j_{1u}}} $ $\frac{ \operatorname{Catchment} }{ \operatorname{mean soil} } = \overline{\theta}_{ev} = \frac{1}{n_{p}} \sum_{j_{1u}}^{n_{c_{1u}}} \overline{\theta}_{i_{1u}} \cdot n_{j_{1u}}} $ $\frac{ \operatorname{Catchment} }{ \operatorname{Soil moisture}} = \overline{\theta}_{ev} = \frac{\sum_{j_{1u}=1}^{n_{1}} \overline{\theta}_{ev_{1}} \cdot m_{1}}{\sum_{j_{1u}=1}^{n_{1}} n_{1}} $ $\frac{ \operatorname{Catchment} }{ \operatorname{Soil moisture}} = \delta_{ev} = \frac{\sum_{j=1}^{n_{1}} \overline{\theta}_{i_{1u}} - \overline{\theta}_{i_{1u},j}} \\ \operatorname{Catchment} = \overline{\theta}_{ev_{1}} \cdot m_{1}} $ $\frac{ \operatorname{Catchment} }{ \operatorname{Soil moisture}} = \delta_{ev} = \sum_{j=1}^{n_{1}} \frac{1}{100} (\overline{\theta}_{fin,1} - \overline{\theta}_{in,j}) \cdot m_{1} $ $\frac{ \operatorname{Catchment} }{ \operatorname{Catchment} } = \delta_{ev} = \sum_{j=1}^{n_{1}} \frac{1}{100} (\overline{\theta}_{fin,1} - \overline{\theta}_{in,j}) \cdot m_{1} $ $\frac{ \operatorname{Catchment} }{ \operatorname{Catchment} } = \delta_{ev} = \sum_{j=1}^{n_{1}} \frac{1}{100} (\overline{\theta}_{fin,1} - \overline{\theta}_{in,j}) \cdot m_{1} $ $\frac{ \operatorname{Catchment} }{ \operatorname{Catchment} } = \delta_{ev} = \sum_{j=1}^{n_{1}} \frac{1}{100} (\overline{\theta}_{fin,1} - \overline{\theta}_{in,j}) \cdot m_{1} $ $\frac{ \operatorname{Catchment} }{ \operatorname{Catchment} } = \frac{ \operatorname{Catchment} }{ \operatorname{Catchment} } = \sqrt{ \operatorname{Catchment} } = \sum_{j=1}^{n_{1}} \operatorname{Catchment} } = \sqrt{ \operatorname{Catchment} } = \frac{ \operatorname{Catchment} }{ \operatorname{Catchment} } = \sqrt{ \operatorname{Catchment} }$	soil moisture	$\overline{\theta}_{i}(t_{ev}) = \frac{1}{2} \sum \theta(x_{i,i}, t_{ev})$		n _i : Number of measurements in
$\begin{array}{llllllllllllllllllllllllllllllllllll$		$n_i \sum_{i=1}^{n} n_i \sum_{i=1}^{n} n_i n_i n_i n_i n_i n_i n_i n_i n_i n_i$		<u>plot j.</u>
$\begin{array}{llllllllllllllllllllllllllllllllllll$	Land use	$-$ 1 $\frac{n_{j_{lu}}}{n_{j_{lu}}}$	<u>(2)</u>	Best estimate for land use classes;
$\begin{array}{c} \begin{array}{c} \mbox{moisture} & n_{\rm hu}_{\rm hu$	mean soil	$\overline{\theta}_{\rm lu}(t_{\rm ev}) = \frac{1}{2} \sum \overline{\theta}_{\rm i}(t_{\rm ev})$		n _{jlu} : Number of meas. in plots of
$\begin{array}{llllllllllllllllllllllllllllllllllll$	moisture	$n_{j_{lu}} \sum_{j_{lu}=1}^{j_{lu}} $		land use <i>lu</i> ; g: grassland, v:
$\begin{array}{llllllllllllllllllllllllllllllllllll$				vineyard, c: cultivated field, f:
$\frac{\text{Catchment}}{\text{mean soil}} = \frac{1}{n_p} \sum_{l_u}^{\infty_{l_u}} \overline{\overline{\theta}}_{l_u} \cdot n_{j_{l_u}} $ $\frac{\text{Formational construction}}{p_{ev}} = \frac{1}{n_p} \sum_{l_u}^{n_u} \overline{\overline{\theta}}_{l_u} \cdot n_{j_{l_u}} $ $\frac{\text{Formational construction}}{p_{ev}} = \frac{1}{n_p} \sum_{l_{u}=1}^{n_u} \overline{\overline{\theta}}_{ev,l} \cdot m_l}{p_{u}} $ $\frac{1}{n_{u}} \sum_{l_{u}=1}^{n_u} \overline{m_l} \sum_{l_{u}=1}^{n_u} \overline{m_l} $ $\frac{1}{n_{u}} \sum_{l_{u}=1}^{n_u} \overline{m_l} \sum_{l_{u}=1}^{n_u} \overline{m_l} \sum_{l_{u}=1}^{n_u} \overline{m_l} $ $\frac{1}{n_{u}} \sum_{l_{u}=1}^{n_u} \overline{m_l} \sum_{l_{u}=$	Catalanaat	n_{c}	(2)	fallow
$\frac{\text{Inter stort}}{\text{moisture}} \qquad \theta_{\text{ev}} = \frac{1}{n_{\text{p}}} \sum_{lu}^{n} \theta_{lu} \cdot n_{l u}} \qquad \qquad h_{\text{p}} \cdot \text{Number of land use classes} \\ n_{\text{clu}} : \text{Number of land use classes} \\ n_{\text{clu}} : \text{Number of land use classes} \\ n_{\text{clu}} = 4) \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{\overline{\theta}}{\theta_{\text{ev}l}} \cdot m_{l}} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} m_{l} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{1}{100} \left(\overline{\overline{\theta}}_{\text{fin,l}} - \overline{\overline{\theta}}_{\text{in,l}} \right) \cdot m_{l} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{1}{100} \left(\overline{\overline{\theta}}_{\text{fin,l}} - \overline{\overline{\theta}}_{\text{in,l}} \right) \cdot m_{l} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{1}{100} \left(\overline{\overline{\theta}}_{\text{fin,l}} - \overline{\overline{\theta}}_{\text{in,l}} \right) \cdot m_{l} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{1}{100} \left(\overline{\overline{\theta}}_{\text{fin,l}} - \overline{\overline{\theta}}_{\text{in,l}} \right) \cdot m_{l} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{1}{100} \left(\overline{\overline{\theta}}_{\text{fin,l}} - \overline{\overline{\theta}}_{\text{in,l}} \right) \cdot m_{l} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{1}{100} \left(\overline{\overline{\theta}}_{\text{fin,l}} - \overline{\overline{\theta}}_{\text{in,l}} \right) \cdot m_{l} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{1}{100} \left(\overline{\overline{\theta}}_{\text{fin,l}} - \overline{\overline{\theta}}_{\text{in,l}} \right) \cdot m_{l} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{1}{100} \left(\overline{\overline{\theta}}_{\text{fin,l}} - \overline{\overline{\theta}}_{\text{in,l}} \right) \cdot m_{l} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{1}{100} \left(\overline{\overline{\theta}}_{\text{fin,l}} - \overline{\overline{\theta}}_{\text{in,l}} \right) \cdot m_{l} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{1}{100} \left(\overline{\overline{\theta}}_{\text{fin,l}} - \overline{\overline{\theta}}_{\text{in,l}} \right) \cdot m_{l} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{1}{100} \left(\overline{\overline{\theta}}_{\text{fin,l}} - \overline{\overline{\theta}}_{\text{in,l}} \right) \cdot m_{l} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{1}{100} \left(\overline{\overline{\theta}}_{\text{fin,l}} - \overline{\overline{\theta}}_{\text{in,l}} \right) \cdot m_{l} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{1}{100} \left(\overline{\overline{\theta}}_{\text{fin,l}} - \overline{\overline{\theta}}_{\text{in,l}} \right) \cdot m_{l} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{1}{100} \left(\overline{\theta}_{l} (t_{ev}) - \overline{\overline{\theta}}_{l} (t_{ev}) \right)^{2} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{1}{100} \sum_{l=1}^{n_{l}} \frac{1}{100} \left(\overline{\theta}_{l} (t_{ev}) - \overline{\theta}_{l} (t_{ev}) \right)^{2} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{1}{100} \sum_{l=1}^{n_{l}} \frac{1}{100} \sum_{l=1}^{n_{l}} \frac{1}{100} \left(\overline{\theta}_{l} (t_{ev}) - \overline{\theta}_{l} (t_{ev}) \right)^{2} \\ \frac{1}{2} \sum_{l=1}^{n_{l}} \frac{1}{100} \sum_{l=1}$	<u>Catchment</u>	$\equiv 1 \sum_{n=1}^{n \leq l_{n}} \equiv$	<u>(3)</u>	Best estimate at catchment scale; $m \in N_{\text{turber of plots}}(m = 11)$;
$\frac{\text{Intrustate}}{\text{solute}} = \frac{\sum_{i=1}^{n_{1}} \overline{\overline{\theta}}_{ev}}{1u} + \frac{\sum_{i=1}^{n_{1}} \overline{\overline{\theta}}_{ev}}{1u} + \frac{m_{1}}{\sum_{i=1}^{n_{1}} m_{1}}} = \frac{\pi_{i_{1}} \overline{\overline{\theta}}_{ev}}{1u} + \frac{m_{1}}{\sum_{i=1}^{n_{1}} m_{1}}} = \frac{\pi_{i_{1}} \overline{\overline{\theta}}_{ev}}{1u} + \frac{m_{1}}{\sum_{i=1}^{n_{1}} m_{1}}} = \frac{\pi_{i_{1}} \overline{\overline{\theta}}_{ev}}{1u} + \frac{\pi_{i_{1}}}{\sum_{i=1}^{n_{1}} m_{1}}} = \frac{\pi_{i_{1}}}{100} \left(\overline{\overline{\theta}}_{fin,1} - \overline{\overline{\theta}}_{in,1}\right) + m_{1}} = \frac{\pi_{i_{1}}}{100} \left(\overline{\overline{\theta}}_{i_{1}} + \overline{\overline{\theta}}_{i_{1}} + \overline{\overline{\theta}}_{i_{1}}\right) + m_{1}} = \frac{\pi_{i_{1}}}{100} \left(\overline{\overline{\theta}}_{i_{1}} + \overline{\overline{\theta}}_{i_{1}}\right) + \pi_{i_{1}}} = \frac{\pi_{i_{1}}}{100} \left(\overline{\overline{\theta}}_{i_{1}} + \overline{\overline{\theta}}_{i_{1}}\right) + \pi_{i_{1}}} = \frac{\pi_{i_{1}}}{100} \left(\overline{\overline{\theta}}_{i_{1}} + \overline{\overline{\theta}}_{i_{1}}\right) + \pi_{i_{1}}} = \frac{\pi_{i_{1}}}{100} \left(\overline{\overline{\theta}}_{i_{1}} + \overline{\overline{\theta}}_{i_{1}} + \overline{\overline{\theta}}_{i_{1}}\right) + \pi_{i_{1}}} = \frac{\pi_{i_{1}}}{100} \left(\overline{\overline{\theta}}_{i_{1}} + \overline{\overline{\theta}}_{i_{1}} + \overline{\overline{\theta}}_{i_{1}}\right) + \pi_{i_{1}}} = \frac{\pi_{i_{1}}}{100} \left(\overline{\overline{\theta}}_{i_{1}$	moisture	$\theta_{\rm ev} = \frac{1}{n_{\rm ev}} \sum_{\rm l} \theta_{\rm lu} \cdot n_{\rm j_{\rm lu}}$		$n_{\rm p}$: Number of plots ($n_{\rm p} = 11$);
Profile mean soil moisture soil moisture soil moisture storage change $\Delta S = \sum_{l=1}^{n_1} \frac{1}{100} \left(\overline{\overline{\theta}}_{\text{fin},l} - \overline{\overline{\theta}}_{\text{ini},l} \right) \cdot m_l$ (4) Best estimate at catchment scale, integrated over the soil profile; $n_{l,i}$ number of layers $(n_l = 3)$; $m_{l,i}$ thickness of layer P_i . (5) Soil water retention during events; $\overline{\overline{\theta}}_{\text{fin},l,i} = \overline{\overline{\theta}}_{\text{ini},l,i}$ final and initial soil moisture in layer l . (6) Estimate of spatial variability at the plot scale. (7) Estimate of spatial variability at the catchment scale. (6) Estimate of spatial variability at the plot scale. (7) Estimate of spatial variability at the catchment scale. (8) Estimate of inter plot variability in the grasslands and vineyards. (9) Estimate variability between the grasslands and vineyards.	moisture	lu lu		n_{clu} : Number of fand use classes
$\begin{array}{llllllllllllllllllllllllllllllllllll$	D ("1	_		$(n_{c_{lu}} = 4)$
$\frac{\text{Soft moisture}}{\text{Soft moisture}} \hat{\theta}_{ev} = \frac{\Delta_{i=1}^{n_{i=1}} - E_{i=1}^{n_{i}}}{\Sigma_{i=1}^{n_{i}} m_{i}} \qquad $	Profile mean	$\sum_{i=1}^{n_1} \overline{\overline{\theta}}_{i+1} \cdot m_1$	<u>(4)</u>	Best estimate at catchment scale,
$\sum_{l=1}^{n_l} m_l \qquad \qquad$	son moisture	$\ddot{\theta}_{\rm ev} = \frac{\omega_{\rm l=1} v_{\rm ev, \rm l}}{\Sigma^{n_{\rm l}}}$		integrated over the soli profile, $n_{\rm l}$:
$\frac{\text{Soil moisture}}{\frac{\text{storage}}{\text{change}}} \Delta S = \sum_{i=1}^{n_{1}} \frac{1}{100} \left(\overline{\overline{\theta}}_{\text{fin,l}} - \overline{\overline{\theta}}_{\text{ini,l}} \right) \cdot m_{1} $ $\frac{\text{Inner plot sd}}{\sigma_{j}^{\text{inner}}(t_{\text{ev}})} = \sqrt{\frac{1}{n_{i} - 1} \sum_{i=1}^{n_{i}} (\theta(x_{i,j}, t_{\text{ev}}) - \overline{\theta}_{j}(t_{\text{ev}}))^{2}} $ $\frac{\text{Inter plot sd}}{(\text{catchment})} \sigma_{\text{cat.}}^{\text{inter}}(t_{\text{ev}}) = \sqrt{\frac{1}{n_{p} - 1} \sum_{j=1}^{n_{p}} (\overline{\theta}_{j}(t_{\text{ev}}) - \overline{\overline{\theta}}_{j}(t_{\text{ev}}))^{2}} $ $\frac{\text{Inter plot sd}}{(g/v)^{b}} \sigma_{lu}^{\text{inter}}(t_{\text{ev}}) = \sqrt{\frac{1}{n_{j} - 1} \sum_{j=1}^{n_{j}} (\overline{\theta}_{j}(t_{\text{ev}}) - \overline{\overline{\theta}}_{lu}(t_{\text{ev}}))^{2}} $ $\frac{\text{Between land}}{use sd} \sigma^{\text{betw}}(t_{\text{ev}}) = \left[\frac{1}{n_{p} - 1} \sum_{j=1}^{n_{i}} (\overline{\theta}_{j}(t_{\text{ev}}) - \overline{\overline{\theta}}_{\text{ev}})^{2}} \right] $ $\frac{(5)}{(6)} \frac{\text{Soil water retended during events;}}{(6)} \frac{\text{Estimate of spatial variability at the plot soll water retended during events;}}{(7)} \frac{\text{Estimate of spatial variability at the catchment scale.}} $ $\frac{(7)}{(7)} \frac{\text{Estimate of spatial variability at the catchment scale.}}{(8)} \frac{\text{Estimate of inter plot variability in the grasslands and vineyards.}} $		$\Delta_{l=1}m_l$		thickness of laver J^{a}
$\Delta S = \sum_{l=1}^{n} \frac{1}{100} \left(\overline{\overline{\theta}}_{in,l} - \overline{\overline{\theta}}_{in,l} \right) \cdot m_{l}$ $\frac{1}{100} \left(\overline{\overline{\theta}}_{in,l} - \overline{\overline{\theta}}_{in,l} \right) \cdot m_{l$	Soil moisture	n_1 $()$	(5)	Soil water retention during events:
$\frac{change}{change} = \sum_{i=1}^{n} 100 \left(-\min_{i} - \min_{i} \right)^{-1} \qquad \qquad$	storage	$\Delta S = \sum \frac{1}{1} \left(\overline{\overline{\theta}}_{\text{fin}1} - \overline{\overline{\theta}}_{\text{ini}1} \right) \cdot m_1$	<u>(-)</u>	Ξ Ξ
$\frac{\text{Inner plot sd}}{\left[\frac{1}{n_{i}-1}\sum_{i=1}^{n_{i}}(\theta(x_{i,j},t_{ev})-\overline{\theta}_{j}(t_{ev}))^{2}\right]} \qquad (6) \qquad \frac{\text{moisture in layer } I}{\text{Estimate of spatial variability at the plot scale.}}$ $\frac{\text{Inter plot sd}}{\left[\frac{(\text{catchment})}{(\text{catchment})}\right]} \qquad \sigma_{\text{cat.}}^{\text{inter}}(t_{ev}) = \sqrt{\frac{1}{n_{p}-1}\sum_{j=1}^{n_{p}}(\overline{\theta}_{j}(t_{ev})-\overline{\overline{\theta}}_{l}(t_{ev}))^{2}} \qquad (7) \qquad \frac{\text{Estimate of spatial variability at the catchment scale.}}{(1)}$ $\frac{\text{Inter plot sd}}{\left[\frac{(g/v)^{b}}{(g/v)^{b}}\right]} \qquad \sigma_{\text{lu}}^{\text{inter}}(t_{ev}) = \sqrt{\frac{1}{n_{j u}-1}\sum_{j=1}^{n_{j u}}(\overline{\theta}_{j}(t_{ev})-\overline{\overline{\theta}}_{lu}(t_{ev}))^{2}}; \qquad (8) \qquad \frac{\text{Estimate of inter plot variability in the grasslands and vineyards.}}{\sigma_{\text{lu}}^{\text{betw}}(t_{ev})} = \sqrt{\frac{1}{n_{j u}-1}\sum_{j=1}^{n_{c}}(\overline{\theta}_{lu}(t_{ev})-\overline{\overline{\theta}}_{lu}(t_{ev}))^{2}; \qquad (9) \qquad \frac{\text{Estimate variability between the grasslands and vineyards.}}{\sigma_{\text{stand}}^{\text{betw}}(t_{ev})} = \sqrt{\frac{1}{n_{p}-1}\sum_{j=1}^{n_{c}}(\overline{\theta}_{lu}(t_{ev})-\overline{\overline{\theta}}_{ev})^{2}} \qquad (9) \qquad \frac{\text{Estimate variability between the grasslands and vineyards.}}{\sigma_{\text{stand}}^{\text{betw}}(t_{ev})} = \sqrt{\frac{1}{n_{p}-1}\sum_{j=1}^{n_{c}}(\overline{\theta}_{lu}(t_{ev})-\overline{\overline{\theta}}_{ev})^{2}} \qquad (9) \qquad \frac{\text{Estimate variability between the grasslands and vineyards.}}{\sigma_{\text{stand}}^{\text{betw}}(t_{ev})} = \sqrt{\frac{1}{n_{p}-1}\sum_{j=1}^{n_{c}}(\overline{\theta}_{lu}(t_{ev})-\overline{\theta}_{ev})^{2}} \qquad (9) \qquad \frac{\text{Estimate variability between the grasslands and vineyards.}}{\sigma_{\text{stand}}^{\text{betw}}(t_{ev})} = \sqrt{\frac{1}{n_{p}-1}\sum_{j=1}^{n_{c}}(\overline{\theta}_{lu}(t_{ev})-\overline{\theta}_{ev})^{2}}} \qquad (1) \qquad$	change	$\sum_{l=1}^{l} 100 \left(\sum_{l=1}^{l} \frac{100}{l} \right)$		$\theta_{\text{fin},l}$; $\theta_{\text{ini},l}$: final and initial soil
$\frac{\text{Inner plot sd}}{\sigma_{j}^{\text{inner}}(t_{ev})} = \sqrt{\frac{1}{n_{i}-1} \sum_{i=1}^{n_{i}} (\theta(x_{i,j}, t_{ev}) - \overline{\theta}_{j}(t_{ev}))^{2}} $ $\frac{\text{Inter plot sd}}{\frac{(\text{catchment})}{(\text{catchment})}} \sigma_{\text{cat.}}^{\text{inter}}(t_{ev}) = \sqrt{\frac{1}{n_{p}-1} \sum_{j=1}^{n_{p}} (\overline{\theta}_{j}(t_{ev}) - \overline{\overline{\theta}}_{l}(t_{ev}))^{2}} $ $\frac{\text{Inter plot sd}}{\frac{(g/v)^{b}}{(g/v)^{b}}} \sigma_{\text{lu}}^{\text{inter}}(t_{ev}) = \sqrt{\frac{1}{n_{ju}-1} \sum_{ju=1}^{n_{ju}} (\overline{\theta}_{j}(t_{ev}) - \overline{\overline{\theta}}_{lu}(t_{ev}))^{2}} $ $\frac{\text{Between land}}{\frac{\text{use sd}}{(g_{ev})}} \sigma_{\text{betw}}^{\text{inter}}(t_{ev}) = \sqrt{\frac{1}{n_{p}-1} \sum_{ju=1}^{n_{c}} (\overline{\theta}_{j}(t_{ev}) - \overline{\overline{\theta}}_{ev})^{2}} $ $\frac{(6)}{(g/v)^{b}} \frac{\text{Estimate of spatial variability at the catchment scale.}}{(7)} \frac{\text{Estimate of spatial variability at the catchment scale.}}{(8)} \frac{\text{Estimate of inter plot variability in the grasslands and vineyards.}}{(9)} \frac{\text{Estimate of inter plot variability between the grasslands and vineyards.}}{(9)} \frac{\text{Estimate variability between the grasslands and vineyards.}}}{(9)} \frac{\text{Estimate variability between the grasslands and vineyards.}}}{(9)} \frac{\text{Estimate variability between the grasslands and vineyards.}}}{(9)} \frac{(9)}{(9)} \frac{\text{Estimate variability between the grasslands and vineyards.}}{(9)} \frac{(9)}{(9)} (9)$	x 1 . 1			<u>moisture in layer <i>l</i>.</u>
$\sigma_{j}^{\text{inner}}(t_{ev}) = \sqrt{\frac{1}{n_{i} - 1} \sum_{i=1}^{n} (\theta(x_{i,j}, t_{ev}) - \overline{\theta}_{j}(t_{ev}))^{2}} \qquad $	Inner plot sd	$\frac{n_i}{1}$	<u>(6)</u>	Estimate of spatial variability at the
$\frac{\text{Inter plot sd}}{(\text{catchment})} \qquad $		$\sigma_{\rm i}^{\rm inner}(t_{\rm ev}) = \left \frac{1}{1} \sum \left(\theta(x_{\rm i}, t_{\rm ev}) - \overline{\theta}_{\rm i}(t_{\rm ev}) \right)^2 \right ^2$		plot scale.
$\frac{\text{Inter plot sd}}{(\text{catchment})} \sigma_{\text{cat.}}^{\text{inter}}(t_{\text{ev}}) = \sqrt{\frac{1}{n_{\text{p}} - 1} \sum_{j=1}^{n_{\text{p}}} (\overline{\theta}_{j}(t_{\text{ev}}) - \overline{\overline{\theta}}(t_{\text{ev}}))^{2}} \frac{(7)}{2} \frac{\text{Estimate of spatial variability at the catchment scale.}}{(7)}$ $\frac{\text{Inter plot sd}}{(g/v)^{b}} \sigma_{\text{lu}}^{\text{inter}}(t_{\text{ev}}) = \sqrt{\frac{1}{n_{j u} - 1} \sum_{j=1}^{n_{j u}} (\overline{\theta}_{j}(t_{\text{ev}}) - \overline{\overline{\theta}}_{lu}(t_{\text{ev}}))^{2}}; \frac{(8)}{2} \frac{\text{Estimate of inter plot variability in the grasslands and vineyards.}}{(1)}$ $\frac{\text{Between land}}{(use sd)} \sigma_{\text{betw}}(t_{\text{ev}}) = \sqrt{\frac{1}{n_{j u} - 1} \sum_{j=1}^{n_{c u}} (\overline{\overline{\theta}}_{lu}(t_{\text{ev}}) - \overline{\overline{\theta}}_{\text{ev}})^{2}} \frac{(9)}{2} \frac{\text{Estimate variability between the grasslands and vineyards.}}{(1)}$		$n_i - 1 \sum_{i=1}^{n_i} (n_i - 1) \sum_{i=1}^{n_i} (n_i -$		
$\frac{(\text{catchment})}{(\text{catchment})} \sigma_{\text{cat.}}^{\text{inter}}(t_{\text{ev}}) = \sqrt{\frac{1}{n_{\text{p}} - 1} \sum_{j=1}^{n_{\text{p}}} (\overline{\theta}_{j}(t_{\text{ev}}) - \overline{\overline{\theta}}(t_{\text{ev}}))^{2}} (\overline{\theta}_{j}(t_{\text{ev}}))^{2} (\overline{\theta}_{j}(t_{\text{ev}}))^{2}} (\overline{\theta}_{j}(t_{\text{ev}}))^{2} (\overline{\theta}_{j}(t_{\text{ev}}))^{2}; (\overline{\theta}_{j}(t_{\text{ev}}))^{2}; $	Inter plot sd		(7)	Estimate of spatial variability at the
$\sigma_{\text{cat.}}^{\text{inter}}(t_{\text{ev}}) = \sqrt{\frac{n_{\text{p}} - 1}{\sum_{j=1}^{j=1}^{j}} (\theta_{j}(t_{\text{ev}}) - \theta(t_{\text{ev}}))^{2}}$ $\frac{\text{Inter plot sd}}{(\underline{g/v})^{b}} \qquad \sigma_{\text{lu}}^{\text{inter}}(t_{\text{ev}}) = \sqrt{\frac{1}{n_{j_{\text{lu}}} - 1} \sum_{j_{\text{lu}}=1}^{n_{j_{\text{lu}}}} (\overline{\theta}_{j}(t_{\text{ev}}) - \overline{\overline{\theta}}_{l_{u}}(t_{\text{ev}}))^{2}}; \qquad \frac{(8)}{\underline{\text{Estimate of inter plot variability in the grasslands and vineyards.}}}$ $\frac{\text{Between land}}{\underline{\text{use sd}}} \qquad \sigma^{\text{betw}}(t_{\text{ev}}) = \sqrt{\frac{1}{n_{1}} \sum_{j=1}^{n_{c_{\text{lu}}}} (\overline{\overline{\theta}}_{l_{u}}(t_{\text{ev}}) - \overline{\overline{\theta}}_{\text{ev}})^{2}} \qquad \frac{(9)}{\underline{\text{Estimate variability between the grasslands and vineyards.}}}$	(catchment)	$1 \sum_{i=1}^{n_p} = 1$	<u>, , , , , , , , , , , , , , , , , , , </u>	catchment scale.
$\frac{\text{Inter plot sd}}{(\underline{g/v})^{b}} \qquad \sigma_{lu}^{(\operatorname{inter}}(t_{ev}) = \sqrt{\frac{1}{n_{j_{lu}} - 1} \sum_{j_{lu}=1}^{n_{j_{lu}}} (\overline{\theta}_{j}(t_{ev}) - \overline{\overline{\theta}}_{lu}(t_{ev}))^{2}}; \qquad \frac{(8)}{\underline{\text{Estimate of inter plot variability in}}} \\ \frac{Between land}{\underline{\text{use sd}}} \\ \sigma^{\text{betw}}(t_{ev}) = \sqrt{\frac{1}{n_{j_{lu}} - 1} \sum_{j_{lu}=1}^{n_{c_{lu}}} (\overline{\overline{\theta}}_{lu}(t_{ev}) - \overline{\overline{\overline{\theta}}}_{ev})^{2}} \qquad \frac{(9)}{\underline{\text{Estimate variability between the}}} \\ \frac{1}{\underline{\text{grasslands and vineyards.}}} \\ \frac{1}{n_{j_{lu}} - 1} \sum_{j_{lu}=1}^{n_{c_{lu}}} (\overline{\overline{\theta}}_{lu}(t_{ev}) - \overline{\overline{\overline{\theta}}}_{ev})^{2}} \qquad \frac{(9)}{\underline{\text{Estimate variability between the}}} \\ \frac{1}{\underline{\text{grasslands and vineyards.}}} \\ \frac{1}{n_{j_{lu}} - 1} \sum_{j_{lu}=1}^{n_{c_{lu}}} (\overline{\overline{\theta}}_{lu}(t_{ev}) - \overline{\overline{\theta}}_{ev})^{2}} \\ \frac{1}{\underline{\text{grasslands and vineyards.}}} \\ \frac{1}{\underline{\text{grasslands and vineyards.}} \\ \frac{1}{\underline{\text{grasslands and vineyards.}}} \\ \frac{1}{\underline{\text{grasslands and vineyards.}}} \\ \frac{1}{\underline{\text{grasslands and vineyards.}} \\ $		$\sigma_{\text{cat.}}^{\text{inter}}(t_{\text{ev}}) = \left \frac{1}{n_{\text{ev}} - 1} \sum_{i=1}^{n_{\text{ev}}} (\theta_{i}(t_{\text{ev}}) - \theta(t_{\text{ev}}))^{2} \right $		
$\frac{\text{Inter plot sd}}{(\underline{g/v})^{b}} \qquad \sigma_{lu}^{\text{inter}}(t_{ev}) = \sqrt{\frac{1}{n_{j_{lu}} - 1} \sum_{j_{lu}=1}^{n_{j_{lu}}} (\overline{\theta}_{j}(t_{ev}) - \overline{\overline{\theta}}_{lu}(t_{ev}))^{2}}; \qquad \frac{(8)}{\underline{\text{Estimate of inter plot variability in}}} \\ \frac{Between land}{\underline{\text{use sd}}} \\ \sigma^{\text{betw}}(t_{ev}) = \sqrt{\frac{1}{n_{-1}} \sum_{j_{lu}=1}^{n_{c_{lu}}} (\overline{\overline{\theta}}_{j}(t_{ev}) - \overline{\overline{\theta}}_{ev})^{2}} \qquad \frac{(9)}{\underline{\text{Estimate variability between the}}} \\ \frac{(9)}{\underline{\text{grasslands and vineyards.}}} \\ \frac{(9)}{\underline{\text{grasslands and vineyards.}} \\ \frac{(9)}{\underline{\text{grasslands and vineyards.}}} \\ \frac{(9)}{\underline{\text{grasslands and vineyards.}} \\ \frac{(9)}{\underline{\text{grasslands and vineyards.}}} \\ \frac{(9)}{\underline{\text{grasslands and vineyards.}} \\ \frac{(9)}{\text{gra$		$\sqrt{\frac{1}{j=1}}$		
$\frac{(g/v)^{b}}{use sd} \qquad \sigma_{lu}^{inter}(t_{ev}) = \sqrt{\frac{1}{n_{j_{lu}} - 1} \sum_{j_{lu}=1}^{\infty} (\overline{\theta}_{j}(t_{ev}) - \overline{\overline{\theta}}_{lu}(t_{ev}))^{2}}; \qquad \frac{the grasslands and vineyards.}{(9)}$ $\frac{Between land}{use sd} \qquad \sigma^{betw}(t_{ev}) = \sqrt{\frac{1}{n_{-1}} \sum_{j=1}^{n_{c_{lu}}} (\overline{\overline{\theta}}_{lu}(t_{ev}) - \overline{\overline{\theta}}_{ev})^{2}} \qquad \frac{(9)}{grasslands and vineyards.}$	Inter plot sd	njlu	<u>(8)</u>	Estimate of inter plot variability in
$\frac{Between \ land}{use \ sd} \qquad \sigma^{betw}(t_{ev}) = \sqrt{n_{j_{lu}} - 1 \sum_{j_{lu}=1}^{n} (\overline{\theta}_{lu}(t_{ev}) - \overline{\theta}_{ev})^2} \qquad \qquad (9) \qquad \underbrace{Estimate \ variability \ between \ the}_{grasslands \ and \ vineyards.}$	$(g/v)^{b}$	$\sigma^{\text{inter}}(t) = \left \frac{1}{\sqrt{2}} \sum_{i=1}^{\infty} (\overline{\theta}_i(t_i)) - \overline{\theta}_i(t_i) \right ^2$		the grasslands and vineyards.
$\frac{\text{Between land}}{\text{use sd}} \sigma^{\text{betw}}(t_{\text{ev}}) = \sqrt{\frac{1}{n_{c_{\text{lu}}} - 1}} \sum_{i=1}^{n_{c_{\text{lu}}}} (\overline{\overline{\theta}}_{iu}(t_{\text{ev}}) - \overline{\overline{\theta}}_{\text{ev}})^2} \qquad \qquad$		$n_{j_{1}u} - 1 \sum_{i_{1}=1}^{i_{1}} (v_{j}(v_{ev}) - v_{j_{1}}(v_{ev})),$		
$\frac{1}{use sd} \sigma^{\text{betw}}(t_{\text{ev}}) = \left \frac{1}{n - 1} \sum_{i=1}^{n_{c_{\text{lu}}}} (\overline{\overline{\theta}}_{lu}(t_{\text{ev}}) - \overline{\overline{\theta}}_{\text{ev}})^2 \right \frac{1}{\frac{1}{2} \text{summer variability between the}} \frac{1}{\frac{1}{2} \text{grasslands and vineyards.}} \right $	Ratwaan land	$\sqrt{\int J_{lu}=1}$	(0)	Estimate variability between the
$\sigma^{\text{betw}}(t_{\text{ev}}) = \left \frac{1}{n - 1} \sum_{v=1}^{\infty} (\overline{\theta}_{\text{lu}}(t_{\text{ev}}) - \overline{\theta}_{\text{ev}})^2 \right ^2$	use sd	$1 \xrightarrow{n_{c_{lu}}} - =$	<u>(9)</u>	grasslands and vinevards
n = 17	<u>use su</u>	$\sigma^{\text{betw}}(t_{\text{ev}}) = \left \frac{1}{1} \sum (\overline{\theta}_{\text{lu}}(t_{\text{ev}}) - \overline{\theta}_{\text{ev}})^2 \right $		grussiands and vincydrus.
$\sqrt{n_{c_{lu}} - 1}$		$\sqrt{n_{c_{lu}}-1}$		
		1		

Table 24: Spatial variability of soil moisture at the plot scale (mean of all events calculated for all plots: mean $\sigma_j^{\text{inner}}(t_{\text{ev}})$, for the grassland plots: mean $\sigma_{j\in g}^{\text{inner}}(t_{\text{ev}})$ and the vineyard plots: mean $\sigma_{j\in v}^{\text{inner}}(t_{\text{ev}})$ calculated with Eq. 6 in Table 1) and at the catchment scale (mean inter plot variability of the grassland plots: mean $\sigma_g^{\text{inter}}(t_{\text{ev}})$ and the vineyard plots mean $\sigma_v^{\text{inter}}(t_{\text{ev}})$ calculated with Eq. 8 in Table 17 as well as between land use class variability mean $\sigma^{\text{betw.}}(t_{\text{ev}})$ calculated with Eq. 7 in Table 18) determined at different depth with the two measuring schemes.

	Initial stat	es			Final state	Final states				
	0-5 cm	10 cm	25 cm	40 cm	0-5 cm	10 cm	25 cm	40 cm		
Mean $\sigma_{\rm i}^{\rm inner}(t_{\rm ev})$	2.62	2.77	1.85	NA	2.91	2.71	1.88	NA		
Mean $\sigma_{ieg}^{inner}(t_{ev})$	2.77	2.18	1.77	NA	2.80	2.18	1.78	NA		
Mean $\sigma_{iev}^{inner}(t_{ev})$	2.49	3.15	2.24	NA	2.84	3.15	2.24	NA		
Mean $\sigma_{\rm g}^{\rm inter}(t_{\rm ev})$	3.48	4.49	2.07	4.40	3.70	4.26	2.04	4.30		
Mean $\sigma_{\rm v}^{\rm inter}(t_{\rm ev})$	2.63	2.26	5.36	6.81	2.11	2.39	5.31	6.65		
Mean $\sigma^{\text{betw.}}(t_{\text{ev}})$	2.12	2.20	1.14	1.71	3.78	1.98	1.10	1.55		

Table 2: Differences between land use classes: results of Student t-tests under significance level $\alpha = 0.05$. The upper panel gives the ratio of the number of rejections of the null hypothesis (true difference in means is equal to zero) to the total number of tests conducted. The lower panel states whether the found differences are systematic (T) or not (F)

	Grassland	Vineyard	Cultivated	Fallow
Grassland		15/26	17/24	12/17
Vineyard	F		11/24	8 / 17
Cultivated	Ŧ	F		4/17
Fallow	F	F	F	

Table 3: Rainfall, soil moisture and discharge characteristics of selected rain events in autumn 2013 in the Gazel catchment: beginning of the rain event, cumulative precipitation (P_{cum}), maximum 20-min rain intensity ($I_{max,20}$), mean intensity (I_{μ}), initial soi moisture ($\tilde{\theta}_{ini}$), final soil moisture ($\tilde{\theta}_{fin}$), standard error of the catchment mean ($SEM_{cat.}^{inter}$) during initial and final stage, soil storage change at depth 0 – 50 cm (ΔS), peak discharge (Q_p), cumulative total discharge ($Q_{tot,cum}$), cumulative event discharge ($Q_{ev,cum}$) and₇ event-based runoff coefficient calculated with the recursive digital filter method (ϕ_{ev}) and maximum signal in electric conductivity (ΔEC_{max}).

	Rainfall				Soil M	oisture				Discharge		
Ev.	Beg. Rain	$P_{\rm cum}$	I _{max,20}	I_{μ}	$ ilde{ heta}_{ m ini}$	$SEM_{cat.}^{inter}$	$ ilde{ heta}_{ m fin}$	$SEM_{cat.}^{inter}$	ΔS	$Q_{ m p}$	$Q_{ m ev,cum}$	ϕ_{ev}
#	DD-MM hh:mm	mm	mm h ⁻¹	mm h ⁻¹	vol%	vol%	vol%	vol%	mm	1 s ⁻¹	mm	-
27	06-09 16:06	41.59	33.30	0.85	27.99	1.73	32.69	1.07	28.17	25	0.34	0.01
28	15-09 00:23	4.82	24.00	0.60	31.40	1.31	31.34	1.38	0.00	14	0.20	0.04
30	28-09 17:09	71.73	59.70	2.39	29.80	1.57	35.20	1.45	32.43	324	1.73	0.02
34	04-10 15:39	27.20	25.50	3.89	34.33	1.29	36.43	1.25	12.62	58	1.03	0.04
36	12-10 05:12	12.86	4.50	0.71	34.55	1.15	35.91	1.02	8.12	18	0.07	0.01
37	15-10 03:46	11.29	9.00	0.81	35.76	1.07	36.54	1.07	4.72	23	0.21	0.02
39	20-10 02:41	82.20	53.40	3.04	35.65	1.10	38.12	1.41	14.83	8660	27.49	0.33
40	23-10 01:01	39.31	60.00	1.79	37.34	1.44	38.40	1.54	6.33	30096	38.91	0.99
41	27-10 03:37	14.64	37.20	0.73	37.17	1.46	38.09	1.52	5.54	361	2.07	0.14
49	18-12 06:52	33.09	8.70	0.57	34.26	1.20	37.90	1.32	21.82	402	5.30	0.16
50	21-12 03:11	8.73	4.20	0.27	38.39	1.23	37.78	1.12	0.00	175	0.95	0.11
52	24-12 01:15	46.33	7.50	0.99	37.09	1.17	39.83	1.08	16.46	1047	12.38	0.27
53	28-12 04:51	18.70	32.70	1.04	37.52	1.40	39.25	1.04	10.41	654	4.25	0.23

	Rainfall				Soil M	oisture				Discharge		
Ev.	Beg. Rain	P _{cum}	I _{max,20}	I_{μ}	$ ilde{ heta}_{ m ini}$	$SEM_{cat.}^{inter}$	$ ilde{ heta}_{ m fin}$	$SEM_{cat.}^{inter}$	ΔS	$Q_{ m p}$	$Q_{ m ev,cum}$	ϕ_{ev}
#	DD-MM hh:mm	mm	$mm h^{-1}$	$mm h^{-1}$	vol%	vol%	vol%	vol%	mm	$m^3 s^{-1}$	mm	-
27	06-09 15:41	43.26	33.3	0.88	27.00	1.38	32.62	1.16	29.05	0.19	0.38	0.01
28	15-09 00:23	3.61	26.7	0.45	31.09	1.32	31.00	1.32	0.00	NA	NA	NA
30	28-09 17:04	77.71	63	2.59	29.09	1.49	34.49	1.24	29.93	16.39	3.16	0.04
34	04-10 15:33	28.09	27	4.01	33.48	1.53	35.46	1.44	11.39	0.59	0.66	0.02
36	12-10 05:12	12.81	4.5	0.71	33.58	1.53	35.20	1.46	8.42	0.13	0.10	0.01
37	15-10 03:46	11.51	9	0.82	34.98	1.55	35.67	1.52	4.34	0.16	0.19	0.02
39	20-10 02:41	83.67	53.4	3.10	34.83	1.63	37.54	1.51	15.33	54.64	20.82	0.25
40	23-10 01:01	51.01	60	2.32	36.92	1.69	38.75	1.60	11.37	60.76	36.37	0.93
41	27-10 03:37	22.06	52.5	1.10	37.62	1.67	38.47	1.64	4.40	12.08	5.44	0.37
49	18-12 06:52	67.91	8.7	1.17	34.40	1.61	38.22	1.44	21.59	NA	NA	NA
50	21-12 03:11	10.98	4.2	0.34	38.70	1.61	38.40	1.63	0.00	NA	NA	NA
52	24-12 01:15	51.24	7.5	1.09	37.70	1.68	40.29	1.58	14.30	NA	NA	NA
53	28-12 04:51	19.65	32.7	1.09	38.09	1.66	39.68	1.60	9.56	NA	NA	NA

Table 4: Rainfall, soil moisture and discharge characteristics of selected rain events in autumn 2013 in the Claduègne catchment, abbreviations as in Table 3.