

Interactive comment on “Evapotranspiration monitoring based on thermal infrared data over agricultural landscapes: comparison of a simple energy budget model and a SVAT model” by Guillaume Bigeard et al.

Anonymous Referee #2

Received and published: 18 December 2018

General comments:

The paper explores the application of the Two-Source Energy Balance (TSEB) using TIR data and a SVAT model (SEtHyS) over experimental agricultural sites in Morocco and France, hence different climate and management practices. With regards to application of TSEB, in the model description, it appears they are using most if not all of the original formulations of the Norman et al (1995) model, for example Eq. (9) for partitioning net radiation (R_n) for the soil and canopy elements. However later they state that they adopt a more physically-based R_n divergence model of Kustas and

[Printer-friendly version](#)

[Discussion paper](#)



Norman (1999). Yet in the sensitivity analysis (Table 1) this extinction coefficient for Eq. (9) is retained and evaluated later in Figure 9 which is not consistent with what is stated in the text. While in the references they appear to cite papers that have included new formulations being implemented in the TSEB since the original 1995 paper, they are not included in this paper. The TSEB model has undergone several modifications since it was first presented by Norman et al. (1995). Changes include refinements to the algorithm estimating soil aerodynamic resistance and shortwave and longwave transmittance through the canopy (as they mention in their paper; Kustas and Norman, 2000) and additionally a means for adjusting the Priestley–Taylor formulation for canopy transpiration (Kustas and Norman 1999). Further improvements include incorporating rigorous treatment of radiation modeling for strongly clumped row crops, accounting for shading effects on soil heat flux (Colaizzi et al. 2012a, 2016a,b), and incorporating alternative formulations for computing the canopy transpiration such as Penman–Monteith (PM) or light-use efficiency (LUE) parameterizations (see Colaizzi et al. 2012b, 2014, 2016c; Anderson et al. 2008). The later two canopy transpiration formulations are mentioned but not applied in this paper. Alternatively, the SETHyS is a SVAT model with 22 parameters and so it is unclear why such a comparison is actually being made between a relatively simple but fairly robust thermal-based model and a SVAT having a large number of tunable parameters. It's also unclear why this comparison does not include application of a newly developed and presumably more robust two-source model SPARSE developed by one of the co-authors (Boulet et al., 2015). Additionally, for the sensitivity analysis, the authors do not appear to be aware of the several studies that have already performed sensitivity analyses for key inputs to TSEB. These include two of the papers mentioned in this manuscript. . . Timmermans et al (2007) and Zhan et al. (1996). There is also Li et al (2005) mentioned in the manuscript and then there is the paper by Kustas and Norman (1997) and Kustas et al. (2012). In summary it appears they conduct an analysis with a dated TSEB model without some of the more current refinements and comparing it to a SVAT that has a number of tunable parameters that would be difficult to prescribe over a large area

[Printer-friendly version](#)

[Discussion paper](#)



without detailed ground information. There are a significant number of analyses performed making it a long paper and is somewhat diffuse in its focus. While I think the paper has some unique findings, it does not consider some of the main advances in TSEB when evaluating model performance for these agricultural sites. Early season conditions when the canopy is small, the soil is playing a major role in the energy exchange, and there is no discussion of soil roughness effects on the TSEB formulation that has been discussed in the literature (Kustas et al., 2016). Errors in TSEB during senescence will largely depend on how well the green fraction is determined. . . however it should be pointed out that these later stages of vegetation condition are not as important to capture the ET as during the main growing season. While I consider this work as having some merit, particularly the analyses performed with SETHyS, it seems the authors do not consider to any degree of the advances/refinements made in the TSEB model since Norman et al (1995) and therefore I question how relevant is their analyses and conclusions using the 20+ year old formulations evaluated here in comparison to the more current parameterizations. Based on these shortcomings I do not find the paper suitable for publication in its current form.

Specific comments:

Page 9: It appears the leaf area and green fraction data are very local and may not reflect conditions viewed by the radiometer. This can be a major issue. Is there any indication where they sampled is representative of the radiometer field of view?

Page 9: Eq (15). What values are assumed in the Penman-Monteith equation for computing LE_{pot} ?

Page 10: How is the calibration of SETHyS carried out and what level of calibration is shown in Figure 2 for the SETHyS model?

Page 10: So the TSEB performance is “sought in its out-of-the box configuration presented in Norman et al (1995)” suggests none of the refinements over the last 20 years are incorporated in this analysis.

Page 10. The 3 parameters identified for study are the Priestley-Taylor coefficient, the net radiation extinction parameter and the fraction of soil net radiation for estimating soil heat flux, G . There is some interdependency here between the amount of canopy net radiation interception and the value of the Priestley-Taylor parameter (Kustas and Norman, 2000). Also for G , refinements of the TSEB include time varying formulation proposed by Santanello and Friedl (2003).

Page 12 line (10): TSEB could be provided albedo inputs from remote sensing. This is something easily done in the model if made available.

Page 12 (line 15): The authors do not seem to be aware of the soil resistance formulation that is sensitive to soil roughness which is discussed in refinements to the TSEB model (Kustas et al., 2016).

Page 12 (Line 30): Its unclear what version of SEtHyS model (1-4 from page 10) is being used in these comparisons.

Page 13 (line 5): The Crow et al (2008) paper actually showed the utility of TSEB in providing an indicator of plant stress for assimilation in a water balance model.

Page 15 Sensitivity analysis to meteorological inputs: It has been long recognized that to apply TSEB regionally requires a way of reducing the need for accurate absolute surface-air temperature differences. This was the motivation for the development of time differencing modeling schemes (Anderson et al., 1997; Norman et al., 2000).

Page 15 Sensitivity analysis to vegetation forcing inputs: The use of micrometeorological measurements close to the canopy height is ill-advised in general due to roughness sublayer effects and so comes as no surprise for the TSEB since the aerodynamic resistances are key to the TSEB calculations. This should be removed

Page 17: Sensitivity analysis to radiative temperature for TSEB: This is well documented and the reason why time differences in radiative temperatures were developed early in the TSEB applications (see Anderson et al., 2004)

[Printer-friendly version](#)

[Discussion paper](#)



Page 17-18: sensitivity analysis to water inputs and soil water content for SEtHyS: This is a major issue with SVAT models. That is why approaches like Crow et al (2008) of combining water balance with remote sensing energy balance is appealing. Moreover, for regional analysis it will be very difficult to acquire irrigation information in a timely manner.

Page 22 (figure 9): These results are related to some extent on the radiation partitioning which the authors appear to have adopted the original formulation of Norman et al (1995) for net radiation extinction and without any clumping effects which row crops tend to have (Anderson et al., 2005).

Page 25 (figure 11): Did the authors consider the fact that extinction of diffuse light through a canopy is quite different from direct and perhaps that is another factor affecting the Priestley-Taylor value?

References:

Anderson MC, Norman JM, Diak GR, Kustas WP, Mecikalski JR. (1997) A two-source time-integrated model for estimating surface fluxes using thermal infrared remote sensing. *Remote Sens Environ* 60:195–216.

Anderson, M.C., Norman, J.M., Mecikalski, J.R., Torn, R.D., Kustas, W.P., & Basara, J.B. (2004). A multi-scale remote sensing model for disaggregating regional fluxes to micrometeorological scales. *J. Hydromet*, 5, 343–363.

Anderson MC, Norman JM, Kustas WP, Li F, Prueger JH, Mecikalski JR (2005) Effects of vegetation clumping on two-source model estimates of surface energy fluxes from an agricultural landscape during SMACEX. *J Hydromet* 6(6):892–909

Anderson MC, Norman JM, Kustas WP, Houborg JM, Starks PJ, Agam N (2008) thermal-based remote sensing technique for routine mapping of land-surface carbon, water and energy fluxes from field to regional scales. *Remote Sens Environ* 112:4227–4241

[Printer-friendly version](#)

[Discussion paper](#)



Boulet, G., Mougenot, B., Lhomme, J.-P., Fanise, P., Lili-Chabaane, Z., Olioso, A., Bahir, M., Rivalland, V., Jarlan, L., Merlin, O., Coudert, B., Er-Raki, S., and Lagouarde, J.-P. (2015) The SPARSE model for the prediction of water stress and evapotranspiration components from thermal infra-red data and its evaluation over irrigated and rainfed wheat, *Hydro Earth Sys. Sci.* 19:4653–4672

Colaizzi PD, Evett SR, Howell TA, Li F, Kustas WP, Anderson MC (2012a) Radiation model for row crops: I. Geometric view factors and parameter optimization. *Agron J* 104:225–240

Colaizzi PD, Kustas WP, Anderson MC, Agam N, Tolk JA, Evett SR, Howell TA, Gowda PH, O’Shaughnessy SA (2012b) Two-source energy balance model estimates of evapotranspiration using component and composite surface temperatures. *Adv Water Resour* 50:134–151

Colaizzi PD, Agam N, Tolk JA, Evett SR, Howell TA, Gowda PH, O’Shaughnessy SA, Kustas WP, Anderson MC (2014) Two-source energy balance model to calculate E, T, and ET: comparison of Priestley–Taylor and Penman–Monteith formulations and two time scaling methods. *Trans ASABE* 57(2):479–498

Colaizzi PD, Evett SR, Agam N, Schwartz RC, Kustas WP (2016a) Soil heat flux calculation for sunlit and shaded surfaces under row crops: 1. Model development and sensitivity analysis. *Agric For Meteorol* 216:115–128

Colaizzi PD, Evett SR, Agam N, Schwartz RC, Kustas WP, Cosh MH, McKee LG (2016b) Soil heat flux calculation for sunlit and shaded surfaces under row crops: 2. Model test. *Agric For Meteorol* 216:129–140

Colaizzi PD, Agam N, Tolk JA, Evett SR, Howell TA, O’Shaughnessy SA, Gowda PH, Kustas WP, Anderson MC (2016c) Advances in a two-source energy balance model: partitioning of evaporation and transpiration for cotton using component and composite surface temperatures. *Trans ASABE* 59(1):181–197. <https://doi.org/10.13031/trans>

Printer-friendly version

Discussion paper



.59.11215

Crow, W. T., Kustas, W. P., and Prueger, J. H. (2008) Monitoring root-zone soil moisture through the assimilation of a thermal remote sensing-based soil moisture proxy into a water balance model, *Remote Sens. Environ.* 112:1268–1281.

Kustas WP, Norman JM. (1997) A two-source approach for estimating turbulent fluxes using multiple angle thermal infrared observations. *Water Resour Res* 33:1495–1508.

Kustas WP, Norman JM (1999) Evaluation of soil and vegetation heat flux predictions using a simple two-source model with radiometric temperatures for partial canopy cover. *Agric For Meteorol* 94:13–29

Kustas W, Norman JM (2000) A two-source energy balance approach using directional radiometric temperature observations for sparse canopy covered surfaces. *Agron J* 92(5):847–854

Kustas WP, Alfieri JG, Anderson MC, Colaizzi PD, Prueger JH, Evett SR, Neale CMU, French AN, Hipps LE, Chávez JL, Copeland KS, Howell TA. (2012) Application of a time differencing technique using local thermal observations in a strongly advective irrigated agricultural area. *Adv Water Resour* 50:120–33.

Kustas WP, Nieto H, Morillas L, Anderson MC, Alfieri JG, Hipps LE, Villagarcía L, Domingo F, García M (2016) Revisiting the paper “using radiometric surface temperature for surface energy flux estimation in mediterranean drylands from a two-source perspective. *Remote Sens Environ* 184:645–653

Li F, Kustas WP, Prueger JH, Neale CMU, Jackson TJ. (2005) Utility of remote sensing based two-source energy balance model under low and high vegetation cover conditions. *J Hydrometeorol* 6:878–91.

Norman JM, Kustas WP, Prueger JH, Diak GR. (2000) Surface flux estimation using radiometric temperature: a dual temperature difference method to minimize measurement error. *Water Resour Res* 36:2263–74.

HESSD

Interactive
comment

Printer-friendly version

Discussion paper



Santanello J Jr, Friedl M (2003) Diurnal covariation in soil heat flux and net radiation. *J Appl Meteorol* 42(6):851–862

Timmermans, WJ, Kustas WP, Anderson MC, French AN. (2007) An intercomparison of the surface energy balance algorithm for land (SEBAL) and the two-source energy balance (TSEB) modeling schemes. *Remote Sens Environ* 108:369–84.

Zhan, X., W. P. Kustas, and K. S. Humes, (1996) An intercomparison study on models of sensible heat flux over partial canopy surfaces with remotely sensed surface temperature. *Remote Sens. Environ.*, 58, 242–256.

Interactive comment on Hydrol. Earth Syst. Sci. Discuss., <https://doi.org/10.5194/hess-2018-295>, 2018.

Printer-friendly version

Discussion paper

