

The authors addressed the reviewers' comments well overall. The paper could, however, still gain from using a more precise terminology (examples below).

The authors thank Bettina Schaeffli for the positive comments and advice, which helped to improve the manuscript. Detailed responses to all comments are given below (blue).

Detailed comments:

- "Definition 1 allows assessing the annual impact of glaciers and snow cover (...)"'. Correct would probably be "(..) impact of glacier melt and snow melt".

The sentence was corrected to:

p.11 l.11-12: "Definition 1 allows assessing the annual impact of glacier melt and snow melt on the water production".

- Section 4.1.3 does still not indicate for which period the geodetic mass balances have been estimated; it is only in the figure legend; I suggest adding a note in the text

The periods for the geodetic mass balances were added to section 4.1.3:

p.18 l.1-5: "Figure 10 compares the simulated mean annual glacier mass balances obtained with the different model configurations with mass balances determined with geodetic methods between 1999 and 2015 (Bolch et al., 2012; Gardelle et al., 2013; Nuimura et al., 2015; King et al., 2016; Brun et al., 2017). These geodetic mass balances range from -0.67 ± 0.45 m w.e.yr⁻¹ (2000-2008) (Nuimura et al., 2015) to -0.32 ± 0.09 m w.e.yr⁻¹ (2000-2015) (Brun et al., 2017)."

- Even if your definitions are correct, I would still be as precise as possible everywhere. Thus the sentence should read "of the outflow as coming from water infiltrated in glaciers and more than 20% from subsurface and groundwater flows (NEW:) generated outside the glacier covered area"

The sentence was modified as suggested (P.20 l.21)

- Figure 13: I would remove the discharge from the top figure since the figure mixes inputs (melt etc.) and outputs (streamflow) and it wrongly implies that you can label how much water comes from melt at a certain time step (which your approach cannot)

We modified Figure 13 and removed the discharge from the top panel.

- Terminology:

- "Total production of runoff" is misleading since this is not water that creates runoff but the water available in the system (the definition is correct but I suggest changing the terminology).

"On average, we find that the outflow is mainly produced by meltwater (...)"'; this certainly holds but in fact, you assess the input from these sources, not the outflow. The same problem repeats in the conclusion: "major contributions from glaciers and snow (...) of the annual outflow produced by ice melt".

Figure 14: here it is extremely obvious that a renaming of “runoff production” is necessary (see later): someone who reads figure 14, necessarily thinks that you assessed how much runoff comes from melt and rain. Why not call it “Total water input”?

We changed the terminology and replaced “runoff production” by “water input” in the text, tables and figures.

For example, p.20 l.16-17: “On average, we find that the outflow is mainly produced by meltwater as 46 % of the annual water input is due to ice melt and 41 % to snow melt (definition 1).”

- V stands usually for Volume; it is however a flux. I suggest changing the symbol and give the units [L/T]

Equation 8 should read: $d\text{Storage}/dt = V(t) - Q(t) - ET(t)$, only if dS/dt (in the catchment) is zero, we have $Q(t) = V(t) - ET(t)$ (ET is subtracted not added). This avoids saying that dS/dt might not be zero. Writing all previous equations as storage change = input – output would probably avoid confusion

Thank you for pointing out this error in the equation. Section “3.2.5 Quantification of the flow components” was modified accordingly

- Italic is only used for variables, not for standard functions such as “max”; units are not in italic;

- Textual subscripts or superscripts should not be italic (e.g. x_{max} , T_{min} where “max” and “min” stand for maximum and minimum, respectively).

The equations of the sections 3.2.2 and 3.2.5 were modified to take into account the two previous remarks.

Detailed responses to all comments are given below (blue).

While the manuscript certainly has been improved, I am a bit disappointed that the authors focused rather on the ‘simple’ revisions than the more substantial ones. As stated in my previous review, I think that model changes and new computations would be needed. To repeat, my concerns regarding the snow redistribution remain:

- 1) Only avalanches as redistribution
- 2) Only cardinal directions

We agree that there are limitations regarding the snow redistribution in our model and we addressed the concerns of the reviewer already during the discussion phase. In short, we concede that a more sophisticated module has the potential to better reflect real conditions for the redistribution of snow. However, processes driving this redistribution like blowing snow require detailed input data concerning the spatially distributed wind direction and speed (Freudiger et al, 2017). Wind fields for the entire catchment derived with the available data (wind speed was measured at two stations with substantial gaps in the data series) would be prone to large and potentially systematic errors. Therefore, we decided to implement at this stage only a simplified module for the snow redistribution, which is also coherent with the current overland flow routing algorithm of DHSVM-GDM considering four directions (Wigmota et al., 1994). We are well aware of these limitations, which are, thus, discussed in detail in the section “5.2 Representation of the cryospheric processes in the model”.

- 3) Description in a rejected manuscript

We removed the citation from the manuscript.

- 1) The same applies to the issue of literature values, ad hoc decisions and ‘tricks’, such as ... My concern regarding lit values has not been addressed

The referee asked for a sensitivity analysis for the most sensitive model parameters. While the objective of this study was certainly not to perform a full sensitivity analysis of all model parameters for a well-established model like DHSVM, we performed sensitivity analyses concerning the newly implemented cryospheric processes concerning for example the glacier outline, the melt coefficient for debris-covered glaciers, and the avalanche parameters. Furthermore, the model sensitivity to the soil parameterization was analyzed. Due to the sparse vegetation and the dominating cryospheric processes in the study catchment we believe that these are the most parameters for a sensitivity analysis for our model set-up. The results of the sensitivity analysis are presented in chapter 5.3.

- 2) 2 m soil under glaciers is just unrealistic and I find it unfortunate that in a manuscript which aims at more realistic simulations such tricks are used to compensate for missing processes in the model. Honestly, I am also not fully happy by the uncertainty analyses.

It is unclear to us to why the referee refers to “tricks” concerning our manuscript. Obviously, any model is an imperfect representation of the real world. Is using a certain parameterization or certain values for model parameters a “trick”? Unfortunately, not all information one would like to have for the selected catchment is available. Nevertheless, we believe that doing simulations for such a region

brings new results for the scientific community and advances our understanding. If observations were not available we performed simulations with parameter values according to the best of our knowledge. These choices are described in the manuscript. While the referee obviously disagrees with (some of) our choices, we do not understand what justifies to qualify these choices as “tricks”. Concerning the soil depth we refer to the study of Pelletier et al. (Global 1-km Gridded Thickness of Soil, Regolith, and Sedimentary Deposit Layers. ORNL DAAC, Oak Ridge, Tennessee, USA.

<http://dx.doi.org/10.3334/ORNLDAA/1304>, 2016), which indicates for more than 50% of the Pheriche catchment area soil depths ranging between 1 and 2 m. While it remains unclear why the referee claims that a soil depth of 2 m, which is used for most model runs, is “unrealistic”, we agree that the soil depth is highly uncertain especially for areas with glaciers. Therefore, we tested the model sensitivity with respect to the soil depth. The results are summarized in chapter 5.3.2.

As a minor comment, I also find the refs to a thesis in French and a submitted manuscript not convincing.

These references are not used in the manuscript and were only used in our response to the referee in order to illustrate other aspects of our work concerning the characterization of the uncertainties in our study. Although written partly in French, the thesis has been evaluated and reviewed by a Ph.D. jury and is published here: <https://tel.archives-ouvertes.fr/tel-01863806>. The manuscript is currently under revision after submission to the Journal of Hydrometeorology in July 2018 (Mimeau et al., Impact of precipitation uncertainty on the simulated hydrological response of a small glacierized Himalayan catchment, JHM-D-18-0157).

Perhaps I am missing something, but I still do not fully understand how the flow components were tracked. Actually, I am now even more confused than before. Did the authors change things in the DHSVM code? If yes, what? If no, how can then the components be tracked with the standard DHSVM?

We only changed the code to have access to internally simulated variables, which are not routinely included in the output of the standard version of DHSVM. No changes were made to the calculation of these variables in the model.

Concerning definition 1, flow components were computed using the water balance equations 5 to 8described in section “3.2.5 Quantification of the flow components”. All variables used for the computation are part of standard output or internal variables of DHSVM.

Concerning the contributions estimated with definition 2, the method used for the computation was added in a supplementary material.

Quantification of different flow components in a high-altitude glacierized catchment (Dudh Koshi, Nepalese Himalaya)

Louise Mimeau¹, Michel Esteves¹, Isabella Zin¹, Hans-Werner Jacobi¹, Fanny Brun¹, Patrick Wagnon¹, Devesh Koirala², and Yves Arnaud¹

¹Université Grenoble Alpes, IRD, Grenoble INP, CNRS, IGE, Grenoble, France

²Nepal Academy of Science and Technology, NAST, Kathmandu, Nepal

Correspondence to: Michel Esteves (michel.esteves@ird.fr)

Abstract. In a context of climate change and water demand growth, understanding the origin of water flows in the Himalayas is a key issue for assessing the current and future water resources availability and planning the future uses of water in downstream regions. This study estimates the relative contributions of rainfall, glacier and snow melt to the Khumbu River streamflow (Upper Dudh Koshi, Nepal, 146 km², 43 % glacierized, elevation range from 4260 to 8848 m a.s.l.), as well as their seasonal,

5 daily, and sub-daily variability during the period 2012-2015, by using the physically based glacio-hydrological model DHSVM-GDM (Distributed Hydrological Soil Vegetation Model - Glaciers Dynamics Model). One of the main issues in high elevated and glacierized catchments hydrology is the limited representation of cryospheric processes, which control the evolution of ice and snow, in distributed hydrological models. Here, the impact of different snow and glacier parameterizations was tested by modifying the original DHSVM-GDM snow albedo parameterization, by adding an avalanche module, and by adding a
10 reduction factor for the melt of debris covered glaciers. The validation of the snow, glacier and hydrological processes was established using three types of validation data (MODIS images, glacier mass balances and in situ discharge measurements). Results show that ~~this the~~ new version of DHSVM-GDM improves the simulation of the ~~snow covered~~ snow-covered area and the glacier mass balances, ~~thus improving the reliability of the overall hydrological simulation.~~ The simulated hydrological contributions differ not only depending on the used models and implemented processes, but also due to different definitions of
15 the estimated flow components. Therefore, we apply here two different definitions based on the water inputs and contributing areas for the diverse flow components. In the presented case study, ~~results show that~~ ice and snow melt contribute each more than 40 % to the annual ~~outflow~~ water inputs and ~~that~~ 69 % of the outflow originates from glacierized areas. ~~Our simulations also highlight that winter flows are mainly controlled by the release from the englacial water storage.~~ The ~~However, the~~ choice of a given parameterization for ~~the~~ snow and glacier processes has a significant impact on the simulated water balance: the
20 different ~~parametrizations tested in this study lead to an ice melt contribution to the outflow~~ tested parameterizations led to ice melt contributions ranging from 45 to 70 %. ~~Our simulations also highlight that winter flows are mainly controlled by the release from the englacial water storage.~~ The sensitivity of the model to the ~~glaciers~~ glacier inventory was tested, demonstrating that the uncertainty related to the glacierized surface leads to an uncertainty of 20 % ~~on for~~ the simulated ice melt component.

1 Introduction

The Himalayan mountain range is known for being the water tower of Central and South Asia (Immerzeel et al., 2010). Its high elevated glaciers and snow cover play an important role in the regional hydrological system (Kaser et al., 2010; Racoviteanu et al., 2013) and provide water resources for the population living in the surrounding countries (Viviroli et al., 2007; Singh et al., 2016; Pritchard, 2017).

In the Hindu Kush-Himalaya (HKH) region climate change is expected to cause shrinkage of the snow and ice cover (Bolch et al., 2012; Benn et al., 2012; Kraaijenbrink et al., 2017). Changes in glacier and snow cover runoff are likely to have a significant impact on the hydrological regime (Akhtar et al., 2008; Immerzeel et al., 2012; Lutz et al., 2014; Nepal, 2016). Development of tourism is also affecting the accessibility to water during the peak of tourist season. In the Everest region in 10 Nepal water needs have increased within the past decades due to higher demand in water supply for tourists and hydroelectricity production, leading to water shortage during months with low flows (winter and spring) (McDowell et al., 2013). Understanding the past and present hydrological regime and more particularly estimating the seasonal contribution of ice melt, snow melt, and rainfall to outflows is thus a key issue for managing water resources within the next decades. Indeed, the quantification of the ice melt contribution enables to assess the proportion of water currently available which is coming from a long term 15 accumulation in the glaciers, and thus to assess the annual decrease of the basin water storage due to glacier melt. Moreover, knowing the fraction of snow melt, ice melt, and rainfall to the river outflow, and understanding their hydrological pathways can give insights into how much water is currently seasonally delayed and how the seasonal outflow and the overall water balance might be impacted in the future when this delay changes or if the ratio snowfall to rainfall changes (Berghuijs et al., 2014).

20 Recent studies have estimated present glacier and snow melt contributions to the outflow in Nepalese Himalayan catchments (e.g., Andermann et al., 2012; Savéan et al., 2015; Ragettli et al., 2015) and simulated future hydrological regimes using glacio-hydrological models (Rees and Collins, 2006; Nepal, 2016; Soncini et al., 2016). Results have demonstrated large differences in the estimates of the contribution of glaciers to the annual outflows of the Dudh Koshi catchment in Nepal, which range from 4 to 60 % (Andermann et al., 2012; Racoviteanu et al., 2013; Nepal et al., 2014; Savéan et al., 2015).

25 One of the main sources of uncertainty in modelling the outflow of Himalayan catchments is the representation of cryospheric processes, which control the evolution of ice and snow-covered surfaces in hydrological models. For instance, the representation of the debris covered glaciers in glacio-hydrological models is a challenge. Debris-covered glaciers represent about 23 % of all glaciers in the Himalaya-Karakoram region (Scheler et al., 2011). The debris layers have been expanding during the last decades due to the glacier recession (Shukla et al., 2009; Bhambri et al., 2011; Benn et al., 2012) and are expected to keep 30 expanding in the near future (Rowan et al., 2015). Since the study of Østrem (1959) it is known that the debris thickness has a strong impact on the meltwater generation, which means that a good representation of the debris-covered glaciers in glacio-hydrological models is essential for estimating the amount of meltwater generated in glacierized catchments in the Himalayas. Many other cryospheric processes such as the liquid water storage and transfer through glaciers, snow transport by avalanches or wind, glacial lake dynamics and snow albedo evolution are either very simplified or not at all represented by the models

(Chen et al., 2017). It is therefore important to estimate the impact of such simplified representations of cryospheric processes on modelling results.

Delineation of the glacierized areas is another key entry-element to the glacio-hydrological model. Glacier inventories are commonly used as forcing data to delineate glacierized areas in glacio-hydrological modelling studies. There are three global 5 major glacier inventories such as the World Glacier Inventory (Cogley, 2009), GlobGlacier (Paul et al., 2009) and the Randolph Glacier Inventory (Pfeffer et al., 2014), and several regional glacier inventories in the HKH region (ICIMOD (Bajracharya et al., 2010), Racoviteanu et al. (2013)), showing substantial differences. These can be due to the definition of the glacierized area 10 itself (Paul et al., 2013; Brun et al., 2017) as well as to the characteristics of the satellite image (date, resolution, spectral properties) used for the delineation (Kääb et al., 2015), and to difficulties related to the interpretation of satellite images for 15 outlying the glaciers, especially when they are debris covered (Bhambri et al., 2011; Racoviteanu et al., 2013; Robson et al., 2015). Thus, the question whether the glacier delineation has a significant impact on the model results needs to be addressed.

These issues of the representation of cryospheric processes and of glaciers delineation in the hydrological modelling are addressed in the present study by (i) adapting the parameterization of the snow albedo evolution of DHSVM-GDM, in order to improve the simulation of the snow cover dynamics; (ii) implementing an avalanche module; (iii) introducing a melting 15 factor for debris covered glaciers and (iv) testing the sensitivity of simulated outflows and flow components with respect to these modifications as well as to glacier delineation for three different outlines coming from different glacier inventories. Both in-situ measurements and satellite data were used for evaluating the outflow simulations as well as snow cover and glacier evolutions focusing on a small headwater catchment.

An uncertainty on the estimation of the glacier contribution also results from how the contributions to the outflow are 20 defined. There are indeed several ways to define the glacier contribution to runoff (Radić and Hock, 2014) : it can be either considered as the total outflow coming from glacierized areas, the outflow produced by the glacier itself (snow, firn and ice melt) or the outflow produced only by the ice melt. The definition of the glacial contribution is dependent to the hydrological model (distributed or lumped, representation of glaciers and snow in the model) and cannot always be chosen. In the Dudh Koshi basin, Andermann et al. (2012); Racoviteanu et al. (2013); Savéan et al. (2015) estimated the fraction of the outflow 25 produced by ice melt, whereas Nepal et al. (2014) defined the glacier contribution as the fraction of the outflow coming from glacierized areas. Here, flow components were estimated using two different definitions of the hydrological contributions ~~in order~~ to control and make the best evaluation of all the terms of the water balance. Finally, the model results are analyzed at the annual, monthly, daily and sub-daily scale in order to explain the origin of the water flows and their seasonal and daily variations.

30 2 Study area

This study focuses on the Pheriche sub-catchment of the Dudh Koshi basin (outlet at coordinates 27.89° N, 86.82° E) located in Nepal on the southern slopes of Mt. Everest in the Sagarmatha National Park (SNP) (Fig. 1). The catchment's area is 146 km² and its elevation extends from 4260 to 8848 m a.s.l.

Local climate is mainly controlled by the Indian summer monsoon (Bookhagen and Burbank, 2006) and is characterized by four different seasons: a cold dry winter from December to March with limited precipitation, a warm and moist summer with most of the annual precipitation occurring during the monsoon from June until September, and two transition seasons: the pre-monsoon season in April and May and the post-monsoon season in October and November (Shrestha et al., 2000). At 5000 m, the annual precipitation is around 600 mm and the mean monthly temperature ranges from -8.4°C in January to 3.5°C in July, according to temperature and precipitation data from the Pyramid EvK2 station (Fig. 2 and Table 1). The hydrological regime follows the precipitation cycle with high flows during the monsoon season, when most of the annual precipitation occurs, complemented by the melting of snow and ice, and low flows during winter.

Due to high elevation, the vegetation in the catchment is scarce. The basin area is mainly covered by rocks and moraines (43 %) (Bajracharya et al., 2010) and glaciers (43 %) (Racoviteanu et al., 2013). Only 14 % of the basin area is covered by grasslands and shrublands. Glaciers belong to the summer-accumulation type (Wagnon et al., 2013) and are partially fed by avalanches (King et al., 2016; Sherpa et al., 2017). In the Pheriche catchment, 60 % of the glaciers are located between 5000 and 6000 m a.s.l.. Debris-covered glaciers are found at low elevations mainly on the ablation tongues of the glaciers (Fig. 3). According to the Racoviteanu et al. (2013) glacier inventory, they represent 30 % of the glacierized area with smaller melting rates at similar elevations than debris-free glaciers due to the insulating effect of the debris layer (Vincent et al., 2016).




Figure 1. Study area : (a) Location map of Pheriche catchment (black) in the Sagarmatha National Park (green) in Nepal. Characteristics of the meteorological stations are summarized in Table 1. (b) Hypsometric curve of the Pheriche catchment.




Figure 2. Daily minimal and maximal air temperature and daily precipitation measured at the Pyramid station.

3 Data and model setup

10 3.1 Database

To describe the topography of the study area, an ASTER DEM originally at 30 m resolution was resampled to a 100 m resolution. The SOTER Nepal soil classification (Dijkshoorn and Huting, 2009) and a landcover classification from ICIMOD (Bajracharya, 2014) were used for the soil and landcover description.

Meteorological data were available at hourly time steps from three automatic weather stations (AWS) located at Pangboche (3950 m a.s.l.), Pheriche (4260 m a.s.l.) and Pyramid (5035 m a.s.l.) (Table 1). Since December 2012, the precipitation has been recorded at the Pheriche and Pyramid AWS by two Geonor T-200 sensors designed to measure both liquid and solid 15 precipitations. Data were corrected for potential undercatch following the method used by Lejeune et al. (2007) and Sherpa et al. (2017). Precipitation at the Pangboche station was recorded with a tipping bucket. Air temperature, wind speed, relative 20 humidity short-wave radiation and long-wave radiation measurements at Pheriche and Pyramid were provided by the EvK2-CNR stations.

Discharge measurements of the Khumbu River at Pheriche were obtained using a pressure water level sensor at a 30 minutes time step since October 2010.

Glacier outline	Area	Satellite imagery used for delineation	Acquisition dates	Spatial resolution of the satellite images used for delineation
Racoviteanu et al. (2013)	Dudh Koshi, Langtang	ASTER, IKONOS-2	2003-2008	1 - 90 m
GAMDAM (Nuimura et al., 2015)	Asian glaciers	SRTM, LANDSAT	1999-2003	30 - 120 m
ICIMOD (Bajracharya et al., 2010)	Nepal	IKONOS, LANDSAT, ASTER	1992-2006	1 - 120 m

Table 2. Glacier outlines characteristics

N°	Name	Elevation (m)	Lat (°)	Lon (°)	Measured parameters	Manager
1	Pangboche	3950	27.857	86.794	T, P	IRD
2	Pheriche	4260	27.895	86.819	T, P, WS, RH, SWin	EvK2-CNR, IRD
3	Pyramid	5035	27.959	86.813	T, P, WS, RH, SWin, SWout, LWin	EvK2-CNR, IRD
4	Changri Nup	5363	27.983	86.779	SWin, SWout	GLACIOCLIM

Table 1. Location of measurements. **T** air temperature, **P** precipitation, **WS** wind speed, **RH** relative humidity, **SWin** incoming shortwave radiation, **SWout** outgoing shortwave radiation, **LWin** incoming longwave radiation.

The MODImLab algorithm developed by Sirguey et al. (2009) was applied to MODIS reflectances data to obtain daily albedo and snow fraction satellite images for the period 2010-2015. We used the Sirguey et al. (2009) algorithm rather than 25 the MOD10A1 500 m snow products because it generates daily regional snow cover images at 250 m resolution and applies corrections on atmospheric and topographic effects which makes the snow cover maps more realistic on mountainous areas. 27 cloud free Landsat8 images were used to generate snow maps at 30 m resolution between 1 November 2014 and 31 December 2015. A NDSI (Normalized-Difference Snow Index) threshold of 0.15 was taken to separate snow free and snow covered pixels on Landsat8 data as proposed by Zhu and Woodcock (2012). Daily snow cover maps were then retrieved from the MODImLab 5 snow fraction product: areas with a snow fraction above 0.15 were defined as snow covered areas so that the MODImLab Snow cover area (SCA) matches the Landsat8 SCA on the common dates. For the rest of this study we call MODIS data albedo and snow cover data obtained with the MODImLab algorithm. We also used snow albedo data from in-situ measurements at Pyramid and Changri Nup (Table 1).

For describing the glacierized area in the basin we compared three different glacier outlines available as vector layers for the Khumbu region: the glacier delineation proposed by Racoviteanu et al. (2013) specifically set up for the Dudh Koshi basin; the GAMDAM inventory covering the entire Himalayan range (Nuimura et al., 2015); and the ICIMOD inventory (Bajracharya et al., 2010) (Fig. 3). The three outlines have been derived on different grids, from different datasets at different spatial resolutions and covering different temporal periods (see Table 2), thus leading at different results.

Mass balances estimated by Sherpa et al. (2017) for the clean-ice West Changri Nup and Pokalde glaciers located in the 5 Pheriche basin (Fig. 3) were used as reference, as well as mean annual glacier mass balances calculated over the Pheriche basin area for the period 2000-2016 by Brun et al. (2017).




Figure 3. Glacier outlines in the Pheriche catchment. (a) Clean glaciers and debris-covered glaciers from Racoviteanu et al. (2013) and location of the clean ice West Changri Nup and Pokalde glaciers (b) GAMDAM (red) and ICIMOD (blue) glacier outlines

3.2 Glacio-hydrological modelling

3.2.1 General description of the model

The glacio-hydrological model DHSVM-GDM (Distributed Hydrological Soil Vegetation Model - Glaciers Dynamics Model) 10 was used for simulating outflows at Pheriche. DHSVM is a physically based, spatially distributed model which was developed for mountain basins with rain and snow hydrological regimes (Wigmota et al., 1994; Nijssen et al., 1997; Beckers and Alila, 2004). A glacier dynamics module was recently implemented in DHSVM by Naz et al. (2014) to simulate glacier mass balance and the runoff production in catchments with glaciers, thus extending the application to ice dominated hydrological regimes. The resulting DHSVM-GDM simulates the spatial distribution and the temporal evolution of the principal water balance terms 15 (soil moisture, evapotranspiration, sublimation, glacier mass balance, snow cover, and runoff) at hourly to daily time scales. It uses a two-layer energy and mass balance module for simulating snow cover evolution and a single layer energy and mass balance module for glaciers (Andreadis et al., 2009; Naz et al., 2014) and has been applied in a number of studies for snow and cold regions hydrology (e.g., Leung et al., 1996; Leung and Wigmota, 1999; Westrick et al., 2002; Whitaker et al., 2003; Zhao et al., 2009; Bewley et al., 2010; Cristea et al., 2014; Frans et al., 2015). Distributed meteorological data (air temperature, 20 precipitation, relative humidity, wind speed, and shortwave and longwave incoming radiation) are requested as input, as well as distributed geographical information (elevation, soil type, landcover, soil depth, and ice thickness).

3.2.2 Snow albedo parameterization

In the original DHSVM-GDM version, the snow albedo α_s [-] is set to its maximum value $\alpha_{s\max}$ (to be fixed either by calibration or from observed albedo values), after a snowfall event and then decreases with time according to the following

25 equations (Wigmasta et al., 1994):

$$\begin{aligned}\alpha_s &= \alpha_{s\max} (\lambda_a)^{N^{0.58}} && \text{if } T_s < 0 \\ \alpha_s &= \alpha_{s\max} (\lambda_m)^{N^{0.46}} && \text{if } T_s \geq 0\end{aligned}\quad (1)$$

Where N is the number of days since the last snowfall, λ_a [-] and λ_m [-] correspond to 0.92 and 0.70 for the accumulation season and the melt season, respectively, and T_s is the snow surface temperature [$^{\circ}\text{C}$].

MODIS albedo images and the albedo measurements from Pyramid and Changri Nup were used to analyse the decrease of snow albedo with age in various locations of our study area. Figure 4 compares the observed albedo decay as a function of time for snow events with at least three consecutive days without clouds after the snowfall with the albedo parameterization in DHSVM-GDM. Since the observed values are not well represented by the standard albedo decrease, the parameterization was replaced by Eq. 2, with a decay of the albedo when there is no new snowfall inspired by the ISBA model albedo

5 parameterization (Douville et al., 1995) and with the fresh snow albedo modified as a function of the amount of snowfall:

$$\begin{aligned}\alpha_s &= (\alpha_{st-1} - \alpha_{s\min}) \exp(-cN) + \alpha_{s\min} && \text{if } i_{\text{snowfall}} = 0 \text{ mm/h} \\ \alpha_s &= \max(0.6, \alpha_{st-1}) && \text{if } 0 \text{ mm/h} < i_{\text{snowfall}} \leq 1 \text{ mm/h} \\ \alpha_s &= \max(0.6, \alpha_{st-1}) + (\alpha_{s\max} - \max(0.6, \alpha_{st-1})) \frac{i_{\text{snowfall}} - 1}{3 - 1} && \text{if } 1 \text{ mm/h} < i_{\text{snowfall}} \leq 3 \text{ mm/h} \\ \alpha_s &= \alpha_{s\max} && \text{if } i_{\text{snowfall}} > 3 \text{ mm/h}\end{aligned}\quad (2)$$

Where α_{st-1} is the albedo from the previous time step, $\alpha_{s\min}$ is the minimal snow albedo of 0.3 (estimated using the mean minimal albedo values observed at the station and on MODIS images), N is the number of days since the last 10 snowfall, c is the coefficient of the exponential decrease [days^{-1}], and i_{snowfall} the snowfall intensity [mm/h]. Since the observed decrease is dependent on elevation, the coefficient c is calculated as a function of elevation according to Eq. 3:

$$c = 20 \exp(-0.001 Z) \quad (3)$$

Where Z is the elevation of the cell in m a.s.l.

The new function for the decrease of the snow albedo is also shown in Fig. 4.




Figure 4. Original and modified parameterization of the snow albedo evolution in DHSVM-GDM and comparison with observed albedo data (2010-2015) in Pheriche, Pyramide and Changri Nup.

15 3.2.3 Avalanches parameterization

The transport of snow by avalanches is not represented in the original version of DHSVM-GDM. The absence of avalanches in the model can lead to an unrealistic accumulation of snow in steep high elevated cells, where the air temperature remains below 0 °C, and to a deficit of snow in the lower areas, where snow melt occurs during the melting season. The simulated water balance directly depends on the snow cover, thus not considering avalanches can lead to significant errors. In order to address these discrepancies, an avalanche module [inspired by Wortmann et al. \(2016\)](#) was implemented in DHSVM-GDM. The avalanche model transfers snow to downslope neighbour cells under the following conditions:

- if the terrain slope is steeper than 35 ° and the amount of dry snow water equivalent (total snow water equivalent minus liquid water content) is higher than 30 cm: 5 cm of snow water equivalent remains in the cell and the rest is removed by avalanches;
- if the terrain slope is ~~less steep than below~~ 35 ° but the difference in snow water equivalent compared to the downslope neighbour cells is larger than 50 cm: 95 % of the difference is removed by avalanches.

The transfer of snow by avalanches is based on the surface runoff routing in DHSVM-GDM: at every time step starting from 10 the highest cell of the DEM to the lowest, each cell can transfer snow to its closest downslope neighbour cells (between 1 and 4 cells). Within the same time step, the amount of snow in the receiving cells is actualised and the avalanches propagate downslope until the conditions cited above are no longer respected.

3.2.4 Glacier parameterization

The distributed ice thickness is derived from the terrain slope following the method described in Haeberli and Högl (1995).

15 Since the standard DHSVM-GDM model does not take into account the impact of the debris layer on melting of the glaciers, the insulating effect of the debris layer is not represented. Here, we implemented a reduction factor for ice melt generated in grid cells with debris-covered glaciers (see Sect. 3.3).

Moreover, in the original DHSVM-GDM version, glacier melt is instantaneously transferred to the soil surface which is parameterized as bedrock under glaciers (Naz et al., 2014). This significantly underestimates the transfer time through glaciers.
 20 In this study we modified the soil parameterization in glacierized areas by increasing the soil depth to 2 m under glaciers. This modification of the soil depth under glaciers enables to compensate the absence of representation of the englacial liquid water storage in the model. This new parametrization also implies to change the values of three soil parameters under glaciers: the vertical and the lateral conductivities, as well as the porosity (Table A2). These were fixed by optimizing the recession shape of the hydrographs.

25 **3.2.5 Quantification of the flow components**

Quantifying the relative contributions of ice melt, snow melt, and rainfall in the river discharge at different time scales is a difficult task because hydrological models usually do not track the origin of water during transfer within the catchment (Weiler et al., 2018). There are also different ways of defining the origins of the streamflow. Weiler et al. (2018) lists three types of contributions: 1) contributions from the source areas i.e. from each class of landcover, 2) contributions from the runoff generation (overland flow, subsurface flow, and groundwater flow), and 3) input contributions (ice melt, ~~snowmelt~~^{snow melt}, and rain).

In this study, two different definitions were used to estimate the hydrological contributions. First, we estimate the ~~input contributions (ice melt~~^($V_{icemelt}$) ~~contributions of ice melt~~^($V_{icemelt}$), ~~snow melt~~^($V_{snowmelt}$), and net rainfall ($V_{rainNet}$) to the total 5 ~~production of runoff~~^(V_{runoff}) ~~(water input (definition 1))~~ according to the following ~~equations: water balance equations (all the terms are fluxes expressed in [L/T])~~:

$$\underline{V_{runoff}} \underline{\text{Input}} = \underline{V_{icemelt}} \underline{\text{Icemelt}} + \underline{V_{snowmelt}} \underline{\text{Snowmelt}} + \underline{V_{rainNet}} \underline{\text{RainNet}} \quad (4)$$

$$\underline{V_{icemelt}} \frac{dI_{wq}}{dt} = \underline{V_{glAcc}} \underline{\text{GlAcc}} - \underline{S_{ice}} \underline{\text{Icemelt}} - \frac{dI_{wq}}{dt} \underline{\text{SublIce}} \quad (5)$$

$$\underline{V_{snowmelt}} \frac{dS_{wq}}{dt} = \underline{P_{solid}} \underline{\text{solid}} - \underline{S_{snow}} \underline{\text{Snowmelt}} - \underline{V_{glAcc}} \underline{\text{SublSnow}} - \frac{dS_{wq}}{dt} \underline{\text{GlAcc}} \quad (6)$$

$$10 \quad \underline{V_{rainNet}} \underline{\text{RainNet}} = \underline{P_{liquid}} \underline{\text{liquid}} - \underline{E_{int}} \underline{\text{int}} \quad (7)$$

$$\underline{Q} \frac{d\text{Storage}}{dt} = \underline{V_{runoff}} + \underline{ET} \underline{\text{Input}} - \underline{Q} - \underline{E_T} \quad (8)$$

Where $\underline{V_{glAcc}} \frac{dI_{wq}}{dt}$ and $\frac{dS_{wq}}{dt}$ are the variations of the ice and snow storages, $\underline{\text{GlAcc}}$ is the amount of snow that is transferred to the ice layer by compaction on glaciers (Naz et al., 2014), $\underline{S_{ice}}$ and $\underline{S_{snow}}$ $\underline{\text{SublIce}}$ and $\underline{\text{SublSnow}}$ are the amounts of sublimation from the ice and snow layers, $\frac{dI_{wq}}{dt}$ and $\frac{dS_{wq}}{dt}$ are the variations of the ice and snow storages, $\underline{P_{solid}}$ and $\underline{P_{liquid}}$ 15 are the $\underline{P_{solid}}$ and $\underline{P_{liquid}}$ are the amounts of solid and liquid precipitation, and $\underline{E_{int}}$ $\underline{E_{int}}$ is the amount of evaporation from

intercepted water stored in the canopy. It is worth noting that the sum of these contributions V_{runoff} (Input) is not equal to the outflow at the catchment outlet Q as it represents all liquid water reaching the soil surface (before infiltration and potential storage in the soils and glaciers ($\frac{d\text{Storage}}{dt}$) and before evapotranspiration ET). Moreover, at daily or monthly time scale, V_{runoff} may not be equal to $Q - ET$ as liquid water can be stored by or evacuated from the soil or glaciers (E_T).

20 In order to evaluate the seasonal components of the outflow at the catchment's outlet, we also define the hydrological contributions as fractions of the outflow coming from the different contributing areas (definition 2):

- direct glacier contribution: direct runoff from glacierized areas,
- delayed glacier contribution: resurging melt water stored inside glaciers,
- direct snow contribution: direct outflow from snow covered non-glacierized areas,
- 25 – direct runoff: direct runoff from areas without snow and glaciers,
- subsurface and groundwater contribution: resurging water from the soil in non-glacierized areas resulting from infiltrated rainfall, snow melt, as well as upstream lateral subsurface flows.

These contributions are obtained from the amount of water reaching the soil surface simulated by DHSVM-GDM (see supplementary material). On each grid cell, this volume is a mixture of ice melt, snowmelt and rainfall and can either infiltrate into the soil or produce runoff. The definition 2 combines contributions from source areas (glacierized and non-glacierized areas) and contributions from runoff generation (direct runoff, englacial contribution, and soil contribution).

5 Figure 5 illustrates the two definitions of the different contributions to outflows. Definition 1 allows assessing the annual impact of glaciers and snow cover glacier melt and snow melt on the water production, while Definition 2 describes the intra-annual routing of the water within the catchment. Moreover, using the two definitions allows to directly compare our the results with other hydrological modelling studies in the Dudh Koshi basin, which have estimated glaciers contributions either from effective ice melt (Savéan et al., 2015; Ragettli et al., 2015; Soncini et al., 2016) or runoff from glacierized areas (Immerzeel 10 et al., 2012; Nepal et al., 2014). Further, we assessed the impact of the definition of hydrological components on the estimated glaciers contribution.

Flow components were estimated for the period 2012-2015 at annual scale, on the basis of the glaciological year (from 1 December to 30 November), as well as monthly, daily, and sub-daily scales, in order to have a better understanding of the seasonal variation of the estimated hydrological contributions.




Figure 5. Definitions of the flow components.

3.3 Experimental set-up

Simulations were run with a 1 h time step and a spatial resolution of 100 m for the period from 1 November 2012 to 27 November 2015 corresponding to the period with most available meteorological and discharge data.

A soil depth map was derived from the DEM using the method proposed in the DHSVM-GDM documentation (Wigmsta 5 et al., 1994). As a result, soil depth outside glacierized areas ranges between 0.5 and 1 m. Under the glaciers, the soil depth was set to 2 m. All parameter values retained for the simulations (with no calibration) are summarized in Appendix A.

In order to test the impact of the representation of the cryospheric processes on the hydrological modelling, we performed simulations with the four following configurations:

- v0: original DHSVM-GDM snow and glacier parameterization;
- 10 – v1: modified snow albedo parameterization;
- v2: modified snow albedo parameterization and avalanche module;
- v3: modified snow albedo parameterization, avalanche module and melt coefficient for debris covered glaciers.

All four configurations were run with the Racoviteanu et al. (2013) glaciers outline. Concerning the melt of the debris-covered glaciers, we use a reduction factor of 0.4 as estimated by Vincent et al. (2016) from a study on uncovered and debris covered
15 areas of the Changri Nup glacier.

Using configuration v3, we also tested the impact of using different glaciers outlines (the GAMDAM and ICIMOD inventories were also considered for simulations). The debris-covered glacier melt reduction factor estimated in Konz et al. (2007), Nepal et al. (2014) and Shea et al. (2015) are respectively equal to 0.3, 0.33 and 0.47. Thus, values between 0.3 and 0.5 were
5 also considered (in addition to the reference of 0.4) in order to evaluate the sensitivity of the model to the debris covered glacier reduction factor.

3.3.1 Model forcing

Meteorological data from the Pheriche and Pyramid stations (Table 1) were spatialized over the basin by an inverse distance interpolation method. Altitudinal lapse rates of precipitation and temperature were calculated at 1 h time step from data collected at Pangboche (3950 m a.s.l.), Pheriche (4260 m a.s.l.) and Pyramid (5035 m a.s.l.) (Fig. 6). Only significant lapse rates with R^2 values higher than 0.75 were retained for precipitation (43 % of the dataset). For smaller R^2 , the lapse rate is considered as not significant and thus set to 0.

In this study, the precipitation lapse rates show a large seasonal variability with daily lapse rates ranging from -41 to
9 mm km $^{-1}$. Precipitation decreases with elevation during the monsoon season and increases with elevation in winter: during
5 the simulation period, we found 450 days (40 %) with no precipitation, 83 days (8 %) with a strictly negative lapse rate and 165 days (15 %) with a strictly positive lapse rate. Concerning temperatures, daily lapse rates range from -0.009 to +0.006 °C m $^{-1}$. We found only 10 days (1 %) showing a temperature inversion with a positive daily lapse rate.




Figure 6. Daily temperature and precipitation lapse rates. Discarded precipitation lapse rates (with a $R^2 < 0.75$) are represented in orange.

3.3.2 Model evaluation

A multi criteria evaluation was made considering simulated outflows, SCA and glacier mass balances. Discharge measurements at Pheriche station were used as reference for the evaluation of simulated outflows. A 15 % confidence interval was retrieved as representative of the uncertainty of measured discharge. Nash-Sutcliffe Efficiency (NSE) (Nash and Sutcliffe, 1970) and Kling-Gupta Efficiency (KGE) (Gupta et al., 2009) were chosen as objective functions and applied to daily discharges. The simulated SCA was evaluated in comparison to daily SCA derived from MODIS images. Because a large number of MODIS images suffer from cloud coverage, we only compared the simulated and observed SCA during days with less than 5 % of cloud cover on the catchment. The simulated glacier mass balances were evaluated at basin scale by a comparison with published regional geodetic mass balances and at local scale using available stake measurements on the clean ice West Changri Nup and the Pokalde glaciers (Sherpa et al., 2017).

4 Results

4.1 Impact of the snow and glacier parameterizations on the simulated results

20 This section presents the simulation results obtained with the different configurations of the model DHSVM-GDM (configurations v0, v1, v2, and v3, see ch. 3.3) and the analysis of the impact of the snow and glacier parameterizations on the [simulated](#) annual outflow, the daily SCA, annual glacier mass balances.

4.1.1 Annual outflow

Figure 7 represents the annual outflow and flow components (definition 1) simulated with the different model configurations, indicating the impact of each modification of the snow and glacier parameterization on the simulated annual outflow and [the glacier contribution to the runoff](#) [flow components](#). Configuration v1 leads to a drastically increased outflow due to an enhanced ice melt component. Implementing the avalanche module (v2) reduces the ice melt component and increases the snow melt component by 21 %. Configuration v3 including debris-covered glaciers further reduces the ice melt, resulting in a simulated annual outflow close to the observations.

Figure 7 shows that configuration v2, which does not consider the debris-covered glaciers, overestimates the outflow at Pheriche with a mean bias of +32 % compared to the annual observed outflow. Without the debris layer, the ice melt component represents 817 mm, which is nearly twice the amount of ice melt obtained with v3 that includes debris-covered glacier melt.

The configuration with all three modifications (v3) gives results similar to the original parameterization of DHSVM-GDM (v0) in terms of glacier mass balance, improving slightly the annual outflow. The ice melt factor for debris covered glaciers and the avalanches compensate the increase of ice melt caused by the new snow albedo parameterization, but the modifications implemented in v3 impact the results for the flow components: on average, less ice melt and more snow melt are generated. Moreover, the configuration v3 modifies the seasonal variation of the outflow by increasing winter discharges and reducing monsoon discharges (not illustrated here) which improves the daily NSE and KGE (Table 3).




Figure 7. Simulated annual hydrological contributions (Definition 1) to Pheriche outflow for 3 glaciological years from 12/2013 to 11/2015.

	v0	v1	v2	v3
NSE	0.87	0.53	0.74	0.91
KGE	0.83	0.5	0.65	0.88

Table 3. NSE and KGE values calculated on the daily discharges on the period 2012-2015 for each model configuration.

4.1.2 Snow cover dynamics

Figures 8 and 9 compare the simulated snow cover area (SCA) and duration obtained with the configurations v0, v1, v2, and v3 to data derived from MODIS images. The SCA is strongly overestimated using the original parameterization v0: Figure 8 shows that after full coverage it does not decrease fast enough compared to the MODIS data. Figure 9 demonstrates that the snow cover duration is over-estimated for the entire catchment area. This indicates that in the simulations snow does not melt fast enough using the original parameterization. Configuration v1 with the modified snow albedo parameterization (Eq. 2) accelerates the snow melt and improves the SCA simulation (Fig. 8). The RMSE between the simulated and observed SCA decreases from 29 % using v0 to 14 % using v1 and v2. Figure 9 shows that with configuration v1 in some areas located at high elevation the snow cover duration is underestimated. This bias is rectified in configuration v2 since the avalanche module transfers snow from high elevated and sloping cells downward and corrects the lack of snow simulated with configuration v1 at the edges of the permanent snow cover (Fig. 9). The results for the SCA and snow cover duration using the configurations v2 and v3 are the same since only the ice melt rate for debris covered glaciers is modified.




Figure 8. Comparison of the MODIS SCA and the simulated daily SCA with the four modelling configurations (v0, v1, v2 and v3) for the Pheriche catchment.




Figure 9. Difference between the mean annual snow cover duration simulated with DHSVM-GDM and derived from MODIS images (in days) for the Pheriche catchment (top panels), with a focus on West Changri Nup (medium panels) and Pokalde glaciers (bottom panels).

4.1.3 Glacier mass balances

Figure 10 compares the simulated mean annual glacier mass balances obtained with the different model configurations with mass balances determined with geodetic methods [between 1999 and 2015](#) (Bolch et al., 2012; Gardelle et al., 2013; Nuimura et al., 2015; King et al., 2016; Brun et al., 2017). These geodetic mass balances range from $-0.67 \pm 0.45 \text{ m w.e.yr}^{-1}$ [\(2000-2008\)](#) (Nuimura et al., 2015) to $-0.32 \pm 0.09 \text{ m w.e.yr}^{-1}$ [\(2000-2015\)](#) (Brun et al., 2017).

Our results show that the snow parameterization has a significant impact on the simulated glacier mass balance. The mass balance simulated with v0 is on average $-0.82 \text{ m w.e.yr}^{-1}$ and decreases to $-2.02 \text{ m w.e.yr}^{-1}$ with the corrected snow albedo (v1) since the modified snow albedo parameterization accelerates the snow melt which leads to more uncovered ice and stronger glacier melt. The avalanche module (v2) adds snow on glaciers and increases the accumulation and, thus, reduces the glacier melt to $-1.69 \text{ m w.e.yr}^{-1}$. Nevertheless the mass balance remains too negative compared to geodetic mass balances, which suggests that the model produces too much ice melt. The implementation of debris-covered glaciers (v3) gives a mean annual glacier mass balance of $-0.84 \pm 0.14 \text{ m w.e.yr}^{-1}$, which is within the intervals of uncertainty and, thus, in good agreement with the results from geodetic methods.




Figure 10. Mean annual glacier mass balances simulated with configurations v0, v1, v2 and v3. The error bar for configuration v3 represents the uncertainty related to the debris layer coefficient melt varying between 0.3 and 0.5.

15 We also evaluated the mass balance at the point scale. Figure 11 shows the simulated mass balances with parameterizations v0, v1, v2 and v3 versus the observed mass balances of the two debris-free glaciers West Changri Nup and Pokalde measured in-situ for the three glaciological years (2012-2015). Here, the configuration v3 gives the same results as the configuration

v2 because in configuration v3 only the ice melt rate on debris-covered glaciers is modified. The simulated mass balances vary according to the model configuration. With configuration v0, the model overestimates the point mass balances because of small snow melt rates (see also section 4.1.2). With configuration v1, the model overestimates the ice melt on the West Changri Nup glacier due to a lack of accumulation in the western part of the catchment and a too strong accumulation on the Pokalde 5 glacier (Fig. 9). The configuration v2 improves the simulated mass balance by transferring snow due to avalanches on the West Changri Nup glacier and by removing exceeding snow accumulation on the Pokalde glacier. For the Pokalde glacier, the mass balances simulated with configuration v3 show a larger variability than the mass balances simulated with configuration v0, but the point mass balances are spread around the diagonal axis which leads to a bias ten times smaller (mean bias of 1 m with v0 and 0.1 m with v3).

10 The results at basin scale and point scale show that the snow parameterization has a strong impact on the simulated glacier mass balance and that the new snow albedo parameterization and the avalanching module clearly improve the simulated glacier mass balance on debris-free glaciers. Nevertheless, regarding point mass balances, the agreement is far from being perfect, due either to simulation errors (including errors depending on the interpolated input fields and errors induced by the representation of slopes and expositions by the DEM) and/or from a lack of representativeness of the measurements.




Figure 11. Annual simulated and measured point mass balances on West Changri Nup (left panel) and Pokalde (right panel) glaciers; also shown is the 1:1 line.

15 4.2 Simulated outflows and flow components

This section presents the outflows and flow components simulated in the Pheriche basin during the period 2012-2015 with the modified version of DHSVM-GDM (configuration v3). The simulation results are analysed using two different definitions of the flow components (definitions 1 and 2, see ch. 3.2.5).

4.2.1 Annual simulated outflow and hydrological contributions

20 The annual outflows simulated with the new parametrization of the model (configuration v3) are in good agreement with the annual observed outflows since they remain within the 15 % interval of estimated error (Fig. 12 and Table 4).

The results show an inter-annual variability of the [hydrological contributions to the overall outflow](#) [flow components](#). During the period 2013-2015, the ice melt component ranged from 41 to 50 %, the snow melt component from 37 to 47 % and the net rainfall component from 12 to 16 %. These variations are related to the meteorological annual variability. The amount of rainfall decreased from 2013 to 2015 and explains the decrease of the net rainfall components from 155 mm in 2013 to 88 mm in 2015. The snow melt component is higher in 2013 because of warmer pre-monsoon and monsoon seasons. The ice melt component is mainly controlled by the amount of winter snowfall. In 2014 a low amount of snowfall was observed, so 5 the snowpack melted more rapidly and the glaciers started melting earlier. In contrast, 2015 was a year with a lot of winter snowfall, which delayed the beginning of the glacier melt and explains the lower ice melt component. The losses by evaporation and sublimation are rather constant through the simulation period ranging from 140 to 150 mm/yr.

The runoff coefficients (ratio between the annual outflow and annual precipitation) were on average equal to 1.4, which means that a considerable amount of water is withdrawn each year from the catchment through ice melt (eventually in the form 10 of a delayed groundwater flow).

On average, we find that the outflow is mainly produced by meltwater as 46 % of the annual [outflow](#) [water input](#) is due to ice melt and 41 % to snow melt (definition 1). The contributions estimated according to definition 2 show the importance of infiltration and subsurface flows in the water balance since more than 40 % of the outflow was coming from water infiltrated in glaciers and more than 20 % from subsurface and groundwater flows [generated outside the glacier covered area](#).

The choice of the definition of the hydrological components leads to different perceptions of the glacier contribution to the outflow. The glacier contribution to the total outflow is 69 % if the contribution from the entire glacierized area (i.e. 5 contributions of ice melt, snow melt and net rainfall) is considered like in definition 2. However, the contribution from ice melt alone, included in definition 1, corresponds to only 46 % of the [outflow](#) [water input](#).




Figure 12. Simulated annual hydrological contributions to Pheriche outflow for the two definitions of the flow components (definition 1 and definition 2) and for 3 the glaciological years from December 2013 to November 2015.

	2012-2013	2013-2014	2014-2015
Total precipitation (mm)	708	644	683
Snowfall (mm)	501	492	561
Q_{obs} (mm)	994	1081	786
Q _{obs} ±15 % (mm)	845 - 1143	919 - 1244	668 - 904
Q_{sim} (mm)	999	933	729
Bias (%)	+1	-14	-7
Evapotranspiration (mm)	61	48	43
Sublimation (mm)	91	96	97
Flow components (Definition 1)			
Net rainfall (mm)	141 <ins>164</ins>	106 <ins>117</ins>	84 <ins>93</ins>
Snow melt (mm)	375 <ins>420</ins>	336 <ins>368</ins>	322 <ins>362</ins>
Ice melt (mm)	422 <ins>476</ins>	443 <ins>496</ins>	284 <ins>317</ins>
Flow components (Definition 2)			
Direct glacier contribution (mm)	293	244	163
Delayed glacier contribution (mm)	414	420	303
Direct runoff (mm)	21	14	9
Direct snow contribution (mm)	51	55	43
Subsurface and groundwater contribution (mm)	220	200	211

Table 4. Annual hydrological balance simulated with configuration v3 for the 3 glaciological years from December 2013 to November 2015.

4.2.2 Seasonal variations of the flow components

Figure 13 presents the daily simulated discharges simulated with configuration v3 and the flow components estimated with the two different definitions. Daily discharges were well simulated for 2012-2013 and 2014-2015 by the model, with NSE equal to 0.91 and KGE equal to 0.88. However, the outflow was under-estimated by the model during the monsoon season in 2014.

The simulated total ~~runoff~~water input (i.e. the sum of snow melt, ice melt and net rainfall) is always higher than the simulated outflow at the catchment outlet before the monsoon season (from February to June) and lower during post-monsoon and winter seasons. This is mainly due to glacier melt water stored inside the glaciers during pre-monsoon and monsoon seasons and continuing surging during winter, as well as to changes in the soil water storage (Fig. 13b and 14b).

Figure 14 shows the mean monthly flow components averaged over the simulation period. From February to May-June, the ~~runoff~~production ~~water input~~ is entirely controlled by snow and ice melt (snow melt between 50 and 60 % ice melt between 40 and 48 %) (Fig. 14a). The net rainfall, snow melt and ice melt absolute contributions are at their maxima in July and August during the monsoon season. During these two months, 24 % of the runoff is generated by net rainfall, 37 % by snow melt,

and 38 % by ice melt. From October to January, the runoff is produced by ice melt (up to 80 % in December) and snow melt (between 20 and 30 %). Groundwater and englacial water represent a significant fraction of the monthly outflow as they contribute more than 50 % of the outflow during the monsoon season and can contribute up to 90 % during winter (Figure 14b). Direct contributions from glacierized areas, snow areas, and direct runoff are highest during the monsoon season, when the englacial and soil storage is saturated.




Figure 13. Daily discharges and flow components simulated with configuration v3: (a) production of ice melt, snow melt and net rainfall (note that the sum of the flow components represent the total runoff water input and is not equal to the discharge at the catchment outlet, see definition 1, ch. 3.2.5) (b) hydrological contributions to the outflow (definition 2, ch. 3.2.5). Observed discharges are represented by the black line with a 15 % interval of error.




Figure 14. Average monthly contributions to the [runoff production-water input](#) (definition 1, ch. 3.2.5) (a) and hydrological contributions (definition 2, ch. 3.2.5) (b) simulated with configuration v3 for the years 2012-2015.

5 4.2.3 Diurnal cycle

Figure 15 presents the diurnal cycles of precipitation and hydrological components averaged for each considered season (winter, pre-monsoon, monsoon, and post-monsoon) obtained with configuration v3. During winter, pre-monsoon, and post-monsoon, the observed outflow is rather constant during the day, with a weak peak around noon when the temperature is at its maximum. During this period, almost all of the precipitation is in the form of snowfall leading to no direct response for the outflow. The peak around noon can be explained by snow melt or the melting of small frozen streams. During the monsoon season, there is a strong diurnal cycle of the precipitation with a maximum occurring during late afternoon or at night causing a peak in the discharge around midnight.

The model simulates ice and snow melt during day time with a maximum at noon as expected. Except for the monsoon season, it seems to simulate accurately the baseflow during night without melt production: the discharge is rather controlled by the release of the glacier and soil storage. However, the model simulates a peak of discharge around 14 h originating mostly from glacierized areas, two hours after the maximum of ice and snow melt, which does not correspond to observed discharges. At daily and longer time scales the water balance is correctly simulated. However, at a sub-daily scale the model responds too quickly to the snow and ice melt production.




Figure 15. Mean hourly precipitation, discharge and flow components simulated with configuration v3 and averaged for the winter (DJFM), pre-monsoon (AM), monsoon (JJAS), and post-monsoon (ON) seasons. Note different y-axis scales for each season.

10 5 Discussion

5.1 Simulation of the discharge and flow components

Overall, the comparison between the two definitions of the hydrological contributions shows that contributions must be explicitly specified in order to allow inter-comparison between models, especially for catchments with a large glacierized area. Moreover, the use of two different definitions allows to get complementary information on the origin of the outflow (processes 15 at the origin of the runoff, types of flow generation, contributive zones). A perspective to improve the quantification of the hydrological contributions to the outflow is to track the ice melt, snowmelt and rainfall component pathways in the model as suggested in Weiler et al. (2018). This would enable to quantify the fractions of the three components contributing to subsurface and groundwater flow, which is not possible with the current version of DHSVM-GDM.

Soncini et al. (2016) studied flow components in the Pheriche catchment for the period 2013-2014 and estimated an annual ice melt component of 55 % and a snow melt component of 20 % of the annual outflow. The ice melt components are thus quite similar in terms of relative contributions to outflow, which is not the case for the snow melt components. We think that the main reason of such a difference is that we use different precipitation input. Indeed, precipitation data are measured here by Geonor sensors, while in Soncini et al. (2016) precipitation data come from tipping buckets. At the Pyramid station, where both sensors are installed, the Geonor sensor measures 60 % more precipitation than the tipping bucket over the period 2013-2015 and the main differences are in terms of solid precipitation (309 mm of mean annual snowfall measured by the Geonor sensor versus 83 mm measured by the tipping bucket, which is known to badly perform with solid precipitation).

Concerning the seasonal contributions to the outflow, our results are consistent with the results from Soncini et al. (2016), who found a main contribution of snow melt during the pre-monsoon season, mixed contributions of rainfall, snow melt and ice melt during the monsoon season and mixed contributions of snow melt and ice melt during post-monsoon and winter season. The studies of Ragettli et al. (2015) and Racoviteanu et al. (2013) concerning the Upper Langtang and the Dudh Koshi basin respectively, showed that most of the winter outflow surges from soil, channel, surface, and englacial storage changes, which is consistent with our results. However, the estimated flow components presented in this study, particularly the soil and englacial contributions, are estimations which strongly depends on the model set-up. Figure 13 shows that the main part of the soil infiltrated water resurges within a day, whereas liquid water can be stored for several months within the glaciers. This difference between the response of the soil storage and the englacial storage results from the soil and glacier parameterization (see sensitivity analysis in ch. 5.3.2).

At hourly scale, the results show that the model cannot represent the diurnal cycle of the outflow correctly as the simulated hydrological response is anticipated. Irvine-Fynn et al. (2017) found that on the Khumbu glacier the presence of supraglacial ponds buffers the runoff by storing diurnally more than 20 % of the discharge. This could explain the longer transfer time observed on the measured outflows which are not represented by the model. This shows that the current representation of the glacier and soil storage in DHSVM-GDM does not allow to reproduce accurately the diurnal variations of discharge and further studies are needed in order to improve the model.

5.2 Representation of the cryospheric processes in the model

One of the main difficulties for hydrological modelling of highly glacierized catchments is to correctly simulate at the same time the outflows, the dynamics of the snow cover, and the glacier mass balances.

The results showed that two different representations of the cryospheric processes in the model (v0 and v3) can lead to similar simulated annual outflows but different estimations of the ice melt and snow melt contributions to the outflow. This is particularly true for the glaciological year 2014-2015, when the ice melt contribution decreases from 59 % with v0 to 41 % with v3 and the snow melt contribution increases from 29 % to 47 % (Fig. 7). This can be explained by the fact that 2014-2015 was a year with a high amount of snowfall (Fujita et al., 2017), therefore, the representation of snow processes in the model has a larger impact on the simulated runoff production than the two other years. This highlights the importance of a correct

representation of snow processes in the model. This also shows the need to use as much validation data as possible to assess the coherence between the ice, snow and hydrological processes and reduce the uncertainty on the flow components estimation.

20 The results also showed that the modification of one specific hydrological process (here, the representation of the snow albedo evolution) can have a significant impact on the simulated hydrological response of the catchment and requires improving other processes (here, considering specific representation of avalanches and debris-covered glaciers).

Further modifications of the model could also lead to different model results and it is also not excluded that different model errors are compensating each other. For example, the results showed that the original model version leads to a correct 25 simulation of the river discharges because the non-representation of the insulation effect for debris covered glaciers on the ice melt was compensated by the incorrect representation of the snow albedo decrease. Due to the complexity of the model and the represented processes, no guarantee can be given that similar compensating effects still occur in the model. In this study, the validation of the model output was extended beyond the annual river discharge to discharges at different time scales, the snow cover area, and glacier mass balances in order to validate the simulations of the snow cover, glacier melt and discharges 30 separately. The results demonstrate that the new version of the model performs well for all three signals. Moreover, the new parametrizations of the snow albedo and ice melt under debris were based on observed data (MODIS and in situ albedo measurements, and coefficient for ice melt under debris from Vincent et al. (2016)) and do not result from a calibration in order to avoid compensation effects. Therefore, it is very likely that the new implementation improved the quality of the represented processes.

5 The results presented in this study also indicate possible forthcoming works for increasing the simulations reliability and reducing uncertainties, especially at short time steps. Indeed, at daily and longer scales, the different hydrological components seem to be well reproduced by the model. However, an analysis of the diurnal cycle (Fig. 15) showed that DHSVM-GDM responds too rapidly to the ice melt production and that the representation of the water storage within the glaciers needs to be improved. Further improvements should be based on studies that analyze the mechanisms of glaciers drainage systems 10 in the Khumbu region and their influence on glaciers outflow (e.g., Gulley et al., 2009; Benn et al., 2017). These studies show that englacial conduits and supraglacial channels, ponds and lakes play a key role in the response of glaciers: DHSVM-GDM could thus be upgraded by implementing a parameterization of such systems and delay the response of glacierized areas, as successfully proposed, for instance, in the model developed by Flowers and Clarke (2002). Other processes such 15 as supraglacial ponds and ice cliffs melting, transport of snow by wind or variation of temperature in the ice pack are not considered in DHSVM-GDM and their impact on the hydrological modelling should also be studied.

The avalanche routine implemented in this study is simplified and only considers 4 directions for the snow redistribution. A perspective of this study is to improve the representation of the avalanches in DHSVM-GDM by considering eight directions for the snow redistribution and considering other parameters such as the age of the snow cover, the snow density and the type of land-cover as it was proposed in Frey and Holzmann (2015).

Glaciers inventory	Basin glacier area		Glacier MB m w.e.yr ⁻¹	Qsim		Flow components (mm)		
	km ²	%		mm	Bias	Net rainfall	Snow melt	Ice melt
Racoviteanu et al. (2013)	60	43	-0.84	887	-7%	118-125 (13%)	363-383 (41%)	406-430 (46%)
GAMDAM	38	24	-1.17	824	-13%	117-123 (14%)	359-387 (44%)	348-370 (42%)
ICIMOD	44	30	-0.89	811	-15%	118-121 (14%)	365-397 (46%)	328-345 (40%)

Table 5. Mean annual glaciers mass balance (MB), outflow and flow components simulated with different glaciers inventories (configuration v3)

20 5.3 Uncertainties and other open-ended questions

5.3.1 Sensitivity to the glacier outline

The three inventories used in this study result in very different estimates of the glacierized area: between 43 % and 24 % of the Pheriche basin with the inventories proposed by Racoviteanu et al. (2013) and GAMDAM (Fig. 3). Table 5 presents the average annual glacier mass balances, outflows, and flow components for the configuration v3 using the three inventories. The GAMDAM inventory leads to a more negative glacier mass balance than the two others inventories with -1.17 m w.e.yr⁻¹ compared to -0.84 and -0.89 m w.e.yr⁻¹ for the Racoviteanu et al. (2013) and ICIMOD inventories. This is due to smaller glacier accumulation areas in the GAMDAM inventory. The amount of snowfalls collected over those areas is lower, leading to more

5 negative mass balances: glaciers receive less snowfall for accumulation, which lowers the mass balance value. Concerning the simulated outflow and flow components, the GAMDAM and ICIMOD inventories lead to fewer ice melt than the Racoviteanu et al. (2013) inventory due to their smaller areas in ablation zones, which leads to a smaller simulated annual outflow. From these results we estimate an uncertainty of 20 % (407-430 mm with the Racoviteanu et al. (2013) inventory versus 327-345 mm with the ICIMOD inventory, cf. Table 5) on the simulated annual ice melt volume related to the glaciers outline. The glacier 10 outline mainly affects the simulated outflow during the monsoon season, when the ice melt contribution to the outflow is more important and leads to an uncertainty of 8 % (154 mm with the Racoviteanu et al. (2013) inventory versus 141 mm with the ICIMOD inventory) on the monthly discharges during monsoon season. This result shows that the choice of the glacier inventory as an input data of the glacio-hydrological model contributes to the uncertainty on the simulation results. Here, the Racoviteanu et al. (2013) inventory gives the best results in terms of glacier mass balance and the smallest bias with respect 15 to the annual outflow. As its area is significantly larger than the other inventories, it gives the largest amount of ice melt. This potentially compensates a lack of precipitation due to a poor knowledge of the precipitation distribution over the catchment, specifically in the areas above 5000 m a.s.l. which constitute more than three quarters of the total area and for which no observations exist. It is worth noting that the glacier mass balances obtained with the Racoviteanu et al. (2013) and ICIMOD inventories are very similar but the amounts of simulated ice melt are different, which shows that a consistent mass balance 20 can lead to errors on the simulated glacier contributions and total outflow.


5.3.2 Sensitivity to the **soil-and-glaciers** parametrization

A major limitation to the estimation of the contributions from the runoff generation (overland flow, subsurface flow, and groundwater flow) is the representation of the groundwater and englacial flows in the model. In this study, we selected standard soil parameters for the simulations since there is very little information about soil properties in high-mountain environments.

5 Moreover, data to validate subsurface and groundwater flows do not exist making it very difficult to adapt the parameter values to the study area. Here, three different values of soil depth under glaciers were tested in order to assess the sensitivity of the soil parametrization on the simulated discharges. Figure 16a shows the simulated discharges and flow components (definition 2) simulated with configuration v3 for a soil depth under glaciers equals to 1 m, 2 m and 5 m. The soil depth under glaciers does not impact the simulated annual outflow and the total annual contributions from glacierized and non-glacierized areas. The soil depth under glaciers only impacts the partitioning between direct and delayed contributions (soil water and englacial water contributions): when the soil depth under glaciers ranges between 1 m and 5 m, the direct glacier contribution ranges from 34 to 20 %, and the delayed glacier contribution ranges from 35 to 47 %. At seasonal scale, an increase of the soil depth under glaciers leads to a delay of the outflow as there is more infiltration simulated during the pre-monsoon and monsoon seasons, but this has a limited impact on the NSE and KGE values (respectively ranging from 0.92 to 0.94 and 0.83 to 0.91).

5 Another limitation of our model lies in the application of a uniform reduction factor for ice melt under debris covered glaciers. Figure 16b shows the sensitivity of the model to the ice melt reduction factor on debris covered glaciers. When the reduction factor varies between 0.3 and 0.5, the simulated annual outflow is modified by $\pm 7\%$ and the mean ice melt flow component ranges from 42 to 50 %. This shows that the results are sensitive to the representation of the debris cover in the model. In order to have a more realistic representation of the debris, the reduction factor could be spatially distributed, at least 10 following elevation or slope exposition, and eventually taken as time-variant. As an example, Ragettli et al. (2015) considered a distributed debris thickness in their glacio-hydrological model and obtained a mean reduction of ice melt under debris of 84 %.

A



B




Figure 16. Annual discharges and flow components simulated with configuration v3 with a) three different soil depths under glaciers (1 m, 2 m, and 5 m) (definition 2), b) three different values for the debris-covered ice melt reduction factor (0.3, 0.4, and 0.5) (definition 1).

We also tested the model sensitivity with respect to the avalanche parameters by changing the values of the thresholds for the slope (35, 40, and 45 °) and the snow height (15, 30 and 60 cm) for steep terrains and for the snow water equivalent difference between two neighboring cells (25, 50, and 100 cm) in case of a flat terrain. In total, 11 combinations of these parameters were tested. The results confirmed that higher thresholds for triggering avalanches and their propagation led to less accumulation on the downslope parts of the glaciers and, thus, to more overall ice melt. Nevertheless, the results also show that the simulated outflow is not very sensitive to the avalanche parameterization since we derived a variation of 3 % of the mean annual outflow. However, the modification of the avalanche parameters leads to an uncertainty of 12 % on the estimation of the ice melt and snow melt components: the simulated contribution of ice melt to the water input varied between 46 and 52 % and the contribution of snow melt varied from 36 to 41 %.

5.3.3 Validation and forcing data uncertainties

A main issue is related to the availability of data for validating the glacio-hydrological modelling parameterizations and outputs. The lack of in-situ measurements at high elevations and the uncertainty related to the available data prevent from assessing the 25 performance of the model in simulating the different cryospheric processes we consider. They only allow to evaluate integrated variables such as the annual glacier mass balance, the seasonal snow cover area dynamics and the outflow at the catchment outlet with significant uncertainties that impact the estimation of the different flow components. For instance, concerning the validation of the simulated glacier mass balances, only the order of magnitude of the simulated and geodetic mass balances used in this study can be compared because the considered areas are not the same in the different studies and the considered 30 time periods differ as well. Indeed, four of the geodetic mass balances were derived for the Khumbu-Changri glacier, while mass balances from this study and from Brun et al. (2017) represent the mean mass balance for all glaciers located in the Pheriche basin. Moreover, the mean annual glacier mass balance is estimated here for the three simulated years (2012-2015), whereas the geodetic mass balances correspond to longer (5 to 15 years) as well as earlier periods beginning between 1999 and 2002 (see Figure 10) and do not take into account the inter-annual variability of the glacier mass balances. Nevertheless, on figure 10, the variability of the simulated glacier mass balances is much larger than the variability of the geodetic mass balances, showing the significant impact of the snow and glacier parameterization on the simulation results. It is also worth noting that the snow cover distribution evaluation is particularly challenging on the Pheriche catchment, as clouds cover more 5 than 50 % of the catchment during more than 150 days per year on average (and almost all the time during the monsoon season).

Finally, a major source of uncertainty lies in the lack of meteorological data at high elevation (because of the inaccessibility on the terrain) and in the measurement errors when observations are available, due to extreme meteorological conditions. Indeed, precipitation is known to be underestimated due to the difficulty of measuring solid precipitation with rain gauges 10 (Wolff et al., 2015). Precipitation fields provided by different atmospheric models and satellites show also high discrepancies in this region of the Himalayas (Andermann et al., 2011; Palazzi et al., 2013; Ceglar et al., 2017): a perspective of this study is to test the sensitivity of the model to different precipitation forcing data sets (in-situ, reanalysis, and satellite) and analyze the impact of different precipitation amounts and spatial distributions on the simulated discharges and flow components.

6 Conclusions

In this study we used a distributed physically-based glacio-hydrological model (DHSVM-GDM) to simulate the outflow of a small catchment in the Everest region and to estimate the different contributions to streamflows, which can be useful for water resources and water-related risks management. Some improvements on the cryospheric processes parameterization in DHSVM-GDM were proposed in order to better represent the snowcover dynamics, the ice melt under debris-covered glaciers, 5 and avalanches. Simulated SCA were compared with MODIS images and calculated glacier mass balances with local in situ and geodetic measurements.

Results showed that the representation of the cryospheric processes in the model has a significant impact on the simulated outflow and flow components. Despite some outstanding issues that have been discussed, we can argue that the most satisfactory snow and glacier parameterizations proposed in this study (model configuration v3) enabled to simulate the snow cover spatial distribution and the glacier mass balance more accurately than the original version of DHSVM-GDM, by increasing the glacier accumulation and reducing ice melt. Major contributions from glaciers and ~~and~~ snow to the outflow were found, with 46 %
5 of the ~~annual outflow~~ ~~annual water input~~ produced by ice melt and 41 % by snow melt. Winter flows are mainly controlled by the release of englacial and soil water storage (up to 78 % in December), which corroborates other studies (Racoviteanu et al., 2013; Ragettli et al., 2015). We estimate an uncertainty related to the ice melt reduction factor by debris (ranging from 0.3 to 0.5) of ± 0.14 m w.e.yr⁻¹ for the annual glacier mass balance and ± 7 % for the annual outflow. The glacier inventories used to outline the glacierized areas have also an important impact on the simulation results. The three inventories tested in this study
10 give estimations of the glacierized area ranging from 26 to 43 % of the basin area and a corresponding uncertainty of 20 % for the ice melt production.

This study also reminds that glacial and snow contributions to the streamflow must be clearly defined, as considering glacial contribution as the total outflow from the glacierized area or as outflow produced by ice melt can lead to very different estimations.

Name	Unit	Value(s)	Reference
Ground Roughness	m	0.04	Brutsaert (2005)
Reference Height	m	2	-
LAI Multiplier for rain interception	-	0.0005	Brutsaert (2005)
LAI Multiplier for snow interception	-	0.00005	Andreadis et al. (2009)
Tree Height	m	2	-
Vegetation Density	-	0.25	-
Distance from bank to canopy	m	2	-
Snow			
Snow Roughness	m	0.001	Brock et al. (2006)
Rain Threshold	°C	0	L'hôte et al. (2005)
Snow Threshold	°C	2	L'hôte et al. (2005)
Snow Water Capacity	-	0.05	Singh (2001)
Minimum Intercepted snow	m	0.005	-
Maximum Snow Albedo	-	0.85	MODImLab
Glaciers			
Glacier Albedo	-	0.3	MODImLab
Melt coefficient for debris-covered glacier	-	0.4	Vincent et al. (2016)

Table A1. Global parameter values used for the glacio-hydrological simulation with DHSVM-GDM

Name	Unit	Value(s)	Reference
Soil type	-	Regosol	
Lateral Conductivity	m/s	0.0053	Clapp and Hornberger (1978)
Exponential Decrease	-	2	Niu et al. (2005)
Depth Threshold	m	10	-
Capillary Drive	-	0.0756	Morel-Seytoux and Nimmo (1999)
Maximum Infiltration	m/s	6.94E-06	FAO
Surface Albedo	-	0.35	ModimLab
Number of Soil Layers	-	3	-
Porosity	-	0.6	calibrated
Pore Size Distribution	-	0.43	Rawls et al. (1982)
Bubbling Pressure	-	0.302	Rawls et al. (1982)
Field Capacity	-	0.31	Meyer et al. (1997)
Wilting Point	-	0.23	Meyer et al. (1997)
Vertical Conductivity	m/s	5.30E-05	Clapp and Hornberger (1978)
Thermal Conductivity layer 1	W/m.°C	7.114	Burns (2012)
Thermal Conductivity layer 2 et 3	W/m.°C	6.923	Burns (2012)
Thermal Capacity	J/m3.°C	1.40E+06	Burns (2012)

Table A2. Soil parameter values used for the glacio-hydrological simulation with DHSVM-GDM. Under glaciers, the values of the vertical and lateral conductivities were adjusted to 0.0003 m/s and the porosity value to 0.8.

Name	Unit	Value(s)			Reference
Vegetation type	-	Shrubland	Grassland	Agriculture	Bare
Overstory Present	-	FALSE	FALSE	FALSE	FALSE
Understory Present	-	TRUE	TRUE	TRUE	TRUE
Impervious Fraction	-	0	0	0	0
Height	m	1	0.3	1	0.15
Maximum	s/m	600	600	600	600
Resistance					Wigmota et al. (1994)
Minimum Resistance	s/m	200	200	120	120
Moisture Threshold	-	0.6	0.6	0.33	0.8
Vapor Pressure	Pa	2880	2880	4000	2000
Deficit					
R _{pc}	-	10	10	10	10
Number of Root	-	3	3	3	3
Zones					
Root Zone Depths 1	m	0.06	0.1	0.06	0.045
Root Zone Depths 2	m	0.13	0.05	0.13	0.025
Root Zone Depths 3	m	0.2	0.05	0.2	0.025
Understory Root	-	0.4	0.4	0.4	0.4
Fraction 1					
Understory Root	-	0.6	0.6	0.6	0.6
Fraction 2					
Understory Root	-	0	0	0	0
Fraction 3					
Understory Monthly	-	5.0 5.0 5.0	0.8 0.9 1.0	3.0 3.0 3.0	1.0 1.0 1.0
LAI		5.0 5.0 5.0	1.1 1.8 3.7	3.0 3.0 3.0	1.0 1.0 1.0
		5.0 5.0 5.0	4.8 4.2 2.0	3.0 3.0 3.0	1.0 1.0 1.0
		5.0 5.0 5.0	1.2 1.0 0.9	3.0 3.0 3.0	1.0 1.0 1.0
Understory Albedo	-	0.2	0.1	0.2	0.2
					Wigmota et al. (1994)

Table A3. Vegetation parameter values used for the glacio-hydrological simulation with DHSVM-GDM.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This work has been supported by the French National Research Agency (ANR) through the ANR-13-SENV-0005-04/05-PRESHINE project and by a grant from Labex OSUG@2020 (investissement d'avenir, ANR 10 LABX56). It has benefited ~~of data from from data of~~ the French Service de l'Observation GLACIOCLIM (<https://glacioclim.osug.fr/>) and from Ev-K2-CNR (www.evkc2cner.org). The DHSVM-GDM model was made available by Chris Frans (Department of Civil and Environmental Engineering, University of Washington). The works of Yves Lejeune (CNRM-GAME) for correcting precipitation data in the GLACIOCLIM database and Ahmed Alatrash for processing MODIS data have been particularly helpful. Authors had fruitful discussions with Pierre Chevalier, Francois Delclaux, Luc Neppel and Judith Eeckman (HSM).

References

25 Akhtar, M., Ahmad, N., and Booij, M.: The impact of climate change on the water resources of Hindu Kush–Karakorum–Himalaya region under different glacier coverage scenarios, *Journal of hydrology*, 355, 148–163, 2008.

Andermann, C., Bonnet, S., and Gloaguen, R.: Evaluation of precipitation data sets along the Himalayan front, *Geochemistry, Geophysics, Geosystems*, 12, 2011.

Andermann, C., Longuevergne, L., Bonnet, S., Crave, A., Davy, P., and Gloaguen, R.: Impact of transient groundwater storage on the 30 discharge of Himalayan rivers, *Nature Geoscience*, 5, 127, 2012.

Andreadis, K. M., Storck, P., and Lettenmaier, D. P.: Modeling snow accumulation and ablation processes in forested environments, *Water Resources Research*, 45, 2009.

Bajracharya, B., Uddin, K., Chettri, N., Shrestha, B., and Siddiqui, S. A.: Understanding land cover change using a harmonized classification system in the Himalaya: a case study from Sagarmatha National Park, Nepal, *Mountain Research and Development*, 30, 143–156, 2010.

35 Bajracharya, S. R.: Glacier status in Nepal and decadal change from 1980 to 2010 based on Landsat data, *International Centre for Integrated Mountain Development*, 2014.

Beckers, J. and Alila, Y.: A model of rapid preferential hillslope runoff contributions to peak flow generation in a temperate rain forest watershed, *Water Resources Research*, 40, 2004.

Benn, D., Bolch, T., Hands, K., Gulley, J., Luckman, A., Nicholson, L., Quincey, D., Thompson, S., Toumi, R., and Wiseman, S.: Response of debris-covered glaciers in the Mount Everest region to recent warming, and implications for outburst flood hazards, *Earth-Science Reviews*, 114, 156–174, 2012.

5 Benn, D. I., Thompson, S., Gulley, J., Mertes, J., Luckman, A., and Nicholson, L.: Structure and evolution of the drainage system of a Himalayan debris-covered glacier, and its relationship with patterns of mass loss, *The Cryosphere*, 11, 2247, 2017.

Berghuijs, W., Woods, R., and Hrachowitz, M.: A precipitation shift from snow towards rain leads to a decrease in streamflow, *Nature Climate Change*, 4, 583, 2014.

Bewley, D., Alila, Y., and Varhola, A.: Variability of snow water equivalent and snow energetics across a large catchment subject to Mountain 10 Pine Beetle infestation and rapid salvage logging, *Journal of Hydrology*, 388, 464–479, 2010.

Bhambri, R., Bolch, T., Chaujar, R. K., and Kulshreshtha, S. C.: Glacier changes in the Garhwal Himalaya, India, from 1968 to 2006 based on remote sensing, *Journal of Glaciology*, 57, 543–556, 2011.

Bolch, T., Kulkarni, A., Kääb, A., Huggel, C., Paul, F., Cogley, J. G., Frey, H., Kargel, J. S., Fujita, K., Scheel, M., et al.: The state and fate of Himalayan glaciers, *Science*, 336, 310–314, 2012.

15 Bookhagen, B. and Burbank, D. W.: Topography, relief, and TRMM-derived rainfall variations along the Himalaya, *Geophysical Research Letters*, 33, 2006.

Brock, B. W., Willis, I. C., and Sharp, M. J.: Measurement and parameterization of aerodynamic roughness length variations at Haut Glacier d’Arolla, Switzerland, *Journal of Glaciology*, 52, 281–297, 2006.

Brun, F., Berthier, E., Wagnon, P., Kääb, A., and Treichler, D.: A spatially resolved estimate of High Mountain Asia glacier mass balances 20 from 2000 to 2016, *Nature Geoscience*, 2017.

Brutsaert, W.: *Hydrology: an introduction*, Cambridge University Press, 2005.

Burns, P. J.: Glacier change in a basin of the Peruvian Andes and implications for water resources, Ph.D. thesis, Oregon State University, 2012.

25 Ceglar, A., Toreti, A., Balsamo, G., and Kobayashi, S.: Precipitation over Monsoon Asia: A Comparison of Reanalyses and Observations, *Journal of Climate*, 30, 465–476, 2017.

Chen, Y., Li, W., Fang, G., and Li, Z.: Hydrological modeling in glacierized catchments of central Asia-status and challenges, *Hydrology and Earth System Sciences*, 21, 669, 2017.

Clapp, R. B. and Hornberger, G. M.: Empirical equations for some soil hydraulic properties, *Water resources research*, 14, 601–604, 1978.

Cogley, J. G.: A more complete version of the World Glacier Inventory, *Annals of Glaciology*, 50, 32–38, 2009.

30 Cristea, N. C., Lundquist, J. D., Loheide, S. P., Lowry, C. S., and Moore, C. E.: Modelling how vegetation cover affects climate change impacts on streamflow timing and magnitude in the snowmelt-dominated upper Tuolumne Basin, Sierra Nevada, *Hydrological processes*, 28, 3896–3918, 2014.

Dickinson, R. E., Henderson-Sellers, A., Rosenzweig, C., and Sellers, P. J.: Evapotranspiration models with canopy resistance for use in climate models, a review, *Agricultural and Forest Meteorology*, 54, 373–388, 1991.

35 Dijkshoorn, K. and Huting, J.: Soil and Terrain database for Nepal, 2009.

Douville, H., Royer, J.-F., and Mahfouf, J.-F.: A new snow parameterization for the Meteo-France climate model, *Climate Dynamics*, 12, 21–35, 1995.

Flowers, G. E. and Clarke, G. K.: A multicomponent coupled model of glacier hydrology 2. Application to Trapridge Glacier, Yukon, Canada, *Journal of Geophysical Research: Solid Earth*, 107, 2002.

Frans, C., Istanbulluoglu, E., Lettenmaier, D. P., Naz, B. S., Clarke, G. K., Condom, T., Burns, P., and Nolin, A. W.: Predicting glacio-hydrologic change in the headwaters of the Zongo River, Cordillera Real, Bolivia, *Water Resources Research*, 51, 9029–9052, 2015.

5 Frey, S. and Holzmann, H.: A conceptual, distributed snow redistribution model, *Hydrology and Earth System Sciences*, 19, 4517, 2015.

Fujita, K., Inoue, H., Izumi, T., Yamaguchi, S., Sadakane, A., Sunako, S., Nishimura, K., Immerzeel, W. W., Shea, J. M., Kayastha, R. B., et al.: Anomalous winter-snow-amplified earthquake-induced disaster of the 2015 Langtang avalanche in Nepal, *Natural Hazards and Earth System Sciences*, 17, 749, 2017.

Gardelle, J., Berthier, E., Arnaud, Y., and Kaab, A.: Region-wide glacier mass balances over the Pamir-Karakoram-Himalaya during 1999–2011 (vol 7, pg 1263, 2013), *Cryosphere*, 7, 1885–1886, 2013.

10 Gulley, J., Benn, D., Screatton, E., and Martin, J.: Mechanisms of englacial conduit formation and their implications for subglacial recharge, *Quaternary Science Reviews*, 28, 1984–1999, 2009.

Gupta, H. V., Kling, H., Yilmaz, K. K., and Martinez, G. F.: Decomposition of the mean squared error and NSE performance criteria: Implications for improving hydrological modelling, *Journal of Hydrology*, 377, 80–91, 2009.

15 Haeberli, W. and Hölzle, M.: Application of inventory data for estimating characteristics of and regional climate-change effects on mountain glaciers: a pilot study with the European Alps, *Annals of glaciology*, 21, 206–212, 1995.

Immerzeel, W. W., Van Beek, L. P., and Bierkens, M. F.: Climate change will affect the Asian water towers, *Science*, 328, 1382–1385, 2010.

Immerzeel, W. W., Van Beek, L., Konz, M., Shrestha, A., and Bierkens, M.: Hydrological response to climate change in a glacierized catchment in the Himalayas, *Climatic change*, 110, 721–736, 2012.

20 Irvine-Fynn, T. D., Porter, P. R., Rowan, A. V., Quincey, D. J., Gibson, M. J., Bridge, J. W., Watson, C. S., Hubbard, A., and Glasser, N. E.: Supraglacial ponds regulate runoff from Himalayan debris-covered glaciers, *Geophysical Research Letters*, 2017.

Kääb, A., Treichler, D., Nuth, C., and Berthier, E.: Brief Communication: Contending estimates of 2003–2008 glacier mass balance over the Pamir–Karakoram–Himalaya, *The Cryosphere*, 9, 557–564, 2015.

Kaser, G., Großhauser, M., and Marzeion, B.: Contribution potential of glaciers to water availability in different climate regimes, *Proceedings of the National Academy of Sciences*, 107, 20 223–20 227, 2010.

King, O., Quincey, D. J., Carrivick, J. L., and Rowan, A. V.: Spatial variability in mass change of glaciers in the Everest region, central Himalaya, between 2000 and 2015, *The Cryosphere Discussions*, pp. 1–35, 2016.

Konz, M., Uhlenbrook, S., Braun, L., Shrestha, A., and Demuth, S.: Implementation of a process-based catchment model in a poorly gauged, highly glacierized Himalayan headwater, *Hydrology and Earth System Sciences Discussions*, 11, 1323–1339, 2007.

Kraaijenbrink, P., Bierkens, M., Lutz, A., and Immerzeel, W.: Impact of a global temperature rise of 1.5 degrees Celsius on Asia's glaciers, *Nature*, 549, 257–260, 2017.

Lejeune, Y., Bouilloud, L., Etchevers, P., Wagnon, P., Chevallier, P., Sicart, J.-E., Martin, E., and Habets, F.: Melting of snow cover in a tropical mountain environment in Bolivia: Processes and modeling, *Journal of Hydrometeorology*, 8, 922–937, 2007.

Leung, L., Wigmosta, M., Ghan, S., Epstein, D., and Vail, L.: Application of a subgrid orographic precipitation/surface hydrology scheme to a mountain watershed, *Journal of Geophysical Research: Atmospheres*, 101, 12 803–12 817, 1996.

Leung, L. R. and Wigmosta, M. S.: Potential climate change impacts on mountain watersheds in the Pacific Northwest, *JAWRA Journal of the American Water Resources Association*, 35, 1463–1471, 1999.

L'hôte, Y., Chevallier, P., Coudrain, A., Lejeune, Y., and Etchevers, P.: Relationship between precipitation phase and air temperature: comparison between the Bolivian Andes and the Swiss Alps/Relation entre phase de précipitation et température de l'air: comparaison entre les Andes Boliviennes et les Alpes Suisse, *Hydrological Sciences Journal*, 50, 2005.

Lutz, A., Immerzeel, W., Shrestha, A., and Bierkens, M.: Consistent increase in High Asia's runoff due to increasing glacier melt and precipitation, *Nature Climate Change*, 4, 587–592, 2014.

McDowell, G., Ford, J. D., Lehner, B., Berrang-Ford, L., and Sherpa, A.: Climate-related hydrological change and human vulnerability in remote mountain regions: a case study from Khumbu, Nepal, *Regional Environmental Change*, 13, 299–310, 2013.

Meyer, P., Rockhold, M., and Gee, G.: Uncertainty analyses of infiltration and subsurface flow and transport for SDMP sites, Tech. rep., Nuclear Regulatory Commission, Washington, DC (United States). Div. of Regulatory Applications; Pacific Northwest National Lab., Richland, WA (United States), 1997.

Morel-Seytoux, H. J. and Nimmo, J. R.: Soil water retention and maximum capillary drive from saturation to oven dryness, *Water resources research*, 35, 2031–2041, 1999.

Nash, J. E. and Sutcliffe, J. V.: River flow forecasting through conceptual models part I—A discussion of principles, *Journal of hydrology*, 10, 282–290, 1970.

Naz, B. S., Frans, C., Clarke, G., Burns, P., and Lettenmaier, D.: Modeling the effect of glacier recession on streamflow response using a coupled glacio-hydrological model, *Hydrology and Earth System Sciences*, 18, 787–802, 2014.

Nepal, S.: Impacts of climate change on the hydrological regime of the Koshi river basin in the Himalayan region, *Journal of Hydro-environment Research*, 10, 76–89, 2016.

Nepal, S., Krause, P., Flügel, W.-A., Fink, M., and Fischer, C.: Understanding the hydrological system dynamics of a glaciated alpine catchment in the Himalayan region using the J2000 hydrological model, *Hydrological Processes*, 28, 1329–1344, 2014.

Nijssen, B., Haddeland, I., and Lettenmaier, D. P.: Point evaluation of a surface hydrology model for BOREAS, *Journal of Geophysical Research: Atmospheres*, 102, 29 367–29 378, 1997.

Niu, G.-Y., Yang, Z.-L., Dickinson, R. E., and Gulden, L. E.: A simple TOPMODEL-based runoff parameterization (SIMTOP) for use in global climate models, *Journal of Geophysical Research: Atmospheres*, 110, 2005.

Nuimura, T., Sakai, A., Taniguchi, K., Nagai, H., Lamsal, D., Tsutaki, S., Kozawa, A., Hoshina, Y., Takenaka, S., Omiya, S., et al.: The
25 GAMDAM glacier inventory: a quality-controlled inventory of Asian glaciers, *The Cryosphere*, 9, 849–864, 2015.

Østrem, G.: Ice melting under a thin layer of moraine, and the existence of ice cores in moraine ridges, *Geografiska Annaler*, 41, 228–230,
1959.

Palazzi, E., Hardenberg, J. v., and Provenzale, A.: Precipitation in the Hindu-Kush Karakoram Himalaya: Observations and future scenarios,
Journal of Geophysical Research: Atmospheres, 118, 85–100, 2013.

30 Paul, F., Kääb, A., Rott, H., Shepherd, A., Strozzi, T., Volden, E., et al.: GlobGlacier: a new ESA project to map the world's glaciers and ice
caps from space, *EARSeL eProceedings*, 8, 11–25, 2009.

Paul, F., Barrand, N. E., Baumann, S., Berthier, E., Bolch, T., Casey, K., Frey, H., Joshi, S., Konovalov, V., Bris, R. L., et al.: On the accuracy
of glacier outlines derived from remote-sensing data, *Annals of Glaciology*, 54, 171–182, 2013.

Pfeffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Cogley, J. G., Gardner, A. S., Hagen, J.-O., Hock, R., Kaser, G., Kienholz, C., et al.: The
35 Randolph Glacier Inventory: a globally complete inventory of glaciers, *Journal of Glaciology*, 60, 537–552, 2014.

Pritchard, H. D.: Asia's glaciers are a regionally important buffer against drought, *Nature*, 545, 169–174, 2017.

Racoviteanu, A. E., Armstrong, R., and Williams, M. W.: Evaluation of an ice ablation model to estimate the contribution of melting glacier
ice to annual discharge in the Nepal Himalaya, *Water Resources Research*, 49, 5117–5133, 2013.

Radić, V. and Hock, R.: Glaciers in the Earth's hydrological cycle: assessments of glacier mass and runoff changes on global and regional
scales, *Surveys in geophysics*, 35, 813–837, 2014.

Ragettli, S., Pellicciotti, F., Immerzeel, W., Miles, E., Petersen, L., Heynen, M., Shea, J., Stumm, D., Joshi, S., and Shrestha, A.: Unraveling
5 the hydrology of a Himalayan catchment through integration of high resolution in situ data and remote sensing with an advanced simulation
model, *Advances in Water Resources*, 78, 94–111, 2015.

Rawls, W. J., Brakensiek, D., and Saxton, K.: Estimation of soil water properties, *Transactions of the ASAE*, 25, 1316–1320, 1982.

Rees, H. G. and Collins, D. N.: Regional differences in response of flow in glacier-fed Himalayan rivers to climatic warming, *Hydrological
processes*, 20, 2157–2169, 2006.

10 Robson, B. A., Nuth, C., Dahl, S. O., Hölbling, D., Strozzi, T., and Nielsen, P. R.: Automated classification of debris-covered glaciers
combining optical, SAR and topographic data in an object-based environment, *Remote Sensing of Environment*, 170, 372–387, 2015.

Rowan, A. V., Egholm, D. L., Quincey, D. J., and Glasser, N. F.: Modelling the feedbacks between mass balance, ice flow and debris transport
to predict the response to climate change of debris-covered glaciers in the Himalaya, *Earth and Planetary Science Letters*, 430, 427–438,
2015.

15 Savéan, M., Delclaux, F., Chevallier, P., Wagnon, P., Gonga-Saholiariliva, N., Sharma, R., Neppel, L., and Arnaud, Y.: Water budget on the
Dudh Koshi River (Nepal): uncertainties on precipitation, *Journal of Hydrology*, 531, 850–862, 2015.

Shea, J., Immerzeel, W., Wagnon, P., Vincent, C., and Bajracharya, S.: Modelling glacier change in the Everest region, Nepal Himalaya, *The
Cryosphere*, 9, 1105–1128, 2015.

Sherpa, S. F., Wagnon, P., Brun, F., Berthier, E., Vincent, C., Lejeune, Y., Arnaud, Y., Kayastha, R. B., and Sinisalo, A.: Contrasted surface
20 mass balances of debris-free glaciers observed between the southern and the inner parts of the Everest region (2007–15), *Journal of
Glaciology*, pp. 1–15, 2017.

Shrestha, A. B., Wake, C. P., Dibb, J. E., Mayewski, P. A., et al.: Precipitation fluctuations in the Nepal Himalaya and its vicinity and
relationship with some large scale climatological parameters, *International Journal of Climatology*, 20, 317–327, 2000.

Shukla, A., Gupta, R., and Arora, M.: Estimation of debris cover and its temporal variation using optical satellite sensor data: a case study in
25 Chenab basin, Himalaya, *Journal of Glaciology*, 55, 444–452, 2009.

Singh, P.: *Snow and glacier hydrology*, vol. 37, Springer Science & Business Media, 2001.

Singh, S., Kumar, R., Bhardwaj, A., Sam, L., Shekhar, M., Singh, A., Kumar, R., and Gupta, A.: Changing climate and glacio-hydrology in
30 Indian Himalayan Region: a review, *Wiley Interdisciplinary Reviews: Climate Change*, 7, 393–410, 2016.

Sirguey, P., Mathieu, R., and Arnaud, Y.: Subpixel monitoring of the seasonal snow cover with MODIS at 250 m spatial resolution in the
35 Southern Alps of New Zealand: Methodology and accuracy assessment, *Remote Sensing of Environment*, 113, 160–181, 2009.

Soncini, A., Bocchiola, D., Confortola, G., Minora, U., Vuillermoz, E., Salerno, F., Viviano, G., Shrestha, D., Senese, A., Smiraglia, C., et al.:
Future hydrological regimes and glacier cover in the Everest region: The case study of the upper Dudh Koshi basin, *Science of the Total
Environment*, 565, 1084–1101, 2016.

Vincent, C., Wagnon, P., Shea, J. M., Immerzeel, W. W., Kraaijenbrink, P., Shrestha, D., Soruco, A., Arnaud, Y., Brun, F., Berthier, E., and
Sherpa, S.: Reduced melt on debris-covered glaciers: investigations from Changri Nup Glacier, Nepal, *The Cryosphere*, 10, 1845, 2016.

Viviroli, D., Dürr, H. H., Messerli, B., Meybeck, M., and Weingartner, R.: Mountains of the world, water towers for humanity: Typology,
mapping, and global significance, *Water resources research*, 43, 2007.

760 Wagnon, P., Vincent, C., Arnaud, Y., Berthier, E., Vuillermoz, E., Gruber, S., Ménégoz, M., Gilbert, A., Dumont, M., Shea, J., et al.: Seasonal
and annual mass balances of Mera and Pokalde glaciers (Nepal Himalaya) since 2007, *The Cryosphere*, 7, 1769–1786, 2013.

Weiler, M., Seibert, J., and Stahl, K.: Magic components—why quantifying rain, snowmelt, and icemelt in river discharge is not easy,
Hydrological Processes, 32, 160–166, 2018.

Westrick, K. J., Storck, P., and Mass, C. F.: Description and evaluation of a hydrometeorological forecast system for mountainous watersheds,
Weather and Forecasting, 17, 250–262, 2002.

765 Whitaker, A., Alila, Y., Beckers, J., and Toews, D.: Application of the distributed hydrology soil vegetation model to Redfish Creek, British
Columbia: model evaluation using internal catchment data, *Hydrological processes*, 17, 199–224, 2003.

Wigmsta, M. S., Vail, L. W., and Lettenmaier, D. P.: A distributed hydrology-vegetation model for complex terrain, *Water resources research*,
30, 1665–1679, 1994.

770 Wolff, M., Isaksen, K., Petersen-Øverleir, A., Ødegaard, K., Reitan, T., and Brækkan, R.: Derivation of a new continuous adjustment function
for correcting wind-induced loss of solid precipitation: results of a Norwegian field study, *Hydrology and Earth System Sciences*, 19, 951,
2015.

Wortmann, M., Bolch, T., Krysanova, V., and Buda, S.: Bridging glacier and river catchment scales: an efficient representation of
glacier dynamics in a hydrological model, *Hydrol. Earth Syst. Sci. Discuss.*, 2016, 1–37, doi:10.5194/hess-2016-272, <http://www.hydrol-earth-syst-sci-discuss.net/hess-2016-272/>, 2016.

775 Zhao, Q., Liu, Z., Ye, B., Qin, Y., Wei, Z., and Fang, S.: A snowmelt runoff forecasting model coupling WRF and DHSVM, *Hydrology and
Earth System Sciences*, 13, 1897–1906, 2009.

Zhu, Z. and Woodcock, C. E.: Object-based cloud and cloud shadow detection in Landsat imagery, *Remote Sensing of Environment*, 118,
83–94, 2012.