



New profiling and mooring records help to assess variability of Lake Issyk-Kul and reveal unknown features of its thermohaline structure

5 Peter O. Zavialov¹, Alexander S. Izhitskiy¹, Georgiy B. Kirillin², Valentina M. Khan³, Boris V. Konovalov¹, Peter N. Makkaveev¹, Vadim V. Pelevin¹, Nikolay A. Rimskiy-Korsakov¹, Salmor A. Alymkulov^{4,5}, and Kubanychbek M. Zhumaliev⁴

¹Shirshov Institute of Oceanology, Moscow, 117997, Russia

²Leibnitz Institute of Freshwater Ecology and Inland Fisheries, Berlin, 12587, Germany

10 ³State Hydrometeorological Research Center, Moscow, 123242, Russia

⁴International University of Innovative Technologies, Bishkek, 119991, Kyrgyzstan

⁵Institute of Physical and Technical Problems and Material Science, Bishkek, 720071, Kyrgyzstan

Correspondence to: Peter O. Zavialov (peter@ocean.ru)

Abstract. This article reports the results of 3 field campaigns conducted in Lake Issyk-Kul in 2015, 2016, and 2017. Vertical CTD profiles were obtained and water samples collected at the total of 75 stations. Some biogeochemical and thermohaline parameters at the lake surface were also mapped at high horizontal resolution along the ship's track. In addition, thermistor chains were deployed at 3 mooring stations in the eastern littoral region of the lake, yielding 147 days long records of temperature data. The measurements revealed that while the thermal state of the active layer, as well as some biogeochemical characteristics, were subject to significant interannual variability mediated by atmospheric forcing, the haline structure of the entire lake was remarkably stable at the interannual scale. Our data do not confirm the reports of progressive warming of the deep Issyk-Kul waters as suggested in some previous publications. However, they do indicate positive trend of salinity in the lake's interior over the last 3 decades. A noteworthy newly found feature is a weak but persistent salinity maximum below the thermocline at the depth of 70-120 m, from where salinity slightly decreased downwards. The data from the moored thermistor chains support previously published hypothesis about significant role of the submerged ancient riverbeds on the eastern shelf in advecting littoral waters into the deep portion of the lake during differential cooling period. We hypothesize that the less saline littoral water penetrating into the deep layers due to this mechanism is responsible for the abovementioned features of salinity profile, and substantiate this hypothesis by estimates based on simple model assumptions.

1 Introduction

30 With its volume of 1,730 km³, Lake Issyk-Kul is the tenth largest by volume lake on Earth overall, and the second largest mountain lake. It is also the sixth deepest lake of the World. Amazingly enough, even until recently, the information on the maximum depth of the lake in the available literature has been contradictory. For example, the official navigation map published by the USSR General Staff in 1978 indicated 663 m, Hoffer et al. (2002) suggested 650 m, the value of 668 m was given by Giralt et al. (2004) and a number of other publications, while some authors (e.g., Alamatov and Mikkola, 2011), as well as assorted sources on the Internet, including the Russian-language Wikipedia, give value of 702 m. However, the most plausible is the maximum depth of 668 m, as it was established based on high quality echo sounding data (De Batist et al., 2001).



As an enclosed terminal water body, Lake Issyk-Kul is sensible to climate change in its catchment area. Some summarizing estimates of the water budget components can be found in (Romanovsky, 2002; Kulenbekov and Merkel, 2012), for example, however, there are large uncertainties in all components. From 1927 to 1997, the Issyk-Kul water level decreased by 3.4 m, and then increased by about 1 m from 1997 on (e.g., Romanovsky et al., 2013). The effects of climate change on thermohaline structure of the lake are less well documented because the data are sparse, especially those for the last 3 decades. Romanovsky et al. (2013) reported a positive trend of temperature in the deep layers of almost 0.5°C for the period 1983–2003, along with substantial increase of surface temperature and river discharges. The interannual and interdecadal variability of salinity in the Issyk-Kul is largely unknown.

The Issyk-Kul is an oligotrophic lake (and, according to the classification of some authors – even ultra-oligotrophic). The level of primary production in it and the concentration of phytoplankton biomass are 10–40 times lower than those characteristic of the World Ocean on average (e.g., Antal, 2000). This affects all parts of the food chain and is reflected in the relatively low biodiversity (Mikkola, 2012). The potential of biological productivity of aquatic objects is often quantified by the ratio of the amount of incoming nutrients to the total volume of water. Since the Issyk-Kul water volume is very large, and the supply of nutrients is restricted to the inflow from relatively small rivers, the oligotrophy of the lake is easily explained from this point of view. However, this argumentation is valid only if the nutrients are distributed throughout the entire water mass of the lake; if, however, most of the nutrients are retained in the upper euphotic layer, the expectancy of biological productivity improves significantly. Thus, the nutrient budgets and biological productivity of the Issyk-Kul are intimately connected with the mixing regimes of the lake. It has been shown by tracer analysis methods that vertical mixing of the Issyk-Kul at the interannual scales is very intense and deep waters renew rapidly, with residence time of only about 11 years or less (Hoffer et al., 2002). The question about the mechanisms of such a fast renewal remains open.

An important role in the horizontal and vertical exchanges of the Issyk-Kul waters may belong to peculiar features of the bottom relief, in particular, the submerged beds of paleo rivers on the extensive eastern shelf of the lake. It is well established that the Issyk-Kul has experienced regressions and transgressions of up to 400 m in the past (e.g., Gebhardt et al. 2017). In particular, during late Pleistocene, the Issyk-Kul level was about 100 m below the current standing. At that time, today's shallow areas in the eastern and western parts of the lake were land. The rivers presently known as the Jyrgalan and the Tyup have left channels, or canyons, on today's lake bottom, buried 20–70 m in relation to the surrounding flat landscape. These channels are well preserved and distinctly visible over many kilometers. At the shelf break, the channels connect to the deep central part of the lake. According to the hypothesis put forward by (Peeters et al. 2003), mixing of the lake at seasonal time scales begins in autumn with differential cooling, when the water in the shoals cools to lower temperatures than that in the deep part of the Issyk-Kul. Further, due to density convection, this cold water descends into the submerged river beds on the shelf, along which it then flows downslope to the deep part of the lake and sinks there to its isopycnic level in the abyssal layers. This hypothesis, however, is yet to be confirmed by direct measurements.

Lake Issyk-Kul is of general scientific interest also because its bottom sediment deposits potentially hold unique information on climate change in continental Eurasia for millions of years. The possibility of organizing deep drilling in the Issyk-Kul in order to collect cores providing a climate reconstruction for 200–300 thousand years has been widely discussed (e.g., Oberhansli and Molnar, 2012). In this context, in order to determine the most informative locations for prospective drilling and correctly interpret the results, it is necessary to understand well the features of the Issyk-Kul hydrology and the processes of suspended matter transport in the lake's water mass.

Lake Issyk-Kul has attracted researchers for centuries, and the most intense studies were underway in the decades of the 1950s through the 1980s. Much of the data obtained at the times were summarized in monographs by Kadyrov



80 (1986) focused on hydrochemical issues, and by Romanovsky (1991) centered on the hydrology of the lake. Both
books, however, were published in Russian language in small series, and now became bibliographic rarities. The
amount of data collected from the Issyk-Kul decreased dramatically in the 1990s after the disintegration of the USSR.
In the 2000s and 2010s, a handful of field campaigns in the lake took place within some international projects.
However, overall, the data coverage remains very sparse. This especially refers to salinity data. During the Soviet era,
85 salinity was determined mainly analytically from water samples, either as a sum of ions or through the dry residue
method. This approach yielded a wealth of robust data, but their precision and spatial resolution were insufficient,
especially considering that salinity variations in the Issyk-Kul, both spatial and temporal, are generally small.
Nowadays, CTD profiling is used, but converting conventional CTD data to true salinity is not straightforward because
of the ionic composition of the Issyk-Kul water, which is significantly different from that of the ocean water. An
90 empirical formula derived from high accuracy laboratory measurements was proposed by Peeters et al. (2003) who
conducted two rather extensive CTD surveys of the lake in the early 2000s, but the conversion has not been widely
used since then.

In the present paper, we report principal findings of three recent field campaigns, focusing on thermohaline structure
and its variability. We use the abovementioned empirical formula to obtain and investigate the true salinity fields. We
95 also address hydrochemical parameters. In a separate section, we analyze water temperature records from thermistor
chains deployed inside and outside the submerged riverbeds on the shelf during the cooling period in an attempt to
quantify the role of the river canyons in advecting the cold water on the shelf towards the deep basin.

2 Data

The data used in this study were collected in three consecutive cruises of R/V Moltur. The first survey took place on
100 June 23 through June 27, 2015, the second on October 31 through November 3, 2016, and the third on June 25 through
July 1, 2017. A total of 75 stations were occupied (Fig. 1). At each station, CTD profiling and water sampling were
performed. A shallow water CTD instrument YSI6600 was used in 2015, so profiling in this survey was confined to
the uppermost portion (70 m) of the water column. In the surveys of 2016 and 2017, a SeaBird's SBE19plus Seacat
instrument allowing for deep profiling was used, but because of rough weather and drift of the vessel leading to
105 inclination of the line, the maximum depth achieved were 570 m in 2016 and 481 m in 2017. Water samples were
taken using HydroBios 5-l Niskin bottles in all surveys at all stations at different depth levels from surface to the
bottom with the depth increment varying from 20 m to 50 m. The samples were analyzed in laboratory using standard
techniques (Winkler method for oxygen; potentiometric method for pH; titration method for alkalinity; colorimetric
assay method for phosphates, silicates, nitrates, and nitrites).

110 In addition to the measurements at stations and moorings, continuous measurements of chlorophyll-“a” and total
suspended matter using ultraviolet fluorescent LiDAR UFL-9 were conducted along the ship's track. The details of
the LiDAR and methodology of this kind of measurements providing horizontal resolution of about 10 m were
described by Palmer et al. (2013), Pelevin et al. (2017), Zavialov et al. (2017). In the cruise of 2016, the continuous
along-track measurements also included soundings of temperature and electric conductivity in the uppermost layer
115 using a pump-through system and the shipboard YSI6600 CTD instrument, which allowed for horizontal resolution
in the along-track direction of the order of 100 m. As mentioned above, in all cases we used the empirical formula
proposed by Peeters et al. (2003) to infer salinity from temperature, electric conductivity, and pressure.

Moreover, 3 mooring stations equipped with thermistor chains were deployed on November 1, 2016, and then
recovered on March 27, 2017. Hence, the stations functioned for 147 days, registering water temperature at different



120 depth levels as 1 min averages. At the southern mooring station deployed in the paleo-bed of the Jyrgalan River where
the total depth was 50.7 m, thirteen RBR TR-1060 temperature sensors were installed at the depth levels of 1.4 m,
5.7 m, 11.8 m, 15.8 m, 19.9 m, 23.9 m, 29.0 m, 32.0 m, 38.1 m, 42.1 m, 46.1 m, 48.2 m, and 49.2 m, and one RBR
TDR-2050 temperature and pressure sensor at 50.6 m depth. At the northern mooring in the Tyup River paleo-bed at
125 the depth 50.2 m, four RBR TR-1060 thermistors were fixed only in the lower portion of the water column at the
levels 19.6 m, 29.7 m, 39.7 m, and 50.0 m. Finally, at the reference mooring station between the two channels at the
depth of 23.0 m, three RBR TR-1060 temperature sensors were installed at 3.1 m, 9.9 m, 16.7 m, and one RBR TDR-
2050 sensor at 23.0 m depth. Prior to the deployment of the moorings, bathymetry mapping was conducted using
FortXXI-Scat300 two-beam echosounder to obtain detailed bottom topography.

3 Results and discussion

130 3.1 Vertical thermohaline structure and its variability

The vertical structure of temperature and salinity as observed on October 31, 2016, and June 29, 2017, in the deepest
area of the Issyk-Kul is shown in Fig. 2. As mentioned above, in 2016, the instrument reached the depth of 570 m,
while in 2017, we were only able to obtain the profiles to the depth of 481 m. Prior to plotting the salinity data, the
data were smoothed using 6 m wide boxcar filtering to remove spikes.

135 Also shown in Fig. 2 are the respective historical data from literature sources. The newly obtained data indicate that
significant interdecadal warming of deep waters of the lake reported by Romanovsky et al. (2013) as about 0.25°C per
decade for the period 1983-2003 is apparently no longer there, as the present-day temperature at 500 m depth is exactly
same as it was in 2003 (namely, 4.44°C). We did observe positive trend of salinity in deep waters, where it has
increased by about 0.05 g kg⁻¹ since 1984, and about 0.02 g kg⁻¹ since 2001. However, according to the literature data
140 (Kadyrov 1986), in 1947 the near-bottom salinity was as high as 6.11 g/kg, which is about 0.09 g kg⁻¹ higher than it
is now.

The recent temperature profiles (red curves in Fig. 2a) include rather steep thermocline where temperature dropped
from about 12°C in 2016 and 17°C in 2017 near the surface to about 5°C at the depth of 80 m and 60 m, respectively.
Farther towards the bottom, the temperature decreases only slightly to 4.44°C at the depth 480 m and 4.37°C at the
145 depth 570 m. The differences between the profiles of 2016 and 2017 practically vanished at the depths below 180 m.
The salinity profiles (blue curves in Fig. 2a) are somewhat more complex than the temperature ones, although the
range of salinity change in the vertical direction is quite small – all values lie within the interval smaller than 0.07
g/kg between 5.960 g kg⁻¹ and 6.026 g kg⁻¹. In 2016, salinity increased from 5.960 g kg⁻¹ at surface to 6.026 g kg⁻¹ at
81 m and then decreased towards the bottom to 6.015 g kg⁻¹ at 570 m. In 2017, salinity first decreased from about
150 5.996 g/kg at the surface to a minimum of 5.990 g/kg at the upper edge of the thermocline at the depth of 15.5 m, and
then increased downwards slightly until it attained the maximum value of 6.019 g/kg at the depth about 125 m (see
the inset panel in Fig. 2). From there and down to the bottom, the salinity was almost constant, but showed extremely
slight decrease to 6.015 g kg⁻¹ at the depth of 481 m.

Therefore, we report here the existence of very weak but observable salinity maximum at intermediate depths below
155 the thermocline. This maximum is too small in magnitude to have been detected by techniques used in the Soviet
period of the lake's exploration, so it has never been reported at the times. However, it appears to be a robust and
persistent feature – a similar behavior of salinity was evident in the data of 2001 presented by Peeters et al. (2003),
although these authors did not discuss the intermediate salinity maximum explicitly. In their observations, the depth



of the maximum varied from 200 to 300 m, and then below this layer the salinity decreased slightly by about 0.002 g/kg towards the bottom.

The TS diagrams for waters below the thermocline for 2016 and 2017 shown in Fig. 2b also clearly indicate the intermediate salinity maxima at 5.5°C, 6.026 g kg⁻¹ for 2016, and 4.8°C, 6.02 g kg⁻¹ in 2017. It is interesting that in the deepest layer, there is a hint of salinity increase below 4.38°C, that is, below 540 m.

The observed salinity inversions are very small in magnitude and do not have any significant effect on density as compared to the effect of temperature stratification, so the water column remains convectively stable. The possible dynamical origin of the intermediate salinity maximum beneath the thermocline will be discussed below in the subsection dedicated to winter cooling on the eastern shelf.

Vertical profile of turbidity obtained in June, 2017, is exhibited in Fig. 2c. It shows distinct maximum at the depth of about 40 m at the upper edge of the thermocline, likely associated with the maximum concentration of phytoplankton (see also Fig. 13). Under the thermocline, turbidity decreased downwards and was nearly constant below 200 m, although some spike-like positive anomalies were seen in the lower portion of the water column.

3.2 Horizontal thermohaline structure and its variability

The salinity and temperature distributions at the surface derived from continuous pump-through CTD measurements of November, 2016, are shown in Fig. 3. The salinity map is virtually the most detailed ever obtained for the Issyk-Kul. In fact, the horizontal variability of the salinity field is remarkably small – with the exception of the areas immediately adjacent to river mouths, the total salinity span over the entire lake does not exceed mere few hundredths of g kg⁻¹ (more precisely, about 0.04 g kg⁻¹). This is indicative of intense mixing processes taking place in the lake.

However, the discharges from the Jyrgalan and the Tyup rivers, bringing to the lake respective 28 m³s⁻¹ and 12 m³s⁻¹ of fresh water on long-term average (Romanovsky, 1990), do exhibit distinct signature in salinity on the eastern shelf, forming a frontal zone at about 78°E. The haline structure in this area is depicted in Fig. 4 based on CTD stations shown by the green bullets in Fig. 1. As it can be seen from Fig. 4, the plume of relatively low salinity water associated with the Jyrgalan River runoff extends for over 20 km from the river inlet, mainly confined to the upper 20 m portion of the water column. Some near-coast regions of freshwater influence from smaller rivers are also evident at the southern and the northern extremities of the Issyk-Kul.

Other notable feature is represented by slightly elevated salinity values in the central part of the lake, which is consistent with the traditional notion of the cyclonic basin-scale circulation resulting in a dome-like structure in the central region where saltier waters below the thermocline come closer to the surface (e.g., Romanovsky, 1991). It is interesting that this area of higher salinity seems to split in a bimodal structure, with one maximum in the central northeastern and the other one in the central southwestern parts of the lake. This gives a hint of two sub-basin scale cyclonic gyres superimposed on the general cyclonic circulation. Such sub-basin gyres are rather common for enclosed seas (e.g., Tuzhilkin, 2010), but have never been established for the Issyk-Kul.

For 2015 and 2017, the available salinity data are restricted to hydrographic stations, so the information about the horizontal distributions is less detailed. Generally, the interpolated surface salinity fields exhibited patterns similar to those characteristic of 2016, namely, very limited (about 0.06 g kg⁻¹) range of spatial variability across the lake area (except in the region of freshwater influence on the eastern shelf), and slight but distinct salinity maximum in the central part of the lake. Figure 5 presents surface salinity field in June of 2017. We omit here similar plot for June of 2015, and instead show in Fig. 6 instead the differences in salinity between June of 2017 and June of 2015. Overall, the observed changes in surface salinity over the 2 years were small, they never exceeded a few hundredths of g kg⁻¹,



and totaled to essentially zero on the areal average. Interestingly, salinity increased in the eastern part of the lake more strongly affected by river discharges, but decreased in the western part. The maximum difference slightly over 0.04 g kg^{-1} was observed in the southeastern sector, namely, the Pokrovsky Bay adjacent to the mouths of the Dzhooku and the Kichene-Kyzylsu Rivers.

The interannual differences observed in the lake surface temperature were more significant than those in salinity. The temperature fields for June of 2017 and June of 2015 are not shown here, while their differences are highlighted in Fig. 7. All over the lake, the surface water temperature in June of 2017 was warmer than that in June of 2015. While amounting to about 0.9°C on the areal average, this misfit, however, was distributed unevenly over the lake area, with maximum changes taking place near the southwestern coast (Fig. 7).

Figure 8 shows ERA-Interim atmospheric reanalysis data (Dee et al., 2011) which indicate that from June 15 through June 24 (i.e., the period immediately preceding the measurements in the surveys), the daily mean near-surface air temperature in the study region was higher in 2017 than that in 2015 by $1\text{--}4^\circ\text{C}$. It also follows from the same reanalysis data that the total heat flux received by the lake surface from the atmosphere (including solar irradiance) for this period in 2017 was equal to about 161% of that in 2015. Therefore, the observed interannual differences in lake temperature are likely to reflect response to different conditions of atmospheric forcing.

3.3 Biogeochemical structure and its variability

3.3.1 Oxygen

During all surveys, the entire water column down to the bottom was fully ventilated. The content of dissolved oxygen varied from 6.1 to 8.4 ml l^{-1} , and its saturation from 69 to 123% (Fig. 9a). Also shown in the figure is the respective range for oxygen saturation for the period 1961-1985, which was from 79 to 117% (Kadyrov, 1986). No significant change in oxygenation can be substantiated, although in the present, both maximum and minimum values extend beyond the “old” range. The highest content of oxygen is usually situated at the upper edge of the thermocline. For surface water, the oxygen content ranged from 6.38 to 7.67 ml l^{-1} . The greatest interannual changes in the content of dissolved oxygen were observed at the upper layer exposed to the influence of continental runoffs and activity of the aquatic biota. Below 200 m, the content of oxygen and its saturation manifested no significant interannual differences.

3.3.2 Alkalinity and pH

Overall, during the three surveys, the total alkalinity ranged from 1.04 to 5.47 mgEq l^{-1} (Fig. 9b). However, low alkalinity values were confined exclusively to the areas adjacent to river mouths, while for the bulk of the lake waters, the span of alkalinity values was between 5.37 and 5.44 mgEq l^{-1} with the mean value 5.41 mgEq l^{-1} , which constitutes noteworthy increase with respect to the values reported by Kadyrov (1986). The mean pH was 8.68, nearly same as 8.71 in the 1980s. Small change in the alkalinity and pH along the vertical, and practically no change from one survey to another once again suggest that the entire water column in the lake mixes actively. In the central part of the lake, a slight increase in pH (by about 0.05 with respect to the average) and total alkalinity (by about 0.05 mgEq l^{-1} with respect to the average) at the surface was evident in horizontal distributions (not shown here). Most likely, this feature observed in all 3 years, as well as the abovementioned salinity maximum in the central area of the lake, is associated with the rise of waters due to cyclonic basin-scale circulation.



3.3.3 Phosphates

The content of phosphates in the lake waters was determined only in 2016 and 2017. The concentration of phosphates varied from the analytical zero (i.e., below the detection limit by the method) to 1.16 μM . The maximum value was detected in the surface waters at the eastern extremity of the lake, most likely due to the influence of continental discharge from the Jergalan and the Tyup rivers. Slightly elevated concentrations of phosphates were also observed in the central region, consistent with similar feature in salinity, pH, and alkalinity. All over the lake area, relatively high content of dissolved inorganic phosphorus was observed in the upper water layer down to about 50 m (Fig. 9c). This may be associated with the influx of phosphorus by runoffs from the coast, including sewage in urban areas, and/or with the oxidation of organic matter formed during the period of active photosynthesis. It is interesting that in the bottom layer, where the organic oxidation processes should take place, we did not observe any significant increase in the content of inorganic phosphorus nor noticeable decrease in dissolved oxygen and pH. Again, this is suggestive of efficient mixing of the water column.

During the November survey of 2016, the phosphate concentration in the layer from 200 m to the bottom varied from an analytical zero to 0.1 μM , which is lower than the existing (although sparse) historical data for the decade of the 1980s when the concentration typically ranged from 0.19 to 0.38 μM below 200 m (Kadyrov, 1986). However, in June of 2017, the phosphate content below the thermocline was significantly higher and totaled to an average of 0.4 μM . Most likely, this was due to oxidation of organic matter coming from the upper active layer into the deep waters of the lake following plankton bloom (see below).

3.3.4 Silicates

The content of dissolved silicates in the lake water varied from 64.8 to 96.1 μM (Fig. 9d). The maximum range of silicate content was observed in surface waters. Below 100 m depth, the average silicate content was about 79 μM , with a certain increase towards the bottom.

The silicates content profile observed in 2017 was close to that documented in 2015. In 2016, the silicate concentrations were higher by about 10% throughout the column. This may reflect seasonal changes, although a longer series of observations is needed to confirm it. However, the concentrations registered in all 3 surveys were notably lower than those reported by (Kadyrov, 1986) whose averages ranged from 86 to 95 μM .

The horizontal distributions of nutrients at the surface observed in all surveys generally looked alike and were characterized by distinct maxima on the eastern shelf in the areas adjacent to the Jyrgalan and Tyup estuaries, and smaller concentrations rather uniformly distributed elsewhere. An example for silicate concentration is shown in Fig. 10. The position of the frontal zone separating the low and high concentration zones is nearly coincident with the salinity front seen in Fig. 3. The observed distributions indicate that the discharges from the Jyrgalan and the Tyup are the principal if not the only source of nutrients supply to the lake.

3.3.5 Nitrates and nitrites

The content of nitrates in the water of the lake varied from 1.2 to 16.1 μM , and that of nitrites from analytical zero to 0.54 μM (Fig. 11), with the largest variability observed in the surface layer. Below the thermocline, nitrate concentration ranged from 1.2 to 12.9 μM . The respective span for nitrites was from 0 to 0.23 μM .

The values of nitrate content obtained during different surveys diverged quite significantly. In 2017, the concentrations were lower than those in 2015 and 2016, virtually throughout the column. In addition, in 2015 and 2016, a moderate



275 increase of nitrate content towards the bottom was manifested in the layers below 100 m, while in 2017, this pattern
was no longer there. As to nitrites, in 2015 and 2016, their concentrations below 100 m depth were very low (especially
in 2016, when they were close to analytical zero) and nearly uniform. In 2017, however, the content of nitrates in the
deep waters of the lake increased to 0.1 μM . In all 3 surveys, the highest content of nitrites was observed at the
“confluence” of the Tyup and Jyrgalan estuaries. In 2017, high content was observed not only in the eastern, but also
in the central part of the lake.

280 According to Kadyrov (1986), in the 1980s, the concentration of nitrates ranged from 2.6 to 16 μM , and that of nitrites
varied between 0.05 and 0.15 μM . Hence, no significant change since then is evident.

3.3.6 Chlorophyll-“a” and total suspended matter

The measurements of chlorophyll-“a” concentration and the content of the total suspended matter (hereinafter TSM)
in the surface layer are summarized in Fig. 12. The inset table in the upper part of the figure refers to the LiDAR
285 measurements encompassing total of 169,166 data points distributed all over the lake along the ship’s tracks, whilst
the bullets in the plot depict data from water samples collected at the stations. The samples were analyzed in laboratory
and the concentrations were determined through the procedures described by Konovalov et al. (2014). Also shown in
Fig. 12 is the so-called pigment index showing essentially the “age” or the life stage of the plankton (e.g., Margalef,
1967, Wozniak and Dera, 2007).

290 The LiDAR data indicate that while the maximum concentrations of TSM are observed in the eastern part of the Issyk-
Kul as it may have been expected because this part of the lake is stronger exposed to river discharges, the chlorophyll
concentrations demonstrate opposite pattern with maximum values pertaining to the western and especially the
northwestern regions. One striking feature evident in both the LiDAR and the in situ data is the fact that overall
chlorophyll concentrations in June of 2017 were much higher than those in June of 2015 by a factor of 5 on the
295 average, and in the southeastern part of the lake most exposed to river discharges even by a factor of 7.3. This
somewhat surprising difference cannot be explained by seasonal cycling, as the two surveys took place almost on the
same days of year. However, the pigment index may give some insights into this matter, showing that at the time of
the observations in 2015, the planktonic communities were represented mainly by “old”, post-bloom plankton, while
those in 2017 (as well as in November of 2016) – mainly by “young” plankton. This indicates that the spring bloom
300 of 2015 occurred earlier than that of 2017, and timing of plankton blooms in the Issyk-Kul is significantly variable
from year to year, perhaps depending on weather conditions and supply of nutrients by rivers. Differences that are
even more significant are documented in TSM whose average concentration in June 2017 exceeded that in June 2015
by a factor of 10.1. On the one hand, this indicates that the suspended matter in the Issyk-Kul is tightly connected with
planktonic communities, and at least a part of the TSM is of biogenic origin. On the other hand, this may point on
305 elevated river discharges bringing both TSM and nutrients immediately before and during the plankton bloom.

Vertical and longitudinal distribution of chlorophyll-“a” along the Jyrgalan canyon based on the sampling conducted
on November 1, 2016 (green bullets in Fig. 1) is shown in Fig. 13. The maximum concentration is observed near the
shore in proximity of the river mouth. To the west of the estuary, vertical structure of chlorophyll exhibits two maxima,
one at the upper boundary of the thermocline at the depth of about 20 m, and the other one near the bottom. The upper
310 maximum of plankton activity just below the mixed layer is characteristic for the entire lake and generally coincides
with the maximum concentration of nutrients. The near-bottom maximum is likely related to downslope advection of
chlorophyll-rich, cold waters from the eastern shelf during cooling period as discussed in the next subsection.



315 3.4 Seasonal cooling on eastern shelf and its possible implications

The data from the moored thermistor chains deployed on November 1 of 2016 and recovered on March 27 of 2017 gave some insights into details of how winter cooling progressed on the eastern shelf of the lake. As mentioned above, two moorings were installed into the paleo beds of the Jyrgalan and the Tyup Rivers, and one on the shelf between them. Fig. 14 shows bathymetric profile across the Jyrgalan and Tyup canyons at 77.9°E obtained by echo-soundings on June 25, 2015. The thermistor chains were deployed on a similar terrain east of this section.

The complete set of data from the thermistor chains is exhibited in Fig. 15, demonstrating gradual general cooling from about 13°C in early November to about 3°C in late February – early March, followed by the onset of spring warming in the mid-March. Also shown in the figure is the surface air temperature from the ERA-Interim reanalysis. Fig. 16 presents a “zoom” of the Jyrgalan canyon record for November, 2016. Both figures exhibit a number of distinct cooling events. Some of these episodes originate from the surface and then propagate downwards, presumably due to convective mixing, such as the events of the late December following abrupt decrease of the air temperature (Fig. 15), or the events of November 7, 17, and 22, clearly seen in Fig. 16, for example. In contrast, other events (November 5, 14, 26) appear in the bottom layer with no sign of equally cold water in the upper parts of the column, and, therefore, can only be attributed to lateral advection.

Closer look at the temperature records reveals that the general cooling was mediated by a large number of relatively short individual events. To investigate their statistics and evaluate the relative roles of advective and convective cooling, we performed the following analysis.

Hereinafter, we refer to the term “bottom cooling event” to identify the situation where temperature at the bottom sensor decreased continuously during a series of sequential readings. Because the sampling rate of the sensors was set to 1 min, only events lasting 2 min or more were taken into account. Furthermore, the event was considered “convective” if at the beginning of it, temperature at the bottom sensor was higher than that at the next sensor of the chain counting from the bottom, and “advective” otherwise. Because the observed temperatures in all cases were above the maximum density temperature, which is about 2.75°C for the Issyk-Kul water (Romanovsky 1991), and salinity stratification in the bottom layer was virtually negligible, the situation of temperature inversion implies convective instability. On the other hand, the absence of such inversion means that the observed decrease of temperature with time cannot be explained by vertical heat exchanges and hence must be attributed to lateral advection.

The entire set of data recorded by three thermistor chains has been analyzed accordingly. The resulting statistics is depicted in Table 1. The total duration of cooling at the bottom over the measurements period was different for the three locations and ranged from 60780 min (42.2 days) for the Jyrgalan canyon to 104981 min (72.9 days) for the Tyup canyon. The total duration of bottom cooling at the reference mooring station situated on the flat shelf between the canyons was 95783 min (66.5 days), which is higher than that for the Tyup canyon and only slightly lower than that for the Jyrgalan canyon. However, while on the shelf outside the canyons the convective and advective mechanisms were responsible for approximately equal number of cooling events, within the canyons the advective mechanism clearly prevailed, accounting for about two thirds of all bottom cooling events in the Tyup canyon, and almost 90% of bottom cooling events in the Jyrgalan canyon. It also appears that the individual advective events within the canyons tended to last longer than those on the flat shelf (maximum duration of 41 min versus 19 min) and produce stronger temperature drop (up to 1.36°C versus 0.29°C considering the maximum values, the average figures are much smaller but seem to be in similar relation). Thus, in this sense, the newly obtained data support the hypothesis



put forward by Peeters et al. (2001) that during the cooling period, the paleo river beds on the eastern shelf of the Issyk-Kul lake may serve as “pathways” guiding relatively cold and dense shelf water formed as a result of differential cooling towards the deep basin. The signature of this water is also seen in the distribution of chlorophyll-“a” concentration along the Jyrgalan canyon recorded on November 1, 2016 already presented above in Fig. 13. The vertical profile shown there is characterized by two maxima of chlorophyll. The first, centered at the depths 20-30 m just below the upper mixed layer, has been commonly observed in the Issyk-Kul lake (Kadyrov, 1986). But the stronger maximum at the bottom, beyond the euphotic layer where plankton normally develops, is likely to originate from surface water in the shallow parts of the shelf sinking to the bottom because of cooling and propagating westward down the slope within the canyon.

Basing on these findings and assuming that the advection of cold shelf waters is a dominant mechanism controlling winter cooling within the submerged river beds, we now can approximately estimate the scale v of the downslope velocity mediating this process. Indeed, in this case,

$$\frac{\partial T}{\partial t} \sim v \frac{\partial T}{\partial x}, \quad (1)$$

where T is the temperature of water in the canyon, t is time, and x is the coordinate along the canyon. The values of $\frac{\partial T}{\partial t}$ and $\frac{\partial T}{\partial x}$ as for the conditions of November, 2016, can be estimated from the data presented above in Figs. 4, 15 and 16 as $\frac{\partial T}{\partial t} \sim 2^\circ\text{C month}^{-1} \sim 10^{-6}^\circ\text{C/s}$, and $\frac{\partial T}{\partial x} \sim 1^\circ\text{C}/10\text{ km} \sim 10^{-4}^\circ\text{C/m}$. It follows then from the equation above that $v \sim 10^{-2}\text{ m s}^{-1}$. Furthermore, given that the cross-sectional area of the Jyrgalan canyon equals is of the order of 10^5 m^2 (Fig. 14), this suggests that the total volume of water that passed through the canyon towards the deep part of the lake during one month of autumn cooling is of the order of 0.1 km^3 .

Admixture of the cold and relatively fresh shelf water, originally at average salinity of about 5.8 g kg^{-1} (Fig. 4), which moves downslope and settles at its isopycnal level in the bulk of the water column, to nearly 6 g kg^{-1} salinity ambient water must result in certain decrease of salinity in the affected part of the column. If we assume that the inflowing water is distributed horizontally over the entire area of the deep floor of the lake ($\sim 4 \cdot 10^9\text{ m}^2$) and vertically throughout the column from the bottom to the depth of the intermediate salinity maximum discussed above, i.e., over $\sim 500\text{ m}$ thick layer, a simple balance estimate demonstrates that the resulting salinity drop in the lower portion of the water column should be of the order of 10^{-3} g kg^{-1} . This effect should be amplified by the presence of the inflow not only through the Jyrgalan canyon, but also through the Tyup canyon and, possibly, flat shelf in a similar process, besides, the duration of the cooling period is almost 4 months (Fig. 15). Therefore, one should expect that the salinity drop in the bottom part of the column due to admixture of cold shelf waters propagating through the Tyup and Jyrgalan canyons would be of the order of a few thousandths of $^\circ\text{C}$ – which is, indeed, evident in the observations.

Hence, our newly obtained data reveal previously unknown details on winter temperatures in the eastern littoral region and suggest a connection of the deep water renewal to horizontal salinity gradients in the Issyk-Kul. The lowest water temperatures observed in February near the lake bottom were in the range $2.4\text{--}3.0^\circ\text{C}$, which falls into the range of the temperatures of maximum density (TMD) for salinities found in Issyk-Kul. According to Romanovsky (1991), the TMD for the Issyk-Kul is 2.75°C . At the outer edge of the cold nearshore areas, a frontal zone of “thermal bar” (e.g., Zilitinkevich et al., 1992) forms in many lakes, dividing water masses at temperatures above and below of the TMD and characterized by strong downward convective flows. An indirect evidence of such flows in the Issyk-Kul is the transient appearance of cold waters with temperatures close to the TMD at the lake bottom with warmer waters above (cold tongues at the lake bottom in February, Fig. 15d). In contrast to large freshwater lakes (Malm et al., 1993;



Shimaraev et al., 1993; Ullman et al., 1998), in the Issyk-Kul, the movement of the thermal bar towards the center of the lake will be restricted by the background lateral salinity gradients formed by freshwater inflows. However, the same salinity gradients together with the bottom slope will ensure volume contraction by mixing of the colder and fresher waters sinking into the warmer and saltier deep pelagial, resulting in the prominent cabbeling effect (Fedorov, 1981, McDougall, 1987). By this, the vertical convective flow formed within the thermal bar will follow the bottom slope towards the warmer lake center with positive vertical acceleration due to the dependence of the thermal expansion coefficient on temperature. This suggested mechanism of deep water renewal differs from that reported in deep freshwater lakes, in particular, Lake Baikal (Killworth et al., 1996; Schmid et al., 2008), where downward vertical acceleration is created by the effect of thermobaricity (dependence of the thermal expansion coefficient on pressure). Still, the latter mechanism cannot be a priori assumed negligible in the ventilation of the Issyk-Kul waters: Killworth (1977) argued that thermobaricity is even more important than cabbeling for deep water formation due to downslope cold freshwater flows, since it provides faster increase of density with depth of fresher waters. In this sense, the Issyk-Kul may be illustrative of a number of physical mechanisms crucial for formation of deep water masses in the ocean, and thus represents a natural “intermediate-scale” laboratory for investigating the cabbeling and thermobaricity effects on deep circulation.

4 Conclusions

As one of the World's largest and deepest lakes, the lake Issyk-Kul reflects the ongoing climate change trends, and, reciprocally, exercises influence on climate, at least, at regional scales. It is, therefore, important to be aware of the variability of the lake. The physical and chemical regimes of the Issyk-Kul have been monitored on a regular basis during the Soviet era, although the information collected then was restricted to measurement techniques and equipment available at the times. For this reason, details of the structure and variability of salinity fields were poorly known. The amount of data obtained from the lake reduced strongly in the early 1990s, with only a few observational studies conducted since then. The aim of the present paper was to present new in situ data collected in three campaigns of 2015-2017, complemented by long-term deployments of moored thermistor chains.

The newly obtained data indicate that significant interdecadal warming of deep waters of the lake reported by Romanovsky et al. (2013) as almost 0.5°C for the period 1983-2003 is apparently no longer there, as the present-day temperature at 500 m depth is exactly same as it was in 2003 (namely, about 4.44°C). However, in deep waters, there did exist positive trend of salinity, which has increased by about 0.05 g kg⁻¹ since 1984, and about 0.02 g kg⁻¹ since 2001.

A pump-through CTD system and use of empirical formula for converting CTD data to true salinity allowed obtaining high resolution salinity maps with some features not resolved in previous measurements, such as a bimodal structure of elevated salinity pattern in the central part of the lake suggesting sub-basin scale cyclonic gyres. Generally, the Issyk-Kul's salinity is amazingly close to constant, even in the surface layer exposed to river discharges. In fact, all of its horizontal variability over the entire lake area (except the river mouths) observed during the cruises of 2015, 2016, and 2017 fitted into the interval of about 0.3 g kg⁻¹, vertical variability was mere 0.03 g kg⁻¹, and the maximum temporal variability at fixed points at the interannual scale was only 0.04 g kg⁻¹. This conservative behavior of salinity is unusual for inland water bodies and points on very efficient horizontal and vertical mixing. However, we did observe some slight redistributions of salinity fields from one year to another.



In contrast to the salinity, the temperature of the surface layer is subject to significant interannual variability controlled by atmospheric forcing conditions. This leads also to changes in timing of phytoplankton blooming in early summer, as expressed in the concentration of chlorophyll-“a”.

The new data revealed no significant long-term changes in nutrients and other hydrochemical constituents, except for silicates whose present-day concentrations throughout the water column are notably lower than those reported for the decades of 1970s and 1980s, and alkalinity, which appears to have increased by about 0.2 mEq l⁻¹.

Finally, we investigated in detail the development of seasonal cooling on the eastern shelf and the role of the paleo river beds in this process. Generally, cooling of the bottom layer occurred through a large number of individual cooling events, some of which originated from convective mixing from above, and others were due to lateral advection along the slope. The measurements demonstrated that while on the flat shelf outside the river canyons the advective and convective cooling events were equally significant, the canyons themselves were dominated by advection, resulting in more efficient cooling. We can conclude, therefore, that the newly obtained data implicitly support the hypothesis by Peeters et al. (2003) that advection through the paleo river beds following differential cooling on the eastern shelf plays an important role in modulating vertical structure in the deep part of the lake.

In this respect, we emphasize as an important finding of this study the intermediate salinity maximum observed below the thermocline, from where the salinity slightly decreased to the bottom. Its depth varied from year to year between 70 and 125 m. This maximum is so weak (0.004-0.009 g kg⁻¹) that it could not possibly have been detected using the techniques of the Soviet era, so it has never been reported at the times. However, the intermediate salinity maximum seems to be a robust and persistent feature. Noteworthy, the opposite signs of temperature and salinity contributions to the density gradients beneath the salinity maximum suggest potential development of double-diffusive convection, which, at salinity decreasing with depth, should take form of salt fingers (Schmitt, 1994) and may additionally contribute to the vertical mixing at water depths below ~100 m. The dynamical origins of the intermediate salinity maximum are likely to be related to the intrusion of lower salinity, lower temperature water from the eastern littoral during the differential cooling period. The observed salinity drop seems to be consistent with the volume of such intrusion estimated independently from thermistor chain measurements.

Acknowledgments

The field work of 2017 and the respective data analyses were funded by the Russian Foundation for Basic Research (Grant No. 17-05-41043). The field campaigns of 2015-2016 were funded by the Russian Science Foundation (Grant No. 14-50-00095). The measurements at the long-term mooring stations and the respective data analyses were supported by the Russian Ministry of Education and Science (Agreement No. 14.W03.31-0006 and State Assignment Theme No. 0149-2018-0002).

References

- Alamatov, A. and Mikkola, H.: Is biodiversity friendly fisheries management possible on Issyk-Kul lake in the Kyrgyz Republic? *AMBIO: A Journal of the Human Environment*, 40(5): 479-495, 2011.
- Antal, T.K.: A study of primary production and phytoplankton characteristics using a submersible fluorescent probe. Ph.D. Thesis, Moscow State University, Department of Biology, Moscow, Russia, 2000.



- De Batist, M., Imbo, Y., Vermeesch, P., Klerkx, J., Giralt, S., Delvaux, D., Lignier, V., Beck, C., Kalugin, I., and
480 Abdrakhmatov, K.E.: Bathymetry and sedimentary environments of Lake Issyk-Kul, Kyrgyz Republic (Central Asia):
A large, high-altitude, tectonic lake. IN: Klerkx, J. and Imanackunov, B. (Eds.), *Lake Issyk-Kul: Its Natural
Environment*. Dordrecht (NATO Science Series, Kluwer), 101–123, 2001.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A.,
Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani,
485 R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler,
M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P.,
Tavolato, C., Thépaut, J.-N. and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data
assimilation system. *Q.J.R. Meteorol. Soc.*, 137: 553–597. doi: 10.1002/qj.828, 2011.
- Fedorov, K. N.: The role of compaction during mixing in the dynamics of oceanic fronts. *Doklady Akademii Nauk
490 SSSR*, 261: 985–988, 1981.
- Gebhardt, A.C., Naudts L., De Mol L., Klerkx J Abdrakhmatov., K., Sobel E.R., and De Batist M.: High-amplitude
lake-level changes in tectonically active Lake Issyk-Kul (Kyrgyzstan) revealed by high-resolution seismic reflection
data. *Clim. Past.*, 13: 73–92, doi:10.5194/cp-13-73-2017, 2017.
- Giralt, S., Klerkx, J., Riera, S., Julia, R., Lignier, V., Beck, C., De Batist, M., and Kalugin, I.: Recent
495 paleoenvironmental evolution of Lake Issyk-Kul. In Klerkx, J., and Imanackunov, B. (Eds.), IN: *Lake Issyk-Kul: Its
Natural Environment*, Dordrecht (NATO Science Series, Kluwer), 125–145, 2002.
- Hofer, M., Peeters F., Aeschbach-Hertig W., Brennwald M., Holocher J., and Livingstone D.M.: Rapid deep-water
renewal in Lake Issyk-Kul (Kyrgyzstan) indicated by transient tracers. *Limnol. Oceanogr.* 47: 1210 – 1216, 2002.
- Kadyrov, V.K.: *Hydrochemistry of Lake Issyk-Kul and its Catchment*. Frunze, Ilim, 212 pp., 1986 [in Russian].
- 500 Killworth, P. D.: Mixing of the Weddell Sea continental slope. *Deep Sea Research*, 24: 427–448, 1977.
- Killworth, P.D., Carmack E.C., Weiss R.F., and Matear R.: Modeling deep-water renewal in Lake Baikal. *Limnol.
and Oceanogr.*, 41: 1521–1538, 1996.
- Konovalov, B.V., Kravchishina M.D., Belyaev N.A., and Novigatsky A.N.: Determination of the concentration of
mineral particles and suspended organic substance based on their spectral absorption. *Oceanology*, 54: 5, 660–
505 667, 2014.
- Kulembekov, Zh. and Merkel B.J.: Investigation of the natural uranium content in the Issyk-Kul lake, Kyrgyzstan.
Freiberg Online Geology, 33: 3–45, 2012.
- Lyons, W. B., Welch K. A., Bonzongo J.-C., Graham E. Y., Shabunin G., Gaudette H. E., and Poretta R. J.: A
preliminary assessment of the geochemical dynamics of Issyk-Kul Lake, Kirghizstan. *Limnol. Oceanogr.* 46: 713–
510 718, 2001.
- Malm, J., Grahn L., Mironov D., and Terzhevik A.: Field investigation of the thermal bar in Lake Ladoga, spring
1991. *Hydrology Research*, 24: 339–358, 1993.
- Margalef, R.: Some concepts relative to the organization of plankton. *Oceanogr. Mar. Biol. Ann. Rev.*, 5: 257–289,
1967.
- 515 McDougall, T.J.: Thermobaricity, cabbeling, and water-mass conversion. *Journal of Geophysical Research*, 92(C5):
5448–5464, 1987.
- Mikkola, H.: Implication of alien species introduction to loss of fish biodiversity and livelihoods on Issyk-Kul Lake
in Kyrgyzstan. IN: *Biodiversity Enrichment in a Diverse World*, G.A. Lameed (Ed.), ISBN 978-953-51-0718-7, 395–
419, 2012.



- 520 Oberhansli, H. and Molnar P. Climate evolution in Central Asia during the past few million years: A case study from Issyk-Kul. *Scientific Drilling*, 13: doi: 10.2204/iodp/sd.13.09.2011, 2012.
- Palmer, S.C.J., Pelevin V.V., Goncharenko I., Kovács A.W., Zlinszky A., Présing M., Horváth H., Nicolás-Perea V., Balzter H., and Tóth V.R.: Ultraviolet fluorescence LiDAR (UFL) as a measurement tool for water quality parameters in turbid lake conditions. *Remote Sens.* 5: 4405-4422; doi:10.3390/rs5094405, 2013.
- 525 Pelevin V., Zlinszky, A., Khimchenko, E., Toth, V.: Ground truth data on chlorophyll-a, chromophoric dissolved organic matter and suspended sediment concentrations in the upper water layer as obtained by UFL LiDAR at high spatial resolution. *International Journal of Remote Sensing*, 38: 7, 1967-1982, doi: 10.1080/01431161.2016.1274446, 2017.
- Peeters, F., Finger D., Hofer M., Brennwald M., Livingstone D.M. and Kipfer R.: Deep-water renewal in Lake Issyk-Kul driven by differential cooling. *Limnol. Oceanogr.* 48: 1419- 1431, 2003.
- 530 Romanovsky, V.V.: Lake Issyk-Kul as natural complex. Ilim, Frunze, 164 pp, 1991 [in Russian].
- Romanovsky, V.V.: Water level variations and water balance of Lake Issyk-Kul. In: J. Klerkx. and B. Imanackunov (Editors). *Lake Issyk-Kul: its natural environment*. NATO Science Series, Kluwer Academic Publishers, Netherlands, Earth and Environmental Sciences 13: 45 – 58, 2002.
- 535 Romanovsky, V.V., Tashbaeva S., Creteaux J.-F., Calmant S., and Drolon V.: The closed lake Issyk-Kul as an indicator of global warming in Tien-Shan. *Natural Science*, 5: 5, 608-623, doi 10.4236/ns.2013.55076, 2013.
- Shimaraev, M. N., Granin N.G., and Zhdanov A.A.: Deep ventilation of Lake Baikal waters due to spring thermal bars. *Limnol. Oceanogr.*, 38: 1068-1072, 1993.
- Schmid, M., Budnev N.M., Granin N.G., Sturm M., Schurter M., and Wüest A.: Lake Baikal deepwater renewal mystery solved. *Geophysical Research Letters*, 35, doi 10.1029/2008GL033223, 2008.
- 540 Schmitt, R.W.: Double diffusion in oceanography. *Annual Review of Fluid Mechanics*, 26(1): 255-285, 1994.
- Tuzhilkin ,V.: Some features of the Black Sea seasonal thermohaline variability: Modern view. *Geography, Environment, Sustainability*. 3(2):42-50. DOI:10.24057/2071-9388-2010-3-2-42-50, 2010.
- Ullman, D., Brown, J., Cornillon, P., and Mavor, T.: Surface temperature fronts in the Great Lakes. *Journal of Great Lakes Research*, 24: 753-775, 1998.
- 545 Wozniak, B., and Dera, J.: *Light Absorption in Sea Water*. Springer, 454 pp., 2007.
- Zavialov, P.O., Pelevin, V.V., Belyaev, N.A., Izhitskiy, A.S., Konovalov, B.V., Kremenskiy, V.V., Goncharenko, I.V., Osadchiv, A.A., Soloviev, D.M., Garcia, C.A.E., Pereira, E.S., Sartorato, L., Moller Jr., , O.O.: High resolution LiDAR measurements reveal fine internal structure and variability of sediment-carrying coastal plume, *Estuarine, Coastal and Shelf Science*, doi:10.1016/j.ecss.2018.01.008, 2018.
- 550 Zilitinkevich, S. S., Kreiman, K. D., and Terzhevik, A. Y.: The thermal bar. *Journal of Fluid Mechanics*, 236: 27-42, 1992.

555

560



565

570




Figure 1: Map of Lake Issyk-Kul with schematized bathymetry contours 200, 300, 400, and 500 m, and hydrographic stations occupied during the field campaigns of June, 2015, November, 2016, and June, 2017. Most stations were occupied during all surveys (orange bullets). Green bullets show stations occupied only once in 2016, and the magenta bullet depicts the station occupied only once in 2017. The red bullets mark the locations of the mooring stations. The dashed lines in the eastern part of the lake indicate the paleo-beds of the Jyrgalan and the Tyup rivers, see the text for details.

580

585




Figure 2: (a) - Vertical profiles of temperature (red) and salinity (blue) observed on October 31, 2016 at 42.524°N, 77.190°E (solid), and on June 29, 2017 at 42.397°N, 77.100°E (dashed), and a summary of available historical data for different years based on (Kadyrov, 1986; Romanovsky, 1991; Peeters et al., 2003; Romanovsky et al., 2013); (b) – TS diagrams for water below the thermocline for October, 2016 (black) and June, 2017 (magenta); (c) – Vertical profile of turbidity in June, 2017.




Figure 3: (a) - Salinity (a) and temperature (b) distributions at the lake surface in November, 2016, based on pump-through CTD measurements.




Figure 4: Distribution of temperature ($^{\circ}\text{C}$, upper panel) and salinity (g kg^{-1} , lower panel) at longitudinal section following the ancient bed of the Jyrgalan River on the eastern shelf (green bullets in Fig. 1). November 1, 2016




Figure 5. Salinity at lake surface in June of 2017, based on data from hydrographic stations.




Figure 6: Difference of salinities at surface observed in June of 2017 and June of 2015.




Figure 7: Difference of lake surface temperatures observed in June of 2017 and June of 2015.




Figure 8: Air temperature at 2 m height in June, 2015, and June, 2017, from ERA-Interim reanalysis. The data correspond to the point 42°20'N, 77°30'E in the central part of the lake.




Figure 9: Vertical distributions of oxygen saturation (a); alkalinity (b), phosphates (c), and silicates (d) in different surveys. Black boxes indicate the typical range for waters below 200 m in 1961-1985 (Kadyrov 1986), except for oxygen where the box indicates the overall range. The vertical lines in the boxes show average values (where available).





Figure 10: Concentration of silicates (μM) at the lake surface in November of 2016.

670



675

Figure 11: Vertical distributions of concentration of nitrates (a) and nitrites (b) during the surveys of 2015, 2016, and 2017. Black box indicates the typical range for waters below 200 m in 1961-1985 (Kadyrov, 1986). The vertical lines in the boxes show average values.

680

685




Figure 12: Chlorophyll-“a” concentration and pigment index for June 2015 (blue bullets), November 2016 (green bullets), and June 2017 (red bullets). The straight lines represent linear regressions. The concentrations of chlorophyll (mg m^{-3}) and total suspended matter (g m^{-3}) are also exhibited in the inset table as the minimum and the maximum values and averages by 4 parts of the lake, delimited here by the 42.45°N parallel and the 77.4°E meridian.





Figure 13: Vertical distribution of chlorophyll-“a” concentration (mg m^{-3}) at longitudinal section following the ancient bed of the Jyrgalan River on the eastern shelf. November 1, 2016.



705

710



715 **Figure 14:** Bathymetric profile at meridional section across the Jyrgalan and the Tyup canyons at 77.9°E obtained by echo-soundings of June 25, 2015.

720





Figure 15: (a) - Air temperature at 2 m from ERA-Interim reanalysis data corresponding to the point 42°20'N, 77°30'E in the central part of the lake; (b) - Water temperature record from thermistor chain in Tyup canyon; (c) - Water temperature record from thermistor chain on the shelf between the Tyup and the Jyrgalan canyons; (d) - Water temperature record from thermistor chain in Jyrgalan canyon.



735 **Figure 16: Water temperature variability in the Jyrgalan canyon in November, 2016, as obtained from thermistor chain data.**

740

745

Table 1: Basic statistics of observed cooling events.

Site	Total duration of cooling, min	Number of cooling events		Duration of cooling event, min (maximum/mean)		ΔT over event, °C (maximum/mean)	
		Convective	Advection	Convective	Advection	Convective	Advection
1. Jergalan canyon, depth 51 m.	60780	1442, 10.7%	11960, 89.3%	15/3.1	41/3.5	0.59/0.01	0.56/0.01
2. Tyup canyon, depth 50 m.	104981	8743, 35.2%	16086, 64.8%	28/3.1	32/3.3	0.79/0.02	1.36/0.03
3. Shelf between the canyons, depth 23 m.	95783	13188, 54.2%	11164, 45.8%	24/3.0	19/3.1	0.25/0.01	0.29/0.01