

1 **Response to reviewers: “Watershed classification for the Canadian Prairies”**

2  
3 *Please note that we have changed the manuscript title to: “A WATERSHED CLASSIFICATION*  
4 *APPROACH THAT LOOKS BEYOND HYDROLOGY: APPLICATION TO A SEMI-ARID,*  
5 *AGRICULTURAL REGION IN CANADA”.*

6  
7 Page and line numbers are shown for each change in reference to the marked-up version.

8  
9 **Response to Editor**

10  
11 On P. 10, line 263 of the marked up version, it says "as the name suggests". The name is no longer  
12 given, just the symbol, so perhaps this statement should be removed.

13 *We have removed this statement as per the Editor's suggestion (10, 263).*

14  
15 On P. 9, line 246, wetland density seems to be defined as the number of wetlands within the watershed,  
16 but should be defined as number of wetlands per unit area (i.e. P. 21, line 611).

17 *We thank the Editor for this comment. We have adjusted the sentence to include reference to unit*  
18 *area: “the number of wetlands within the watershed per unit area (i.e., wetland density (km<sup>-2</sup>)).”*  
19 *(9, 246).*

20  
21 Typo on P. 26, line 751 in the spelling of Baulch.

22 *Thank you and we have made the correction.*

23  
24 In the acknowledgements, you might mention the Global Water Futures Program, for which the CFREF  
25 grant was awarded, and since both CFREF and Prairie Water are noted here.

26 *We have included the acknowledgement of the Global Water Futures program in this section.*

27  
28 Note that the map in Fig. 1 includes a north arrow pointing up, but the direction of north varies in the map  
29 projections for both the main and inset maps. (The same may apply to Fig. 5.)

30 *The projection for both the main and inset maps is the same. However, to improve clarity, we*  
31 *have added a north arrow and scale bar for the inset map. Fig. 5 does not have an inset map, so*  
32 *the map remains unchanged.*

33  
34

1 **A WATERSHED CLASSIFICATION APPROACH THAT LOOKS BEYOND**  
2 **HYDROLOGY: APPLICATION TO A SEMI-ARID, AGRICULTURAL REGION IN**  
3 **CANADA**

4 Jared D. Wolfe<sup>1\*</sup>, Kevin R. Shook<sup>2</sup>, Chris Spence<sup>3</sup>, Colin J. Whitfield<sup>1,4</sup>

5  
6 <sup>1</sup>Global Institute for Water Security, University of Saskatchewan, Saskatoon, Saskatchewan,  
7 Canada

8 <sup>2</sup>Centre for Hydrology, Saskatoon, Saskatchewan, Canada

9 <sup>3</sup>National Hydrology Research Centre, Environment and Climate Change Canada, Saskatoon,  
10 Saskatchewan, Canada

11 <sup>4</sup>School of Environment and Sustainability, University of Saskatchewan, Saskatoon,  
12 Saskatchewan, Canada

13

14

15 \*corresponding author: jared.wolfe@usask.ca

Formatted: Section start: Continuous

16 **ABSTRACT**

17       Classification and clustering approaches provide a means to group watersheds according  
18 to similar attributes, functions, or behaviours, and can aid in managing natural resources.  
19 Although they are widely used, approaches based on hydrological response parameters restrict  
20 analyses to regions where well-developed hydrological records exist, and overlook factors  
21 contributing to other management concerns, including biogeochemistry and ecology. In the  
22 Canadian Prairie, hydrometric gauging is sparse and often seasonal. Moreover, large areas are  
23 endorheic and the landscape is highly modified by human activity, complicating classification  
24 based solely on hydrological parameters. We compiled climate, geological, topographical, and  
25 land cover data from the Prairie and conducted a classification of watersheds using a hierarchical  
26 clustering of principal components. Seven classes were identified based on the clustering of  
27 watersheds, including those distinguishing southern Manitoba, the pothole region, river valleys,  
28 and grasslands. Important defining variables were climate, elevation, surficial geology, wetland  
29 distribution, and land cover. In particular, three classes occur almost exclusively within regions  
30 that tend not to contribute to major river systems, and collectively encompass the majority of the  
31 study area. The gross difference in key characteristics across the classes suggests that future  
32 water management and climate change may carry with them heterogeneous sets of implications  
33 for water security across the Prairie. This emphasizes the importance of developing management  
34 strategies that target sub-regions expected to behave coherently as current human-induced  
35 changes to the landscape will affect how watersheds react to change. The study provides the first  
36 classification of watersheds within the Prairie based on climatic and biophysical attributes, with  
37 the framework used being applicable to other regions where hydrometric data are sparse. Our  
38 findings provide a foundation for addressing questions related to hydrological, biogeochemical,  
39 and ecological behaviours at a regional level, enhancing the capacity to address issues of water  
40 security.

41 **A WATERSHED CLASSIFICATION APPROACH THAT LOOKS BEYOND**  
42 **HYDROLOGY: APPLICATION TO A SEMI-ARID, AGRICULTURAL REGION IN**  
43 **CANADA**

44

45 **1. INTRODUCTION**

46

47 Watershed classification methods provide a means of grouping watersheds according to  
48 similar attributes, or behaviours, and can identify sub-regions that are expected to exhibit  
49 coherent responses. This strategy can identify how catchment characteristics are similar, or  
50 dissimilar, among groups of watersheds and thus might influence hydrologic behaviour  
51 (McDonnell and Woods, 2004). Classifying watersheds can be useful for developing predictions  
52 in ungauged basins (Peters et al., 2012), and moreover, classification can be used to inform how  
53 changes to key traits (e.g., climate and land management) may affect system function.  
54 Establishing these links between watershed function and biophysical structure, including  
55 hydroclimate, is an opportunity of watershed classification (Wagener et al., 2007). Accordingly,  
56 the regionalization of hydrological response through watershed classifications has been used to  
57 inform natural resource management (Detenbeck et al., 2000; Jones et al., 2014).

58 Many different approaches to watershed classification have been employed to date,  
59 including non-linear dimension reduction techniques (Kanishka and Eldho, 2017), decision trees  
60 (Bulley et al. 2008), and independent component analysis (Mwale et al., 2011), among others.  
61 Hydrological characteristics (e.g., statistical properties of streamflow regime) are widely used to  
62 inform classification owing to their potential linkages between watershed features and  
63 hydrologic responses (Brown et al., 2014; Sivakumar et al., 2013; Spence and Saso, 2005). Other  
64 classification exercises have included a wider number of characteristics, including biophysical  
65 attributes along with streamflow response, to differentiate watershed classes (e.g., Sawicz et al.,  
66 2014; Burn, 1990). Ecoregions, which incorporate historical aspects of climate, topography, and  
67 vegetation regimes, have also served as a method of differentiation for eco-hydrological studies  
68 (Masaki and Rosenberry, 2002; Loveland and Merchant, 2004). In select cases, classification is  
69 performed independently of streamflow response factors (Knoben et al., 2018). In arid or poorly  
70 gauged regions of the world, these types of approaches to classification that are independent  
71 from or not strongly dependent on hydrological indices (streamflow response), are needed,

72 although few such classifications have been performed. The need for new approaches to  
73 watershed classification can also be true of regions undergoing strong pressures from climate  
74 change and land-use, where historical streamflow records may not reflect current behaviour,  
75 particularly if a regime shift has occurred.

76 In Canada, watershed classification has been applied in many regions (e.g., Cavadias et  
77 al., 2001; Ouarda et al., 2002; Spence and Saso, 2005). To date, most have focused on larger  
78 basins, and none have covered in detail the semi-arid Canadian Prairie, which spans nearly 5 x  
79 10<sup>5</sup> km<sup>2</sup> in western Canada, from the Rocky Mountain foothills in the west to Lake Winnipeg in  
80 the east (Fig. 1). This is despite its importance as a major food producing region of the world and  
81 one that faces numerous water security challenges (Gober and Wheeler, 2014; Spence et al.,  
82 2018). Earlier work by Durrant and Blackwell (1959) grouped large Prairie watersheds based on  
83 flood regimes. A recent classification that included the Prairie region focused on stream  
84 hydrology (e.g., MacCulloch and Whitfield, 2012) but was broader and included watersheds  
85 from mountainous and forested regions to the west and north, respectively. In the Canadian  
86 Prairie, and similar regions elsewhere, extrapolating catchment-scale field and modelling studies  
87 presents challenges. It is inherently difficult to explain or predict different responses among  
88 basins, as poorly developed stream networks with intermittent or seasonal flow do not easily lend  
89 themselves to classification methods featuring streamflow response. MacCulloch and Whitfield  
90 (2012), who found a single streamflow class across the Canadian Prairie, raised the question as  
91 to whether a single grouping is appropriate, and suggested the need to expand classifications to  
92 include a greater diversity of biological, physical and chemical properties.

93 Like many of the world's agricultural regions, the Canadian Prairie has undergone vast  
94 environmental change co-incident with the green revolution. Predominant agricultural practices  
95 have changed over the decades, and each is known to influence water cycling and storage,  
96 including tillage practices, summer fallowing, and cropping type (Awada et al., 2014; Van der  
97 Kamp et al., 2003; Shook et al., 2015). Significant warming over the last 70 years, especially in  
98 winter (Coles et al., 2017; DeBeer et al., 2016) has resulted in more rain at the expense of snow  
99 (Vincent et al., 2015), and multiple-day rainfall events have been increasing in frequency relative  
100 to shorter events in some regions (Dumanski et al., 2015; Shook and Pomeroy, 2012). These  
101 observed changes in precipitation have reduced the predictability of runoff derived from  
102 snowmelt, and add uncertainty to water management and agricultural decision-making.

Field Code Changed

Field Code Changed

103           Disentangling the relative impacts of climate and land-use changes on water quantity and  
104 quality is complex, particularly as their effects are heterogeneous across spatial extent and scale.  
105 For the Prairie and elsewhere, new approaches to classification that can distinguish sub-regional  
106 and, importantly, sub-hydrometric station variability, are needed. Further, because land  
107 management decisions in agricultural regions are intrinsically linked to system function, there is  
108 a need for classifications that can inform decision-makers at a relevant scale. Indeed, stable  
109 isotope-based investigations of runoff from small lake catchments in the Boreal Plains (north of  
110 the Prairie) emphasize the need for local-scale characterization of watershed behaviour (Gibson  
111 et al., 2010, 2016), while streamflow dynamics for the Prairie and nearby Boreal Plain are linked  
112 to local surface geology and land cover (Devito et al., 2005; Mwale et al., 2011), suggesting an  
113 opportunity for a new approach to watershed classification in the region. Another potential  
114 advantage of a more comprehensive approach is that by de-emphasizing available hydrometric  
115 observations for larger and well-studied or monitored basins and including other environmental  
116 characteristics, the risk of overlooking other functions (e.g., ecology, biogeochemistry) that may  
117 be equally important to the management of a watershed's natural resources can be reduced. A  
118 system-based watershed classification for the Prairie that avoids the prejudice of classifying only  
119 those watersheds where a reasonably robust understanding of hydrology or streamflow exists can  
120 serve as a template for other regions of the world where streamflow-based classification is not  
121 viable.

122           The objective of the present work is to develop a watershed classification system based  
123 on hydrologically and ecologically significant traits for the Canadian Prairie. In this region,  
124 assessment of localized hydrological response to change is challenged by limited spatial  
125 resolution of observed streamflow data, and higher order streamflow being unrepresentative of  
126 local response due to a poorly-developed drainage network. In establishing such an approach, we  
127 seek to advance our understanding of watershed hydrology and broader watershed behaviour  
128 within the Prairie whilst also providing a framework for similar classification exercises in other  
129 regions where streamflow-based methods are not ideal. Our approach avoids the limitations of  
130 classifying according to known hydrologic response, and increases the spatial resolution of  
131 watershed classification relative to many existing approaches. We compile physio-~~ge~~ographic  
132 characteristics, including geology, wetland distribution, and land cover, of watersheds  
133 approximately 100 km<sup>2</sup> to achieve the classification. This framework will identify those areas

134 that are climatically and ~~physio~~-geographically similar, and thus might be expected to respond in  
135 a hydrologically coherent manner to climate and land management changes. Additionally, it  
136 provides a foundation on which to base prediction of watershed hydrologic, biogeochemical and  
137 ecological responses to these stressors.

## 138 139 **2. DATA COLLECTION & COMPILATION**

### 140 141 *2.1. Region domain and description*

142 The Canadian Prairie (Prairies ecozone) spans the provinces of Alberta, Saskatchewan,  
143 and Manitoba, and is part of the Nelson Drainage Basin (Fig. 1). Climate is semi-arid, with mean  
144 annual precipitation ranging between 350 and 610 mm (1970–2000) increasing from west to east.  
145 Mean annual temperature was 1–6°C over the same period with warmer conditions towards the  
146 southwest (Mekis and Vincent, 2011; Vincent et al., 2012;  
147 [http://climate.weather.gc.ca/climate\\_normals/index\\_e.html](http://climate.weather.gc.ca/climate_normals/index_e.html)). Much of the region deglaciated  
148 during the Late Pleistocene approximately 10,000 years before present, resulting in an often  
149 hummocky landscape with numerous depressions. Combined with the dry climate, the relatively  
150 short post-glaciation history has prevented maturing of a ubiquitous drainage network, and many  
151 headwaters remain disconnected from higher order streams (Shook et al., 2015). Depressions in  
152 the hummocky landscape, and the wetlands that form within them, are important features for  
153 Prairie hydrology (Van der Kamp et al., 2016) and often facilitate groundwater recharge (i.e.,  
154 depression-focused recharge) (Van der Kamp and Hayashi, 2009). The location of wetlands and  
155 their size, relative to the watershed outlet controls hydrologic gate-keeping (e.g., Spence and  
156 Woo, 2003), and thus the potential to contribute streamflow to higher-order watersheds  
157 (Leibowitz et al., 2016; Shaw et al., 2012; Shook et al., 2013). The size distribution of wetlands  
158 within a watershed and their spatial arrangement also dictate biogeochemical function and  
159 provide habitat and foraging for biota (Evenson et al., 2018). Terrestrial vegetation is typically  
160 open grassland, with aspen parkland ecotone along the northern edges of the ecozone boundary  
161 (Ecological Stratification Working Group, 1995).

### 162 163 *2.2. Watershed boundaries*

164 The focus of this study was on those watersheds that drain a distinctively prairie  
165 landscape, with watersheds defined according to topographic delineation. Thus, we constrained  
166 our study to the Canadian Prairie ecozone ( $4.7 \times 10^5 \text{ km}^2$ ) ~~and watersheds occurring therein;~~  
167 ~~watershed areas of larger exotic streams in the region originating in the Rocky Mountains to the~~  
168 ~~west were not included.~~ Delineations of candidate study watersheds were obtained from the  
169 HydroSHEDS global dataset (Lehner and Grill, 2013). Watershed boundaries within this dataset  
170 were based on Shuttle Radar Topographic Mission (SRTM) digital elevation model (DEM)  
171 calculated at a 15 arc-second resolution. The resolution is equivalent to for example  
172 approximately 285 m east-west and 464 m north-south at Saskatoon, SK. As with other SRTM  
173 products, the HydroSHEDs dataset may be prone to errors in regions with low relief due to  
174 elevation precision of 1 m. However, the dataset provided ~~an objective watershed~~ delineations  
175 over the geographic region of interest and at a fine enough scale (i.e.,  $100 \text{ km}^2$ ), and thus, it was  
176 sufficient based on data availability for purpose of the current study.

177 Only those watersheds completely within the Canadian Prairie ecozone (~~Fig. 1~~) were  
178 extracted ( $n = 4729$ ) from the HydroSHEDs dataset. Those watersheds that were very large  
179 ( $>4000 \text{ km}^2$ ) or small ( $<5 \text{ km}^2$ ) were removed from analysis (see Table S1). Because  
180 HydroSHEDs includes the basins of larger water bodies, including lakes, watersheds consisting  
181 of a majority of water were removed as the study only concerns the uplands of these systems.  
182 Finally, highly urbanized areas (i.e., watersheds with cover being  $>40\%$  urban) were removed.  
183 After considering these criteria, 4175 watersheds remained for use in subsequent analyses,  
184 covering a total area of  $4.2 \times 10^5 \text{ km}^2$ . Mean watershed area for this subset was  $99.8 \pm 58.7 \text{ km}^2$ .  
185

### 186 2.3. ~~Physio-geographic~~ data collection

187 The physio-~~geographic~~ watershed variables were assembled from Canadian provincial  
188 and federal governments and non-governmental agency datasets (see Table S2 for a full list of  
189 variables and their sources). Variables were derived from climatic, hydrologic, geological,  
190 geographic, and land cover data, and details are described briefly below. Spatial processing and  
191 statistical analyses were conducted in ArcGIS version 10.5 and R version 3.4.3 (R Core Team,  
192 2018), respectively.

#### 193 194 2.3.1. Climate

195 Mean annual precipitation and temperature data were derived from the Canadian Gridded  
196 Temperature and Precipitation Anomalies (CANGRD) dataset spanning (ECCC, 2017).  
197 CANGRD is the only gridded climate product available for the region that uses adjusted and  
198 homogenized station data, and was picked for this reason (Mekis and Vincent, 2011; Vincent et  
199 al., 2012). The 1970–2000 period was chosen because the number of stations with adjusted and  
200 homogenized data used to derive CANGRD significantly diminished after 2000 (Laudon et al.,  
201 2017). Mean annual values over the 30-year period were constructed from 50 km resolution  
202 gridded cells ( $n = 626$ ) within and surrounding the Prairie ecozone, and interpolated to a higher  
203 spatial resolution raster by kriging using a spherical semivariogram. Values were clipped  
204 according to the watershed boundaries, and averaged over the watersheds to obtain mean annual  
205 precipitation and temperature for each watershed. Mean annual potential evapotranspiration  
206 (PET) was derived as a measure of dryness across the region. To maintain consistency among  
207 climate data, and use the same temperature data as described above, options were limited with  
208 which to calculate PET. The Thornthwaite equation (Thornthwaite, 1948) was applied using the  
209 R package *SPEI* (Vicente-Serrano et al., 2010). A disadvantage of the Thornthwaite approach is  
210 that it ~~assumes~~ calculates PET solely as a function of air temperature and latitudinal position,  
211 and it assumes a fixed  $\alpha$ - $\epsilon$  correlation between temperature and radiative forcing. As such, it  
212 integrates effects of other factors directly or indirectly influencing radiation or latent heat, like  
213 advection, vegetation, and humidity. ~~and~~The calculation adjusts for any lag in this relationship  
214 using corrections for latitude and month; however, it likely does not represent the full annual and  
215 seasonal variability in PET across a landscape, given regional heterogeneity of the  
216 aforementioned factors. Despite the limitations, the simplicity of this method is ideal for  
217 application across the wide geographic area of interest with limited data required as input,  
218 allowing for approximation of mean annual PET for the study area.

219

### 220 2.3.2. Wetland traits

221 Large regions within the Canadian Prairie have been designated as being “non-effective”,  
222 where they do not contribute flow to the stream network, at least one year in two (Godwin and  
223 Martin, 1975). The location of these regions are shown in Figure 1. This definition stems from  
224 work by Agriculture and Agri-Food Canada where prairie drainage areas were divided into *gross*  
225 and *effective* drainage areas, whereby the former describes the area within a topographic divide

226 that is expected to contribute under highly wet conditions, and the latter is the area that  
227 contributes runoff during a mean annual runoff event (Mowchenko and Meid, 1983). Thus, at its  
228 simplest, the non-effective area is the difference between the gross and effective drainage area;  
229 however, the exact area contributing runoff is dynamic and the controls complex, which include  
230 antecedent storage capacity and climatic conditions (Shaw et al., 2012; Shook and Pomeroy,  
231 2015). Briefly, the “non-effective” regions are caused by the intermittent connectivity of runoff  
232 among the landscape depressions, which trap runoff, and prevent it from contributing to  
233 downstream flow when the depressions are not connected. Trapped surface water can form  
234 wetlands (hereafter, inclusively referring to water area ponded in these depressions). These  
235 depressions can store water, and are indicative of water storage of the basin. Thus the non-  
236 effective portion of a basin is an index of its lack of contribution and is an important quality  
237 when considering the hydrological dynamics of this region (Shook et al., 2012).

238 The Global Surface Water dataset (Pekel et al., 2016) provides a geographically  
239 comprehensive layer of any ~30 m x 30 m pixel that was inundated at least once between 1984  
240 and 2015, as identified from the Landsat constellation of satellites. It was assumed that the  
241 dataset was indicative of potential maximum wetland coverage, as this period spanned several  
242 wet [climate periods](#). As such, “wetland” in this context can include some seasonal ponds (i.e.,  
243 prairie potholes) as well as larger or more permanent shallow water bodies (but see Section 2.2  
244 and Table S1). Using the R package *raster* (Hijmans, 2017), wetland variables were calculated  
245 for each study watershed, including fractional wetland area, and the number of wetlands within  
246 the watershed per unit area (i.e., wetland density ( $\text{km}^{-2}$ )). The ratio of the area of the largest  
247 wetland to total wetland area in the watershed was also used as a metric (i.e.,  $W_L$ ). Further, we  
248 used the ratio of the linear distance of the largest wetland’s centroid to the watershed outlet ( $L_W$ ),  
249 to the maximum watershed boundary distance to the outlet ( $L_O$ ) to represent a centroid fraction  
250 ( $L_W/L_O$ ; i.e., the relative location of the largest wetland to watershed outlet). The basin outlet was  
251 defined as the point of lowest elevation on the watershed boundary. Both  $W_L$  and  $L_W/L_O$  can be  
252 used to evaluate the relative importance of hydrological gate-keeping; for example, larger  
253 wetland depressions located closer to the outlet control the likelihood of the watershed  
254 contributing flow downstream and attenuating peakflow (Shook and Pomeroy, 2011; Ameli and  
255 Creed, 2019).

256 To estimate wetland size distribution, it was assumed that they followed a Generalized  
257 Pareto Distribution (GPD) defined according to (Shook et al., 2013):

258

$$F(z) = GPD(\mu, \beta, \xi) = 1 - \left[ 1 + \xi \left( \frac{z - \mu}{\beta} \right)^{-1/\xi} \right] \quad (1)$$

259

260 Where  $z$  is wetland area, and  $\mu$  is the location parameter (i.e., the minimum size for which the  
261 distribution was fitted and has units of  $m^2$ ), and the scale ( $\beta$ ) and shape ( $\xi$ ) parameters are  
262 determined for each watershed. The  $\beta_{scale}$  parameter is an index of the dispersion of the  
263 distribution, similar to the standard deviation, with the same units as the data being fitted (in this  
264 case  $m^2$ ). The  $\xi_{shape}$  parameter is dimensionless and governs the shape of the fitted distribution.  
265 Hosking and Wallace (1987) plot the effect of variation in the shape parameter on the GPD. The  
266 scale and shape parameters were used to quantify the size distribution of wetlands and thus to  
267 describe the wetland frequency distributions for the cluster analyses (see 3.2). Note that because  
268 the sizes of the water bodies were taken from infrequent remote-sensing measurements (i.e., the  
269 Landsat data have a minimum revisit time of 8 or 16 days), they also are biased against short-  
270 lived water bodies.

271

### 272 2.2.3. Topographical parameters

273 Topographic variables including the mean elevation, mean and coefficient of variation of  
274 slope, and stream density were also calculated for each watershed. Because of the hummocky  
275 nature of many regions in the domain, it is possible for a basin to have some fraction of its area  
276 located at an elevation below that of the outlet. As such, the fraction of area below the basin  
277 outlet ( $A_{BO}$ ) was calculated for each basin. The elevation and slope variables were based on a  
278 DEM generated from the SRTM dataset. Stream vectors were obtained from the Hydrographic  
279 features CanVec (1:50000) series available from Natural Resources Canada  
280 (<https://open.canada.ca/data/en/dataset?q=canvec&sort=&collection=fgp>). The total length of  
281 streams within a watershed was calculated, and divided by the watershed area to produce the  
282 stream density. Additionally, the dimension shape factor (DSF) was used to describe watershed  
283 shape, as it has been found important for hydrological responses in previous Canadian catchment  
284 classification exercises (Spence and Saso, 2005). The DSF ( $km^{-1}$ ) was calculated as follows:

285

$$DSF = \frac{(0.28 \cdot P)}{A} \quad (2)$$

286

287 Where  $P$  (km) and  $A$  (km<sup>2</sup>) are the watershed perimeter and area, respectively, and derived from  
288 the HydroSHEDS global dataset (Lehner and Grill, 2013).

289

Geographical parameters of surficial geology, local surface landforms, soil particle size  
290 classes (sand, silt, clay), and soil zone were included in the analysis. Surficial geology polygons  
291 were derived by compiling provincial government data sources for Alberta (Atkinson, 2017),  
292 Saskatchewan (Simpson, 2008), and Manitoba (Matile, 2006). Due to the different geological  
293 classification schemes for each province, more detailed classes were grouped to broader  
294 categories related to depositional environment and surficial materials using those from the  
295 Geological Survey of Canada (2014), which provided for comparison across provincial  
296 boundaries. Local surface form (i.e., areas categorized by slope, relief, and morphology) and soil  
297 zone data were obtained from the Soil Landscape dataset (AAFC, 2013). The soil zones in the  
298 Canadian Prairie, used in the analyses were black, dark brown, brown, gray, and dark gray. The  
299 zones incorporate characteristics of colour and organic content, which are influenced by regional  
300 climate and vegetation. Clay, silt, and sand content were collected from the Detailed Soil Survey  
301 of Canada (AAFC, 2015). Mean catchment values of each of surficial geology, local surface  
302 landform, soil zone, and particle size class were determined by areal weighting of soil polygons  
303 within the watershed boundaries.

304

#### 305 2.3.4. Land cover and cropland practice

306

Fractional areas of land-use types were derived from the Agriculture and Agri-Food  
307 Canada's 2016 Annual Crop Inventory (AAFC, 2016). These raster data define land-use and land  
308 cover. Variables used in our analysis were standardized to watershed area and included  
309 unmanaged grasslands, forests (i.e., the sum of coniferous, deciduous, and mixed forest areas),  
310 pasture, and cropland (sum of cropped land areas). Predominant cropland practice was defined  
311 according to the fractional area of tillage by agricultural region sub-division (e.g., normalized to  
312 the area prepared for seed within that division by year). Averaged areas over the years 2011 and  
313 2016 for each practice, including zero-till, conservation till (leaving crop residue on soil surface),

314 and conventional till (incorporating residues into soil) (Statistics Canada, 2016), were used to  
315 describe these activities, and normalized as a fraction of the watershed.

316

### 317 2.3.5. Hydrological variable calculation

318 The relatively sparse hydrometric stream gauging in the domain, and the resulting paucity  
319 of data, presents two notable challenges to hydrologic response-based watershed classification.

320 The first is that the basin network is biased to stations on higher-order (and often exotic) streams  
321 traversing the region (i.e., larger river basins), and thus there are a limited number of  
322 hydrometric gauges on streams draining solely Prairie watersheds, particularly at the spatial  
323 resolution of our study watersheds (~100 km<sup>2</sup>). Further, only a subset of these are considered  
324 reference stations (i.e., gauging unmanaged flows). Second, in the more arid and/or cold regions  
325 of the Prairie, some of these hydrometric stations are operated only seasonally, presenting  
326 additional challenges in using these records for classification exercises (e.g., MacCulloch and  
327 Whitfield, 2012).

328 As a result, mean annual runoff (Q2) and 1:100 year flood (Q100) magnitudes were  
329 estimated for the 4175 watersheds using relationships defined from canonical correlation  
330 analysis (CCA) to correlate gauged data to multivariate climatic and physio-~~geo~~graphic data  
331 according to procedures given by Spence and Saso (2005). According to Spence and Saso  
332 (2005), expected uncertainty using these methods approached 50% but exhibited biases of less  
333 than 15% (n = 34). Hydrological stations used were those identified in MacCulloch and  
334 Whitfield (2012) and within the Prairie region (n = 11), and data were obtained from archived  
335 databases of the Water Survey of Canada ([https://wateroffice.ec.gc.ca/search/historical\\_e.html](https://wateroffice.ec.gc.ca/search/historical_e.html))  
336 between 1990–2014. We note that greater uncertainty than that reported by Spence and Saso  
337 (2005) may result when using the CCA approach with a smaller sample size. Multivariate  
338 geophysio-~~geo~~graphic data were collected as outlined in the above sections according to the  
339 watershed boundaries for the hydrological stations. Due to the fact that many watersheds within  
340 the HydroSHEDS dataset are likely to drain internally and do not consistently connect to a  
341 higher-order stream network, these streamflow data were interpreted as “runoff”, meaning the  
342 amount of water accumulated within the watershed polygon that drains to its lowest point  
343 annually.

344 Briefly, CCA correlates the streamflow record of gauged basins to physico-climatic  
345 characteristics of watersheds by representing ~~the original~~these variables as a reduced set of  
346 canonical variables. The analysis results in two canonical variable sets: one for the physico-  
347 climatic variables (i.e., V1 and V2) and another for the hydrological variables (i.e., W1 and W2).  
348 These canonical variables are constructed from linear combinations of the ~~original~~-variable sets  
349 such that the correlation of the canonical variables are maximized. Canonical variables plotting  
350 similarly on X-Y plots (W1-W2 and V1-V2), indicate good correlation (Spence and Saso, 2005).  
351 Where canonical correlations ( $\lambda_1, \lambda_2$ ) were above 0.75 (Cavadias et al., 2001), that set of  
352 physico-climatic variables was deemed useful for estimating hydrological variables. Those  
353 physico-climatic variables passing this threshold were included as variables in a multiple  
354 regression to develop a predictive equation for Q2. Analyses were performed using the R  
355 package *vegan* (Oksanen et al., 2018).

### 357 3. DATA ANALYSIS

#### 359 3.1. Pre-processing compositional datasets

360 Principal components analysis (PCA) was used as a pre-processing step to reduce the  
361 dimensionality associated with compositional datasets (e.g., topographical and land cover  
362 parameters) (Fig. S1). Using this approach, the principal components (PC) that could  
363 cumulatively explain 80% of the variation in a subset of compositional data were included in the  
364 subsequent cluster analysis. This procedure identified the major data patterns and aided in  
365 reducing the number of zero-weighted variables. Where necessary, variables that were not  
366 transformed into PCs were log-transformed to reduce data skewness. Variable unit ranges were  
367 also scaled during the PCA to reduce the impact of certain variables exhibiting a large range of  
368 values on the subsequent cluster analysis.

#### 370 3.2. Agglomerative hierarchical clustering of principal components and watershed classification

371 Clustering analysis was performed on the suite of physio-~~geo~~graphic variables, which  
372 included PC variables derived from pre-processing (Table S2; Table SS3). Agglomerative  
373 hierarchical clustering of principal components (HCPC) was used to define clusters of  
374 watersheds using the *HCPC* function in the R package *FactoMineR* (Lê et al., 2008; Husson et

375 al., 2009) to apply a PCA on the standardized multivariate dataset of watershed attributes and  
376 was the basis for clustering. The majority of physio-geographic variables were included as active  
377 variables in the PCA and thus influenced the arrangements of the PCs. In contrast, watershed  
378 area, DSF, latitude, and longitude were used only as supplementary variables, and thus did not  
379 explicitly affect the clustering analysis. These variables did, however, aid in watershed class  
380 characterization and interpretation. The first set of PCs that together explained 50% of the  
381 variation in the dataset ( $n=6$ ) was retained for agglomerative clustering. Retaining these first  
382 PCs at a threshold of 50% allowed for clearer focus on main trends in the data and reduced the  
383 impact of noise on subsequent analyses, which might occur if subsequent, less influential, PCs  
384 were retained.

385 The agglomerative hierarchical clustering was performed using the Euclidean distances  
386 (from the PCA) and Ward's criterion for aggregating clusters. Ward's criterion decomposes the  
387 total inertia of clusters into between and within-group variance, and this method dictates merging  
388 for clusters (or watersheds) such that the growth in within-group inertia is minimal (Husson et  
389 al., 2010). The total inertia is partitioned into within- and between-group inertias. Within-group  
390 inertia represented the homogeneity, or similarity, of watershed within a cluster. Consequently,  
391 watersheds located close to each other in PC-space were deemed to be similar in their attributes.  
392 ~~This approach decomposes the total variability, or inertia, into within and between group~~  
393 ~~inertias.~~ Watersheds are grouped according to pairs that minimize within-group inertia (Begou et  
394 al., 2015), and are differentiated based on between-group inertia gained by adding clusters. The  
395 variables contributing to cluster characteristics were determined by  $t$ -tests (Husson et al., 2009),  
396 which assessed whether the cluster mean for a given variable was significantly ( $\alpha=p < 0.05$ )  
397 greater or smaller than the overall mean.

398

### 399 3.3. Comparing class-specific observed and simulated wetland depression data

400 To compare how well the GPD parameters predicted the observed wetland area  
401 distributions from the Global Surface Water (GSW) dataset, wetland size distributions were  
402 simulated for each class. Wetland area for select watershed class-specific percentiles (i.e., 25<sup>th</sup>,  
403 50<sup>th</sup>, and 75<sup>th</sup> percentiles) derived from the simulated data were then compared to the wetland  
404 areas for corresponding watershed class-specific percentiles of the observed watershed data to

405 [assess the potential usefulness of using these parameters in representing wetland size](#)  
406 [distribution.](#)

407 For this comparison, the fitted wetland area distributions were constrained in their  
408 minimum and maximum values by the Global Surface Water dataset spatial resolution (i.e., the  
409 30 m pixel size) and the median area of the largest wetland observed for each watershed class,  
410 respectively. The median area of the distribution of largest wetlands for each watershed class  
411 gave an indication of the maximum sizes of the water bodies in those watersheds, and thus  
412 provided a maximum value for simulating wetland areas using the GPD. Wetland areas were  
413 simulated using the R package *SpatialExtremes* (Ribatet, 2018).

#### 415 3.4. Resampling and re-classifying procedure

416 The robustness of the HCPC procedure on characterizing Prairie watersheds was tested  
417 using additional hierarchical clustering on ten subsets of the entire [set of 4175](#). For each  
418 iteration, ten percent of watersheds were removed from the original dataset ( $n = 4175$ ) without  
419 replacement, and the remaining watersheds ( $n = 3757$ ) were then re-analyzed according to the  
420 HCPC outlined above (Fig. S1). The number of potential classes allowed was set at seven ( $k =$   
421 7), for consistency with the complete analysis. The resulting classifications were then compared  
422 to the classification performed on the complete dataset, with the watersheds being assessed on  
423 the percentage of iterations in which they were assigned to the same class as the complete  
424 classification. The proportion of membership agreement was calculated and visualized to assess  
425 the likelihood of classing watersheds consistently.

## 427 4. RESULTS

### 429 4.1. Geographical data processing

#### 430 4.1.1 Dimension reduction: [Compositional datasets and ~~Variable~~ principal components analysis](#)

431 Variation in geology and soil was best explained by two or three principal components  
432 (Table 1; Fig. S2). Two PCs captured over 80% of the variation in surficial geology, with PC1  
433 (proportion explained: 73%) positively relating to glacial till deposits and negatively with  
434 glaciolacustrine deposits, and PC2 (14%) positively related to riverine or erosive deposits, such  
435 as glaciofluvial, alluvial, and eolian deposits. Particle size class data were explained by the first

436 two PCs, where PC1 (75%) was positively associated with sand and negatively associated with  
437 silt and clay, while PC2 (14%) was related negatively to silt. Positive PC1 (55%) scores defined  
438 the dominance of black soils, and PC2 (43%) described dominance of brown or dark brown soils  
439 on positive or negative scores, respectively. Three PCs described the local surface form dataset.  
440 PC1 (55%) captured the change from greater portion of hummocky forms to undulating forms,  
441 and PC2 (24%) was negatively associated with higher river-incised landscape fraction. The  
442 portion of level surface form was negatively related to PC3 (12%).

443 Three PCs were needed to explain over 80% of the variation in land cover (Table 1; Fig.  
444 S2). Land cover PC1 (37%) was positively associated with higher cropland and negatively with  
445 unmanaged grassland; whereas PC2 (25%) was negatively associated with higher pasture and  
446 forest cover. PC3 was associated with greater fallow and pasture areal proportion (21%).  
447 Cropland practice was described by PC1 (90%), with zero-till practices being negatively  
448 associated to this component. Although it only explained 9%, PC2 was also retained to describe  
449 the change between conventional and conservation till practices, with the practices exhibiting a  
450 positive and negative relationship, respectively.

451

#### 452 4.1.2 Canonical correlation analysis

453 The canonical coefficients from the CCA were acceptably high at  $\lambda_1$  0.97 and  $\lambda_2$  0.77,  
454 respectively, indicating that the physico-climatic variables exhibited influence on the  
455 hydrological variables (Cavadias et al., 2001; Spence and Saso, 2005). ~~Mean-e~~ Canonical  
456 correlation values between the hydrological variables and W2 were greater than those with W1  
457 (Table 2); ~~and because both values of  $\lambda$  were acceptably large (Cavadias et al., 2001) thus,~~ the  
458 physico-climatic variables strongly associated to second canonical correlation (i.e., V2) were  
459 used in the multiple regressions. These variables were watershed area, DSF, areal fraction of  
460 rock, and areal fraction of natural area. Plots of observed and predicted runoff Q2 ( $R^2=0.45$ ) and  
461 Q100 ( $R^2=0.48$ ) show moderate agreement at lower flow values (Fig. 2). There is a negative  
462 bias estimated between 26 and 29%, which is greater than that documented by Spence and Saso  
463 (2005) using comparable extrapolation methods, but this is not unexpected because of the  
464 smaller sample size in the current study. As Q2 and Q100 exhibited high collinearity, only Q2  
465 was included in subsequent cluster analyses to:

466

$$\log(Q2) = 0.130*\log(A) - 0.077*\log(N) + 0.117*\log(R) - 0.141*\log(DSF) - 0.620 \quad (3)$$

467  
 468 Where  $A$  was the watershed area,  $N$  was the natural area fraction and the sum of grasslands and  
 469 forest,  $R$  was the rock fraction area, and  $DSF$  was the dimensional shape factor of the watershed.  
 470 The equation was then used to calculate  $Q2$  for each watershed included in the clustering  
 471 analysis.

472  
 473 *4.2. Watershed classification*

474 *4.2.1. Principal component analysis*

475 In total, 29 watershed attributes, including the PCs from compositional datasets ([see](#)  
 476 [Table 1](#)), were used in the clustering analysis as active variables, and four were included as  
 477 supplementary (Table 3). In the pre-clustering PCA, the first six PCs explained 54.3% of data  
 478 variation, and were retained for the HCPC analysis (Fig. 3). The influence of subsequent PCs  
 479 declined dramatically, and eleven PCs were required to explain >80%. [Variable importance in](#)  
 480 [the classification was not related to the log-transformed range exhibited by that variable \(data not](#)  
 481 [shown\), and impact was mitigated by scaling the ranges of input variables in the PCA.](#)

482 Principal components 1 and 2 captured changes in physical, land cover, and wetland  
 483 characteristics (Fig. 3). PC1 was strongly associated with physical and land cover characteristics,  
 484 such as elevation, wetland density, and the land cover PCs. PC2 was strongly related to metrics  
 485 characterising the hydrological landscape, including river and wetland density, non-effective area  
 486 fraction, landscape surface form, and size of the largest wetland ( $W_L$ ). Subsequent PCs explained  
 487 less variation and were more specialized in the variables associated with them. Generally, these  
 488 PCs were associated with differences in soil zone and texture class, surficial geology, and  
 489 varying surface land form. A more detailed account of associations of the variables with the PCs  
 490 is provided below.

491 PC1 was positively associated with elevation, mean slope, land cover PC2, and surface  
 492 form PC3, and negatively with, total annual precipitation, soil zone PC1, wetland density, land  
 493 practice PC1, land cover PC1, and longitude (Table 3; Fig. 3). PC2 was associated with non-  
 494 effective area fraction, wetland density,  $\beta$ , and surface form PC2, and negatively related to land  
 495 practice PC1,  $W_L$ , and river density. PC3 was positively related to wetland fraction,  $W_L$ ,  $\zeta$ , soil  
 496 texture PC2, and DSF. Watershed area and runoff were negatively associated with PC3.

497 Variable correlations were ~~less-strong~~weaker for the remaining three PCs (Table 3). PC4  
498 was mainly associated with soil texture PC1, surficial geology PC1 and surface landform PC1,  
499 characteristic of sandier soil areas featuring glacial till deposits and higher hummocky surface  
500 forms, as well as higher mean slope. PC4 was negatively related to land cover PC2. PC5 was  
501 related positively to PET, fraction below outlet, and soil zone PC2, and negatively to land cover  
502 PC1, river density, and slope CV. Finally, PC6 was mainly associated with soil texture PC2 and  
503 land cover PC3, and negatively with surface landform PC2.

504

#### 505 4.2.1. Agglomerative hierarchical cluster analysis

506 Seven clusters were identified from the hierarchical cluster analysis based on the  
507 between-group inertia gained by increasing cluster number ( $k$ ). The HCPC analysis suggested  
508 three clusters resulted in the greatest reduction of within-group inertia while minimally  
509 increasing  $k$  (Fig. 4). Further increasing  $k$  refined the separation and differentiation of clusters up  
510 to seven ( $k=7$ ). Minimal added separation was observed up to  $k=9$ , and increasing  $k > 9$   
511 resulted in little inertia gained between clusters. Thus, seven clusters, or classes, were manually  
512 selected based on these observations (Fig. 4).

513

#### 514 4.2.3. Class characteristics and interpretation

515 Our methodology yields sub-regional watershed classes according to climatic,  
516 physiographic, wetland, and land cover variables. The seven classes (Fig. 5), are defined by  
517 multivariate sets of attributes (Table 4). Influential classifying variables in all classes were mean  
518 elevation, total annual precipitation, land practice, surface forms, and wetland density. Other  
519 variables influential to class differentiation included fraction of non-effective area, land cover,  
520 and soil variables. Climate and elevation gradients are likely responsible for the west to east  
521 watershed clustering pattern. Moreover, we observe strong spatial concordance among some  
522 classes (Fig. 5), which is likely due to the hierarchical nature of the analysis. For simplicity, we  
523 interpret classes based on the variables where large, significant differences in class mean versus  
524 the overall mean of the dataset were observed. The classes can be assigned as follows: Southern  
525 Manitoba (C1); a Prairie Pothole region (C2, C3); Major River Valleys (C4); and Grasslands  
526 (C5, C6, and C7).

527

528 *Southern Manitoba (C1)*

529 The majority of Class 1 (C1; n = 365) watersheds occurred in the eastern prairie south of  
530 Lake Winnipeg (Fig. 5) and thus “Southern Manitoba” is used as the class name. Distinguishing  
531 characteristics associated with this class included soil zone PC1 (predominantly black soils) and  
532 cropland practice PC1 (predominantly conventional till) (Table 4). Southern Manitoba had a high  
533 incidence of glaciolacustrine and alluvial deposits, as indicated by moderately negative and  
534 positive relationships with surficial geology PC1 and PC2, respectively, and the class also had  
535 low mean elevation. Topography tended to be level, with mild slopes and strong association with  
536 land surface form PC3 (Table 4). Notably, these watersheds exhibited both high annual  
537 precipitation and PET compared to other classes, and this class was the only one to have no mean  
538 moisture deficit (i.e., precipitation – PET > 0) (Fig. 6). Southern Manitoba watersheds also  
539 exhibited smaller fractions of non-effective areas and grasslands than other classes (Fig. 7).

540

541 *Prairie Potholes (C2 and C3)*

542 The Prairie Pothole group, consisting of Class 2 (C2; n = 879), or Pothole Till, and Class  
543 3 (C3; n = 681), Pothole Glaciolacustrine, represents the largest class of watersheds spatially,  
544 spanning the northern part of the Alberta prairie to the southeastern part of Saskatchewan (Fig.  
545 5). Mean annual precipitation was relatively high for the study area, contributing to a slightly  
546 negative moisture deficit (Fig. 6). These watersheds contained large fractions of non-effective  
547 area (~75%) (Fig. 7a), and they exhibited positive scores on land cover PC1 (Table 4) indicating  
548 high cropland cover (~70%), whereas unmanaged grassland cover was typically very low  
549 (<20%) (Fig. 7b-c). On average, Pothole watersheds had high wetland densities (wetlands km<sup>-2</sup>),  
550 with C2 exhibiting the greatest density of all classes (Fig. 8a).

551 Surficial geology differentiated classes C2 and C3. Overall, glacial till and hummocky  
552 landforms dominated the pothole region; however, C2 was more associated with these  
553 characteristics, scoring greater mean values on PC1 of local surface form and surficial geology.  
554 In contrast, glaciolacustrine deposits were more common in C3, and soils had a higher incidence  
555 of clay and silt, where C2 watersheds were sandier (Table 4). Although both classes contain  
556 many wetlands, C2 watersheds had the smallest values of W<sub>L</sub>, indicating lower areal water extent  
557 was contained in the largest wetland (Fig. 8b).

558

559 *Major River Valleys (C4)*

560 Class 4 (C4; n = 536) watersheds were associated with river valleys, and as such, extend  
561 across the prairie region (Fig. 5) and generally coincide with major rivers (e.g., North and South  
562 Saskatchewan, Qu'Appelle) and large lakes. These watersheds had the greatest value of the  
563 fraction of water area in the largest depression ( $W_L$ ) (Fig. 8b), as well as high slope CV, wetland  
564 fraction, and fractions of black soil (i.e., higher soil zone PC1 scores) (Table 4). These  
565 watersheds were also associated with soil texture PC1 and surficial geology PC2, suggestive of  
566 higher incidence of sandy riverine deposits (e.g., alluvial and glaciofluvial deposits). The Major  
567 River Valleys class tended to have large "wetland" area, which is interpreted as the area of water  
568 of these rivers.

569 Taken together, these watersheds were related to parameters typical of fluvial  
570 environments, including glaciofluvial or alluvial deposits, and sandier soils. Large values of  
571 mean and CV of slope were also typical of river valley watersheds. About half the basin area  
572 tends to be non-effective in these watersheds, compared to the much greater fractions in the  
573 pothole regions (Fig. 7a) that surround many of the Major River Valleys watersheds. Being river  
574 valleys, C4 watersheds were generally narrow and small in area. Higher DSF (i.e., narrower  
575 watersheds) and smaller areas were generally associated with lower Q2 values (Table 2). Thus,  
576 although these watersheds have a high likelihood of contributing to streamflow of major rivers,  
577 the watershed Q2 contributions were predicted to be small (Table 4).

578

579 *Grasslands (C5, C6, and C7)*

580 The southwestern Canadian Prairie, which includes the majority of southern Alberta and  
581 western Saskatchewan between the South Saskatchewan River and the Cypress Hills, was  
582 occupied by classes C5, C6, and C7. These watersheds tended to have large fractions of  
583 unmanaged grasslands (negative land cover PC1) and mean elevation (Table 4). Compared to the  
584 rest of the Prairie, this sub-region tended to be arid, with a strong moisture deficit (Fig. 6). As a  
585 result, these classes exhibited relatively low wetland density (Fig. 8a).

586 Classes 5 (C5; n = 635), Interior Grasslands, and 6 (C6; n = 702), High-Elevation  
587 Grasslands, were characteristic of the grasslands in southeastern Alberta. These watersheds had  
588 the greatest values of mean fractional grassland area, with cropland and grassland fractions being  
589 comparable (35–40%) (Fig. 7). Distinguishing features of Interior Grasslands were greater values

590 of the fraction of area below the basin outlet,  $A_{BO}$ , and a notably large non-effective area fraction  
591 (Fig. 7a). High scores on land cover PC2 and PC3 indicate large fractions of fallow and pasture.  
592 These watersheds also scored higher on soil zone PC2, suggesting more common occurrences of  
593 brown soils. Small magnitudes of mean slope and stream densities were observed, suggesting  
594 that the wetlands within the Interior Grasslands are relatively disconnected from the drainage  
595 network. This characteristic might explain why these watersheds have relatively large wetlands  
596 (Fig. 8c). In contrast, High Elevation Grasslands were characterized by greater mean elevation  
597 and slope values, and smaller non-effective fractions (Table 4; Fig. 7). These watersheds also  
598 had greater stream densities and smaller wetland densities.

599 Class 7 (C7;  $n = 377$ ), Sloped Incised, watersheds are characterized by dissected, river-  
600 incised landscapes, as indicated by positive associations with local surface form PC3 (Table 4).  
601 Like High Elevation Grasslands (C6), Sloped Incised watersheds followed the Bow, Red Deer,  
602 as well as the Milk River valleys, suggesting a similar function to those of the Major River  
603 Valleys class. Wetland density is smallest in Sloped Incised watersheds, owing to their steepness,  
604 resulting in surface water reaching stream networks rather than collecting on the landscape (Fig.  
605 8).

606

#### 607 4.3. Predicting wetland size distributions from class parameters

608 Simulated wetland area distributions by class were compared to observed size  
609 distributions from study watersheds to evaluate the concordance of the approximate class-  
610 specific distribution to that of the observed distributions of watersheds, collectively. The median  
611 wetland density was greatest in C2, followed by C3, C1, and C5 (Fig. 8a). The median wetland  
612 densities in C6 and C7 were less than  $1 \text{ km}^{-2}$ . C4 had the greatest areal fraction of water in the  
613 largest wetland ( $W_L$ ), which was over 40% (Fig. 8b), while C2 had the smallest value at ~10%.  
614 For the rest of the classes, this value was between 28% and 34%. The simulated wetland area  
615 distributions slightly overestimated those of the observed values, especially at the 25<sup>th</sup> percentile.  
616 However, the patterns of wetland area in the quartiles was generally consistent among all classes  
617 (Fig. 8c). The area of the smallest 25% of the wetlands appears quite consistent across the  
618 classes, with more variation occurring at higher percentiles. The largest difference among classes  
619 in wetland size was in the 75<sup>th</sup> percentile, with the greatest range being in C5 and the smallest in  
620 C1.

621

#### 622 4.4. Resampling and re-classifying procedure

623 The HCPC and watershed classification was repeated with ten random subsets of 3757  
624 watersheds. The majority of watershed were removed from at least one iteration, with only 50  
625 watersheds being removed a total of 4-6 times (Fig. S3). This resulted in ten unique watershed  
626 subsets to test clustering and agreement to the seven classes, outlined above.

627 Percent membership agreement of a watershed varied by class, with the majority of  
628 classes exhibiting high agreement even after resampling. Classes exhibiting high membership  
629 agreement were Pothole Till (C2), Interior Grasslands (C5), High Elevation Grasslands (C6), and  
630 Sloped Incised (C7), with a large proportion having more than 90% agreement with the seven  
631 classes from the complete classification (Fig. 9; Table S4). Although a large mean agreement  
632 was observed overall, a few watershed classes exhibited low agreement and inconsistent  
633 classification. Southern Manitoba (C1) exhibited a bimodal distribution, where most were  
634 generally classed as C1 over 75% of the time and a second set only ~60% agreement (Fig. 9).  
635 This was due to a new class appearing (Fig. 10). Hereafter, this class is referred to as “Eastern  
636 Manitoba”. Briefly, Eastern Manitoba was association with large fraction of conventional tillage  
637 practice (i.e., positive association with land practice PC1 and land practice PC2) and large  
638 fractional effective areas (data not shown). The Major River Valleys class was the only one that  
639 did not include a watershed that achieved 100% agreement across the ten iterations; this class  
640 exhibited a peak of total agreement at approximately 60% (Fig. 9). Where Major River Valleys  
641 watersheds were classified inconsistently, the most common alternative classification were  
642 Pothole Glaciolacustrine (C3) or secondarily High Elevation Grasslands (C6) (Fig. 10). The loss  
643 of Major River Valleys occurred for iterations when the Eastern Manitoba class (C8) became  
644 apparent.

645

## 646 5. DISCUSSION

647

### 648 5.1. Classifying Prairie watersheds

#### 649 5.1.1. Hydrological approaches

650 Our classification procedure grouped watersheds of approximately 100 km<sup>2</sup> into seven  
651 classes. Few studies [anywhere](#) have classified watersheds [at this granularity, and our](#)

652 ~~investigation gives specifically within the Canadian Prairie with~~ particular attention to ~~these~~  
653 characteristics that ~~control~~ influence hydrological ~~and ecological~~ behaviour. Many previous  
654 studies ~~in the region~~ spanned larger areas, and this often results in the Prairie being identified as  
655 a homogenous region due to relatively low streamflow and atypical geology and surface  
656 topography (MacCulloch and Whitfield, 2012; Mwale et al., 2011). ~~Our results are novel in that~~  
657 ~~they characterize in greater detail, and at small watershed scales, the potential for different~~  
658 ~~hydrological behaviour of watersheds within the region.~~ The only ~~similar~~ example that was  
659 found in the literature was by Durrant and Blackwell (1959), whose findings parallel those of this  
660 study, ~~but at a larger watershed scale.~~ Durrant and Blackwell (1959) described broad regions of  
661 Saskatchewan and Manitoba based on mean annual flood, distinguishing five sub-regions  
662 including southwestern Saskatchewan, north and central Saskatchewan, and southern Manitoba  
663 near the Red River and Assiniboine River confluence. In the current study, surficial geology and  
664 land surface form strongly influenced how grasslands were separated into three classes, which  
665 reinforces the role of local topography on hydrological response, as seen elsewhere (Mwale et  
666 al., 2011). Likewise, surficial geology was particularly important for distinguishing the Pothole  
667 (Till and Glaciolacustrine) classes. Similarities to the work of Durrant and Blackwell (1959)  
668 based on streamflow in larger basins suggest that our approach, with consideration of factors  
669 important to watershed behaviour, can yield classification with relevance to hydrologic function,  
670 despite the use of few hydrologic indices in our analysis (Fig. 5). This approach holds potential  
671 for use in other regions of the world that are dry, ungauged, or feature low effective areas, and  
672 thus cannot rely on streamflow characteristics as a primary means of classification according to  
673 functional behaviour.

674 ~~The~~ Our classification grouped Prairie watersheds using geological, biophysical, and  
675 hydroclimatic attributes. In their review of classification approaches, Sivakumar et al. (2013)  
676 indicate that solely using ~~physiographic~~ geographic data is advantageous when there are limited  
677 hydrological data; however, the relationship between physical attributes and hydrologic  
678 behaviour is not necessarily definitive in all regions. For these reasons, it was important to  
679 include traits indicative of structural hydrological connectivity, such as Q2 estimates and wetland  
680 parameters. It is important to note that while Q2 emerged as a defining feature for several of the  
681 classes, it was ~~always~~ consistently one of many variables important for characterization of that  
682 class (Table 4), suggesting that while it provides value added, it does not stand out as a major

683 driving factor in the classification. In particular, the immature drainage network and relatively  
684 high depressional water storage capacity make prairie hydrology relatively distinct (Jones et al.,  
685 2014; Shook et al., 2013, 2015). Notably, three classes (i.e., Pothole Till, Pothole  
686 Glaciolacustrine, and Interior Grasslands) occur almost exclusively within regions that tend not  
687 to contribute to major river systems, and collectively encompass the majority of the study area  
688 (Table 4; Fig. 5). It is therefore expected that hydrological response will be very different  
689 between classes that exhibit higher hydrological connectivity (i.e., potentially lower wetland to  
690 stream densities and non-effective area fractions), such as the Major River Valleys or Sloped  
691 Incised watersheds, than those that do not, such as Pothole classes.

692

### 693 *5.1.2. Ecoregions and human impacts*

694 Ecoregions are commonly used to characterize landscapes according to geographical or  
695 ecological similarity (~~Masaki and Rosenberry, 2002;~~ Omernik and Griffith, 2014). Similar to our  
696 approach, ecoregion classifications are often hierarchical in nature, allowing for differing levels  
697 of detail, spatial extent, and thus defining characteristics depending on the scale of interest  
698 (Loveland and Merchant, 2004). Ecoregion classifications used in the United States (Omernik  
699 and Griffith, 2014) and Canada (Ecological Stratification Group, 1995) employ a “top-down”  
700 approach, where broad categories are partitioned into smaller, more specialized units. In contrast,  
701 our approach provides a bottom-up, agglomerative approach where similar watersheds are  
702 merged. Assumptions are inherent in either approach; however, the latter was applicable to the  
703 current study to allow for grouping of watersheds given similarities in ~~physio-geographic~~  
704 characteristics. This approach does not limit class ~~membershipification~~ to the geographic extent  
705 of a higher level class, allowing for ~~class-membership~~ to ~~potentially~~ span ~~a large the~~ geographic  
706 extent of the Canadian Prairie domain (Fig. 5).

707 Despite the differing methods for distinguishing similarities (or differences),  
708 arrangements of watershed classes in some cases exhibited similar ranges to ecoregion  
709 boundaries. The boundaries of Lake Manitoba Plain and Mixed Grassland ecoregions  
710 (Ecological Working Group, 1995) correspond roughly to those of the broader Southern  
711 Manitoba (C1) and Grasslands (C5, C6, and C7) classes, respectively (Fig. S4). Mwale et al.  
712 (2011) also found that annual hydrological regimes based on data from 200 stations and physical  
713 attributes in Alberta linked closely with provincial ecoregions. Our emphasis on inclusion of

714 hydrologically relevant characteristics, such as wetland traits and effective areas that are likely  
715 important contributors to function, has proven useful for further distinguishing among the  
716 Grassland classes as well as the Pothole classes (C2 and C3) (Fig. 5; Fig. S4). Due to the  
717 fundamental differences in effective areas and in wetland versus river dominated systems (Table  
718 4; ~~Fig. 8; Fig. 8~~), we expect different hydrological behaviour between these classes. This is an  
719 advantage of the HCPC classification approach in that it allows for identifying the potential  
720 similarity at relatively fine spatial scales, and does not require similar watersheds to be  
721 physically adjacent to one another. This confers the opportunity to further investigate these  
722 systems, such as (e.g., through hydrological modelling and contrasting resulting responses under  
723 climate and land-use scenarios).

724 The highly managed prairie landscape reinforces the importance of considering  
725 anthropogenic alteration in hydrological understanding. Crop rotation and the ways in which  
726 fields are managed for winter affect the accumulation and redistribution of snow (Fang et al.,  
727 2010; Harder et al., 2018; Van der Kamp et al., 2003). Spring snowmelt and consequent runoff  
728 are imperative to summer surface water availability (Dumanski et al., 2015; Shook et al., 2015),  
729 and depression-focused recharge of snowmelt into groundwater facilitates storage and mitigates  
730 flood impacts (Hayashi et al., 2016). Thus, classifying procedures in the Prairie must consider  
731 the human influence on the water cycle.

732 An example of the complexities introduced by human land management activities can be  
733 shown by the C1 (Southern Manitoba) watersheds, where the land practice variable was a strong  
734 class descriptor. Agricultural activity is high everywhere in the Prairie; however, only C1 was  
735 associated with low zero-till practices, instead favouring conventional tillage (Table 4).  
736 Manitoba has seen less coherent adoption of zero-till practices since the early 1990s in compared  
737 to Alberta and Saskatchewan, with conventional or other conservation till practices remaining  
738 common in Manitoba (reviewed in Awada et al., 2014). Sustained use of conventional tillage  
739 practice within this region may increase the risk of soil erosion, which can negatively affect  
740 downstream water bodies (Cade-Menun et al., 2013). This practice, combined with landscape  
741 modifications, such as artificial drainage networks, serve to facilitate removal of water and may  
742 contribute to concurrent nutrient export from agricultural lands (Weber et al., 2017).

743 These management practices can be viewed as a trade-off, where high numbers of  
744 wetlands and level topography can pose flood risk during wet periods as wetlands fill and merge

745 (Leibowitz et al., 2016), inundating tracts of adjacent land. Conversely, where landscape  
746 modification to enhance water export occurs, local, field-scale flood risk may be reduced, while  
747 heightening the risk of downstream flooding. Land-use and land management are important  
748 factors in understanding the connectivity and chemical transport in prairie landscapes (Leibowitz  
749 et al., 2018). In southern Manitoba, where artificial drainage has been used to increase the area of  
750 arable land, beneficial management practices in the form of agricultural reservoirs have been  
751 implemented as a means of reducing nutrient export and improving downstream water quality  
752 while also mitigating the risk of downstream flooding (Gooding and Baulch, 2017). These  
753 factors illustrate the complexities when classifying and understanding hydrological response of  
754 watershed embedded in highly managed landscapes, and underscore that necessity of considering  
755 the human influence on the water cycle in such approaches.

756

## 757 5.2. HCPC as a clustering and classification framework

### 758 5.2.1. Using the HCPC approach and limitations

759 The HCPC method provides a procedure for integrating multiple physio-~~geo~~geographic  
760 attributes and describes resulting clusters by sets of significant variables (Husson et al., 2009).  
761 As discussed above, an advantage of the method is that it groups individual watersheds based on  
762 similarities, ~~and T~~therefore, it lends itself well ~~as a~~ setting a foundation for investigating  
763 hydrological behaviour ~~to be applied to~~through modelling efforts. In the case of the current  
764 study, modelling efforts can be applied at a 100 km<sup>2</sup> scale to evaluate responses to environmental  
765 changes. An additional advantage is that that one may select variables or sets of variables of  
766 interest to inform the clustering of watersheds, such as those based only on topographic  
767 parameters or those dictating local hydrology. For example, climate variables may be excluded if  
768 the goal of the classification is ~~informing application~~parameterizing of a hydrological model, as  
769 these variables could instead be ~~part of~~described by ~~model parameterization~~local climate forcing.  
770 The relative ease with which different sets of variables can be added to or excluded from the  
771 analysis to consider different permutations of the classification is a real strength of the approach.  
772 Although this may result in differing cluster results, assessment of how these classes change with  
773 addition or removal of certain datasets can identify the variables that control class definition as  
774 well as elucidate spatial patterning of classes.

775           There are a few considerations when using this method. First, the linear restrictions of  
776 this method are challenging when working with environmental data, which often do not conform  
777 to assumptions of normality. Non-linear PCA methods and self-organizing maps have been  
778 applied successfully to classify watersheds in Ontario and to regionalize streamflow metrics  
779 (Razavi and Coulibaly, 2013, 2017). Although these methods might be logical next steps for the  
780 current study, we chose to focus on conventional PCA due to its smaller computational cost  
781 when classifying the large number of watersheds in our study.

782           Second, the current analysis weighs all variables equally. This can bias the analysis  
783 towards attributes that exhibit greater variability, as these can overshadow other more  
784 constrained variables. For example, the location of the largest pond relative to the watershed  
785 outlet (coded as Lw/Lo) is important to controlling local prairie hydrology and hydrological gate-  
786 keeping potential (i.e., the likelihood of releasing surface water to the next order watershed)  
787 (Shook et al., 2013, 2015) and water quality (Hansen et al., 2018). Despite its hydrological  
788 importance, this variable had little influence on the clustering procedure overall, and was only a  
789 minor descriptor in certain classes, such as C5 and C6 (Table 4).

790           ~~The classes resulting from the HCPC are ultimately dependent on the types of data  
791 included. The availability of data and its geographic coverage determined the environmental  
792 parameters included in our analyses. Ideally, a more detailed estimate of runoff for each  
793 watershed, as well as a soil moisture dataset would have been include. A comprehensive wetland  
794 inventory or an index of wetland drainage activity that is comparable across the three Provinces  
795 does not currently exist. These would be valuable additions to future efforts to classify Prairie  
796 watersheds given the important role of land modification on watershed functions.~~

797           Class membership in our approach is also determinate. In reality, there can be large  
798 variability in attributes within a class (e.g., Fig. 7), and membership is determined by the  
799 collective similarity of watershed attributes. Previous studies have used fuzzy c-means and  
800 Bayesian approaches that can assign a likelihood of membership to classes (Jones et al., 2014;  
801 Rao and Srinivas, 2006; Sawicz et al., 2011). An advantage to this approach is that it allows for  
802 fuzzy boundaries between classes where a gradient of features likely exists (Loveland and  
803 Merchant, 2004). Our re-classifying analysis supports the proposition that boundaries among  
804 classified regions are fuzzy and some watershed might flicker among class memberships (Fig.  
805 10). Such approaches, which are also un-supervised and are probabilistic in nature and will

806 eliminate the subjectivity due to the researcher pre-defining the number of classes. Our future work  
807 thus should consider these fuzzy boundaries and potential for watersheds to exhibit partial  
808 membership to multiple classes. ~~will include applying a fuzzy cluster Bayesian framework to~~  
809 ~~assess the current classification framework.~~

### 811 *5.2.2. Data quality and availability*

812 The classes resulting from the HCPC are also ultimately dependent on the types of data  
813 included. The availability of data and its geographic coverage determined the environmental  
814 parameters included in our analyses. Ideally, a more detailed estimate of runoff for each  
815 watershed would be a valuable contribution. In the current study, we used the CCA and eleven  
816 reference stations to approximate runoff values for the clustering watersheds. Given the number  
817 of watersheds included in the analyses, the diversity of physical characteristics and potential  
818 hydrological behaviour is likely not completely represented in the small sample size of available  
819 hydrometric stations, and is a limitation of our approach. Soil moisture would be important to  
820 consider in future studies given its role in influencing vegetation community composition, PET,  
821 and over all water balance (Hayashi et al., 2003; Shook et al., 2015). Where data is available,  
822 future work should consider variables related to snow formation and melt, as well the proportion  
823 of annual precipitation as snowfall. These variables are likely influential when describing  
824 hydrological behaviour of the watersheds and classes in the current study, and other cold regions  
825 (Knoben et al., 2018; Shook and Pomeroy, 2012). Furthermore, a comprehensive wetland  
826 inventory or an index of wetland drainage activity that is comparable across the three Provinces  
827 does not currently exist. These would be valuable additions to future efforts to classify Prairie  
828 watersheds given the important role of land modification on watershed functions.

829 One consideration with the Global Surface Water dataset is that the pixel size (30 m) is  
830 quite coarse and will miss numerous smaller wetlands, underestimating the number of wetlands  
831 observed. Consequently, it is likely that the analysis omitted some ephemeral wetlands for which  
832 persistence is short and size is small. Despite their known important ecological functions  
833 (Calhoun et al., 2017; Van Meter and Basu, 2015), their size and transient nature is a challenge  
834 to their inclusion in comprehensive datasets spanning large geographic areas. This may  
835 inadvertently result in the role of smaller wetlands being under-represented in our analysis, or  
836 others that rely on this dataset.

837 Use of the  $\zeta$  and  $\beta$  parameters as indices of the wetland area frequency distributions were  
838 shown to estimate classes area distributions reasonably well (Fig. 8c). Although for consistency,  
839 we restricted our simulated dataset to the spatial resolution of the surface water raster, one could  
840 use these parameters to estimate the frequencies of smaller wetlands in watersheds, which would  
841 otherwise be missed by satellite measurements, assuming conformity to a Generalized Pareto  
842 Distribution (Shook et al., 2013). Our analysis supports this application as simulated wetland  
843 areas generally approximated those seen across the observed data (Fig. 8c). Nonetheless, in  
844 regions where wetland drainage has been undertaken, it is expected that wetland area distribution  
845 has been altered via preferential loss of smaller water bodies (Evenson et al., 2018; Van Meter  
846 and Basu, 2015). This is exacerbated by the fact that remote sensed satellite data tends to omit  
847 smaller, ephemeral ponds. A more robust characterization of the size and permanence of  
848 wetlands in our study watersheds would be expected to improve the current dataset and enhance  
849 the clustering and classification analyses.

850

### 851 5.3. Management implications

852 Classification frameworks help to define sub-regions with potentially similar  
853 characteristics or behaviours. For example, climatic zones can be delineated, specifically the dry  
854 Grassland watersheds in the southwest and the wet Potholes in the northeast and in Manitoba  
855 (Fig. 5). In some cases, this may be related to local wetland densities, with large densities  
856 observed corresponding with low moisture deficits (Fig. 6b) (Liu and Schwartz, 2012). Climate  
857 variation may divide watersheds with seemingly similar ~~physio~~-geography into differing classes,  
858 as is the case with Major River Valleys and Sloped Incised watersheds. Both sets of watersheds  
859 tended to follow river valleys, but the former exhibit greater precipitation and smaller PET  
860 (Table 4). These divisions can be used to give context to regions we might expect to behave  
861 similarly, whether hydrologically, or ecologically, based solely on physical attributes, and echoes  
862 other methods, such as ecodistricts (Ecological Stratification Working Group, 1995) to classify  
863 landscapes. For example, areas that are geologically similar may differ in terrestrial or aquatic  
864 community assemblages, which should influence how each area might be managed (Jones et al.,  
865 2014; Wagner et al., 2007). If classifications are used to inform management, the resulting  
866 decisions for a given location will depend on the strength of the delineation, the scale at which  
867 management is applied, relationships among management practices and the attributes used to

868 define that area, and the relationship of those attributes to the response variable of concern  
869 (Wagner et al., 2007)(~~Wagener et al., 2007~~).

870 This set of analyses was unique among watershed classification exercises in Canada in  
871 that it considered a suite of wetland variables. The arrangement of wetlands or landscape  
872 depressions and their size distribution define the hydrological behavior of Prairie watersheds  
873 (Shook et al., 2015; Shook and Pomeroy, 2011). The storage capacity and subsequent spilling or  
874 merging controls wetland connectivity, and thus the quantity of water available to move from  
875 one watershed to another (Leibowitz et al., 2016; Shaw et al., 2012; Shook et al., 2015). In turn,  
876 a wetland or depression's hydrological gate-keeping potential, or its likelihood to prevent  
877 connectivity to the downstream watershed, is a function of both its storage capacity and  
878 landscape position. Large wetlands near an outlet have a great gate-keeping potential, as they  
879 block much of the watershed from connecting, and it takes a great deal of water to fill them  
880 before permitting flow to the next order watershed (Shook and Pomeroy, 2011). Simulated  
881 frequency distributions of wetland areas indicate that the depressional storages of the classes are  
882 very different (Fig. 8). It may be that wetland management practices will have different  
883 influences between each pothole class, and possibly among all the classes. This has implications  
884 for managing salinizing soils (Goldhaber et al., 2014), biodiversity (Balas et al., 2012), and  
885 floodings potential (Evenson et al., 2018; Golden et al., 2017).

886 Wetland drainage and wetland consolidation change hydrological connectivity and  
887 therefore the transport of nutrients and their loading into receiving water bodies (Brown et al.,  
888 2017; Vanderhoof et al., 2017). More positive values of the moisture deficit (i.e., where  $P \geq$   
889  $PET$ ) were associated with greater wetland densities (Fig. 6b) (Liu and Schwartz, 2012), and  
890 these areas were generally associated with greater fractions of cropland, such as Pothole Till,  
891 Pothole Glaciolacustrine, and Southern Manitoba watersheds. In these regions wetland drainage  
892 is widely practiced, historically or at present, and conflict over available arable land and wetland  
893 conservation is high (Breen et al., 2018).

894 Extensive drainage in combination with agricultural activity is known to increase the risk  
895 of agricultural nutrient mobility (Kerr, 2017) from the landscape to receiving water bodies.  
896 Increased connectivity also reduces water residence time and thus tends to decrease wetland  
897 nutrient retention (Marton et al., 2015). Over time, zero-till practices can promote nutrient  
898 stratification in soils, where concentrations (especially phosphorus) accumulate at the surface,

899 which can increase nutrient loading when surface runoff is generated (Cade-Menun et al., 2013).  
900 The cropland-wetland interface might also have important implications for pesticide mobility in  
901 Pothole Till and northern Pothole Glaciolacustrine watersheds. These areas coincide with  
902 extensive use of canola, which has been linked to high application rates of neonicotinoid  
903 pesticides which are known to have high persistence in small, temporary wetlands (Main et al.,  
904 2014). Watersheds in the Pothole Till class appear to have more hummocky landscapes than the  
905 Pothole Glaciolacustrine classification and smaller, more numerous wetlands (Fig. 8). Moreover,  
906 the water area fraction occupied by the largest wetland differs between the classes. The  
907 landscape biogeochemical functionality of pothole wetlands is known to vary considerably  
908 according to pothole character (Evenson et al., 2018; Van Meter and Basu, 2015). As such, our  
909 classification may highlight contrasting biogeochemical functioning, including nutrient retention,  
910 between these classes. Thus, although water quality risks are common within the region, the  
911 classes may respond very differently to environmental and land management stresses.

912

## 913 **6. CONCLUSION**

914

915 This study provides an overview of a classification framework that can be applied in  
916 regions with limited understanding of or data describing streamflow. The HCPC procedure offers  
917 a flexible analysis to elucidate the spatial arrangement of watershed classes given a large number  
918 of units to classify and a diverse set of attributes to inform the classification. In contrast to  
919 classifications based solely on hydrological function, using physio-geographic data allows for  
920 classifying small basins, which are unlikely to be gauged, and confers advantages over alternate  
921 procedures that rely heavily on observations of hydrological parameters, namely statistics  
922 describing streamflow.

923 Use of the classification approach for small Canadian Prairie watersheds identified  
924 regions of similar climatic and physio-geographic features and, potentially, of hydrological  
925 response (Fig. 5). This yielded watershed classes that consider not only drainage patterns, but  
926 also land cover, and land-use, and the underlying geology. In the Prairie region, wetland  
927 variables incorporate the hydrologic gate-keeping potential of wetlands as well as parameters  
928 indicative of wetland size distributions. With the classification based on a large and diverse set of  
929 attributes, a diversity of behaviours is captured. This represents a major step forward for

930 classification of Prairie watersheds that have to-date offered only a much more homogenized  
931 depiction of watershed function in the region. The watershed classification framework presented  
932 promises to be useful in other dry or semi-arid regions, and those that are poorly gauged. Given  
933 the inclusive nature of the classification approach, which incorporates landscape controls on  
934 hydrology as well as those influencing biogeochemistry and ecology, it also provides a  
935 foundation to evaluate the efficacy of land and watershed management practices in the context of  
936 a changing climate.

937  
938  
939  
940  
941  
942  
943

944 **Author contributions**

945 JDW, CJW, and CS conceived the study, and JDW collected data and performed analyses. KRS  
946 wrote code to analyze basin and wetland data. JDW wrote the manuscript with input from all co-  
947 authors.

948

949 **Acknowledgements**

950 This work was pursued under the Prairie Water project and funded by the Global Water Futures  
951 program, which was supported by the Canada First Research Excellence Fund. The authors  
952 would like to thank John Pomeroy for his valuable input on the scoping and approach to the  
953 study. We would also like to thank Wouter Knoben and two anonymous reviewers for their  
954 insightful comments on the manuscript. Finally, we would like to thank the Prairie Water team  
955 and the Global Institute for Water Security for ongoing support. The authors declare that they  
956 have no conflict of interest.

957

958

959 **REFERENCES**

- 960
- 961 AAFC: Annual Crop Inventory, Agriculture and Agri-Food Canada, Government of Canada.  
962 Available from: <https://open.canada.ca/data/en/dataset/ba2645d5-4458-414d-b196-6303ac06c1c9>, 2016.
- 963
- 964 AAFC: Detailed Soil Surveys, Agriculture and Agri-Food Canada, Government of Canada..  
965 Available from: <https://open.canada.ca/data/en/dataset/7ed13bbe-fbac-417c-a942-ea2b3add1748>, 2015.
- 966
- 967 AAFC: Soils of Canada, Derived. Soil Landscapes of Canada and Detailed Soil Surveys, version  
968 3.2. Canadian Soil Information Service, Agriculture and Agri-Food Canada, Government of  
969 Canada. Available from: <https://open.canada.ca/data/en/dataset/8f496e3f-1e54-4dbb-a501-a91eccf616b8>, 2013.
- 970
- 971 Atkinson, N., Utting, D. J., and Pawley, S. M.: Surficial geology of west-central Alberta (GIS  
972 data, polygon features); Alberta Energy Regulator, AER/AGS Digital Data 2017-0031.  
973 Available from: [http://ags.aer.ca/publications/DIG\\_2017\\_0031.html](http://ags.aer.ca/publications/DIG_2017_0031.html), 2017.
- 974 Awada, L., Lindwall, C. W. and Sonntag, B.: The development and adoption of conservation  
975 tillage systems on the Canadian Prairies, *Int. Soil Water Conserv. Res.*, 2, 47–65,  
976 doi:10.1016/S2095-6339(15)30013-7, 2014.
- 977 Balas, C. J., Euliss, N. H. and Mushet, D. M.: Influence of conservation programs on amphibians  
978 using seasonal wetlands in the prairie pothole region, *Wetlands*, 32, 333–345,  
979 doi:10.1007/s13157-012-0269-9, 2012.
- 980 Begou, J., Bazie, P. and Afouda, A.: Catchment classification: multivariate statistical analyses  
981 for physiographic similarity in the Upper Niger Basin, *J. Eng. Res. Appl.*, 5, 60–68, 2015.
- 982 Breen, S.-P. W., Loring, P. A. and Baulch, H.: When a Water Problem Is More Than a Water  
983 Problem: Fragmentation, Framing, and the Case of Agricultural Wetland Drainage, *Front.*  
984 *Environ. Sci.*, 6, 1–8, doi:10.3389/fenvs.2018.00129, 2018.
- 985 Brown, R., Zhang, Z., Comeau, L. P. and Bedard-Haughn, A.: Effects of drainage duration on  
986 mineral wetland soils in a Prairie Pothole agroecosystem, *Soil Tillage Res.*, 168, 187–197,  
987 doi:10.1016/j.still.2016.12.015, 2017.
- 988 Brown, S. C., Lester, R. E., Versace, V. L., Fawcett, J. and Laurenson, L.: Hydrologic landscape  
989 regionalisation using deductive classification and random forests, *PLoS One*, 9,  
990 doi:10.1371/journal.pone.0112856, 2014.
- 991 Bulley, H.N.N., Marx, D.B., Merchant, J.W., Holz, J.C., and Holz, A.A.: A comparison of  
992 Nebraska reservoir classes estimated from watershed-based classification models and  
993 ecoregions, *J. Environ. Informatics*, 11, 90–102, doi:10.3808/jei.200800114, 2008.
- 994 Burn, D.: Cluster analysis as applied to regional flood frequency, *J. Water Resour. Plan. Manag.*,  
995 115, 567–582, 1990.
- 996 Cade-Menun, B. J., Bell, G., Baker-Ismail, S., Fouli, Y., Hodder, K., McMartin, D. W., Perez-  
997 Valdivia, C. and Wu, K.: Nutrient loss from Saskatchewan cropland and pasture in spring  
998 snowmelt runoff, *Can. J. Soil Sci.*, 93, 445–458, doi:10.4141/cjss2012-042, 2013.
- 999 Calhoun, A. J. K., Mushet, D. M., Bell, K. P., Boix, D., Fitzsimons, J. A. and Isselin-Nondedeu,  
1000 F.: Temporary wetlands: challenges and solutions to conserving a “disappearing” ecosystem,  
1001 *Biol. Conserv.*, 211, 3–11, doi:10.1016/j.biocon.2016.11.024, 2017.

1002 Cavadias, G. S., Ouarda, T. B. M. J., Bobee, B. and Girard, C.: A canonical correlation approach  
1003 to the determination of homogeneous regions for regional flood estimation of ungauged  
1004 basins, *Hydrol. Sci. J.*, 46, 499–512, doi:10.1080/02626660109492846, 2001.

1005 Coles, A. E., McConkey, B. G. and McDonnell, J. J.: Climate change impacts on hillslope runoff  
1006 on the northern Great Plains, 1962–2013, *J. Hydrol.*, 550, 538–548,  
1007 doi:10.1016/j.jhydrol.2017.05.023, 2017.

1008 DeBeer, C. M., Wheeler, H. S., Carey, S. K. and Chun, K. P.: Recent climatic, cryospheric, and  
1009 hydrological changes over the interior of western Canada: A review and synthesis, *Hydrol.*  
1010 *Earth Syst. Sci.*, 20, 1573–1598, doi:10.5194/hess-20-1573-2016, 2016.

1011 Detenbeck, N. E., Batterman, S. L., Brady, V. J., Brazner, J. C., Snarski, V. M., Taylor, D. L.,  
1012 Thompson, J. A. and Arthur, J. W.: a Test of Watershed Classification Systems for  
1013 Ecological Risk Assessment, *Environ. Toxicol. Chem.*, 19, 1174, doi:10.1897/1551-  
1014 5028(2000)019<1174:ATOWCS>2.3.CO;2, 2000.

1015 Devito, K., Creed, I., Gan, T., Mendoza, C., Petrone, R., Silins, U. and Smerdon, B.: A  
1016 framework for broad-scale classification of hydrologic response units on the Boreal Plain: Is  
1017 topography the last thing to consider?, *Hydrol. Process.*, 19, 1705–1714,  
1018 doi:10.1002/hyp.5881, 2005.

1019 Dumanski, S., Pomeroy, J. W. and Westbrook, C. J.: Hydrological regime changes in a Canadian  
1020 Prairie basin, *Hydrol. Process.*, 29, 3893–3904, doi:10.1002/hyp.10567, 2015.

1021 Durrant, E. F. and Blackwell, S. R.: The magnitude and frequency of floods in the Canadian  
1022 Prairies, *Proc. Of Symposium No. 1, Spillway Design Floods*, sub-committee on hydrology,  
1023 National Research Council Associate Committee on Geodesy and Geophysics. The Queens  
1024 Printer, Ottawa, 1959.

1025 ECCC: Hydatt series. Water Survey of Canada, Environment and Climate Change Canada.  
1026 Government of Canada, 2016. Available from: [https://ec.gc.ca/rhc-  
1027 wsc/default.asp?n=9018B5EC-1](https://ec.gc.ca/rhc-wsc/default.asp?n=9018B5EC-1), 2016.

1028 ECCC: Canadian Gridded Temperature and Precipitation Anomalies. Environment and Climate  
1029 Change Canada. Government of Canada. Accessed November 2017. Available from:  
1030 <https://open.canada.ca/data/dataset/3d4b68a5-13bc-48bb-ad10-801128aa6604>, 2017.

1031 Ecological Stratification Working Group: A National Ecological Framework for Canada.  
1032 Agriculture and Agri-Food Canada, Research Branch, Centre for Land and Biological  
1033 Resources Research and Environment Canada, State of the Environment Directorate,  
1034 Ecozone Analysis Branch, Ottawa/Hull. Report and national map at 1:7500 000 scale, 1995

1035 Evenson, G. R., Golden, H. E., Lane, C. R., McLaughlin, D. L. and D’Amico, E.: Depressional  
1036 wetlands affect watershed hydrological, biogeochemical, and ecological functions, *Ecol.*  
1037 *Appl.*, 28, 953–966, doi:10.1002/eap.1701, 2018.

1038 Fang, X., Pomeroy, J. W., Westbrook, C. J., Guo, X., Minke, A. G. and Brown, T.: Prediction of  
1039 snowmelt derived streamflow in a wetland dominated prairie basin, *Hydrol. Earth Syst. Sci.*,  
1040 14, 991–1006, doi:10.5194/hess-14-991-2010, 2010.

1041 [Geological Survey of Canada: Surficial geology of Canada / Géologie des formations  
1042 superficielles du Canada. Canadian Geoscience Map, Geological Survey of Canada,  
1043 Government of Canada. 195, <https://doi.org/10.4095/295462>, 2014](https://doi.org/10.4095/295462)

1044 Gibson, J. J., Birks, S. J., Jeffries, D. S., Kumar, S., Scott, K. A., Aherne, J. and Shaw, D. P.:  
 1045 Site-specific estimates of water yield applied in regional acid sensitivity surveys across  
 1046 western Canada, *J. Limnol.*, 69, 67–76, doi:10.3274/JL10-69-S1-08, 2010.

1047 Gibson, J. J., Yi, Y. and Birks, S. J.: Isotope-based partitioning of streamflow in the oil sands  
 1048 region, northern Alberta: Towards a monitoring strategy for assessing flow sources and  
 1049 water quality controls, *J. Hydrol. Reg. Stud.*, 5, 131–148, doi:10.1016/j.ejrh.2015.12.062,  
 1050 2016.

1051 Gober, P., and Wheeler, H.S. Socio-hydrology and the science-policy interface: a case study of  
 1052 the Saskatchewan River basin. *Hydrol. Earth Syst. Sci.*, 18, 1413–1422, doi: 10.5194/hess-  
 1053 18-1413-2014, 2014.

1054 Godwin RB, Martin FRJ. Calculation of gross and effective drainage areas for the Prairie  
 1055 Provinces. In: Canadian Hydrology Symposium - 1975 Proceedings, 11-14 August 1975,  
 1056 Winnipeg, Manitoba. Associate Committee on Hydrology, National Research Council of  
 1057 Canada, pp. 219–223. 1975.

1058 Golden, H. E., Creed, I. F., Ali, G., Basu, N. B., Neff, B. P., Rains, M. C., McLaughlin, D. L.,  
 1059 Alexander, L. C., Ameli, A. A., Christensen, J. R., Evenson, G. R., Jones, C. N., Lane, C. R.  
 1060 and Lang, M.: Integrating geographically isolated wetlands into land management decisions,  
 1061 *Front. Ecol. Environ.*, 15, 319–327, doi:10.1002/fee.1504, 2017.

1062 Goldhaber, M. B., Mills, C. T., Morrison, J. M., Stricker, C. A., Mushet, D. M. and LaBaugh, J.  
 1063 W.: Hydrogeochemistry of prairie pothole region wetlands: Role of long-term critical zone  
 1064 processes, *Chem. Geol.*, 387, 170–183, doi:10.1016/j.chemgeo.2014.08.023, 2014.

1065 Gooding, R. M. and Baulch, H. M.: Small reservoirs as a beneficial management practice for  
 1066 nitrogen removal, *J. Environ. Qual.*, 46, 96-104, doi:10.2134/jeq2016.07.0252, 2017.

1067 Hansen, A. T., Dolph, C. L., Fofoula-Georgiou, E. and Finlay, J. C.: Contribution of wetlands  
 1068 to nitrate removal at the watershed scale, *Nat. Geosci.*, 11, 127–132, doi:10.1038/s41561-  
 1069 017-0056-6, 2018.

1070 Harder, P., Helgason, W. D. and Pomeroy, J. W.: Modeling the Snowpack Energy Balance  
 1071 during Melt under Exposed Crop Stubble, *J. Hydrometeorol.*, 19, 1191–1214,  
 1072 doi:10.1175/JHM-D-18-0039.1, 2018.

1073 ~~Hayashi, M. and Rosenberry, O. D.: Effects of ground water exchange on the hydrology and~~  
 1074 ~~ecology of surface water, *Groundwater*, 40, 309–316, 2002.~~

1075 Hijmans, R. J.: raster: Geographic data analysis and modeling, R package version 2.6-7,  
 1076 <https://CRAN.R-project.org/package=raster>, 2017.

1077 Hosking, J.R. and Wallis, J.R.: Parameter and quantile estimation for the generalized Pareto  
 1078 distribution. *Technometrics*, 29, 3, 339-349, 1987.

1079 Husson, F., Josse, J., Lê, S., and Mazet, J.: FactoMineR: Multivariate Exploratory Data Analysis  
 1080 and Data Mining with R. R package version 1.12. Available from: <http://factominer.free>,  
 1081 2009

1082 Jones, N. E., Schmidt, B. J., Melles, S. J. and Brickman, D.: Characteristics and distribution of  
 1083 natural flow regimes in Canada: a habitat template approach, *Can. J. Fish. Aquat. Sci.*, 71,  
 1084 1616–1624, doi:10.1139/cjfas-2014-0040, 2014.

1085 Kanishka, G., and Eldho, T.I.: Watershed classification using isomap technique and  
1086 hydrometeorological attributes, *J. Hydrol. Eng.*, 22, 04017040, doi:  
1087 10.1061/(ASCE)HE.1943-5584.0001562, 2017.

1088 Knoben, W. J. M., Woods, R. A., & Freer, J. E.: A quantitative hydrological climate  
1089 classification evaluated with independent streamflow data, *Water Resources Research*, 54,  
1090 5088–5109, <https://doi.org/10.1029/2018WR022913>, 2018.

1091 Kerr, J. G.: Multiple land use activities drive riverine salinization in a large, semi-arid river basin  
1092 in western Canada, *Limnol. Oceanogr.*, 62, 1331–1345, doi:10.1002/lno.10498, 2017.

1093 Lê, S., Josse, J., and Husson, F.: FactoMineR: An R Package for Multivariate Analysis, *Journal*  
1094 *of Statistical Software*, 25, 1-18, 2008

1095 Lehner, B., and Grill, G.: Global river hydrography and network routing: baseline data and new  
1096 approaches to study the world’s large river systems, *Hydrol. Process.*, 27, 2171–2186. Data  
1097 is available at [www.hydrosheds.org](http://www.hydrosheds.org), 2013

1098 Leibowitz, S. G., Mushet, D. M. and Newton, W. E.: Intermittent Surface Water Connectivity:  
1099 Fill and Spill Vs. Fill and Merge Dynamics, *Wetlands*, 36, 323–342, doi:10.1007/s13157-  
1100 016-0830-z, 2016.

1101 Leibowitz, S. G., Wigington, P. J., Schofield, K. A., Alexander, L. C., Vanderhoof, M. K. and  
1102 Golden, H. E.: Connectivity of Streams and Wetlands to Downstream Waters: An Integrated  
1103 Systems Framework, *J. Am. Water Resour. Assoc.*, 54, 298–322, doi:10.1111/1752-  
1104 1688.12631, 2018.

1105 Liu, G. and Schwartz, F. W.: Climate-driven variability in lake and wetland distribution across  
1106 the Prairie Pothole Region: From modern observations to long-term reconstructions with  
1107 space-for-time substitution, *Water Resour. Res.*, 48, 1–11, doi:10.1029/2011WR011539,  
1108 2012.

1109 Loveland, T. R. and Merchant, J. M.: Ecoregions and Ecoregionalization: Geographical and  
1110 Ecological Perspectives, *Environ. Manage.*, 34(S1), S1–S13, doi:10.1007/s00267-003-5181-  
1111 x, 2004.

1112 MacCulloch, G. and Whitfield, P. H. H.: Towards a Stream Classification System for the  
1113 Canadian Prairie Provinces, *Can. Water Resour. J.*, 37, 311–332, doi:10.4296/cwrj2011-905,  
1114 2012.

1115 Main, A. R., Headley, J. V., Peru, K. M., Michel, N. L., Cessna, A. J. and Morrissey, C. A.:  
1116 Widespread use and frequent detection of neonicotinoid insecticides in wetlands of Canada’s  
1117 prairie pothole region, *PLoS One*, 9, doi:10.1371/journal.pone.0092821, 2014.

1118 Marton, J. M., Creed, I. F., Lewis, D. B., Lane, C. R., Basu, N. B., Cohen, M. J. and Craft, C. B.:  
1119 Geographically isolated wetlands are important biogeochemical reactors on the landscape,  
1120 *Bioscience*, 65, 408–418, doi:10.1093/biosci/biv009, 2015.

1121 Matile, G.L.D., and Keller, G.R.: Surficial geology of the Norway House map sheet (NTS 63H),  
1122 Manitoba. Surficial Geology Compilation Map Series SG-63H, scale 1:250 000. Manitoba  
1123 Science, Technology, Energy and Mines, Manitoba Geological Survey. Available from:  
1124 <https://www.gov.mb.ca/iem/info/libmin/SG-63H.zip>, 2006.

1125 McDonnell, J. J. and Woods, R.: On the need for catchment classification, *J. Hydrol.*, 299, 2–3,  
1126 doi:10.1016/j.jhydrol.2004.09.003, 2004.

1127 Mekis, É. and Vincent, L. A.: An overview of the second generation adjusted daily precipitation  
1128 dataset for trend analysis in Canada, *Atmos. - Ocean*, 49(2), 163–177,  
1129 doi:10.1080/07055900.2011.583910, 2011.

1130 Mowchenko M, Meid PO.: The Determination of Gross and Effective Drainage Areas in the  
1131 Prairie Provinces, Regina SK (ed). *Agriculture - Prairie Farm Rehabilitation - Engineering*  
1132 Branch: Canada, 22, 1983.

1133 Mwale, D., Gan, T. Y., Devito, K. J., Silins, U., Mendoza, C. and Petrone, R.: Regionalization of  
1134 Runoff Variability of Alberta, Canada, by Wavelet, Independent Component, Empirical  
1135 Orthogonal Function, and Geographical Information System Analyses, *J. Hydrol. Eng.*, 16,  
1136 93–107, doi:10.1061/(ASCE)HE.1943-5584.0000284, 2011.

1137 NRC: Hydro features (1:50000). National Hydro Network, CanVec series. Earth Sciences Sector,  
1138 Natural Resources Canada. Government of Canada. Available from:  
1139 <http://open.canada.ca/en>, 2016.

1140 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R.,  
1141 O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E. and Wagner, H.:  
1142 vegan: Community Ecology Package. R package version 2.5-2. [https://CRAN.R-](https://CRAN.R-project.org/package=vegan)  
1143 [project.org/package=vegan](https://CRAN.R-project.org/package=vegan), 2018.

1144 Omernik, J. M. and Griffith, G. E.: Ecoregions of the Conterminous United States: Evolution of  
1145 a Hierarchical Spatial Framework, *Environ. Manage.*, 54(6), 1249–1266,  
1146 doi:10.1007/s00267-014-0364-1, 2014.

1147 Ouarda, T. B., Hache, M., Bruneau, P. and Bobee, B.: Regional Flood Peak and Volume  
1148 Estimation in Northern Canadian Basin, 14, 176–191, 2002.

1149 Pekel, J.-F., Cottam, A., Gorelick, N. and Belward, A. S.: High-resolution mapping of global  
1150 surface water and its long-term changes, *Nature*, 540, 418–422, doi:10.1038/nature20584,  
1151 2016.

1152 Peters, D.L., Boon, S., Huxter, E.H., Spence, C., van Meerveld, H.J., and Whitfield, P.H.: Zero  
1153 Flow: A PUB (Predic- tion in Ungauged Basins) Workshop on Temporary Streams:  
1154 Summary of workshop discussions and future directions, *Canadian Water Resources Journal*  
1155 37, 425–431, 2012.

1156 Rao, A. R. and Srinivas, V. V.: Regionalization of watersheds by fuzzy cluster analysis, *J.*  
1157 *Hydrol.*, 318, 57–79, doi:10.1016/j.jhydrol.2005.06.004, 2006.

1158 Razavi, T. and Coulibaly, P.: Classification of Ontario watersheds based on physical attributes  
1159 and streamflow series, *J. Hydrol.*, 493, 81–94, doi:10.1016/j.jhydrol.2013.04.013, 2013.

1160 Razavi, T. and Coulibaly, P.: An evaluation of regionalization and watershed classification  
1161 schemes for continuous daily streamflow prediction in ungauged watersheds, *Can. Water*  
1162 *Resour. J.*, 42, 2–20, doi:10.1080/07011784.2016.1184590, 2017.

1163 R Core Team. R: A language and environment for statistical computing. R Foundation for  
1164 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>, 2018.

1165 Ribatet, M.: SpatialExtremes: Modelling Spatial Extremes. R package version 2.0-7.  
1166 <https://CRAN.R-project.org/package=SpatialExtremes>, 2018.

1167 Sawicz, K., Wagener, T., Sivapalan, M., Troch, P. A. and Carrillo, G.: Catchment classification:  
1168 Empirical analysis of hydrologic similarity based on catchment function in the eastern USA,  
1169 *Hydrol. Earth Syst. Sci.*, 15, 2895–2911, doi:10.5194/hess-15-2895-2011, 2011.

1170 Seekell, D. A. and Pace, M. L.: Does the Pareto distribution adequately describe the size-  
 1171 distribution of lakes?, *Limnol. Oceanogr.*, 56, 350–356, doi:10.4319/lo.2011.56.1.0350,  
 1172 2011.

1173 Shaw, D. A., van der kamp, G., Conly, F. M., Pietroniro, A. and Martz, L.: The Fill-Spill  
 1174 Hydrology of Prairie Wetland Complexes during Drought and Deluge, *Hydrol. Process.*, 26,  
 1175 3147–3156, doi:10.1002/hyp.8390, 2012.

1176 Shook, K. and Pomeroy, J.: Changes in the hydrological character of rainfall on the Canadian  
 1177 prairies, *Hydrol. Process.*, 26, 1752–1766, doi:10.1002/hyp.9383, 2012.

1178 Shook, K., Pomeroy, J. W., Spence, C. and Boychuk, L.: Storage dynamics simulations in prairie  
 1179 wetland hydrology models: Evaluation and parameterization, *Hydrol. Process.*, 27, 1875–  
 1180 1889, doi:10.1002/hyp.9867, 2013.

1181 Shook, K., Pomeroy, J. and van der Kamp, G.: The transformation of frequency distributions of  
 1182 winter precipitation to spring streamflow probabilities in cold regions; case studies from the  
 1183 Canadian Prairies, *J. Hydrol.*, 521, 394–409, doi:10.1016/j.jhydrol.2014.12.014, 2015.

1184 Shook, K. R. and Pomeroy, J. W.: Memory effects of depression storage in Northern Prairie  
 1185 hydrology, *Hydrol. Process.*, 25, 3890–3898, doi:10.1002/hyp.8381, 2011.

1186 Simpson, M.A.: Surficial Geology Map of Saskatchewan (250k\_surficial, vector digital data).  
 1187 Original maps published 1984 to 1988; merged digital version made available 2008,  
 1188 compiled by M.A. Simpson, Environment Branch, Saskatchewan Research Council and  
 1189 Saskatchewan Ministry of the Economy. Available from:  
 1190 <https://gisappl.saskatchewan.ca/Html5Ext/index.html?viewer=GeoAtlas>, 2008.

1191 Sivakumar, B., Singh, V. P., Berndtsson, R. and Khan, S. K.: Catchment Classification  
 1192 Framework in Hydrology: Challenges and Directions, *J. Hydrol. Eng.*, 20,  
 1193 130426211354007, doi:10.1061/(ASCE)HE.1943-5584.0000837, 2013.

1194 Spence, C. and Saso, P.: A hydrological neighbourhood approach to predicting streamflow in the  
 1195 Mackenzie valley, *Predict. Ungauged Basins Approaches Canada's Cold Reg.*, 21–44, 2005.

1196 Statistics Canada: Table 32-10-0162-01. Selected land management practices and tillage  
 1197 practices used to prepare land for seeding, historical data. Census of Agriculture. Statistics  
 1198 Canada, Government of Canada. Available from:  
 1199 <https://www150.statcan.gc.ca/t1/tbl1/en/cv.action?pid=3210016201>, 2016.

1200 Ulrich, A. E., Malley, D. F. and Watts, P. D.: Lake Winnipeg Basin: Advocacy, challenges and  
 1201 progress for sustainable phosphorus and eutrophication control, *Sci. Total Environ.*, 542,  
 1202 1030–1039, doi:10.1016/j.scitotenv.2015.09.106, 2016.

1203 Van der Kamp, G. and Hayashi, M.: Groundwater-wetland ecosystem interaction in the semiarid  
 1204 glaciated plains of North America, *Hydrogeol. J.*, 17, 203–214, doi:10.1007/s10040-008-  
 1205 0367-1, 2009.

1206 Van der Kamp, G., Hayashi, M. and Gallén, D.: Comparing the hydrology of grassed and  
 1207 cultivated catchments in the semi-arid Canadian prairies, *Hydrol. Process.*, 17, 559–575,  
 1208 doi:10.1002/hyp.1157, 2003.

1209 Van der Kamp, G., Hayashi, M., Bedard-Haughn, A. and Pennock, D.: Prairie Pothole Wetlands  
 1210 – Suggestions for Practical and Objective Definitions and Terminology, *Wetlands*, 36, 229–  
 1211 235, doi:10.1007/s13157-016-0809-9, 2016.

1212 Vanderhoof, M. K., Christensen, J. R. and Alexander, L. C.: Patterns and drivers for wetland  
1213 connections in the Prairie Pothole Region, United States, *Wetl. Ecol. Manag.*, 25, 275–297,  
1214 doi:10.1007/s11273-016-9516-9, 2017.

1215 Van Meter, K. J. and Basu, N. B.: Signatures of human impact: size distributions and spatial  
1216 organization of wetlands in the Prairie Pothole landscape, *Ecol. Appl.*, 25, 451–465,  
1217 doi:10.1890/14-0662.1, 2015.

1218 Vincent, L. A., Zhang, X., Brown, R. D., Feng, Y., Mekis, E., Milewska, E. J., Wan, H. and  
1219 Wang, X. L.: Observed trends in Canada’s climate and influence of low-frequency  
1220 variability modes, *J. Clim.*, 28, 4545–4560, doi:10.1175/JCLI-D-14-00697.1, 2015.

1221 Vicente-Serrano S.M., Beguería, S., and López-Moreno, J.I.: A Multi-scalar drought index  
1222 sensitive to global warming: The Standardized Precipitation Evapotranspiration Index –  
1223 SPEI, *J. Clim.*, 23, 1696–1718, doi: 10.1175/2009JCLI2909.1, 2010.

1224 Wagener, T., Sivapalan, M., Troch, P., and Woods, R.: Catchment classification and hydrologic  
1225 similarity, *Geogr. Compass*, 1, 901–931, doi: 10.1111/j.1749-8198.2007.00039.x, 2007.

1226 Wagner, T., Bremigan, M. T., Cheruvilil, K. S., Soranno, P. A., Nate, N. A. and Breck, J. E.: A  
1227 multilevel modeling approach to assessing regional and local landscape features for lake  
1228 classification and assessment of fish growth rates, *Environ. Monit. Assess.*, 130, 437–454,  
1229 doi:10.1007/s10661-006-9434-z, 2007.

1230 Weber, D., Sadeghian, A., Luo, B., Waiser, M. J. and Lindenschmidt, K. E.: Modelling  
1231 Scenarios to Estimate the Potential Impact of Hydrological Standards on Nutrient Retention  
1232 in the Tobacco Creek Watershed, Manitoba, Canada, *Water Resour. Manag.*, 31, 1305–  
1233 1321, doi:10.1007/s11269-017-1578-9, 2017.

1234

1235 **Tables-TABLES AND FIGURES and Figures**

1236 **Table 1** – Pre-processing of compositional data PCA results. Shown are the respective subsets,  
 1237 the number of initial fractional area variables before dimensional reduction, the number of  
 1238 principal components retains to reach over 80% of subset variation (except for tillage practice),  
 1239 and the proportion of variation explained by each component.

Variable subset	Number of initial variables	Number of principal components	Total variation explained by component
Surficial geology	6	2	1: 72.8% 2: 14.4%
Particle size class	3	2	1: 74.8% 2: 15.6%
Soil zone	5	2	1: 54.6% 2: 42.7%
Local surface form	5	3	1: 54.5% 2: 24.2% 3: 11.9%
Land cover	5	3	1: 36.8% 2: 25.2% 3: 20.6%
Tillage practice	3	2	1: 90.9% 2: 8.5%

1240

1241 **Table 2** – Canonical correlation coefficients for watershed attribute and hydrological variables of  
 1242 hydrological research stations from the canonical correlation analysis. Those variables used in  
 1243 multiple regression equations are denoted with a ‘\*’.

	Correlation	
	V1	V2
<b>Watershed attributes</b>		
Area*	0.36	-0.83
DSF*	-0.26	0.90
Fraction rock*	-0.64	0.61
Fraction natural area*	-0.26	0.71
Stream density	-0.27	0.37
Mean annual precipitation	-0.14	-0.30
Fraction water area	0.53	-0.19
<b>Hydrological variables</b>	<b>W1</b>	<b>W2</b>
Q2	-0.82	-0.58
Q100	-0.22	-0.98
Canonical $\lambda$	0.97	0.77

1244

1245

1246 **Table 3** – Correlation of study watershed attributes to principal components (PC). The values for  
 1247 the six PCs used in the cluster analysis are shown.

Variable	Abbreviation	PC1	PC2	PC3	PC4	PC5	PC6
Mean elevation	elevation	0.81	0.34	-0.14	0.09	-0.16	-0.17
Mean slope	slope.mean	0.61	-0.23	0.06	0.37	-0.10	0.11
Slope CV	slope.CV	0.30	-0.38	0.22	0.14	-0.41	-0.09
Total precipitation	precip	-0.85	-0.12	0.13	0.16	-0.07	0.30
Potential evapotranspiration	PET	0.31	-0.33	-0.33	-0.06	0.47	0.13
Non-effective area	NE.area	0.02	0.70	0.31	0.10	0.01	-0.15
Areal fraction below outlet ( $A_{BO}$ )	A_BO	0.14	0.25	0.27	-0.17	0.42	-0.01
Stream density	stream.density	0.08	-0.42	-0.39	0.03	-0.41	0.08
Wetland density	wetland.density	-0.63	0.46	0.11	-0.04	0.12	0.24
Wetland fraction	wetland.frac	-0.30	0.19	0.66	-0.36	0.02	0.11
Water area in largest wetland to total in watershed ( $W_L$ )	W_L	0.31	-0.44	0.51	-0.32	-0.06	-0.12
Location of largest wetland to outlet ( $L_W/L_O$ )	L_W/L_O	-0.01	-0.06	-0.22	0.09	-0.07	-0.07
Beta ( $\beta$ )	beta	0.17	0.49	-0.02	0.01	0.09	0.05
Xi ( $\xi$ )	xi	0.21	-0.23	0.57	-0.31	-0.10	-0.17
Runoff ( $Q_2$ )	Q2	-0.13	0.35	-0.47	0.00	-0.33	0.10
Soil texture PC1	Text.PC1	-0.07	-0.04	0.28	0.55	0.19	-0.32
Soil texture PC2	Text.PC2	0.02	-0.32	0.43	0.03	-0.31	0.54
Soil zone PC1	Soil.PC1	-0.65	-0.29	-0.07	0.19	-0.10	-0.24
Soil zone PC2	Soil.PC2	0.27	-0.12	-0.06	-0.11	0.40	0.25
Land cover PC1	LC.PC1	-0.44	0.38	-0.21	-0.26	-0.43	0.12
Land cover PC2	LC.PC2	0.42	0.22	-0.17	-0.53	0.15	0.03
Land cover PC3	LC.PC3	0.21	0.30	0.15	0.25	0.11	0.46
Surficial geology PC1	SF.PC1	0.06	0.21	-0.19	0.50	0.17	-0.09
Surficial geology PC2	SF.PC2	0.06	-0.38	0.24	0.47	0.11	-0.03
Surface form PC1	LL.PC1	-0.16	0.20	0.17	0.47	0.26	0.26
Surface form PC2	LL.PC2	-0.20	0.44	0.12	-0.03	0.04	-0.55
Surface form PC3	LL.PC3	0.41	0.38	0.20	0.21	-0.27	0.27
Land practice PC1	LP.PC1	-0.54	-0.58	-0.13	-0.10	0.32	-0.09
Land practice PC2	LP.PC2	0.14	-0.16	-0.24	-0.22	0.29	0.30
<i>Supplementary variables</i>							
Latitude	Lat	-0.15	0.24	0.26	-0.01	-0.33	-0.41
Longitude	Long	-0.73	-0.24	0.06	0.10	0.16	0.39
Area	Area	-0.05	0.27	-0.44	0.09	-0.15	-0.03
DSF	DSF	-0.02	-0.25	0.42	-0.05	0.12	0.01

1248

1249 **Table 4** – Classes and distinguishing variables of prairie watersheds. The v-test statistics, based  
 1250 on Ward’s criterion, are shown. Variables with v-test values greater or less than 10 and –10,  
 1251 respectively, are bolded to emphasize defining features of each class. All variables are significant  
 1252 to  $p < 0.001$ . *Classes: Southern Manitoba (1), Pothole Till (2), Pothole Glaciolacustrine (3),*  
 1253 *Major River Valleys (4), Interior Grasslands (5), High Elevation Grasslands (6), Sloped Incised*  
 1254 *(7).*

Class 1 (n=365)		Class 2 (n=879)		Class 3 (n=681)		Class 4 (n=536)	
Variable	v-test	Variable	v-test	Variable	v-test	Variable	v-test
<b>LP.PC1</b>	<b>48.11</b>	<b>wetland.density</b>	<b>28.23</b>	<b>LC.PC1</b>	<b>22.60</b>	<b>SF.PC2</b>	<b>19.83</b>
<b>precip</b>	<b>30.33</b>	<b>LL.PC1</b>	<b>24.81</b>	<b>wetland.frac</b>	<b>12.74</b>	<b>slope.CV</b>	<b>19.35</b>
<b>Soil.PC1</b>	<b>23.60</b>	<b>precip</b>	<b>22.74</b>	<b>Q2</b>	<b>12.63</b>	<b>xi</b>	<b>16.05</b>
<b>LP.PC2</b>	<b>14.74</b>	<b>SF.PC1</b>	<b>21.74</b>	<b>NE.area</b>	<b>11.12</b>	<b>W_L</b>	<b>15.39</b>
<b>PET</b>	<b>13.10</b>	<b>LC.PC1</b>	<b>17.19</b>	LL.PC2	9.45	<b>Text.PC2</b>	<b>15.07</b>
wetland.density	7.39	<b>LL.PC2</b>	<b>16.42</b>	wetland.density	8.05	<b>Text.PC1</b>	<b>14.40</b>
DSF	6.81	<b>Q2</b>	<b>15.77</b>	LC.PC2	6.70	<b>Soil.PC1</b>	<b>14.01</b>
SF.PC2	6.53	<b>Soil.PC1</b>	<b>15.76</b>	LL.PC3	6.53	<b>DSF</b>	<b>11.76</b>
stream.density	4.61	<b>NE.area</b>	<b>15.72</b>	xi	5.89	<b>precip</b>	<b>10.97</b>
LC.PC1	-3.37	<b>area</b>	<b>13.15</b>	W_L	4.58	<b>wetland.frac</b>	<b>10.92</b>
A_BO	-4.22	<b>Text.PC1</b>	<b>12.00</b>	precip	3.47	slope.mean	7.29
area	-5.46	LC.PC3	6.76	A_BO	-3.79	LP.PC1	3.52
slope.CV	-6.49	beta	5.31	slope.CV	-4.97	A_BO	-3.83
Q2	-8.47	L_W/L_O	4.20	L_W/L_O	-5.17	wetland.density	-4.41
SF.PC1	-8.90	LL.PC3	3.93	LP.PC2	-7.11	SF.PC1	-4.56
LC.PC2	-9.21	SF.PC2	-3.97	LC.PC3	-9.71	LC.PC1	-5.13
<b>LL.PC2</b>	<b>-14.18</b>	LP.PC1	-4.87	<b>LP.PC1</b>	<b>-12.38</b>	soil.PC2	-6.93
<b>slope.mean</b>	<b>-16.17</b>	stream.density	-5.92	<b>Soil.PC2</b>	<b>-13.01</b>	beta	-7.60
<b>beta</b>	<b>-16.88</b>	elevation	-7.15	<b>Text.PC1</b>	<b>-14.58</b>	elevation	-8.03
<b>LC.PC3</b>	<b>-18.13</b>	A_BO	-7.86	<b>slope.mean</b>	<b>-15.92</b>	<b>area</b>	<b>-11.04</b>
<b>NE.area</b>	<b>-28.97</b>	Text.PC2	-9.15	<b>SF.PC2</b>	<b>-17.03</b>	<b>LP.PC2</b>	<b>-11.44</b>
<b>LL.PC3</b>	<b>-36.59</b>	DSF	-9.93	<b>LL.PC1</b>	<b>-17.83</b>	<b>Q2</b>	<b>-13.27</b>
<b>elevation</b>	<b>-47.42</b>	<b>LP.PC2</b>	<b>-10.88</b>	<b>SF.PC1</b>	<b>-18.83</b>	<b>PET</b>	<b>-13.98</b>
		<b>Soil.PC2</b>	<b>-12.00</b>	<b>PET</b>	<b>-23.29</b>	<b>LC.PC2</b>	<b>-20.86</b>
		<b>PET</b>	<b>-13.15</b>				
		<b>slope.mean</b>	<b>-13.50</b>				
		<b>slope.CV</b>	<b>-16.26</b>				
		<b>LC.PC2</b>	<b>-16.29</b>				
		<b>xi</b>	<b>-21.49</b>				
		<b>W_L</b>	<b>-32.96</b>				


1255

1256 **Table 4** – (cont'd)

Class 5 (n=635)		Class 6 (n=702)		Class 7 (n=377)	
Variable	v-test	Variable	v-test	Variable	v-test
<b>A_BO</b>	<b>34.10</b>	<b>elevation</b>	<b>29.29</b>	<b>Text.PC2</b>	<b>27.65</b>
<b>LC.PC2</b>	<b>21.53</b>	<b>PET</b>	<b>20.16</b>	<b>LL.PC3</b>	<b>25.69</b>
<b>Soil.PC2</b>	<b>20.81</b>	<b>slope.CV</b>	<b>17.67</b>	<b>slope.mean</b>	<b>22.32</b>
<b>LC.PC3</b>	<b>17.44</b>	<b>slope.mean</b>	<b>16.12</b>	<b>LC.PC3</b>	<b>14.84</b>
<b>NE.area</b>	<b>16.22</b>	<b>stream.density</b>	<b>14.55</b>	<b>stream.density</b>	<b>13.82</b>
<b>beta</b>	<b>15.96</b>	<b>LC.PC2</b>	<b>14.09</b>	<b>Soil.PC2</b>	<b>13.09</b>
<b>elevation</b>	<b>13.31</b>	<b>W_L</b>	9.47	<b>elevation</b>	<b>12.42</b>
<b>PET</b>	<b>11.47</b>	<b>L_W/L_O</b>	6.80	<b>PET</b>	<b>11.47</b>
<b>LL.PC2</b>	8.11	<b>LP.PC2</b>	5.73	<b>SF.PC2</b>	6.80
<b>LP.PC2</b>	7.67	<b>area</b>	3.72	<b>LP.PC2</b>	6.39
<b>LL.PC3</b>	7.31	<b>LL.PC2</b>	3.62	<b>slope.CV</b>	5.87
<b>wetland.frac</b>	5.77	<b>LP.PC1</b>	-3.60	<b>W_L</b>	4.63
<b>LL.PC1</b>	5.50	<b>Q2</b>	-3.94	<b>precip</b>	-4.75
<b>SF.PC2</b>	-4.74	<b>DSF</b>	-4.91	<b>A_BO</b>	-5.65
<b>area</b>	-4.86	<b>A_BO</b>	-9.47	<b>LC.PC1</b>	-7.62
<b>L_W/L_O</b>	-7.11	<b>Soil.PC1</b>	<b>-10.17</b>	<b>Text.PC1</b>	-8.34
<b>Q2</b>	-9.34	<b>LL.PC3</b>	<b>-10.62</b>	<b>LP.PC1</b>	<b>-11.42</b>
<b>LP.PC1</b>	-9.96	<b>LC.PC3</b>	<b>-13.17</b>	<b>NE.area</b>	<b>-13.33</b>
<b>Text.PC2</b>	<b>-11.36</b>	<b>NE.area</b>	<b>-14.11</b>	<b>wetland.frac</b>	<b>-13.64</b>
<b>LC.PC1</b>	<b>-11.38</b>	<b>LL.PC1</b>	<b>-15.44</b>	<b>wetland.density</b>	<b>-16.27</b>
<b>slope.CV</b>	<b>-12.42</b>	<b>Text.PC2</b>	<b>-15.78</b>	<b>Soil.PC1</b>	<b>-16.43</b>
<b>precip</b>	<b>-20.86</b>	<b>LC.PC1</b>	<b>-17.15</b>	<b>LL.PC2</b>	<b>-39.41</b>
<b>Soil.PC1</b>	<b>-23.58</b>	<b>wetland.frac</b>	<b>-21.48</b>		
<b>stream.density</b>	<b>-26.34</b>	<b>wetland.density</b>	<b>-29.58</b>		
		<b>precip</b>	<b>-37.27</b>		

1257

1258




1259

1260 **Figure 1** – Map of the study area spanning the Prairies ecozone in western Canada (inset). Large  
1261 cities in each of the three provinces are shown for reference, while the region characterized as  
1262 not contributing runoff (2-year) is also shown. Prairie ecozone based on the region classified by  
1263 the Ecological Stratification Group (1995).


1264

1265



1266


1267 **Figure 2** – Observed versus predicted estimates for (a) Q2, and (b) Q100. The dashed grey line  
1268 depicts the linear regression between observed and predicted flow values, and the black, solid  
1269 line shows a 1:1 relationship.



1274 **Figure 3** – Principal components analysis for candidate variables for classification. Active and  
1275 supplementary variables are shown as solid black, and dashed blue arrows, respectively.  
1276 Eigenvalues for PC axes are provided (inset), with black bars denoting the six PCs used in the  
1277 hierarchical clustering analysis.

1278


1279



1293 **Figure 4** – Dendrogram resulting from the hierarchical cluster analysis of principal components.  
1294 The blue, dashed line indicates the cut in the tree, resulting in seven clusters. The amount of  
1295 inertia gained by increasing the number of clusters ( $k$ ) is depicted in the inset panel.

1296





1297



1298

1299 **Figure 5** – Classification of Prairie ecozone watersheds. Watershed delineations are from Lehner  
1300 and Grills (2013), available at [www.hydrosheds.org](http://www.hydrosheds.org). See text for detailed interpretation of the  
1301 seven clusters.


1302



1303



1304 **Figure 6** – Climatic variation among watershed classes. (a) Boxplots of total annual precipitation  
 1305 (grey) and potential evapotranspiration (PET) (white) for each watershed cluster. Lower, middle,  
 1306 and upper limits of boxes show the 25<sup>th</sup>, 50<sup>th</sup>, and 75<sup>th</sup> quantiles, respectively. (b) Wetland  
 1307 density to moisture deficit (Precipitation – PET). *Classes: Southern Manitoba (1), Pothole Till*  
 1308 *(2), Pothole Glaciolacustrine (3), Major River Valleys (4), Interior Grasslands (5), High*  
 1309 *Elevation Grasslands (6), Sloped Incised (7).*

1310



1311

1312 **Figure 7** – Boxplots of select variables by watershed class: (a) fraction of non-effective area; (b)  
1313 fraction of cropland; and (c) fraction of grassland. *Classes: (1) Southern Manitoba, (2) Pothole*  
1314 *Till, (3) Pothole Glaciolacustrine, (4) Major River Valleys, (5) Interior Grassland, (6) High*  
1315 *Elevation Grasslands, and (7) Sloped Incised.*




1316

1317 **Figure 8** – Wetland variables and simulated size distributions. Median (a) density of wetlands  
 1318 and (b) fraction of total watershed water area in the largest wetland (W<sub>L</sub>) are depicted by class.  
 1319 Panel (c) shows observed (dark) and simulated (light) percentiles of wetland areas. Predicted  
 1320 values are based on a generalized Pareto distribution and using median parameters of  $\beta$  and  $\zeta$   
 1321 for each cluster. Simulated data were restricted to the raster pixel resolution of observed data from  
 1322 the Global Surface Water dataset. *Classes: Southern Manitoba (1), Pothole Till (2), Pothole*  
 1323 *Glaciolacustrine (3), Major River Valleys (4), Interior Grasslands (5), High Elevation*  
 1324 *Grasslands (6), Sloped Incised (7).*

1325

1326

1327




1328

1329 **Figure 9** – Density distributions of percent agreement of watersheds to the classification in Fig.  
1330 5 by watershed class. *Classes: Southern Manitoba (1), Pothole Till (2), Pothole Glaciolacustrine*  
1331 *(3), Major River Valleys (4), Interior Grasslands (5), High Elevation Grasslands (6), Sloped*  
1332 *Incised (7).*

1333

1334

1335



1336

1337 **Figure 10** – Agreement of assigned watershed classification from the (original) complete  
1338 analysis, with class assignments from the iterative approach using re-sampling. Classes are  
1339 coloured according to that shown in Fig. 5, with those identified under a new class (C8) depicted  
1340 in black. Watersheds that were removed from the subsets analyzed are in white. *Classes:*  
1341 *Southern Manitoba (1), Pothole Till (2), Pothole Glaciolacustrine (3), Major River Valleys (4),*  
1342 *Interior Grasslands (5), High Elevation Grasslands (6), Sloped Incised (7).*