

1  
1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15

**A WATERSHED CLASSIFICATION APPROACH THAT LOOKS BEYOND  
HYDROLOGY: APPLICATION TO A SEMI-ARID, AGRICULTURAL REGION IN  
CANADA**

Jared D. Wolfe<sup>1\*</sup>, Kevin R. Shook<sup>2</sup>, Chris Spence<sup>3</sup>, Colin J. Whitfield<sup>1,4</sup>

<sup>1</sup>Global Institute for Water Security, University of Saskatchewan, Saskatoon, Saskatchewan, Canada

<sup>2</sup>Centre for Hydrology, Saskatoon, Saskatchewan, Canada

<sup>3</sup>National Hydrology Research Centre, Environment and Climate Change Canada, Saskatoon, Saskatchewan, Canada

<sup>4</sup>School of Environment and Sustainability, University of Saskatchewan, Saskatoon, Saskatchewan, Canada

\*corresponding author: [jared.wolfe@usask.ca](mailto:jared.wolfe@usask.ca)

16 **ABSTRACT**

17           Classification and clustering approaches provide a means to group watersheds according  
18 to similar attributes, functions, or behaviours, and can aid in managing natural resources.  
19 Although they are widely used, approaches based on hydrological response parameters restrict  
20 analyses to regions where well-developed hydrological records exist, and overlook factors  
21 contributing to other management concerns, including biogeochemistry and ecology. In the  
22 Canadian Prairie, hydrometric gauging is sparse and often seasonal. Moreover, large areas are  
23 endorheic and the landscape is highly modified by human activity, complicating classification  
24 based solely on hydrological parameters. We compiled climate, geological, topographical, and  
25 land cover data from the Prairie and conducted a classification of watersheds using a hierarchical  
26 clustering of principal components. Seven classes were identified based on the clustering of  
27 watersheds, including those distinguishing southern Manitoba, the pothole region, river valleys,  
28 and grasslands. Important defining variables were climate, elevation, surficial geology, wetland  
29 distribution, and land cover. In particular, three classes occur almost exclusively within regions  
30 that tend not to contribute to major river systems, and collectively encompass the majority of the  
31 study area. The gross difference in key characteristics across the classes suggests that future  
32 water management and climate change may carry with them heterogeneous sets of implications  
33 for water security across the Prairie. This emphasizes the importance of developing management  
34 strategies that target sub-regions expected to behave coherently as current human-induced  
35 changes to the landscape will affect how watersheds react to change. The study provides the first  
36 classification of watersheds within the Prairie based on climatic and biophysical attributes, with  
37 the framework used being applicable to other regions where hydrometric data are sparse. Our  
38 findings provide a foundation for addressing questions related to hydrological, biogeochemical,  
39 and ecological behaviours at a regional level, enhancing the capacity to address issues of water  
40 security.

41 **A WATERSHED CLASSIFICATION APPROACH THAT LOOKS BEYOND**  
42 **HYDROLOGY: APPLICATION TO A SEMI-ARID, AGRICULTURAL REGION IN**  
43 **CANADA**

44

45 **1. INTRODUCTION**

46

47 Watershed classification methods provide a means of grouping watersheds according to  
48 similar attributes, or behaviours, and can identify sub-regions that are expected to exhibit  
49 coherent responses. This strategy can identify how catchment characteristics are similar, or  
50 dissimilar, among groups of watersheds and thus might influence hydrologic behaviour  
51 (McDonnell and Woods, 2004). Classifying watersheds can be useful for developing predictions  
52 in ungauged basins (Peters et al., 2012), and moreover, classification can be used to inform how  
53 changes to key traits (e.g., climate and land management) may affect system function.

54 Establishing these links between watershed function and biophysical structure, including  
55 hydroclimate, is an opportunity of watershed classification (Wagener et al., 2007). Accordingly,  
56 the regionalization of hydrological response through watershed classifications has been used to  
57 inform natural resource management (Detenbeck et al., 2000; Jones et al., 2014).

58 Many different approaches to watershed classification have been employed to date,  
59 including non-linear dimension reduction techniques (Kanishka and Eldho, 2017), decision trees  
60 (Bulley et al. 2008), and independent component analysis (Mwale et al., 2011), among others.  
61 Hydrological characteristics (e.g., statistical properties of streamflow regime) are widely used to  
62 inform classification owing to their potential linkages between watershed features and  
63 hydrologic responses (Brown et al., 2014; Sivakumar et al., 2013; Spence and Saso, 2005). Other  
64 classification exercises have included a wider number of characteristics, including biophysical  
65 attributes along with streamflow response, to differentiate watershed classes (e.g., Sawicz et al.,  
66 2014; Burn, 1990). Ecoregions, which incorporate historical aspects of climate, topography, and  
67 vegetation regimes, have also served as a method of differentiation for eco-hydrological studies  
68 (Loveland and Merchant, 2004). In select cases, classification is performed independently of  
69 streamflow response factors (Knoben et al., 2018). In arid or poorly gauged regions of the world,  
70 these types of approaches to classification that are independent from or not strongly dependent  
71 on hydrological indices (streamflow response), are needed, although few such classifications

72 have been performed. The need for new approaches to watershed classification can also be true  
73 of regions undergoing strong pressures from climate change and land-use, where historical  
74 streamflow records may not reflect current behaviour, particularly if a regime shift has occurred.

75 In Canada, watershed classification has been applied in many regions (e.g., Cavadias et  
76 al., 2001; Ouarda et al., 2002; Spence and Saso, 2005). To date, most have focused on larger  
77 basins, and none have covered in detail the semi-arid Canadian Prairie, which spans nearly 5 x  
78 10<sup>5</sup> km<sup>2</sup> in western Canada, from the Rocky Mountain foothills in the west to Lake Winnipeg in  
79 the east (Fig. 1). This is despite its importance as a major food producing region of the world and  
80 one that faces numerous water security challenges (Gober and Wheeler, 2014; Spence et al.,  
81 2018). Earlier work by Durrant and Blackwell (1959) grouped large Prairie watersheds based on  
82 flood regimes. A recent classification that included the Prairie region focused on stream  
83 hydrology (e.g., MacCulloch and Whitfield, 2012) but was broader and included watersheds  
84 from mountainous and forested regions to the west and north, respectively. In the Canadian  
85 Prairie, and similar regions elsewhere, extrapolating catchment-scale field and modelling studies  
86 presents challenges. It is inherently difficult to explain or predict different responses among  
87 basins, as poorly developed stream networks with intermittent or seasonal flow do not easily lend  
88 themselves to classification methods featuring streamflow response. MacCulloch and Whitfield  
89 (2012), who found a single streamflow class across the Canadian Prairie, raised the question as  
90 to whether a single grouping is appropriate, and suggested the need to expand classifications to  
91 include a greater diversity of biological, physical and chemical properties.

92 Like many of the world's agricultural regions, the Canadian Prairie has undergone vast  
93 environmental change co-incident with the green revolution. Predominant agricultural practices  
94 have changed over the decades, and each is known to influence water cycling and storage,  
95 including tillage practices, summer fallowing, and cropping type (Awada et al., 2014; Van der  
96 Kamp et al., 2003; Shook et al., 2015). Significant warming over the last 70 years, especially in  
97 winter (Coles et al., 2017; DeBeer et al., 2016) has resulted in more rain at the expense of snow  
98 (Vincent et al., 2015), and multiple-day rainfall events have been increasing in frequency relative  
99 to shorter events in some regions (Dumanski et al., 2015; Shook and Pomeroy, 2012). These  
100 observed changes in precipitation have reduced the predictability of runoff derived from  
101 snowmelt, and add uncertainty to water management and agricultural decision-making.

102           Disentangling the relative impacts of climate and land-use changes on water quantity and  
103 quality is complex, particularly as their effects are heterogeneous across spatial extent and scale.  
104 For the Prairie and elsewhere, new approaches to classification that can distinguish sub-regional  
105 and, importantly, sub-hydrometric station variability, are needed. Further, because land  
106 management decisions in agricultural regions are intrinsically linked to system function, there is  
107 a need for classifications that can inform decision-makers at a relevant scale. Indeed, stable  
108 isotope-based investigations of runoff from small lake catchments in the Boreal Plains (north of  
109 the Prairie) emphasize the need for local-scale characterization of watershed behaviour (Gibson  
110 et al., 2010, 2016), while streamflow dynamics for the Prairie and nearby Boreal Plain are linked  
111 to local surface geology and land cover (Devito et al., 2005; Mwale et al., 2011), suggesting an  
112 opportunity for a new approach to watershed classification in the region. Another potential  
113 advantage of a more comprehensive approach is that by de-emphasizing available hydrometric  
114 observations for larger and well-studied or monitored basins and including other environmental  
115 characteristics, the risk of overlooking other functions (e.g., ecology, biogeochemistry) that may  
116 be equally important to the management of a watershed's natural resources can be reduced. A  
117 system-based watershed classification for the Prairie that avoids the prejudice of classifying only  
118 those watersheds where a reasonably robust understanding of hydrology or streamflow exists can  
119 serve as a template for other regions of the world where streamflow-based classification is not  
120 viable.

121           The objective of the present work is to develop a watershed classification system based  
122 on hydrologically and ecologically significant traits for the Canadian Prairie. In this region,  
123 assessment of localized hydrological response to change is challenged by limited spatial  
124 resolution of observed streamflow data, and higher order streamflow being unrepresentative of  
125 local response due to a poorly-developed drainage network. In establishing such an approach, we  
126 seek to advance our understanding of watershed hydrology and broader watershed behaviour  
127 within the Prairie whilst also providing a framework for similar classification exercises in other  
128 regions where streamflow-based methods are not ideal. Our approach avoids the limitations of  
129 classifying according to known hydrologic response, and increases the spatial resolution of  
130 watershed classification relative to many existing approaches. We compile physiographic  
131 characteristics, including geology, wetland distribution, and land cover, of watersheds  
132 approximately 100 km<sup>2</sup> to achieve the classification. This framework will identify those areas

133 that are climatically and geographically similar, and thus might be expected to respond in a  
134 hydrologically coherent manner to climate and land management changes. Additionally, it  
135 provides a foundation on which to base prediction of watershed hydrologic, biogeochemical and  
136 ecological responses to these stressors.

137

## 138 **2. DATA COLLECTION & COMPILATION**

139

### 140 *2.1. Region domain and description*

141 The Canadian Prairie (Prairies ecozone) spans the provinces of Alberta, Saskatchewan,  
142 and Manitoba, and is part of the Nelson Drainage Basin (Fig. 1). Climate is semi-arid, with mean  
143 annual precipitation ranging between 350 and 610 mm (1970–2000) increasing from west to east.  
144 Mean annual temperature was 1–6°C over the same period with warmer conditions towards the  
145 southwest (Mekis and Vincent, 2011; Vincent et al., 2012;  
146 [http://climate.weather.gc.ca/climate\\_normals/index\\_e.html](http://climate.weather.gc.ca/climate_normals/index_e.html)). Much of the region deglaciated  
147 during the Late Pleistocene approximately 10,000 years before present, resulting in an often  
148 hummocky landscape with numerous depressions. Combined with the dry climate, the relatively  
149 short post-glaciation history has prevented maturing of a ubiquitous drainage network, and many  
150 headwaters remain disconnected from higher order streams (Shook et al., 2015). Depressions in  
151 the hummocky landscape, and the wetlands that form within them, are important features for  
152 Prairie hydrology (Van der Kamp et al., 2016) and often facilitate groundwater recharge (i.e.,  
153 depression-focused recharge) (Van der Kamp and Hayashi, 2009). The location of wetlands and  
154 their size, relative to the watershed outlet controls hydrologic gate-keeping (e.g., Spence and  
155 Woo, 2003), and thus the potential to contribute streamflow to higher-order watersheds  
156 (Leibowitz et al., 2016; Shaw et al., 2012; Shook et al., 2013). The size distribution of wetlands  
157 within a watershed and their spatial arrangement also dictate biogeochemical function and  
158 provide habitat and foraging for biota (Evenson et al., 2018). Terrestrial vegetation is typically  
159 open grassland, with aspen parkland ecotone along the northern edges of the ecozone boundary  
160 (Ecological Stratification Working Group, 1995).

161

### 162 *2.2. Watershed boundaries*

163 The focus of this study was on those watersheds that drain a distinctively prairie  
164 landscape, with watersheds defined according to topographic delineation. Thus, we constrained  
165 our study to the Canadian Prairie ecozone ( $4.7 \times 10^5 \text{ km}^2$ ) and watersheds occurring therein.  
166 Delineations of candidate study watersheds were obtained from the HydroSHEDS global dataset  
167 (Lehner and Grill, 2013). Watershed boundaries within this dataset were based on Shuttle Radar  
168 Topographic Mission (SRTM) digital elevation model (DEM) calculated at a 15 arc-second  
169 resolution. The resolution is equivalent to for example approximately 285 m east-west and 464 m  
170 north-south at Saskatoon, SK. As with other SRTM products, the HydroSHEDs dataset may be  
171 prone to errors in regions with low relief due to elevation precision of 1 m. However, the dataset  
172 provided watershed delineations over the geographic region of interest and at a fine enough scale  
173 (i.e.,  $100 \text{ km}^2$ ), and thus, it was sufficient based on data availability for purpose of the current  
174 study.

175 Only those watersheds completely within the Canadian Prairie ecozone were extracted ( $n$   
176 = 4729) from the HydroSHEDs dataset. Those watersheds that were very large ( $>4000 \text{ km}^2$ ) or  
177 small ( $<5 \text{ km}^2$ ) were removed from analysis (see Table S1). Because HydroSHEDs includes the  
178 basins of larger water bodies, including lakes, watersheds consisting of a majority of water were  
179 removed as the study only concerns the uplands of these systems. Finally, highly urbanized areas  
180 (i.e., watersheds with cover being  $>40\%$  urban) were removed. After considering these criteria,  
181 4175 watersheds remained for use in subsequent analyses, covering a total area of  $4.2 \times 10^5 \text{ km}^2$ .  
182 Mean watershed area for this subset was  $99.8 \pm 58.7 \text{ km}^2$ .

183

### 184 *2.3. Physiographic data collection*

185 The physiographic watershed variables were assembled from Canadian provincial and  
186 federal governments and non-governmental agency datasets (see Table S2 for a full list of  
187 variables and their sources). Variables were derived from climatic, hydrologic, geological,  
188 geographic, and land cover data, and details are described briefly below. Spatial processing and  
189 statistical analyses were conducted in ArcGIS version 10.5 and R version 3.4.3 (R Core Team,  
190 2018), respectively.

191

#### 192 *2.3.1. Climate*

193 Mean annual precipitation and temperature data were derived from the Canadian Gridded  
194 Temperature and Precipitation Anomalies (CANGRD) dataset spanning (ECCC, 2017).  
195 CANGRD is the only gridded climate product available for the region that uses adjusted and  
196 homogenized station data, and was picked for this reason (Mekis and Vincent, 2011; Vincent et  
197 al., 2012). The 1970–2000 period was chosen because the number of stations with adjusted and  
198 homogenized data used to derive CANGRD significantly diminished after 2000 (Laudon et al.,  
199 2017). Mean annual values over the 30-year period were constructed from 50 km resolution  
200 gridded cells ( $n = 626$ ) within and surrounding the Prairie ecozone, and interpolated to a higher  
201 spatial resolution raster by kriging using a spherical semivariogram. Values were clipped  
202 according to the watershed boundaries, and averaged over the watersheds to obtain mean annual  
203 precipitation and temperature for each watershed. Mean annual potential evapotranspiration  
204 (PET) was derived as a measure of dryness across the region. To maintain consistency among  
205 climate data, and use the same temperature data as described above, options were limited with  
206 which to calculate PET. The Thornthwaite equation (Thornthwaite, 1948) was applied using the  
207 R package *SPEI* (Vicente-Serrano et al., 2010). A disadvantage of the Thornthwaite approach is  
208 that it calculates PET solely as a function of air temperature and latitudinal position, and it  
209 assumes a fixed correlation between temperature and radiative forcing. As such, it integrates  
210 effects of other factors directly or indirectly influencing radiation or latent heat, like advection,  
211 vegetation, and humidity. The calculation adjusts for any lag in this relationship using  
212 corrections for latitude and month; however, it likely does not represent the full annual and  
213 seasonal variability in PET across a landscape, given regional heterogeneity of the  
214 aforementioned factors. Despite the limitations, the simplicity of this method is ideal for  
215 application across the wide geographic area of interest with limited data required as input,  
216 allowing for approximation of mean annual PET for the study area.

217

### 218 2.3.2. *Wetland traits*

219 Large regions within the Canadian Prairie have been designated as being “non-effective”,  
220 where they do not contribute flow to the stream network, at least one year in two (Godwin and  
221 Martin, 1975). The location of these regions are shown in Figure 1. This definition stems from  
222 work by Agriculture and Agri-Food Canada where prairie drainage areas were divided into *gross*  
223 and *effective* drainage areas, whereby the former describes the area within a topographic divide



224 that is expected to contribute under highly wet conditions, and the latter is the area that  
225 contributes runoff during a mean annual runoff event (Mowchenko and Meid, 1983). Thus, at its  
226 simplest, the non-effective area is the difference between the gross and effective drainage area;  
227 however, the exact area contributing runoff is dynamic and the controls complex, which include  
228 antecedent storage capacity and climatic conditions (Shaw et al., 2012; Shook and Pomeroy,  
229 2015). Briefly, the “non-effective” regions are caused by the intermittent connectivity of runoff  
230 among the landscape depressions, which trap runoff, and prevent it from contributing to  
231 downstream flow when the depressions are not connected. Trapped surface water can form  
232 wetlands (hereafter, inclusively referring to water area ponded in these depressions). These  
233 depressions can store water, and are indicative of water storage of the basin. Thus the non-  
234 effective portion of a basin is an index of its lack of contribution and is an important quality  
235 when considering the hydrological dynamics of this region (Shook et al., 2012).

236 The Global Surface Water dataset (Pekel et al., 2016) provides a geographically  
237 comprehensive layer of any ~30 m x 30 m pixel that was inundated at least once between 1984  
238 and 2015, as identified from the Landsat constellation of satellites. It was assumed that the  
239 dataset was indicative of potential maximum wetland coverage, as this period spanned several  
240 wet climate periods. As such, “wetland” in this context can include some seasonal ponds (i.e.,  
241 prairie potholes) as well as larger or more permanent shallow water bodies (but see Section 2.2  
242 and Table S1). Using the R package *raster* (Hijmans, 2017), wetland variables were calculated  
243 for each study watershed, including fractional wetland area, and the number of wetlands within  
244 the watershed per unit area (i.e., wetland density ( $\text{km}^{-2}$ )). The ratio of the area of the largest  
245 wetland to total wetland area in the watershed was also used as a metric (i.e.,  $W_L$ ). Further, we  
246 used the ratio of the linear distance of the largest wetland’s centroid to the watershed outlet ( $L_W$ ),  
247 to the maximum watershed boundary distance to the outlet ( $L_O$ ) to represent a centroid fraction  
248 ( $L_W/L_O$ ; i.e., the relative location of the largest wetland to watershed outlet). The basin outlet was  
249 defined as the point of lowest elevation on the watershed boundary. Both  $W_L$  and  $L_W/L_O$  can be  
250 used to evaluate the relative importance of hydrological gate-keeping; for example, larger  
251 wetland depressions located closer to the outlet control the likelihood of the watershed  
252 contributing flow downstream and attenuating peakflow (Shook and Pomeroy, 2011; Ameli and  
253 Creed, 2019).

254 To estimate wetland size distribution, it was assumed that they followed a Generalized  
255 Pareto Distribution (GPD) defined according to (Shook et al., 2013):

256

$$F(z) = GPD(\mu, \beta, \xi) = 1 - \left[ 1 + \xi \left( \frac{z - \mu}{\beta} \right)^{-1/\xi} \right] \quad (1)$$

257

258 Where  $z$  is wetland area, and  $\mu$  is the location parameter (i.e., the minimum size for which the  
259 distribution was fitted and has units of  $m^2$ ), and the scale ( $\beta$ ) and shape ( $\xi$ ) parameters are  
260 determined for each watershed. The  $\beta$  parameter is an index of the dispersion of the distribution,  
261 similar to the standard deviation, with the same units as the data being fitted (in this case  $m^2$ ).  
262 The  $\xi$  parameter is dimensionless and governs the shape of the fitted distribution. Hosking and  
263 Wallace (1987) plot the effect of variation in the shape parameter on the GPD. The scale and  
264 shape parameters were used to quantify the size distribution of wetlands and thus to describe the  
265 wetland frequency distributions for the cluster analyses (see 3.2). Note that because the sizes of  
266 the water bodies were taken from infrequent remote-sensing measurements (i.e., the Landsat data  
267 have a minimum revisit time of 8 or 16 days), they also are biased against short-lived water  
268 bodies.

269

### 270 2.2.3. Topographical parameters

271 Topographic variables including the mean elevation, mean and coefficient of variation of  
272 slope, and stream density were also calculated for each watershed. Because of the hummocky  
273 nature of many regions in the domain, it is possible for a basin to have some fraction of its area  
274 located at an elevation below that of the outlet. As such, the fraction of area below the basin  
275 outlet ( $A_{BO}$ ) was calculated for each basin. The elevation and slope variables were based on a  
276 DEM generated from the SRTM dataset. Stream vectors were obtained from the Hydrographic  
277 features CanVec (1:50000) series available from Natural Resources Canada  
278 (<https://open.canada.ca/data/en/dataset?q=canvec&sort=&collection=fgp>). The total length of  
279 streams within a watershed was calculated, and divided by the watershed area to produce the  
280 stream density. Additionally, the dimension shape factor (DSF) was used to describe watershed  
281 shape, as it has been found important for hydrological responses in previous Canadian catchment  
282 classification exercises (Spence and Saso, 2005). The DSF ( $km^{-1}$ ) was calculated as follows:

283

$$DSF = \frac{(0.28 \cdot P)}{A} \quad (2)$$

284

285 Where  $P$  (km) and  $A$  (km<sup>2</sup>) are the watershed perimeter and area, respectively, and derived from  
286 the HydroSHEDS global dataset (Lehner and Grill, 2013).

287 Geographical parameters of surficial geology, local surface landforms, soil particle size  
288 classes (sand, silt, clay), and soil zone were included in the analysis. Surficial geology polygons  
289 were derived by compiling provincial government data sources for Alberta (Atkinson, 2017),  
290 Saskatchewan (Simpson, 2008), and Manitoba (Matile, 2006). Due to the different geological  
291 classification schemes for each province, more detailed classes were grouped to broader  
292 categories related to depositional environment and surficial materials using those from the  
293 Geological Survey of Canada (2014), which provided for comparison across provincial  
294 boundaries. Local surface form (i.e., areas categorized by slope, relief, and morphology) and soil  
295 zone data were obtained from the Soil Landscape dataset (AAFC, 2013). The soil zones in the  
296 Canadian Prairie, used in the analyses were black, dark brown, brown, gray, and dark gray. The  
297 zones incorporate characteristics of colour and organic content, which are influenced by regional  
298 climate and vegetation. Clay, silt, and sand content were collected from the Detailed Soil Survey  
299 of Canada (AAFC, 2015). Mean catchment values of surficial geology, local surface landform,  
300 soil zone, and particle size class were determined by areal weighting of soil polygons within the  
301 watershed boundaries.

302

#### 303 *2.3.4. Land cover and cropland practice*

304 Fractional areas of land-use types were derived from the Agriculture and Agri-Food  
305 Canada's 2016 Annual Crop Inventory (AAFC, 2016). These raster data define land-use and land  
306 cover. Variables used in our analysis were standardized to watershed area and included  
307 unmanaged grasslands, forests (i.e., the sum of coniferous, deciduous, and mixed forest areas),  
308 pasture, and cropland (sum of cropped land areas). Predominant cropland practice was defined  
309 according to the fractional area of tillage by agricultural region sub-division (e.g., normalized to  
310 the area prepared for seed within that division by year). Averaged areas over the years 2011 and  
311 2016 for each practice, including zero-till, conservation till (leaving crop residue on soil surface),

312 and conventional till (incorporating residues into soil) (Statistics Canada, 2016), were used to  
313 describe these activities, and normalized as a fraction of the watershed.

314

### 315 *2.3.5. Hydrological variable calculation*

316 The relatively sparse hydrometric stream gauging in the domain, and the resulting paucity  
317 of data, presents two notable challenges to hydrologic response-based watershed classification.  
318 The first is that the basin network is biased to stations on higher-order (and often exotic) streams  
319 traversing the region (i.e., larger river basins), and thus there are a limited number of  
320 hydrometric gauges on streams draining solely Prairie watersheds, particularly at the spatial  
321 resolution of our study watersheds (~100 km<sup>2</sup>). Further, only a subset of these are considered  
322 reference stations (i.e., gauging unmanaged flows). Second, in the more arid and/or cold regions  
323 of the Prairie, some of these hydrometric stations are operated only seasonally, presenting  
324 additional challenges in using these records for classification exercises (e.g., MacCulloch and  
325 Whitfield, 2012).

326 As a result, mean annual runoff (Q2) and 1:100 year flood (Q100) magnitudes were  
327 estimated for the 4175 watersheds using relationships defined from canonical correlation  
328 analysis (CCA) to correlate gauged data to multivariate climatic and physiographic data  
329 according to procedures given by Spence and Saso (2005). According to Spence and Saso  
330 (2005), expected uncertainty using these methods approached 50% but exhibited biases of less  
331 than 15% (n = 34). Hydrological stations used were those identified in MacCulloch and  
332 Whitfield (2012) and within the Prairie region (n = 11), and data were obtained from archived  
333 databases of the Water Survey of Canada ([https://wateroffice.ec.gc.ca/search/historical\\_e.html](https://wateroffice.ec.gc.ca/search/historical_e.html))  
334 between 1990–2014. We note that greater uncertainty than that reported by Spence and Saso  
335 (2005) may result when using the CCA approach with a smaller sample size. Multivariate  
336 geographic data were collected as outlined in the above sections according to the watershed  
337 boundaries for the hydrological stations. Due to the fact that many watersheds within the  
338 HydroSHEDS dataset are likely to drain internally and do not consistently connect to a higher-  
339 order stream network, these streamflow data were interpreted as “runoff”, meaning the amount  
340 of water accumulated within the watershed polygon that drains to its lowest point annually.

341 Briefly, CCA correlates the streamflow record of gauged basins to physico-climatic  
342 characteristics of watersheds by representing these variables as a reduced set of canonical

343 variables. The analysis results in two canonical variable sets: one for the physico-climatic  
344 variables (i.e., V1 and V2) and another for the hydrological variables (i.e., W1 and W2). These  
345 canonical variables are constructed from linear combinations of the variable sets such that the  
346 correlation of the canonical variables are maximized. Canonical variables plotting similarly on  
347 X-Y plots (W1-W2 and V1-V2), indicate good correlation (Spence and Saso, 2005). Where  
348 canonical correlations ( $\lambda_1, \lambda_2$ ) were above 0.75 (Cavadias et al., 2001), that set of physico-  
349 climatic variables was deemed useful for estimating hydrological variables. Those physico-  
350 climatic variables passing this threshold were included as variables in a multiple regression to  
351 develop a predictive equation for Q2. Analyses were performed using the R package *vegan*  
352 (Oksanen et al., 2018).

353

### 354 **3. DATA ANALYSIS**

355

#### 356 *3.1. Pre-processing compositional datasets*

357 Principal components analysis (PCA) was used as a pre-processing step to reduce the  
358 dimensionality associated with compositional datasets (e.g., topographical and land cover  
359 parameters) (Fig. S1). Using this approach, the principal components (PC) that could  
360 cumulatively explain 80% of the variation in a subset of compositional data were included in the  
361 subsequent cluster analysis. This procedure identified the major data patterns and aided in  
362 reducing the number of zero-weighted variables. Where necessary, variables that were not  
363 transformed into PCs were log-transformed to reduce data skewness. Variable unit ranges were  
364 also scaled during the PCA to reduce the impact of certain variables exhibiting a large range of  
365 values on the subsequent cluster analysis.

366

#### 367 *3.2. Agglomerative hierarchical clustering of principal components and watershed classification*

368 Clustering analysis was performed on the suite of physiographic variables, which  
369 included PC variables derived from pre-processing (Table S2; Table S3). Agglomerative  
370 hierarchical clustering of principal components (HCPC) was used to define clusters of  
371 watersheds using the *HCPC* function in the R package *FactoMineR* (Lê et al., 2008; Husson et  
372 al., 2009) to apply a PCA on the standardized multivariate dataset of watershed attributes and  
373 was the basis for clustering. The majority of physiographic variables were included as active

374 variables in the PCA and thus influenced the arrangements of the PCs. In contrast, watershed  
375 area, DSF, latitude, and longitude were used only as supplementary variables, and thus did not  
376 explicitly affect the clustering analysis. These variables did, however, aid in watershed class  
377 characterization and interpretation. The first set of PCs that together explained 50% of the  
378 variation in the dataset ( $n = 6$ ) was retained for agglomerative clustering. Retaining these first  
379 PCs at a threshold of 50% allowed for clearer focus on main trends in the data and reduced the  
380 impact of noise on subsequent analyses, which might occur if subsequent, less influential, PCs  
381 were retained.

382         The agglomerative hierarchical clustering was performed using the Euclidean distances  
383 (from the PCA) and Ward's criterion for aggregating clusters. Ward's criterion decomposes the  
384 total inertia of clusters into between and within-group variance, and this method dictates merging  
385 for clusters (or watersheds) such that the growth in within-group inertia is minimal (Husson et  
386 al., 2010). The total inertia is partitioned into within- and between-group inertias. Within-group  
387 inertia represented the homogeneity, or similarity, of watershed within a cluster. Consequently,  
388 watersheds located close to each other in PC-space were deemed to be similar in their attributes.  
389 Watersheds are grouped according to pairs that minimize within-group inertia (Begou et al.,  
390 2015), and are differentiated based on between-group inertia gained by adding clusters. The  
391 variables contributing to cluster characteristics were determined by v-tests (Husson et al., 2009),  
392 which assessed whether the cluster mean for a given variable was significantly ( $p < 0.05$ ) greater  
393 or smaller than the overall mean.

394

### 395 *3.3. Comparing class-specific observed and simulated wetland depression data*

396         To compare how well the GPD parameters predicted the observed wetland area  
397 distributions from the Global Surface Water (GSW) dataset, wetland size distributions were  
398 simulated for each class. Wetland area for select watershed class-specific percentiles (i.e., 25<sup>th</sup>,  
399 50<sup>th</sup>, and 75<sup>th</sup> percentiles) derived from the simulated data were then compared to the wetland  
400 areas for corresponding watershed class-specific percentiles of the observed watershed data to  
401 assess the potential usefulness of using these parameters in representing wetland size  
402 distribution.

403         For this comparison, the fitted wetland area distributions were constrained in their  
404 minimum and maximum values by the Global Surface Water dataset spatial resolution (i.e., the

405 30 m pixel size) and the median area of the largest wetland observed for each watershed class,  
406 respectively. The median area of the distribution of largest wetlands for each watershed class  
407 gave an indication of the maximum sizes of the water bodies in those watersheds, and thus  
408 provided a maximum value for simulating wetland areas using the GPD. Wetland areas were  
409 simulated using the R package *SpatialExtremes* (Ribatet, 2018).

410

### 411 *3.4. Resampling and re-classifying procedure*

412 The robustness of the HCPC procedure on characterizing Prairie watersheds was tested  
413 using additional hierarchical clustering on ten subsets of the entire set of 4175. For each  
414 iteration, ten percent of watersheds were removed from the original dataset ( $n = 4175$ ) without  
415 replacement, and the remaining watersheds ( $n = 3757$ ) were then re-analyzed according to the  
416 HCPC outlined above (Fig. S1). The number of potential classes allowed was set at seven ( $k =$   
417  $7$ ), for consistency with the complete analysis. The resulting classifications were then compared  
418 to the classification performed on the complete dataset, with the watersheds being assessed on  
419 the percentage of iterations in which they were assigned to the same class as the complete  
420 classification. The proportion of membership agreement was calculated and visualized to assess  
421 the likelihood of classing watersheds consistently.

422

## 423 **4. RESULTS**

424

### 425 *4.1. Geographical data processing*

#### 426 *4.1.1 Dimension reduction: Compositional datasets and principal components analysis*

427 Variation in geology and soil was best explained by two or three principal components  
428 (Table 1; Fig. S2). Two PCs captured over 80% of the variation in surficial geology, with PC1  
429 (proportion explained: 73%) positively relating to glacial till deposits and negatively with  
430 glaciolacustrine deposits, and PC2 (14%) positively related to riverine or erosive deposits, such  
431 as glaciofluvial, alluvial, and eolian deposits. Particle size class data were explained by the first  
432 two PCs, where PC1 (75%) was positively associated with sand and negatively associated with  
433 silt and clay, while PC2 (14%) was related negatively to silt. Positive PC1 (55%) scores defined  
434 the dominance of black soils, and PC2 (43%) described dominance of brown or dark brown soils  
435 on positive or negative scores, respectively. Three PCs described the local surface form dataset.

436 PC1 (55%) captured the change from greater portion of hummocky forms to undulating forms,  
437 and PC2 (24%) was negatively associated with higher river-incised landscape fraction. The  
438 portion of level surface form was negatively related to PC3 (12%).

439 Three PCs were needed to explain over 80% of the variation in land cover (Table 1; Fig.  
440 S2). Land cover PC1 (37%) was positively associated with higher cropland and negatively with  
441 unmanaged grassland; whereas PC2 (25%) was negatively associated with higher pasture and  
442 forest cover. PC3 was associated with greater fallow and pasture areal proportion (21%).  
443 Cropland practice was described by PC1 (90%), with zero-till practices being negatively  
444 associated to this component. Although it only explained 9%, PC2 was also retained to describe  
445 the change between conventional and conservation till practices, with the practices exhibiting a  
446 positive and negative relationship, respectively.

447

#### 448 *4.1.2 Canonical correlation analysis*

449 The canonical coefficients from the CCA were acceptably high at  $\lambda_1$  0.97 and  $\lambda_2$  0.77,  
450 respectively, indicating that the physico-climatic variables exhibited influence on the  
451 hydrological variables (Cavadias et al., 2001; Spence and Saso, 2005). Canonical correlation  
452 values between the hydrological variables and W2 were greater than those with W1 (Table 2);  
453 thus, the physico-climatic variables strongly associated to second canonical correlation (i.e., V2)  
454 were used in the multiple regressions. These variables were watershed area, DSF, areal fraction  
455 of rock, and areal fraction of natural area. Plots of observed and predicted runoff Q2 ( $R^2 = 0.45$ )  
456 and Q100 ( $R^2 = 0.48$ ) show moderate agreement at lower flow values (Fig. 2). There is a  
457 negative bias estimated between 26 and 29%, which is greater than that documented by Spence  
458 and Saso (2005) using comparable extrapolation methods, but this is not unexpected because of  
459 the smaller sample size in the current study. As Q2 and Q100 exhibited high collinearity, only  
460 Q2 was included in subsequent cluster analyses to:

461

$$\log(Q2) = 0.130*\log(A) - 0.077*\log(N) + 0.117*\log(R) - 0.141*\log(DSF) - 0.620 \quad (3)$$

462

463 Where  $A$  was the watershed area,  $N$  was the natural area fraction and the sum of grasslands and  
464 forest,  $R$  was the rock fraction area, and  $DSF$  was the dimensional shape factor of the watershed.



465 The equation was then used to calculate Q2 for each watershed included in the clustering  
466 analysis.

467

## 468 *4.2. Watershed classification*

### 469 *4.2.1. Principal component analysis*

470 In total, 29 watershed attributes, including the PCs from compositional datasets (see  
471 Table 1), were used in the clustering analysis as active variables, and four were included as  
472 supplementary (Table 3). In the pre-clustering PCA, the first six PCs explained 54.3% of data  
473 variation, and were retained for the HCPC analysis (Fig. 3). The influence of subsequent PCs  
474 declined dramatically, and eleven PCs were required to explain >80%. Variable importance in  
475 the classification was not related to the log-transformed range exhibited by that variable (data not  
476 shown), and impact was mitigated by scaling the ranges of input variables in the PCA.

477 Principal components 1 and 2 captured changes in physical, land cover, and wetland  
478 characteristics (Fig. 3). PC1 was strongly associated with physical and land cover characteristics,  
479 such as elevation, wetland density, and the land cover PCs. PC2 was strongly related to metrics  
480 characterising the hydrological landscape, including river and wetland density, non-effective area  
481 fraction, landscape surface form, and size of the largest wetland ( $W_L$ ). Subsequent PCs explained  
482 less variation and were more specialized in the variables associated with them. Generally, these  
483 PCs were associated with differences in soil zone and texture class, surficial geology, and  
484 varying surface land form. A more detailed account of associations of the variables with the PCs  
485 is provided below.

486 PC1 was positively associated with elevation, mean slope, land cover PC2, and surface  
487 form PC3, and negatively with, total annual precipitation, soil zone PC1, wetland density, land  
488 practice PC1, land cover PC1, and longitude (Table 3; Fig. 3). PC2 was associated with non-  
489 effective area fraction, wetland density,  $\beta$ , and surface form PC2, and negatively related to land  
490 practice PC1,  $W_L$ , and river density. PC3 was positively related to wetland fraction,  $W_L$ ,  $\zeta$ , soil  
491 texture PC2, and DSF. Watershed area and runoff were negatively associated with PC3.

492 Variable correlations were weaker for the remaining three PCs (Table 3). PC4 was  
493 mainly associated with soil texture PC1, surficial geology PC1 and surface landform PC1,  
494 characteristic of sandier soil areas featuring glacial till deposits and higher hummocky surface  
495 forms, as well as higher mean slope. PC4 was negatively related to land cover PC2. PC5 was

496 related positively to PET, fraction below outlet, and soil zone PC2, and negatively to land cover  
497 PC1, river density, and slope CV. Finally, PC6 was mainly associated with soil texture PC2 and  
498 land cover PC3, and negatively with surface landform PC2.

499

#### 500 *4.2.1. Agglomerative hierarchical cluster analysis*

501 Seven clusters were identified from the hierarchical cluster analysis based on the  
502 between-group inertia gained by increasing cluster number ( $k$ ). The HCPC analysis suggested  
503 three clusters resulted in the greatest reduction of within-group inertia while minimally  
504 increasing  $k$  (Fig. 4). Further increasing  $k$  refined the separation and differentiation of clusters up  
505 to seven ( $k = 7$ ). Minimal added separation was observed up to  $k = 9$ , and increasing  $k > 9$   
506 resulted in little inertia gained between clusters. Thus, seven clusters, or classes, were manually  
507 selected based on these observations (Fig. 4).

508

#### 509 *4.2.3. Class characteristics and interpretation*

510 Our methodology yields sub-regional watershed classes according to climatic,  
511 physiographic, wetland, and land cover variables. The seven classes (Fig. 5), are defined by  
512 multivariate sets of attributes (Table 4). Influential classifying variables in all classes were mean  
513 elevation, total annual precipitation, land practice, surface forms, and wetland density. Other  
514 variables influential to class differentiation included fraction of non-effective area, land cover,  
515 and soil variables. Climate and elevation gradients are likely responsible for the west to east  
516 watershed clustering pattern. Moreover, we observe strong spatial concordance among some  
517 classes (Fig. 5), which is likely due to the hierarchical nature of the analysis. For simplicity, we  
518 interpret classes based on the variables where large, significant differences in class mean versus  
519 the overall mean of the dataset were observed. The classes can be assigned as follows: Southern  
520 Manitoba (C1); a Prairie Pothole region (C2, C3); Major River Valleys (C4); and Grasslands  
521 (C5, C6, and C7).

522

#### 523 *Southern Manitoba (C1)*

524 The majority of Class 1 (C1;  $n = 365$ ) watersheds occurred in the eastern prairie south of  
525 Lake Winnipeg (Fig. 5) and thus “Southern Manitoba” is used as the class name. Distinguishing  
526 characteristics associated with this class included soil zone PC1 (predominantly black soils) and

527 cropland practice PC1 (predominantly conventional till) (Table 4). Southern Manitoba had a high  
528 incidence of glaciolacustrine and alluvial deposits, as indicated by moderately negative and  
529 positive relationships with surficial geology PC1 and PC2, respectively, and the class also had  
530 low mean elevation. Topography tended to be level, with mild slopes and strong association with  
531 land surface form PC3 (Table 4). Notably, these watersheds exhibited both high annual  
532 precipitation and PET compared to other classes, and this class was the only one to have no mean  
533 moisture deficit (i.e., precipitation – PET > 0) (Fig. 6). Southern Manitoba watersheds also  
534 exhibited smaller fractions of non-effective areas and grasslands than other classes (Fig. 7).

535

#### 536 *Prairie Potholes (C2 and C3)*

537 The Prairie Pothole group, consisting of Class 2 (C2; n = 879), or Pothole Till, and Class  
538 3 (C3; n = 681), Pothole Glaciolacustrine, represents the largest class of watersheds spatially,  
539 spanning the northern part of the Alberta prairie to the southeastern part of Saskatchewan (Fig.  
540 5). Mean annual precipitation was relatively high for the study area, contributing to a slightly  
541 negative moisture deficit (Fig. 6). These watersheds contained large fractions of non-effective  
542 area (~75%) (Fig. 7a), and they exhibited positive scores on land cover PC1 (Table 4) indicating  
543 high cropland cover (~70%), whereas unmanaged grassland cover was typically very low  
544 (<20%) (Fig. 7b-c). On average, Pothole watersheds had high wetland densities (wetlands km<sup>-2</sup>),  
545 with C2 exhibiting the greatest density of all classes (Fig. 8a).

546 Surficial geology differentiated classes C2 and C3. Overall, glacial till and hummocky  
547 landforms dominated the pothole region; however, C2 was more associated with these  
548 characteristics, scoring greater mean values on PC1 of local surface form and surficial geology.  
549 In contrast, glaciolacustrine deposits were more common in C3, and soils had a higher incidence  
550 of clay and silt, where C2 watersheds were sandier (Table 4). Although both classes contain  
551 many wetlands, C2 watersheds had the smallest values of W<sub>L</sub>, indicating lower areal water extent  
552 was contained in the largest wetland (Fig. 8b).

553

#### 554 *Major River Valleys (C4)*

555 Class 4 (C4; n = 536) watersheds were associated with river valleys, and as such, extend  
556 across the prairie region (Fig. 5) and generally coincide with major rivers (e.g., North and South  
557 Saskatchewan, Qu'Appelle) and large lakes. These watersheds had the greatest value of the

558 fraction of water area in the largest depression ( $W_L$ ) (Fig. 8b), as well as high slope CV, wetland  
559 fraction, and fractions of black soil (i.e., higher soil zone PC1 scores) (Table 4). These  
560 watersheds were also associated with soil texture PC1 and surficial geology PC2, suggestive of  
561 higher incidence of sandy riverine deposits (e.g., alluvial and glaciofluvial deposits). The Major  
562 River Valleys class tended to have large “wetland” area, which is interpreted as the area of water  
563 of these rivers.

564 Taken together, these watersheds were related to parameters typical of fluvial  
565 environments, including glaciofluvial or alluvial deposits, and sandier soils. Large values of  
566 mean and CV of slope were also typical of river valley watersheds. About half the basin area  
567 tends to be non-effective in these watersheds, compared to the much greater fractions in the  
568 pothole regions (Fig. 7a) that surround many of the Major River Valleys watersheds. Being river  
569 valleys, C4 watersheds were generally narrow and small in area. Higher DSF (i.e., narrower  
570 watersheds) and smaller areas were generally associated with lower Q2 values (Table 2). Thus,  
571 although these watersheds have a high likelihood of contributing to streamflow of major rivers,  
572 the watershed Q2 contributions were predicted to be small (Table 4).

573

#### 574 *Grasslands (C5, C6, and C7)*

575 The southwestern Canadian Prairie, which includes the majority of southern Alberta and  
576 western Saskatchewan between the South Saskatchewan River and the Cypress Hills, was  
577 occupied by classes C5, C6, and C7. These watersheds tended to have large fractions of  
578 unmanaged grasslands (negative land cover PC1) and mean elevation (Table 4). Compared to the  
579 rest of the Prairie, this sub-region tended to be arid, with a strong moisture deficit (Fig. 6). As a  
580 result, these classes exhibited relatively low wetland density (Fig. 8a).

581 Classes 5 (C5;  $n = 635$ ), Interior Grasslands, and 6 (C6;  $n = 702$ ), High-Elevation  
582 Grasslands, were characteristic of the grasslands in southeastern Alberta. These watersheds had  
583 the greatest values of mean fractional grassland area, with cropland and grassland fractions being  
584 comparable (35–40%) (Fig. 7). Distinguishing features of Interior Grasslands were greater values  
585 of the fraction of area below the basin outlet,  $A_{BO}$ , and a notably large non-effective area fraction  
586 (Fig. 7a). High scores on land cover PC2 and PC3 indicate large fractions of fallow and pasture.  
587 These watersheds also scored higher on soil zone PC2, suggesting more common occurrences of  
588 brown soils. Small magnitudes of mean slope and stream densities were observed, suggesting

589 that the wetlands within the Interior Grasslands are relatively disconnected from the drainage  
590 network. This characteristic might explain why these watersheds have relatively large wetlands  
591 (Fig. 8c). In contrast, High Elevation Grasslands were characterized by greater mean elevation  
592 and slope values, and smaller non-effective fractions (Table 4; Fig. 7). These watersheds also  
593 had greater stream densities and smaller wetland densities.

594 Class 7 (C7; n = 377), Sloped Incised, watersheds are characterized by dissected, river-  
595 incised landscapes, as indicated by positive associations with local surface form PC3 (Table 4).  
596 Like High Elevation Grasslands (C6), Sloped Incised watersheds followed the Bow, Red Deer,  
597 as well as the Milk River valleys, suggesting a similar function to those of the Major River  
598 Valleys class. Wetland density is smallest in Sloped Incised watersheds, owing to their steepness,  
599 resulting in surface water reaching stream networks rather than collecting on the landscape (Fig.  
600 8).

601

#### 602 *4.3. Predicting wetland size distributions from class parameters*

603 Simulated wetland area distributions by class were compared to observed size  
604 distributions from study watersheds to evaluate the concordance of the approximate class-  
605 specific distribution to that of the observed distributions of watersheds, collectively. The median  
606 wetland density was greatest in C2, followed by C3, C1, and C5 (Fig. 8a). The median wetland  
607 densities in C6 and C7 were less than 1 km<sup>-2</sup>. C4 had the greatest areal fraction of water in the  
608 largest wetland (W<sub>L</sub>), which was over 40% (Fig. 8b), while C2 had the smallest value at ~10%.  
609 For the rest of the classes, this value was between 28% and 34%. The simulated wetland area  
610 distributions slightly overestimated those of the observed values, especially at the 25<sup>th</sup> percentile.  
611 However, the patterns of wetland area in the quartiles was generally consistent among all classes  
612 (Fig. 8c). The area of the smallest 25% of the wetlands appears quite consistent across the  
613 classes, with more variation occurring at higher percentiles. The largest difference among classes  
614 in wetland size was in the 75<sup>th</sup> percentile, with the greatest range being in C5 and the smallest in  
615 C1.

616

#### 617 *4.4. Resampling and re-classifying procedure*

618 The HCPC and watershed classification was repeated with ten random subsets of 3757  
619 watersheds. The majority of watershed were removed from at least one iteration, with only 50

620 watersheds being removed a total of 4-6 times (Fig. S3). This resulted in ten unique watershed  
621 subsets to test clustering and agreement to the seven classes, outlined above.

622         Percent membership agreement of a watershed varied by class, with the majority of  
623 classes exhibiting high agreement even after resampling. Classes exhibiting high membership  
624 agreement were Pothole Till (C2), Interior Grasslands (C5), High Elevation Grasslands (C6), and  
625 Sloped Incised (C7), with a large proportion having more than 90% agreement with the seven  
626 classes from the complete classification (Fig. 9; Table S4). Although a large mean agreement  
627 was observed overall, a few watershed classes exhibited low agreement and inconsistent  
628 classification. Southern Manitoba (C1) exhibited a bimodal distribution, where most were  
629 generally classed as C1 over 75% of the time and a second set only ~60% agreement (Fig. 9).  
630 This was due to a new class appearing (Fig. 10). Hereafter, this class is referred to as “Eastern  
631 Manitoba”. Briefly, Eastern Manitoba was association with large fraction of conventional tillage  
632 practice (i.e., positive association with land practice PC1 and land practice PC2) and large  
633 fractional effective areas (data not shown). The Major River Valleys class was the only one that  
634 did not include a watershed that achieved 100% agreement across the ten iterations; this class  
635 exhibited a peak of total agreement at approximately 60% (Fig. 9). Where Major River Valleys  
636 watersheds were classified inconsistently, the most common alternative classification were  
637 Pothole Glaciolacustrine (C3) or secondarily High Elevation Grasslands (C6) (Fig. 10). The loss  
638 of Major River Valleys occurred for iterations when the Eastern Manitoba class (C8) became  
639 apparent.

640

## 641 **5. DISCUSSION**

642

### 643 *5.1. Classifying Prairie watersheds*

#### 644 *5.1.1. Hydrological approaches*

645         Our classification procedure grouped watersheds of approximately 100 km<sup>2</sup> into seven  
646 classes. Few studies anywhere have classified watersheds at this granularity, and our  
647 investigation gives particular attention to characteristics that influence hydrological and  
648 ecological behaviour. Many previous studies in the region spanned larger areas, and this often  
649 results in the Prairie being identified as a homogenous region due to relatively low streamflow  
650 and atypical geology and surface topography (MacCulloch and Whitfield, 2012; Mwale et al.,

2011). Our results are novel in that they characterize in greater detail, and at small watershed scales, the potential for different hydrological behaviour of watersheds within the region. The only similar example that was found in the literature was by Durrant and Blackwell (1959), whose findings parallel those of this study, but at a larger watershed scale. Durrant and Blackwell (1959) described broad regions of Saskatchewan and Manitoba based on mean annual flood, distinguishing five sub-regions including southwestern Saskatchewan, north and central Saskatchewan, and southern Manitoba near the Red River and Assiniboine River confluence. In the current study, surficial geology and land surface form strongly influenced how grasslands were separated into three classes, which reinforces the role of local topography on hydrological response, as seen elsewhere (Mwale et al., 2011). Likewise, surficial geology was particularly important for distinguishing the Pothole (Till and Glaciolacustrine) classes. Similarities to the work of Durrant and Blackwell (1959) based on streamflow in larger basins suggest that our approach, with consideration of factors important to watershed behaviour, can yield classification with relevance to hydrologic function, despite the use of few hydrologic indices in our analysis (Fig. 5). This approach holds potential for use in other regions of the world that are dry, ungauged, or feature low effective areas, and thus cannot rely on streamflow characteristics as a primary means of classification according to functional behaviour.

Our classification grouped Prairie watersheds using geological, biophysical, and hydroclimatic attributes. In their review of classification approaches, Sivakumar et al. (2013) indicate that solely using geographic data is advantageous when there are limited hydrological data; however, the relationship between physical attributes and hydrologic behaviour is not necessarily definitive in all regions. For these reasons, it was important to include traits indicative of structural hydrological connectivity, such as Q2 estimates and wetland parameters. It is important to note that while Q2 emerged as a defining feature for several of the classes, it was consistently one of many variables important for characterization of that class (Table 4), suggesting that while it provides value added, it does not stand out as a major driving factor in the classification. In particular, the immature drainage network and relatively high depressional water storage capacity make prairie hydrology relatively distinct (Jones et al., 2014; Shook et al., 2013, 2015). Notably, three classes (i.e., Pothole Till, Pothole Glaciolacustrine, and Interior Grasslands) occur almost exclusively within regions that tend not to contribute to major river systems, and collectively encompass the majority of the study area (Table 4; Fig. 5). It is

682 therefore expected that hydrological response will be very different between classes that exhibit  
683 higher hydrological connectivity (i.e., potentially lower wetland to stream densities and non-  
684 effective area fractions), such as the Major River Valleys or Sloped Incised watersheds, than  
685 those that do not, such as Pothole classes.

686

### 687 *5.1.2. Ecoregions and human impacts*

688 Ecoregions are commonly used to characterize landscapes according to geographical or  
689 ecological similarity (Omernik and Griffith, 2014). Similar to our approach, ecoregion  
690 classifications are often hierarchical in nature, allowing for differing levels of detail, spatial  
691 extent, and thus defining characteristics depending on the scale of interest (Loveland and  
692 Merchant, 2004). Ecoregion classifications used in the United States (Omernik and Griffith,  
693 2014) and Canada (Ecological Stratification Group, 1995) employ a “top-down” approach,  
694 where broad categories are partitioned into smaller, more specialized units. In contrast, our  
695 approach provides a bottom-up, agglomerative approach where similar watersheds are merged.  
696 Assumptions are inherent in either approach; however, the latter was applicable to the current  
697 study to allow for grouping of watersheds given similarities in physiographic characteristics.  
698 This approach does not limit class membership to the geographic extent of a higher level class,  
699 allowing for membership to potentially span the geographic extent of the Canadian Prairie  
700 domain (Fig. 5).

701 Despite the differing methods for distinguishing similarities (or differences),  
702 arrangements of watershed classes in some cases exhibited similar ranges to ecoregion  
703 boundaries. The boundaries of Lake Manitoba Plain and Mixed Grassland ecoregions  
704 (Ecological Working Group, 1995) correspond roughly to those of the broader Southern  
705 Manitoba (C1) and Grasslands (C5, C6, and C7) classes, respectively (Fig. S4). Mwale et al.  
706 (2011) also found that annual hydrological regimes based on data from 200 stations and physical  
707 attributes in Alberta linked closely with provincial ecoregions. Our emphasis on inclusion of  
708 hydrologically relevant characteristics, such as wetland traits and effective areas that are likely  
709 important contributors to function, has proven useful for further distinguishing among the  
710 Grassland classes as well as the Pothole classes (C2 and C3) (Fig. 5; Fig. S4). Due to the  
711 fundamental differences in effective areas and in wetland versus river dominated systems (Table  
712 4; Fig. 8), we expect different hydrological behaviour between these classes. This is an



713 advantage of the HCPC classification approach in that it allows for identifying the potential  
714 similarity at relatively fine spatial scales, and does not require similar watersheds to be  
715 physically adjacent to one another. This confers the opportunity to further investigate these  
716 systems, such as through hydrological modelling and contrasting resulting responses under  
717 climate and land-use scenarios.

718         The highly managed prairie landscape reinforces the importance of considering  
719 anthropogenic alteration in hydrological understanding. Crop rotation and the ways in which  
720 fields are managed for winter affect the accumulation and redistribution of snow (Fang et al.,  
721 2010; Harder et al., 2018; Van der Kamp et al., 2003). Spring snowmelt and consequent runoff  
722 are imperative to summer surface water availability (Dumanski et al., 2015; Shook et al., 2015),  
723 and depression-focused recharge of snowmelt into groundwater facilitates storage and mitigates  
724 flood impacts (Hayashi et al., 2016). Thus, classifying procedures in the Prairie must consider  
725 the human influence on the water cycle.

726         An example of the complexities introduced by human land management activities can be  
727 shown by the C1 (Southern Manitoba) watersheds, where the land practice variable was a strong  
728 class descriptor. Agricultural activity is high everywhere in the Prairie; however, only C1 was  
729 associated with low zero-till practices, instead favouring conventional tillage (Table 4).  
730 Manitoba has seen less coherent adoption of zero-till practices since the early 1990s in compared  
731 to Alberta and Saskatchewan, with conventional or other conservation till practices remaining  
732 common in Manitoba (reviewed in Awada et al., 2014). Sustained use of conventional tillage  
733 practice within this region may increase the risk of soil erosion, which can negatively affect  
734 downstream water bodies (Cade-Menun et al., 2013). This practice, combined with landscape  
735 modifications, such as artificial drainage networks, serve to facilitate removal of water and may  
736 contribute to concurrent nutrient export from agricultural lands (Weber et al., 2017).

737         These management practices can be viewed as a trade-off, where high numbers of  
738 wetlands and level topography can pose flood risk during wet periods as wetlands fill and merge  
739 (Leibowitz et al., 2016), inundating tracts of adjacent land. Conversely, where landscape  
740 modification to enhance water export occurs, local, field-scale flood risk may be reduced, while  
741 heightening the risk of downstream flooding. Land-use and land management are important  
742 factors in understanding the connectivity and chemical transport in prairie landscapes (Leibowitz  
743 et al., 2018). In southern Manitoba, where artificial drainage has been used to increase the area of

744 arable land, beneficial management practices in the form of agricultural reservoirs have been  
745 implemented as a means of reducing nutrient export and improving downstream water quality  
746 while also mitigating the risk of downstream flooding (Gooding and Baulch, 2017). These  
747 factors illustrate the complexities when classifying and understanding hydrological response of  
748 watershed embedded in highly managed landscapes, and underscore that necessity of considering  
749 the human influence on the water cycle in such approaches.

750

## 751 *5.2. HCPC as a clustering and classification framework*

### 752 *5.2.1. Using the HCPC approach and limitations*

753 The HCPC method provides a procedure for integrating multiple physiographic attributes  
754 and describes resulting clusters by sets of significant variables (Husson et al., 2009). As  
755 discussed above, an advantage of the method is that it groups individual watersheds based on  
756 similarities. Therefore, it lends itself well as a foundation for investigating hydrological  
757 behaviour through modelling efforts. In the case of the current study, modelling efforts can be  
758 applied at a 100 km<sup>2</sup> scale to evaluate responses to environmental changes. An additional  
759 advantage is that that one may select variables or sets of variables of interest to inform the  
760 clustering of watersheds, such as those based only on topographic parameters or those dictating  
761 local hydrology. For example, climate variables may be excluded if the goal of the classification  
762 is parameterizing a hydrological model, as these variables could instead be described by local  
763 climate forcing. The relative ease with which different sets of variables can be added to or  
764 excluded from the analysis to consider different permutations of the classification is a real  
765 strength of the approach. Although this may result in differing cluster results, assessment of how  
766 these classes change with addition or removal of certain datasets can identify the variables that  
767 control class definition as well as elucidate spatial patterning of classes.

768 There are a few considerations when using this method. First, the linear restrictions of  
769 this method are challenging when working with environmental data, which often do not conform  
770 to assumptions of normality. Non-linear PCA methods and self-organizing maps have been  
771 applied successfully to classify watersheds in Ontario and to regionalize streamflow metrics  
772 (Razavi and Coulibaly, 2013, 2017). Although these methods might be logical next steps for the  
773 current study, we chose to focus on conventional PCA due to its smaller computational cost  
774 when classifying the large number of watersheds in our study.

775           Second, the current analysis weighs all variables equally. This can bias the analysis  
776 towards attributes that exhibit greater variability, as these can overshadow other more  
777 constrained variables. For example, the location of the largest pond relative to the watershed  
778 outlet (coded as  $L_w/L_o$ ) is important to controlling local prairie hydrology and hydrological gate-  
779 keeping potential (i.e., the likelihood of releasing surface water to the next order watershed)  
780 (Shook et al., 2013, 2015) and water quality (Hansen et al., 2018). Despite its hydrological  
781 importance, this variable had little influence on the clustering procedure overall, and was only a  
782 minor descriptor in certain classes, such as C5 and C6 (Table 4).

783           The original set of watersheds in the clustering analysis can affect the final classification;  
784 however, there was a high degree of agreement between classified subsets of the original dataset,  
785 and the classification generated using the complete set of watersheds ( $n = 4175$ ) (Fig. 9). Overall,  
786 watersheds designated as part of the Pothole and Grassland classes were classified consistently,  
787 with most exhibiting over 90% agreement. Major River Valleys exhibited the weakest agreement  
788 (Fig. 9), due to the appearance of a unique (new) class consistent with the Lake Manitoba Plain  
789 ecoregion (Fig. S4) for some of the subsets. In these cases, those watersheds previously  
790 classified as Major River Valleys were re-distributed to mainly High Elevation Grasslands or  
791 Pothole classes depending on dominate watershed features (Fig. 10). Although we do not include  
792 a detailed account of the new Eastern Manitoba class that emerged during this exercise, defining  
793 characteristics included a high fraction of effective area (i.e., the most eastern portion of the  
794 Prairie in Fig. 1), low relief, and lower use of zero-till agriculture (as reviewed in Awada et al.,  
795 2014). Since this new class would not be expected to translate to notable differences in  
796 management outcomes. Moreover, previous reviews on the usefulness of ecoregion  
797 classifications agree that strict geographic boundaries are unlikely, and are instead more likely  
798 “fuzzy” (Loveland and Merchant, 2004; Omernik and Griffiths, 2014).

799           Class membership in our approach is also determinate. In reality, there can be large  
800 variability in attributes within a class (e.g., Fig. 7), and membership is determined by the  
801 collective similarity of watershed attributes. Previous studies have used fuzzy c-means and  
802 Bayesian approaches that can assign a likelihood of membership to classes (Jones et al., 2014;  
803 Rao and Srinivas, 2006; Sawicz et al., 2011). An advantage to this approach is that it allows for  
804 fuzzy boundaries between classes where a gradient of features likely exists (Loveland and  
805 Merchant, 2004). Our re-classifying analysis supports the proposition that boundaries among

806 classified regions are fuzzy and some watershed might flicker among class memberships (Fig.  
807 10). Such approaches are also un-supervised and probabilistic in nature and will eliminate the  
808 subjectivity due to the researcher pre-defining the number of classes. Future work thus should  
809 consider these fuzzy boundaries and potential for watersheds to exhibit partial membership to  
810 multiple classes.

811

### 812 *5.2.2. Data quality and availability*

813 The classes resulting from the HCPC are also ultimately dependent on the types of data  
814 included. The availability of data and its geographic coverage determined the environmental  
815 parameters included in our analyses. Ideally, a more detailed estimate of runoff for each  
816 watershed would be a valuable contribution. In the current study, we used the CCA and eleven  
817 reference stations to approximate runoff values for the clustering watersheds. Given the number  
818 of watersheds included in the analyses, the diversity of physical characteristics and potential  
819 hydrological behaviour is likely not completely represented in the small sample size of available  
820 hydrometric stations, and is a limitation of our approach. Soil moisture would be important to  
821 consider in future studies given its role in influencing vegetation community composition, PET,  
822 and over all water balance (Hayashi et al., 2003; Shook et al., 2015). Where data is available,  
823 future work should consider variables related to snow formation and melt, as well the proportion  
824 of annual precipitation as snowfall. These variables are likely influential when describing  
825 hydrological behaviour of the watersheds and classes in the current study, and other cold regions  
826 (Knoben et al., 2018; Shook and Pomeroy, 2012). Furthermore, a comprehensive wetland  
827 inventory or an index of wetland drainage activity that is comparable across the three Provinces  
828 does not currently exist. These would be valuable additions to future efforts to classify Prairie  
829 watersheds given the important role of land modification on watershed functions.

830 One consideration with the Global Surface Water dataset is that the pixel size (30 m) is  
831 quite coarse and will miss numerous smaller wetlands, underestimating the number of wetlands  
832 observed. Consequently, it is likely that the analysis omitted some ephemeral wetlands for which  
833 persistence is short and size is small. Despite their known important ecological functions  
834 (Calhoun et al., 2017; Van Meter and Basu, 2015), their size and transient nature is a challenge  
835 to their inclusion in comprehensive datasets spanning large geographic areas. This may

836 inadvertently result in the role of smaller wetlands being under-represented in our analysis, or  
837 others that rely on this dataset.

838 Use of the  $\xi$  and  $\beta$  parameters as indices of the wetland area frequency distributions were  
839 shown to estimate classes area distributions reasonably well (Fig. 8c). Although for consistency,  
840 we restricted our simulated dataset to the spatial resolution of the surface water raster, one could  
841 use these parameters to estimate the frequencies of smaller wetlands in watersheds, which would  
842 otherwise be missed by satellite measurements, assuming conformity to a Generalized Pareto  
843 Distribution (Shook et al., 2013). Our analysis supports this application as simulated wetland  
844 areas generally approximated those seen across the observed data (Fig. 8c). Nonetheless, in  
845 regions where wetland drainage has been undertaken, it is expected that wetland area distribution  
846 has been altered via preferential loss of smaller water bodies (Evenson et al., 2018; Van Meter  
847 and Basu, 2015). This is exacerbated by the fact that remote sensed satellite data tends to omit  
848 smaller, ephemeral ponds. A more robust characterization of the size and permanence of  
849 wetlands in our study watersheds would be expected to improve the current dataset and enhance  
850 the clustering and classification analyses.

851

### 852 *5.3. Management implications*

853 Classification frameworks help to define sub-regions with potentially similar  
854 characteristics or behaviours. For example, climatic zones can be delineated, specifically the dry  
855 Grassland watersheds in the southwest and the wet Potholes in the northeast and in Manitoba  
856 (Fig. 5). In some cases, this may be related to local wetland densities, with large densities  
857 observed corresponding with low moisture deficits (Fig. 6b) (Liu and Schwartz, 2012). Climate  
858 variation may divide watersheds with seemingly similar geography into differing classes, as is  
859 the case with Major River Valleys and Sloped Incised watersheds. Both sets of watersheds  
860 tended to follow river valleys, but the former exhibit greater precipitation and smaller PET  
861 (Table 4). These divisions can be used to give context to regions we might expect to behave  
862 similarly, whether hydrologically, or ecologically, based solely on physical attributes, and echoes  
863 other methods, such as ecodistricts (Ecological Stratification Working Group, 1995) to classify  
864 landscapes. For example, areas that are geologically similar may differ in terrestrial or aquatic  
865 community assemblages, which should influence how each area might be managed (Jones et al.,  
866 2014; Wagner et al., 2007). If classifications are used to inform management, the resulting

867 decisions for a given location will depend on the strength of the delineation, the scale at which  
868 management is applied, relationships among management practices and the attributes used to  
869 define that area, and the relationship of those attributes to the response variable of concern  
870 (Wagner et al., 2007).

871 This set of analyses was unique among watershed classification exercises in Canada in  
872 that it considered a suite of wetland variables. The arrangement of wetlands or landscape  
873 depressions and their size distribution define the hydrological behavior of Prairie watersheds  
874 (Shook et al., 2015; Shook and Pomeroy, 2011). The storage capacity and subsequent spilling or  
875 merging controls wetland connectivity, and thus the quantity of water available to move from  
876 one watershed to another (Leibowitz et al., 2016; Shaw et al., 2012; Shook et al., 2015). In turn,  
877 a wetland or depression's hydrological gate-keeping potential, or its likelihood to prevent  
878 connectivity to the downstream watershed, is a function of both its storage capacity and  
879 landscape position. Large wetlands near an outlet have a great gate-keeping potential, as they  
880 block much of the watershed from connecting, and it takes a great deal of water to fill them  
881 before permitting flow to the next order watershed (Shook and Pomeroy, 2011). Simulated  
882 frequency distributions of wetland areas indicate that the depressional storages of the classes are  
883 very different (Fig. 8). It may be that wetland management practices will have different  
884 influences between each pothole class, and possibly among all the classes. This has implications  
885 for managing salinizing soils (Goldhaber et al., 2014), biodiversity (Balas et al., 2012), and  
886 flooding potential (Evenson et al., 2018; Golden et al., 2017).

887 Wetland drainage and wetland consolidation change hydrological connectivity and  
888 therefore the transport of nutrients and their loading into receiving water bodies (Brown et al.,  
889 2017; Vanderhoof et al., 2017). More positive values of the moisture deficit (i.e., where  $P \geq$   
890 PET) were associated with greater wetland densities (Fig. 6b) (Liu and Schwartz, 2012), and  
891 these areas were generally associated with greater fractions of cropland, such as Pothole Till,  
892 Pothole Glaciolacustrine, and Southern Manitoba watersheds. In these regions wetland drainage  
893 is widely practiced, historically or at present, and conflict over available arable land and wetland  
894 conservation is high (Breen et al., 2018).

895 Extensive drainage in combination with agricultural activity is known to increase the risk  
896 of agricultural nutrient mobility (Kerr, 2017) from the landscape to receiving water bodies.  
897 Increased connectivity also reduces water residence time and thus tends to decrease wetland

898 nutrient retention (Marton et al., 2015). Over time, zero-till practices can promote nutrient  
899 stratification in soils, where concentrations (especially phosphorus) accumulate at the surface,  
900 which can increase nutrient loading when surface runoff is generated (Cade-Menun et al., 2013).  
901 The cropland-wetland interface might also have important implications for pesticide mobility in  
902 Pothole Till and northern Pothole Glaciolacustrine watersheds. These areas coincide with  
903 extensive use of canola, which has been linked to high application rates of neonicotinoid  
904 pesticides which are known to have high persistence in small, temporary wetlands (Main et al.,  
905 2014). Watersheds in the Pothole Till class appear to have more hummocky landscapes than the  
906 Pothole Glaciolacustrine classification and smaller, more numerous wetlands (Fig. 8). Moreover,  
907 the water area fraction occupied by the largest wetland differs between the classes. The  
908 landscape biogeochemical functionality of pothole wetlands is known to vary considerably  
909 according to pothole character (Evenson et al., 2018; Van Meter and Basu, 2015). As such, our  
910 classification may highlight contrasting biogeochemical functioning, including nutrient retention,  
911 between these classes. Thus, although water quality risks are common within the region, the  
912 classes may respond very differently to environmental and land management stresses.

913

## 914 **6. CONCLUSION**

915

916 This study provides an overview of a classification framework that can be applied in  
917 regions with limited understanding of or data describing streamflow. The HCPC procedure offers  
918 a flexible analysis to elucidate the spatial arrangement of watershed classes given a large number  
919 of units to classify and a diverse set of attributes to inform the classification. In contrast to  
920 classifications based solely on hydrological function, using physiographic data allows for  
921 classifying small basins, which are unlikely to be gauged, and confers advantages over alternate  
922 procedures that rely heavily on observations of hydrological parameters, namely statistics  
923 describing streamflow.

924 Use of the classification approach for small Canadian Prairie watersheds identified  
925 regions of similar climatic and geographic features and, potentially, of hydrological response  
926 (Fig. 5). This yielded watershed classes that consider not only drainage patterns, but also land  
927 cover, land-use, and the underlying geology. In the Prairie region, wetland variables incorporate  
928 the hydrologic gate-keeping potential of wetlands as well as parameters indicative of wetland

929 size distributions. With the classification based on a large and diverse set of attributes, a diversity  
930 of behaviours is captured. This represents a major step forward for classification of Prairie  
931 watersheds that have to-date offered only a much more homogenized depiction of watershed  
932 function in the region. The watershed classification framework presented promises to be useful  
933 in other dry or semi-arid regions, and those that are poorly gauged. Given the inclusive nature of  
934 the classification approach, which incorporates landscape controls on hydrology as well as those  
935 influencing biogeochemistry and ecology, it also provides a foundation to evaluate the efficacy  
936 of land and watershed management practices in the context of a changing climate.

937  
938  
939  
940  
941  
942  
943

#### 944 **Author contributions**

945 JDW, CJW, and CS conceived the study, and JDW collected data and performed analyses. KRS  
946 wrote code to analyze basin and wetland data. JDW wrote the manuscript with input from all co-  
947 authors.

948

#### 949 **Acknowledgements**

950 This work was pursued under the Prairie Water project and funded by the Global Water Futures  
951 program, which was supported by the Canada First Research Excellence Fund. The authors  
952 would like to thank John Pomeroy for his valuable input on the scoping and approach to the  
953 study. We would also like to thank Wouter Knoben and two anonymous reviewers for their  
954 insightful comments on the manuscript. Finally, we would like to thank the Prairie Water team  
955 and the Global Institute for Water Security for ongoing support. The authors declare that they  
956 have no conflict of interest.

957  
958



959 **REFERENCES**

- 960
- 961 AAFC: Annual Crop Inventory, Agriculture and Agri-Food Canada, Government of Canada.  
962 Available from: <https://open.canada.ca/data/en/dataset/ba2645d5-4458-414d-b196-6303ac06c1c9>, 2016.
- 963
- 964 AAFC: Detailed Soil Surveys, Agriculture and Agri-Food Canada, Government of Canada..  
965 Available from: <https://open.canada.ca/data/en/dataset/7ed13bbe-fbac-417c-a942-ea2b3add1748>, 2015.
- 966
- 967 AAFC: Soils of Canada, Derived. Soil Landscapes of Canada and Detailed Soil Surveys, version  
968 3.2. Canadian Soil Information Service, Agriculture and Agri-Food Canada, Government of  
969 Canada. Available from: <https://open.canada.ca/data/en/dataset/8f496e3f-1e54-4dbb-a501-a91eccf616b8>, 2013.
- 970
- 971 Atkinson, N., Utting, D. J., and Pawley, S. M.: Surficial geology of west-central Alberta (GIS  
972 data, polygon features); Alberta Energy Regulator, AER/AGS Digital Data 2017-0031.  
973 Available from: [http://ags.aer.ca/publications/DIG\\_2017\\_0031.html](http://ags.aer.ca/publications/DIG_2017_0031.html), 2017.
- 974 Awada, L., Lindwall, C. W. and Sonntag, B.: The development and adoption of conservation  
975 tillage systems on the Canadian Prairies, *Int. Soil Water Conserv. Res.*, 2, 47–65,  
976 doi:10.1016/S2095-6339(15)30013-7, 2014.
- 977 Balas, C. J., Euliss, N. H. and Mushet, D. M.: Influence of conservation programs on amphibians  
978 using seasonal wetlands in the prairie pothole region, *Wetlands*, 32, 333–345,  
979 doi:10.1007/s13157-012-0269-9, 2012.
- 980 Begou, J., Bazie, P. and Afouda, A.: Catchment classification: multivariate statistical analyses  
981 for physiographic similarity in the Upper Niger Basin, *J. Eng. Res. Appl.*, 5, 60–68, 2015.
- 982 Breen, S.-P. W., Loring, P. A. and Baulch, H.: When a Water Problem Is More Than a Water  
983 Problem: Fragmentation, Framing, and the Case of Agricultural Wetland Drainage, *Front.*  
984 *Environ. Sci.*, 6, 1–8, doi:10.3389/fenvs.2018.00129, 2018.
- 985 Brown, R., Zhang, Z., Comeau, L. P. and Bedard-Haughn, A.: Effects of drainage duration on  
986 mineral wetland soils in a Prairie Pothole agroecosystem, *Soil Tillage Res.*, 168, 187–197,  
987 doi:10.1016/j.still.2016.12.015, 2017.
- 988 Brown, S. C., Lester, R. E., Versace, V. L., Fawcett, J. and Laurenson, L.: Hydrologic landscape  
989 regionalisation using deductive classification and random forests, *PLoS One*, 9,  
990 doi:10.1371/journal.pone.0112856, 2014.
- 991 Bulley, H.N.N., Marx, D.B., Merchant, J.W., Holz, J.C., and Holz, A.A.: A comparison of  
992 Nebraska reservoir classes estimated from watershed-based classification models and  
993 ecoregions, *J. Environ. Informatics*, 11, 90–102, doi:10.3808/jei.200800114, 2008.
- 994 Burn, D.: Cluster analysis as applied to regional flood frequency, *J. Water Resour. Plan. Manag.*,  
995 115, 567–582, 1990.
- 996 Cade-Menun, B. J., Bell, G., Baker-Ismail, S., Fouli, Y., Hodder, K., McMartin, D. W., Perez-  
997 Valdivia, C. and Wu, K.: Nutrient loss from Saskatchewan cropland and pasture in spring  
998 snowmelt runoff, *Can. J. Soil Sci.*, 93, 445–458, doi:10.4141/cjss2012-042, 2013.
- 999 Calhoun, A. J. K., Mushet, D. M., Bell, K. P., Boix, D., Fitzsimons, J. A. and Isselin-Nondedeu,  
1000 F.: Temporary wetlands: challenges and solutions to conserving a “disappearing” ecosystem,  
1001 *Biol. Conserv.*, 211, 3–11, doi:10.1016/j.biocon.2016.11.024, 2017.

1002 Cavadias, G. S., Ouarda, T. B. M. J., Bobee, B. and Girard, C.: A canonical correlation approach  
1003 to the determination of homogeneous regions for regional flood estimation of ungauged  
1004 basins, *Hydrol. Sci. J.*, 46, 499–512, doi:10.1080/02626660109492846, 2001.

1005 Coles, A. E., McConkey, B. G. and McDonnell, J. J.: Climate change impacts on hillslope runoff  
1006 on the northern Great Plains, 1962–2013, *J. Hydrol.*, 550, 538–548,  
1007 doi:10.1016/j.jhydrol.2017.05.023, 2017.

1008 DeBeer, C. M., Wheeler, H. S., Carey, S. K. and Chun, K. P.: Recent climatic, cryospheric, and  
1009 hydrological changes over the interior of western Canada: A review and synthesis, *Hydrol.*  
1010 *Earth Syst. Sci.*, 20, 1573–1598, doi:10.5194/hess-20-1573-2016, 2016.

1011 Detenbeck, N. E., Batterman, S. L., Brady, V. J., Brazner, J. C., Snarski, V. M., Taylor, D. L.,  
1012 Thompson, J. A. and Arthur, J. W.: a Test of Watershed Classification Systems for  
1013 Ecological Risk Assessment, *Environ. Toxicol. Chem.*, 19, 1174, doi:10.1897/1551-  
1014 5028(2000)019<1174:ATOWCS>2.3.CO;2, 2000.

1015 Devito, K., Creed, I., Gan, T., Mendoza, C., Petrone, R., Silins, U. and Smerdon, B.: A  
1016 framework for broad-scale classification of hydrologic response units on the Boreal Plain: Is  
1017 topography the last thing to consider?, *Hydrol. Process.*, 19, 1705–1714,  
1018 doi:10.1002/hyp.5881, 2005.

1019 Dumanski, S., Pomeroy, J. W. and Westbrook, C. J.: Hydrological regime changes in a Canadian  
1020 Prairie basin, *Hydrol. Process.*, 29, 3893–3904, doi:10.1002/hyp.10567, 2015.

1021 Durrant, E. F. and Blackwell, S. R.: The magnitude and frequency of floods in the Canadian  
1022 Prairies, *Proc. Of Symposium No. 1, Spillway Design Floods*, sub-committee on hydrology,  
1023 National Research Council Associate Committee on Geodesy and Geophysics. The Queens  
1024 Printer, Ottawa, 1959.

1025 ECCC: Hydat series. Water Survey of Canada, Environment and Climate Change Canada.  
1026 Government of Canada, 2016. Available from: [https://ec.gc.ca/rhc-  
1027 wsc/default.asp?n=9018B5EC-1](https://ec.gc.ca/rhc-wsc/default.asp?n=9018B5EC-1), 2016.

1028 ECCC: Canadian Gridded Temperature and Precipitation Anomalies. Environment and Climate  
1029 Change Canada. Government of Canada. Accessed November 2017. Available from:  
1030 <https://open.canada.ca/data/dataset/3d4b68a5-13bc-48bb-ad10-801128aa6604>, 2017.

1031 Ecological Stratification Working Group: A National Ecological Framework for Canada.  
1032 Agriculture and Agri-Food Canada, Research Branch, Centre for Land and Biological  
1033 Resources Research and Environment Canada, State of the Environment Directorate,  
1034 Ecozone Analysis Branch, Ottawa/Hull. Report and national map at 1:7500 000 scale, 1995

1035 Evenson, G. R., Golden, H. E., Lane, C. R., McLaughlin, D. L. and D’Amico, E.: Depressional  
1036 wetlands affect watershed hydrological, biogeochemical, and ecological functions, *Ecol.*  
1037 *Appl.*, 28, 953–966, doi:10.1002/eap.1701, 2018.

1038 Fang, X., Pomeroy, J. W., Westbrook, C. J., Guo, X., Minke, A. G. and Brown, T.: Prediction of  
1039 snowmelt derived streamflow in a wetland dominated prairie basin, *Hydrol. Earth Syst. Sci.*,  
1040 14, 991–1006, doi:10.5194/hess-14-991-2010, 2010.

1041 Geological Survey of Canada: Surficial geology of Canada / Géologie des formations  
1042 superficielles du Canada. Canadian Geoscience Map, Geological Survey of Canada.  
1043 Government of Canada. 195, <https://doi.org/10.4095/295462>, 2014

1044 Gibson, J. J., Birks, S. J., Jeffries, D. S., Kumar, S., Scott, K. A., Aherne, J. and Shaw, D. P.:  
 1045 Site-specific estimates of water yield applied in regional acid sensitivity surveys across  
 1046 western Canada, *J. Limnol.*, 69, 67–76, doi:10.3274/JL10-69-S1-08, 2010.

1047 Gibson, J. J., Yi, Y. and Birks, S. J.: Isotope-based partitioning of streamflow in the oil sands  
 1048 region, northern Alberta: Towards a monitoring strategy for assessing flow sources and  
 1049 water quality controls, *J. Hydrol. Reg. Stud.*, 5, 131–148, doi:10.1016/j.ejrh.2015.12.062,  
 1050 2016.

1051 Gober, P., and Wheeler, H.S. Socio-hydrology and the science-policy interface: a case study of  
 1052 the Saskatchewan River basin. *Hydrol. Earth Syst. Sci.*, 18, 1413–1422, doi: 10.5194/hess-  
 1053 18-1413-2014, 2014.

1054 Godwin RB, Martin FRJ. Calculation of gross and effective drainage areas for the Prairie  
 1055 Provinces. In: *Canadian Hydrology Symposium - 1975 Proceedings*, 11-14 August 1975,  
 1056 Winnipeg, Manitoba. Associate Committee on Hydrology, National Research Council of  
 1057 Canada, pp. 219–223. 1975.

1058 Golden, H. E., Creed, I. F., Ali, G., Basu, N. B., Neff, B. P., Rains, M. C., McLaughlin, D. L.,  
 1059 Alexander, L. C., Ameli, A. A., Christensen, J. R., Evenson, G. R., Jones, C. N., Lane, C. R.  
 1060 and Lang, M.: Integrating geographically isolated wetlands into land management decisions,  
 1061 *Front. Ecol. Environ.*, 15, 319–327, doi:10.1002/fee.1504, 2017.

1062 Goldhaber, M. B., Mills, C. T., Morrison, J. M., Stricker, C. A., Mushet, D. M. and LaBaugh, J.  
 1063 W.: Hydrogeochemistry of prairie pothole region wetlands: Role of long-term critical zone  
 1064 processes, *Chem. Geol.*, 387, 170–183, doi:10.1016/j.chemgeo.2014.08.023, 2014.

1065 Gooding, R. M. and Baulch, H. M.: Small reservoirs as a beneficial management practice for  
 1066 nitrogen removal, *J. Environ. Qual.*, 46, 96-104, doi:10.2134/jeq2016.07.0252, 2017.

1067 Hansen, A. T., Dolph, C. L., Foufoula-Georgiou, E. and Finlay, J. C.: Contribution of wetlands  
 1068 to nitrate removal at the watershed scale, *Nat. Geosci.*, 11, 127–132, doi:10.1038/s41561-  
 1069 017-0056-6, 2018.

1070 Harder, P., Helgason, W. D. and Pomeroy, J. W.: Modeling the Snowpack Energy Balance  
 1071 during Melt under Exposed Crop Stubble, *J. Hydrometeorol.*, 19, 1191–1214,  
 1072 doi:10.1175/JHM-D-18-0039.1, 2018.

1073 Hayashi, M., Van der Kamp, G. and Rosenberry, D. O.: Hydrology of Prairie Wetlands:  
 1074 Understanding the Integrated Surface-Water and Groundwater Processes, *Wetlands*, 36, 1–  
 1075 18, doi:10.1007/s13157-016-0797-9, 2016.

1076 Hijmans, R. J.: raster: Geographic data analysis and modeling, R package version 2.6-7,  
 1077 <https://CRAN.R-project.org/package=raster>, 2017.

1078 Hosking, J.R. and Wallis, J.R.: Parameter and quantile estimation for the generalized Pareto  
 1079 distribution. *Technometrics*, 29, 3, 339-349, 1987.

1080 Husson, F., Josse, J., Lê, S., and Mazet, J.: FactoMineR: Multivariate Exploratory Data Analysis  
 1081 and Data Mining with R. R package version 1.12. Available from: <http://factominer.free>,  
 1082 2009

1083 Jones, N. E., Schmidt, B. J., Melles, S. J. and Brickman, D.: Characteristics and distribution of  
 1084 natural flow regimes in Canada: a habitat template approach, *Can. J. Fish. Aquat. Sci.*, 71,  
 1085 1616–1624, doi:10.1139/cjfas-2014-0040, 2014.

1086 Kanishka, G., and Eldho, T.I.: Watershed classification using isomap technique and  
1087 hydrometeorological attributes, *J. Hydrol. Eng.*, 22, 04017040, doi:  
1088 10.1061/(ASCE)HE.1943-5584.0001562, 2017.

1089 Knobon, W. J. M., Woods, R. A., & Freer, J. E.: A quantitative hydrological climate  
1090 classification evaluated with independent streamflow data, *Water Resources Research*, 54,  
1091 5088–5109, <https://doi.org/10.1029/2018WR022913>, 2018.

1092 Kerr, J. G.: Multiple land use activities drive riverine salinization in a large, semi-arid river basin  
1093 in western Canada, *Limnol. Oceanogr.*, 62, 1331–1345, doi:10.1002/lno.10498, 2017.

1094 Lê, S., Josse, J., and Husson, F.: FactoMineR: An R Package for Multivariate Analysis, *Journal*  
1095 *of Statistical Software*, 25, 1-18, 2008

1096 Lehner, B., and Grill, G.: Global river hydrography and network routing: baseline data and new  
1097 approaches to study the world’s large river systems, *Hydrol. Process.*, 27, 2171–2186. Data  
1098 is available at [www.hydrosheds.org](http://www.hydrosheds.org), 2013

1099 Leibowitz, S. G., Mushet, D. M. and Newton, W. E.: Intermittent Surface Water Connectivity:  
1100 Fill and Spill Vs. Fill and Merge Dynamics, *Wetlands*, 36, 323–342, doi:10.1007/s13157-  
1101 016-0830-z, 2016.

1102 Leibowitz, S. G., Wigington, P. J., Schofield, K. A., Alexander, L. C., Vanderhoof, M. K. and  
1103 Golden, H. E.: Connectivity of Streams and Wetlands to Downstream Waters: An Integrated  
1104 Systems Framework, *J. Am. Water Resour. Assoc.*, 54, 298–322, doi:10.1111/1752-  
1105 1688.12631, 2018.

1106 Liu, G. and Schwartz, F. W.: Climate-driven variability in lake and wetland distribution across  
1107 the Prairie Pothole Region: From modern observations to long-term reconstructions with  
1108 space-for-time substitution, *Water Resour. Res.*, 48, 1–11, doi:10.1029/2011WR011539,  
1109 2012.

1110 Loveland, T. R. and Merchant, J. M.: Ecoregions and Ecoregionalization: Geographical and  
1111 Ecological Perspectives, *Environ. Manage.*, 34(S1), S1–S13, doi:10.1007/s00267-003-5181-  
1112 x, 2004.

1113 MacCulloch, G. and Whitfield, P. H. H.: Towards a Stream Classification System for the  
1114 Canadian Prairie Provinces, *Can. Water Resour. J.*, 37, 311–332, doi:10.4296/cwrj2011-905,  
1115 2012.

1116 Main, A. R., Headley, J. V., Peru, K. M., Michel, N. L., Cessna, A. J. and Morrissey, C. A.:  
1117 Widespread use and frequent detection of neonicotinoid insecticides in wetlands of Canada’s  
1118 prairie pothole region, *PLoS One*, 9, doi:10.1371/journal.pone.0092821, 2014.

1119 Marton, J. M., Creed, I. F., Lewis, D. B., Lane, C. R., Basu, N. B., Cohen, M. J. and Craft, C. B.:  
1120 Geographically isolated wetlands are important biogeochemical reactors on the landscape,  
1121 *Bioscience*, 65, 408–418, doi:10.1093/biosci/biv009, 2015.

1122 Matile, G.L.D., and Keller, G.R.: Surficial geology of the Norway House map sheet (NTS 63H),  
1123 Manitoba. Surficial Geology Compilation Map Series SG-63H, scale 1:250 000. Manitoba  
1124 Science, Technology, Energy and Mines, Manitoba Geological Survey. Available from:  
1125 <https://www.gov.mb.ca/iem/info/libmin/SG-63H.zip>, 2006.

1126 McDonnell, J. J. and Woods, R.: On the need for catchment classification, *J. Hydrol.*, 299, 2–3,  
1127 doi:10.1016/j.jhydrol.2004.09.003, 2004.

1128 Mekis, É. and Vincent, L. A.: An overview of the second generation adjusted daily precipitation  
1129 dataset for trend analysis in Canada, *Atmos. - Ocean*, 49(2), 163–177,  
1130 doi:10.1080/07055900.2011.583910, 2011.

1131 Mowchenko M, Meid PO.: The Determination of Gross and Effective Drainage Areas in the  
1132 Prairie Provinces, Regina SK (ed). *Agriculture - Prairie Farm Rehabilitation - Engineering*  
1133 Branch: Canada, 22, 1983.

1134 Mwale, D., Gan, T. Y., Devito, K. J., Silins, U., Mendoza, C. and Petrone, R.: Regionalization of  
1135 Runoff Variability of Alberta, Canada, by Wavelet, Independent Component, Empirical  
1136 Orthogonal Function, and Geographical Information System Analyses, *J. Hydrol. Eng.*, 16,  
1137 93–107, doi:10.1061/(ASCE)HE.1943-5584.0000284, 2011.

1138 NRC: Hydro features (1:50000). National Hydro Network, CanVec series. Earth Sciences Sector,  
1139 Natural Resources Canada. Government of Canada. Available from:  
1140 <http://open.canada.ca/en>, 2016.

1141 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R.,  
1142 O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E. and Wagner, H.:  
1143 vegan: Community Ecology Package. R package version 2.5-2. [https://CRAN.R-](https://CRAN.R-project.org/package=vegan)  
1144 [project.org/package=vegan](https://CRAN.R-project.org/package=vegan), 2018.

1145 Omernik, J. M. and Griffith, G. E.: Ecoregions of the Conterminous United States: Evolution of  
1146 a Hierarchical Spatial Framework, *Environ. Manage.*, 54(6), 1249–1266,  
1147 doi:10.1007/s00267-014-0364-1, 2014.

1148 Ouarda, T. B., Hache, M., Bruneau, P. and Bobee, B.: Regional Flood Peak and Volume  
1149 Estimation in Northern Canadian Basin, 14, 176–191, 2002.

1150 Pekel, J.-F., Cottam, A., Gorelick, N. and Belward, A. S.: High-resolution mapping of global  
1151 surface water and its long-term changes, *Nature*, 540, 418–422, doi:10.1038/nature20584,  
1152 2016.

1153 Peters, D.L., Boon, S., Huxter, E.H., Spence, C., van Meerveld, H.J., and Whitfield, P.H.: Zero  
1154 Flow: A PUB (Prediction in Ungauged Basins) Workshop on Temporary Streams:  
1155 Summary of workshop discussions and future directions, *Canadian Water Resources Journal*  
1156 37, 425–431, 2012.

1157 Rao, A. R. and Srinivas, V. V.: Regionalization of watersheds by fuzzy cluster analysis, *J.*  
1158 *Hydrol.*, 318, 57–79, doi:10.1016/j.jhydrol.2005.06.004, 2006.

1159 Razavi, T. and Coulibaly, P.: Classification of Ontario watersheds based on physical attributes  
1160 and streamflow series, *J. Hydrol.*, 493, 81–94, doi:10.1016/j.jhydrol.2013.04.013, 2013.

1161 Razavi, T. and Coulibaly, P.: An evaluation of regionalization and watershed classification  
1162 schemes for continuous daily streamflow prediction in ungauged watersheds, *Can. Water*  
1163 *Resour. J.*, 42, 2–20, doi:10.1080/07011784.2016.1184590, 2017.

1164 R Core Team. R: A language and environment for statistical computing. R Foundation for  
1165 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>, 2018.

1166 Ribatet, M.: SpatialExtremes: Modelling Spatial Extremes. R package version 2.0-7.  
1167 <https://CRAN.R-project.org/package=SpatialExtremes>. 2018.

1168 Sawicz, K., Wagener, T., Sivapalan, M., Troch, P. A. and Carrillo, G.: Catchment classification:  
1169 Empirical analysis of hydrologic similarity based on catchment function in the eastern USA,  
1170 *Hydrol. Earth Syst. Sci.*, 15, 2895–2911, doi:10.5194/hess-15-2895-2011, 2011.

1171 Seekell, D. A. and Pace, M. L.: Does the Pareto distribution adequately describe the size-  
1172 distribution of lakes?, *Limnol. Oceanogr.*, 56, 350–356, doi:10.4319/lo.2011.56.1.0350,  
1173 2011.

1174 Shaw, D. A., van der kamp, G., Conly, F. M., Pietroniro, A. and Martz, L.: The Fill-Spill  
1175 Hydrology of Prairie Wetland Complexes during Drought and Deluge, *Hydrol. Process.*, 26,  
1176 3147–3156, doi:10.1002/hyp.8390, 2012.

1177 Shook, K. and Pomeroy, J.: Changes in the hydrological character of rainfall on the Canadian  
1178 prairies, *Hydrol. Process.*, 26, 1752–1766, doi:10.1002/hyp.9383, 2012.

1179 Shook, K., Pomeroy, J. W., Spence, C. and Boychuk, L.: Storage dynamics simulations in prairie  
1180 wetland hydrology models: Evaluation and parameterization, *Hydrol. Process.*, 27, 1875–  
1181 1889, doi:10.1002/hyp.9867, 2013.

1182 Shook, K., Pomeroy, J. and van der Kamp, G.: The transformation of frequency distributions of  
1183 winter precipitation to spring streamflow probabilities in cold regions; case studies from the  
1184 Canadian Prairies, *J. Hydrol.*, 521, 394–409, doi:10.1016/j.jhydrol.2014.12.014, 2015.

1185 Shook, K. R. and Pomeroy, J. W.: Memory effects of depressional storage in Northern Prairie  
1186 hydrology, *Hydrol. Process.*, 25, 3890–3898, doi:10.1002/hyp.8381, 2011.

1187 Simpson, M.A.: Surficial Geology Map of Saskatchewan (250k\_surficial, vector digital data).  
1188 Original maps published 1984 to 1988; merged digital version made available 2008,  
1189 compiled by M.A. Simpson, Environment Branch, Saskatchewan Research Council and  
1190 Saskatchewan Ministry of the Economy. Available from:  
1191 <https://gisappl.saskatchewan.ca/Html5Ext/index.html?viewer=GeoAtlas>, 2008.

1192 Sivakumar, B., Singh, V. P., Berndtsson, R. and Khan, S. K.: Catchment Classification  
1193 Framework in Hydrology: Challenges and Directions, *J. Hydrol. Eng.*, 20,  
1194 130426211354007, doi:10.1061/(ASCE)HE.1943-5584.0000837, 2013.

1195 Spence, C. and Saso, P.: A hydrological neighbourhood approach to predicting streamflow in the  
1196 Mackenzie valley, *Predict. Ungauged Basins Approaches Canada's Cold Reg.*, 21–44, 2005.

1197 Statistics Canada: Table 32-10-0162-01. Selected land management practices and tillage  
1198 practices used to prepare land for seeding, historical data. Census of Agriculture. Statistics  
1199 Canada, Government of Canada. Available from:  
1200 <https://www150.statcan.gc.ca/t1/tbl1/en/cv.action?pid=3210016201>, 2016.

1201 Ulrich, A. E., Malley, D. F. and Watts, P. D.: Lake Winnipeg Basin: Advocacy, challenges and  
1202 progress for sustainable phosphorus and eutrophication control, *Sci. Total Environ.*, 542,  
1203 1030–1039, doi:10.1016/j.scitotenv.2015.09.106, 2016.

1204 Van der Kamp, G. and Hayashi, M.: Groundwater-wetland ecosystem interaction in the semiarid  
1205 glaciated plains of North America, *Hydrogeol. J.*, 17, 203–214, doi:10.1007/s10040-008-  
1206 0367-1, 2009.

1207 Van der Kamp, G., Hayashi, M. and Gallén, D.: Comparing the hydrology of grassed and  
1208 cultivated catchments in the semi-arid Canadian prairies, *Hydrol. Process.*, 17, 559–575,  
1209 doi:10.1002/hyp.1157, 2003.

1210 Van der Kamp, G., Hayashi, M., Bedard-Haughn, A. and Pennock, D.: Prairie Pothole Wetlands  
1211 – Suggestions for Practical and Objective Definitions and Terminology, *Wetlands*, 36, 229–  
1212 235, doi:10.1007/s13157-016-0809-9, 2016.

- 1213 Vanderhoof, M. K., Christensen, J. R. and Alexander, L. C.: Patterns and drivers for wetland  
1214 connections in the Prairie Pothole Region, United States, *Wetl. Ecol. Manag.*, 25, 275–297,  
1215 doi:10.1007/s11273-016-9516-9, 2017.
- 1216 Van Meter, K. J. and Basu, N. B.: Signatures of human impact: size distributions and spatial  
1217 organization of wetlands in the Prairie Pothole landscape, *Ecol. Appl.*, 25, 451–465,  
1218 doi:10.1890/14-0662.1, 2015.
- 1219 Vincent, L. A., Zhang, X., Brown, R. D., Feng, Y., Mekis, E., Milewska, E. J., Wan, H. and  
1220 Wang, X. L.: Observed trends in Canada’s climate and influence of low-frequency  
1221 variability modes, *J. Clim.*, 28, 4545–4560, doi:10.1175/JCLI-D-14-00697.1, 2015.
- 1222 Vicente-Serrano S.M., Beguería, S., and López-Moreno, J.I.: A Multi-scalar drought index  
1223 sensitive to global warming: The Standardized Precipitation Evapotranspiration Index –  
1224 SPEI, *J. Clim.*, 23, 1696–1718, doi: 10.1175/2009JCLI2909.1, 2010.
- 1225 Wagener, T., Sivapalan, M., Troch, P., and Woods, R.: Catchment classification and hydrologic  
1226 similarity, *Geogr. Compass*, 1, 901–931, doi: 10.1111/j.1749-8198.2007.00039.x, 2007.
- 1227 Wagner, T., Bremigan, M. T., Cheruvilil, K. S., Soranno, P. A., Nate, N. A. and Breck, J. E.: A  
1228 multilevel modeling approach to assessing regional and local landscape features for lake  
1229 classification and assessment of fish growth rates, *Environ. Monit. Assess.*, 130, 437–454,  
1230 doi:10.1007/s10661-006-9434-z, 2007.
- 1231 Weber, D., Sadeghian, A., Luo, B., Waiser, M. J. and Lindenschmidt, K. E.: Modelling  
1232 Scenarios to Estimate the Potential Impact of Hydrological Standards on Nutrient Retention  
1233 in the Tobacco Creek Watershed, Manitoba, Canada, *Water Resour. Manag.*, 31, 1305–  
1234 1321, doi:10.1007/s11269-017-1578-9, 2017.
- 1235

1236 **TABLES AND FIGURES**

1237 **Table 1** – Pre-processing of compositional data PCA results. Shown are the respective subsets,  
 1238 the number of initial fractional area variables before dimensional reduction, the number of  
 1239 principal components retains to reach over 80% of subset variation (except for tillage practice),  
 1240 and the proportion of variation explained by each component.

<b>Variable subset</b>	<b>Number of initial variables</b>	<b>Number of principal components</b>	<b>Total variation explained by component</b>
Surficial geology	6	2	1: 72.8% 2: 14.4%
Particle size class	3	2	1: 74.8% 2: 15.6%
Soil zone	5	2	1: 54.6% 2: 42.7%
Local surface form	5	3	1: 54.5% 2: 24.2% 3: 11.9%
Land cover	5	3	1: 36.8% 2: 25.2% 3: 20.6%
Tillage practice	3	2	1: 90.9% 2: 8.5%

1241



1242 **Table 2** – Canonical correlation coefficients for watershed attribute and hydrological variables of  
 1243 hydrological research stations from the canonical correlation analysis. Those variables used in  
 1244 multiple regression equations are denoted with a ‘\*’.

	Correlation	
	V1	V2
<b>Watershed attributes</b>		
Area*	0.36	-0.83
DSF*	-0.26	0.90
Fraction rock*	-0.64	0.61
Fraction natural area*	-0.26	0.71
Stream density	-0.27	0.37
Mean annual precipitation	-0.14	-0.30
Fraction water area	0.53	-0.19
<b>Hydrological variables</b>	<b>W1</b>	<b>W2</b>
Q2	-0.82	-0.58
Q100	-0.22	-0.98
Canonical $\lambda$	0.97	0.77

1245

1246

1247 **Table 3** – Correlation of study watershed attributes to principal components (PC). The values for  
 1248 the six PCs used in the cluster analysis are shown.

Variable	Abbreviation	PC1	PC2	PC3	PC4	PC5	PC6
Mean elevation	elevation	0.81	0.34	-0.14	0.09	-0.16	-0.17
Mean slope	slope.mean	0.61	-0.23	0.06	0.37	-0.10	0.11
Slope CV	slope.CV	0.30	-0.38	0.22	0.14	-0.41	-0.09
Total precipitation	precip	-0.85	-0.12	0.13	0.16	-0.07	0.30
Potential evapotranspiration	PET	0.31	-0.33	-0.33	-0.06	0.47	0.13
Non-effective area	NE.area	0.02	0.70	0.31	0.10	0.01	-0.15
Areal fraction below outlet ( $A_{BO}$ )	A_BO	0.14	0.25	0.27	-0.17	0.42	-0.01
Stream density	stream.density	0.08	-0.42	-0.39	0.03	-0.41	0.08
Wetland density	wetland.density	-0.63	0.46	0.11	-0.04	0.12	0.24
Wetland fraction	wetland.frac	-0.30	0.19	0.66	-0.36	0.02	0.11
Water area in largest wetland to total in watershed ( $W_L$ )	W_L	0.31	-0.44	0.51	-0.32	-0.06	-0.12
Location of largest wetland to outlet ( $L_w/L_o$ )	L_W/L_O	-0.01	-0.06	-0.22	0.09	-0.07	-0.07
Beta ( $\beta$ )	beta	0.17	0.49	-0.02	0.01	0.09	0.05
Xi ( $\xi$ )	xi	0.21	-0.23	0.57	-0.31	-0.10	-0.17
Runoff ( $Q_2$ )	Q2	-0.13	0.35	-0.47	0.00	-0.33	0.10
Soil texture PC1	Text.PC1	-0.07	-0.04	0.28	0.55	0.19	-0.32
Soil texture PC2	Text.PC2	0.02	-0.32	0.43	0.03	-0.31	0.54
Soil zone PC1	Soil.PC1	-0.65	-0.29	-0.07	0.19	-0.10	-0.24
Soil zone PC2	Soil.PC2	0.27	-0.12	-0.06	-0.11	0.40	0.25
Land cover PC1	LC.PC1	-0.44	0.38	-0.21	-0.26	-0.43	0.12
Land cover PC2	LC.PC2	0.42	0.22	-0.17	-0.53	0.15	0.03
Land cover PC3	LC.PC3	0.21	0.30	0.15	0.25	0.11	0.46
Surficial geology PC1	SF.PC1	0.06	0.21	-0.19	0.50	0.17	-0.09
Surficial geology PC2	SF.PC2	0.06	-0.38	0.24	0.47	0.11	-0.03
Surface form PC1	LL.PC1	-0.16	0.20	0.17	0.47	0.26	0.26
Surface form PC2	LL.PC2	-0.20	0.44	0.12	-0.03	0.04	-0.55
Surface form PC3	LL.PC3	0.41	0.38	0.20	0.21	-0.27	0.27
Land practice PC1	LP.PC1	-0.54	-0.58	-0.13	-0.10	0.32	-0.09
Land practice PC2	LP.PC2	0.14	-0.16	-0.24	-0.22	0.29	0.30
<i>Supplementary variables</i>							
Latitude	Lat	-0.15	0.24	0.26	-0.01	-0.33	-0.41
Longitude	Long	-0.73	-0.24	0.06	0.10	0.16	0.39
Area	Area	-0.05	0.27	-0.44	0.09	-0.15	-0.03
DSF	DSF	-0.02	-0.25	0.42	-0.05	0.12	0.01

1249

1250 **Table 4** – Classes and distinguishing variables of prairie watersheds. The v-test statistics, based  
 1251 on Ward’s criterion, are shown. Variables with v-test values greater or less than 10 and –10,  
 1252 respectively, are bolded to emphasize defining features of each class. All variables are significant  
 1253 to  $p < 0.001$ . *Classes: Southern Manitoba (1), Pothole Till (2), Pothole Glaciolacustrine (3),*  
 1254 *Major River Valleys (4), Interior Grasslands (5), High Elevation Grasslands (6), Sloped Incised*  
 1255 *(7).*


Class 1 (n=365)		Class 2 (n=879)		Class 3 (n=681)		Class 4 (n=536)	
Variable	v-test	Variable	v-test	Variable	v-test	Variable	v-test
<b>LP.PC1</b>	<b>48.11</b>	<b>wetland.density</b>	<b>28.23</b>	<b>LC.PC1</b>	<b>22.60</b>	<b>SF.PC2</b>	<b>19.83</b>
<b>precip</b>	<b>30.33</b>	<b>LL.PC1</b>	<b>24.81</b>	<b>wetland.frac</b>	<b>12.74</b>	<b>slope.CV</b>	<b>19.35</b>
<b>Soil.PC1</b>	<b>23.60</b>	<b>precip</b>	<b>22.74</b>	<b>Q2</b>	<b>12.63</b>	<b>xi</b>	<b>16.05</b>
<b>LP.PC2</b>	<b>14.74</b>	<b>SF.PC1</b>	<b>21.74</b>	<b>NE.area</b>	<b>11.12</b>	<b>W_L</b>	<b>15.39</b>
<b>PET</b>	<b>13.10</b>	<b>LC.PC1</b>	<b>17.19</b>	LL.PC2	9.45	<b>Text.PC2</b>	<b>15.07</b>
wetland.density	7.39	<b>LL.PC2</b>	<b>16.42</b>	wetland.density	8.05	<b>Text.PC1</b>	<b>14.40</b>
DSF	6.81	<b>Q2</b>	<b>15.77</b>	LC.PC2	6.70	<b>Soil.PC1</b>	<b>14.01</b>
SF.PC2	6.53	<b>Soil.PC1</b>	<b>15.76</b>	LL.PC3	6.53	<b>DSF</b>	<b>11.76</b>
stream.density	4.61	<b>NE.area</b>	<b>15.72</b>	xi	5.89	<b>precip</b>	<b>10.97</b>
LC.PC1	-3.37	<b>area</b>	<b>13.15</b>	W_L	4.58	<b>wetland.frac</b>	<b>10.92</b>
A_BO	-4.22	<b>Text.PC1</b>	<b>12.00</b>	precip	3.47	slope.mean	7.29
area	-5.46	LC.PC3	6.76	A_BO	-3.79	LP.PC1	3.52
slope.CV	-6.49	beta	5.31	slope.CV	-4.97	A_BO	-3.83
Q2	-8.47	L_W/L_O	4.20	L_W/L_O	-5.17	wetland.density	-4.41
SF.PC1	-8.90	LL.PC3	3.93	LP.PC2	-7.11	SF.PC1	-4.56
LC.PC2	-9.21	SF.PC2	-3.97	LC.PC3	-9.71	LC.PC1	-5.13
<b>LL.PC2</b>	<b>-14.18</b>	LP.PC1	-4.87	<b>LP.PC1</b>	<b>-12.38</b>	soil.PC2	-6.93
<b>slope.mean</b>	<b>-16.17</b>	stream.density	-5.92	<b>Soil.PC2</b>	<b>-13.01</b>	beta	-7.60
<b>beta</b>	<b>-16.88</b>	elevation	-7.15	<b>Text.PC1</b>	<b>-14.58</b>	elevation	-8.03
<b>LC.PC3</b>	<b>-18.13</b>	A_BO	-7.86	<b>slope.mean</b>	<b>-15.92</b>	<b>area</b>	<b>-11.04</b>
<b>NE.area</b>	<b>-28.97</b>	Text.PC2	-9.15	<b>SF.PC2</b>	<b>-17.03</b>	<b>LP.PC2</b>	<b>-11.44</b>
<b>LL.PC3</b>	<b>-36.59</b>	DSF	-9.93	<b>LL.PC1</b>	<b>-17.83</b>	<b>Q2</b>	<b>-13.27</b>
<b>elevation</b>	<b>-47.42</b>	<b>LP.PC2</b>	<b>-10.88</b>	<b>SF.PC1</b>	<b>-18.83</b>	<b>PET</b>	<b>-13.98</b>
		<b>Soil.PC2</b>	<b>-12.00</b>	<b>PET</b>	<b>-23.29</b>	<b>LC.PC2</b>	<b>-20.86</b>
		<b>PET</b>	<b>-13.15</b>				
		<b>slope.mean</b>	<b>-13.50</b>				
		<b>slope.CV</b>	<b>-16.26</b>				
		<b>LC.PC2</b>	<b>-16.29</b>				
		xi	-21.49				
		<b>W_L</b>	<b>-32.96</b>				

1256

**Table 4 – (cont'd)**

Class 5 (n=635)		Class 6 (n=702)		Class 7 (n=377)	
Variable	v-test	Variable	v-test	Variable	v-test
<b>A_BO</b>	<b>34.10</b>	<b>elevation</b>	<b>29.29</b>	<b>Text.PC2</b>	<b>27.65</b>
<b>LC.PC2</b>	<b>21.53</b>	<b>PET</b>	<b>20.16</b>	<b>LL.PC3</b>	<b>25.69</b>
<b>Soil.PC2</b>	<b>20.81</b>	<b>slope.CV</b>	<b>17.67</b>	<b>slope.mean</b>	<b>22.32</b>
<b>LC.PC3</b>	<b>17.44</b>	<b>slope.mean</b>	<b>16.12</b>	<b>LC.PC3</b>	<b>14.84</b>
<b>NE.area</b>	<b>16.22</b>	<b>stream.density</b>	<b>14.55</b>	<b>stream.density</b>	<b>13.82</b>
<b>beta</b>	<b>15.96</b>	<b>LC.PC2</b>	<b>14.09</b>	<b>Soil.PC2</b>	<b>13.09</b>
<b>elevation</b>	<b>13.31</b>	<b>W_L</b>	9.47	<b>elevation</b>	<b>12.42</b>
<b>PET</b>	<b>11.47</b>	<b>L_W/L_O</b>	6.80	<b>PET</b>	<b>11.47</b>
LL.PC2	8.11	LP.PC2	5.73	SF.PC2	6.80
LP.PC2	7.67	area	3.72	LP.PC2	6.39
LL.PC3	7.31	LL.PC2	3.62	slope.CV	5.87
wetland.frac	5.77	LP.PC1	-3.60	W_L	4.63
LL.PC1	5.50	Q2	-3.94	precip	-4.75
SF.PC2	-4.74	DSF	-4.91	A_BO	-5.65
area	-4.86	A_BO	-9.47	LC.PC1	-7.62
L_W/L_O	-7.11	<b>Soil.PC1</b>	<b>-10.17</b>	Text.PC1	-8.34
Q2	-9.34	<b>LL.PC3</b>	<b>-10.62</b>	<b>LP.PC1</b>	<b>-11.42</b>
LP.PC1	-9.96	<b>LC.PC3</b>	<b>-13.17</b>	<b>NE.area</b>	<b>-13.33</b>
<b>Text.PC2</b>	<b>-11.36</b>	<b>NE.area</b>	<b>-14.11</b>	<b>wetland.frac</b>	<b>-13.64</b>
<b>LC.PC1</b>	<b>-11.38</b>	<b>LL.PC1</b>	<b>-15.44</b>	<b>wetland.density</b>	<b>-16.27</b>
<b>slope.CV</b>	<b>-12.42</b>	<b>Text.PC2</b>	<b>-15.78</b>	<b>Soil.PC1</b>	<b>-16.43</b>
<b>precip</b>	<b>-20.86</b>	<b>LC.PC1</b>	<b>-17.15</b>	<b>LL.PC2</b>	<b>-39.41</b>
<b>Soil.PC1</b>	<b>-23.58</b>	<b>wetland.frac</b>	<b>-21.48</b>		
<b>stream.density</b>	<b>-26.34</b>	<b>wetland.density</b>	<b>-29.58</b>		
		<b>precip</b>	<b>-37.27</b>		


1259




1260

1261 **Figure 1** – Map of the study area spanning the Prairies ecozone in western Canada (inset). Large  
1262 cities in each of the three provinces are shown for reference, while the region characterized as  
1263 not contributing runoff (2-year) is also shown. Prairie ecozone based on the region classified by  
1264 the Ecological Stratification Group (1995).

1265




1268 **Figure 2** – Observed versus predicted estimates for (a) Q2, and (b) Q100. The dashed grey line  
1269 depicts the linear regression between observed and predicted flow values, and the black, solid  
1270 line shows a 1:1 relationship.



1275 **Figure 3** – Principal components analysis for candidate variables for classification. Active and  
 1276 supplementary variables are shown as solid black, and dashed blue arrows, respectively.  
 1277 Eigenvalues for PC axes are provided (inset), with black bars denoting the six PCs used in the  
 1278 hierarchical clustering analysis.

1279

1280



1281


1294 **Figure 4** – Dendrogram resulting from the hierarchical cluster analysis of principal components.

1295 The blue, dashed line indicates the cut in the tree, resulting in seven clusters. The amount of

1296 inertia gained by increasing the number of clusters ( $k$ ) is depicted in the inset panel.

1297






1299

1300 **Figure 5** – Classification of Prairie ecozone watersheds. Watershed delineations are from Lehner  
1301 and Grills (2013), available at [www.hydrosheds.org](http://www.hydrosheds.org). See text for detailed interpretation of the  
1302 seven clusters.


1303



1304


1305 **Figure 6** – Climatic variation among watershed classes. (a) Boxplots of total annual precipitation  
 1306 (grey) and potential evapotranspiration (PET) (white) for each watershed cluster. Lower, middle,  
 1307 and upper limits of boxes show the 25<sup>th</sup>, 50<sup>th</sup>, and 75<sup>th</sup> quantiles, respectively. (b) Wetland  
 1308 density to moisture deficit (Precipitation – PET). *Classes: Southern Manitoba (1), Pothole Till*  
 1309 *(2), Pothole Glaciolacustrine (3), Major River Valleys (4), Interior Grasslands (5), High*  
 1310 *Elevation Grasslands (6), Sloped Incised (7).*

1311



1312

1313 **Figure 7** – Boxplots of select variables by watershed class: (a) fraction of non-effective area; (b)  
1314 fraction of cropland; and (c) fraction of grassland. *Classes: (1) Southern Manitoba, (2) Pothole*  
1315 *Till, (3) Pothole Glaciolacustrine, (4) Major River Valleys, (5) Interior Grassland, (6) High*  
1316 *Elevation Grasslands, and (7) Sloped Incised.*




1317

1318 **Figure 8** – Wetland variables and simulated size distributions. Median (a) density of wetlands  
 1319 and (b) fraction of total watershed water area in the largest wetland ( $W_L$ ) are depicted by class.  
 1320 Panel (c) shows observed (dark) and simulated (light) percentiles of wetland areas. Predicted  
 1321 values are based on a generalized Pareto distribution and using median parameters of  $\beta$  and  $\zeta$  for  
 1322 each cluster. Simulated data were restricted to the raster pixel resolution of observed data from  
 1323 the Global Surface Water dataset. *Classes: Southern Manitoba (1), Pothole Till (2), Pothole*  
 1324 *Glaciolacustrine (3), Major River Valleys (4), Interior Grasslands (5), High Elevation*  
 1325 *Grasslands (6), Sloped Incised (7).*

1326

1327

1328




1329

1330 **Figure 9** – Density distributions of percent agreement of watersheds to the classification in Fig.  
1331 5 by watershed class. *Classes: Southern Manitoba (1), Pothole Till (2), Pothole Glaciolacustrine*  
1332 *(3), Major River Valleys (4), Interior Grasslands (5), High Elevation Grasslands (6), Sloped*  
1333 *Incised (7).*

1334

1335

1336



1337

1338 **Figure 10** – Agreement of assigned watershed classification from the (original) complete  
1339 analysis, with class assignments from the iterative approach using re-sampling. Classes are  
1340 coloured according to that shown in Fig. 5, with those identified under a new class (C8) depicted  
1341 in black. Watersheds that were removed from the subsets analyzed are in white. *Classes:*  
1342 *Southern Manitoba (1), Pothole Till (2), Pothole Glaciolacustrine (3), Major River Valleys (4),*  
1343 *Interior Grasslands (5), High Elevation Grasslands (6), Sloped Incised (7).*