

1 Response to Reviewers

2 **Title:** The Probability Distribution of Daily Precipitation at the Point and Catchment
3 Scales in the United States

4 **Manuscript ID:** hess-2015-92

5 **Authors:** Lei Ye, Lars S. Hanson, Pengqi Ding, Dingbao Wang, Richard M. Vogel
6

7
8 Dear Editor:

9 We greatly appreciate the constructive comments and suggestions provided by the
10 reviewers. We fully agree with the comments on the analysis of complete series of daily
11 precipitation. Thank you for pointing this out, and we have deleted all the results
12 pertaining to the complete rainfall series. Our detailed responses to the comments are
13 listed below.
14

15 **Reviewer #1**
16

17 **Comment 1: It only makes sense to advance a single continuous pdf for the wet**
18 **day case, regardless of where the data arises from, if one adds zeros, the**
19 **Lmoments will always land exactly on the Pearson Type III curve**

20 Response:

21 Thank you so much for pointing this out. In this revised manuscript, we have eliminated
22 all the results relating to the 'all day' or $X \geq 0$ conditions, focusing only on the
23 probability distribution of wet-day precipitation.
24

25 **Comment 2: Lines 363-365: “demonstrating that the parameter Gamma**
26 **distribution cannot describe the tail behavior of full-record series of**
27 **precipitation as has often been assumed in the past.” These lines are just the**
28 **opportunity for commenting on tail issues. Summary shape statistics are of**
29 **course affected by the tail behavior but they are not sufficient to reveal in a**

30 **robust way the behavior of the tail if the whole sample is used (I mean all**
31 **nonzero values) and not values that belong to the tail. For example in the**
32 **paper the authors cite (Papalexiou and Koutsoyiannis, 2016) after the fitting**
33 **using L-moments various measures were proposed in order to compare the**
34 **fitting in the most extreme value, the largest extremes the whole sample etc.**
35 **The author can see that the performance of distributions changed, still the**
36 **GG performed better but the BrXII increased its performance too. I just want**
37 **to say that indeed this approach can favour specific distributions and exclude**
38 **others like the G2 the authors mention, yet this is based judging the whole**
39 **distributional shape properties and it is not really robust to judge on the tail**
40 **when using the whole nonzero sample. Also other global studies indicated the**
41 **sub exponential nature of tails focusing on using only “tail” data (Papalexiou**
42 **et al., 2013; Serinaldi and Kilsby, 2014); the latter was also applied in a**
43 **seasonal basis, which by the way might be also a nice idea, i.e., the authors to**
44 **explore seasonal variation.**

45 Response:

46 Thank you. The sentence has been removed accordingly in the revised manuscript since
47 our analysis of the full-record series of precipitation has been eliminated. The reviewer
48 has suggested a good idea to explore the seasonal variation of the distribution of daily
49 rainfall. The ideas introduced by Papalexiou and Koutsoyiannis (2016) and others
50 concerning the impact of seasons on rainfall distributions could be a future effort. We
51 are aware that the choice of a suitable distribution for modeling rainfall would be quite
52 different if we were to focus our attention on the extreme tail behavior, as is the case
53 for example, when one fits a distribution to the series of annual maxima or peaks above
54 some threshold. However given our interest in wet-day precipitation, most situations of
55 practical relevance and concern are not with extreme rainfall, thus we focus on the
56 complete series of wet-day amounts in this paper, without special attention given to the
57 largest values.

58

59 **Comment 3: The P3 distribution is just the two-parameter Gamma distribution**
60 **(G2) with an additional location parameter which does not affect the shape**
61 **characteristics and thus $\tau\tau3$ and $\tau\tau4$. So the curve of P3 shown in $\tau4-\tau\tau3$ ratio**
62 **plots is the same as the G2. And obviously they have the same tail. The same**
63 **holds for GPA and GP2 and for any other distribution that has one shape**
64 **parameter and additional location parameters are added. Maybe to ease the**
65 **reader, as different formulations can be found in the literature, it would be no**
66 **harm to add a table of the distributions functions used.**

67 Response:

68 Thank you. We have added distribution functions into Tables 2 and 3.

69 **Table 2:** Theoretical probability distributions presented on the L-Kurtosis vs L-Skew
70 L-moment diagram. *Italicized distributions are special cases of other*
71 *distributions.*

| Distribution | Abbreviation | PDF | Parameters |
|------------------------------------|--------------|---|------------|
| Generalized Extreme Value Type III | GEV | $f(x) = \frac{1}{\eta} \left[1 - \left(\frac{x - \omega}{\eta} \right) \beta \right]^{\frac{1}{\beta} - 1} \exp \left\{ - \left[1 - \left(\frac{x - \omega}{\eta} \right) \beta \right]^{\frac{1}{\beta}} \right\}$ | 3 |
| Generalized Logistic | GLO | $f(x) = \frac{be^{-\frac{x-\mu}{\sigma}}}{\sigma \left(1 + e^{-\frac{x-\mu}{\sigma}} \right)^{b+1}}$ | 3 |
| Generalized Pareto | GPA | $f(x) = \frac{1}{\sigma} \left(1 + \frac{\xi(x-\mu)}{\sigma} \right)^{-(1/\xi-1)}$ | 3 |
| Lognormal | LN3 | $f(x) = \frac{1}{\sqrt{2\pi}(x-\alpha)\sigma} \exp \left[-\frac{1}{2\sigma^2} (\ln(x-\alpha) - \mu^2) \right]$ | 3 |
| Pearson Type III | P3 | $f(x) = \frac{p/a^d}{\Gamma(d/p)} x^{d-1} e^{-(x/a)^p}$ | 3 |
| <i>Exponential</i> | E | $f(x) = \begin{cases} \lambda e^{-\lambda x}, & x \geq 0 \\ 0, & x < 0 \end{cases}$ | 2 |
| <i>Gumbel</i> | G | $f(x) = \frac{1}{\beta} e^{-(z+e^{-z})}, z = \frac{x-\mu}{\beta}$ | 2 |
| <i>Normal</i> | N | $f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$ | 2 |

| | | | |
|-----------------|---|---|---|
| <i>Logistic</i> | L | $f(x) = \frac{e^{-\frac{x-\mu}{s}}}{s \left(1 + e^{-\frac{x-\mu}{s}}\right)^2}$ | 2 |
| <i>Uniform</i> | U | $f(x) = \begin{cases} \frac{1}{b-a}, & a < x < b \\ 0, & x < a \text{ or } x > b \end{cases}$ | 1 |

72

73 **Table 3:** Theoretical probability distributions presented on the L-Cv vs L-Skew L-
74 moment diagram.


| Distribution | Abbreviation | PDF | Parameters |
|--------------------|--------------|---|------------|
| Gamma | G2 | $f(x) = \frac{\beta^\alpha x^{\alpha-1} e^{-\beta x}}{\Gamma(\alpha)}$ | 2 |
| Generalized Pareto | GP2 | $f(x) = \frac{1}{\sigma} \left(1 + \frac{\xi x}{\sigma}\right)^{(-1/\xi-1)}$ | 2 |
| Lognormal | LN2 | $f(x) = \frac{1}{x\sigma\sqrt{2\pi}} e^{-\frac{(\ln x - \mu)^2}{2\sigma^2}}$ | 2 |
| Weibull | W2 | $f(x) = \frac{k}{\phi} \left(\frac{x}{\phi}\right)^{k-1} \exp\left\{-\left[\frac{k}{\phi}\right]^k\right\}$ | 2 |

75

76 **Comment 4:** The Weibull distribution could also be added in the analysis as a
77 **representative of distributions with stretched exponential tails.**

78 Response:

79 Thanks. We have added the Weibull (W2) distribution in the figure below and found
80 that W2 was not competitive with the three potential candidate pdfs G2, P2, and KAP.
81 Again, we emphasize that our interest is in the entire distribution, without focusing
82 attention on the extreme tail behavior.



83

84

85 **Comment 5: When we use distributions with a location parameter to describe a**
 86 **positive variable like the nonzero precipitation this parameter might end far**
 87 **from zero or even negative indicating a lower bound. So, this distribution**
 88 **cannot be used in stochastic modelling of precipitation as it will result in**
 89 **inconsistent values. It would be interesting the authors to actually show some**
 90 **box plots of the estimated parameters.**

91 Response:

92 Thank you so much for your comment. We agree with you that some parameters can be
 93 very interesting. However, the focus of the paper is on the choice of distribution for
 94 daily precipitation. In order to save the length of the paper, we don't discuss the
 95 parameters here. We will study the parameters in the future work.

96

97 **Comment 6: The principle of parsimony should always be applied. If the authors,**
 98 **generate samples from a 4-parameter distribution like the kappa and try to**
 99 **estimate a posteriori the parameters, even for the sample sizes used here, they**
 100 **will find a huge variability that makes, in my opinion, the operational use of**
 101 **4-parameter distributions quite risky. Of course a 4-parameter distribution**
 102 **like the kappa has a great flexibility, yet this does imply that a better fitting**

103 **to an observed sample is a better choice to extrapolate values for large return**
104 **periods.**

105 Response:

106 We fully agree that in most applications in hydrology, the principle of parsimony is
107 absolutely paramount, due to the short samples available for fitting distributions.
108 However, in this application, with samples sizes in the tens of thousands, concerns over
109 parsimony are not nearly as critical, even when estimating the KAP distribution. This
110 fact has been shown nicely in the recent paper by Blum et al (2017) where they
111 demonstrated, for similarly length samples of daily streamflow the sampling properties
112 of estimated Lmoments from synthetic samples in their Figure 2.

113 We do not think we should be advocating a four parameter distribution (KAP) unless
114 absolutely necessary, because it is a much more complex model than may be needed.
115 We conclude that for representing wet-day precipitation, the Gamma and Pearson-III
116 distributions are comparable with the 4-parameter Kappa distribution for the areal
117 average precipitation, with P3 only slightly better than G2; however, when the point
118 precipitation is of concern, the Kappa distribution could be the distribution of choice.

119


120 **Comment 7: The authors, since this is the first large scale study on catchment**
121 **precipitation, could provide some analysis on the relation of catchment size**
122 **and distributional shape. As the spatial averaging process will tend to make**
123 **the distributions more bell-shaped and with thinner tails. This is the**
124 **explanation of the performance decrease of the heavy-tailed distribution**
125 **shown in Fig. 7 compared to Fig. 6 (commenting on the Wet-day; full-day**
126 **results should be modified).**

127 Response:

128 Thank you for your suggestion. In the revised manuscript, we have explored the relation
129 between catchment size and distribution shape in the Discussion section.

130 Lines 393-403: "Figure 9 displays the PPCC values of P3 and G2 pdfs versus catchment
131 drainage area for areal average wet-day series. The PPCC values are chosen from 0.99-

132 1, approximately 96% of catchments are displayed on the figure; the remaining points
 133 lie outside the plot domains. It can be seen that for most of the catchments, the PPCC
 134 values for G2 and P3 pdfs are very close, with points corresponding to G2 and P3 pdfs
 135 almost overlapping. This is especially true for PPCC values higher than 0.998. The
 136 phenomena clearly indicates that when G2 can well represent the behavior of
 137 catchment-scale wet-day precipitation series, P3 also provides very good performance.
 138 However, for the areas where PPCC values are lower than 0.996, the P3 distribution
 139 outperforms the G2 distribution for most cases, with a very slight improvement.”



140

141 **Figure 9:** the PPCC values of P3 and G2 pdfs versus catchment drainage area for areal average wet-
 142 day series.

143


144 **Comment 8: Some regions of the USA, mainly in Midwest, show quite intense**
 145 **changes (or maybe natural variability) on extremes. The authors could also**
 146 **comment on that or do a quick extra analysis on the daily precipitation.**

147 Response:

148 Thank you very much for you suggestion. We have displayed the best distribution
 149 functions for areal average wet-day series of the catchments by showing the location
 150 on a map.

151 Lines 405-412: “KAP distribution is the best pdf for large proportion of the catchments
 152 especially in the middle of US. P3 distribution occupies the second large proportion of

153 the catchments especially in east-central US. Only a very few catchments can be best
154 represented by G2 distribution. Seen from Figure 10, it seems that the performances of
155 the three pdfs vary greatly. However, as we have seen from previous figures, the
156 differences between the three pdfs for catchments are very small.”



157
158 **Figure 10:** The spatial map of catchments with the corresponding best distribution functions for
159 areal average wet-day series.

160

161 **Comment 9:** I believe the literature should be updated with many other works, e.g.,
162 there are several papers that are using other distributions for daily
163 precipitation, e.g., one that came to mind is the by Wilson and Toumi (2005).

164 Response:

165 Thank you. We have updated the literature review with a focus on the distribution of
166 wet-day precipitation amounts. The following papers have been added in the
167 References.

168 Chen, J., Brissette, F. P., and Leconte, R.. Downscaling of weather generator
169 parameters to quantify hydrological impacts of climate change. *Climate Research*, 51,
170 185-200, 2012.

171 Kigobe, M., McIntyre, N., Wheeler, H., and Chandler, R.. Multi-site stochastic
172 modelling of daily rainfall in Uganda. *Hydrological sciences journal*, 56, 17-33, 2011.

173 Li, Z., Brissette, F., Chen, J.. Finding the most appropriate precipitation probability
174 distribution for stochastic weather generation and hydrological modelling in Nordic
175 watersheds. *Hydrological Processes*, 27: 3718-3729, 2013.

176 Wilson, P. S., and Toumi, R.. A fundamental probability distribution for heavy
177 rainfall. Geophysical Research Letters, 32, L14812, 2005.

178

179 **Reviewer #2**

180

181 **Comment 1: It should probably be explained in the Introduction why**
182 **“Establishing a probability distribution that provides a good fit to daily**
183 **precipitation depths has long been a topic interest”.**

184 Response:

185 Thank you for your comment. We have explained why “Establishing a probability
186 distribution that provides a good fit to daily precipitation depths has long been a topic
187 interest” in the introduction section as follows.

188 Lines 33-36: “Precipitation is paramount in the fields of hydrology, meteorology,
189 climatology, and others. However, long series of precipitation data are not always
190 available; therefore, establishing a probability distribution that provides a good fit to
191 daily precipitation depths has long been a topic interest.”

192

193 **Comment 2: The research objectives are included in the subsection “Precipitation**
194 **trends and changes”, which isn’t really logical. Consider restructuring the**
195 **Introduction, for example, by adding a “Research objectives” subsection.**

196 Response:

197 Thank you. We have reorganized the introduction section by adding a “Research
198 objectives” subsection so that we state the research objectives explicitly at the end of
199 the introduction section.

200

201 **Comment 3: The Introduction is almost half of the paper. Considering shortening**
202 **it or moving the less essential material to a background subsection.**

203 Response:

204 Thank you. We have shortened the introduction section in the revised manuscript to

205 focus on the distribution of wet-day precipitation.

206

207 **Comment 4: Line 267: Regarding “less than “0.01” recordable precipitation,”**
208 **what are the units of the 0.01? Isn’t this threshold too low given the detection**
209 **limit of gauges (approximately 0.25 mm)?**

210 *Response:*

211 It is common practice in the US to report daily precipitation amounts in inches and 0.01
212 inches is commonly considered the detection limit which is approximately equivalent
213 to 0.25 mm. We have explained it in the paper as follows.

214 Lines 198-200: “The wet-day series were constructed by excluding zero and “trace”
215 values (those with less than 0.01 inches (approximately equivalent to 0.25 mm)
216 recordable precipitation).”


217

218 **Comment 5: Can you show some maps of the results to reveal what the spatial**
219 **patterns in the results look like? Are there any striking differences between,**
220 **for example, the temperate southeastern and arid southwestern US?**

221 *Response:*

222 Thank you very much for your suggestion. We have displayed the best distribution
223 functions for areal average wet-day series of the catchments by showing the location
224 on a map.

225 Lines 405-412: “KAP distribution is the best pdf for large proportion of the catchments
226 especially in the middle of US. P3 distribution occupies the second large proportion of
227 the catchments especially in east-central US. Only a very few catchments can be best
228 represented by G2 distribution. Seen from Figure 10, it seems that the performances of
229 the three pdfs vary greatly. However, as we have seen from previous figures, the
230 differences between the three pdfs for catchments are very small.”



231

232 **Figure 10:** The spatial map of catchments with the corresponding best distribution functions for
 233 areal average wet-day series.

234

235 **Comment 6:** A Discussion section is missing from the paper.

236 *Response:*

237 We have added a Discussion section before the Conclusions.

238

239 **Comment 7:** What is the broader significance of the results? Are the results
 240 representative of the rest of the world?

241 *Response:*

242 With increased attention on the impact of climate change, an understanding of the
 243 distribution of daily wet-day precipitation is paramount for modeling such impacts.

244 Importantly, the U.S. continent has extremely broad variations in climatic conditions so
 245 that the results of our study, which employ very large continental datasets, should be
 246 illustrative of most other regions of the globe, with the exception of extremely tropical
 247 and extremely frigid climates, of which there are none in the U.S.

248

249 **Reviewer #3**

250

251 **Comment 1:** The ‘Introduction’ section describes in great detail the vast literature
 252 related to the topics of (1) stochastic precipitation modeling, (2) precipitation

253 frequency analysis, and (3) precipitation trends and climate changes. In this
254 thorough review it is apparent that the Pearson Type III (P3) distribution has
255 not been considered as a candidate distribution to describe wet-day, AMS or
256 PDS daily precipitation series. Yet the consideration of the P3 distribution is
257 largely explored in this paper. Recommend the authors add why they believe
258 the P3 is an appropriate distribution for the extreme values of rainfall.

259

260 Response:

261 The two parameter Gamma distribution is the most widely used distribution of daily
262 rainfall in previous studies. Therefore it is only natural that one should also consider
263 fitting a three parameter version of the Gamma distribution, known as the P3
264 distribution to daily rainfall amounts. Given that hundreds of studies have assumed the
265 Gamma distribution, we were very surprised to find so little attention given to the three
266 parameter version of the Gamma distribution, known as the P3 distribution. This is one
267 of the most important contributions of our paper, bringing this fact to light.

268

269 **Comment 2: Similar to Referee #2, I believe too much detail is presented in the**
270 **‘Introduction’ section. The lengthy discussion doesn’t add to the flow of the**
271 **paper. Recommend reducing the literature review discussion, highlighting the**
272 **important studies related to the topics in this paper and refer the reader to**
273 **Table 1 for a more thorough review of previous studies.**

274 Response:

275 As you said, the introduction section indeed accounts for a relatively large proportion
276 of the manuscript. We have shortened the introduction section significantly and refer
277 the reader to Table 1 for a more thorough review of previous studies.

278

279 **Comment 3: Similar to Referee #2, a ‘Discussion’ section is missing in this paper**
280 **and I recommend it be added.**

281 Response:

282 We have added a Discussion section before the Conclusions.

283

284 **Comment 4: 234-239 is interpretive and describes the findings of this paper. This**
285 **should be moved to the ‘Discussion’ and/or ‘Conclusions’ sections. Similarly,**
286 **the last sentence in the ‘Introduction’ section (lines 243-245) is interpretive**
287 **and should be moved to the ‘Discussion’ and/or ‘Conclusions’ sections.**

288 *Response:*

289 We have moved the contents of lines 234-239 and lines 243-245 to the Conclusions
290 section.

291

The Probability Distribution of Daily Precipitation at the Point and Catchment Scales in the United States

Lei Ye^{1*}, Lars S. Hanson², Pengqi Ding¹, Dingbao Wang³, Richard M. Vogel⁴

¹ School of Hydraulic Engineering, Dalian University of Technology, Dalian, China

² Institute for Public Research, Center for Naval Analyses, Arlington, Virginia, USA.

³ Department of Civil, Environmental, and Construction Engineering, University of Central Florida, Orlando, Florida, USA

⁴ Department of Civil and Environmental Engineering, Tufts University, Medford, Massachusetts, USA

Abstract: Choosing a probability distribution to represent daily precipitation depths is important for precipitation frequency analysis, stochastic precipitation modeling and in climate trend assessments. Early studies identified the 2-parameter Gamma (G2) distribution as a suitable distribution for wet-day precipitation based on traditional goodness of fit tests. Here, probability plot correlation coefficients and L-moment diagrams are used to examine distributional alternatives for the wet-day series of daily precipitation for hundreds of stations at the point and catchment scales in the United States. Importantly, the G2 distribution performs poorly in comparison to either the Pearson Type-III (P3) or Kappa (KAP) distributions particularly for point rainfall. Our analysis indicates that the KAP distribution best describes the distribution of wet-day precipitation at the point scale, whereas the performance of G2 and P3 distributions are comparable for wet-day precipitation at the catchment scale, with P3 generally providing improved goodness of fit over G2. Since the G2 distribution is currently the most widely used probability density function, our findings could be considerably important, especially within the context of climate change investigations.

Key Words: Climate; Rainfall; Weather; L-moment diagram; PPCC; Pearson type III; Kappa; Gamma; Wet-day; Frequency analysis; Trend detection; Stochastic weather models

1. Introduction

Precipitation is paramount in the fields of hydrology, meteorology, climatology, and others. However, long series of precipitation data are not always available; therefore, establishing a probability distribution that provides a good fit to daily precipitation depths has long been a topic interest. Investigations into the probability distribution of daily precipitation can be found in at least three main research areas, namely, (1) stochastic precipitation models, (2) frequency analysis of precipitation, and (3) precipitation trends related to global climate change. Table 1 displays a sampling of the literature related to those three topics including the particular precipitation series and durations under investigation, and the proposed probability distributions recommended. Table 1 is by no means exhaustive; it only attempts to document the widespread interest in the

* Corresponding author. E-mail address: yelei@dlut.edu.cn

43 determination of a suitable distribution for daily precipitation totals in a wide range of
44 studies across a wide range of fields of inquiry.

45 *[Table 1 goes here]*

46 **1.1 Stochastic precipitation models:**

47 Our central goal is to select a suitable generalized probability distribution for modeling
48 daily precipitation depths, thus we are only concerned with the class of “two-part”
49 stochastic daily precipitation models that utilize a probability distribution function to
50 describe precipitation amounts on wet-days, while a probabilistic representation of
51 precipitation occurrences can be separately described using a Markov model or some
52 form of a stochastic renewal process (Buishand, 1978;Geng et al., 1986;Waymire and
53 Gupta, 1981;Watterson, 2005). We only consider the selection of a suitable distribution
54 for modeling wet-day daily rainfall, leaving the stochastic representation of the
55 occurrence of zeros, to others.

56 It is evident from Table 1 that the wet-day precipitation series is the primary
57 series considered within the stochastic precipitation model literature. Thom’s (1951)
58 suggestion of the 2-parameter Gamma (G2) distribution function for wet-day amounts
59 seems to carry considerable weight. Buishand (1978) lent support to the suggestion of
60 the G2 distribution by showing that for the wet-day series at six stations, the empirical
61 Coefficient of Variation to Coefficient of Skewness ratio was quite close to the
62 theoretical value of two for a G2 distribution. Geng et al. (1986) provided a review of
63 other literature supporting the use of the G2 distribution for modeling wet-day rainfall.

64 While the G2 distribution is by far the most commonly advocated distribution for
65 wet-day precipitation amounts, other distributions have also been suggested. Woolhiser
66 and Roldan (1982), Wilks (1998) and Li et al. (2013) suggested the use of a three-
67 parameter mixed exponential distribution instead of G2. Through a variety of goodness
68 of fit tests and log-likelihood analyses, the mixed exponential was preferred to G2 (Wilks,
69 1998).

70 The Weibull (W2) and to a lesser extent the exponential distribution have also
71 been suggested for modeling daily precipitation amounts (Duan et al., 1995;Burgueno et
72 al., 2005). Duan et al. (1995) used a Chi-squared test to demonstrate that synthetic
73 rainfall generated from the W2 and G2 models best match the observed daily rainfall data
74 within each month. Burgueno et al. (2005) used graphical methods and the Kolmogorov-
75 Smirnov test to give support to the W2 and exponential distributions.

76 **1.2 Precipitation frequency analysis:**

77 The second section of Table 1 displays a small portion of the literature related to
78 precipitation frequency analyses. Since extreme rainfall values are of primary
79 importance in these studies, censored series of rainfall (e.g. the Annual Maximum Series
80 (AMS) and Partial Duration Series (PDS)) are often useful in these analyses (Stedinger et
81 al., 1993). Table 1 displays that many of the precipitation frequency investigations of
82 daily precipitation depths have selected the AMS series.

83 For many years, the most common approach to summarizing precipitation
84 frequency analyses in the United States was the work of Hershfield (1961), which is

85 commonly referred to as TP-40. Hershfield (1961) fitted a Gumbel distribution to the
86 AMS of 24-hour precipitation. In the context of a national revision to the TP-40, Bonnin
87 et al., (2006) fitted a generalized extreme value (GEV) distribution to the AMS of rainfall.

88 While the results of Bonnin et al. (2006) apply to the United States, other
89 researchers have found similar results using similar methods in other parts of the world.
90 Pilon et al. (1991) used L-moment goodness-of-fit results to show that the Gumbel
91 distribution should be rejected in the favor of the GEV in Ontario, Canada. In Korea,
92 Park and Jung (2002) successfully used the Kappa distribution (of which the GEV is a
93 special case) to generate extreme precipitation quantile maps. In perhaps the most
94 comprehensive assessment of the distribution of precipitation extremes, Papalexiou and
95 Koutsoyiannis (2013) examined the goodness-of-fit of the GEV distribution to a global
96 dataset of AMS. Analysis of such a large dataset enabled them to conclude that GEV
97 models of AMS of daily precipitation provide a good approximation.

98 Interestingly, while a great deal of attention is given to fitting distributions to the
99 relatively short AMS series of precipitation depths, very few studies directly explore the
100 probability distribution of the complete series of daily precipitation (including zeros) or
101 the wet-day series of daily precipitation (zeros excluded). Shoji and Kitaura (2006)
102 investigated both complete and wet-day daily precipitation series, but included only the
103 normal, lognormal, exponential, and W2 distributions as candidate distributions, and did
104 not employ modern regional hydrologic methods such as the method of L-moments.
105 Deidda and Puliga (2006) investigated the degree of left-censoring of wet-day series
106 needed to fit a Generalized Pareto (GPA) distribution for 200 stations in Italy with a
107 range of modern statistical analysis techniques. Wilson and Toumi (2005) derived a
108 fundamental distribution for heavy rainfall, with a simple expression for rainfall as the
109 product of mass flux, specific humidity and precipitation efficiency. Statistical theory
110 predicted that the tail of the derived rainfall distribution has a stretched exponential form
111 with a shape parameter of 2/3, which was verified by a global daily precipitation data set.

112 Perhaps the most thorough investigations, to date, on the probability distribution
113 of daily precipitation amounts are the global studies by Papalexiou and Koutsoyiannis
114 (2012, 2016). Papalexiou and Koutsoyiannis (2012) derived a generalized Gamma
115 distribution (GG) from Entropy theory, using plausible constraints for wet-day series of
116 daily precipitation series. Together, the two studies by Papalexiou and Koutsoyiannis
117 (2012, 2016) revealed that the GG distribution provides a good approximation to the
118 behavior of observed L-moments of global series of wet-day daily precipitation at 11,519
119 and 14,157 stations, respectively.

120 **1.3 Precipitation trends and changes:**

121 The third section of Table 1 summarizes a small portion of the precipitation trend
122 literature which has become a rather large area of inquiry due to concerns over climate
123 change, as evidenced from recent reviews on the subject (Easterling et al., 2000;
124 Trenberth, 2011; Madsen et al., 2014). Almost universally, the G2 distribution appears to
125 be accepted without serious consideration of alternative distributions. For instance,
126 Groisman et al. (1999) compared maps of the empirical probability of summer 1-day
127 rainfall exceeding 50.4 mm with maps of probabilities determined by a stochastic model

128 using the fitted G2 distribution for the amounts. They found acceptable fits in regions
129 where there are enough observed daily rainfall events greater than 50.4 mm.

130 This is an interesting contrast to the precipitation frequency analysis literature
131 where a G2 distribution is often fit to wet-day series for the purpose of examining
132 extreme rainfalls instead of using the AMS series fitted by a GEV or other distribution.
133 Yoo et al. (2005) explained that conventional frequency analysis (using AMS) cannot
134 expect to predict precipitation changes resulting from climate change; while an
135 examination of the differences in the G2 distribution's parameters (fitted to the whole
136 wet-day record) might predict such changes. They found that modifying the parameters
137 of the daily G2 distribution can explain changes in rainfall quantiles predicted by General
138 Circulation Models under various climate change scenarios.

139 In a national study of precipitation trends, Karl and Knight (1998) employed the
140 G2 distribution to fill in missing precipitation observations. Both Watterson and Dix
141 (2003) and Watterson (2005) assumed a G2 distribution for daily precipitation in the
142 development of stochastic rainfall models for use in evaluating changes in precipitation
143 extremes.

144 **1.4 Research objectives:**

145 In summary, there are a wide variety of previous studies which have explored the
146 probability distribution of daily precipitation for the purposes of precipitation frequency
147 analysis, stochastic precipitation modeling and for trend detection. There seems to be a
148 consensus that annual maxima appear to be well approximated by either a GEV, Gumbel
149 or GPA probability density function (pdf) and that series of wet-day daily precipitation
150 totals are well approximated by a G2, GG, W2 or in some cases a mixed exponential pdf.
151 However, other than the two recent global studies by Papalexiou and Koutsoyiannis
152 (2012, 2016), we are unaware of any studies that have used recent developments in
153 regional hydrologic frequency analysis such as L-moment diagrams or probability plot
154 goodness of fit evaluations to evaluate the probability distribution of very large regional
155 datasets comprised of the wet-day series of daily precipitation.

156 The recent studies by Papalexiou and Koutsoyiannis (2012; 2016) represent
157 perhaps the most comprehensive studies to date. However, their L-moment evaluations
158 only evaluate the relationship between L-Skewness and L-Cv; thus they were unable to
159 fully evaluate the goodness-of-fit of the several relatively new three-parameter pdfs
160 introduced in their studies such as the GG and the generalized Burr type XII pdfs which
161 would require construction of L-Kurtosis versus L-Skew diagrams which are currently
162 unavailable for those pdfs. Analogous to those two studies, this paper uses two large
163 scale national datasets to re-examine the question of which of the commonly used
164 continuous distribution functions which are widely used in the fields of hydrology,
165 meteorology and climate best fit wet-day series of observed daily precipitation data. We
166 focus our research interest on the distribution of wet-day series of precipitation since the
167 pdf of complete series can be derived by a mixed distribution consisting of a combination
168 of the pdf of wet-day series and a stochastic model of the percentage and occurrence of
169 zeros.

170 Instead of considering the GG distribution, the pdf recommended by both
171 Papalexiou and Koutsoyiannis (2012, 2016), which has seen very limited use and for

172 which analytical and/or polynomial relationships for L-Kurtosis are unavailable as they
173 are for most commonly used pdfs in hydrology, we consider the more widely used 3
174 parameter generalization of the G2 distribution known as the Pearson type III (P3)
175 distribution. Our primary objective is to use a very large national spatially distributed
176 dataset at both the point and catchment scales, to determine a suitable probability
177 distribution of wet-day series of daily precipitation using L-moment diagrams and
178 probability plot correlation coefficient goodness of fit statistics.

179 **2. Study area and data**

180 Precipitation depths at the point and catchment scales provide important information in
181 hydrology, meteorology, and other fields, thus our study focuses on both scales. For
182 point precipitation, we employ a data set comprised of daily precipitation depths at 237
183 first-order NOAA stations from 49 U.S. states (Hawaii is excluded due to fundamentally
184 different precipitation behavior). Station locations are shown in Figure 1a. In contrast,
185 the areal average precipitation for 305 catchments in the international Model Parameter
186 Estimation Experiment (MOPEX) data set (Duan et al., 2006) is also selected for analysis.
187 The catchment locations and boundaries are shown in Figure 1b. The data were quality
188 controlled to remove null values. When greater than 6 null values occurred in a given
189 year or greater than 3 in a given month, the full year of data was removed. When fewer
190 than these numbers of null values were present, they were treated as zeroes. The average
191 record length for point precipitation depths for the 237 sites is 24,657 days (67.5 years).
192 The distribution of record lengths corresponding to the 237 first-order NOAA stations is
193 shown in Figure 2. The MOPEX data set consists of 56 years of areal average daily
194 precipitation from 1948 to 2003, corresponding to a fixed record length 20,454 days for
195 each of the 305 catchments shown in Figure 1b.

196 *[Figure 1 goes here]*

197 *[Figure 2 goes here]*

198 The wet-day series were extracted from both data sets. The wet-day series were
199 constructed by excluding zero and “trace” values (those with less than 0.01 inches
200 (approximately equivalent to 0.25 mm) recordable precipitation). Wilks (1990) discussed
201 other ways to treat trace precipitation and left-censored data, but for convenience, they
202 are simply excluded. The mean wet-day record lengths for point and areal average
203 precipitation are 7,219 days (equivalent to nearly 20 years) and 14,043 days (more than
204 38 years), respectively. The distributions of wet-day record length are shown in Figure 3.
205 As expected, the proportion of wet-days in the areal average precipitation data set is
206 higher than that in the point precipitation data set.

207 *[Figure 3 goes here]*

208 **3. Methodology**

209 This section describes the methods of analysis used for assessing the goodness-of-fit of
210 various distributional hypotheses, namely, L-moment diagrams and probability plot
211 correlation coefficients.

212 **3.1 L-Moment Diagrams**

213 L-moment diagrams are now a widely accepted approach for evaluating the goodness of
214 fit of alternative distributions to observations. The theory and application of L-moments
215 introduced by Hosking (1990) is now widely available in the literature (Stedinger et al.,
216 1993; Hosking and Wallis, 1997), hence it is not reproduced here.

217 The distribution of daily precipitation totals is highly skewed due to the large
218 proportion of days with zero or small precipitation. Higher order conventional moment
219 ratios such as skewness and kurtosis are very sensitive to extreme values and can exhibit
220 enormous downward bias even for extremely large sample sizes (Vogel and Fennessey,
221 1993) as is the case here. However, L-moment ratios are approximately unbiased in
222 comparison to conventional moment ratios, thus providing a particularly useful tool for
223 investigating the pdf of daily wet-day precipitation series.

224 L-moment ratio diagrams provide a convenient graphical image to view the
225 characteristics of sample data compared to theoretical statistical distributions. The L-
226 moment diagrams: L-Kurtosis (τ_4) vs L-Skew (τ_3) and L-Cv (τ_2) vs L-Skew (τ_3) enable us
227 to compare the goodness of fit of a range of three-parameter, two-parameter, and one-
228 parameter (or special case) distributions. Table 2 displays distributions analyzed by
229 means of the τ_4 vs τ_3 L-moment ratio diagrams.

230 *[Table 2 goes here]*

231 Table 3 displays distributions analyzed by means of the τ_2 vs τ_3 L-moment ratio
232 diagrams.

233 *[Table 3 goes here]*

234 L-moment ratio diagrams have been used before to examine the distribution of
235 series of annual maximum precipitation data (Pilon et al., 1991; Park and Jung, 2002; Lee
236 and Maeng, 2003; Papalexioiu and Koutsoyiannis, 2013) and left-censored records
237 (Deidda and Puliga, 2006). Other than the two recent global studies by Papalexioiu and
238 Koutsoyiannis (2012, 2016) which examined the agreement between empirical and
239 theoretical relationships between L-Cv and L-Skew, this is the only study we are aware
240 of, in which a set of daily wet-day precipitation records have been subjected to such a
241 comprehensive L-moment goodness-of-fit analysis. L-moment estimators were chosen in
242 this study for a variety of reasons: (1) they are easily computed and nicely summarized
243 by Hosking and Wallis (1997) for all the cases considered in this study, and (2) estimates
244 of L-moments are unbiased and estimates of L-moment ratios are nearly unbiased, and
245 thus for the extremely large sample sizes considered here, sampling variability of
246 empirical L-moment ratios will be extremely small especially when contrasted to the
247 variability among the theoretical L-moment ratios corresponding to the various
248 distributions considered..

249 **3.2 Probability plot correlation coefficient goodness-of-fit evaluation**

250 Probability plots are constructed for each of the wet-day series using L-moment
251 estimators of the distribution parameters (see Hosking and Wallis (1997)) for the
252 distributions indicated in Table 4. A probability plot is constructed in such a manner as
253 to ensure that the observations will appear to create a linear relationship when they arise
254 from the hypothesized distribution assumed for each plot.

255

[Table 4 goes here]

256 The goodness of fit of each probability plot is summarized using a probability plot
257 correlation coefficient (PPCC, or simply, r) which is simply a measure of the linearity of
258 the plot. The PPCC statistic has a maximum value of 1. The PPCC has been shown to be
259 a powerful statistic for evaluating the goodness-of-fit of a wide range of alternative
260 distributional hypotheses (Stedinger et al., 1993) and for performing hypothesis tests of
261 various two parameter distributional alternatives.

262 To construct a probability plot and to estimate a PPCC, requires estimation of a
263 plotting position. There are two classes of plotting positions, those that yield unbiased
264 exceedance probabilities and those that yield unbiased quantile estimates. The Weibull
265 plotting position given by $p=i/(n+1)$ yields an unbiased estimate of exceedance
266 probability regardless of the underlying distribution (see Stedinger et al. (1993)).
267 Alternatively there would be a unique plotting position to use for each probability
268 distribution, and it is now well known that unbiased plotting positions for three parameter
269 distributions require an additional parameter to estimate within the plotting position. For
270 example, Vogel and McMartin (1991) derived an unbiased plotting position for the P3
271 distribution which depends upon the skewness of the distribution, a parameter which adds
272 so much additional uncertainty to the analysis that led Vogel and McMartin (1991), after
273 considerable analysis, to not recommend its use. To put all the distributional alternatives
274 on the same footing, we chose to use the Weibull plotting position for estimation of all
275 PPCC values.

276 **4. Results and analysis**

277 **4.1 L-Moment Diagrams**

278 **4.1.1 L-Cv vs L-Skew**

279 Figure 4 displays empirical and theoretical distributional relationships between L-Cv and
280 L-Skew for point values of daily precipitation (Figure 4a) and areal average values of
281 daily precipitation (Figure 4b). The various curves represent the theoretical relationship
282 between L-Cv and L-Skew for the distributions indicated. Each plotted point represents
283 the empirical relationship between L-Cv and L-Skew for a single precipitation station or
284 catchment. By comparing the empirically derived points with the theoretical curves, it is
285 possible to see the degree to which the distributional tail behavior of the data record
286 matches those of the candidate distributions. We emphasize again, importantly, that the
287 sample sizes are large enough in this study so that one may, approximately, ignore
288 sampling variability in all L-moment diagrams. This phenomenon was nicely illustrated
289 in Figure 2 of Blum et al. (2017), using synthetic data, for record lengths similar to those
290 used here, but corresponding to daily streamflow records.

291

[Figure 4 goes here]

292 In Figure 4a, the L-moment ratios fall primarily within a region bounded by the
293 G2 and GP2 theoretical curves, with the W2 passing through some of the points. In
294 Figure 4b, the L-moment ratios fall primarily in the upper region of the W2 theoretical
295 curve, with the G2 passing through or very close to most of the points. These patterns do
296 not indicate a clearly preferred distribution for point values, especially considering that

297 the large sample sizes associated with these series result in negligible sampling variability.
298 However, Figure 4b documents that the G2 pdf provides a good approximation to the pdf
299 of wet-day series for areal average values.

300 Blum et al. (2017, Figure 2) used L-moment diagrams for complete and synthetic
301 series of daily streamflow observations to demonstrate that the sampling variability in L-
302 moment ratios is negligible for the sample sizes considered in this study. Thus, the
303 scatter shown in Figure 4 is likely due to real distributional differences rather than due to
304 sampling variability as is often the case when one constructs L-moment diagrams for
305 short AMS precipitation and streamflow records, as is the case in most previous studies
306 which have employed L-moment ratio diagrams.

307 **4.1.2 L-Kurtosis vs L-Skew**

308 Figure 5 displays empirical and theoretical distributional relationships between L-
309 Kurtosis vs L-Skew point values of daily precipitation (Figure 5a) and areal average
310 values of daily precipitation (Figure 5b). The empirical relationships of plotted points for
311 both wet-day series are very similar to the theoretical relationship for the P3 distribution.
312 In fact, among the pdfs considered in Figure 5, the P3 pdf seems to be the only 3-
313 parameter distribution that could possibly fit the wet-day record data. Although there is a
314 small proportion of points lying outside the P3 curve, the overall fit is still very striking.

315 It should also be noted that the L-moment ratio estimates for both wet-day series
316 occupy a space that can be well represented by the KAP distribution, which occupies a
317 region of the L-Kurtosis vs L-Skew diagram as shown in Figure A1 of Hosking and
318 Wallis (1997). A complete description of the 4-parameter KAP distribution is referred to
319 Hosking (1994) and Hosking and Wallis (1997).

320 *[Figure 5 goes here]*

321 **4.2 PPCC**

322 **4.2.1 Standard boxplots of PPCC**

323 The L-moment ratio diagrams were useful for identifying several potential candidate
324 distributions for representing the wet-day daily precipitation series at the point and
325 catchment scales. From that analysis we conclude that a four parameter Kappa pdf is
326 needed to approximate the pdf of point wet-day series whereas a G2 and P3 pdf are
327 adequate to approximate the pdf of areal average wet-day series. The PPCC statistic
328 offers another quantitative method for comparing the goodness of fit of different
329 distributions to the daily precipitation observations. Table 5 summarizes the central
330 tendency and spread of the values of PPCC for each of the distributions for the wet-day
331 series of point and catchment scale daily precipitation, respectively. The highest values
332 for the mean, median, 95th percentile, and 5th percentile of the PPCC are shown in bold
333 type. The lowest values of the sample standard deviation of the PPCC values, denoted $\hat{\sigma}$,
334 are also shown in bold. Figure 6 illustrates box-plots of the values of PPCC for
335 distributions fitted to the wet-day series of daily precipitation data at the point and
336 catchment scales.

337 *[Table 5 goes here]*

338 *[Figure 6 goes here]*

339 Figure 6 and Table 5 indicate that for the wet-day-series of point daily
340 precipitation depths, all the distributions have median PPCCs well above 0.9, but only the
341 median PPCCs of G2, P3, and KAP distributions are over 0.99. The same situation
342 appears in the catchment scale precipitation, except that the median PPCCs of the
343 remaining four distributions are significantly lower than the corresponding values for
344 point precipitation.

345 The insets in Figure 6 show detailed views of the boxplots of PPCC values for the
346 G2, P3, and KAP distributions for point and areal average daily precipitation. From
347 Figure 6a, KAP distribution results in the best goodness-of-fit for point precipitation
348 because all of its indices are the best, while the P3 distribution generally performs better
349 than the G2 distribution. However, for catchment-scale precipitation (Figure 6b), the
350 four parameter KAP distribution is no longer competitive, and both the G2 and P3 pdfs
351 will suffice. We are reluctant to advocate the use of a four parameter pdf, such as the
352 KAP distribution, due to its inherent complexity, though such a pdf may be needed for
353 point values as evidenced from our analyses.

354 **4.2.2 Graphical comparison of P3, G2, and KAP**

355 Across all previous comparisons, the P3, G2, and KAP are the best fitting distributions
356 for describing daily precipitation at the point or catchment scales. The insets in Figure 6
357 identify the distributions that exhibit the best fit to the each observed series. However,
358 these inserts do not indicate by how much the best performing distribution outperforms
359 the second or third best. For this purpose, pairwise comparisons of the PPCC values of
360 two highly performing distributions for all the stations and catchments are instructive. A
361 simple graphical method can accomplish this goal.

362 Figure 7 compares the PPCC values of the P3 (vertical axis) and G2 (horizontal
363 axis) distributions for point- and catchment-scale daily precipitation. Approximately
364 98% of stations are displayed on the figure; the remaining points lie outside the plot
365 domains. Points lying above the diagonal line indicate that the P3 distribution has a
366 higher PPCC for that particular station, and points lying below the diagonal line indicate
367 the G2 results in a higher PPCC. Figure 7a shows that in nearly every case, the P3
368 distribution outperforms the G2 distribution. When the G2 does outperform the P3, the
369 PPCCs are both very high and nearly equal. The point-scale precipitation plot shows that
370 the P3 distribution performs significantly better than the G2 distribution in many cases.
371 Thus, we conclude the P3 distribution better represents wet-day daily point precipitation
372 than the more commonly used G2 distribution in nearly every case. Figure 7b compares
373 the PPCC values of P3 and G2 for the catchment-scale precipitation. The results are
374 nearly the same as for the point-scale precipitation in the sense that most points are above
375 the diagonal line; while, for a few catchments where G2 does outperform P3, the points
376 lie on the dividing line, showing only very slight superiority.

377 *[Figure 7 goes here]*

378 Figure 8 displays similar plots comparing the KAP (vertical axis) and P3
379 (horizontal axis) distribution for point- and catchment-scale daily precipitation. It can be
380 seen in Figure 8a that the KAP distribution does not always outperform the P3 pdf, as one
381 might expect given that it has an additional parameter. We are reluctant to advocate the

382 KAP pdf given its additional model complexity combined with the fact that it does not
383 appear to provide a uniform improvement, in either case, over the P3 pdf.

384 *[Figure 8 goes here]*

385

386 **5. Discussion**

387 From the L-moment diagrams and PPCC comparisons we concluded that a KAP pdf is
388 required to fully capture the tail behavior of point wet-day series, though both P3 and G2
389 can provide reasonable approximations in many situations. In contrast, we found that a
390 KAP pdf is not needed to approximate the behavior of areal average wet-day series,
391 where instead, either a P3 or G2 model would suffice. In this section we evaluate the
392 relationship between these findings and the size of the catchments considered.

393 Figure 9 displays the PPCC values of P3 and G2 pdfs versus catchment drainage
394 area for areal average wet-day series. The PPCC values are chosen from 0.99-1,
395 approximately 96% of catchments are displayed on the figure; the remaining points lie
396 outside the plot domains. It can be seen that for most of the catchments, the PPCC values
397 for G2 and P3 pdfs are very close, with points corresponding to G2 and P3 pdfs almost
398 overlapping. This is especially true for PPCC values higher than 0.998. The phenomena
399 clearly indicates that when G2 can well represent the behavior of catchment-scale wet-
400 day precipitation series, P3 also provides very good performance. However, for the areas
401 where PPCC values are lower than 0.996, the P3 distribution outperforms the G2
402 distribution for most cases, with a very slight improvement.

403 *[Figure 9 goes here]*

404 Figure 10 shows the spatial map of catchments with the corresponding best
405 distribution functions for areal average wet-day series. KAP distribution is the best pdf
406 for large proportion of the catchments especially in the middle of U.S. P3 distribution
407 occupies the second large proportion of the catchments especially in east-central U.S.
408 Only a very few catchments can be best represented by G2 distribution. Seen from
409 Figure 10, it seems that the performances of the three pdfs vary greatly. However, as we
410 have seen from previous figures, the differences between the three pdfs for catchments
411 are very small.

412 *[Figure 10 goes here]*

413 **6. Conclusions**

414 This study has demonstrated that L-moment diagrams and probability plot correlation
415 coefficient goodness of fit evaluations can provide new insight into the distribution of
416 very long series of daily wet-day precipitation at both the point and catchment scales.
417 Although previous studies have claimed that the commonly used 2-parameter Gamma
418 distribution performs fairly well on the basis of traditional goodness-of-fit tests, this
419 study reveals, through the use of L-moment diagrams and probability plot correlation
420 coefficient goodness of fit evaluations that very long series of uncensored daily point and
421 areal average precipitation are better approximated by a KAP distribution and a Pearson-
422 III distribution respectively, and importantly, they do not resemble any of the other

423 commonly used distributions. Analogous to the recent study by Papalexiou and
424 Koutsoyiannis (2016), our evaluations yield very different conclusions than previous
425 research on this subject and thus could have important implications in climate change
426 investigations and other studies which employ a pdf of daily precipitation.

427 We conclude that for representing wet-day precipitation, the Gamma and Pearson-
428 III distributions are comparable with the 4-parameter Kappa distribution for the areal
429 average precipitation; however, when the point precipitation is of concern, the Kappa
430 distribution should be the distribution of choice. We also conclude that future
431 investigations should consider comparisons between the generalized Gamma distribution
432 introduced by Papalexiou and Koutsoyiannis (2012, 2016) for wet-day daily precipitation
433 and the G2, Pearson type III and Kappa distributions recommended here.

434 Once analytical and polynomial L-moment relationships and parameter estimation
435 methods become available for the GG distribution, future studies should compare the P3
436 and GG distributions on wet-day series, because on the basis of this study, and
437 Papalexiou and Koutsoyiannis (2016), the P3 and GG distributions appear to have
438 tremendous potential for approximating the distribution of wet-day series.

439

440

441 **Acknowledgements**

442 The first and third authors are partially supported by the National Natural Science
443 Foundation of China (No. 91547116, 51709033, 91647201). Special thanks are given to
444 Dr. Simon M. Papalexiou and other two anonymous reviewers and editors for their
445 constructive remarks.
446

447 **References**

- 448 Blum, A. G., Archfield, S. A., and Vogel, R. M.: On the probability distribution of daily
449 streamflow in the United States, *Hydrology and Earth System Sciences*, 21, 3093, 2017.
- 450 Bonnin, G. M., Martin, D., Lin, B., Parzybok, T., Yekta, M., and Riley, D.: Precipitation-
451 frequency atlas of the United States, NOAA atlas, 14, 2006.
- 452 Buishand, T. A.: Some remarks on the use of daily rainfall models, *Journal of Hydrology*,
453 36, 295-308, 1978.
- 454 Burgueno, A., Martinez, M. D., Lana, X., and Serra, C.: Statistical distributions of the
455 daily rainfall regime in Catalonia (northeastern Spain) for the years 1950-2000,
456 *International Journal of Climatology*, 25, 1381-1403, 2005.
- 457 Chen, J., and Brissette, F. P.: Stochastic generation of daily precipitation amounts: review
458 and evaluation of different models, *Climate Research*, 59, 189-206, 2014.
- 459 Deidda, R., and Puliga, M.: Sensitivity of goodness-of-fit statistics to rainfall data
460 rounding off, *Physics and Chemistry of the Earth, Parts A/B/C*, 31, 1240-1251, 2006.
- 461 Duan, J., Sikka, A. K., and Grant, G. E.: A comparison of stochastic models for
462 generating daily precipitation at the HJ Andrews Experimental Forest, 1995.
- 463 Duan, Q., Schaake, J., Andreassian, V., Franks, S., Goteti, G., Gupta, H. V., Gusev, Y.
464 M., Habets, F., Hall, A., and Hay, L.: Model Parameter Estimation Experiment (MOPEX):
465 An overview of science strategy and major results from the second and third workshops,
466 *Journal of Hydrology*, 320, 3-17, 2006.
- 467 Easterling, D. R., Evans, J., Groisman, P. Y., Karl, T. R., Kunkel, K. E., and Ambenje, P.:
468 Observed variability and trends in extreme climate events: a brief review, *Bulletin of the
469 American Meteorological Society*, 81, 417-425, 2000.
- 470 Geng, S., de Vries, F. W. P., and Supit, I.: A simple method for generating daily rainfall
471 data, *Agricultural and Forest Meteorology*, 36, 363-376, 1986.
- 472 Groisman, P. Y., Karl, T. R., Easterling, D. R., Knight, R. W., Jamason, P. F., Hennessy,
473 K. J., Suppiah, R., Page, C. M., Wibig, J., and Fortuniak, K.: Changes in the probability
474 of heavy precipitation: important indicators of climatic change, in: *Weather and Climate
475 Extremes*, Springer, 243-283, 1999.
- 476 Hershfield, D. M.: Rainfall frequency atlas of the United States for durations from 30
477 minutes to 24 hours and return periods from 1 to 100 years, 1961.

478 Hosking, J. R.: L-moments: analysis and estimation of distributions using linear
479 combinations of order statistics, *Journal of the Royal Statistical Society. Series B*
480 (Methodological), 105-124, 1990.

481 Hosking, J. R.: The four-parameter kappa distribution, *IBM Journal of Research and*
482 *Development*, 38, 251-258, 1994.

483 Hosking, J. R. M., and Wallis, J. R.: *Regional frequency analysis : an approach based on*
484 *L-moments*, Cambridge University Press, 1997.

485 Karl, T. R., and Knight, R. W.: Secular trends of precipitation amount, frequency, and
486 intensity in the United States, *Bulletin of the American Meteorological society*, 79, 231-
487 241, 1998.

488 Kigobe, M., McIntyre, N., Wheeler, H., and Chandler, R.. *Multi-site stochastic modelling*
489 *of daily rainfall in Uganda. Hydrological sciences journal*, 56, 17-33, 2011.

490 Lee, S. H., and Maeng, S. J.: Frequency analysis of extreme rainfall using L moment,
491 *Irrigation and Drainage*, 52, 219-230, 2003.

492 Li, Z., Brissette, F., Chen, J.. *Finding the most appropriate precipitation probability*
493 *distribution for stochastic weather generation and hydrological modelling in Nordic*
494 *watersheds. Hydrological Processes*, 27: 3718-3729, 2013.

495 Madsen, H., Lawrence, D., Lang, M., Martinkova, M., and Kjeldsen, T.: Review of trend
496 analysis and climate change projections of extreme precipitation and floods in Europe,
497 *Journal of Hydrology*, 519, 3634-3650, 2014.

498 Mehrotra, R., Srikanthan, R., and Sharma, A.: A comparison of three stochastic multi-site
499 precipitation occurrence generators, *Journal of Hydrology*, 331, 280-292, 2006.

500 Naghavi, B., and Yu, F. X.: Regional frequency analysis of extreme precipitation in
501 Louisiana, *Journal of Hydraulic Engineering*, 121, 819-827, 1995.

502 Papalexiou, S. M., and Koutsoyiannis, D.: Entropy based derivation of probability
503 distributions: A case study to daily rainfall, *Advances in Water Resources*, 45, 51-57,
504 2012.

505 Papalexiou, S. M., and Koutsoyiannis, D.: Battle of extreme value distributions: A global
506 survey on extreme daily rainfall, *Water Resources Research*, 49, 187-201, 2013.

507 Papalexiou, S. M., and Koutsoyiannis, D.: A global survey on the seasonal variation of
508 the marginal distribution of daily precipitation, *Advances in Water Resources*, 94, 131-
509 145, 2016.

510 Park, J.-S., and Jung, H.-S.: Modelling Korean extreme rainfall using a Kappa
511 distribution and maximum likelihood estimate, *Theoretical and Applied climatology*, 72,
512 55-64, 2002.

513 Pilon, P. J., Adamowski, K., and Alila, Y.: Regional analysis of annual maxima
514 precipitation using L-moments, *Atmospheric Research*, 27, 81-92, 1991.

515 Schoof, J. T., Pryor, S. C., and Surprenant, J.: *Development of daily precipitation*
516 *projections for the United States based on probabilistic downscaling, Journal of*
517 *Geophysical Research: Atmospheres*, 115, D13, 2010.

518

519 Shoji, T., and Kitaura, H.: Statistical and geostatistical analysis of rainfall in central Japan,
520 Computers & Geosciences, 32, 1007-1024, 2006.

521 Srikanthan, R., and McMahon, T.: Stochastic generation of annual, monthly and daily
522 climate data: A review, Hydrology and Earth System Sciences Discussions, 5, 653-670,
523 2001.

524 Stedinger, J. R., R.M. Vogel and E. Foufoula-Georgiou: Frequency analysis of extreme
525 events, Handbook of Hydrology, Chapter 18, McGraw Hill Book Co, D.R. Maidment -
526 editor in chief, 1993..

527 Thom, H. C.: A frequency distribution for precipitation, Bulletin of the American
528 Meteorological Society, 32, 397, 1951.

529 Trenberth, K. E.: Changes in precipitation with climate change, Climate Research, 47,
530 123-138, 2011.

531 Vogel, R. M., and Fennessey, N. M.: L moment diagrams should replace product moment
532 diagrams, Water Resources Research, 29, 1745-1752, 1993.

533 Vogel, R. W., and McMartin, D. E.: Probability Plot Goodness-of-Fit and Skewness
534 Estimation Procedures for the Pearson Type 3 Distribution, Water resources research, 27,
535 3149-3158, 1991.

536 Waggoner, P.E., 1989: Anticipating the frequency distribution of precipitation if climate
537 change alters its mean, Agricultural and Forest Meteorology, 47, 321 – 337.

538 Watterson, I., and Dix, M.: Simulated changes due to global warming in daily
539 precipitation means and extremes and their interpretation using the gamma distribution,
540 Journal of Geophysical Research: Atmospheres, 108, 2003.

541 Watterson, I. G.: Simulated changes due to global warming in the variability of
542 precipitation, and their interpretation using a gamma-distributed stochastic model,
543 Advances in Water Resources, 28, 1368-1381, 2005.

544 Waymire, E., and Gupta, V. K.: The mathematical structure of rainfall representations: 1.
545 A review of the stochastic rainfall models, Water resources research, 17, 1261-1272,
546 1981.

547 Wilby, R. L., and Wigley, T.: Future changes in the distribution of daily precipitation
548 totals across North America, Geophysical Research Letters, 29, 2002.

549 Wilks, D. S.: Maximum likelihood estimation for the gamma distribution using data
550 containing zeros, Journal of Climate, 3, 1495-1501, 1990.

551 Wilks, D. S.: Multisite generalization of a daily stochastic precipitation generation model,
552 Journal of Hydrology, 210, 178-191, 1998.

553 Wilks, D. S., and Wilby, R. L.: The weather generation game: a review of stochastic
554 weather models, Progress in physical geography, 23, 329-357, 1999.

555 **Wilson, P. S., and Toumi, R.: A fundamental probability distribution for heavy rainfall.**
556 **Geophysical Research Letters, 32, L14812, 2005.**

557 Woolhiser, D. A., and Roldan, J.: Stochastic daily precipitation models: 2. A comparison
558 of distributions of amounts, *Water resources research*, 18, 1461-1468, 1982.

559 Yoo, C., Jung, K. S., and Kim, T. W.: Rainfall frequency analysis using a mixed Gamma
560 distribution: evaluation of the global warming effect on daily rainfall, *Hydrological*
561 *Processes*, 19, 3851-3861, 2005.

562

563

564 **Table captions:**

565 **Table 1:** Review of literature pertinent to daily precipitation probability distribution
566 selection.

567 **Table 2:** Table 2: Theoretical probability distributions presented on the L-Kurtosis vs L-
568 Skew L-moment diagram. *Italicized distributions are special cases of other distributions.*

569 **Table 3:** Theoretical probability distributions presented on the L-Cv vs L-Skew L-
570 moment diagram.

571 **Table 4:** Distributions used in probability plot goodness of fit evaluations.

572 **Table 5:** Central tendency and spread of values of PPCC for the 237 precipitation
573 stations 305 areal average precipitation catchments.

574 **Figure captions:**

575 **Figure 1:** Map showing locations of a) 237 precipitation gaging stations, and b) 305
576 catchments.

577 **Figure 2:** Distribution of full record length of point precipitation base on weather stations.

578 **Figure 3:** Distribution of wet-day record length: a) point precipitation; and b) areal
579 average precipitation over watersheds. Days with zero precipitation are removed in the
580 wet-day records

581 **Figure 4:** L-Cv vs L-Skew L-moment ratio diagram of sample L-moments and
582 theoretical distributions: a) point precipitation; and b) areal average precipitation depths.

583 **Figure 5:** L-Skew vs L-Kurtosis L-moment ratio diagram of sample L-moments and
584 theoretical distributions: a) point precipitation; and b) areal average precipitation depths.
585 Logistic (L), Normal (N), Uniform (U), Gumbel (G), and Exponential (E) distributions
586 appear as a single point.

587 **Figure 6:** Standard boxplots of r for all 7 distributions evaluated for wet-day series of a)
588 point precipitation, and b) areal average precipitation depths.

589 **Figure 7:** Comparison of PPCC (r) values for the P3 (vertical axis) and G2 (horizontal
590 axis) distributions for the a) point, and b) areal average precipitation depths series. Points
591 lying above the line represent stations with a higher r for the P3 distribution than G2
592 distribution.

593 **Figure 8:** Comparison of r values for P3 (horizontal axis) and KAP (vertical axis)
594 distributions for the a) point and b) areal average precipitation depths' wet-day series.

595 **Figure 9:** The spatial distribution of best daily precipitation distribution function.

596 **Figure 10:** The spatial map of catchments with the corresponding best distribution
597 functions for areal average wet-day series.

598

599 **Tables**

600

Table 1: Review of literature pertinent to daily precipitation probability distribution selection.

| 1. Stochastic Precipitation Modelling: | | | | | | |
|--|------|----------------------|-------------------|----------------|----------------------|--|
| Author | Year | Stations | Series type | Duration | Distribution | Justification |
| Thom | 1951 | | Wet-day | 1-day | Gamma | |
| Buishand | 1978 | 6 | Wet-day | 1-day | Gamma | Cv-Cs ratio |
| Geng et al | 1986 | 6 | Wet-day, by month | 1-day, monthly | Gamma | Regress. fit: β vs mean wet-day depth |
| Woolhiser and Roldan | 1982 | | Wet-day | 1-day | Mixed Exponential | MLE, Akaike Information Criterion |
| Duan et al | 1995 | 1 | Wet-day, by month | 1-day | Calib. W2, Gamma | MLE, Chi-sq test |
| Wilks | 1998 | 25 | Wet-day | 1-day | Mixed Exponential | MLE, goodness of fit |
| Waterson and Dix | 2003 | | Wet-day | 1-day | Gamma | Literature |
| Burgueno et al | 2005 | 75 | Wet-day | 1-day | Exponential, Weibull | Normalized Rainfall Curve |
| Kigobe et al | 2011 | 110 | Wet-day, by month | | Gamma | |
| Li et al | 2013 | 24 | Wet-day | 1-day | Mixed Exponential | Goodness of fit and Kolmogorov–Smirnov tests |
| Schoof | 2010 | Grided precipitation | Wet-day | 1-day | Gamma | Goodness of fit |

| 2. Precipitation Frequency Analysis | | | | | | |
|-------------------------------------|------|----------|---------------------------|------------------------|-------------------------------------|--|
| Author | Year | Stations | Series type | Duration | Distribution | Justification |
| Hershfield (TP-40) | 1962 | | AMS | 24 hour | Gumbel | |
| Pilon et al | 1991 | 75 | AMS | 5 min - 24 hour | GEV | L-moments |
| Naghavi & Yu | 1995 | 25 | AMS | 1-24 hour | GEV | L-moments, PWMs, Monte Carlo experiments |
| Park and Jung | 2002 | 61 | AMS | 1, 2-day | Kappa(4) | |
| Lee and Maeng | 2003 | 38 | AMS | 1-day | GEV, GLO | L-moments |
| Bonnin et al | 2006 | | AMS | 5 min - 24 hour | GEV | L-moments |
| Shoji and Kitaura | 2006 | 243 | Complete, Wet-day | Hour, Day, Month, Year | Lognormal, Weibull | Goodness of fit |
| Deidda and Puliga | 2006 | 200 | Left Censored Wet-day PDS | 1-day | Generalized Pareto | “Failure-to-reject” method, L-moments |
| Wilson and Toumi | 2005 | 270 | Complete | 1-day | Self-derived | |
| Papalexiou and Koutsoyiannis | 2012 | 11,519 | Wet-Day | 1-day | Generalized Gamma | L-moments |
| Papalexiou and Koutsoyiannis | 2013 | 15,137 | AMS | 1-day | GEV | L-moments |
| Papalexiou and Koutsoyiannis, | 2016 | 14,157 | Wet-Day, by month | 1-day | Generalized Gamma and Burr type XII | L-moments and Goodness-of-fit |

3. *Precipitation Trends and Climate Change*

| Author | Year | Stations | Series type | Duration | Distribution | Justification |
|------------------|-------------|-----------------|--------------------|------------------------|---------------------|--|
| Waggoner | 1989 | 55 | Monthly | 1-month | Gamma | Literature Review |
| Groisman et al | 1999 | 1313 | Summer (wet-day) | 1-day | Gamma | Literature Review, goodness of fit to extreme rainfall quantiles |
| Wilby and Wigley | 2002 | GCM | Seasonal | 1-day | Gamma | Literature Review |
| Yoo et al | 2005 | 31 | Monthly (wet-day) | 1-day | Gamma | Literature Review |
| Watterson | 2005 | GCM | January, July | 1-month (daily forced) | Gamma | Literature Review |

Table 2: Theoretical probability distributions presented on the L-Kurtosis vs L-Skew L-moment diagram. *Italicized distributions are special cases of other distributions.*

| Distribution | Abbreviation | PDF | Parameters |
|------------------------------------|--------------|---|------------|
| Generalized Extreme Value Type III | GEV | $f(x) = \frac{1}{\eta} \left[1 - \left(\frac{x-\omega}{\eta} \right) \beta \right]^{\frac{1}{\beta}-1} \exp \left\{ - \left[1 - \left(\frac{x-\omega}{\eta} \right) \beta \right]^{\frac{1}{\beta}} \right\}$ | 3 |
| Generalized Logistic | GLO | $f(x) = \frac{be^{-\frac{x-\mu}{\sigma}}}{\sigma \left(1 + e^{-\frac{x-\mu}{\sigma}} \right)^{b+1}}$ | 3 |
| Generalized Pareto | GPA | $f(x) = \frac{1}{\sigma} \left(1 + \frac{\xi(x-\mu)}{\sigma} \right)^{(-1/\xi-1)}$ | 3 |
| Lognormal | LN3 | $f(x) = \frac{1}{\sqrt{2\pi}(x-\alpha)\sigma} \exp \left[-\frac{1}{2\sigma^2} (\ln(x-\alpha) - \mu^2) \right]$ | 3 |
| Pearson Type III | P3 | $f(x) = \frac{p/a^d}{\Gamma(d/p)} x^{d-1} e^{-(x/a)^p}$ | 3 |
| <i>Exponential</i> | E | $f(x) = \begin{cases} \lambda e^{-\lambda x}, & x \geq 0 \\ 0, & x < 0 \end{cases}$ | 2 |
| <i>Gumbel</i> | G | $f(x) = \frac{1}{\beta} e^{-(z+e^{-z})}, z = \frac{x-\mu}{\beta}$ | 2 |
| <i>Normal</i> | N | $f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$ | 2 |
| <i>Logistic</i> | L | $f(x) = \frac{e^{-\frac{x-\mu}{s}}}{s \left(1 + e^{-\frac{x-\mu}{s}} \right)^2}$ | 2 |
| <i>Uniform</i> | U | $f(x) = \begin{cases} \frac{1}{b-a}, & a < x < b \\ 0, & x < a \text{ or } x > b \end{cases}$ | 1 |

Table 3: Theoretical probability distributions presented on the L-Cv vs L-Skew L-moment diagram.

| Distribution | Abbreviation | PDF | Parameters |
|--------------------|--------------|--|------------|
| Gamma | G2 | $f(x) = \frac{\beta^\alpha x^{\alpha-1} e^{-\beta x}}{\Gamma(\alpha)}$ | 2 |
| Generalized Pareto | GP2 | $f(x) = \frac{1}{\sigma} \left(1 + \frac{\xi x}{\sigma} \right)^{(-1/\xi-1)}$ | 2 |
| Lognormal | LN2 | $f(x) = \frac{1}{x\sigma\sqrt{2\pi}} e^{-\frac{(\ln x - \mu)^2}{2\sigma^2}}$ | 2 |

| | | | |
|---------|----|---|---|
| Weibull | W2 | $f(x) = \frac{k}{\phi} \left(\frac{x}{\phi}\right)^{k-1} \exp\left\{-\left[\frac{x}{\phi}\right]^k\right\}$ | 2 |
|---------|----|---|---|

Table 4: Distributions used in probability plot goodness of fit evaluations.

| Distribution | Abbreviation | Parameters |
|------------------------------------|--------------|------------|
| Generalized Extreme Value Type III | GEV | 3 |
| Generalized Logistic | GLO | 3 |
| Generalized Pareto | GPA | 3 |
| Lognormal | LN3 | 3 |
| Pearson Type III | P3 | 3 |
| Gamma | G2 | 2 |
| Kappa | KAP | 4 |

Table 5: Central tendency and spread of values of PPCC for the 237 precipitation stations and 305 catchments.

| Distribution | Point precipitation | | | Percentiles | | Areal average precipitation | | | Percentiles | |
|--------------|---------------------|---------------|---------------|---------------|---------------|-----------------------------|---------------|---------------|---------------|---------------|
| | Mean | Median | \hat{s} | 95th | 5th | Mean | Median | \hat{s} | 95th | 5th |
| P3 | 0.9952 | 0.9971 | 0.0063 | 0.9995 | 0.9872 | 0.9977 | 0.9985 | 0.0028 | 0.9996 | 0.9936 |
| GEV | 0.9338 | 0.9375 | 0.0222 | 0.9609 | 0.8944 | 0.8003 | 0.7965 | 0.0474 | 0.8917 | 0.7264 |
| GPA | 0.9793 | 0.9828 | 0.0145 | 0.9949 | 0.9500 | 0.8688 | 0.8687 | 0.0484 | 0.9586 | 0.7894 |
| GLO | 0.9115 | 0.9154 | 0.0235 | 0.9423 | 0.8734 | 0.7800 | 0.7750 | 0.0441 | 0.8669 | 0.7101 |
| LN3 | 0.9838 | 0.9855 | 0.0075 | 0.9924 | 0.9727 | 0.9362 | 0.9373 | 0.0224 | 0.9737 | 0.8983 |
| G2 | 0.9925 | 0.9949 | 0.0079 | 0.9990 | 0.9789 | 0.9974 | 0.9985 | 0.0034 | 0.9996 | 0.9924 |
| KAP | 0.9971 | 0.9985 | 0.0048 | 0.9997 | 0.9915 | 0.9976 | 0.9987 | 0.0026 | 0.9998 | 0.9929 |

Figures




Figure 1: Map showing locations of a) 237 point precipitation gaging stations, and b) 305 MOPEX catchments.




Figure 2: Distribution of length of records of point daily precipitation data for the 237 gaging stations depicted in Figure 1a.




Figure 3: Distribution of wet-day record lengths corresponding to the two datasets: a) point precipitation; and b) areal average precipitation over catchments. Days with zero precipitation are removed in the wet-day records




Figure 4: L-Cv vs L-Skew L-moment ratio diagram of sample L-moments and theoretical distributions for: a) point daily precipitation; and b) areal average daily precipitation depths.




Figure 5: L-Skew vs L-Kurtosis L-moment ratio diagram of sample L-moments and theoretical distributions for: a) point daily precipitation; and b) areal average daily precipitation depths. Note that Logistic (L), Normal (N), Uniform (U), Gumbel (G), and Exponential (E) distributions appear as a single point.





Figure 6: Standard boxplots of r for all 7 distributions evaluated for wet-day series of a) point precipitation, and b) areal average precipitation depths.




Figure 7: Comparison of PPCC (r) values for the P3 (vertical axis) and G2 (horizontal axis) distributions for the a) point, and b) areal average precipitation depths series. Points lying above the line represent stations with a higher r for the P3 distribution than G2 distribution.




Figure 8: Comparison of r values for P3 (horizontal axis) and KAP (vertical axis) distributions for the a) point and b) areal average precipitation depths' wet-day series.




Figure 9: the PPCC values of P3 and G2 pdfs versus catchment drainage area for areal average wet-day series.




Figure 10: The spatial map of catchments with the corresponding best distribution functions for areal average wet-day series.