

Interactive comment on “Multimodel simulation of vertical gas transfer in a temperate lake” by Sofya Guseva et al.

Sofya Guseva et al.

guseva@uni-landau.de

Received and published: 11 November 2019

Dear Referees and Editor,

we highly appreciate your time and work in revising our manuscript. We thank the Referees for the very important and helpful comments which helped us to find the mistakes and improve the modeling results. We went carefully through all the suggestions/comments/corrections and respond to them at our best.

Kind regards, Authors

Interactive comment on Hydrol. Earth Syst. Sci. Discuss., <https://doi.org/10.5194/hess-2019-146>, 2019.

[Printer-friendly version](#)

[Discussion paper](#)



Interactive comment on “Multimodel simulation of vertical gas transfer in a temperate lake” by Sofya Guseva et al.

Sofya Guseva et al.

guseva@uni-landau.de

Received and published: 5 November 2019

1. (1) Concerns arise however about the way one of the models - FLake - was treated in the study. The authors correctly state in the description of the model experiments that FLake “stands aside from the other 1-D models due to the . . . bulk-structure which employs the concept of <temperature profile> self-similarity. . . ”. This high level of parameterization ensures computational efficiency of the model, which was primarily designed for prediction of surface temperatures in global/regional climate models and numerical weather prediction (NWP). On the other hand, the model parameterizations put some constraints on the model application to real lakes. One crucially important feature of FLake is that the model equations are derived in the assumption of preserving the heat capacity or volume of the lake. In this regard, the “baseline” configuration

[Printer-friendly version](#)

[Discussion paper](#)



applied in this study with the maximum lake depth as model input is inappropriate for FLake and would produce a priori incorrect results. Another issue of the parameterized model refers to correct choice of the few “shape factor” parameters determining the stratification pattern. The latter is not resolved in FLake numerically, but parameterized via few “shape factor”-constants related to the spatial integrals over the stratified layer. Several recent publications, including co-authorship of one of the authors of the present study, (Shatwell et al., 2016; Kirillin et al., 2017; Shatwell et al., 2019; Su et al., 2019) discussed the appropriate choice of the shape factor constants in FLake and have demonstrated that the set of constants used in the NWP-version of the model should be amended if the vertical thermal structure is in question apart from the lake surface temperatures. In particular, the unrealistically weak deep stratification and the corresponding high depth of the surface mixed layer, as reported in the present study, are the results of applying the NWP-constants together with the maximum lake depth as the model lake depth. Hence, the FLake-outcomes discussed here are useless or even misleading for the potential FLake-users. I see two ways of possible modification of the study: (i) excluding FLake from the study completely, confining the set of tested models to four models; (ii) re-designing the FLake-experiments in the corrected way with corresponding changes in the results/discussion.

(2) We generally agree with these suggestions of the referee and have re-designed experiments with FLake model testing the sensitivity of the model to one of the constants determining "shape factor" – dimensionless relaxation constant C_{rc} . In the current study, we have got the best agreement with the measured temperature profile using the value $C_{rc} = 0.3$ (in comparison with the standard value of 0.003) and chose this setting for the baseline experiment RefSim. The model now reproduces the temperature profile much better and we conducted other experiments including the one with mean lake depth and varying light attenuation coefficient with $C_{rc} = 0.3$.

(3) Page 7 line 15: The concept of the self-similarity in the FLake model (Sect. 2.2) includes the "shape factor" coefficient C_{θ} , which determines the vertical temperature

[Printer-friendly version](#)

[Discussion paper](#)



profile below the mixed layer. Evolution of this parameter is controlled by a relaxation time scale t_{rc} (Mironov, 2008). This time scale includes the dimensionless relaxation constant C_{rc} having a default value of 0.003 in the model. This constant is a calibration parameter which is generally individual for each lake (Shatwell et al., 2016, Kirillin et al., 2017). We tested the sensitivity of the model to the variation of C_{rc} using values 0.03, 0.3, 0.5, 1, 2, 3, 30 and achieved the best agreement with the measured temperature profile using $C_{rc} = 0.3$ (for details, see Sect. 3.1). In the following sections, this C_{rc} setting is used in RefSim and other experiments with FLake model.

Page 10 line 11: In RefSim we test the model with different values of C_{rc} (see Sect. 2.3 above). The best agreement with observed temperature profile we get using $C_{rc} = 0.3$. In particular, the error (RMSE_c) for surface temperature reduces from 3.4 ° C (using standard value of $c_{relax} = 0.003$) to 2.5 ° and becomes close to other models. At the depth of ~ 6-10 m RMSE_c reduces from 5.7 ° C (using standard value of $c_{relax} = 0.003$) to 5.2 °. In addition, RMSE_c reduces from 3.3-0.7 ° C to 1.8-0.2 ° C at the depths 20-27 m.

We added new figures with new results for FLake model to the main manuscript: Fig. 2, Fig. 3, Fig. 4 ; add new figures to SI: Fig. S2; changed metrics for FLake model performance in Tables: S2, S3, S4.

2. (1) Regarding the biogeochemical part, the study would benefit from an extension of the discussion on possible ways of improving the representation of biogeochemistry, in particular, the deep oxygen (chlorophyll) maximum in oligotrophic lakes, and the vertical distribution of carbon dioxide across the water column.

(2)-(3) Page 19 line 34: Successful representation of maximal O₂ content below mixed layer during summer may be an important modelling skill for simulating correctly CH₄ in a lake because the former acts as a sink region for the latter. Furthermore, realistic oxygen concentration reduction during periods of stable stratification means that the respective CO₂ production from aerobic organic matter decomposition is reproduced

Printer-friendly version

Discussion paper



reasonably as well.

Page 21 line 21: As stated in the previous section, a realistic decay of oxygen content during stratified periods in ALBM and LAKE models suggests that CO₂ amount produced by aerobic decomposition of organic matter both in water column and in the top part of sediments is reasonably simulated as well. Satisfactory agreement of computed oxygen in the mixed layer and below with observations implies that photosynthesis minus respiration rate is fairly captured as well, and so do the models for CO₂ gain or loss due to these processes. Hence, the primary drawback of the models used in respect to DIC simulation is likely to be not explicitly simulating transport of carbon species from catchment to a water body. Thus, modelling approaches coupling the catchment and a lake presented recently (Futter et al., 2008; Duffy et al., 2018; McCullough et al., 2018) should be elaborated and wider used.

3. (1) Note also that the temperature profile within the ice cover in FLake is not assumed to be linear, but parameterized via a time-varying shape-function with a linear asymptotic.

(2) Yes, that is correct.

(3) Page 5 line 20: In FLake model, the temperature profile within the ice cover is parameterized via a time-varying shape-function having a linear asymptotic.

References: 1. Duffy C.J., Dugan H.A., Hanson P.C.: The age of water and carbon in lake-catchments: A simple dynamical model, *Limnol. Oceanogr. Lett.*, Vol. 3, No. 3, pp. 236–245, <https://doi.org/10.1002/lol2.10070>, 2018.

2. Futter M.N., Starr M., Forsius M., Holmberg M.: Modelling the effects of climate on long-term patterns of dissolved organic carbon concentrations in the surface waters of a boreal catchment, *Hydrol. Earth Syst. Sci.*, Vol. 12, No. 2, pp. 437–447, <https://doi.org/10.5194/hess-12-437-2008>, 2008.

3. Kirillin, G., Wen, L., Shatwell, T.: Seasonal thermal regime and climatic trends

in lakes of the Tibetan highlands, *Hydrol. Earth Syst. Sci.*,21(4), 1895-1909, <https://doi.org/10.5194/hess-21-1895-2017>, 2017.

4. McCullough I.M., Dugan H.A., Farrell K.J., Morales-Williams A.M., Ouyang Z., Roberts D., Scordo F., Bartlett S.L., Burke S.M., DoubekJ.P., Krivak-Tetley F.E., Skaff N.K., Summers J.C., Weathers K.C., Hanson P.C.: Dynamic modeling of organiccarbon fates in lake ecosystems, *Ecol. Modell.* Vol. 386. pp. 71–82, <https://doi.org/10.1016/j.ecolmodel.2018.08.009>, 2018.

5. Mironov, D. V.: Parameterisation of lakes in numerical weather prediction. Description of a lake model, COSMO Technical Report 11,Deutscher Wetterdienst, Offenbach am Main, Germany, 2008. 6. Shatwell, T., Adrian, R., and Kirillin, G.: Planktonic events may cause polymictic-dimictic regime shifts in temperate lakes., *Sci. Rep.-UK*,256, 24361, <https://doi.org/10.1038/srep24361>, 2016.

Additional comment: Attached file contains all changes in the manuscript and SI mentioned above.

Please also note the supplement to this comment:

<https://www.hydrol-earth-syst-sci-discuss.net/hess-2019-146/hess-2019-146-AC1-supplement.zip>

Interactive comment on *Hydrol. Earth Syst. Sci. Discuss.*, <https://doi.org/10.5194/hess-2019-146>, 2019.

Printer-friendly version

Discussion paper



Interactive comment on “Multimodel simulation of vertical gas transfer in a temperate lake” by Sofya Guseva et al.

Sofya Guseva et al.

guseva@uni-landau.de

Received and published: 5 November 2019

General comments

1. (1) It is stated that the study is a continuity of LakeMIP exercises accounting for biogeochemical processes comparisons. Although the first part of the paper is dedicated to thermal stratification and ice cover study, and involves the five models, the fact that only two of them have the possibility to simulate O₂ and CO₂ dynamics, indicate that this is not a real LakeMIP type exercise, and therefore constitutes a limitation in my view to be considered as a true intercomparison model experiment for vertical gas transfer.

(2) Dissolved gas transport throughout the water column is mainly carried out by turbu-

[Printer-friendly version](#)

[Discussion paper](#)



lence. Despite the fact that only two of the models have an ability directly to reproduce the concentration the gases (oxygen and carbon dioxide) our numerical experiments with a passive tracer (which can be seen as a "prototype" of gas or other constituent) involving four models out of five showed the potential simulation of gases governed by seasonal stratification and ice cover. We can consider this experiment as the inter-comparison model experiment in respect to primary physical controls of gas dynamics. We consider the intercomparison experiment involving full biogeochemical models as a next step of this study.

(3) No comments/corrections are added to the manuscript.

2. (1) In the first LakeMIP exercise (Stepanenko et al., 2010) the sensitivity of lake depth has been studied and the experiment setup accounted for simulations with maximum depth, local depth and average depth. And this is crucial especially (maybe only) for FLake model as it was demonstrated that an average lake depth was necessary for FLake simulation in order to be conservative in terms of energy. In the current setup, the maximum lake depth is used for all models and this is contradictory with a correct use of FLake which should be run using the LDSim configuration. It could be interesting to compare in the same graph RefSim and LDSim at least for FLake.

(2) The main focus of our study is the vertical diffusion which is the main driver for the gas distribution. Of major interest are such gases as carbon dioxide and methane which are primarily produced in lake sediments. Setting the maximum lake depth becomes crucial in terms of the total vertical distance required for transport of bottom-originated gases to the atmosphere. For this reason, we conduct the baseline experiment with the maximum depth. We added new figures to the manuscript with results from the LDSim experiment – FLake showed the largest sensitivity to the variation of the lake depth among the other models: e.g. $RMSE_c$ for surface temperature as well as for the whole temperature profile reduced almost twofold (see Table S4, Fig. S3, S4). This suggests potential limitation of FLake for correctly simulating gas dynamics in deep lakes (given the biogeochemical block is added to the model).

[Printer-friendly version](#)

[Discussion paper](#)



(3) Page 10 line 24: Remarkably, FLake demonstrates the largest sensitivity among all models to the lake depth – $RMSE_c$ is reduced approximately twofold (from $2.5\text{ }^{\circ}\text{C}$ to $1.28\text{ }^{\circ}\text{C}$) (see SI, Table S4, Fig. S3, S4) when using a mean depth of 13.32 m instead of the maximal depth 37.5 m. The thickness of mixed layer depth reduces up to 4.5 m and have a better agreement with the observations in comparison with RefSim experiment. We added the mean depth of the mixed layer for all models in the LDSim experiment to the Table S2 in SI. We also put two additional figures for LDSim simulation to SI: Fig. S3, S4.

3. (1) As a consequence, the sensitivity test on light extinction for FLake is not relevant since the thermal profile cannot be well simulated, and therefore should be conducted with a depth of 13.32 m.

(2) We redid all FLake simulations using calibrated shape factor, and temperature profile is now much better reproduced compared to results presented in the first version of the manuscript. Correspondingly, the sensitivity test on light extinction coefficient with FLake demonstrates now result similar to other models (see new Fig. 4).

(3) Page 11 line 32: This leads to temperature change at respective depths up to $4\text{ }^{\circ}\text{C}$ in FLake model and $8\text{ }^{\circ}\text{C}$ in ALBM and LAKEoneD. Fig. 4 contains only 3 models with different types of turbulent closure because the effect of varying the light attenuation coefficient is very similar in models on the same type (e.g. $k-\epsilon$ models LAKEoneD and LAKE).

We also added the result of ExtMaxSim simulation for FLake in Fig. 4 in the manuscript.

Specific comments

4. (1) Looking at the bathymetry indicates strong gradients from the shoreline to the point of maximum depth. There are probably 3d-circulations that take place when dense waters flow along the bottom slopes. And these circulations are not accounted for in the 1d simulations. What is in your view the potential impact on these circulations

[Printer-friendly version](#)

[Discussion paper](#)



on the thermal stratification and ice cover? And on modeled vertical transport? It would be interesting to add a discussion on that particular point.

(2) 1D models have certain limitations when it comes to such a 3-dimensional system as lake. They do not capture all the lake mixing mechanisms such as density driven currents, which can be important under certain conditions. In general, these currents are very slow (only a few cm s^{-1}) (Bengtsson, 2012) and are not important in terms of the turbulence production. The strong contribution of these flows in the vertical exchange of deep and near-surface water was found for deep lakes (> 100 m) with an extended littoral zone, e.g. Lake Geneva and Lake Van (Fer et al., 2001, Fer et al. 2002, Kaden et al., 2010). For Harp Lake, in particular, this process may not be important during summer due to a smaller depth and a limited area of shallow part. In winter, a large-scale convective circulation (up to $3\text{-}5 \text{ cm s}^{-1}$) due to heat exchange at the sediment-water interface may develop under ice (Kirillin et al., 2015). However, the importance of this circulation in terms of gas transfer have not been studied yet. At the moment, this type of the lake circulation is a topic of the ongoing research and it is not included into 1D models. Modeling such 3-d phenomena requires an application of 2D or 3D models.

(3) Added to Page 24 line 22 : - 1D models have certain limitations when applying to such a 3-dimensional system as lake. They do not capture all the lake mixing mechanisms such as density driven currents, which can be important under certain conditions (Samolyubov,1999). It was found that for deep lakes with extended littoral zone, such as Lake Geneva (Fer et al., 2001, 2002) and Lake Van (Kaden et al., 2010), these flows can be significant in terms of the vertical heat and gas transfer. In particular, for Harp Lake it may not be important due to a smaller depth and a not large shallow area. In winter, a large-scale convective circulation (up to $3\text{-}5 \text{ cm s}^{-1}$) due to heat exchange at the sediment-water interface may develop under ice (Kirillin et al., 2015). However, the importance of this circulation in terms of gas transfer have not been studied yet. So far, to the best of our knowledge, these currents are not parameterized in 1D models.

5. (1) The abstract mentions the need to improve biogeochemical processes in lake models to enhance weather prediction and climate projection capabilities. I'm not convinced improving biogeochemical processes will improve weather prediction. What is crucial in weather prediction or climate modeling is to simulate a correct surface temperature and fluxes because these are the variables that will be used in the coupling to the atmosphere. That's true that in climate simulations the knowledge of carbon dioxide or methane emissions are of high interest, however to my knowledge only the LAKE model offers the capability to simulate CO₂, O₂ and CH₄ dynamics and be coupled to a climate model. Please add a discussion on that point to also highlight the difficulty to increase the complexity of lake models and ensure a correct coupling to an atmospheric climate model.

(2)-(3) As in this study we do not focus on methane, due to absence of respective measurement data, and discuss models performance for CO₂ content, we added a comment on issues which will arise when implementing lacustrine CO₂ emissions in climate models (Page 21 line 29): The important role of catchment processes in building up the DIC levels in lakes introduces an extra difficulty for implementing lacustrine CO₂ emissions in the Earth system models. This is caused by the necessity to provide regional or global data on lake's catchments geometric, physical and biogeochemical properties, which are not currently available. In relation to this, we can also note a faster progress on a roadmap to introduce lake CH₄ dynamics in climate models, where simulations passed from site-levels studies to regional estimates (Tan et al., 2015), whereas CO₂ modeling is currently confined to individual lakes (Stepanenko et al., 2016; Tan et al., 2017; Kiuru et al., 2018).

6. (1) Ice cover is a key variable for vertical transfer of gases. It has been shown that freezeup or brake-up presented delays of several weeks potentially. Don't you think more effort should be put on the representation of ice and snow over ice in lake models, especially when working in NWP and/or climate contexts?

(2)-(3) We agree and added the following comment at Page 23 line 18: This implies

[Printer-friendly version](#)[Discussion paper](#)

more efforts of a community should be spent on elaborating ice-snow schemes in lake models, where currently vertical homogeneity and temporal invariance of optical and thermodynamic properties are typically assumed which does not correspond to a bulk of existing knowledge (Lepparanta et al., 2015). In a recent study, (Tan et al., 2018) one possible direction to improve the simulation of ice-cover is shown. They demonstrated that including the conversion of snow to white or slush ice when the weight of ice and snow exceeds the buoyancy of the ice cover, can significantly improve the ice simulation results. In special case of saline lakes, effects of salts trapping in ice cover become important and should be adequately parameterized as well (Stepanenko et al., 2019).

7. (1) It would also be of interest to have a comparison of surface temperatures observed at 10cm to the model simulations to be sure that the daily cycle of temperature is well reproduced. This is a key feature for any further coupling to an atmospheric model. Could you add such a graph in the revised manuscript?

(2) The diurnal cycle of the surface temperature has a limited relevance to the problem of the vertical diffusion of gases originated from bottom. Below there is a graph showing the diurnal course of temperature for 1 month, July, averaged over all 5 years. There is a systematic difference between the results of models and observation data (e.g. up to 4 °C for ALBM model), however, in this study we do not discuss the reason of these errors given the main focus of the study. (see attached figure: Fig.1)

8. (1) A comparison of methane profiles for ALBM and LAKE would also be interesting (climate change context, ...) even if no observation is available.

(2) The difference between these two models in methane simulation is significant: the maximum bottom concentration is 2.8 ppm and 1.1 ppm for ALBM and LAKE, respectively. The mean bottom methane concentration in the LAKE model (0.03 ppm) is an order of magnitude smaller than in the ALBM model (0.24 ppm). In general, the modeled concentration of methane in the lake is small, because of a high oxygen con-

[Printer-friendly version](#)

[Discussion paper](#)



centration. However, there are no observations available and we cannot consider one or another model as more realistic in this respect. Hence, we do not include this result into the main manuscript. (see attached figures: Fig.2, Fig.3)

Technical comments

1. (1) Page 4 line 12: change – by of
(2) We could not find these corrections.
2. (1) Page 6 line 5: summarized (2)-(3) Page 6 line 5: changed to "summarized"

References:

1. Fer, I., Lemmin, U., Thorpe, S. A.: Cascading of water down the sloping sides of a deep lake in winter, *Geophys. Res. Lett.*, 28(10), 2093-2096, <https://doi.org/10.1029/2000GL012599>, 2001.
2. Fer, I., Lemmin, U., Thorpe, S. A.: Winter cascading of cold water in Lake Geneva, *J. Geophys. Res.-Oceans*, 107(C6), 13-1, <https://doi.org/10.1029/2001JC000828>, 2002.
Forbes G. S. and Meritt J. H.: Mesoscale vortices over
3. Kaden, H., Peeters, F., Lorke, A., Kipfer, R., Tomonaga, Y., Karabiyikoglu, M.: Impact of lake level change on deep-water renewal and oxic conditions in deep saline Lake Van, Turkey, *Water Resour. Res.*, 46(11), <https://doi.org/10.1029/2009WR008555>, 2010.
4. Kirillin, G. B., Forrest, A. L., Graves, K. E., Fischer, A., Engelhardt, C., Laval, B. E.: Axisymmetric circulation driven by marginal heating in ice-covered lakes, *Hydrol. Geophys. Res. Lett.*, 42(8), 2893-2900, <https://doi.org/10.1002/2014GL062180>, 2015.
5. Kiuru, P., Ojala, A., Mammarella, I., Heiskanen, J., Kämäräinen, M., Vesala, T., Huttula, T.: Effects of climate change on CO₂ concentration and efflux in a humic boreal lake: A modeling study, *J. Geophys. Res.-Biogeo.*, 123, 2212– 2233, <https://doi.org/10.1029/2018JG004585>, 2018.

6. Lepparanta M. Freezing of lakes and the evolution of their ice cover: Springer Heidelberg New York Dordrecht London, 301 p.,doi:10.1007/978-3-642-29081-7, 2015.
7. Samolyubov, B. I.: Bottom stratified currents, Nauchny Mir, 464, 1999 (in Russian).
8. Stepanenko, V., Mammarella, I., Ojala, A., Miettinen, H., Lykosov, V., and Vesala, T.: LAKE 2.0: a model for temperature, methane, carbon dioxide and oxygen dynamics in lakes, Geosci. Model Dev., 9(5), 1977-2006, <https://doi.org/10.5194/gmd-9-1977-2016>, 2016.
9. V. M. Stepanenko, I. A. Repina, G. Ganbat, and G. Davaa. Numerical simulation of ice cover of saline lakes. Izv. Atmos. Ocean Phy.,55(1):129–138, <http://dx.doi.org/10.1134/S0001433819010092>, 2019.
10. Tan, Z., Zhuang, Q., and Walter Anthony, K.: Modeling methane emissions from arctic lakes: Model development and site level study, J. Adv. Model Earth Sy., 7(2), 459-483, <https://doi.org/10.1002/2014MS000344>, 2015.
11. Tan, Z., Zhuang, Q., Shurpali, N. J., Marushchak, M. E., Biasi, C., Eugster, W., and Walter Anthony, K.: Modeling CO2 emissions from Arctic lakes: Model development and site-level study, J. Adv. Model Earth Sy., 9(5), 2190-2213, <https://doi.org/10.1002/2017MS001028>, 2017.
12. Tan, Z., Yao, H., Zhuang, Q.: A small temperate lake in the 21st century: Dynamics of water temperature, ice phenology, dissolved oxygen and chlorophyll-a. Water Resour. Res., 54, <https://doi.org/10.1029/2017WR022334>, 2018.

Additional comment: Attached file contains all changes in the manuscript and SI mentioned above.

Please also note the supplement to this comment:

<https://www.hydrol-earth-syst-sci-discuss.net/hess-2019-146/hess-2019-146-AC2-supplement.zip>

Printer-friendly version

Discussion paper



Interactive comment on Hydrol. Earth Syst. Sci. Discuss., <https://doi.org/10.5194/hess-2019-146>, 2019.

HESSD

Interactive
comment

Printer-friendly version

Discussion paper






Fig. 1. The diurnal cycle of the surface water temperature (in July), averaged over 5 years.

Printer-friendly version

Discussion paper






Fig. 2. Vertical distribution of methane concentration in two models: ALBM, LAKE

Printer-friendly version

Discussion paper






Fig. 3. Vertical distribution of methane concentration (logarithmic scale) in two models: ALBM, LAKE

Printer-friendly version

Discussion paper



List of additional changes

1. SI Page 5 Table S4: correction of two numbers for model ALBM for LDSim (typo in the first version)
2. Page 10 line 16: text for FLake model analysis was rearranged .

Multimodel simulation of vertical gas transfer in a temperate lake

Sofya Guseva¹, Tobias Bleninger², Klaus Jöhnk³, Bruna Arcie Polli², Zeli Tan⁴, Wim Thiery^{5,6}, Qianlai Zhuang⁷, James Anthony Rusak⁸, Huaxia Yao⁸, Andreas Lorke¹, and Victor Stepanenko^{9,10}

¹Institute for Environmental Sciences, University of Koblenz- Landau, Landau, Germany. *

²Graduate Program on Water Resources and Environmental Engineering, Federal University of Paraná, Curitiba, Brazil.

³CSIRO Land and Water, Black Mountain, Canberra ACT 2601, Australia.

⁴Pacific Northwest National Laboratory, Richland, Washington, USA.

⁵Institute for Atmospheric and Climate Science, ETH Zurich, Zurich, Switzerland.

⁶Department of Hydrology and Hydraulic Engineering, Vrije Universiteit Brussel, Brussels, Belgium.

⁷Department of Earth, Atmospheric, and Planetary Sciences, Purdue University, West Lafayette, Indiana, USA.

⁸Dorset Environmental Science Centre, Ontario Ministry of Environment, Conservation and Parks, Dorset, Ontario, POA 1E0, Canada.

⁹Laboratory for Supercomputer Modeling of Climate System Processes, Research Computing Center, Lomonosov Moscow State University, Moscow, Russia.

¹⁰Department of Meteorology and Climatology, Faculty of Geography, Lomonosov Moscow State University, Moscow, Russia.

Correspondence to: Sofya Guseva (guseva@uni-landau.de)

Abstract.

In recent decades, several lake models of varying complexity have been developed and incorporated in numerical weather prediction systems and climate models. To foster enhanced forecasting ability and verification, improvement of these lake models remains essential. This especially applies to the limited simulation capabilities of biogeochemical processes in lakes and greenhouse gas exchanges with the atmosphere. Here we present multi-model simulations of physical variables and dissolved gas dynamics in a temperate lake (Harp Lake, Canada). The five models (ALBM, FLake, LAKE, LAKEoneD, MTCR-1) considered within this most recent round of the Lake Model Intercomparison Project (LakeMIP) all captured the seasonal temperature variability well. In contrast, none of the models is able to reproduce the exact dates of ice-cover and ice-off, leading to considerable errors in the simulation of eddy diffusivity around those dates. We then conducted an additional modeling experiment with a diffusing passive tracer to isolate the effect of the eddy diffusivity on gas concentration. Remarkably, sophisticated $k - \varepsilon$ models do not demonstrate a significant difference in the vertical diffusion of a passive tracer compared to models with much simpler turbulence closures. All models simulate less intensive spring overturn compared to autumn. Reduced mixing in the models consequently leads to the accumulation of the passive tracer distribution in the water column. The lake models with a comprehensive biogeochemical module, such as ALBM and LAKE, predict dissolved oxygen dynamics adequate to the observed data. However, for the surface carbon dioxide concentration the correlation between modeled (ALBM, LAKE) and observed data is weak (~ 0.3). Overall our results indicate the need to improve the representation of physical and biogeochemical processes in lake models, thereby contributing to enhanced weather prediction and climate projection capabilities.

*Previous affiliation of S.Guseva: Department of Geography, Lomonosov Moscow State University, Moscow, Russia.

1 Introduction

The past two decades have seen a renewed interest in lake modeling. Due to the ever-increasing spatial resolution in numerical weather prediction systems and climate models, many lakes became resolvable on the model grid. Lakes are responsible for changing local atmospheric conditions by modifying turbulent fluxes of heat, moisture and momentum relative to adjacent land (Forbes and Meritt, 1984; Mahrt, 2000; Long et al., 2007; Thiery et al., 2015). This effect can be substantial over regions where the total coverage of inland water bodies is high, e.g. in Finland or Sweden ($\approx 10\%$, 9% lakes of the total area, respectively (Lehner and Döll, 2004)), or where large lakes dominate (Docquier et al., 2016; Thiery et al., 2015, 2016, 2017). Additionally, lakes are increasingly recognized as a considerable source of greenhouse gases, such as carbon dioxide (CO_2) and methane (CH_4), to the atmosphere (Tranvik et al., 2009; Bastviken et al., 2011; Raymond et al., 2013; Tan and Zhuang, 2015; Wik et al., 2016).

With the development of multiple lake models of varying complexity, it becomes necessary to compare them and examine their merits and drawbacks in the context of atmospheric and limnological applications. The scientific community has addressed this need through the Lake Model Intercomparison Project (LakeMIP) launched in 2008 (<http://netfam.fmi.fi/Lake08/>). Since then, a methodology for lake model comparison has been developed (Stepanenko et al., 2010) and lake model intercomparison experiments have been conducted for a range of limnological and climatic conditions. In each experiment, input parameters were unified as much as possible, including identical meteorological forcing, optical properties of water, ice, and snow, initial temperature profiles, and lake depth and/or morphometry. This allowed for a detailed inter-model comparison in respect of limnophysical processes representation.

Previous LakeMIP studies primarily investigated the ability of the lake models to simulate the thermal regime of the water bodies that cover a wide range of size, depth and mixing regimes in different latitudes. In particular, the major effort has been spent on the simulation of the evolution of the vertical temperature profile along with the surface temperature and energy fluxes to the atmosphere. At the same time, biogeochemical processes were not considered in the model intercomparison studies.

LakeMIP simulations have been performed using seven one-dimensional lake models: 1) CLM4-LISSS (Hostetler and Bartlein, 1990; Subin et al., 2012), 2) FLake (Mironov, 2003), 3) Hostetler model (Hostetler et al., 1993), 4) LAKE (Stepanenko et al., 2011, 2016), 5) LAKEoneD (Jöhnk and Umlauf, 2001; Jöhnk et al., 2008), 6) MINLAKE96/2012 (Fang and Stefan, 1996), and 7) SimStrat (Goudsmit et al., 2002). They targeted deep lakes: monomictic Lake Geneva (Switzerland/France), dimictic Lake Michigan (USA), meromictic equatorial Lake Kivu (Central Africa), and for shallow lakes: Lake Sparkling (USA), Lake Großer Kossenblatter See (Germany), Lake Valkea-Kotinen (Finland). The lake models in general simulated reasonable seasonal variability of temperature and thermocline characteristics (Perroud et al., 2009; Stepanenko et al., 2010). However, the simulated and observed evolution of the mixed layer thickness and the bottom temperature tend to disagree. The $k - \varepsilon$ models (SimStrat, LAKE, LAKEoneD) demonstrate intensive mixing of the lakes during summer, emphasizing their applicability especially to shallow lakes (Stepanenko et al., 2010, 2014). Nevertheless, this type of lake models can also be appropriate for studying physical processes in deep lakes (Thiery et al., 2014a). Furthermore, lake models using the Henderson-Sellers parameterisation of vertical mixing (e.g. CLM4-LISSS, Hostetler lake model, (Henderson Sellers, 1985)) represent the

thermodynamic state of a water body adequately, but can underestimate mixolimnion temperatures (Thiery et al., 2014a). The FLake model has demonstrated less satisfactory agreement with observations in terms of summer stratification in previous studies, yet simulates surface temperatures well.

This paper presents a logical continuation of LakeMIP by advancing intercomparisons to the study of biogeochemical processes in lakes. It primarily focuses on the simulation of key physical factors affecting the vertical transport of greenhouse gases, such as thermal stratification, vertical diffusion of gases, and ice cover. The ice cover is of special interest for greenhouse gas dynamics, as it can cause the depletion of oxygen, which favors CH₄ accumulation until spring ice breakup (Phelps et al., 1998; Karlsson et al., 2013; Jammot et al., 2015).

Our study aims at identifying the major merits and shortcomings of different lake model formulations, as well as uncertainties in simulating the limnophysical controls on greenhouse gas distribution and emissions to the atmosphere. Simulations were run for Harp Lake, a lake in Ontario, Canada with long-term high-frequency monitoring data for meteorology, water temperature, CO₂ and O₂. Five lake models were used in this study to simulate thermal dynamics and turbulent diffusivity in the lake: 1) Arctic Lake Biogeochemistry Model, ALBM (Tan et al., 2015; Tan and Zhuang, 2015; Tan et al., 2017, 2018) 2) FLake (Mironov, 2003) 3) LAKE (Stepanenko et al., 2011, 2016) 4) LAKEoneD (Jöhnk and Umlauf, 2001; Jöhnk et al., 2008) 5) Modelagem do Transporte de Calor no Reservatório, MTCR-1 (Polli and Bleninger, 2015, 2018). ALBM and LAKE models include comprehensive biogeochemical modules for calculation of dissolved gas concentrations, which were then tested in their ability to reproduce CO₂ and oxygen O₂ dynamics. The study is structured as follows: Sect. 2 includes the description of the study site, observational data, lake models, and experimental setup. Section 3 is devoted to the simulation of the lake's thermal regime, i.e. the temperature patterns and eddy diffusivity (ED). To assess the pure effect of the ED on the gas distribution and dynamics in the lake, additional experiments have been conducted, where the fate of passive tracers emitted from the bottom of the lake was analyzed. In a final section, the models ALBM and LAKE are compared in terms of the CO₂ and O₂ concentration in Harp Lake.

2 Materials and methods

2.1 Study site and data

2.1.1 Study area

The object of the current study is Harp Lake (45° 22' N , 79° 07' W, 327 m above sea level), a dimictic temperate lake located in south-central Ontario, Canada. The lake and its surrounding forested catchment are representative of the southern Canadian Shield landscape (Cox, 1978; Dillon et al., 1987) (Fig. 1). It has a 0.71 km² surface area, a volume of 0.0095 km³, a maximum depth of 37.5 m and mean depth of 13.32 m. The lake is oligotrophic, characterized by average total phosphorus and chlorophyll-a concentrations of 6.7 and 3.5 µg l⁻¹ (Molot and Dillon, 1991). The water is slightly acidic (pH 6.3) (Dillon et al., 1987). Harp Lake has six inflows and one outflow, with a long-time average discharge of 0.067 m³ s⁻¹ for the inflows and 0.091 m³ s⁻¹ for the outflow. The difference is due to small ungauged inflows and overland flow. Its average residence time is 3.1 years.

Due to this relatively long residence time, the effect of the inflows on temperature can be neglected. The lake has only small water level fluctuations (maximum ~ 0.2 m).



Figure 1. The location and morphometry of Harp Lake (HP3 to HP6A are the six inflows, HPOF is the lake outflow (Yao et al., 2014)). Red circle indicates the position of the high-frequency monitoring buoy ($45^{\circ} 22' 44.88''$ N, $79^{\circ} 8' 7.77''$ W).

2.1.2 Observational data

The lake has been monitored by the Dorset Environmental Science Centre (DESC) for more than 40 years, thus providing a comprehensive database supporting the model experiments. The high-frequency and continuous observations used in this study have been collected during 5 years (14.07.2010-19.10.2015) with 10-minute resolution on a monitoring buoy in the lake center (Fig. 1). The meteorological variables collected are: air temperature, humidity at the height of 1.75 m above the water surface; wind speed at 1.75 m, and downwelling shortwave and longwave radiation. Pressure and precipitation are available from a land-based meteorological station 500 m away from the western lake shore. The high-frequency dataset further includes the a vertical temperature profile (up to the depth of 27.1 m prior to the summer of 2012 and to a depth of 9.85 m thereafter, aggregated to time step of 1 h), O_2 concentrations (at two depths: 1 m, 18 m, aggregated to a time step – 1 h) and CO_2 concentration (at a depth of 0.39 m, for the period 12.03.2012 – 19.10.2015, aggregated to a time step of 1 h). In addition, traditional temperature and O_2 profiles were collected to the depth of 35 m, except during the winter time period, at time step of approximately 2 weeks, and Secchi depth was also measured fortnightly or monthly during these 5 years (for more details see SI Table S6, S7).

The meteorological conditions of the region during the period under study are typical for continental climate of mid-latitudes (Table 1). This region is subject to the influence of the Azores High during summer, and is located in between the Icelandic Low, and the Canadian High forming in the central part of the continent during winter. Cold arctic air usually invades this area during winter, leading to the formation of stable freezing weather. However, it should be considered that winters are usually characterized by several abrupt rises in air temperature above $0^{\circ}C$ (see SI, Fig. S1). This phenomenon causes the 2012 winter to rank as the warmest winter during the 5 measurement years. Average annual wind speed is relatively low (≈ 2 m s^{-1}), likely

caused by wind-shading effects imposed by the surrounding forest. Predominant wind directions are WSW (230° - 280°) and SE (130°).

Table 1. Hydrometeorological conditions during the period 14.07.2010 – 19.10.2015.

	Max	Min	Average	Standard deviation, σ^*
Air temperature, T °C	31.4	-37.5	winter: -9.2 summer: + 18	12.4
Lake surface temperature, T_s °C ¹	28.4	-1.5	10.9	9.42
$\Delta T = T_s - T_{27.1m}$, °C	24.1	-4.3	5.6	9
Wind speed, W m s ⁻¹	9	0	2	1.28
Precipitation, R mm h ⁻¹	39.7	0	0.12	0.71

a. ¹ Water temperature measured at 0.1 m depth.

2.2 Lake models

In this study, a suite of 1-D lake models was applied to Harp Lake. Two classes of finite difference models can be identified based on different approaches of calculating the vertical ED: Henderson-Sellers-based models (Henderson Sellers, 1985; Hostetler and Bartlein, 1990) which are computationally inexpensive, and the more sophisticated models based on $k - \varepsilon$ turbulence closure scheme. The Henderson-Sellers approach computes the ED coefficient as a function of the Richardson number and considers wind-driven turbulence during stable or neutral stratification. A separate procedure is considered for convective mixing and assumes instantaneous mixing of unstably stratified depth layers. The $k - \varepsilon$ formulation involves prognostic equations for turbulent kinetic energy and its dissipation. Both wind-induced and convective turbulence are taken into account in this approach. Within the first class there are recently developed models ALBM and MTCR-1; while $k - \varepsilon$ models include LAKE and LAKEoneD.

The FLake model stands aside from the other 1-D models due to the two-layer bulk structure which employs the concept of self-similarity to estimate the temperature profile in the mixed layer, and thermocline, respectively (Mironov, 2008). In the mixed layer the temperature is assumed to be constant, whereas below it is parameterized as a function of non-dimensional depth. ED is not explicitly specified in this model. Of all models, FLake has the lowest computational cost (Thiery et al., 2014a).

Ice formation is one of the main controlling factors for the dissolved gas dynamics in a lake during winter. Models like LAKEoneD and MTCR-1 involve a simple ice parameterisation, in which ice-cover is governed only by air temperature and lake temperature is not considered (Ashton, 1980, 2011). **In FLake model, the temperature profile within the ice cover is parameterized via a time-varying shape-function having a linear asymptotic.** The ALBM model incorporates a more comprehensive ice parameterisation developed by Fang and Stefan (1994), taking into account the physical processes of freezing of the surface water, radiation and heat penetration through ice and snow. The LAKE model takes into account the same physical processes

and uses a multilayer ice parameterisation (Stepanenko and Lykossov, 2005; Stepanenko et al., 2011) based on an unsteady heat transfer equation. Light extinction in all models follows the Beer–Lambert law with a constant light attenuation coefficient.

ALBM and LAKE incorporate the most comprehensive biogeochemical modules, including the sources, sinks and transport of CO₂, O₂, CH₄ (and nitrogen in ALBM) (see SI, Table S1). LAKEoneD model calculates O₂ concentration only. Parameters and characteristics of all models are summarized in Table 2. All models use finite-difference numerical schemes along vertical coordinates, except for FLake which is based on the two-layer representation of lake temperature profile (see above). The vertical grid of these models is fixed in time, while the number of numerical layers varies from 51 to 105 between models. The time step varies within the range of 20-3600 s between the models. ALBM uses a fourth-order adaptive Runge-Kutta-Fehlberg solver and a variable time step. If there are abrupt changes of boundary conditions (such as air temperature, solar radiation and wind), the time step in this model will be reduced to avoid numerical instability.

Table 2. Characteristics of lake models relevant to the study.

Lake model	Number of layers	Time step (s)	Turbulent mixing parameterisation	Gases considered in biogeochemical module / analyzed in experiments	Ice module
ALBM	51	1-1800	Henderson-Sellers-based	CO ₂ , O ₂ , CH ₄ , N ₂ / CO ₂ , O ₂	bulk model (Fang and Stefan, 1994)
MTCR-1	95	3600	Henderson-Sellers-based	–	simple model (Ashton, 1980, 2011)
FLake	2	3600	two-layer bulk model, assuming self-similar temperature profile	–	bulk model, assuming self-similar temperature profile
LAKE	105	20	$k - \varepsilon$	CO ₂ , O ₂ , CH ₄ / CO ₂ , O ₂	multilayer model (Stepanenko and Lykossov, 2005)
LAKEoneD	75	240	$k - \varepsilon$	O ₂	simple model (Ashton, 1980, 2011)

2.3 Experimental setup

In this study, a set of numerical experiments were conducted to compare model simulation outputs assuming different morphometries, light extinction coefficients, and for the biogeochemical model parts initial CO₂ concentrations (Table 3):

- 1) a reference simulation in which the depth is set to a maximum value (37.5 m), light attenuation coefficient is set to a mean measured value ($\mu = 0.4 \text{ m}^{-1}$) and the morphometry is not taken into account (dependence of horizontal cross-section area on depth (hypsometric curve) is not included). [RefSim];
- 2) model simulations testing the sensitivity of models to variations in:
 - lake depth, setting it to mean value: $h_{\text{ave}} = 13.32 \text{ m}$ [LDSim],
 - the light attenuation coefficient, prescribing $\mu_{\text{min}} = 0.28 \text{ m}^{-1}$, and $\mu_{\text{max}} = 0.68 \text{ m}^{-1}$ which are minimal and maximal values measured, respectively [ExtMinSim, ExtMaxSim],
- 3) a model simulation testing the effect of the vertical diffusion on the distribution of the passive tracer [PassTrSim].
- 4) a model simulation with biogeochemical modules activated including the calculation of the O₂ and CO₂ concentration [GasSim]. Measured mean CO₂ concentration and O₂ observations are used as initial profiles. Time step of model output in all experiments is 1 h.

Table 3. Parameters of numerical experiments of LakeMIP-Harp.

Experiment (14.07.2010 - 19.10.2015) ¹	Light attenuation coefficient, μ , m^{-1}	Lake depth, m	Hypsometric curve	Initial CO ₂ concentration, $\text{mol}\cdot\text{m}^{-3}$	Participating models
RefSim	0.4	37.5	–	–	all
LDSim	0.4	13.32	–	–	all
ExtMinSim, ExtMaxSim	0.28/0.68	37.5	–	–	all
PassTrSim	0.4	37.5	–	–	ALBM, MTCR-1, LAKE, LAKEoneD
GasSim	0.4	37.5	+	0.098	ALBM, LAKE

a. ¹ For more details see the simulation protocol in SI. Some of the experiments are not included into our study.

- 15 **The concept of the self-similarity in the FLake model (Sect. 2.2) includes the "shape factor" coefficient C_θ , which determines the vertical temperature profile below the mixed layer. Evolution of this parameter is controlled by a relaxation time scale t_{rc} (Mironov, 2008). This time scale includes the dimensionless relaxation constant C_{rc} having a default value of 0.003 in the model. This constant is a calibration parameter which is generally individual for each lake (Shatwell et al., 2016; Kirillin et al., 2017). We tested the sensitivity of the model to the variation of C_{rc} using values 0.03, 0.3, 0.5, 1, 2, 3, 30 and achieved the**

best agreement with the measured temperature profile using $C_{rc} = 0.3$ (for details, see Sect. 3.1). In the following sections, this C_{rc} setting is used in RefSim and other experiments with FLake model.

The meteorological forcing is identical for all models and experiments. This, however, does not imply that heat fluxes to the atmosphere are the same in models, as models use different surface flux schemes. Some of the models do not include bathymetry (e.g. FLake); thus, for a more consistent comparison, in the RefSim experiment all models are integrated assuming homogeneous depth. In the experiment GasSim, lake morphometry is considered because the lake's carbon budget cannot be accurately calculated without including sediment oxygen demand along a sloping bottom.

The effect of the tributaries is neglected due to the large residence time of Harp Lake. Likewise, water level variations are not considered in the models given the small temporal fluctuations.

The choice of the lake depth in the one-dimensional model without bathymetry may be crucial for the modeling results, not only for the temperature regime (Balsamo et al., 2010) but also for the gas dynamics. The gas dynamics is affected by lake depth as the latter controls the amount of bottom-originated gas to be biochemically-transformed in the water column (e.g. oxidation of CH_4). Thus, it is important to examine the sensitivity of models to this parameter. In all experiments except LDSim, the maximal depth is used to compare the output with all available measurements. To test the effect of the chosen depth on model performance, the depth is set to mean depth (13.32 m) in the LDSim experiment, as such the modeled lake volume is equal to the real lake volume.

The light attenuation coefficient is an important parameter, regulating the thermal regime of a lake (Heiskanen et al., 2015) and biogeochemical processes including photosynthesis. Secchi depth measurements during the 5-year study period on Harp Lake are irregular but enable calculation of mean, maximum and minimum light attenuation coefficient using the Poole and Atkins formula (Poole and Atkins, 1929).

The motivation for conducting experiments with perturbed input parameters (depth and light attenuations coefficient) and initial conditions (dissolved gases) is that these properties, especially light attenuation coefficient, become considerably uncertain in global applications. While a first global lake depth database recently became available (Choulga et al., 2014; Kourzeneva, 2010), such data set currently does not exist for water transparency.

To understand the role of ED coefficient in gas distribution and dynamics, an additional experiment solving the pure vertical diffusive transport equation for a passive tracer governed by eddy diffusivities resulting from different lake models was conducted. Details on this experiment are presented in Sec. 3.3.

Harp Lake is a deep lake suitable for studying the transport of O_2 and CO_2 as well as their biogeochemical transformations due to long-term regular measurements of these gases. In this respect, the additional numerical experiment [GasSim] was conducted with ALBM and LAKE, the lake models with most sophisticated biogeochemical modules. To harmonize the experimental setup, the contribution of CO_2 and O_2 fluxes through a sloping bottom to the lake carbon budget was switched on in the LAKE model. It should be noted that ALBM model includes CO_2 sink/source due to outlets and inlets, whereas LAKE does not take this effect into account. In the ALBM model, a set of biogeochemical parameters was calibrated, specifically: maximum chlorophyll-specific photosynthetic rate, maximum metabolic loss potential, aquatic dissolved organic matter (DOM) microbial degradation rate, terrestrial DOM microbial degradation rate, groundwater dissolved organic carbon (DOC)

concentration (Tan et al., 2017). No parameter calibration was performed for the LAKE model (a complete set of default values of model parameters is given in (Stepanenko et al., 2016)).

The model skill scores used in this study are listed in Table 4.

Table 4. Statistical measures of model performance.

Symbol	Property	Definition
f_n, r_n	Model and observed data	
σ_f, σ_r	Standard deviations of f and r, respectively	
DM	Difference between the mean values of the model and the observed data	$DM = \bar{f} - \bar{r}$
R	Correlation coefficient	$R = \frac{\frac{1}{N} \sum_{n=1}^N (f_n - \bar{f})(r_n - \bar{r})}{\sigma_f \sigma_r}$
RMSE _c	Centered root mean square error	$RMSE_c = \left(\frac{1}{N} \sum_{n=1}^N [(f_n - \bar{f}) - (r_n - \bar{r})]^2 \right)^{\frac{1}{2}}$

3 Results and Discussion

3.1 Temperature and ice

All lake models simulate seasonal water temperature dynamics adequately (Fig. 2): the epilimnion is warming throughout the first half of summer, accompanied by thermocline deepening, eventually leading to overturn in autumn. During winter, the lake temperature is at the freezing point immediately below the ice-cover, causing stable stratification. During spring, the lake starts to mix due to the penetration of shortwave radiation. The ice formation and ice melt dates play a key role in various processes: the interaction of the lake surface with the atmosphere, the overturning during spring, biogeochemical processes, release of gases that have accumulated during wintertime, e.g. CO₂ or CH₄. Yet none of the models is able to precisely predict these dates. The difference between the model results and observations become even larger towards the end of simulation, with differences exceeding 2 weeks in winter 2014-2015 (see SI, Table S3).

FLake model. In RefSim we test the model with different values of C_{rc} (see Sect. 2.3 above). The best agreement with observed temperature profile we get using $C_{rc} = 0.3$. In particular, the error (RMSE_c) for surface temperature reduces from 3.4 °C (using standard value of $c_{relax} = 0.003$) to 2.5 ° and becomes close to other models. At the depth of ~ 6-10 m RMSE_c reduces from 5.7 °C (using standard value of $c_{relax} = 0.003$) to 5.2 °. In addition, RMSE_c reduces from 3.3-0.7 °C to 1.8-0.2 °C at the depths 20-27 m.

FLake demonstrates the largest error among other models for the depth of thermocline (Fig. 3 (a)), (RMSE_c up to 5.2 °C). In contrast, the model performance in simulation of the lake surface temperature is comparable with other models and the error is only 2.5 °C (see SI, Fig. S2 (a)), which points to the fact that it includes similar formulations for the surface heat balance as the other models.

As highlighted in former studies (e.g. Stepanenko et al. (2010); Thiery et al. (2014b)), this model overestimates the surface mixed layer depth during summer, even under ice (Fig. 2 (b)). The mixed-layer depth in the FLake model is 1.5 times larger than in observed data (5.6 m versus 3.6 m). Predictions of ice-cover and ice-off are characterized by large errors (see SI, Table S3), with modeled dates more than two weeks off from the observations.

Remarkably, FLake demonstrates the largest sensitivity among all models to the lake depth – RMSE_c is reduced approximately twofold (from 2.5 °C to 1.28 °C) (see SI, Table S4, Fig. S3, S4) when using a mean depth of 13.32 m instead of the maximal depth 37.5 m. The thickness of mixed layer depth reduces up to 4.5 m and have a better agreement with the observations in comparison with RefSim experiment.

FLake showed similar response in the sensitivity model test to variation of the light attenuation coefficient as other models. However, the difference in temperature is smaller than in other models: in RefSim and both ExtMinSim, ExtMaxSim is less than 4 °C in comparison with 8 °C (Fig. 4). It can be related to the weaker temperature gradient in the thermocline in comparison with other models (Fig. 2).

ALBM model. The warmest model in terms of the epilimnion temperature was ALBM model (Fig. 2 (a)). The maximum water temperature reached in the model is 32.4°C (28.4°C – in observations). However, ALBM demonstrates the lowest RMSE_c (1.53° C) and the highest correlation (0.98) with the observed data for the temperature of the water column (except

the surface) during the entire simulation period among all models (Fig. 3 (a), and see SI, Fig. S2, Table S4). It can be noted visually, that the shape of the thermocline is better reproduced by this model than by other model-participants (Fig. 3 (a)). In addition, the correlation coefficient of ALBM temperature decreases more slowly with depth than in all other models (Fig. 3 (b)). In contrast, $RMSE_c$ for the surface temperature in ALBM is almost 1.5 larger than in $k - \epsilon$ models (see SI, Fig. S2 (a)).

5 The ALBM model successfully reproduces autumnal overturn, although the homogeneous temperature distribution in this model occurs approximately 7-10 days later compared to the measurements. The likely reason for this offset is the excessive accumulation of heat in the mixed layer during summer, which delays the cooling to the temperature of maximum density and subsequent vertical mixing.

The ice-cover and ice-off dates are quite close to observations (e.g. 24.12.2011 vs 28.12.2011, see SI, Table S3). Differences
10 between model and observations ice-cover dates by of more than two weeks happen three times. The large errors mainly occur for ice-cover, likely due to the excessive heat accumulated in summer.

MTCR-1 model. In terms of temperature, the Henderson-Sellers-based MTCR-1 model shows contrasting performance compared to ALBM (Fig. 2 (e)). It is generally the coldest model (the temperature reaches a maximum of 24.2 °C in the epilimnion) and results in the deepest mixed layer (5.9 m versus 3.6 m observed). This significant difference in mixed layer
15 depth (see SI, Table S2) is possibly associated with different schemes of convective mixing.

Similar to the FLake model, the surface temperature in MTCR-1 is sensitive to changes in lake depth. The surface temperature DM reduced 4 times compared to the reference model simulation when decreasing the lake depth from 37.5 to 13.32 m (see SI, Table S4).

While MTCR-1 and LAKEoneD use similar parameterisation for ice formation and melt, their results are similar only for
20 the ice-off dates. The ice-cover dates greatly vary between these models with differences reaching up to 1 month (winter 2011-2012). Interestingly, LAKEoneD produces ice melt periods in winters 2011-2012 and 2012-2013 (see SI, Table S3) when MTCR-1 does not show it at all (there is no observed ice melt during these winters).

LAKEoneD and LAKE models. Even though LAKE and LAKEoneD use the same turbulence closure, they represent the temperature dynamics in the lake with considerable differences (Fig. 2 (c,d)). In particular, the mean depth of the mixed layer
25 as simulated by LAKE is 1 m deeper compared to LAKEoneD results (4.8 m and 3.9 m, respectively). LAKE demonstrates greater heating of the epilimnion (up to 30 °C) than LAKEoneD (maximum value of 27.3 °C). The homogeneous distribution of temperature during autumn mixing predicted by LAKEoneD reaches the depth of 15 m in most years, as opposed to LAKE (up to 35 m) and the observational data (at least up to 25 m, where the deepest sensor is located). Remarkably, only in these two models periods of unstable ice-cover occur. For example, the ice in LAKE is thin and often comes off during winter (more
30 than 5 times in winter 2011-2012), leading to wind-driven surface layer mixing (see SI, Table S3).

The sensitivity experiments with varying light attenuation coefficient (see SI, Table S4) generally result in a similar response in all models. Increasing the light attenuation coefficient (i.e. reducing water transparency) leads to an upward shift of the thermocline position (Fig. 4), and conversely, decreasing the attenuation coefficient causes a downward move of the thermocline.

This leads to temperature change at respective depths up to 4 °C in FLake model and 8 °C in ALBM and LAKEoneD. (Fig. 4)




Figure 2. Time-depth pattern of temperature in Harp Lake (14.07.2010-19.10.2015), RefSim and observed data. The grey boxes represent duration of the ice-cover period, the white patterns denote the absence of data.




Figure 3. (a) Centered root-mean-square error of modeled temperature calculated for each individual depth; (b) the correlation coefficient between the simulated and observed temperature at each depth.




Figure 4. Temperature difference between the experiments ExtMaxSim and RefSim in three lake models: (a) ALBM; (b) FLake; (c) LAKEoneD.

contains only 3 models with different types of turbulent closure because the effect of varying the light attenuation coefficient is very similar in models on the same type (e.g. $k-\epsilon$ models LAKEoneD and LAKE).

It is important to note that all models have a maximum error, $RMSE_c$ for temperature (compared to observations) in the thermocline (Fig. 3 (a)), which agrees with a previous study (Stepanenko et al., 2014). Conversely, the correlation between simulated and observed temperature decreases with depth in all models (Fig. 3 (b)): starting from 0.9-1.0 near the surface to 0.1-0.7 close to the bottom.

5 3.2 Eddy diffusivity

Eddy diffusivity (ED), K , is the variable controlling the vertical turbulent transport in lake models. This section examines the results of simulated ED values, thereby focusing on discrepancies between models caused by different turbulent closure schemes.

The simulated time-depth distribution of ED is shown in Fig. 5. ED in ALBM, and MTCR-1 do not reproduce spring and autumn whole-water-column mixing as the Henderson-Sellers diffusivity is not valid for unstable stratification. In contrast, $k - \varepsilon$ models correctly predict the deepening of the mixed layer in the ED pattern during summertime towards a complete mixing during autumn.

Even models employing the same type of turbulence parameterisation differ in simulated eddy diffusivities in the thermocline by orders of magnitude. This is caused by the different background values of ED added to those estimated by turbulence closures and mimicking mixing mechanisms not directly resolved in 1D models such as internal waves breaking (e.g. Hondzo and Stefan (1993)). Apparently, the background diffusivity is switched off in MTCR-1 model during ice-cover.

The increase of ED near the bottom during the stratified period in the ALBM can be explained by the change from a stable stratification in the thermocline to a quasi-neutral stratification beneath.

A peculiar feature of LAKE is that it features high-intensity turbulence during much of the ice-cover period in the lower part of the water column that is apparently related to residual flow after the autumn overturn. In contrast, in LAKEoneD, turbulence dissipates shortly after the ice-cover. Both of these models demonstrate a subsurface region of high ED values in winter, which is not reproduced by other models. Production of turbulent kinetic energy (TKE) below the ice may be caused by both momentum transfer through partial ice-cover (implemented in LAKE) and shortwave radiation penetration causing under-ice convection. The conspicuous difference between LAKE and LAKEoneD results is that springtime deep mixing does not develop in LAKEoneD. This may be linked to different constants and stability functions used in its $k - \varepsilon$ closures, or different buoyancy and momentum fluxes at the lake-atmosphere interface during this season.

It is worthy to note that the significant difference between the modeled and observed dates of ice-cover and ice-off (see SI, Table S3) leads to errors in the simulation of ED as water opens and interacts with the atmosphere.

The main features of ED distribution over seasons described in this section hold in experiment LDSim with different lake depth value (13.32 m).




Figure 5. Time-depth distribution of the ED in Harp Lake (14.07.2010-19.10.2015), RefSim model experiment.

3.3 Effects of eddy diffusivity on the vertical distribution of a passive tracer

The vertical distribution of ED in lake models is one of the major drivers of vertical transport of dissolved gases and their emission to the atmosphere. To isolate the effect of the ED parameterisation in the different models, we solve the diffusion equation for a passive tracer concentration C :

$$5 \quad \frac{\partial C}{\partial t} = \frac{\partial}{\partial z} K \frac{\partial C}{\partial z}. \quad (1)$$

The emission rate of the passive tracer at the bottom is set constant:

$$K \left. \frac{\partial C}{\partial z} \right|_{z=h} = F_{bot}, F_{bot} = const > 0, \quad (2)$$

while at the surface the flux is assumed to be large enough to make the surface concentration negligible compared to deep-water concentration, so that the boundary condition here becomes:

$$10 \quad C|_{z=0} = 0. \quad (3)$$

Setting zero concentration at the surface is justified by the fact, that surface concentration of gases of interest (CO_2 , CH_4) is typically orders of magnitude less than bottom concentration. The total (molecular plus turbulent) diffusivity K is taken from different models, so that the solution C of Eq. (1)-(3) is individual for each lake model.

Calculation of tracer diffusion was carried out using simulated values $K(z, t)$ from the two Henderson-Sellers-based models – ALBM, MTCR-1, and two $k - \varepsilon$ models – LAKE and LAKEoneD, covering the period of 5 years. In order to take into account vertical mixing under unstable stratification performed in ALBM and MTCR-1 and not expressed by Henderson-Sellers diffusivity, a value $K = 10^5 \text{ m}^2/\text{s}$ was applied for the convective layer in such cases.

The results of the numerical tracer experiment are presented in Fig. 6. All models demonstrate seasonal variability of the passive tracer concentration, including the near-bottom accumulation during stratified periods. Three models (ALBM, MTCR-1, and LAKE) perform the full overturn of the water column during autumn and spring, leading to the emission of almost the whole tracer storage to the atmosphere and forming a homogeneous concentration profile. In LAKEoneD, the vertical mixing does not reach the lake bottom during spring and autumn in certain years (e.g. 2011 and 2014). Hence, not all tracer amount is removed from the water column during overturn periods (see SI, Fig. S5 (b)) in this model. As a result, not only the concentration of the tracer near the bottom, but also the total tracer amount in water column increases in LAKEoneD throughout the simulation period. However this effect is a result of the assumption of a bathtub shaped lake bathymetry and would be less when including the depth dependence of lake volume.

The MTCR-1 model demonstrates short overturn periods similar to those of LAKEoneD, but the mixing reaches deeper. Hence, the passive tracer does not accumulate over time. Furthermore, ALBM exhibits a higher near-bottom concentration than MTCR-1 because the ED simulated by ALBM is one order of magnitude lower compared to MTCR-1 at 30-35 m depth (see Fig. 6).




Figure 6. Vertical distribution of passive tracer concentration in Harp Lake simulations, according to different models

To assess the effect of different eddy diffusivities on the surface tracer flux, we use the ratio of the surface flux, F_{surf} , to the bottom flux, F_{bot} , where the former is given as:

$$F_{surf} = K_{surf} \cdot \left(\frac{C_w - C_{eq}}{\Delta z} \right), \quad (4)$$

with $C_{eq} = 0$ the surface concentration, C_w the concentration at the first model output level below surface (0.3 m), both in $[\text{mol m}^{-3}]$, $\Delta z = 0.3 \text{ m}$, and K_{surf} is surface diffusivity coefficient.

The Henderson-Sellers-based models produce short periods with high fluxes, primarily during autumnal mixing (see SI, Fig. S5 (a)) and the drops of the integral concentration in the water column at the same time (see SI, Fig. S5 (b)). The flux spikes are several orders of magnitude higher compared to $k - \varepsilon$ models. A reasonable explanation is that instantaneous convective mixing implemented in ALBM, MTCR-1 models leads to almost the whole tracer content inside the convective layer to be released to the atmosphere during a single convective event. All models demonstrate much weaker tracer fluxes during spring overturn compared to those resulting from autumnal convection.

Lake models with $k-\varepsilon$ closure produce substantial tracer fluxes during the stratified period in particular years (e.g. 2013 and 2014). The likely reason for this effect is rapid changes in mixed-layer depth, due to sudden wind forcing, entraining higher concentrations from the hypolimnion into the mixed layer.

The tracer diffusion experiment was also conducted at the reduced depth of 13.32 m (see SI, Fig. S5 (c,d)). Due to the lower
5 depth, seasonal vertical mixing now extends over the whole water column in all models, providing more effective release of the substance. ALBM and MTCR-1 demonstrate episodic high fluxes during summer along with $k-\varepsilon$ models. The spring mixing in LAKEoneD now reaches the bottom in some years, hence the integral tracer amount evolves here in a similar way compared to all other models. The time-averaged tracer emission in LAKEoneD is thus the most sensitive to depth reduction. The mean
10 surface flux in this model increases up to approximately 54 %, while for ALBM, MTCR-1, LAKE – this increase is 4 %, 19 %, and 8 %, respectively. This can be primarily explained by the contribution of the increased fluxes during summer stratification and of the autumn mixing now reaching the bottom.

3.4 Harp Lake biogeochemistry

3.4.1 Oxygen

The evolution of observed vertical distribution of O₂ concentration in Harp Lake is shown in Fig. 7 c excluding winter periods and data with higher sampling frequency (1 h) collected at 1, 18 m depth (Fig. 8). During spring (end of April - beginning of May) after ice-off, convective mixing occurs, leading to a homogenized O₂ profile. The similar process happens in autumn. During summer, O₂ depletes in the hypolimnion due to its consumption by organic matter degradation in sediments and deep water layers. The main source of O₂ is located in the photic layer, where it forms as a result of photosynthesis. Below the mixed layer (at 5-10 m depth), there is a maximum of O₂ concentration due to weak turbulence, so that the produced O₂ is not transported to the surface and eventually emitted to the atmosphere. Emergence of this maximum is consistent to observations on a number of other freshwater lakes (Camacho, 2006; Cantin et al., 2011). In particular, the O₂ maximum is remarkable during the summer of 2012: ($C_{O_2\max}$ increases up to 15 mg l⁻¹) at 5 m depth. During winter, the O₂ concentration is usually high, indicating relatively low O₂ consumption rate from organic matter degradation in the lake. In summer, surface oxygen content decreases following reduction of temperature-controlled solubility. The measurements thus indicate common trends observed in most deep oligotrophic lakes at mid-latitudes (Mitchell and Prepas, 1990).

Both models depict a reasonable representation of O₂ dynamics at different depths (Fig. 7 (a, b)). ALBM includes the O₂ model described in Tan et al. (2017), whereas LAKE employs the representation of O₂ sinks and source from (Stefan and Fang, 1994). Even though they use different oxygen and turbulent parameterizations, both models show similar spatio-temporal pattern. In particular, they reproduce the maximum of O₂ concentration below the mixed layer, its quasi-linear in time decrease in the hypolimnion (18 m) during stratified periods, and the effect of mixing events during spring and autumn, marked by rapid changes of O₂ concentration at 18 m depth. However, the O₂ concentration as computed by LAKE is on average 1-2 mg l⁻¹ higher than the observed values. A plausible explanation for this positive bias is that photosynthesis in LAKE yields excessive production of O₂ (parameters of biogeochemical parameterisations in this model have not been calibrated in the study). Another deficiency of LAKE simulations is that during ice-cover near-surface O₂ decays linearly, and then exhibits significant peaks at thinning of ice-cover and ice-off, whereas in ALBM and measurements, oxygen content holds almost constant during these periods, which is more consistent with measurements.

The time series of O₂ concentration from ALBM and LAKE correlate reasonably to observed data, with linear correlation coefficient ranging between 0.5-0.6 at 1 m, 0.6-0.8 at 18 m depth, and with RMSE_c 1.27-1.56 mg l⁻¹ at 1 m, 1.19-1.52 at 18 m depth (see SI, Fig. S6). For both depths, ALBM demonstrates higher correlation with observations. Both models correlate better with observations at large depth. Decrease of the temperature in the ExtMaxSim simulation (Fig. 4) due to the upward shift of thermocline leads to the reduction of O₂ concentration at the respective depths due to temperature dependence of photosynthesis.

Overall, the oxygen content is highly dependent on physical controls which are temperature, radiation, ED and ice cover. Given that these factors are reasonably reproduced by ALBM and LAKE, O₂ spatiotemporal pattern is captured as well.

Successful representation of maximal O₂ content below mixed layer during summer may be an important modelling skill

for simulating correctly CH_4 in a lake because the former acts as a sink region for the latter. Furthermore, realistic oxygen concentration reduction during periods of stable stratification means that the respective CO_2 production from aerobic organic matter decomposition is reproduced reasonably as well.




Figure 7. Time-depth profiles of O_2 in Harp Lake (14.07.2010-19.10.2015), model experiment including sediments and morphometry (Gas-Sim). White patterns in (c) indicate absence of data.




Figure 8. Time series of O_2 concentration at depths 1 m and 18 m, observed and simulated

3.4.2 Carbon Dioxide

The vertical distribution of CO₂ concentration simulated by ALBM and LAKE are shown in Fig. 9. ALBM simulates higher concentrations of CO₂ compared to LAKE. This is likely because, contrary to LAKE, ALBM considers dissolved inorganic carbon (DIC) inflow from the catchment and explicitly treats DOC and particulate organic carbon (POC) dynamics (Tan et al., 2017). Parameter calibration applied in ALBM brought CO₂ content closer to measurements in this model. Correlation with the observational data at depth of 0.39 m is relatively small for the both models (0.3) and RMSE_c is 4.5, 2.7 mg l⁻¹ for LAKE and ALBM, respectively.

Both models reproduce the seasonality of the CO₂ concentrations: during summer CO₂ accumulates in the hypolimnion, while CO₂ is consumed in the mixed layer due to photosynthesis. For the same reason, the depth of the O₂ maximum (Fig. 7 (c)) is consistent with the metalimnetic CO₂ minimum.

Vertical turbulent diffusion (Sec. 3.3) greatly affects CO₂ patterns in models. In particular, the mixing periods clearly seen in Fig. 9 coincide with those in Fig. 6. Accumulation of CO₂ below mixed layer during stratified periods is clearly demonstrated by both models. However, the vertical CO₂ distribution in the thermocline is very different, i.e. in LAKE, concentration attains its maximum near the bottom, whereas in ALBM the maximum is located at 15-20 m depth. This can be related to the fact that in ALBM, DOC and DIC from surface water and ground water can be injected in the middle of the water column.

In contrast to oxygen, measured CO₂ content is much less correlated to simulated series. The wintertime increase of surface concentration seen in calculated series is hardly observable in measured data. In summer, simulated CO₂ series are smooth, whereas there are significant fluctuations in empirical data. There is some correspondence between seasonal-mixing-caused peaks in modeled concentration and measured one, though with time lags. All this leads to conclusion, that, compared to oxygen, carbon dioxide is more controlled by biogeochemical processes misrepresented in lake models, than by physical factors. As stated in the previous section, a realistic decay of oxygen content during stratified periods in ALBM and LAKE models suggests that CO₂ amount produced by aerobic decomposition of organic matter both in water column and in the top part of sediments is reasonably simulated as well. Satisfactory agreement of computed oxygen in the mixed layer and below with observations implies that photosynthesis minus respiration rate is fairly captured as well, and so do the models for CO₂ gain or loss due to these processes. Hence, the primary drawback of the models used in respect to DIC simulation is likely to be not explicitly simulating transport of carbon species from catchment to a water body. Thus, modelling approaches coupling the catchment and a lake presented recently (Futter et al., 2008; Duffy et al., 2018; McCullough et al., 2018) should be elaborated and wider used.

The important role of catchment processes in building up the DIC levels in lakes introduces an extra difficulty for implementing lacustrine CO₂ emissions in the Earth system models. This is caused by the necessity to provide regional or global data on lake's catchments geometric, physical and biogeochemical properties, which are not currently available. In relation to this, we can also note a faster progress on a roadmap to introduce lake CH₄ dynamics in climate models, where simulations passed from site-levels studies to regional estimates (Tan et al., 2015), whereas CO₂ modeling is currently confined to individual lakes (Stepanenko et al., 2016; Tan et al., 2017; Kiuru et al., 2018).




Figure 9. Time-depth profiles of CO_2 concentration in Harp Lake (17.08.2012-20.04.2015), experiment including modules of sediments and morphometry (GasSim)

4 Conclusions

A lake model intercomparison study focusing on physical controls of biogeochemical species (i.e., dissolved gas concentration) is performed, making a step forward in the LakeMIP exercises, which have been previously focused on thermodynamic modeling only. To evaluate model performance with respect to biogeochemical processes we had to first evaluate their performance in hydro/thermodynamic simulations and variations due to uncertainty in major driving parameters, such as temperature and turbulent diffusivity.

The participating lake models (ALBM, FLake, LAKE, LAKEoneD, MTCR-1) were evaluated regarding their ability to reproduce the thermodynamic regime of Harp lake. All models capture the seasonal variability of the temperature profile in the deep boreal dimictic lake. A substantial discrepancy was found between models in representation of thermocline evolution – consistent with previous studies – and the correlation coefficient for calculated and observed temperature was found to decrease from surface to bottom (from 0.9-1 to 0.1-0.6). The sensitivity of the simulation results to the light attenuation coefficient and the lake depth was assessed, given that in global applications, such as numerical weather prediction, they are external parameters known with high uncertainty.

All considered lake models do not accurately reproduce the dates of ice-cover and off, similar to the results from a previous simulation study for the same lake which used other lake models and an earlier time period (Yao et al., 2014). The difference between the model results and the observations exceeds 2 weeks in particular years. The unrealistic representation of ice formation and ice melt dates leads to untimely changes in eddy diffusivity (ED) within the water column, which in turn strongly affects the modeled distribution of gases and their emission to the atmosphere. **This implies more efforts of a community should be spent on elaborating ice-snow schemes in lake models, where currently vertical homogeneity and temporal invariance of optical and thermodynamic properties are typically assumed which does not correspond to a bulk of existing knowledge (Lepparanta, 2015). In a recent study, (Tan et al., 2018) one possible direction to improve the simulation of ice-cover is shown. They demonstrated that including the conversion of snow to white or slush ice when the weight of ice and snow exceeds the buoyancy of the ice cover, can significantly improve the ice simulation results. In special case of saline lakes, effects of salts trapping in ice cover become important and should be adequately parameterized as well (Stepanenko et al., 2019).**

To study the effect of turbulent transport on the vertical distribution of gases, we conducted passive tracer simulations using simulated ED from different models.

This experiment reveals that all lake models demonstrate more intense mixed layer deepening during autumn than during spring, with spring overturn in some cases not even reaching the lake bottom (e.g. in MTCR-1, and LAKEoneD).

Moreover, less intensive mixing during spring and autumn leads to an accumulation of passive tracer concentrations in deep water at a multiyear timescale.

In addition, the representation of an equivalent lake depth in 1-dimensional models is found to be a crucial parameter regulating both passive tracer concentration and its flux to the atmosphere: when reducing the depth of a lake approximately by 50 %, the average tracer flux increased by 4 %-54 %.

The lake models with most comprehensive biogeochemical modules such as ALBM, LAKE demonstrated reasonable reproduction of the O₂ concentration profile and its seasonal evolution. For instance, both models predict an oxygen concentration maximum and corresponding CO₂ minimum below the mixed layer in summer, elevated surface concentration and linear decay of deep-water content under ice-cover, vertical homogenization during spring and autumn overturns.

- 5 The main difference between the models in the CO₂ simulation is that the ALBM model involves additional source of DIC from the catchment. Due to this, the average CO₂ concentration in ALBM is higher than in LAKE and closer to measurements. Results from both models are weakly correlated to measurements of surface CO₂ concentration. This leads to conclusion, that, compared to oxygen, carbon dioxide is much more controlled by biogeochemical processes misrepresented in lake models, than by physical factors.
- 10 As a result of this study, we emphasize the following issues to be addressed by the lake modeling community in respect to dissolved gases simulation:
- simplified lake modeling approaches especially employing parameterized temperature profile and representing reasonably the surface temperature may fail in calculating vertical temperature distribution with potentially significant effect on biogeochemical processes, if the respective module is coupled to thermodynamic compartment;
 - 15 – for deep lakes, models differ in the depth of autumn and especially spring overturn; if this depth reaches bottom, the gas content originated from sediments and accumulated in the hypolimnion during the preceding stratified period is almost completely released to the atmosphere, otherwise these gases may continuously accumulate in deep water at a multi-year timescale; however, more research is needed to establish the effect of lake morphometry on these processes;
 - oxygen content in both surface and deep water is efficiently controlled by physical and biogeochemical factors, successfully simulated by lake models, whereas measured surface CO₂ exhibits significant temporal variability not captured by the models, calling for more research in biogeochemical mechanisms and parameterisations of carbon dioxide dynamics.
 - 20 – 1D models have certain limitations when applying to such a 3-dimensional system as lake. They do not capture all the lake mixing mechanisms such as density driven currents, which can be important under certain conditions (Samolyubov, 1999). It was found that for deep lakes with extended littoral zone such as Lake Geneva (Fer et al., 2001, 2002), Lake
 - 25 Van (Kaden et al., 2010) these flows can be significant in terms of vertical heat, gas transfer. In particular, for Harp Lake it maybe not important due to smaller depth and not large shallow area. In winter, a large-scale convective circulation (up to 3-5 cm s⁻¹) due to heat exchange at the sediment-water interface may develop under ice (Kirillin et al., 2015). However, the importance of this circulation in terms of gas transfer have not been studied yet. So far, to the best of our knowledge, these currents are not parameterized in 1D models.

Model/Data	Code and data availability
ALBM	Available upon request to Zeli Tan, e-mail: tanzeli1982@gmail.com
MTCR-1	Available upon request to Bruna Arcie Polli, e-mail: brunapolli@gmail.com
FLake	Publicly available at the website: http://www.flake.igb-berlin.de/
LAKEoneD	Available upon request to Klaus Jöhnk, e-mail: science@limnophysics.de
LAKE	Publicly available at the website: http://tesla.parallel.ru/Viktor/LAKE/wikis/LAKE-model
Observational data	Available upon request to Huaxia Yao, e-mail: huaxia.yao@ontario.ca

5 Code and data availability

6 Author contribution

V.S., W.T., K.J., Z.T., H.Y. conceptually designed the experiments. S.G., B.A.P., T.B., W.T., K.J., Z.T., Q.Z. carried them out and provided the results of the simulations. J.A.R. and H.Y. provided observational data. S.G. made a formal analysis and combined all the data, visualized them under the supervision of V.S. S.G. prepared the manuscript under the supervision of A.L., V.S. with significant contributions from all co-authors. The authors declare that they have no conflict of interest.

Acknowledgements. Simulations performed with LAKE model have been supported by Russian Science Foundation (grant 17-17-01210). The research is partially supported by a USGS project (G17AC00276) to Q.Z. Z.T. is supported by the U.S. DOE's Earth System Modeling program through the Energy Exascale Earth System Model (E3SM) project. The supercomputing resource for ALBM model simulations is provided by Rosen Center for Advanced Computing at Purdue University. W.T. is supported by an ETH Zurich postdoctoral fellowship (Fel-45 15-1). The Uniscientia Foundation and the ETH Zurich Foundation are acknowledged for their support to this research. Thanks are given to DESC staff (e.g. R. Ingram, C. McConnell, T. Field) for contributing to data collection and to the Ontario Ministry of the Environment and Climate Change for funding the traditional and high-frequency lake monitoring activities. Simulations of MTCR-1 were supported by the brazilian funding agencies CNPQ (grant number: 308758/2017-0) and CAPES .

References

- Ashton, G. D.: Freshwater ice growth, motion and decay, in: Dynamics of Snow and Ice Masses, edited by: Colbeck, S., Elsevier, 261-304, 1980.
- Ashton, G. D.: River and lake ice thickening, thinning and snow ice formation, *Cold. Reg. Sci. Technol.*, 68, 3-19, 5 <https://doi.org/10.1016/j.coldregions.2011.05.004>, 2011.
- Balsamo, G., Dutra, E., Stepanenko, V. M., Viterbo, P., Miranda, P., and Mironov, D.: Deriving an effective lake depth from satellite lake surface temperature data: a feasibility study with MODIS data, *Boreal Environ. Res.*, 15(2), <https://doi.org/10.21957/be525ccu>, 2010.
- Bastviken, D., Tranvik, L. J., Downing, J. A., Crill, P. M. and Enrich-Prast, A.: Freshwater methane emissions offset the continental carbon sink, *Science*, 331(6013), 50, <https://doi.org/10.1126/science.1196808>, 2011.
- 10 Choulga, M., Kourzeneva, E., Zakharova, E., and Doganovsky, A.: Estimation of the mean depth of boreal lakes for use in numerical weather prediction and climate modeling, *Tellus A*, 66(1), 21295, <https://doi.org/10.3402/tellusa.v66.21295>, 2014.
- Camacho, A.: On the occurrence and ecological features of deep chlorophyll maxima (DCM) in Spanish stratified lakes, *Limnetica*, 25: 453–478, 2006.
- Cantin, A., Beisner, B. E., Gunn, J. M., Prairie, Y. T., Winter, J. G.: Effects of thermocline deepening on lake plankton communities, *Can. J. Fish. Aquat.*, 68: 260–276, 2011.
- 15 Cox, E. T.: Counts and Measurements of Ontario Lakes: Watershed Unit Summaries Based on Maps of Various Scales by Watershed Unit, Ontario Ministry of Natural Resources, 1978.
- Dillon, P. J., Reid, R. A., and De Grosbois, E.: The rate of acidification of aquatic ecosystems in Ontario, Canada, *Nature*, 329, 45-48, 1987.
- Docquier, D., Thiery, W., Lhermitte, S., and Van Lipzig, N.: Multi-year wind dynamics around Lake Tanganyika, *Clim. Dynam.*, 47(9-10), 20 3191-3202, <https://dx.doi.org/10.1007/s00382-016-3020-z>, 2016.
- Duffy C.J., Dugan H.A., Hanson P.C.: The age of water and carbon in lake-catchments: A simple dynamical model, *Limnol. Oceanogr. Lett.*, Vol. 3, No. 3, pp. 236–245, <https://doi.org/10.1002/lo12.10070>, 2018.**
- Fang, X., and Stefan, H. G.: Temperature and dissolved oxygen simulations for a lake with ice cover, U.S. Environmental Protection Agency, Environmental Research Laboratory, Duluth, Minn, 365(65), 1994.
- 25 Fang, X., and Stefan, H. G.: Long-term lake water temperature and ice cover simulations/measurements, *Cold. Reg. Sci. Technol.*, 24(3), 289-304, 1996.
- Fer, I., Lemmin, U., Thorpe, S. A.: Cascading of water down the sloping sides of a deep lake in winter, *Geophys. Res. Lett.*, 28(10), 2093-2096, <https://doi.org/10.1029/2000GL012599>, 2001.**
- Fer, I., Lemmin, U., Thorpe, S. A.: Winter cascading of cold water in Lake Geneva, *J. Geophys. Res.-Oceans*, 107(C6), 13-1, 30 <https://doi.org/10.1029/2001JC000828>, 2002.**
- Forbes G. S. and Meritt J. H.: Mesoscale vortices over the Great Lakes in wintertime, *Mon. Weather Rev.*, 112(2), 377-381, 1984.
- Futter M.N., Starr M., Forsius M., Holmberg M.: Modelling the effects of climate on long-term patterns of dissolved organic carbon concentrations in the surface waters of a boreal catchment, *Hydrol. Earth Syst. Sci.*, Vol. 12, No. 2, pp. 437–447, <https://doi.org/10.5194/hess-12-437-2008>, 2008.**
- 35 Goudsmit, G. H., Burchard, H., Peeters, F. and Wüest, A.: Application of $k-\epsilon$ turbulence models to enclosed basins: The role of internal seiches, *J. Geophys. Res-oceans*, 107(C12), <https://dx.doi.org/10.1029/2001JC000954>, 2002.

- Goyette, S., McFarlane, N. A. and Flato, G. M.: Application of the Canadian Regional Climate Model to the Laurentian Great Lakes region: Implementation of a lake model, *Atmos. Ocean* 38(3), 481-503, <https://doi.org/10.1080/07055900.2000.9649657>, 2000.
- Heiskanen, J. J., Mammarella, I., Ojala, A., Stepanenko, V., Erkkilä, K. M., Miettinen, H., and Vesala, T.: Effects of water clarity on lake stratification and lake-atmosphere heat exchange, *J. Geophys. Res-Atmos.*, 120(15), 7412-7428, <https://doi.org/10.1002/2014JD022938>, 2015.
- Henderson-Sellers, B.: New formulation of eddy diffusion thermocline models, *Appl. Math. Model*, 9(6), 441-446, 1985.
- Hondzo, M., and Stefan, H. G.: Lake water temperature simulation model, *J. Hydraul. Eng.*, 119(11), 1251-1273, 1993.
- Hostetler, S. W., and Bartlein, P. J.: Simulation of lake evaporation with application to modeling lake level variations of Harney-Malheur Lake, Oregon, *Water Resour. Res.*, 26(10), 2603-2612, 1990.
- 10 Hostetler, S. W., Bates, G. T. and Giorgi, F.: Interactive coupling of a lake thermal model with a regional climate model, *J. Geophys. Res.-Atmos.*, 98(D3), 5045-5057, 1993.
- Jammet, M., Crill, P., Dengel, S., and Friborg, T.: Large methane emissions from a subarctic lake during spring thaw: Mechanisms and landscape significance, *J. Geophys. Res-Bioge.*, 120(11), 2289-2305, <https://doi.org/10.1002/2015JG003137>, 2015.
- Jöhnk K. D. and Umlauf L.: Modeling the metalimnetic oxygen minimum in a medium sized alpine lake, *Ecol. Model.*, 136(1), 67-80, 15 [https://doi.org/10.1016/S0304-3800\(00\)00381-1](https://doi.org/10.1016/S0304-3800(00)00381-1), 2001.
- Jöhnk, K. D., Huisman, J. E. F., Sharples, J., Sommeijer, B. E. N., Visser, P. M., and Stroom, J. M.: Summer heatwaves promote blooms of harmful cyanobacteria, *Glob. Change Biol.*, 14(3), 495-512, <https://doi.org/10.1111/j.1365-2486.2007.01510.x>, 2008.
- Kaden, H., Peeters, F., Lorke, A., Kipfer, R., Tomonaga, Y., Karabiyikoglu, M.: Impact of lake level change on deep-water renewal and oxic conditions in deep saline Lake Van, Turkey, *Water Resour. Res.*, 46(11), <https://doi.org/10.1029/2009WR008555>, 2010.
- 20 Karlsson, J., Giesler, R., Persson, J., and Lundin, E.: High emission of carbon dioxide and methane during ice thaw in high latitude lakes, *Geophys. Res. Lett.*, 40(6), 1123-1127, <https://doi.org/10.1002/grl.50152>, 2013.
- Kirillin, G. B., Forrest, A. L., Graves, K. E., Fischer, A., Engelhardt, C., Laval, B. E.: Axisymmetric circulation driven by marginal heating in ice-covered lakes, *Hydrol. Geophys. Res. Lett.*, 42(8), 2893-2900, <https://doi.org/10.1002/2014GL062180>, 2015.
- Kirillin, G., Wen, L., Shatwell, T.: Seasonal thermal regime and climatic trends in lakes of the Tibetan highlands, *Hydrol. Earth Syst. Sci.*, 25 21(4), 1895-1909, <https://doi.org/10.5194/hess-21-1895-2017>, 2017.
- Kiuru, P., Ojala, A., Mammarella, I., Heiskanen, J., Kämäräinen, M., Vesala, T., Huttula, T.: Effects of climate change on CO₂ concentration and efflux in a humic boreal lake: A modeling study, *J. Geophys. Res.-Bioge.*, 123, 2212– 2233, <https://doi.org/10.1029/2018JG004585>, 2018.
- Kourzeneva, E.: External data for lake parameterisation in Numerical Weather Prediction and climate modeling, *Boreal Environ. Res.*, 15, 30 165-177, 2010.
- Lehner, B. and Döll, P.: Development and validation of a global database of lakes, reservoirs and wetlands, *J. Hydrol.*, 296(1), 1-22, <https://doi.org/10.1016/j.jhydrol.2004.03.028>, 2004.
- Lepparanta M. Freezing of lakes and the evolution of their ice cover: Springer Heidelberg New York Dordrecht London, 301 p., [doi:10.1007/978-3-642-29081-7](https://doi.org/10.1007/978-3-642-29081-7), 2015.
- 35 Long, Z., Perrie, W., Gyakum, J., Caya, D., and Laprise, R.: Northern lake impacts on local seasonal climate, *J. Hydrometeorol.*, 8(4), 881-896, <https://doi.org/10.1175/JHM591.1>, 2007.
- Mahrt L.: Surface heterogeneity and vertical structure of the boundary layer, *Bound-Lay. Meteorol.*, 96(1), 33-62, <https://doi.org/10.1023/A:1002482332477>, 2000.

- McCullough I.M., Dugan H.A., Farrell K.J., Morales-Williams A.M., Ouyang Z., Roberts D., Scordo F., Bartlett S.L., Burke S.M., Doubek J.P., Krivak-Tetley F.E., Skaff N.K., Summers J.C., Weathers K.C., Hanson P.C.: Dynamic modeling of organic carbon fates in lake ecosystems, *Ecol. Modell.* Vol. 386. pp. 71–82, <https://doi.org/10.1016/j.ecolmodel.2018.08.009>, 2018.
- Mironov, D. V.: Parameterisation of lakes in numerical weather prediction. Part 1: Description of a lake model. *Available from the author; Dmitrii. Mironov@dwd.de.*, 2003.
- Mironov, D. V.: Parameterisation of lakes in numerical weather prediction. Description of a lake model, COSMO Technical Report 11, Deutscher Wetterdienst, Offenbach am Main, Germany, 2008.
- Atlas of Alberta lakes, edited by: Mitchell, P. and Prepas, E. E., University of Alberta, 1990.
- Molot, L. A., and Dillon, P. J.: Nitrogen/phosphorus ratios and the prediction of chlorophyll in phosphorus-limited lakes in central Ontario. *Can. J. Fish. Aquat. Sci.*, 48, 140-145, 1991.
- Perroud, M., Goyette, S., Martynov, A., Beniston, M. and Anneville, O.: Simulation of multiannual thermal profiles in deep Lake Geneva: a comparison of one-dimensional lake models, *Limnol. Oceanogr.-Meth.*, 54(5), 1574-1594, <https://doi.org/10.4319/lo.2009.54.5.1574>, 2009.
- Phelps, A. R., Peterson, K. M., and Jeffries, M. O.: Methane efflux from high-latitude lakes during spring ice melt, *J. Geophys. Res.- Atmos.*, 103(D22), 29029-29036, <https://doi.org/10.1029/98JD00044>, 1998.
- Polli, B. A. and Bleninger, T.: Modelagem do transporte de calor no reservatório Vossorooca. *21 Simpósio Brasileiro De Recursos Hídricos*, 1-8, 2015.
- Polli, B. A. and Bleninger, T.: Reservoir 1D heat transport model, *J. Applied Water Engineering and Research*, <https://doi.org/10.1080/23249676.2018.1497560>, 2018.
- Poole, H. H., and Atkins, W. R. G.: Photo-electric measurements of submarine illumination throughout the year, *J. Mar. Biol. Assoc. UK*, 16(1), 297-324, 1929.
- Raymond, P. A., Hartmann, J., Lauerwald, R., Sobek, S., McDonald, C., Hoover, M., ... and Kortelainen, P.: Global carbon dioxide emissions from inland waters, *Nature*, 503(7476), 355, <https://doi.org/10.1038/nature12760>, 2013.
- Samolyubov, B. I.: Bottom stratified currents, *Nauchny Mir*, 464, 1999 (in Russian).
- Shatwell, T., Adrian, R., and Kirillin, G.: Planktonic events may cause polymictic-dimictic regime shifts in temperate lakes., *Sci. Rep.-UK*, 6, 24361, <https://doi.org/10.1038/srep24361>, 2016.
- Stefan, H. G., and Fang, X.: Dissolved oxygen model for regional lake analysis, *Ecol. Model.*, 71(1-3), 37-68, [https://doi.org/10.1016/0304-3800\(94\)90075-2](https://doi.org/10.1016/0304-3800(94)90075-2), 1994.
- Stepanenko, V. M., and Lykossov, V. N.: Numerical modeling of heat and moisture transfer processes in a system lake-soil, *Russ. Meteorol. Hydrol.*, 3, 95-104, 2005.
- Stepanenko, V. M., Goyette, S., Martynov, A., Perroud, M., Fang, X. and Mironov, D.: First steps of a Lake Model Intercomparison Project: LakeMIP, *Boreal Environ. Res.*, 15, 191-202, 2010.
- Stepanenko, V. M., Machulskaya, E. E., Glagolev, M. V., and Lykossov, V. N.: Numerical modeling of methane emissions from lakes in the permafrost zone, *Izv. Atmos. Ocean Phys.*, 47(2), 252-264, <https://doi.org/10.1134/S0001433811020113>, 2011.
- Stepanenko, V. M., Martynov, A., Jöhnk, K. D., Subin, Z. M., Perroud, M., Fang, X., Beyrich, F., Mironov, D. and Goyette, S.: A one-dimensional model intercomparison study of thermal regime of a shallow, turbid midlatitude lake, *Geosci. Model Dev.*, 6(4), 1337-1352, <https://doi.org/10.5194/gmd-6-1337-2013>, 2013.

- Stepanenko, V., Jöhnk, K. D., Machulskaya, E., Perroud, M., Subin, Z., Nordbo, A., Mammarella, I. and Mironov, D.: Simulation of surface energy fluxes and stratification of a small boreal lake by a set of one-dimensional models, *Tellus A*, 66, <https://doi.org/10.3402/tellusa.v66.21389>, 2014.
- Stepanenko, V., Mammarella, I., Ojala, A., Miettinen, H., Lykosov, V., and Vesala, T.: LAKE 2.0: a model for temperature, methane, carbon dioxide and oxygen dynamics in lakes, *Geosci. Model Dev.*, 9(5), 1977-2006, <https://doi.org/10.5194/gmd-9-1977-2016>, 2016.
- V. M. Stepanenko, I. A. Repina, G. Ganbat, and G. Davaa. Numerical simulation of ice cover of saline lakes. *Izv. Atmos. Ocean Phy.*, 55(1):129–138, <http://dx.doi.org/10.1134/S0001433819010092>, 2019.
- Subin, Z. M., Riley, W. J. and Mironov, D.: An improved lake model for climate simulations: model structure, evaluation, and sensitivity analyses in CESM1, *J. Adv. Mod. Earth Sys.*, 4(1), M02001, <https://doi.org/10.1029/2011MS000072>, 2012.
- 10 Tan, Z., Zhuang, Q., and Walter Anthony, K.: Modeling methane emissions from arctic lakes: Model development and site level study, *J. Adv. Model Earth Sy.*, 7(2), 459-483, <https://doi.org/10.1002/2014MS000344>, 2015.
- Tan, Z. and Zhuang, Q.: Arctic lakes are continuous methane sources to the atmosphere under warming conditions, *Environ. Res. Lett.*, 10(5), 054016, <https://doi.org/10.1088/1748-9326/10/5/054016>, 2015.
- Tan, Z., Zhuang, Q., Shurpali, N. J., Marushchak, M. E., Biasi, C., Eugster, W., and Walter Anthony, K.: Modeling CO₂ emissions from Arctic lakes: Model development and site-level study, *J. Adv. Model Earth Sy.*, 9(5), 2190-2213, <https://doi.org/10.1002/2017MS001028>, 2017.
- 15 Tan, Z., Yao, H., & Zhuang, Q.: A small temperate lake in the 21st century: Dynamics of water temperature, ice phenology, dissolved oxygen and chlorophyll-a. *Water Resour. Res.*, 54, <https://doi.org/10.1029/2017WR022334>, 2018.
- Thiery, W., Stepanenko, V. M., Fang, X., Jöhnk, K. D., Li, Z., Martynov, A., Perroud, M., Subin, Z. M., Darchambeau, F., Mironov, D. ... Van Lipzig, N. P.: LakeMIP Kivu: evaluating the representation of a large, deep tropical lake by a set of one-dimensional lake models, *Tellus A*, 66(1), 21390, <https://doi.org/10.3402/tellusa.v66.21390>, 2014.
- 20 Thiery, W., Martynov, A., Darchambeau, F., Descy, J. P., Plisnier, P. D., Sushama, L., and van Lipzig, N. P.: Understanding the performance of the FLake model over two African Great Lakes, *Geosci. Model Dev.*, 7(1), 317-337, <https://doi.org/10.5194/gmd-7-317-2014>, 2014.
- Thiery, W., Davin, E. L., Panitz, H.-J., Demuzere, M., Lhermitte, S., Lipzig, N.: The Impact of the African Great Lakes on the Regional Climate, *J. Clim.*, 28(10), 4061-4085, <https://doi.org/10.1175/JCLI-D-14-00565.1>, 2015.
- 25 Thiery, W., Davin, E. L., Seneviratne, S. I., Bedka, K., Lhermitte, S., and van Lipzig, N. P.: Hazardous thunderstorm intensification over Lake Victoria, *Nat. Commun.*, 7, 12786, <https://doi.org/10.1038/ncomms12786>, 2016.
- Thiery, W., Gudmundsson, L., Bedka, K., Semazzi, F. H., Lhermitte, S., Willems, P., ... and Seneviratne, S. I.: Early warnings of hazardous thunderstorms over Lake Victoria, *Environ. Res. Lett.*, 12(7), 074012, <https://doi.org/10.1088/1748-9326/aa7521>, 2017.
- 30 Tranvik, L. J., Downing, J. A., Cotner, J. B., Loiselle, S. A., Striegl, R. G., Ballatore, T. J., Dillon, P., Finlay, K., Fortino, K., Knoll, L. B. ... and Kortelainen, P. L.: Summarizing multiple aspects of model performance in a single diagram, *Limnol. Oceanogr.*, 54(6 part 2), 2298-2314, <https://doi.org/10.1029/2000JD900719>, 2009.
- Wik, M., Varner, R. K., Anthony, K. W., MacIntyre, S. and Bastviken, D.: Climate-sensitive northern lakes and ponds are critical components of methane release, *Nat. Geosci.*, 9(2), 99, <https://doi.org/10.1038/ngeo2578>, 2016.
- 35 Yan, N. and Pawson.: Changes in the crustacean zooplankton community of Harp Lake, Canada, following invasion by *Bythotrephes cederstroemi*, *Freshwater Biol.*, 37(2), 409-425, <https://doi.org/10.1046/j.1365-2427.1997.00172.x>, 1997.

Yao, H., Samal, N. R., Joehnk, K. D., Fang, X., Bruce, L. C., Pierson, D. C., ... and James, A.: Comparing ice and temperature simulations by four dynamic lake models in Harp Lake: past performance and future predictions, *Hydrol. Process.*, 28(16), 4587-4601, <https://doi.org/10.1002/hyp.10180>, 2014.

Yeates, P.S. and Imberger, J.: Pseudo two-dimensional simulations of internal and boundary fluxes in stratified lakes and reservoirs, *International Journal of River Basin Management*, 1(4), 297-319, <https://doi.org/10.1080/15715124.2003.9635214>, 2003.

5

Supporting information to the paper by
Guseva, S., Bleninger, T., Jöhnk, K., Polli, B.A., Tan, Z., Thiery, W., Zhuang, Q.,
Rusak, J., Yao, H., Lorke, A. and Stepanenko, V. "Multimodel simulation of verti-
cal gas transfer in a temperate lake"

5 1 Figures




Figure S1. Time-series of meteorological variables: air temperature, wind speed and precipitation. Black line indicates the moving average with a window of half a month.




Figure S2. Taylor diagram for the water temperature at (a) surface (b) all depths in RefSim. Grey lines indicate $RMSE_c$ ($^{\circ}C$), the radial distance from the origin is the standard deviation ($^{\circ}C$), an azimuthal position denote the correlation coefficient (Taylor et al., 2001).




Figure S3. Time-depth pattern of temperature in Harp Lake (14.07.2010-19.10.2015), LDSim and observed data.




Figure S4. Taylor diagram for the water temperature at (a) surface (b) all depths in LDSim. Grey lines indicate $RMSE_c$ ($^{\circ}C$), the radial distance from the origin is the standard deviation ($^{\circ}C$), an azimuthal position denote the correlation coefficient.




Figure S5. Time-series of the surface flux, normalized by bottom flux, and the total tracer amount in the water column: (a, b) RefSim, 37.5 m, (c, d) LDSim, 13.32 m.




Figure S6. Taylor diagram for the modeled O_2 concentration at depths of (a) 1 m (b) 18 m in GasSim.

2 Tables

Table S1. Biogeochemical processes represented in ALBM, LAKE models.

	ALBM	LAKE
Gas exchange at the air-water interface	Surface renewal model (Heiskanen et al., 2014)	Surface renewal model (MacIntyre et al., 2010; Heiskanen et al., 2014)
Sedimentary oxygen demand	(Hanson et al., 2004; Stefan and Fang, 1994)	(Walker and Snodgrass, 1986)
Photosynthesis	(Tian, 2006; Hipsey et al., 2008; Li et al., 2010)	(Megard et al., 1984; Stefan and Fang, 1994)
Respiration	(Hanson et al., 2004)	(Stefan and Fang, 1994)
Biochemical oxygen demand	(Hipsey et al., 2008; Hanson et al., 2011)	(Stefan and Fang, 1994)
DIC from inflow	(Hanson et al., 2004)	-
DOC and POC	(Hanson et al., 2004)	-

Table S2. Mean depth of mixed layer in models in RefSim and LDSim.

Mean depth of mixed layer, m ¹	ALBM	FLake	LAKE	LAKEoneD	MTCR-1	Observed
RefSim	2.4	5.6	4.8	3.9	5.9	3.6
LDSim	2.1	4.5	4.9	3.7	5.8	3.6

a. ¹ is calculated as a mean depth of a maximum for the buoyancy frequency: $N^2 = \frac{g}{\rho_0} \frac{\partial \rho(T)}{\partial z}$.

b. The period of averaging is starting from spring ice-off and ending at August, 31

Table S3. Dates of ice formation and ice melt in RefSim.

	Ice-	ALBM	LAKE	LAKEoneD	MTCR-1	FLake	Observations
2010-2011	cover	24.12.2010**	15.12.2010	06.12.2010	05.01.2011**	22.01.2011**	10.12.2010
	off	01.05.2011	03.04.2011	02.05.2011	04.05.2011*	09.04.2011**	27.04.2011
2011-2012	cover	24.12.2011	28.12.2011	10.12.2011*	29.12.2011	15.01.2012**	28.12.2011
	off	02.04.2012	12.03.2012**	23.03.2012	25.03.2012	21.03.2012*	30.03.2012
2012-2013	cover	05.01.2013**	27.12.2012**	28.11.2012**	24.12.2012*	23.01.2013**	12.12.2012
	off	22.04.2013	29.03.2013**	01.05.2013	04.05.2013*	26.03.2013**	25.04.2013
2013-2014	cover	13.12.2013*	12.12.2013*	10.12.2013*	13.12.2013*	01.01.2014**	01.12.2013
	off	29.04.2014	14.04.2014**	11.05.2014*	13.05.2014*	21.04.2014*	30.04.2014
2014-2015	cover	29.12.2014**	22.12.2014**	17.11.2014**	12.12.2014*	05.01.2015**	02.12.2014
	off	17.04.2015	02.04.2015**	04.05.2015**	07.05.2015**	15.04.2015	21.04.2015

a. Difference between the model and observations:* > 1 week ; ** > 2 weeks

b. **10.12.2011** dates denoted by bold indicate multiple freezing-melting events after the date

Table S4. Statistics of the surface temperature of the lake in RefSim, LDSim, ExtMinSim, ExtMaxSim.

Experiment		ALBM	FLake	LAKE	LAKEoneD	MTCR-1
RefSim	$h_{max} = 37.5$ m	2.13 ;	0.36 ;	1.11 ;	0.15 ;	0.68 ;
	DM (°C) ; RMSE _c (°C)	2.17	2.5	1.52	1.32	2.62
LDSim	$h_{ave} = 13.32$ m	1.1 ;	0.28 ;	1.03 ;	0.21 ;	0.16 ;
	DM (°C) ; RMSE _c (°C)	3.5	1.28	1.51	1.25	2.15
ExtMinSim	$\mu = 0.28$ m ⁻¹	2.08 ;	0.37 ;	1.15 ;	0.08 ;	0.7 ;
	DM (°C) ; RMSE _c (°C)	2.12	2.7	1.68	1.74	3.07
ExtMaxSim	$\mu = 0.68$ m ⁻¹	2.1 ;	0.38 ;	1.13 ;	0.2 ;	0.7 ;
	DM (°C) ; RMSE _c (°C)	2.25	2.48	1.51	1.02	2.03

3 Appendix

Brief description of field data at Harp Lake, Ontario, Canada

Oct. 8, 2015 (by Huaxia Yao)

1. Site and lake facts

Table S5. Chemical status of lake water.

	mg L ⁻¹		mg L ⁻¹		mg L ⁻¹
Salinity	15	DOC	1.8	DIC	1.5
Cl	2.8	TP	4.4	NO ₃	0.1
SO ₄	6.6	Ca	2.6		
silica	1.8	Na	2.2		

5 2. High frequency data from a raft buoy (10 minute intervals)

Table S6. Meteorological measurements.

	Height, m	Period of measurements	Instrument
Air temperature	1.75	July 14, 2010 – present	Vaisala HMP45C air temperature and relative humidity sensor
Relative humidity	1.75	July 14, 2010 – present	Vaisala HMP45C air temperature and relative humidity sensor
Wind speed	2	July 14, 2010 – present	RM Young wind monitor model 05103
Wind direction	2	July 14, 2010 – present	RM Young wind monitor model 05103
Atmospheric longwave radiation (earlier data can be got from another land-based station)		Nov. 5, 2012 – present	Kipp & Zonen CGR3 Pyrgeometer
Short-wave radiation (earlier data can be got from another land-based station)		Nov. 5, 2012 – present	Kipp & Zonen CGR3 Pyrgeometer
Precipitation			not measured on the raft, but available from neighboring station
Atmospheric pressure			not measured on the raft, but available from neighboring station

Table S7. Measurements in lake water.

	Depth, m	Period of measurements	Instrument
Chlorophyll-a	1	Sept. 7, 2011 –present	YSI 6025 Chlorophyll sensor
CO2 concentration	0.39	Mar. 12, 2012 –present	Vaisala GMP343 CO2 Probe with sintered PTFE filter
Electrical conductivity	1, 18	Sept. 11, 2011 –present	YSI 6560 temperature and conductivity sensor
DO concentration	1, 18	June 30, 2011 – present	YSI 6150 optical oxygen sensor
Water temperature	0.1 to 10.1 (every 0.25 m intervals between 0.1 to 10.1 m)	July 14, 2010 –present	PME thermistor T-chain with pressure transducer
Water temperature	0.1 to 27.1 (1m interval)	Mar. 12, 2012 - present	Campbell Scientific custom thermistor string of model 109 temperature probe

3. Low frequency lake data (regular monitoring) (weekly or bi-weekly for ice-free seasons)

Lake profile data: temperature and DO, at 1m interval from 0.1 m to 35m. Started late 1970s. For our selected years 2010-2015, these weekly profile measurements do not cover the winter or ice seasons. Secchi disk depth (bi-weekly or monthly): since 1978

Ice date: 1978 to 2015 Ice thickness (monthly): 1978-1993; and 2013-2015

5 Regular meteorology data (daily or hourly): land-based stations, 1978-2015.

4. Hydrology data

Stream flow discharge (daily): at 5 inlet streams (HP3, HP3a, HP4, HP5, HP6a), and the outlet (HP0), long-term data Stream water chemistry (weekly or bi-weekly): at all streams

5. Gap filling, QA/QC

10 The data are basically from raw data of the raft/buoy sensors, with a simple quality check and gap filling.

For our LMIP project, the data have been edited and gap-filled. Missing data (gaps) in atmospheric forcing are infilled mainly by using another dataset of a neighbouring climate station (backyard of DESC office site). Except for air temperature, any missed hourly data at the raft is provided by the values at the neighbouring station. The missings in air temperature are filled by regressions between the two stations which are established for each month (using available hourly data). Any doubt and concern with data are kindly reported to participants and

15 Huaxia Yao for a solution.

References

- Hanson, P. C., Pollard, A. I., Bade, D. L., Predick, K., Carpenter, S. R., and Foley, J. A.: A model of carbon evasion and sedimentation in temperate lakes, *Glob. Change Biol.*, 10(8), 1285-1298, 2004.
- 5 Hanson, P. C., Hamilton, D. P., Stanley, E. H., Preston, N., Langman, O. C., and Kara, E. L.: Fate of allochthonous dissolved organic carbon in lakes: a quantitative approach, *Plos One*, 6(7), e21884, 2011.
- Heiskanen, J. J., Mammarella, I., Haapanala, S., Pumpanen, J., Vesala, T., MacIntyre, S., and Ojala, A.: Effects of cooling and internal wave motions on gas transfer coefficients in a boreal lake, *Tellus B*, 66, 22827, doi:10.3402/tellusb.v66.22827, 2014.
- Hipsey, M. R., and Hamilton, D. P.: Computational aquatic ecosystem dynamic model: CAEDYM ver. 3. Science Manual, Cent. for Water Res. Rep., Univ. of West. Aust., Nedlands, Australia., 2008.
- 10 Li, Q. P., Franks, P. J., Landry, M. R., Goericke, R., and Taylor, A. G.: Modeling phytoplankton growth rates and chlorophyll to carbon ratios in California coastal and pelagic ecosystems, *J. Geophys. Res.-Biogeo.*, 115(G4), 2010.
- MacIntyre, S., Jonsson, A., Jansson, M., Aberg, J., Turney, D. E., and Miller, S. D.: Buoyancy flux, turbulence, and the gas transfer coefficient in a stratified lake, *Geophys. Res. Lett.*, 37, L24604, doi:10.1029/2010GL044164, 2010.
- 15 Megard, R. O., Tonkyn, D. W., and Senft, W. H.: Kinetics of oxygenic photosynthesis in planktonic algae, *J. Plankton Res.*, 6(2), 325-337, 1984.
- Taylor, K. E.: Summarizing multiple aspects of model performance in a single diagram, *J. Geophys. Res-Atmos.*, 106 (D7), 7183-7192, 2001.
- Tian, R. C.: Toward standard parameterizations in marine biological modeling, *Ecol. Model.*, 193(3-4), 363-386, 2006.
- Stefan, H. G., and Fang, X.: Dissolved oxygen model for regional lake analysis, *Ecol. Model.*, 71(1-3), 37-68, 1994.
- 20 Walker, R. R. and Snodgrass, W. J.: Model for Sediment Oxygen Demand in Lakes, *J. Environ. Eng.*, 112, 25-43, doi:10.1061/(ASCE)0733-9372(1986)112:1(25), 1986.