

## Responses to Referee #1

We highly appreciate your time in reviewing our manuscript for publication in HESS. The relevant comments have improved the quality of the manuscript. We now revised the manuscript accordingly. The detail responses are listed below. The modifications in the article are marked in the annotated manuscript.

Q1. ... It remains a difficult issue to evaluate results of the analysis of large spatial datasets (like airborne EM) with individual boreholes, and I would suggest that the evaluation of the method focusses some more on the geological plausibility of the results.

**Re:** We now enhanced the description of geological background of palaeovalley system in both “Study area” (Line 71-85 in annotated manuscript) and “training data generation” (Line 144-166), to provide a more detail geological background to support the predicted palaeovalley system.

Q2. Line 32: Both (x,y) and z?

**Re:** It here means x-y plane, which is now clarified in Line 32.

Q3. Line 52: Gunnink and Siemomn, 2015 did not use ANN to classify AEM

**Re:** This is now deleted.

Q4. Line 82: Is 15m arbitrary chosen? Are there no clays in the alluvial setting to be expected?

**Re:** The value of 15m is the observed maximum thickness of eolian sediments (now Line 79). Because it is difficult to differentiate from the borehole logs the eolian from the alluvial sands, sediments with less than 15 m of sands are classified as transition between alluvium and Aeolian sands.

Analysis of the borehole logs shows that clays present in the alluvial deposits. In our analysis of borehole logs, alluvium is defined on the basis of the relative thickness of coarse to fine-grained sediments within the borehole. If coarse sediments represent more than 50% of the total depth, then the entire borehole depth is considered as alluvium. If, on the other hand, coarse sediments exist in less than 50% of the borehole, the entire borehole depth is considered as transition between alluvium, Aeolian and bedrock. We now rephrased Line 90-94.

Q5. Line 94: Justify the dimensions of the spatial resolution.

**Re:** The constraint on the lateral resolution of the AEM data is determined by the line spacing of the survey. In the case of the APY lands survey with a survey line spacing of 2 km, it was gridded to a cell size of one fifth of the line spacing – hence the 400m value, to maintain a high fidelity. This is a common approach taken in geophysics. The depth interval defined is somewhat arbitrary, but is commonly between 5-10 m increasing exponentially with depth (Because AEM is a diffusive technology). In the APY lands a vertical interval starting at 10 m is selected and remains unchanged in the first 100 m depth, to avoid generating too many interval conductivity slices (Now Line 103-109).

Q6. Line 95: From this I conclude that the analysis will basically be 2D, per slice. If so, that should be mentioned here explicitly.

**Re:** This is now clarified in Line 107-109.

Q7. Figure1 caption: what does dS mean?

**Re:** dS or deciSiemens is a unit of electrical conductivity and often used in groundwater science. But it seems not necessary in this article and is now deleted.

Q8. Line 104: supply reference.

**Re:** References are now added in Line 119.

Q9. Line 106: You should also mention the EC differences between clay / shales and sands / sandstone. It is not just groundwater that controls EC!

**Re:** Indeed. This is now mentioned in Line 119. Since both Archie's Equation and the weighted average method applied to calculate bulk EC in this study considered the EC in both water and matrix (solids), we now deleted the sentence in Line 121 to avoid confusion.

Q10. Line 110: What about the spatial variability of EC in the paleo-valley itself, as mentioned on page 4: "variable water quality from upper reaches (lower salinity) to down-valley (higher salinity)"

**Re:** The description in page 4 is rather a general trend in palaeovalleys across entire Australia (this statement is now deleted). In this specific study area, we found only a limited number of water salinity measurements, from which an approximate EC range in palaeovalleys was obtained. These measurements didn't show a particular trend of EC variation within the palaeovalley. We assume a uniform distribution of EC in both palaeovalleys and non-valley areas (now Line 190-191), respectively, without considering the spatial variation of EC in palaeovalleys.

Q11. Line 114: These changes might also be due to salinity changes in the groundwater in the paleo valley.

**Re:** Agree. This is now added in Line 129.

Q12. Line 129-130: This is a rather strong statement. Although in general true, you need to elaborate more on the validity in this specific case. E.g. what about larger discharges in the past that might have caused larger dimensions? Or a much lower erosional base, causing deeper incisions instead of broad valleys? Can climatic and geomorphological arguments help in justifying this statement? Line 132: Are the current "topographical lows" oversized, given the current river discharges? If so, this might be an additional argument for the statement that the current river valley network is an analogue for the paleo-system.

**Re:** We have rewritten this paragraph to accommodate these comments (see Line 144-166).

Q13. Line 136: why 80km x 80 km; any special reason?

**Re:** There is no special reason. It was mainly selected on the basis of the AEM-survey data availability.

Q14. Line 137: How was the DEM downscaled from 30x30 to 100x100; averging, min, max, moving window?

**Re:** The downscaling here is completed by a bicubic interpolation, which is mentioned now in Line 158-160.

Q15. Why was cropping to 50x50 necessary?

**Re:** The large training image represents a rather specific feature of the valley system. However, if we crop it into small training images and rotate by different angles, the combination of these small images allows to create more valley patterns. In this manner, the training image pool can be more representative. This is now described in Line 164-166.

Q16. Line 164: Was a statistical distribution of the EC values assumed or was the random sampling from a uniform distribution?

**Re:** They were assumed to follow the uniform distribution (now Line 190-191).

Q17. it was stated before that the bulk EC in the non-valley was  $< 2.5$  mS/m.

**Re:** This is now explained in Line 192-197. We added the overlap in EC values between valley and non-valley in two sequential steps: first step is to enlarge the EC range in bedrocks, and then to enhance the overlap by downscaling and interpolation. The first step is to add noise relating to several uncertainties (e.g. land cover, local weathering of bed rocks, etc.) in AEM survey and interpretation, while the subsequent smoothing is to address the spreading of this noise. The noise added in training images would enhance the prediction capability of the neural network, as the neural network has the opportunity to learn to identify these noises. But if too much noise is added in training image, the neural network may overestimate the noise and make the palaeovalley disconnected. The selection of the size of the overlap and its influence on the output is discussed in Appendix A1 in Line 490-531.

Q18. Line 169: I assume that the size of the images is the same, only the pixels get larger (smaller number of pixels with same image size leads to larger pixels). If so, I do not understand how nearest neighbor interpolation leads to blurred images. In my experience, nearest neighbor creates more crisp images, because gradients in the original images are not expressed in the nearest neighbor images.

**Re:** We now rephrased this sentence in Line 198-202. The downscaling was operated by the nearest neighbour method, while the downscaled images were interpolated back to their original size by bicubic interpolation.

Q19. Line 173: Why using pre-interpolation images?

**Re:** We try to make the training images representative of multiple degrees of smoothing.

Q20. Line 201: Is this a probability?

**Re:** It is not a true probability, although a probability distribution function is created from the values 0-1. It is an index indicating presence of a palaeovalley when the value approaches 1.0 and non-valley when values approaches 0.0. This is now added in Line 232-234.

Q21. Eq. 6 What does e mean?

**Re:** It represents an exponential function, we now modified this equation.

Q22. Line 205: How are these filter sizes determined and how do they influence the results? Line 207: Finally, spatial correlation is mentioned! It should have been mentioned before and taken into account in the generation of the training images. How is the spatial correlation length determined?

**Re:** We now rephrased Line 239-248 to address these two issues.

The filter size is originally selected according to the classical SRCNN structure proposed by Dong et al 2016, and then adjusted by trial-and-error test as explained in Appendix A3 (Line 555-588), and also the training image size as discussed in Appendix A2 (Line 532-553).

The spatial correlation length is an unknown, and is not explicitly calculated in the convolution neural network. We use a large filter to guarantee that the spatial correlation in the images can be accounted for and the accuracy of resulting binary palaeovalley is sufficiently high. A two-layer example is used in Line 242-246 to demonstrate the relationship between filter size and correlation scale incorporated in the neural network.

Q23. Line 211: What is meant by "width"?

**Re:** It is the number of images contained in each layer (now line 239).

Q24. Line 267: what is the exact value? How does the choice of this value influence the results?

**Re:** A threshold value of 0.5 is used here. As the palaeovalley index follows a bimodal distribution with values grouped near 1.0 and 0.0, the selection of a threshold value in the range 0.2 to 0.8 doesn't really affect the classification of the palaeovalley significantly (Now Line 302-305).

Q25. Fig. 5: It seems from histogram (e) that the number of pixels with value  $> 0.8$  is much smaller than number of pixels between 0 and 0.2. But, Figure (d) and (f) suggests otherwise....

**Re:** The histograms represent statistics of all 4000 images (now in caption of Fig.5), and many images have a very narrow palaeovalley.

Q26. Line 287 This seems a small overlap. What happens when the overlap is larger, as is often the case in "real life"?

**Re:** We here arbitrarily use an overlap of 2.5% to test if the trained neural network can identify the palaeovalley pattern properly. After smoothing, the overlapping size could exceed 2.5%. For example, in the training images, the artificial overlap size is given as 5%, but after smoothing, as shown in the histogram of Fig. 2d, the EC value in non-valley areas in training images can even reach 40 mS/m and in the valley zone can reach 1 mS/m. The maximum overlapping size addressed in training images can reach 50%. We discuss the influence of overlapping size used in training and test images on the output in Appendix A1 in Line 490-531.

Q27. Fig. 8c Here again, it seems that the proportion  $<$  and  $> 0.5$  is not visible in the map; From (a) to (b) seems an increase in Ec value, is that correct and if so, is that what you want?

**Re:** The threshold value of 0.5 is now marked in Fig. 8c. There is a normalization operation from EC values ranging from 1 to 80 mS/m (a) to values in 0 to 1.0 (b).

Q28. Line 338: this implies that there is no trend in depth. Is that reasonable to assume?

**Re:** We now rephrased this sentence in Line 374-376.

As the training images were generated from a DEM, training, validation and prediction of the neural network in this study were based on 2D images, and the trained neural network is employed layer by layer to generate a quasi-3D image of the palaeovalley. Since training images include palaeovalleys with varying geometries (e.g. width, curviness, length...), the palaeovalley pattern with varying geometry in each layer can be generated, determined by the EC values.

However, one limitation is that the vertical relationship between EC and palaeovalley pattern cannot be addressed in such 2D image-based neural network. This is now mentioned in Line 474-480, and also future works that potentially overcome this problem.

Q29. Fig. 9. There seems to be quite a lot of paleo valleys. Is that realistic?

**Re:** The palaeodrainage system in arid zone of Australia is very developed. The more detail geological background on the formation of these palaeovalleys are now described in Line 71-85.

## Responses to referee #2

Thanks a lot for your suggestions on this manuscript, which definitely helped us improve the article. We now modified the manuscript accordingly, and the detail responses are listed below.

Q1. The geolithological description is very synthetical. More details should be given and can you provide the image of a bore log for coherence with the described lithological classes?

Re: We now added in the article the history of palaeovalley formation in the arid zone of Australia and described the general lithology facies of sedimentary infills in the palaeovalley (now Line 71-85 in annotated manuscript, and Fig. S1 in the attachment). Generally, the coarse sediments were deposited at the bottom of the palaeovalley in a fluvial environment, overlain by fine-grained sediments deposited in lacustrine environment. In the Quaternary, these palaeovalley sediments were partly covered by aeolian sediments with a maximum thickness 15 m. Although the borehole logs show a similar trend, those fine and coarse sediments often interbedded with each other, and it is also difficult to differentiate between the aeolian sands and fluvial sands without particle size analysis. We thus classify borehole logs according to the portion of coarse sediments (i.e. sands) and fine (i.e. silts and clays) sediments, and also the thickness. More detail principles for classification are now given in Line 89-94 in the annotated manuscript. We hope that these new explanations can make the interpretation of borehole logs more sound.




Fig. s1. Typical sedimentary facies in palaeovalley in arid zone of Australia (modified according to Magee 2009)

Q2: My main concern is about the scale effect of the AEM measurement to small resolution data, such as bore scale. Can you discuss more about the uncertainty in the bicubic algorithm for image scaling and the calibration/validation of the methodology with real data?

Re: In this study, the bicubic algorithm is employed on images of electrical conductivity, which are the input of the neural network. The binary palaeovalleys, as output, are then generated by the neural network. We could use other interpolation methods on the electrical conductivity image, e.g. kriging. However, as long as the interpolation method used in the training image generation is consistent with the prediction, the quality of the output binary palaeovalley is mainly determined by the structure of the trained neural network (e.g. filter size, crop size, depth and width), rather than by the interpolation method, as the neural network has learned to filter

out the errors induced by the interpolation. The sensitivity of output binary palaeovalley to these parameters is discussed in Appendix A of the manuscript in Line 555-588 of annotated manuscript.

To make sure that the trained neural network works well, we first tested it based on synthetic examples.

Regarding the scale effect, it is illustrated in section 4.3 (Line 320-355) that the methodology used in this study allows upscaling of the original electrical conductivity image to binary palaeovalleys by ten-times in resolution, without significant loss in accuracy.

When the method is applied to the study area, we don't have many hard information to validate our results on, apart from the borehole logs. The AEM-derived electrical conductivity images were classified into binary palaeovalleys with a spatial resolution in horizontal plane of 40 m × 40 m. For validation, it is assumed that the control area of each borehole is also 40 m × 40 m. This is a reasonable assumption as the short-distance lateral stratigraphic heterogeneity is considered small over a few tens of meters.

Q3: Instead of summarizing the results, the conclusion should focus more on the novelty of the approach compared to other traditional methods. I would suggest to insert the part 'Future work' inside the conclusion to critically discuss the limit of the approach and the proposed future studies to improve it.

Re: We rewrote the conclusion and merged the limitations and future work in the conclusions section (now Line 456-482 in annotated manuscript).

# High-resolution palaeovalley classification from airborne electromagnetic imaging and deep neural network training using digital elevation model data

Zhenjiao Jiang<sup>1,2\*</sup>, Dirk Mallants<sup>2</sup>, Luk Peeters<sup>3</sup>, Lei Gao<sup>2</sup>, Camilla Soerensen<sup>3</sup>, Gregoire Mariethoz<sup>4</sup>

5 <sup>1</sup>Key Laboratory of Groundwater Resources and Environment, Ministry of Education, College of Environment and Resources, Jilin University, Changchun, 130021, China

<sup>2</sup>CSIRO Land & Water, Locked Bag 2, Glen Osmond, SA 5064, Australia

<sup>3</sup>CSIRO Mineral Resources, Locked Bag 2, Glen Osmond, SA 5064, Australia

10 <sup>4</sup>University of Lausanne, Faculty of Geosciences and Environment, Institute of Earth Surface Dynamics, Lausanne, Switzerland

*Correspondence to:* Zhenjiao Jiang (jiangzhenjiao@hotmail.com)

**Abstract.** Palaeovalleys are buried ancient river valleys that often form productive aquifers, especially in the semi-arid and arid areas of Australia. Delineating their extent and hydrostratigraphy is however a challenging task in groundwater system characterization. This study developed a methodology based on the deep learning super-resolution convolutional neural network (SRCNN) approach, to convert electrical conductivity (EC) estimates from an airborne electromagnetic (AEM) survey in South Australia to a high-resolution binary palaeovalley map. The SRCNN was trained and tested with a synthetic training dataset, where valleys were generated from readily available digital elevation model (DEM) data from the AEM survey area. Electrical conductivities typical of valley sediments were generated by Archie's Law, and subsequently blurred by down-sampling and bicubic interpolation to represent noise from the AEM survey, inversion and interpolation. After a model training step, the SRCNN successfully removed such noise, and reclassified the low-resolution, converted unimodal but skewed EC values into a high-resolution palaeovalley index following a bimodal distribution. The latter allows distinguishing valley from non-valley pixels. Furthermore, a realistic spatial connectivity structure of the palaeovalley was predicted when compared with borehole lithology logs and valley bottom flatness indicator. Overall the methodology permitted to better constrain the three-

15

20



dimensional palaeovalley geometry from AEM images that are becoming more widely available for groundwater prospecting.

25 **Keywords:** Deep learning; Convolutional neural network; Palaeovalley; Airborne electromagnetics; Electrical conductivity.

## 1 Introduction

A palaeovalley is the remnant of an inactive ancient river valley filled by unconsolidated, semi-consolidated or lithified sediments, which often have a higher porosity and permeability than the surrounding rocks (Jackson, 2005). Palaeovalleys are important in mineral exploration as they may contain remobilized gold, uranium and heavy minerals (Hou et al., 2008) and in groundwater exploration, as they often form productive aquifers (Samadder et al., 2011; Mulligan et al., 2007; Knight et al., 2018). However, delineating the geometry and connectivity of palaeovalleys at the regional scale (tens to hundreds of kilometers) with a high resolution (tens of meters in horizontal plane) is challenging (Holzschuh, 2002; Lane, 2002). This is mainly because surface geophysical surveys and borehole data often do not yield the required spatial resolution and coverage to reliably and cost-effectively map connected palaeovalleys at a regional scale.

35 Aerial geophysical surveys, such as airborne electromagnetic (AEM) surveys, can be inverted to provide depth profiles of electrical conductivity (EC) at regional scale over tens to hundreds of kilometers (Fitterman et al., 1991). Their horizontal resolution depends on the distance between flight lines (typically between 250 m to 30 km), which can be tailored to the problem at hand, while vertical resolution ranges from meters to tens of meters. Classification of geophysical properties into palaeovalleys and non-valley zones is most often done manually, although several methods have been developed to automate the identification of lithofacies from electrical conductivity estimates. Most of these methods assume a simplified petrophysical relationship between electrical conductivity and hydraulic parameters (e.g. porosity and permeability) (Vilhelmsen et al., 2014; Marker et al., 2015; Pollock and Cirpka, 2010). Using synthetic borehole data, Christensen et al. (2017) converted AEM data to lithofacies at a scale of kilometers by use of Markov Chain Monte Carlo and sequential indicator simulation methods.

45 Electrical conductivity values estimated from AEM surveys are subject to uncertainties introduced by variations in land cover during surveys, inversion processes, and the interpolation of EC values to the required resolution (Viezzoli et al., 2008; Robinson et al., 2008). Consequently, the relationship between EC and lithofacies is complex and difficult to identify. In this

paper, we introduce a deep-learning (neural network)-based methodology (including training dataset generation, and neural network construction and training) for automatic classification of high resolution binary palaeovalley maps from AEM-derived EC data with noise.

50 Artificial neural networks (ANN), which can express the complex and nonlinear relationship between input and outputs, were previously applied for the inversion of EC values from original AEM data (Ahl, 2003) and to classify lithology from AEM-derived EC data (Gunnink et al., 2012). However, the large number of weights involved in ANN make it difficult to train the network and often leads to overfitting problems (Tu, 1996). Deep learning approaches based on convolutional neural networks with sharing weights were established in 2006 (Gu et al., 2017), and are now well accepted in the field of visual recognition,  
55 speech recognition and language processes. They provide efficient high-dimensional interpolators that cope with multiple scales and heterogeneous information (Marcais and de Dreuzy, 2017), and have been applied in geoscience for earthquake detection based on seismic monitoring (Perol et al., 2018), object and disaster recognition from remote sensing data (Långkvist et al., 2016; Amit et al., 2016), and mineral prospectivity evaluation by the fusing of different geophysical datasets (Granek, 2016; Meller et al., 2013). Furthermore, a super-resolution convolutional neural network (SRCNN) approach composed merely  
60 of convolutional layers was established to directly capture the relationship between low- and high-resolution images (Dong et al., 2016). The SRCNN was found to be accurate, robust and fast for removing noise from low-resolution images and reconstructing a super-resolution image (Hao et al., 2018; Tuna et al., 2018; Luo et al., 2017).

In this study, concepts from the SRCNN approach are used to identify palaeovalleys at high spatial resolution from a regional scale AEM survey. The objective is to develop a methodology based on SRCNN to generate a high-resolution, regional scale  
65 map of palaeovalleys from low-resolution AEM-derived EC data, that (1) reproduces palaeovalley connectivity and (2) accounts for noise in the EC data. The method is applied to an arid region of South Australia to identify palaeovalleys at depths up to 100 m, i.e. the depth up to which the AEM-derived EC has a sufficient signal to noise level. The paper is organized as follows; Section 2 presents the data availability in our study area. Sections 3 introduces the methodology, which is followed by performance analyses in Section 4. Section 5 concludes the major findings.

Australian landscapes are ancient, featuring the product of subdued tectonics, long-term subaerial exposure and an extremely limited extent of Quaternary glaciation. This often manifests itself in an extensive palaeovalley network with deep weathering profiles and thick accumulation of unconsolidated alluvium and colluvium. The widespread palaeovalley networks in today's arid landscape are remnants of the Early Cenozoic inset valleys with Tertiary sedimentary infill and a thin and variable Quaternary cover (Magee, 2009). In the intra-continental Cenozoic sedimentary basins, palaeovalley infill sediments typically consist of Eocene sediments overlain by finer-grained sediments of Oligocene to Miocene age. The Eocene sediments are dominantly coarse-grained fluvial sands and basal gravels, deposited under wet climatic conditions. The Oligocene to Miocene sediments were deposited by relatively lower-energy drainage systems under drier climatic conditions. During the Quaternary, eolian sediments with maximum observed thickness of 15 m covered portions of the palaeovalleys at a time when fluvial or lacustrine deposition had ceased (Magee, 2009).

This study focuses on the Anangu Pitjantjatjara Yankunytjatjara (APY) Lands, which are part of the Musgrave Province in northern South Australia (Fig. 1). This area features an arid climate with very low and unreliable rainfall averaging about 230 mm/year (Jones et al., 2009). However, an extensive palaeovalley system with sedimentary faces aligning with the Cenozoic sedimentary basins above represents a shallow dynamic groundwater system exhibiting reliable water resources for local communities and mining (English et al., 2012; Munday et al., 2013).

Within the study area 128 bores, drilled between 1970 and 2018, with lithological information were retrieved from the South Australia Government Waterconnect database ([www.waterconnect.sa.gov.au](http://www.waterconnect.sa.gov.au)). Three lithological classes were derived from the logs:

- (1) bedrock: basement at surface or covered with in-situ weathering products (clays, broken basement fragments);
- (2) alluvium (sediments in palaeovalleys): basement covered with more than 15 m of unconsolidated sediments consisting of sand and gravel with minor silt and clay, showing indication of alluvial sediment transport;
- (3) transition: basement covered with up to 15 m of eolian sands or lacustrine sediments consisting of silt and clay with minor amounts of sand or gravel, showing limited indication of transport.

95 While the information content in these logs was often limited, they provided independent lithological data to verify the predicted palaeovalley network in this study (see further).

Two AEM surveys were flown in the APY Lands in 2016, covering a total area of 33,500 km<sup>2</sup> and featuring a line spacing of 2 km in the north-south direction (Soerensen et al., 2016). An area 80 km by 80 km in a central-east section of the APY Lands is selected and used to test mapping palaeovalleys based on a SRCNN analysis of electrical conductivity (EC) generated from the AEM survey (Fig. 1). In this area, the AEM survey was undertaken using the helicopter borne SkyTEM<sup>312FAST</sup> system (Soerensen et al., 2016). The averaging trapezoidal filter was used to reduce the noise in low- and high- moment amplitude response data. Aarhus Workbench software was used to invert AEM data to obtain EC (Auken et al., 2009; Auken et al., 2014). In a final step, ordinary kriging was used to interpolate EC values to a spatial resolution of 400 m × 400 m in the horizontal plane and 10 m in the vertical cross section (Ley-Cooper and Munday, 2013; Soerensen et al., 2016). The constraint on the lateral resolution of the AEM data was determined by the line spacing of the survey (2 km). In the APY lands, it was gridded to a cell size of fifth of the line spacing (i.e. 400 m), to maintain the fidelity. The depth interval is commonly between 5-10 meters increasing exponentially with depth, because AEM is a diffusive technology (Yang et al., 2013; Spies, 1989). In the APY lands the vertical resolution is 10 m for the first 100 m depth interval to avoid generating too many interval conductivity slices. Only the EC values in the first 10 depth slices, up to 100m depth, are used in this study to construct the binary palaeovalley pattern per slice, which are then stacked up to a quasi-3D image of palaeovalley. ~~The first 10 depth slices, up to 100m depth, are used in this study.~~

100

105

110




Figure 1: (a) Electrical conductivity at 100 m depth ranging from 1 to 80 mS/m ( $1 \text{ mS/m} = 0.01 \text{ dS/m}$ ), as interpreted from airborne electromagnetic surveys in the Anangu Pitjantjatjara Yankunytjatjara (APY) Lands, Australia; (b) inset shows detail of EC map at a spatial resolution of  $400 \times 400 \text{ m}$  (Soerensen et al., 2016).

115

Bulk electrical conductivity of the subsurface depends on both the solid phase (i.e., the rock mass) and the liquid phase (i.e., soil water and groundwater). It is further influenced by the porosity, tortuosity of the pore space, and degree of water saturation. Unweathered rocks are generally a poor electrical conductor with EC values typically less than 1 mS/m for igneous and metamorphic rock, and 1 to 1000 mS/m for regolith (e.g. gravel, sand, silt and clay) (Lane, 2002); saline groundwater with a similar salinity level as seawater has an EC of around 3000-5000 mS/m, while freshwater EC is up to 150 mS/m (Lane, 2002; Rhoades et al., 1976; Purvance and Andricevic, 2000). As a result, variations in bulk EC are usually controlled by groundwater quantity and salinity.

120

Previous hydrogeological characterization studies in the APY Lands study area indicated that palaeovalley porosity values are relatively high (from 10 % to 30 %) with the mean salinity of the pore fluid reaching 4500 mg/L (700 mS/m) EC; the surrounding rocks (fractured granites and gneiss) have a much lower porosity (< 1%) and water salinity values (< 1000 mg/L,

125

160 mS/m ~~EC~~) (Varma, 2012; Taylor et al., 2015). It is reasonable to assume that a clear distinction exists in this study area between EC values of the valley and non-valley lithologies, and thus only EC is used to distinguish palaeovalleys from surrounding basement. However, due to the data smoothing methods used during inversion of the AEM data and EC interpolation, and the continuous variation of water salinity near the interface between palaeovalley and fractured bedrocks, the resulting EC values vary continuously (Fig. 1b), which makes the boundary between valley and non-valley lithologies rather diffuse. Our novel methodology allows to automatically identify the boundaries between valley and non-valley lithologies based on convolutional neural networks.

### 3 Methodology

The method developed in the present study to identify palaeovalleys is comprised of three key steps: (1) A deep neural network training dataset is generated by creating synthetic palaeovalley networks from a digital elevation model (DEM) of the study area; the palaeovalley network is converted to EC values by applying Archie's Law (see further) to the water bearing formations, while EC values for the non-valley zone composed of fractured bedrock are obtained as a volume weighted average of EC values of rock and fluid components; (2) the SRCNN is trained and validated using the synthetic EC and corresponding palaeovalleys to remove noise and establish a non-linear relationship between EC image and palaeovalley image; (3) the SRCNN is then applied to predict the palaeovalley in the APY Lands based on measured AEM data. The algorithm of training dataset generation and SRCNN, and the performance metrics to evaluate SRCNN palaeovalley classification are described in detail below.

#### 3.1 Synthetic training data generation

Australia is well-known for its relative tectonic stability and is a stable continent located in an intraplate position. The palaeovalley networks are coherent, dominantly dendritic, and largely concordant with modern topographic expression (Magee, 2009). Although palaeovalleys in the arid zone are partly covered by Quaternary eolian deposits, the topographic expression of the palaeovalley pattern is still evident in high-resolution digital elevation model (DEM) data (Magee, 2009). In the APY

150 Lands, crustal architecture has been preserved since the Cenozoic, and it is considered to have been unaffected by later tectonic events (Drexel and Preiss, 1995). Previous studies in the study area have considered that the palaeovalleys are coincident with topographic lows that characterize the contemporary landscape, with AEM images being particularly useful for locating the position of the deeper portions of the older valley system (Munday et al., 2013). It is thus assumed that the present-day valley pattern indicated by topographic lows in the study area is comparable to the palaeovalley pattern according to the principle of uniformitarianism (Simpson, 1970), but shifts in valley width, orientation and connectivity between present-day valley and palaeovalley are allowed. Following this principle, we generate synthetic palaeovalley image based on digital elevation model (DEM).

155 First, a DEM of the study area with a resolution of  $30 \times 30$  m (<https://earthexplorer.usgs.gov/>) is used to generate 15 sets of palaeovalley images, mimicking palaeovalleys of various spatial densities and width over an area of  $80 \times 80$  km based on the hydrological analysis in ArcGIS (Fig. 2a) (details in Maidment and Morehouse, 2002). For convenience in the subsequent neural network operation, each resulting valley image is downscaled by the bicubic interpolation method to contain  $800 \times 800$  pixels with spatial resolution of 100 m. Valley widths range from 1 to 10 pixels (i.e. 100 m to 1000 m). The 15 images generated from the DEM were rotated between zero and 360 degrees and randomly cropped into 20,000 small training images with a size of  $50 \times 50$  pixels (Fig. 2b). Thus, the potential differences in the width and orientation of present-day valley and palaeovalley induced by several uncertain factors, e.g. variation of the river discharge and geomorphology, can be addressed in the training images. The re-combination of small training images allows recreation of valley patterns beyond those 15 full-size images generated from DEM data. A broad range of likely palaeovalley patterns at varying principle orientations, widths and connectivity are available in the SRCNN training image pool.

165 The properties in the porous palaeovalley sediments are then converted to EC values using Archie's Law (Archie, 1942):

$$R = R_0 \theta^{-m}, \quad (1)$$

170 where  $R$  is the electrical resistivity of the water-bearing formation (ohm.m),  $R_0$  is the electrical resistivity of the pore water relating to water salinity (ohm.m),  $\theta$  is the porosity,  $m$  is a constant relating to the lithology (with value ranging from 1.8 to 2.0) (Worthington, 1993). Electrical conductivity values are calculated as the inverse of resistivity values (i.e.  $EC=1/R$ ). In the


present study area,  $R_0$  is considered to range from 1.4 to 1.7 ohm.m, corresponding to water salinities of 3,000 to 6,000 mg/L (Varma, 2012), while  $\theta$  is considered to range from 10% to 30% (Taylor et al., 2015;Varma, 2012). As a result, palaeovalley EC values are estimated to be within the range 6 to 80 mS/m, which are in the range of AEM-derived EC values in Fig. 1.

175 In contrast, the non-palaeovalley zone is predominantly fractured rock with solid phase EC values  $< 1$  mS/m, characteristic porosity of  $< 1$  % and fluid salinity values of  $< 150$  mS/m EC (1000 mg/L) (Olhoeft, 1981;Parkhomenko, 2012). The bulk EC values in the non-palaeovalley zones were estimated as volume weighted average of EC in fractured rock and fluid, following:

$$EC = EC_s \cdot (1 - \varphi) + EC_f \cdot \varphi, \quad (2)$$

where  $EC$  is the bulk electrical conductivity,  $EC_s$  is the EC value of rocks,  $EC_f$  is the EC of fluid, and  $\varphi$  is the ratio of fracture void volume to total volume. The resulting bulk EC values are lower than 2.5 mS/m. Again, these synthetic EC values are similar to the AEM-derived values for the presumed fractured bedrock areas (Fig. 1).





**Figure 2: Workflow diagram of methodology used to generate training and validation datasets. (a) Synthetic palaeovalley networks generated from DEM data of the study area; (b) Rotation and cropping to randomly generate 20,000 sub-images from 15 initial**

palaeovalley networks; (c) Conversion of valley images to EC values using Archie's Law in valley and weighted averaging in non-valley zone; (d) Down-sampling of resulting  $50 \times 50$  pixel EC spatial distributions to  $10 \times 10$  pixels resolution (20,000),  $20 \times 20$  pixels (20,000),  $30 \times 30$  pixels (20,000) and  $40 \times 40$  pixels (20,000), respectively and reconstructed to obtain 80,000 blurred EC images. A total of 70,000 EC images are randomly selected from original EC images (c) and reconstructed EC images (d), forming 70,000 image pairs including 20,000 binary valley images (with some EC images corresponding to the same valley image) to train the SRCNN.

Furthermore, to represent the effects from data smoothing and inherent noise associated with the AEM survey, inversion and data interpolation, artificial noise is generated by randomly sampling EC values in the non-palaeovalley zones (following a uniform distribution ranging from 1 to 10 mS/m) and in the palaeovalley zones (following a uniform distribution ranging from 6 and 80 mS/m). It is also noted that the upper limit EC values in fractured bedrock areas are enlarged artificially from 2.5 mS/m to 10 mS/m, to assure that in the training images palaeovalley and non-palaeovalley zones overlap in EC by 4 mS/m (5% of the total range of EC values between 1 to 80 mS/m) (Fig. 2c). The SRCNN can then learn to identify this overlap in EC between palaeovalley and non-valley zones. However, Appendix A1 shows that setting the overlapping size in EC too large results in the trained SRCNN overestimating the extent of the non-valley zones, which make the predicted palaeovalleys disconnected.


The overlap in EC near the boundary between palaeovalley and non-valley is further enhanced by data smoothing: the resultant EC images of  $50 \times 50$  pixels is first downsampled into images with a smaller number of pixels, i.e.  $40 \times 40$  (20,000 images),  $30 \times 30$  (20,000 images),  $20 \times 20$  (20,000 images) and  $10 \times 10$  (20,000 images) pixels, respectively, by nearest neighbor interpolation. These resulting 80,000 images are then upsampled by bicubic interpolation to yield blurred images with the original resolution of  $50 \times 50$  pixels (Fig. 2d). In this manner, the EC values in the palaeovalley and non-palaeovalley zones are smoothed and the boundary between palaeovalley and non-palaeovalley becomes blurred.

We then randomly selected 70,000 EC images from a total of 100,000 images, composed of 20,000 pre-interpolation EC images (Fig. 2c) and 80,000 reconstructed blurred EC images (Fig. 2d) with a size of  $50 \times 50$  pixels, as input to the neural network

(see further); with the original synthetic palaeovalley images (pixel code 1) and non-palaeovalley (pixel code 0) pixels (Fig. 2b), as output. From the random set of 70,000 images, 60,000 pairs of EC (Fig. 2c and 2d, as input) and palaeovalley images (Fig. 2b, as output) are used as “training dataset” for training the SRCNN. A total of 6,000 pairs are used as “validation dataset” for validation and another 4,000 are used as “testing dataset” to demonstrate the performance of the trained SRCNN in removing the noise in EC images and lithofacies (palaeovalley and non-palaeovalley) classification.

### 3.2 SRCNN algorithm

To quantify the relationship between EC images and palaeovalley images, the super-resolution convolutional neural network (SRCNN) algorithm is employed. Neural networks are regression models that provide a general way of identifying nonlinear relationships between two sets of variables (Bishop, 1996;Moyses et al., 2003), where one set of variables is considered to be the input (herein electrical conductivity) and another is a network output (binary palaeovalley). The SRCNN algorithm can directly train the relationship between a low-resolution (input) and a high-resolution image (output) (Dong et al., 2016). A typical SRCNN is composed of three convolution layers (Fig. 3), representing patch extraction and representation, nonlinear mapping and reconstruction, respectively.



**Figure 3: Algorithm of converting (a) low-resolution EC image to (b) high-resolution palaeovalley image based on (c) the super-resolution convolutional neural network. (d) Convolutional processes of data from an input image to an output image by a filter with size of 2, moving through the input image by 1 pixel at the time.**

In the patch extraction and representation layer, the input is a normalized  $50 \times 50$  pixel EC image, which is operated by a convolution process:

$$\mathbf{H}^1(\mathbf{X}) = \max(0, \langle \mathbf{X}, \mathbf{W}^1 \rangle + \mathbf{b}^1), \quad (3)$$

where  $\mathbf{H}$  represents the output images,  $\langle \rangle$  is the convolution operator,  $\mathbf{X}$  represents the input EC image,  $\mathbf{W}$  and  $\mathbf{b}$  represent the weight filter and bias, respectively.  $\mathbf{W}^1$  corresponds to  $n_1$  filters with size of  $f_1 \times f_1$  and  $\mathbf{b}^1$  is an  $n_1$ -dimensional vector. After convolution,  $\mathbf{H}^1$  contains  $n_1$  generated  $50 \times 50$  pixel images that are input into the nonlinear mapping layer. It is then convoluted by:

$$\mathbf{H}^2(\mathbf{H}^1) = \max(0, \langle \mathbf{H}^1, \mathbf{W}^2 \rangle + \mathbf{b}^2) \quad (4)$$

to generate  $\mathbf{H}^2$  composed by  $n_2$   $50 \times 50$  pixel images, where  $\mathbf{W}^2$  contains  $n_2$  filters with size of  $n_1 \times f_2 \times f_2$  and  $\mathbf{b}^2$  is a  $n_1 \times n_2$  matrix.

Finally, an output palaeovalley index (~~with values 0 to 1~~ with values approaching zero indicating a non-valley pixel and values approaching unity indicating a palaeovalley pixel) can be reconstructed from  $\mathbf{H}^2$  by:

$$\mathbf{H}^3(\mathbf{H}^2) = G(\langle \mathbf{H}^2, \mathbf{W}^3 \rangle + \mathbf{b}^3). \quad (5)$$

235  $\mathbf{H}^3$  contains one  $50 \times 50$  pixel palaeovalley index image, and  $\mathbf{W}^3$  contains one filter with size of  $n_2 \times f_3 \times f_3$  and  $\mathbf{b}^3$  is a  $n_2 \times 1$  matrix.  $G(\cdot)$  is a sigmoid function to assist the palaeovalley classification and accelerate the training processes, which is written as:

$$G(\cdot) = \exp(\cdot) / [1 + \exp(\cdot)] \quad (6)$$

In this study,  $f_1, f_2$  and  $f_3$  are referred to as filter size with values of 9, 1 and 5, respectively, and  $n_1$  and  $n_2$  are the layer width (the number of images contained in each layer) with values of 64 and 32, respectively, following the classical structure of SRCNN used in Dong et al (2016). The influence of the filter size and width on the quality of the output images was investigated in Appendix A3. The filter size in the SRCNN controls the spatial correlation length of EC values that can be considered in the neural network operator. As illustrated in Fig. 3d, in each calculation, the EC values in the filter are convoluted to form a value at a single pixel in the output image. An EC image convoluted by the filter with the size of 2 and stride of 1 (i.e. filter moving 1 pixel at the time) and 1 hidden layer, leads to a palaeovalley index at one pixel of the output image that relates to EC values from  $3 \times 3$  pixels in the input image. In this example, the spatial correlation scale able to be addressed is equal to 3 pixels multiplied by the size of each pixel (meter). In addition, the width of each layer determines the degree of the nonlinear relationship between input and output, while the depth of the network affects both the spatial correlation length and the nonlinearity (See further in Appendix A3).

240

245

The initial weight values are randomly generated, following a standard normal distribution, while initial bias values are given as 0.1. Both weight and bias values for each of the three convolutional neural network layers are optimized simultaneously using the adaptive moment estimation algorithm (Kingma and Ba, 2014) to minimize the loss function,  $L$ , which is defined as

250

the mean sum of squared residuals:

$$L(\mathbf{W}^1, \mathbf{W}^2, \mathbf{W}^3, \mathbf{b}^1, \mathbf{b}^2, \mathbf{b}^3) = \frac{1}{N} \sum_{i=1}^N \|\mathbf{H}^3 - \mathbf{Y}\|^2, \quad (7)$$

where  $\mathbf{Y}$  is the known binary palaeovalley pattern (0 represents non-palaeovalley, 1 corresponds to the palaeovalley) in the training data, and  $N$  is the number of image pixels in each training.

### 255 3.3 Performance metrics of the SRCNN algorithm

To verify the performance of the SRCNN, the following image quality indices are calculated:

(1) Peak signal to noise ratio (PSNR) (Wang and Bovik, 2002):

$$PSNR = -10 \log_{10} \left[ \frac{1}{N} \sum_{i=1}^N (\tilde{Y}_i - Y_i)^2 \right], \quad (8)$$

where  $Y$  represents the synthetic binary palaeovalley index generated from the DEM (0 for non-palaeovalley and 1 for palaeovalley) (Fig. 2b) and  $\tilde{Y}$  is the calculated palaeovalley index (Eq. 5) from SRCNN using EC images as input, the term

260 between brackets is the mean square error. PSNR is a traditional approach to image quality assessment. A high PSNR represents a high-quality palaeovalley generation, e.g. a PSNR = 20 value is equivalent to a mean square error of 0.01.

(2) Structure similarity index (SSIM) (Wang et al., 2004):

$$SSIM = \frac{2\mu_Y\mu_{\tilde{Y}} + \varepsilon}{\mu_Y^2 + \mu_{\tilde{Y}}^2 + \varepsilon} \cdot \frac{2cov(Y, \tilde{Y}) + \varepsilon}{\sigma_Y^2 + \sigma_{\tilde{Y}}^2 + \varepsilon}, \quad (9)$$

where  $\mu$  is the mean,  $\sigma^2$  the variance, and  $cov(\cdot)$  the covariance of the synthetic or calculated palaeovalley index and  $\varepsilon$  is a small number ( $10^{-6}$ ). SSIM is complementary to PSNR, but focuses on structural similarity between a reference and distorted

265 image. It ranges theoretically from 0 to 1.0. The higher the SSIM, the higher the resolution of the palaeovalley network being reconstructed.

(3) Connectivity function (Pardo-Igúzquiza and Dowd, 2003; Renard and Allard, 2013):

$$\tau(\mathbf{h}) = \frac{N(u \leftrightarrow u+h | u, u+h \in S)}{N(u, u+h \in S)}, \quad (10)$$

where  $N(u, u+h \in S)$  is the number of palaeovalley pixels in a certain direction within the distance  $h$ , while  $N(u \leftrightarrow u+h | u, u+h \in S)$  is the number of connected palaeovalley pixels in this direction. It ranges from 0 to 1.0, and high values

270 indicate a strong spatial connectivity.


## 4 Results and discussion

We here (1) monitor both PSNR and SSIM between the palaeovalley index generated from SRCNN and DEM for 60,000 training and 6,000 validation datasets to test for the overfitting problem, (2) generate palaeovalley index maps from synthetic EC images in 4,000 testing datasets to demonstrate the performance of SRCNN in identifying the noise in EC images, 275 classification and recreate the connectivity of the palaeovalley, and (3) infer binary palaeovalley maps by applying the trained SRCNN to the AEM-based EC values in the study area, and (4) compare the resulting palaeovalley image with borehole lithology logs and existing palaeovalley indicators, i.e. multiple resolution valley bottom flatness.

### 4.1 Training and preliminary testing

The training dataset composed of 60,000 pairs of EC and valley images in Fig. 2 is divided into 1,200 batches (inner number 280 of iterations) with each batch containing 50 images. The epoch (outer number of iterations) is put to 5, and the 60,000 training image pairs are resorted at the beginning of each epoch. In this scenario, weights and biases in the SRCNN are updated by 6,000 iterations ( $5 \times 1200$ ), according to the loss function calculated based on 50 pairs of images in each batch.

After each iteration, the PSNR (Eq. 8) and SSIM (Eq. 9) for 50 training images in each batch are calculated (Fig. 4a and 4b). Moreover, the PSNR and SSIM for 6,000 validation images are calculated for every 50 iterations. It is illustrated that PNSR 285 for each training batch fluctuates near 18 (which corresponds to a mean square error of 0.015 based on Eq. 8), while the SSIM stabilizes at 0.96. The PSNR and SSIM values for the validation images agree well with those of the training images. This suggests that the SRCNN is sufficiently trained to recreate the palaeovalley with a high accuracy without overfitting problems, and importantly, preserving structural similarity.



290

**Figure 4: (a) PSNR and (b) SSIM values between palaeovalley index generated from SRCNN and DEM recorded for both training (60,000 images) and validation (6000 images) dataset. For SRCNN training 50 images are used per iteration.**

#### 4.2 Performance of SRCNN for noise removal, lithofacies classification and recreating connectivity


295

The trained SRCNN is then applied to generate palaeovalley images based on 4,000 testing EC images; we here randomly selected four images to demonstrate the ability of SRCNN. The synthetic palaeovalley images from DEM and their corresponding blurred EC images are illustrated in Fig. 5a and 5b, respectively. The histogram of EC values for all 4000 images (each containing  $50 \times 50$  pixels) in the testing dataset follows a unimodal, right skewed distribution (Fig. 5c). It is not trivial to define an EC threshold value from such unimodal distribution that can be used to distinguish the palaeovalley and non-palaeovalley cells from Fig. 5b. After calibration of the SRCNN, a palaeovalley index map is obtained (Fig. 5d). However, the histogram of the resultant palaeovalley index displays a bimodal behavior, with peaks centered at 0 and 1 (Fig. 5e). By selecting a threshold palaeovalley index value of 0.5~~between 0.2 and 0.8~~, the palaeovalley and the non-palaeovalley data can be

300



305 differentiated and converted to a binary palaeovalley map (Fig. 5f). The resultant palaeovalleys compare well with the reference (i.e. synthetic) palaeovalleys in Fig. 5a. The selection of a threshold palaeovalley index in the range 0.2 to 0.8 does not have a significant influence on the resultant binary palaeovalley pattern.




310 Figure 5: (a) DEM-generated synthetic palaeovalley used as reference image in testing the SRCNN; (b) normalized electrical conductivities corresponding to the palaeovalleys following (c) a skewed distribution (based on 4000 images in the test dataset). (d) Palaeovalley index generated through processing EC images via the SRCNN; (e) bimodal distribution of palaeovalley index (based on 4000 images in the test dataset); (f) by arbitrarily selecting the palaeovalley index threshold as 0.5, the palaeovalley index map converted into a binary palaeovalley map.

315 Moreover, the resultant palaeovalley index is less noisy in both palaeovalley and non-palaeovalley parts (Fig. 5d). The SRCNN is able to create connected palaeovalley networks from the poorly connected EC values generated by bicubic interpolation (Fig. 5b), which is one of the most challenging features in geostatistics. Fig. 5 demonstrates three advantages of applying

SRCNN: (1) it removes the noise in EC values, (2) it recreates the connectivity of the palaeovalleys, and (3) it classifies the palaeovalley and non-palaeovalley components, which allows the selection of a threshold index to define palaeovalley and non-palaeovalley zones.


#### 320 4.3 SRCNN performance under different image resolutions

The next synthetic example considers 400-m wide synthetic palaeovalleys generated in ArcGIS from the DEM in the zone about 60 km southwestern to the study area in Fig. 1a. The total extent of each synthetic palaeovalley image is 80 km by 80 km, with the resolutions ranging from  $200 \times 200$  to  $2000 \times 2000$  pixels. The palaeovalley image (Fig. 6e) with  $200 \times 200$  pixels is converted to EC values based on Archie's Law (Fig. 6a), with EC values overlapping by 2.5% between palaeovalley and non-palaeovalley zones. This low-resolution EC image is upscaled to a high-resolution EC image by bicubic interpolation (Fig. 6b), which is then cropped to images of  $50 \times 50$  pixels and used as input image for the SRCNN. Subsequently, the palaeovalley index and histogram at different resolutions are obtained (Fig. 6c). Following the histogram of the palaeovalley index, it is easy to select an arbitrary threshold in the range 0.25-0.8 to convert the palaeovalley index (Fig. 6c) to a binary palaeovalley (Fig. 6d). The choice of threshold in this range does not affect the resultant binary palaeovalley pattern, as after SRCNN processing, the palaeovalley index is already well grouped. The calculation of the palaeovalley index at the resolution of  $2000 \times 2000$  pixels takes 52s.



**Figure 6: Workflow for generating a binary palaeovalley map: (a) upscaling the 200×200 pixel electrical conductivity image to (b) a 2000×2000 pixel image using bicubic interpolation; (c) SRCNN processing; (d) generating a binary palaeovalley at resolution of 2000×2000 pixels; (e) binary palaeovalley with comparable characteristics to the original synthetic palaeovalley.**

It is worth noting that as the resolution of the resultant palaeovalley increases, the PSNR and SSIM goodness of fit metrics and connectivity do not change significantly (Fig. 7). Both PSNR and SSIM increase with the resolution from 200 x 200 to 800 x 800 pixels, because the bicubic interpolation smoothed the EC values and reduces the noise in EC values. When the image resolution further increases from 800 to 2000, PSNR degrades weakly from 18.48 to 17.09 (corresponding to an increase in mean square error from 0.014 to 0.019) and, similarly, SSIM decreases from 0.8919 to 0.8522.




345 **Figure 7: The PSNR, SSIM and connectivity of palaeovalleys generated by SRCNN for different resolutions of upscaling the low-resolution EC image to a high-resolution binary palaeovalley.**

Because each image has a fixed extent of  $80 \times 80$  km, as the resolution increases, the distance between pixels and the real geological scale of  $50 \times 50$  pixels images reduces. When the resolution increases from  $200 \times 200$  to  $2000 \times 2000$  pixels, the distance between pixels reduces from 400 m to 40 m and the real scale of each training image reduces from  $20 \times 20$  km to  $2 \times 2$  km. When training the SRCNN, the distance between pixels was not accounted for. The training images in the training dataset include images without any palaeovalley to images being fully occupied by the palaeovalleys, with the narrowest palaeovalley occupying merely one pixel. These palaeovalley patterns are unrelated to the real scale of the training image, i.e. across the range from  $20 \times 20$  km to  $2 \times 2$  km. Thus, the trained SRCNN works well to infer palaeovalleys across different resolutions and scales.

#### 4.4 Application to APY Lands AEM data

Following the training and testing of the SRCNN method based on synthetic DEM-derived palaeovalley networks, we now

apply the trained network to an area in the APY lands to convert EC values at a spatial resolution of 400 x 400 m to identify palaeovalleys at a resolution of 40 x 40 m in an area of 80 x 80 km. The methodology was first applied to a single depth AEM image (i.e. 100 m) to illustrate the procedure and discuss main findings. In a second step we will apply the methodology to the AEM images from all ten depths to extract specific information on the depth structure of the palaeovalley network.



**Figure 8: Steps to derive a binary palaeovalley network in an 80 × 80 km region in APY Lands, Australia. (a) Raw EC map at a depth of 100 m; (b) EC map after bicubic interpolation; (c) palaeovalley channel index map after application of the trained SRCNN method; (d) binary palaeovalley map.**

Fig. 8 summarizes how the previously trained SRCNN successfully converts the low-resolution EC values resulting from an AEM survey to a binary map composed of palaeovalleys and non-palaeovalley areas (Fig. 8a). First, the bicubic interpolation

370 generates a high-resolution EC image characterized by a right skewed distribution of normalized EC values (Fig. 8b). This map does not yet allow a clear differentiation of the palaeovalley from the surrounding fractured rocks. However, once we apply the SRCNN, a palaeovalley index map with the palaeovalley index following a binomial distribution is produced (Fig. 8c). Selecting an appropriate index (0.5 here) separates palaeovalley from non-palaeovalley pixels (Fig. 8d).

375 Inversion of AEM derived EC maps at ten depths within the first 100 m below the land surface (at 10 m intervals) is shown in Fig. 9a. EC values available at 10 layers are converted to binary palaeovalley images by SRCNN, based on the premise that both the palaeovalley pattern and bulk electrical conductivity from the 100-m depth interval can be represented in training images for a single depth SRCNN training. As shown in Fig. 9a, normalized EC values derived from AEM survey are characterized by a right skewed distribution. However, once we apply the SRCNN, the resulting palaeovalley index map (Fig. 9b) displays a binomial distribution of palaeovalley indices. Selecting an appropriate index (0.5 here) generates a regional-  
380 scale 3D binary palaeovalley image with a horizontal resolution of 40m and vertical resolution of 10 m (Fig. 9c).





Figure 9: (a) Rescaled AEM-derived EC map and corresponding histogram of normalized EC values within the depth interval of 100 m in an 80 × 80 km region in APY Lands, Australia; (b) palaeovalley indices map and corresponding histogram after application of the trained SRCNN; (c) binary palaeovalley map.

385

In the subsequent discussion we first test the derived palaeovalley map with independent, yet limited, borehole data and auxiliary land surface maps. Next we extract further information from Fig. 9c about the depth structure of the palaeovalleys to better constrain the areas for groundwater prospection.



390

**Figure 10: (a) SRCNN palaeovalley index by aggregating the binary palaeovalley in the vertical direction within 100 m, (b) multiple resolution valley bottom flatness (MRVBF) indicating the position of alluvium sediment accumulation. The black lines showing the boundary between palaeovalley and non-palaeovalley interpreted from (b), while the white lines representing the contour lines of SRCNN palaeovalley index from (a). The box-plot of SRCNN palaeovalley index (c) and MRVBF (d) with respect to the boreholes logs showing bedrock, alluvium and transition between bedrock and alluvium.**

395



To compare the palaeovalley map (Fig. 9c) with borehole logs and an alternative indicator of the location of valley in the land surface (i.e. the Multiple Resolution Valley Bottom Flatness index, Gallant and Dowling, 2003), we aggregated the ten depth slices of Fig. 9c into a 2D palaeovalley index map, with values ranging from zero (i.e., no palaeovalley within the ten depth layers) to ten (i.e., palaeovalley detected across all depth layers) (Fig. 10a).

400 The resulting palaeovalley index map in Fig. 10a is first compared to the Multiple Resolution Valley Bottom Flatness (MRVBF) index in Fig. 10b which was originally calculated by Gallant and Dowling (2003) based on a digital elevation model with a spatial resolution of 100 m. High MRVBF values indicate a high probability of deposition of alluvium sediments. It was used by Munday et al. (2013), together with field observations of regolith, to obtain a hydrofacies map (black line in Fig. 10a). A comparison of the contours of the SRCNN palaeovalley index 10 and 6 with the MRVBF index shows the emergence of similar  
405 patterns (Fig. 10a and 10b). While this confirms that the SRCNN palaeovalley index map is not inconsistent with the MRVBF index, the latter contains insufficient information for testing the palaeovalley map.

The degree to which the MRVBF index can be used to identify the main three hydrofacies (bedrock, alluvium and transition material) is discussed on the basis of Fig. 10d. High MRVBF values correspond to bores with both alluvial lithology and transition material lithology, while a large number of bedrock boreholes also show high MRVBF values. In other words, the  
410 alluvial (i.e. palaeovalley) and bedrock/transition material (non-palaeovalley) lithology classes could not be fully identified by the MRVBF index.


In contrast, the AEM survey and the automatic neural network based palaeovalley classification in this study has improved capability to identify position of palaeovalleys. The box-plot of Fig. 10c shows that the boreholes classified as ‘alluvium’ correspond to a higher median SRCNN palaeovalley index of 4, compared to the two other lithology classes of median  
415 palaeovalley index of 0 and 2, respectively. For 128 boreholes identified in the study area, (i) those drilled in bedrock (66 boreholes) had the smallest SRCNN palaeovalley index (median of 0), (ii) those drilled in alluvium (57 boreholes) had the largest SRCNN index (median of 4), and (iii) those drilled in transition zones (5 boreholes) had the next largest SRCNN index (median of 2). Despite the relatively small dataset of borehole logs (3 per 100 km<sup>2</sup>), there is a clear trend that bores in alluvial sediments correspond to the areas with the highest SRCNN index. It is reasonable to assume that these alluvial

420 represent palaeovalleys, although the lithological classification did not provide this level of detail. However, for 11 alluvial boreholes, only a low corresponding palaeovalley index of  $< 2$  was identified. This may be due to the limited lithological and sedimentary information captured by the downhole logs, which were mainly recorded in the 1970s with limited description of the subsurface environment. The same is true for the boreholes in bedrock and transition zones, which may have been misclassified due to insufficient data.

425 The palaeovalley network shown in Fig. 10a is based on an analysis of ten depth layers and hence gives greater confidence about the location of deep palaeovalleys than the analysis of a single-depth palaeovalley map (Fig. 8). A significant proportion of the image has a maximum index of 10, meaning that a palaeovalley has been detected throughout the full investigation depth. This is thus an area with a high certainty (i.e. all pixels with index 10 have 10 layers identified as palaeovalley) that at least a 100-m deep palaeovalley is present. For the subsequent indices, e.g. 8, 6, 4, etc., at least 8, 6, and 4 depth layers with a  
430 palaeovalley were identified, respectively.

Moreover, the burial depth of the palaeovalley (defined by the vertical distance between the uppermost parts of the palaeovalley to the land surface) is calculated based on the 3D binary palaeovalley. It is shown in Fig. 11a that a wide range of the palaeovalleys are buried up to a depth of 10 to 20 m, which cannot be observed directly from the land surface, but can be revealed by the methodology proposed in this work based on geophysical prospecting (here 3D AEM data).

435



**Figure 11: (a) Burial depth and thickness (>40 m) of the alluvium sediments in palaeovalleys inferred from the 3D binary valley image of Fig. 9c with a 10-m vertical resolution. The hollow zone in (a) represents no identified palaeovalleys within the depth of 100 m in the study area.**

We finally calculate the thickness of palaeovalley layers (potentially representing the thickness of an alluvium aquifer) from the ten depth layers with palaeovalley indices. As a result, the thickness of the palaeovalley calculated by the distance between bottom (lowest part) and top (uppermost part) of the palaeovalley (Fig. 11b) is identical to the palaeovalley index (Fig. 10b) multiplied by the layer thickness of 10 m. This indicates that except for those pixels that were shown to have a 10 to 20 m cover of non-palaeovalley sediments (see burial depth in Fig. 11a), all other pixels had uninterrupted palaeovalley layers starting from the land surface. In those palaeovalley zones without surface sediment cover, the SRCNN palaeovalley index 8, 6, and 4 of Fig. 10a are representative for uninterrupted palaeovalley sediments in the depth intervals 0-80 m, 0-60 m, and 0-40 m, respectively.

Note that in Fig. 10a at any pixel with given palaeovalley index  $n$  (from 0 to 10), the probability of finding  $n$  consecutive palaeovalley layers can be inferred; in our test case area this was 100% everywhere - except for the buried pixels with 10 to 20 m cover of non-palaeovalley sediments - as no interruption was detected in the sequence of palaeovalley layers identified. This demonstrates that despite expected vertical lithological heterogeneities within palaeovalleys (Knight et al., 2018), AEM images combined with our SRCNN methodology are able to identify and differentiate a broad series of sediments that make up a palaeovalley from the surrounding bedrock. The SRCNN palaeovalley index map thus provides an improved tool for groundwater prospectivity.

## 5 Conclusions

The super-resolution convolutional neural network (SRCNN) approach is one of many deep learning methods developed recently to sharpen image quality and to extract particular features from images. This study applied as one of the first the SRCNN approach to resolve a long-standing challenge in the earth sciences: how to generate high-resolution binary palaeovalley maps from low-resolution electrical conductivity data derived from airborne electromagnetic surveys. The training images were generated using present-day valley patterns derived from DEM data as analogues to the palaeovalley patterns at different depths, together with Archie's equation and bicubic interpolation to generate the corresponding electrical conductivity images. The large training image dataset featured the responses of airborne electromagnetics (AEM) data to the

465 palaeovalley system with noise. Following a supervised learning, SRCNN successfully removed noise from AEM-derived  
electrical conductivity (EC) data and classified EC values into two separate palaeovalley index groups: one close to zero (the  
non-palaeovalley areas) and another one near unity (the palaeovalley areas). The resultant bimodal histogram of palaeovalley  
index was then used to select threshold values to convert palaeovalley indices to a binary palaeovalley and non-palaeovalley  
image. SRCNN can accommodate the spatial correlation between EC and palaeovalley index by moving filters to recreate the  
connectivity of the palaeovalley network. Moreover, the high-resolution of palaeovalley patterns can be inferred from low-  
470 resolution EC images via SRCNN, as long as their relationship is addressed in the training image dataset.  
However, there are several limitations to the method that require more work. In applying the SRCNN methodology, only EC  
images were used here to identify the palaeovalley network. In those area where palaeovalley and non-palaeovalley zones  
contain fluid with similar salinity, leading to similar bulk EC values, more geophysical information, e.g. gravity and magnetics,  
can be used as inputs in SRCNN to distinguish the position of palaeovalley. To generate a large training image pool, SRCNN  
475 was based on 2D training images derived from DEM data. The trained SRCNN were employed at different depth slices  
independently, where they were stacked up into a quasi-3D palaeovalley image. However, the vertical relationship between  
EC and palaeovalley index could not be addressed. In the future, the 3D palaeovalley patterns in the training dataset could be  
generated by process-based methods (e.g. sedimentary processes modelling) or multiple geostatistical approaches, and 3D  
images could be used in SRCNN to address horizontal and vertical correlations between EC and palaeovalley index  
480 simultaneously. In addition, when applying the SCRNN methodology to a new study area, the training images need to be  
updated according to the factors influencing the relationship between target geobody and electrical conductivities (i.e. porosity,  
water content, and sediment components in Archie's Equation).

### **Appendix: Robustness testing of the SRCNN methodology**

The neural network settings used in this study were as follows:  $f_1=9, f_2=1$  and  $f_3=5, n_1=64, n_2=32$  and  $n_3=1$ , where  $f$  represents  
485 the filter size and  $n$  represents number of output images from layer 1, layer 2 and layer 3, respectively; size of input image is  
50; and overlapping size of EC values between palaeovalley and non-palaeovalley zone is 5 %. We now modify each of these

parameters individually while fixing the others to investigate the robustness of the SRCNN as quantified by the performance metrics OSNR, SSIM and connectivity.

### A1. Overlapping size

490 In this study, an overlap in EC values between palaeovalley and non-palaeovalley zones is induced to reflect impact by factors such as noise and smoothing in the AEM data interpretation and interpolation; the maximum overlapping size discussed is 5% of the range of EC values (1-80 mS/m).




Figure A1: (a) PSNR and (b) SSIM calculated by SRCNN based on testing datasets in 400 iterations.

495

As shown in Fig. A1, the EC overlap between non-palaeovalley and palaeovalley zones in the training dataset only alters the speed at which the metrics PSNR and SSIM stabilize, but it does not affect the final PSNR and SSIM values. When the overlap size in the training dataset is comparable to that in the testing dataset (i.e. 5%), the SRCNN can be trained to generate images with a similar accuracy. Furthermore, a cross-test in Fig. A2 illustrated that the trained SRCNN can identify the palaeovalley in the testing dataset with overlap size smaller than that of the training image. However, if a small overlap size was employed in the training dataset (e.g. overlap size of 1%), the trained SRCNN failed to identify the palaeovalley cells in the testing dataset that had a larger overlap size (e.g. 5%).

500

This indicates that the SRCNN can be trained to remove noise in EC and identify the palaeovalley cells based on training

505 datasets, despite a certain degree of overlap in EC values between palaeovalley and non-palaeovalley. As a general rule, for the SCRNN to be successful, the overlap size in the training dataset should be larger than that in the testing dataset.





Figure A2: Effect of different degrees of EC overlap between palaeovalley and non-palaeovalley cells on model performance using (a) PSNR and (b) SSIM metrics.

510



**Figure A3: (a) Connectivity function in the northwest to southeast direction when applying the trained SRCNN to generate synthetic palaeovalleys. Resultant palaeovalley patterns using trained SRCNN with various degrees of overlap (1% (b) and 5% (c)) in comparison with real palaeovalley (d).**

515

However, this does not mean that the larger degree of overlapping in a training dataset is always expected. As shown in Fig. A3, when compared to the synthetic palaeovalley, the connectivity of palaeovalleys resulting from SRCNN decays with the increase of the degree of overlapping in training dataset. This is because when a large degree of overlapping is contained in the training dataset, SRCNN considers more pixels with similar EC in both palaeovalley and non-palaeovalley zones as noise.

520

After training, SRCNN removes too much noise and the resultant palaeovalleys are disconnected (Fig. A3-c). In contrast, when the degree of overlapping in the training dataset is low, the resulting image can contain noise in both palaeovalleys and non-palaeovalleys zones (Fig. A3-b), but a better palaeovalley connectivity is obtained. This suggests that although SRCNN can be trained to identify palaeovalleys from EC images with a certain degree of overlapping, it is still desirable to constrain the degree of overlapping EC between palaeovalley and non-palaeovalley zones based on field data, e.g. the groundwater salinity, porosity and major minerals in rocks.

525

Moreover, the overlapping EC values here do not indicate that palaeovalley and non-palaeovalleys cells have the same EC; otherwise, the AEM data will not contain enough information to separate the palaeovalley and non-palaeovalley zones. Furthermore, we need additional geophysical data, e.g. seismic velocity or gravity, to further constrain the palaeovalley position.

530

The inherent flexibility in the SRCNN methodology allows adding more geophysical dataset, e.g. gravity and seismic velocity to the input image, to obtain an improved training of the relationship between the binary palaeovalley image and multiple geophysical datasets. Demonstrating the information content of such data sets is beyond the scope of this paper.

## **A2. Input image size**

The EC and binary palaeovalley images with a size of  $30 \times 30$  to  $100 \times 100$  pixels are used to train the weights in the SRCNN. Although a larger input image size results in a higher PSNR metric, it does not significantly affect the SSIM metric (Fig. A4).



535 Given the same number of iterations (6,000) and batch size (50), the loss function is calculated at more pixels per iteration based on the larger input image. Consequently, longer computation times are required to train the SRCNN. Considering 6,000 iterations takes merely 51 min to train the  $30 \times 30$  pixels images, but 766 min are required to train the  $100 \times 100$  pixels images (Fig. A5). Using large input images to train the SRCNN with fewer iterations has the same effects as using small input image with more iterations.

540





Figure A4: Performance criteria PSNR (a) and SSIM (b) calculated by the SRCNN for the testing dataset under varying input image sizes (30, 50 and 100).



545

**Figure A5: The connectivity function of the palaeovalley generated by SRCNN, with the weight and bias values learned from the training images with size of 30, 50 and 100, respectively.**


550

However, as is evident from Fig. A5, the connectivity of SRCNN-generated palaeovalleys decreases for input images of  $30 \times 30$  pixels. This is because the correlation scale of EC and palaeovalley index exceeds the input image size. In other words, the small-size training image limits the ability of SRCNN to address the spatial correlation of EC values and to recreate spatial connectivity. When the image size exceeds  $50 \times 50$  pixels, the connectivity of generated palaeovalleys corresponds well with the synthetic palaeovalley. Further increasing the image size does not significantly affect the resultant palaeovalley pattern.

### A3. SRCNN depth, width and filter size


555

A larger filter size and network depth means more weights to be updated in the network, which potentially enhances the ability of the SRCNN in reproducing the palaeovalley and non-palaeovalley feature at each pixel. However, there is no strict criterion to determine the number of weights that yield a successful SRCNN model. It is reasonable to select the number of weights (i.e., unknowns) close to the size of training datasets (i.e., 60,000 knowns). Less weights could limit the capability of the SRCNN, while too many weights could cause overfitting risks in the SRCNN.



**Figure A6: Model performance PSNR (a) and SSIM (b) calculated for the test dataset with varying SRCNN filter depths and filter sizes. Numbers are as follows: 5-1-5 in 5(64)-1(32)-5(1) represents the filter size in layer 1, 2 and 3, respectively, and (64)-(32)-(1) represents the number of output images of layer 1, 2 and 3.**

In the three-layer network with filter size of 5-1-5, and output images of 64-32-1, the number of weights is 6592. When the filter size in the first layer increases to 9 and the depth of the network increases to 5, the number of weights becomes 59,328. Both are less than the size of training dataset (60,000). While the increase in filter size and depths of SRCNN yield slightly higher PSNR and SSIM (Fig. A6 and A7), the drawback is that longer computation times are required (Fig. A7). With the total number of weights getting close to the size of the training dataset, the rate at which PSNR improves with increasing network depth slows down (Fig. A6). On the other hand, a too deep network may remove too much noise from the palaeovalley part, which makes the palaeovalleys disconnected and the connectivity of the calculated palaeovalley (green line in Fig. A7) diverts from the reference (black line in Fig. A7).



575

**Figure A7: Connectivity of palaeovalley generated by SRCNN with multiple depths and filter sizes. Computation cost is the time taken to train the SRCNN.**

580

The filter size determines the spatial correlation length of EC values accounted for. Since we increase the filter size in the second layer to 5, a peak in PSNR and SSIM values and connectivity function are obtained in the full-size synthetic test (Fig. A7), although the number of weights in the network structure of 9(64)-5(32)-5(1) is not the largest among the five networks discussed. This suggests that a larger filter size is desirable to better address the spatial correlation of the EC values for palaeovalley cells. However, it is also noted that to maintain the size of output image to be the same as that of the input image, part of the filter covers the zone outside the input image, where EC values of zero are used. This may cause errors in palaeovalley index calculation, which is referred to as edge effect and can increase with filter size.

585

The depth of the network can also increase the correlation scale that is accounted for; the degree of this influence is determined by the filter size in each layer. In contrast, the width of each layer is unrelated to the correlation scale of EC and palaeovalleys, it merely alters the degree of nonlinearity of the network by affecting the number of weights.

## 6 Acknowledgement

590 Funding support for this study was provided by the National Key R&D Program of China (2018YFC0604305-03), the National Natural Science Foundation of China (No. 41572215 and No. 41502222), and the International Postdoctoral Exchange Fellowship Program (2017) from China Postdoctoral Council in combination with CSIRO funding through the Land and Water Business Unit and the Future Science Platform *Deep Earth Imaging*.

## 7 Reference

- 595 Ahl, A.: Automatic 1D inversion of multifrequency airborne electromagnetic data with artificial neural networks: discussion and a case study, *Geophysical Prospecting*, 51, 89-98, 2003.
- Amit, S. N. K. B., Shiraishi, S., Inoshita, T., and Aoki, Y.: Analysis of satellite images for disaster detection, *Geoscience and Remote Sensing Symposium (IGARSS)*, 2016 IEEE International, 2016, 5189-5192,
- Archie, G. E.: The electrical resistivity log as an aid in determining some reservoir characteristics, *Transactions of the AIME*,  
600 146, 54-62, 1942.
- Auken, E., Christiansen, A. V., Westergaard, J. H., Kirkegaard, C., Foged, N., and Viezzoli, A.: An integrated processing scheme for high-resolution airborne electromagnetic surveys, the SkyTEM system, *Exploration Geophysics*, 40, 184-192, 2009.
- Auken, E., Christiansen, A. V., Kirkegaard, C., Fiandaca, G., Schamper, C., Behroozmand, A. A., Binley, A., Nielsen, E.,  
605 Effersø, F., and Christensen, N. B.: An overview of a highly versatile forward and stable inverse algorithm for airborne, ground-based and borehole electromagnetic and electric data, *Exploration Geophysics*, 46, 223-235, 2014.
- Bishop, C. M.: *Neural networks for pattern recognition*, Oxford University Press, 1996.
- Christensen, N. K., Minsley, B. J., and Christensen, S.: Generation of 3-D hydrostratigraphic zones from dense airborne electromagnetic data to assess groundwater model prediction error, *Water Resources Research*, 53, 1019-1038, 2017.
- 610 Dong, C., Loy, C. C., He, K., and Tang, X.: Image super-resolution using deep convolutional networks, *IEEE transactions on*

pattern analysis and machine intelligence, 38, 295-307, 2016.

Drexel, J. F., and Preiss, W. V.: The Geology of South Australia, Geological Survey of South Australia Bulletin, 1995.

English, P., Lewis, S., Bell, J., Wischusen, J., Woodgate, M., Bastrakov, E., Macphail, M., and Kilgour, P.: Water for  
Australia's arid zone—Identifying and assessing Australia's palaeovalley groundwater resources: Summary report,  
615 Waterlines Report Series, 2012.

Fitterman, D. V., Menges, C. M., Al Kamali, A. M., and Jama, F. E.: Electromagnetic mapping of buried paleochannels in  
eastern Abu Dhabi Emirate, UAE, *Geoexploration*, 27, 111-133, 1991.

Gallant, J. C., and Dowling, T. I.: A multiresolution index of valley bottom flatness for mapping depositional areas, *Water  
resources research*, 39, 2003.

620 Granek, J.: Application of machine learning algorithms to mineral prospectivity mapping, University of British Columbia,  
2016.

Gu, J., Wang, Z., Kuen, J., Ma, L., Shahroudy, A., Shuai, B., Liu, T., Wang, X., Wang, G., and Cai, J.: Recent advances in  
convolutional neural networks, *Pattern Recognition*, 2017.

Gunnink, J., Bosch, J., Siemon, B., Roth, B., and Auken, E.: Combining ground-based and airborne EM through Artificial  
625 Neural Networks for modelling glacial till under saline groundwater conditions, *Hydrology and Earth System Sciences*,  
16, 3061, 2012.

Hao, S., Wang, W., Ye, Y., Li, E., and Bruzzone, L.: A Deep Network Architecture for Super-Resolution-Aided Hyperspectral  
Image Classification With Classwise Loss, *IEEE Transactions on Geoscience and Remote Sensing*, 56, 4650-4663, 2018.

Holzschuh, J.: Low-cost geophysical investigations of a paleochannel aquifer in the Eastern Goldfields, Western Australia,  
630 *Geophysics*, 67, 690-700, 2002.

Hou, B., Frakes, L., Sandiford, M., Worrall, L., Keeling, J., and Alley, N.: Cenozoic Eucla Basin and associated palaeovalleys,  
southern Australia—climatic and tectonic influences on landscape evolution, sedimentation and heavy mineral  
accumulation, *Sedimentary Geology*, 203, 112-130, 2008.

Jackson, J. A.: Glossary of geology, *Glossary of Geology*, by JA Jackson. 2005 Approx. 900 p. 5th revised and enlarged ed.

- 635 ISBN 3-540-27951-2. Berlin: Springer, 2005., 5th, 2005.
- Jones, D. A., Wang, W., and Fawcett, R.: High-quality spatial climate data-sets for Australia, *Australian Meteorological and Oceanographic Journal*, 58, 233, 2009.
- Kingma, D. P., and Ba, J.: Adam: A method for stochastic optimization, arXiv preprint arXiv:1412.6980, 2014.
- Knight, R., Smith, R., Asch, T., Abraham, J., Cannia, J., Viezzoli, A., and Fogg, G.: Mapping aquifer systems with airborne  
640 electromagnetics in the Central Valley of California, *Groundwater*, 2018.
- Långkvist, M., Kiselev, A., Alirezaie, M., and Loutfi, A.: Classification and segmentation of satellite orthoimagery using convolutional neural networks, *Remote Sensing*, 8, 329, 2016.
- Lane, R.: Ground and airborne electromagnetic methods, *Geophysical and Remote Sensing Methods for Regolith Exploration*. CRCLEME Open File Report, 144, 2002.
- 645 Ley-Cooper, A., and Munday, T.: Groundwater Assessment and Aquifer Characterization in the Musgrave Province, South Australia: Interpretation of SPECTREM Airborne Electromagnetic Data, Goyder Institute for Water Research Technical Report Series, 2013.
- Luo, Y., Zhou, L., Wang, S., and Wang, Z.: Video Satellite Imagery Super Resolution via Convolutional Neural Networks, *IEEE Geoscience and Remote Sensing Letters*, 14, 2398-2402, 2017.
- 650 Magee, J. W.: Palaeovalley groundwater resources in arid and semi-arid Australia: A literature review, *Geoscience Australia*, 2009.
- Maidment, D. R., and Morehouse, S.: Arc Hydro: GIS for water resources, ESRI, Inc., 2002.
- Marcais, J., and de Dreuzy, J. R.: Prospective Interest of Deep Learning for Hydrological Inference, *Groundwater*, 55, 688-692, 10.1111/gwat.12557, 2017.
- 655 Marker, P. A., Foged, N., He, X., Christiansen, A., Refsgaard, J., Auken, E., and Bauer-Gottwein, P.: Performance evaluation of groundwater model hydrostratigraphy from airborne electromagnetic data and lithological borehole logs, *Hydrology and Earth System Sciences*, 19, 3875, 2015.
- Meller, C., Genter, A., and Kohl, T.: The application of a neural network to map clay zones in crystalline rock, *Geophysical*

Journal International, 196, 837-849, 2013.

- 660 Moysey, S., Caers, J., Knight, R., and Allen-King, R. M.: Stochastic estimation of facies using ground penetrating radar data, *Stochastic Environmental Research And Risk Assessment*, 17, 306-318, 10.1007/s00477-003-0152-6, 2003.
- Mulligan, A. E., Evans, R. L., and Lizarralde, D.: The role of paleochannels in groundwater/seawater exchange, *Journal of hydrology*, 335, 313-329, 2007.
- Munday, T., Abdat, T., Ley-Cooper, Y., and Gilfedder, M.: Facilitating Long-term Outback Water Solutions (G-FLOWS  
665 Stage-1: Hydrogeological Framework, Goyder Institute for Water Research Technical Report Series, 2013.
- Olhoeft, G. R.: Electrical properties of granite with implications for the lower crust, *Journal of Geophysical Research: Solid Earth*, 86, 931-936, 1981.
- Pardo-Igúzquiza, E., and Dowd, P. A.: CONNEC3D: a computer program for connectivity analysis of 3D random set models, *Computers & geosciences*, 29, 775-785, 2003.
- 670 Parkhomenko, E. I.: *Electrical properties of rocks*, Springer Science & Business Media, 2012.
- Perol, T., Gharbi, M., and Denolle, M.: Convolutional neural network for earthquake detection and location, *Science Advances*, 4, e1700578, 2018.
- Pollock, D., and Cirpka, O. A.: Fully coupled hydrogeophysical inversion of synthetic salt tracer experiments, *Water Resources Research*, 46, 2010.
- 675 Purvance, D. T., and Andricevic, R.: On the electrical-hydraulic conductivity correlation in aquifers, *Water Resources Research*, 36, 2905-2913, 2000.
- Renard, P., and Allard, D.: Connectivity metrics for subsurface flow and transport, *Advances in Water Resources*, 51, 168-196, 2013.
- Rhoades, J., Raats, P., and Prather, R.: Effects of liquid-phase electrical conductivity, water content, and surface conductivity  
680 on bulk soil electrical conductivity 1, *Soil Science Society of America Journal*, 40, 651-655, 1976.
- Robinson, D., Binley, A., Crook, N., Day-Lewis, F., Ferré, T., Grauch, V., Knight, R., Knoll, M., Lakshmi, V., and Miller, R.: Advancing process-based watershed hydrological research using near-surface geophysics: A vision for, and review of,



electrical and magnetic geophysical methods, *Hydrological Processes*, 22, 3604-3635, 2008.

685 Samadder, R. K., Kumar, S., and Gupta, R. P.: Paleochannels and their potential for artificial groundwater recharge in the western Ganga plains, *Journal of Hydrology*, 400, 154-164, 2011.

Simpson, G. G.: Uniformitarianism. An inquiry into principle, theory, and method in geohistory and biohistory, in: *Essays in evolution and genetics in honor of Theodosius Dobzhansky*, Springer, 43-96, 1970.

690 Soerensen, C. C., Munday, T. J., Ibrahimi, T., Cahill, K., and Gilfedder, M.: Musgrave Province, South Australia: processing and inversion of airborne electromagnetic (AEM) data: Preliminary results, in, 1839-2725, Goyder Institute for Water Research Technical Report Series, 57, 2016.

Spies, B. R.: Depth of investigation in electromagnetic sounding methods, *Geophysics*, 54, 872-888, 1989.

Taylor, A., Pichler, M., Olifent, V., Thompson, J., Bestland, E., Davies, P., Lamontagne, S., Suckow, A., Robinson, N., and Love, A.: Groundwater Flow Systems of North-eastern Eyre Peninsula (G-FLOWS Stage-2): Hydrogeology, geophysics and environmental tracers, Goyder Institute for Water Research Technical Report Series, 2015.

695 Tu, J. V.: Advantages and disadvantages of using artificial neural networks versus logistic regression for predicting medical outcomes, *Journal of clinical epidemiology*, 49, 1225-1231, 1996.

Tuna, C., Unal, G., and Sertel, E.: Single-frame super resolution of remote-sensing images by convolutional neural networks, *International Journal of Remote Sensing*, 39, 2463-2479, 2018.

700 Varma, S.: Hydrogeological review of the Musgrave Province, South Australia, Goyder Institute for Water Research Technical Report Series, 2012.

Viezzoli, A., Christiansen, A. V., Auken, E., and Sørensen, K.: Quasi-3D modeling of airborne TEM data by spatially constrained inversion, *Geophysics*, 73, F105-F113, 2008.

Vilhelmsen, T. N., Behroozmand, A. A., Christensen, S., and Nielsen, T. H.: Joint inversion of aquifer test, MRS, and TEM data, *Water Resources Research*, 50, 3956-3975, 2014.

705 Wang, Z., and Bovik, A. C.: A universal image quality index, *IEEE signal processing letters*, 9, 81-84, 2002.

Wang, Z., Bovik, A. C., Sheikh, H. R., and Simoncelli, E. P.: Image quality assessment: from error visibility to structural

similarity, IEEE transactions on image processing, 13, 600-612, 2004.

Worthington, P. F.: The uses and abuses of the Archie equations, 1: The formation factor-porosity relationship, Journal of Applied Geophysics, 30, 215-228, 1993.

710 Yang, D., Oldenburg, D. W., and Haber, E.: 3-D inversion of airborne electromagnetic data parallelized and accelerated by local mesh and adaptive soundings, Geophysical Journal International, 196, 1492-1507, 2013.