

Hess-2019-242-RC1

Reply to Hessel Winsemius' comments

Emery et al. propose a new data assimilation scheme (AEnKF), to be used for assimilation of wide-swath altimeter information from the upcoming Surface Water and Ocean Topography (SWOT) mission. Given the large-scale nature and long-time scale revisit times, using an asynchronous scheme, that to the best degree possible utilizes time and space varying availability of information seems like a logical and useful choice, compared to more synchronous approaches such as classical EnKF. I consider this to be a useful contribution to HESS and very much in scope, and useful as a preparation for using SWOT in large-scale hydraulic simulations and forecasts.

*The authors would like to thank referee Hessel Winsemius for his detailed and very helpful comments, which helped us to better articulate our paper. Please find below our replies and associated modifications to the manuscript in order to address these concerns. To easily address all comments, we numbered each comment: they are referenced by a "M" for the major comments or by a "m" for the minor comments.*

*We kept the reviewer's comments and question in bold while our replies are in italic. When the associated modifications in the manuscript remain small, we inserted the modified paragraph in our reply in plain text: the black text corresponds to the original unmodified text, the crossed text corresponds to deleted text and the blue text corresponds to new text.*

*The abstract, the introduction (the latest paragraphs) and the discussions sections have been heavily modified to take into account several remarks from the three reviewers. Therefore, we refer the reviewers directly to the corrected annotated manuscript. In the annotated manuscript, we used a color code to differentiate which reviewer made the comment and suggested a modification: comments from reviewer 1 (Hessel Winsemius), 2 (Claire Michailovsky) and 3 (Paul Bates) are in purple, orange and green respectively. Each modification is also referenced by a code in bracket indicating the reviewer (R#\*\*\*) and the type and index of the comment (M/m#\*\*\*) such as: "[R#3-M#1]".*

**I do have a number of comments that lead to my verdict that this paper requires major revisions.**

## **MAJOR COMMENTS**

**M#1.** My largest comments are a) the choice to update Manning's  $n$  (rather than a state); and related to this, b) the choice to only evaluate the assimilation performance on the basis of water levels (or depth). In most applications, the user will require a good estimate of the river flow (besides water levels), because river flow (a volume in time) controls availability of water for some process that is to be predicted by the model used, not just the water level. A hydrology-hydraulic model-cascade could for instance be used to provide inputs to water allocation predictions for the forthcoming weeks/months (requiring an amount of flow over a given time span), or an upstream boundary condition for a flood simulation of a downstream river stretch. For all such applications, an accurate volume per unit of time is required, not just a stage (except for a flood simulation in a

steep area, where floodplain storage is negligible to event accumulated flow, but these are generally small streams, definitely not comparable with the size of the Amazon and its tributaries). I consider this a large (and unnecessary) weakness in the approach, with the additional risk that the Manning roughness will change to physically highly illogical values (which it in fact does in this study!), because e.g. either the water balance of the underlying hydrological simulations of ISBA are biased, or the channel dimensions are poorly defined. To become useful for typical applications, data assimilation should be as much as possible aimed at correcting the amount of water in channel sections, so that predictions after state updating can be made useful and reliable. This is now not proven. I find it a pity that the authors decided to apply this AEnKF on parameters with (As far as I can find in the text) the sole reason being that other authors already used it for state estimation experiments. This makes the study purely theoretical, as I don't really see how the experiments would ever be applied in a real-world case. The authors should at least show river flow as an additional benchmark variable and show how the 3 experiments affect the accuracy of river flow and discuss this result. My logical feeling is that discharge will be quite heavily impacted especially in experiment 3 where a bias is introduced.

*We agree with the reviewer on the fact that river flow/discharge is one of the main variable of interest for river modeling applications and, following that, state estimation are well adapted for hydrology data assimilation applications. However, we would like to emphasize that we already dedicated another study focused on the correction of the state and the discharge in the context of remotely-sensed hydrology products, see Emery et al. (2018). Moreover, as SWOT is a scientific mission with a nominal lifespan of 3 years (we indicate now this point in the abstract), an important application of SWOT data would be to calibrate hydrological models in order to get better simulations over past and future periods. Therefore, the present parameter estimation study works as a complementary study focusing on correcting crucial parameters for hydrological applications that are still not well known. To set this framework clearer, we added those remarks at the end of the introduction. Still, it is possible to apply the current framework (e.g. twin experiments with SWOT-like observations) to correct either water depths or river discharges (state estimation). We added this point to the perspectives and future works in the manuscript's conclusions.*

*The choice of the Manning coefficient as control variable is directly linked to the results of a sensitivity analysis (SA) of the ISBA-CTRIP river outputs to its routing parameters (Emery et al 2016). This SA showed that, among all CTRIP parameters and their tested ranges, simulated water depths are essentially sensitive to the Manning coefficients. Then, as SWOT water elevations product is closest to ISBA-CTRIP's water depths, it was chosen to build a framework based on the assimilation of SWOT-like water depths to correct the model most sensitive input parameters.*

*Besides, the reviewer also made very relevant points regarding the consequence of forcing/LSM bias on the roughness coefficient value. Indeed, in a real-case framework, if these types of bias are not considered, the assimilation scheme will try to change Manning coefficient values to compensate water elevations variations due to these biases. We realized that this point was not discussed in the initial manuscript, so we dedicated a new paragraph in the Discussions section where we acknowledge this limitation of our current approach and suggest solutions to handle it in future developments.*

Moreover, we would like to clarify that it is not the direct value of the Manning coefficient that is corrected but a multiplying factor applied to the Manning distribution (see our reply to your M#2). Therefore, the high values for the control variable displayed in Figures 7 and 9 correspond to these multiplying factors which remain within a range of 0.5 and 1.5, while the corresponding Manning coefficients are within physical value between 0.02 and 0.07.

Finally, as suggested by the reviewer, the model and data assimilation performances on water depth and discharge were easily calculated. Given the idealized framework of the OSSE, the statistics were exceptionally good for both water depth and discharge and did not bring essential new information to the manuscript. These tables were then added to the manuscript in a dedicated appendix.

- *Modified introduction >> see annotated manuscript P.3 L.29-33 + P.3 L.35 – P.4 L.4 + P.4 L.11-14*
- *Modified manuscript in the study's perspectives, P.21 L.32, L.35 – P.22 L.1-3:*

These experiments offer several perspectives. They mainly consist in going towards more realistic data assimilation experiments that take into account more sources of uncertainties between the model and the observations, (such as correlated observation errors or [uncertainties in the forcing and LSM surface and sub-surface runoff](#)). To test the performances' limitations regarding the DEM/bathymetry bias issue, one can use simulated water surface elevations referenced to a geoid instead of water depths from the model or even assimilate water depths from another model where the bathymetry is different. [As most applications generally require a good estimate of the river flow and river water volume, another lead of investigation could maintain the SWOT-based OSSE framework but to correct the simulated water storage and/or discharge, either as a single state estimation framework or as a dual state-parameter estimation framework \(similarly to dual discharge-bathymetry inference methods developed by Oubanas et al., 2018 and Brisset et al. 2018 for some hydraulic models\)](#). Moreover, along with observations of water surface elevations, SWOT will also provide two-dimensional maps of river widths and surface slopes. One can also study the possibility of assimilating such product to ~~correct the corresponding parameters in ISBA-CTRIIP such as the model river width or maybe~~ constrain other parameters such as the bankful depth that controls the model flooding scheme.

- *Modified discussions >> see annotated manuscript P.19 L.24-29*
- *Added appendix D*

**M#2. Second point: I don't fully understand the zonal approach to updating Manning's n. To me it would make more sense to use an upstream-downstream relation in manning coefficients (e.g. update manning coefficient at location, as well as upstream and downstream) which could easily introduce a logical covariance between n values (rather than assuming everything to be independent). In fact, the full zonal approach with areas that may have very little relationship to each other suggests, that there is a 100% covariance across the zone. Why was this selected in this way (or would things become overcomplicated if done in a different way)? Consider discussing this in the last sections of the paper.**

*The zonal approach is applied to the control variables only. As previously mentioned in the reply to M#1, the control variable are not directly the Manning coefficients but rather a set of multiplying factors applied to the Manning coefficient distribution. It is those factors that are set constant over the 9 hydro-geomorphologic areas while the Manning coefficient distribution remains spatially-distributed at the grid-cell scale and built on an upstream-downstream relationship (specified in Eq. 1 of the manuscript). The interest of using such zonal approach to correct the Manning coefficient distribution was indeed to maintain this upstream-downstream relationship between the grid-cell (which was not the case when each Manning coefficient was individually updated in Pedinotti et al.*

2014). We realized that this aspect might not be clear enough in the manuscript as reviewer #2 also raised similar questions. Therefore, we modified the manuscript adequately in Section 3.2.2 to better present the definition of the control variable and recall it in Sections 4 and 6.

- *Modified manuscript in section 3.2.2: P.8 L.18-24*

Following the conclusions from the ISBA-CTrip sensitivity analysis to its routing parameters in Emery et al. (2016), we determined that assimilating water-depth-like observations would be efficient to correct the distribution of the river Manning coefficients. These coefficients are spatially-distributed at the grid-cell scale. However, from Pedinotti et al. (2014), equifinality issues were raised by correcting the distribution at this scale while also affecting its upstream-to-downstream spatial distribution. Thus, we chose to correct it by applying multiplying factors defined at a coarser scale, namely at the scale of the 9 hydro-geomorphological areas defined in Section 2.3 and illustrated in Figure 1b. Within a same area, the Manning coefficients of all grid-cells are identically updated by being multiplied by the same correcting factor. Thus, data assimilation will focus on directly adjusting these multiplying factors. Therefore, the control vector is composed of the  $n_x$  multiplying factors  $N_{mult,i}$ ,  $i = 1 : : n_x$ , applied to correct the distribution of the river Manning coefficient:

$$x_k = [N_{mult,1}, \dots, N_{mult,n_x}]^T, \quad (8)$$

giving  $n_x = 9$ .

- *Modified manuscript in section 4: P.12 L.17, L.19*

In the incoming experiments, the true control variables  $x_t$  are:

... (17)

and their background a priori values at for the first assimilation cycle  $x_b$  are:

... (18)

- *Modified manuscript in section 6: P.16 L.12-13*

We present now the results from the data assimilation experiments presented in Table 2 and in Section 4.4.2. Recall that these experiments aim at correcting a set of 9 multiplying factors applied to the manning coefficient distribution and constant over 9 hydro-geomorphological zones the spatially varying Manning coefficient in the nine zones covering the Amazon basin.

**M#3. The English writing and sentence constructions are not everywhere up to standards. Please make sure the manuscript is reviewed by a (near-)native English person.**

*Reviewer #2 had a similar remark. Therefore, while preparing the replies to the reviewers, we submitted the manuscript to an independent English-speaking proofreader to improve the English.*

## **MINOR COMMENTS**

**m#1. Introduction: there are many “however”s in the text. Some or many of these can be removed.**

*Thank you for noticing this. We read through the introduction and modified it to use different linking words.*

**m#2. P. 3, l. 32 “at a coarser scale”, please just describe the scale.**



Here, the “coarser resolution” relates to the zonal distribution, in opposition to the grid-cell finer resolution. As the zonal distribution is not introduced yet in the manuscript, it is true that it might be confusing. Therefore, we modified the sentence in the introduction to be more explicit.

- Modified manuscript in Introduction P.4 L.22-23

For the current study, it was decided to update the Manning coefficient distribution, not a the grid-cell resolution but at a coarser zonal resolution, by applying multiplying correcting factors constant over each zones, at a coarser regional scale identical to the one used in Emery et al. (2016).

### **m#3. p. 4. l. 13: “gravitational drainage”, do you mean groundwater outflow?**

Not exactly. When we use the term “gravitational drainage”, we are describing the LSM where the water flows toward the deep soil and feeds CTRIP’s groundwater reservoir (denoted G). The term “groundwater outflow” is used for CTRIP and represents the flow from the groundwater reservoir G into CTRIP’s river/surface reservoir S. It is now specified in the manuscript.

- Modified manuscript in section 2.1: P.5 L.8

In particular, ISBA gives a diagnostic of the surface runoff (QISBA,sur) and the gravitational drainage (QISBA,sub, , i.e. water percolating to the deep layers of the soil ) later used as forcing inputs for the RRM denoted CTRIP.

### **m#4. p. 4. l. 20. Replace “empties” by “spills”**

Thank you this suggestion. The modification was made into the manuscript.

### **m#5. p. 4, l. 29, Replace “fixes” by “results in”**

Thank you this suggestion. The modification was made into the manuscript.

### **m#6. p. 5 l. 2 (p.5): “. . .values between 0. 75 and 1 for smaller and mountainous tributaries. . .” I guess you mean 0.075 and 0.1 s m-1/3. The values you mention are ridiculously high!**

Yes, you are absolutely right. This is a typing error, thank you for noticing it. The manuscript has been corrected.

### **m#7. p. 5. Eq. 1. Why is SOmax not simply 1 as it is only a way to scale values?**

Actually, there is another typing error here. SO is the stream order taking values ranging from 1 at source cells (grid cells without any upstream grid cells, according to the river network) to a maximal SO associated to the outlet grid cell (depending on the depth of the river network). It is Nmin and Nmax that takes values of 0.04 and 0.06, following the configuration from Decharme et al (2012). These definitions were corrected in the manuscript.

- Modified manuscript in section 2.2: P.6 L.3-6

with  $SO$  being the stream size relative measure at the current cell;  $SO_{max} = 0.06$  is the same measure at the river mouth (which value depends on the depth of the river network) and  $SO_{min} = 0.041$  the measure at source cells (namely cells without any upstream cells according to the river network) (Decharme et al., 2012). The Manning coefficient is then set to be constant in time while its spatial values decrease as the cells approaches the river outlet, taking values between  $N_{min}=0.04$  and  $N_{max}=0.06$  (Decharme et al 2012). (following the river network).

**m#8. p. 5, l. 10 replace “as the cells approaches” for “towards”**

*Thank you this suggestion. The modification was made into the manuscript.*

**m#9. p. 5, l 11.  $V(t)$  is not the surface flow, but the average cross-sectional flow velocity.**

*You are right. By “surface”, we meant the flow velocity in CTRIP’s surface reservoir  $S$ . The confusion is now corrected in the manuscript.*

- *Modified manuscript in section 2.2: P.6 L.7-8*

All these parameters are eventually essential to estimate the spatially- and time-varying surface average cross-sectional flow velocity in the surface reservoir  $v(t)$  following the Manning formula.

**m#10. p. 5, eq. 3.  $S$  is not defined**

*Actually, it was introduced p4, l.17, but it is true that it is hidden within the text and does not clearly appear as a variable. Therefore, we added it in p.5*

- *Modified manuscript in section 2.2: P.6 L.10*

where  $h_S$  is the river water depth estimated from the river storage  $S$  by

**m#11. p. 6, l. 3. “forcings are considered perfect”. This is my point above. They never are and the assimilation should work to correct these forcings. In the case of ISBA this concerns errors in the water balance, and in CTRIP errors in the transport of mass and momentum through the channel network.**

*We agree with the reviewer that the forcings are never perfect. By “perfect” here, we meant that the forcing uncertainties are not included in the generation of the ensemble, but they should definitely be included in future studies. First, in the manuscript, we withdraw the use of the term “perfect” when presenting the forcing (section 2.4) but instead write it is considered “as such” in the data assimilation framework (section 3.4.3). Then, following a similar remark from reviewer #2, we dedicate a paragraph to the “perfect” forcing assumption in the discussions. Finally, the objective to directly correct these forcings with assimilation is here out of the scope of the study but it is pointed out in the discussions.*

- *Modified manuscript in section 2.4.*

~~In the entire study, those forcing are considered perfect.~~

- *Modified manuscript in section 3.4.3: P.11 L.14-17*

To generate the background control ensemble, we solely stochastically perturb the variables within the control vector. Note that, by only perturbing the variables in the control vectors to generate the ensemble, we assume that all other features of the forward model, e.g. the atmospheric forcings, the LSM structure and therefore the surface and sub-surface runoff, are perfect. While this is the case for purely OSSE, such features are never perfect in real-case experiment. This assumption is further discussed in the Section 6.

- *Modified discussions P.19 L.21-24*

**m#12. p. 6, l. 26. “white noise”, is this a reasonable assumption? And if reality is different, how would it affect your results? Discuss this in Section 7.**

*When the study was developed, there was, at the time, no large-scale SWOT simulator allowing considering correlated SWOT-like errors so the white noise assumption was the most reasonable, in default of having a better error model. Nevertheless, reviewer #2 had a similar comment inviting us to discuss more those observation errors in the Discussions; therefore we added a dedicated paragraph in the Discussions.*

- *Modified discussions P.20 L.17-27*

**m#13. p. 11, eq 17 and 18. Are these the selected zonal Manning;s n values? These are unrealistically high. Why are these selected in such a strange domain?**

*No, those are the multiplying factors of the Manning coefficient distribution. This explains why their values are around 1.0. Following the modifications from M#2, this is now explicitly specified in the Section.*

**m#14. p. 11, 19. Describe briefly what your expected results are (i.e. why these experiments) on both water levels and flows!**

*Following the reviewer’s suggestion, we added a few sentences in the Section 4.2.*

- *Modified manuscript in section 4.2: P.13 L.8-10*

The first experiment, denoted as PE1, is configured from the aforementioned sensitivity test outcome. The parameters defining the experiment (spinup, starting date, ensemble size, control error) will be those giving the best results in the sensitivity tests in Table 2. Also, the reference level between the observed and simulated water depths is the same. In other words, there is no bias in the observation. This first idealized experiment serves as proof-of-concept *as the observations nature matches exactly the type of the simulated variables. Consequently, with this experiment, we expect to retrieve the true value of the control variables and hence the correct water depths and discharges.*

- *Modified manuscript in section 4.2: P.13 L.26-30*

First, in experiment PE2, there will still be no bias between the observed and simulated river bathymetry to see how the assimilation of anomalies performs. *Similarly to PE1, we expect this experiment to be able to retrieve the true control and state variables.* Finally, the last experiment PE3, which introduces a constant relative bias between CTRIP and SWOT, will be carried out. *For this experiment, we anticipate that the assimilation will still be able to retrieve the model state variables. The use of anomalies as observations should limit the impact of the inserted bias however, we do not exclude that it may be slightly echoed on the control variables.*

**m#15. Section 5.1. Describe what you I hydrological sense expect from the spinup time experiment. You can relate the expected required spinup to the time of concentration of the considered basin.**

*Thank you for this suggestion. We added the following sentences at the end of section 5.1.*

- *Additional remark in section 5.1: P.14 L.26-27*

*This period corresponds to the basin concentration time or, in other terms, the required time for the river network to totally empty.*

**m#16. p. 13. L. 9, you only show spatial average results. Why not spatial patterns? That may reveal the locations where things go right / wrong.**

*You are right. Initially, we chose to display the results averaged over the basin to keep the figure easy to read and limit the number of figures within the manuscript. However, these plots can definitively be generated. Therefore, we added the corresponding figure in a new appendix.*

- *Added Appendix C*

**m#17. p. 15, l. 16-28. I find this paragraph not very clear, it is not clear why the results behave so differently from zone 1,2,3. I was wondering if there is not simply too little water in the system to get correct results in this part of the domain? If you only change manning's n, you can never introduce new water or take water out of the system (see main comment).**

*You are right. We focused on the sensitivity analysis results to explain our results but it is true that, during the low flow period, there might be just too little water entering the system and the assimilation is unable to perform efficiently probably due to a forecast ensemble that is not spread enough. The section was slightly modified to include this comment and to simplify the interpretation.*

- *Modified manuscript in section 6.1: P.17 P.12-15*

Subsequently, zones 6, 7 and 8 correspond to right-bank tributaries, namely the Juruá and Purus rivers (zone 6), the Madeira river (zone 7) and the Tapajós and Xingu rivers (zone 8). These right-bank tributary zones are characterized by a strong seasonal cycle (see Figure 8, zones 6-8). Then, by comparing the corresponding plots in Figures 7 and 8, we notice that the period when the analysis control variable spreads from the truth corresponds to the low flow season in these zones. According to the global sensitivity analysis results, water depths in these zones are ~~not~~ less sensitive to the Manning coefficient in low flow conditions. *Additionally, there is very little water in the zones during this period. Consequently, the background control ensemble is not spread enough for the EnKF to be efficient.* Meanwhile, the EnKF still “sees” ~~a positive discrepancy between the model and the observations (i.e. that the observations are higher than the model predictions),~~ (as seen *with the positive innovation in these zones shown in the time evolution per zone of the innovation* in Figure A1). Therefore, in order to increase the simulated water depths, the EnKF corrects the Manning coefficient so that its value gets higher (a higher Manning coefficient means a slower flow velocity and then a higher simulated water depth). ~~However, the water being insensitive to the Manning coefficient during this period, the correction is not transferred to the simulated water depth and the Manning coefficient value keeps being increased during the low flow season.~~ Finally, once the low flow season ends, the analysis Manning coefficient converges back to the truth (see the last assimilation cycles).

**m#18. p. 17, l. 20. Around this part you should definitely discuss the state updating versus parameter updating, and the water storage errors that you can never resolve with parameter updating.**

*Following the reviewer's suggestion, we completed the Discussions paragraph on the forcing and LSM bias.*

- *Modified discussions P.19 L.24-29*

**m#19. p. 17, l. 31. I am very curious what kind of exceptional hydrological event you mean here**

*Maybe the term was not the most adapted here. By "exceptional", we meant intense flooding for example. We also changed the term by "extreme".*

- *Modified discussions P.18 L.29-30*

**m#20. Some of the figures have too small fonts, please make the figures readable throughout the text.**

*Following the reviewer's suggestion and a similar comment from reviewer #2, we re-worked the figures, namely Figures 7 to 10.*

Hess-2019-242-RC2

Reply to Claire Michailovsky's comments

### General Comments

This paper presents a data assimilation experiment using synthetic observations from the planned Surface Water Ocean Topography (SWOT) mission to update the Manning roughness parameters of a large scale routing model. The assimilation method used in the Asynchronous Ensemble Kalman Filter (AEnKF) with a 21-day time window (one orbit repeat cycle) which allows for measurements at different locations acquired at different times to be included in a single assimilation step. The application of the AEnKF is logical considering the specificities of the data and this is a useful contribution to the work preparing for SWOT.

*The authors would like to thank referee Claire Michailovsky for her detailed and very helpful comments. She raises very pertinent points that helped us better articulate our purpose. Please find below our replies and associated modifications to the manuscript with which we hope to address her concerns. To easily address all comments, we numbered each comment: they are referenced with a "M" for the major comments and with a "m" for the minor comments.*

*We kept the reviewer's comments and question in bold while our replies are in italic. When the associated manuscript's modifications remain small, we inserted the modified paragraph in our reply in plain text: the black text corresponds to the original unmodified text, the crossed text corresponds to deleted text and the blue text corresponds to new text.*

*The abstract, the introduction (the latest paragraphs) and the discussions sections have been heavily modified to take into account several remarks from the three reviewers. Therefore, we refer the reviewers directly to the corrected annotated manuscript. In the annotated manuscript, we used a color code to differentiate which reviewer made the comment and suggested a modification: comments from reviewer 1 (Hessel Winsemius), 2 (Claire Michailovsky) and 3 (Paul Bates) are in purple, orange and green respectively. Each modification is also referenced by a code in bracket indicating the reviewer (R#\*\*\*) and the type and index of the comment (M/m#\*\*\*) such as: "[R#3-M#1]".*

**I recommend publication with major revisions, including a thorough review for improving language.**

*Reviewer #1 had a similar remark. Therefore, while preparing the replies to the reviewers, we submitted the manuscript to an independent English-speaking proofreader to improve the English after taking into account your own corrections.*

### MAJOR COMMENTS

**My main comment relates to the specificity of the study as a synthetic experiment and with how some of the simplifications/assumptions are presented. I recommend further discussion on the impact of these on a study using real data, and on the estimates of uncertainties which are crucial to any assimilation experiment.**



*Thank you for this important remark. Reviewer #1 raised similar issues and also asked for more discussions regarding the impact of the OSSE's assumptions going towards more realistic experiments. Therefore, the "Discussions" section will be profoundly updated to take into account all the points raised by the reviewers. Please find below our more detailed reply.*

**These are specifically:**

### **M#1 - The assimilation of depth rather than elevation**

*We agree that SWOT will provide water surface elevations and not water depths. The assimilation of water depths in the first experiment serves as a reference or benchmark, as we expect it to perform well. Then, going towards more realistic experiments and directly use water elevations as observations, the first challenge would be to deal with the change of reference between the water depths simulated by the model and the water elevations measured by the satellites. To do so, we need to know the elevation of the bottom of the river bed referenced to the same geoid as the observations. Yet, this information is generally not well known. Therefore, we chose to use water surface elevation anomalies as observations because this type of variable does not depend on a reference or a bathymetry.*

*This essential point (use elevation anomalies to overcome the issue of unknown bathymetry) constitutes one of the main motivation for this study. Still, we realized it was not clearly stated neither in the abstract nor the introduction. Therefore, we modified these two sections.*

- *Modified abstract: P.1 L.7-10 + P.1 L.12 + P.1 L.17*
- *Modified introduction: P.4 L.6 + P.4 L.7-8 + P.4 L.10-11*

### **M#2 - The fact that the truth is generated by the same model:**

**- no model structure error**

**- there is a "real" Manning to converge to, which might not be the case with real data**

**- even in PE3, the bathymetry is not significantly changed, only the river bed elevation.**

*To our knowledge, the model structure error is still a challenging error to estimate and most of data assimilation studies assume no model structure error. However, when using ensemble-based model, a possibility to deal with this error is to enrich the background ensemble by considering more uncertainties from variables that are not necessarily included in the control vector (including errors in forcings or parameters from both the LSM and the RRM). The capacity of such ensemble to deal with the model structure error can then be tested in a framework where we still use synthetic observations but generated from a different model. We added those remarks to the Discussions.*

*We agree that bathymetry errors are more complex than just errors on river bankful depth Errors on widths and also representativeness errors due to the use of a simplified bathymetry are also important. With the presence of such errors, one could expect to generate updated Manning coefficient distribution with unrealistic values that translate the incoherence between the simulated and observed bathymetry. We added a paragraph in the Discussions dedicated to these errors.*

- *Modified Discussions: P.19 L.12-13, P.19 L.19-20, P.20 L.3-11*

### **M#3 - Assumption of perfect forcing**

Reviewer #1 had a very similar remark. We are aware that the “perfect” forcing assumption is a very strong simplifying assumption as the forcing control the amount of water entering the system. We do not plan on correcting the forcing with data assimilation here but rather calibrating the model given the current forcing. However, to smooth the effect of forcing errors on the assimilation, it is possible to include them in the generation of the background ensemble. A dedicated paragraph in the Discussions section has been added to the manuscript to discuss these aspects.

- Modified Discussions: P.19 L.21-29

## **MINOR COMMENTS**

**m#1 - P6, l.21:** this is brought up later when the anomalies are assimilated, but more focus should be placed here on the fact that depth is assimilated while SWOT will produce elevations. The conversion from level to depth is one of the big issues with using altimetry in hydrological studies. Does the SWOT simulator directly produce depths? You are assuming known river bed elevation, and this should be clearly specified. (I can see this is mentioned p11, l.26 but that is too late in the paper).

*You are right. This aspect is not clearly set when describing the observations variables. We modified the corresponding paragraph accordingly.*

- Modified manuscript in section 3.2.1: P.7 L.16-26

In the present study, the observed variables are water depths issued from a [simplified SWOT simulator](#). [Note that this simulator will produce water depths while the real SWOT satellite will provide water elevations](#). As in Biancamaria et al. (2011) and Pedinotti et al. (2014), [our](#) this SWOT simulator replicates SWOT spatio-temporal coverage. At a given date, the simulator selects the ISBA-CTrip cells contained (at least 50% of their area) in the SWOT ground tracks. Figure 2 shows some selected ISBA-CTrip cells under the real swaths over the Amazon basin. The true run is used as a basis to get the true water depths  $Y_{tk}$ . Then, to generate the observation vector  $y_{ok}$  from the extracted true water depths, each of them is randomly perturbed by adding a white noise characterized by a standard deviation  $\sigma_o$  such that:

Equation (6)

[Using water depths observation is a strong simplification of the real SWOT product](#). Therefore, in order to take into account that SWOT will provide water elevations and not directly water depths, this study will look at the assimilation of both water depths and water anomalies. The method to generate the anomalies will be further detailed in Section 4.2.

**m#2 - P7, l.9:** similar issue as previous comment, elevation and not depth is required at 10cm accuracy. The depth accuracy will be much lower. You assume no representativeness error due to scale, but how about level vs. depth?

*It is true that the 10cm vertical accuracy corresponds to water elevations measurements. We should expect higher errors on water depths, as we do not know the bathymetry. Still, it is complex to quantify such bathymetry errors and they only apply to water depths assimilation. As water depth assimilation is considered as a benchmark, we are still considering 10 cm vertical accuracy in our study for water depth. For water elevation anomalies, the error might even be lower, but as we are only considering white noise, we keep 10 cm for anomalies. We added a few remarks to acknowledge this aspect in the manuscript.*

- *Modified manuscript in section 3.2.1: P.8 L.5-10*

The observation error is the addition of the measurement error and the representativeness error. The first is associated to inherent instrumental errors when processes are observed and the second represents the error introduced when the observed and simulated variables are not exactly the same (in nature or scale). Following the SWOT uncertainty requirements (Esteban Fernandez, 2017), SWOT-like water surface elevation measurements have a vertical accuracy of 10 cm (when averaged over a water area of 1 km<sup>2</sup>). This uncertainty accounts for measurement errors due to the remotely-sensed acquisition such as instrumental thermal noise, speckle, troposphere and ionosphere effects. Moreover, we omit error correlations along the swath so that observation errors follow a white noise model. Accounting for spatially-correlated observation errors is an active research area in the field of data assimilation (Guillet et al., 2018) that is beyond the scope of demonstrating the feasibility of assimilating SWOT-type data. ~~Finally,~~ In the framework of OSSE, observed and simulated water depths have the same scale as the ISBA-CTRIIP model is used to generate both. Therefore, in the following, we assume there is no representativeness error [related to scale](#) in the system. [However, it is worth acknowledging that we should expect higher errors on water depths, compared to water elevations, as we do not know the bathymetry. Assimilation of water depths is performed as a benchmark, against which assimilation of water anomalies will be compared to.](#) Ultimately,  $\sigma^o$  is chosen equal to 10 cm for all observed variables (i.e. both water depths and water elevation anomalies).

**m#3 - P8, l11: Would this be necessary if you had a better representation of your measurement error (re: previous comments)? What is the magnitude of this additional error?**

*Not necessarily. In our study, we are using a stochastic version of the EnKF method that is, each member of the background ensemble is updated using the analysis equation 12 (in opposition to deterministic version like the Ensemble Transform Kalman Filter, where the observations randomization is not necessary as the analysis equation is only applied to the ensemble mean and the covariance matrix is estimated from a transformation of the prior covariance matrix). According to Burgers et al. (1998), the randomization of the observation vector should be used so that all the variables within the EnKF are random. Without this, the observation vector is deterministic, which conflicts with the EnKF analysis scheme where the ensemble covariance is used instead of the error covariance. Moreover, we know that the stochastic EnKF tends to under-estimate the analysis error. Therefore, we generate different realization of the observation vector so that the analysis ensemble retains enough variability from one assimilation cycle to another and avoid “ensemble collapse” (when all ensemble members are identical).*

*This randomization is specific to the algorithm itself, not the data. Still, the magnitude of the randomization is generally fixed by the observation error (hence, the randomization will be minimal if the observations are very accurate). Therefore, the observation randomization is done by adding to each observation a random perturbation following a white noise with a standard deviation  $\sigma^o=10\text{cm}$ . This is now specified in the manuscript.*

- *Modified manuscript in section 3.3: P.9 L.18-20*

To avoid ensemble collapse, [the observation vector in Eq. \(5\) is randomized by adding a supplementary white noise with the same observation error standard deviation  \$\sigma^o=10\text{ cm}\$  \(Burgers et al., 1998\) such that](#)

$$\forall j = 1 \dots n_y, \forall l = 1 \dots n_e, y_{k,j}^{o,[l]} = y_{k,j}^o + \epsilon_j^{o,[l]}, \epsilon_j^{o,[l]} \sim N(0, \sigma^o).$$

An observation ensemble is generated

$$Y_{e,k}^o = \left[ y_k^{o,[1]}, y_k^{o,[2]}, \dots, y_k^{o,[n_e]} \right].$$

**m#4 - At some points vague language is used (f.ex: “without really converging”), please be more precise.**

*Thank you for this advice. We carefully went through the manuscript and focused on correcting this language.*

**m#5 - P7, l.25: It is not clearly explained if  $x_k$  are the Mannings coefficients themselves or the multiplying factors as described above. This issue is repeated elsewhere in the paper and while I understand  $x_k$  does refer to the factors, please make sure this is clear throughout.**

*Reviewer #1 had a similar comment. Therefore, we modified the manuscript to clarify this aspect.*

- *Modified manuscript in section 3.2.2: P.8 L.21-24*

**m#6 - Figure 8: the black lines cannot really be seen, perhaps increase the width?**

*Thank you for this suggestion. We modified Figures 8 and 10 such that the different curves can be seen more easily.*

## **ADDITIONAL COMMENTS FROM SUPPLEMENT**

**m#7 – P2, l.9: it is not unavoidable that reductions in structure uncertainty are linked to increase in parameterization.**

*Following the reviewer’s recommendation, we re-wrote this sentence.*

**m#8 – P2, l.27-28: This sentence feels odd right after the mention of important vertical errors.**

*Yes, this is true. In the previous sentence, we implies that the signal with high vertical error was then ignored, diminishing more the spatio-temporal coverage of the observations. To be clearer, the part “or with important vertical error” was deleted.*

**m#9 – P5, l.3: in time I agree, but is it not usually spatially variable?**

*To our knowledge, there are models where a spatially-constant Manning coefficient is applied such as Biancamaria et al (2009), Beighley et al.,(2009) and other with a spatially-distributed Manning coefficients such as Decharme et al. (2012). However, it is true that the most recent model developments use a spatially-distributed coefficient. Therefore, this sentence was added to the manuscript.*

**m#10 – P10, l.10: this is confusing, is this study a standalone PE? This paragraph is unnecessarily complicated.**

*You are right; the term “standalone” is confusing. We withdrew it from the manuscript.*

**m#11 – P11, l.26: this should be mentioned way earlier in the text.**

*Please see reply to m#1.*

**m#12 – P12, l.8: isn't it river bed elevations you are talking about?**

*Yes, this is right. We changed the term in the manuscript.*

**m#13 – P15, l.1-5: I don't see a clear distinction between the behavior of these 2 groups in the figures**

*This is true. Besides, for the interpretation of the results in the next paragraph, zones 1 and 9 are interpreted along with zones 2, 3, 4 and 5. Therefore, this item is deleted and zones 1 and 9 are now cited in the first item.*

**m#14 – P16, l.1-2: It is overstating it that the bathymetries are different, the shape itself remains the same, only the bed elevation is shifted.**

*This is true. We re-adjusted this sentence in the corrected manuscript and worked on the paragraph.*

- *Modified manuscript in section 6.2: P.17 L.22-23:*

*In the PE2 experiment, there is no difference of bathymetry between the simulated and observed water anomalies while the river bankful depth is different in the PE3 experiment.*

**m#15 – p17, l.2-6: I find it odd that this is how you begin the discussion. If that is why assimilation is introduced, then the natural conclusion would be to update the river bed elevation.**

*It is true that this sentence was confusing. This part has been rewritten.*

- *Modified discussions P.18 L.24-26 + P.19 L.6-8*

**m#16 – P17, l.14-15: This will help only if width, bank slope etc are also correct or corrected**

*You are right. Following previous reviewers' comments, this point was covered elsewhere in the Discussions and the current sentence does not appear anymore in the manuscript.*

#### ***Additional references***

*Sylvain Biancamaria, Paul Bates, Aaron Boone, Nelly Mognard. Large-scale coupled hydrologic andhydraulic modelling of the Ob river in Siberia. Journal of Hydrology, Elsevier, 2009, 379 (1-2), pp.136-150. doi 10.1016/j.jhydrol.2009.09.054*

*Beighley, R. E., Eggert, K. G., Dunne, T., He, Y., Gummadi, V., and Verdin, K. L.: Simulating hydrologic and hydraulic processes throughout the Amazon River Basin, Hydrol. Process., 23, 1221–1235, doi:10.1002/hyp.7252, 2009*



Hess-2019-242-RC3

Reply to Paul Bates' comments

### General Comments

In addition to the comments of the previous referees, I would add that I think the paper also needs to better articulate what is the new contribution to knowledge made by the work. Previous papers have established that SWOT data is very likely going to allow us to better constrain parameters and states in a variety of environmental models, but I was missing what in addition to this that we learn from the present work. There are a number of specific technical differences here to the previous papers by Pedinotti et al (2014) and Emery et al (2016) which are described in the last paragraph of the introduction, but I am not sure these changes lead to very different conclusions.

*The authors would like to thank referee Paul Bates for his helpful comments. Please find below our replies with which we hope to address his concerns. To easily address all comments, we numbered each comment: they are referenced with a "M" for the major comments and with a "m" for the minor comments.*

*The abstract, the introduction (the latest paragraphs) and the discussions sections have been heavily modified to take into account several remarks from the three reviewers. Therefore, we refer the reviewers directly to the corrected annotated manuscript. In the annotated manuscript, we used a color code to differentiate which reviewer made the comment and suggested a modification: comments from reviewer 1 (Hessel Winsemius), 2 (Claire Michailovsky) and 3 (Paul Bates) are in purple, orange and green respectively. Each modification is also referenced by a code in bracket indicating the reviewer (R#\*\*\*) and the type and index of the comment (M/m#\*\*\*) such as: "[R#3-M#1]".*

**M#1** - Overall the abstract states that the assimilation scheme "is able to retrieve the true value of the Manning coefficients within one assimilation cycle most of the time", but isn't this ability already well established, at least in general terms? What do we learn about the likely potential of the SWOT data that we did not know before? Alternatively, what do we learn about the kinds of assimilation schemes that are going to work well with the SWOT data? In addressing this issue the comments of the other referees about the rather specific nature of the numerical experiment are pertinent: if the experiment does not reflect the assimilation problem that will be faced in practice when ISBA-CTRIP is confronted with SWOT data then the conclusions drawn may not be easy to generalize.

**I think a significant effort needs to be made to re-draft the abstract introduction to more clearly identify what is the new contribution to knowledge made by the paper. I'm sure there is one, so this is more a question of properly assembling the arguments in order to bring this out.**

*Beyond the objective to show the capability of assimilating SWOT-like data to correct the model's parameters, we wanted to introduce the assimilation of water surface anomalies as observed variables. Indeed, hydrological models usually produce water depths referenced to the bottom of the modeled bathymetry while SWOT will provide water surface elevations referenced to a reference surface (a geoid or an ellipsoid). Then, one of the biggest challenges in assimilating satellite-based water surface elevations remain in how they can be compared to simulations without introducing a large bias in the system. Therefore, we propose a method where we avoid the comparison between the model and observations reference by using the water surface anomalies that can be both*

*extracted from the model or the observations and compared. The use of anomalies offers then a method to handle SWOT data when we do not have any information on the bathymetry. However, it was required to show that assimilating water elevation anomalies would provide comparable results than assimilating water depths. This is the purpose of our study.*

*By gathering the comments from all reviewers, we realized this aspect was not enough emphasized in our manuscript. Therefore, we modified the abstract and the introduced to better assert this aspect. Moreover, the water surface anomalies are also introduced earlier in the manuscript: they were initially presented in Section 4.2 describing the assimilation strategy and they are now mentioned in Section 3.2.1 with the description of the observation variable. Finally, following a comment from reviewer #2, the manuscript has been cleaned to avoid the confusion between water depths and water surface elevations.*

*Our methodology was also modified in light of Pedinotti et al. (2014)'s results. We are now using a more complex assimilation scheme, the asynchronous EnKF over a 21-day assimilation window and we adopted a zonal approach to correct the Manning coefficient distribution.*

*Moreover, we are aware of limitations of our experiments. Going towards more realistic experiments will require some modifications to our OSSE. For example, more realistic model errors are needed, such more complex bathymetry errors, forcing and LSM parameters errors, model structure errors. More realistic SWOT observation errors must also be taken into account in future works. In our current study, we partially look at errors on the riverbed bankful depth. The other types of uncertainties are of a different nature and would require more processing than just the use of anomalies as observations. As this represents a first test with water surface anomalies, we limited our experiments to the proof-of-concept*

*and the test of the river bankful depth error. However, following the reviewers' recommendations, we intensively reworked the Discussions section to discuss the other types of uncertainties: how they would affect the assimilation performances and how they can be handled.*

- *Modified abstract, P.1 L.9-10, P.1 L.18-19*
- *Modified introduction, P.4 L.8-10*

## **List of all relevant changes to the manuscript (following the annotated manuscript outlines)**

- **P.1, L.4 [R#1-M#1]:** added the lifetime of SWOT
- **P.1, L.7-10 [R#2-M#1]+[R#3-M#1]:** additionnal remarks to better contextualize the study and its scientific contribution, i.e. the assimilation of water surface anomalies
- **P.1, L.12 + P.1, L. 17 [R#2-M#1]:** additional remarks to clearly state that the study will look at the assimilation of water surface anomalies
- **P.2, L.10, L.14 [R#1-m#1]:** rewording
- **P.2, L.16 [R#2-m#7]:** rewording
- **P.1, L.19 [R#3-M#1]:** additional remark to show the contribution of the study
- **P.3, L. 29 – P.4, L.4 [R#1-M#1]:** additional remark to better contextualize the study and justify the chosen parameter estimation framewok
- **P.4, L.7-8 [R#2-m#11]:** additionnal remark to clearly differentiate simulated water depths from observed water surface elevations and how it impacts the study
- **P.4, L.8-10 [R#3-M#1]:** additional remark to better define the study's objectives
- **P.4, L.10-11 [R#2-M#1]:** additional remark to better contextualize the study
- **P.4, L.12-14 [R#1-M#1]:** additional remark to better contextualize the study and justify the chosen parameter estimation framewok
- **P.4, L.22-23 [R#1-m#2]:** rewording to better explain the different variable scales
- **P.5, L.8 [R#1-m#3]:** additional definition
- **P.5, L.28 [R#1-m#6]:** corrected typing error
- **P.5, L.29-30 [R#2-m#8]:** modified statement on the use of the Manning coefficient in global scale hydrologic model
- **P.6, L.3-6 [R#1-m#7]+[R#1-m#8]:** rewording and corrected typing error to better explain Equation 1
- **P.6, L.7-8 [R#1-m#9]:** rewording to better define the velocity variable
- **P.6, L.10 [R#1-m#10]:** added missing definition of river storage S

- **P.7, L.16-17 [R#2-m#1]:** additional remark to clearly differentiate simulated water depths from observed water surface elevations
- **P.7, L.24-26 [R#2-m#1]:** additional remark to introduce water anomalies
- **P.8, L.5-10 [R#2-m#2]:** additional remark to discuss the observation error model
- **P.8, L.18-24 [R#1-M#2]:** additional paragraph to better justify the choice of control variables
- **P.9, L.18-20 [R#2-m#3]:** rewording to better present the observation randomization process
- **P.11, L.14-17 [R#1-m#11]:** additional remark on OSSE's assumptions
- **P.13, L.8-10 [R#1-m#14]:** additional remarks on what we expect the assimilation results to be
- **P.13, L.26-30 [R#1-m#14]:** additional remarks on what we expect the assimilation results to be
- **P.14, L.26-28 [R#1-m#15]:** additional remark on the physical meaning of the spinup time
- **P.16, L.12-13 [R#1-M#2]:** additional remark to better explain how the control variables are defined
- **P.16, L.19-20 [R#1-M#1]:** additional remark to introduce a new figure in appendix
- **P.17, L.12-15 [R#1-m#17]:** additional remark on the physical meaning of the assimilation results
- **P.17, L.22-23 [R#2-m#14]:** rewording
- **P.17, L.25-26 [R#1-M#1]:** additional remark to introduce new tables in appendix
- **P.18, L.4-9 [R#2-m#4]:** rewording
- **P.18, L.24-26 [R#2-m#15]:** rewording
- **P.18, L.29-30 [R#1-m#19]:** rewording
- **P.19, L.6-8 [R#2-m#15]:** rewording
- **P.19, L.12-13 [R#2-M#2]:** rewording

- **P.19, L.19-20 [R#2-M#2]:** additionnal discussions on the impact of bathymetry error on the corrected control variables
- **P.19, L.21-29 [R#1-M#1]+[R#1-m#18]:** additionnal paragraph on the impact of forcing and LSM errors
- **P.20, L.3-11 [R#2-M#2]:** additionnal paragraph on the impact of model structure error
- **P.20, L.17-27 [R#1-m#12]:** additionnal paragraph on more realistic observation error model
- **P.21, L.31-32 [R#1-M#1]:** added more study perspectives
- **P.21, L.35 – P.22, L.1-3 [R#1-M#1]:** additionnal perspectives on dual state-parameter estimation perspective
- **P.23, L.2-5:** new appendix in which the table explaining the spinup sensitivity test is moved to unload the main manuscript.
- **P.23, L.6-7 [R#1-m#16]:** new appendix with additionnal figure
- **P.23, L.8-16 [R#1-M#1]:** new appendix with additional tables showing the assimilation performances in terms of water depth and discharge RMSEn



# Assimilation of wide-swath altimetry observations to correct large-scale river routing model parameters

Charlotte Marie Emery<sup>1,2</sup>, Sylvain Biancamaria<sup>1</sup>, Aaron Boone<sup>3</sup>, Sophie Ricci<sup>4</sup>, Mélanie C. Rochoux<sup>4</sup>, Vanessa Pedinotti<sup>5</sup>, and Cédric H. David<sup>2</sup>

<sup>1</sup>LEGOS, 16 Avenue Edouard Belin, 31400 Toulouse, France.

<sup>2</sup>Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

<sup>3</sup>CNRM-GAME, Meteo-France, 42 Avenue Gaspard Coriolis, 31000 Toulouse, France

<sup>4</sup>CECI, Université de Toulouse, CERFACS, CNRS, 42 Avenue Gaspard Coriolis, 31057 Toulouse cedex 1, France

<sup>5</sup>Magellium, 1 Rue Ariane, 31520 Ramonville-Saint-Agne, France

**Correspondence:** Charlotte Marie Emery (charlotte.emery@jpl.nasa.com)

**Abstract.** Land surface models combined with river routing models are widely used to study the continental part of the water cycle. They give global estimates of water flows and storages but they are not without non-negligible uncertainties among which inexact input parameters play a significant part. The incoming Surface Water and Ocean Topography (SWOT) satellite mission, with a launch scheduled for 2021 and with a required lifetime of at least three years [R#1-M#1], will be dedicated to the measuring of water surface elevations, widths and surface slopes of rivers wider than 100 meters, at a global scale. SWOT will provide a significant amount of new observations for river hydrology and maybe combined, through data assimilation, with global-scale models in order to correct their input parameters and reduce their associated uncertainty. Comparing simulated water depths with measured water surface elevations remains however a challenge and can introduce within the system large bias. A promising alternative for assimilating water surface elevations consists of assimilating water surface elevation anomalies which do not depend on a reference surface [R#2-M#1][R#3-M#1]. The objective of this study is to present a data assimilation platform based on the asynchronous ensemble Kalman filter (AEnKF) that can assimilate synthetic SWOT observations of water depths and water elevation anomalies [R#2-M#1] to correct the input parameters of a large scale hydrologic model over a 21-day time window. The study is applied to the ISBA-CTRIP model over the Amazon basin and focuses on correcting the spatial distribution of the river Manning coefficients. The data assimilation algorithm, tested through a set of Observing System Simulation Experiments (OSSE), is able to retrieve the true value of the Manning coefficients within one assimilation cycle much of the time (basin-averaged Manning coefficients RMSEn is reduced from 33% to [1%-10%] after one assimilation cycle) and shows promising perspectives with assimilating water anomalies [R#2-M#1] (basin-averaged Manning coefficients RMSEn is reduced from 33% to [1%-2%] when assimilating water surface elevation anomalies over one year) which allows us to overcome the issue of unknown bathymetry [R#3-M#1].

## 1 Introduction

Global Hydrological Models (GHM) are extensively exploited to study the continental component of the global water cycle (Doll et al., 2015; Sood and Smakhtin, 2015). Such models have been extensively developed over the past two decades in order to quantify freshwater flows and storage changes over continental surfaces (Bierkens, 2015). They are based on the coupling of a Land Surface Model (LSM) with a River Routing Model (RRM). As an example, the ISBA-CTRIP (Decharme et al., 2019) hydrologic model results from the coupling of the ISBA LSM (Noilhan and Planton, 1989) and the TRIP RRM (Oki and Sud, 1998). LSMs simulate the energy and water balance at the soil-atmosphere-vegetation interface while RRM emulate the lateral transfer of freshwater toward the continent-ocean interface. The current study focuses on the river component of the terrestrial water cycle simulated by RRM.

GHMs give a global view of the state of the water flow and storage at model spatial and temporal resolutions. Nonetheless [R#1-m#1], they suffer from multiple sources of uncertainties which are related to the model structure, the external forcing and the input parameters (Liu and Gupta, 2007; Renard et al., 2010). Model structure uncertainties initially arose from a lack of knowledge in the hydrologic processes or from simplifying assumptions made to limit simulation computational cost. Still [R#1-m#1], with the increase of computational power, models are more and more complex (Liu and Gupta, 2007; Melsen et al., 2016): they run at finer spatial resolution, they include new physical processes and use an increasing number of fully distributed forcing and parameter datasets (Liu et al., 2012). This has led to an increase [R#2-m#7] in the number of model input parameters and potentially inflates the model uncertainty in those parameters. Input parameters express the spatial and/or temporal properties of the system. The spatial scale of parameters measurable on the field may differ from the model scale, while other conceptual parameters are not directly observable and measurable on the field (Moradkhani et al., 2005; Melsen et al., 2016) and are inferred using geomorphological empirical formula and/or indirect methods such as calibration (Gupta et al., 1998; Beven, 2012).

Another way in which to study the terrestrial water cycle is to use direct observations of the system. Most parts of the terrestrial water cycle are currently observed and measured from in situ or remote techniques (Sanoo et al., 2011; Vinukollu et al., 2011; Rodell et al., 2015). For the observations of rivers, in situ techniques measure river water elevations at gauge stations. In situ measurements are commonly very accurate and also frequent (i.e. sub-daily) but their main limitation is their spatially sparse sampling and their decreasing number over recent decades at a global scale (International Association of Hydrological Sciences Ad Hoc Group on Global Water Sets et al., 2001). Coincidentally, remotely-sensed data provided by satellite missions have increased quite significantly since the 90's and deliver effective river observations. The most common instrument operating to assess river water levels remains the nadir altimeter. Nadir altimetry gives localized water elevation measurements along the satellite ground track. Initially, altimeters were designed to monitor ocean topography but their application has broadened to the observation of lakes (Cretaux et al., 2009), floodplains (Birkett et al., 2002) and later on, rivers (Silva et al., 2010). Yet [R#1-m#1], their main limitation remains their limited spatial and temporal samplings: generally several days between two consecutive measurements at a limited number of locations. Besides, over continental surfaces, the signal is not always

retrievable. Current river observations therefore provide a more accurate view of the river system than models but they are quite limited by their sparse availability in space and time.

The incoming Surface Water and Ocean Topography (SWOT) mission, jointly developed by NASA, CNES, CSA and UKSA and scheduled for a launch in 2021, will be dedicated to the observation of continental free surface water with a better spatial and temporal coverage than the current nadir missions (such as EnviSat, the JASON series or also Sentinel-3A/B). SWOT main payload called KaRIn, for Ka-band Radar INterferometer (Fjórtoft et al., 2014), will observe surfaces under two swaths of 50 km each separated by a nadir-gap of 20 km and will have a near-global coverage. For hydrology, SWOT will observe rivers wider than 100 meters as well as lakes and wetlands larger than  $250 \times 250 \text{ m}^2$  within the latitudes  $78^\circ$  South and  $78^\circ$  North and with a revisit time of 21 days. SWOT will provide two-dimensional images of water surface elevations with a vertical accuracy of 10 cm when averaged over  $1 \text{ km}^2$  of water area. Along with water surface elevation measurements in rivers, SWOT will also provide observations of river width, surface slope and estimates of discharge based on SWOT observations. SWOT will provide a significant amount of new data for surface hydrology. It will give an ensemble of constraints that will allow a better depiction of surface water in hydrological models. This new data could be combined or integrated into global-scale hydrological models in order to correct them and improve their performances and forecasting capabilities.

Data assimilation techniques are a set of mathematical methods which combine a physical model and related external measurements taking their relative uncertainties into account. Data assimilation aims at improving the model ability to forecast and/or emulate the physical system's evolution. For this purpose, data assimilation methods are built to correct either the model's outputs (state estimation) or the model's input parameters (parameter estimation or PE), sometimes both simultaneously. Data assimilation for state estimation has been widely applied in meteorology and oceanography, and is more and more developed for large-scale terrestrial hydrology (Clark et al., 2008; Michailovsky et al., 2013; Paiva et al., 2013; Emery et al., 2018). Data assimilation for PE in hydrology has been initially developed as a dynamic alternative to model calibration (Montzka et al., 2011; Panzeri et al., 2013; Ruiz et al., 2013; Shi et al., 2015). In most models, parameters are assumed to be constant in time whereas, in reality, they may vary seasonally or under evolving climate and/or anthropogenic conditions. Sequential data assimilation can therefore help track model parameters variations in time (Kurtz et al., 2012; Deng et al., 2016; Pathiraja et al., 2016). PE is also used to retrieve conceptual parameters of hydrologic models such as friction coefficients (Pedinotti et al., 2014; Oubanas et al., 2018; Hafliger et al., 2019) or residence time of quick- and slow-flow reservoirs and partition of runoff excess (Vrugt et al., 2012; Pathiraja et al., 2016) which can not be directly measured.

Before launch, in the preparatory phase, Observing System Simulation Experiments (OSSE) can be performed in order to assess the benefits of assimilating SWOT data into a hydrological model and to evaluate the most adapted methodologies to assimilate this data into models. Several studies assimilating synthetic and/or simplified SWOT like data have been published so as to evaluate the correction of river model state namely river depth (Andreadis et al., 2007; Biancamaria et al., 2011), river storages (Munier et al., 2015) and river discharges (Andreadis and Schumann, 2014) at various scales [R#1-M#1]. But also, several studies focused on the possibility of using SWOT data to retrieve critical river parameters such as river bathymetry (Durand et al., 2008; Yoon et al., 2012; Mersel et al., 2013) and/or riverbed roughness/friction coefficient (Pedinotti et al., 2014; Oubanas et al., 2018; Hafliger et al., 2019). Indeed, SWOT is a scientific mission with a three year nominal lifetime. Therefore,

SWOT observations will help to better calibrate hydrological models and to improve their performances even over time periods beyond its lifetime. Moreover, other studies using real remote-sensing data have also been published and give insight into the challenges related to the assimilation of space-borne products such as Michailovsky et al. (2013); Michailovsky and Bauer-Gottwein (2014) and Emery et al. (2018) which assimilate nadir radar altimetry data [R#1-M#1].

5 In the present study, a data assimilation framework is used to correct input parameters of the large-scale ISBA-CTRIP model. More specifically, synthetic SWOT observations of water surface depths and anomalies [R#2-M#1] are assimilated in order to correct the spatially-distributed riverbed friction coefficients (or Manning coefficients). As SWOT will not directly measure water depths (it provides water elevation and the bathymetry is required to derive water depth) [R#2-m#11], the purpose of this study is to evaluate the possibility of assimilating water elevation anomalies to correct the model's parameters and to assess how the assimilation performances are impacted, compared to the direct assimilation of water depths [R#3-M#1]. Assimilating water elevation anomalies is done to overcome a potential lack of bathymetry data [R#2-M#1].

This study is presented as a complementary study to that of Emery et al. (2018) which is dedicated to the state estimation (river storage and discharge) of the same ISBA-CTRIP model, using real satellite-based discharge products [R#1-M#1]. The choice of the roughness coefficient as control variable was made following the results from the ISBA-CTRIP sensitivity analysis in Emery et al. (2016). In this preliminary study, the sensitivity of the simulated water depths and also anomalies to several river input parameters (such as riverbed width, depth, slope and also friction coefficient) was evaluated. The results showed that the highest sensitivity was in the Manning coefficient.

This study is, furthermore, also built on the conclusions from the work of Pedinotti et al. (2014). In our study, an Ensemble Kalman Filter (EnKF) is used (instead of the Extended Kalman Filter in Pedinotti et al. (2014)) to better account for the nonlinearities of the system and to better estimate the model errors. Also, Pedinotti et al. (2014) chose to update the Manning coefficients distribution at the grid cell scale and the question of equifinality arose (Beven and Freer, 2001) in their results. For the current study, it was decided to update the Manning coefficient distribution not at the grid-cell resolution, but at a coarser zonal resolution, by applying multiplying correcting factors uniformly over each zone [R#1-m#2], identical to the one used in Emery et al. (2016). Finally, Pedinotti et al. (2014) used an assimilation window of 2 days. This configuration resulted in updated Manning coefficient time series displaying "unrealistic jumps" with a frequency of about 20 days associated to the orbit repeat cycle (longer than the 2-day window). To avoid this phenomenon, the present study uses an assimilation window of 21 days corresponding to the current SWOT orbit repeat cycle.

Section 2 will first give a description of the ISBA-CTRIP model used for this study. Section 3 will present the particular data assimilation method developed for this study and finally, after presenting the assimilation strategy in Section 4, Sections 5 and 6 will give the data assimilation results.

## 2 Model

### 2.1 The ISBA-CTRIIP large-scale hydrological model

The ISBA model (Noilhan and Planton, 1989) is a LSM defined at global-scale on a  $0.5^\circ \times 0.5^\circ$  regular mesh grid that establishes the energy and water budget over continental surfaces. This study operates the ISBA-3L version based on a three-layer soil (Boone et al., 1999). The budget equations are solved separately on each grid cell. Still, larger-scale spatial patterns in the radiative and precipitation forcing, the soil composition and the vegetation cover ensure spatial correlations between those cells. (for more details, see Decharme et al., 2012, 2019). In particular, ISBA gives a diagnostic of the surface runoff ( $Q_{\text{ISBA,sur}}$ ) and the gravitational drainage ( $Q_{\text{ISBA,sub}}$ , i.e. water percolating to the deep layers of the soil [R#1-m#3]) later used as forcing inputs for the RRM denoted CTRIP.

The CTRIP model (Decharme et al., 2010, 2012, 2019), is defined on the same mesh grid as ISBA and follows a river network to laterally transfer water from one cell to another, down to the interface with the ocean (Oki and Sud, 1998). The study is based on the CTRIP version from Decharme et al. (2012) with three reservoirs, as illustrated in Figure 1a. The water mass [kg] stored in a groundwater reservoir  $G$  and a floodplain reservoir  $F$  interacts with the water mass in the surface reservoir  $S$  representing the river. Only the surface reservoir  $S$  is related to the river network and fills with the surface runoff  $Q_{\text{ISBA,sur}}$ , the outflow from upstream cells and the delayed drainage  $Q_{\text{ISBA,sub}}$  by means of the groundwater reservoir. Occasionally, when the amount of water in the river exceeds a given threshold (defined by the water level in the reservoir), the river spills [R#1-m#4] into the floodplains.

### 2.2 CTRIP parameters

Within a  $0.5^\circ \times 0.5^\circ$  cell, the surface reservoir is a unique river channel that may gather multiple real river branches. Its rectangular cross-section is described by its slope  $s$  [-], its width  $W$  [m], its bankful depth  $H_c$  [m], its length  $L$  [m] and finally a Manning or friction coefficient  $N$  [ $\text{s m}^{-1/3}$ ] that assesses the reach resistance at the bottom of the river.

Each cell's elevation is deduced from the STN-30p Digital Elevation Model ([http://daac.ornl.gov/ISLSCP\\_II/islscpii.shtml](http://daac.ornl.gov/ISLSCP_II/islscpii.shtml)). These elevations are then compared to determine the riverbed slope  $s$ . Global empirical geomorphologic relationships are used to define the river width  $W$  and bankful depth  $H_c$ . The arc length between grid cell centers, inflated by a meandering factor  $\mu$ , results in [R#1-m#5] the river reach length  $L$ . More details on these parameters can be found in Oki and Sud (1998) and Decharme et al. (2012).

The Manning coefficient  $N$  is generally more complicated to estimate. Following Maidment (1993), it should take values between 0.025 and 0.03 for natural streams and values between 0.075 and 0.1 [R#1-m#6] for smaller and mountainous tributaries and also floodplains. Global studies can apply either a constant (Beighley et al., 2009; Biancamaria et al., 2009) or a spatially-distributed (Decharme et al., 2012) Manning coefficient [R#2-m#8]. However, it is ordinarily accepted that this parameter should vary in space and even in time across the river catchment. Consequently, CTRIP uses a spatially-distributed Manning coefficient based on a simple linear relationship between the relative stream size in the current cell, denoted  $SO$ , and

the size at the river mouth and the source cells, so that:

$$N = N_{min} + (N_{max} - N_{min}) \frac{SO_{max} - SO}{SO_{max} - SO_{min}}, \quad (1)$$

- $SO$  is the stream size relative measure at the current cell;  $SO_{max}$  (whose value depends on the network depth) the same measure at the river mouth and  $SO_{min} = 1$  the measure at source cells (namely, cells without any upstream cells, according to the river network). The Manning coefficient is then set to be constant in time while its spatial values decrease towards the river outlet (following the river network), with values between  $N_{min} = 0.04$  and  $N_{max} = 0.06$  [R#1-m#7][R#1-m#8].

All these parameters are eventually essential to estimate the spatially- and time-varying average cross-sectional flow velocity in the surface reservoir  $v(t)$  following the Manning formula (Manning, 1891) [R#1-m#9]:

$$v(t) = \frac{s^{\frac{1}{2}}}{N} \left( \frac{Wh_S(t)}{W + 2h_S(t)} \right)^{\frac{2}{3}}, \quad (2)$$

- where  $h_S$  is the river water depth estimated from the river storage  $S$  [R#1-m#10] by

$$h_S = \frac{S}{\rho WL}, \quad (3)$$

and  $\rho$  the water density. The flow velocity is ultimately used to estimate the discharge leaving the CTRIP cell:

$$Q_{out}^S(t) = \frac{v(t)}{L} S(t). \quad (4)$$

- As the definition of most of these parameters is based on empirical relationships, we have to be aware that they inevitably have substantial uncertainties.

### 2.3 CTRIP implementation over the Amazon basin

In this study, we present an OSSE test case over the Amazon river basin whose hydrology is carefully described in Molinier et al. (1993); Wisser et al. (2010). This choice was motivated as the present work follows and complements studies over the same domain (Emery et al., 2016, 2018).

- For ISBA-CTRIP, the Amazon basin is composed of a total number of 2028 cells. Based on the basin geomorphology and hydrology (Meade et al., 1991), the basin has been split into 9 spatial regions. These zones, illustrated in Figure 1b, were initially introduced in (Emery et al., 2016) and will be re-exploited here within the application of data assimilation. For a detailed description of the zones, the reader can refer to (Emery et al., 2016).

### 2.4 ISBA-CTRIP forcing

- For the present study, ISBA-CTRIP needs external atmospheric forcing in order to run. Similarly to Emery et al. (2016), such data is provided by the Global Soil Wetness Project 3 (GSWP3, <http://hydro.iis.u-tokyo.ac.jp/GSWP3>) at a 3-hourly time resolution.



### 3 Method: Synthetic parameter estimation on ISBA-CTrip

#### 3.1 OSSE framework

In OSSE, we introduce beforehand a reference configuration for the model input parameters that we will consider thereafter as the *truth*. From those true parameters, we directly deduce the *true run* from a ISBA-CTrip model integration. The synthetic observations used for data assimilation are obtained from perturbing the true observables (variables that are used as observations) using an error model that is representative of the real observation errors. The control variables (the model variables to be corrected with data assimilation) first guess are obtained by directly perturbing the true control variables. Control error also has to be chosen to be representative of the real modeling errors. OSSEs are prerequisite tests to ensure that the implementation of the EnKF algorithm is correct and adapted to the hydrologic problem under consideration (temporal/spatial length-scales, sources of uncertainty, observation operator...).

#### 3.2 Data assimilation variables

##### 3.2.1 Observation variables and their errors

The observation vector, denoted  $\mathbf{y}_k^o$  at the assimilation cycle  $k$ , is composed of the  $n_y$  available observations at cycle  $k$ :

$$\mathbf{y}_k^o = \left[ y_{k,1}^o, y_{k,2}^o, \dots, y_{k,n_y}^o \right] \quad (5)$$

where  $y_{k,j}^o$ ,  $j = 1 \dots n_y$ , is the  $j$ -th observation among the  $n_y$  at cycle  $k$ .

In the present study, the observed variables are water depths issued from a simplified SWOT simulator. **Note that this simulator will produce water depths while the real SWOT satellite will provide water elevations [R#2-m#1]**. As in Biancamaria et al. (2011) and Pedinotti et al. (2014), this SWOT simulator replicates SWOT spatio-temporal coverage. At a given date, the simulator selects the ISBA-CTrip cells contained (at least 50% of their area) in the SWOT ground tracks. Figure 2 shows some selected ISBA-CTrip cells under the real swaths over the Amazon basin. The true run is used as a basis to get the true water depths  $\mathbf{y}_k^t$ . Then, in order to generate the observation vector  $\mathbf{y}_k^o$  from the extracted true water depths, each of them is randomly perturbed by adding a white noise characterized by a standard deviation  $\sigma^o$  so that:

$$\forall j = 1 \dots n_y, y_{k,j}^o = y_{k,j}^t + \epsilon_j^o, \epsilon_j^o \simeq \mathcal{N}(0, \sigma^o). \quad (6)$$

**Using water depth observations is a strong simplification of the real SWOT product. Therefore, in order to take into account that SWOT will provide water elevations and not directly water depths, this study will look at the assimilation of both water depths and water anomalies. The method for generating these anomalies will be further detailed in Section 4.2 [R#2-m#1].**

The observation error is the addition of the measurement error and the representativeness error. The first is associated with inherent instrumental errors when processes are observed and the second represents the error introduced when the observed and simulated variables are not exactly the same (in nature or scale). Following the SWOT uncertainty requirements (Esteban Fernandez, 2017), SWOT-like water surface elevation measurements have a vertical accuracy of 10 cm (when averaged over a

water area of 1 km<sup>2</sup>). This uncertainty accounts for measurement errors due to the remotely-sensed acquisition such as instrumental thermal noise, speckle, troposphere and ionosphere effects. Moreover, we omit error correlations along the swath so that observation errors follow a white noise model. Accounting for spatially-correlated observation errors is an active research area in the field of data assimilation (Guillet et al., 2018) which is beyond the scope of demonstrating the feasibility of assimilating

5 SWOT-type data. In the framework of OSSE, observed and simulated water depths have the same scale as the ISBA-CTrip model which is used to generate both. In the following, we assume therefore that there is no representativeness error related to the scale in the system. However, it is worth acknowledging that we should expect higher errors on water depths, compared to water elevations, as we do not know the bathymetry. Assimilation of water depths is performed as a benchmark, against which assimilation of water anomalies will be compared. Ultimately,  $\sigma^o$  is chosen as being equal to 10 cm for all observed variables

10 (i.e. both water depths and water elevation anomalies) [R#2-m#2].

### 3.2.2 Control, observation space and their errors

The control vector is denoted by  $\mathbf{x}_k \in \mathbb{R}^{n_x}$ . It includes the  $n_x$  uncertain variables to be estimated through the  $k$ -th data assimilation cycle. The choice of the control variables determines the observation operator  $\mathcal{H}_k$ :

$$\mathbf{y}_k = \mathcal{H}_k(\mathbf{x}_k), \quad (7)$$

15 where  $\mathbf{y}_k$  are the simulated observables, in other words,  $\mathcal{H}_k$  maps the control variables onto the observation space. They are then compared to the measured observations  $\mathbf{y}_k^o$  during the data assimilation experiment. This difference is referred to as the innovation vector.

Following the conclusions from Emery et al. (2016), we determined that assimilating water-depth-like observations would be efficient for the correction of the distribution of the river Manning coefficients. These coefficients are spatially-distributed

20 at the grid-cell scale. However, from Pedinotti et al. (2014), equifinality issues were raised through the correcting of the distribution at this scale. They also affected its upstream-to-downstream spatial distribution. We chose to correct it therefore by applying multiplying factors defined at a coarser scale, namely at the scale of the 9 hydro-geomorphological areas defined in Section 2.3 and illustrated in Figure 1b. Within the same area, the Manning coefficient values are all identically modified by being multiplied by the same factor [R#1-M#2][R#1-m#5].

25 The control vector is composed therefore of the  $n_x$  multiplying factors  $N_{\text{mult},i}$ ,  $i = 1 \dots n_x$ , applied to the correcting of the spatial distribution of the river Manning coefficient:

$$\mathbf{x}_k = [N_{\text{mult},1} \dots N_{\text{mult},n_x}]^T, \quad (8)$$

giving  $n_x = 9$ .

The observation operator  $\mathcal{H}_k$  maps the control variables (Manning coefficients dimensionless multiplying factors) into the

30 observables (river water depths in meters) as follows:

1. first, apply the multiplying factors ( $\mathbf{x}_k$ ) to the Manning coefficients distribution;

2. then, apply the ISBA-CTrip model  $\mathcal{M}_{[k-1,k]}$  over the assimilation window  $[k-1, k]$  to determine the model states that correspond to the perturbed Manning coefficients distribution;
3. after, turn the CTrip surface water storage into equivalent water depths following Eq. 3 (we denote by  $\mathcal{Z}_k$  the diagnostic operator turning the surface storage variable into the water depth variable);
- 5 4. finally, select the simulated water depths under the SWOT swath mask (we denote by  $\mathcal{S}_k$  this operator).

The observation operator is therefore the composition of three operators:

$$\mathbf{y}_k = (\mathcal{S}_k \circ \mathcal{Z}_k \circ \mathcal{M}_{[k-1,k]})(\mathbf{x}_k) = \mathcal{H}_k(\mathbf{x}_k). \quad (9)$$

Such a non-linear observation operator  $\mathcal{H}_k$  is difficult to formulate explicitly which is why we use an EnKF algorithm to estimate the Kalman gain in a statistical way.

### 10 3.3 The EnKF general formulation

In the EnKF framework, the model  $\mathcal{M}_{[k-1,k]}$  and observation  $\mathcal{H}_k$  operators are generally not linear. The main assumption for the EnKF is to use stochastic ensembles to represent first- and second-order moments (namely the means and the covariances) of the control variable errors (Evensen, 1994, 2003). Indeed, it is assumed that the distribution of the ensemble is similar to that of the error of the control vector and it is also assumed that the Probability Density Function (PDF) of the error is gaussian, thus well described by its first and second moments. The background control variables  $\mathbf{x}_k^b$  (the first guess) are therefore represented by an ensemble of  $n_e$  members:

$$\mathbf{X}_{e,k}^b = \begin{bmatrix} \mathbf{x}_k^{b,[1]} & \mathbf{x}_k^{b,[2]} & \dots & \mathbf{x}_k^{b,[n_e]} \end{bmatrix}, \quad (10)$$

To avoid the collapsing of the ensemble, the observation vector in Eq. 5 is randomized by adding a supplementary white noise with the same observation error standard deviation  $\sigma^o$  (Burgers et al., 1998) [R#2-m#3] so that

$$20 \quad \forall j = 1 \dots n_y, \forall l = 1 \dots n_e, y_{k,j}^{o,[l]} = y_{k,j}^o + \epsilon_j^{o,[l]}, \epsilon_j^o \simeq \mathcal{N}(0, \sigma^o). \quad (11)$$

An observation ensemble is generated:

$$\mathbf{Y}_{e,k}^o = \begin{bmatrix} \mathbf{y}_k^{o,[1]} & \mathbf{y}_k^{o,[2]} & \dots & \mathbf{y}_k^{o,[n_e]} \end{bmatrix}. \quad (12)$$

Note that alternatives exist to the observation randomization chosen here. However, for the present study, we choose to use a full stochastic filter.

25 Finally, the EnKF analysis step is applied to each member of the ensemble so that

$$\forall l = 1 \dots n_e, \mathbf{x}_k^{a,[l]} = \mathbf{x}_k^{b,[l]} + \mathbf{K}_{e,k} \left( \mathbf{y}_k^{o,[l]} - \mathcal{H}_k(\mathbf{x}_k^{b,[l]}) \right), \quad (13)$$

where  $\mathbf{K}_{e,k}$  is the the Kalman gain. It is built from the control and observation error covariance matrices  $\mathbf{P}$  and  $\mathbf{R}$  and the linearized observation operator  $\mathbf{H}$  so that (see Appendix A for more details):

$$\mathbf{K}_{e,k} = [\mathbf{P}\mathbf{H}^T]_{e,k} \left( [\mathbf{H}\mathbf{P}\mathbf{H}^T]_{e,k} + \mathbf{R}_k \right)^{-1}. \quad (14)$$

Figure 3 summarizes the general OSSE framework used for the present study. The figure reads from top to bottom and from left to right. An assimilation cycle  $[k-1, k]$  includes a forecast step where an ensemble of ISBA-CTRIP simulations is integrated, each member having a different spatially-distributed Manning coefficient; an analysis step where the ensemble of Manning coefficients is corrected using synthetic observations through the Kalman filter update in Eq. 13; and a cycling step where the ISBA-CTRIP model is re-run with these analysis estimates to obtain updated model states.

### 3.4 SWOT-based data assimilation special feature

#### 3.4.1 Choice of the assimilation window

We use a 21-day assimilation window corresponding to a SWOT orbit revisit period. During one assimilation window, every pixel under the observation mask is therefore observed at least once. However, it implies that new observations are available at times which differ from the update time. Such a case has already been addressed in several studies. The Ensemble Kalman Smoother (EnKS) for example, introduced by Evensen and Leeuwen (2000), is a direct extension of the EnKF. It consists of generating an update of the control variables taking into consideration the present and all past observations when a new observation is available. The EnKS is actually a sequential version of the Ensemble Smoother (Leeuwen and Evensen, 1996). The latter takes into consideration all past and future observations but turned out to be less effective than the EnKF and the EnKS (Evensen, 1997; Evensen and Leeuwen, 2000). Alternatively, Hunt et al. (2004) developed the 4D-EnKF (4D as in the 4D-VAR variational assimilation methods; Talagrand and Courtier, 1987), which also assimilates observations available at different time-steps. In the 4D-EnKF, all model observations are expressed as a linear combination of the model observations at analysis time and the problem is transformed into a classic EnKF problem. Similarly, Hunt et al. (2007) also presented an asynchronous version of the Local Ensemble Transform Kalman Filter (Bishop et al., 2001; Ott et al., 2004). In the framework of the present study, we apply an Asynchronous Ensemble Kalman Filter (AEnKF) as described by Sakov et al. (2010) and Rakovec et al. (2015). The principle is to increase the dimension of the state in order to consider observations at past and analysis times. This increases the dimension of the matrices which contain covariances between observations available at different times. To our knowledge, the AEnKF has not been used for parameter estimation, as Sakov et al. (2010) and Rakovec et al. (2015) described the method for state estimation experiments.

#### 3.4.2 The Asynchronous EnKF

To start with,  $k$  represents the assimilation cycle index but it needs to be distinguished from the day index (within the assimilation cycle) which is the time unit for the observations. We will then denote by  $k(i)$ ,  $i = 1 \dots 21$ , the  $i$ -th day in the current assimilation cycle.

On the  $i$ -th day of the  $k$ -th assimilation cycle, the  $n_{y,k(i)}$  observations are gathered in the vector  $\mathbf{y}_{k(i)}^o$ . The overall observation vector at cycle  $k$ ,  $\mathbf{y}_k^o$ , then concatenates the 21 daily observation vectors  $\mathbf{y}_{k(i)}^o$  so that:

$$\mathbf{y}_k^o = \left[ \mathbf{y}_{k(1)}^{o,T} \cdots \mathbf{y}_{k(21)}^{o,T} \right]^T \in \mathbb{R}^{n_{y,k}}, \quad n_{y,k} = \sum_{i=1}^{21} n_{y,k(i)}. \quad (15)$$

Similarly to the observation vector, the overall observation operator at the  $k$ -th cycle,  $\mathcal{H}_k$ , is the concatenation of the daily observation operators  $\mathcal{H}_{k(i)}$ , defined from Eq. 9, but by considering the operator  $\mathcal{M}$  integrating the model between  $k(0)$  ( $= (k-1)(21)$ ) and  $k(i)$  as well as the diagnostic and selection operators a time step  $k(i)$ ,  $\mathcal{Z}_{k(i)}$  and  $\mathcal{S}_{k(i)}$ .

The observation error covariance matrix  $\mathbf{R}_k$  is the concatenation of the daily observation error covariance matrices:

$$\mathbf{R}_k = \begin{pmatrix} \mathbf{R}_{k(1)} & & & & \\ & \mathbf{R}_{k(2)} & 0 & & \\ & 0 & \ddots & & \\ & & & \ddots & \\ & & & & \mathbf{R}_{k(21)} \end{pmatrix} \quad \text{with } \mathbf{R}_{k(i)} = (\sigma^o)^2 \mathbf{I}_{n_{y,k(i)}}, \quad (16)$$

where  $\mathbf{I}_{n_{y,k(i)}}$  is the identity matrix of size  $n_{y,k(i)} \times n_{y,k(i)}$ . It turns out that:

$$\mathbf{R}_k = (\sigma^o)^2 \mathbf{I}_{n_y}. \quad (17)$$

Following the same equations as the EnKF, the AEnKF generates, for each member  $l = 1 \dots n_e$  of the control ensemble, an analysis control vector  $\mathbf{x}_k^{a,[l]}$ .

### 3.4.3 Generation of the ensemble

To generate the background control ensemble, we solely stochastically perturb the variables within the control vector. Note that it amounts to the assumption that all other features of the forward model, e.g. the atmospheric forcings, the LSM structure and therefore the surface and sub-surface runoff, are perfect. While this applies for OSSEs, such features are never perfect in real-case experiments. This assumption is further discussed in Section 7 [R#1-m#11].

The ensemble of background control vectors  $\mathbf{X}_{e,k}^b$ , of size  $n_x \times n_e$ , is generated so that  $\mathbf{x}_k^{b,[l]}$ ,  $l = 1 \dots n_e$ , follows a gaussian law of mean  $\overline{\mathbf{x}_k^b}$  and covariance matrix  $\mathbf{P}_{e,k}^b$ . For the first assimilation cycle, the control variables mean value  $\overline{\mathbf{x}_1^b}$  is arbitrarily chosen as the openloop run input parameter (the openloop or free run is the model run without assimilation) and the background error covariance matrix  $\mathbf{P}_{e,1}^b$  is a diagonal matrix defined as

$$\mathbf{P}_{e,1}^b = ((\sigma^b)^2 \mathbb{I}_{n_x})$$

with  $\mathbb{I}_{n_x}$  the identity matrix of size  $n_x \times n_x$  and  $\sigma^b$  the vector that gathers the initial control variable error standard deviation.

Once the analysis ensemble  $\mathbf{X}_{e,k}^a$  is determined, the next step is to propagate the correction in time. In a PE framework, it is necessary to re-run the ensemble model runs during the current assimilation window with the analysis parameters as inputs. Then, the contribution of the updated parameters is propagated through the model, up to the end of the current assimilation window and put into the model initial condition for the next assimilation cycle.

For the next assimilation cycles, the background mean estimate is set equal to the analysis mean estimate from the previous cycle:

$$\overline{\mathbf{x}}_k^b = \overline{\mathbf{x}}_{k-1}^a.$$

There are different ways of defining  $\mathbf{P}_{e,k}^b$ . One could choose to stochastically estimate  $\mathbf{P}_{e,k-1}^a$  from the analysis ensemble at the previous cycle and use it as  $\mathbf{P}_{e,k}^b$ . Contrary to state estimation experiments where the analysis error covariance matrix is propagated in time using the model along with the control variables, parameter estimation experiments use it directly as the background error covariance matrix as there is no dynamical model for the Manning coefficient. The issue with this approach is that the analysis ensemble variance can be strongly reduced and provide too small an ensemble spread to have efficient AEnKF updates in time. To ensure that enough uncertainty is maintained in the ensemble, one can maintain the initial background error covariance matrix through all cycles or impose a minimal value for the variance elements (see Section 5.4).

The background error cross-covariance matrix  $[\mathbf{PH}^T]_{e,k}$  and covariances matrix  $[\mathbf{HPH}^T]_{e,k}$  are directly built from the definition suggested by Evensen (2004), Moradkhani et al. (2005) and Durand et al. (2008), see Appendix A for more details. The matrices are of size  $n_x \times n_{y,k}$  and  $n_{y,k} \times n_{y,k}$  respectively. The elements in the error cross-covariance matrices result directly from the characterization of the background ensemble, namely the parameter uncertainties accounted for for the generating of the control matrix  $\mathbf{X}_{e,k}^b$  and  $\mathcal{H}(\mathbf{X}_{e,k}^b)$ .

## 4 Assimilation strategy

In the incoming experiments, the true control variables  $\mathbf{x}^t$  are:

$$\mathbf{x}^t = \begin{bmatrix} 1.65 & 0.85 & 0.85 & 0.95 & 0.90 & 0.95 & 0.90 & 1.30 & 1.40 \end{bmatrix} \quad (18)$$

and the a priori values  $\overline{\mathbf{x}}_1^b$  at the first assimilation cycle are:

$$\overline{\mathbf{x}}_1^b = \begin{bmatrix} 1.50 & 0.50 & 0.50 & 0.50 & 0.50 & 0.50 & 0.50 & 1.50 & 1.50 \end{bmatrix}. \quad (19)$$

We increase the Manning value in mountainous zones (zones 1 in the Andes and zones 8 and 9 over the shields) and lower the Manning value over the other zones with the lowest values (in zones 2 and 3) corresponding to the main stem. Both true and background values were chosen accordingly.

### 4.1 Sensitivity tests

During one EnKF assimilation cycle, the analysis potentially depends on the following parameters: model spin up, time period (high/low flow), size of the ensemble, control error. Note that the observation error also has an impact on the analysis but its value is already fixed for all subsequent experiments (see Section 5).

A first set of experiments (either model runs or data assimilation runs) will serve as sensitivity tests for the data assimilation platform with respect to the above features. During these sensitivity tests, the different features are tested individually. Table 1 details the range of variations for each tested feature.

## 4.2 Assimilation tests

Following the sensitivity tests, a set of three data assimilation experiments will be run and is presented in Table 2. The data assimilation experiments are divided into two categories: the first one uses water depths as observations, the second considers water depth anomalies. All experiments are run across a year, corresponding to 17 assimilation cycles of 21 days.

5 The first experiment, denoted as PE1, is configured from the aforementioned sensitivity test outcome. The parameters defining the experiment (spinup, starting date, ensemble size, control error) will be those which provide the best results in the sensitivity tests in Table 2. The reference level between the observed and simulated water depths is also the same. In other words, there is no bias in the observation. This first idealized experiment serves as proof-of-concept as the observations type matches exactly the type of the simulated variables. Consequently, with this experiment, we expect to retrieve the true value of  
10 the control variables and hence the correct water depths and discharges [R#1-m#14].

The next step is to head towards more realistic experiments by including new sources of uncertainties in the data assimilation system and seeing how to address them. In this context, two additional experiments denoted as PE2 and PE3 will be carried out. As an example of new uncertainties, SWOT will in fact observe water elevations (water surface elevation as referenced to a geoid or an ellipsoid) whilst CTRIP produces water depths (water surface elevation as referenced to the bottom of the river  
15 bed). To perform data assimilation, one needs to convert CTRIP water depths ( $h_S^{\text{CTRIP}}$ ) into CTRIP water elevations ( $H_{\text{alti}}^{\text{CTRIP}}$ ) or inversely for SWOT. It is highly plausible that this operation induces a bias between the modeled and observed water elevations. A simplified example of such situation is illustrated in Figure 4. In this case, SWOT catches the right water elevation dynamic (as  $h_S^{\text{SWOT}}$  and  $h_S^{\text{CTRIP}}$  are equal) but the direct assimilation of SWOT water elevation  $H_{\text{alti}}^{\text{SWOT}}$  will induce a bias as the elevations of the river bed ( $H_{\text{bed}}^{\text{SWOT}}$  and  $H_{\text{bed}}^{\text{CTRIP}}$ ) are different between CTRIP and SWOT.

20 A solution for the handling of this issue is to assimilate water depth anomalies instead of water depths. The next data assimilation experiments, denoted as PE2 and PE3, will therefore test the feasibility of assimilating anomalies. In these experiments, the water depth anomalies are generated by subtracting a time-averaged reference water depth from the current water depth. For all runs (true, openloop or analysis), this time-averaged reference water depth is computed as the mean (true, openloop or analysis) water depth over the year before the start of the assimilation window. It is therefore different for each member of  
25 the ensemble. Firstly, in experiment PE2, there will still be no bias between the observed and simulated river bathymetry to observe how the assimilation of anomalies performs. Similarly to PE1, we expect this experiment to be able to retrieve the true control and state variables [R#1-m#14]. Finally, the last experiment PE3, which introduces a constant relative bias between CTRIP and SWOT, will be carried out. For this experiment, we anticipate that the assimilation will still be able to retrieve the model states variables. The use of anomalies as observations should limit the impact of the inserted bias. We do not exclude  
30 however that it may be slightly echoed on the control variables [R#1-m#14].

## 5 Assimilation sensitivity tests

### 5.1 Model spinup sensitivity tests

The objective of the spinup sensitivity tests is to evaluate the minimum spinup period required by the model before applying data assimilation. For this purpose, the model is run several times across two years, from December 19th, 2006 to December 22nd, 2008, corresponding to 35 windows of 21 days (735 days).

A first simulation is run using the true Manning spatial distribution (see Eq. 18) over the two-year time period. We then run 18 additional simulations over the same period with a varying length of the spinup period (see Table 1). Initially, the simulation setup corresponds to the openloop configuration with the openloop Manning spatial distribution (see Eq. 19). At a given time during the first year, the Manning spatial distribution is instantaneously changed to the true distribution (see Eq. 18) and the model is run until the end of the two years with the true Manning spatial distribution. Table A1 summarizes for each run, the date when the Manning coefficients are changed. The spinup period (expressed as a number of windows of 21 days) corresponds to the period between when the Manning distribution is changed and the start of the second year, i.e. January 1st, 2008.

To evaluate the spinup impact, the relative difference between the reference run and the test runs is evaluated over the second year of simulation (from January 1st to December 22nd, 2008) and averaged over every window of 21 days. Figure 5 presents the results for the spinup sensitivity test. Each test run (on the x-axis) is identified by its corresponding spinup period length (expressed as the number of windows of 21 days, see Table A1). We then count (on the y-axis) the number of 21-day windows during which the relative error between the test run and the reference run is higher than a given threshold. We assume that the spinup period is long enough when this number is equal to 0. This number is evaluated from the basin-averaged relative difference and from the relative difference at the downstream station of Obidos, both in terms of water depth and discharge. Note that in Figure 5, when the number of spinup windows of 21 days is equal to 4 on the x-axis, the change from the openloop Manning distribution to the true one is imposed on October 9th, 2007. Similarly, when this number is equal to 10, the change is imposed on June 5th, 2007. Note also that two thresholds are considered, 0.01 and 0.001. Basin-averaged results are not sensitive to this threshold. There are some differences at Obidos; however, we retain the basin-averaged results to evaluate the required model spinup period.

From all of the results we conclude that a minimum spinup period of 4 windows of 21 cycles, i.e. 84 days, is required. **This period corresponds to the basin concentration time or, in other terms, the required time for the river network to totally empty [R#1-m#15].** In the following sensitivity tests, the model runs start on October 9th, 2007, to be consistent with these results.

### 5.2 Data assimilation starting date sensitivity tests

To evaluate the impact of the starting date, a set of 17 one-cycle-long data assimilation experiments is carried out over the second running year. All experiments have the same general configuration except for the initial date starting from January 1st, 2008 and shifted by 21 days until December 22nd, 2008. This means that the last experiment starts on December 2nd, 2008. The performance of each experiment is evaluated by simply evaluating the spatial average difference between the analysis and



the true Manning coefficients. Results presented in Figure 6a-b indicate that there are no significant differences between the data assimilation experiments (for all 17 experiments, the error of the updated Manning coefficients with respect to the true value of the coefficients is below 5 %).

Note that the results in Figure 6a-b show a slight increase in the errors for the experiment starting at the end of the year, presently from September 9th to December 2nd, 2008. This period corresponds to the low-flow season in the Amazon hydrological cycle. Concerning the sensitivity analysis results (Emery et al., 2016), the water depths showed a very low sensitivity to the Manning coefficient during the low flow season. As a consequence of data assimilation, the EnKF is less effective in low flow seasons in the correcting of the Manning coefficient. The analysis relative error (in Figure 6a) and the analysis ensemble dispersion (in Figure 6b) is therefore higher in the low flow season.

For all of the following sensitivity tests, we are only considering therefore one-assimilation-cycle experiments, which will start on January 1st, 2008.

### 5.3 Ensemble size sensitivity tests

The next sensitivity test is dedicated to the ensemble size  $n_e$ , a critical parameter of any EnKF algorithm. This parameter has to be high enough so as to accurately estimate the Kalman gain matrix but low enough to limit the computational cost (the higher  $n_e$ , the more model runs are required over each data assimilation window to obtain the analysis estimate of the Manning coefficients).

We consider different ensemble sizes through a one-assimilation-cycle experiment,  $n_e$  varies between 10 and 200. Figure 6c compares the analysis Manning coefficient relative error for each ensemble size  $n_e$  as in Figure 6a. Results show that the analysis relative error decreases when the ensemble size  $n_e$  increases. For an ensemble size  $n_e$  equal to 20, the analysis error is below 5 %. Also, for an ensemble size  $n_e$  higher than 50, the analysis relative error has converged to a constant value while the analysis ensemble dispersion showed in Figure 6d stabilises.

These results indicate that the ensemble size for future data assimilation experiments should be at least equal to 20; we consider  $n_e = 25$  in the present study to limit computational time.

### 5.4 Model error standard deviation sensitivity tests

In this study, we only consider parameter estimation, implying that the background error covariance matrix is associated with the parameter space. We assume that the errors in the Manning coefficients are independent so that the background error covariance matrix is initially specified as a diagonal matrix, where all diagonal elements correspond to the error variances in the spatially-varying Manning coefficients and are equal to the same variance  $(\sigma^b)^2$ , where  $\sigma^b$  is the background error standard deviation. Similarly to previous sensitivity tests, we study here the sensitivity of the data assimilation results to the value of  $\sigma^b$  through a one-assimilation-cycle experiment. Figure 6c shows, in logarithmic-scale on the x-axis, the relative error of the updated Manning coefficient with respect to the true coefficient. The analysis error curve shows a decreasing behavior until  $\sigma^b$  is in the order of 0.4. It then increases again.

Note that the actual Manning coefficient error before data assimilation is equal to 0.33 (see the blue curve in Figure 6c showing the openloop Manning coefficient error). Consistently, the best data assimilation results are obtained when  $\sigma^b$  provides a good approximation of the real error standard deviation. Note also that when  $\sigma^b$  becomes too small, data assimilation is less effective. The EnKF algorithm is known to be under-dispersive. Therefore, for future data assimilation experiments, when updating the error covariance matrix from one cycle to another, we will need to make sure that the ensemble dispersion is high enough to cover possible model behavior over the forecast time window by imposing a minimum value for the error variance. Given the sensitivity test results, the minimum value for  $\sigma^b$  is set to 0.005. Thus, in the following data assimilation experiments, we use the analysis error covariance matrix as the background error covariance matrix for the next assimilation cycle, while applying the minimum threshold value on the matrix diagonal terms.

## 6 Data assimilation results

We now present the results from the data assimilation experiments described in Table 2 and in Section 4.4.2. Recall that these experiments aim at correcting a set of 9 multiplying factors applied to the Manning coefficients distribution and constant s over 9 hydro-geomorphological zones dividing the Amazon basin [R#1-M#2].

### 6.1 Assimilation of water depths (PE1)

Figure 7 gives, for each zone, the time evolution of the mean analysis control variable (red) with its dispersion (even though it is very narrow) compared to the truth (black) and the first guess (blue). Similarly, Figure 8 shows, for each zone, the time evolution of the analysis water depth (red) compared to the truth (black) and the openloop (blue). To generate one plot per zone, we use, for each time step, the ensemble of water depths over all cells in the zone and estimate the median value, the first decile and the ninth decile. Furthermore, zone-averaged normalized Root Mean Square Error (RMSEn) statistics are given in Tables A2 and A3 in Appendix D [R#1-M#1].

In general, the PE1 experiment gives very good results as the analysis mean for each zone retrieves the true value with a very low dispersion. However, the data assimilation algorithm features spatially-dependent behavior, see Figure 7):

- Firstly, the control variable for the zones 1, 2, 3, 4, 5 and 9 converges instantaneously (in only one assimilation cycle) toward the true values and remains at these true values for all following cycles.
- A similar behavior can be observed for zones 6, 7 and 8 from the first cycle to around the ninth cycle. For the remaining cycles, we notice an increase in the mean analysis estimate, along with the ensemble dispersion, until around the thirteenth cycle and after, a decrease back to the true value.

These observations can be explained with the global sensitivity analysis results for water depths in Emery et al. (2016) and using Figure 8:

- Firstly, zones 1, 2 and 3 correspond to the river main stem, whilst zones 4, 5 and 9 correspond to the main left-bank tributaries, namely the Caquetá/Japurá river (zone 4) and the Negro river (zone 5). In these zones, as the Manning

coefficient is directly corrected in the first assimilation cycle, the analysis water depths (red) overlap the true water depth (black). Furthermore, Figure 8 for these zones shows that the openloop (blue) and true (black) water depths have a very similar variability in time but differ by a constant bias. The global sensitivity analysis results in these zones showed a constant first-order sensitivity in time to the Manning coefficient all year long. This first-order sensitivity means that the contribution of the Manning coefficient to the water depth is linear. Correcting the Manning coefficient in these zones equates therefore to correcting the bias between the openloop and the true water depths.

- Subsequently, zones 6, 7 and 8 correspond to right-bank tributaries, namely the Juruá and Purus rivers (zone 6), the Madeira river (zone 7) and the Tapajós and Xingu rivers (zone 8). These right-bank tributary zones are characterized by a strong seasonal cycle (see Figure 8, zones 6-8). By comparing the corresponding plots in Figures 7 and 8, we notice that the period when the analysis control variable spreads from the truth corresponds to the low flow season in these zones. According to the global sensitivity analysis results, water depths in these zones are less sensitive to the Manning coefficient in low flow conditions. Additionally, there is very little water in the zones during this period and consequently, the background control ensemble is not spread out enough for the EnKF to be efficient [R#1-m#17]. Meanwhile, the EnKF still sees that the observations are higher than the model predictions (as seen with the positive innovations in these zones shown in Figure A2) [R#1-m#17]. In order to increase the simulated water depth, the EnKF therefore corrects the Manning coefficient so that its value rises (a higher Manning coefficient means a slower flow velocity and then a higher simulated water depth). Finally, once the low flow season ends, the analysis Manning coefficient converges back to the truth (see the last assimilation cycles).

## 6.2 Assimilation of water anomalies (PE2 and PE3)

The assimilation of anomalies has been tested over two experiments denoted PE2 and PE3 (see Table 2). Note that the observation error standard deviation  $\sigma^o$  remains equal to 10 cm as, with these experiments, we only aim at testing the feasibility of assimilating water depth anomalies. In the PE2 experiment, there is no difference of bathymetry between the simulated and observed water anomalies whilst the river bankful depth is different in the PE3 experiment [R#2-m#14]. Figure 9 gives, for each zone, the time evolution of the mean analysis control variable for PE2 (orange) and PE3 (purple) with their dispersion compared to the truth (black) and the first guess (blue). Again, zone-averaged normalized Root Mean Square Error (RMSEn) statistics for these experiments are given in Tables A2 and A3 in Appendix D [R#1-M#1].

The general configuration of the experiments PE1 and PE2 is the same. The only difference between the two experiments is the nature of the observations: water depths for PE1 and water anomalies for PE2. Like experiment PE1, experiment PE2 (the orange line in Figure 9) gives very good results. All control variables converge toward the true values more or less rapidly. The control variable for the zones 4 to 8 instantaneously (in only one assimilation cycle) converges toward the true value while the convergence is slower for the remaining zones as around five assimilation cycles are needed to retrieve the true value. This slower convergence for these zones can be explained by the fact that the magnitude of the observed water anomalies is generally smaller, compared to the water depths assimilated in PE1. The ratio between the observation error and

the observations themselves is also then smaller, resulting in a smaller EnKF gain. The control variable correcting increment is smaller for the anomalies therefore than for the water depth and more cycles are needed to converge.

As for the PE3 experiment results - the purple line in Figure 9 - the assimilation still gives good results but not as good as in previous experiments. The control variables still instantaneously converge toward the truth in the zones 4, 5 and 6. Concerning the other zones, there is no clear convergence towards the true value. Instead, the analysis control variables either get closer but remain distinct from the true value (zones 2 and 3) or temporarily deviate from the truth during the experiment (zones 1, 7, 8 and 9). Still, despite the control variables not clearly converging towards the truth, the simulated water depths using the analysis control variables, presented in Figure 10, display a very low deviation from the truth, confirming the general good performance of the data assimilation procedure [R#2-m#4].

Comparing the control variables and water depth time variations, it appears that the control variables are deviating from the truth mainly when water depths are decreasing, in between the high flow and low flow seasons. During this period, the model goes from a state where floods occur to a state where there is no flood, particularly in the zones 2-3 and 7-8 with a clear seasonal cycle. On the other hand, no flood event was spotted in zones 4, 5 and 6 where the best results were obtained. In ISBA-CTRIP, the activation/deactivation of the flooding scheme is triggered by the simulated water depth exceeding/becoming lower than the river bankful depth. Yet, in the experiment PE3, this river bankful depth differs between the model and the observation because we artificially inserted a bias between the simulated and observed water depths. More specifically, the river bankful depth is lower in the model than in the observations. Therefore, the control variables deviating from the true value when the water depth is decreasing are indicative of the simulated water depths presenting floods when there is no flood in the observations. The activation of the flood scheme changes the dynamics of the water depth in the river. As part of the water in the river is spilled into the floodplains, water level variations in the river are slower. The flooded model needs then a stronger variation of the Manning coefficient in order to catch the non-flooded observed water level. Ultimately, the stronger variations of the estimated Manning coefficient allow the retrieval of the true water depths.

## 7 Discussions

The results presented here are preliminary investigations into the assimilation of SWOT water surface elevations product into a large-scale hydrological model. This study focused on the correction of a critical river parameter, here the river Manning coefficient [R#1-m#15].

For all simulations, the Manning coefficient distribution is set to be constant in time. For each grid cell, one value of the Manning coefficient is used for the entire simulation. However, in reality, it is commonly accepted that this parameter could vary in time, depending on the seasonal cycle or also some extreme hydrological event such as large flooding events, which can even modify the bathymetry itself [R#1-m#19]. The results showed that, for this OSSE, the data assimilation is able to converge quite quickly towards the true value. For example, for the left-bank tributaries zones, namely zones 4 and 5, in every experiment, the associated control variable converges toward the true value in only one assimilation cycle. In a real-case experiment, we could expect to retrieve the temporal variations of the Manning coefficient from one assimilation cycle to

another. The good performances of the assimilation platform are mainly related to the fact that, in the ISBA-CTRIIP model, the water depth diagnostic variables are sensitive to the Manning coefficient (Emery et al., 2016). Simulated water depths are not then that sensitive to the Manning coefficient (e.g. in the right-bank tributary zones during the low flow season), the data assimilation performances slightly degrade. These results are specific to the ISBA-CTRIIP model. To apply the same method to another model and even another region, one needs to first study the sensitivity of the (other) model to the (other) study region.

Secondly, the study investigates the potential of assimilating water surface anomalies instead of direct water surface elevations. The use of water surface anomalies is driven by the need to avoid potential bias between the control and the observed variables [R#1-m#15]. Indeed, a bias will likely be introduced from a discrepancy between the elevation of the river bed in the model and in the observations with respect to a reference surface such as a geoid or an ellipsoid (see Figure 4). Under the assumption that the water variations are the same between the model and the observations, the use of anomalies as observed variables should prevent this bias from affecting the results.

Another likely bathymetry error corresponds to errors on the river bankful depth, the river width and more generally, representativeness errors due to the use of a simplified bathymetry [R#2-M#2]. This type of error was artificially introduced by perturbing the model bankful depth in PE3. Specifically to ISBA-CTRIIP, the river bankful depth controls when the model floods, which has a direct impact on the water depth dynamics. Background anomalies and observed anomalies may therefore present different dynamics where either the observed variables flood while the model variables do not or inversely. The experiment PE3 illustrated the effect of this bias on the variations of Manning coefficients. Instead of being maintained at their true value, their value slightly varied around the true values to account for the difference in dynamics between the model and the observations. However, one could expect even more variations in the updated control variables around the true value to increase if more and different errors in the bathymetry exist (which will likely happen with more realistic experiments) [R#2-M#2].

Furthermore, real-case experiments may suffer from another type of bias originating from errors in the atmospheric forcing and in the surface and sub-surface runoff provided by the LSM (i.e. ISBA). Both control the amount of water entering the river system. A basic idea to attenuate this issue would be to consider their uncertainties when generating the background ensemble [R#1-m#11]. This approach may become limited when the errors in the forcing are very large. it may also lead to unrealistic, even non-physical, updated Manning coefficient values. Besides, when correcting the model's parameters, we only re-distribute the water volume within the basin whilst such types of errors could actually require adding/withdrawing water to/from the system. The potential solution would then be to include such forcing or LSM variables in the control vector or, to update variables closer to the observations, including CTRIIP's state variables such as the water storage. This would change the current framework to a dual state-parameter estimation approach [R#1-M#1][R#1-m#18][R#2-M#3].

Noting this, there may be an additional advantage in assimilating water anomalies instead of the direct water depths. Comparing the Kalman gain between the PE1 experiment (which assimilated direct water depths) and the PE2/PE3 experiments (which assimilated water anomalies), the gain magnitude for the water anomalies is lower than the water depth gain magnitude. This is to be expected as the Kalman gain is stochastically estimated from an ensemble of model runs and the magnitude of the simulated water anomalies is lower than the simulated water depth magnitude. The consequence of this lower gain is

that the correction applied to the control variable is also lower. If the convergence towards the true value takes more than one assimilation cycle, the divergence from it in the presence of bias is also diminished.

Beyond the bias issues, real-data assimilation configuration will raise the question of the unknown true parameter value, if it exists. Firstly, there will be no true estimates of the control variables with which to compare the assimilated simulations.

5 The assimilation will be evaluated against the observed variables directly. Then, with the real data, model structure error will be introduced. To our knowledge, the model structure error is still a challenging error to estimate and most data assimilation studies assume no model structure error. However, when using an ensemble-based model, a possibility for dealing with such structure error is to enrich the background ensemble by considering more uncertainties from variables that are not necessarily in the control vector (including errors in the forcing or parameters from both the LSM and the RRM). The capacity of such ensembles to tackle model structure errors can be tested using synthetic observations based on a different hydrological model [R#2-M#2].

15 Additionally, the real SWOT data will have a finer resolution than the synthetic SWOT data currently used. Still, the coarser resolution observations are found to provide information to constrain the model and to improve the value of the spatially-varying Manning coefficients. Then, when moving to real-data assimilation experiments, we can consider averaging the fine-scale SWOT product over a coarse grid cell corresponding to an ISBA-CTRIP cell so that the resolution of the observations and the model matches.

20 Ultimately, heading towards more realistic experiments also implies more realistic representations of the observation errors. But more complex errors should be expected for the real SWOT product. Some correlated errors along the swath should be expected due to the instrument but also to the motion of the satellite and delays due to propagation of the electromagnetic waves in the ionosphere and atmosphere. Nevertheless, as part of the mission science requirements, the sum of all errors should not exceed 10 cm when the measured data is averaged over 1 km<sup>2</sup>. There should also be additional errors affecting the observations that can be described as "detectability errors" such as "dark water", "layover" and "false positive". "Dark water" pixels will result in missing data and will not be included in the assimilation, "layover" pixels will have a higher vertical error due to surrounding vegetation and topography, but should also be flagged (Biancamaria et al., 2016). Eventually, "false positive" pixels (i.e. pixels classified as water, whereas they correspond to land) will be the most complicated to anticipate. With these additional errors taken into account in the assimilation framework, one could expect a slower convergence of the control variables. Note that these aspects of the measurement errors are related to water surface elevation products [R#1-m#12].

## 8 Conclusions

30 This study presents a series of OSSE that assimilates SWOT-like synthetic observations of water depths and anomalies into the large-scale hydrological model ISBA-CTRIP in order to correct the spatially-distributed Manning coefficient. The study is applied over the Amazon river basin. Prior to the actual data assimilation experiments, a series of sensitivity tests was conducted to study the sensitivity of the data assimilation performance to the different features of the EnKF, in particular the size of the

ensemble. Then, three full-year data assimilation experiments were run based on the outcomes of the sensitivity tests. For all three experiments, the assimilation was able to track back the true value of the Manning coefficient distribution.

The sensitivity tests successively studied the sensitivity of the data assimilation platform to model spinup period, the experiment starting date through the hydrological year, the size of the ensemble for the EnKF and the initial control variable standard deviation. These tests showed first of all that a spin up of four windows of 21 days is sufficient for the transitional period due to a sudden change in the Manning coefficient distribution in the model. The second sensitivity test then demonstrated that the data assimilation performance is not clearly sensitive to the period of the hydrological year when the experiment is done. The next sensitivity test informed us that an ensemble of 25 members was enough to obtain good EnKF performances. Finally, the last sensitivity test studied the effect of the control variable error standard deviation and the best performances were obtained for prior standard deviation between 0.05 and 0.75, which corresponds to the order of magnitude of the actual error between the true and openloop control variables.

Using these results, we run three data assimilation experiments over approximately one year (the year 2008). The first experiment (PE1) assimilated direct pseudo-observations of water depths. Results showed the capability of the data assimilation algorithm to converge very quickly toward the true value, generally in only one assimilation cycle. Still, during the low flow season, the assimilation was less effective in the zones with a clear seasonal cycle. This was explained by the fact that during this period, water depths are less sensitive to the Manning coefficient.

The two other experiments (PE2 and PE3) introduced and tested the assimilation of water surface anomalies. The anomalies were obtained by subtracting a yearly-averaged water depth from the current water depth in both the model and the observations. The first water anomalies assimilation experiment (PE2) provided very good results with all the control variables also converging towards their associated true values. However, the convergence was slightly slower than during the assimilation of the water depth (between 1 and 5 assimilation cycles). This is explained by a lower Kalman gain when updating the Manning coefficient.

The last experiment also assimilated water anomalies (PE3). For this particular experiment however a bias was artificially introduced in the river bathymetry. For this experiment, the assimilation was still able to get closer to the true value but, for the some zones like the mainstream zones, there was no convergence as the control variables kept varying around the true value. This phenomenon was explained by the detection of floods in the model but not in the observations. Still, the statistics of the Manning coefficient distribution and the simulated water depths after assimilation remain as improved compared to the openloop simulations. Ultimately, these two experiments demonstrated the feasibility of assimilating water surface anomalies to correct the Manning coefficient.

These experiments offer several perspectives. They mainly consist of approaching more realistic data assimilation experiments which take into account more sources of uncertainties between the model and the observations, such as correlated observation errors or uncertainties in the forcing and the LSM surface and sub-surface runoff [R#1-M#1]. To test the platform's limitations regarding the DEM/bathymetry bias issue, one can use simulated water surface elevations referenced to a geoid instead of water depths from the model or even assimilate water depths from another model where the bathymetry differs. As most applications generally require a good estimate of the river flow and river water volume, another lead of investigation could

maintain the SWOT-based OSSE framework but correct the simulated water storage and/or discharge, either as a single state estimation framework or as a dual state-parameter estimation framework (similarly to dual discharge-bathymetry inference methods developed by Oubanas et al., 2018 and Brisset et al., 2018 for some hydraulic models) [R#1-M#1]. Moreover, along with observations of water surface elevations, SWOT will also provide two-dimensional maps of river widths and surface slopes. One can also study the possibility of assimilating such products to constrain other parameters such as the bankful depth that controls the model flooding scheme.

*Code and data availability.* The CTRIP code is open source and is available as a part of the surface modelling platform called SURFEX, which can be downloaded at <http://www.cnrm-game-meteo.fr/surfex/>. SURFEX is updated approximately every 3 to 6 months and the CTRIP version presented in this paper is from SURFEX version 7.3. If more frequent updates are needed, please follow the procedure informing you of how to obtain a SVN or Git account in order to access real-time modifications of the code (see the instructions in the previous link). The ISBA-CTRIP model is coupled to the DA codes via the OpenPalm coupler available at [http://www.cerfacs.fr/globc/PALM\\_WEB/](http://www.cerfacs.fr/globc/PALM_WEB/). To get the DA routines coupled to ISBA-CTRIP with OpenPalm, please directly contact C. Emery ([charlotte.emery@jpl.nasa.gov](mailto:charlotte.emery@jpl.nasa.gov)) or S. Biancamaria ([sylvain.biancamaria@legos.obs-mip.fr](mailto:sylvain.biancamaria@legos.obs-mip.fr)). To obtain the GSWP3 forcings, please refer to the following url: [http://search.diasjp.net/en/dataset/GSWP3\\_EXP1\\_Forcing](http://search.diasjp.net/en/dataset/GSWP3_EXP1_Forcing) (<https://doi.org/10.20783/DIAS.501>).

## 15 Appendix A: Definition of error covariance matrices

The background error cross-covariance matrices  $[\mathbf{PH}^T]_{e,k}$  and  $[\mathbf{HPH}^T]_{e,k}$  are defined based on Evensen (2004); Moradkhani et al. (2005); Durand et al. (2008) so that:

$$[\mathbf{PH}^T]_{e,k} = (n_e - 1)^{-1} \left( \mathbf{X}_{e,k}^b - \overline{\mathbf{X}_{\bullet,k}^b} \cdot \mathbf{1}_{n_e}^T \right) \left( \mathcal{H}(\mathbf{X}_{e,k}^b) - \overline{\mathcal{H}(\mathbf{X}_{\bullet,k}^b)} \cdot \mathbf{1}_{n_e}^T \right)^T, \quad (\text{A1})$$

and

$$[\mathbf{HPH}^T]_{e,k} = (n_e - 1)^{-1} \left( \mathcal{H}(\mathbf{X}_{e,k}^b) - \overline{\mathcal{H}(\mathbf{X}_{\bullet,k}^b)} \cdot \mathbf{1}_{n_e}^T \right) \left( \mathcal{H}(\mathbf{X}_{e,k}^b) - \overline{\mathcal{H}(\mathbf{X}_{\bullet,k}^b)} \cdot \mathbf{1}_{n_e}^T \right)^T. \quad (\text{A2})$$

In those definitions,  $\mathbf{X}_{e,k}^b$  is the control matrix storing the  $n_e$  control vectors  $\mathbf{x}_k^{b,[l]}$ ,  $l = 1 \dots n_e$ , from the background ensemble such that

$$\mathbf{X}_{e,k}^b = \begin{bmatrix} \mathbf{x}_k^{b,[1]} & \dots & \mathbf{x}_k^{b,[N_e]} \end{bmatrix}$$

Next,  $\mathcal{H}(\mathbf{X}_{e,k}^b)$  represents the same control matrix but mapped into the observation space:

$$\mathcal{H}(\mathbf{X}_{e,k}^b) = \begin{bmatrix} \mathcal{H}(\mathbf{x}_k^{b,[1]}) & \dots & \mathcal{H}(\mathbf{x}_k^{b,[n_e]}) \end{bmatrix}.$$

Also,  $\overline{\mathbf{X}_{\bullet,k}^b}$  and  $\overline{\mathcal{H}(\mathbf{X}_{\bullet,k}^b)}$  are the corresponding ensemble expectations such that

$$\overline{\mathbf{X}_{\bullet,k}^b} = \frac{1}{n_e} \sum_{l=1}^{n_e} \mathbf{x}_k^{b,[l]} \quad \overline{\mathcal{H}(\mathbf{X}_{\bullet,k}^b)} = \frac{1}{n_e} \sum_{l=1}^{n_e} \mathcal{H}(\mathbf{x}_k^{b,[l]}).$$



These vectors dimension are  $n_x$  and  $n_{y,k}$  respectively. Finally,  $\mathbf{1}_{n_e}$  is a vector of size  $n_e$  containing only 1s.

## Appendix B: Spinup sensitivity test additional tables

Table A1 summarizes for each run, the date when the Manning coefficients are changed. The spinup period (expressed as a number of windows of 21 days) corresponds to the period between when the Manning distribution is changed and the start of the second year, i.e. January 1st, 2008.

## Appendix C: Sensitivity tests results per zones [R#1-m#16]

Figure A1 displays the sensitivity tests results (as in Figure 6) but for each zone separately [R#1-m#16].

## Appendix D: Assimilation performances at the zones scales [R#1-M#1]

At each grid-cell of the study domain, we estimated the normalized Root Mean Square Error (RMSEn) before and after assimilation by comparing the openloop and the mean analysis simulations respectively to the true simulation, for both the simulated water depth and discharge:

$$\text{RMSEn}_i = \frac{\sqrt{\frac{1}{N} \sum_{n=1}^N (V_{n,i}^* - V_{n,i}^t)^2}}{\overline{V_{.,i}^t}}, \quad (\text{D1})$$

where the state variable  $V$  is either the discharge or the water depth,  $n$  is the time indice,  $i$  is the grid-cell indice, the  $t$  superscript represents the "truth" and the  $*$  superscript represents either the openloop or the analysis ensemble average.

Table A2-A3 gives these statistics for all experiments averaged over each control zone. Table A2 shows the water depths zone-averaged RMSEn and Table A3 shows the discharge zone-averaged RMSEn.

## Appendix E: Assimilation results additional figures

Figure A2 displays the evolution along the assimilation cycles of the averaged innovations. The sign of the innovation will drive the direction of the correction brought by the assimilation:

- A positive innovation means that the observations are higher than the model. Physically, the simulated flow is too fast and the water leaves the river reservoir too quickly. This means that the river Manning coefficient needs to be increased to slow the flow,
- A negative innovation means that the observations are lower than the model. Physically, the simulated flow is too slow and the water remains in the river reservoir. This means that the river Manning coefficient needs to be increased to accelerate the flow.

*Author contributions.* Charlotte M. Emery designed and carried out the experiments under the supervision of Sylvain Biancamaria and Aaron Boone, as part of her PhD project. Sylvain Biancamaria provided the SWOT-based observation mask used to generate the synthetic observations and Aaron Boone provided support for the use of the ISBA-CTRIP hydrologic model. Charlotte M. Emery prepared the manuscript with contribution from all co-authors.

5 *Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* This work was supported by the CNES, through a grant from Terre-Océan-Surfaces Continentales-Atmosphère (TOSCA) committee attributed to the project entitled "Towards an improved understanding of the global hydrological cycle using SWOT measurements". C. M. Emery received doctoral research support from CNES/région Midi-Pyrénées grant. C. M. Emery and C. H. David received support from the Jet Propulsion Laboratory, California Institute of Technology, under a contract with NASA; including grants from the

10 SWOT Science Team and the Terrestrial Hydrology Program.

## References


- Andreadis, K. M. and Schumann, G. J. P.: Estimating the impact of satellite observations on the predictability of large-scale hydraulic models, *Advances in Water Resources*, 73, 44–54, <https://doi.org/10.1016/j.advwatres.2014.06.006>, 2014.
- Andreadis, K. M., Clark, E. A., Lettenmaier, D. P., and Alsdorf, D. E.: Prospects for river discharge and depth estimation through assimilation  
5 of swath-altimetry into a raster-based hydrodynamics model, *Geophysical Research Letter*, 34, <https://doi.org/10.1029/2007GL029721>, 2007.
- Beighley, R. E., Eggert, K. G., Dunne, T., He, Y., Gummadi, V., and Verdin, K. L.: Simulating hydrologic and hydraulic processes throughout the Amazon basin, *Hydrological Processes*, 23, 1221–1235, <https://doi.org/10.1002/hyp.7252>, 2009.
- Beven, K. and Freer, J.: Equifinality, data assimilation, and uncertainty estimation in mechanistic modelling of complex environmental  
10 systems using GLUE methodology, *Journal of Hydrology*, 249, 11–29, [https://doi.org/10.1016/S0022-1694\(01\)00421-8](https://doi.org/10.1016/S0022-1694(01)00421-8), 2001.
- Beven, K. J.: *Rainfall-Runoff modelling*, chap. Chap 1: Down to basics: runoff processes and the modelling of processes, pp. 1–22, John Wiley and Sons, West Sussex, UK, 2012.
- Biancamaria, S., Bates, P., Boone, A., and Mognard, N.: Large-scale coupled hydrologic and hydraulic modelling of the Ob river in Siberia, *Journal of Hydrology*, 379, 136–150, <https://doi.org/10.1016/j.jhydrol.2009.09.054>, 2009.
- 15 Biancamaria, S., Durant, M., Andreadis, K. M., Bates, P. D., Boone, A., Mognard, N. M., Rodriguez, E., Alsdorf, D. E., Lettenmaier, D. P., and Clark, E. A.: Assimilation of virtual wide swath altimetry to improve Arctic river modeling, *Remote Sensing of Environment*, 115, 373–381, <https://doi.org/10.1016/j.rse.2010.09.008>, 2011.
- Biancamaria, S., Lettenmaier, D. P., and Pavelsky, T. M.: The SWOT mission and its capabilities for land hydrology, *Journal of Geophysics*, 37, <https://doi.org/10.1007/s10712-015-9346-y>, 2016.
- 20 Bierkens, M. F. P.: Global hydrology 2015: State, trends, and directions, *Water Resources Research*, 51, 4923–4947, <https://doi.org/10.1002/2015WR017173>, 2015.
- Birkett, C. M., Mertes, L. A. K., Dunne, T., Costa, M. H., and Jasinski, M. J.: Surface water dynamics in the Amazon basin: Application of satellite radar altimetry, *Journal of Geophysical Research*, 107, <https://doi.org/10.1029/2001JD000609>, 2002.
- Bishop, C. H., Etherton, B. J., and Majumbar, S. J.: Adaptive sampling with the ensemble transform Kalman filter. Part I: Theoretical  
25 aspects, *Monthly Weather Review*, 129, 420–436, [https://doi.org/10.1175/1520-0493\(2001\)129<0420:ASWTET>2.0.CO;2](https://doi.org/10.1175/1520-0493(2001)129<0420:ASWTET>2.0.CO;2), 2001.
- Boone, A., Calvet, J.-C., and Noilhan, J.: Inclusion of a Third Soil Layer in a Land Surface Scheme Using the Force-Restore Method, *Journal of Hydrometeorology*, 38, 1611–1630, [https://doi.org/10.1175/1520-0450\(1999\)038<1611:IOATSL>2.0.CO;2](https://doi.org/10.1175/1520-0450(1999)038<1611:IOATSL>2.0.CO;2), 1999.
- Brisset, P., Monnier, J., Garambois, P.-A., and Roux, H.: On the assimilation of altimetry data in 1D Saint-Venant river models, *Advances in Water Resources*, 119, 41–59, <https://doi.org/10.1016/J.advwatres.2018.06.004>, 2018.
- 30 Burgers, G., Leeuwen, P. J. V., and Evensen, G.: Analysis Scheme in the Ensemble Kalman Filter, *Monthly Weather Review*, 126, 1719–1724, [https://doi.org/10.1175/1520-0493\(1998\)126<1719:ASITEK>2.0.CO;2](https://doi.org/10.1175/1520-0493(1998)126<1719:ASITEK>2.0.CO;2), 1998.
- Clark, M. P., Rupp, D. E., Woods, R. A., Zheng, X., Ibbitt, R. P., Slater, A. G., Schmidt, J., and Uddstrom, M. J.: Hydrological data assimilation with the ensemble Kalman filter: Use of streamflow observations to update states in a distributed hydrological model, *Advances in Water Resources*, 31, 1309–1324, <https://doi.org/10.1016/j.advwatres.2008.06.005>, 2008.
- 35 Cretaux, J.-F., Calmant, S., Romanoski, V., Shabunin, A., Lyard, F., Berge-Nguyen, M., Cazenave, A., Hernandez, F., and Perosanz, F.: An absolute calibration site for radar altimeters in the continental domain: Lake Issykkul in Central Asia, *Journal Of Geodesy*, 83, 723–735, <https://doi.org/10.1007/s00190-008-0289-7>, 2009.

- Decharme, B., Alkama, R., Douville, H., Becker, M., and Cazenave, A.: Global Evaluation of the ISBA-TRIP Continental Hydrological System. Part II: Uncertainties in River Routing Simulation Related to Flow Velocity and Groundwater Storage, *J. Hydrometeor.*, 11, 601–617, <https://doi.org/10.1175/2010JHM1212.1>, 2010.
- Decharme, B., Alkama, R., Papa, F., Faroux, S., Douville, H., and Prigent, C.: Global off-line evaluation of the ISBA-TRIP flood model, *Climate Dynamics*, 38, 1389–1412, <https://doi.org/10.1007/s00382-011-1054-9>, 2012.
- Decharme, B., C., D., Minvielle, M., J., C., J.-P., V., D., A. A. S.-M., R., S., S., S., and A., V.: Recent changes in the ISBA-CTrip land surface system for use in CNRM-CM6 climate model and global off-line hydrological applications., *Journal of Advances in Modeling Earth Systems*, 11, <https://doi.org/10.1029/2018MS001545>, 2019.
- Deng, C., Liu, P., Guo, S., Li, Z., and Wang, D.: Identification of hydrological model parameter variation using ensemble Kalman filter, *Hydrology and Earth System Sciences*, 20, 4949–4961, <https://doi.org/10.5194/hess-20-4949-2016>, 2016.
- Doll, P., Douville, H., Güntner, A., Schmied, H. M., and Wada, Y.: Modelling Freshwater Resources at the Global Scale: Challenges and Propects, *Surveys in Geophysics*, 37, <https://doi.org/10.1007/s10712-015-9343-1>, 2015.
- Durand, M., Andreadis, K., Alsdorf, D., Lettenmaier, D., Moller, D., and Wilson, M.: Estimation of bathymetric depth and slope from data assimilation of swath altimetry into a hydrodynamic model, *Geophysical Research Letters*, 35, <https://doi.org/10.1029/2008GL034150>, 2008.
- Emery, C. M., Biancamaria, S., Boone, A., Garambois, P.-A., Ricci, S., Rochoux, M. C., and Decharme, B.: Temporal variance-based sensitivity analysis of the river routing component of the large scale hydrological model ISBA-TRIP: Application on the Amazon Basin, *Journal of Hydrometeorology*, 17, 3007–3027, <https://doi.org/10.1175/JHM-D-16-0050.1>, 2016.
- Emery, C. M., Paris, A., Biancamaria, S., Boone, A., Calmant, S., Garambois, P.-A., and Silva, J. S. D.: Large scale hydrological model river storage and discharge correction using satellite altimetry-based discharge product, *Hydrology and Earth System Sciences*, 22, 2135–2162, <https://doi.org/10.5194/hess-22-2135-2018>, 2018.
- Esteban Fernandez, D.: SWOT Project, Mission performance and error budget, Tech. rep., Jet Propulsion Laboratory, 2017.
- Evensen, G.: Sequential data assimilation with a nonlinear quasi-geostropic model using Monte Carlo methods to forecast error statistics, *Journal of Geophysical Research*, 99, 10 143–10 162, <https://doi.org/10.1029/94JC00572>, 1994.
- Evensen, G.: Advanced data assimilation for strongly nonlinear dynamics, *Monthly Weather Review*, 125, 1342–1354, [https://doi.org/10.1175/1520-0493\(1997\)125<1342:ADAFSN>2.0.CO;2](https://doi.org/10.1175/1520-0493(1997)125<1342:ADAFSN>2.0.CO;2), 1997.
- Evensen, G.: The Ensemble Kalman Filter: theoretical formulation and practical implementation, *Ocean Dynamics*, 53, 343–367, <https://doi.org/10.1007/s10236-003-0036-9>, 2003.
- Evensen, G.: Sampling strategies and square root analysis schemes for the EnKF, *Ocean Dynamics*, 54, 539–560, <https://doi.org/10.1007/s10236-004-0099-2>, 2004.
- Evensen, G. and Leeuwen, P. V.: An ensemble kalman smoother for nonlinear dynamics, *Monthly Weather Review*, 128, 1852–1867, [https://doi.org/10.1175/1520-0493\(2000\)128<1852:AEKSFN>2.0.CO;2](https://doi.org/10.1175/1520-0493(2000)128<1852:AEKSFN>2.0.CO;2), 2000.
- Fjórtoft, R., Gaudin, J.-M., Pourthie, N., Lalaurie, J.-C., Mallet, A., Nouvel, J.-F., Martinot-Lagarde, J., Oriot, H., Borderies, P., Ruiz, C., and Daniel, S.: KaRIn on SWOT: Characteristics of near-nadir Ka-band interferometric SAR imagery, *IEEE Transactions on Geoscience and Remote Sensing*, 52, 2172–2185, <https://doi.org/10.1109/TGRS.2013.2258402>, 2014.
- Guillet, O., Weaver, A., Vasseur, X., Michel, M., Gratton, S., and Gürol, S.: Modelling spatially correlated observation errors in variational data assimilation using a diffusion operator on an unstructured mesh, *Quarterly Journal of the Royal Meteorological Society*, in revision, 2018.

- Gupta, H. V., Sorooshian, S., and Yapo, P. O.: Toward improved calibration of hydrological models: multiple and noncommensurable measures of information, *Water Resources Research*, 34, 751–763, <https://doi.org/10.1029/97WR03495>, 1998.
- Hafliker, V., Martin, E., Boone, A., Ricci, S., and Biancamaria, S.: Assimilation of synthetic SWOT river depths in a regional hydrometeorological model, *Water*, 11, <https://doi.org/10.3390/w11010078>, 2019.
- 5 Hunt, B., Kalnay, E., Kostelich, E. J., Ott, E., Patil, D. T., Sauer, T., Szunyogh, I., Yorke, J. A., and Zimin, A. V.: Four-dimensional ensemble Kalman filtering, *Tellus*, 56, 273–277, <https://doi.org/10.1111/j.1600-0870.2004.00066.x>, 2004.
- Hunt, B. R., Kostelich, E. J., and Szunyogh, I.: Efficient data assimilation for spatiotemporal chaos: A local ensemble transform Kalman filter, *Physica D*, 230, 112–126, <https://doi.org/10.1016/j.physd.2006.11.008>, 2007.
- International Association of Hydrological Sciences Ad Hoc Group on Global Water Sets, Vörösmarty, C., Askew, A., Grabs, W., Barry, R. G., Birkett, C., Döll, P., Goodison, B., Hall, A., Jenne, R., Kitaev, L., Landwehr, J., Keeler, M., Leavesley, G., Schaake, J., Strzepek, K., Sundarvel, S. S., Takeuchi, K., and Webster, F.: Global water data: a newly endangered species, *Eos. Trans. AGU*, 82, 54–58, <https://doi.org/10.1029/01EO00031>, 2001.
- 10 Kurtz, W., Hendricks-Frassen, H.-J., and Vereecken, H.: Identification of time-variant river bed properties with Ensemble Kalman Filter, *Water Resources Research*, 48, <https://doi.org/10.1029/2011WR011743>, 2012.
- 15 Leeuwen, P. V. and Evensen, G.: Data assimilation and inverse methods in terms of a probabilistic formulation, *Monthly Weather Review*, 124, 2898–2913, [https://doi.org/10.1175/1520-0493\(1996\)124<2898:DAAIMI>2.0.CO;2](https://doi.org/10.1175/1520-0493(1996)124<2898:DAAIMI>2.0.CO;2), 1996.
- Liu, Y. and Gupta, H. V.: Uncertainty in hydrological modeling: Towards an integrated data assimilation framework, *Water Resources Research*, 43, <https://doi.org/10.1029/2006WR005756>, 2007.
- Liu, Y., Weerts, A. H., Clark, M., Hendricks Franssen, H.-J., Kumar, S., Moradkhani, H., Seo, D.-J., Schwanenberg, D., Smith, P., van Dijk, A. I. J. M. N., van Velzen, N., He, M., Lee, H., Noh, S. J., Rakovec, O., and Restrepo, P.: Advancing data assimilation in operational hydrologic forecasting: progress, challenges, and emerging opportunities, *Hydrology and Earth System Sciences*, 16, 3863–3887, <https://doi.org/10.5194/hess-16-3863-2012>, 2012.
- 20 Maidment, D. R.: *Handbook of hydrology*, McGraw Hill Professional, 1993.
- Manning, R.: On the flow of water in open channels and pipes, *Institution of Civil Engineers of Ireland*, 20, 161–207, 1891.
- 25 Meade, R., Rayol, J., Conceição, S. D., and Natividade, J.: Backwater Effects in the Amazon River Basin of Brazil, *Environmental Geology and Water Sci.*, 18, 105–114, <https://doi.org/10.1007/BF01704664>, 1991.
- Melsen, L., Teuling, A., Torfs, P., Zappa, M., Mizukami, N., Clark, M., and Uijlenhoet, R.: Representation of spatial and temporal variability in large-domain hydrological models: case study for a mesoscale pre-Alpine basin, *Hydrology and Earth System Sciences*, 20, 2207–2226, <https://doi.org/10.5194/hess-20-2207-2016>, 2016.
- 30 Mersel, M. K., Smith, L. C., Andreadis, K. M., and Durand, M. T.: Estimation of river depth from remotely-sensed hydraulic relationship, *Water Resources Research*, 49, 3165–3179, <https://doi.org/10.1002/wrcr.20176>, 2013.
- Michailovsky, C. I. and Bauer-Gottwein, P.: Operational reservoir inflow forecasting with radar altimetry: the Zambezi case study, *Hydrology and Earth System Sciences*, 18, 997–1007, <https://doi.org/10.5194/hess-18-997-2014>, 2014.
- Michailovsky, C. I., Milzow, C., and Bauer-Gottwein, P.: Assimilation of radar altimetry to a routing model of the Brahmaputra river, *Water Resources Research*, 49, 4807–4816, <https://doi.org/10.1002/wrcr.20345>, 2013.
- 35 Molinier, M., Guyot, J.-L., Orstom, B., Guimarães, V., de Oliveira, E., and Dnaee, B.: Hydrologie du bassin de l’Amazone, in: *Grands Bassins Fluviaux Périalantiques*, edited by Editions, O., pp. 335–345, PEGI-INSA-CNRS-ORSTOM, Paris, 1993.


- Montzka, C., Moradkhani, H., Weihermüller, L., Hendricks-Franssen, H.-J., Canty, M., and Vereecken, H.: Hydraulic parameter estimation by remotely-sensed top soil moisture observations with the particle filter, *Journal of Hydrology*, 399, 410–421, <https://doi.org/10.1016/j.jhydrol.2011.01.020>, 2011.
- Moradkhani, H., Hsu, K.-L., Gupta, H., and Sorooshian, S.: Uncertainty assessment of hydrologic model states and parameters: Sequential data assimilation using particle filter, *Water Resources Research*, 41, <https://doi.org/10.1029/2004WR003604>, 2005.
- Munier, S., Polebistki, A., Brown, C., Belaud, G., and Lettenmaier, D. P.: SWOT data assimilation for operational reservoir management on the upper Niger river basin, *Water Resources Research*, 51, 554–575, <https://doi.org/10.1002/2014WR016157>, 2015.
- Noilhan, J. and Planton, S.: A simple parameterization of land surface processes for meteorological models, *Mon. Wea. Rev.*, 117, 536–549, [https://doi.org/10.1175/1520-0493\(1989\)117<0536:ASPOLS>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<0536:ASPOLS>2.0.CO;2), 1989.
- 10 Oki, T. and Sud, Y. C.: Design of Total Integrating Pathways (TRIP)-A Global River Channel Network, *Earth Interactions*, 2, 1–36, [https://doi.org/10.1175/1087-3562\(1998\)002<0001:DOTRIP>2.3.CO;2](https://doi.org/10.1175/1087-3562(1998)002<0001:DOTRIP>2.3.CO;2), 1998.
- Ott, E., Hunt, B. R., Szunyogh, I., Kostelich, A. V. Z. A. J., Corazza, M., Kalnay, E., Patil, D. J., and Yorke, J. A.: A local ensemble Kalman filter for atmospheric data assimilation, *Tellus A*, 56, 415–428, <https://doi.org/10.1111/j.1600-0870.2004.00076.x>, 2004.
- Oubanas, H., Gejadze, I., Malaterre, P.-O., Durand, M., Wei, R., Frasson, R. P. M., and Domeneghetti, A.: Discharge estimation in ungauged basins through variational data assimilation: the potential of the SWOT mission, *Water Resources Research*, 54, <https://doi.org/10.1002/2017WR021735>, 2018.
- 15 Paiva, R. C. D., Buarque, D. C., Collischonn, W., Bonnet, M.-P., Frappart, F., Calmant, S., and Mendes, C. A. B.: Large scale hydrological and hydrodynamic modeling of the Amazon River basin, *Water Resources Research*, 49, 1226–1243, <https://doi.org/10.1002/wrcr.20067>, 2013.
- 20 Panzeri, M., Riva, M., Guadagnini, A., and Neuman, S. P.: Data assimilation and parameter estimation via ensemble kalman filter coupled with stochastic moment equations of transient groundwater flow, *Water Resources Research*, 49, 1334–1344, <https://doi.org/10.1002/wrcr.20113>, 2013.
- Pathiraja, S., Marshall, L., Sharma, A., and Moradkhani, H.: Detecting non-stationar hydrologic model parameters in a paired catchment system using data assimilation, *Advances in Water Resources*, 94, 103–119, <https://doi.org/10.1016/j.advwatres.2016.04.021>, 2016.
- 25 Pedinotti, V., Boone, A., Ricci, S., Biancamaria, S., and Mognard, N.: Assimilation of satellite data to optimize large-scale hydrological model parameters: a case study for the SWOT mission, *Hydrology and Earth System Science*, 18, 4485–4507, <https://doi.org/10.5194/hess-18-4485-2014>, 2014.
- Rakovec, O., Weerts, A. H., Sumihar, J., and Uijlenhoet, R.: Operational aspects of asynchronous filtering for flood forecasting, *Hydrology and Earth System Sciences*, 19, 2911–2924, <https://doi.org/10.5194/hess-19-2911-2015>, 2015.
- 30 Renard, B., Kavetski, D., Kuczera, G., Thyer, M., and Franks, S. W.: Understanding predictive uncertainty in hydrologic modeling: the challenge of identifying input and structural errors, *Water Resources Research*, 46, <https://doi.org/10.1029/2009WR008328>, 2010.
- Rodell, M., Beaudoin, H. K., L’Ecuyer, T. S., Olson, W. S., Famiglietti, J. S., Houser, P. R., Adler, R., Bosilovich, M. G., Clayson, C. A., Chambers, D., Clark, E., Fetzer, E. J., Gao, X., Gu, G., Hilburn, K., Huffman, G. H., Lettenmaier, D. P., Liu, W. T., Robertson, F. R., Schlosser, C. A., Sheffield, J., and Wood, E. F.: The observed state of the water cycle in the early twenty-first century, *Journal of Climate*, 28, 8289–8318, <https://doi.org/10.1175/JCLI-D-14-00555.1>, 2015.
- 35 Ruiz, J. J., Pulido, M., and Miyoshi, T.: Estimating model parameters with ensemble-based data assimilation, *Journal of the Meteorological Society of Japan*, 91, 79–99, <https://doi.org/10.2151/jmsj.2013-201>, 2013.

- Sakov, P., Evensen, G., and Bertino, L.: Asynchronous data assimilation with the EnKF, *Tellus*, 62, 24–29, <https://doi.org/10.1111/j.1600-0870.2009.00417.x>, 2010.
- Sanoo, A. K., Pan, M., Troy, T. J., Vinukollu, R. K., Sheffield, J., and Wood, E. F.: Reconciling the global terrestrial water budget using satellite remote sensing, *Remote Sensing of Environment*, 115, 1850–1865, <https://doi.org/10.1016/j.rse.2011.03.009>, 2011.
- 5 Shi, Y., Davis, K. J., Zhang, F., Duffy, C. J., and Yu, X.: Parameter estimation of physically-based land surface model hydrologic model using an ensemble Kalman filter: a multivariate real-data experiment, *Advances in Water Resources*, 83, 421–427, <https://doi.org/10.1016/j.advwatres.2015.06.009>, 2015.
- Silva, J. S. D., Calmant, S., Seyler, F., Filho, O. C. R., Cochonneau, G., and Mansur, W. J.: Water levels in the Amazon basin derived from the ERS 2 and ENVISAT radar altimetry missions, *Remote Sensing of Environment*, 114, 2160–2181, <https://doi.org/10.1016/j.rse.2010.04.020>, 2010.
- 10 Sood, A. and Smakhtin, V.: Global hydrological models: a review, *Hydrological Sciences Journal*, 60, 549–565, <https://doi.org/10.1080/02626667.2014.950580>, 2015.
- Talagrand, O. and Courtier, P.: Variational assimilation of meteorological observations with the adjoint vorticity equation, Part 1: Theory, *Quarterly Journal of the Royal Meteorological Society*, 113, 1311–1328, <https://doi.org/10.1002/qj.49711347812>, 1987.
- 15 Vinukollu, R. K., Meynadier, R., Sheffield, J., and Wood, E. F.: Multi-model, multi-sensor estimates of global evapotranspiration : climatology, uncertainties and trends, *Hydrological Processes*, 25, 3993–4010, <https://doi.org/10.1002/hyp.8393>, 2011.
- Vrugt, J. A., ter Braak, C. J. F., Diks, C. G. H., and Shoups, G.: Hydrological data assimilation using particle Markov chain Monte Carlo simulation: Theory, concepts and applications, *Advances in Water Resources*, 51, 457–478, <https://doi.org/10.1016/j.advwatres.2012.04.002>, 2012.
- 20 Wisser, D., Feketa, B., Vörösmarty, C. J., and Schumann, A. H.: Reconstructing 20th century global hydrography: a contribution to the Global Terrestrial Network- Hydrology (GTN-H), *Hydrology and Earth System Sciences*, 14, 1–24, <https://doi.org/10.5194/hess-14-1-2010>, 2010.
- Yoon, Y., Durand, M., Merry, C. J., Clark, E. A., Andreadis, K. M., and Alsdorf, D. E.: Estimating river bathymetry from data assimilation of synthetic SWOT measurements, *Journal of Hydrology*, 464–465, 363–375, <https://doi.org/10.1016/j.jhydrol.2012.07.028>, 2012.




**Figure 1.** (a) The ISBA-CTrip system for a given grid cell. ISBA surface runoff ( $Q_{\text{ISBA,sur}}$ ) flows into the river/surface reservoir  $S$ , ISBA gravitational drainage ( $Q_{\text{ISBA,sub}}$ ) feeds groundwater reservoir  $G$ . The surface water is transferred from one cell to another following the TRIP river routing network. (b) Hydro-geomorphological areas of the Amazon basin from Emery et al. (2016) with the gauge of Óbidos located by the white circle at the entry of the zone 3.







**Figure 2.** SWOT swaths at ISBA-CTrip resolution over the Amazon basin.




**Figure 3.** Data assimilation framework over the assimilation cycle  $[k - 1, k]$  including (1) a forecast step to integrate the ensemble of ISBA-CTRIP simulations, each member having a different spatially-distributed Manning coefficient, (2) an analysis step to correct this ensemble of Manning coefficients using synthetic observations through the Kalman filter equation and (3) re-run the ISBA-CTRIP model with these analysis estimates to obtain the updated model states.




**Figure 4.** Illustration of a bias case between the water elevation as observed by SWOT ( $H_{\text{alti}}^{\text{SWOT}}$ ) and the one simulated by CTRIP ( $H_{\text{alti}}^{\text{TRIP}}$ ) because of a bias between model and true river beds.




**Figure 5.** Results for the spinup sensitivity test. Each test run is represented along the  $x$ -axis and referenced by its number of spinup windows. The  $y$ -axis displays the number of windows during which the relative difference between the true run and the openloop run in which the Manning spatial distribution is modified is above the chosen threshold. These statistics are obtained for the discharge (a,c) and the water depth (b,d) and evaluated over the entire basin (a,b) and at the downstream station of Óbidos (c,d). Note that the vertical dashed line corresponds to the minimum model spinup period retained in this study.




**Figure 6.** (Top) Relative error (to the truth) and (Bottom) dispersion of the analysis control ensemble (averaged over all control variables) for the sensitivity tests to (a-b) the data assimilation starting date, (c-d) the ensemble size  $n_e$ , (e-f) the background error standard deviation  $\sigma^b$ . For each test, a set of one-cycle-long data assimilation experiments is run.



**Figure 7.** Control variables assimilation results for the PE1 experiment: evolution of the ensemble-averaged analysis control variable (red line) for each zone (one zone per subplot) with respect to the assimilation cycle and compared to the corresponding true value (black line) and the openloop value (blue line).




**Figure 8.** Water depths assimilation results for the PE1 experiment: daily evolution of the ensemble-averaged analysis water depth (red lines) compared to the true water depths (black lines) and the openloop water depths (blue line). For each zone (one per subplot), the median (full line), the first decile (dotted line) and the ninth decile (dashed line) of water depth ensemble over all grid cells in the zone are represented.



**Figure 9.** Control variables assimilation results for the PE2 and PE3 experiments: evolution of the ensemble-averaged analysis control variable for the PE2 experiment (orange line) and the PE3 experiment (purple line) for each zone (one zone per subplot) with respect to the assimilation cycle and compared to the corresponding true value (black line) and the openloop value (blue line).






**Figure 10.** Water depths assimilation results for the PE3 experiment: daily evolution of the ensemble-averaged analysis water depth (red lines) compared to the true water depths (black lines) and the openloop water depths (blue line). For each zone (one per subplot), the median (full line), the first decile (dotted line) and the ninth decile (dashed line) of water depth ensemble over all grid cells in the zone are represented.

Parameter	Nb run	Range
Spinup	18	From 0 window to 17 windows of 21 days
Starting date	17	Starting January 1st, 2008 and on, every 21 days
$n_e$	9	[ 10 20 30 40 50 75 100 150 200 ]
$\sigma^b$	13	[ 0.01 0.02 0.05 0.1 0.2 0.3 0.4 0.5 0.75 1.0 1.2 1.5 1.75 ]


**Table 1.** Tested data assimilation parameters in the sensitivity tests.

Simulation name	Observation variables	Bathymetry bias
PE1	Water depths	No
PE2	Water depth anomalies	No
PE3	Water depth anomalies	Yes

**Table 2.** List of data assimilation experiments. All experiments are run over approximately one year (17 cycles of 21 days) starting on January 1st, 2008. The ensemble size is  $n_e = 25$ , the observation error standard deviation is  $\sigma^o = 0.1$  m and the initial control variable error standard deviation is  $\sigma^b = 0.3$ .



**Figure A1.** (Top) Relative error (to the truth) and (Bottom) dispersion of the analysis control ensemble for each zones for the sensitivity tests to (a-b) the data assimilation starting date, (c-d) the ensemble size  $n_e$ , (e-f) the background error standard deviation  $\sigma^b$ . For each test, a set of one-cycle-long data assimilation experiments is run. (Top only) The relative error in zone 1 (dark blue line), zone 2 (orange line), zone 3 (yellow line), zone 4 (purple line), zone 5 (green line), zone 6 (light blue line), zone 7 (burgundy red line), zone 8 (pink line) and zone 9 (gray line) are compared to the basin-averaged openloop relative error (black line).



**Figure A2.** Evolution of the EnKF innovations (" $\mathbf{y}_k^{o,[l]} - \mathcal{H}_k(\mathbf{x}_k^{b,[l]})$ ") term in Eq. 13) with respect to the assimilation cycle for PE1 (red line), PE2 (orange line) and PE3 (purple line). For each zone (one zone per subplot), the displayed innovation is the averaged of all the innovations in the corresponding zones.

Run	Starting date	Manning distr. change	Spinup length (in wdws of 21 days)
Reference	Dec 16, 2006	-	18 (=378 days)
1	Dec 16, 2006	Jan 09, 2007	17 (=357 days)
2	Dec 16, 2006	Jan 30, 2007	16 (=336 days)
3	Dec 16, 2006	Feb 20, 2007	15 (=315 days)
4	Dec 16, 2006	Mar 13, 2007	14 (=294 days)
5	Dec 16, 2006	Apr 03, 2007	13 (=273 days)
6	Dec 16, 2006	Apr 24, 2007	12 (=252 days)
7	Dec 16, 2006	May 15, 2007	11 (=231 days)
8	Dec 16, 2006	Jun 05, 2007	10 (=210 days)
9	Dec 16, 2006	Jun 26, 2007	9 (=189 days)
10	Dec 16, 2006	Jul 17, 2007	8 (=168 days)
11	Dec 16, 2006	Aug 07, 2007	7 (=147 days)
12	Dec 16, 2006	Aug 28, 2007	6 (=126 days)
13	Dec 16, 2006	Sept 18, 2007	5 (=105 days)
14	Dec 16, 2006	Oct 09, 2007	4 (=84 days)
15	Dec 16, 2006	Oct 30, 2007	3 (=63 days)
16	Dec 16, 2006	Nov 20, 2007	2 (=42 days)
17	Dec 16, 2006	Dec 11, 2007	1 (=21 days)
18	Dec 16, 2006	Jan 01, 2008	0 (=0 days)

**Table A1.** Spinup sensitivity test set up: each run consists in an approximately-two-year-long ISBA-CTRIP run starting on Dec 16th, 2006 (column 2) and ending on Dec 22nd, 2008. After a given number of 21-day windows during the first year (column 3), the Manning distribution is changed to replicate an assimilation update step while the reference run (row 2) used the same Manning for the entire run. The period between the instant when the Manning distribution is changed and the beginning of the second year of simulation corresponds to the spinup period (column 4). The simulated water depth/discharge during the second year of run are then compared to the reference run in order to evaluate the impact of the spin up.

Zones	1	2	3	4	5	6	7	8	9
Openloop	7.57	28.84	30.28	33.51	33.41	37.15	38.83	11.48	5.23
PE1	0.48	0.27	0.37	0.34	0.53	0.88	0.76	1.11	1.58
PE2	1.34	0.30	1.12	0.90	0.31	1.49	0.91	0.06	1.40
PE3	2.23	3.52	8.31	1.48	0.57	1.39	1.56	1.67	1.46

**Table A2.** Zone-averaged RMSEn for the openloop water depths (row 2) and the ensemble-averaged analysis water depths (rows 3-5) compared to the true water depths.

Zones	1	2	3	4	5	6	7	8	9
Openloop	2.73	4.65	6.46	3.89	5.57	4.63	9.46	3.50	3.26
PE1	0.35	0.38	0.34	0.15	0.25	0.53	0.29	0.30	0.59
PE2	0.74	0.14	0.22	0.08	0.04	0.34	0.85	0.12	0.45
PE3	1.20	4.52	7.52	0.15	0.30	0.35	2.21	1.65	2.14

**Table A3.** Zone-averaged RMSEn for the openloop discharges (row 2) and the ensemble-averaged analysis discharges (rows 3-5) compared to the true discharges.