

RESPONSE TO COMMENTS from 2nd Referee Report

Review of the revised manuscript entitled “Assessment of near 0°C temperature and precipitation characteristics across Canada” by Mekis et al. 2019. The authors have addressed most of my comments in the revised manuscript. However, there are still a few minor issues in the revised manuscript that should be addressed.

1. Lines 120-124. The country is divided into 6 broad regions. Does this study show any results for these 6 broad regions?

Line in question: “Canada is a vast country with many different climatic regions (Gullett et al., 1992) and ecozones (Zhang et al., 2001). For easier evaluation of the results, similar to Vincent et al. (2018), the country was divided into six broad regions based on the 13 provincial/territorial boundaries (Fig. 2). In addition to the Canada-wide analysis, nine stations were chosen for further analysis (Fig. 2). These stations, having high quality consistent observing practices, represent contrasting climatic conditions across the country.”

Response: No, we did not compute the results for the six broad regions. All the results are shown on the individual station level. We used these names for grouping the results and patterns for easier evaluation of the results. Throughout the paper, these names had been used extensively.

New/Revised Text: There is no change in the text.

2. Lines 167-169. The information regarding the average annual surface temperature lines is out of context. Is this provided on fig. 2?

Line in question: “The average annual surface temperature contour lines was computed from the 1981-2010 climate normal period (http://climate.weather.gc.ca/climate_normals/index_e.html, ECCC, 2019c). Kriging with a linear Variogram model and a grid spacing of 50 km was applied to create the interpolated surface temperature map from the 1619 stations for Canada.”

Response: Since the contour lines computed from the 1981-2010 climate normals are new results (never been used or published); the brief description of the applied method is appropriate to be in the method section. It is true that these lines first appeared on Fig. 2, but it was subsequently used in Figures 3-5 and multiple places in the analysis. Additional text was added (bold) to better link it into the chapter.

New/Revised Text: **Given the significance of near 0°C isotherms location in this study, the -2, 0, +2 °C average annual surface temperature contour lines were computed from the 1981-2010 climate normal period (http://climate.weather.gc.ca/climate_normals/index_e.html, ECCC, 2019c) to be used later in the analysis.** Kriging with a linear Variogram model and a grid spacing of 50 km was applied to create the interpolated surface temperature map from the 1619 stations for Canada.

3. Table 2. The caption is incomplete. Average annual frequencies of the number of days, hours, events and maximum duration when $-2^{\circ}\text{C} \leq T \leq 2^{\circ}\text{C}$ (in brackets) with the 31 year...

Line in question: “Table 2: Average annual frequencies of the selected near 0°C indicators along (in brackets) with the 31 year trend values based on hourly surface temperature over the 1981-2011 period (minimum 90% of data). The numbers in bold indicate significant changes at 5% level.”

Response: As requested, addition text is added to Table 2 caption (bold).

New/Revised Text: Table 2: Average annual frequencies of the **number of the days, hours, events and maximum duration when $-2^{\circ}\text{C} \leq T \leq 2^{\circ}\text{C}$** along (in brackets) with the 31 year trend values based on hourly surface temperature over the 1981-2011 period (minimum 90% of data). The numbers in bold indicate significant changes at 5% level.

4. Table 3. Should not provide ‘(no)’ for no significant change but instead mention in the caption that no brackets means no significant change.

Response: Thank you. The “(no)” is deleted from the table and a sentence is added to its title. This change indeed made the table cleaner.

New/Revised Text: Sentence added to the title (shown in bold): “No bracket means no significant change.”

Table 3 before:

	Whitehorse	Vancouver	Calgary	Cambridge Bay	Winnipeg	Churchill	Toronto	Montreal	St. John's	Units
Rain	16.3 (no)	25.2 (no)	12.7 (no)	29.9 (no)	23.1 (no)	60.7 (no)	49.9 (no)	56.3 (no)	83.9 (no)	hours
Rain Showers	4.2 (>)	9.6 (no)	4.6 (no)	15.1 (>)	4.1 (no)	9.1 (<)	6.8 (no)	5.7 (no)	34.5 (<)	hours
Drizzle	2.0 (no)	3.7 (no)	6.5 (no)	30.2 (no)	15.1 (no)	41.8 (<)	21.4 (<)	16.2 (no)	132.2 (no)	hours
Freezing Rain	n/a	2.2 (rare)	n/a	4.9 (no)	3.6 (no)	10.0 (no)	8.3 (no)	13.4 (no)	41.0 (no)	hours
Freezing Drizzle	n/a	n/a	4.2 (no)	10.9 (no)	5.9 (no)	15.3 (<)	5.2 (no)	6.3 (<)	85.7 (<)	hours
Snow	111.8 (no)	41.2 (no)	96.2 (no)	91.9 (no)	79.6 (no)	163.3 (no)	115.1 (no)	152.0 (no)	173.2 (>)	hours
Snow Grains	n/a	1.4 (rare)	1.8 (no)	8.6 (no)	2.8 (no)	5.7 (no)	6.0 (no)	2.4 (no)	12.8 (<)	hours
Ice Crystal	n/a	n/a	0.0 (rare)	n/a	n/a	0.1 (no)	n/a	n/a	n/a	hours
Ice Pellets	n/a	n/a	2.4 (rare)	4.5 (<)	3.6 (no)	3.9 (<)	7.6 (no)	7.9 (no)	16.8 (no)	hours
Ice Pellet Showers	n/a	n/a	n/a	n/a	0.8 (rare)	1.9 (rare)	0.9 (no)	0.9 (no)	1.2 (no)	hours
Snow Showers	29.1 (no)	16.7 (no)	36.5 (<)	68.6 (>)	11.9 (no)	44.4 (<)	53.3 (no)	40.2 (no)	174.2 (<)	hours
Snow Pellets	0.9 (no)	n/a	2.5 (no)	3.4 (no)	1.5 (no)	2.6 (no)	2.7 (no)	1.5 (no)	3.0 (no)	hours
Any of the 12 types	154.7 (no)	85.6 (no)	151.6 (no)	250.5 (no)	136.3 (no)	323.2 (<)	252.6 (no)	276.2 (no)	686.1 (no)	hours

After: Table 3: Average number of hours with the 12 different precipitation type occurrences along in bracket with the indication of changes over the 1981-2011 period. **No bracket means no significant change.** “>” indicates statistically significant increasing change; “<” indicates statistically significant decreasing change; “no” indicates no significant change; “rare” indicates that the given precipitation type rarely occurs and the change is not computed; “n/a” indicates that there is not enough data to compute the change. The statistical significance of the trends were assessed at 5% level.

	Whitehorse	Vancouver	Calgary	Cambridge Bay	Winnipeg	Churchill	Toronto	Montreal	St. John's	Units
Rain	16.3	25.2	12.7	29.9	23.1	60.7	49.9	56.3	83.9	hours
Rain Showers	4.2 (>)	9.6	4.6	15.1 (>)	4.1	9.1 (<)	6.8	5.7	34.5 (<)	hours
Drizzle	2	3.7	6.5	30.2	15.1	41.8 (<)	21.4 (<)	16.2	132.2	hours
Freezing Rain	n/a	2.2 (rare)	n/a	4.9	3.6	10	8.3	13.4	41	hours
Freezing Drizzle	n/a	n/a	4.2	10.9	5.9	15.3 (<)	5.2	6.3 (<)	85.7 (<)	hours
Snow	111.8	41.2	96.2	91.9	79.6	163.3	115.1	152	173.2 (>)	hours
Snow Grains	n/a	1.4 (rare)	1.8	8.6	2.8	5.7	6	2.4	12.8 (<)	hours
Ice Crystal	n/a	n/a	0.0 (rare)	n/a	n/a	0.1	n/a	n/a	n/a	hours
Ice Pellets	n/a	n/a	2.4 (rare)	4.5 (<)	3.6	3.9 (<)	7.6	7.9	16.8	hours
Ice Pellet Showers	n/a	n/a	n/a	n/a	0.8 (rare)	1.9 (rare)	0.9	0.9	1.2	hours
Snow Showers	29.1	16.7	36.5 (<)	68.6 (>)	11.9	44.4 (<)	53.3	40.2	174.2 (<)	hours
Snow Pellets	0.9	n/a	2.5	3.4	1.5	2.6	2.7	1.5	3	hours
Any of the 12 types	154.7	85.6	151.6	250.5	136.3	323.2 (<)	252.6	276.2	686.1	hours

5. Table 4. What are “Onset” and “Cessation” trends? Are these expressions necessary? They need to be defined.

Lines in question: “Further insight can be gained by examining shorter time scales. For example, Fig. 10 shows annual cycles of near 0°C conditions and associated precipitation type occurrences at four stations (Vancouver, Cambridge Bay, Toronto and St. John’s) chosen to illustrate a range of variation. Near 0°C conditions do not occur during summer at all southern stations (Figs. 10a, c, d) but they can occur in any other month. In contrast, these conditions only occur in summer at Cambridge Bay (Fig. 10b). The most frequent occurrence can be seen at St John’s (Fig. 10d). Furthermore, the occurrence of any precipitation

type tends to take place towards the ‘winter’ side at southern stations, although there are exceptions. Such occurrences on the ‘winter’ side are probably linked with storms passing over the stations with associated precipitation, whereas occurrences on the ‘summer’ side normally just reflect the diurnal cycle.

It would be expected that, with overall warming (Bush and Lemmen, 2019), the onset for near 0°C would occur later in autumn and earlier in spring. As shown in Table 3 and Fig. 10c, both of these trends are significant at only one location, Toronto. Whitehorse shows significant earlier spring cessation (Table 4) and St. John’s shows significant later autumn onset (Fig. 10d and Table 4). Cambridge Bay experiences near 0°C conditions in every ‘warm season’ month (Fig. 10b and Table 4) so the onset of near 0°C in the spring and its cessation in the autumn were considered; neither showed statistically significant trends. No analysis was conducted for Churchill because near 0°C conditions occurred in every month (Table 4).”

Response: There had been no specific comments by reviewers questioning the use of ‘onset’ and ‘cessation’. However, Reviewers 2 and 4 comments asked for clarification about this section of the article. We thought we had addressed these but obviously not as well as necessary. As suggested, we now clearly define these two terms.

We feel that the analysis using these two terms is critical. The article is concerned with patterns of near 0°C conditions including their occurrence over the annual cycle. Onset and cessation are defined to allow us to quantify when near 0°C conditions started and stopped over the annual cycle. For most of the 9 selected stations, onset refers to the first occurrence of near 0°C conditions in the autumn and cessation refers to the last occurrence of these conditions in the spring. For Cambridge Bay, onset refers to the first occurrence of near 0°C conditions in the spring and cessation refers to the last occurrence of these conditions in the autumn.

New/Revised Text: This issue is referred to in 3 paragraphs as follows.

Paragraph 1: No change in the paragraph starting with “Further insight...”

Paragraph 2: New Paragraph: This new paragraph comes right after the one shown above starting with “Further insight...”

“Important features of near 0°C conditions are its first and last occurrences over the annual cycle (Fig. 10). For stations with ‘warm season’ surface temperatures continually well above 0°C, onset is defined as the first occurrence of near 0°C conditions in the autumn and cessation is defined to the last occurrence of these conditions in the spring. For stations such as Cambridge Bay with surface temperatures near 0°C in every ‘warm season’ month and below this value in every ‘cold season’ month, onset is defined as the first occurrence of these conditions in the spring and cessation is defined as the last occurrence of these conditions in the autumn. For stations such as Churchill, near 0°C conditions can occur every month so neither onset nor cessation can be defined.”

Paragraph 3: Revised Paragraph: This revised paragraph comes right after the one just above starting with “Important features...”. It is a revised version of the current one. Deleted text is shown with a strike out and inserted text is in bold.

“It would be expected that, with overall warming (Bush and Lemmen, 2019), the onset for near 0°C would occur later in autumn and earlier in spring at most locations. As shown in Table 4 and Fig. 10c, both of these trends are significant at only one location, Toronto. Whitehorse shows significant earlier spring cessation (Table 4) and St. John’s shows significant later autumn onset (Table 4 and Fig. 10d). ~~Cambridge Bay experiences near 0°C conditions in every ‘warm season’ month (Fig. 9b and Table X) so the onset of near 0°C in the spring and its cessation in the autumn were considered; neither showed statistically significant trends.~~ **Neither spring onset nor autumn cessation illustrated a statistically significant trend at Cambridge Bay (Table 4 and Fig. 10b).** No analysis was conducted for Churchill because near 0°C conditions occurred in every month (Table 4).”

6. Fig. 2 caption. “Blue ellipses and red crosses”: blue crosses instead?

Line in question: “Figure 2: The 92 stations used in the analysis (see text for details). Blue ellipses and red crosses show the 9 selected stations across Canada.”

Response: Thank you, it is corrected now. The words “ellipses and red” is removed.

New/Revised Text: Figure 2: The 92 stations used in the analysis (see text for details). Blue crosses show the 9 selected stations across Canada.

7. Fig. 8. Why is fig. 8 not in agreement with Table 2? For example, St. Johns shows an increase of 7 hours in maximum duration in Table 2 while fig. 8 shows a decrease. Toronto shows a decrease of 26.9 hours in Table 2 while fig. 8 show a decrease of about 400 hours.

Line in question: Table 2 and Figure 8

	Whitehorse	Vancouver	Calgary	Cambridge Bay	Winnipeg	Churchill	Toronto	Montreal	St. John's	Units
Number of Days	139.5 (100.4)	62.3 (7.2)	161.0 (46.1)	67.9 (-110.2)	108.7 (0)	92.6 (-143.6)	121.6 (-316.9)	107.0 (-69)	155.0 (90.4)	days
Number of Hours	1229.9 (-10.3)	530.3 (1.8)	1149.1 (-2.6)	854.3 (-2)	902.3 (-8.9)	1025.7 (-3.1)	1201.8 (-26.9)	979.5 (-10.3)	1744.5 (-3.6)	hours
Number of Events	194.3 (-18.5)	93.9 (3.1)	279.8 (-16.9)	66.5 (0)	145.8 (9.3)	99.7 (0.8)	142.1 (-35.8)	133.7 (-15.5)	163.6 (-15.5)	events
Maximum Duration	58.3 (22.9)	37.2 (-5.2)	39.5 (-6.3)	89.5 (-36.6)	53.8 (8.9)	84.1 (-27.9)	58.0 (-25.4)	52.2 (6.2)	91.0 (7)	hours

Response: Thank you for noticing this. Because of this comment, we found an error in the revised document. Due to Reviewer 3 and 4 comments, Table 2 was updated and modified. In the effort to follow the order in Table 1 the order of lines (1 vs 2) was switched; the average was changed but not the trend in bracket. The Table 2 numbers were correct in the originally submitted document, but not in the revised version. Here is the original Table 2 from the first version of the submitted paper:

	Number of Hours		Number of Days		Number of Events		Maximum Duration	
	Aver	Sign	Aver	Sign	Aver	Sign	Aver	Sign
Whitehorse	100.4	no	-10.3	no	-18.5	no	22.9	no
Vancouver	7.2	no	1.8	no	3.1	no	-5.2	no
Calgary	46.1	no	-2.6	no	-16.9	no	-6.3	no
Cambridge Bay	-110.2	no	-2.0	no	0.0	no	-36.6	no
Winnipeg	0.0	no	-8.9	no	9.3	no	8.9	no
Churchill	-143.6	no	-3.1	no	0.8	no	-27.9	no
Toronto	-316.9	↻	-26.9	↻	-35.8	↻	-25.4	↻
Montreal	-69.0	no	-10.3	no	-15.5	↻	6.2	no
St. John's	90.4	no	-3.6	no	-15.5	no	7.0	no

In the analysis section we used the original trend values; so no modification is required.


This update also explains the second example, the difference found in Toronto. The first minor difference observed at St John's is due to the fact, that on Figure 8 trend lines were created using a simple linear regression versus the more sophisticated trend computation described in Sect. 3.2. This new figure was created due to the request of Reviewer 2 and served as a visualization time-series and major trend directions. The new updated Figure 8 trend line values are now matching the trend results listed in Table 2.

New/Revised Text:

Revised Table 2:

	Whitehorse	Vancouver	Calgary	Cambridge Bay	Winnipeg	Churchill	Toronto	Montreal	St. John's	Units
Number of Days	139.5 (-10.3)	62.3 (1.8)	161.0 (-2.6)	67.9 (-2)	108.7 (-8.9)	92.6 (-3.1)	121.6 (-26.9)	107.0 (-10.3)	155.0 (-3.6)	days
Number of Hours	1229.9 (100.4)	530.3 (7.2)	1149.1 (46.1)	854.3 (-110.2)	902.3 (0)	1025.7 (-143.6)	1201.8 (-316.9)	979.5 (-69)	1744.5 (90.4)	hours
Number of Events	194.3 (-18.5)	93.9 (3.1)	279.8 (-16.9)	66.5 (0)	145.8 (9.3)	99.7 (0.8)	142.1 (-35.8)	133.7 (-15.5)	163.6 (-15.5)	events
Maximum Duration	58.3 (22.9)	37.2 (-5.2)	39.5 (-6.3)	89.5 (-36.6)	53.8 (8.9)	84.1 (-27.9)	58.0 (-25.4)	52.2 (6.2)	91.0 (7)	hours

Revised Figure 8:



8. Fig. 11. The temperature pathways and corresponding text need more explanation. Is this section and figure really needed?

Line in question at end of Sect. 4, lines 426-431: “It is recognized that there are four pathways through which surface temperatures enter and leave near 0°C conditions (Fig. 11). These pathways are events (1) from above to below, (2) from below to above, (3) from above back to above, and (4) from below back to below. As shown in Fig. 11, pathway 3 is most common at all selected stations, Calgary experiences this pathway more than anywhere else (94 events/year), and this pathway dominates at Vancouver. Except for Vancouver and Calgary, the least common pathways are 1 and 2 (which have very similar values at all stations) with the overall lowest value being at Cambridge Bay (< 4 events/year).”

Response: First, the temperature pathways section was developed in response to, in particular, Reviewer 4, Major Comment 2. This comment states “...an event that drops into the window but never reaches zero might produce the same count as an event that warms into the window without ever exceeding zero, but have quite different precipitation characteristics.” Second, we had mentioned such issues in our original submission but we had not carried through a quantitative analysis.

This analysis is very appropriate. The article is concerned with patterns of near 0°C conditions including the temporal ones leading to their formation. This material is important because it addresses the manner through which near 0°C conditions arise. To our knowledge, no comparable analysis has ever been carried out before. The term ‘pathway’ is used and it refers to the surface temperature evolution through which it enters and leaves near 0°C conditions. There are only four possibilities as we have shown.

New/Revised Text: Here is the revised paragraph with changes shown. New text is shown in bold and deleted text is crossed out.

~~It is recognized that there are four pathways through which surface temperatures enter and leave near 0°C conditions (Fig. 11).~~ **It is also recognized that surface temperatures can enter and leave near 0°C conditions in different manners. These are referred to as pathways and there are only four possibilities (Fig. 11).** These pathways are events **with surface temperatures passing** (1) from above to below **near 0°C conditions**, (2) from below to above, (3) from above back to above, and (4) from below back to below. As shown in Fig. 11, pathway 3 is most common at all selected stations, Calgary experiences this pathway more than anywhere else (94 events/year), and this pathway dominates at Vancouver. Except for Vancouver and Calgary, the least common pathways are 1 and 2 (which have very similar values at all stations) with the overall lowest value being at Cambridge Bay (< 4 events/year).

There is no change in other, later paragraphs in which the term ‘pathways’ is used.

A REQUEST FROM THE AUTHORS

We noticed that we made a mistake in responding to initial comments by Reviewer 4. We would like to revise the deletion of a sentence.

The comment in question (repetition from previous review):

Reviewer 4 / Comment 27 / Lines 246-248: This text warrants literature support. Once a lake is frozen it no longer serves as a heat sink.

Lines 246-248 text: “For some stations in the North and the Atlantic region, the development and melting of sea ice also acts to maintain temperatures near 0°C; inland stations must experience a similar effect from lake ice. Most stations are similarly affected by melting and freezing of snow on the surface.”

Response: The text has been modified and references have been added to address the reviewer’s comment.

Revised Text: “Many of the oceans surrounding Canada are close to 0°C, especially in the cold season (Phillips, 1990; Larouche and Galbraith, 2016). For some stations in the North and Atlantic regions, the development and melting of sea ice also must act to maintain temperatures near 0°C; inland stations in close proximity to large lakes likely experience a similar effect from the development and melting of lake ice. For example, Larouche and Galbraith (2016) showed that water temperatures in parts of the Great Lakes are close to 0°C during the cold season. Most stations are similarly affected by melting and freezing of snow on the surface (Takeuchi et al., 2002).”

The requested revision

Our mistake: The last sentence in the above paragraph (Lines 248-249) was not referred to by Reviewer 4. This sentence is as follows in the original submission. “*Many stations are also affected by frozen soil; its freezing in the autumn and its melting in the spring would again act to maintain temperatures near 0°C (Oke, 1987).*” We subsequently found that the Oke (1987) reference was not suitable so we deleted the whole sentence. We have now found an appropriate reference, which is Boone et al. (2000).

We request that the paragraph include the original last sentence with the new reference in the paper.

New/Revised Text: The revised paragraph should read as follows with the sentence in question in bold:

“Many of the oceans surrounding Canada are close to 0°C, especially in the cold season (Phillips, 1990; Larouche and Galbraith, 2016). For some stations in the North and Atlantic regions, the development and melting of sea ice also must act to maintain temperatures near 0°C; inland stations in close proximity to large lakes likely experience a similar effect from the development and melting of lake ice. For example, Larouche and Galbraith (2016) showed that water temperatures in parts of the Great Lakes are close to 0°C during the cold season. Most stations are similarly affected by melting and freezing of snow on the surface (Takeuchi et al., 2002). **Many stations are also affected by frozen soil; its freezing in the autumn and its melting in the spring would again act to maintain temperatures near 0°C (Boone et al., 2000).**”

New reference:

Boone, A., Masson, V., Meyers, T., and Noilhan, J.: The influence of the inclusion of soil freezing on simulations by a soil–vegetation–atmosphere transfer scheme. *Journal of Applied Meteorology*, 39, 1544-1569. doi: 10.1175/1520-0450(2000)0392.0.CO;2.hal-02269591, 2000.

Manuscript showing all revisions

Dated February 27, 2020

Assessment of Near 0°C surface Temperature and Precipitation type Characteristics patterns across Canada

Eva Mekis¹, Ronald E. Stewart², Julie M. Theriault³, Bohdan Kochtubajda⁴, Barrie R. Bonsal⁵ and Zhuo Liu²

¹Meteorological Research Division, Environment and Climate Change Canada, Toronto, M3H5T4, Canada

²Department of Environment and Geography, University of Manitoba, Winnipeg, R3T2N2, Canada

³Centre ESCER, Department of Earth and Atmospheric Sciences, Université du Québec à Montréal, H2X3Y7, Canada

⁴Meteorological Service of Canada, Environment and Climate Change Canada, Edmonton, T6B1K5, Canada

⁵Watershed Hydrology and Ecology Research Division, Environment and Climate Change Canada, Saskatoon, S7N3H5, Canada

Correspondence to: Eva Mekis (eva.mekis@canada.ca)

Abstract. The 0°C temperature threshold is critical to many meteorological and hydrological processes driven by melting and freezing in the atmosphere, surface and sub-surface and by the associated precipitation varying between rain, freezing rain, wet snow and snow. This threshold ~~,linked with freeze-thaw,~~ is especially important in cold regions such as Canada ~~because it is linked with freeze-thaw, snowmelt and permafrost.~~ This study develops a Canada-wide perspective on near 0°C conditions using hourly surface temperature and precipitation type observations from 92 climate stations for the period 1981 to 2011 ~~with a particular focus on the occurrence of its associated precipitation. Since this analysis requires hourly values of surface temperature and precipitation type observations, it was limited to 92 stations over the 1981-2011 period.~~ In addition, nine stations ~~representative of~~ various climatic regions are selected for further analysis. Near 0°C conditions are defined as periods when the surface temperature is between -2°C and 2°C. Near 0°C conditions occur often across all regions of the country although the annual number of days and hours and the duration of these events varies dramatically. Various ~~forms~~-types of precipitation (~~including for example~~ rain, freezing rain, wet snow and ice pellets) ~~sometimes occur with these temperatures. The near 0°C conditions and the reported precipitation type occurrences tends to be higher in Atlantic Canada, although high values also occur in other regions. are sometimes linked with these temperatures with highest fractions tending to occur in Atlantic Canada.~~ Trends of most temperature-based and precipitation-based indicators show little or no change despite a systematic warming in annual surface temperatures over Canada. Over the annual cycle, near 0°C temperatures and precipitation often exhibit a pattern with short durations near summer driven by the diurnal cycle, while longer durations tend to occur more towards winter associated with storms. There is also a tendency for near 0°C surface temperatures to occur more often than expected relative to other temperature windows at some stations; due at least in part to diabatic cooling and heating occurring with melting and freezing, respectively, in the atmosphere and at the surface.

1 Introduction

In cold regions such as Canada, ~~numerous both~~ environmental processes and socio-economic activities are significantly impacted by temperatures near 0°C. At larger spatial and temporal scales, the seasonal arrival and retreat of 0°C temperatures influence snow melt/accumulation, hydrologic processes (spring freshet, freshwater ice duration ~~that affects open water evaporation~~, flooding), permafrost thaw and related slumping, transportation (e.g., ice roads), growing season length, and animal hibernation (e.g., Bonsal and Prowse, 2003). At smaller scales, periodic transitional episodes from below to above 0°C (or vice versa) can have adverse effects including mid-winter ice-jams and related flooding (e.g., Beltaos et al., 2006; Lindenschmidt et al., 2016), ~~animal starvation,~~

freeze-thaw damage to infrastructure (e.g., Kraatz et al., 2019), unseasonal frosts (e.g., McKenney et al., 2014), and recreation impacts (skiing, avalanches) (e.g., Moen and Fredman, 2007; Laute and Beylich, 2018). Furthermore, if these periods are associated with precipitation (e.g., freezing rain/ice-storms), severe and sometimes life-threatening impacts are possible including damaged electrical transmission infrastructure, air traffic disruptions and hazardous road conditions. If near 0°C occurrences change, this will have subsequent impacts on all these issues.

Many regions of ~~Canada the country~~ have experienced major impacts from near 0°C events. For example, a prolonged period of freezing rain, drizzle, and ice pellets severely affected a region stretching from southeastern Ontario to southwestern Quebec, as well as northeastern United States in early January 1998 (Henson et al., 2007; Henson et al., 2011). Some areas experienced in excess of 100 mm of freezing precipitation. Transportation was shut down and damage to the electrical infrastructure, trees and farms was catastrophic. There were an estimated 47 fatalities in Canada and the United States and \$4 billion U.S. in total losses (Lecomte et al., 1998; Kerry et al., 1999; Milton and Bourque, 1999; Klaassen et al., 2003). More recently, ice storms over northeast New Brunswick in January 2017 (McFadden and Theriault, 2018) and the Fraser Valley of British Columbia in December 2017 (Mendoza and Schmunk, 2017) severely impacted electrical power grids, snapping poles, downing lines and leaving thousands of people without electricity for days.

Near 0°C conditions occur over many areas of the world. Over land areas, even high terrain experiences such conditions. Some of the most critical questions associated with our changing climate are associated with temperature change and the hydrological cycle. This certainly applies to the future occurrence of near 0°C conditions and its associated precipitation.

There naturally continues to be many studies documenting global, regional and local surface temperatures ranging from climate change synthesis reports (Intergovernmental Panel on Climate Change, 2013) to studies of temperature at individual weather stations. Many of these of course include 0°C but few have focused on this temperature, although, for example, Wexler et al. (1954), Fujibe (2001) and Takeuchi et al. (2002) all showed that melting and/or freezing can affect its occurrence and/or duration.

In contrast, several studies have examined the varying precipitation types near 0°C. Examples of studies examining the occurrence of rain and snow in relation to temperature and, in some cases, relative humidity include U.S. Army Corps of Engineers (1956), Auer (1974), Matsuo and Sasyo (1981a,b), Matsuo et al. (1981) and Kienzle (2008). In general, these studies found a steady increase in the probability of rain as temperatures increase from near 0°C to higher values and they also pointed out that lower atmospheric moisture generally leads to higher temperatures needed for rain to occur. Dai (2008) built on this work to examine global distributions of rain and snow although freezing rain was just included within the snow category in this study. Sims and Liu (2015) developed an algorithm for use with remotely sensed observations to discriminate different types of surface precipitation; they emphasized the importance of atmospheric moisture. Jennings et al. (2018) examined the threshold temperature at which rain and snowfall with equal frequency, mapped this parameter over the Northern Hemisphere, and illustrated its strong dependence on atmospheric humidity. Overall, many of these studies illustrated that the variation in atmospheric moisture directly affects the temperature threshold needed for rain to occur with lower values acting to increase required temperatures. In addition, atmospheric humidity can also influence the characteristics of the melting particles and consequently the threshold. For example, under saturated conditions aloft with rising air, cloud droplets can be produced and later be captured by falling precipitation particles. The ensuing rimed, more dense particles require greater fall distances and/or higher temperatures to melt (Stewart et al., 2019).

There have been previous Canadian studies related to aspects of the 0°C temperature threshold including associated precipitation. At large scales, Bonsal and Prowse (2003) assessed 20th century trends and variability in spring and autumn 0°C dates across the

country and found significant trends toward earlier springs (particularly in western areas) but little change during autumn. Regionally, Stewart and Yiu (1993) examined near 0°C conditions including their horizontal scales and associated precipitation over southern Ontario. In terms of associated precipitation, MacKay and Thompson (1969) published the first climatology of freezing precipitation for Canada and this was later updated by Stuart and Isaac (1999) and Wang (2006). Many case study analyses of heavy precipitation and/or freezing rain events have been carried out to investigate storm structure and associated precipitation production mechanisms (for example, Henson et al., 2007; Henson et al., 2011). As well, two recent articles, one focused on western and northern Canada (Kochtubajda et al., 2017) and one on eastern regions (Bresson et al., 2017), have collectively documented some of the hazardous freezing conditions occurring within the country.

A number of studies have examined climatological characteristics of freezing precipitation across North America, Europe and Asia (see for example, Carriere et al., 2000; Cortinas, 2000; Changnon, 2003; Houston and Changnon, 2007; Groisman et al., 2016; Kämäräinen et al., 2016). These investigations used a variety of observational and model datasets, found wide variations in its freezing precipitation occurrence and, in some cases, related these findings to contributing factors such as mountain barriers and coastlines. A recent study in Europe also identified the impacts of extreme weather on critical infrastructure in Europe (Groenemeijer et al., 2017). One of the results of interviews with infrastructure and emergency managers was that the impacts of freezing precipitation, snowfall, snow loading and snow storms were of most concern.

Although studies have assessed various aspects of the 0°C conditions, there has been no Canada-wide assessment describing characteristics and trends in their historical occurrence. A preliminary assessment in the frequency of near 0°C surface temperatures across Canada (Fig. 1) shows that they occur at stations at which average annual surface, with the exception of Vancouver, there is a relatively high frequency of these temperatures range between below, near, and substantially above 0°C in all regions of the country despite wide ranging climates. Given the aforementioned importance of this threshold, additional information is necessary to better understand its spatial and temporal characteristics across the country Canada. This includes the associated precipitation that often results in the greatest impacts.

The objective of this study is therefore to quantify and improve understanding develop a Canada-wide perspective on of near 0°C conditions with a particular focus on its associated temperature and precipitation type patterns across Canada. Data and methods are described first, followed by an assessment of climatology and trends in key identified variables. Next, representative selected stations are examined in more depth followed by a discussion of several critical near 0°C features and by concluding remarks.

2 Data and methods

2.1 Data

A combination of automated systems and human observations comprise Environment and Climate Change Canada's (ECCC) surface weather networks. These measurements are subject to subsequent manual and automated quality control procedures and are available from ECCC's National Climate Data and Information Archive Historical Climate Data portal at various temporal scales ranging from hourly to annual (<http://www.climate.weatheroffice.ec.gc.ca>, ECCC, 2019a). Since this study focuses on the identification of conditions near 0°C at the surface along with associated precipitation, hourly surface temperature and the manual precipitation type observations across Canada were retrieved from the archive.

Selecting appropriate stations for further analysis was determined by data availability and is fraught with several complicating issues including missing values and changes to the observing program. The initial period considered was 1953-2016, with the criterion of a minimum 25 years of record. Although 227 stations satisfied this criterion for both hourly surface temperature and precipitation type information, many were not operating 24 hours a day whereas others changed their daily-observing practices over the period of interest (for example, from 24 hours per day to fewer or vice versa). It was therefore decided that only stations operating 24 hours a day would be used, and these had to have at least 90% hourly surface dry bulb temperature data availability (equivalent to an average requirement of 21 hours per day). For precipitation type observations, the only condition was the existence of the hourly observing program during the study period.

Consequently, to maximize the number of stations but still maintain a sufficiently long period for climatological studies, the 31 year period of 1981-2011 was chosen. This latter date was influenced by the dramatic drop in the number of stations archiving information after 2012 (Mekis et al., 2018). This resulted in 92 stations being used for analyses (Fig. 2) that provide reasonable coverage over the country. For these stations, the following 12 manual weather (precipitation) types observations are considered: rain, rain showers, drizzle, freezing rain, freezing drizzle, snow, snow grains, ice crystals, ice pellets, ice pellet showers, snow showers and snow pellets. Reporting was carried out according to World Meteorological Organization standards that are described in the Manual of Surface Weather Observation Standards (MANOBS <https://www.canada.ca/en/environment-climate-change/services/weather-manuals-documentation/manobs-surface-observations.html>, ECCC, 2019b). Precipitation intensity is characterized using four distinct values based on visibility or the rate of rainfall ranging from absent to 'heavy', but, for the purposes of this study, only the presence or absence of precipitation types was considered. Definitions are described in American Meteorological Society (2018), ECCC (2019b) and WMO (2017).

Canada is a vast country with many different climatic regions (Gullett et al., 1992) and ecozones (Zhang et al., 2001). For easier evaluation of the results, similar to Vincent et al. (2018), the country was divided into six broad regions based on the 13 provincial/territorial boundaries (Fig. 2). In addition to the Canada-wide analysis, nine representative stations were chosen for further analysis (Fig. 2). These stations, having high quality consistent observing practices, represent contrasting climatic conditions across the country.

2.2 Methods

This analysis identified key near 0°C characteristics and threshold events during the study period. There is no obvious precise, physically -based criterion that can be used to characterize conditions near 0°C although we did want to focus on conditions in which embedded change in precipitation types is common. We note that WMO Solid Precipitation Intercomparison Experiment (SPICE) broke down precipitation into three categories based on surface temperature (T) with those being snow < -2°C, mixed precipitation -2°C ≤ T ≤ 2°C, and rain > 2°C (Nitu et al., 2018); this approach was similar to that used by Yang et. (1995, 1998). As well, Matsuo et al. (1981) found that almost all of the precipitation near rain-snow transitions in Japan is rain if the surface temperature is ≥ 2°C and relative humidity is ≥ 90% and we also note that Kochtubajda et al. (2017) found that 75% of freezing rain across the Canadian Prairies and northern Canada fell at surface temperatures ≥ -2°C. To provide a reasonable symmetric (dry bulb) temperature (defined as T) window straddling 0°C with embedded large fractions of overall occurrences of varying precipitation types, we defined near 0°C, these conditions were defined as -2 ≤ T ≤ 2°C throughout the paper.

A total of 21 indicators were considered (Table 1). The first four indicators are associated with the near 0°C temperature condition (measured as hourly surface dry bulb temperature) without any consideration of precipitation. ~~They are the average-number of days per year having temperatures in this range; average-number of hours per year with temperature in this range; average-number of events per year; and annual maximum duration of the events within the study period. The event is defined as the number of consecutive hourly observations within the $-2 < T < 2^{\circ}\text{C}$ range. Note that to be considered a single event, there could be no more than three continuous hours of missing data.~~ To assess precipitation during near 0°C conditions, a further thirteen precipitation type indicators were computed. In addition, the combination of temperature and precipitation type provides an additional four indicators. ~~Four temperature-based indices were calculated from this information (Table 1). They are the average number of days per year having temperatures in this range; average number of hours per year with temperature in this range; average number of events per year; and annual maximum duration of the events within the study period. The event is defined as the number of consecutive hourly observations within the $-2 \leq T \leq 2^{\circ}\text{C}$ range. Note that to be considered a single event, there could be no more than three continuous hours of missing data.~~

~~The first set of indicators are associated with the near 0°C condition without the assurance of any precipitation occurrence. To assess precipitation during near 0°C conditions, further indicators combining the temperature and precipitation type conditions were computed (Table 1).~~ These include the annual average hours with any of the 12 aforementioned precipitation type conditions; the annual average hours with only freezing rain; the percentage of time in which any precipitation (from the 12 types) occurred; and the percentage of time that freezing rain alone occurred. Freezing rain is highlighted, ~~these rather than other frozen precipitation~~, since its occurrence often results in major impacts.

To characterize these variables, the 1981-2011 average was determined for each station and mapped across the country. The assessment of the trend can be challenging for time series with low occurrences and repetitive (tied) values (Frei and Schar, 2001 and Keim and Cruise, 1998) such as the annual number of days with certain precipitation types. The Chi-square test was applied to the data to determine whether the data is following normal or Poisson distribution. Depending on the results, two different approaches were used to estimate the trends. For the data with normal distribution (most of the surface temperature related indicators), the estimated magnitude of the trend is based on the slope estimator of Sen (1968), and the statistical significance of the trend is based on the nonparametric Kendall's tau-test (Kendal, 1955). This same approach was recently used to assess trends in Canada's climate (Vincent et al., 2015) and in surface temperature and precipitation indices (Vincent et al., 2018). ~~In addition, nonparametric linear trends were estimated using the approach by Sen (1968) with statistical significance based on the nonparametric Kendall's test (Kendall, 1955).~~ This test is less sensitive to the non-normality of the data distribution, and less affected by missing values and outliers as compared to the frequently used least squares method. Since serial correlation is often present in climatological time series, the method also involves an iterative procedure that takes into account the first lag autocorrelation (Zhang et al., 2000). For the data following the parametric Poisson distribution (mainly the precipitation type indicators), the logistic model was applied to transform the time-series (McCullagh and Nelder, 1989). This case only the direction (positive and negative trends) and the significance of the fitted curve were used in the analysis. The statistical significance of the trends were assessed at the 5% level. Given the aforementioned aspect of missing data and similar to earlier works (Mekis et al., 2015; Vincent et al., 2018), ~~it was decided that, for trend computation, a minimum of 90% of the values (29 out of 31 years) was required.~~, the trends were calculated if there were at least 90% of the values (29 out of 31 years) were available, otherwise the trends were set to missing.

180 In addition to the temperature and precipitation type occurrence information, hourly sky cover was also extracted over the nine ~~representative~~-selected stations. Sky conditions are reported in units of octas according to World Meteorological Organization standards that are described in the Manual of Surface Weather Observation Standards (ECCC, 2019b). If multiple cloud layers are observed, then the octas of the layers are summed. This information was only examined for the longest duration near 0°C conditions at these stations.

185 Given the significance of the 0°C reference line in this study, the -2, 0 +2 °C average annual surface temperature contour lines was computed from the 1981-2010 climate normal period (http://climate.weather.gc.ca/climate_normals/index_e.html, ECCC, 2019c) to be used later in the analysis. Kriging with a linear Variogram model and a grid spacing of 50 km was applied to create the interpolated surface temperature map from the 1619 stations for Canada.

3 Climatology and Trends

190 3.1 The 31-Year Climatology for 1981-2011

Figure 3a shows the ~~average~~-number of days per year when surface temperatures ~~were-was~~ between -2°C and +2°C. In relation to the average annual 0°C surface temperature line, it is apparent that high values generally occur at stations that are near or above 0°C. There are distinct regional patterns with the largest values (120 to 200 days) concentrated in three main areas. Highest occurrences are found in interior British Columbia and southern Alberta extending into southern Saskatchewan with maximum values within or on the leeward side of the western ~~e~~Cordillera. ~~It is likely that factors such as chinooks contribute to these high occurrences by occasionally bring warm air into the region that, during the cold season, results in temperatures near 0°C.~~ The second region with high values is in Atlantic Canada where temperatures often fluctuate around 0°C during the cold season due to the Maritime influence. Southern Ontario also has a relatively high number of occurrences likely due to its more southerly location and resultant influxes of warmer southern air masses during the cold season. Mid-range values (80 to 120 days) occur in the
195
200 continental interior stretching from the Yukon through central Canada to Quebec and Labrador. This area is colder than the previously mentioned regions with ~~less-fewer~~ incursions of warm air during the cold season. Lowest (40 to 80) values are in the North due to even fewer warm air incursions. Low values also occur in southwestern British Columbia where temperatures seldom dip to values below 0°C.

The preceding indicates that, on average, near 0°C conditions can occur over 50% of the days in regions with the highest values in
205 Fig. 3a. Even in the most northerly locations, such conditions occur approximately once per week on average.

Figure 3b shows the average number of hours per year with surface temperatures between -2°C and +2°C. Unlike the number of days in Fig. 3a, the number of hours shows less dependence on the location of the average annual 0°C surface temperature line. Overall, the spatial distribution is similar to that of Fig. 3a, but ~~some~~-differences are apparent. The same three general regions of high values still occur but the western one is more localized and does not extend east of the Canadian Rockies. This may indicate
210 that such conditions are short-lived east of the Canadian Rockies so they show up in the number of days but not ~~in-as~~ an extended number of hours.

High values are even more pronounced in Fig. 3b than in Fig. 3a at one ~~near~~-northern coastal British Columbia station (Terrace) where maximum values of 1800-2000 hours represent approximately 80 days per year ~~are also evident~~. This northern British Columbia station ~~next to~~near the ocean may share many characteristics to ones in the Atlantic region. Note that northern coastal

215 British Columbia maximum values (1800 - 2000 hours) represent approximately 80 days per year. In contrast to Fig 3a, there, here
are now three ~~general~~-distinct areas of low values (< 800 hours) in the average number of hours near 0°C (Fig. 3b). These areas
are the far North; southwestern British Columbia; and- The northern region is split into two with one being in the far North and the
~~second being in~~ northern British Columbia, ~~and Alberta as well as in the Yukon and western Nunavut~~and the western Northern
Territories. This latter region experiences warm summer conditions when temperatures seldom reach this low and cold winter
220 conditions when temperatures seldom reach this high. ~~Stations to the east more often experience near 0°C temperatures in the
warm season.~~

Figure 3c shows the average number of events per year. Similar to Fig. 3a, almost all high values occur at stations with average
annual surface temperatures near or above 0°C. Spatial patterns are also similar to those in Fig. 3a with maximum values (180-
280) in the west extending into southern Saskatchewan. High values are observed within the previously mentioned areas of southern
225 ~~Ontario and~~ the Atlantic region, but also occur on the north shore of Lake Superior. The number of events in Atlantic Canada is
comparable to the eastern Prairies even though there are far more hours in the Atlantic region. The number of events at some
stations in southern Ontario (120 - 140) is also comparable to the number in the southern Northwest Territories even though, again,
there are far more hours in southern Ontario.

The annual maximum duration of ~~the annual events lengths in hours~~, characterizing the occasional persistence of such events, is
230 shown in Fig. ~~ure~~ 3d. Maximum duration of events tend to show little dependence on the average annual 0°C surface temperature
line across the entire country. The spatial pattern differs from those in Figs. 3a-c with longest durations in the Atlantic region and
some interior stations in British Columbia including the northern coastal region. Other large values occur near coastlines in the
North. These values range up to 130 hours or ~5 days. Lowest maximum durations (10 - 20 hours) occur in the lee of the Rocky
Mountains as well as in southern British Columbia. ~~This may be related to the occurrence of chinooks within which temperatures
can quickly pass from below to above 0°C (Brinkman and Ashwell 1968).~~

Figures 4a to 4d provide climatologies of precipitation types associated with near 0°C conditions. First, Fig. 4a shows the average
annual number of hours with any of the 12 reported precipitation types listed in Table 1. There is no strong dependence in the
occurrence of any weather type in relation to the position of the average annual 0°C surface temperature line. Maximum values
(up to 800 hours) are primarily concentrated in the eastern half of the country although, as with the number of hours in Fig. 3b,
240 northern coastal British Columbia again is associated with high values. Such precipitation types are rare in most of western Canada
(except northern coastal British Columbia) with the lowest value (40 - 80 hours) occurring on Vancouver Island. ~~The magnitude
of variation is enormous.~~ There is more than an order of magnitude difference between the lowest and highest values across the
country. There is even a huge variation between the two farthest north stations (120 - 160 hours and 360-800 hours).

Figure 4b shows the percentage of near 0°C conditions with associated precipitation types, where the number hour reported any
245 12 weather type events are divided with all hours with near 0°C conditions. As in Fig. 4a, there is little dependence on average
annual surface temperature. The spatial pattern is quite similar to that in Fig. 4a but with a few exceptions. Highest values are again
in eastern Canada and northern coastal British Columbia. The map reveals that over 40% of near 0°C conditions are associated
with precipitation types in these regions but it is only of order 10 ~~to~~ 20% in western Canada.

Figure 4c shows the average annual number of hours of freezing rain with surface temperatures near 0°C. In contrast to Figs. 4a-
250 b, highest values of freezing rain primarily occur at stations with average annual surface temperature above 0°C. Maximum values
by far are in the Atlantic region (36 - 42 hours) and there is a regional maximum near Montreal (20 - 24 hours). Low-~~Small~~ values

(0 - 4 hours) occur in other regions of the country, especially in the North and much of the west. In fact, freezing rain is rarely ~~ever~~ reported on the western side of the Prairies and Northern areas.

Figure 4d shows the percentage of near 0°C hours with freezing rain. Consistent with the freezing rain hours in Fig. 4c, highest values of freezing rain percent also tend to occur at stations with average annual surface temperature above 0°C. The spatial pattern is also similar to that in Fig. 4c. This includes highest values (1.8 - 2.5%) being in the Atlantic region with a secondary maximum near Montreal. One isolated, high value (1.0 - 1.2%) does occur near Vancouver however.

3.2 Trends

Figure 5a shows trends in the number of days with temperatures near 0°C ~~over~~ during the 1981-2011 period significant at 5% level. There is no apparent relationship in these trends to the average annual 0°C surface temperature line. The majority of stations are characterized by a non-significant decrease with few having a non-significant increase (mainly in central Alberta and parts of British Columbia). Only two stations exhibited significant change; Toronto and the most northerly station Eureka which experienced a decrease of 27 and 19 days over the 31 year period respectively.

Figure 5b shows trends in the annual number of hours with near 0°C conditions. As in Fig. 5a, there is no simple relationship to the average annual 0°C surface temperature line, although more decreasing trends are evident in areas above 0°C. As in Fig. 5a, ~~most~~ Most stations also experienced a decrease over the 1981-2011 period. This is most pronounced over southern Ontario but also over other large areas of the country. Newfoundland, several stations in the North, as well as southern British Columbia experienced increases. Significant decrease were observed in many southern Ontario stations and again in Eureka. Only one station (western Newfoundland) showed a significant increase. The maximum changes with over 300 hours over the 31-year period (the equivalent of over 12 days) were found in three locations in Ontario, namely Toronto, Wiarton, which is north of Toronto, and London in the extreme southwestern part of the province.

Figure 5c shows trends in annual number of events with near 0°C conditions. No apparent pattern in relation to the 0°C surface temperature line was found. The pattern is similar to those in Figs. 5a and 5b, however, both increase and decreases are often observed in the same general areas. One exception is southern Ontario, which again shows consistent decreases. Only Toronto, Eureka, Sioux Lookout in northwestern Ontario, and Montreal show statistically significant change with the decrease of 36, 31, 20 and 16 number of events, respectively, over the 31 year period.

Figure 5d shows trends of the maximum duration. As in Figs. 5a-c, there is no simple relation to average annual surface temperature. There is a mix of decreasing and increasing patterns as well as many stations showing little change. Southern Ontario is completely characterized by decreases while other regions have a mixed pattern. Stations experiencing a statistically significant change are scattered across the country and exhibit both increases and decreases. Only four locations show significant change, decreasing in Toronto and Eureka with 25 and 18 hours per event and increasing in Yarmouth in Nova Scotia and Thompson in Manitoba with 34 and 19 hour per event, respectively.

4 Representative-Selected stations

The preceding analyses have illustrated Canada-wide conditions but it is also critical to examine individual stations in more detail. To address this, nine stations representing contrasting climatic conditions were chosen (Fig. 1). Annual average frequencies for

the four surface temperature and 13 precipitation type related indicators are available in Table 2 and 3 at each case study locations. The greatest annual number of days and number of hours with near 0°C conditions occurred at St John's with 155 days and 1744 hours respectively. The maximum annual number of independent event occurred at Calgary (280 events) while the maximum duration are the coaster stations St John's, Cambridge Bay and Churchill (91, 89 and 84 hours respectively). This is consistent with regional patterns shown in Figs. 3 and 4.

An important characteristic of near 0°C conditions is the duration of events ~~is studied further~~. As shown in Fig. ~~ure~~ 6, this distribution shows wide variation at each of the selected stations as well as between stations. For example, almost half of the events at Calgary were less than 2 hours but at Cambridge Bay this value was 7 hours; 90% of the events at Calgary were less than 12 hours but at Cambridge Bay this value was 43 hours. In parallel, Calgary experienced the largest number of events; Cambridge Bay the ~~least~~fewest.

While Table 3 contains the average number of hours ~~Figure 7 shows the distribution in the occurrence~~ of the 12 precipitation types, Fig. 7 shows the distribution in the occurrence ~~included in this study~~ based on the same input at the 9 stations over the 1981-2011 period. Snow dominates at all stations, except at St. John's, and it is most prevalent over Whitehorse and Calgary. Freezing rain is minimal at most western stations but steadily increases eastward. There is also a large variation in precipitation type occurrences between the two northern stations with, for example, drizzle and freezing drizzle being minimal at Whitehorse but not at Cambridge Bay.

As discussed in Sect. 3.2, many stations exhibited some change but few were ~~statistically~~ significant (Fig. 8 and Table 2). This characteristic is also prevalent for the 9 case studies with only two experiencing significant change. In particular, Toronto showed significant decreases in all 4 temperature-related indicators while Montreal had a decrease in the number of events per year. ~~These regional patterns are also somewhat reflected in dependence on average annual surface temperature; high values generally occur at stations with near or above 0°C conditions, although long durations also often occur at stations with below 0°C conditions.~~

The trend summary for the 12 weather types is shown in Table 3. Only Churchill exhibited a statistically significant trend (decrease) in the occurrence of any of these 12 types (last column). This arises from ~~five~~ significant decreasing types, namely rain showers, drizzle, freezing drizzle, ice pellets and snow showers. St. John's experienced five significant changes (in rain showers, freezing drizzle, snow, snow grains and snow showers precipitation types) but, due to the shift from snow grains and snow showers to snow, the overall changes for all types are not significant.

Patterns on sub-annual time scales are also examined. ~~The average monthly occurrences of near 0°C conditions were calculated and these values were compared with average monthly surface temperatures.~~ As shown in Fig. ~~ure~~ 9a, there is a strong dependence of average near 0°C occurrences on average monthly ~~surface~~ temperature over the 1981-2011 period; this pattern is largely independent of station. Largest occurrences naturally take place when average temperatures are close to 0°C. By $\pm 10^\circ\text{C}$, values have fallen to of order 25-35% of those at the peaks ~~with higher values generally associated with lower temperatures~~. Although rare, near 0°C conditions sometimes occurred with average monthly ~~surface~~ temperatures with more than 20°C away from the 0°C line (Fig. 9b). The five coldest differences occurred at Churchill and Cambridge Bay, and the five warmest ones occurred at Toronto, Winnipeg and Montreal.

Further insight can be gained by examining ~~the even~~ shorter time scales. ~~For example, Figure~~ Fig. 10 shows annual cycles of near 0°C ~~conditions~~ and associated precipitation type occurrences at four ~~of the nine representative~~ stations (Vancouver, Cambridge Bay, Toronto and St. John's) chosen to illustrate a range of variation. Near 0°C conditions do not occur during summer at all

southern stations (Figs. 10a, c, d) but they can occur in any other month. In contrast, these conditions only occur in summer at Cambridge Bay (Fig. 10b). The most frequent occurrence can be seen at St John's (Fig. 10d).

Furthermore, the occurrence of any precipitation type tends to take place towards the 'winter' side at southern stations, although there are exceptions. Such occurrences on the 'winter' side are probably linked with storms passing over the stations with associated precipitation, whereas occurrences on the 'summer' side normally just reflect the diurnal cycle.

Important features of near 0°C conditions are its first and last occurrences over the annual cycle (Fig. 10). For stations with 'warm season' surface temperatures continually well above 0°C, onset is defined as the first occurrence of near 0°C conditions in the autumn and cessation is defined to the last occurrence of these conditions in the spring. For stations such as Cambridge Bay with surface temperatures near 0°C in every 'warm season' month and below this value in every 'cold season' month, onset is defined as the first occurrence of these conditions in the spring and cessation is defined as the last occurrence of these conditions in the autumn. For stations such as Churchill, near 0°C conditions can occur every month so neither onset nor cessation can be defined.

~~It is evident that there is large inter-annual variability and this can mask expected systematic trends. For example, it would be expected that, with overall warming (Bush and Lemmen, 2019), the onset for near 0°C would occur later in autumn and earlier in spring. Although not as shown in Table 3 and, these patterns were generally not statistically significant. Only Toronto (Fig. 105c,) shows both of these trends to be statistically significant at only one location, Toronto. Whitehorse shows significant earlier spring cessation (Table 4) and St. John's shows significant later autumn onset (Fig. 109d and Table 4). Neither spring onset nor autumn cessation illustrated a statistically significant trend at Cambridge Bay (Table 4 and Fig. 10b). Cambridge Bay experiences near 0°C conditions in every 'warm season' month (Fig. 9b) so the onset of near 0°C in the spring and its cessation in the autumn were considered; neither showed statistically significant trends.~~ No analysis was conducted for Churchill because near 0°C conditions occurred in every month (Table 4).

It is also recognized that surface temperatures can enter and leave near 0°C conditions in different manners. These are referred to as pathways and there are only four possibilities (Fig. 11). These pathways are events (1) from above to below, (2) from below to above, (3) from above back to above, and (4) from below back to below. As shown in Fig. 11, pathway 3 is most common at all selected stations, Calgary experiences this pathway more than anywhere else (94 events/year), and this pathway dominates at Vancouver. Except for Vancouver and Calgary, the least common pathways are 1 and 2 (which have very similar values at all stations) with the overall lowest value being at Cambridge Bay (< 4 events/year).

Pathways with the highest and lowest fractions of any type of precipitation vary. The highest fractional occurrence of any precipitation is associated with pathway 1 at St. John's (88%), and this pathway is also associated with the highest fraction at Calgary and Cambridge Bay. Pathway 3 is associated with the highest fraction at Vancouver; pathway 4 is associated with the highest fraction at the remaining five stations. The lowest fractional occurrence is associated with pathway 2 at Calgary (19%), and this pathway is associated with the lowest fraction at all stations. St. John's experiences the maximum occurrence of freezing rain for all pathways. The pathway most often associated with freezing rain events varies between stations. For example, it is pathway 1 at Churchill; pathway 2 at Toronto, Montreal and St. John's; and pathway 4 at Winnipeg.

5 Characterizations and interpretations

5.1 ~~Country~~Canada-wide patterns and governing factors

The Canada-wide plots exhibit a number of patterns in the temperature indicators. First, there are three general regions in terms of high occurrences of near 0°C conditions (Fig. 3a-c). These are in central British Columbia and sometimes stretching to Saskatchewan, southern Ontario and the Atlantic region. The size of the regions varies with indicator. Regions with generally low occurrences are in the north, where such temperatures are not often reached, and in the lower southwestern part of British Columbia, where temperatures seldom reach 0°C. In terms of duration (Fig. 3d), highest values tend to be in the Atlantic region, eastern Northern region as well as northern coastal British Columbia. Lowest values are in southwestern British Columbia and stations just east of the Rocky Mountains in Alberta. These regional patterns are also somewhat reflected in dependence on the average annual 0°C surface temperature line with higher values generally occur at stations near or above 0°C. However, long durations also often occur at stations with below 0°C conditions.

There are numerous factors contributing to these patterns. One is the proximity to coastlines. Many of the oceans surrounding Canada are close to 0°C, especially in the cold season (Phillips, 1990; Larouche and Galbraith, 2016), and this acts to maintain station temperatures near this value. For some stations in the North and the Atlantic regions, the development and melting of sea ice also must acts to maintain temperatures near 0°C; inland stations in close proximity to large lakes likely must experience a similar effect from the development and melting of lake ice. For example, Larouche and Galbraith (2016) showed that water temperatures in parts of the Great Lakes, are close to 0°C during the cold season. Most stations are similarly affected by melting and freezing of snow on the surface (Takeuchi et al., 2002). Many stations are also affected by frozen soil; its freezing in the autumn and its melting in the spring would again act to maintain temperatures near 0°C (Oke, 1987; Boone et al., 2000).

Mountains can also be a contributor to near 0°C conditions. Circulations such as chinooks (American Meteorological Society, 2012) and valley/mountain flows are continually shifting temperatures and these can pass through 0°C in the cold season (Longley, 1967). Chinooks are common at Calgary. Longley (1967) found an average of 27 days per season (December-February) over the 1931-1965 period whereas Nkemdirim (1996) found 50 per season (November-February) over the 1951-1990 period. Associated temperature changes are typically rapid and are sometimes associated with several passes through near 0°C in a day (Brinkman and Ashwell 1968). Temperature changes with during chinooks can be dramatic. Nkemdirim (1997) pointed out that temperature increases of more than 25°C in less than 24 hours are typical, and Gough (2008) indicated that the largest observed temperature change at Calgary was associated with a 4-hour event in 1883 during which the temperature increased 30°C (from -17°C to +13°C). vary between approximately 0.6°C/h and 8.3°C/h with an average of 2.1°C/h. During chinook episodes, large temperature swings can also occur. For example, Brinkman and Ashwell (1968) showed that, at Calgary on Feb. 16, 1965, there were 4 near 0°C events (two 1-h periods, one 2-h, and one 5-hour) and no precipitation at all. Weather conditions were precipitation free this day.

These chinook effects undoubtedly contribute to the findings at Calgary. As shown in Sects. 3 and 4, it experiences many days and events with near 0°C conditions but relatively few hours since the events are short and it also experiences few hours of associated precipitation.

Long duration periods of near 0°C conditions based on surface temperature were further examined by identifying the longest duration events periods at the 9 representative selected stations (Table 45). The longest period was found in Cambridge Bay (197 hours) and the shortest in Vancouver (68 hours) which is still almost 3 days. These events occurred in every season with the longest being in June (Cambridge Bay). An analysis of the hourly sky conditions during these events was carried out by identifying the associated frequency of clear or mainly clear sky conditions (≤ 2 octas of clouds). All of the events Using the hourly observations

395 ~~at these stations, it is evident that all the events~~ were dominated by cloudy conditions; ~~which—that were often—sometimes~~
accompanied with ~~persistent~~ precipitation (Table 5). Such sky ~~cover—conditions~~ would contribute to reducing temperature swings
associated with daytime heating and nighttime cooling (see, for example, Ahrens et al., 2015). Many of the mainly or completely
clear reports were linked with temperatures initially passing into or finally passing out of these long duration near 0°C conditions.
There are also patterns with the occurrence of the associated precipitation types. ~~The country—Canada~~ is almost split in two between
400 west (low values) and east (high values) (Fig. 4). ~~This is likely a reflection of moisture access with eastern regions receiving warm,~~
~~humid tropical and subtropical air much more often than western regions and, in association, raising temperatures through 0°C~~
~~(Hare, 1997).~~ One exception in the west is northern coastal British Columbia (Terrace near the coast and nearby Smithers more
inland). Some Northern stations, particularly in the eastern portion, experience more occurrences than do many in British Columbia
and the Prairies. ~~The regional patterns related to the occurrence of any precipitation type are not directly reflected in dependence~~
405 ~~on the average annual 0°C surface temperature line. This is not the case for freezing rain where high values generally occur at~~
~~stations with above 0°C conditions.~~

The fractional occurrence of precipitation types (Fig. 7) can generally be explained as follows. In some of the western regions
(such as Calgary and Whitehorse), the atmosphere is normally dry which means that melting of snow aloft is reduced since the wet
bulb temperature is lowered. Over the temperature window studied here of 2°C, more of the snow will not have melted (~~Matsuo et~~
410 ~~al., 1981~~). In contrast, the Atlantic region is ~~buffeted by—subject to~~ large storms coming from the south as discussed by, for example,
~~Stewart et al. (1987) and Stewart et al. (1995a)~~. These vigorous storms almost always ~~mean—associated with~~ surface temperatures
passing from below to above 0°C with near saturated conditions (~~Stewart and Patenaude, 1988~~); they also ~~have—associated with~~
strong warm air advection aloft which often leads to inversions and freezing rain (~~Stewart et al., 1987~~).

An examination of the four pathways through which surface temperatures enter and leave near 0°C conditions revealed additional
415 patterns (Fig. 11). Regardless of whether the average annual surface temperature is above, near or below 0°C, the pathway
associated with initially above 0°C temperatures and back again (pathway 3) is associated with the most number of events at all
nine selected stations. In contrast, this most common pathway is associated with a variable fraction of precipitation occurrence
from low values, perhaps simply reflecting the diurnal cycle, to high values, possibly reflecting the passage of weather systems.
Freezing rain is most likely in events with temperatures passing from below to above 0°C (pathway 2) at several locations; this
420 pathway follows the temperature evolution during the passage of a warm front, which commonly leads to freezing rain (see, for
example, ~~Stewart et al., 1995b~~).

5.2 Enhanced occurrence of near 0°C conditions

As shown in Fig. 1, near 0°C conditions are prominent in several of the ~~representative—selected~~ stations. At four locations, this is
the most common temperature band despite wide variations in their whole span of temperature, ~~average annual temperature~~ as well
425 as geographic location. These stations are Whitehorse, Churchill, Toronto and St. John's. As well, there is a secondary peak near
0°C at Winnipeg. Cambridge Bay and Montreal did not display dramatic change near 0°C although there is a jump in occurrence
from colder temperatures. At Vancouver and Calgary, no obvious enhancement is apparent.

The enhanced occurrence of near 0°C conditions is similar to the pattern found in Japan by Fujibe (2001). This study attributed the
enhancement to the melting of falling snow, which cooled ~~surface~~ temperatures towards 0°C, as previously noted by, for example,
430 ~~Wexler et al. (1954). This is likely occurring in Canada—b~~But other factors are also critical as discussed in Sect. 5.1. For example,

the only mention by Fujibe (2001) of the role played by snowcover by Fujibe (2001) was as a factor leading to stable atmospheric conditions, which would reduce mixing that acts to eliminate isothermal layers near 0°C but there was no mention of the cooling effect on the atmosphere of the melting snowpack and refreezing of meltwater (Takeuchi et al., 2002). As well, the freezing of freezing rain drops at the surface acts to raise temperatures towards 0°C (Stewart, 1985); rain reaching the surface that subsequently freezes with lowering temperatures would have the same effect of the melting snowpack itself.

5.3 Factors affecting change

A question that arises is whether the observed warming over most Canada during the last few decades (Vincent et al., 2015, 2018), has impacted the occurrence of near 0°C conditions. Results from this study indicate a general lack of statistically significant change in the frequency and maximum duration of near 0°C conditions (Sect. 3.2). This is consistent with the Canada-wide assessment of annual freeze-thaw days (defined as the number of days with daily minimum temperature $\leq 0^\circ\text{C}$ and daily maximum temperature $\geq 0^\circ\text{C}$) for the period 1948-2016 that found a slight decrease in these events when averaged over the entire country (Vincent et al., 2018). However, regional differences were apparent including minor increases in the Prairies and Ontario.

The lack of significant change also holds for the four temperature indicators in this study including the onset and cessation of near 0°C conditions. The major exception is Toronto with significant declines in all four temperature indicators, as well as in the delayed onset and cessation of near 0°C conditions in the autumn, and earlier spring cessation of these conditions (Table 2 and Table 4). Montreal only showed significant declines in the number of near 0°C events (Table 2); Whitehorse showed significantly earlier spring cessation and St. John's showed significantly later autumn onset (Table 4). Several stations illustrated significant change (increase and/or decrease) in the occurrence of at least one precipitation type but only Churchill experienced a significant decrease in the occurrence of any precipitation type (Table 3).

It is not surprising that significant trends are not always evident. This may simply reflect a relatively short observational period (31 years) for stations with large inter-annual variability. As well, and as discussed in Sect. 5.1 and 5.2, many factors contribute to the occurrence of near 0°C conditions and these can counter each other. At many stations, temperatures in mid-winter are far below 0°C (Table 4). Even with the recent warming, that is still largely the case. For such stations, overall warming simply leads to a shift in near 0°C timing towards winter so the total number of occurrences does not necessarily change. In contrast, some stations, such as in southern Ontario (for example, Toronto), are normally not far below 0°C in mid-winter (Table 4). With the observed overall warming, these stations experience a shift in timing of near 0°C occurrence towards winter but more instances of mid-winter temperatures above 2°C will probably also occur, so they should would likely experience an overall decrease in near 0°C occurrences. In addition, the melting and freezing factors contributing to near 0°C conditions would continue to be active despite warming (Sect. 5.1). Snowcover and sea ice formed and melted; ground frozen and thawed. One exception may again be in southern Ontario where the ground may has not have frozen as much or be covered by as much snow (Vincent et al., 2015; Bush and Lemmen, 2019); these factors may have contributed to some of this region's significant decreases in occurrence related indicators.

These findings represent a basis for examining how near 0°C conditions may change in the future. Some studies of future freezing rain conditions have been carried out over North America (e.g., Lambert and Hansen, 2011; Jeong et al. 2019) and southern Quebec (Matte et al., 2018) but none has focused specifically on near 0°C conditions.

One can anticipate more indications of significant trends in near 0°C conditions. Even though this is not generally apparent so far, there should eventually be, for example, widespread delays in occurrence in autumn and earlier cessation in spring at southern locations. But the total number of occurrences may be countered by warmer mid-winters, although at locations such as in Toronto (Table 4), mid-winter temperatures ~~mid-winter~~ may be substantially more often sustained above 2°C (and thus no longer as often near 0°C). This latter factor may not occur for a long time at locations such as Winnipeg, where temperatures are normally far below 0°C in mid-winter (Table 4). Such ~~expected~~ changes would be expected to affect the occurrence of the near 0°C conditions shown in Fig. 1. For example, these conditions should become less prominent at Toronto but not necessarily at stations such as Winnipeg.

6 Concluding remarks

Temperatures near 0°C represent an important issue all across Canada. A comprehensive characterization of near 0°C conditions and the occurrence of associated precipitation types has been carried out. To accomplish this, the study had to carefully address which stations had good quality hourly data for a sufficient period; 92 locations were finally used for the 1981-2011 period. The period's last year 2011 was determined by the shrinking manual observation program required for the precipitation type observations. The analysis was completed for four temperature related indicators, 12 precipitation types and four combined temperature and precipitation type indicators. With the $-2 \leq T \leq 2^\circ\text{C}$ criterion, important insight was gained from the 31 year climatological and trend assessments for all locations and from in depth analysis of the nine case study locations. Key findings include:

- ~~The entire country~~Canada is characterized by highly variable near 0°C occurrences, events, durations, and associated precipitation types that have been quantified for the first time.
- There are systematic preferred regions in the occurrence of near 0°C conditions and the associated precipitation types that can be explained by large to regional scale conditions. Stations near oceans, for example, tend to have the largest values due to the moderating effects of near 0°C oceanic temperatures.
- A distinctive pattern related to the occurrence of several precipitation-related indicators is an east-west divide roughly down the center of the country. ~~This is likely the reflection of moisture access with eastern regions being subject to more moisture from warmer sub-tropical oceanic sources whereas much of western Canada does not have this moisture source.~~
- As expected, the monthly near 0°C occurrences peak when average monthly temperatures are near 0°C and there is a sharp fall off as averages move away from this value although there are rare occurrences at much colder or warmer temperatures.
- The occurrence of near 0°C values is influenced by numerous factors including ~~daily, radiatively-driven temperature variation the solar heating~~ and the annual temperature cycle. As well, ~~precipitation, snowcover, precipitation, ground conditions~~ and sea/lake ice can be important; all of these ~~simultaneously can~~ melt and freeze with effects always being to tip temperatures towards 0°C. In addition, ~~ocean~~ sea surface temperatures ~~near-around~~ Canada tend to ~~often~~ be near 0°C during the cold season especially ~~near some of Canada's coastlines~~ (Larouche and Galbraith, 2016). This is especially evident in Atlantic Canada and the Arctic; parts of the Great Lakes have comparable values in some seasons as well.

When such conditions occur, SSTs are generally slightly above 0°C although in some small areas they are below. SST values near 0°C would act to bring the temperature of the overlying air towards similar values.

- The longest duration events at the 9 selected stations are associated with persistent cloud conditions which act to reduce diurnally-driven temperature swings ~~are associated with prolonged cloud cover.~~

The aforementioned processes lead to near 0°C temperatures often being the most common occurring temperatures during the year. This observation is evident over wide geographic and climatic areas.

- Four pathways into and out of near 0°C conditions occur. At all of the nine selected stations across Canada, the pathway in which surface temperatures never becomes lower than -2°C is associated with the most number of events, and the one in which surface temperatures pass from below to above 0°C is associated with the lowest fraction of events with any precipitation except at Vancouver
- Even though surface temperatures have generally increased over the 1981-2011 period (Bush and Lemmen, 2019), the occurrences of near 0°C conditions have not trended in a similar fashion. This arises at least in part because increased temperatures in the warm season lead to fewer conditions but this is largely balanced by more in the cold season. In addition, the processes acting to maintain near 0°C conditions have generally continued to occur even as overall temperatures have increased. One exception is Toronto, which always had cold season temperatures not too far below 0°C.
- There has been no significant change in the frequency of occurrence of any of the 12 precipitation types or of freezing rain at most stations. However, the period of 31 years is relatively short to detect statistically significant changes. Using different selection criteria and period, Wang (2006) found that some areas of Canada experienced an increasing trend of freezing rain events over the 1953-2004 period.

Although our $-2 \leq T \leq 2^\circ\text{C}$ window for defining near 0°C conditions is justified as discussed in Sect. 2.2, it is recognized that other surface temperature windows could have been used. Overall findings would undoubtedly be similar, although quantitative values would change. For example, although not shown, near 0°C conditions largely exhibit the same occurrence patterns at the selected stations (Fig. 1) whether a $-1 \leq T \leq 1^\circ\text{C}$ window or a $-3 \leq T \leq 3^\circ\text{C}$ window is used instead. As well, a narrower (wider) surface temperature window would lead to smaller (larger) fractions of precipitation type occurrences simply being rain and snow. Future research should nonetheless thoroughly investigate the implications of applying different definitions to the study of near 0°C conditions. Related to trend computation, this study followed the statistical approach used in several recent Canadian studies (Vincent et al., 2015; Vincent et al., 2018), but the use of "significance/non-significance" terms for trends analysis can be restrictive. Additional research is needed to examine the strengths and probabilities of each relationships as discussed by, for example, Wasserstein et al. (2019).

~~Although this analysis has provided valuable insight regarding near 0°C temperatures across Canada, further research is required to obtain a better understanding of these conditions. For example, this study did not address the directional change in near 0°C occurrence, which impacts whether the region is going from a cold to warm state or vice versa. Cold to warm versus warm to cold would be linked with the two branches of the diurnal cycle as well as with warm and cold fronts, respectively; these factors should affect duration and precipitation types. In addition, near 0°C conditions are closely related to freeze thaw cycles that can have numerous economic and environmental impacts. This is also a critical area of future research.~~ The findings of this study may be applicable to many other regions. On examining a global map of the location of the average annual 0°C isotherm (Ahrens et al., 2016), one can appreciate that it slices through large expanses of land; areas at least occasionally passing through 0°C will be

massive. The recent WMO international project focusing on solid precipitation measurement (Nitu et al., 2018), utilized observational sites across North America, Europe and Asia and concerned, in part, with the $-2^{\circ}\text{C} \leq T \leq 2^{\circ}\text{C}$ so called “mixed” precipitation range examined here. And, Dai (2008) and Jennings et al. (2018) illustrated rain/snow issues over many regions of the world. Given the importance of near 0°C conditions and the large areas of the world subject to them, analyses carried out in this study should be conducted elsewhere.

In summary, this study can be considered an important step in the better understanding of near 0°C conditions and associated precipitation types across Canada and possibly many other regions.

~~This study can be considered a first step in the better understanding of near 0°C conditions and associated precipitation across Canada.~~

Data availability. The dataset used in this article is available through Environment Climate Change Canada.

Author Contribution. EM was the lead author who obtained and analyzed most of the data as well as writing several sections of the article. RES wrote several sections and carried out some of the analyses. JMT and ZL carried out analyses and contributed to the manuscript. BK and BRB contributed to the manuscript. All authors contributed scientifically by providing comments and suggestions.

Competing interests. There are no competing interests.

Special issue statement. This article is part of the special issue “Understanding and predicting Earth system and hydrological change in cold regions”. It is not associated with a conference.

Acknowledgements. The authors would like to thank David Hudak from Environment and Climate Change Canada, and XX anonymous reviewers for their useful comments and suggestions that have helped to improve the manuscript. The research was supported by the Changing Cold Regions Network supported by the Natural Sciences and Engineering Research Council (NSERC); the Global Water Futures study supported by Canada First Research Excellence Fund; the NSERC Discovery grants of RES and JMT.

References

Ahrens, C. D., Jackson, P. L. and Jackson, C. E. J.: *Meteorology Today: An introduction to weather, climate, and the environment: Second Canadian edition*, Nelson Education, ISBN-13: 978-0176530792, 2015.

Auer Jr, A. H.: The rain versus snow threshold temperatures, *Weatherwise*, 27(2), 67–67, 1974.

American Meteorological Society: Chinook. Glossary of Meteorology. Retrieved from <http://glossary.ametsoc.org/wiki/Chinook>, 2012.

American Meteorological Society: Glossary of Meteorology, available at: http://glossary.ametsoc.org/wiki/Main_Page (last access: October 3, 2019), 2018.

Beltaos, S., Prowse, T., Bonsal, B., Mackay, R., Romolo, L., Pietroniro, A. and Toth, B.: Climatic effects on ice-jam flooding of the Peace-Athabasca Delta, *Hydrological Processes*, 20(19), 4031–4050, doi:10.1002/hyp.6418, 2006.

- Bonsal, B. R. and Prowse, T. D.: Trends and variability in spring and autumn 0°C-isotherm dates over Canada, *Climatic Change*, 57(3), 341–358, doi:10.1023/a:1022810531237, 2003.
- Boone, A., Masson, V., Meyers, T., and Noilhan, J.: The influence of the inclusion of soil freezing on simulations by a soil–vegetation–atmosphere transfer scheme. *Journal of Applied Meteorology*, 39, 1544–1569. doi: 10.1175/1520-0450(2000)0392.0.CO;2.hal-02269591, 2000.
- Bresson, É., Laprise, R., Paquin, D., Thériault, J. and Elía, R. D.: Evaluating the ability of CRCM5 to simulate mixed precipitation, *Atmosphere-Ocean*, 55(2), 79–93, doi:10.1080/07055900.2017.1310084, 2017.
- Brinkmann, W. and Ashwell, I.: The structure and movement of the chinook in Alberta, *Atmosphere*, 6(2), 1–10, doi:10.1080/00046973.1968.9676548, 1968.
- Bush, E. and Lemmen, D.S. (Eds.): *Canada’s changing climate report*, Government of Canada, Ottawa, Ontario, 444 pp., 2019.
- Carrière, J.-M., Lainard, C., Bot, C. L. and Robart, F.: A climatological study of surface freezing precipitation in Europe, *Meteorological Applications*, 7(3), 229–238, doi:10.1017/s1350482700001560, 2000.
- Changnon, S. A.: Characteristics of ice storms in the United States, *Journal of Applied Meteorology*, 42(5), 630–639, doi:10.1175/1520-0450(2003)042<0630:coisit>2.0.co;2, 2003.
- Cortinas, J.: A climatology of freezing rain in the Great Lakes region of North America, *Monthly Weather Review*, 128(10), 3574–3588, doi:10.1175/1520-0493(2001)129<3574:acofri>2.0.co;2, 2000.
- Dai, A.: Temperature and pressure dependence of the rain-snow phase transition over land and ocean, *Geophysical Research Letters*, 35(12) [online] Available from: <http://onlinelibrary.wiley.com/doi/10.1029/2008GL033295/full>, 2008.
- ECCC (Environment and Climate Change Canada): Historical Climate Data portal, available at: <http://www.climate.weatheroffice.ec.gc.ca>, (last access: October 3, 2019), 2019a.
- ~~Environment Canada: MANOBS (Manual of Surface Weather Observations), (7th ed.), ISBN: 978-1-100-25445-6, Environment Canada, Downsview, Ontario, 2015.~~
- ECCC (Environment and Climate Change Canada): MANOBS (Manual of Surface Weather Observation Standards), 8th Edition, available at: <https://www.canada.ca/en/environment-climate-change/services/weather-manuals-documentation/manobs-surface-observations.html> (last access: October 3, 2019), 2019b.
- ECCC (Environment and Climate Change Canada): Canadian Climate Normals, available at: http://climate.weather.gc.ca/climate_normals/index_e.html (last access: 24 October 2019), 2019c.
- Frei, C., & Schär, C. (2001). Detection probability of trends in rare events: Theory and application to heavy precipitation in the Alpine region. *Journal of Climate*, 14, 1568–1584.
- Fujibe, F.: On the near-0°C. frequency maximum in surface air temperature under precipitation: A statistical evidence for the melting effect. *Journal of the Meteorological Society of Japan*, 79 (3), 731–739, 2001.
- Groenemeijer, P., Becker, N., Djidara, M., Gavin, K., Hellenberg, T., Holzer, A. M., Juga, I., Jokinen, P., Jylhä, K., Lehtonen, I., Mäkelä, H., Napoles, O., Nissen, K., Paprotny, D., Prak, P., Púčik, T., Tijssen L. and Vajda, A.: Past Cases of Extreme Weather

- 605 Impact on Critical Infrastructure in Europe. Office of Critical Infrastructure Protection and Emergency Preparedness. FP7-SEC-2013-1 , 129 pp., 2017.
- Groisman, P. Y., Bulygina, O. N., Yin, X., Vose, R. S., Gulev, S. K., Hanssen-Bauer, I. and Førland, E.: Recent changes in the frequency of freezing precipitation in North America and Northern Eurasia, *Environmental Research Letters*, 11(4), 045007, doi:10.1088/1748-9326/11/4/045007, 2016.
- 610 Hare, K.: *Canada's climate: An overall perspective*. In "The Surface Climates of Canada". McGill-Queen's University Press, 3-20, 1997.
- Henson, W., Stewart, R. and Kochtubajda, B.: On the precipitation and related features of the 1998 Ice Storm in the Montréal area, *Atmospheric Research*, 83(1), 36–54, doi:10.1016/j.atmosres.2006.03.006, 2007.
- Henson, W., Stewart, R., Kochtubajda, B. and Thériault, J.: The 1998 Ice Storm: Local flow fields and linkages to precipitation, 615 *Atmospheric Research*, 101(4), 852–862, doi:10.1016/j.atmosres.2011.05.014, 2011.
- Houston, T. G. and Changnon, S. A.: Freezing rain events: a major weather hazard in the conterminous US, *Natural Hazards*, 40(2), 485–494, doi:10.1007/s11069-006-9006-0, 2007.
- Intergovernmental Panel on Climate Change: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T. F., Qin, D., Plattner, G.-K., 620 Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V. and Midgley, P. M. (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp., doi:10.1017/CBO9781107415324, 2013.
- Jennings, K. S., Winchell, T. S., Livneh, B. and Molotch, N. P.: Spatial variation of the rain–snow temperature threshold across the Northern Hemisphere, *Nature Communications*, 9(1), doi:10.1038/s41467-018-03629-7, 2018.
- Jeong, D. I., Cannon, A. J. and Zhang, X.: Projected changes to extreme freezing precipitation and design ice loads over North 625 America based on a large ensemble of Canadian regional climate model simulations, *Natural Hazards and Earth System Sciences Discussions*, 1–23, doi:10.5194/nhess-2018-395, 2019.
- Kämäräinen, M., Hyvärinen, O., Jylhä, K., Vajda, A., Neiglick, S., Nuottokari, J. and Gregow, H.: A method to estimate freezing rain climatology from ERA-Interim reanalysis over Europe, *Natural Hazards and Earth System Sciences Discussions*, 1–29, doi:10.5194/nhess-2016-225, 2016.
- 630 Kendall, M. B.: *Rank Correlation Methods*. Hafner Publishing Company, New York, 1955.
- Kerry, M., Kelk, G., Etkin, D., Burton, I., Kalhok, S.: Glazed over: Canada copes with the Ice Storm of 1998. *Environment*, 41, 6–11, 1999.
- Kienzie, S. W.: A new temperature based method to separate rain and snow, *Hydrological Processes*, 22(26), 5067–5085, doi:10.1002/hyp.7131, 2008.
- 635 Klaassen, J., Cheng, C. S., Auld, H., Li, Q., Ros, E., Geast, M., Li, G. and Lee, R.: Estimation of Severe Ice Storm Risks for South-Central Canada. Report prepared for Office of Critical Infrastructure Protection and Emergency Preparedness. Meteorological Service of Canada, Environment Canada, Toronto, Canada, 2003.

- Keim, B. D., & Cruise, J. F. (1998). A technique to measure trends in the frequency of discrete random events. *Journal of Climate*, 11, 848–855.
- 640 Kochtubajda, B., Mooney, C. and Stewart, R.: Characteristics, atmospheric drivers and occurrence patterns of freezing precipitation and ice pellets over the Prairie Provinces and Arctic Territories of Canada: 1964–2005, *Atmospheric Research*, 191, 115–127, doi:10.1016/j.atmosres.2017.03.005, 2017.
- Kraatz, S., Jacobs, J. M. and Miller, H. J.: Spatial and temporal freeze-thaw variations in Alaskan roads, *Cold Regions Science and Technology*, 157, 149–162, doi:10.1016/j.coldregions.2018.10.006, 2019.
- 645 Lambert, S. J. and Hansen, B. K.: Simulated changes in the freezing rain Climatology of North America under global warming using a coupled climate model, *Atmosphere-Ocean*, 49(3), 289–295, doi:10.1080/07055900.2011.607492, 2011.
- Larouche, P. and Galbraith, P. S.: Canadian coastal seas and Great Lakes Sea Surface Temperature climatology and recent trends, *Canadian Journal of Remote Sensing*, 42(3), 243–258, doi:10.1080/07038992.2016.1166041, 2016.
- Laute, K. and Beylich, A. A.: Potential effects of climate change on future snow avalanche activity in western Norway deduced from meteorological data, *Geografiska Annaler: Series A, Physical Geography*, 100(2), 163–184, doi:10.1080/04353676.2018.1425622, 2018.
- 650 Lecomte, E. L., Pang, A. W., Russell, J. W.: Ice Storm '98. Institute for Catastrophic Loss Reduction, Toronto, Ontario, Research Paper Series 1, 37 pp., 1998.
- Lindenschmidt, K.-E., Das, A., Rokaya, P. and Chu, T.: Ice-jam flood risk assessment and mapping, *Hydrological Processes*, 30(21), 3754–3769, doi:10.1002/hyp.10853, 2016.
- 655 Longley, R. W.: The frequency of winter chinooks in Alberta, *Atmosphere*, 5(4), 4–16, doi:10.1080/00046973.1967.9676538, 1967.
- MacKay, G. A. and Thompson, H. A.: Estimating the hazard of ice accretion in Canada from climatological data, *Journal of Applied Meteorology*, 8(6), 927–935, doi:10.1175/1520-0450(1969)008<0927:ethoia>2.0.co;2, 1969.
- 660 Matsuo, T., Sasyo, Y. and Sato, Y.: Relationship between types of precipitation on the ground and surface meteorological elements, *Journal of the Meteorological Society of Japan. Ser. II*, 59(4), 462–476, doi:10.2151/jmsj1965.59.4_462, 1981.
- Matsuo, T. and Sasyo, Y.: Empirical formula for the melting rate of snowflakes, *Journal of the Meteorological Society of Japan. Ser. II*, 59(1), 1–9, doi:10.2151/jmsj1965.59.1_1, 1981a.
- Matsuo, T. and Sasyo, Y.: Melting of snowflakes below freezing level in the atmosphere, *Journal of the Meteorological Society of Japan. Ser. II*, 59(1), 10–25, doi:10.2151/jmsj1965.59.1_10, 1981b.
- 665 Matte, D., Thériault, J. M. and Laprise, R.: Mixed precipitation occurrences over southern Québec, Canada, under warmer climate conditions using a regional climate model, *Climate Dynamics*, <https://doi-org.uml.idm.oclc.org/10.1007/s00382-018-4231-2>, 2018.
- McCullagh, P., & Nelder, J. A. (1989). *Generalized linear models*. 2nd ed. Monogr. on Statistics and Appl. Probability, No. 37, Chapman and Hall. London, UK.
- 670

- McFadden, V. and J. M. Thériault, J. M.: Simulations of the ice storm in the Maritime Provinces on 24-26 January 2017, Canadian Meteorological and Oceanographic Society Congress, Halifax, Nova Scotia, 2018.
- McKenney, D. W., Pedlar, J. H., Lawrence, K., Papadopol, P., Campbell, K. and Hutchinson, M. F.: Change and evolution in the plant hardiness zones of Canada, *BioScience*, 64(4), 341–350, doi:10.1093/biosci/biu016, 2014.
- 675 Mekis, E., Donaldson, N., Reid, J., Zucconi, A., Hoover, J., Li, Q., Nitu, R. and Melo, S.: An overview of surface-based precipitation observations at Environment and Climate Change Canada, *Atmosphere-Ocean*, 56(2), 71–95, doi:10.1080/07055900.2018.1433627, 2018.
- Mendoza, G.-P. and Schmunk, R.: Fraser Valley ice storm creates slippery roads, leaves thousands without power | CBC News, CBCNews [online] Available from: [https://www.cbc.ca/news/canada/british-columbia/fraser-valley-winter-storm-december-2017-](https://www.cbc.ca/news/canada/british-columbia/fraser-valley-winter-storm-december-2017-1.4467154)
680 1.4467154, 2017.
- Milton, J. and Bourque, A.: A climatological account of the January 1998 Ice Storm in Québec. Atmospheric Sciences and Environmental Issues Division, Environment Canada, Québec Region, 87 pp., 1999.
- Moen, J. and Fredman, P.: Effects of climate change on alpine skiing in Sweden, *Journal of Sustainable Tourism*, 15(4), 418–437, doi:10.2167/jost624.0, 2007.
- 685 Nitu, R., Roulet, Y.A., Wolff, M., Earle, M., Reverdin, A., Smith, C., Kochendorfer, J., Morin, S., Rasmussen, R., Wong, K., Alastrué, J., Arnold, L., Baker, B., Buisán, S., Collado, J.L., Colli, M., Collins, B., Gaydos, A., Hannula, H.R., Hoover, J., Joe, P., Kontu, A., Laine, T., Lanza, L., Lanzinger, E., Lee, G.W., Lejeune, Y., Leppänen, L., Mekis, E., Panel, J.M., Poikonen, A., Ryu, S., Sabatini, F., Theriault, J., Yang, D., Genthon, C., van den Heuvel, F., Hirasawa, N., Konishi, H., Nishimura, K. and Senese, A.: WMO Solid Precipitation Intercomparison Experiment (SPICE) (2012-2015), World Meteorological Organization
690 Instruments and Observing Methods Report No. 131, Geneva, 2018.
- Nkemdirim, L. C.: Canada's chinook belt, *International Journal of Climatology*, 16(4), 441–462, doi:10.1002/(sici)1097-0088(199604)16:4<441::aid-joc21>3.0.co;2-t, 1996.
- ~~Oke, T. R.: Boundary Layer Climates, Second Edition, Routledge, New York, NY, 435 pp., 1987.~~
- Phillips, D.: The Climates of Canada. Supply and Services Canada, Ottawa, 176 pp., 1990.
- 695 Sen, P. K.: Estimates of the regression coefficient Based on Kendall's tau, *Journal of the American Statistical Association*, 63(324), 1379–1389, doi:10.1080/01621459.1968.10480934, 1968.
- Sims, E. M. and Liu, G.: A parameterization of the probability of snow–rain transition, *Journal of Hydrometeorology*, 16(4), 1466–1477, doi:10.1175/jhm-d-14-0211.1, 2015.
- Stewart, R.E.: Precipitation types in winter storms. *Pure and Applied Geophysics*, 123, 597-609, 1985.
- 700 Stewart, R. E., Isaac, G. A. and Shaw, R. W.: Canadian Atlantic Storms Program: The meteorological field project, *Bulletin of the American Meteorological Society*, 68(4), 338–345, doi:10.1175/1520-0477(1987)068<0338:casptm>2.0.co;2, 1987.
- Stewart, R. and Patenaude, L.: Rain-snow boundaries and freezing precipitation in Canadian East Coast winter storms, *Atmosphere-Ocean*, 26(3), 377–398, doi:10.1080/07055900.1988.9649309, 1988.

- Stewart, R. E. and Yiu, D. T.: Distributions of precipitation and associated parameters across precipitation type transitions over southern Ontario, *Atmospheric Research*, 29(3-4), 153–178, doi:10.1016/0169-8095(93)90002-6, 1993.
- Stewart, R., Bachand, D., Dunkley, R., Giles, A., Lawson, B., Legal, L., Miller, S., Murphy, B., Parker, M., Paruk, B. and Yau, M.: Winter storms over Canada, *Atmosphere-Ocean*, 33(2), 223–247, doi:10.1080/07055900.1995.9649533, 1995a.
- Stewart, R. E. and D. T. Yui and K. K. Chung and D. R. Hudak and E. P. Lozowski and M. Oleskiw and B. E. Sheppard and K. K. Szeto, 1995: Weather conditions associated with the passage of precipitation type transition regions over Eastern Newfoundland, *Atmos. Ocean*, 33, 25-53, 1995b.
- Stuart, R. and Isaac, G.: Freezing precipitation in Canada, *Atmosphere-Ocean*, 37(1), 87–102, doi:10.1080/07055900.1999.9649622, 1999.
- Stewart, R. E., Szeto, K. K., Bonsal, B. R., Hanesiak, J. M., Kochtubajda, B., Li, Y., Thériault, J. M., Debeer, C. M., Tam, B. Y., Li, Z., Liu, Z., Bruneau, J. A., Duplessis, P., Marinier, S. and Matte, D.: Summary and synthesis of Changing Cold Regions Network (CCRN) research in the interior of western Canada – Part 1: Projected climate and meteorology, *Hydrology and Earth System Sciences*, 23(8), 3437–3455, doi:10.5194/hess-23-3437-2019, 2019.
- Takeuchi, Y., Kodama, Y. and Ishikawa, N.: The thermal effect of melting snow/ice surface on lower atmospheric temperature, *Arctic, Antarctic, and Alpine Research*, 34(1), 20, doi:10.2307/1552504, 2002.
- Vincent, L. A., Zhang, X., Brown, R. D., Feng, Y., Mekis, E., Milewska, E. J., Wan, H. and Wang, X. L.: Observed trends in Canada’s climate and influence of low-frequency variability modes, *Journal of Climate*, 28(11), 4545–4560, doi:10.1175/jcli-d-14-00697.1, 2015.
- Vincent, L. A., Zhang, X., Mekis, E., Wan, H. and Bush, E. J.: Changes in Canada’s climate: Trends in indices based on daily temperature and precipitation data, *Atmosphere-Ocean*, 56(5), 332–349, doi:10.1080/07055900.2018.154579, 2018.
- U.S. Army Corps of Engineers: Snow hydrology: Summary report of the snow investigations, North Pacific Division, Portland, Ore. 437 pp., 1956.
- Wang, X. L. (2006), Climatology and trends in some adverse and fair weather conditions in Canada, 1953–2004, *J. Geophys. Res.*, 111, D09105, doi:10.1029/2005JD006155.
- Wasserstein, R. L., Schirm, A. L. and Lazar, N. A.: Moving to a world beyond “ $p < 0.05$ ”. *The American Statistician*. 73, 1-19, 2019.
- Wexler, R., Reed, R. J. and Honig, J.: Atmospheric cooling by melting snow, *Bulletin of the American Meteorological Society*, 35(2), 48–51, doi:10.1175/1520-0477-35.2.48, 1954.
- World Meteorological Organization - International Cloud Atlas Manual on the Observation of Clouds and Other Meteors. (WMO-No. 407), available at: <https://cloudatlas.wmo.int/home.html> (last access: October 3, 2019), 2017.
- Yang, D., Goodison, B. E., Metcalfe, J. R., Golubev, V. S., Elomaa, E., Gunther, T., Bates, R., Pangburn, T., Hanson, C., Emerson, D., Copaciu, V. and Milkovic, J.: Accuracy of Tretyakov precipitation gauge: Result of WMO intercomparison. *Hydrological Processes*, 9(8), 877-895. doi:10.1002/hyp.3360090805, 1995.

Yang, D., Goodison, B. E., Metcalfe, J. R., Golubev, V. S., Bates, R., Pangburn, T. and Hanson, C. L.: Accuracy of NWS 8" standard nonrecording precipitation gauge: Results and application of WMO Intercomparison. *Journal of Atmospheric and Oceanic Technology*, 15(1), 54-68. doi:10.1175/1520-0426(1998)0152.0.co;2, 1998.

740 Zhang, X., Vincent, L. A., Hogg, W. and Niitsoo, A.: Temperature and precipitation trends in Canada during the 20th century, *Atmosphere-Ocean*, 38(3), 395–429, doi:10.1080/07055900.2000.9649654, 2000.

Tables

745 **Table 1: Table of indicators in the $-2 \leq T \leq 2^{\circ}\text{C}$ range. The “T” refers to hourly surface dry bulb temperature. Definitions were obtained from American Meteorological Society (2018), ECCC (2019b) and WMO (2017).**

~~Table of indicators in the $-2 \leq T \leq 2^{\circ}\text{C}$ range.~~

DEFINITIONS	UNIT
Temperature related	
Annual number of days with $-2^{\circ}\text{C} \leq T_{\text{drybulb}} \leq +2^{\circ}\text{C}$	days
Annual number of hours with $-2^{\circ}\text{C} \leq T_{\text{drybulb}} \leq +2^{\circ}\text{C}$	hours
Annual number of independent events (continuous hours) with $-2^{\circ}\text{C} \leq T_{\text{drybulb}} \leq +2^{\circ}\text{C}$	events
Annual Maximum Lengths with $2^{\circ}\text{C} \leq T_{\text{drybulb}} \leq +2^{\circ}\text{C}$	hours
Precipitation types related (occurrences; 12e): Annual number of hours with	
... Rain	hours
... Rain Showers	hours
... Drizzle	hours
... Freezing Rain	hours
... Freezing Drizzle	hours
... Snow	hours
... Snow Grains	hours
... Ice Crystals	hours
... Ice Pellets	hours
... Ice Pellet Showers	hours
... Snow Showers	hours
... Snow Pellets	hours
... all 12 weather types above	hours
Combination of temperature and precipitation type	
Annual number of hours with freezing rain and $-2^{\circ}\text{C} \leq T_{\text{drybulb}} \leq +2^{\circ}\text{C}$	hours
Annual number of hours with any of the 12 precip types and $-2^{\circ}\text{C} \leq T_{\text{drybulb}} \leq +2^{\circ}\text{C}$	hours
The fraction of $[-2^{\circ}\text{C} \leq T_{\text{drybulb}} \leq +2^{\circ}\text{C}]$ conditions associated with Freezing Rain	%
The fraction of $[-2^{\circ}\text{C} \leq T_{\text{drybulb}} \leq +2^{\circ}\text{C}]$ conditions with any of the 12 precip types	%

DEFINITIONS	UNIT
Surface temperature related	
Annual number of days with $-2^{\circ}\text{C} \leq T \leq +2^{\circ}\text{C}$	days
Annual number of hours with $-2^{\circ}\text{C} \leq T \leq +2^{\circ}\text{C}$	hours
Annual number of independent events (continuous hours) with $-2^{\circ}\text{C} \leq T \leq +2^{\circ}\text{C}$	events
Annual Maximum Lengths with $2^{\circ}\text{C} \leq T \leq +2^{\circ}\text{C}$	hours
Precipitation types related (occurrences): Annual number of hours with	
... Rain: Liquid water drops having diameters $> 0.5\text{ mm}$	hours
... Rain Showers: Rainfall where intensity can be variable and may change rapidly	hours
... Drizzle: Liquid water droplets having diameters between $100\text{ }\mu\text{m}$ and 0.5 mm	hours
... Freezing Rain: Rain that freezes upon contact forming a layer of ice on the ground or on exposed objects	hours
... Freezing Drizzle: Drizzle that freezes upon contact forming a layer of ice on the ground or on exposed objects	hours
... Snow: Precipitation of ice crystals singly or agglomerated into snowflakes	hours
... Snow Grains: Very small opaque white particles of ice having diameters $< 1\text{ mm}$	hours
... Ice Crystals: Crystalline forms in which ice appears including hexagonal columns, hexagonal platelets, dendritic crystals, ice needles and combinations of these forms.	hours
... Ice Pellets: Transparent ice particles usually spheroidal or irregular and rarely conical having diameters $< 5\text{ mm}$.	hours
... Ice Pellet Showers: Ice pellets falling where intensity can be variable and may change rapidly	hours
... Snow Showers: Snowfall where intensity can be variable and may change rapidly	hours
... Snow Pellets: White and opaque ice particles, generally conical or rounded having diameters as large as 5 mm	hours
... all 12 precipitation types above	hours
Combination of surface temperature and precipitation type	
Annual number of hours with freezing rain and $-2^{\circ}\text{C} \leq T \leq +2^{\circ}\text{C}$	hours
Annual number of hours with any of the 12 precipitation types and $-2^{\circ}\text{C} \leq T \leq +2^{\circ}\text{C}$	hours
The fraction of $[-2^{\circ}\text{C} \leq T \leq +2^{\circ}\text{C}]$ conditions associated with Freezing Rain	%
The fraction of $[-2^{\circ}\text{C} \leq T \leq +2^{\circ}\text{C}]$ conditions with any of the 12 precipitation types	%

Table 2: Average annual frequencies of the number of the days, hours, events and maximum duration when $-2^{\circ}\text{C} \leq T \leq 2^{\circ}\text{C}$ along (in brackets) with the 31 year trend values based on hourly surface temperature over the 1981-2011 period (minimum 90% of data). The numbers in bold indicate significant changes at 5% level. ~~Table 2: Near 0°C trends based on surface temperature for the 1981-2011 period (minimum 90% of data). Average (Aver) trend values over the 1981-2011 period. An arrow indicates a statistically significant (sign) increase (pointed up) or decrease (pointed down). Stations are arranged from west to east.~~

	Whitehorse	Vancouver	Calgary	Cambridge Bay	Winnipeg	Churchill	Toronto	Montreal	St. John's	Units
Number of Days	139.5 (-10.3)	62.3 (1.8)	161.0 (-2.6)	67.9 (-2)	108.7 (-8.9)	92.6 (-3.1)	121.6 (-26.9)	107.0 (-10.3)	155.0 (-3.6)	days
Number of Hours	1229.9 (100.4)	530.3 (7.2)	1149.1 (46.1)	854.3 (-110.2)	902.3 (0)	1025.7 (-143.6)	1201.8 (-316.9)	979.5 (-69)	1744.5 (90.4)	hours
Number of Events	194.3 (-18.5)	93.9 (3.1)	279.8 (-16.9)	66.5 (0)	145.8 (9.3)	99.7 (0.8)	142.1 (-35.8)	133.7 (-15.5)	163.6 (-15.5)	events
Maximum Duration	58.3 (22.9)	37.2 (-5.2)	39.5 (-6.3)	89.5 (-36.6)	53.8 (8.9)	84.1 (-27.9)	58.0 (-25.4)	52.2 (6.2)	91.0 (7)	hours

	Number of Hours		Number of Days		Number of Events		Maximum Duration	
	Aver	Sign	Aver	Sign	Aver	Sign	Aver	Sign
Whitehorse	100.4	no	-10.3	no	-18.5	no	22.9	no
Vancouver	7.2	no	1.8	no	3.1	no	-5.2	no
Calgary	46.1	no	-2.6	no	-16.9	no	-6.3	no
Cambridge Bay	-110.2	no	-2.0	no	0.0	no	-36.6	no
Winnipeg	0.0	no	-8.9	no	9.3	no	8.9	no
Churchill	-143.6	no	-3.1	no	0.8	no	-27.9	no
Toronto	-316.9	↗	-26.9	↘	-35.8	↗	-25.4	↗
Montreal	-69.0	no	-10.3	no	-15.5	↗	6.2	no
St. John's	90.4	no	-3.6	no	-15.5	no	7.0	no

760 Table 3: Average number of hours with the 12 different precipitation type occurrences along in bracket with the indication of changes
 over the 1981-2011 period. No bracket means no significant change. “>” indicates statistically significant increasing change; “<” indicates
 statistically significant decreasing change; “no” indicates no significant change; “rare” indicates that the given precipitation type rarely
 occurs and the change is not computed; “n/a” indicates that there is not enough data to compute the change. The statistical significance
 of the trends were assessed at 5% level. ~~Table 3: Near 0°C trends based on 12 precipitation type occurrence information for the 1981-
 2011 period (minimum 10 years) Average (Aver) trend values over the 1981-2011 period are shown. An arrow indicates a statistically
 significant (sign) increase (pointed up) or decrease (pointed down) and the last columns refers to the average trend of any of the 12
 precipitation (precip) types and whether these are statistically significant. Stations are arranged from west to east.~~

	Whitehorse	Vancouver	Calgary	Cambridge Bay	Winnipeg	Churchill	Toronto	Montreal	St. John's	Units
Rain	16.3	25.2	12.7	29.9	23.1	60.7	49.9	56.3	83.9	hours
Rain Showers	4.2 (>)	9.6	4.6	15.1 (>)	4.1	9.1 (<)	6.8	5.7	34.5 (<)	hours
Drizzle	2	3.7	6.5	30.2	15.1	41.8 (<)	21.4 (<)	16.2	132.2	hours
Freezing Rain	n/a	2.2 (rare)	n/a	4.9	3.6	10	8.3	13.4	41	hours
Freezing Drizzle	n/a	n/a	4.2	10.9	5.9	15.3 (<)	5.2	6.3 (<)	85.7 (<)	hours
Snow	111.8	41.2	96.2	91.9	79.6	163.3	115.1	152	173.2 (>)	hours
Snow Grains	n/a	1.4 (rare)	1.8	8.6	2.8	5.7	6	2.4	12.8 (<)	hours
Ice Crystal	n/a	n/a	0.0 (rare)	n/a	n/a	0.1	n/a	n/a	n/a	hours
Ice Pellets	n/a	n/a	2.4 (rare)	4.5 (<)	3.6	3.9 (<)	7.6	7.9	16.8	hours
Ice Pellet Showers	n/a	n/a	n/a	n/a	0.8 (rare)	1.9 (rare)	0.9	0.9	1.2	hours
Snow Showers	29.1	16.7	36.5 (<)	68.6 (>)	11.9	44.4 (<)	53.3	40.2	174.2 (<)	hours
Snow Pellets	0.9	n/a	2.5	3.4	1.5	2.6	2.7	1.5	3	hours
Any of the 12 types	154.7	85.6	151.6	250.5	136.3	323.2 (<)	252.6	276.2	686.1	hours

	Rain		Rain Showers		Drizzle		Freezing Rain		Freezing Drizzle		Snow		Snow Grains		Ice Crystal		Ice Pellets		Ice Pellet Showers		Snow Showers		Snow Pellets		All 12 Prec Types	
	Aver	Sign	Aver	Sign	Aver	Sign	Aver	Sign	Aver	Sign	Aver	Sign	Aver	Sign	Aver	Sign	Aver	Sign	Aver	Sign	Aver	Sign	Aver	Sign	Aver	Sign
Whitehorse	-1.8	no	5.6	⬆️	-1.0	no	-	n/a	-	n/a	-45.1	no	-	n/a	-	n/a	-	n/a	-	n/a	24.8	no	-1.7	no	-6.7	no
Vancouver	-8.2	no	3.4	no	0.0	no	-	n/a	-	n/a	-6.4	no	-	n/a	-	n/a	-	n/a	-	n/a	-10.4	no	-	n/a	-10.9	no
Calgary	5.2	no	-3.2	no	0.0	no	-	n/a	0.0	no	44.4	no	0.0	no	-	n/a	-	n/a	-	n/a	-69.3	⬆️	-2.6	no	-5.2	no
Cambridge Bay	7.6	no	30.2	⬆️	-18.6	no	0.0	no	-5.9	no	16.4	no	-2.1	no	-	n/a	-4.0	⬆️	-	n/a	78.1	⬆️	-0.8	no	85.6	no
Winnipeg	-0.6	no	0.0	no	-5.4	no	0.0	no	-1.1	no	-10.5	no	1.1	no	-	n/a	0.4	no	-	n/a	8.5	no	0.0	no	8.1	no
Churchill	-1.1	no	-9.4	⬆️	-70.6	⬆️	0.0	no	-20.7	⬆️	12.7	no	-17.5	no	0.0	no	-7.8	⬆️	-	n/a	-54.9	⬆️	-2.8	no	-144.5	⬆️
Toronto	-13.5	no	-6.7	no	-21.7	⬆️	-3.9	no	-4.4	no	-9.7	no	1.3	no	-	n/a	-2.2	no	0.0	no	-14.2	no	-4.6	no	-95.6	no
Montreal	0.0	no	-4.9	no	-6.9	no	-7.4	no	-8.9	⬆️	0.0	no	-11.3	no	-	n/a	-2.8	no	0.0	no	-17.7	no	-3.3	no	-24.8	no
St. John's	15.5	no	-48.1	⬆️	-31.0	no	-18.9	no	-66.6	⬆️	112.7	⬆️	-20.8	⬆️	-	n/a	-0.1	no	0.0	no	-14.6	⬆️	-3.9	no	-51.1	no

770


Table 4: Average monthly surface temperatures (°C) and trends (days/31 years) of the onset of near 0°C conditions in autumn and cessation in spring for the period 1981-2011. For Cambridge Bay (*), average monthly values refer to June, July and August and trends refer to the cessation of near 0°C conditions in autumn and onset in spring. No calculations were carried out at Churchill. Average monthly surface temperatures were obtained for the 1981-2010 period from ECCC (2019c). The numbers in bold indicate trends significant at the 5% level.

Station	Average Monthly Temperature (°C)			Autumn Onset Trend (days/31 years)	Spring Cessation Trend (days/31 years)
	December	January	February		
Whitehorse	-12.5	-15.2	-12.7	6.4	-11.8
Vancouver	3.6	4.1	4.9	8.3	-4.1
Calgary	-6.8	-7.1	-5.4	5.4	1.1
Cambridge Bay*	2.7	8.9	6.8	23.8	0.2
Winnipeg	-13.2	-16.4	-13.2	4	-2.4
Churchill	-21.9	-26	-24.5	-	-
Toronto	-2.2	-5.5	-4.5	25.1	-15.4
Montreal	-5.4	-9.7	-7.7	9.6	-0.8
St. John's	-1.5	-4.5	-4.9	32.3	-10.9

Table 45: The longest duration events at the 9 ~~representative-selected~~ stations. Columns indicate maximum duration, start time, and hours (and fraction of duration) with mainly or completely clear ~~skies~~. ~~Times are UTC.~~sky conditions, hours with any of the 12 reported precipitation types (and fraction of duration). Times are local standard.

Station	Duration	Start Time	Mainly/Completely Clear		Precipitation	
	(h)	(YYYY-MM-DD-HH)	(h)	(%)	(h)	(%)
Whitehorse	110	1998-10-06-21	3.0	2.7	12.0	10.9
Vancouver	68	2005-01-06-04	1.0	1.5	44.0	64.7
Calgary	73	2003-05-05-10	0.0	0.0	60.0	82.2
Cambridge Bay	197	1987-06-11-08	3.0	1.5	63.0	32.0
Winnipeg	128	1983-03-01-15	0.0	0.0	106.0	82.8
Churchill	141	1986-10-18-12	1.0	0.1	98.0	69.5
Toronto	158	1986-12-26-16	0.0	0.0	13.0	8.2
Montreal	114	2007-12-24-12	8.0	7.0	34.0	29.8
St. John's	148	1983-04-01-06	14.0	9.5	75.0	50.7

Station	Duration	Start Time	Mainly/Completely Clear	
	(h)	(YYYY-MM-DD-HH)	(h)	(%)
Whitehorse	110	1998-10-06-21	3.0	2.7
Vancouver	68	2005-01-06-04	1.0	1.5
Calgary	73	2003-05-05-10	0.0	0.0
Cambridge Bay	197	1987-06-11-08	3.0	1.5
Winnipeg	128	1983-03-01-15	0.0	0.0
Churchill	141	1986-10-18-12	1.0	0.1
Toronto	158	1986-12-26-16	0.0	0.0
Montreal	114	2007-12-24-12	8.0	7.0
St. John's	148	1983-04-01-06	14.0	9.5








Figure 1: The average annual surface temperature distribution (4°C bins) from 1981 to 2011 for 9 representative stations across Canada as shown in Fig. 2. The vertical black line indicates the average annual surface temperature using information from ECCC (2019a). The red bar identifies near 0°C surface temperatures defined as $-2 \leq T \leq 2^\circ\text{C}$. Stations are arranged from west to east. Details on the temperature data are in Sect. 2.




795 **Figure 2: The 92 stations used in the analysis (see text for details). Blue ellipses and red crosses show the 9 representative selected stations**
across Canada. British Columbia region includes all stations in British Columbia (BC); Prairies region - all stations in Alberta (AB),
Saskatchewan (SK) and Manitoba (MB); Ontario region - all stations in Ontario (ON); Quebec region - all stations in Quebec (QC);
Atlantic region - all stations in New Brunswick (NB), Prince Edward Island (PE), Nova Scotia (NS), Newfoundland and Labrador (NL-
L); and Northern region - all stations in Yukon (YK), Northwest Territories (NT) and Nunavut (NU). **The average annual 0°C, -2°C and**
800 **2°C surface temperature lines, computed from 1981-2010 climate normals of 1619 locations (ECCC, 2019c), are also shown.**


a) Number of days




b) Number of hours




c) Number of events




d) Maximum duration




a) Number of days



b) Number of hours



c) Number of events



d) Maximum duration






Figure 3: Average near 0°C ($-2 \leq T \leq 2^\circ\text{C}$) conditions over the 1981-2011 period for (a) number of days per year, (b) number of hours per year, (c) number of events per year, and the (d) annual maximum (max) duration of events. The red line indicates the annual average 0°C surface temperature computed from 1981-2010 climate normals (ECCC, 2019c).


a) Any 12 weather type hours




b) Any 12 weather type percent




c) Freezing Rain hours




d) Freezing Rain percent




a) Any 12 weather type hours



b) Any 12 weather type percent



c) Freezing Rain hours



d) Freezing Rain percent






Figure 4: Precipitation type occurrences during near 0°C ($-2 \leq T \leq 2^\circ\text{C}$) conditions over the 1981-2011 period, where (a) average annual number of hours with reported precipitation types (any of the 12), (b) percentage of near 0°C conditions associated with reported precipitation types (any of the 12), (c) average annual number of hours with freezing rain, and (d) percentage of near 0°C conditions associated with freezing rain. The red line indicates the annual average 0°C surface temperature computed from 1981-2010 climate normals (ECCC, 2019c).

a) Number of days




b) Number of hours



c) Number of events



d) Maximum duration







Figure 5: Trends in near 0°C ($-2 \leq T \leq 2^\circ\text{C}$) conditions over the 1981-2011 period. (a) annual average number of days, (b) annual average hours, (c) annual number of events, and (d) annual maximum duration. A solid triangle indicates statistical significance at 5% level. The red line indicates the annual average 0°C surface temperature computed from 1981-2010 climate normals (ECCC, 2019c).



820





Figure 6: Cumulative distribution of events (%) as a function of duration (h) of near 0°C ($-2 \leq T \leq 2^\circ\text{C}$) events at the 9 representative stations across Canada over the 1981-2011 period arranged from west to east. The total number of events is shown in brackets also indicated and duration is plotted on a logarithmic scale.






Figure 7: The distribution of precipitation type occurrence with near 0°C conditions at each of the 9 stations over the 1981-2011 period. Stations are arranged from west to east.





Figure 8: The 1981-2011 values of average (a) annual number of days, (b) annual number of hours, (c) annual number of events and (d) annual maximum duration (in hours) during near 0°C ($-2 \leq T \leq 2^\circ\text{C}$) conditions for the nine selected locations in Canada. Linear trend lines are superimposed in red.








Figure 98: a) Average monthly occurrence (hours) of near 0 $^{\circ}\text{C}$ conditions as a function of average monthly surface temperature for the nine representative-selected stations over the 1981-2011 period. The open diamond symbol indicates no occurrence. Filled symbols indicate occurrence and open or unfilled symbols indicate no occurrence. b) An expanded view to better illustrate low values of average monthly occurrence.





845 **Figure 109:** The occurrence of near 0°C conditions and any (of the 12) associated precipitation types at (a) Vancouver, (b) Cambridge Bay (c) Toronto and (d) St. John's over the 1981-2011 period. Shading refers to the weekly fraction (%) of near 0°C hours with (or without) precipitation, the '0+' symbol refers to at least one hour of precipitation whereas the gray '0' means no precipitation even if the near 0°C criterion was met. Blank areas indicate no occurrence of near 0°C conditions. The vertical, dashed line indicates January 1. Blue and red bars indicate hourly occurrences of temperature and associated precipitation types (any of the 12), respectively. The vertical, dashed line indicates January 1.

850




Figure 11: Average annual number of near 0°C events at the 9 selected stations organized by surface temperature pathways. The four pathways are shown schematically below the figure. They are represented by bars at each station from pathway 1 to 4 (left to right). No shading refers to the occurrence of near 0°C conditions, dark shading refers to the occurrence of freezing rain near 0°C, and light grey refers to the occurrence of all other types of precipitation near 0°C.