



BESS-STAIR: a framework to estimate daily, 30-meter, and all-weather crop evapotranspiration using multi-source satellite data for the U.S. Corn Belt

Chongya Jiang^{1,2}, Kaiyu Guan^{1,2,3}, Ming Pan⁴, Youngryel Ryu⁵, Bin Peng^{1,3}, Sibow Wang³

5 ¹College of Agricultural, Consumer and Environmental Sciences, University of Illinois at Urbana Champaign, Urbana, Illinois, USA

²Center for Advanced Bioenergy and Bioproducts Innovation, University of Illinois at Urbana Champaign, Urbana, Illinois, USA

³National Center of Supercomputing Applications, University of Illinois at Urbana Champaign, Urbana, Illinois, USA

10 ⁴Department of Civil and Environmental Engineering, Princeton University, New Jersey, USA

⁵Department of Landscape Architecture and Rural Systems Engineering, Seoul National University, Seoul, Republic of Korea

Correspondence to: Chongya Jiang (chongya.jiang@email.com)

15 **Abstract.** With increasing crop water demands and drought threats, mapping and monitoring of cropland evapotranspiration (ET) at high spatial and temporal resolutions becomes increasingly critical for water management and sustainability. However, estimating ET from satellite for precise water resources management is still challenging due to the limitations in both existing ET models and satellite input data. Specifically, the process of ET is complex and difficult to model, and existing satellite remote sensing data could not fulfill high resolutions in both space and time. To address the above two

20 issues, this study presented a new high spatiotemporal resolution ET mapping framework, i.e., BESS-STAIR, which integrates a satellite-driven water-carbon-energy coupled biophysical model BESS (Breathing Earth System Simulator) with a generic and fully-automated fusion algorithm STAIR (SaTallite dAta IntegRation). In this framework, STAIR provides daily 30-meter multispectral surface reflectance by fusing Landsat and MODIS satellite data to derive fine-resolution leaf area index and visible/near-infrared albedo, all of which, along with coarse-resolution meteorological and CO₂ data, are used

25 to drive BESS to estimate gap-free 30-m resolution daily ET. We applied BESS-STAIR from 2000 through 2017 in six areas across the U.S. Corn Belt, and validated BESS-STAIR ET estimations using flux tower measurements over 12 sites (85 site-years). Results showed that BESS-STAIR daily ET achieved an overall $R^2 = 0.75$, with RMSE = 0.93 mm d⁻¹ and relative error = 27.9% when benchmarked with the flux measurements. In addition, BESS-STAIR ET estimations well captured the spatial patterns, seasonal cycles, and interannual dynamics in different sub-regions. The high performance of the BESS-

30 STAIR framework is primarily resulted from: (1) the implementation of coupled constraints on water, carbon, and energy in BESS, (2) high-quality daily 30-m data from STAIR fusion algorithm, and (3) BESS's applicability under all-sky conditions. BESS-STAIR is calibration-free and has great potentials to be a reliable tool for water resources management and precision agriculture applications for the U.S. Corn Belt, and even for worldwide given the global coverage of its input data.



1 Introduction

35 Accurate field-level management of water resources urgently demands reliable estimations of evapotranspiration (ET) at high spatial and temporal resolutions. ET is the sum of water loss from soil surface through evaporation and that from plant components through transpiration, and ET at cropland is usually considered as crop water needs (Allen et al., 1998). ET consumes up to 90% of total water inputs (precipitation plus irrigation) in agro-ecosystems in the Western and Midwestern United States (Irmak et al., 2012). In the U.S. Corn Belt, increasing vapor pressure deficit (VPD) and drought sensitivity has
40 been recognized as severe threats to future crop security (Lobell et al., 2014; Ort and Long, 2014). The vulnerability to drought in this region is further exacerbated by elevated rates of grass-to-crop conversion and expansion of irrigated areas (Brown and Pervez, 2014; Wright and Wimberly, 2013). Furthermore, precision water resources management requires capacity to account for spatial heterogeneity and to guide real-time decision-making (USDA, 1997). Accordingly, reliable tools are urgently needed to estimate, map and monitor the total amount and spatial and temporal variations of cropland ET.

45 One critical requirement for the accurate estimations of ET at high spatiotemporal resolutions is reliable and advanced satellite-based models. This is challenging because the process of ET is complex and difficult to model. ET results from competitions between atmospheric water demand and soil water supply, and it is also regulated by plants through canopy development and stomatal behaviors in order to optimize their water, carbon and energy use strategies (Katul et al., 2012; Wang and Dickinson, 2012). A large number of satellite-based ET estimation methods have been developed based on different theories and techniques. In general, they can be grouped into many categories: statistical or machine learning methods (Jung et al., 2010; Lu and Zhuang, 2010), water balance methods (Pan et al., 2012; Wan et al., 2015), energy balance methods (Anderson et al., 1997; Su, 2002), triangular or trapezoid space methods (Jiang and Islam, 1999; Li et al., 2009), Priestley–Taylor methods (Fisher et al., 2008; Miralles et al., 2011), and Penman–Monteith methods (Mu et al., 2011; Yebra et al., 2013). Kalma et al. (2008), Li et al. (2009) and Zhang et al. (2016) have provided detailed reviews on the pros
55 and cons of different remote sensing approaches.

Given the complexity of the ET process, we argue that a reliable ET model should include both necessary biophysical processes and high-quality multi-source observations to constrain ET estimations (Loew et al., 2016). While remote sensing-
60 based approaches tend to focus on constraints from various satellite data, land surface models (LSMs) are proficient to include processes that account for interactions between environment and plant structure and functions. Given the gaps between remote sensing and LSMs, a distinct ET model, the Breathing Earth System Simulator (BESS), was developed (Jiang and Ryu, 2016; Ryu et al., 2011). Different from the above-mentioned remote sensing models, BESS is a biophysical model, which adopts modules commonly-implemented in land surface models but uses various satellite remote sensing data
65 as direct inputs. Specifically, BESS is a two-leaf water-carbon-energy coupled model driven by environmental and vegetation variables derived from multi-source satellite data. As the energy cycle, carbon cycle and water cycle are jointly modeled and mutually constrained in BESS, it has produced a series of high-quality global long-term (2000–2017) products,



including the 5-km resolution global radiation (R_g) and photosynthetically active radiation (PAR) and diffuse PAR products (Ryu et al., 2018), and 1-km resolution gross primary productivity (GPP) and ET products (Jiang and Ryu, 2016), which
70 enables tracking crop growth and yields too (Huang et al., 2018). In particular, the 1-km resolution BESS ET product is able to capture the total amount and spatial and temporal variations in arid/semi-arid areas like Australia (Whitley et al., 2016, 2017), California (Baldocchi et al., 2019) and Northwestern China (Wei et al., 2019). The fidelity of coarse-resolution BESS ET product suggests its potential at fine resolutions.

75 The other critical requirement for accurate estimations of ET at high spatiotemporal resolutions is satellite input data at high resolutions in both space and time. This is challenging because existing satellite missions cannot satisfy the two conditions simultaneously. Data fusion techniques, which take multi-sensor data to generate fusion data with high resolutions in both space and time, provide a possible and scalable solution. Several such algorithms have been developed over the past decade (Gao et al., 2006; Houborg and McCabe, 2018; Zhu et al., 2010), and they have been successful for localized applications
80 (Gao et al., 2017; Gómez et al., 2016; Wu et al., 2015). Notably, energy balance and thermal-based ET models such as ALEXI/DisALEXI and SEBS have been combined with the fusion algorithm such as STARFM and ESTARFM to generate daily 30-m ET estimations with favourable performance at several sites (Anderson et al., 2018; Cammalleri et al., 2013; Li et al., 2017; Ma et al., 2018).

85 Here we propose and present a new ET estimation framework that combines BESS with a novel fusion algorithm SaTallite dAta IntegRation (STAIR) (Luo et al., 2018), for accurate ET estimation at both high resolution in time and space. BESS has demonstrated its high performance in estimating ET at medium to coarse resolutions, but the major obstacle of moving BESS's ET estimation to finer resolutions is the lack of key vegetation status variables at higher spatial resolutions, including leaf area index (LAI), and visible and near-infrared albedo (α_{VIS} and α_{NIR}). In BESS, these surface information are
90 critical to resolving spatial heterogeneity, while environmental information such as radiation, temperature, humidity and CO₂ concentration are relatively homogeneous. To cope with the absence of high spatiotemporal resolution vegetation data, we propose to couple STAIR with BESS. STAIR is a genetic and fully-automated fusion algorithm to generate cloud-/gap-free surface reflectance product in high spatiotemporal resolution (Luo et al., 2018). Instead of manually selecting image pairs adopted by most other data fusion algorithms, STAIR automatically takes full advantage of time-series of daily coarse-
95 resolution images and fine-resolution but less frequent images. Moreover, STAIR's high efficiency in computation allows scalability for large scale productions, which enable this new framework to deliver daily 30-m ET at regional and decadal scales.


The objective of this study is to address a fundamental issue in agro-ecological science and applications: lack of high
100 spatiotemporal gap-free ET data for decision-making. We implemented a new ET estimation framework BESS-STAIR and tested it at six study areas across the U.S. Corn Belt from 2000 to 2017. This is the first attempt to couple a satellite-driven



105 LSM with data fusion technique to provide daily 30m-resolution ET estimations at regional and decadal scales. While existing frameworks retrieve clear-sky ET from satellite-observed LST and fill ET gaps for cloudy-sky days, BESS-STAIR simulates all-sky ET and LST as a result of crop biophysical properties. This manner has more referential significance for crop modeling studies and has potential of breaking a new path to agro-ecological science and applications. We conducted comprehensive evaluation on the BESS-STAIR ET estimations with regards to the overall performance, spatial patterns, seasonal cycles and interannual dynamics, benchmarked on the ET observations from 12 eddy-covariance flux towers across the U.S. Corn Belt. The paper also discussed on the performance, advantages, limitations and potential improvements of the BESS-STAIR ET framework.

110

2 Materials and methods



115 Figure 1. The BESS-STAIR framework. The BESS ET estimation model and the STAIR data fusion algorithm are highlighted in green boxes. Blue boxes are satellite data, yellow boxes are ancillary data, and red boxes are key inputs to BESS. The output of BESS-STAIR is the 30-m resolution daily ET highlighted in white box.

120 BESS-STAIR estimates cropland ET at 30-m resolution at daily interval (Figure 1). BESS is driven by environmental variables (radiation, temperature, humidity, and CO₂ concentration), plant structural variables (LAI, α_{VIS} and α_{NIR}), and plant functional variables (peak maximum carboxylation rate at 25 °C (peak V_{cmax25}) and Ball-Berry coefficients, for C3 and C4 plants, respectively). Among these key inputs, LAI, α_{VIS} and α_{NIR} characterize crop canopy structure, which are usually very



heterogeneous. In the global BESS ET product (Jiang and Ryu, 2016), these vegetation variables are derived from MODIS satellite data at 1-km resolution; while in BESS-STAIR, they are derived from 30-m resolution surface reflectance fused from high spatial resolution Landsat data and high temporal resolution MODIS data by STAIR.

125 2.1 The ET estimation model: BESS

BESS is a sophisticated satellite-driven water-carbon-energy coupled biophysical model designed to continuously monitor and map water and carbon fluxes (Jiang and Ryu, 2016; Ryu et al., 2011). It is a simplified land surface model, including an atmosphere radiative transfer module (Kobayashi and Iwabuchi, 2008; Ryu et al., 2018), a two-leaf canopy radiative transfer module (De Pury and Farquhar, 1997), and an integrated carbon assimilation – stomatal conductance – energy balance
130 module. Specifically, the Farquhar model for C3 and C4 plants (Collatz et al., 1992, 1991), the Ball-Berry model (Ball et al., 1987), and the quadratic form of the Penman-Monteith equation (Paw U and Gao, 1988) are used for the simulation of carbon assimilation, stomatal conductance and energy balance, respectively. This carbon-water integrated module employs an iterative procedure to solve intercellular CO₂ concentration, stomatal conductance and leaf temperature for sunlit and shade canopy. Instantaneous sunlit/shade GPP and sunlit/shade/soil ET and net radiation at Terra and Aqua overpass times
135 are simultaneously estimated, followed by a temporal upscaling procedure to derive daily GPP and ET using semi-empirical cosine functions (Ryu et al., 2012). The Priestley-Taylor equation is used to compute daily potential ET (PET) based on daily net radiation and meteorological data.

A unique feature of BESS is that BESS takes full advantages of atmospheric and land products derived from multi-source
140 satellite data. By using MOD/MYD 04 aerosol products (Sayer et al., 2014), MOD/MYD 06 cloud products (Baum et al., 2012), MOD/MYD 07 atmospheric profile products (Seemann et al., 2003), along with gap-free atmospheric data provided by MERRA-2 reanalysis products (Gelaro et al., 2017), BESS calculates direct/diffuse visible/near-infrared radiation components at 0.05° resolution. By coupling CO₂ concentration derived from SCIAMACHY and GOSAT satellite data (Dils et al., 2014) with those from OCO-2 satellite data (Hammerling et al., 2012), as well as NOAA long-term field observations
145 (www.esrl.noaa.gov/gmd/ccgg/trends/), BESS derives long-term continuous monthly CO₂ concentration maps. Finally, in this study BESS uses air temperature and dew point temperature provided by ERA5 reanalysis products at 0.1° resolution (Hersbach and H., 2016). In addition to these environmental variables, BESS also highly relies on vegetation structural and functional variables. By using satellite-derived LAI, α_{VIS} and α_{NIR} , BESS quantifies the absorption of ultraviolet/visible/near infrared radiation by sunlit/shaded canopy through a canopy radiative transfer model. This model also upscales leaf level
150 (V_{cmax25}) to sunlit/shade canopy, which is used in the Farquhar photosynthesis model. V_{cmax25} is a parameter depending on the plant functional type (Bonan et al., 2011; Kattge et al., 2009), and its seasonal variation is empirically parameterized by LAI (Ryu et al., 2011).



2.2 The data fusion algorithm: STAIR

155 STAIR is a generic and fully-automated method for fusing multi-spectral satellite data to generate high spatiotemporal
resolution and cloud-/gap-free data (Luo et al., 2018). It fully leverages the complementary strengths in the high temporal
resolution MCD43A4 nadir reflectance (daily but 500 m resolution) (Schaaf et al., 2002) and the high spatial resolution
Landsat L2 nadir reflectance (30-m resolution but 16-day revisiting frequency) (Masek et al., 2006) time series data. STAIR
first imputes the missing pixels using an adaptive-average correction procedure, and then employs a local interpolation
160 model to capture finer spatial information provided by Landsat data, followed by a time-series refinement step that
incorporates the temporal patterns provided by MODIS data. This strategy allows higher efficiency in missing-data
interpolation as well as greater robustness against concurrently missed MODIS and Landsat observation, which is a common
situation during continuous cloudy/snowy days.

165 The algorithm starts from the imputation of the missing pixels (due to cloud cover or Landsat 7 Scan Line Corrector failure)
in satellite images. For MODIS images, a Savitzky-Golay filter is first applied to reconstruct continuous time series. For
Landsat images, a two-step approach is employed using both temporal and spatial information from clear-sky observations.
First, a temporal interpolation through a linear regression is applied as the initial gap-filling, based on the whole time series
of images throughout a year. Second, an adaptive-average correction procedure is applied to remove inharmonic spatial
170 patterns between gap-filled and original data. The target image is partitioned into multiple segments, each of which contains
one type of homogeneous pixels. The relative difference between a gap pixel and neighbourhood pixels of it within the same
segment is calculated using clear-sky observations acquired in several dates close to the target image acquisition date. Based
on the assumption that the relative difference remains roughly the same across different dates in a short time period (e.g., <
2–3 weeks), such difference is used to correct the filled values of the gap pixel derived from temporal interpolation so that
175 the spatial relationship between the gap-filled pixel and its neighbourhood pixels within the same segment is consistent with
those in clear-sky observations.

The STAIR fusion algorithm fully exploits the spatial and temporal information in the time series of gap-filled MODIS and
Landsat images throughout the growing season (April - October). A nearest neighbour sampling is conducted for all the
180 MODIS images to achieve the same image size, pixel resolution and projected coordinate system with Landsat images.
Difference image is calculated for each pair of Landsat and resampled MODIS images, and a linear interpolation is applied
to reconstruct the difference image for any given date when no Landsat image is available. Such difference image is used to
correct the resampled MODIS image on that date and to generate a fused Landsat image. In this manner, the fused image
captures the most informative spatial information provided by the high spatial resolution Landsat data and incorporates the
185 temporal patterns provided by the high temporal resolution MODIS data without any user interference. The fusion algorithm
is applied to the six Landsat bands: blue, green, red, near-infrared (nir), shortwave infrared 1 (swir1), and shortwave
infrared-2 (swir2).



2.3 Derivation of BESS inputs from STAIR data

190 At global scale, LAI, α_{VIS} and α_{NIR} can be obtained from MODIS and other satellite data, but for field-scale agricultural applications high spatial resolution data are needed to account for the spatial heterogeneity between fields or within a field. At this point, we employed two approaches to estimate 30-m resolution daily LAI from STAIR fused surface reflectance data: an empirical approach based on linear relationship with vegetation indices (VIs) and a mechanic approach based on inversion of a canopy radiative transfer model (RTM).

195

First, we estimated LAI using the empirical approach, because of availability of field LAI measurements in the study area. We calculated four VIs calculated from STAIR-derived spectral reflectance: Wide Dynamic Range Vegetation Index (WDRVI), Green Wide Dynamic Range Vegetation Index (GWDRVI), Enhanced Vegetation Index (EVI), and Land Surface Water Index (LSWI) for corn and soybean, respectively (Eq. (1) – (3)). These four VIs were chosen because they utilized
200 information from different band combinations.

$$WDRVI = \frac{0.1\rho_N - \rho_R}{0.1\rho_N + \rho_R} \quad (1)$$

$$GWDRVI = \frac{0.1\rho_N - \rho_G}{0.1\rho_N + \rho_G} \quad (2)$$

$$EVI = 2.5 \frac{\rho_N - \rho_R}{\rho_N + 6\rho_R - 7.5\rho_B + 1} \quad (3)$$

$$LSWI = \frac{\rho_N - \rho_{SW1}}{\rho_N + \rho_{SW2}} \quad (4)$$

where ρ_B , ρ_G , ρ_R , ρ_N , and ρ_{SW1} refer to the surface reflectance at blue, green, red, near-infrared, the first shortwave-infrared band, respectively. Subsequently, we used field measured LAI data collected using destructive method at Mead, Nebraska from 2001 through 2007 to build VI-LAI relationships (Gitelson et al., 2007). For each of the four VIs we build a linear regression between time series of VI and LAI for corn, soybean, and the combination of corn and soybean, respectively. At
205 this point, the equation derived from the combination of corn and soybean was used for vegetation cover other than corn and soybean. Although this might cause bias for forest LAI estimation, it is not a concern in this study as we focused on crop ET only. We applied linear regressions to four VIs separately and averaged the four derived LAI as the final LAI estimation, with the expectation that such average would reduce uncertainty caused by individual VI-LAI relationship.

210 Second, we inversed PROSAIL RTM using a look-up table (LUT) method. PROSAIL is an efficient and widely-used model to simulate canopy reflectance given a set of sun-object-view geometry, canopy structure, leaf biochemical, and soil optical parameters (Jacquemoud et al., 2009). It is a combination of the PROSPECT leaf hyperspectral properties model (Jacquemoud et al., 1996; Jacquemoud and Baret, 1990) and the SAIL canopy bidirectional reflectance model (Verhoef,



1984, 1985). PROSAIL is particularly suitable for grasslands and croplands (Darvishzadeh et al., 2008; Xu et al., 2019), and
215 therefore used in this study. LUT is a robust and easy method to retrieve model parameters from observed canopy reflectance
(Verrelst et al., 2018). It is based on the generation of simulated canopy reflectance database for a number of plausible
combinations of model parameter value ranges, and the identification of parameter values in the database leading to the best
agreement between simulated and observed canopy reflectance. LUT is particularly suitable for big data processing (Myneni
et al., 2002), and therefore used in this study.

220

We established a database by running PROSAIL with sampled parameter values listed in Table 1. For computation
efficiency, we only sampled varied values for four parameters while others were fixed. These four free parameters, including
LAI (10 values), fraction of vegetation cover (6 values), soil brightness (5 values) and chlorophyll content (4 values), were
chosen because they have been identified as the most sensitive parameters in canopy radiative transfer models (Bacour et al.,
225 2002; Mousivand et al., 2014). Leaf inclination distribution function is also sensitive but we set fixed types "spherical",
"planophile" and "plagiophile" for corn, soybean, and other biomes, respectively (Nguy-Robertson et al., 2012; Pisek et al.,
2013). The fixed values of other parameters were set according to literature (Baret et al., 2007; Feret et al., 2008;
Jacquemoud et al., 2009). Solar zenith angle at satellite overpass time can be calculated so we did not set it as a free
parameter. Instead, we built a set of databases with solar zenith angle values (°) of 20, 25, 30, 35, 40, 45 and 50, respectively,
230 representing the range during growing season in the study area. In PROSAIL, specific absorption coefficients and refractive
index of leaf material are pre-measured hyperspectral data from 400 to 2500 nm with 1 nm interval (Feret et al., 2008), we
averaged them over wavelengths to match Landsat 7 bands and assumed differences of spectral ranges between Landsat 5,
Landsat 7 and Landsat 8 have marginal influence on LAI retrieval. We did not use default soil spectrum in PROSAIL, but
spatiotemporally averaged all cropland pixels spectral reflectance in April when no crop is planted across the study area to
235 derive representative soil spectral reflectance.

Table 1. Parameter values needed to establish the canopy reflectance database by PROSAIL.

Parameters	Values
LAI (m ² m ⁻²)	0, 0.5, 1, 2, 3, 4, 5, 6, 7, 8
Fraction of vegetation cover (m ² m ⁻²)	0, 0.2, 0.4, 0.6, 0.8, 1
Soil brightness (a.u.)	0.01, 0.4, 0.8, 1.2, 1.6
Chlorophyll content (ug cm ⁻²)	0, 20, 40, 60
Leaf inclination distribution function	spherical for corn, planophile for soybean, plagiophile for others
Structure coefficient (a.u.)	1.75
Carotenoid content (ug cm ⁻²)	0
Equivalent Water Thickness (cm)	0.015
Leaf Mass per area (g cm ⁻²)	0.0075



Brown pigment content (a.u.)	0
Hot spot parameter (a.u.)	0.1
View zenith angle (°)	0
Azimuth angle (°)	0

To retrieve LAI, we compared STAIR-derived surface reflectance (R_{STAIR}) with records in the canopy reflectance database simulated by PROSAIL ($R_{PROSAIL}$) pixel by pixel. We used root mean square error (RMSE) as the cost function which was defined as:

$$RMSE = \sqrt{\frac{1}{l} \sum_{\lambda=1}^l [R_{STAIR}(\lambda) - R_{PROSAIL}(\lambda)]^2}, \quad (5)$$

where $\lambda = 1, 2, \dots, l$ indicates band number and $l = 6$ for STAIR. Ideally, the simulated reflectance in the database yielding the smallest RMSE can be considered as the best simulation, and the corresponding LAI value can be considered as the solution for the satellite pixel. However, in reality the solution might not be unique, because different parameter combinations could derive similar reflectance simulations and errors in both satellite and model could further amplify this problem (Verrelst et al., 2018). For this reason, we chose top 10% small RMSE simulations in the database and considered the average of corresponding LAI values as the final solution. The threshold 10% was decided by evaluating LUT-retrieved LAI against field-measured LAI at three Mead sites, and it was within a reasonable range from top 50 records to top 20% records suggested by previous studies (Duan et al., 2014; Weiss et al., 2000).

250

We further employed semi-empirical equations to calculate α_{VIS} and α_{NIR} (Liang, 2001) from STAIR-derived spectral reflectance in six Landsat bands:

$$\alpha_{VIS} = 0.443\rho_B + 0.317\rho_G + 0.240\rho_R \quad (6)$$

$$\alpha_{NIR} = 0.693\rho_N + 0.212\rho_{SW1} + 0.116\rho_{SW2} - 0.003 \quad (7)$$

where ρ_{SW2} is the surface reflectance at the second shortwave-infrared band. CI was set 0.75 for herbaceous and 0.70 for woody plants according to the global mean value of different plant functional types (He et al., 2012). For C3 crops/grasses, forests, and C4 crops/grasses, peak V_{cm^2} values were set 180, 60 and 45, respectively (Kattge et al., 2009; Zhang et al., 2014). Ball-Berry slope and intercept are another two important parameters used in the stomatal conductance model, and their values were set 13.3 and 0.02 for C3 crops/grasses, 9.5 and 0.005 for forests, and 5.8 and 0.04 for C4 crops, respectively (Miner et al., 2017). Distributions of C3 and C4 crops were obtained from Crop Data Layer (CDL) data (Boryan et al., 2011).


260

2.4 Evaluation of BESS-STAIR ET

The BESS-STAIR ET estimations were evaluated against flux tower ET measurements in the U.S. Corn Belt. The U.S. Corn Belt (Figure 2) generally refers to a region in the Midwestern United States that has dominated corn and soybean production



265 in the United States (Green et al., 2018), which currently produces about 45% and 30% of the global corn and soybean, respectively (USDA, 2014). The region is characterized by relatively flat land, deep fertile soils, and a high soil organic matter (Green et al., 2018). Most part of the U.S. Corn Belt has favorable growing conditions of temperature and rainfall. A majority of the croplands in the U.S. Corn Belt are rainfed, with a small portion in the west part relying on irrigation.



270 Figure 2. Study areas. Red dots indicate 12 flux tower sites scattered in six areas across the U.S. Corn Belt. The background map indicates the percent each state contribute to the total national corn and soybean plantation area (USDA, 2018).

A total of 12 cropland sites scattered in six areas across the U.S. Corn Belt are registered in the AmeriFlux or FLUXNET network with publicly-available ET data (Figure 2 and Table 2). These sites include both corn only and corn/soybean rotation sites and both rainfed and irrigated sites, covering typical cropping patterns in the U.S. Corn Belt. All of them were used in this study to ensure the representativeness of the validation for the precision agriculture applications in this region. For six sites: US-Bo1 (Meyers and Hollinger, 2004), US-Bo2 (Bernacchi et al., 2005), US-Br1 (Prueger et al., 2003), US-Br3 (Prueger et al., 2003), US-Ro2 (Turner et al., 2016) and US-SFP (Wilson and Meyers, 2007), level 2 half-hourly data were downloaded from the AmeriFlux website (<http://ameriflux.lbl.gov/>). For three sites: US-IB1 (Matamala et al., 2008), US-Ro1 (Griffis et al., 2010) and US-Ro3 (Griffis et al., 2010), standardized gap-filled level 4 daily mean data were downloaded from the Carbon Dioxide Information Analysis Center data archive website ([https://mirrors.asun.co/climate-mirror/](https://mirrors.asun.co/climate-mirror/cdiac.ornl.gov/pub/ameriflux/)
280 [cdiac.ornl.gov/pub/ameriflux/](https://mirrors.asun.co/climate-mirror/cdiac.ornl.gov/pub/ameriflux/)). For the other three sites (Suyker et al., 2004): US-Ne1, US-Ne2 and US-Ne3, standardized high-quality gap-filled daily mean data were downloaded from the FLUXNET2015 website (<http://fluxnet.fluxdata.org/data/fluxnet2015-dataset/>).

285

Table 2. Information of 12 flux tower sites used for validation.



ID	Site	Location	Latitude	Longitude	Plant	Irrigation	Years	Source
1	US-Bo1	Bondville, IL	40.0062	-88.2904	corn/soybean	N	2000-2008	AmeriFlux L2
2	US-Bo2	Bondville, IL	40.0090	-88.2900	corn/soybean	N	2004-2006	AmeriFlux L2
3	US-Br1	Brooks Field, IA	41.9749	-93.6906	corn/soybean	N	2005-2011	AmeriFlux L2
4	US-Br3	Brooks Field, IA	41.9747	-93.6936	corn/soybean	N	2005-2011	AmeriFlux L2
5	US-IB1	Fermilab, IL	41.8593	-88.2227	corn/soybean	N	2005-2007	AmeriFlux L4
6	US-Ne1	Mead, NE	41.1651	-96.4766	corn	Y	2001-2012	FLUXNET2015
7	US-Ne2	Mead, NE	41.1649	-96.4701	corn/soybean	Y	2001-2012	FLUXNET2015
8	US-Ne3	Mead, NE	41.1797	-96.4397	corn/soybean	N	2001-2012	FLUXNET2015
9	US-Ro1	Rosemount, MN	44.7143	-93.0898	corn/soybean	N	2004-2006	AmeriFlux L4
10	US-Ro2	Rosemount, MN	44.7288	-93.0888	corn/soybean/clover	N	2008-2016	AmeriFlux L2
11	US-Ro3	Rosemount, MN	44.7217	-93.0893	corn/soybean	N	2004-2006	AmeriFlux L4
12	US-SFP	Sioux Falls, SD	43.2408	-96.9020	corn	N	2007-2009	AmeriFlux L2

By comparing with eddy covariance ET, we evaluated three ET estimations: BESS-STAIR with VIs-based LAI, BESS-STAIR with RTM-based LAI, and BESS-STAIR with MODIS LAI. MODIS LAI refers to MCD15A3H 500m resolution 4-day composite LAI product downloaded from <https://lpdaac.usgs.gov/tools/data-pool/>. Since eddy covariance technique used by flux towers provides water flux observations in term of latent heat (LE) rather than ET, evaluations were conducted by comparing BESS-STAIR daily LE estimates with flux tower measurements. At this point, water flux can be simply converted from energy unit (LE, MJ m⁻² d⁻¹) to water unit (ET mm d⁻¹) by dividing latent heat of vaporization which is a function of daily temperature (Henderson-Sellers, 1984). Flux tower measurements usually have an irregular and dynamic footprint at scales from 100-m to 1-km (Fu et al., 2014), but for simplicity, only 30-m resolution BESS-STAIR pixels containing the flux tower were used for the direct comparison. With regard to flux towers, measurements data were directly used without energy closure adjustment. For AmeriFlux level 2 data, half-hourly data were averaged to daily LE only if no gaps exist during the day to avoid sampling bias caused by missing data. For AmeriFlux level 4 data and FLUXNET2015 data, gap-filled daily LE were used directly.

3 Results

3.1 Performance of STAIR LAI

LAI is the key input of BESS. The accuracy of high-resolution LAI estimations determine the validity of high-resolution ET estimations. We evaluated VIs-based LAI and RTM-based LAI estimations derived from 30-m resolution STAIR fused surface reflectance data against field measurements. We also compared them with 500-m resolution MODIS LAI. Overall, STAIR-derived LAI agree well with measured LAI, with $R^2 > 0.85$, $RMSE < 0.8$ and mean bias error (MBE) ≈ 0 (Figure 3). The RTM-based method which is calibration free yields same performance with VIs-based method which requires substantial field measurements to build empirical relationships. Misclassification of CDL data between corn and soybean is



310 an important uncertainty source since both methods rely on crop types. During 2001 – 2007, 4 out of 21 site years (19%)
over the three Mead sites were misclassified. By using the correct classification (not shown), the accuracy of LAI
estimations reach $R^2 = 0.90$ and $RMSE = 0.62$ for VIs-based method and $R^2 = 0.89$ and $RMSE = 0.68$ for RTM-based
method. By comparison, coarse-resolution MODIS LAI has relatively large errors, especially a negative bias ($R^2 = 0.55$,
 $RMSE = 1.68$ and $MBE = -0.97$).

315





Figure 3. Scatter plots between LAI measurements and LAI estimations. LAI measurements are destructively collected at
three Mead sites. (a) – (b) STAIR-derived daily 30m-resolution LAI using VIs-based method and RTM-based method,
respectively. (c) 500m-resolution MODIS LAI.


320

3.2 Performance of BESS-STAIR ET

BESS-STAIR daily LE estimations are in a highly aligned agreement with ground truth from the 12 flux-tower
measurements (Figure 4). Across all of the 12 sites, BESS-STAIR LE with RTM-based LAI achieves an overall coefficient
of determination (R^2) of 0.75, root-mean-square error (RMSE) of $2.29 \text{ MJ m}^{-2} \text{ d}^{-1}$, relative error ($E(|X_{\text{estimation}} -$
325 $X_{\text{measurement}}|)/E(X_{\text{measurement}})$, RE) of 27.9%, and no overall bias. Figure 5 further exhibits its performance over all of the 12
flux tower sites. R^2 values range from 0.68 to 0.94 for corn, and 0.65 to 0.81 for soybean, highlighting the robustness of
BESS-STAIR ET in the U.S. Corn Belt. BESS-STAIR LE with VIs-based LAI has similar performance ($R^2 = 0.75$, $RMSE =$
 $2.24 \text{ MJ m}^{-2} \text{ d}^{-1}$, and $RE = 27.4\%$). Considering relatively small difference between BESS-STAIR using RTM-based LAI
and that using VIs-based LAI, only the former one which is calibration-free is demonstrated in the following parts of this
330 paper. By comparison, BESS LE with MODIS LAI shows larger errors ($R^2 = 0.65$, $RMSE = 2.50 \text{ MJ m}^{-2} \text{ d}^{-1}$, and $RE =$
 30.2%) comparing to BESS-STAIR.



335 Figure 4. Density scatter plots between LE measurements and LE estimations. LE measurements are from eddy covariance data collected at 12 flux towers. (a) and (b) BESS-STAIR LE with VIs-based LAI and RTM-based LAI, respectively. (c) 500m-BESS LE with MODIS LAI.




340 Figure 5. Site-by-site R^2 between flux tower measured and BESS-STAIR estimated daily LE for corn and soybean, respectively. Crop type is from CDL data.

345 Figure 6 shows the comparison between BESS-STAIR daily LE estimations and flux tower measurements over site years with least data gaps in measurements. Across all of the 12 sites, BESS-STAIR well captures the seasonal characteristics of LE observation from flux towers, as they exhibit generally consistent variations over the growing season. During the peak growing season (June, July and August), the radiation displays a dominant impact on measured daily LE, and it is reasonably




350 estimated by BESS-STAIR LE as well. In most cases, measured daily LE do not show strong and fast response to precipitation and/or irrigation, possibly due to the plentiful water storage in soil. Two exceptions are US-IB1 (2006) and US-Ne3 (2012). In case of US-IB1, no precipitation is available in August and little in July. As a result, daily LE measurements drop slightly quicker in August than other cases. Such anomaly is also depicted by BESS-STAIR LE. In case of US-Ne3, the severe drought in 2012 summer causes much lower LE values than the two adjacent irrigation sites (US-Ne1 and US-Ne2). BESS-STAIR LE also captures this considerable reduction, although a slight bias is observed in July. Figure 7 further demonstrates that the seasonal cumulative ET at three Mead sites calculated for both the flux tower measurements and BESS-STAIR estimations overall agree well throughout the peak growing season (June – September).






355 Figure 6. Seasonal time series of flux tower measured and BESS-STAIR estimated daily LE for 12 selected site years. Daily radiation and precipitation/irrigation are overlaid except for US-Br3.



360 Figure 7. Multi-year mean cumulative ET for flux tower measurements and BESS-STAIR estimations at three Mead sites from 2001 through 2012.

3.3 Spatiotemporal variations of BESS-STAIR ET



365 Figure 8. Daily LE ($\text{MJ m}^{-2} \text{d}^{-1}$) derived from (a) BESS-STAIR and (b) BESS-MODIS at Mead ($41.1^\circ\text{N} - 41.2^\circ\text{N}$, $96.4^\circ\text{W} - 96.5^\circ\text{W}$) on August 1, 2012.

BESS-STAIR daily ET demonstrates prominent spatial variations within the $0.1^\circ \times 0.1^\circ$ area near the Mead site in Nebraska (Figure 8). Because of the impact of drought, central pivot irrigated fields characterized by round-shaped plots generally display higher values than surrounding croplands, and croplands have much higher values than grasslands. Variabilities of

370



ET between different crop fields and within individual crop fields are also observable. Such variabilities might be attributed to different irrigation strategies, varieties and/or other management. By comparison, though 500m-resolution BESS-MODIS ET is able to capture the general spatial pattern, it has many mixed pixels and is unable to demonstrate gradients across field boundaries.

375





Figure 9. Monthly mean BESS-STAIR ET at Brooks Field (41.9°N – 42.0°N, 96.65°W – 96.75°W) during the growing season of 2000, along with a CDL land cover map. The last subplot shows the average time series of corn, soybean and grass.

380

Reasonable seasonal cycles for different land cover types are revealed by BESS-STAIR monthly ET averaged from gap-free daily estimations. An example time series of monthly ET maps at Brooks Field during the growing season of 2000 is shown in Figure 9. BESS-STAIR ET clearly captures the temporal dynamics throughout the growing season. All vegetation show low values (e.g., <math> < 2 \text{ mm d}^{-1}</math>) in April, May, September and October, but high values in June, July and August (JJA), with their peaks in July. Different seasonal cycles for corn, soybean and grass are also captured. Grass has the highest ET among the three vegetation types from April through June. Corn has higher ET than soybean in June and July, and decreases quickly since August. Soybean has the lowest ET from April through June, but has the highest ET in August.

385



390 Figure 10. Monthly mean BESS-STAIR ET in July at Rosemount (44.65°N – 44.75°N, 93.05°W – 93.15°W) from 2004
throughout 2017, along with a scatter plot between regional-averaged monthly mean precipitation and ET in July over the 14
years. Monthly precipitation maps are from PRISM (<http://www.prism.oregonstate.edu/historical/>).

BESS-STAIR is also able to produce long-term ET estimations due to its high computational efficiency. Figure 10 shows an
395 example time series of monthly ET in July at Rosemount from 2004 throughout 2017. The interannual variability is
observable, although much smaller than seasonal variability (Figure 10). In this case, time-series of BESS-STAIR ET is in
line with that of precipitation, as indicated by a significant linear correlation $r = 0.73$ ($p < 0.005$). It is also noted that ET is
relatively steady given the high interannual variation in precipitation, and BESS-STAIR enables the investigation of such
response at field scales.

400

4 Discussions

4.1 Performance of BESS-STAIR ET

In this study, we have presented BESS-STAIR, a new framework for estimating croplands ET at field and daily scale, and
we have demonstrated its high performance in the U.S. Corn Belt. The process-based biophysical model BESS, driven by
405 30-m resolution vegetation-related variables derived from STAIR fused surface spectral reflectance data (Figure 3) and
medium resolution environmental inputs derived from MODIS and other satellite data (Figure 1), is able to produce gap-free



ET and PET estimations on field-scale and at daily interval across space and time (Figure 4-7). Over the 12 sites across the U.S. Corn Belt (Figure 2), BESS-STAIR explains 75% variations in flux tower measured daily LE (Figure 4), with an overall RMSE of $2.29 \text{ MJ m}^{-2} \text{ d}^{-1}$ (equivalent to 0.93 mm d^{-1} or 26 W m^{-2}), a 27.9% relative error, and stable performance across sites (Figure 5), as well as consistent seasonal dynamics with respect to flux tower measurements (Figure 6-7).

The error statistics of BESS-STAIR are commensurate with previous high resolution croplands ET mapping studies. Typical RMSE values include 25 W m^{-2} by TSEB-DTD (Guzinski et al., 2014), 35 W m^{-2} by METRIC (Irmak et al., 2011), 62 W m^{-2} by SEBS (McCabe and Wood, 2006), 0.60 mm d^{-1} by SSEBop (Senay et al., 2016), and 1.04 mm d^{-1} by SEBAL (Singh et al., 2008). Nevertheless, it is worth mentioning that those studies used original Landsat data and therefore suffered from considerably large data gaps. In contrast, BESS-STAIR uses daily Landsat-MODIS fusion data free from any gaps, which leads to temporally continuous ET estimation at the field level, thus can meet the requirements of precision agriculture. In addition, it is worth mentioning that BESS-STAIR is calibration-free and therefore is scalable. It also indicates that the accuracy of BESS-STAIR ET is likely to further improve by using locally optimized driving force or parameter values.

BESS-STAIR is also comparable to other croplands ET mapping studies utilizing data fusion techniques. For example, DisALEXI-STARFM daily ET estimates were validated against the flux tower measurements over the three Mead sites (Yang et al., 2018). They reported error statistics around 1.2 mm d^{-1} RMSE and 29% relative error. BESS-STAIR's performance at these three same sites shows an average of 0.89 mm d^{-1} RMSE and 25.3% relative error (Figure 11). At monthly scale, the average RMSE and relative errors are only 0.48 mm d^{-1} and 14.3% (Figure A1). In addition, BESS-STAIR has a potential to apply to any croplands around the world back to 1984 when both high spatial resolution data (e.g., Landsat/TM) and high temporal resolution data (e.g., NOAA/AVHRR) were available.




Figure 11. Scatter plots between LE measurements and LE estimations at three sites US-Ne1, US-Ne2 and US-Ne3.



4.2 Scientific advantages of BESS-STAIR ET

The efficacy of BESS-STAIR lies in several aspects. First, BESS is a water-carbon-energy coupled biophysical model. BESS employs atmospheric and canopy radiative transfer modules, carbon assimilation module, stomatal conductance module, and energy balance module (Jiang and Ryu, 2016; Ryu et al., 2011). BESS integrates the simulation of carbon cycle, water cycle and energy cycle in the same framework. Such carbon-water-energy coupling strategy realistically and coherently simulates plant physiology and their response to the environment, specifically the carbon uptake and water loss by plants have been simulated synchronously through environmental constraints on stomatal conductance, with further constraints by available energy (Baldocchi and Meyers, 1998; Leuning et al., 1995). Many land surface models have already adopted such strategy and have successfully simulated the evolution of terrestrial ecosystems (Ju et al., 2006; Sellers, 1997; Tian et al., 2010). However, this is not the case in commonly-used remote sensing models. Empirical methods, water balance methods, and Priestley–Taylor methods only focus on the water cycle. Energy balance methods, triangular space methods, and Penman–Monteith methods couple water cycle and energy cycle and consider ET in the context of energy partitioning. BESS, unlike these remote sensing models, constrains ET with regards to both energy requirement and carbon requirement, thanks to explicit modeling of radiative transfer and stomatal behavior processes. For above reasons, BESS-STAIR ET does not only achieve high accuracy (Figure A1 – A3), but also accurately capture responses to GPP, radiation, temperature, and humidity at daily scale (Table 3). Thus, BESS-STAIR has the potential to advance the understanding of crop responses to climate change through bridging remote sensing data and land surface models, which was first suggested by Sellers et al. (1997) more than 20 years ago.

Table 3. BESS-STAIR captures the correct response of daily LE to GPP, radiation (R_g), temperature (T_a) and humidity (VPD) as compared to flux tower measurements over the three Mead sites from 2001 through 2012. The linear equation slopes and correlation coefficients between LE and other factors are similar in flux tower measurements and BESS-STAIR estimations, for both the whole growing season (April – October) or only peak growing season (June, July and August). For “flux tower” columns, R_g, T_a and VPD are from site measurements, while for “BESS-STAIR” columns, they are from satellite-derived coarse resolution inputs.

Time period	Relationship	Flux tower		BESS-STAIR	
		Equation	Correlation	Equation	Correlation
Growing season (April – October)	LE (MJ m ⁻² d ⁻¹) ~ GPP (gC m ⁻² d ⁻¹)	y = 0.44x + 3.19	r = 0.86 (p < 0.01)	y = 0.65x + 2.60	r = 0.90 (p < 0.01)
	LE (MJ m ⁻² d ⁻¹) ~ R _g (MJ m ⁻² d ⁻¹)	y = 0.33x - 0.07	r = 0.56 (p < 0.01)	y = 0.29x + 0.28	r = 0.51 (p < 0.01)
	LE (MJ m ⁻² d ⁻¹) ~ T _a (°C)	y = 0.44x - 1.60	r = 0.67 (p < 0.01)	y = 0.43x - 1.64	r = 0.54 (p < 0.01)
	LE (MJ m ⁻² d ⁻¹) ~ VPD (hPa)	y = 0.32x + 4.10	r = 0.31 (p < 0.01)	y = 0.44x - 2.50	r = 0.68 (p < 0.01)
JJA (June, July, August)	LE (MJ m ⁻² d ⁻¹) ~ GPP (gC m ⁻² d ⁻¹)	y = 0.35x + 4.85	r = 0.76 (p < 0.01)	y = 0.39x + 6.36	r = 0.63 (p < 0.01)
	LE (MJ m ⁻² d ⁻¹) ~ R _g (MJ m ⁻² d ⁻¹)	y = 0.36x + 1.79	r = 0.58 (p < 0.01)	y = 0.36x + 3.40	r = 0.77 (p < 0.01)
	LE (MJ m ⁻² d ⁻¹) ~ T _a (°C)	y = 0.52x - 2.20	r = 0.46 (p < 0.01)	y = 0.15x + 9.40	r = 0.28 (p < 0.01)



$$\text{LE (MJ m}^{-2} \text{ d}^{-1}) \sim \text{VPD (hPa)} \quad y = 0.28x + 7.61 \quad r = 0.32 \text{ (} p < 0.001 \text{)} \quad y = 0.28x + 4.37 \quad r = 0.30 \text{ (} p < 0.001 \text{)}$$

460 The second strength is that BESS-STAIR is designed most sensitive to the variables which can be well-quantified from
remote sensing data. BESS ET is most sensitive to solar radiation, followed by LAI (Ryu et al., 2011), as BESS ET is mainly
constrained by net radiation and GPP. In most cases, solar radiation is the predominant component of net radiation, while
LAI determines the capacity of radiation absorption and subsequently determines GPP. BESS explicitly computes radiation
components in high accuracy by driving an atmosphere radiative transfer model FLiES using MODIS cloud, aerosol and
atmospheric profile products. Globally, BESS-estimated solar radiation has its R^2 about 0.85 and 0.95 for MODIS snapshots
465 and 4-day averages, respectively (Ryu et al., 2018). On the other hand, BESS-STAIR calculates high spatiotemporal
resolution LAI and albedo from fused surface reflectance data. Since Landsat and MODIS surface reflectance products are
publicly-available and highly-reliable (Claverie et al., 2015; Masek et al., 2006), spatial heterogeneity and temporal
dynamics of crop growing conditions are well captured (Figure 8). This study only uses reflectance data fused from Landsat
and MODIS, but STAIR can be easily extended to further incorporate other types of data, such as Sentinel-2 (10 m
470 resolution) and Planet Lab CubeSats (3 m resolution) (McCabe et al., 2017). By incorporating more high resolution
observations, the relevance of reconstructed high resolution image series can be further improved.

The third strength is that BESS-STAIR is able to perform under all-weather conditions. BESS-STAIR fills data gaps in
surface reflectance, which has a smooth day-to-day variation even with changes in sky conditions (Liu et al., 2017). Based
475 on filtered surface reflectance, LAI and albedo time series are well-reconstructed, and subsequently BESS-STAIR could
directly work under all-weather condition. In this manner, BESS-STAIR has no need to fill cloudy-sky ET using clear-sky
ET estimations, which is error prone because the empirically-filled ET estimations usually lack sophisticated process-level
model constraints and thus can have large uncertainties. Figure 12 shows that the estimation errors of BESS-STAIR ET do
not change significantly under different sky conditions, with low to high “sky clearness index” referring more cloudy to
480 more clear sky conditions.




Figure 12. BESS-STAIR estimated daily LE has similar performance with varying sky clearness index (the ratio of incoming radiation on surface to that on top-of-atmosphere). The lower and upper boundaries of boxes refer to the first and third quartile of error statistics. The bars inside boxes refer to median values. The whiskers indicate 1.5 times of distance between the first and third quartiles.

4.3 Limitations and future improvements of BESS-STAIR ET

In this study, several inputs used by BESS have some limitations in terms of generality and accessibility. First, three plant functional parameters, peak $V_{\text{cmax}25}$, Ball-Barré slope and intercept are obtained from literatures, assuming constant given C3 or C4 plant type. Other land surface models tend to use the similar strategy by assigning fixed values to a given plant functional type (PFT) (Bonan et al., 2011; Kattge et al., 2009; Miner et al., 2017). The drawback of this strategy is the overlook of within-PFT variations and the feedback mechanisms between vegetation and its environment (Van Bodegom et al., 2014). These limitations might be mitigated by incorporating innovative leaf trait estimation techniques emerged in recent years, such as imaging spectroscopy (Serbin et al., 2015), sun-induced fluorescence (Zhang et al., 2018), and plant optimization theory (Walker et al., 2017; Wang et al., 2017). Second, BESS-STAIR in this study uses CDL data which is only available in United States. Fortunately, BESS does not require specific crop types but only C3/C4 distributions, and the separation of the major C4 crop maize from other crops is practical using time-series satellite data (Cai et al., 2018; Zhong et al., 2016). It is noted that misclassification of C3 and C4 crops are likely to cause large bias in GPP, but relatively small bias in ET (Fig. A1 – A3).

Though BESS-STAIR is able to capture water stress impact on ET in the U.S. Corn Belt where atmospheric demands play a major role, its applicability to regions where soil supply dominates needs further investigation. Some studies suggest that optical signal as an indicator of drought performs at a longer time scale than thermal signal does (Otkin et al., 2017). Drought first decreases soil moisture content due to enhanced ET induced by high atmospheric demand, then decreases ET



due to low soil moisture content, and finally causes damage to plants which changes surface reflectance. Accordingly, LAI may not serve as a relevant early warning of droughts. Furthermore, severe soil moisture stress may cause physiological deterioration in addition to structural damage that has been reflected in LAI. To address this issues for dry regions, we acknowledge that LST observations may provide essential adding values. At this point, the capacity of BESS-STAIR in
510 estimating LST leads to a possibility of optimizing BESS-STAIR using satellite-derived LST. Recent advances of innovative thermal observation platforms such as ECOSTRESS (Hulley et al., 2017), GOES-R (Schmit et al., 2017), and Sentinel-3 (Zheng et al., 2019) have provide great opportunity to integrate satellite-derived LST with the BESS-STAIR.

The BESS model itself in essence estimates instantaneous ET. The ratio of snapshot potential solar radiation to daily
515 potential solar radiation is adopted as a scaling factor for the temporal upscaling of ET (Ryu et al., 2012). In this study, BESS runs two times per day, utilizing radiation components derived from Terra/MODIS (around 11:00 AM) and Aqua/MODIS (around 1:00 PM) data, respectively. The two instantaneous ET estimates are separately upscaled to daily estimates and averaged. In spite of robustness of the upscaling algorithm (Ryu et al., 2012), bias cannot be avoided if the sky
520 cloud. Since BESS is a time-independent model and can perform at any time during daytime, adding more snapshots to account for the diurnal variations of radiation can solve this problem. Unfortunately, fine-resolution polar-orbiting satellite usually have similar overpass times (10:00 AM – 11:00 AM and 1:00 PM – 2:00 PM), so even adding more satellites is likely to bring redundant information only. Reanalysis radiation data covering diurnal cycle have limited accuracy and
525 coarse resolution (Babst et al., 2008; Zhang et al., 2016b), so they may be unable to provide much added values as well. Next-generation geostationary satellites, acquiring data with both high spatial and high temporal resolutions such as GOES-R and GaoFen-4 (Goodman et al., 2012; Xu et al., 2017), are expected to enable BESS-STAIR ET in hourly or sub-hourly interval and subsequently generate more realistic daily ET estimates.

5 Conclusions

530 In this study we presented BESS-STAIR, a new framework to estimate high spatiotemporal resolution ET that can be used for field-level precision water resources management. BESS-STAIR couples a satellite-driven water-energy-carbon coupled biophysical model BESS with a generic and fully-automated fusion algorithm STAIR to generate gap-free 30-m resolution daily ET estimations. Comprehensive evaluation of BESS-STAIR ET estimations revealed: 1) reliable performance over 12 flux tower sites across the U.S. Corn Belt, and 2) reasonable spatial patterns, seasonal cycles and interannual dynamics. The
535 proposed BESS-STAIR framework has demonstrated its ability to provide significant advancements with regard to daily field-level estimations of ET at regional and decadal scales. We expect BESS-STAIR to become a solid tool for precision water resources management and other precision agriculture applications for the U.S. Corn Belt as well as other agricultural areas around the world, thanks to the global coverage of input data.



540 Data availability. The data generated in this study are available upon request.

Competing interests. The authors declare that they have no conflict of interest.


545 Author contribution. C.J. and K.G. designed the study, C.J. conducted the modeling and analysis, M.P. and B.P. provided research inputs during the analysis, Y.R. and S.W. provided guidance in the BESS model and the STAIR algorithm, respectively. All the authors contributed to the writing of the manuscript.

Acknowledgements


550 C.J. and K.G. were funded by the DOE Center for Advanced Bioenergy and Bioproducts Innovation (U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research under Award Number DE-SC0018420). Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the views of the U.S. Department of Energy. K.G., B. P., and S.W. are funded by NASA awards (NNX16AI56G and 80NSSC18K0170) and USDA National Institute of Food and Agriculture (NIFA) Foundational Program award (2017-67013-26253, 2017-68002-26789, 2017-67003-28703). The development of the BESS model was mainly
555 supported by The National Research Foundation of Korea (NRF-2014R1A2A1A11051134). K.G. and C.J. also acknowledge the support from Blue Waters Professorship from National Center for Supercomputing Applications of UIUC. This research is part of the Blue Waters sustained-petascale computing project, which is supported by the National Science Foundation (awards OCI-0725070 and ACI-1238993) and the state of Illinois. Blue Waters is a joint effort of the University of Illinois at Urbana-Champaign and its National Center for Supercomputing Applications. We thank the U.S. Landsat project
560 management and staff at USGS Earth Resources Observation and Science (EROS) Center South Dakota for providing the Landsat data free of charge. We also thank NASA freely share the MODIS products.



Appendix



565 Figure A1. Time series of monthly mean LE from flux tower measurements and BESS-STAIR estimations.



570 Figure A2. Time series of monthly mean GPP from flux tower measurements and BESS-STAIR estimations. Significant underestimations in 2003 and 2011 for Ne1, in 2005 for Ne2, and in 2005 for Ne3 are due to misclassification of corn as soybean in CDL. Significant overestimations in 2006 for Ne2 are due to misclassification of soybean as corn in CDL.




Figure A3. Time series of monthly mean Rn from flux tower measurements and BESS-STAIR estimations.

References

- 575 Allen, R. G., Pereira, L. S., Raes, D. and Smith, M.: Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56, Italy: Rome. [online] Available from: https://s3.amazonaws.com/academia.edu.documents/40878584/Allen_FAO1998.pdf?AWSAccessKeyId=AKIAIWOWYYGZ2Y53UL3A&Expires=1527634764&Signature=GkL44Ekv1zkaqSV04%2F4EwT99LEk%3D&response-content-disposition=inline%3B filename%3DAllen_FAO1998.pdf, 1998.
- 580 Anderson, M., Gao, F., Knipper, K., Hain, C., Dulaney, W., Baldocchi, D., Eichelmann, E., Hemes, K., Yang, Y., Medellin-Azuara, J. and Kustas, W.: Field-Scale Assessment of Land and Water Use Change over the California Delta Using Remote Sensing, *Remote Sens.*, 10(10), doi:10.3390/rs10060889, 2018.
- Anderson, M. C., Norman, J. M., Diak, G. R. and Mecikalski, J. R.: A Two-Source Time-Integrated Model for Estimating Surface Fluxes Using Thermal Infrared Remote Sensing, *Remote Sens. Environ.*, 60, 195–216, 1997.
- 585 Babst, F., Mueller, R. W. and Hollmann, R.: Verification of NCEP reanalysis shortwave radiation with mesoscale remote sensing data, *IEEE Geosci. Remote Sens. Lett.*, 5(1), 34–37, doi:10.1109/LGRS.2007.907537, 2008.
- Bacour, C., Jacquemoud, S., Tourbier, Y., Dechambre, M. and Frangi, J.: Design and analysis of numerical experiments to compare four canopy reflectance models, *Remote Sens. Environ.*, 79, 72–83, 2002.
- Baldocchi, D. and Meyers, T.: On using eco-physiological, micrometeorological and biogeochemical theory to evaluate carbon dioxide, water vapor and trace gas fluxes over vegetation: A perspective, *Agric. For. Meteorol.*, 90(1–2), 1–25, doi:10.1016/S0168-1923(97)00072-5, 1998.
- 590



- Baldocchi, D., Dralle, D., Jiang, C. and Ryu, Y.: How Much Water is Evaporated Across California?: A Multi-Year Assessment Using a Biophysical Model Forced with Satellite Remote Sensing Data, *Water Resour. Res.*, 2018WR023884, doi:10.1029/2018WR023884, 2019.
- 595 Ball, J. T., Woodrow, I. E. and Berry, J. A.: A Model Predicting Stomatal Conductance and its Contribution to the Control of Photosynthesis under Different Environmental Conditions, in *Progress in Photosynthesis Research*, pp. 221–224, Springer Netherlands, Dordrecht., 1987.
- Baret, F., Hagolle, O., Geiger, B., Bicheron, P., Miras, B., Huc, M., Berthelot, B., Niño, F., Weiss, M., Samain, O., Roujean, J. L. and Leroy, M.: LAI, fAPAR and fCover CYCLOPES global products derived from VEGETATION. Part 1: Principles
600 of the algorithm, *Remote Sens. Environ.*, 110, 275–286, doi:10.1016/j.rse.2007.02.018, 2007.
- Bernacchi, C. J., Hollinger, S. E. and Meyers, T.: The conversion of the corn/soybean ecosystem to no-till agriculture may result in a carbon sink, *Glob. Chang. Biol.*, 11(11), 1867–1872, doi:10.1111/j.1365-2486.2005.01050.x, 2005.
- Van Bodegom, P. M., Douma, J. C. and Verheijen, L. M.: A fully traits-based approach to modeling global vegetation distribution., *Proc. Natl. Acad. Sci. U. S. A.*, 111(38), 13733–8, doi:10.1073/pnas.1304551110, 2014.
- 605 Bonan, G. B., Lawrence, P. J., Oleson, K. W., Levis, S., Jung, M., Reichstein, M., Lawrence, D. M. and Swenson, S. C.: Improving canopy processes in the Community Land Model version 4 (CLM4) using global flux fields empirically inferred from FLUXNET data, *J. Geophys. Res.*, 116(G2), G02014, doi:10.1029/2010JG001593, 2011.
- Boryan, C., Yang, Z., Mueller, R. and Craig, M.: Monitoring US agriculture: The US department of agriculture, national agricultural statistics service, cropland data layer program, *Geocarto Int.*, 26(5), 341–358,
610 doi:10.1080/10106049.2011.562309, 2011.
- Brown, J. F. and Pervez, M. S.: Merging remote sensing data and national agricultural statistics to model change in irrigated agriculture, *Agric. Syst.*, 127, 28–40, doi:10.1016/j.agsy.2014.01.004, 2014.
- Cai, Y., Guan, K., Peng, J., Wang, S., Seifert, C., Wardlow, B. and Li, Z.: A high-performance and in-season classification system of field-level crop types using time-series Landsat data and a machine learning approach, *Remote Sens. Environ.*,
615 210(March), 35–47, doi:10.1016/j.rse.2018.02.045, 2018.
- Cammalleri, C., Anderson, M. C., Gao, F., Hain, C. R. and Kustas, W. P.: A data fusion approach for mapping daily evapotranspiration at field scale, *Water Resour. Res.*, 49(8), 4672–4686, doi:10.1002/wrcr.20349, 2013.
- Claverie, M., Vermote, E. F., Franch, B. and Masek, J. G.: Evaluation of the Landsat-5 TM and Landsat-7 ETM+ surface reflectance products, *Remote Sens. Environ.*, 169, 390–403, doi:10.1016/j.rse.2015.08.030, 2015.
- 620 Collatz, G., Ribas-Carbo, M. and Berry, J.: Coupled Photosynthesis-Stomatal Conductance Model for Leaves of C 4 Plants, *Aust. J. Plant Physiol.*, 19(5), 519, doi:10.1071/PP9920519, 1992.
- Collatz, G. J., Ball, J. T., Grivet, C. and Berry, J. a: Physiological and environmental regulation of stomatal conductance, photosynthesis and transpiration: a model that includes a laminar boundary layer, *Agric. For. Meteorol.*, 54(2–4), 107–136, doi:10.1016/0168-1923(91)90002-8, 1991.
- 625 Darvishzadeh, R., Skidmore, A., Schlerf, M. and Atzberger, C.: Inversion of a radiative transfer model for estimating



- vegetation LAI and chlorophyll in a heterogeneous grassland, *Remote Sens. Environ.*, 112, 2592–2604, doi:10.1016/j.rse.2007.12.003, 2008.
- Dils, B., Buchwitz, M., Reuter, M., Schneising, O., Boesch, H., Parker, R., Guerlet, S., Aben, I., Blumenstock, T., Burrows, J. P., Butz, A., Deutscher, N. M., Frankenberg, C., Hase, F., Hasekamp, O. P., Heymann, J., De Mazière, M., Notholt, J.,
630 Sussmann, R., Warneke, T., Griffith, D., Sherlock, V. and Wunch, D.: The greenhouse gas climate change initiative (GHG-CCI): Comparative validation of GHG-CCI SCIAMACHY/ENVISAT and TANSO-FTS/GOSAT CO₂ and CH₄ retrieval algorithm products with measurements from the TCCON, *Atmos. Meas. Tech.*, 7(6), 1723–1744, doi:10.5194/amt-7-1723-2014, 2014.
- Duan, S. B., Li, Z. L., Wu, H., Tang, B. H., Ma, L., Zhao, E. and Li, C.: Inversion of the PROSAIL model to estimate leaf
635 area index of maize, potato, and sunflower fields from unmanned aerial vehicle hyperspectral data, *Int. J. Appl. Earth Obs. Geoinf.*, 26(1), 12–20, doi:10.1016/j.jag.2013.05.007, 2014.
- Feret, J. B., François, C., Asner, G. P., Gitelson, A. a., Martin, R. E., Bidel, L. P. R., Ustin, S. L., le Maire, G. and Jacquemoud, S.: PROSPECT-4 and 5: Advances in the leaf optical properties model separating photosynthetic pigments, *Remote Sens. Environ.*, 112(6), 3030–3043, doi:10.1016/j.rse.2008.02.012, 2008.
- 640 Fisher, J. B., Tu, K. P. and Baldocchi, D. D.: Global estimates of the land-atmosphere water flux based on monthly AVHRR and ISLSCP-II data, validated at 16 FLUXNET sites, *Remote Sens. Environ.*, 112, 901–919, doi:10.1016/j.rse.2007.06.025, 2008.
- Fu, D., Chen, B., Zhang, H., Wang, J., Black, T. A., Amiro, B. D., Bohrer, G., Bolstad, P., Coulter, R., Rahman, A. F., Dunn, A., McCaughey, J. H., Meyers, T. and Verma, S.: Estimating landscape net ecosystem exchange at high spatial–temporal
645 resolution based on Landsat data, an improved upscaling model framework, and eddy covariance flux measurements, *Remote Sens. Environ.*, 141, 90–104, doi:10.1016/j.rse.2013.10.029, 2014.
- Gao, F., Masek, J., Schwaller, M. and Hall, F.: On the blending of the landsat and MODIS surface reflectance: Predicting daily landsat surface reflectance, *IEEE Trans. Geosci. Remote Sens.*, 44(8), 2207–2218, doi:10.1109/TGRS.2006.872081, 2006.
- 650 Gao, F., Anderson, M. C., Zhang, X., Yang, Z., Alfieri, J. G., Kustas, W. P., Mueller, R., Johnson, D. M. and Prueger, J. H.: Toward mapping crop progress at field scales through fusion of Landsat and MODIS imagery, *Remote Sens. Environ.*, 188, 9–25, doi:10.1016/j.rse.2016.11.004, 2017.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W.,
655 Kim, G. K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M. and Zhao, B.: The modern-era retrospective analysis for research and applications, version 2 (MERRA-2), *J. Clim.*, 30(14), 5419–5454, doi:10.1175/JCLI-D-16-0758.1, 2017.
- Gitelson, A. A., Wardlow, B. D., Keydan, G. P. and Leavitt, B.: An evaluation of MODIS 250-m data for green LAI estimation in crops, *Geophys. Res. Lett.*, 34(20), doi:10.1029/2007GL031620, 2007.



- 660 Gómez, C., White, J. C. and Wulder, M. A.: Optical remotely sensed time series data for land cover classification: A review, *ISPRS J. Photogramm. Remote Sens.*, 116, 55–72, doi:10.1016/j.isprsjprs.2016.03.008, 2016.
- Goodman, S. J., Gurka, J., De Maria, M., Schmit, T. J., Mostek, A., Jedlovec, G., Siewert, C., Feltz, W., Gerth, J., Brummer, R., Miller, S., Reed, B. and Reynolds, R. R.: The goes-R proving ground: Accelerating user readiness for the next-generation geostationary environmental satellite system, *Bull. Am. Meteorol. Soc.*, 93(7), 1029–1040, doi:10.1175/BAMS-D-11-00175.1, 2012.
- 665 Griffis, T. J., Baker, J. M., Sargent, S. D., Erickson, M., Corcoran, J., Chen, M. and Billmark, K.: Influence of C4 vegetation on ^{13}C discrimination and isoforcing in the upper Midwest, United States, *Global Biogeochem. Cycles*, 24(4), n/a-n/a, doi:10.1029/2009GB003768, 2010.
- Guzinski, R., Nieto, H., Jensen, R. and Mendiguren, G.: Remotely sensed land-surface energy fluxes at sub-field scale in heterogeneous agricultural landscape and coniferous plantation, *Biogeosciences*, 11(18), 5021–5046, doi:10.5194/bg-11-5021-2014, 2014.
- 670 Hammerling, D. M., Michalak, A. M. and Kawa, S. R.: Mapping of CO_2 at high spatiotemporal resolution using satellite observations: Global distributions from OCO-2, *J. Geophys. Res.*, 117(D6), D06306, doi:10.1029/2011JD017015, 2012.
- He, L., Chen, J. M., Pisek, J., Schaaf, C. B. and Strahler, A. H.: Global clumping index map derived from the MODIS BRDF product, *Remote Sens. Environ.*, 119, 118–130, doi:10.1016/j.rse.2011.12.008, 2012.
- 675 Hersbach, H. and H.: The ERA5 Atmospheric Reanalysis., *Am. Geophys. Union, Fall Gen. Assem. 2016*, Abstr. id. NG33D-01 [online] Available from: <http://adsabs.harvard.edu/abs/2016AGUFMNG33D..01H> (Accessed 18 July 2019), 2016.
- Houborg, R. and McCabe, M. F.: Remote Sensing of Environment A Cubesat enabled Spatio-Temporal Enhancement Method (CESTEM) utilizing Planet , Landsat and MODIS data, *Remote Sens. Environ.*, 209(January), 211–226, doi:10.1016/j.rse.2018.02.067, 2018.
- 680 Huang, Y., Ryu, Y., Jiang, C., Kimm, H., Kim, S., Kang, M. and Shim, K.: BESS-Rice: A remote sensing derived and biophysical process-based rice productivity simulation model, *Agric. For. Meteorol.*, 256–257(March), 253–269, doi:10.1016/j.agrformet.2018.03.014, 2018.
- Hulley, G., Hook, S., Fisher, J. and Lee, C.: ECOSTRESS, A NASA Earth-Ventures Instrument for studying links between the water cycle and plant health over the diurnal cycle, in *International Geoscience and Remote Sensing Symposium (IGARSS)*, vol. 2017-July, pp. 5494–5496, IEEE., 2017.
- 685 Irmak, A. ;, Ratcliffe, I. ;, Ranade, P. ;, Hubbard, K. G., Singh, R. K. and Kamble: ESTIMATION OF LAND SURFACE EVAPOTRANSPIRATION WITH A SATELLITE REMOTE SENSING PRO, *Gt. Plains Res. Spring*, 21(1), 73–88, doi:10.2307/23779969, 2011.
- 690 Irmak, S., Kabenge, I., Skaggs, K. E. and Mutibwa, D.: Trend and magnitude of changes in climate variables and reference evapotranspiration over 116-yr period in the Platte River Basin, central Nebraska-USA, *J. Hydrol.*, 420–421, 228–244, doi:10.1016/j.jhydrol.2011.12.006, 2012.
- Jacquemoud, S. and Baret, F.: PROSPECT: A model of leaf optical properties spectra, *Remote Sens. Environ.*, 34(2), 75–91,



- doi:10.1016/0034-4257(90)90100-Z, 1990.
- 695 Jacquemoud, S., Ustin, S. L., Verdebout, J., Schmuck, G., Andreoli, G. and Hosgood, B.: Estimating leaf biochemistry using the PROSPECT leaf optical properties model, *Remote Sens. Environ.*, 56(3), 194–202, doi:10.1016/0034-4257(95)00238-3, 1996.
- Jacquemoud, S., Verhoef, W., Baret, F., Bacour, C., Zarco-Tejada, P. J., Asner, G. P., François, C. and Ustin, S. L.: PROSPECT + SAIL models: A review of use for vegetation characterization, *Remote Sens. Environ.*, 113(SUPPL. 1), S56–
700 S66, doi:10.1016/j.rse.2008.01.026, 2009.
- Jiang, C. and Ryu, Y.: Multi-scale evaluation of global gross primary productivity and evapotranspiration products derived from Breathing Earth System Simulator (BESS), *Remote Sens. Environ.*, 186, 528–547, doi:10.1016/j.rse.2016.08.030, 2016.
- Jiang, L. and Islam, S.: A methodology for estimation of surface evapotranspiration over large areas using remote sensing
705 observations, *Geophys. Res. Lett.*, 26(17), 2773–2776, doi:10.1029/1999GL006049, 1999.
- Ju, W., Chen, J. M., Black, T. A., Barr, A. G., Liu, J. and Chen, B.: Modelling multi-year coupled carbon and water fluxes in a boreal aspen forest, *Agric. For. Meteorol.*, 140, 136–151, doi:10.1016/j.agrformet.2006.08.008, 2006.
- Jung, M., Reichstein, M., Ciais, P., Seneviratne, S. I., Sheffield, J., Goulden, M. L., Bonan, G., Cescatti, A., Chen, J., de Jeu, R., Dolman, a J., Eugster, W., Gerten, D., Gianelle, D., Gobron, N., Heinke, J., Kimball, J., Law, B. E., Montagnani, L., Mu,
710 Q., Mueller, B., Oleson, K., Papale, D., Richardson, A. D., Rouspard, O., Running, S., Tomelleri, E., Viovy, N., Weber, U., Williams, C., Wood, E., Zaehle, S. and Zhang, K.: Recent decline in the global land evapotranspiration trend due to limited moisture supply., *Nature*, 467(7318), 951–954, doi:10.1038/nature09396, 2010.
- Kalma, J. D., McVicar, T. R. and McCabe, M. F.: Estimating land surface evaporation: A review of methods using remotely sensed surface temperature data, *Surv. Geophys.*, 29(4–5), 421–469, doi:10.1007/s10712-008-9037-z, 2008.
- 715 Kattge, J., Knorr, W., Raddatz, T. and Wirth, C.: Quantifying photosynthetic capacity and its relationship to leaf nitrogen content for global-scale terrestrial biosphere models, *Glob. Chang. Biol.*, 15(4), 976–991, doi:10.1111/j.1365-2486.2008.01744.x, 2009.
- Katul, G. G., Oren, R., Manzoni, S., Higgins, C. and Parlange, M. B.: Evapotranspiration: a process driving mass transport and energy exchange in the soil-plant-atmosphere-climate system, *Rev. Geophys.*, 50(RG3002), RG000366: 1-25,
720 doi:10.1029/2011RG000366.1.INTRODUCTION, 2012.
- Kobayashi, H. and Iwabuchi, H.: A coupled 1-D atmosphere and 3-D canopy radiative transfer model for canopy reflectance, light environment, and photosynthesis simulation in a heterogeneous landscape, *Remote Sens. Environ.*, 112(1), 173–185, doi:10.1016/j.rse.2007.04.010, 2008.
- Leuning, R., Kelliher, F. M., DePury, D. G. G. and Schulze, E. -D.: Leaf nitrogen, photosynthesis, conductance and
725 transpiration: scaling from leaves to canopies, *Plant. Cell Environ.*, 18(10), 1183–1200, doi:10.1111/j.1365-3040.1995.tb00628.x, 1995.
- Li, Y., Huang, C., Hou, J., Gu, J., Zhu, G. and Li, X.: Mapping daily evapotranspiration based on spatiotemporal fusion of



- ASTER and MODIS images over irrigated agricultural areas in the Heihe River Basin, Northwest China, *Agric. For. Meteorol.*, 244–245(June), 82–97, doi:10.1016/j.agrformet.2017.05.023, 2017.
- 730 Li, Z. L., Tang, R., Wan, Z., Bi, Y., Zhou, C., Tang, B., Yan, G. and Zhang, X.: A review of current methodologies for regional Evapotranspiration estimation from remotely sensed data, *Sensors*, 9(5), 3801–3853, doi:10.3390/s90503801, 2009.
- Liang, S.: Narrowband to broadband conversions of land surface albedo I Algorithms, *Remote Sens. Environ.*, 76, 213–238, doi:10.1016/S0034-4257(00)00205-4, 2001.
- Liu, Y., Wang, Z., Sun, Q., Erb, A. M., Li, Z., Schaaf, C. B., Zhang, X., Román, M. O., Scott, R. L., Zhang, Q., Novick, K.
735 A., Sydonia Bret-Harte, M., Petroy, S. and SanClements, M.: Evaluation of the VIIRS BRDF, Albedo and NBAR products suite and an assessment of continuity with the long term MODIS record, *Remote Sens. Environ.*, 201(September), 256–274, doi:10.1016/j.rse.2017.09.020, 2017.
- Lobell, D. B., Roberts, M. J., Schlenker, W., Braun, N., Little, B. B., Rejesus, R. M. and Hammer, G. L.: Greater Sensitivity to Drought Accompanies Maize Yield Increase in the U.S. Midwest, *Science (80-.)*, 344(6183), 516–519,
740 doi:10.1126/science.1251423, 2014.
- Loew, A., Peng, J. and Borsche, M.: High-resolution land surface fluxes from satellite and reanalysis data (HOLAPS v1.0): Evaluation and uncertainty assessment, *Geosci. Model Dev.*, 9(7), 2499–2532, doi:10.5194/gmd-9-2499-2016, 2016.
- Lu, X. and Zhuang, Q.: Evaluating evapotranspiration and water-use efficiency of terrestrial ecosystems in the conterminous United States using MODIS and AmeriFlux data, *Remote Sens. Environ.*, 114(9), 1924–1939, doi:10.1016/j.rse.2010.04.001,
745 2010.
- Luo, Y., Guan, K. and Peng, J.: STAIR: A generic and fully-automated method to fuse multiple sources of optical satellite data to generate a high-resolution, daily and cloud-/gap-free surface reflectance product, *Remote Sens. Environ.*, 214(March), 87–99, doi:10.1016/j.rse.2018.04.042, 2018.
- Ma, Y., Liu, S., Song, L., Xu, Z., Liu, Y., Xu, T. and Zhu, Z.: Estimation of daily evapotranspiration and irrigation water efficiency at a Landsat-like scale for an arid irrigation area using multi-source remote sensing data, *Remote Sens. Environ.*,
750 216(December 2016), 715–734, doi:10.1016/j.rse.2018.07.019, 2018.
- Masek, J. G., Vermote, E. F., Saleous, N. E., Wolfe, R., Hall, F. G., Huemmrich, K. F., Gao, F., Kutler, J. and Lim, T.: A Landsat Surface Reflectance Dataset, *IEEE Geosci. Remote Sens. Lett.*, 3(1), 68–72, doi:10.1109/LGRS.2005.857030, 2006.
- Matamala, R., Jastrow, J. D., Miller, R. M. and Garten, C. T.: Temporal changes in C and N stocks of restored prairie: Implications for C sequestration strategies, *Ecol. Appl.*, 18(6), 1470–1488, doi:10.1890/07-1609.1, 2008.
755
- McCabe, M. F. and Wood, E. F.: Scale influences on the remote estimation of evapotranspiration using multiple satellite sensors, *Remote Sens. Environ.*, 105(4), 271–285, doi:10.1016/j.rse.2006.07.006, 2006.
- McCabe, M. F., Rodell, M., Alsdorf, D. E., Miralles, D. G., Uijlenhoet, R., Wagner, W., Lucieer, A., Houborg, R., Verhoest, N. E. C., Franz, T. E., Shi, J., Gao, H. and Wood, E. F.: The future of Earth observation in hydrology, *Hydrol. Earth Syst. Sci.*, 21(7), 3879–3914, doi:10.5194/hess-21-3879-2017, 2017.
760
- Meyers, T. P. and Hollinger, S. E.: An assessment of storage terms in the surface energy balance of maize and soybean,



- Agric. For. Meteorol., 125(1–2), 105–115, doi:10.1016/j.agrformet.2004.03.001, 2004.
- Miner, G. L., Bauerle, W. L. and Baldocchi, D. D.: Estimating the sensitivity of stomatal conductance to photosynthesis: a review, *Plant Cell Environ.*, 40(7), 1214–1238, doi:10.1111/pce.12871, 2017.
- 765 Miralles, D. G., Holmes, T. R. H., De Jeu, R. A. M., Gash, J. H., Meesters, A. G. C. A. and Dolman, A. J.: Global land-surface evaporation estimated from satellite-based observations, *Hydrol. Earth Syst. Sci.*, 15(2), 453–469, doi:10.5194/hess-15-453-2011, 2011.
- Mousivand, A., Menenti, M., Gorte, B. and Verhoef, W.: Global sensitivity analysis of the spectral radiance of a soil-vegetation system, *Remote Sens. Environ.*, 145, 131–144, doi:10.1016/j.rse.2014.01.023, 2014.
- 770 Mu, Q., Zhao, M. and Running, S. W.: Improvements to a MODIS global terrestrial evapotranspiration algorithm, *Remote Sens. Environ.*, 115(8), 1781–1800, doi:10.1016/j.rse.2011.02.019, 2011.
- Myneni, R. B., Hoffman, S., Knyazikhin, Y., Privette, J. L., Glassy, J., Tian, Y., Wang, Y., Song, X., Zhang, Y., Smith, G. R., Lotsch, a., Friedl, M., Morisette, J. T., Votava, P., Nemani, R. R. and Running, S. W.: Global products of vegetation leaf area and fraction absorbed PAR from year one of MODIS data, *Remote Sens. Environ.*, 83, 214–231, doi:10.1016/S0034-775 4257(02)00074-3, 2002.
- Nguy-Robertson, A., Gitelson, A., Peng, Y., Viña, A., Arkebauer, T. and Rundquist, D.: Green leaf area index estimation in maize and soybean: Combining vegetation indices to achieve maximal sensitivity, *Agron. J.*, 104(5), 1336–1347, doi:10.2134/agronj2012.0065, 2012.
- Ort, D. R. and Long, S. P.: Limits on Yields in the Corn Belt, *Science (80-.)*, 344(6183), 484–485, doi:10.1126/science.1253884, 2014.
- 780 Otkin, J. A., Svoboda, M., Hunt, E. D., Ford, T. W., Anderson, M. C., Hain, C. and Basara, J. B.: Flash Droughts: a Review and Assessment of the Challenges Imposed By Rapid Onset Droughts in the United States, *Bull. Am. Meteorol. Soc.*, (MAY), BAMS-D-17-0149.1, doi:10.1175/BAMS-D-17-0149.1, 2017.
- Pan, M., Sahoo, A. K., Troy, T. J., Vinukollu, R. K., Sheffield, J. and Wood, A. E. F.: Multisource estimation of long-term terrestrial water budget for major global river basins, *J. Clim.*, 25(9), 3191–3206, doi:10.1175/JCLI-D-11-00300.1, 2012.
- 785 Paw U, K. T. and Gao, W.: Applications of solutions to non-linear energy budget equations, *Agric. For. Meteorol.*, 43(2), 121–145, doi:10.1016/0168-1923(88)90087-1, 1988.
- Pisek, J., Sonnentag, O., Richardson, A. D. and Möttus, M.: Is the spherical leaf inclination angle distribution a valid assumption for temperate and boreal broadleaf tree species?, *Agric. For. Meteorol.*, 169, 186–194, doi:10.1016/j.agrformet.2012.10.011, 2013.
- 790 Prueger, J. H., Kustas, W. P., Hipps, L. E., Hatfield, J. L., Cahill, A., Williams, C. A., Albertson, J. D., Eichinger, W. E., Cooper, D. I., Brunsell, N. and others: Spatial variability of turbulent fluxes across a corn/soybean production region in central Iowa’, in *AMS 17th Conference on Hydrology*, Long Beach, CA, USA., 2003.
- De Pury, D. G. G. and Farquhar, G. D.: Simple scaling of photosynthesis from leaves to canopies without the errors of big-leaf models, *Plant, Cell Environ.*, 20(5), 537–557, doi:10.1111/j.1365-3040.1997.tb00466.x, 1997.
- 795



- Ryu, Y., Baldocchi, D., Kobayashi, H., Van Ingen, C., Li, J., Black, T. A., Beringer, J., Van Gorsel, E., Knohl, A., Law, B. and Rouspard, O.: Integration of MODIS land and atmosphere products with a coupled-process model to estimate gross primary productivity and evapotranspiration from 1 km to global scales, *Global Biogeochem. Cycles*, 25(GB4017), GB4017, doi:10.1029/2011GB004053, 2011.
- 800 Ryu, Y., Baldocchi, D. D., Black, T. A., Detto, M., Law, B. E., Leuning, R., Miyata, A., Reichstein, M., Vargas, R., Ammann, C., Beringer, J., Flanagan, L. B., Gu, L., Hutley, L. B., Kim, J., McCaughey, H., Moors, E. J., Rambal, S. and Vesala, T.: On the temporal upscaling of evapotranspiration from instantaneous remote sensing measurements to 8-day mean daily-sums, *Agric. For. Meteorol.*, 152(1), 212–222, doi:10.1016/j.agrformet.2011.09.010, 2012.
- Ryu, Y., Jiang, C., Kobayashi, H. and Detto, M.: MODIS-derived global land products of shortwave radiation and diffuse
805 and total photosynthetically active radiation at 5 km resolution from 2000, *Remote Sens. Environ.*, 204(January 2018), 812–825, doi:10.1016/j.rse.2017.09.021, 2018.
- Sayer, A. M., Munchak, L. A., Hsu, N. C., Levy, R. C., Bettenhausen, C. and Jeong, M. J.: MODIS Collection 6 aerosol products: Comparison between Aqua’s e-Deep Blue, Dark Target, and “merged” data sets, and usage recommendations, *J. Geophys. Res. Atmos.*, 119(24), 13965–13989, doi:10.1002/2014JD022453, 2014.
- 810 Schmit, T. J., Griffith, P., Gunshor, M. M., Daniels, J. M., Goodman, S. J. and Lebar, W. J.: A closer look at the ABI on the goes-r series, *Bull. Am. Meteorol. Soc.*, 98(4), 681–698, doi:10.1175/BAMS-D-15-00230.1, 2017.
- Seemann, S. W., Li, J., Gumley, L. E., Strabala, K. I. and Menzel, W. P.: Operational retrieval of atmospheric temperature, moisture, and ozone from MODIS infrared radiances, in *Third International Asia-Pacific Environmental Remote Sensing Remote Sensing of the Atmosphere, Ocean, Environment, and Space*, edited by W. P. Menzel, W.-J. Zhang, J. Le Marshall,
815 and M. Tokuno, pp. 168–176, International Society for Optics and Photonics., 2003.
- Sellers, P. J.: Modeling the Exchanges of Energy, Water, and Carbon Between Continents and the Atmosphere, *Science* (80-.), 275(5299), 502–509, doi:10.1126/science.275.5299.502, 1997.
- Sellers, P. J., Dickinson, R. E., Randall, D. A., Betts, A. K., Hall, F. G., Berry, J. A., Collatz, G. J., Denning, A. S., Mooney, H. A., Nobre, C. A., Sato, N., Field, C. B. and Henderson-Sellers, A.: Modeling the exchanges of energy, water, and carbon
820 between continents and the atmosphere, *Science* (80-.), 275(5299), 502–509, doi:10.1126/science.275.5299.502, 1997.
- Senay, G. B., Friedrichs, M., Singh, R. K. and Velpuri, N. M.: Evaluating Landsat 8 evapotranspiration for water use mapping in the Colorado River Basin, *Remote Sens. Environ.*, 185, 171–185, doi:10.1016/j.rse.2015.12.043, 2016.
- Serbin, S. P., Singh, A., Desai, A. R., Dubois, S. G., Jablonski, A. D., Kingdon, C. C., Kruger, E. L. and Townsend, P. a.: Remotely estimating photosynthetic capacity, and its response to temperature, in vegetation canopies using imaging
825 spectroscopy, *Remote Sens. Environ.*, 167, 78–87, doi:10.1016/j.rse.2015.05.024, 2015.
- Singh, R. K., Irmak, A., Irmak, S. and Martin, D. L.: Application of SEBAL Model for Mapping Evapotranspiration and Estimating Surface Energy Fluxes in South-Central Nebraska, *J. Irrig. Drain. Eng.*, 134(3), 273–285, doi:10.1061/(ASCE)0733-9437(2008)134:3(273), 2008.
- Su, Z.: The surface energy balance system (SEBS) for estimation of turbulent heat fluxes, *Hydrol. Earth Sys. Science*, 6(1),



- 830 85–99, 2002.
- Suyker, A. E., Verma, S. B., Burba, G. G., Arkebauer, T. J., Walters, D. T. and Hubbard, K. G.: Growing season carbon dioxide exchange in irrigated and rainfed maize, *Agric. For. Meteorol.*, 124(1–2), 1–13, doi:10.1016/j.agrformet.2004.01.011, 2004.
- Tian, H., Chen, G., Liu, M., Zhang, C., Sun, G., Lu, C., Xu, X., Ren, W., Pan, S. and Chappelka, A.: Model estimates of net
835 primary productivity, evapotranspiration, and water use efficiency in the terrestrial ecosystems of the southern United States during 1895–2007, *For. Ecol. Manage.*, 259(7), 1311–1327, doi:10.1016/j.foreco.2009.10.009, 2010.
- Turner, P. A., Baker, J. M., Griffis, T. J. and Venterea, R. T.: Impact of Kura Clover Living Mulch on Nitrous Oxide Emissions in a Corn–Soybean System, *J. Environ. Qual.*, 45(5), 1782, doi:10.2134/jeq2016.01.0036, 2016.
- USDA: National Engineering Handbook: Irrigation Guide, USA: Washington DC., 1997.
- 840 USDA: 2012 census of agriculture, USDA National Agricultural Statistics Service Washington, DC. [online] Available from: <https://www.agcensus.usda.gov/Publications/2012/>, 2014.
- USDA: Acreage. [online] Available from: <http://usda.mannlib.cornell.edu/usda/current/Acre/Acre-06-29-2018.pdf>, 2018.
- Verhoef, W.: Light scattering by leaf layers with application to canopy reflectance modeling: The SAIL model, *Remote Sens. Environ.*, 16, 125–141, doi:10.1016/0034-4257(84)90057-9, 1984.
- 845 Verhoef, W.: Earth observation modeling based on layer scattering matrices, *Remote Sens. Environ.*, 17(1931), 165–178, doi:10.1016/0034-4257(85)90072-0, 1985.
- Verrelst, J., Malenovsky, Z., van der Tol, C., Camps-Valls, G., Gastellu-Etchegorry, J. P., Lewis, P., North, P. and Moreno, J.: Quantifying Vegetation Biophysical Variables from Imaging Spectroscopy Data: A Review on Retrieval Methods, *Surv. Geophys.*, 1–41, doi:10.1007/s10712-018-9478-y, 2018.
- 850 Walker, A. P., Quaipe, T., van Bodegom, P. M., De Kauwe, M. G., Keenan, T. F., Joiner, J., Lomas, M. R., MacBean, N., Xu, C., Yang, X. and Woodward, F. I.: The impact of alternative trait-scaling hypotheses for the maximum photosynthetic carboxylation rate (V_{cmax}) on global gross primary production, *New Phytol.*, 215(4), 1370–1386, doi:10.1111/nph.14623, 2017.
- Wan, Z., Zhang, K., Xue, X., Hong, Z., Hong, Y. and Gourley, J. J.: Water balance-based actual evapotranspiration
855 reconstruction from ground and satellite observations over the conterminous United States, *Water Resour. Res.*, 51(8), 6485–6499, doi:10.1002/2015WR017311, 2015.
- Wang, H., Prentice, I. C., Davis, T. W., Keenan, T. F., Wright, I. J. and Peng, C.: Photosynthetic responses to altitude: an explanation based on optimality principles, *New Phytol.*, 213(3), 976–982, doi:10.1111/nph.14332, 2017.
- Wang, K. and Dickinson, R. E.: A review of global terrestrial evapotranspiration: observation, modelling, climatology, and
860 climatic variability, *Rev. Geophys.*, 50(2011), 1–54, doi:10.1029/2011RG000373.1.INTRODUCTION, 2012.
- Wei, J., Chen, Y., Gu, Q., Jiang, C., Ma, M., Song, L. and Tang, X.: Potential of the remotely-derived products in monitoring ecosystem water use efficiency across grasslands in Northern China, *Int. J. Remote Sens.*, 00(00), 1–21, doi:10.1080/01431161.2019.1587208, 2019.



- Weiss, M., Baret, F., Myneni, R., Pragnère, A. and Knyazikhin, Y.: Investigation of a model inversion technique to estimate canopy biophysical variables from spectral and directional reflectance data, *Agronomie*, 20(1), 3–22, doi:10.1051/agro:2000105, 2000.
- Whitley, R., Beringer, J., Hutley, L. B., Abramowitz, G., De Kauwe, M. G., Duursma, R., Evans, B., Haverd, V., Li, L., Ryu, Y., Smith, B., Wang, Y.-P., Williams, M. and Yu, Q.: A model inter-comparison study to examine limiting factors in modelling Australian tropical savannas, *Biogeosciences*, 13(11), 3245–3265, doi:10.5194/bg-13-3245-2016, 2016.
- Whitley, R., Beringer, J., Hutley, L. B., Abramowitz, G., De Kauwe, M. G., Evans, B., Haverd, V., Li, L., Moore, C., Ryu, Y., Scheiter, S., Schymanski, S. J., Smith, B., Wang, Y.-P., Williams, M. and Yu, Q.: Challenges and opportunities in modelling savanna ecosystems, *Biogeosciences*, 14, 4711–4732, doi:10.5194/bg-2016-190, 2017.
- Wilson, T. B. and Meyers, T. P.: Determining vegetation indices from solar and photosynthetically active radiation fluxes, *Agric. For. Meteorol.*, 144(3–4), 160–179, doi:10.1016/j.agrformet.2007.04.001, 2007.
- Wright, C. K. and Wimberly, M. C.: Recent land use change in the Western Corn Belt threatens grasslands and wetlands, *Proc. Natl. Acad. Sci.*, 110(10), 4134–4139, doi:10.1073/pnas.1215404110, 2013.
- Wu, P., Shen, H., Zhang, L. and Göttsche, F. M.: Integrated fusion of multi-scale polar-orbiting and geostationary satellite observations for the mapping of high spatial and temporal resolution land surface temperature, *Remote Sens. Environ.*, 156, 169–181, doi:10.1016/j.rse.2014.09.013, 2015.
- Xu, J., Liang, Y., Liu, J. and Huang, Z.: Multi-frame super-resolution of gaofen-4 remote sensing images, *Sensors*, 17(9), doi:10.3390/s17092142, 2017.
- Xu, M., Liu, R., Chen, J. M., Liu, Y., Shang, R., Ju, W., Wu, C. and Huang, W.: Retrieving leaf chlorophyll content using a matrix-based vegetation index combination approach, *Remote Sens. Environ.*, 224(February), 60–73, doi:10.1016/j.rse.2019.01.039, 2019.
- Yang, Y., Anderson, M. C., Gao, F., Wardlow, B., Hain, C. R., Otkin, J. A., Alfieri, J., Yang, Y., Sun, L. and Dulaney, W.: Field-scale mapping of evaporative stress indicators of crop yield: An application over Mead, NE, USA, *Remote Sens. Environ.*, 210(January), 387–402, doi:10.1016/j.rse.2018.02.020, 2018.
- Yebra, M., Van Dijk, A., Leuning, R., Huete, A. and Guerschman, J. P.: Evaluation of optical remote sensing to estimate actual evapotranspiration and canopy conductance, *Remote Sens. Environ.*, 129, 250–261, doi:10.1016/j.rse.2012.11.004, 2013.
- Zhang, K., Kimball, J. S. and Running, S. W.: A review of remote sensing based actual evapotranspiration estimation, *Wiley Interdiscip. Rev. Water*, 3(6), 834–853, doi:10.1002/wat2.1168, 2016a.
- Zhang, X., Liang, S., Wang, G., Yao, Y., Jiang, B. and Cheng, J.: Evaluation of the reanalysis surface incident shortwave radiation products from NCEP, ECMWF, GSFC, and JMA using satellite and surface observations, *Remote Sens.*, 8(3), doi:10.3390/rs8030225, 2016b.
- Zhang, Y., Guanter, L., Berry, J. a, Joiner, J., van der Tol, C., Huete, A., Gitelson, A., Voigt, M. and Köhler, P.: Estimation of vegetation photosynthetic capacity from space-based measurements of chlorophyll fluorescence for terrestrial biosphere



- models., *Glob. Chang. Biol.*, 20(12), 3727–42, doi:10.1111/gcb.12664, 2014.
- 900 Zhang, Y., Guanter, L., Joiner, J., Song, L. and Guan, K.: Spatially-explicit monitoring of crop photosynthetic capacity through the use of space-based chlorophyll fluorescence data, *Remote Sens. Environ.*, 210(June 2017), 362–374, doi:10.1016/j.rse.2018.03.031, 2018.
- Zheng, Y., Ren, H., Guo, J., Ghent, D., Tansey, K., Hu, X., Nie, J. and Chen, S.: Land Surface Temperature Retrieval from Sentinel-3A Sea and Land Surface Temperature Radiometer, Using a Split-Window Algorithm, *Remote Sens.*, 11(6), 650, doi:10.3390/rs11060650, 2019.
- 905 Zhong, L., Hu, L., Yu, L., Gong, P. and Biging, G. S.: Automated mapping of soybean and corn using phenology, *ISPRS J. Photogramm. Remote Sens.*, 119, 151–164, doi:10.1016/j.isprsjprs.2016.05.014, 2016.
- Zhu, X., Chen, J., Gao, F., Chen, X. and Masek, J. G.: An enhanced spatial and temporal adaptive reflectance fusion model for complex heterogeneous regions, *Remote Sens. Environ.*, 114(11), 2610–2623, doi:10.1016/j.rse.2010.05.032, 2010.