Response to comments by Anonymous Referee #1:

General Comments:

The manuscript by Groh et al. aims primarily at evaluating if, and to what extent, changes in local climatic conditions among some German sites affect crop production and water use efficiency. The evaluation is carried out through a "space-for-time" (SFT) framework by moving soil monoliths contained in lysimeters in two locations subjected to different aridity index. Among the various outcomes of this study, the authors claim that a more efficient crop water use occurs under less optimal soil moisture conditions. The text reads well and is properly organized, although some parts are too wordy or seem going astray in describing the moving of the lysimeters. Figures and tables are satisfactory, but I suggest that the readability of Figs. 2 and 3 should be improved. As I will specify below, I have some concerns about the approach and modeling tool used, and the discussion of some results. Therefore, while the topic is of current interest for the HESS readership, this paper should not be accepted in its present form, requires major revisions or should be rejected altogether.

Response: The authors thank Referee #1 for reviewing our paper.

We improved the readability of the text and figures (see line 281 Fig. 2 and line 305 Fig. 3, tick mark labels) in the revised manuscript.

We included a clearer description of the transfer of lysimeter and if necessary, a shortening of the text (see Line 135 until 150).

The main concerns of Referee #1 on our manuscript refer to the space-for-time approach in the set-up and the used modeling approach. We think that our approach can be well justified and hope that our answers to the specific questions are convincing. To the best of our knowledge, we provide for the first time observations on water fluxes, crop yield, biomass, N-content of yield for different soil types, but each under different climatic conditions (wet & dry) following a modified space-for-time substitution.

Specific Comments:

1a) About the SFT approach. The authors employ the SFT approach in the context of moving the lysimeters from one location to another in Germany. SFT is not new, actually, and mostly used in Ecology, but some examples can be seen in the hydrologic literature (e.g. Scanlon et al., AWR 28:291-302; Troch et al., HESS 17:2209-2217). However, the way the authors have employed this approach does not

seem to follow a standard (I guess), and therefore I think that an evaluation is required to test its soundness in the context of the submitted paper.

Response: We modified the text from Line 134 – 151 to:

Local excavated lysimeters (i.e. intact soil monoliths) were transferred between the stations to subject them to different climate regimes so as to generate a crossed soilclimate setup according to the space for time (SFT) substitution approach. It should be noted, that we did not follow the SFT substitution as used in ecological (e.g. Pickett, 1989; Blois et al., 2013; Wogan and Wang, 2018) or hydrological studies (e.g. Scanlon et al., 2005; Troch et al., 2013). Typically such SFT studies assume that spatial and temporal variations are equivalent (Pickett, 1989). By translocating soils from one test site to another while keeping some of the lysimeters at their original site, we actually account for unsuspected effects from the past. In this way we eliminate effects caused by past local events such as disturbances, pedogenesis, or site management. This is in contrast to the standard SFT approach. The spatial transfer of intact soil monoliths in the lysimeters followed an assumed direction of climatic changes of increased temperature and precipitation. For this study, we considered all arable-land lysimeter at the central sites Bad Lauchstädt and Selhausen of the TERENO-SOILCan lysimeter network. Each central experimental site contains three replicates of soils from different locations: Bad Lauchstädt (BL; Haplic Chernozems, loess), Dedelow (Dd; Calcic Luvisols and Haplic Luvisols, glacial till), Sauerbach (Sb; Colluvic Regosols; colluvial deposits), and Selhausen (Se; Haplic Luvisols, loess) allowing to investigate the response of the corresponding soil type under different climates. The transfer of soils between the research stations imitates a change in climatic conditions and compares for identical soils the effects of different climatic conditions on crop yield and soil water fluxes with those at the original location. By transferring lysimeters between stations, the "climatic shift" is abrupt such that we are not able to follow the gradual changes of the soil ecosystem over time as suggested in standard SFT approaches. Instead, crop yield and fluxes for the same soil under different climatic conditions are compared.

1b) The "long-term" concept exerts a key role when applying the SFT approach, but in this study only thirty years of weather data are exploited (just a minimum from a meteorological viewpoint) and then only six years are considered for the analysis (from 2012 to 2017). In view of this, I suggest that questions about "climate change" should be left out of this paper, whereas at least the authors might deal with possible changes, if any, in weather seasonality (e.g. a prolonged dry period or wet period, namely anomalies with respect to what observed during the 30 years of records).

Response: We agree that the available observation period is too short for questions about climate change and we clarified this within the revised manuscript and refer it

to change in climatic conditions or weather seasonality. However, we will also note that moving soils in regions with different climatic conditions gives us a perspective to evaluate the impact of changing climatic conditions and the longer the observation period the better. TERENO-SOILCan is an ongoing project to monitor the soil-plantatmosphere-continuum. Considering that lysimeter operations are expensive, require relatively high maintenance, such that sustainable quality lysimeter data are still limited.

1c) If longer time series of weather data were available (but 30 years could be used just like a threshold), plots of "Standardized Precipitation Index, SPI" or "Standardized Precipitation Evapotranspiration Index, SPEI" would definitely help. By the way, Walter and Lieth's climate diagrams for only six years is not a good practice.

Response: The SPI or SPEI would help to quantify the extent of the drought in 2015. However, several studies across Europe, including our region, have shown the extent of the drought in 2015. Thus we don't think that plots of SPI or SPEI would add much information to our investigation. We used the Walter and Lieth's diagram to compare between the climate conditions at the two sites. Although 6-7 years is a relatively short time, the diagram shows relative differences between both sites during the observation period. Nevertheless, we have added two additional subplots to Fig. 1 in the revised manuscript (see line 171) that describe the longer term climate conditions (1988-2017) according to Walter and Lieth.

2) About the modeling tool. My view is that the topic coved in the paper is addressed more effectively if one looks at the derivative of the system dynamics and not at its integral behavior. In order to translate my comment in modeling terms, what I am suggesting is the use of a Richards-based model instead of the bucketing type approach expressed by Eq.(1). Giving a look at the paper by Pütz et al. (2016), I see that the lysimeters are fully equipped with soil water content and potential sensors, together with other sensing devices. Therefore, I am wondering why the authors did not exploit the potential of this information to use the Richards equation.

Response: We don't agree with the Referee #1 in this point and we think that this is a misunderstanding of our intention of the manuscript. We actually provide a first comprehensive data set covering a nearly seven-year period, which can be used in a next step to model the soil-plant atmosphere system. Equation 1 describes how we obtain the soil water storage change directly from the raw data and the evapotranspiration was obtained using the water balance (Equation 4). We do not consider this as a modeling tool because it does not require any assumptions on soil hydraulic properties as would be the case when using Richards' equation. It should

be noted, that although a considerable amount of instruments was installed in the lysimeters, a considerable uncertainty is involved when trying to derive hydraulic properties from these measurements as previously reported by Groh *et al.* (2018). For the questions addressed in this paper we are directly using the measured data to calculate the water balance, thereby avoiding any uncertainties that would be introduced by model assumptions.

3) Concerns about determining ET My comment in this point 3) is linked somehow to the previous point 2). The use of ET, instead of making the partitioning of this variable in actual evaporation (Ea) and actual transpiration (Ta), can be something that may strongly limit the understanding of the ongoing processes and might yield erroneous outcomes. The use of the bucket model of Eq.(1) does not account for the important aspect of evaluating the possible onset of stress conditions in the crops and hence the computations of water use efficiency. The concept of "available water" or "readily available water" (as implied by Eq.(1), if I understood well) is definitely not adequate for the objectives of this paper. The plant can be under stress conditions due to the atmospheric demand even if a good amount of soil water is in the soil profile. Water transport resistances into the plant also play a key role. Moreover, what about possible physiological reactions of the vegetation when moving the lysimeter from one location to another? Did the authors check this aspect? Usually, vegetation shows some sort of resilience to its moving, at least during the initial stage of this moving. Can the authors comment on those points?

Response: We did not mention that we used the concept of "available water" or "readily available water". But we agree that the partitioning between E and T would be helpful to clarify further findings from our investigation, but this was beyond the scope of our study. The components E and T cannot be separated by Equation 1, which is used here only to determine the changes in soil water storage from the lysimeter mass data. We are aware that there are different methods to determine water use efficiency and discussed this in line 86 to 92 (e.g. use of T instead of ET), but we think that E is always related to the crop specific development and management and hence represents a kind of "cropping system" water use efficiency rather than plant water use efficiency at the leaf level. Although we did not monitor the crop stress status, the yield data provided an indication of the cumulative stress that was experienced by the crop. The Referee #1 mentioned possible physiological reactions of the vegetation when moving the lysimeter from one to another location. Please note that the soil monoliths were bare during the transfer and we used annual crops. The crop rotation was identical at both sites.

Groh, J., Stumpp, C., Lücke, A., Pütz, T., Vanderborght, J., Vereecken, H., 2018. Inverse Estimation of Soil Hydraulic and Transport Parameters of Layered Soils from Water Stable Isotope and Lysimeter Data. Vadose Zone Journal 17.

Response to comments by Anonymous Referee #2

General Comments:

The manuscript by Groh et al. presents results from the lysimeter network SOILCan. The focus of this study is on the effects of different weather and soil texture conditions on crop water use efficiency using a space-for-time approach. Hereby, weighable lysimeters with soils from four sites were moved and monitored at two of the sites with a drier and wetter climate, respectively. Instead of assessing changes in soil water storage as a residual of the water balance components the changes in lysimeter weights were used to avoid an accumulation of errors. One of the main outcomes was that the water-use-efficiency was improved (due to lower evaporation loss from soils) under drier soil moisture conditions not following a linear function. Further, the effects of drought were still visible in the following season and even beyond that especially on finer-textured soils. Overall, the manuscript reads very well with a logical structure. The manuscript deals with the very relevant topic of changing climate conditions on agricultural productivity. The combination of weighable lysimeters in a space-for-time approach investigating four different soils with data over seven years provides valuable and interesting insights on how crop production may be affected. One of the strengths of this MS is that the authors present a comprehensive data set covering a seven-year period. The measurement data can be used for model development, calibration and validation. I recommend that the authors present such a model study in a follow-up paper. I recommend the acceptance of the manuscript upon minor revisions.

<u>Response</u>: The authors thank Referee #2 for reviewing our paper and their positive feedback/ comments concerning our manuscript. We are currently conducting a study and use lysimeter data for the calibration of different crop models.

Specific Comments:

M&M section information about the soil texture of all four soil would be helpful as it later becomes an important part in the discussion (coarser vs finer textured soils)

<u>Response</u>: We added as suggested a soil profile description in the supplement of the revised manuscript (see Table A1; line 486).

Figure 2 Please improve readability. Tick mark labels are very small

<u>Response</u>: We changed the tick mark labels to improve the readability of Figure 2 and 3 (see line 281 and 305 in the revised manuscript)

L24 & L123 'monitored from April 2011 until December 2018' versus 'lysimeter data from April 2011 until December 2017' Please clarify.

Response: We changed December 2018 to December 2017 in line 23

L244-264 Could this be related to a higher infiltration capacity of the coarser textured soil allowing for a more rapid recharge? It would be interesting if the authors made any observations on silting, cracking etc. of the soil surfaces especially of the finer-textured soils which might explain deficiencies in soil water recharge.

Response: The infiltration capacity is dependent on the conductivity at the soil surface. Silting, which more often occur at the soil surface of fine textured soil, affects the macropore structure (destruction of soil aggregates) and reduce the infiltration. No surface runoff was observed during the observation period. Thus we don't think that the annual carry –over of soil moisture deficits are related to a different infiltration capacity of the soil.

Some qualitative observations were made during the harvest time, but the soil surface has been modified by tillage, and the topsoil organic matter content and the plant roots are counteracting silting and cracking. We included this information and discussion in the revised paper (see line 268 - 274).

L410 '. . .net fluxes were observed. . .'

<u>Response</u>: We changed "net fluxes observed" to "net fluxes were observed" in the revise manuscript (see line 438 in the revised manuscript)

Responses of soil water storage and crop water use efficiency to changing climatic conditions: A lysimeter-based space-for-time approach

Jannis Groh^{1, 2}, Jan Vanderborght², Thomas Pütz², Hans-Jörg Vogel³, Ralf Gründling³, Holger Rupp³, Mehdi Rahmati⁴, Michael Sommer^{5, 6}, Harry Vereecken², Horst H. Gerke¹

¹Research Area 1 "Landscape Functioning", Working Group "Hydropedology", Leibniz Centre for Agricultural Landscape Research (ZALF), Müncheberg, 15374, Germany
 ²Institute of Bio- and Geoscience IBG-3: Agrosphere, Forschungszentrum Jülich GmbH, Jülich, 52425, Germany
 ³Department of Soil Physics, Helmholtz Centre for Environmental Research — UFZ, Halle (Saale), 06120, Germany

⁴Department of Soil Science and Engineering, Faculty of Agriculture, University of Maragheh, Maragheh, Iran ⁵Research Area 1 "Landscape Functioning", Working Group "Landscape Pedology", Leibniz Centre for Agricultural Landscape Research (ZALF), Müncheberg, 15374, Germany ⁶Institute of Environmental Science and Geography, University of Potsdam, Potsdam, 14476, Germany

Correspondence to: Jannis Groh (groh@zalf.de or j.groh@fz-juelich.de)

15

5

Abstract. Future crop production will be affected by climatic changes. In several regions, the projected changes in total rainfall and seasonal rainfall patterns will lead to lower soil water storage (SWS) which in turn affects crop water uptake, crop yield, water use efficiency, grain quality and groundwater recharge. Effects of climate change on those variables depend on the soil properties and were often estimated based on model simulations. The objective of this study was to investigate the

20 response of key variables in four different soils and for two different climates in Germany with different aridity index: 1.09 for the wetter (range: 0.82 to 1.29) and 1.57 for the drier climate (range: 1.19 to 1.77), by using high-precision weighable lysimeters. According to a "space-for-time" concept, intact soil monoliths that were moved to sites with contrasting climatic conditions have been monitored from April 2011 until December 201<u>7</u>.

Evapotranspiration was lower for the same soil under the relatively drier climate whereas crop yield was significantly higher,

- 25 without affecting grain quality. Especially 'non-productive' water losses (evapotranspiration out of the main growing period) were lower which led to a more efficient crop water use in the drier climate. A characteristic decrease of the SWS for soils with a finer texture was observed after a longer drought period under a drier climate. The reduced SWS after the drought remained until the end of the observation period which demonstrates carry-over of drought from one growing season to another and the overall long term effects of single drought events. In the relatively drier climate, water flow at the soil profile
- 30 bottom showed a small net upward flux over the entire monitoring period as compared to downward fluxes (ground water recharge) or drainage in the relatively wetter climate and larger recharge rates in the coarser- as compared to finer-textured soils. The large variability of recharge from year to year and the long lasting effects of drought periods on SWS imply that long term monitoring of soil water balance components is necessary to obtain representative estimates. Results confirmed a

more efficient crop water use under less optimal soil moisture conditions. Long-term effects of changing climatic conditions

35 on the SWS and ecosystem productivity should be considered when trying to develop adaptation strategies in the agricultural sector.

1 Introduction

The amount of water stored within the root zone of the soil and the vadose zone is a central and characteristic component of terrestrial ecosystems. Soil water storage (SWS) is important for provisioning (e.g., crop production, water balance, and
plant available nutrients) as well as regulating and supporting ecosystem services (e.g. water, nutrients, climate, flood, drought; Adhikari and Hartemink, 2016; Vereecken et al., 2016). The SWS capacity (SWSC) depends on soil texture, organic matter content, bulk density, and soil structure and is related to the effective field capacity, which can be derived from the soil water retention function (Vereecken et al., 2010). The knowledge on magnitude and temporal variation of the SWS is essential for understanding ecological and hydrological processes and to manage ecosystems (Cao et al., 2018).
Climate change will modify the temporal availability of soil water, increase the frequency and duration of droughts, affecting the quantity and quality of aquifer recharge and might affect crop production. Thus future ecosystem productivity (e.g. crop yield) is expected to respond to changes in weather (short-term) and climate (long-term), because it will alter the crop water balance components, such as SWS, evapotranspiration (ET) and drainage (Yang et al., 2016). How to produce more crop yield with less water is a major challenge in agriculture, because i) water is a limiting factor for crop production in many

50 regions of the world, and ii) predictions of future climate indicate an increasing water limitation for crop production caused by reduced rainfall and changing seasonal rainfall distribution (Lobell and Gourdji, 2012). Several studies have been conducted to investigate the impact of global climate change on crop water balance components (Sebastiá, 2007; Wu et al., 2015) and crop or grain yield (Ewert et al., 2002; Zhao et al., 2016; Schauberger et al., 2017;

Asseng et al., 2019). Understanding the impact of weather signals on the agricultural productivity is of crucial importance

- 55 for managing future crop production, since variations in weather conditions could explain much of the yield variability (Frieler et al., 2017). Temperature rise and changing seasonal rainfall patterns could alter the probability of droughts and affect freshwater resources (Gudmundsson and Seneviratne, 2016; Gudmundsson et al., 2017). Negative impacts of rising temperature on the yield of major crops at the global scale (Asseng et al., 2014; Zhao et al., 2017) are highlighting the potential vulnerability of agricultural productivity to climate change. Schauberger et al. (2017) showed a consistent negative
- 60 response of US crops under rainfed conditions being mainly related to water stress induced by higher temperatures. In addition to the direct effects of a temperature rise, an elevated atmospheric CO₂-concentration, and changes in rainfall amounts on crop yield (Ewert et al., 2002; Asseng et al., 2014; Gammans et al., 2017; Scheelbeek et al., 2018), the higher temperatures could affect crop yields indirectly. Indirect effects caused by increasing the atmospheric water demand, limiting ET due to water stress and reducing the SWS, could in turn lead to a decrease in crop yield (Zhao et al., 2016; Zhao et al.,

- 65 2017). Thus investigating the response of crop water balance components and yield to climate change is important to develop suitable adaptation and mitigation strategies (Albert et al., 2017; Rogers et al., 2017). Previous studies reported estimates of crop water balance components and crop yield mostly based on either manipulative experiments or observational studies to predict the ecological response of crops to climate change (Yuan et al., 2017). Wu et al. (2015) showed that the inter-annual variation of the SWS at northern middle and high latitudes increased under a warmer
- 70 climate with higher values during the wetter and lower values of the SWS during the drier season. In this case, the frequency of water logging events or soil crack formation will increase and probably alter soil properties such as macroporosity and SWSC and thus affect vadose zone hydrology at different scales (Robinson et al., 2016; Hirmas et al., 2018). Robinson et al. (2016) showed for a manipulative long-term experiment that intense summer droughts altered the soil water retention characteristic and lowered the SWSC.
- 75 Nevertheless, current knowledge on changes of SWS are still limited mostly to the analysis of soil moisture observations related to restricted soil volumes and soil moisture ranges (Mei et al., 2019; Yost et al., 2019). As an alternative method, weighable lysimeters allow the direct observation of SWS by monitoring the temporal changes of the total soil mass in mostly cylindrical containers. However, the use of weighable lysimeters was often limited in the past to the quantitative determination of the water balance components of precipitation (P), evapotranspiration (ET), and subsurface inflow (Q_{in}) and
- 80 outflow (Q_{out} ; e.g. drainage); the change of SWS was obtained as residual of the water balance components (e.g. Herbrich et al., 2017; Groh et al., 2018b). This approach accumulated all possible errors introduced by other components into the SWS, causing a relatively low precision. The direct derivation of SWS from lysimeter mass changes could provide a new perspective on the use of lysimeter data as an additional model calibration variable and for lysimeters that are large enough to fully capture the complete soil profile with the relevant soil horizons and intact soil structures to be representative for the 85 pedon scale.
- The water use efficiency (WUE) links the carbon and water exchange between vegetated soil and the atmosphere (Niu et al., 2011). Several definitions have been used to describe the WUE at the leaf or ecosystem level (for more details see Zhou et al., 2017). At the ecosystem level, WUE defined as the ratio between grain yield or total biomass and the water lost to the atmosphere by ET_(Fan et al., 2018) is one possibility to quantify the impact of changes in environmental conditions and of management decisions (e.g. irrigation) on agricultural productivity. The use of ET instead of transpiration (T) only for calculating WUE represents a water use efficiency at the ecosystem rather than leaf level because it accounts for evaporation, which is also depending on crop specific development and soil management. The WUE provides insights to better manage and understand the productivity and ecological functioning of agricultural ecosystems (Zhang et al., 2015). The prevailing general hypothesis for WUE is that plant productivity increases with increasing water use (ET; Hatfield and Dold, 2019), which implies that WUE efficiency is a linear function of the water used by a crop to produce grain yield or the total above ground biomass. But several studies have shown that crop WUE was negatively correlated with annual rainfall and plants
 - 3

achieved their maximum crop WUE under less favourable soil water availability (Zhang et al., 2010; Ponce-Campos et al., 2013; Xiao et al., 2013; Zhang et al., 2015). The last statement might imply that plants are able to adapt their water use

during drought conditions by improving their WUE or that there is simply less non-productive water losses by evaporation.

- 100 Nevertheless, temperature above a certain threshold (extremely high temperature) especially during the reproductive period (Gourdji et al., 2013) or due to drought and heat stress reduce yield. However, such investigations are often focused on one specific environmental variable (e.g. *P* or temperature) in manipulation experiments. This basically ignores joint effects of several climate variables on the crop WUE in climate impact research studies. The impact of altered climatic conditions on different agricultural ecosystems within manipulative experiments has not been thoroughly studied yet; due to problems to
- 105 either realistically manipulate the climatic conditions at a specific site or to move an intact soil to another site with contrasting climatic conditions.

Here, we hypothesize that WUE will not increase for drier climate; because a change in plant productivity will simultaneously alter the water use (ET) and thus describe WUE as a linear function between both variables. In addition we wanted to test if observed lysimeter mass changes can be used to monitor the long-term change of SWS, which might be in

- 110 addition to water flux observation a useful dataset for the calibration of vadose zone models. We used observations from a German soil-climate crossed factorial experiment (TERENO-SOILCan; Pütz et al., 2016). The lysimeter network of TERENO-SOILCan has been initiated to assess effects of climatic changes on arable and grassland soil ecosystems including the water balance components (ET, SWS, net drainage) and crop characteristics including yield, yield quality and WUE. As part of this project, arable-land lysimeters filled with four different soils were transferred within and between
- 115 TERENO observatories (space-for-time; see details in Pütz et al., 2016) to expose soils from originals sites to other climatic conditions. The space-for-time approach means that soils are translocated in space instead of waiting at the same location for changes in climatic conditions in time. The concept initially intended to evaluate the impact of climate on agricultural ecosystems (Pütz et al., 2016). It represents basically a crossed soil type and climate experimental setup that could allow quantifying changes in the soil water balance and the crop production as response to imposed variations in climatic
- 120 conditions. Results from this experimental setup can primarily be used to evaluate models that predict changes in response to possible future climatic conditions.

Our objectives were: i) to develop an approach to obtain time series of changes in SWS directly from lysimeter data, ii) to determine the other soil water balance components (*P*, ET, inflow and drainage) of soils each exposed to two different climates, iii) to compare the net flux (inflow and drainage)/SWS dynamics for the same soils in relatively dry and wet climates and iv) to test the hypothesis that WUE of crops remains constant under changing climatic conditions in these

crossed soil type and climate experiment. The analysis was based on lysimeter data from April 2011 until December 2017.

2 Material and Methods

2.1 Site descriptions

- 130 The study was conducted at the experimental field sites Selhausen (50°52′7′N, 6°26′58′E) and Bad Lauchstädt (51°23′37′N, 11°52′41′E), which are part of the Eifel/Lower Rhine Valley and the Harz/Central German Lowland Observatory of TERENO in Germany (Wollschläger et al., 2016; Bogena et al., 2018), respectively. The TERENO-SOILCan lysimeter network was established at several experimental stations across a rainfall and temperature gradient. Local excavated lysimeters (i.e. intact soil monoliths) were transferred between the stations to subject them to different
- 135 climate regimes so as to generate a crossed soil-climate setup according to the space for time <u>(SFT)</u> substitution approach. It should be noted, that we did not follow the SFT substitution as used in ecological (e.g. Pickett, 1989; Blois et al., 2013; Wogan and Wang, 2018) or hydrological studies (e.g. Scanlon et al., 2005; Troch et al., 2013). Typically such SFT studies assume that spatial and temporal variations are equivalent (Pickett, 1989). By translocating soils from one test site to another while keeping some of the lysimeters at their original site, we actually account for unsuspected effects from the past. In this
- 140 way we eliminate effects caused by past local events such as disturbances, pedogenesis, or site management. This is in contrast to the standard SFT approach. The spatial transfer of intact soil monoliths in the lysimeters followed an assumed direction of climatic changes of increased temperature and precipitation. For this study, we considered all arable-land lysimeter at the central sites Bad Lauchstädt and Selhausen of the TERENO-SOILCan lysimeter network. Each central experimental site contains three replicates of soils from different locations: Bad Lauchstädt (BL; Haplic Chernozems, loess),
- 145 Dedelow (Dd; Calcic Luvisols and Haplic Luvisols, glacial till), Sauerbach (Sb; Colluvic Regosols; colluvial deposits), and Selhausen (Se; Haplic Luvisols, loess) allowing to investigate the response of the corresponding soil type under different climates. The transfer of soils between the research stations imitates a change in climatic conditions and compares for identical soils the effects of different climatic conditions on crop yield and soil water fluxes with those at the original location. By transferring lysimeters between stations, the "climatic shift" is abrupt such that we are not able to follow the
- 150 gradual changes of the soil ecosystem over time as suggested in standard SFT approaches. Instead, crop yield and fluxes for the same soil under different climatic conditions are compared. Further information on soil texture and the transfer of soil monoliths from the TERENO-observatories to the central sites can be taken from Table A1 to A2 (see appendix). The transferred eroded Luvisol soil monoliths from Dedelow have a varying soil depth to the clay illuviation horizon (B_t) and to the marly, illitic glacial till (*C*-horizon). They represent part of the erosion gradient typically observed in agricultural
- 155 landscapes of hummocky ground moraines (Sommer et al., 2008; Rieckh et al., 2012; Herbrich et al., 2017). Detailed information about the lysimeter design and general experimental-set up of TERENO-SOILCan can be found in Pütz et al. (2016). The climatic conditions of the central sites from 1 January 2012 to 31 December 2017 (complete years) are shown in Fig. 1 according to Walter and Lieth (1967). Although the patterns in average monthly temperature values are relatively similar at both sites (Fig. 1), a more pronounced amplitude of the temperature variations over the year could be found in Bad
- 160 Lauchstädt (representing a more continental climate) as compared to the more temperate and humid climate (sub-oceanic or

sub-Atlantic) in Selhausen (Fig. 1). The average annual grass reference evapotranspiration (ET₀) calculated with the FAO56 Penman-Monteith method (Allen et al., 2006) is slightly higher at Bad Lauchstädt (710 mm) than at Selhausen (694 mm). Larger differences are shown in the annual rainfall and the rainfall distribution over the year (Fig. 1). The lower annual *P* in Bad Lauchstädt (458 mm) than in Selhausen (644 mm) corresponds with a higher aridity index (AI = ET₀ P^{-1} , see data

165 repository) of 1.57 for Bad Lauchstädt than for Selhausen (1.09). The rainfall distribution over the year was more uniform in Selhausen whereas the probability of relatively dry periods in spring (April) and late summer (September) was higher in Bad Lauchstädt. Thus, the climatic conditions at the SOILCan experimental sites can be defined as drier for Bad Lauchstädt and wetter at Selhausen, which corresponds well to long-term weather station data, from stations at Bernburg / Saale (Nord; German weather service) and at Forschungszentrum Juelich (see Fig.1 c and d).

170



subplot a) indicate months were frost is likely to occur.

Figure 1: Climate diagrams according to Walter and Lieth (1967) for Bad Lauchstädt (a), and Selhausen (b) for 2012 to 2017 and Bernburg/Saale (Nord) (c), and Forschungszentrum Juelich (d) for 1981 to 2017. Data were obtained from the SOILCan weather stations at Selhausen and a climate station at Bad Lauchstädt above sea level (asl.). The long-term weather data were taken from the weather stations of the German weather service at Bernburg / Saale (Nord) and the Forschungszentrum Juelich. The blue bars at the bottom of

175

2.2 Soil water storage (SWS)

Monthly changes in SWS (Δ SWS) were calculated from lysimeter observations as:

$$\Delta SWS = \Delta W + \Delta L_{yscor} \tag{1}$$

180 where ΔW is the monthly lysimeter mass change, and ΔL_{yscor} corresponds to mass changes by maintenance, harvesting, or other disturbances that occur accidently (e.g. erroneous load cells) or naturally (e.g., animals). The variable ΔW was directly obtained by analysing lysimeter mass data (average value: 12°AM until 2°AM) defined as:

$$\Delta W = W_{i+1} - W_i \tag{2}$$

where *W* is the lysimeter mass at the beginning of month *i*. The variable ΔL_{yscor} was determined from monthly changes of lysimeter mass during maintenance work. Less than 0.6 % of Δ SWS values could not be calculated, because lysimeter mass data at the beginning of the corresponding month were missing. A linear regression model obtained for the entire time series between Δ SWS of the soils was used for interpolation to fill the gaps. This was first based on Δ SWS from surrounding lysimeters of the same soil type and if not available, then the average values of Δ SWS obtained from all available lysimeters at the respective station were used.

190 2.3 Crop water use efficiency (WUE), grain yield and yield quality

In total 12 arable land lysimeters (three replicates of four soil types) with a surface area of 1 m^2 and a depth of 1.5 m were embedded within larger fields at the respective central experimental site at Selhausen (250 m^2) and Bad Lauchstädt (720 m^2). The same crops were grown and identical tillage and crop management procedures were carried out at both sites and in the field around the lysimeters. The lysimeters were cultivated with peas (Pisum sativum L.; cultivar: Mascara), winter barley

- (Hordeum vulgare L.; cultivar: Lomerit), winter canola (Brassica napus L.; cultivar: Adriana), oat (avena sativa L.; cultivar: Max G), winter wheat (Triticum aestivum L.; cultivar: Glaucus), winter barley (Hordeum vulgare L., cultivar: Antonella) and winter rye (Secale cereal L.; cultivar: SU Santini), whereas the applications of seasonal plant protection, crop growth regulators and nitrogen-fertilizer (see appendix Table <u>A3</u>) have been adapted to local farmer conditions at the respective experimental site. Dry mass of the yield and plant residual matter were gravimetrically determined with a precision balance
- 200 (Selhausen: EMS 6K0.1, KERN, Balingen-Frommern, Germany; Bad Lauchstädt: LC 6200 D, Satorius, Göttingen, Germany) after drying at 75°C for 24 hours (Bad Lauchstädt) and at 60°C for >24 hours (Selhausen; until reaching a constant weight). The determination of total nitrogen of the dry yield and plant residual material was obtained with an elementary analyser (VarioelCube, elementar, Langenselbold, Germany).

The following Eq. (3) was used to calculate the crop WUE (kg m^{-3}):

$$205 \quad \text{WUE} = \frac{\gamma}{\text{ET}} \tag{3}$$

where Y is the grain yield (kg m⁻²), and ET (m³ m⁻²) is the measure of the consumed water during the growing season of the corresponding crop (Katerji et al., 2008). The growing periods of the crops were defined as the time between sowing and

harvest (see appendix Table <u>A3</u>). The required ET during the growing season was estimated based on the monthly water balance equation and observed precipitation (P) in mm per month as:

210 ET = $P - \Delta SWS - Q_{net} - \Delta L_{ysvol}$

(4)

where Q_{net} is the monthly sum of net water flux across the lysimeter bottom ($Q_{net} > 0$: drainage; $Q_{net} < 0$: capillary rise) and ΔL_{vsvol} is mass change determined from monthly soil water sampling volume. P was measured with a tipping bucket rain gauge (15189, Lambrecht, Göttingen, Germany) at Bad Lauchstädt (experimental station Bad Lauchstädt), and with a weighing rain gauge (Ott Pluvio2 L, Ott, Kempten, Germany) at Selhausen (Se_BDK_002). Data of the latter station is 215 available at TERENO data portal (http://teodoor.icg.kfa-juelich.de/ddp/index.jsp). The Ott rain gauge was installed in April 2013; data before April 2013 was estimated by linear regression models and P data from surrounding climate stations of the TERENO data portal (station names: SE_BDK_002; RU_BCK_003; RU_K_001; ME_BCK_001), which can be used to interpolate between the given data points. We used the R software (R-Core-Team, 2016) and the function Im of the package stats (R-Core-Team, 2016) to set-up linear regressions. The coefficient of determination (R^2) was used to determine the goodness of fit of the linear regression. A stepwise gap-filling approach was used to gap-fill missing P data after April 2013, 220 which consisted of an analysis of data from other meteorological stations that were operating and missing values, were filled based on the observation which had the highest R^2 . Monthly Q_{net} values were obtained from mass changes of the leachate from the lysimeters, collected with a weighable reservoir tank. The lysimeter bottom boundary pressure head condition was imposed by a pumping mechanism, which enabled either outflow or inflow according to differences in pressure head values

- at 1.4 m depth between lysimeter and surrounding field soil. This control of the bottom boundary allowed imitating the upward and downward water fluxes and representation of ET processes in lysimeters (Groh et al., 2016) more realistically and comparable to the intact soil profile. More technical details can be found in Pütz et al. (2016). Missing data in the time series of Q_{net} were filled for small gaps of about one minute by linear interpolation and for gaps between >1 and 10 minutes by using a moving average with a window width of 30 minutes. Larger gaps in the time series were filled by average water
- 230 flux values from other lysimeters of the same soil type. Nearly 5% of monthly ET values were found not plausible perhaps due to water loss by leaking during periods with water-saturated conditions at the lysimeter bottom. These conditions occurred mainly in winter, when monthly ET fluxes were in general relatively low as compared to summer conditions, so that potential error was low and easily detectable. A linear regression based on either single or average ET values from other non-affected lysimeters with similar soils were used for interpolation to fill the gaps. Detailed information on the monthly
- water balance data and missing data can be taken from the TERENO data portal (see section Data availability).

3 Results and Discussion

3.1 Soil water storage change

For the observation period (April 2011- January 2018), evapotranspiration (ET) and cumulative soil water storage change (Δ SWS) differed at both stations, Selhausen and Bad Lauchstädt, in amount and temporal development between transferred

- soils and those from the original site (Fig. 2). The variability in terms of the standard deviation of Δ SWS and ET of the three replicate soils were small and ranged across the different soils for Δ SWS between 4.3 and 7.4 mm and for ET between 3.7 and 5.5 mm. This clearly demonstrates that the differences in Δ SWS and ET between the same soils at original and new locations are larger than their scatter (Fig. 2). This suggests that uncertainty in the calculation of water fluxes were in general smaller than effects of transferring soils between the test sites. Larger deviations in Δ SWS between origin and transferred
- soils were visible for the crop winter canola after date of harvest in summer 2013 (soils from BL, Sb, and Se Fig. 2b, 2d, 2h) and winter barley 2016 (all soils). Largest depletions of SWS during the entire observation period could be observed for all soils during the spring-summer period (March and July) in 2015. At Bad Lauchstädt, the aridity index (AI = $ET_0 P^{-1}$) of 2.7 for March-July 2015 was larger as compared to the average AI value of 2.0 calculated for all March and July periods between 2012 and 2017. Also the value of the AI for Selhausen was with 2.0 slightly larger as compared to the average AI
- value of 1.6 for all March-July periods. The SWS depletion in 2015 was larger at both sites for soils from Bad Lauchstädt (BL; Fig. 2b) and Sauerbach (Sb; Fig. 2d) as compared to that of the other two soils from Dedelow (Dd; Fig. 2f) and Selhausen (Se; Fig. 2h). The Sb and BL soils were strongly desiccated by the winter wheat crop in 2015, which can be seen from ET June 2015 for BL and Sb of about 125 175 mm/month (Figs. 2a and 2c) was larger than for Dd and Se soils of about 100 125 mm/month (Figs. 2e and 2g) even for the soils exposed to the drier climate in Bad Lauchstädt. For the BL
- 255 (Fig. 2b) and Sb (Fig. 2d) soils, the amount of rainfall after the growing season of 327 mm (August 2015 April 2016) in Bad Lauchstädt was not sufficient to compensate for ET and drainage such that the soil profile did not return to a SWS capacity (i.e., typical spring moisture) at the end of the winter period characterized by a value close to 0 of the cumulative Δ SWS. The soil moisture deficit from 2015 was carried over to the growing seasons of 2016 and even of 2017. For the Dd and Se soils (Figs. 2f and 2h, the SWS deficit during the 2015 growing season under the climate of Bad Lauchstädt was less
- and the amount of precipitation after the growing season was sufficient for the soils to return to a typical SWS value although this value was reached later and not before the next spring. The AI of 1.77 at BL in 2015 (January-December) was considerably higher than the average AI for the 5 year period at BL (1.57). For the same year 2015, the AI was 1.13 at Se, and thus only slightly higher than the 5-years average AI-value of 1.09. For all soils in Se (blue lines in Figs 2b, 2d, 2f, 2h), the amount of precipitation after the growing season of 501 mm of 2015 (August 2015 April 2016) was sufficient for the
- lysimeters to return to their 'typical' SWS value at the end of the winter. These results indicate soil type dependent changes in SWS during drought periods. The annual carry-over of soil moisture deficits demonstrates the vulnerability towards drought risks even for finer-textured soils, despite having an overall larger SWSC than coarser-textured soils. This might be related to a higher infiltration capacity of the coarser-textured soil, which allows for a more rapid recharge. The infiltration capacity is dependent on the conductivity at the soil surface. Silting and cracking, which more often occur at the soil surface
- 270 of fine textured soils, affects the macropore structure (destruction of soil aggregates) and change the infiltration. However, no surface runoff was observed during the observation period and qualitative observations on cracking were made during the harvest time, but the soil surface has been modified by tillage, and the topsoil organic matter content and the plant roots are counteracting silting and cracking. This suggests that the annual carry-over of soil moisture deficits were not related to a

different infiltration capacity of the soil. The observed stronger depletion of soil water corresponds with soil drying reports from larger scale observations on the occurrence of a severe drought during the summer 2015, where effects of the drought has been observed from a climatological (Ionita et al., 2016) and hydrological (Laaha et al., 2016) perspective. The carryover of soil moisture deficits to the time after the drought at the local scale in Bad Lauchstädt agrees well with the results from Laaha et al. (2016), which showed for several stations in Europe that soil water storage (catchment scale) at the end of the study period (November 2015) has not totally recovered from the summer drought in 2015.





Figure 2: Monthly evapotranspiration (ET) and cumulative monthly changes in soil water storage (Δ SWS) from April 2011 until January 2018 at the lysimeter stations in Selhausen and in Bad Lauchstädt for soils from Bad Lauchstädt (a, b), Sauerbach (c, d), Dedelow (e, f), and Selhausen (g, h); mean values (dots) and standard deviations (error bars) are from 3 individual lysimeter monoliths of each soil. The background colour corresponds with the cropping periods at the TERENO-SOILCan lysimeters: bare soil (white) and crops (green).

285

290

Furthermore changing climatic conditions and a more frequent occurrence of drought could alter the SWSC because of the increasingly unavailable pore spaces due to different sources, including physical e.g. swelling and shrinking processes (te Brake et al., 2013; Herbrich and Gerke, 2017), biological e.g. vegetation induced soil desiccation that enhanced soil cracking (Robinson et al., 2016), biochemical e.g. enhanced organic matter mineralization, due to increasingly oxidation of the organic horizons during dry periods (Robinson et al., 2016), which will consequently result in a degradation of organic soil structure, or change in the soil wettability (Ellerbrock et al., 2005).

3.2 Net drainage

- 295 The water fluxes across the suction rake system at the lysimeter bottom in 1.5 m depth were cumulated to monthly net drainage fluxes (Q_{Mnet}) . The time series' of Q_{Mnet} for all soils at Se, the site with relatively wet climate, were in general directed downward during the winter months and upward (capillary rise) during spring and summer (Fig. 3). However, the magnitude of monthly fluxes Q_{Mnet} differed between the soil types (e.g. soils in Se for 2012 or 2013 see Fig.3); Q_{Mnet} for lysimeters with the coarser-textured soils from Dd (Fig. 3c) was mostly larger (e.g., drainage during bare fallow 2014) than
- 300
- for those with the finer-textured soils from BL (Fig. 3a), Sb (Fig. 3b), and Se (Fig. 3d). For the same soils under the relatively dry climate in BL, time series' of Q_{Mnet} were rather similar, with the largest values of upward fluxes for the soil from Dd (Fig. 3c). The magnitude of Q_{Mnet} for soils under BL climate was mostly smaller for drainage and larger for upward directed fluxes as compared to the Q_{Mnet} values for the soils under the wet climate in Selhausen.



305

Figure 3: Monthly net water fluxes across the lysimeter bottom in 1.5 m soil depth from April 2011 until January 2018 at the stations Selhausen and Bad Lauchstädt for soils from a) Bad Lauchstädt, b) Sauerbach, c) Dedelow, and d) Selhausen; mean values (dots) and standard deviation (error bars). Positive values are defined to drainage and negative values to upward direct water flux from capillary rise. Error bars indicate the variability of storage changes between individual lysimeters of each soil group. The background colour corresponds to different crops lysimeter cover types: bare soil (white) and different crops (green).

310

The Q_{Mnet} time series' (Fig. 3) demonstrate that weather conditions in 2015 impacted the soil water fluxes in the following years: Under the dry climate in BL, hardly any drainage was observed for all soils after 2015. This indicates that the soils remained so dry during the winter period that downward water percolation or groundwater drainage was limited. The lack of 315 water recharge during winter also affected the upward directed Q_{Mnet} flux rates in the following years, which generally decreased after 2015, especially for soils from BL and Sb. The nearly unchanged Q_{Mnet} values for the soils at BL after 2015 indicate that soil water saturation and dynamics is limited throughout the soil profile.

The annual net water fluxes (Q_{Anet}) at the bottom (in 1.5 m) of the same soils under the dry and wet climates are compared in

- form of scatterplots (Fig. 4). The scatterplots clearly show that fluxes were in general directed upward (i.e., negative values 320 of Q_{Anet} for soils under a dry climate in BL; positive values of Q_{Anet} (i.e. drainage) were only observed for 2011 and 2014 (Fig. 4). The larger values of Q_{Anet} for 2014 could be due to the lower ET after an earlier harvesting of the oat crop and a longer bare soil period without crop transpiration. The coarser-textured soils from Dedelow showed the largest range of Q_{Anet} values (from -78 mm to +164 mm) at the site with a relatively dry climate (BL) during the observation period of 2011-2017.
- This range could be explained by variation in soil water storage capacities between Dd soils, which depended on the 325 thickness of the upper soil horizons that were modified by soil erosion (Herbrich et al., 2017). The long-term average values of Q_{Anet} for all soils in the dry climate were negative and varied only in a small range (from -18 mm to -28 mm; see appendix Table A2). Long term negative groundwater recharge is only possible at sites where groundwater can be replenished, for instance, by lateral subsurface water flow. Whether the Q_{Anet} flux under the BL climate will continue to be negative for all
- 330 soils would require a longer time series. Nevertheless, a low and even negative groundwater recharge has not only an impact on the groundwater quantity, but it will also affect the groundwater quality. In case of a small net recharge, the concentrations of solutes from agricultural fertilizers, pesticides, and those of dissolved minerals and salts in the water-filled soil pores will become relatively high, and soil water movement still remains negligibly small. Thus under conditions of relatively small leaching rates, solutes including plant nutrients will largely be retained within the soil's root zone. Under
- long term conditions of net negative leaching, soils and soil horizons may accumulate carbonates (e.g., BL soil Haplic 335 Chernozems), or if leaching is small such that the carbonates from the topsoil horizons precipitate already in the subsoil within the 1.5 m soil monoliths like in the Ccv horizons in Dd subsoil of Calcic Luvisols (see soil profile descriptions in Herbrich and Gerke, 2017) and eventually salts.
- Q_{Anet} values under a relatively wet climate (in Se) were for all soils positive, indicating in general downward directed 340 drainage fluxes (Fig. 4). The long-term average Q_{Anet} values ranged between 49 to 119 mm (see appendix Table A2) depended on the soil type. The Q_{Anet} value was larger for the coarser-textured soil from Dd (Fig. 4c) as compared to the other soils. For 2013 (Winter Canola crop), the Q_{Anet} fluxes were negative for all finer-textured soils (i.e. Bad Lauchstädt, Sauerbach, and Selhausen, Fig. 4a, b, d), which might be related to the deeper reaching root system of the crop canola (Breuer et al., 2003) in and a consequently larger plant water uptake in comparison to other crops. Upward directed Q_{Anet}
- 345

values were observed during the year 2017 for the soils from Bad Lauchstädt under winter rye crop (Fig. 4a) and during 2015 for the soils from Sauerbach under winter wheat (Fig. 4b).



Figure 4: Comparison of net fluxes for the same soils at two sites: Annual observed net soil water flux at 1.5 m soil depth of soils from a)
 Bad Lauchstädt, b) Sauerbach, c) Dedelow, and d) Selhausen under a dry climate (Bad Lauchstädt) and wet climate (Selhausen) for the years between 2011-2017; average values (symbols) and standard deviations (error bars) for observations from the same soil.

3.3 Crop yield and Water Use Efficiency

The grain yields were in general larger for a dry climate at Bad Lauchstädt than for a wet climate at Selhausen except for the peas (Fig. 5a). The pea crop had in comparison to the other cereal crops a relatively short vegetation period and depends more on conditions during germination in early spring than on differences in climatic conditions in late spring and summer. For the other crops the spread of fungal pathogens under a more humid climate (Talley et al., 2002; Agam and Berliner, 2006) and frequent occurrence of dew formation (Xiao et al., 2009; Groh et al., 2018a; Brunke et al., 2019; Groh et al., 2019) could explain the generally lower yield of grain crops for soils under a wet climate in Selhausen. However, an appropriate 360 crop management with one to three applications of fungicides during the growing season (see appendix Table <u>A3</u>), except for pea crop in 2011 (BL and Se) and winter rye 2017 (Se) should have prevented the spread of fungal diseases and their impact on crop yield such that other reasons have to be considered. The yield varied for the most crops among the soil replicates at a certain site, which can be described by the coefficient of variation (CV), below a CV value of 28%, except for pea, which showed for all soils a high value, for winter canola grown on finer-textured soils in Se (BL, Se see appendix Table A4), and for winter barley (Dd and Sb in 2012, Sb in 2016) cropped at Se. For winter canola this might be related to a higher loss of rapeseeds during manual harvesting, natural pod shattering, cleaning and threshing (Alizadeh et al., 2007; Kuai et al., 2015). The CV value of the observed yield variability between each soil type corresponds to values reported between 5 to 27 % by Joernsgaard and Halmoe (2003) and Wallor et al. (2018). The yield of winter wheat (7.8 t ha⁻¹ see appendix Table A4) for the soil from BL at BL agreed well with observations on yields from a long term fertilization experiment at the BL site (Merbach and Schulz, 2013), which demonstrates the high yield potential of the soil from BL.



Figure 5: Comparison of annual crop yield- and ET-related parameters for the same soils from Bad Lauchstädt, Dedelow, Sauerbach, and Selhausen (three lysimeters each origin) at the two sites with relatively dry (Bad Lauchstädt) and wet climate (Selhausen); average values
(symbols) and standard deviation (error bars) between observations from the same soils for (a) observed yield, (b) total biomass, (c) harvest index, (d) evapotranspiration, (e) water use efficiency (WUE) from yield, (f) WUE from total biomass, and (h) nitrogen (*N*) content in the grain yield, and (g) the relationship between grain yield and evapotranspiration of all soils and crops during the years 2011-2017.

- 380 The scatterplot of the total biomass (Fig. 5b) shows that most crops produced relatively similar amounts of total above ground biomass at both sites with the exception of winter barley in years 2012 and 2016. The crops could probably use comparable amounts of solar radiation during the observation period (average annual radiation from 2011 to 2017, obtained from the weather stations; BL: 1181.4 kWh m⁻² and Se: 1180 kWh m⁻²). Despite a similar amount of radiation received by the crops the harvest index, which is defined as the ratio of yield to the total biomass, was found to be larger under a dry 385 climate than under a wet climate (Fig. 5c). This means that crops under a dry climate were more productive with respect to
- crop yield than under a wet climate. The crop ET (i.e., ET related to the vegetation period) was larger under the wet than under the dry climate (Fig. 5d), and the corresponding crop water use efficiency (WUE) was larger at the site with the

relatively dry (BL) as compared to the wet (Se) climate (Fig. 5e). These results demonstrated that plants were more efficient to produce yield at a site with a suboptimal water supply. The present results are in line with earlier findings from Zhang et

- 390 al. (2015), who showed that the WUE reached a maximum under warm and dry and a stable minimum under warm-wet climatic conditions. Also when WUE was calculated based on the total aboveground biomass, a higher WUE was observed for the corresponding crop under a dry than under a wet climate (Fig. 5f), which demonstrated that climatic conditions were not only beneficial for the grain yield but also for that of the straw. However, differences in fertilizer application (see appendix Table <u>A3</u>) with lower nitrate application in the wet site could be another reason for the differences in yield and
- 395 biomass production.

400

405

The lower WUE under a wet climate might be related to a higher soil evaporation and plant canopy interception evaporation. Kunrath et al. (2018) found for the crop tall fescue that limiting nitrogen-supply conditions negatively affected WUE values by a reduced leaf area index, leaf photosynthesis and radiation efficiency, which hence increased the ratio of soil evaporation to transpiration. Thus, we further compared the ET during periods when ET was either transpiration (ET_T) or evaporation (ET_E) dominated. The transpiration-dominated period was defined from the beginning of April, which corresponds well with the temporal increase of the monthly ET, until the time when plants reached the growth stage of ripening /maturity of their fruit or seeds about a month before harvest (see appendix Table <u>A3</u>). The rest of the vegetation period was defined as the evaporation-dominated period are shown in Table 1. The differences for ET_E between all soils in the dry and wet climate from 359 mm to 576 mm was larger than the differences for ET_T (range: -72 mm to 199 mm). Especially the

larger soil evaporation (ET_E) at Selhausen contributed to the lower WUE under wet climate. The relationship between yield and ET was reported to correspond with the productivity function of crops (grain yield vs. ET) and often assumed to be linear (Tolk and Howell, 2009; Wichelns, 2014). However, for our present data, a quadratic

productivity function (Fan et al., 2018) of the winter barley and pea crops (Fig. 5g) rather than a linear one could explain the

- 410 observed larger WUE of soils under a dry climate at Bad Lauchstädt. The crop winter canola could be best described by a linear productivity function with a negative slope (Fig. 5g). The other crops, winter rye and winter wheat, could neither be described by a linear nor a quadratic function. Longer time series' with more crop yield observations under different climatic conditions would be necessary to confirm the assumed quadratic productivity function for these crops.
- Grain yield quality in terms of the nitrogen content of the grains is an additional important variable to characterize the quality of legume and cereal crops (Kemanian et al., 2007). The scatterplot of the nitrogen content in the yield compares results from the same soils in the dry and wet climate (Fig. 5h). The comparison showed no effect of climatic conditions or of the fertilization on the crop grain quality. Larger deviations from the 1:1 line were only visible for the soils from Dedelow and the crop pea under a dry climate and for soils from Bad Lauchstädt and crop winter rye under a wet climate (Fig. 5h). Nuttall et al. (2017) remarked that heat stress during the time of flowering and higher temperatures during the post-anthesis
- 420 period of crops impact grain-size and milling yield. The impact of rising temperatures and increasing CO_2 concentrations in the atmosphere on yield quality could affect the nutritional quality and end-use value (Asseng et al., 2019). The grain yield

quality was reported to be influenced mainly by genetics, crop management, and environmental conditions (Nuttall et al., 2017). Since in the present study, the crop management was similar and the same cultivars were used, the altered climatic conditions seemed not to affect the quality of the yield in our crossed soil-climate experiment.

425

430

Table 1: Average values (of 3 lysimeters each) of cumulative evapotranspiration ($\sum ET$) for the whole observation period (2011-2017), and cumulative transpiration ($\sum ET_T$) and evaporation ($\sum ET_E$) for periods dominated by evaporation (E) or transpiration (T), for soils from Bad Lauchstädt (BL), Sauerbach (Sb), Dedelow (Dd), and Selhausen (Se) under a dry climate at BL and a wet climate at Se. The ET_T values were defined from the beginning of the vegetation period (April) until ripening/maturity of the fruit or seeds; the data for $\sum ET_E$ comprised the values from rest of the season. The differences of the cumulative values for the same soils between the sites BL and Se are denoted by $\Delta \sum ET_E$ and $\Delta \sum ET_T$.

Location		Se	BL	Se	BL	Se	BL	Se	BL	
Soil		BL	BL	Sb	Sb	Dd	Dd	Se	Se	
ΣET	(mm)	4090.1	3490.8	4121.0	3406.8	3593.9	3316.7	3985.0	3323.0	
$\sum ET_E$	(mm)	2102.5	1616.9	2110.3	1595.1	1941.7	1593.1	2228.2	1668.0	
$\sum ET_{T}$	(mm)	1987.5	1873.9	2010.7	1811.7	1652.1	1723.7	1756.8	1655.0	
ΔΣΕΤ	(mm)	599.3		714.2		277.1		661.9		
$\Delta \sum ET_E$	(mm)	485.7		515.2		348.7		560.2		
$\Delta \Sigma ET_{T}$	(mm)	113.6	113.6		199.0			101.8		

4 Conclusion

Lysimeter data from a German-wide lysimeter network (TERENO-SOILCan), where intact soil monoliths were moved to sites with contrasting climatic conditions, were used to analyse effects of soil and climate on agricultural ecosystems in a

- 435 soil-climate crossed factorial design. In the wet climate, there was a net drainage which was larger for the coarser- than for the finer-textured soils. In the dry climate, a small negative net drainage (upward flux) was obtained when observing the long-term average for the whole period 2011-2017. In the wet climate, drainage dominated for all soils. When looking at shorter periods, negative values of monthly net fluxes <u>were</u> observed during the summer months at both sites.
- During winter months, the soil water storage (SWS) returned to a typical value and drainage occurred when this value was reached. In the dry climate, this critical SWS was not reached in two soils after the growing season of 2015 in which the SWS was strongly depleted. The resulting insufficient refilling of the soil water storage capacity after a drought suggests that the precipitation during the following winter months was not sufficient to refill the soil so that no drainage took place. This lack of drainage had consequences for the upward water fluxes in the following growing seasons. Future studies about the impact of climate change, which in general are expected to increase the frequency and duration of droughts, on agroecosystem water balances and crop development should consider the long lasting impact of droughts on the soil water balance and soil water fluxes that are carried over to following years. Results indicate that direct observation on SWS will
 - 16

become increasingly important in environmental climate change studies, where changing climatic conditions could affect the SWSC. Longer term monitoring data are needed to observe effects of impacts on soil properties.

Crops were more productive in terms of grain yield and used less water under drier climatic conditions. Plant development

- 450 and a higher crop water use efficiency demonstrated that less plant available soil water did not go along with a decline of grain yield, because plants used the available soil water resources under such conditions more efficiently (e.g. by reduced soil evaporation). Results revealed in contrast to our hypothesis of a linear productivity function for some crops a quadratic productivity function and thus showed that plants can maximize their grain yield under an intermediate ET range in rainfed agriculture. However, longer time series are necessary to confirm the latter hypothesis of a quadratic productivity function of
- 455 the corresponding crop. Our results suggest that despite the higher grain yield (quantity) climatic conditions seemed not to affect the quality of the yield, which might reflect a positive effect of the regional drier climatic conditions for crop production. The results of this study so far confirmed that typical soil water balance components, crop water use and especially the soil water storage dynamics undergo a substantial change when exposed to different climatic conditions. We could show that:
- 460 1) The result further suggests that a new approach based on lysimeter mass data can enable the long-term monitoring of SWS changes at the pedon scale.
 - 2) SWS dynamics were vulnerable to droughts and led to an insufficient refilling of the soil water storage capacity.
 - 3) Crossed soil-climate experiments are useful to determine the impact of changing climatic conditions on the ecosystem water balances.
- 465 4) Crop water use efficiencies <u>increased with reduced water supply</u>.

The results herald the need to account for potential changes in soil water storage and plant reactions due to changes in climatic conditions and variability when trying to develop adaptation strategies in the agricultural sector.

Data availability

All data for the specific lysimeter and weather station (raw data) can be freely obtained from the TERENO data portal

470 (https://teodoor.icg.kfa-juelich.de/ddp/index.jsp, lysimeter station Bad Lauchstädt and Selhausen: SE_Y_03, SE_Y_04). Climate data for the experimental station Bad Lauchstädt can be acquired upon request from Ralf Gründling. The processed data to support the findings of this study can be acquired also from the TERENO data portal (https://hdl.handle.net/20.500.11952/butt.metadata.handle/000000010).

Author contribution

475 TP conceived the experiments. JG, JV, HHG, and TP had the idea and designed the study. JG and RG provided the data for the corresponding lysimeter stations. JG performed the data analysis and wrote the manuscript with equal contributions from all co-authors.

Acknowledgements

We acknowledge the support of TERENO and SOILCan, which were funded by the Helmholtz Association (HGF) and the
 Federal Ministry of Education and Research (BMBF). We thank the colleagues at the corresponding lysimeter station for
 their kind support: Sylvia Schmögner, Petra Petersohn and Ines Merbach (Bad Lauchstädt), Jörg Haase (Dedelow), Werner
 Küpper, Ferdinand Engels, Philipp Meulendick, Rainer Harms, and Leander Fürst (Selhausen). Thanks to Gernot Verch
 (Dedelow-ZALF) for helpful discussions and constructive comments.

Competing interests

485 The authors declare that they have no conflict of interest.

Appendix data:

Table A1: Basic soil texture information for soils from Bad Lauchstädt (BL), Sauerbach (Sb), Dedelow (Dd), and Selhausen (Se). Please note the soil monoliths from Dedelow have a varying soil depth to the clay illuviation horizon (Bt) and to the marly, illitic glacial till (C-horizon). The profile for Dedelow describes the soil profile from soil monolith Dd_1.

<u>Soil</u>	<u>Horizon</u>	Upper depth	Lower depth	Sand	<u>Silt</u>	<u>Clay</u>
		<u>m</u>	<u>m</u>	<u>%</u>	<u>%</u>	<u>%</u>
BL	<u>Ap2</u>	<u>0</u>	<u>0.23</u>	<u>5</u>	<u>73</u>	<u>22</u>
	<u>Axh1-2</u>	<u>0.23</u>	<u>0.33</u>	<u>4</u>	<u>83</u>	<u>13</u>
	<u>Axh3-2</u>	<u>0.33</u>	<u>0.70</u>	<u>4</u>	<u>82</u>	<u>14</u>
	<u>C1-1</u>	<u>0.70</u>	<u>1.10</u>	<u>6</u>	<u>67</u>	<u>27</u>
	<u>C2-2</u>	<u>1.10</u>	<u>1.45</u>	<u>5</u>	<u>72</u>	<u>23</u>
	<u>C3-1</u>	<u>1.45</u>	<u>>1.50</u>	<u>65</u>	<u>16</u>	<u>19</u>
<u>Sb</u>	<u>Ap</u>	<u>0</u>	<u>0.26</u>	<u>13</u>	<u>62</u>	<u>25</u>
	<u>M1</u>	<u>0.26</u>	<u>0.75</u>	<u>15</u>	<u>70</u>	<u>15</u>
	<u>M2</u>	<u>0.75</u>	<u>1.04</u>	<u>11</u>	<u>74</u>	<u>15</u>
	<u>M3</u>	<u>1.04</u>	<u>1.5</u>	<u>6</u>	<u>75</u>	<u>19</u>
Dd	<u>Ap</u>	0.00	<u>0.30</u>	<u>54</u>	<u>31</u>	<u>15</u>
	<u>Al+Bt</u>	<u>0.30</u>	<u>0.42</u>	<u>51</u>	<u>34</u>	<u>15</u>
	<u>Bt</u>	<u>0.42</u>	<u>0.90</u>	<u>52</u>	<u>23</u>	<u>25</u>
	elCcv	0.80	<u>1.50</u>	<u>59</u>	<u>29</u>	<u>12</u>
<u>Se</u>	<u>Ap</u>	<u>0</u>	0.23	<u>15</u>	<u>68</u>	<u>17</u>
	<u>Al-Bv</u>	<u>0.23</u>	<u>0.68</u>	<u>15</u>	<u>63</u>	<u>22</u>
	<u>II-Btv</u>	<u>0.68</u>	<u>>1.50</u>	<u>17</u>	<u>61</u>	<u>22</u>

490 Table <u>A2</u>: Average observed soil water flux at 1.5m soil depth of soils from Bad Lauchstädt (BL), Sauerbach (Sb), Dedelow (Dd), and Selhausen (Se) under a dry climate Bad Lauchstädt (BL) and wet climate Selhausen (Se).

Origin	Transfer	2011*	2012	2013	2014	2015	2016	2017	Average [#]	
		mm	mm	mm	mm	mm	mm	mm	mm	
BL-	·	-8	-91	-91	119	-16	-35	-53	-28	
Sb	BL	17	-18	-97	93	-38	-32	-12	-17	
Dd	BL	33	-78	-63	164	-34	-52	-44	-18	
Se	BL	17	-19	-99	96	-27	-41	-19	-18	
BL	Se	31	61	-57	206	7	94	-14	50	
Sb	Se	19	28	-28	132	-8	177	46	58	
Dd	Se	48	66	76	284	41	223	27	119	
Se-	·	28	30	-5	142	20	68	40	49	
* April-December; # 2012 – 2017										

Table <u>A3</u>: Site management information on seasonal crop type, sowing and harvesting date, crop growth length N-fertilizer and number of fungicide applications at Bad Lauchstädt and Selhausen. Calcium ammonium nitrate (KAS) was mainly us as *N*-fertilizer. Except for 2013 in Selhausen, where ammonium sulphate nitrate (ASS) instead of KAS was used.

Crop	Bad Lauc	hstädt				Selhausen						
	Sowing	Harvest	Duration	N-Fertilizer	Number of	Sowing	Harvest	Duration	N-Fertilizer	Number of		
	YY/mm	YY/mm	[days]	(KAS)	fungicide	YY/mm/	YY/mm/dd	[days]	(KAS)	fungicide		
	/dd	/dd		kg N ha ⁻¹	applications	dd			kg N ha⁻¹	applications		
Pea	11/	11/	99		0	11/06/01	11/08/25	85		0		
	05/04	08/11										
Winter	11/	12/	278	145	3	11/10/14	12/07/10	270	50	§		
Barley	09/30	07/12										
Winter	12/	13/	330	210	2	12/09/18	13/07/25	310	130#	1		
Canola	08/27	07/23										
Oat	14/	14/	82	60	0	14/03/05	14/06/03	90	60	0		
	03/13	06/03										
Winter	14/	15/	288	60	2	14/ 10/07	15/07/21	279	90	2		
Wheat	10/13	07/28										
Winter	15/ 09/	16/	282	100	3	15/ 10/07	16/07/08	275	80	2		
Barley	22	06/30										
Winter	16/	17/	285	100	2	16/10/11	17/07/21	283	78	0		
Rye	10/05	07/17										

ASS; § no data available

Test site		2	2011		2012		2013		2015	2015			2017	2017	
Origin Transfer	Pea		Winter barley		Winte	Winter canola		Winter wheat		Winter barley		Winter rye			
		t ha ⁻¹	CV %	t ha ⁻¹	CV %	t ha ⁻¹	CV %	t ha ⁻¹	t ha ⁻¹	CV %	t ha ⁻¹	CV %	t ha ⁻¹	CV %	
BL	BL	2.20	20	9.30	11	7.01	23	#	7.81	5	8.46	3	7.35	18	
		(±0.44)		(±1.01)		(±1.60)			(±0.42)		(±0.23)		(±1.35)		
Dd	BL	1.11	43	8.79	8	5.64	9	#	5.90	4	9.42	9	7.74	13	
		(±0.48)		(±0.71)		(±0.50)			(±0.26)		(±0.81)		(±0.97)		
Sb	BL	2.44	38	10.28	24	4.65	13	#	8.33	4	11.36	4	6.83	13	
		(±0.92)		(±2.47)		(±0.61)			(±0.37)		(±0.46)		(±0.92)		
Se	BL	0.99	17	7.76	19	5.07	16	#	6.36	6	8.42	8	5.45	10	
		(±0.17)		(±1.50)		(±0.82)			(±0.36)		(±0.66)		(±0.53)		
BL	Se	1.87	24	8.43	15	1.32	62	#	5.66	16	5.14	1	4.77	9	
		(±0.45)		(±1.29)		(±0.80)			(±0.88)		(±0.05)		(±0.44)		
Dd	Se	1.13	57	4.02	57	2.21	17	#	4.73	18	3.51	1	4.77	9	
		(±0.64)		(±2.31)		(±0.38)			(±0.83)		(±2.31)		(±0.44)		
Sb	Se	1.90	12	5.06	70	3.66	28	#	5.67	2	5.01	40	6.37	9	
		(±0.22)		(±3.56)		(±1.03)			(±0.14)		(±2.02)		(±0.57)		
Se-	•	2.10	25	7.93	8	3.94	77	#	4.61	15	3.49	15	4.22	23	
		(±0.53)		(±0.61)		(±3.05)			(±0.71)		(±0.51)		(±0.97)		

Table <u>A4</u>: Average and coefficient of variation (CV) of yield for soil from Bad Lauchstädt (BL), Dedelow (Dd), Sauerbach (Sb), Selhausen (Se) under dry (BL)500and wet climate (Se). Origin describes the location, where the soil was taken from and the location where the soil was transferred to, where the soil was
transferred to. The value in the brackets describes the variability of yield for each soil type (standard deviation from three replicates).

Crop was not harvested but biomass was cut and removed in June. Manually tilled so that soil was bare fallow during summer

References

10.1016/S0304-3800(03)00274-6, 2003.

- 505 Adhikari, K., and Hartemink, A. E.: Linking soils to ecosystem services A global review, Geoderma, 262, 101-111, 10.1016/j.geoderma.2015.08.009, 2016.
 Agam, N., and Berliner, P. R.: Dew formation and water vapor adsorption in semi-arid environments A review, Journal of Arid Environments, 65, 572-590, 10.1016/j.jaridenv.2005.09.004, 2006.
 Albert, M., Nagel, R.-V., Nuske, R. S., Sutmöller, J., and Spellmann, H.: Tree Species Selection in the Face of Drought Risk—Uncertainty
- 510 in Forest Planning, Forests, 8, 363, 10.3390/f8100363, 2017.
 Alizadeh, M. R., Bagheri, I., and Payman, M. H.: Evaluation of a Rice Reaper Used for Rapeseed Harvesting, American-Eurasian J. Agric. & Environ. Sci., 2, 388-394, 2007.
 Allen, R. G., Pruitt, W. O., Wright, J. L., Howell, T. A., Ventura, F., Snyder, R., Itenfisu, D., Steduto, P., Berengena, J., Yrisarry, J. B., Smith, M., Pereira, L. S., Raes, D., Perrier, A., Alves, I., Walter, I., and Elliott, R.: A recommendation on standardized surface resistance
- 515 for hourly calculation of reference ETo by the FAO56 Penman-Monteith method, Agricultural Water Management, 81, 1-22, 10.1016/j.agwat.2005.03.007, 2006.
- Asseng, S., Ewert, F., Martre, P., Rötter, R. P., Lobell, D. B., Cammarano, D., Kimball, B. A., Ottman, M. J., Wall, G. W., White, J. W., Reynolds, M. P., Alderman, P. D., Prasad, P. V. V., Aggarwal, P. K., Anothai, J., Basso, B., Biernath, C., Challinor, A. J., De Sanctis, G., Doltra, J., Fereres, E., Garcia-Vila, M., Gayler, S., Hoogenboom, G., Hunt, L. A., Izaurralde, R. C., Jabloun, M., Jones, C. D., Kersebaum,
- 520 K. C., Koehler, A. K., Müller, C., Naresh Kumar, S., Nendel, C., O'Leary, G., Olesen, J. E., Palosuo, T., Priesack, E., Eyshi Rezaei, E., Ruane, A. C., Semenov, M. A., Shcherbak, I., Stöckle, C., Stratonovitch, P., Streck, T., Supit, I., Tao, F., Thorburn, P. J., Waha, K., Wang, E., Wallach, D., Wolf, J., Zhao, Z., and Zhu, Y.: Rising temperatures reduce global wheat production, Nature Climate Change, 5, 143, 10.1038/nclimate2470, 2014.
- Asseng, S., Martre, P., Maiorano, A., Rötter, R. P., O'Leary, G. J., Fitzgerald, G. J., Girousse, C., Motzo, R., Giunta, F., Babar, M. A.,
 Reynolds, M. P., Kheir, A. M. S., Thorburn, P. J., Waha, K., Ruane, A. C., Aggarwal, P. K., Ahmed, M., Balkovič, J., Basso, B., Biernath, C., Bindi, M., Cammarano, D., Challinor, A. J., De Sanctis, G., Dumont, B., Eyshi Rezaei, E., Fereres, E., Ferrise, R., Garcia-Vila, M.,
 Gayler, S., Gao, Y., Horan, H., Hoogenboom, G., Izaurralde, R. C., Jabloun, M., Jones, C. D., Kassie, B. T., Kersebaum, K.-C., Klein, C.,
 Koehler, A.-K., Liu, B., Minoli, S., Montesino San Martin, M., Müller, C., Naresh Kumar, S., Nendel, C., Olesen, J. E., Palosuo, T.,
 Porter, J. R., Priesack, E., Ripoche, D., Semenov, M. A., Stöckle, C., Stratonovitch, P., Streck, T., Supit, I., Tao, F., Van der Velde, M.,
- Wallach, D., Wang, E., Webber, H., Wolf, J., Xiao, L., Zhang, Z., Zhao, Z., Zhu, Y., and Ewert, F.: Climate change impact and adaptation for wheat protein, Global Change Biology, 25, 155-173, 10.1111/gcb.14481, 2019.
 Blois, J. L., Williams, J. W., Fitzpatrick, M. C., Jackson, S. T., and Ferrier, S.: Space can substitute for time in predicting climate-change effects on biodiversity, Proceedings of the National Academy of Sciences, 110, 9374-9379, 10.1073/pnas.1220228110, 2013.
 Bogena, H. R., Montzka, C., Huisman, J. A., Graf, A., Schmidt, M., Stockinger, M., von Hebel, C., Hendricks-Franssen, H. J., van der
- 535 Kruk, J., Tappe, W., Lücke, A., Baatz, R., Bol, R., Groh, J., Pütz, T., Jakobi, J., Kunkel, R., Sorg, J., and Vereecken, H.: The TERENO-Rur Hydrological Observatory: A Multiscale Multi-Compartment Research Platform for the Advancement of Hydrological Science, Vadose Zone Journal, 17, 10.2136/vzj2018.03.0055, 2018. Breuer, L., Eckhardt, K., and Frede, H.-G.: Plant parameter values for models in temperate climates, Ecological Modelling, 169, 237-293,
- 540 Brunke, M., Groh, J., Vanderborght, J., Vereecken, H., and Pütz, T.: Tau ein bedeutsamer Bestandteil des Wasserhaushaltes eines Grünlandstandortes Eine lysimeterbasierte Studie (In German, with English abstract.) 18. Gumpensteiner Lysimetertagung 2019, 49 58, 2019.

Cao, R., Jia, X., Huang, L., Zhu, Y., Wu, L., and Shao, M. a.: Deep soil water storage varies with vegetation type and rainfall amount in the Loess Plateau of China, Scientific Reports, 8, 12346, 10.1038/s41598-018-30850-7, 2018.

- 545 Ellerbrock, R. H., Gerke, H. H., Bachmann, J., and Goebel, M.-O.: Composition of Organic Matter Fractions for Explaining Wettability of Three Forest Soils, Soil Science Society of America Journal, 69, 57-66, 10.2136/sssaj2005.0057, 2005.
 Ewert, F., Rodriguez, D., Jamieson, P., Semenov, M. A., Mitchell, R. A. C., Goudriaan, J., Porter, J. R., Kimball, B. A., Pinter, P. J., Manderscheid, R., Weigel, H. J., Fangmeier, A., Fereres, E., and Villalobos, F.: Effects of elevated CO2 and drought on wheat: testing crop simulation models for different experimental and climatic conditions, Agriculture, Ecosystems & Environment, 93, 249-266, 10.1016/S0167-8809(01)00352-8, 2002.
 - Fan, Y., Wang, C., and Nan, Z.: Determining water use efficiency of wheat and cotton: A meta-regression analysis, Agricultural Water Management, 199, 48-60, 10.1016/j.agwat.2017.12.006, 2018.

Frieler, K., Schauberger, B., Arneth, A., Balkovič, J., Chryssanthacopoulos, J., Deryng, D., Elliott, J., Folberth, C., Khabarov, N., Müller, C., Olin, S., Pugh, T. A. M., Schaphoff, S., Schewe, J., Schmid, E., Warszawski, L., and Levermann, A.: Understanding the weather signal in national crop-vield variability. Earth's Future. 5, 605-616, 10.1002/2016EF000525, 2017.

- in national crop-yield variability, Earth's Future, 5, 605-616, 10.1002/2016EF000525, 2017.
 Gammans, M., Mérel, P., and Ortiz-Bobea, A.: Negative impacts of climate change on cereal yields: statistical evidence from France, Environmental Research Letters, 12, 054007, 10.1088/1748-9326/aa6b0c, 2017.
 Gourdji, S. M., Sibley, A. M., and Lobell, D. B.: Global crop exposure to critical high temperatures in the reproductive period: historical trends and future projections, Environmental Research Letters, 8, 024041, 10.1088/1748-9326/8/2/024041, 2013.
- 560 Groh, J., Vanderborght, J., Pütz, T., and Vereecken, H.: How to Control the Lysimeter Bottom Boundary to Investigate the Effect of Climate Change on Soil Processes?, Vadose Zone Journal, 15, 10.2136/vzj2015.08.0113, 2016. Groh, J., Slawitsch, V., Herndl, M., Graf, A., Vereecken, H., and Pütz, T.: Determining dew and hoar frost formation for a low mountain range and alpine grassland site by weighable lysimeter, Journal of Hydrology, 563, 372-381, 10.1016/j.jhydrol.2018.06.009, 2018a. Groh, J., Stumpp, C., Lücke, A., Pütz, T., Vanderborght, J., and Vereecken, H.: Inverse Estimation of Soil Hydraulic and Transport
- 565 Parameters of Layered Soils from Water Stable Isotope and Lysimeter Data, Vadose Zone Journal, 17, 10.2136/vzj2017.09.0168, 2018b. Groh, J., Pütz, T., Gerke, H. H., Vanderborght, J., and Vereecken, H.: Quantification and prediction of nighttime evapotranspiration for two distinct grassland ecosystems, Water Resources Research, 10.1029/2018wr024072, 2019. Gudmundsson, L., and Seneviratne, S. I.: Anthropogenic climate change affects meteorological drought risk in Europe, Environmental Research Letters, 11, 044005, 10.1088/1748-9326/11/4/044005, 2016.
- 570 Gudmundsson, L., Seneviratne, S. I., and Zhang, X.: Anthropogenic climate change detected in European renewable freshwater resources, Nature Climate Change, 7, 813, 10.1038/nclimate3416, 2017.
 Hatfield, J. L., and Dold, C.: Water-Use Efficiency: Advances and Challenges in a Changing Climate, Frontiers in Plant Science, 10, 10.3389/fpls.2019.00103, 2019.
- Herbrich, M., and Gerke, H. H.: Scales of Water Retention Dynamics Observed in Eroded Luvisols from an Arable Postglacial Soil 575 Landscape, Vadose Zone Journal, 10.2136/vzj2017.01.0003, 2017.

Herbrich, M., Gerke, H. H., Bens, O., and Sommer, M.: Water balance and leaching of dissolved organic and inorganic carbon of eroded Luvisols using high precision weighing lysimeters, Soil and Tillage Research, 165, 144-160, 10.1016/j.still.2016.08.003, 2017.
Hirmas, D. R., Giménez, D., Nemes, A., Kerry, R., Brunsell, N. A., and Wilson, C. J.: Climate-induced changes in continental-scale soil macroporosity may intensify water cycle, Nature, 561, 100-103, 10.1038/s41586-018-0463-x, 2018.

580 Ionita, M., Tallaksen, L. M., Kingston, D. G., Stagge, J. H., Laaha, G., Van Lanen, H. A. J., Chelcea, S. M., and Haslinger, K.: The European 2015 drought from a climatological perspective, Hydrol. Earth Syst. Sci. Discuss., 2016, 1-32, 10.5194/hess-2016-218, 2016. Joernsgaard, B., and Halmoe, S.: Intra-field yield variation over crops and years, European Journal of Agronomy, 19, 23-33, 10.1016/S1161-0301(02)00016-3, 2003.

Katerji, N., Mastrorilli, M., and Rana, G.: Water use efficiency of crops cultivated in the Mediterranean region: Review and analysis, 585 European Journal of Agronomy, 28, 493-507, 10.1016/j.eja.2007.12.003, 2008.

Kemanian, A. R., Stöckle, C. O., and Huggins, D. R.: Estimating Grain and Straw Nitrogen Concentration in Grain Crops Based on Aboveground Nitrogen Concentration and Harvest Index, Agronomy Journal, 99, 158-165, 10.2134/agronj2006.0090, 2007.

Kuai, J., Sun, Y., Zuo, Q., Huang, H., Liao, Q., Wu, C., Lu, J., Wu, J., and Zhou, G.: The yield of mechanically harvested rapeseed (Brassica napus L.) can be increased by optimum plant density and row spacing, Scientific Reports, 5, 18835, 10.1038/srep18835, 2015.

- 590 Kunrath, T. R., Lemaire, G., Sadras, V. O., and Gastal, F.: Water use efficiency in perennial forage species: Interactions between nitrogen nutrition and water deficit, Field Crops Research, 222, 1-11, 10.1016/j.fcr.2018.02.031, 2018. Laaha, G., Gauster, T., Tallaksen, L. M., Vidal, J. P., Stahl, K., Prudhomme, C., Heudorfer, B., Vlnas, R., Ionita, M., Van Lanen, H. A. J., Adler, M. J., Caillouet, L., Delus, C., Fendekova, M., Gailliez, S., Hannaford, J., Kingston, D., Van Loon, A. F., Mediero, L., Osuch, M.,
- Romanowicz, R., Sauquet, E., Stagge, J. H., and Wong, W. K.: The European 2015 drought from a hydrological perspective, Hydrol. Earth
 Syst. Sci. Discuss., 2016, 1-30, 10.5194/hess-2016-366, 2016.

Lobell, D. B., and Gourdji, S. M.: The Influence of Climate Change on Global Crop Productivity, Plant Physiology, 160, 1686-1697, 10.1104/pp.112.208298, 2012.

Mei, X., Ma, L., Zhu, Q., Li, B., Zhang, D., Liu, H., Zhang, Q., Gou, Q., and Shen, M.: The variability in soil water storage on the loess hillslopes in China and its estimation, CATENA, 172, 807-818, 10.1016/j.catena.2018.09.045, 2019.

600 Merbach, I., and Schulz, E.: Long-term fertilization effects on crop yields, soil fertility and sustainability in the Static Fertilization Experiment Bad Lauchstädt under climatic conditions 2001–2010, Archives of Agronomy and Soil Science, 59, 1041-1057, 10.1080/03650340.2012.702895, 2013.

Niu, S., Xing, X., Zhang, Z. H. E., Xia, J., Zhou, X., Song, B., Li, L., and Wan, S.: Water-use efficiency in response to climate change: from leaf to ecosystem in a temperate steppe, Global Change Biology, 17, 1073-1082, 10.1111/j.1365-2486.2010.02280.x, 2011.

605 Nuttall, J. G., O'Leary, G. J., Panozzo, J. F., Walker, C. K., Barlow, K. M., and Fitzgerald, G. J.: Models of grain quality in wheat—A review, Field Crops Research, 202, 136-145, 10.1016/j.fcr.2015.12.011, 2017. Pickett, S. T. A.: Space-for-Time Substitution as an Alternative to Long-Term Studies, in: Long-Term Studies in Ecology: Approaches and Alternatives, edited by: Likens, G. E., Springer New York, New York, NY, 110-135, 1989. Ponce-Campos, G. E., Moran, M. S., Huete, A., Zhang, Y., Bresloff, C., Huxman, T. E., Eamus, D., Bosch, D. D., Buda, A. R., Gunter, S.

A., Scalley, T. H., Kitchen, S. G., McClaran, M. P., McNab, W. H., Montoya, D. S., Morgan, J. A., Peters, D. P. C., Sadler, E. J., Seyfried, M. S., and Starks, P. J.: Ecosystem resilience despite large-scale altered hydroclimatic conditions, Nature, 494, 349, 10.1038/nature11836, 2013.

Pütz, T., Kiese, R., Wollschläger, U., Groh, J., Rupp, H., Zacharias, S., Priesack, E., Gerke, H. H., Gasche, R., Bens, O., Borg, E., Baessler, C., Kaiser, K., Herbrich, M., Munch, J.-C., Sommer, M., Vogel, H.-J., Vanderborght, J., and Vereecken, H.: TERENO SOILCan: a lysimeter-network in Germany observing soil processes and plant diversity influenced by climate change, Environmental

- Earth Sciences, 75, 1-14, 10.1007/s12665-016-6031-5, 2016.
 Rieckh, H., Gerke, H. H., and Sommer, M.: Hydraulic properties of characteristic horizons depending on relief position and structure in a hummocky glacial soil landscape, Soil and Tillage Research, 125, 123-131, 10.1016/j.still.2012.07.004, 2012.
 Robinson, D. A., Jones, S. B., Lebron, I., Reinsch, S., Domínguez, M. T., Smith, A. R., Jones, D. L., Marshall, M. R., and Emmett, B. A.:
- Experimental evidence for drought induced alternative stable states of soil moisture, Scientific Reports, 6, 20018, 10.1038/srep20018, 2016.

Rogers, B. M., Jantz, P., and Goetz, S. J.: Vulnerability of eastern US tree species to climate change, Global Change Biology, 23, 3302-3320, 10.1111/gcb.13585, 2017.

Scanlon, T. M., Caylor, K. K., Manfreda, S., Levin, S. A., and Rodriguez-Iturbe, I.: Dynamic response of grass cover to rainfall variability: implications for the function and persistence of savanna ecosystems, Advances in Water Resources, 28, 291-302, , 2005.

Schauberger, B., Archontoulis, S., Arneth, A., Balkovic, J., Ciais, P., Deryng, D., Elliott, J., Folberth, C., Khabarov, N., Müller, C., Pugh, T. A. M., Rolinski, S., Schaphoff, S., Schmid, E., Wang, X., Schlenker, W., and Frieler, K.: Consistent negative response of US crops to high temperatures in observations and crop models, Nature Communications, 8, 13931, 10.1038/ncomms13931, 2017.

Scheelbeek, P. F. D., Bird, F. A., Tuomisto, H. L., Green, R., Harris, F. B., Joy, E. J. M., Chalabi, Z., Allen, E., Haines, A., and Dangour,
A. D.: Effect of environmental changes on vegetable and legume yields and nutritional quality, Proceedings of the National Academy of Sciences, 115, 6804-6809, 10.1073/pnas.1800442115, 2018.
Sebastiá, M.-T.: Plant guilds drive biomass response to global warming and water availability in subalpine grassland, Journal of Applied Ecology, 44, 158-167, 10.1111/j.1365-2664.2006.01232.x, 2007.

Sommer, M., Gerke, H. H., and Deumlich, D.: Modelling soil landscape genesis — A "time split" approach for hummocky agricultural landscapes, Geoderma, 145, 480-493, 10.1016/j.geoderma.2008.01.012, 2008.

Talley, S. M., Coley, P. D., and Kursar, T. A.: The effects of weather on fungal abundance and richness among 25 communities in the Intermountain West, BMC Ecol, 2, 7-7, 10.1186/1472-6785-2-7, 2002.

te Brake, B., van der Ploeg, M. J., and de Rooij, G. H.: Water storage change estimation from in situ shrinkage measurements of clay soils, Hydrol. Earth Syst. Sci., 17, 1933-1949, 10.5194/hess-17-1933-2013, 2013.

- 640 Tolk, J. A., and Howell, T. A.: Transpiration and Yield Relationships of Grain Sorghum Grown in a Field Environment 1 The mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture, Agronomy Journal, 101, 657-662, 10.2134/agronj2008.0079x, 2009.
- Troch, P. A., Carrillo, G., Sivapalan, M., Wagener, T., and Sawicz, K.: Climate-vegetation-soil interactions and long-term hydrologic partitioning: signatures of catchment co-evolution, Hydrol. Earth Syst. Sci., 17, 2209-2217, 10.5194/hess-17-2209-2013, 2013.
- Vereecken, H., Weynants, M., Javaux, M., Pachepsky, Y., Schaap, M. G., and Genuchten, M. T. v.: Using Pedotransfer Functions to Estimate the van Genuchten–Mualem Soil Hydraulic Properties: A Review All rights reserved. No part of this periodical may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information storage and retrieval system, without permission in writing from the publisher, Vadose Zone Journal, 9, 795-820, 10.2136/vzj2010.0045, 2010.
- Vereecken, H., Schnepf, A., Hopmans, J. W., Javaux, M., Or, D., Roose, T., Vanderborght, J., Young, M. H., Amelung, W., Aitkenhead, M., Allison, S. D., Assouline, S., Baveye, P., Berli, M., Brüggemann, N., Finke, P., Flury, M., Gaiser, T., Govers, G., Ghezzehei, T., Hallett, P., Hendricks Franssen, H. J., Heppell, J., Horn, R., Huisman, J. A., Jacques, D., Jonard, F., Kollet, S., Lafolie, F., Lamorski, K., Leitner, D., McBratney, A., Minasny, B., Montzka, C., Nowak, W., Pachepsky, Y., Padarian, J., Romano, N., Roth, K., Rothfuss, Y.,
- 655 Rowe, E. C., Schwen, A., Šimůnek, J., Tiktak, A., Van Dam, J., van der Zee, S. E. A. T. M., Vogel, H. J., Vrugt, J. A., Wöhling, T., and Young, I. M.: Modeling Soil Processes: Review, Key Challenges, and New Perspectives, Vadose Zone Journal, 15, 10.2136/vzj2015.09.0131, 2016.
 Wallor, E., Kersebaum, K.-C., Ventrella, D., Bindi, M., Cammarano, D., Coucheney, E., Gaiser, T., Garofalo, P., Giglio, L., Giola, P.,

Wallor, E., Kersebaum, K.-C., Ventrella, D., Bindi, M., Cammarano, D., Coucheney, E., Gaiser, T., Garolalo, P., Giglio, L., Giola, P., Hoffmann, M. P., Iocola, I., Lana, M., Lewan, E., Maharjan, G. R., Moriondo, M., Mula, L., Nendel, C., Pohankova, E., Roggero, P. P., Traka, M., and Trambi, G.: The reapones of process based agra accounter models to within field variability in site conditions. Field

Trnka, M., and Trombi, G.: The response of process-based agro-ecosystem models to within-field variability in site conditions, Field Crops Research, 228, 1-19, 10.1016/j.fcr.2018.08.021, 2018.
Walter, H., and Lieth, H. H. F.: Klimadiagramm-Weltatlas, Fischer Verlag, Jena, 1967.
Wichelns, D.: Do Estimates of Water Productivity Enhance Understanding of Farm-Level Water Management?, Water, 6, 778-795, 10.3390/w6040778 2014.

- Wogan, G. O. U., and Wang, I. J.: The value of space-for-time substitution for studying fine-scale microevolutionary processes, Ecography, 41, 1456-1468, 10.1111/ecog.03235, 2018.
 Wollschläger, U., Attinger, S., Borchardt, D., Brauns, M., Cuntz, M., Dietrich, P., Fleckenstein, J. H., Friese, K., Friesen, J., Harpke, A., Hildebrandt, A., Jäckel, G., Kamjunke, N., Knöller, K., Kögler, S., Kolditz, O., Krieg, R., Kumar, R., Lausch, A., Liess, M., Marx, A., Merz, R., Mueller, C., Musolff, A., Norf, H., Oswald, S. E., Rebmann, C., Reinstorf, F., Rode, M., Rink, K., Rinke, K., Samaniego, L.,
- Vieweg, M., Vogel, H.-J., Weitere, M., Werban, U., Zink, M., and Zacharias, S.: The Bode hydrological observatory: a platform for integrated, interdisciplinary hydro-ecological research within the TERENO Harz/Central German Lowland Observatory, Environmental Earth Sciences, 76, 29, 10.1007/s12665-016-6327-5, 2016.

Wu, W.-Y., Lan, C.-W., Lo, M.-H., Reager, J. T., and Famiglietti, J. S.: Increases in the annual range of soil water storage at northern middle and high latitudes under global warming, Geophysical Research Letters, 42, 3903-3910, 10.1002/2015GL064110, 2015.

- 675 Xiao, G., Zheng, F., Qiu, Z., and Yao, Y.: Impact of climate change on water use efficiency by wheat, potato and corn in semiarid areas of China, Agriculture, Ecosystems & Environment, 181, 108-114, 10.1016/j.agee.2013.09.019, 2013. Xiao, H., Meissner, R., Seeger, J., Rupp, H., and Borg, H.: Effect of vegetation type and growth stage on dewfall, determined with high precision weighing lysimeters at a site in northern Germany, Journal of Hydrology, 377, 43-49, 10.1016/j.jhydrol.2009.08.006, 2009. Yang, Y., Liu, D. L., Anwar, M. R., O'Leary, G., Macadam, I., and Yang, Y.: Water use efficiency and crop water balance of rainfed
- wheat in a semi-arid environment: sensitivity of future changes to projected climate changes and soil type, Theoretical and Applied Climatology, 123, 565-579, 10.1007/s00704-015-1376-3, 2016.
 Viet J, L. Huang, L. and Hartamink, A. E. Spatial temporal analysis of soil water storage and deep drainage under irrigated potatoes in
- Yost, J. L., Huang, J., and Hartemink, A. E.: Spatial-temporal analysis of soil water storage and deep drainage under irrigated potatoes in the Central Sands of Wisconsin, USA, Agricultural Water Management, 217, 226-235, 10.1016/j.agwat.2019.02.045, 2019.
- Yuan, Z. Y., Jiao, F., Shi, X. R., Sardans, J., Maestre, F. T., Delgado-Baquerizo, M., Reich, P. B., and Peñuelas, J.: Experimental and observational studies find contrasting responses of soil nutrients to climate change, eLife, 6, e23255, 10.7554/eLife.23255, 2017.
- Zhang, B., Liu, W., Chang, S. X., and Anyia, A. O.: Water-deficit and high temperature affected water use efficiency and arabinoxylan concentration in spring wheat, Journal of Cereal Science, 52, 263-269, 10.1016/j.jcs.2010.05.014, 2010.
- Zhang, J., Ren, W., An, P., Pan, Z., Wang, L., Dong, Z., He, D., Yang, J., Pan, S., and Tian, H.: Responses of Crop Water Use Efficiency to Climate Change and Agronomic Measures in the Semiarid Area of Northern China, PLOS ONE, 10, e0137409, 10.1371/journal.pone.0137409, 2015.
- Zhao, C., Piao, S., Huang, Y., Wang, X., Ciais, P., Huang, M., Zeng, Z., and Peng, S.: Field warming experiments shed light on the wheat yield response to temperature in China, Nature Communications, 7, 13530, 10.1038/ncomms13530, 2016. Zhao, C., Liu, B., Piao, S., Wang, X., Lobell, D. B., Huang, Y., Huang, M., Yao, Y., Bassu, S., Ciais, P., Durand, J.-L., Elliott, J., Ewert,

F., Janssens, I. A., Li, T., Lin, E., Liu, Q., Martre, P., Müller, C., Peng, S., Peñuelas, J., Ruane, A. C., Wallach, D., Wang, T., Wu, D., Liu,

- 695 Z., Zhu, Y., Zhu, Z., and Asseng, S.: Temperature increase reduces global yields of major crops in four independent estimates, Proceedings of the National Academy of Sciences, 114, 9326-9331, 10.1073/pnas.1701762114, 2017. Zhou, S., Yu, B., Schwalm, C. R., Ciais, P., Zhang, Y., Fisher, J. B., Michalak, A. M., Wang, W., Poulter, B., Huntzinger, D. N., Niu, S., Mao, J., Jain, A., Ricciuto, D. M., Shi, X., Ito, A., Wei, Y., Huang, Y., and Wang, G.: Response of Water Use Efficiency to Global Environmental Change Based on Output From Terrestrial Biosphere Models, Global Biogeochemical Cycles, 31, 1639-1655,
- 700 10.1002/2017gb005733, 2017.

Dear Editor Markus Weiler,

Thank you for your response on our replies. We added the proposed changes and discussed the points in detail which were suggested by the two reviewers in the revised manuscript. We also include briefly in the revised manuscript that the measurements errors in determining the water fluxes were in general much smaller than the effect of moving the soils in the revised manuscript. We hope that revision of our manuscript could answer the open questions from the replies and the Editor.

Kind regards

Jannis Groh