

Response to reviews on the manuscript hess-2019-432 “Ability of an Australian reanalysis dataset to characterise sub-daily precipitation” by Suwash Chandra Acharya et al.

We would like to thank the two anonymous referees for their constructive comments and suggestions on our paper.

In the following sections we provide a detailed response to all the remarks and suggestions made by the referees to improve the manuscript. The reviewers’ comments (in black), our corresponding reply (in blue), and revisions (*italicised*).

Response to Referee #1

General comments

The manuscript “Ability of an Australian reanalysis dataset to characterize sub-daily precipitation” by Acharya et al. evaluates sub-daily precipitation from a gridded reanalysis dataset, BARRA, covering Australia, against gauge observations and radar data. The aim is to assess the performance of BARRA, specifically for the use in catchment hydrology applications. The aim is clearly stated and the analysis is indeed relevant, as the need for continuous precipitation data of high spatial and temporal resolution is obvious. The manuscript is well structured and has a clear language.

We thank Referee #1 for acknowledging the relevance of the analyses presented and overall positive remarks on the manuscript.

On a general term I have some minor suggestions to improve the manuscript as listed below:

For non-Australians it would be useful to have more information about the climate and topography in Australia, specifically the rainfall climate in the selected areas. Also, please discuss results in light of regional differences in rainfall climate. I think there is room for one more analysis and figure.

We agree that including a description of Australian climate and topography will help non-Australian readers to generalise the conclusions of the paper for other studies. In Figure 1, we present the climate classification across Australia, based on the Köppen-Geiger classification which is not adequately discussed within the text. To address this comment, we have revised the study area and data sources section (Lines: 95-102) as follows:

*The pluviometer observations in this study encompass three of the broad climatic zones across Australia, namely tropical, temperate and arid (**Error! Reference source not found.**). Rainfall in Australia is highly variable across space and time. Rainfall is concentrated during summer in the tropical north, whereas rainfall is more prevalent during winter in the temperate south and southwest. The southeast region experiences a more consistent rainfall throughout the year. The central arid region receives the least total rainfall, and the eastern coast the highest average rainfall. The spatial evaluation across the entire Australia is impeded by the availability of high-resolution observation-based dataset. Therefore, the spatial evaluation is undertaken by using radar-based datasets at four*

city-centred regions (Brisbane, Darwin, Melbourne and Sydney). These study areas are located in the tropical and temperate zones with varying rainfall climates.

In the manuscript subsection “5.3 Performance dependence on spatial location”, we discuss the results focusing on the location of selected areas. This discussion implicitly considers the differences in climate of the selected areas. However, after describing the climate classification and rainfall climate of the selected areas, we have revised the discussion section (Lines: 329-332) as follows:.

*Both the bias analysis of quantiles and the FSS evaluation show that the performance of BARRA gradually improves as we move from northern to southern regions (**Error! Reference source not found.** and **Error! Reference source not found.**). This variation in performance can partly be attributed to the different rainfall climate at these regions. The convective precipitation during summer season is dominant at northern (low) latitudes, whereas winter (or uniform) non-convective rainfall are dominant at southern (high) latitudes.*

Since the target application is catchment hydrology, I would like to see more emphasize on the evaluation of areal precipitation. For instance, a case study analysing the evolution of high-impact events over catchments would be interesting.

We also recognise the importance of areal precipitation in the catchment hydrological applications. The evaluation and use of areal rainfalls require a substantial assessment in itself, and indeed we are in the final stages of preparing a paper that explores the evaluation and design applications of areal rainfall estimates. We have revised the conclusion section (Lines: 384, 388) as follows:.

The ability of BARRA to provide areal rainfall estimates is useful in catchment hydrological applications.

...

Accordingly, future work will be directed towards evaluation and design applications of areal rainfall estimates.

It might be beneficial to the reader to have other titles on sub-chapters, particularly in the Results section. For instance, “4.3 Fractions Skill Score (FSS)” could be renamed to describe what FSS actually evaluates.

We have re-named the sub-chapters to be more descriptive. The sub-chapters in the Methodology and corresponding Results section are revised as follows:

3.1 “Frequency distribution of rainfall” to “Frequency distribution of sub-daily rainfall”

3.2 “Spatio-temporal distribution” to “Neighbourhood-based diurnal patterns at point locations”

3.3 “Fractions skill score (FSS)” to “Neighbourhood-based spatial evaluation”

Specific comments

Although you state that a spatial resolution of 12 km is considered high, I would argue that the parameterization of convection is a major limitation when studying sub-daily rainfall. Please discuss this in more detail.

Our statement that the spatial resolution of BARRA is “high” is based on its comparison to other available reanalysis datasets for Australia. However, we agree that the spatial resolution is not fine enough for parameterisation of convection.

The Unified Model’s parameterised sub-grid convection scheme (the NWP model used in BARRA) which works independently at each grid point, produces a bias towards widespread precipitation (Clark et al., 2016; Su et al., 2019). The parameterisation scheme (for detail, see: Su et al., 2019) adopted for sub-grid convection is limiting in resolving convective rainfall and affects the locations dominated by such rainfall (especially tropics). This is observed in daily evaluation of BARRA by (Acharya et al., 2019) where the performance was better in temperate than tropical regions. We have revised the study area and data sources section (Lines: 137-138), and discussion section (Lines: 329-332) as follows:

This parameterisation scheme adopted for sub-grid convection is limiting in resolving convective rainfall and affects the locations dominated by such rainfall (especially tropics).

...

*Both the bias analysis of quantiles and the FSS evaluation show that the performance of BARRA gradually improves as we move from northern to southern regions (**Error! Reference source not found.** and **Error! Reference source not found.**). This variation in performance can partly be attributed to the different rainfall climate at these regions. The convective precipitation during summer season is dominant at northern (low) latitudes, whereas winter (or uniform) non-convective rainfall are dominant at southern (high) latitudes.*

I can’t see that you address the uncertainty in the observation based data. Please discuss these, and perhaps make an attempt to quantify them and include in the figures.

In our study we use two datasets as a benchmark for evaluation: gauge and blended radar. In absence of any other alternative of high-quality datasets, these two datasets represent the best available point and spatial estimates of rainfall and thus provide an appropriate basis for comparison. We acknowledge the uncertainty arising due to comparison of point rainfall against BARRA grid and have discussed it accordingly. However, without suitable reference data sets it is not possible to calculate the uncertainty in these sub-daily observations and there is no published study on this which is relevant to the regions we studied. Similarly, the radar datasets are prone to various error sources which are discussed in the “2. Study area and data sources” section of the manuscript. The Bureau of Meteorology have blended radar estimates with gauged data and have estimated the associated uncertainties however, this information has not been published and is not publicly available.

p81228: You state that BARRA tends to overestimate light rain events. Please add a reference or show this in a figure.

A reference has been added and the sentence reads (Lines:240-241) *This difference is partly due to the tendency of BARRA to overestimate light rain events (Su et al., 2019).*

p101292: You state that point precipitation is generally higher than areal rainfall at 12 km scale. Could you suggest a way to overcome this? Could you consider using an areal reduction factor? If not, why? In many studies lately there has been a focus on quantifying the contribution of changes in intensity and changes in the frequency to trends in (heavy) precipitation. Could you please relate your results to the how well BARRA represents intensity and frequency?

Despite an apparent mismatch in spatial resolution/representation, we use point rainfall as a benchmark for evaluating BARRA rainfall because it is one of the best available datasets for evaluation. Accordingly, we discuss the variation in performance in light of the difference in spatial scale. Addressing the performance variation arising from differences in the spatial representation of point and areal rainfalls is not straightforward. In “design rainfall” applications (ie those based on probabilistic representations of rainfall intensities for design purposes), areal reduction factors are applied to scale point rainfalls to areal rainfalls, however, we note that 1) such factors could lack the actual properties of large rainfall events and result in mis-estimation of flood risks (Wright et al., 2014), and 2) there is limited robust evidence for areal reduction factors at this small scale (Podger et al., 2015; Stensmyr et al., 2015). In hydrological modelling applications, any constant scaling applied to point rainfall would not hold true for individual events as they are derived to preserve statistical characteristics of the entire time series. One way of addressing such differences could be to evaluate BARRA rainfall against high-quality reference spatial datasets, but such data sets would require considerable effort to derive.

The focus of our study is to present an assessment of sub-daily rainfall from BARRA at point and spatial scales, where the period of evaluation is limited to a six-year period (2010-2015) based on the availability of observed (benchmark) datasets. Given the limited time period of the BARRA data available at the time of this study, it is difficult to make any meaningful inferences regarding trends in the intensity and frequency of rainfalls. However, it is hoped that future assessments will consider the contribution of changes in rainfall intensity and frequency to trends in (heavy) precipitation.

Figures

Figure 1: Please thicken the line marking the four study regions.

Addressed in the revised manuscript.

Figure 2: Although I like this figure, it is a bit hard to see the colors etc due to the small maps. Could you split the maps and the boxplots into two figures?

We agree that the figure as currently presented is slightly difficult to read. Since the maps and boxplots are linked together, following reviewer #2’ suggestion, we have changed the orientation of the plot to make it bigger in the revised manuscript.

Figure 5: I might have missed something, but I do not understand why you here only study precipitation up to 6 hours, while up to 24 hours in Figure 2.

Figure 2 and Figure 5 are results from two different evaluation approaches: non-conditional frequency distribution at a point, and rainfall events over an area. For the former, we explore the bias in rainfall

intensity at various frequencies and temporal accumulations up to 24 hours. Our attempt to compute the frequencies up to 24 hours was to understand the variation in rainfall frequencies at different temporal accumulations. As discussed in introduction section, this assessment could be useful for developing intensity-frequency curves for design applications.

With the areal analysis, we use Fractions Skill Scores (FSS) to understand the representativeness of spatial patterns of rainfall from BARRA at sub-daily scales. Our analysis of temporal accumulations of 3 and 6 hours was undertaken to assess the utility of the areal rainfalls at sub-daily temporal scales that are relevant to hydrological modelling. The evaluation of multiple sub-daily aggregations (1, 3 and 6 hours) allows us to determine a suitable temporal aggregation for hydrological modelling purposes. While extending this accumulation to 24 hours will definitely show improved metrics, the resulting time series would have more limited application to hydrological modelling.

In response to this comment, we have revised the Methodology section (Lines: 157-158, 229-231) as follows:

The analysis of frequencies corresponding to different accumulations (up to 24 h) is selected to be of relevance to design rainfall applications.

...

It is worth noting that FSS metric only provides information on variation of performance with increasing spatial scale. The timing errors at finer temporal scales can be indirectly discerned by analysing variation of FSS with time accumulations. Therefore, the FSS is evaluated at temporal aggregations of 3 h and 6 h.

Technical corrections:

p2132-33: This sentence should be rephrased. p2138: Remove “a”.

p2152: “developing use cases”? Please rephrase.

p91258: Add “the” before “two datasets”

p101304: Do you mean “mixed result between locations”?

These technical corrections have been addressed in the revised manuscript.

Response to Referee #2

General comments

The article is sound and makes a modest contribution (by examining sub-daily time scales) to body of literature on the evaluation of reanalysis rainfall data. I think it should be published subject to some clarification.

We thank Referee #2 for their positive remarks on the paper, and for their constructive comments for improving the manuscript.

Specific comments

Some items for correction / clarification:

L19: quantile should be replaced with percentile.

Agreed. We have made corrections in the revised manuscript.

L47-48: what does it mean that BARRA is driven by ERA-Interim?

The initial and boundary conditions required for BARRA is obtained from ERA-Interim. We have revised the sentence (Lines: 46-49) to make it clearer.

It is driven with initial and boundary conditions from the global ERA-Interim reanalysis (~79 km) and provides estimates over the Australasian region from 1990 to 2018.

L104 - 115: this seems an overly critical view of radar rainfall. In the assessment of the sources of error in radar retrievals, are the authors referring to their own gauge correction procedure or other published work? Further, I understand that the authors use the radar data to evaluate the spatial distribution of rainfall, but how does aggregating the 1.5km data to 12km (which I assume they did for a fair evaluation) change the interpretation of the spatial patterns for such small regions of Australia. At 12km resolution, an evaluation against satellite retrievals (e.g. GPM IMERG final product) may provide the same information but for the whole country.

Our assessment of radar was based on the review of various studies mentioned in the paper. The blended radar data were made available from Bureau of Meteorology, Australia. The blended radar data had already undergone gauge correction procedures and any assessment regarding such corrections is beyond the scope of the paper.

We acknowledge that the radar dataset is the best available spatial dataset and provides an accurate estimate of spatio-temporal distribution of rainfall. In addition, we apply area-weighted approach to re-grid BARRA (~12km) to radar grid (~1.5). It can be expected that the re-gridding of BARRA will underestimate the intensity of rainfall at a finer scale. To address this, we apply a percentile-based threshold while calculating Fractions Skill Score (FSS) to evaluate the spatial distribution of the rainfall field. Nonetheless, any results from FSS obtained for a spatial scale less than <12km needs to be interpreted carefully.

Similarly, we agree that evaluating BARRA across the whole country would provide valuable information. Such evaluation, however, would be limited by the availability of high-quality and high-resolution benchmark datasets. As mentioned in the comment by Reviewer #2, satellite retrievals such as the IMERG final product could possibly be used for such a large-scale evaluation. Assessments of the IMERG final product have been shown to perform better than TRMM or the IMERG initial run (Beck et al., 2019; Wang et al., 2017), however, a majority of such evaluations are limited to daily scales (Beck et al., 2019; Wang and Yong, 2020). As our assessment focuses on sub-daily rainfall from the BARRA reanalysis dataset it is necessary to use benchmark data that is accurate at that temporal scale. However, there are currently no comprehensive assessments of the IMERG final run at sub-daily scales for Australian continent, and this forces us to focus our evaluation on more accurate radar datasets at selected locations. A more detailed spatial assessment of BARRA would be possible once further comprehensive assessments of high-resolution satellite datasets are available.

In the revised manuscript, we have clarified the rationale for the choice of study areas and reference datasets used in the current study. We have further discussed our current limitations in evaluating over entire Australia and provided comment on possible directions for future assessments. The manuscript is revised as follows (Lines: 99-101, 367-371):

The spatial evaluation across the entire Australia is impeded by the availability of high-resolution observation-based datasets. Therefore, the spatial evaluation is undertaken by using high quality radar-based datasets at four city-centred regions (Brisbane, Darwin, Melbourne and Sydney).

...

One of the limitations of this study is that the spatial assessment of BARRA is restricted to a few locations where radar-based datasets are available and the performance of BARRA across Australia is generalised based on evaluation at those locations. Currently, the lack of high-quality and high-resolution benchmark datasets limits our ability to fully understand the performance of BARRA across the entire continent. However, a more detailed spatial assessment of BARRA would be possible once such benchmark datasets are available.

Box plots in Fig. 2 are very difficult to see on the printed version. Perhaps a landscape layout for figure 2 might help?

We agree that the figure, currently, is difficult to read. We have changed the orientation of the plot in the revised manuscript.

References

- Acharya, S. C., Nathan, R., Wang, Q. J., Su, C.-H. and Eizenberg, N.: An evaluation of daily precipitation from a regional atmospheric reanalysis over Australia, *Hydrol. Earth Syst. Sci.*, 23(8), 3387–3403, doi:10.5194/hess-23-3387-2019, 2019.
- Beck, H. E., Pan, M., Roy, T., Weedon, G. P., Pappenberger, F., van Dijk, A. I. J. M., Huffman, G. J., Adler, R. F. and Wood, E. F.: Daily evaluation of 26 precipitation datasets using Stage-IV gauge-radar data for the CONUS, *Hydrol. Earth Syst. Sci.*, 23(1), 207–224, doi:10.5194/hess-23-207-2019, 2019.
- Clark, P., Roberts, N., Lean, H., Ballard, S. P. and Charlton-Perez, C.: Convection-permitting models: A step-change in rainfall forecasting, *Meteorol. Appl.*, 23(2), 165–181, doi:10.1002/met.1538, 2016.
- Ebert, E. E., Janowiak, J. E. and Kidd, C.: Comparison of near-real-time precipitation estimates from satellite observations and numerical models, *Bull. Am. Meteorol. Soc.*, 88(1), 47–64, doi:10.1175/BAMS-88-1-47, 2007.
- de Leeuw, J., Methven, J. and Blackburn, M.: Evaluation of ERA-Interim reanalysis precipitation products using England and Wales observations, *Q. J. R. Meteorol. Soc.*, 141(688), 798–806, doi:10.1002/qj.2395, 2015.
- Podger, S., Green, J., Jolly, C., Beesley, C. and others: Creating long duration areal reduction factors, in 36th Hydrology and Water Resources Symposium: The art and science of water, p. 39, Engineers Australia., 2015.
- Stensmyr, P., Babister, M., Adam, M. and others: Short duration areal reduction factors: Sydney,

Melbourne and Brisbane, in 36th Hydrology and Water Resources Symposium: The art and science of water, p. 121., 2015.

Su, C.-H., Eizenberg, N., Steinle, P., Jakob, D., Fox-Hughes, P., White, C. J., Rennie, S., Franklin, C., Dharssi, I. and Zhu, H.: BARRA v1.0: the Bureau of Meteorology Atmospheric high-resolution Regional Reanalysis for Australia, *Geosci. Model Dev.*, 12(5), 2049–2068, doi:10.5194/gmd-12-2049-2019, 2019.

Wang, H. and Yong, B.: Quasi-Global Evaluation of IMERG and GSMaP Precipitation Products over Land Using Gauge Observations, *Water*, 12(1), 243, doi:10.3390/w12010243, 2020.

Wang, Z., Zhong, R., Lai, C. and Chen, J.: Evaluation of the GPM IMERG satellite-based precipitation products and the hydrological utility, *Atmos. Res.*, 196, 151–163, doi:10.1016/j.atmosres.2017.06.020, 2017.

Wright, D. B., Smith, J. A. and Baeck, M. L.: Critical Examination of Area Reduction Factors, *J. Hydrol. Eng.*, 19(4), 769–776, doi:10.1061/(ASCE)HE.1943-5584.0000855, 2014.

Ability of an Australian reanalysis dataset to characterise sub-daily precipitation

Suwash Chandra Acharya¹, Rory Nathan¹, Quan J Wang¹, Chun-Hsu Su^{1,2}, Nathan Eizenberg²

¹Department of Infrastructure Engineering, the University of Melbourne, Australia

5 ²Bureau of Meteorology, Melbourne, Australia

Correspondence to: Suwash Chandra Acharya (suwasha@student.unimelb.edu.au)

Abstract. The high spatio-temporal variability of precipitation is often difficult to characterise due to limited measurements. The available low-resolution global reanalysis datasets inadequately represent the spatio-temporal variability of precipitation relevant to catchment hydrology. The Bureau of Meteorology Atmospheric high-resolution Regional Reanalysis for Australia (BARRA) provides a high-resolution atmospheric reanalysis dataset across the Australasian region. For hydrometeorological applications, however, it is essential to properly evaluate the sub-daily precipitation from this reanalysis. In this regard, this paper evaluates the sub-daily precipitation from BARRA for a period of 6 years (2010-2015) over Australia against point observations and blended radar products. We utilise a range of existing and bespoke metrics for evaluation at point and spatial scales. We examine bias in quantile estimates and spatial displacement of sub-daily rainfall at a point scale. At a spatial scale, we use the Fractions Skill Score as a spatial evaluation metric. The results show that the performance of BARRA precipitation depends on spatial location with poorer performance in tropical relative to temperate regions. A possible spatial displacement during large rainfall is also found at point locations. This displacement, evaluated by comparing the distribution of rainfall within a day, could be quantified by considering the neighbourhood grids. On spatial evaluation, hourly precipitation from BARRA are found to be skilful at a spatial scale of less than 100km (150km) for a threshold of 75%-th percentile quantile (90%-th quantilepercentile) at most of the locations. The performance across all the metrics improves significantly at time resolutions higher than 3 h. Our evaluations illustrate that the BARRA precipitation, despite discernible spatial displacements, serves as a useful dataset for Australia, especially at sub-daily resolutions. Users of BARRA are recommended to properly account for possible spatio-temporal displacement errors, especially for applications where the spatial and temporal characteristics of rainfall are deemed very important.

25 1 Introduction

Precipitation is highly variable across both space and time, especially at spatial and temporal scales relevant to catchment hydrology (Michaelides et al., 2009). An understanding of the spatio-temporal pattern of precipitation is vital for many scientific and operational applications such as hydro-climatic modelling and the forecasting of floods (Golding et al., 2016; Kucera et al., 2013; Paschalis et al., 2014). This understanding relies on access to high-resolution precipitation datasets.

30 However, the availability of fine-scale precipitation products (e.g. spatial resolution less than around 0.25° at hourly time scale) are limited. The general sources of precipitation data are gauge measurements, ground-based radars, satellites and atmospheric reanalysis models (Michaelides et al., 2009). Gauge measurements, which are localised in nature, are hindered~~hindered by sparse measurement and~~by the sparse and uneven ~~density distribution~~ of gauge network, ~~—~~ whereas the coverage of ground-based radars is limited. Global reanalysis datasets (e.g. NCEP-CFSR, Saha et al., 2010; ERA-Interim, Dee et al., 2011; JRA-55, Kobayashi et al., 2015) and satellite products (e.g. TMPA, Huffman et al., 2007; IMERG, Huffman et al., 2018) provide a continuous and consistent estimate at varying spatial (0.05° to 2.5°) and temporal resolution (hourly to daily).

An atmospheric reanalysis merges observations and models to provide a four-dimensional earth system data at a regular spatial and temporal resolution over ~~a~~ long time period, often years and decades (Parker, 2016). The variables in the reanalysis (such as precipitation, cloud and soil moisture) are related to one another through modelled physical relations and with the analysed observations. By undertaking modelling over a limited area, a regional atmospheric reanalysis can provide precipitation estimates at finer spatial and temporal scale than a global reanalysis. It can incorporate more observations at a finer scale to better constrain the evolution of a higher-resolution model, and thus can account for the effects of mountains, coastlines, and mesoscale atmospheric circulations in greater detail. Such analyses can thereby provide precipitation estimates with greater spatial relevance to local fine-scale studies than coarser-scale models. BARRA (Bureau of Meteorology atmospheric high-resolution regional reanalysis for Australia) is one such high-resolution (~ 12 km) regional reanalysis (BARRA-R). It is driven with initial and boundary conditions from ~~by~~ the global ~~~ 79 km~~ ERA-Interim reanalysis (~ 79 km) and provides estimates over the Australasian region from 1990 to 2018 (Jakob et al., 2017; Su et al., 2019).

50 BARRA-R (referred to here as simply BARRA) provides hourly precipitation estimates at 12 km horizontal resolution (Jakob et al., 2017; Su et al., 2019). Rainfall observations are not assimilated in this reanalysis and precipitation is estimated by model physics and parameterisation. Evaluating its suitability for scientific studies and ~~developing use cases~~its use in hydrological applications can be facilitated by identifying its strengths and limitations through a quantitative assessment. An initial assessment of daily precipitation from BARRA showed that it was able to reproduce the precipitation statistics and large precipitation at point (gauged locations) and grid scale of 5 km (Acharya et al., 2019; Su et al., 2019). However, precipitation data at sub-daily temporal resolutions are essential to support the application of flood modelling (Chiaravalloti et al., 2018) and analysis of precipitation extremes in convective systems. Therefore, it is useful to assess the performance of BARRA at sub-daily resolution along with its ability to represent the spatial structure of rainfall events.

60 The performance of BARRA at a daily scale was summarised using continuous metrics and categorical metrics at both point and grid scales (Acharya et al., 2019). However, at sub-daily scales, such one-on-one evaluations can be misleading for several reasons (Jerney and Renshaw, 2016). First, sub-daily precipitation is dominated by zero values and has a highly skewed distribution, which makes it difficult to interpret the correlation statistics. In addition, these metrics doubly penalise a modelled

rainfall field that is displaced in space or time: once as a missed observation, and again as a false alarm. This situation is popularly known as “double-penalty problem” (Rossa et al., 2008), and to mitigate this issue it is necessary to adopt an evaluation approach which considers likely displacements in space and time (Gilleland et al., 2009; Jerney and Renshaw, 2016; Thiemig et al., 2012). For example, when evaluating heavy rainfall events at point scale, Thiemig et al. (2012) adopted an error metric that explicitly considers possible time-lags in the gridded dataset. Similarly, Jerney and Renshaw (2016) considered temporal displacement while evaluating the precipitation by grouping the events into seven synthetic categories with varying temporal shift and bias structure. Besides these methods which look at temporal displacement, a possible shift across space can be considered when evaluating high-resolution precipitation datasets. However, such evaluations may be limited by the availability of suitable gridded observation data for comparison, and it is thus preferable to undertake assessments using multiple evaluation techniques (Jerney and Renshaw, 2016).

The metrics used to undertake the evaluation need to match the prospective use of the dataset. For example, it may be sufficient for some urban and stormwater design problems to only evaluate the non-conditional frequency distribution of precipitation at a point. However, for most hydrological modelling purposes, it is necessary to evaluate the representation of spatio-temporal characteristics of rainfall events over the catchment of interest. The spatial and temporal resolution of the prospective applications also play a critical role in selection of the performance measures, so it is useful to assess the efficacy of the reanalysis over a range of different temporal and spatial accumulations.

The objective of this paper is to present an evaluation of sub-daily BARRA precipitation over temporal and spatial resolutions that are relevant to catchment hydrology applications. The evaluation is undertaken using ground measurements and radar-based rainfall observations across Australia over a six-year period spanning 2010-2015. Following the assessment of BARRA at daily scale (Acharya et al., 2019), this study further explores BARRA precipitation at different time accumulations ranging from hourly to daily. In addition, we compare the temporal distribution of gauged rainfalls with neighbouring grids, specifically for large rainfall events. We also employ a spatial evaluation metric (Roberts and Lean, 2008) to evaluate the ability of the reanalysis dataset to represent the spatial distribution of rainfalls against a gauge-corrected radar dataset. Overall, our aim is to assess the ability of BARRA to capture the behaviour of sub-daily precipitation at the catchment scale, particularly for large events. To this end, we use a combination of existing and bespoke metrics to evaluate performance over different temporal and spatial accumulations in different climatic zones.

2 Study area and data sources

The primary sources of reference data used to evaluate the performance of BARRA data were derived from pluviometer gauges and radar-based products. Ground-based observations are not assimilated in BARRA and thus these data serve as an independent dataset for evaluation. [The pluviometer observations in this study encompass three of the broad climatic zones](#)

across Australia, namely tropical, temperate and arid (Figure 1). Rainfall in Australia is highly variable across space and time. Rainfall is concentrated during summer in the tropical north, whereas rainfall is more prevalent during winter in the temperate south and southwest. The southeast region experiences a more consistent rainfall throughout the year. The central arid region receives the least total rainfall, and the eastern coast the highest average rainfall. The spatial evaluation across the entire Australia is impeded by the availability of high-resolution observation-based dataset. Therefore, the spatial evaluation is undertaken by using radar-based datasets at four city-centred regions (Brisbane, Darwin, Melbourne and Sydney). These study areas are located in the tropical and temperate zones with varying rainfall climates.

~~The study areas for the analysis of pluviometer observations encompass three of the broad climatic zones across Australia, namely tropical, temperate and arid. The radar-based evaluation is undertaken at four city-centred regions that are located in these three zones (Brisbane, Darwin, Melbourne and Sydney).~~

Hourly rainfall observations from pluviometer gauges were obtained from the Australian Bureau of Meteorology. Observations from a total of 441 gauges were selected, covering a common period of record between 2010 and 2015 (Figure 1). The observed rainfall is used to approximate precipitation as most precipitation in Australia is in the form of rainfall.

The blended radar precipitation dataset was also obtained from the Australian Bureau of Meteorology for regions surrounding the four city centres (see Figure 1). The spatial resolution of the radar data is ~1.5 km, and it is only available from 2014. The radar fields have been blended with the observed rainfall using conditional merging (Sinclair and Pegram, 2005). The approach is in principle close to a copula-based approach, which exhibits less bias and yields a smaller error metric compared to non-parametric radar rainfall estimation (Hasan et al., 2016). The blended radar estimates are still likely to be erroneous and biased due to reasons that include: errors in reflectivity measurement (e.g., radar beam overshoot, terrain blocking, clutter), inaccurate radar reflectivity and rain rate relationship, tendency of the radar to underestimate rainfall at distance, and quality control algorithms rejecting gauge data used for bias adjustment (Chumchean et al., 2006; Seo et al., 2010). Further, there is a fundamental difference in representativity between radar measurements and modelled precipitation, where radar infers precipitations at height over a cubic kilometre in size. However, we follow prior studies (e.g. Mittermaier et al., 2013; Roberts and Lean, 2008) and use radar rainfall to compare spatial rainfall at sub-daily time steps assuming that the spatial distribution of rainfall is preserved. In addition, we apply quantile-based thresholds in order to remove the potential bias in the daily rainfall totals, although we note the strategy remains limited by the fact that the bias is spatially varying.

BARRA utilises the Unified Model (UM, Davies et al., 2005) and its 4D variational data assimilation system (4D Var). BARRA extends spatially over 65.0° to 196.9° east, -65.0° to 19.4° north at a spatial resolution of 0.11° (approximately 12 km) and with 70 levels up to 80 km into the atmosphere. The model includes a comprehensive set of parametrisations, including a modified boundary layer scheme, mixed phase cloud microphysics, a mass flux convection scheme, and a radiation scheme. The model parameterisation in BARRA is mainly inherited from the UKMO Global Atmosphere (GA) 6.0 configurations as

130 described in Walters et al. (2017). Observations from land surface stations, ships, buoys, aircrafts, radiosondes, and satellites are assimilated in BARRA, conducted 4 times a day with a 6-hour analysis window centred at time $t_0 = 0, 6, 12$ and 18 UTC. Surface and satellite rainfall observations are not assimilated, and the precipitation fields are determined by the modelled dynamics. In particular, they are estimated from the 12-hour (h) forecast runs of the UM from $t_0 - 3h$, using the microphysics scheme based on Wilson and Ballard (1999) and the mass flux convective parameterisation scheme of Gregory and Rowntree
135 (1990). The former describes the atmospheric processes that transfer water between the various states of water to remove moisture resolved on the grid scale. At 12 km horizontal resolution, BARRA requires the convection scheme to model sub-grid scale convection using an ensemble of cumulus clouds as a single entraining-detraining plume (Clark et al., 2016). The scheme prevents unstable growth of cloud structures on the grid and explicit vertical circulations, and can only predict an area-average rainfall instead of a spectrum of rainfall rates. This parameterisation scheme adopted for sub-grid convection is limiting
140 in resolving convective rainfall and affects the locations dominated by such rainfall (especially tropics). Further details on parameterisation and assimilation schemes in BARRA are provided by Su et al. (2019).

3 Methodology

Hourly BARRA precipitation estimates at 12 km resolution are evaluated at varying spatial and temporal scales. A range of point-to-grid and grid-to-grid evaluations are undertaken using gauged rainfall and blended radar estimates over various
145 temporal and spatial scales.

3.1 Frequency distribution of sub-daily rainfall

A point-to-grid analysis is undertaken to evaluate the sub-daily frequency distribution of BARRA precipitation. At each gauge location as shown in Figure 1, corresponding BARRA estimates are obtained using the nearest neighbour interpolation. The basis of comparison requires some thought as the fraction of zero values in sub-daily rainfall data is high (the 95% quantiles
150 at hourly temporal resolution are zero). This issue could be addressed by selecting suitable conditional thresholds, though different thresholds would need to be adopted for each temporal accumulation. For example, a threshold of 0.1mm/h could be adopted for hourly data and 1 mm/day for daily. However, the slight problem with this approach is that any trends in the performance metric with temporal accumulation will be confounded by the somewhat arbitrary choice of thresholds for intermediate temporal accumulations (3h, 6h, and 12h). Accordingly, we derived quantiles for sub-daily rainfalls that occurred
155 only on the rain days for each dataset individually, where, a “rain day” is defined based on a threshold of 1 mm/day (Ebert et al., 2007).

In investigating the frequency of rainfall, we compute the various quantiles (80, 90, 95, 99%) at different time accumulations (1 h, 3 h, 6 h, 12 h and 24 h). The analysis of frequencies corresponding to different accumulations (up to 24 h) is selected to
160 be of relevance for design rainfall applications. We then estimate percentage bias using $(m-o)/o * 100$, where m and o denote

the reanalysis and observed precipitation corresponding to the selected quantiles. This evaluation is further stratified across three broad climatic zones (arid, tropical, and temperate) as defined by the Köppen-Geiger classification (Peel et al., 2007; Figure 1). It is worth noting that comparisons of point (gauged) and areal (gridded) rainfall are generally biased as the average precipitation over a grid cell is lower than rainfall recorded at a particular point. It would also be expected that sub-daily point rainfalls are more variable than those averaged over a grid cell area.

3.2 Neighbourhood-based diurnal patterns at point locations ~~Spatio-temporal distribution~~

A direct comparison of precipitation, especially in higher resolution datasets, is difficult as the conventional metrics are not able to penalise intensity and location errors in a desirable manner (Rossa et al., 2008). Performance metrics which compare point observations with model estimates averaged over a grid cell are heavily influenced by errors in the spatial pattern and location of rainfall events, even if the average rainfall depths over the local domain are the same. This is particularly the case with high-resolution precipitation datasets, and with the analysis of sub-daily periods.

To mitigate this problem in a gauge-based evaluation, we adopt an approach that allows for possible timing and displacement differences in rainfall occurrence. The approach is similar to the neighbourhood (or “fuzzy”) approach used with single observation-neighbourhood forecasts (Ebert, 2008), in which gridded observations were evaluated against gridded forecasts. This method accounts for situations where an event defined on the basis of gauge measurements may miss the nearby grid cell resulting in spatial error, and/or appear non-coincident at the nearby grid cell signifying a temporal error. For the former case, an evaluation that considers neighbouring grid cells can account for spatial errors. For cases involving a time displacement, a moving storm may appear at a neighbouring grid cell at the time-step under consideration, approaching towards the nearby grid cell.

To account for these different types of errors, we employ an analysis that explicitly considers the occurrence and timing of rainfalls at neighbouring grid cells. This evaluation is undertaken by selecting large rainfall events (greater than 10mm/day) in the gauge dataset. The cumulative fraction of rainfall over the day is computed for the gauge and the nearest neighbouring grid (Eq. 1). The squared difference in the fraction of cumulative rainfall occurring in each hour between the gauge and reanalysis rainfalls is then averaged to provide an error score (Eq. 2).

$$F_h = \sum_{i=1}^h f_i, \quad 1 \leq h \leq 24 \quad (1)$$

where, f_i is the fraction of daily rainfall occurring at i^{th} hour.

$$\text{error score} = \frac{1}{23} \sum_{h=1}^{23} (F_{h,model} - F_{h,gauge})^2 \quad (2)$$

Note that 23 is used in the denominator as $F_{24, model} = F_{24, gauge} = 1$ and are not included in the computation of the error score. The temporal error is considered by using cumulative precipitation at a daily scale, which penalises large temporal errors more than small ones. Similarly, to account for possible spatial displacement, we analyse the temporal distribution of precipitation by searching over neighbourhood space to find the best performing grid cell. The error score is computed for both the nearest grid and the neighbourhood grid cells equidistant from the nearest grid. The minimum error score is selected and then averaged across all rainfall events at a location.

The average error score varies between 0 and 1. A score of 1 represents the worse possible situation in which rainfall occurs in the first hour in one dataset and the last hour in the other. Conversely, a score of 0 indicates a perfect match between observations and model estimates. Scores between 0 and 1 indicate differing degrees of temporal error, where for example, a score of 0.33 indicates that rainfall occurs in either the first or last hour in one data set and is distributed uniformly throughout the day in the other.

3.3 Fractions skill score (FSS) Neighbourhood-based spatial evaluation

Finally, a spatial evaluation of reanalysis precipitation against blended radar estimates is undertaken. For each spatial domain, the largest 25 rainfall events to have occurred over the 3-year period are selected based on domain-averaged daily precipitation. Radar precipitation, which is available as 30-minute accumulations, is aggregated to hourly to match the temporal resolution of BARRA. Similarly, a common spatial resolution is adopted. The precipitation from BARRA (~12km) is re-gridded to the resolution of the radar grid (~1.5km) using area-weighted re-gridding, which means that single BARRA precipitation estimates are distributed over 8×8 radar grid cells.

Different metrics have been developed for undertaking spatially variable evaluations (Ebert, 2008, 2009). This study adopts the FSS from Roberts and Lean (2008), as it measures the variation of skill across increasing spatial scales and hence indicates the minimum spatial scale at which the model is skilful.

The FSS metric is based on the likelihood that rainfall over a given threshold occurs somewhere within the neighbouring window of grid cells. A common threshold rainfall rate is selected for both the observed and modelled grid cells. An increasing window of size $n \times n$ centred on a particular grid is selected (which yields N windows over the whole domain). For each window i , a fraction of grid points exceeding the threshold in observed rainfall ($p_{o,i}$) and modelled rainfall ($p_{m,i}$) are computed. Then, the FSS is calculated as:

$$FSS = 1 - \frac{\frac{1}{N} \sum_{i=1}^N (p_{m,i} - p_{o,i})^2}{\frac{1}{N} \sum_{i=1}^N p_{m,i}^2 + \frac{1}{N} \sum_{i=1}^N p_{o,i}^2} \quad (3)$$

The FSS score varies between 0 and 1. A score of 0 represents a complete mismatch between two rainfall fields and a score of 1 represents a perfect match. Usually, FSS is computed for varying size of spatial windows and results are plotted as a function of window size. A random score (FSS_{random}) is the FSS that would be achieved, on average, by a random field with the same fraction of observed events (p_o) over the domain. A benchmark score, a target or ‘uniform’ skill (FSS_{uniform}), is given to a uniform field with a probability of occurrence equal to p_o at each grid cell (Roberts and Lean, 2008). FSS_{uniform} is expressed as $0.5 + p_o/2$, which is halfway between a perfect score (1) and a random score ($FSS_{\text{random}} = p_o$). This FSS_{uniform} can be approximated to 0.5 when p_o is small, as in the case of larger precipitation thresholds.

225

The thresholds used to calculate FSS are based on rainfall quantiles in observed and reanalysis data which are derived for each time step. A rainfall intensity greater than 0.2 mm/h is used as a threshold to define a rainy grid cell for which quantile-based thresholds are computed. This ensures that the model and observed rainfall fields have an identical fraction of rain events for each threshold value. The application of quantile based thresholds aims to remove the impact of any bias in rainfall amount and focus solely on spatial accuracy of modelled precipitation (Mittermaier et al., 2013; Roberts and Lean, 2008; Skok and Roberts, 2018). It is worth noting that FSS metric only provides information on variation of performance with increasing spatial scale. The timing errors at finer temporal scales can be indirectly discerned by analysing variation of FSS with time accumulations. Therefore, the FSS is evaluated at temporal aggregations of 3 h and 6 h.

230

4 Results

235

4.1 Frequency distribution of sub-daily rainfall

240

Figure 2 illustrates the spatial distribution of quantile bias and its summary across climatic zones. Overall, the spatial distribution of bias across all time accumulations and quantiles exhibit a similar pattern. The biases in the northern region are higher and spatially more variable than those in the southern region. The variation in bias across quantiles is the highest for hourly rainfall, especially in arid and tropical climatic zones. For the 80% quantile of hourly rainfalls, all stations in the arid and tropical zones exhibit a higher positive bias than those in the temperate zone. This difference is partly due to the tendency of BARRA to overestimate light rain events (Su et al., 2019). Despite considering wet days only in comparing quantiles, hourly rainfall at 80% quantile for arid and tropical region dominated by small precipitation amount which results in a high positive bias. At higher quantiles, the BARRA estimates are all lower than the gauged point rainfalls. This negative bias is largest in the tropical zone, followed by the arid zone. Although the nature of the high hourly bias varies with location, a step reduction

245 in bias is observed when the accumulation time periods increase from 1 to 3 and 6 hours. This is partly due to reduced inherent bias arising from the adoption of longer temporal accumulations which reduces the potential for differences in timing between observations and model estimates.

The spatial distribution of biases associated with the different quantiles exhibits a similar pattern for all temporal accumulations considered. At shorter accumulations (up to 3 hours), the biases change from positive to negative and gradually increase in magnitude with increasing quantile. At higher accumulations (6, 12 and 24 hours), the bias is negative but decreases in magnitude with increasing quantiles. At higher quantiles (95% and 99%), the bias is the least at the temperate zone. The BARRA estimates are under-predicted at all time accumulations, but this improves with increasing temporal accumulations.

4.2 Neighbourhood-based diurnal patterns at point locations ~~Spatio-temporal patterns~~

255 Figure 3 shows how the minimum error score changes with varying neighbourhood grid size. It is seen that the error score decreases significantly when a neighbourhood of 3×3 grid cells (about 35×35 km) is considered instead of a single nearest neighbour only. The error score continues to decrease slightly as the size of the neighbourhood increases, though the adoption of a larger neighbourhood increases the likelihood that rainfall events unrelated to the gauge observations are being considered. The results show that the temporal distribution of the BARRA precipitation estimates within a wet day are representative yet displaced in terms of location. The error scores are all less than 0.33, which indicates that the temporal distribution of sub-daily BARRA precipitation is on average superior to a uniform distribution of estimates derived by simply disaggregating daily rainfalls over a day.

4.3 Neighbourhood-based spatial evaluation ~~Fractions Skill Score (FSS)~~

265 Figure 4 shows the hourly precipitation rates corresponding to different quantile thresholds in the Brisbane, Darwin, Melbourne, and Sydney regions, for the largest 25 events that occurred between 2014 to 2016. The Melbourne and Sydney regions are similar than other domains in terms of the frequency distribution of hourly precipitation and the discrepancy between radar and BARRA precipitation.

270 Overall, the rainfall magnitude corresponding to quantiles is higher for the radar estimates than for BARRA, and this can be attributed to the difference in spatial scale and the area-weighted re-gridding scheme. The difference between the two datasets is greater at higher quantiles. However, at around 99% quantiles, the precipitation from BARRA is close to or greater than radar precipitation.

275 FSS is calculated for varying quantile thresholds (50, 75, 90, 95, and 99%) at different accumulations across time (1h, 3h and 6h) and presented in Figure 5. As expected, FSS increases with increasing neighbourhood size and decreases with increasing

threshold. If there is no frequency bias, then the FSS curve is expected to asymptote to 1 as the neighbourhood size is increased. In Figure 5, the maximum FSS of 1 is not achieved even at the neighbourhood size of 300 km, which signifies a frequency bias in BARRA.

280 The FSS scores vary with location, where the regions in increasing order of performance are: Darwin, Brisbane, Melbourne, and Sydney. The results for hourly precipitation and 75% quantile threshold suggest that the skilful spatial scale L ($FSS > 0.5 + p_o/2$) is less than 100 km for all the locations except Darwin. The skilful spatial scale for 90% and 95% quantile thresholds increases to 150 km. For the 99% quantile threshold, BARRA estimates only exhibit useful skill over spatial scales larger than around 250 km.

285

The FSS metric only provides information on how performance varies with increasing spatial scale. It does not account for timing errors associated with events that might be initiated at different times and/or evolve at different rates. An indication of such timing errors may be discerned by assessing how the FSS varies with increasing time accumulations. The results shown in Figure 5 indicate that FSS improves with increasing time accumulations. For example, at Darwin, BARRA is able to provide skilful estimates of 50% threshold of hourly rainfalls only at a scale of 200 km; however, at 3 h and 6 h accumulations, the corresponding spatial scale reduces to 125 km and 100 km, respectively. The accuracy of the BARRA estimates decreases with increasing rainfall severity, and even at the longest time accumulations, the spatial scale at which rainfalls above the 99% quantiles are skilful extends out to 300 km.

290

5 Discussion

295 To understand the performance of BARRA precipitation at sub-daily scales, a range of evaluation methods are employed to ascertain its spatial and temporal characteristics. Key insights arising from the results are discussed below.

5.1 Performance based on wet day quantiles

The unconditional evaluation of precipitation frequency in terms of wet day quantiles examines the representativeness of sub-daily climatology (Figure 2). The minimum bias across all time accumulations and quantiles in the temperate zone suggests that BARRA provides unbiased estimates of temperate rainfalls over the observed range of events. In addition, an improved performance is observed across all regions when temporal accumulations are considered. This includes the arid and tropical zones where BARRA performs poorly at an hourly scale. A slight negative bias is observed across all locations in most of the quantiles. This underestimation is however expected due to the mismatch in spatial scale between point observations and grid average precipitation (Maraun, 2013). The point precipitation, in general, is expected to be higher than areal rainfall at 12 km spatial scale.

305

5.2 Spatio-temporal representation of rainfall

A potential displacement of precipitation field in space and time is expected when evaluating high-resolution precipitation datasets, especially when performance is assessed at an hourly timescale and at a point (or single grid cell) location (Rossa et al., 2008). The BARRA estimates exhibit such displacement errors, as evidenced by improved performance at neighbourhood analysis and FSS analysis. An assessment of hourly temporal patterns shows an improvement when neighbourhood grid cells are considered (Figure 3). This suggests that when using precipitation from BARRA, users could benefit from considering spatial and temporal displacement in the precipitation field, especially during large events. This further suggests an opportunity to utilise hourly distribution of rainfall for disaggregating daily totals.

At the catchment scale, the accuracy of the spatial distribution of rainfall is important for flood modelling. The evaluation of the spatial performance of BARRA against radar data for selected large events showed a mixed result ~~based on~~between locations. BARRA rainfalls for 90% quantile at 3 h accumulation achieves the target FSS at a spatial scale of range 100-140 km for all domains except Darwin (250 km) (Figure 5). Achieving useful skill only at a large spatial scale can be partly attributed to the spatial error in the precipitation fields. The higher quantile thresholds are related to small-scale features which are even more likely to be subject to spatial error. Therefore, it is difficult to achieve a skilful spatial scale for these more extreme rainfalls. This has also been pointed out by Roberts and Lean (2008) based on the skill of NWP model outputs at various quantile thresholds (75, 90, 95 and 99%). The nature of such spatial displacement errors should be considered if BARRA estimates of precipitation are intended to be used for hydrological modelling.

This spatial evaluation is conducted at selected locations to take advantage of the availability of high-resolution blended radar datasets. Across the whole Australian continent there is a dearth of such high-resolution observations of precipitation, and BARRA could be relied upon to provide estimates of sub-daily precipitation at selected spatial scales in regions where there is a paucity of gauging data.

5.3 Performance dependence on spatial location

The overall performance of BARRA varies with location, and this similar trend was evident across all evaluation methods. ~~Both the bias analysis of quantiles and the FSS evaluation show that BARRA performs more poorly in the tropical regions (Figure 2 and Figure 5). Both the bias analysis of quantiles and the FSS evaluation show that the performance of BARRA gradually improves as we move from northern to southern regions (Figure 2 and Figure 5). This variation in performance can partly be attributed to the different rainfall climate at these regions. The convective precipitation during summer season is dominant at northern (low) latitudes, whereas winter (or uniform) non-convective rainfall are dominant at southern (high) latitudes.~~ The poorer performance in the tropics, where convective precipitation is dominant, reflects the limitations of the parameterisation scheme to describe sub-grid scale rainfall (Su et al., 2019). This is consistent with the finding of Ebert et al.

(2007) that focuses on general performance of numerical weather prediction models. The variation in performance of sub-daily BARRA precipitation across spatial location and different climatologies is consistent with the daily evaluation performed in Acharya et al. (2019).
340

5.4 Performance as a function of temporal resolution

Availability of hourly rainfall observations (based on both pluviometer and radar products) enable BARRA estimates of hourly rainfalls to be evaluated. The uneven and sparse distribution of pluviometer gauges across Australia and the limited availability of radar products is not sufficient for an overall assessment of sub-daily precipitation across the whole of Australia, yet given the wide range of climatologies considered, the analyses provide a comprehensive evaluation of BARRA. Our unconditional (Figure 2) and direct comparison (Figure 5) both illustrate a similar dependency of performance on temporal resolution. The performance shows a significant improvement when estimates are accumulated from 1h to 3h. The increased performance for coarser temporal resolution can also be linked to the smoothing effect of rainfall at larger accumulation times and reduction of inherent bias.
345

350

The trade-off between performance and temporal resolution needs to be considered in combination with the objectives to which the precipitation estimates are used. However, in general terms the increased performance at 3 h and 6 h accumulation suggests that it is better to use these accumulations from BARRA. Given that the performance of BARRA does vary with location, it is expected that shorter (3 h) accumulations may be appropriate for use in temperate locations and longer accumulations (6 h) in the tropical and arid regions.
355

6 Conclusions

An accurate representation of spatial and sub-daily temporal characteristics of precipitation fields is important for many hydro-meteorological applications. BARRA is a regional reanalysis dataset that provides long-term high-resolution estimates of atmospheric variables over the Australian continent. In this study, the spatio-temporal characteristics of sub-daily BARRA precipitation estimates are assessed using various metrics to evaluate its performance at various spatial and temporal scales. Based on the results, we conclude:
360

1. Sub-daily precipitation exhibits negative bias at higher quantiles. The magnitude of bias decreases with increasing temporal accumulation and is the lowest at daily accumulations.
2. The spatial displacement of rainfall estimates is evident, especially for rainfall corresponding to higher quantiles.
- 365 3. The performance of BARRA varies spatially, and more accurate estimates are provided in southern and eastern Australia (temperate zone) than in northern Australia (tropical zone). These spatial trends in model performance are evident

in evaluations undertaken using both gauged point rainfalls and blended radar observations, and are consistent with the evaluation of BARRA estimates of daily rainfall (Acharya et al., 2019).

4. Performance increases with time aggregation, which is expected due to smoothing in accumulated rainfall time series.

370 The performance is reasonably skilful at most of the locations for temporal accumulations of 3h and greater.

One of the limitations of this study is that the spatial assessment of BARRA is restricted to a few locations where radar-based datasets are available and the performance of BARRA across Australia is generalised based on evaluation at those locations. Currently, the lack of high-quality and high-resolution benchmark datasets limits our ability to fully understand the performance of BARRA across the entire continent. However, a more detailed spatial assessment of BARRA would be possible once such benchmark datasets are available.

375

A natural interest of the users of BARRA is the potential application of the data in hydro-meteorology. Information on sub-daily rainfalls across the Australian continent is limited due to sparse gauge measurements, the availability of radar at only few locations, spatially coarser global reanalysis products, and few satellite products (~25km, going back to only 1998).

380 BARRA stands out as one of few available datasets at hourly temporal resolution. In that respect, it serves as a useful dataset for the applications requiring sub-daily precipitation. Further, it can be used to help characterise rainfall behaviour in regions where gauges are sparse or non-existent. One of the potential applications of BARRA could be for probabilistic design rainfalls, that is for those engineering applications which utilise information on the relationship between rainfall magnitude and its exceedance probability, either at a point or over an area; such applications are probabilistic in nature and are less sensitive to event-based errors in temporal and spatial behaviour. For applications in flood modelling, where the spatial and temporal distribution of precipitation is important, a proper account needs to be given to the likely spatio-temporal displacement errors in the BARRA precipitation.

385

-The ability of BARRA to provide areal rainfall estimates is useful in catchment hydrological applications. In addition to direct reliance on absolute estimates of precipitation, it may be that BARRA is well suited to providing information on sub-daily patterns of areal rainfalls that can be scaled to match more reliable estimates of daily rainfalls. The extent to which such estimates might provide a better understanding of areal sub-daily rainfall behaviour than is possible from gauged point rainfalls warrants further investigation. Accordingly, future work will be directed towards evaluation and design applications of areal rainfall estimates.

390

Acknowledgements

395 The authors gratefully acknowledge the financial support provided by Seqwater and the Bureau of Meteorology to partially fund SCA's PhD scholarship. We would like to thank Susan Rennie, Kevin Cheong, and Alan Seed (Bureau of Meteorology) for their advice on the use of the radar rainfall products, as retrieved from the Rainfields Archiving System provided by the


Bureau of Meteorology. Also, Dörte Jakob and Peter Steinle (Bureau of Meteorology) contributed valuable discussions on general methodology and early results.

400 **References**

- Acharya, S. C., Nathan, R., Wang, Q. J., Su, C.-H. and Eizenberg, N.: An evaluation of daily precipitation from a regional atmospheric reanalysis over Australia, *Hydrol. Earth Syst. Sci.*, 23(8), 3387–3403, doi:10.5194/hess-23-3387-2019, 2019.
- Chiaravalloti, F., Brocca, L., Procopio, A., Massari, C. and Gabriele, S.: Assessment of GPM and SM2RAIN-ASCAT rainfall products over complex terrain in southern Italy, *Atmos. Res.*, 206(December 2017), 64–74, doi:10.1016/j.atmosres.2018.02.019, 2018.
- 405 Chumchean, S., Sharma, A. and Seed, A.: An integrated approach to error correction for real-time radar-rainfall estimation, *J. Atmos. Ocean. Technol.*, 23(1), 67–79, doi:10.1175/JTECH1832.1, 2006.
- Davies, T., Cullen, M. J. P., Malcolm, A. J., Mawson, M. H., Staniforth, A., White, A. A. and Wood, N.: A new dynamical core for the Met Office’s global and regional modelling of the atmosphere, *Q. J. R. Meteorol. Soc.*, 131(608), 1759–1782, doi:10.1256/qj.04.101, 2005.
- 410 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J. N. and Vitart, F.: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, 137(656), 553–597, doi:10.1002/qj.828, 2011.
- Ebert, E. E.: Fuzzy verification of high-resolution gridded forecasts: A review and proposed framework, *Meteorol. Appl.*, 15(1), 51–64, doi:10.1002/met.25, 2008.
- Ebert, E. E.: Neighborhood Verification: A Strategy for Rewarding Close Forecasts, *Weather Forecast.*, 24(6), 1498–1510, doi:10.1175/2009WAF2222251.1, 2009.
- 420 Ebert, E. E., Janowiak, J. E. and Kidd, C.: Comparison of near-real-time precipitation estimates from satellite observations and numerical models, *Bull. Am. Meteorol. Soc.*, 88(1), 47–64, doi:10.1175/BAMS-88-1-47, 2007.
- Gilleland, E., Ahijevych, D., Brown, B. G., Casati, B. and Ebert, E. E.: Intercomparison of Spatial Forecast Verification Methods, *Weather Forecast.*, 24(5), 1416–1430, doi:10.1175/2009WAF2222269.1, 2009.
- 425 Golding, B., Roberts, N., Leoncini, G., Mylne, K. and Swinbank, R.: MOGREPS-UK Convection-Permitting Ensemble Products for Surface Water Flood Forecasting: Rationale and First Results, *J. Hydrometeorol.*, 17(5), 1383–1406, doi:10.1175/JHM-D-15-0083.1, 2016.
- Hasan, M. M., Sharma, A., Johnson, F., Mariethoz, G. and Seed, A.: Merging radar and in situ rainfall measurements: An assessment of different combination algorithms, *Water Resour. Res.*, 52(10), 8384–8398, doi:10.1002/2015WR018441, 2016.

- 430 Huffman, G. J., Bolvin, D. T., Nelkin, E. J., Wolff, D. B., Adler, R. F., Gu, G., Hong, Y., Bowman, K. P. and Stocker, E. F.: The TRMM Multisatellite Precipitation Analysis (TMPA): Quasi-Global, Multiyear, Combined-Sensor Precipitation Estimates at Fine Scales, *J. Hydrometeorol.*, 8(1), 38–55, doi:10.1175/JHM560.1, 2007.
- Huffman, G. J., Bolvin, D. T., Braithwaite, D., Hsu, K., Joyce, R., Kidd, C., Nelkin, E. J., Sorooshian, S., Tan, T. and Xie, P.: NASA Global Precipitation Measurement (GPM) Integrated Multi-satellitE Retrievals for GPM (IMERG), Algorithm Theor.
- 435 Basis Doc., (February), 1–31, doi:https://pmm.nasa.gov/resources/documents/gpm-integrated-multi-satellite-retrievals-gpm-imerg-algorithm-theoretical-basis-, 2018.
- Jakob, D., Su, C.-H., Eizenberg, N., Kociuba, G., Steinle, P., Fox-Hughes, P. and Bettio, L.: An atmospheric high-resolution regional reanalysis for Australia, *B. Aus. Meteorol. Oceanog. Soc.*, 30(3), 16–23, 2017.
- Jermey, P. M. and Renshaw, R. J.: Precipitation representation over a two-year period in regional reanalysis, *Q. J. R. Meteorol.*
- 440 Soc., 142(696), 1300–1310, doi:10.1002/qj.2733, 2016.
- Kobayashi, S., Ota, Y., Harada, Y., Ebata, A., Moriya, M., Onoda, H., Onogi, K., Kamahori, H., Kobayashi, C., Endo, H., Miyaoka, K. and Takahashi, K.: The JRA-55 Reanalysis: General Specifications and Basic Characteristics, *J. Meteorol. Soc. Japan. Ser. II*, 93(1), 5–48, doi:10.2151/jmsj.2015-001, 2015.
- Kucera, P. A., Ebert, E. E., Turk, F. J., Levizzani, V., Kirschbaum, D., Tapiador, F. J., Loew, A. and Borsche, M.: Precipitation
- 445 from space: Advancing earth system science, *Bull. Am. Meteorol. Soc.*, 94(3), 365–375, doi:10.1175/BAMS-D-11-00171.1, 2013.
- Maraun, D.: Bias Correction, Quantile Mapping, and Downscaling: Revisiting the Inflation Issue, *J. Clim.*, 26(6), 2137–2143, doi:10.1175/JCLI-D-12-00821.1, 2013.
- Michaelides, S., Levizzani, V., Anagnostou, E., Bauer, P., Kasparis, T. and Lane, J. E.: Precipitation: Measurement, remote
- 450 sensing, climatology and modeling, *Atmos. Res.*, 94(4), 512–533, doi:10.1016/j.atmosres.2009.08.017, 2009.
- Mittermaier, M., Roberts, N. and Thompson, S. A.: A long-term assessment of precipitation forecast skill using the Fractions Skill Score, *Meteorol. Appl.*, 20(2), 176–186, doi:10.1002/met.296, 2013.
- Parker, W. S.: Reanalyses and Observations: What’s the Difference?, *Bull. Am. Meteorol. Soc.*, 97(9), 1565–1572, doi:10.1175/BAMS-D-14-00226.1, 2016.
- 455 Paschalis, A., Fatichi, S., Molnar, P., Rimkus, S. and Burlando, P.: On the effects of small scale space–time variability of rainfall on basin flood response, *J. Hydrol.*, 514, 313–327, doi:10.1016/j.jhydrol.2014.04.014, 2014.
- Peel, M. C., Finlayson, B. L. and McMahon, T. A.: Updated world map of the Köppen-Geiger climate classification, *Hydrol. Earth Syst. Sci.*, 11(5), 1633–1644, doi:10.5194/hess-11-1633-2007, 2007.
- Roberts, N. M. and Lean, H. W.: Scale-Selective Verification of Rainfall Accumulations from High-Resolution Forecasts of
- 460 Convective Events, *Mon. Weather Rev.*, 136(1), 78–97, doi:10.1175/2007MWR2123.1, 2008.
- Rossa, A., Nurmi, P. and Ebert, E.: Overview of methods for the verification of quantitative precipitation forecasts, in *Precipitation: Advances in Measurement, Estimation and Prediction*, pp. 419–452, Springer Berlin Heidelberg, Berlin, Heidelberg., 2008.

- Saha, S., Moorthi, S., Pan, H. L., Wu, X., Wang, J., Nadiga, S., Tripp, P., Kistler, R., Woollen, J., Behringer, D., Liu, H.,
465 Stokes, D., Grumbine, R., Gayno, G., Wang, J., Hou, Y. T., Chuang, H. Y., Juang, H. M. H., Sela, J., Iredell, M., Treadon, R.,
Kleist, D., Van Delst, P., Keyser, D., Derber, J., Ek, M., Meng, J., Wei, H., Yang, R., Lord, S., Van Den Dool, H., Kumar, A.,
Wang, W., Long, C., Chelliah, M., Xue, Y., Huang, B., Schemm, J. K., Ebisuzaki, W., Lin, R., Xie, P., Chen, M., Zhou, S.,
Higgins, W., Zou, C. Z., Liu, Q., Chen, Y., Han, Y., Cucurull, L., Reynolds, R. W., Rutledge, G. and Goldberg, M.: The NCEP
climate forecast system reanalysis, *Bull. Am. Meteorol. Soc.*, 91(8), 1015–1057, doi:10.1175/2010BAMS3001.1, 2010.
- 470 Seo, D.-J., Seed, A. and Delrieu, G.: Radar and multisensor rainfall estimation for hydrologic applications, *Geophys. Monogr. Ser.*, 191, 79–104, doi:10.1029/2010GM000952, 2010.
- Sinclair, S. and Pegram, G.: Combining radar and rain gauge rainfall estimates using conditional merging, *Atmos. Sci. Lett.*, 6(1), 19–22, doi:10.1002/asl.85, 2005.
- Skok, G. and Roberts, N.: Estimating the displacement in precipitation forecasts using the Fractions Skill Score, *Q. J. R. Meteorol. Soc.*, 144(711), 414–425, doi:10.1002/qj.3212, 2018.
- 475 Su, C.-H., Eizenberg, N., Steinle, P., Jakob, D., Fox-Hughes, P., White, C. J., Rennie, S., Franklin, C., Dharssi, I. and Zhu, H.: BARRA v1.0: the Bureau of Meteorology Atmospheric high-resolution Regional Reanalysis for Australia, *Geosci. Model Dev.*, 12(5), 2049–2068, doi:10.5194/gmd-12-2049-2019, 2019.
- Thiemig, V., Rojas, R., Zambrano-Bigiarini, M., Levizzani, V. and De Roo, A.: Validation of Satellite-Based Precipitation
480 Products over Sparsely Gauged African River Basins, *J. Hydrometeorol.*, 13(6), 1760–1783, doi:10.1175/JHM-D-12-032.1, 2012.
- Walters, D., Boutle, I., Brooks, M., Melvin, T., Stratton, R., Vosper, S., Wells, H., Williams, K., Wood, N., Allen, T., Bushell, A., Copsey, D., Earnshaw, P., Edwards, J., Gross, M., Hardiman, S., Harris, C., Heming, J., Klingaman, N., Levine, R., Manners, J., Martin, G., Milton, S., Mittermaier, M., Morcrette, C., Riddick, T., Roberts, M., Sanchez, C., Selwood, P., Stirling, A., Smith, C., Suri, D., Tennant, W., Vidale, P. L., Wilkinson, J., Willett, M., Woolnough, S. and Xavier, P.: The Met Office Unified Model Global Atmosphere 6.0/6.1 and JULES Global Land 6.0/6.1 configurations, *Geosci. Model Dev.*, 10(4), 1487–1520, doi:10.5194/gmd-10-1487-2017, 2017.



490

Figure 1: Study area with locations of pluviometer gauges (points) and the radar data (box surrounding city centres). The locations of radar datasets are Darwin, Brisbane, Sydney and Melbourne regions. The climatic classification is adapted from Peel et al. (2007).




Figure 2: Percentage bias in precipitation at quantiles (columns) and accumulations (rows). The boxplots represent the summary of percentage bias across climatic zones for respective rows and columns.





Figure 3: Boxplots showing the distribution of minimum error score across stations in each neighbourhood grid size. The box represents 25-75th quantile values, horizontal line in the box represents median, and whiskers represent 5-95th quantiles.



505 Figure 4: Hourly precipitation rates corresponding to quantiles for BARRA and blended radar data at four different locations.




Figure 5: Mean FSS as a function of neighbourhood distance for rainfall above quantile thresholds (indicated by colours) at various locations (rows) and accumulations (columns). The dashed horizontal lines indicate the target or uniform (FSS_{target} or FSS_{uniform}) skill for each threshold as specified by Roberts and Lean (2008)

510