

Effects of climate anomalies on warm-season low flows in Switzerland

Marius G. Floriancic^{1,2}, Wouter R. Berghuijs², Tobias Jonas³, James W. Kirchner^{2,4}, Peter Molnar¹

¹ Institute of Environmental Engineering, ETH Zurich, 8093 Zürich, Switzerland

² Department of Environmental Systems Science, ETH Zurich, 8092 Zürich, Switzerland

³ WSL Institute for Snow and Avalanche Research SLF, 7260 Davos Dorf, Switzerland

⁴ Swiss Federal Research Institute WSL, 8903 Birmensdorf, Switzerland

Corresponding author: Marius G. Floriancic (floriancic@ifu.baug.ethz.ch)

Keywords: low flow, hydrological drought, precipitation, evapotranspiration

Short summary: Low river flows affect societies and ecosystems. Here we study how precipitation and potential evapotranspiration shape annual warm-season low flows across a network of 380 Swiss catchments. Low flows in these rivers typically result from below-average precipitation and above-average potential evapotranspiration. The lowest low flows result from long periods of the combined effects of both drivers.

Abstract. Switzerland has faced extended periods of low river flows in recent years (2003, 2011, 2015, and 2018), with major economic and environmental consequences. Understanding the origins of events like these is important for water resources management. In this work we provide data illustrating the individual and joint contributions of precipitation and evapotranspiration to low flows in both typical and dry years. To quantify how weather drives low flows, we explore how deviations from mean seasonal climate conditions (i.e., climate anomalies) of precipitation and potential evapotranspiration correlate with the occurrence and magnitude of annual 7-day lowest flows (Q_{\min}) during the warm season (May through November) across 380 Swiss catchments from 2000 through 2018. Most warm-season low flows followed periods of below-average precipitation and above-average potential evapotranspiration, and the lowest low flows resulted from both of these drivers acting together. Low-flow timing was spatially variable across Switzerland in all years, including the driest (2003, 2011, 2015, and 2018). Low flows in these driest years were associated with much longer-lasting climate anomalies than the ≤ 2 -month anomalies which preceded typical warm-season low flows in other years. We found that snow water equivalent and winter precipitation totals only slightly influenced the magnitude and timing of warm-season low flows in low-elevation catchments across Switzerland. Our results provide insight into how precipitation and potential evapotranspiration jointly shape warm-season low flows across Switzerland, and potentially aid in assessing low-flow risks in similar mountain regions using seasonal weather forecasts.

1. Introduction

In recent decades, Europe has experienced several severe droughts (Van Lanen et al., 2016). Their impacts, such as dry river reaches and high water temperatures, have a range of adverse effects on society and river ecology (e.g., Poff et al., 1997; Bradford & Heinonen, 2008; Price et al., 2011; Rolls et al., 2012; van Vliet et al., 2012). Severe low flows in the years 2003, 2011, 2015 and 2018 led to substantial economic losses by limiting water availability for households, industry, irrigation and hydropower, as well as impacting river transportation (Stahl et al., 2016; Munich Re, 2019). Such effects are expected to become more severe and frequent as water demand rises, and as droughts are anticipated to increase in frequency and intensity in the future (e.g., De Stefano et al., 2012; Wada et al., 2013), leading to calls for improved understanding and management of droughts and their effects on low flows across Europe (e.g., Seneviratne et al., 2012a; Van Lanen et al., 2016; WMO, 2008).

In temperate climates, annual low flows typically occur in two distinct seasons, i.e. during late summer and autumn in warmer regions and during winter in colder regions (Fiala et al., 2010; Smakhtin, 2001). This typical low-flow seasonality has been reported for many regions of the world, including, for example, Austria (Laaha & Blöschl, 2006; Van Loon & Laaha, 2015), the Rhine river basin (Demirel et al., 2013; Tongal et al., 2013), and North America (Cooper et al., 2018; Dierauer et al., 2018; Wang, 2019). Switzerland also has two low-flow seasons, where the distinction between warm-season low flows and winter low flows is strongly connected to elevation (Wehren et al., 2010; Weingartner & Aschwanden, 1992). Low flows tend to occur in late summer and early autumn (August through October) in low-elevation Swiss catchments, and during the winter (January through March) in high-elevation catchments.

Catchment properties shape low flows by controlling the storage and release of water (e.g., Stoelzle et al., 2014; Van Lanen et al., 2013; Van Loon & Laaha, 2015; Staudinger et al., 2017), but the landscape itself does not cause low flows. Instead, the drivers of low flows are meteorological conditions that dry out catchments (e.g., Fleig et al., 2006; Haslinger et al., 2014; Smakhtin, 2001). Warm-season low flows are typically caused by sustained periods of high evapotranspiration and low precipitation, whereas winter low flows often follow sustained periods of sub-freezing temperatures (e.g.; Laaha et al., 2013; Van Loon, 2015). The duration of these anomalous weather conditions is critical in shaping the annual lowest flows. Their timing varies between years and is largely driven by climate seasonality. In this paper we refer to weather conditions that deviate from the seasonal norm as ‘climate anomalies’ regardless of the magnitude of this departure.

The two main climatic factors controlling water storage and release in a catchment are precipitation and temperature (through its influence on snow processes and evapotranspiration). Therefore precipitation (P) and potential evapotranspiration (E_p) anomalies are expected to be important drivers of warm-season low flows across Switzerland. Precipitation controls the amount of water that is available for runoff in a catchment, and sustained periods with little precipitation will inevitably reduce storage and thereby limit streamflow. Because there is a time lag between low precipitation and low streamflow, meteorological droughts (i.e., precipitation deficits) result in hydrological droughts and/or low flows if they persist for long enough (e.g., Peters et al., 2006; Tallaksen & Van Lanen, 2004; Van Loon, 2015). In Switzerland, there is limited precipitation seasonality, but precipitation can still vary substantially within seasons or from year to year. However, precipitation is expected to become increasingly seasonal with changing climatic conditions in the future, with less precipitation during summer and more precipitation in winter. In addition, anticipated changes in snowfall and snow packs may also alter river flows (CH2018, 2018).

High temperatures can be an indicator of high E_p , and thus high potential for depletion of soil moisture storage, reducing aquifer recharge and streamflow (e.g., Jaeger & Seneviratne, 2011; Vidal et al., 2010). Temperature extremes can be amplified

when low soil moisture limits evapotranspiration, leading to lower relative humidity and higher air temperatures, which further
75 increase E_p (Granger, 1989). Furthermore, vegetation decreases the amount of water available for streamflow by increasing
transpiration during periods of high vapor pressure deficits. Although these mechanisms are known, the effects of
evapotranspiration on river low flows have received relatively little attention compared to precipitation effects.
Seneviratne et al. (2012) reported that low flows across Switzerland in 2003 more likely resulted from excess
evapotranspiration than from spring precipitation deficits, and Teuling et al. (2013) have documented the depletion of soil
80 water storage by high evapotranspiration during past European low flows. Woodhouse et al. (2016) reported that temperatures
rather than precipitation explained the interannual streamflow variations of the Colorado river. More recently Cooper et al.
(2018) reported that summer low flows in the maritime Western US are largely driven by summer E_p , rather than by winter
precipitation or snow water equivalent. Mastrotheodoros et al. (2020) modeled how increasing evapotranspiration strongly
reduced streamflow across the European Alps during the summer of 2003. Future E_p is projected to increase along with
85 increases in incoming longwave radiation (Roderick et al., 2014), with uncertain consequences for future low flows. In
Switzerland, temperatures are expected to rise even quicker than the global average in the next decades (CH2018, 2018),
potentially influencing low-flow dynamics.

Future climate changes will also affect low flows in mountain regions by altering snow accumulation and melt. Multiple studies
90 have examined how winter precipitation and snow water equivalent affect summer low flows in high-elevation catchments.
For example, Godsey et al. (2014) found that shrinking snowpacks in the Sierra Nevada of California led to smaller low flows
in the following summers. Jenicek et al. (2016) reported that maximum snow accumulation strongly affected summer low
flows across several Swiss mountainous catchments. Dierauer et al. (2018) found that warmer winters with less snow
accumulation led to lower summer low flows in mountainous catchments of the Western United States. Recently, Wang (2019)
95 reported that climate warming might increase aquifer conductivity and thereby streamflow in cold region watersheds. Future
climate warming in both warm and cold seasons will most likely impact summer low flows through different mechanisms. In
summer, higher temperatures increase potential evapotranspiration, whereas in winter they reduce snowpacks (e.g., Déry et al.,
2009; Diffenbaugh et al., 2015; Musselman et al., 2017).

100 The effects of precipitation, temperature and evapotranspiration on low flows have been investigated for individual events or
individual catchments and regions in the literature. Previous studies have largely focused on how signatures of low flows
(averaged across many events) relate to catchment and climate characteristics (e.g., Fangmann & Haberlandt, 2019; Hannaford,
2015; Laaha & Blöschl, 2006; Van Loon & Laaha, 2015). To our knowledge, however, no studies have systematically assessed
the direct impact of temperature and precipitation during periods immediately preceding individual annual low-flow events
105 across many catchments in a topographically diverse region.

Here we explore how precipitation and E_p deviations from their seasonal norms (here termed “climate anomalies”) jointly
shape the occurrence and magnitude of annual warm-season low flows across a network of 380 Swiss catchments. Because
the annual lowest flow is an atypical flow condition, we expect it to follow atypical weather conditions, rather than reflecting
110 climate seasonality alone. Therefore, we hypothesize that annual lowest flows will typically occur after anomalous weather
conditions, that is, weather conditions that deviate from the seasonal average. Understanding how anomalous weather drives
low flows may help to reveal the processes at work, and also support low-flow forecasting. Switzerland is an interesting study
region because gauging and climate data are available from a dense station network spanning a wide range of elevations,
climates, and topographies. We investigate (a) how precipitation and E_p anomalies separately and jointly shape the occurrence

115 and magnitude of warm-season low flows across Switzerland, (b) which durations of these anomalies have the strongest impact
on low-flow occurrence and magnitude, both in typical and in exceptionally dry years, and (c) how winter precipitation and
snow packs influence the magnitude and timing of warm-season low flows. Understanding these connections is important for
anticipating how streamflows are likely to respond as the exceptionally dry years of today are expected to become more typical
in a future warmer climate.

120

2. Data and methodology

2.1. Streamflow and climate data

We compiled daily streamflows for 380 gauging stations across Switzerland for a 19-year period (2000-2018), using data
collected by the Swiss Federal Office of the Environment (FOEN) and the Swiss Cantonal authorities. This data set excludes
125 catchments with obvious anthropogenic influences on the hydrograph, e.g., from major dams or hydropeaking operations. Low
flows were defined as the lowest 7-day average streamflow for each year (Q_{\min}). We calculated the magnitude and timing of
 Q_{\min} in each catchment for each year from 2000 to 2018. Not all catchments had continuous data for all 19 years; in total we
could calculate low-flow magnitude and timing for 6237 station-years. This data set included years when the lowest annual
flows were much higher than typical low flows (e.g., in especially wet years and years without distinct dry periods). We
130 removed all annual low flows above the threshold of 2.5 mm d^{-1} , which is the 25th percentile of daily discharges across all
catchments, because flows above this threshold cannot be considered truly low flows. This resulted in the removal of
approximately 2% of all low flows, leaving a total of 6124 station-years for our analysis. We split the dataset of annual low
flows into cold-season low flows occurring between December and April and warm-season low flows occurring between May
and November. In total, we observed 2122 cold-season low flows and 4002 warm-season low flows across the 380 catchments
135 within the 19-year time period.

We determined catchment area and mean catchment elevation for each gauging station based on a 2-m DEM
(SwissAlti3D 2016, Swisstopo), using functions provided in the ArcGIS “Spatial Analyst” toolbox. The catchments range in
size from 1 to 519 km^2 , vary in mean elevation from 309 to 2930 m, and have diverse landcovers and climates. Daily gridded
140 precipitation and temperature data ($\sim 2 \times 2 \text{ km}$ cells; Meteoswiss products “RhiresD” and “TabsD”) were used to derive
catchment-averaged weather and climate conditions. Daily potential evapotranspiration (E_p) was estimated following the
method of Hargreaves & Samani (1985). A gridded dataset of snow water equivalent (SWE) on March 1st of each year was
used to estimate catchment-average SWE. The SWE product was based on data from 320 Swiss snow monitoring stations that
were assimilated into a distributed snow cover model (Magnusson et al., 2014; Griessinger et al., 2016). We use SWE on
145 March 1st instead of April 1st because our focus is on warm-season low flows in lower-elevation catchments, most of which
have no substantial snow left by April 1st (Steger et al., 2013; Winstral et al., 2019; Lüthi et al., 2019).

2.2. Anomalies of climate variables

To infer climate conditions preceding annual low flows, we selected the annual 7-day minimum streamflow events (Q_{\min}) in
150 each catchment for each year from 2000 to 2018. We then calculated precipitation and potential evapotranspiration for time
windows of different lengths prior to each annual low flow. We hypothesize that severe low flows will usually follow periods
in which precipitation and potential evapotranspiration significantly deviate from their seasonal averages. Thus, we define
climate anomalies as deviations of precipitation and potential evapotranspiration from their climatic norms, defined as their
long-term averages on the same day of the year. For example, we quantify precipitation anomalies (in mm) by:

$$\sum_{t=d_l-d_t}^{d_l} (P(t) - \bar{P}(t)) \quad Eq.(1)$$

where $P(t)$ is daily precipitation (mm) at day t , $\bar{P}(t)$ is the climatic mean precipitation on day t averaged across all of the years of record, d_l is the time period over which anomalies are calculated for each annual low flow, and d_l is the day of the low flow. We vary the time period d_l from one week to half a year (7, 14, 30, 60, 90, 120, 182 days), with the endpoint always being the date of the low flow. For example, the 30-day precipitation anomaly for a low flow that happened on 30 September 2018 is calculated using the sum of precipitation from 1st to 30th September 2018 minus the mean of precipitation for all 1st to 30th September periods from 2000 to 2018. We calculate E_p anomalies in the same way.

2.3. Statistical tests and quantification of process importance

We first report the spatial distribution of the timing of the annual lowest flows across Switzerland for 2000 until 2018. We then show the magnitude of 30-day climate anomalies before each annual low flow as a function of elevation. The mechanisms involved in generating annual cold-season low flows and warm-season low flows are different, thus we split our dataset into cold-season and warm-season low flows. From this point on, we report results only for warm-season low flows. To quantify the relationship between the magnitudes of climate anomalies and the magnitudes of warm-season annual low flows, we use Spearman rank correlation coefficients (r_s) as a robust estimator (Legates & McCabe, 1999). We report these rank correlations across all catchments in a histogram. To test the regional significance of the r_s coefficients we use the sign test.

We assess the impact of the length of climate anomalies preceding the annual warm-season low flows by comparing the magnitude of P and E_p anomalies for the different time windows (7, 14, 30, 60, 90, 120, 182 days) between the four driest years and the more typical years. We also correlate the magnitude of warm-season Q_{min} with the number of days that P and E_p exceed certain thresholds. The threshold that defines low precipitation is the 20th percentile of 10-day running averages of precipitation over the entire period of record. Similarly, the threshold that defines high E_p is the 80th percentile of 10-day running averages of E_p over the entire period of record. We report the distribution of rank correlations calculated for each catchment based on the 19 years of data in histograms. The magnitudes of the annual low flows are shown as boxplots for each individual year. The horizontal line in the boxplots indicates the median, the box represents the interquartile range, and the whiskers extend to 1.5 times the interquartile range above and below the box; the dots are outliers.

To quantify the individual and joint importance of the magnitude of P and E_p anomalies, we first calculated the bivariate Spearman rank correlation between the individual anomalies and Q_{min} for the different time windows (30, 60, 90, 120, 182 days) for all years (2000-2018) and for the years with the lowest low flows (2003, 2011, 2015 and 2018). For this analysis we reduced the original dataset to only those catchments where at least 5 years of Q_{min} data were available, as suggested in WMO (2008). In a next step we used the joint anomalies of P and E_p for all durations 30, 60, 90, 120, 182 days to predict Q_{min} with a multivariate stepwise generalized linear model (GLM) fitted by minimizing RMSE. We then computed the fraction of the GLM's R^2 attributable to the individual precipitation and E_p anomalies for each duration, to assess the relative contribution of each anomaly to the prediction of Q_{min} . We compared the results for all years to those for the lowest-flow years (2003, 2011, 2015 & 2018) to assess whether the relations between climate anomalies and Q_{min} differed during the driest years.

To test how warm-season low flows are influenced by precipitation and snow processes in the preceding winter, we calculated the Spearman rank correlations between the total December-March precipitation sum and the following warm-season Q_{min} ,

and between SWE on 1st March and the following warm-season Q_{\min} . We again report these rank correlations across multiple catchments in histograms, and test the significance of these distributions of correlations by the sign test.

Finally, we assess whether the correlations we obtained between P and E_p anomalies and warm-season Q_{\min} are influenced by the extent of human impact in each catchment. We quantify human impact by the fraction of human affected land cover in each catchment. As a proxy for human activity we use the Corine landcover dataset (CLC, 2018) and calculate the fraction of catchment area with “Artificial surfaces”. We then show histograms of the rank correlations between P , E_p , and Q_{\min} in the 20% of catchments with the most human-influenced land use, and the 20% of catchments with the least human-influenced land use, compared to the distribution across all catchments. We assessed the significance of the differences between the obtained distributions by the Student t-Test.

3. Results

3.1. Spatial patterns of low-flow timing

During the dry years of 2003, 2011, 2015, and 2018, low-flow conditions occurred across large parts of Europe (Laaha et al., 2017; Van Lanen et al., 2016). Annual low flows did not occur simultaneously across Switzerland, but instead occurred primarily during winter in the Alpine regions and summer and autumn in the Swiss Plateau (Fig. 1). In addition, within these two sub-regions, the timing of low flows was still spatially variable, indicating that annual low flows may be surprisingly asynchronous across Switzerland, even in unusually dry years. Within the Swiss Plateau, low-flow timing is more spatially consistent during some years without severe low flows (e.g., 2009, 2013, 2016), than during others (e.g., 2000, 2002, 2004, 2010, 2017).




Figure 1: The timing of occurrence of annual low flows across Switzerland for the years 2000 to 2018 in the two main regions: the Swiss Plateau and Swiss Alps (roughly the northern and southern halves of the country, respectively). Low-flow timing tended to be spatially heterogeneous, even in years when large parts of Europe simultaneously experienced severe low flows (2003, 2011, 2015, 2018).

220 **3.2. Climate anomalies control low-flow timing and magnitude**

The occurrence of low flows is linked to periods of below-average P and above-average E_p (Fig. 2a&b). However, distinct site-to-site differences exist: at elevations below approximately 1500 m asl, almost all annual low flows occur after periods of anomalously high potential evapotranspiration and anomalously low precipitation (Fig. 2a&b). At higher elevations, by contrast, E_p anomalies have no systematic effect and precipitation anomalies become less important with increasing elevation.

225 This reduced importance of anomalies at these higher elevations is probably because low flows here result primarily from freezing temperatures (or periods of snow accumulation), rather than precipitation or E_p patterns. Low flows at higher elevations occur during the winter months when there is a lack of liquid water inputs to catchments, due to precipitation mostly accumulating as snow, and little snowmelt. These processes are mainly driven by sustained below-zero temperatures. Thus, the main determining factor in winter low flows at high elevations (or in cold environments) will likely be the length of the

230 snow accumulation period, rather than what the exact temperatures were, or how much precipitation occurred.




Figure 2: Altitudinal variation in 30-day anomalies of precipitation (a) and potential evapotranspiration (b) preceding (warm and cold season) annual low flows from 2000 through 2018. Blue and red horizontal bars indicate the range between the minima and maxima of these anomalies at each catchment across the 19 years of this study. Yellow bars show moving averages of these climate anomalies for bins of 10 catchments ordered by elevation. Note that the elevation scale is not linear. Low flows are associated with below-average precipitation (a) and above-average potential

evapotranspiration (b). Histograms of rank correlations between anomalies of precipitation (c) and potential evapotranspiration (d) and low-flow magnitudes for warm-season (May through November) low flows across Swiss catchments. Results for cold-season low flows can be found in the supplementary Fig. S2.

More severe climate anomalies lead to lower low flows (Fig. 2c&d). Spearman rank correlations of magnitudes of the climate anomalies to magnitudes of Q_{\min} (shown for the months May through November) indicate that lower precipitation in the 30 days prior to Q_{\min} usually results in smaller Q_{\min} (median $r_s=0.28$). Similarly, higher potential evapotranspiration usually results in smaller Q_{\min} (median $r_s=-0.44$). This indicates that the magnitudes of both precipitation and E_p anomalies affect low-flow magnitudes (p-values < 0.001 according to the sign test), but with substantial site-to-site variability. The r_s between 30-day climate anomalies and Q_{\min} does not show distinct spatial patterns across Switzerland (see supplementary Fig.S3). The r_s between the 30-day precipitation anomaly and Q_{\min} is not correlated with mean catchment elevation ($R^2 = 0.08$), and the r_s between the 30-day E_p anomaly and Q_{\min} is weakly correlated with mean catchment elevation ($R^2 = 0.33$).

3.3. Combined effects of climate anomalies on warm-season low flows

The results shown in Fig. 2 indicate that both P and E_p can affect low flows. However, most low flows are not caused by only one driver, but instead result from the combined effects of below-average P and above-average E_p during the same time period. Warm-season low flows usually follow periods of below-average precipitation and above-average potential evapotranspiration (72.2% of low flows fall in the top left quadrant of Fig. 3a). Less than a quarter of the annual low flows occur after periods of below-average precipitation and below-average potential evapotranspiration (20.5% lower left quadrant – Fig. 3a). Only very few annual low flows (7.3%) occur after periods of above-average precipitation. Thus, precipitation anomalies appear to be the most important driver for warm-season low flows in Switzerland, and potentially also in other regions with distinct warm-season low flows. While potential evapotranspiration appears to be less important than precipitation, more than 70% of low flows are caused by a combination of both drivers. The combined effect of above-average E_p thus more than triples the chance of an annual low flow (compared to when precipitation is below average, but there is below-average E_p).

In particular, the most severe low flows occur through the combined effects of low precipitation and high potential evapotranspiration. For example, 96% of low flows during the most severe low-flow year (2003, shown by green markers in Fig. 3a) follow periods of both below-average precipitation and above-average potential evapotranspiration. This behavior is not unique to the 2003 event, but was also observed for other years with severe annual low flows such as 2011, 2015 and 2018 (Fig. 3b&c).




Figure 3: Anomalies in precipitation and potential evapotranspiration 30 days prior to each annual warm-season (May through November) low-flow period in each catchment (grey dots); annual cold-season low flows were excluded (a). The most severe low-flow year during the study (2003) is highlighted in green. Almost all (92.7%) annual low flows occurred following below-average precipitation (the left half of the figure), and 72.2% of all low flows occurred following a combination of below-average precipitation and above-average potential evapotranspiration (the upper left quadrant of the figure). Boxplots of warm-season 7-day minimum flows for the Swiss study catchments (b) and the catchment distribution of the signs of precipitation and evapotranspiration anomalies that preceded these low flows (c). The most severe low-flow years (2003, 2011, 2015, and 2018) were characterized by negative precipitation anomalies and positive E_p anomalies for the large majority of catchments, as indicated by the light grey bars in (c).

3.4. Duration of climate anomalies

The magnitudes of low flows are also related to the durations of the preceding precipitation and evapotranspiration anomalies. Longer periods of below-threshold P and above-threshold E_p tend to lead to lower low flows in most of our catchments (Fig. 4). The duration of high E_p is more strongly correlated with low-flow magnitudes than the duration of low precipitation is (mean Spearman correlations r_s of -0.27 and -0.11 respectively; median r_s values differ from 0 at $p < 0.001$ by sign test; Fig. 4). The weaker correlation with the duration of below-threshold precipitation probably arises because precipitation is more erratic through the years than E_p . A single precipitation event may exceed the precipitation threshold (according to the criterion outlined in Sect. 2.3), but be insufficient to end the low flow in the stream. Low-flow magnitudes are less strongly correlated with the duration of below-threshold precipitation than with the intensity of 30-day precipitation anomalies (compare Fig. 4 with Fig. 1; mean r_s of -0.11 and 0.26, respectively). Similarly, low-flow magnitudes are less strongly correlated with the duration of above-threshold E_p than with the intensity of 30-day E_p anomalies (compare Fig. 4 with Fig. 1; mean r_s of -0.27 and -0.41, respectively).




Figure 4: Histograms of rank correlations between the magnitudes of warm-season low flows and the lengths of the preceding intervals with below-threshold precipitation (a) or above-threshold E_p (b). Longer periods of high E_p are associated with lower low flows, whereas a weaker association is seen between lower low flows and longer periods with low precipitation.

Summing P and E_p anomalies over time windows ranging from one week to half a year indicates that most low flows can be well explained by anomalies of up to 60 days (Fig. 5h). This is because in the typical Swiss climate, precipitation and E_p anomalies usually last for 60 days or less. This is depicted by the grey cloud of points in Fig. 5, as well as the mean anomalies (indicated by the dotted lines in Figs. 5a-g) which remain approximately stable for periods exceeding 60 days. Thus, while longer precipitation and E_p anomalies would lead to lower flows, most low flows result from anomalies of up to 60 days. This is because most anomalies peak at around that 60-day time scale, which is also indicated by the means of the precipitation and E_p anomalies as functions of timescale (dashed lines in Figs. 5h and 5i).

The severe low flows in 2003, 2011, 2015 and 2018, however, are associated with P and E_p anomalies that grow for much longer, and thus become much larger, than the roughly 60-day anomalies that are typical in this climate (colored symbols in Fig. 5). Long periods of above-average E_p appear to be an important factor for these severe low flows; the colored points in Figs. 5e-g expand more on the y-axis than the x-axis for timescales >60 days. Thus, severe low flows result from longer-lasting (and thus larger) P and E_p anomalies, whereas more typical low flows result from climate anomalies that end after roughly 60 days, as illustrated by Figs. 5h&i.





Figure 5: Cumulative anomalies of precipitation and potential evapotranspiration over 7, 14, 30, 60, 90, 120 and 182 days prior to every annual warm-season low flow in each catchment (a-g), and the evolution of the mean anomalies over the different time windows (h & i). Each grey dot represents the combination of precipitation and E_p anomalies before one low-flow event at one site. Low-flow anomalies in the most severe low-flow years are indicated by different colors (2003 in green, 2011 in yellow, 2015 in cyan and 2018 in orange). The dotted lines indicate the mean precipitation and E_p anomalies. The mean anomalies (dotted lines in all panels) clearly increase within the first 60 days prior to low flows, but show no clear trend over longer time windows. During the most severe low-flow years, however, the mean anomalies continue to increase across all of the time windows examined here. In particular, the E_p anomalies during the severe low-flow years grow well beyond the range that is observed during more typical years.

3.5. The relative importance of P and E_p anomalies for warm-season low-flow magnitudes

We further assessed the relative importance of each of the climate drivers and their duration in predicting the magnitude of annual low flows by calculating the bivariate Spearman rank correlation between each climate driver and Q_{\min} as one value for all stations and years together (Fig. 6). The results also include the site-to-site variability in Q_{\min} ; thus the overall r_s correlations are weaker than those shown in Fig. 1c&d. Typical low flows across all years of the observation period (2000-2018) are more

330 strongly correlated to precipitation anomalies than to E_p anomalies (see also Fig. 1), and this correlation becomes slightly stronger at longer durations. However, during the driest years of our dataset (2003, 2011, 2015 and 2018), the correlation between precipitation anomalies and Q_{\min} drops to roughly zero, suggesting that under these extreme conditions low precipitation alone cannot explain the variation in annual low-flow magnitudes. Instead, in these dry years, E_p anomalies retain their predictive power for Q_{\min} , suggesting a relatively more important role of E_p in dry years.


335



340 **Figure 6: Bivariate Spearman rank correlation coefficients between precipitation (blue) and E_p anomalies (red) and Q_{\min} of warm-season (May to November) low flows for durations of 30 to 182 days, across all stations and years. The overall explanatory power of the climate anomalies in a bivariate regression framework is low, although precipitation anomalies are slightly better correlated to Q_{\min} than E_p anomalies in the whole dataset (a). In the four driest years (b) the overall explanatory power of precipitation anomalies is much smaller, and the explanatory power of E_p anomalies is slightly greater, than in all years combined.**

To quantify how much of the maximum predictive power lies in individual anomalies, we first used a multivariate stepwise generalized linear model (GLM) to predict Q_{\min} as a function of all precipitation and E_p anomalies for all durations of 30, 60, 90, 120 and 182 days. In Fig. 7 we show the fraction of the model R^2 explained by individual P and E_p anomalies for the different durations. Across all stations and years of the observation period (2000-2018), warm season Q_{\min} is best predicted by precipitation anomalies with increasing duration (Fig. 7a), which shows the cumulative effect of low precipitation. However, in the years with the lowest annual warm season low flows (2003, 2011, 2015 and 2018) the picture reverses, and instead E_p explains most of the variability in Q_{\min} . This is true across a wider range of durations, starting even at 30 days. Thus, although precipitation anomalies are a good predictor for typical low flows, low-flow magnitudes in the driest years are more strongly related to E_p anomalies when precipitation is also very low.

350



355 **Figure 7: The fraction of multivariate R^2 (calculated by a stepwise generalized linear regression model that explains warm-season low-flow magnitudes using all climate variables and durations) that can be explained by a precipitation (blue) or E_p (red) anomaly of the specified duration. Precipitation anomalies explain most of the variation in Q_{\min} when looking at all stations and all years (a). However, precipitation anomalies are not good predictors for low flows that occurred in the driest years (2003, 2011, 2015, 2018), when E_p anomalies are instead much better predictors of Q_{\min} variability (b).**

360

3.6. The impact of winter precipitation and snow on warm-season low flows

Previous studies indicate that winter snowpack and snowfall can influence the timing and magnitude of summer low flows in some regions (e.g., Dierauer et al., 2018; Jenicek et al., 2016; Godsey et al., 2014). If this holds true for our study catchments, more winter precipitation (December through March), or higher SWE on March 1st, should lead to larger and later warm-season low flows. To test for this effect, we calculated Spearman rank correlations between winter precipitation totals and subsequent warm-season low-flow magnitudes and timing. The correlations between winter precipitation and the magnitude and timing of Q_{\min} (mean absolute $r_s < 0.11$ for both; grey bars in Fig. 8) are weaker than those between low-flow magnitudes and climate anomalies directly preceding low flows (Figs. 1c&d), and they do not vary systematically with elevation. We also calculated the Spearman rank correlations between March 1st SWE and subsequent low-flow magnitudes and timing, and found no strong relationship (Fig. 8, green bars; mean absolute $r_s < 0.17$ for both).

370




Figure 8: Histograms of the rank correlations between winter precipitation (December through March; grey bars) and March 1st snow water equivalent (SWE; green bars) and the magnitude (a) and timing (b) of warm-season low flows. Winter precipitation is weakly associated with higher, and later, warm-season low flows, as indicated by the positive r_s for the majority of catchments; however, overall correlations are weak, with considerable site-to-site variability.

4. Discussion

4.1 Climate anomalies control low-flow timing and magnitude

Anomalies of precipitation and potential evapotranspiration affect the magnitude of low flows, but their influence decreases with elevation (Fig. 2a&b). This pattern is probably not unique to Switzerland, and we expect precipitation and E_p anomalies to also be relatively unimportant in other cold regions where low flows primarily occur in winter (e.g., Dierauer et al., 2018; Laaha & Blöschl, 2006; Van Loon et al., 2015; Wang, 2019), driven by extended freezing periods. However, warm-season low flows are more common globally (e.g., Dettinger & Diaz, 2000; Eisner et al., 2017), suggesting that summer climate anomalies are likely to be important not only for the lower-elevation catchments in Switzerland, but also across many other regions of the world.

We found that the combined effects of P and E_p anomalies shape the occurrence and magnitude of low flows, with the more extreme low flows being driven by longer-duration anomalies. Typical warm-season low flows result from climate anomalies of up to 60 days (Fig.5). In Switzerland, typical low flows result from relatively short climate anomalies, probably because precipitation does not have a strong seasonal signature. In climates that typically have frequent precipitation events, short periods (e.g., one to two months) with less precipitation than normal will most likely precede the annual low flow. Similarly short E_p deviations from the norm often precede typical annual low flows. In the years with the lowest low flows (2003, 2011, 2015 and 2018), the durations of climate anomalies were significantly longer, and especially the impacts of E_p anomalies were larger (Figure 5). This highlights precipitation and evapotranspiration as combined drivers of severe low flows, consistent with findings in several experimental catchments during the 2003 low-flow year (Teuling et al., 2013). Our results suggest that the

magnitude and duration of these precipitation and E_p anomalies are generally important controls on low flows in a large, diverse sample of mesoscale catchments across Switzerland. These compound effects of E_p and precipitation anomalies might also be important for low flows across larger regions (e.g., Stahl et al., 2010), as the climate conditions in Switzerland are comparable to those in other densely populated regions of the world. However, we only analyze these processes on timescales of up to half a year, so long-term memory effects in low-flow generation may not be fully captured by this approach.

The pronounced effect of E_p in the years with the lowest low flows might also reflect the coupling of P and E_p during dry and warm periods. Low precipitation and high air temperature might lead to soil moisture depletion, forcing plants to reduce transpiration. Lower latent heat fluxes and greater sensible heat fluxes from the surface increase air temperature and thus increase E_p while reducing actual evapotranspiration. This complementary relationship between evapotranspiration and E_p can amplify the apparent effect of E_p during (extended) dry periods (Granger, 1989). Conversely, in locations where transpiration is not limited by water availability (e.g., at higher elevations), high temperatures and larger vapor pressure deficits (i.e., high E_p) may drive increases in transpiration rates, accelerating the depletion of catchment water stores and thereby reducing runoff. For example, Mastrotheodoros et al. (2020) showed how increased evapotranspiration at higher elevations systematically amplified runoff deficits during severe low flows in 2003 across the European Alps. These processes are especially relevant in view of potential future climatic changes. In Switzerland, climate change is expected to increase temperatures by more than the global average, resulting in warmer summers with less warm-season precipitation (CH2018, 2018). Similar trends are also expected in many other regions. This highlights the effects of water removal through evapotranspiration, especially during extended dry periods, which are expected to become more severe with changing climate conditions.

A small fraction of all warm-season low flows in the period 2000 to 2018 followed periods of above-average precipitation and below-average E_p (4% in lower right quadrant of Fig. 3a). These anomalies are expected to lead to above-average flow conditions, but can nonetheless lead to annual low flows for at least two reasons. First, these low flows occur in years that are relatively wet, with relatively high annual low flows (Fig. 3b). Second, flow conditions in most Swiss catchments are highly seasonal (Wehren et al., 2010; Weingartner & Aschwanden, 1992), meaning that the seasonality of the flow regime can, in some years, outweigh the effects of shorter-term weather.

4.2 The influence of winter precipitation and snow on warm-season low flows


Previous work in several Swiss catchments has suggested that the snow-water equivalent (SWE) accumulated in the winter snowpack strongly affects summer low-flow magnitudes (Jenicek et al., 2016). Our more complete dataset of Swiss catchments indicates that winter precipitation and SWE (on March 1st) are only weakly related to the magnitude and timing of the following warm-season low flows. In addition, these weak correlations did not significantly increase at higher-elevation catchments, suggesting that even at the higher-elevation sites, SWE is not a major control on warm-season low flows. We caution, however, that this analysis excludes many of the highest-elevation catchments, in which the annual low flow occurs during the winter. Thus the discrepancy between our results and those of Jenicek et al., 2016 probably arises from differences between our respective definitions of low flows. We studied annual 7-day minima, and included only the annual low flows that occur between May and November (thus excluding many high-elevation sites where annual low flows occur in the winter instead), whereas Jenicek et al. (2016) studied 7-day summer minima regardless of whether they are annual minima. Thus, winter precipitation and SWE do affect summer streamflow in Alpine catchments (Jenicek et al., 2016), but our results suggest that for most of the rest of Switzerland, projected changes in winter snowpacks (e.g., Harpold et al., 2017; Mote et al., 2018) might only slightly affect the magnitude and timing of annual low flows that occur during the warm season.

440 **4.3 Human impacts on warm-season low-flow statistics**

Almost every catchment in Switzerland, and elsewhere where dense gauging data exist, is to some extent affected by human activity (e.g., Grill et al., 2019, Lehner et al., 2011). This could be through, for example, water management operations, water abstractions, hydropower operations, and wastewater treatment plant return flows. Especially in Central Europe, almost no pristine catchments exist and quantitative information capturing all potential human influences on streamflow at catchment scale is unavailable. As described in the methods, we removed any catchments with any obvious anthropogenic influences on streamflow (e.g., from hydropowering or dams), however some regulation effects may still be present in the dataset.

To assess the impact of human influence on the results, we recalculated Fig. 2c & d for the 20% of catchments with the largest fraction of human-affected land use, and the 20% of catchments with the smallest fraction of human-affected land use. We thereby tested whether the relationships between the 30-day anomalies of precipitation and E_p and the magnitude of warm-season Q_{min} are significantly different in catchments with a lot of human activity compared to catchments with relatively little human activity (Fig. 9). The results were broadly similar with no significant differences between the strongly affected and weakly affected catchments ($p > 0.2$ by Student's t-test).

The consistency of the results may be due to the fact that, although human water use during low flows will change their absolute magnitudes (and thus may affect site-to-site differences in low flows, which are not considered here), it may have a smaller effect on their relative magnitudes from year to year at any given site. Thus human influences may not greatly alter the rankings of annual low flows throughout the observation period; drier years are still expected to have lower low flows and wetter years are still expected to have higher low flows, largely independent of human influences. Therefore the Spearman rank correlation coefficient is likely to be a relatively robust index for assessing the effects of climate anomalies on the timing and magnitude of annual low flows. Recent studies across US catchments have also found limited effects of human influence on low flows compared to climate drivers (Ferrazzi et al., 2019; Sadri et al., 2016). Nevertheless, the unexplained variance in our established relationships suggests that human-induced shifts in the Q_{min} ranking may have an effect on low-flow behaviors in some catchments.



465

Figure 9: Histograms of rank correlations between low-flow magnitudes and anomalies of precipitation (a & b) and potential evapotranspiration (c & d) for warm-season low flows across Swiss catchments. The left side (a & c) shows the distributions for the 20% of catchments with the least human impact (blue & red) on top of the distributions for all data (grey). The right side (b & d) shows the distributions for the 20% of catchments with the most human impact (blue & red), again plotted on top of the distributions for all data (grey). The observed distributions of correlations between the 30-day climate anomalies and the magnitudes of low flows are similar in catchments with the most and the least human activity.

470

475

4.4 Broader implications

Our overall results are largely consistent with previously discussed drivers of low flows (e.g., Teuling et al., 2013; Woodhouse et al., 2016; Hannaford, 2015). Our work builds upon past research by studying a large dataset which shows the variability and consistency in low-flow/climate relationships among many catchments. We also quantify the effect of the duration of climate anomalies and analyze the interplay of P and E_p as drivers. Our work thereby emphasizes how both

precipitation and E_p anomalies are important drivers of low flows, especially during severe low flows. This is in line with a growing literature on severe events arising from the interplay of multiple drivers (e.g., Zscheischler et al., 2018). Our study also highlights that the relevant properties of low-flow drivers are multidimensional: their magnitudes, timings, and durations all matter. For example, in a lower-elevation catchment, a precipitation anomaly in spring will not have the same impact as a similar anomaly in autumn. Likewise, periods of above-average E_p will have different implications for streamflow in May than they would in September. Thus, antecedent catchment conditions matter. It is not sufficient to look at climate anomalies alone as drivers of low flows, since they may have different implications at different times of the year. Although our study is based on a network of Swiss catchments, we expect our findings to be more broadly applicable to climatically similar regions as well. We see similar patterns in low-flow seasonality in other regions of the world (e.g., Laaha & Blöschl, 2006; Demirel et al., 2013; Dettinger & Diaz, 2000), suggesting that the effects of climate anomalies in these other regions may also be largely similar. For example, the severe summer low flows in California in recent years have been driven by below-average precipitation magnified by above-average temperatures and thus potential evapotranspiration (Diffenbaugh et al., 2015). Van Loon et al. (2015) and Van Loon & Laaha (2015) reported similar driving mechanisms for low flows in Austria and Norway. Thus, our approach for assessing the effects of multiple dimensions of climate impacts (i.e., timing, duration and magnitude) on low flows could potentially be used to derive insight into low flows in other regions.

5. Conclusions

Annual low flows in Switzerland typically occur in two distinct seasons: in winter at higher elevations due to sub-freezing temperatures, and in summer and autumn at lower elevations, following periods of above-average potential evapotranspiration and below-average precipitation (Figs. 2a&b). The magnitudes of these climate anomalies strongly affect the magnitudes of annual low flows across our network of catchments (Figs. 2c&d). Almost all (about 92%) of our catchments' annual low flows follow periods of unusually low precipitation, and many (about 70%) also follow periods of unusually high potential evapotranspiration (Fig. 3a). Thus, most low flows arise from the combined effects of precipitation and E_p anomalies. Severe low flows, such as in the years 2003, 2011, 2015 and 2018, almost exclusively occurred after anomalies in both precipitation and E_p (Fig. 3a). During these especially dry years, low flows occurred simultaneously across large parts of Europe, but their timing was highly variable across Switzerland (Fig. 1). Longer periods of below-threshold precipitation and above-threshold E_p generally led to lower low flows (Fig. 4). Anomalies preceding low flows typically acted over timescales of up to 60 days, while precipitation and E_p anomalies in unusually dry years (2003, 2011, 2015 and 2018) grew for much longer, and thus became much larger (Fig 5). Long periods of above-average E_p appear to be especially important drivers of the most severe low flows (Fig. 5). Typical low flows were mainly driven by precipitation anomalies; however, the low flows in the driest years (2003, 2011, 2015 and 2018) were more related to E_p anomalies (Fig. 6 & 7). Total winter precipitation (and SWE) affected the magnitude and timing of warm-season low flows (Fig. 8), but was less important than the climate anomalies in the month prior to the low-flow period (Figs. 1c&d). Our results describe how the timing, magnitude and duration of precipitation and E_p anomalies drive warm-season low flows across Switzerland. In combination with seasonal weather forecasts, these results could help in predicting and managing low flows.

Data availability: The data that support the findings of this study are available in the ETH library open-access repository. Discharge time series can be obtained from FOEN (Swiss Federal Office of the Environment) and Swiss Cantonal Authorities; meteorological data can be obtained from MeteoSwiss, geodata from Swisstopo (Swiss Federal Office of Topography). Contact information for these agencies is provided in the Supplementary Material.

Author contribution: MF, WB, PM designed the study; MF performed the analyses and wrote the first draft, all authors discussed the results and edited the manuscript.

Competing interests: The authors declare that they have no competing interests.

525

Acknowledgements: The project was funded by the Swiss Federal Office of the Environment (FOEN). We thank FOEN and the Cantons Aargau, Basel Landschaft, Bern, Geneva, Graubünden, Luzern, Neuchatel, Schaffhausen, Solothurn, St.Gallen, Thurgau, Ticino, Vaud and Zürich for providing discharge data, the Swiss Federal Office of Meteorology (Meteoswiss) for providing gridded climate data and the Swiss Federal Office of Topography (Swisstopo) for providing geodata. We thank Anna Costa for additional analyses during the revision process, as well as the editor Kerstin Stahl, the four anonymous reviewers and Ryan Teuling and his students for their comments that greatly improved the original manuscript.

530

References

- 535 Bradford, M. J. and Heinonen, J. S.: Low Flows, Instream Flow Needs and Fish Ecology in Small Streams, *Can. Water Resour. J. Rev. Can. Ressour. Hydr.*, 33(2), 165–180, doi:10.4296/cwrj3302165, 2008.
- CH2018: CH2018 – Climate Scenarios for Switzerland, Technical Report, National Centre for Climate Services, Zurich. [online] Available from: <https://www.nccs.admin.ch/nccs/en/home/klimawandel-und-auswirkungen/schweizer-klimaszenarien/technical-report.html> (Accessed 12 February 2020), 2018.
- 540 CLC: CLC 2018 — Copernicus Land Monitoring Service, [online] Available from: <https://land.copernicus.eu/pan-european/corine-land-cover/clc2018> (Accessed 12 February 2020), 2018.
- Cooper, M. G., Schaperow, J. R., Cooley, S. W., Alam, S., Smith, L. C. and Lettenmaier, D. P.: Climate Elasticity of Low Flows in the Maritime Western U.S. Mountains, *Water Resour. Res.*, 54(8), 5602–5619, doi:10.1029/2018WR022816, 2018.
- De Stefano, L., Duncan, J., Dinar, S., Stahl, K., Strzepek, K. M. and Wolf, A. T.: Climate change and the institutional resilience of international river basins, *J. Peace Res.*, 49(1), 193–209, doi:10.1177/0022343311427416, 2012.
- 545 Demirel, M. C., Booij, M. J. and Hoekstra, A. Y.: Impacts of climate change on the seasonality of low flows in 134 catchments in the River Rhine basin using an ensemble of bias-corrected regional climate simulations, *Hydrol. Earth Syst. Sci.*, 17(10), 4241–4257, doi:10.5194/hess-17-4241-2013, 2013.
- Déry, S. J., Stahl, K., Moore, R. D., Whitfield, P. H., Menounos, B. and Burford, J. E.: Detection of runoff timing changes in pluvial, nival, and glacial rivers of western Canada, *Water Resour. Res.*, 45(4), doi:10.1029/2008WR006975, 2009.
- 550 Dettinger, M. D. and Diaz, H. F.: Global Characteristics of Stream Flow Seasonality and Variability, *J. Hydrometeorol.*, 1(4), 289–310, doi:10.1175/1525-7541(2000)001<0289:GCOSFS>2.0.CO;2, 2000.
- Dierauer, J. R., Whitfield, P. H. and Allen, D. M.: Climate Controls on Runoff and Low Flows in Mountain Catchments of Western North America, *Water Resour. Res.*, 54(10), 7495–7510, doi:10.1029/2018WR023087, 2018.
- 555 Diffenbaugh, N. S., Swain, D. L. and Touma, D.: Anthropogenic warming has increased drought risk in California, *Proc. Natl. Acad. Sci.*, 112(13), 3931–3936, doi:10.1073/pnas.1422385112, 2015.
- Eisner, S., Flörke, M., Chamorro, A., Daggupati, P., Donnelly, C., Huang, J., Hundecha, Y., Koch, H., Kalugin, A., Krylenko, I., Mishra, V., Piniewski, M., Samaniego, L., Seidou, O., Wallner, M. and Krysanova, V.: An ensemble analysis of climate change impacts on streamflow seasonality across 11 large river basins, *Clim. Change*, 141(3), 401–417, doi:10.1007/s10584-016-1844-5, 2017.
- 560 Fangmann, A. and Haberlandt, U.: Statistical approaches for identification of low-flow drivers: temporal aspects, *Hydrol. Earth Syst. Sci.*, 23(1), 447–463, doi:10.5194/hess-23-447-2019, 2019.
- Ferrazzi, M., Vivian, R. and Botter, G.: Sensitivity of Regulated Streamflow Regimes to Interannual Climate Variability, *Earth's Future*, 7(11), 1206–1219, doi:10.1029/2019EF001250, 2019.
- 565 Fiala, T., Ouarda, T. B. M. J. and Hladný, J.: Evolution of low flows in the Czech Republic, *J. Hydrol.*, 393(3), 206–218, doi:10.1016/j.jhydrol.2010.08.018, 2010.
- Fleig, A. K., Tallaksen, L. M., Hisdal, H. and Demuth, S.: A global evaluation of streamflow drought characteristics, *Hydrol. Earth Syst. Sci.*, 10(4), 535–552, doi:10.5194/hess-10-535-2006, 2006.
- Godsey, S. E., Kirchner, J. W. and Tague, C. L.: Effects of changes in winter snowpacks on summer low flows: case studies in the Sierra Nevada, California, USA, *Hydrol. Process.*, 28(19), 5048–5064, doi:10.1002/hyp.9943, 2014.
- 570 Granger, R. J.: A complementary relationship approach for evaporation from nonsaturated surfaces, *J. Hydrol.*, 111(1–4), 31–38, doi:10.1016/0022-1694(89)90250-3, 1989.
- Griessinger, N., Seibert, J., Magnusson, J. and Jonas, T.: Assessing the benefit of snow data assimilation for runoff modeling in Alpine catchments, *Hydrol. Earth Syst. Sci.*, 20(9), 3895–3905, doi:10.5194/hess-20-3895-2016, 2016.

- 575 Grill, G., Lehner, B., Thieme, M., Geenen, B., Tickner, D., Antonelli, F., Babu, S., Borrelli, P., Cheng, L., Crochetiere, H., Macedo, H. E., Filgueiras, R., Goichot, M., Higgins, J., Hogan, Z., Lip, B., McClain, M. E., Meng, J., Mulligan, M., Nilsson, C., Olden, J. D., Opperman, J. J., Petry, P., Liermann, C. R., Sáenz, L., Salinas-Rodríguez, S., Schelle, P., Schmitt, R. J. P., Snider, J., Tan, F., Tockner, K., Valdujo, P. H., Soesbergen, A. van and Zarfl, C.: Mapping the world's free-flowing rivers, *Nature*, 569(7755), 215–221, doi:10.1038/s41586-019-1111-9, 2019.
- 580 Hannaford, J.: Climate-driven changes in UK river flows: A review of the evidence, *Prog. Phys. Geogr. Earth Environ.*, 39(1), 29–48, doi:10.1177/0309133314536755, 2015.
- Hargreaves, G. H. and Samani, Z. A.: Reference Crop Evapotranspiration from Temperature, *Appl. Eng. Agric.*, 1(2), 96, doi:10.13031/2013.26773, 1985.
- 585 Harpold, A. A., Kaplan, M. L., Klos, P. Z., Link, T., McNamara, J. P., Rajagopal, S., Schumer, R. and Steele, C. M.: Rain or snow: hydrologic processes, observations, prediction, and research needs, *Hydrol. Earth Syst. Sci.*, 21(1), 1–22, doi:10.5194/hess-21-1-2017, 2017.
- Haslinger, K., Koffler, D., Schöner, W. and Laaha, G.: Exploring the link between meteorological drought and streamflow: Effects of climate-catchment interaction, *Water Resour. Res.*, 50(3), 2468–2487, doi:10.1002/2013WR015051, 2014.
- 590 Jaeger, E. B. and Seneviratne, S. I.: Impact of soil moisture–atmosphere coupling on European climate extremes and trends in a regional climate model, *Clim. Dyn.*, 36(9–10), 1919–1939, doi:10.1007/s00382-010-0780-8, 2011.
- Jenicek, M., Seibert, J., Zappa, M., Staudinger, M. and Jonas, T.: Importance of maximum snow accumulation for summer low flows in humid catchments, *Hydrol Earth Syst Sci*, 20(2), 859–874, doi:10.5194/hess-20-859-2016, 2016.
- Laaha, G. and Blöschl, G.: Seasonality indices for regionalizing low flows, *Hydrol. Process.*, 20(18), 3851–3878, doi:10.1002/hyp.6161, 2006.
- 595 Laaha, G., Gauster, T., Tallaksen, L. M., Vidal, J.-P., Stahl, K., Prudhomme, C., Heudorfer, B., Vlnas, R., Ionita, M., Van Lanen, H. A. J., Adler, M.-J., Caillouet, L., Delus, C., Fendekova, M., Gailliez, S., Hannaford, J., Kingston, D., Van Loon, A. F., Mediero, L., Osuch, M., Romanowicz, R., Sauquet, E., Stagge, J. H. and Wong, W. K.: The European 2015 drought from a hydrological perspective, *Hydrol Earth Syst Sci*, 21(6), 3001–3024, doi:10.5194/hess-21-3001-2017, 2017.
- 600 Legates, D. R. and McCabe, G. J.: Evaluating the use of “goodness-of-fit” Measures in hydrologic and hydroclimatic model validation, *Water Resour. Res.*, 35(1), 233–241, doi:10.1029/1998WR900018, 1999.
- Lehner, B., Liermann, C. R., Revenga, C., Vörösmarty, C., Fekete, B., Crouzet, P., Döll, P., Endejan, M., Frenken, K., Magome, J., Nilsson, C., Robertson, J. C., Rödel, R., Sindorf, N. and Wisser, D.: High-resolution mapping of the world's reservoirs and dams for sustainable river-flow management, *Front. Ecol. Environ.*, 9(9), 494–502, doi:10.1890/100125, 2011.
- 605 Magnusson, J., Gustafsson, D., Hüsler, F. and Jonas, T.: Assimilation of point SWE data into a distributed snow cover model comparing two contrasting methods, *Water Resour. Res.*, 50(10), 7816–7835, doi:10.1002/2014WR015302, 2014.
- Mastrotheodoros, T., Pappas, C., Molnar, P., Burlando, P., Manoli, G., Parajka, J., Rigon, R., Szeles, B., Bottazzi, M., Hadjidoukas, P. and Fatichi, S.: More green and less blue water in the Alps during warmer summers, *Nat. Clim. Change*, 10(2), 155–161, doi:10.1038/s41558-019-0676-5, 2020.
- 610 Mote, P. W., Li, S., Lettenmaier, D. P., Xiao, M. and Engel, R.: Dramatic declines in snowpack in the western US, *Npj Clim. Atmospheric Sci.*, 1(1), 2, doi:10.1038/s41612-018-0012-1, 2018.
- Munich Re: NatCatSERVICE - Natural catastrophe statistics online, München; Germany. [online] Available from: <https://natcatservice.munichre.com> (Accessed 29 August 2019), 2019.
- 615 Musselman, K. N., Clark, M. P., Liu, C., Ikeda, K. and Rasmussen, R.: Slower snowmelt in a warmer world, *Nat. Clim. Change*, 7(3), 214–219, doi:10.1038/nclimate3225, 2017.
- Peters, E., Bier, G., van Lanen, H. A. J. and Torfs, P. J. J. F.: Propagation and spatial distribution of drought in a groundwater catchment, *J. Hydrol.*, 321(1–4), 257–275, doi:10.1016/j.jhydrol.2005.08.004, 2006.

- Poff, N. L., Allan, J. D., Bain, M. B., Karr, J. R., Prestegard, K. L., Richter, B. D., Sparks, R. E. and Stromberg, J. C.: The Natural Flow Regime, *BioScience*, 47(11), 769–784, doi:10.2307/1313099, 1997.
- 620 Price, K., Jackson, C. R., Parker, A. J., Reitan, T., Dowd, J. and Cyterski, M.: Effects of watershed land use and geomorphology on stream low flows during severe drought conditions in the southern Blue Ridge Mountains, Georgia and North Carolina, United States, *Water Resour. Res.*, 47(2), doi:10.1029/2010WR009340, 2011.
- Roderick, M. L., Sun, F., Lim, W. H. and Farquhar, G. D.: A general framework for understanding the response of the water cycle to global warming over land and ocean, *Hydrol. Earth Syst. Sci.*, 18(5), 1575–1589, doi:10.5194/hess-18-1575-2014, 625 2014.
- Rolls, R. J., Leigh, C. and Sheldon, F.: Mechanistic effects of low-flow hydrology on riverine ecosystems: ecological principles and consequences of alteration, *Freshw. Sci.*, 31(4), 1163–1186, doi:10.1899/12-002.1, 2012.
- Sadri, S., Kam, J. and Sheffield, J.: Nonstationarity of low flows and their timing in the eastern United States, *Hydrol. Earth Syst. Sci.*, 20(2), 633–649, doi:10.5194/hess-20-633-2016, 2016.
- 630 Seneviratne, S. I., Nicholls, N., Easterling, D., Goodess, C., Kanae, S., Kossin, J., Luo, Y., Marengo, J., McInnes, K., Rahimi, M. and et al.: Changes in Climate Extremes and their Impacts on the Natural Physical Environment. A special report of working groups I and II of the Intergovernmental Panel on Climate Change (IPCC)., IPCC, Cambridge; UK. [online] Available from: <https://www.ipcc.ch/report/managing-the-risks-of-extreme-events-and-disasters-to-advance-climate-change-adaptation/changes-in-climate-extremes-and-their-impacts-on-the-natural-physical-environment/> (Accessed 19 April 2019a), 635 2012.
- Seneviratne, S. I., Lehner, I., Gurtz, J., Teuling, A. J., Lang, H., Moser, U., Grebner, D., Menzel, L., Schroff, K., Vitvar, T. and Zappa, M.: Swiss prealpine Rietholzbach research catchment and lysimeter: 32 year time series and 2003 drought event, *Water Resour. Res.*, 48(6), doi:10.1029/2011WR011749, 2012b.
- Smakhtin, V. U.: Low flow hydrology: a review, *J. Hydrol.*, 240(3–4), 147–186, doi:10.1016/S0022-1694(00)00340-1, 2001.
- 640 Stahl, K., Hisdal, H., Hannaford, J., Tallaksen, L. M., van Lanen, H. A. J., Sauquet, E., Demuth, S., Fendekova, M. and Jódar, J.: Streamflow trends in Europe: evidence from a dataset of near-natural catchments, *Hydrol. Earth Syst. Sci.*, 14(12), 2367–2382, doi:10.5194/hess-14-2367-2010, 2010.
- Stahl, K., Kohn, I., Blauhut, V., Urquijo, J., De Stefano, L., Acácio, V., Dias, S., Stagge, J. H., Tallaksen, L. M., Kampragou, E., Van Loon, A. F., Barker, L. J., Melsen, L. A., Bifulco, C., Musolino, D., de Carli, A., Massarutto, A., 645 Assimacopoulos, D. and Van Lanen, H. A. J.: Impacts of European drought events: insights from an international database of text-based reports, *Nat. Hazards Earth Syst. Sci.*, 16(3), 801–819, doi:10.5194/nhess-16-801-2016, 2016.
- Staudinger, M., Stoelzle, M., Seeger, S., Seibert, J., Weiler, M. and Stahl, K.: Catchment water storage variation with elevation, *Hydrol. Process.*, 31(11), 2000–2015, doi:10.1002/hyp.11158, 2017.
- Stoelzle, M., Stahl, K., Morhard, A. and Weiler, M.: Streamflow sensitivity to drought scenarios in catchments with different 650 geology, *Geophys. Res. Lett.*, 41(17), 6174–6183, doi:10.1002/2014GL061344, 2014.
- Tallaksen, L. M. and Van Lanen, H. A. J.: Hydrological Drought – Processes and Estimation Methods for Streamflow and Groundwater, *Eur. Drought Cent.* [online] Available from: <http://europeandroughtcentre.com/resources/hydrological-drought-1st-edition/> (Accessed 22 April 2019), 2004.
- Teuling, A. J., Van Loon, A. F., Seneviratne, S. I., Lehner, I., Aubinet, M., Heinesch, B., Bernhofer, C., Grünwald, T., 655 Prasse, H. and Spank, U.: Evapotranspiration amplifies European summer drought, *Geophys. Res. Lett.*, 40(10), 2071–2075, doi:10.1002/grl.50495, 2013.
- Tongal, H., Demirel, M. C. and Booij, M. J.: Seasonality of low flows and dominant processes in the Rhine River, *Stoch. Environ. Res. Risk Assess.*, 27(2), 489–503, doi:10.1007/s00477-012-0594-9, 2013.
- Van Lanen, H. A. J., Wanders, N., Tallaksen, L. M. and Van Loon, A. F.: Hydrological drought across the world: impact of 660 climate and physical catchment structure, *Hydrol. Earth Syst. Sci.*, 17(5), 1715–1732, doi:10.5194/hess-17-1715-2013, 2013.

- Van Lanen, H. A. J., Laaha, G., Kingston, D. G., Gauster, T., Ionita, M., Vidal, J.-P., Vlnas, R., Tallaksen, L. M., Stahl, K., Hannaford, J., Delus, C., Fendekova, M., Mediero, L., Prudhomme, C., Rets, E., Romanowicz, R. J., Gailliez, S., Wong, W. K., Adler, M.-J., Blauhut, V., Caillouet, L., Chelcea, S., Frolova, N., Gudmundsson, L., Hanel, M., Haslinger, K., Kireeva, M., Osuch, M., Sauquet, E., Stagge, J. H. and Van Loon, A. F.: Hydrology needed to manage droughts: the 2015 European case, *Hydrol. Process.*, 30(17), 3097–3104, doi:10.1002/hyp.10838, 2016.
- 665 Van Loon, A. F.: Hydrological drought explained, *Wiley Interdiscip. Rev. Water*, 2(4), 359–392, doi:10.1002/wat2.1085, 2015.
- Van Loon, A. F. and Laaha, G.: Hydrological drought severity explained by climate and catchment characteristics, *J. Hydrol.*, 526, 3–14, doi:10.1016/j.jhydrol.2014.10.059, 2015.
- 670 Van Loon, A. F., Ploum, S. W., Parajka, J., Fleig, A. K., Garnier, E., Laaha, G. and Van Lanen, H. A. J.: Hydrological drought types in cold climates: quantitative analysis of causing factors and qualitative survey of impacts, *Hydrol. Earth Syst. Sci.*, 19(4), 1993–2016, doi:10.5194/hess-19-1993-2015, 2015.
- Vidal, J.-P., Martin, E., Franchistéguy, L., Habets, F., Soubeyroux, J.-M., Blanchard, M. and Baillon, M.: Multilevel and multiscale drought reanalysis over France with the Safran-Isba-Modcou hydrometeorological suite, *Hydrol. Earth Syst. Sci.*, 14(3), 459–478, doi:10.5194/hess-14-459-2010, 2010.
- 675 van Vliet, M. T. H., Yearsley, J. R., Ludwig, F., Vögele, S., Lettenmaier, D. P. and Kabat, P.: Vulnerability of US and European electricity supply to climate change, *Nat. Clim. Change*, 2, 676, 2012.
- Wada, Y., Beek, L. P. H. van, Wanders, N. and Bierkens, M. F. P.: Human water consumption intensifies hydrological drought worldwide, *Environ. Res. Lett.*, 8(3), doi:10.1088/1748-9326/8/3/034036, 2013.
- 680 Wang, S.: Freezing Temperature Controls Winter Water Discharge for Cold Region Watershed, *Water Resour. Res.*, 55(12), 10479–10493, doi:10.1029/2019WR026030, 2019.
- Wehren, B., Weingartner, R., Schädler, B. and Viviroli, D.: General characteristics of Alpine waters, in *Alpine waters*, edited by U. Bundi, pp. 17–58, Springer, Berlin., 2010.
- Weingartner, R. and Aschwanden, H.: *Abflussregimes als Grundlage zur Abschätzung von Mittelwerten des Abflusses*, Geographisches Institut der Universität Bern, Bern., 1992.
- 685 WMO: *Manual on Low-flow Estimation and Prediction.*, 2008.
- Woodhouse, C. A., Pederson, G. T., Morino, K., McAfee, S. A. and McCabe, G. J.: Increasing influence of air temperature on upper Colorado River streamflow, *Geophys. Res. Lett.*, 43(5), 2174–2181, doi:10.1002/2015GL067613, 2016.
- Zscheischler, J., Westra, S., van den Hurk, B. J. J. M., Seneviratne, S. I., Ward, P. J., Pitman, A., AghaKouchak, A., Bresch, D. N., Leonard, M., Wahl, T. and Zhang, X.: Future climate risk from compound events, *Nat. Clim. Change*, 8(6), 469–477, doi:10.1038/s41558-018-0156-3, 2018.
- 690