

1 **The influence of water table depth on evapotranspiration in the Amazon arc of deforestation**

2 Authors: John O'Connor¹, Maria J. Santos², Karin T. Rebel¹ and Stefan C. Dekker^{1,3}

3 ¹Copernicus Institute of Sustainable Development, Department Environmental Sciences, Utrecht University, The
4 Netherlands

5 ²University Research Priority Program in Global Change and Biodiversity and Department of Geography,
6 University of Zürich, Switzerland

7 ³Faculty of Management, Science and Technology, Open University, Heerlen, The Netherlands
8

9 *Correspondence to:* John C. O Connor (j.c.oconnor@uu.nl)

10

11 Abstract:

12

13 The Amazon rainforest evapotranspiration (ET) flux provides climate regulating and moisture provisioning
14 ecosystem services through a moisture recycling system. The dense complex canopy and deep root system
15 creates an optimum structure to provide large ET fluxes to the atmosphere forming the source for precipitation.
16 Extensive land use and land cover change (LULCC) from forest to agriculture in the arc of deforestation breaks
17 this moisture recycling system. Crops such as soybean are planted in large homogeneous monocultures and the
18 maximum rooting depth of these crops is far shallower than forest. This difference in rooting depth is key as
19 forests can access deep soil moisture and show no signs of water stress during the dry season while in contrast
20 crops are highly seasonal with a growing season dependant on rainfall. As access to soil moisture is a limiting
21 factor in vegetation growth, we hypothesised that if crops could access soil moisture they would undergo less
22 water stress and therefore would have higher evapotranspiration rates than crops which could not access soil
23 moisture.

24

25 We combined remote sensing data with modelled groundwater table depth (WTD) to assess whether vegetation
26 in areas with a shallow WTD had higher ET than vegetation in deep WTD areas. We randomly selected areas of
27 forest, savanna and crop with deep and shallow WTD and examined whether they differ on MODIS
28 Evapotranspiration (ET), Land Surface Temperature (LST) and Enhanced Vegetation Index (EVI), from 2001 to
29 2012, annually and during transition periods between the wet and dry season. As expected, we found no
30 differences in ET, LST, and EVI for forest vegetation between deep and shallow WTD, which because of their
31 deep roots could access water and maintain evapotranspiration for moisture recycling during the entire year. We
32 found significantly higher ET and lower LST in shallow WTD crop areas than in deep WTD during the dry
33 season transition, suggesting that crops in deep WTD undergo higher water stress than crops in shallow WTD
34 areas.

35

36 The differences found between crop in deep and shallow WTD, however, are of low significance with regards
37 the moisture recycling system as the difference resulting from conversion of forest to crop has an overwhelming
38 influence (ET in forest is $\approx 2 \text{ mm day}^{-1}$ higher than that in crops) and has the strongest impact on energy balance
39 and ET. However, access to water during the transition between wet and dry seasons may positively influence
40 growing season length in crop areas.

41

42

43 **Introduction**

44
45 The Amazon rainforest has been reduced to 80% of its original size due to deforestation over the past few decades
46 (Davidson et al., 2012). Land use and land cover change (LULCC) from forest and savanna to agricultural land
47 disrupts the Amazonian water cycle due to changes in evapotranspiration, infiltration, and runoff (Fearnside, 1997;
48 Lawrence and Vandecar, 2014). Changes in evapotranspiration result in major changes to the water energy balance,
49 as forest vegetation has high evapotranspiration rates and is replaced with agricultural vegetation with lower
50 evapotranspiration which results in a lower latent heat flux and higher sensible heat flux (Swann et al., 2015). In
51 addition, a decline in evapotranspiration reduces the available atmospheric moisture which can reduce rainfall.

52 Differences in vegetation structure are suggested to be the main drivers affecting the evapotranspiration rates.
53 Three major land cover classes can be identified at the Amazon arc of deforestation; forest, savanna (Brazilian
54 cerrado, here we use savanna to keep terms equal with the land cover classification used) and agriculture. Forest
55 vegetation has the highest total leaf surface area while savanna has a lower leaf surface areas owing to its mixed
56 structure of grasses shrubs and trees with a more open canopy and agricultural vegetation usually have a lower
57 leaf area (Asner et al., 2003; Costa et al., 2007). This difference in leaf area lowers the potential surface area for
58 both interception evaporation and transpiration. In addition, the rooting depth of forest savanna and agricultural
59 vegetation differs greatly (Costa and Foley, 2000). Forest vegetation have deep roots which facilitate access to
60 deep soil moisture maintaining their supply of water necessary for photosynthesis even during the dry season.
61 Therefore, forest evapotranspiration remains high throughout the year, unaffected by periods of low rainfall
62 (Maeda et al., 2017; Staal et al., 2018). While the rooting depth of savanna tree species have been shown to be
63 deep, the savanna landscape also contains more open shallow rooted shrubs and grasses. Following LULCC from
64 forest or savanna to agriculture the new vegetation cover lacks deep roots and therefore no longer accesses deeper
65 soil moisture. Over the past few decades, the developing agricultural industry driven by international demand
66 encouraged extensive LULCC (Brando et al., 2014; Foley et al., 2007; Sampaio et al., 2007) concentrated along
67 the southern and eastern edge of the Amazon in an area known as the arc of deforestation (Costa and Pires, 2010;
68 Malhi et al., 2008). LULCC negatively impacts the ecosystem service provision of the Amazon including highly
69 valuable services such as carbon storage and sequestration and moisture recycling and regulation. However, little
70 is known whether LULCC that occurred in areas with a shallow WTD facilitates access to water and leads to
71 higher vegetation productivity and evapotranspiration compared to areas with a deep WTD. Understanding the
72 effect that LULCC has on evapotranspiration is important as the loss of evapotranspiration impacts both climate
73 and precipitation on local and regional scales.

74
75 Local climate can be impacted by LULCC due to changes in the energy balance as loss of evapotranspiration
76 reduces latent heat and increases sensible heat. Studies in the Amazon have shown that temperatures increase on
77 average 1.4 °C with a max of 7 °C following conversion to crop (Badger and Dirmeyer, 2015). The seasonal impact
78 of LULCC is particularly strong during the dry season as crop evapotranspiration is at its lowest, latent heat flux
79 can be reduced by 78% and the sensible heat flux can increase by 85% relative to forest (Ponte De Souza et al.,
80 2011). The loss of evapotranspiration impacts rainfall both locally and on the continental scale. Evapotranspiration
81 returns water to the atmosphere where it can precipitate again either in situ or be carried further downwind (Eltahir
82 and Bras, 1994). Large forests like the Amazon, because of their density and extent create large evapotranspiration
83 fluxes, leading to underpressure over land and the pressure differences draw moisture towards land (Makarieva

84 and Gorshkov, 2007; Sheil, 2014). As high as 70% of rainfall in the Amazon and southern Brazil is a result of
85 Amazonian evapotranspiration (van der Ent and Savenije, 2011). This evapotranspiration precipitation cycle is
86 highly important in both maintaining the forest itself but also providing precipitation to non forested areas. LULCC
87 reduces the evapotranspiration and breaks this moisture recycling system resulting in lower rainfall locally and
88 downwind. The seasonal loss of evapotranspiration in crop areas during the dry season is of great significance,
89 evidence already suggests that LULCC has resulted in a lengthening of the dry season (Costa and Pires, 2010;
90 Debortoli et al., 2017). Model simulations predict that if deforestation continues by 2050 the loss of
91 evapotranspiration will result in a negative effect further reducing forest cover and evapotranspiration (Foley et
92 al., 2007; Spracklen et al., 2012). The conversion of forest and savanna to agricultural land in Brazil is driven by
93 an increasing demand for agricultural production which has almost doubled since 2000 (Zalles et al., 2019);
94 however, losses in evapotranspiration could lead to subsequent losses in agricultural productivity as rainfall is
95 reduced and the growing season is shortened (Oliveira et al., 2013).

96

97 [Crops in the Amazon arc of deforestation are predominantly rainfed and as such impacted by the high seasonality](#)
98 [in rainfall unseen in forest vegetation](#). Forest vegetation provides an optimum structure for evapotranspiration due
99 to its tall complex, dense canopy and deep root systems which can access deep soil moisture stores and maintain
100 high transpiration rates even during periods of low rainfall (Nepstad et al., 1994; Sheil, 2014). [Savanna has a mixed](#)
101 [composition, with both trees and grass layers, more open canopy and lower leaf area. Savanna trees can have a](#)
102 [deep rooting depth \(> 10 m\) facilitating access to deep soil water](#) (Canadell et al., 1996). Agricultural crops, are
103 known to contribute much less to evapotranspiration as a result of their shorter canopy and simpler structure
104 (Fearnside, 1997). In addition, agricultural crops lack the deep root systems of forest which are credited for
105 maintaining evapotranspiration throughout the dry season (Nepstad et al., 1994). In theory, if vegetation continues
106 to access the water table within the root zone then this vegetation will continue to transpire during periods of
107 reduced rainfall. Thus access to soil moisture is an important limiting factor for photosynthesis and transpiration.
108 Shallow water table depths across South America are widely distributed and correspond to an area of
109 approximately 36% of the Amazon (Fan and Miguez-Macho, 2010). We hypothesize that areas of shallow water
110 table depth (WTD) allow shallow rooted vegetation to access soil moisture, potentially facilitating vegetation
111 productivity and higher evapotranspiration when compared to areas of deep WTD. Experimental manipulation of
112 WTD using sub irrigation systems of soybean demonstrated that shallow WTD benefitted productivity and
113 increased yield (Kahlow et al., 2005; Mejia et al., 2000). [In the Amazonian arc of deforestation, irrigation of](#)
114 [crops is relatively uncommon](#) (Lathuilière et al., 2012) and [increases in agricultural productivity have been](#)
115 [achieved primarily by increasing the area of crops](#) (Oliveira et al., 2013). [If agricultural vegetation can access soil](#)
116 [moisture in these shallow WTD areas it could potentially increase the growing season length and productivity](#)
117 [without the need for investment in irrigation systems. In turn, less land would be required to achieve the same](#)
118 [agricultural output](#). During the wet season, soybean can reach rates of evapotranspiration similar to that of forest
119 (Costa and Foley, 2000). Some studies have suggested that the difference in annual ET between forest and
120 agricultural crops is primarily due to access to water during the dry season (Costa et al., 2007).

121

122 In this study, [we use a number of freely available remote sensing products in combination with modelled water](#)
123 [table depth](#) to investigate if naturally occurring shallow water table depth could increase evapotranspiration

124 compared to deep water table depth. We expect the greatest influence to be seen in crop areas as they have the
125 shallowest rooting depth and are most dependent on precipitation. As reported in other studies the influence of
126 WTD should not be visible for deep rooted vegetation (Nepstad et al., 1994) like forest and some savanna species.
127 As savanna has mixed vegetation and rooting depths, we expect that to find some differences in ET as a result of
128 deep and shallow WTD. We expect that the differences as a result of WTD will be greater in the transition periods
129 between wet and dry seasons as rainfall as a water source is limited. In areas of shallow WTD, the saturated zone
130 is closer to the root zone of vegetation. In these locations we, therefore, expect crop vegetation to be buffered
131 against the reduction in rainfall during the dry season transition and experience drought conditions later, thus
132 delaying the decline of transpiration due to the dry season. Similarly, during the wet season transition (WST), we
133 expect that areas of shallow WTD will have higher productivity as crop vegetation may access the shallow WTD
134 to supplement their demand when rainfall is low, therefore growing sooner than areas with deep WTD, effectively
135 shortening the dry season. Finally, we discuss whether differences found in ET between deep and shallow WTD
136 are important for moisture recycling, vegetation productivity and what are the implications for future LULCC.

137 **2 Methods**

138 **2.1 Study Area**

139 The study area is located in the southern Amazon, mostly in the northern region of Mato Grosso and incorporating
140 the border area with Pará (Figure 1). Mato Grosso is classified into three major biomes with rainforest in the North,
141 cerrado (a vegetation type that resembles savanna) in the central region and wetlands in the southwest (Kastens et
142 al., 2017; Lathuillière et al., 2012). The climate has two seasons, the wet season in the austral winter and the dry
143 season in austral summer, the dry season lasts around 5 months with an annual average rainfall of 2000 mm and
144 monthly mean temperatures between 22 - 26 °C (Arvor et al., 2014). This precipitation level is within the natural
145 range supporting both savanna (700 to 2000 mm/year), and forest (1000-2500 mm/year). Mean elevation over the
146 study area is 345 m ± 100 m with a max of 700 m and a min of 100 m. Runoff in the Amazon basin is usually low
147 with groundwater convergence accounting for as high as 90% of streamflow (Miguez-Macho and Fan, 2012).
148 Mean WTD of the study area is 12 m with approximately 20 % shallow (< 2 m). The max WTD is 60 - 70 m. This
149 region is well-known as a dynamic agricultural frontier – the arc of deforestation – with high rates of LULCC,
150 where forest and savanna are converted for extensive agriculture, mostly cattle ranching and soy production
151 (Kastens et al., 2017). Mato Grosso is the leading producer of agricultural crops such as soybean in Brazil (Gusso
152 et al., 2014). We chose a 750 km x 750 km study area which is centrally located in the arc of deforestation and has
153 large areas of primary forest (73 %), savanna (19 %) and crops (3 %).



154
155
156 **Figure 1: Study area on the arc of deforestation the Amazon, in Northern Mato Grosso. Inlayed image shows MODIS**
157 **land cover classification map (2001) for the three land cover classes analysed. Forest – Green, Savanna – Beige, Crop –**
158 **Yellow and Other - Grey. Due to the sinusoidal projection of MODIS satellite data, the study area looks distorted.**

159

160 2.2 Datasets

161 2.2.1 Remote sensing data

162 Remote sensing offers excellent tools for monitoring changes in vegetation over large regions as it provides full
163 geographic coverage, high temporal frequency at spatial scales relevant to most Earth system processes (Chambers
164 et al., 2007). Here we use three separate products from the Moderate Resolution Imaging Spectrometer (MODIS),
165 namely MODIS Evapotranspiration (MOD16A2), MODIS Land Surface Temperature (MOD11A2), and MODIS
166 Enhanced Vegetation Index (MOD13A2), to assess the influence of WTD on evapotranspiration. MODIS remote
167 sensing products were used as they offer a moderate spatial resolution and a high temporal resolution which is
168 ideal for examination of seasonal processes. We chose to perform the analysis for the currently available MODIS
169 land cover archive using data from 2001 to 2012. In addition, this period represents a time with high variability of
170 precipitation extremes in which the Amazon experienced droughts, floods and could depict the variability the
171 system experiences (Nobre et al., 2016). Data was downloaded from the NASA data sharing portal
172 (earthdata.nasa.gov). Data was rescaled to 1 km resolution, no additional post-processing was conducted.

173
174 MODIS Evapotranspiration (hereafter ET) data (Mu et al., 2011) provides 8 day accumulated evapotranspiration
175 at 500 m resolution (rescaled to 1 km). The ET dataset is one of the best available datasets due to its relatively
176 high spatial and temporal resolution as such it has been widely used to investigate the effect of land use change on
177 evapotranspiration in the Amazon (Loarie et al., 2011; Neill et al., 2013; Vergopolan and Fisher, 2016). The
178 baseline algorithm to the MODIS ET product is based on the Penman-Monteith equation, and combines parameters
179 such as land cover, leaf area index (LAI), Albedo and Fraction of Photosynthetically Active Radiation (FPAR)
180 directly observed with or modelled from MODIS data, with reanalysis data on Radiation, Air Temperature and
181 Humidity from the Global Modelling and Assimilation Office (Mu et al., 2011). The MODIS ET products were
182 previously tested over the Amazon by comparing its outputs with eddy covariance tower data, showing that the
183 product is more accurate over longer temporal scales (monthly timesteps) and larger areas (e.g. drainage basin)
184 (Ruhoff et al., 2013; Velpuri et al., 2013). While MODIS ET product is known to be underperforming at fine
185 temporal resolutions and newer novel methods show promising results at nine flux sites across the Amazon (Xu et
186 al., 2019), we believe that the application of the new method for our question on the influence of WTD and our
187 time series analysis was beyond the scope of this study. This is also the reason why we chose to also analyse the
188 effects of WTD on satellite retrieved EVI and LST. As with these additional products differences might be
189 detectable, and potentially show a signal to the effects of WTD on the water cycle.

190 MODIS Land Surface Temperature (hereafter LST) provides an 8 day mean day time land surface temperature in
191 degrees Kelvin at 1 km resolution. LST data are produced by detection of thermal infrared radiation between 3 –
192 15 μm spread across 15 bands of the thermal sensor on board the MODIS satellite system and temperatures are
193 modelled based on land cover classification with a clear sky accuracy of 1 degree K (Wan, 2014). MODIS LST
194 data was converted to degrees Celsius. Despite low albedo in the Amazon and high net radiation, the strong
195 evapotranspiration results in a net cooling effect (Bonan, 2008). Therefore, areas with lower LST will be observed
196 in areas of higher ET (Eltahir and Bras, 1994).

197 MODIS Enhanced Vegetation Index (hereafter EVI) provides an observation on vegetation greenness at a
198 frequency of 16 days and 500 m resolution (rescaled to 1 km). EVI is a vegetation index that measures greenness
199 as a proxy for productivity (Huete et al., 2002). It was developed to improve upon the normalized difference

200 vegetation index (NDVI), as it is less sensitive to saturation in highly dense canopies as those in the Amazon, and
201 EVI also corrects for canopy background effects and atmospheric aerosol effects (Huete et al., 2002). This MODIS
202 product offers an observation of vegetation productivity as it measures “greenness” and is correlated to
203 photosynthesis/evapotranspiration (Mu et al., 2011). Thus vegetation with adequate access to water near their root
204 zone will have a comparatively higher EVI than vegetation which is water stressed. This higher EVI, in turn, would
205 correspond to areas of higher ET.

206 In addition, we also used the MODIS land cover product for selection of our analysis sites (see below). MODIS
207 land cover (hereafter land cover) provides a classification of global land cover at 1 km resolution, and it is annually
208 updated. For this study, we only used pixels that were classified as the same land cover type during the entire study
209 period 2001 - 2012. The study area chosen provides a sufficient number of representative pixels for random
210 selection of each land cover type. The use of stable land cover classes was necessary to determine and describe
211 the patterns of ET, LST, and EVI over time and assess the effects of WTD on such trends without the confounding
212 effect of land cover change. Further, we used MODIS land cover as it is the same land cover classification map as
213 used for the MODIS ET product (Friedl et al., 2010) to avoid effects of land cover classification errors from
214 different maps.

215 Over the Amazon cloud cover and shadows are an issue, especially in the wet season. Pixels with high cloud cover
216 were excluded from the analysis. The high seasonal difference in cloud cover is clear, at each time step we used a
217 spatial mean of only available pixels, due to our large sample size we still enough pixels for the analysis (see figure
218 SI.10.1). We compared the cloud cover per land cover class, and found no bias or significant differences between
219 deep and shallow areas.

220 Finally, topography might influence the MODIS data in a number of ways. Elevation can influence
221 meteorological forcing (i.e. temperature and vapor pressure) which is used to calculate ET. Topography can also
222 influence water availability on a pixel due to slope and catchment size of the surrounding area, impacting water
223 available to vegetation therefore influencing ET and EVI. Serious errors due to topography are filtered by MODIS
224 quality control dataset and these pixels were excluded from our analysis. We used SRTM (Shuttle Radar
225 Topography Mission) data to examine elevation and calculate the topographic wetness index (an integrated
226 measure of water accumulation) of our studied pixels. No significant differences were found between elevation of
227 deep and shallow WTD areas of forest or savanna. Crop elevation was found to be significantly different between
228 deep and shallow areas for half of our randomisations. However, the difference in mean elevation was only 10 m
229 leading us to believe that this will not have a strong impact on the meteorological forcing data or ET. We found
230 no significant differences in the topographic wetness index between deep and shallow land covers (see figure
231 SI.9.1-SI.9.4).

232 Finally, Tropical Rainfall Measuring Mission (hereafter TRMM) 3B42 provides daily precipitation at a resolution
233 of 0.25° (downloaded from earthdata.nasa.gov). We calculated daily mean rainfall of our study area using the
234 TRMM data which was then applied to an accumulation to calculate the seasonality of rainfall, i.e. start of the dry
235 season and the wet season across the study area and not per pixel (see below for further details).

236 2.2.2 Water table depth

237 Water table depth (WTD) values were extracted from the Fan et al. (2010) equilibrium WTD model of South
238 America at 30 arc seconds (~ 1 km). The model was created as a long term mean water table depth using a


239 combination of literature reported depths and national databases of groundwater table depth most of which are
 240 from drinking water wells from areas of high population. This data is interpolated using a groundwater model
 241 forced by present day climate, terrain, and sea level. We used the output of the model to obtain WTD data, which
 242 was projected to the same sinusoidal projection of the MODIS data. The equilibrium WTD model is intended for
 243 use in dynamic simulations, and although our study is not the intended use of the WTD model, it is the best
 244 currently available. As the WTD model output is in “equilibrium” it gives a better indication of the annual average
 245 WTD compared to interpolated WTD measurements which may be biased depending on when they were recorded.
 246 The authors compared their WTD calculations with values reported in the literature and found good agreement for
 247 shallower WTD; however, the model over estimated deep WTD. We selected two broad WTD classes in order to
 248 further reduce some of the uncertainty around this key parameter: Shallow <2 m and Deep >8 m (and we will refer
 249 to these as such from hereafter). Figure 2. shows a theoretical graphical representation of the difference between
 250 forest (deep rooting depth), savanna (mix rooting depth), and crop (shallow rooting depth) land cover classes.
 251 These depths were selected as they represented rooting depth values for crop and forest vegetation from literature
 252 (Fan et al., 2016; Moreira et al., 2000; Nepstad et al., 1994; Setiyono et al., 2008).

253 2.3 Sampling design

254 2.3.1 Spatial sampling

255 We chose to avoid pixels which experienced LULCC during the study period as we wanted to use the full time
 256 series for each pixel. We used MODIS land cover to identify pixels of each land cover class which remained
 257 unchanged between years and used these for analysis. We combined three land cover classes with the two water
 258 table depths and analysed the following classes: Forest Deep, Forest Shallow, Savanna Deep, Savanna Shallow,
 259 Crop Deep, and Crop Shallow.

260 For each class, we randomly selected 1000 pixels and performed this random selection 20 times to account for the
 261 effect of the randomization process in the results. This random selection method increased computational
 262 efficiency by limiting the number of total pixels examined and producing comparable group sizes for statistical
 263 analysis. During the wet season the number of usable pixels was as low as 200 – 300 pixels per class for some
 264 time steps while in the dry season the number of usable pixels was above 900 (see supplemental information fig
 265 SI.8.1).



267
 268 **Figure 2: Diagram showing that forest (A) root depth can reach until the saturated zone in both shallow (< 2 m) and**
 269 **deep (> 8 m) WTD, savanna (B) has a mixed rooting depth with only tree roots reaching deep WTD and crop (C)**
 270 **vegetation have a low maximum rooting depth (crops having a maximum rooting depth of 2 m and savanna having a**
 271 **maximum rooting depth > 10 m (Canadell et al., 1996). Shallow roots can reach the saturated zone in shallow WTD (<**
 272 **2 m); however, they cannot reach the saturated zone in deep WTD (> 8 m).**

273 2.3.2 Data analysis


274 The Amazon arc of deforestation is located in a region that has two major seasons defined by the difference in
275 rainfall, the wet season from October to March (approximately 1500 mm) and the dry season from April to
276 September (approximately 400 mm). The difference in rainfall can have significant impacts as the area can be
277 prone to both seasonal flooding and droughts. In recent years the Amazon arc of deforestation has undergone an
278 increased frequency of extreme weather events with drought in 2005, 2010 and flooding in 2009, 2012 (Nobre et
279 al., 2016). These extreme climatic conditions can have a large influence on ET, and vegetation distribution as
280 waterlogging of soils can lead to anoxia in the root zone. Due to the selection of only consistently classified pixels
281 the influence of waterlogging can be avoided as over time these areas will fall under different classifications.
282 Investigation into the drivers of these extreme variations and how each land cover class is influenced is however
283 beyond the scope of this study.

284
285 Analysis of the data was conducted using three primary time periods. At first, we compared mean daily values of
286 ET, EVI and LST between deep and shallow WTD as this gives an indication of the influence of WTD on our land
287 cover classes without considering the seasonal variation. Secondly, we compared ET, EVI and LST of our land
288 cover classes during the dry and wet season transition periods.

289
290 For each year we calculated the dry and wet season transitions using mean daily precipitation of our study area
291 from TRMM with the anomalous accumulation method of Liebmann et al., (2007). This method uses the following
292 equation:

$$293 \quad A(n) = \sum_{n=1}^{day} [R(n) - \bar{R}]$$

294 Where $R(n)$ is daily precipitation and \bar{R} is the average daily precipitation. Calculation of the anomalous
295 accumulation begins at the driest month of the year, when the difference between daily precipitation and annual
296 average is summed to a running total of the anomalous accumulation (A). The wet season onset is defined as the
297 beginning of the longest period where the anomalous accumulation remains positive while the dry season onset is
298 defined as the day after this anomalous accumulation reaches its maximum (Figure 3). These onset points of the
299 dry and wet seasons were applied to find the closest time stamp from each MODIS product in the time series. We
300 then considered the dry season transition to last on average 8 repeats in the MODIS record (5 for EVI due to the
301 lower frequency of the product) and the wet season transition 7 repeats (4 for EVI). We used an average value for
302 each remote sensing product over these transition periods to assess the difference between shallow and deep WTD
303 on evapotranspiration.



305

306 **Figure 3** Mean annual precipitation of the study area calculated from TRMM with 7 day average for graphical
 307 smoothing. wet season transition (WST) and dry season transition (DST) periods are represented in lighter blue. The
 308 vertical lines represent the average start and end dates, however exact dates were calculated per year between July 2001
 309 and July 2012. Red line represents anomalous accumulation method A(n) from Liebmann et al. (2007).

310 These seasonal periods were selected as LULCC in the arc of deforestation is correlated with a lengthening of the
 311 dry season in particular delays in the WST (Butt et al., 2011; Dubreuil et al., 2012; Fu et al., 2013). Recently,
 312 evapotranspiration has been shown to draw moist air over the Amazon triggering the wet season before migration
 313 of the ICTZ (Wright et al., 2017). In this study, we focus our analysis on differences in the DST and the WST.
 314 During the DST, there is already significant drydown (anomalous accumulation is at a maximum, and precipitation
 315 already went down before, see figure 3) which should be apparent in vegetation without access to deeper water
 316 sources. Further into the dry season, other factors may cause a decline in transpiration as well, like heat stress.
 317 During the WST, we focus on the recovery of the vegetation, which should be faster when they have access to
 318 deeper water sources, like deep roots or a shallow WTD.. Thus shallow rooted, vegetation in shallow WTD areas
 319 may have higher access to water as their root zone is closer to the water table this will likely produce higher ET,
 320 EVI and lower LST during the dry season transition than shallow rooted vegetation in deep WTD areas. This is
 321 because the WTD is much deeper and further from the vegetation rooting zone, which leads to a lack of access to
 322 water and the vegetation will likely be stressed. Similarly, during the wet season transition, shallow rooted
 323 vegetation in shallow WTD may exhibit higher ET, EVI and lower LST than that in deep WTD because vegetation
 324 cannot yet be sustained by precipitation alone.. We do not expect these differences with deeply rooted vegetation
 325 We tested whether ET, LST, and EVI followed a normal distribution using the Kolmogorov–Smirnov test. This
 326 test served two purposes, to assess whether parametric statistics could be used and also indicate whether the WTD
 327 influences the frequency distribution of ET, LST, and EVI. Since a large number of response variables were not
 328 normally distributed, we chose to use non-parametric methods. Therefore, Wilcoxon rank sum test was used to

329 test whether there was a significant difference in median ET, LST, and EVI due to the deep and shallow water
330 table.

331 We further examined the frequency distribution of deep and shallow WTD of each of the datasets using the
332 methodology of Wilcox (2012) where the lower and upper quantiles of the distribution are compared. Wilcox's
333 method utilises bootstrapping in order to compare the distribution of the 10th and 90th quantile using the Wilcoxon–
334 Mann–Whitney test. Due to our large sample size, 100 bootstrapped datasets were used.

335 Statistical analysis between each deep and shallow land cover pair was performed separately each year for all 20
336 randomisations e.g. differences in forest ET was tested for significance 12 years * 20 randomisations. **For one**
337 **year, the difference in ET, EVI or LST was considered statistically significant** when more than 66.7% of
338 randomisations were significant and an overall significance was determined if the majority (>50 %) of the years
339 were significant. Statistical analysis was performed using Matlab R2018a (The MathWorks Inc., Natick, USA)
340 statistical toolbox and Wilcox (2012) quantile distribution tool.


341 **3. Results**

342 The following results section is split into three subsections, one for each of the MODIS products used in the
343 analysis. Each of the subsections and accompanying figures follows the same structure. Each figure uses three
344 panels for the three time periods on the analysis A) annual daily mean, B) daily mean during DST, C) daily mean
345 during WST. Each panel has three pairs of box plots which represent the deep and shallow WTD data for forest,
346 savanna and crop.

347 **3.1 Effect of ground water depth on Evapotranspiration**

348 None of the three land cover classes had significant differences in the average daily evapotranspiration (ET_{daily})
349 between deep and shallow WTD areas (Figure 4A). However, while we did not find consistent significant
350 differences, in both forest and crop ET_{daily} we do see a trend towards higher ET_{daily} in shallow WTD areas for both
351 (average \pm standard deviation: Forest Deep = 3.953 ± 0.08 mm day⁻¹, Forest Shallow 3.967 ± 0.09 mm day⁻¹; Crop
352 Deep = 1.697 ± 0.07 mm day⁻¹, Crop Shallow = 1.713 ± 0.08 mm day⁻¹). Interestingly, we found significant
353 differences for Savanna at the extremes of the distributions, depicted by the arrows in Figure 4A. Both the 10th and
354 90th quantiles of ET_{daily} were significantly higher in deep WTD areas than in shallow (difference of 10th = 0.017mm
355 day⁻¹, difference of 90th = 0.02 mm day⁻¹, see supplemental information table S.2.4 for all the quantile analyses).

356



357

358 **Figure 4. (A) Average daily evapotranspiration (ET) annually ET_{daily} , (B) during the dry season transition period ET_{DST} ,**
 359 **(C) during the wet season transition ET_{WST} . Red boxes represent deep WTD Blue boxes represent shallow WTD.**
 360 **Significant results are shown by the green filled boxes if significance was found with both Wilcoxon Rank (WR) and**
 361 **Kolmogorov–Smirnov (KS). Significant differences in 10th and 90th quantile are depicted by the arrows.**

362

363 Clear differences in seasonality occur between the different land cover types (see supplemental information figures
 364 SI.3.1, SI.3.2 and SI.3.3). During the wet season mean ET of all land cover types can be above 4 mm day⁻¹. Both
 365 crop and savanna show clear suppression of ET during the dry season.

366

367 Crop ET during the dry season transition (hereafter ET_{DST}) was significantly higher in shallow than deep WTD
 368 areas (average \pm standard deviation ET: Deep = 2.196 ± 0.11 mm day⁻¹, Shallow = 2.26 ± 0.12 mm day⁻¹, see the
 369 green filled boxes in Fig 4B). Again we observed significant differences at the extremes of the distribution for
 370 savanna, on average the 10th quantile of ET_{DST} was higher in shallow (average difference = 0.003 mm day⁻¹) and
 371 on average the 90th quantile of ET_{DST} was higher in shallow (average difference = 0.005 mm day⁻¹).

372 During the wet season transition (WST), while on average ET_{WST} was higher in shallow WTD areas than in deep
 373 WTD areas (average difference: Forest = 0.01 mm day⁻¹; Savanna = 0.01 mm day⁻¹; Crop = 0.06 mm day⁻¹) this
 374 difference was not significant (Figure 4C).


375

376 3.2 Effect of ground water depth in Land Surface Temperature

377 We found that the distribution of the average land surface temperature (LST_{daily}) was significantly different only
 378 for savanna and the 90th quantile of crop. Deep WTD areas of savanna showed a distribution skewed towards lower
 379 temperatures (average \pm standard deviation LST: Deep = 31.705 °C \pm 0.38, Shallow = 31.848 °C \pm 0.37), see yellow
 380 filled boxes in Figure 5A. The 90th quantile of crop LST_{daily} deep WTD areas was on average 0.1 °C higher than in
 381 shallow WTD areas. Although this is only part of the distribution, it indicates that the warmest crop areas are found
 382 in deep WTD.

383

384



385


386 **Figure 5. (A) Average daily land surface temperature (LST) annually LST_{daily} , (B) during the dry season transition**
387 **period LST_{DST} , (C) during the wet season transition LST_{WST} . Red boxes represent deep WTD Blue boxes represent**
388 **shallow WTD. Yellow filled boxes represent a statistical difference in skewness, calculated by Kolmogorov–Smirnov,**
389 **and green filled boxes represent statistical differences by both Wilcoxon-rank and Kolmogorov–Smirnov. Significant**
390 **differences in 10th and 90th quantile are depicted by the arrows.**

391

392 LST shows clear seasonal differences between the different land covers. Crop LST has the highest range in LST
393 with the warmest period coming towards the end of the dry season (August/September). (Supplemental
394 information figure S.5.1, S.5.2 and S.5.3). During the dry season transition, we found that crop in deep WTD areas
395 had a significantly higher LST than in shallow WTD areas (average \pm standard deviation LST: Deep = $31.256 \pm$
396 0.29 °C, Shallow = 30.864 ± 0.31 °C, green filled boxes in Figure 5B). In addition, the 10th quantile of the crop
397 distributions was significantly higher by 0.42 °C in deep WTD areas than in shallow. During these periods we
398 found again a significant difference in the distribution of savanna, where deep savanna distribution was skewed
399 towards lower LST values. No significant differences were found during the wet season transition (Figure 5C).

400 3.3 Effect of ground water depth on Enhanced Vegetation Index

401 We found significant differences in daily average EVI (EVI_{daily}) between deep and shallow only in crop (average
402 \pm standard deviation EVI: Deep = 0.352 ± 0.01 ; Shallow = 0.357 ± 0.01), with shallow WTD areas EVI being
403 higher than that of deep WTD areas (Figure 6A green filled boxes).



405
 406 **Figure 6. (A) Average enhanced vegetation index (EVI) annually EVI_{daily} , (B) during the dry season transition period**
 407 **EVI_{DST} , (C) during the wet season transition EVI_{WST} . Red boxes represent deep WTD Blue boxes represent shallow**
 408 **WTD. Green filled boxes represent statistical differences by both Wilcoxon-rank and Kolmogorov-Smirnov tests.**
 409 **Significant differences in 90th quantile are depicted by the arrows.**

410
 411 Seasonality in EVI is shown in Supplemental information figure S.7.1, S.7.2 and S.7.3. Crop EVI shows the highest
 412 variation between land cover types. When looking at the DST (May/June) of crop EVI it seems that the response
 413 is delayed in shallow WTD compared to deep WTD; for the WST (October/November) it seems that EVI in
 414 shallow areas increases faster than in deep WTD areas.

415 Mean EVI during the dry season transition (EVI_{DST}) for crop showed a trend towards higher EVI in shallow WTD
 416 areas; however, this difference was only significant in 5 of the 11 years and therefore is not considered consistent
 417 enough to be statistically significant (average \pm standard deviation EVI: Deep = 0.352 ± 0.01 , Shallow = $0.3656 \pm$
 418 0.01 . Figure 6B, Table S.6.8). The 90th quantile EVI of crop was significantly higher in shallow WTD areas than
 419 deep. During the wet season transition (EVI_{WST}), crop was the only different class where EVI was significantly
 420 higher in shallow WTD areas than in deep WTD areas (average \pm standard deviation EVI: Deep = 0.364 ± 0.01 ,
 421 Shallow = 0.378 ± 0.02 , green filled boxes in Figure 6C).

422 4 Discussion

423 In this study, we tested the hypothesis that areas of shallow water table depth (WTD) would have higher
 424 evapotranspiration when compared to areas of deep WTD. As crop vegetation has the shallowest roots (< 2 m) we
 425 expect to see the largest influence of WTD. In areas of deep WTD the root zone is far from the saturated zone
 426 resulting in less uptake of deep soil water, while in areas of shallow WTD the root zone is close to the saturated
 427 zone therefore providing the crops access to ground water. However, we found no support for this as the annual
 428 daily mean ET was not different between crop in deep and shallow WTD areas. One potential explanation is that
 429 since crops experience high seasonality, this annual variability may override differences between deep and shallow

430 WTD areas in the daily average values of ET. For example, average crop ET reaches a high of 3.5 mm day⁻¹ in the
431 wet season while the dry season ET reaches a low of 0.4 mm day⁻¹. Interestingly we found significant differences
432 in annual mean LST and EVI for crop. For LST, we found that the upper 90th quantile was 0.11 °C higher in deep
433 than in shallow WTD areas. While this difference is only found in the 90th quantile of the distribution it does
434 indicate that LST in deep WTD areas can reach higher temperatures than shallow WTD areas. In addition, we
435 found that crops in shallow WTD areas had a significantly higher EVI than in deep WTD. Crop EVI in shallow
436 WTD areas is 1.2 % higher than in deep WTD. This provides support to our hypothesis that crop would have
437 higher EVI in shallow WTD compared to deep WTD areas. The maximum rooting depth for most crops in the
438 region is 2 m, in shallow WTD areas this means the root zone is close to the WTD and would have access to water
439 while in deep WTD the roots are far from the saturated zone. This access to water in shallow WTD areas could
440 also lead to higher ET and therefore evaporative cooling explaining the cooler temperatures in the 90th quantile.

441
442 The second part of our hypothesis was that the effect of WTD would be most evident during the transition periods
443 between wet and dry seasons when rainfall is reduced and vegetation activity is limited by access to soil moisture.
444 We found support for this hypothesis during the dry season transition. In the DST, crop ET was significantly higher
445 in shallow WTD areas and crop LST was significantly lower in shallow WTD areas, while in crop EVI we saw a
446 trend towards higher EVI in shallow WTD areas (significant differences were only found in 5 of the 11 years).
447 While the difference in crop ET is not large (0.063 mm day⁻¹, 2.9 % higher in shallow); during the dry season
448 transition, the results are important as they indicate that crops in the shallow WTD areas have a delayed response
449 to lower rainfall and have a relatively longer growing season. Further evidence of this delayed response can be
450 seen in the EVI seasonality graphs (see figure SI.7.3) where the response of shallow crop to the DST seems delayed
451 compared to deep areas. Crop LST further supports our hypothesis as LST in deep WTD areas was 0.39 °C higher
452 than in shallow WTD areas, while no significant effects were found in EVI. Therefore cooler temperatures in
453 shallow WTD areas are expected to be the result of higher evaporative cooling from ET. These relatively low
454 differences in ET as measured with MODIS data might also be due to the ET product itself. The ET model used
455 for MODIS is not optimised for comparison over relatively small spatial extents and short temporal scales (Ruhoff
456 et al., 2013). In addition, the ET model does not take into account soil water storage and ET is based largely on
457 atmospheric forcing and global land cover parameterisation. Therefore the differences we found for the dry season
458 transition may be under estimated in the MODIS ET values.

459
460 Ponte De Souza et al. (2011) highlighted that one of the strongest impacts of LULCC from forest to crop was due
461 the simultaneous 85% increase in sensible heat flux and 78% reduction in latent heat recorded during the dry
462 season. Studies examining the change in LST due to LULCC found that LST increased by 6 °C from forest to crop
463 (Silvério et al., 2015) and 1.5 °C from savanna to crop (Loarie et al., 2011). Further global models estimated an
464 increase of 5 °C during the summer season for the Amazon, due to a shift from forest to grass (Brovkin et al., 2009;
465 Dekker et al., 2010). This increase in temperature could be influenced by WTD and land cover change; in shallow
466 WTD areas this may result in a less severe temperature change while in deep WTD it could lead to a greater change
467 in temperature; however, WTD was not used as input for these modelling studies. Our results show a maximum
468 temperature of 30 °C in forest compared to a maximum temperature of 38 °C in crops.

469

470 We also expected that the influence of WTD would be **most** important during the wet season transition, as in this
471 period rainfall is increasing. In areas of shallow WTD, vegetation **with a root zone close to the water table** may
472 still access water to supplement if rainfall is not sufficient. Therefore, vegetation growth may be accelerated in
473 comparison to areas of deep WTD which rely more directly on precipitation. Crop EVI was significantly higher in
474 shallow than deep WTD by about 3.8 %, and this was the only data for which we found a significant difference.
475 **Looking at the seasonality of EVI (figure SI.7.3) during the WST EVI is increase faster in shallow WTD areas**
476 **than in deep WTD.** EVI measures vegetation greenness and could be an indication of more rapid growth in shallow
477 WTD areas. As EVI data is directly observed and not modelled the differences are solely reliant on differences in
478 reflected radiation. It may be that smaller differences between deep and shallow WTD areas are more easily
479 detectable using this data. Along the arc of deforestation observations of a lengthening dry season since the 1970s,
480 are linked to a delay in the WST (Butt et al., 2011; Fu et al., 2013). This delay correlates with LULCC and the
481 large reduction this has on ET (Debortoli et al., 2017). Although the difference in WTD seen in crops does not
482 have a strong influence on ET when compared to the difference in ET between the land cover classes, evidence of
483 earlier or faster growth due to the shallow WTD could be beneficial on a local scale.

484

485 These results are even more relevant when comparing the effects of WTD in crop and forest. As forest has been
486 shown to maintain ET throughout the seasons (Kunert et al., 2017) as its deep roots access deeper groundwater
487 (Gash and Nobre, 1997; Nepstad et al., 1994), we hypothesised that no change should be observed in ET, LST,
488 and EVI. Indeed, we found no significant differences across the three MODIS products, both annually or during
489 the dry and wet season transitions. While this does not directly support our hypothesis about the role of WTD for
490 shallow rooted vegetation, this does help validate that our approach reflects our knowledge of the system for
491 vegetation with deep roots.

492

493 Savanna is a complex land cover type because its natural structure makes it is challenging to classify with remote
494 sensing data (Gibbes et al., 2010). MODIS classification accuracy of savanna is about 40 %, about half of that of
495 forest and crop (90 % and 80 % respectively) (Friedl et al., 2010). Savanna includes both trees and grasses, which
496 through the deep roots of trees may access moisture directly and facilitate moisture uptake via hydraulic
497 redistribution (Oliveira et al., 2005) and large areas of shallow root grasses without trees would be negatively
498 affected by water stress. A number of the findings for savanna were not in line with our proposed hypothesis. The
499 distribution between shallow and deep LST was significantly different, with deep WTD areas having a skewed
500 distribution towards lower temperatures. In our hypothesis, we expected to find lower temperature where shallow
501 WTD occurs or no differences in temperature. A similar trend was found in ET where the 10th and 90th quantiles
502 of the distribution were significantly higher in deep WTD areas. The difference in ET was very small, **less than 1**
503 **% difference** between deep and shallow WTD areas. **Water logging of soils has been shown to be an important**
504 **factor in determining vegetation distribution .** Although we believe that larger flooding event leading to changes
505 in vegetation composition are removed from our study due to the selection of pixels that during the time series
506 were always classified as one land cover type, shorter periods of water logging may occur in shallow WTD areas.
507 **However, much higher spatial and temporal resolution imagery would be needed to identify this possibility.**

508

509 The differences found for crop support our hypothesis that shallow WTD areas may facilitate water uptake
510 compared with areas of deep WTD during the transition between wet and dry seasons. Previous crop production
511 studies have shown that artificially maintaining a shallow WTD through sub irrigation systems can increase the
512 productivity of crops such as soy (Kahlow et al., 2005; Mejia et al., 2000) but this has not been previously shown
513 in the naturally occurring shallow WTD areas of the arc of deforestation in the Amazon. In deep WTD areas, crop
514 vegetation undergoes more severe water stress compared with shallow WTD further reducing evapotranspiration
515 and its potential impact on the moisture recycling system. At the regional scale, the difference between deep and
516 shallow WTD is not that important. The most significant differences in ET are driven by deforestation and strong
517 annual variations in rainfall. Although not analysed specifically in this study, the remote sensing data clearly shows
518 these distinctions between different land cover classes and high seasonal and inter-annual variability. On a local
519 scale, the difference between deep and shallow WTD on crop may be of great importance. During the dry season
520 transition crop areas in shallow WTD maintained higher ET. This difference may be important for overall
521 productivity as the dry season influence is delayed and as a result, is increasing the growing season length. This
522 could facilitate natural double cropping systems without the need for investment in irrigation which is still an
523 uncommon practice in the Amazon arc of deforestation (Lathuilière et al., 2012). Agricultural intensification is a
524 pathway to increasing the sustainability of agriculture in the arc of deforestation if it prevents or reduces
525 deforestation or facilitates reforestation (Oliveira et al., 2013). If agricultural productivity can be increased by
526 focusing on already cleared shallow WTD areas, areas of deep WTD could be reforested or returned to secondary
527 forest. Reforestation of previously degraded or logged forest has been shown to return to near natural levels of ET
528 within a few years (Davidson et al., 2012; Hölscher et al., 1997). *The patterns seen in crop vegetation may be
529 caused by factors not considered in this paper. Spatially explicit details about specific crops or agricultural
530 practices were not known for the study. Planting of soybean is determined by the WST and can vary between
531 September at October (Gusso et al., 2014). It is possible that the differences seen in shallow WTD could be the
532 result of earlier sowing and double cropping systems. However, it may be that these agricultural management
533 decisions are implemented more often in shallow WTD because of the higher availability of soil water.*

534

535 This study is a first approach into gaining a better understanding on the influence of shallow WTD on shallow
536 rooted vegetation and it heavily relies on models and remote sensing data which are most appropriate for analyses
537 at larger spatial and temporal scales.

538 The results presented here are limited by the inherent uncertainty of the data used, both in the WTD model and the
539 remote sensing data. Although we believe that the WTD model used here is the best currently available, due to
540 limited data availability it was created using data located mostly in the coastal regions of the continent with very
541 few observations from near our study site (Fan and Miguez-Macho, 2010). In this study, the authors note that there
542 is an overestimation of deep WTD areas when validated against literature reported values. We believe that by the
543 use of a conservative definition of deep WTD >8 m the model outputs are appropriate for our purposes. As
544 discussed above, the remote sensing data has obvious limitations but does provide some insights into how depth
545 of the water table at a local scale might affect water transfer and evaporative processes. Nonetheless, the second
546 main source of uncertainty is in the MODIS land cover classification. We chose to use this land cover classification
547 as the ET and LST products use this classification in their algorithm. Although the classes used are broad and do
548 not reflect the full complexity and heterogeneity of the arc of deforestation, they are robust enough for our

549 purposes. As the influence of WTD on ET is most relevant on smaller scales, further research in these areas could
550 focus on the smaller spatial scales and validate sites with accurate classification and WTD measurements.

551

552 **5 Conclusion**

553 This study aimed to investigate if naturally occurring shallow water table depth supported higher ET compared to
554 deep WTD. In particular if shallow rooted crop vegetation would have higher ET due to increased access to soil
555 water in shallow WTD areas as the distance from the root zone to the saturated zone is shorter. Comparison of EVI
556 showed evidence to support this hypothesis as daily mean EVI was significantly higher in shallow WTD crop
557 areas. However, the difference between deep and shallow WTD is overshadowed by the clear differences between
558 land cover classes. Although not the focus of this study, differences in ET, LST and EVI were largest between
559 land cover classes. In terms of larger scale processes like moisture recycling, LULCC is far more impactful than
560 WTD differences. The main driver of LULCC is agricultural expansion. So although our results are not directly
561 relevant at regional or continental scales on a local scale show signs that shallow WTD areas are more productive
562 than deep WTD.

563 The influence of WTD on crop vegetation was concentrated during the transition periods between wet and dry
564 seasons. We found higher ET and lower LST during the DST and higher EVI during the WST for crop in shallow
565 WTD areas. This higher vegetation productivity of crops due to the shallow WTD help effectively increases the
566 growing season length. The higher productivity in shallow WTD areas may facilitate natural double cropping
567 increasing the agricultural efficiency of the areas. These local scale effects can become significant when scaled to
568 the level of the Amazon. Deforestation rates grew as high as 28,000 km² year⁻¹ in 2004 (Davidson et al., 2012).
569 Any LULCC which occurs in areas of deep WTD are leading to inefficiencies in agricultural production and higher
570 impacts to the moisture recycling system.

571 The results presented here help to demonstrate that the LULCC impacts can vary spatially due to differences in
572 WTD. Future studies investigating the impact of LULCC should incorporate WTD to help disentangle the full
573 impact on the moisture recycling system.

574

575 **References**

- 576 Arvor, D., Dubreuil, V., Ronchail, J., Simões, M. and Funatsu, B. M.: Spatial patterns of rainfall regimes related
577 to levels of double cropping agriculture systems in Mato Grosso (Brazil), *Int. J. Climatol.*, 34(8), 2622–2633,
578 doi:10.1002/joc.3863, 2014.
- 579 Asner, G. P., Scurlock, J. M. O. and A. Hicke, J.: Global synthesis of leaf area index observations: implications
580 for ecological and remote sensing studies, *Glob. Ecol. Biogeogr.*, 12(3), 191–205, doi:10.1046/j.1466-
581 822X.2003.00026.x, 2003.
- 582 Badger, A. M. and Dirmeyer, P. A.: Climate response to Amazon forest replacement by heterogeneous crop cover,
583 *Hydrol. Earth Syst. Sci.*, 19(11), 4547–4557, doi:10.5194/hess-19-4547-2015, 2015.
- 584 Bonan, G. B.: Forests and climate change: forcings, feedbacks, and the climate benefits of forests., *Science*,
585 320(5882), 1444–1449, doi:10.1126/science.1155121, 2008.
- 586 Brando, P. M., Balch, J. K., Nepstad, D. C., Morton, D. C., Putz, F. E., Coe, M. T., Silverio, D., Macedo, M. N.,
587 Davidson, E. A., Nobrega, C. C., Alencar, A. and Soares-Filho, B. S.: Abrupt increases in Amazonian tree
588 mortality due to drought-fire interactions, *Proc. Natl. Acad. Sci.*, 111(17), 6347–6352,
589 doi:10.1073/pnas.1305499111, 2014.
- 590 Brovkin, V., Raddatz, T., Reick, C. H., Claussen, M. and Gayler, V.: Global biogeophysical interactions between
591 forest and climate, *Geophys. Res. Lett.*, 36(7), 1–5, doi:10.1029/2009GL037543, 2009.
- 592 Butt, N., de Oliveira, P. A. and Costa, M. H.: Evidence that deforestation affects the onset of the rainy season in
593 Rondonia, Brazil, *J. Geophys. Res.*, 116(D11), D11120, doi:10.1029/2010JD015174, 2011.
- 594 Canadell, J., Jackson, R., Ehleringer, J., Mooney, H. A., Sala, O. E. and Schulze, E.-D.: Maximum rooting depth
595 of vegetation types at the global scale, *Oecologia*, 108, 583–595, doi:10.1007/BF00329030, 1996.
- 596 Chambers, J. Q., Asner, G. P., Morton, D. C., Anderson, L. O., Saatchi, S. S., Espírito-Santo, F. D. B., Palace, M.
597 and Souza, C.: Regional ecosystem structure and function: ecological insights from remote sensing of tropical
598 forests, *Trends Ecol. Evol.*, 22(8), 414–423, doi:10.1016/j.tree.2007.05.001, 2007.
- 599 Costa, M. H. and Foley, J. A.: Combined effects of deforestation and doubled atmospheric CO₂ concentrations on
600 the climate of Amazonia, *J. Clim.*, 13(1), 18–34, doi:10.1175/1520-0442(2000)013<0018:CEODAD>2.0.CO;2,
601 2000.
- 602 Costa, M. H. and Pires, G. F.: Effects of Amazon and Central Brazil deforestation scenarios on the duration of the
603 dry season in the arc of deforestation, *Int. J. Climatol.*, 30(13), 1970–1979, doi:10.1002/joc.2048, 2010.
- 604 Costa, M. H., Yanagi, S. N. M., Souza, P. J. O. P., Ribeiro, A. and Rocha, E. J. P.: Climate change in Amazonia
605 caused by soybean cropland expansion, as compared to caused by pastureland expansion, *Geophys. Res. Lett.*,
606 34(7), 2–5, doi:10.1029/2007GL029271, 2007.
- 607 Davidson, E. a., de Araújo, A. C., Artaxo, P., Balch, J. K., Brown, I. F., C. Bustamante, M. M., Coe, M. T., DeFries,

- 608 R. S., Keller, M., Longo, M., Munger, J. W., Schroeder, W., Soares-Filho, B. S., Souza, C. M. and Wofsy, S. C.:
609 The Amazon basin in transition, *Nature*, 481(7381), 321–328, doi:10.1038/nature10717, 2012.
- 610 Debortoli, N. S., Dubreuil, V., Hirota, M., Filho, S. R., Lindoso, D. P. and Nabucet, J.: Detecting deforestation
611 impacts in Southern Amazonia rainfall using rain gauges, *Int. J. Climatol.*, 37(6), 2889–2900,
612 doi:10.1002/joc.4886, 2017.
- 613 Dekker, S. C., de Boer, H. J., Brovkin, V., Fraedrich, K., Wassen, M. J. and Rietkerk, M.: Biogeophysical
614 feedbacks trigger shifts in the modelled vegetation-atmosphere system at multiple scales, *Biogeosciences*, 7(4),
615 1237–1245, doi:10.5194/bg-7-1237-2010, 2010.
- 616 Dubreuil, V., Debortoli, N., Funatsu, B., Nédélec, V. and Durieux, L.: Impact of land-cover change in the Southern
617 Amazonia climate: a case study for the region of Alta Floresta, Mato Grosso, Brazil, *Environ. Monit. Assess.*,
618 184(2), 877–891, doi:10.1007/s10661-011-2006-x, 2012.
- 619 Eltahir, E. A. B. and Bras, R. L.: Precipitation recycling in the Amazon basin, *Q. J. R. Meteorol. Soc.*, 120(518),
620 861–880, doi:10.1002/qj.49712051806, 1994a.
- 621 Eltahir, E. A. B. and Bras, R. L.: Sensitivity of regional climate to deforestation in the Amazon basin, *Adv. Water
622 Resour.*, 17(1–2), 101–115, doi:10.1016/0309-1708(94)90027-2, 1994b.
- 623 van der Ent, R. J. and Savenije, H. H. G.: Length and time scales of atmospheric moisture recycling, *Atmos. Chem.
624 Phys.*, 11(5), 1853–1863, doi:10.5194/acp-11-1853-2011, 2011.
- 625 Fan, J., McConkey, B., Wang, H. and Janzen, H.: Root distribution by depth for temperate agricultural crops, *F.
626 Crop. Res.*, 189, 68–74, doi:10.1016/j.fcr.2016.02.013, 2016.
- 627 Fan, Y. and Miguez-Macho, G.: Potential groundwater contribution to Amazon evapotranspiration, *Hydrol. Earth
628 Syst. Sci.*, 14(10), 2039–2056, doi:10.5194/hess-14-2039-2010, 2010.
- 629 Fearnside, P. M.: Environmental services as a strategy for sustainable development in rural Amazonia, *Ecol. Econ.*,
630 20(1), 53–70, doi:10.1016/S0921-8009(96)00066-3, 1997.
- 631 Foley, J. A., Asner, G. P., Costa, M. H., Coe, M. T., DeFries, R., Gibbs, H. K., Howard, E. A., Olson, S., Patz, J.,
632 Ramankutty, N. and Snyder, P.: Amazonia revealed: forest degradation and loss of ecosystem goods and services
633 in the Amazon Basin, *Front. Ecol. Environ.*, 5(1), 25–32, doi:10.1890/1540-9295(2007)5[25:ARFDAL]2.0.CO;2,
634 2007a.
- 635 Foley, J. A., Asner, G. P., Heil Costa, M., Coe, M. T., DeFries, R., Gibbs, H. K., Howard, E. A., Olson, S., Patz,
636 J., Ramankutty, N. and Snyder, P.: Amazonia revealed: forest degradation and loss of ecosystem goods and
637 services in the Amazon Basin, *Front Ecol Environ.*, 5(1), 25–32, doi:http://doi.wiley.com/10.1890/1540-
638 9295(2007)5[25:ARFDAL]2.0.CO;2, 2007b.
- 639 Friedl, M. A., Sulla-Menashe, D., Tan, B., Schneider, A., Ramankutty, N., Sibley, A. and Huang, X.: MODIS
640 Collection 5 global land cover: Algorithm refinements and characterization of new datasets, *Remote Sens.
641 Environ.*, 114(1), 168–182, doi:10.1016/j.rse.2009.08.016, 2010.

- 642 Fu, R., Yin, L., Li, W., Arias, P. A., Dickinson, R. E., Huang, L., Chakraborty, S., Fernandes, K., Liebmann, B.,
643 Fisher, R. and Myneni, R. B.: Increased dry-season length over southern Amazonia in recent decades and its
644 implication for future climate projection, *Proc. Natl. Acad. Sci.*, doi:10.1073/pnas.1302584110, 2013.
- 645 Gash, J. H. C. and Nobre, C. A.: Climatic Effects of Amazonian Deforestation: Some Results from ABRACOS,
646 *Bull. Am. Meteorol. Soc.*, 78(5), 823–830, doi:10.1175/1520-0477(1997)078<0823:CEOADS>2.0.CO;2, 1997.
- 647 Gibbes, C., Adhikari, S., Rostant, L., Southworth, J. and Qiu, Y.: Application of Object Based Classification and
648 High Resolution Satellite Imagery for Savanna Ecosystem Analysis, *Remote Sens.*, 2(12), 2748–2772,
649 doi:10.3390/rs2122748, 2010.
- 650 Gusso, A., Arvor, D., Ricardo Ducati, J., Veronez, M. R. and Da Silveira, L. G.: Assessing the modis crop detection
651 algorithm for soybean crop area mapping and expansion in the Mato Grosso state, Brazil, *Sci. World J.*, 2014(1),
652 doi:10.1155/2014/863141, 2014.
- 653 Hölscher, D., de A. Sá, T. ., Bastos, T. ., Denich, M. and Fölster, H.: Evaporation from young secondary vegetation
654 in eastern Amazonia, *J. Hydrol.*, 193(1–4), 293–305, doi:10.1016/S0022-1694(96)03145-9, 1997.
- 655 Huete, A., Didan, K., Miura, H., Rodriguez, E. P., Gao, X. and Ferreira, L. F.: Overview of the radiometric and
656 biophysical performance of the MODIS vegetation indices, *Remote Sens. Environ.*, 83, 195–213 [online]
657 Available from: [https://ac-els-cdn-com.sire.ub.edu/S0034425702000962/1-s2.0-S0034425702000962-](https://ac-els-cdn-com.sire.ub.edu/S0034425702000962/1-s2.0-S0034425702000962-main.pdf?_tid=420e9dda-1821-11e8-b53d-00000aacb35d&acdnat=1519339356_ef83dc686b96e75a110fbad8d8dc950a)
658 [main.pdf?_tid=420e9dda-1821-11e8-b53d-](https://ac-els-cdn-com.sire.ub.edu/S0034425702000962/1-s2.0-S0034425702000962-main.pdf?_tid=420e9dda-1821-11e8-b53d-00000aacb35d&acdnat=1519339356_ef83dc686b96e75a110fbad8d8dc950a)
659 [00000aacb35d&acdnat=1519339356_ef83dc686b96e75a110fbad8d8dc950a](https://ac-els-cdn-com.sire.ub.edu/S0034425702000962/1-s2.0-S0034425702000962-main.pdf?_tid=420e9dda-1821-11e8-b53d-00000aacb35d&acdnat=1519339356_ef83dc686b96e75a110fbad8d8dc950a), 2002.
- 660 Kahlow, M. A., Ashraf, M. and Zia-UI-Haq: Effect of shallow groundwater table on crop water requirements and
661 crop yields, *Agric. Water Manag.*, 76(1), 24–35, doi:10.1016/j.agwat.2005.01.005, 2005.
- 662 Kastens, J. H., Brown, J. C., Coutinho, A. C., Bishop, C. R. and Esquerdo, J. C. D. M.: Soy moratorium impacts
663 on soybean and deforestation dynamics in Mato Grosso, Brazil, *PLoS One*, 12(4), 1–21,
664 doi:10.1371/journal.pone.0176168, 2017.
- 665 Kunert, N., Maria Aparecido, L. T., Wolff, S., Higuchi, N., dos, J., Carioca de Araujo, A. and Trumbore, S.: A
666 revised hydrological model for the Central Amazon: The importance of emergent canopy trees in the forest water
667 budget, *Agric. For. Meteorol.*, 239, 47–57, doi:10.1016/j.agrformet.2017.03.002, 2017.
- 668 Lathuillière, M. J., Johnson, M. S. and Donner, S. D.: Water use by terrestrial ecosystems: temporal variability in
669 rainforest and agricultural contributions to evapotranspiration in Mato Grosso, Brazil, *Environ. Res. Lett.*, 7(2),
670 024024, doi:10.1088/1748-9326/7/2/024024, 2012.
- 671 Lawrence, D. and Vandecar, K.: Effects of tropical deforestation on climate and agriculture, *Nat. Clim. Chang.*,
672 5(1), 27–36, doi:10.1038/nclimate2430, 2014.
- 673 Liebmann, B., Camargo, S. J., Seth, A., Marengo, J. A., Carvalho, L. M. V., Allured, D., Fu, R. and Vera, C. S.:
674 Onset and end of the rainy season in South America in observations and the ECHAM 4.5 atmospheric general
675 circulation model, *J. Clim.*, 20(10), 2037–2050, doi:10.1175/JCLI4122.1, 2007.
- 676 Loarie, S. R., Lobell, D. B., Asner, G. P., Mu, Q. and Field, C. B.: Direct impacts on local climate of sugar-cane

- 677 expansion in Brazil, *Nat. Clim. Chang.*, 1(2), 105–109, doi:10.1038/nclimate1067, 2011.
- 678 Maeda, E. E., Ma, X., Wagner, F., Kim, H., Oki, T., Eamus, D. and Huete, A.: Evapotranspiration seasonality
679 across the Amazon basin, *Earth Syst. Dyn. Discuss.*, (January), 1–28, doi:10.5194/esd-2016-75, 2017.
- 680 Makarieva, a M. and Gorshkov, V. G.: Biotic pump of atmospheric moisture as driver of the hydrological cycle
681 on land, *Hydrol. Earth Syst. Sci.*, 11(2), 1013–1033, doi:10.5194/hess-11-1013-2007, 2007.
- 682 Malhi, Y., Roberts, J. T., Betts, R. a, Killeen, T. J., Li, W. and Nobre, C. a: Climate Change, Deforestation, and
683 the Fate of the Amazon, *Science (80-.)*, 319(5860), 169–172, doi:10.1126/science.1146961, 2008.
- 684 Mejia, M. N., Madramootoo, C. A. and Broughton, R. S.: Influence of water table management on corn and
685 soybean yields, *Agric. Water Manag.*, 46(1), 73–89, doi:10.1016/S0378-3774(99)00109-2, 2000.
- 686 Miguez-Macho, G. and Fan, Y.: The role of groundwater in the Amazon water cycle: 1. Influence on seasonal
687 streamflow, flooding and wetlands, *J. Geophys. Res. Atmos.*, 117(15), 1–30, doi:10.1029/2012JD017539, 2012.
- 688 Moreira, M. Z., Sternberg, L. D. L. and Nepstad, D. C.: Vertical patterns of soil water uptake by plants in a primary
689 forest and an abandoned pasture in the eastern Amazon: an isotopic approach, *Plant Soil*, 222(1–2), 95–107,
690 doi:10.1023/A:1004773217189, 2000.
- 691 Mu, Q., Zhao, M. and Running, S. W.: Improvements to a MODIS global terrestrial evapotranspiration algorithm,
692 *Remote Sens. Environ.*, 115(8), 1781–1800, doi:10.1016/j.rse.2011.02.019, 2011.
- 693 Neill, C., Coe, M. T., Riskin, S. H., Krusche, A. V., Elsenbeer, H., Macedo, M. N., McHorney, R., Lefebvre, P.,
694 Davidson, E. A., Scheffler, R., Figueira, A. M. A. S., Porder, S. and Deegan, L. A.: Watershed responses to
695 Amazon soya bean cropland expansion and intensification, *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, 368(1619),
696 2–7, doi:10.1098/rstb.2012.0425, 2013.
- 697 Nepstad, D. C., de Carvalho, C. R., Davidson, E. A., Jipp, P. H., Lefebvre, P. A., Negreiros, G. H., da Silva, E. D.,
698 Stone, T. a., Trumbore, S. E. and Vieira, S.: The role of deep roots in the hydrological and carbon cycles of
699 Amazonian forests and pastures, *Nature*, 372(6507), 666–669, doi:10.1038/372666a0, 1994.
- 700 Nobre, C. A., Sampaio, G., Borma, L. S., Castilla-Rubio, J. C., Silva, J. S. and Cardoso, M.: Land-use and climate
701 change risks in the Amazon and the need of a novel sustainable development paradigm, *Proc. Natl. Acad. Sci.*,
702 201605516, doi:10.1073/pnas.1605516113, 2016a.
- 703 Nobre, C. A., Marengo, J. A., Seluchi, M. E., Cuartas, A. and Alves, L. M.: Some Characteristics and Impacts of
704 the Drought and Water Crisis in Southeastern Brazil during 2014 and 2015 Some Characteristics and Impacts of
705 the Drought and Water Crisis in Southeastern Brazil during 2014 and 2015, *J. od Water Resour. Prot.*, 8(August),
706 252–262, doi:10.4236/jwarp.2016.82022, 2016b.
- 707 Oliveira, L. J. C., Costa, M. H., Soares-Filho, B. S. and Coe, M. T.: Large-scale expansion of agriculture in
708 Amazonia may be a no-win scenario, *Environ. Res. Lett.*, 8(2), 024021, doi:10.1088/1748-9326/8/2/024021, 2013.
- 709 Oliveira, R. S., Dawson, T. E., Burgess, S. S. O. and Nepstad, D. C.: Hydraulic redistribution in three Amazonian

- 710 trees, *Oecologia*, 145(3), 354–363, doi:10.1007/s00442-005-0108-2, 2005.
- 711 Ponte De Souza, P. J. D. O., Ribeiro, A., Da Rocha, E. J. P., Botelho, M. D. N., De Sousa, A. M. L., De Souza, E.
712 B. and Bouças Farias, J. R.: Impacts of soybean expansion on the Amazon energy balance: A case study, *Exp.*
713 *Agric.*, 47(03), 553–567, doi:10.1017/S0014479711000391, 2011.
- 714 Ruhoff, A. L., Paz, A. R., Aragao, L. E. O. C., Mu, Q., Malhi, Y., Collischonn, W., Rocha, H. R. and Running, S.
715 W.: Assessment of the MODIS global evapotranspiration algorithm using eddy covariance measurements and
716 hydrological modelling in the Rio Grande basin, *Hydrol. Sci. J.*, 58(8), 1658–1676,
717 doi:10.1080/02626667.2013.837578, 2013.
- 718 Sampaio, G., Nobre, C., Costa, M. H., Satyamurty, P., Soares-Filho, B. S. and Cardoso, M.: Regional climate
719 change over eastern Amazonia caused by pasture and soybean cropland expansion, *Geophys. Res. Lett.*, 34(17),
720 L17709, doi:10.1029/2007GL030612, 2007.
- 721 Setiyono, T. D., Weiss, A., Specht, J. E., Cassman, K. G. and Dobermann, A.: Leaf area index simulation in
722 soybean grown under near-optimal conditions, *F. Crop. Res.*, 108(1), 82–92, doi:10.1016/j.fcr.2008.03.005, 2008.
- 723 Sheil, D.: How plants water our planet: advances and imperatives, *Trends Plant Sci.*, 19(4), 209–211,
724 doi:10.1016/j.tplants.2014.01.002, 2014.
- 725 Silvério, D. V., Brando, P. M., Macedo, M. N., Beck, P. S. A., Bustamante, M. and Coe, M. T.: Agricultural
726 expansion dominates climate changes in southeastern Amazonia: The overlooked non-GHG forcing, *Environ. Res.*
727 *Lett.*, 10(10), doi:10.1088/1748-9326/10/10/104015, 2015.
- 728 Spracklen, D. V., Arnold, S. R. and Taylor, C. M.: Observations of increased tropical rainfall preceded by air
729 passage over forests, *Nature*, 489(7415), 282–285, doi:10.1038/nature11390, 2012.
- 730 Staal, A., Tuinenburg, O. A., Bosmans, J. H. C., Holmgren, M., van Nes, E. H., Scheffer, M., Zemp, D. C. and
731 Dekker, S. C.: Forest-rainfall cascades buffer against drought across the Amazon, *Nat. Clim. Chang.*, 1,
732 doi:10.1038/s41558-018-0177-y, 2018.
- 733 Swann, A. L. S., Longo, M., Knox, R. G., Lee, E. and Moorcroft, P. R.: Future deforestation in the Amazon and
734 consequences for South American climate, *Agric. For. Meteorol.*, 214–215, 12–24,
735 doi:10.1016/j.agrformet.2015.07.006, 2015.
- 736 Velpuri, N. M., Senay, G. B., Singh, R. K., Bohms, S. and Verdin, J. P.: A comprehensive evaluation of two
737 MODIS evapotranspiration products over the conterminous United States: Using point and gridded FLUXNET
738 and water balance ET, *Remote Sens. Environ.*, 139, 35–49, doi:10.1016/j.rse.2013.07.013, 2013.
- 739 Vergopolan, N. and Fisher, J. B.: The impact of deforestation on the hydrological cycle in Amazonia as observed
740 from remote sensing, *Int. J. Remote Sens.*, 37(22), 5412–5430, doi:10.1080/01431161.2016.1232874, 2016.
- 741 Wan, Z.: New refinements and validation of the collection-6 MODIS land-surface temperature/emissivity product,
742 *Remote Sens. Environ.*, 140, 36–45, doi:10.1016/j.rse.2013.08.027, 2014.

- 743 Wilcox, R. R.: Comparing two independent groups via a quantile generalization of the wilcoxon-mann-whitney
744 test, *J. Mod. Appl. Stat. Methods*, 11(2), doi:10.22237/jmasm/1351742460, 2012.
- 745 Wright, J. S., Fu, R., Worden, J. R., Chakraborty, S., Clinton, N. E., Risi, C., Sun, Y. and Yin, L.: Rainforest-
746 initiated wet season onset over the southern Amazon, *Proc. Natl. Acad. Sci.*, 114(32), 8481–8486,
747 doi:10.1073/pnas.1621516114, 2017.
- 748 Xu, D., Agee, E., Wang, J. and Ivanov, V. Y.: Estimation of Evapotranspiration of Amazon Rainforest Using the
749 Maximum Entropy Production Method, *Geophys. Res. Lett.*, 1402–1412, doi:10.1029/2018gl080907, 2019.
- 750 Zalles, V., Hansen, M. C., Potapov, P. V., Stehman, S. V., Tyukavina, A., Pickens, A., Song, X., Adusei, B., Okpa,
751 C., Aguilar, R., John, N. and Chavez, S.: Near doubling of Brazil’s intensive row crop area since 2000
752 [Sustainability Science], *Proc. Natl. Acad. Sci. U. S. A.*, 116(2), doi:10.1073/pnas.1810301115, 2019.

753