

Revised responses to reviewers - Text in blue is added since our online responses

----- REVIEWER 1 -----

We thank Reviewer 1 for reading our paper and for praising it as well-written. It is unfortunate that the reviewer says it lacks “originality and significance”, which we argue against below, and faults us for providing “little or no methodological innovation”.

We first point out that the themes addressed in our paper align with the following guidance from the HESS web site:

HESS encourages and supports fundamental and **applied research that advances the understanding of hydrological systems, their role in providing water for ecosystems and society**, and the role of the water cycle in the functioning of the Earth system. [*emphasis added*]

and

HESS, therefore, aims to serve not only the hydrological science community but all earth and life scientists, **water engineers, and water managers**, who wish to publish original findings on the interactions and feedbacks between the governing processes of the water cycle and processes governing atmospheric circulation and climate, bio-geochemical cycling, dynamics, and **resilience of ecosystems and socio-economy**.

[.....]

3. the study of interactions with human activity of all the processes, budgets, fluxes, and pathways as outlined above, **and the options for influencing them in a sustainable manner, particularly in relation to floods**, droughts, desertification, land degradation, eutrophication, and other aspects of global change. [*emphasis added*]

We next note that applying methods use previously is not given as a criterion for disqualification,

As for the originality of our paper, although there are examples of similar work (cited in our paper), we have not found such a comprehensive study, either for the Columbia Basin or of other large basins, of **changing flood risk** that accounts for and quantifies key sources of uncertainty (see our ANOVA analysis) and, moreover, describes both the changes in magnitude and seasonality of flood risk and how they change as one travels down a river. Multi-GCM, multi-hydrological model, analyses of changing flood risk across a large area are still very rare (we found one – Thober et al., 2018 – but even they examine only the 1-in-2 year event and they don’t explore the hydrological processes that contribute to variability changes in space) If the reviewer is aware of a study that includes the components of our study, we would be grateful to learn of it.

The significance of the paper lies both in its uniqueness and in its generalizability. It is unique in that it provides key numerical input into international treaty negotiations that are currently underway. Many academic papers conclude with a vague admonition to water managers to pay attention to the results. By contrast, most of the authors of this paper have been deeply involved in

developing the key dataset used by the US Entity, and the paper thus has a deep and integral connection to an important policy process. Second, it is generalizable in that we show how complex the pattern of change (with space and with season) can be in a mixed rain-and-snow basin. Basins of similar size and hydrological response to warming exist on most continents, so our results provide a warning about using that simplistic answers about changing flood risk beyond just the Columbia Basin. [We have added some of this response to the text.](#)

References

Tober, S., Kumar, R., Wanders, N., Marx, A., Pan, M., Rakovec, O., Samaniego, L., Sheffield, J., Wood, E.F. and Zink, M., 2018. Multi-model ensemble projections of European river floods and high flows at 1.5, 2, and 3 degrees global warming. *Environmental Research Letters*, 13(1), p. 014003.

----- REVIEWER 2 -----

We are grateful to Reviewer 2 for a particularly thoughtful review and also for responding to Reviewer 1.

On Page 7, l 193, the authors describe the behavior on the Snake as surprising. I would argue rather that it is the upper Columbia that is surprising. The increase in flood ratio along the Snake is consistent with the shift of snow-dominant to transient basins described in Hamlet and Lettenmaier (2010) and so forth. I found the large flood ratios found in the coldest parts of the Columbia surprising, and this is partly where this study diverges from Tohver et al (2014). I'd like to see more explanation of the Upper Columbia; it's hard to see how the effect of temperature alone could explain this, and there must be an increase in the snowpack at the head waters. The effects of temperature and precipitation could potentially be discerned by comparing changes in discharge timing and intensity (Fig 9) – if intensity changes without a change in timing, precipitation changes are likely the cause. Results at TDA suggest a longer less abrupt melt season in the future in contrast to CRNIC.

[We made some changes to the text but a full analysis of the Upper Columbia will require work beyond the scope of this paper.](#)

I'd like to see some comparison in the results section to Tohver et al (2014) rather than leave it for the conclusions. Their fig 5 (flipped on y-axis) is essentially the same as your fig 3. Results are essentially the same for rain basins, but you find more consistent increases for transient and snow basins. Pointing this out here helps show the continuity and newness of these results.

[We added some text to the results section as suggested.](#)

p 8: This section brings a lot of fresh insight, and I'd like to see a little more. In particular, given the interesting result of high flood ratios [*sic*] in the Upper Columbia, it would be interesting to understand what's different about models like cnrm-cm5 and canesm2 that show the strongest result.

[Analysis of the sort suggested would require detailed analysis of the individual GCM runs, and is beyond the scope of this paper.](#)

At l232 you say "the role of climate grows more important and the role of hydrologic variability less important." The word "role" suggests that in the real world, changes in climate and hydrology have differing effects. Really, it's just the uncertainty in modeling these that map onto uncertainty in the results differently depending on location. So

We changed the text as suggested.

In the conclusions, you make comparisons to Salathé et al (2014), and attribute a lot of the difference to the single scenario used there. Another factor is that the RCM is going to have a lot shorter spatial coherence (ie lack of temporal correlation across small distances) in the precipitation. So, whether it is snow or rain, for a single realization, there will be a few stations that buck the larger (deterministic) trend by chance. A large ensemble would reduce that effect.

Great point. We changed the text accordingly.

----- REVIEWER 3 -----

We are grateful to Reviewer 3 for taking the time to perform this review and for noting the effort required to conduct this thorough assessment. We address below the reviewer's concerns.

"My first concern is the use of GCMs and not RCMs."

We have several responses to this concern. [We have added substantial text in the manuscript, including all the key elements of our responses below, to address this concern.](#)

- 1) RCMs certainly have their place in such work and were used in some previous studies noted in our paper. But this dataset was developed in order to sample a larger climate space than is possible with RCMs, which must be driven by GCMs anyway and are too expensive to run to generate the 40 different climate scenarios used here. As far as we are aware, the most comprehensive published dataset of RCM-GCM combinations is the NARCCAP ensemble which is now nearly a decade old and was conducted using just 6 RCMs, all at 50km spatial resolution.
- 2) Our ANOVA analysis (Figure 8) shows that the climate scenarios contribute a majority of the variation among results for most of the basin. Consequently, it is of great importance to sample the climate scenarios. Using RCMs would constrain us to a much smaller range of climate scenarios.
- 3) RCMs are still at too coarse a resolution to use in our simulations, and thus would have to be further downscaled and most likely bias corrected.
- 4) Since the flood events for the Snake and Columbia have a significant snowmelt component, the value of getting the hydrological processes right probably exceeds the additional value of RCMs in simulating daily rainfall vs the MACA approach for linking large-scale flow to local processes through the constructed analogs approach.

5) The mere fact that an alternative approach exists should not mean that the current approach, which has substantial backing in the literature, is rejected. Using RCMs would be an entirely different study, with (as noted above) its own weaknesses.

“My second concern is the use of an analogue based downscaling approach which may be compromised in its ability to represent unseen extremes”

The MACA dataset has been in wide use for nearly a decade while undergoing improvements, and although other approaches exist, the most recent improvement on MACA (LOCA) also uses constructed analogs. In short, this approach is still the benchmark. Points 4 and 5 are relevant here too. [We have added a new verification section, 2.3, to show the performance of the modeling setup.](#)

Alternately the authors must try to capture some novel question in their analysis that may shed light on processes elsewhere. For instance, a significant portion of the flow in the Columbia comes because of melt. Additionally, it is well known GCM simulations are not very reliable in the context of precipitation. Is there a research question in how one could downscale snow and rain using GCMs in a way snowpack dynamics for the current climate period are well represented? Additionally, how this downscaling would compare [sic] with the higher spatial scale simulations from RCMs over the study region. There may be other questions too that could be of interest. Given the work the authors have already done, I urge them to identify such questions and change their presentation to addressing these instead of reporting overall changes in the basin.

We contend that our approach is novel – the reviewer has not pointed us toward, nor could we find, a paper in the literature that uses such a large climate-hydrological ensemble to characterize the changes in flood magnitude over a basin, let alone while systematically presenting the dependence of results on climate scenario, location/hydrological characteristics, and other factors. Our ANOVA results, and plots distinguishing the variation across climate scenarios and hydrology scenarios, are all unique in the literature as far as we are aware. Moreover, the very purpose of this dataset – to inform international treaty negotiations – sets it apart from standard academic research. The reviewer’s other suggestions to investigate snowpack dynamics or to use RCMs, are considerably beyond the scope of this paper.

In conclusion, we contend that our dataset and our approach are sufficiently state-of-the-science to merit publication, that HESS publishes papers of similar novelty and geographic focus, and that the reviewer has not provided enough evidence either that RCMs are the only acceptable tool for scientific studies such as ours nor that others have effectively done the same kind of study.

----- REVIEWER 4 -----

We are grateful to Reviewer 4 for a constructive and insightful review, and for describing the paper as important and relevant to HESS. Below we outline our plans for heeding the excellent points raised in this review.

1. How does the daily rainfall and resultant flood statistics compare with historic data? How well does the GCM downscaling match up and how well does the 4 hydrologic model generated flood event

compare with the historic data for magnitude and timing. As mentioned in the paper, the previous work was based on annual and monthly flood statistics and the change in modelling timesteps require some sort of validation. Acknowledging that this basin has a lot of flow control, even the comparison of the upper reaches of the river system might be sufficient to gain some level of comparison on how well the four hydrologic model simulations compare with historic data.

We added a new section 2.3 to summarize new analysis we performed in which we compare our large ensemble (hydrologic models driven by free-running GCMs, not observed meteorology) with 'observed' streamflows from a dataset called No Reservoirs No Irrigation (NRNI; RMJOC 2017). We compared return period curves as in Figure 2, without GEV fits, and twin-violin plots as in Figure 9.

2. As addressed in the conclusion of this paper, the question on how much of a contribution does the PRMS model results have on the increase in flood ratios needs to be addressed in some way as this paper suggests an increase in flood risk which is different to many other studies. A possible way that can be considered might be, were PRMS flood predictions higher for both periods of comparison? How do the result change if the results from the PRMS models are not considered?

The analysis in section 2.3 provided reasons to retain the PRMS results.

3 The points raised in the table are good recommendations for clarifying and expanding the text and we addressed them all in our revisions.

River Management Joint Operating Committee (RMJOC). (2017). NRNI Flows 1929-2008 Corrected 04-2017. Bonneville Power Administration. Retrieved from <https://www.bpa.gov/p/Power-Products/Historical-Streamflow-Data/Pages/No-Regulation-No-Irrigation-Data.aspx>

Ubiquitous increases in flood magnitude in the Columbia River Basin under climate change

Laura E. Queen¹, Philip W. Mote¹, David E. Rupp¹, Oriana Chegwiddden², and Bart Nijssen²

¹Oregon Climate Change Research Institute, Oregon State University, Corvallis OR 97331 USA

²Department of Civil and Environmental Engineering, University of Washington Seattle WA 98105 USA

Correspondence to: Laura Queen (lqueen@uoregon.edu)

Abstract. The US and Canada have entered negotiations to modernize the Columbia River Treaty, signed in 1961. Key priorities are balancing flood risk, hydropower production, and improving aquatic ecosystem function while incorporating projected effects of climate change. In support of the US effort, Chegwiddden et al. (2017) developed a large-ensemble dataset of past and future daily flows at 396 sites throughout the Columbia River Basin (CRB) and select other watersheds in western Washington and Oregon, using state-of-the-art climate and hydrologic models. In this study, we use that dataset - the largest now available - to present new analyses of the effects of future climate change on flooding using water year maximum daily flows. For each simulation, flood statistics are estimated from Generalized Extreme Value distributions fit to simulated water year maximum daily flows for 50-year windows of the past (1950-1999) and future (2050-2099) periods. Our results contrast with previous findings: we find that the vast majority of locations in the CRB are estimated to experience an increase in future discharge magnitudes. We show that on the Columbia and Willamette rivers, increases in discharge magnitudes are smallest downstream and grow larger moving upstream. For the Snake River, however, the pattern is reversed, with increases in discharge magnitudes growing larger moving downstream to the confluence with the Salmon River tributary, and then abruptly dropping. We decompose the variation in results attributable to **variability in** climate and hydrologic factors **across the ensemble**, finding that climate contributes more variation in larger basins while hydrology contributes more in smaller basins. Equally important for practical applications like flood control rule curves, the seasonal timing of flooding shifts dramatically on some rivers (e.g., on the Snake, 20th century floods occur exclusively in late spring, but by the end of the 21st century some floods occur as early as December) and not at all on others (e.g. the Willamette).

1 Introduction

Among natural disasters in the Northwest, flooding ranks second behind fire in federal disaster declarations¹ since 1953 despite extensive flood prevention infrastructure. The largest flood in modern times on the Columbia occurred in late spring (May-June) 1948, and obliterated the town of Vanport which lay on an island between Portland, OR and Vancouver, WA, permanently displacing its 18,500 residents². Other disruptive floods in the region include the Heppner flood in 1903, one of the deadliest flash floods in US history (Byrd, 2014); floods on the Chehalis River in both December 2007³ and January 2009⁴ that closed Interstate 5, the main north-south transportation corridor through the Northwest, for several days each time at a cost of several \$m per day to freight movement alone; and floods on the Willamette River in February 1996 and April 2019. The timing of typical floods varies widely across the region: low-elevation basins in western Washington and Oregon typically flood in November through February, whereas the snow-dominant basins east of the Cascades more typically flood in spring, even as late as June (Berghuis et al. 2016).

The Columbia River drains much of the Northwest, with the fourth largest annual flow volume in the US and a drainage that includes portions of seven states plus the Canadian province of British Columbia (BC), an area of 668,000 km² (Fig. 1). Its numerous federal and nonfederal dams provide flood protection, hydropower production, navigation, irrigation, and recreation services. A treaty between the US and Canada, signed in 1961, codified joint management of the river's reservoirs (and funded construction of new reservoirs in BC) primarily to provide flood protection and hydropower production⁵. The US and Canada have entered negotiations to update the treaty; the USA's "key objectives include continued, careful management of flood risk; ensuring a reliable and economical power supply; and improving the ecosystem in a modernized Treaty regime." (*ibid.*) Both countries have expressed an intention to include the effects of climate change on flows, and clearly a key aspect of hydrologic change is to inform the treaty negotiations of the influence of climate change on the magnitude of flooding.

¹ <https://www.fema.gov/data-visualization-summary-disaster-declarations-and-grants> accessed 8/6/2019

² https://www.oregonlive.com/portland/2017/05/vanport_flood_may_30_1948_chan.html accessed 8/6/2019

³ <https://www.seattletimes.com/seattle-news/extensive-flooding-3-confirmed-deaths-hundreds-of-rescues/> accessed 8/6/2019

⁴ <https://www.seattletimes.com/seattle-news/despite-drying-cooling-trend-flooding-and-road-closures-continue/> accessed 8/6/2019

⁵ <https://www.state.gov/columbia-river-treaty/> accessed 8/6/2019

While rising temperatures potentially affect all parts of the hydrologic cycle, in a snowmelt-dominated hydrologic system such as many of the Northwest's river basins, warming directly affects snow accumulation and melt (e.g., Hamlet et al. 2005). Observational studies have shown consistent changes toward lower spring snowpack (Mote et al. 2018), earlier spring flow (Stewart et al. 2005), and lower summer flow (Fritze et al. 2011) since the mid-20th century. Observations of trends in flooding in the US have generally failed to find any consistent trends (Lins and Slack 1999; Douglass et al. 2000; Sharma et al. 2018). Sharma et al. (2018) offer several possible explanations, chiefly "decreases in antecedent soil moisture, decreasing storm extent, and decreases in snowmelt". The detection of trends in floods is complicated by the interaction of extreme events and nonstationarity (Serinaldi and Kilsby, 2015). Moreover, as a result of the substantial alteration of rivers to prevent flooding (e.g., by the construction of dams and levees) during the observational period, the best long-term records - i.e., on streams with the least modifications - are on rivers that were not producing sufficiently disruptive floods to lead decision-makers to construct flood protection structures. That is, as flooding of settlements, infrastructure, or other assets led to the investments in flood protection structures on most rivers, thereby altering the flow regime and dividing any gauged records into pre- and post- modification, the ones that were left unmodified tended to be small and/or remote.

To interpret the ambiguous results from observed trends, Hamlet and Lettenmaier (2007) used the Variable Infiltration Capacity (VIC) hydrologic model forced twice with detrended observed daily weather for the period 1916-2003, with about 1°C of temperature difference between the two. They then compared 20- and 100-year flood quantiles for basins at varying sizes in the western US and found a wide range of changes in flood magnitude ranging from large decreases to large increases (+/- 30%). Broadly, the responses depended somewhat on basin winter temperature, with the coldest basins (<-6°C) showing reductions in flood magnitude owing to reduced snowpack, basins with moderate temperatures exhibiting a wide range of changes, and rain-dominant (>5°C) basins showing little change, though the warm basins in coastal areas of Washington, Oregon, and California showed increased flood magnitude.

Modeling work using state-of-the-art hydrologic models has been applied to understand where and how flood magnitudes may change in the future. Tohver et al (2014) found widespread increases in flood magnitudes, especially in temperature-sensitive basins (mainly on the west side of the Cascades), but their approach used monthly GCM output so changes in daily precipitation would not be represented. Salathé et al. (2014) used a

single global climate model (GCM), the ECHAM5, linked to a regional climate model to obtain high-resolution (in space and time) driving data for VIC over the period 1970-2069. As did Hamlet and Lettenmaier (2007), they compared the ratio of flood change (2050s vs 1980s) against mean historical winter temperature and found a majority of locations with a higher 100-year flood, in some cases by a factor of 2 or more; while they projected increases in every one of the warmer basins ($>0^{\circ}\text{C}$), a substantial fraction of colder locations had decreases in flood magnitude.

As noted above and detailed below, Chegwidden et al. (2019) describe the process used to generate the streamflow ensemble used here. In addition, they used analysis of variance (ANOVA) to analyze the different influences of choices of emissions scenario (as a Representative Concentration Pathway - RCP), GCM, downscaling method, and hydrologic model, and how those influences varied spatially across the domain and also seasonally and by hydrologic variable. They found that the RCP and GCM had the largest influence on the range of annual streamflow volume and timing, and hydrologic model had the largest influence on low flows. The hydrologic variables they considered were snowpack (maximum snow water equivalent and date of maximum SWE), annual streamflow volume, centroid timing (the date at which half the water year's flow has passed), and seasonal streamflow volume; primary focus was on centroid timing, annual volume, and minimum 7-day flow. They did not examine maximum daily flow. The purpose of this paper is to address this important gap in our understanding of the future Northwest hydrology; to do so, we use the largest available ensemble of climate-hydrology scenarios. By using a large ensemble, we ensure a reasonable breadth of climatic and hydrological futures in order to better describe the range of possible future flooding and how it varies across the region with its diverse hydroclimates.

2 Methods

2.1 Hydrologic modeling data set

To assess changing flood magnitudes under climate change, we analyzed changes in water year maximum daily flows in a large ensemble of streamflow simulations at 396 locations in the CRB (Figure MAP) and select watersheds in western Oregon and Washington (Chegwidden et al., 2017). The simulations were constructed from permutations of modeling decisions on forcing datasets and hydrologic modeling. Specifically, choices included two RCPs (RCP4.5 and RCP8.5), ten GCMs, two methods of downscaling the climate model output to the resolution of the hydrologic models, and four hydrologic model implementations, for a total of 160 permutations. For our analysis, we extracted a more tractable dataset of 40 simulations per location, by only considering simu-

lations with RCP 8.5 and the Multivariate Adaptive Constructed Analogs (MACA) downscaling method (Abatzoglou and Brown, 2012).

The rationale for using a subset of the available data is as follows. First, the time-dependent set of greenhouse gas concentrations in RCP4.5 is fully included in RCP8.5, so any concentration of greenhouse gases on the RCP4.5 path can be converted to a point on RCP8.5 (at a different time). We analyzed results for both RCP8.5 and RCP4.5, and found that to first order the changes in flood magnitude in RCP4.5 were approximately 2/3 those in RCP8.5, which is also roughly the ratio of global temperature change over the period considered (IPCC Summary for Policymakers, 2014). For clarity we show only the results for RCP8.5. Second, we considered only simulations using the MACA downscaling method because of the method's ability to capture the daily GCM-simulated meteorology critical for assessing changes in extremes and its skill in topographically complex regions (Lute et al., 2015). The other downscaling approach used by Chegwiddden et al. (2019), the Bias Correction and Statistical Downscaling (BCSD) method (Wood et al. 2004), produces probability distributions of daily precipitation inconsistent with the GCM response to forcings because the method stochastically disaggregates monthly data to daily data based on historical statistical properties of the daily data. This statistical property limits the ability of BCSD to reproduce changes in storm frequency in the future, making it a less attractive choice for daily extreme flow analysis (Hamlet et al. 2010; Guttman et al. 2014).

The GCMs used in this study are the CanESM2, CCSM4, CNRM-CM5, CSIRO-Mk3-6-0, GFDL-ESM2M, HadGEM2-CC, HadGEM2-ES, Inmcm4, IPSL-CM5A-MR, and MIROC5. These ten GCMs were chosen primarily for their ability to accurately reproduce observed climate metrics during the historical period mainly of the Northwest US but also at sub-continental and larger scales as assessed in Rupp et al. (2013) and RMJOC (2018). The four hydrologic model implementations originated from two distinct hydrologic models: the Variable Infiltration Capacity (VIC; Liang et al., 1994) model and the Precipitation Runoff Modeling System (PRMS; Leavesley et al., 1983). VIC and PRMS are process-based, energy balance models and were both run on the same 1/16th degree grid with output saved at a daily time step for the period 1950 to 2099. VIC is a macroscale semi-distributed hydrologic model that solves full water and energy balances, and in these simulations it also included a glacier model (Hamman & Nijssen, 2015). Three unique implementations of VIC were used with independently derived parameter sets (P1, P2, P3) marked by differences in calibrated parameters, calibration methodology, and meteorological and streamflow reference sets. PRMS is a distributed, deterministic hydrologic model which, in contrast to VIC, does not allow for subgrid heterogeneity. See Chegwiddden et al (2019) for details. It is important to note that these hydrologic simulations and calibrations do not include reservoir models.

2.2 Flood magnitude

We assessed changes in flood magnitude in the Columbia River Basin by comparing maximum daily streamflows over a 150-year period (1950-2100). We estimated the 10, 5, 2, and 1% probability of occurrence (commonly referred to as the 10-, 20-, 50-, and 100-year flood, respectively) by fitting generalized extreme value (GEV) probability distributions to simulated water year maximum daily flows for 50-year windows of the past (1950-1999) and future (2050-2099) periods. (We also looked at 30- and 75-year windows, choosing 50 years as a balance between sample size favoring longer periods, and nonstationarity considerations favoring shorter periods.) We used Python's `scipy.stats.genextreme` module (Jones et al., 2001) to fit a Gumbel distribution and estimate flood magnitudes for each return period. We assessed change in flood magnitude as the “discharge ratio” of the estimated future to past floods for a given return period; a ratio greater than 1 indicates an increase in flood magnitudes while a ratio less than 1 indicates a decrease.

We describe how changes in flood magnitude vary by climatic zone across the PNW by using an efficient and internally consistent proxy for climatic zone: the centroid of timing – the day in the water year that half the annual volume of water has passed through the location. The centroid of timing is a metric of snow dominance (e.g., Stewart et al. 2005) which is related to the spatial distribution of temperature and tends to decrease downstream. This temporal proxy of a hydrologic characteristic is effective in the Columbia Basin where most of the precipitation occurs in winter and the relative magnitude and timing of the freshet from the spring thaw is a good indicator of importance of snowmelt to streamflow. An early centroid indicates that rain, which falls predominantly during the cooler, earlier part of the year, is the driver of the peak flows at the location, while a late centroid indicates that snowmelt during later spring months is the prime hydrological driver. We computed the centroid using the 1950-79 simulated years. Note that Chegwiddden et al. (2019) also used the *change* in centroid as a hydrologic variable of interest; below, we discuss our results in the context of their findings.

2.3 Verification

Comparing directly between gauged flows and modeled flows is inadvisable since the flows are substantially altered by regulation. However, a set of streamflows called No Reservoirs No Irrigation (NRNI; RMJOC 2017) has been developed by federal agencies to support practical analysis. The NRNI dataset exists at ~190 sites across the Columbia River Basin for the years 1950-2008, and are adjusted to correct for reservoir management and the diversions and evaporation associated with both the reservoirs and with irrigated agriculture. This dataset is suitable for comparisons with our modeling setup, and we have computed return period curves at all the NRNI locations (not shown). On the lower mainstem Columbia, the return period curves are extremely close

to those computed from NRNI. Individual hydrologic model configurations are not consistently biased across the basin nor across return periods; despite its different provenance, PRMS generally lies within the return period flows of the three VIC configurations rather than being consistently different from all VIC configurations, except on the lower Snake, where PRMS is consistently an outlier on the low end of the distribution. Only at Hills Creek in the Willamette Basin do the modeled return period curves all lie outside NRNI, and only for the longest return periods (>10 years).

We also examined the ensemble performance for 1950-2008 in the distribution of timing of peak daily flow for 28 locations along the Columbia, Snake, and Willamette. At all locations we examined, the median date (as well as earliest and latest quartiles) of annual maximum flow in the ensemble is within 10 days of the observed, from NRNI. The modeled distribution is a bit later than NRNI on the lower Columbia and a bit earlier than NRNI on the Willamette. Note that the GCM simulations used to drive the hydrologic models during this verification period are independent of the observed meteorology, so both the magnitude and the timing of annual maximum flows are computed from first principles, and represent a remarkable agreement with observations. Although the modeled flows are calibrated, the statistical approach to calibrations is not sensitive to the extreme maximum daily flow studied here.

3 Results

3.1 Regional changes in flood ratio

Figure 3 shows the changes in maximum daily discharge for all of the 396 flow locations for different return periods. The horizontal position of each circle represents the centroid of timing. The circles are semi-opaque so overlapping circles lead to a deeper saturation. Points on the same river are ordered from more to less snow dominant (i.e., right to left) traveling downstream; strings of circles in a smooth pattern usually indicate one of the larger rivers, highlighted in Figure 4. Each circle in Figures 3 and 4 represents an average of 40 simulations: 10 GCMs and 4 hydrologic model configurations.

A striking result in Figure 3 is that, in contrast to the results of Tohver et al. (2014), the flood magnitude increases (i.e., the discharge ratio exceeds one) at nearly every flow location and return period (though not for every individual climate scenario, as shown in Figure 5). Broadly, the patterns are similar across all return periods though with slightly higher ratios for longer return periods, and subsequent figures will show only the 10-

and 100-year floods. For the flow locations with centroid <125 or so (i.e. February 2), flood ratios are fairly concentrated about 1.25 for all return periods. For mixed rain-snow basins, roughly delineated by centroids between 125 and 160 (March 8 most years), flood ratios range widely from just below 1 to about 2.4 for the 10-year and 3.2 for 50- and 100-year floods. For the longer return intervals, there is a wide range of projected changes in daily flood at many locations (indicated by the red coloring). This is undoubtedly partly due to the GEV fit extrapolating from 50 to 100 years. Finally, for the basins with flow centroid >160, the ratios have a smaller range, from slightly greater than 1 to a maximum that increases from about 2 for the 10-year, to about 2.75 for 100-year. [Tohver et al. \(2014\)](#) distinguished basins by their DJF temperature, a rough proxy for our snow dominance metric, and found a substantial number of locations where the flood ratio for both 20-year and 100-year flood was as much as 20% lower for the 2040s compared with a historical period. We return to this point in the conclusions.

To understand better how flood magnitude changes along the length of a river, we focus (Figure 4) on a handful of significant rivers in the region: the mainstem Columbia, Willamette (along with major tributaries the McKenzie and Middle Fork Willamette), and Snake, and also on the Chehalis in southwest Washington (see Introduction). Flow locations are listed in Table 3 Rivers in the Appendix. [Many of the larger tributaries also have flow points in our dataset, so we can infer the role of tributaries in changing the flood magnitudes in the future, as discussed below.](#) The Columbia River includes the most snow-dominant basins, with a centroid of >190 days (early to mid April) in the Canadian portion of the basin. The flood ratio decreases almost uniformly along the length of the river, from 1.3 for the 10-year and >1.5 for the 100-year in the Canadian portion to just above 1 at the last few points along the river (The Dalles, Bonneville, and Portland). [Past flood events on the mainstem Columbia are exclusively associated with large spring snowmelt, and the large tributaries \(the Yakima, Snake, and Willamette\) contribute annual flow volume but rarely contribute peak flow at the same time; as shown below, the future flood timing changes but flood magnitudes change little in the lower Columbia owing to the fact that the Columbia integrates such diverse hydroclimates.](#) Like the Columbia, the Willamette also has flood ratios that decrease along the length of the river [as it integrates more diverse hydroclimates](#), from 1.7 to 1.35 for both return periods. The McKenzie River (points 15-17), one of the three tributaries that converge at Eugene to form the Willamette, is a highly [spring-fed](#) river with higher baseflow than is represented in the hydrologic models, though it is unclear how that difference would manifest in the flood statistics.

[In contrast to the Columbia and the Willamette](#), the Snake behaves oppositely: flood ratio increases along the length of the river, until the confluence with the Salmon River, which drains a large mountainous area of central

Idaho. On parts of the Snake the ratios are as high as 1.4 for 10-year and 1.6 for 100-year. Then after the confluence with the Salmon River, which has much lower change in discharge ratio, the ratios on the Snake drop to about 1.2 for 10-year and about 1.3 for the 100-year. Our hypothesis is that in the Snake above the Salmon River, the tributaries shift from snow-dominant to rain-dominant, so that a single storm can drive large rainfall-driven increases (possibly with a snowmelt component) leading to larger synchronous discharges. The Salmon and Clearwater rivers retain less exposure to such shifts, and dilute the effects of single large storms on flooding.

Each circle in Figures 3 and 4 represents an average of 40 simulations: 10 GCMs and 4 hydrologic model configurations. To better understand the range in results, Figure 5 shows the discharge ratio for all 40 simulations at each point on the mainstem Columbia. Although the mean flood ratio at the lowest two points is only barely above 1, several ensemble members have ratios less than one, and a few have ratios >1.5 . Moving upstream, the range in results increases, as shown also by the color of the dots.

3.2 Dependence of results on modeling choices

As in Chegwiddden et al (2019), we separate the results - here for the three largest rivers - into variations across GCM (Figure 6) and variations across hydrologic model configurations (Figure 7). The ranking of flood ratios by GCM changes substantially between basins and even within a basin, and does not correspond to the changes in seasonal precipitation. For the upper Columbia River, the models with the least warming - inmcm4 and GFDL-ESM2M (Rupp et al 2017) - have almost no change in flood magnitude, but the HadGEM2-ES which warms considerably in summer produces a large decrease in flood magnitude. In the Willamette and Snake Rivers, the range of projected flood changes by different GCMs remains large from the headwaters to the mouth of the river, whereas for the Columbia the range diminishes considerably as one moves downriver.

The variation of results depends less on hydrologic model than on GCM (Figure 7), though the differences across hydrological models are still substantial. For the Willamette, lower Snake, and both upper and lower Columbia, the PRMS model predicts substantially larger increases in flooding than the three calibrations of the VIC model. For the upper Snake, it predicts substantially smaller change than any VIC calibration. While it is perhaps not surprising that the three calibrations of VIC are close to each other, it is striking just how different are the projections from PRMS at most locations on these three rivers. Chegwiddden et al. (2019) found that the main contributors to differences in hydrologic variables (except low flows) generally were the climate scenarios (GCM and RCP), consistent with our findings here. (The order of models is similar in the equivalent figure for the 100-year return period, but we elected to show the 10-year figure since the 100-year figure is more difficult to decipher because the symbols overlap with those from other rivers.)

To parse the contributions of climate factors (represented by the GCMs) and hydrologic factors (represented by the hydrologic models), we perform ANOVA on the 40 discharge ratios. The pie charts in Fig. 8 show the proportion of the total variance explained by climate factors and hydrologic factors at different locations. For the Willamette River, the [portion of uncertainty connected to the climate](#) grows more important and the [portion of uncertainty connected to the hydrologic variability](#) less important going from the confluence of the three major tributaries at Eugene to the mouth. For the Snake and Columbia rivers, climate is responsible for virtually all of the variance in projections in the upper reaches, but only about half at the lowest point, similar to the Willamette. The Willamette basin is much smaller, and a large storm can affect the entire basin on the same day, whereas storms typically take a couple of days to move across the Snake and Columbia (and generally move upstream). With larger and more diverse contributing areas, differences in the rates with which the hydrological models transfer precipitation to the point of interest become more important. Unlike Chegwiddden et al. (2019), we did not attempt to isolate the response to anthropogenic forcing from internal climate variability. Though several techniques for separating these two factors have been used (e.g., Hawkins and Sutton, 2009; Rupp et al., 2017; Chegwiddden et al., 2019), these techniques are either infeasible with our dataset or we question their suitability for the application to changes in extreme river flows.

3.3 Change in timing

Although in a broad hydrologic sense a flood is a flood regardless of what time of year it occurs, there are potentially significant ecological differences depending on time of year; for example, scouring salmon redds (Goode et al. 2013). Moreover, water management policies are strongly linked to the calendar year (see Discussion). We computed the probability of flooding for (all 40) past and future simulations at all the points on the three rivers (Figure 4) as a function of day of year (Figure 9). For the Willamette, no significant change in timing occurs; however, for the upper Willamette, a single peak in likelihood in February becomes more diffuse. For the Snake, all locations see a shift toward earlier floods, consistent with the transition to less snow-dominant and more rain-dominant. Whereas floods were historically concentrated in the period of mid-May to mid-July, the projected future flooding period spans December to June. For the Columbia, the mode in the flood timing shifts earlier by half a month in the upper Columbia to about a month in the lower Columbia. The distribution also broadens with an elongated tail towards winter such that there is low, but non-negligible, probability of floods occurring as early as January. Although the magnitude of the 10- and 100-year flood events in the lower Columbia do not increase much (Figures 4-7), the risk of major flood on any given day decreases, and the likelihood of major flooding in May or April (or even February and March) increases.

4 Discussion and conclusions

Our study joins a small number of others in examining large hydroclimate ensembles. Gangrade et al. 2020 used a similar ensemble approach analyzing hydrological projections for the Alabama-Coosa-Tallapoosa River Basin with 11 dynamically downscaled and bias corrected GCMs (10 of which our studies share) and 3 hydrologic models (including VIC and PRMS). While they did not examine extreme daily flows, they did calculate changes in the 95% percentile of daily streamflow (Q95). Perhaps because of the hydroclimatic uniformity of that basin, they found very small differences in Q95 across hydrologic models, which contrasts with our results showing changes in flood magnitudes varying by watershed and distance downstream. Thober et al. (2018) conducted a similar study in some European river basins, but rather than using a climate ensemble they simply imposed uniform warming scenarios on a hydrologic model (i.e. a more straightforward temperature sensitivity analysis rather than an exploration of the range of future climate scenarios).

Returning to the Northwest, our findings contrast with earlier work. Salathe et al. (2014) found decreases in flood magnitude at a substantial number of sites, but our results show increases in flood magnitude at nearly every return period and location, which includes about 100 locations not included in their study. They also noted that directly downscaling the GCM outputs leads to a smaller range of results than when running the regional model as an intermediate step, so we infer that if we had had access to RCM simulations driven by all 40 of our GCMs, our range of results might have been larger. Another important difference may be in the spatiotemporal coherence of extreme precipitation, which in the RCM would be generated directly by the interaction of synoptic-scale storms, topography, and to a small extent by surface water and energy balance; and in our study, by the interaction of the GCM-scale synoptic storms and constructed analogs derived from observations. A large ensemble would reduce the magnitude of that effect. In our study, the MACA statistical downscaling approach preserves much of the daily variability from the GCM, so the primary reason for the difference between our results and theirs is probably the fact that we analyzed 40 scenarios. Some locations, for example the points on the lower Columbia river, had a handful of ensemble members with decreasing flood magnitude. But averaging the entire ensemble nearly always resulted in an increase in flood magnitude. It is possible therefore that their study, repeated with a larger ensemble of hydrologic-climate model combinations, might have found ubiquitous increases in flood magnitude as ours did.

Prior results (Hamlet and Lettenmaier 2007, Tohver et al. 2014, Salathe et al. 2014) suggested a decrease in flood magnitude in snowmelt-dominated basins like the Columbia, since reduced snowpack reduces the store of water available to be released quickly in a spring flood (like the May-June 1948 Vanport flood). In a subbasin of

the Willamette, Surfleet and Tullos (2013) projected decreases in flood magnitude for return periods > 10 years in the Santiam River basin under a high-emissions scenario (SRES A1B, 2070-2099 vs. 1960-2010; 8 GCMs), attributing the decreases to fewer large rain-on-snow events. Our results for the Santiam River show an *increase* of 40% for both 10- and 100-year floods; this result includes rain-on-snow events, since they are represented in VIC, which computes the accumulation of water in the snowpack and determines whether sufficient energy has been provided to create a melt event. Our results point to ubiquitous increases throughout the basin, even on the lower mainstem Columbia. The coldest basins including the headwaters of the Columbia also had some large increases in flood magnitude, suggesting that the former results were missing some key details. It seems likely that any reduction in flood magnitude originating from the warming-induced reduction in spring snowpack is offset by the increased pace of melt (including possibly rain-on-snow events). These results emphasize the necessity of revisiting reservoir rule curves, which are strongly tied to historical hydrographs, and also emphasize that changes in the seasonality of flooding can be dramatically different from the changes in the mean hydrograph. In particular, in the lower Snake and lower Columbia, changes in magnitude of flooding are modest but changes in timing of the earliest quartile of flood events is much larger than the 0.5-1 month shift in the mean hydrograph.

A strength of our study compared with earlier studies is the use of a large ensemble, which samples a wide climate space by using GCMs as opposed to RCMs. Conventional wisdom and evidence from the weather and seasonal climate forecasting realms illustrate the utility of considering ensembles, and that generally the true outcome of a prediction lies near the middle of the ensemble. Our ANOVA analysis (Figure 8) shows that climate scenarios contribute a majority of the variation among results for most of the basin. Consequently, it is of great importance to sample the climate scenarios broadly, which only GCMs can do. Large ensembles of RCMs are rare; the 12-member NARCCAP ensemble (6 RCMs, 4 GCMs; Mearns et al. 2013), some of whose model runs were completed a decade ago, remains the largest, but has a spatial resolution of only 50km. CORDEX North America, similarly now has a comparable-size ensemble, but mostly still at 50 km (some at 0.22°), and was not available in such large numbers when we began our hydrologic simulations. At such spatial resolutions, RCMs would still have to be further downscaled and bias corrected to use in our hydrologic models (~6km spatial resolution). In the tradeoff between breadth of climate scenarios and spatial resolution, these ensembles offer insufficient improvement in spatial resolution relative to our GCM ensemble to justify sacrificing the breadth in climate scenarios represented by choosing just 4 GCMs. While RCMs certainly have their place in such work and were used in some previous studies, using GCMs in this study allowed for a larger climate space to be sampled, thus adding to the robustness of our results.

The spread of results shown in Fig 5 suggests that although the likeliest outcome is little change in flood magnitude in the lower Columbia, a prudent risk management strategy would consider the range of possibilities. However, we view the highest outcomes (>50% increase in peak 100-yr flood) as less likely than other individual scenarios, because they are the product of a hydrologic model that may be less suited to calculating the extreme changes in a much warmer world.

Our findings provide an initial indication of how existing flood risk management could respond to a warming climate. Reservoir management is guided by rule curves which are intended to reflect the changing priorities and risks during the year. For example, reservoirs used for flood control have rule curves that require reservoir levels to be lowered when approaching the time of year when flood likelihood increases, and reservoir levels may be raised as the likelihood decreases. For the Willamette, we found little change in the distribution of timing of flood events, which indicate that with the state of the science today, reservoir rule curves may need to be altered as to magnitude of flooding (which our results indicate will increase by 30-40%) but not timing; a reservoir model would be required for complete understanding of how flood risk (magnitude and timing) will actually change. For the Snake, larger shifts in the timing imply a need to completely rethink the existing rule curves. For the Columbia, the mode in flood timing shifts earlier by half a month in the upper Columbia to about a month in the low Columbia. The distribution also broadens, with an elongated tail towards winter such that there is low, but non-negligible, probability of floods occurring as early as January. These changes in timing imply a need for moderate alteration of rule curves for reservoirs in the Canadian portion of the Columbia Basin.

Our results should not be taken as a precise prediction of flood magnitude change but rather as the best available projections given the current state of the science. Two important factors need to be taken into account in interpreting our results: first, in using RCP8.5, we selected the most extreme emissions scenario. If efforts to stabilize the climate before 2050 are successful, the flood magnitudes shown here will undoubtedly be smaller (our analysis suggests most of the locations would see a change in flood magnitude about 1/3 smaller, for RCP4.5; e.g., a ratio of 1.3 (30% increase) for RCP8.5 would correspond to a ratio of 1.2 for RCP4.5).

The second important factor in interpreting our results is that the actual river system in the Northwest includes many dams, a majority of which have flood control as a primary (or one of a few top) objective. As a result, actual flows (and the changes in flow) at a given point in the river would be altered by reservoir management. Translating these changes in flood magnitude into actual changes would require a reservoir model for the basin

or subbasin of relevance. One could then compute optimal rule curves for the major flood control reservoirs (perhaps time-evolving every couple of decades, to reflect the likely changes in scientific understanding and emissions trajectory). Even without that additional analysis, however, our results stress that the magnitude and/or timing of flood events will change throughout the basin. In other words, what worked for flood control in the past will not work as well in the future.

This study may have some utility in framing and quantifying the possible changes in flood risk as the Columbia River Treaty is in renegotiation, but further work would be needed to assign probabilities to future flood magnitude. Such work includes (a) understanding whether the PRMS projections of much larger change are reliable (our analysis in section 2.3 shows that PRMS performs about as well as the three calibrations of VIC for simulating past peak flows, but more work would be needed to understand the reasons for divergence in future projections), (b) applying different statistical and/or dynamical downscaling methods, and (c) using a more sophisticated approach to evaluating extremes in a nonstationary climate (as advocated by Serinaldi and Kilsby, 2015). The mechanisms of flooding in the upper Columbia and elsewhere are also a key question arising from this work; this and other work is needed to decipher the cause of the discharge ratio patterns we found along the major rivers. Furthermore, a new generation of GCM outputs (CMIP6, Eyring et al. 2016) already has data available from over 25 GCMs; in the near future, it would be feasible to apply a newer multi-model hydrologic modeling approaches (e.g., Clark et al., 2015) to the new generation of GCMs, though perhaps no significant changes would result.

Nonetheless, with current knowledge the fact that very few locations would see a decrease in flood risk under any climate/hydrologic scenario is a strong statement of the need to update all aspects of flood preparation: the definition of N-year (especially 100-year) return period flows, flood plain mapping, and reservoir rule curves, to name a few. Moreover, the challenges that the renegotiated Columbia River Treaty faces in accounting for climate change now appear to include the necessity of incorporating the likely increase in flood risk throughout the region.

Generally, this study shows how complex the spatial and temporal patterns of change can be in a mixed rain-and-snow basin. Basins of similar size and hydrological response to warming exist on most continents, so our results provide a warning against using a small number of climate scenarios or a single hydrologic model to estimate changes in flood risk in other basins.

Code/data availability. The data used here are available at <https://zenodo.org/record/854763>.

Author contribution. L. Queen performed all analyses, wrote portions of the text, and edited the document. P. Mote guided the analysis and wrote much of the text. D. Rupp guided the analysis and edited the document. O. Chegwiddden generated the underlying dataset, guided the analysis, provided assistance with programming, and commented on the text. B. Nijssen generated the underlying dataset and commented on the text.

Competing interests. The authors declare no competing interests.

Acknowledgments. This project originated as a senior honors thesis by the first author, who thanks Hank Childs of the University of Oregon for his mentorship. The research was supported by the NOAA Climate Impacts Research Consortium, under award #NA15OAR4310145.

References

Berghuijs, W.R., R.A. Woods, C.J. Hutton, and M. Sivapalan, Dominant Flood Generating Mechanisms Across the United States. *Geophys. Res. Letts.*, 43, 4382-4390, doi: 10.1002/2016GL068070, 2016.

Byrd, J. G.: Calamity: The Heppner Flood of 1903. University of Washington Press, 2014.

Chegwidden, O. S., B. Nijssen, D.E. Rupp, and P.W. Mote, Hydrologic Response of the Columbia River System to Climate Change [Data set]. Zenodo. doi:10.5281/zenodo.854763, 2017.

Chegwidden, O. S., B. Nijssen, D.E. Rupp, J.R. Arnold, M.P. Clark, J.J. Hamman, S. Kao, et al: How Do Modeling Decisions Affect the Spread Among Hydrologic Climate Change Projections? Exploring a Large Ensemble of Simulations Across a Diversity of Hydroclimates. *Earth's Future*, 7, 623–637, doi: 10.1029/2018EF001047, 2019.

Clark, M. P., Nijssen, B., Lundquist, J. D., Kavetski, D., Rupp, D. E., Woods, R. A., ... & Arnold, J. R. (2015). A unified approach for process-based hydrologic modeling: 1. Modeling concept. *Water Resources Research*, 51(4), 2498-2514.

Do, H. X., F. Zhao, S. Westra, M. Leonard, L. Gudmundsson, J. Chang, P. Ciais, D. Gerten, S.N. Gosling, H.M. Schmied, T. Stacke, B.J.E. Stanislas, and Y. Wada: Historical and Future Changes in Global Flood Magnitude – Evidence from a Model-Observation Investigation. *Hydrol. Earth Syst. Sci. Discuss*, doi: 10.5194/hess-2019-388, in review, 2019.

Douglas, E.M., R.M. Vogel, and C.N. Kroll: Trends in Floods and Low Flows in the United States: Impact of Spatial Correlation. *Journal of Hydrology*, doi: 10.1016/S0022-1694(00)00336-X, 2000.

Eyring, V., S. Bony, G.A. Meehl, C.A. Senior, B. Stevens, R.J. Stouffer, and K.E. Taylor: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) Experimental Design and Organization. *Geosci. Model Dev.*, 9, 1937-1958, doi: 10.5194/gmd-9-1937-2016, 2016.

Fritze, H., I.T. Stewart, and E. J. Pebesma: Shifts in Western North American Snowmelt Runoff Regimes for the Recent Warm Decades. *Journal of Hydrometeorology*, doi: 10.1175/2011JHM1360.1, 2011.

Gangrade, Sudershan & Kao, Shih-Chieh & McManamay, Ryan. (2020). Multi-model Hydroclimate Projections for the Alabama-Coosa-Tallapoosa River Basin in the Southeastern United States. *Scientific Reports*. 10.1038/s41598-020-59806-6.

Goode, J.R., J.M. Buffington, D. Tonina, D.J. Isaak, R.F. Thurow, S. Wenger, D. Nagel, C. Luce, D. Tetzlaff, and C. Soulsby: Potential effects of climate change on streambed scour and risks to salmonid survival in snow-dominated mountain basins. *Hydrological Processes*, 27, 750-765, doi: 10.1002/hyp.9728.

Gutmann, E., T. Pruitt, M. P. Clark, L. Brekke, J.R. Arnold, D. A. Raff, and R.M. Rasmussen: An Intercomparison of Statistical Downscaling Methods Used for Water Resource Assessments in the United States. *Water Resources Research*, 50, 7167–7186, doi: 10.1002/2014WR015559, 2014.

Hamlet, A.F., and D.P. Lettenmaier: Effects of 20th Century Warming and Climate Variability on Flood Risk in the Western U.S. *Water Resour. Res.*, 43, W06427, doi: 10.1029/2006WR005099, 2007.

Hamlet, A.F., P.W. Mote, M.P. Clark, and D.P. Lettenmaier, 2005: Effects of precipitation and temperature variability on snowpack trends in the western United States, *J. Climate*, 18, 4545–4561.

Hamlet, A.F., E.P. Salathé, and P. Carrasco: Statistical Downscaling Techniques for Global Climate Model Simulations of Temperature and Precipitation with Application to Water Resources Planning Studies. Chapter 4 in Final Report for the Columbia Basin Climate Change Scenarios Project, Climate Impacts Group, Center for Science in the Earth System, Joint Institute for the Study of the Atmosphere and Ocean, University of Washington, Seattle, 2010.

Hamman, J., and B. Nijssen: VIC 4.2.glacier. Retrieved from <https://github.com/UW-Hydro/VIC/tree/support/VIC.4.2.glacier>, 2015.

Hawkins, E., and R. Sutton: The potential to narrow uncertainty in regional climate predictions. *Bulletin of the American Meteorological Society*, 90, 1095–1108, doi: 10.1175/2009BAMS2607.1, 2009.

Kundzewicz, Z.W., S. Kanae, S.I. Seneviratne, J. Handmer, N. Nicholls, P. Peduzzi, R. Mechler, L.M. Bouwer, N. Arnell, K. Mach, R. Muir-Wood, G.R. Brakenridge, W. Kron, G. Benito, Y. Honda, K. Takahashi, and B. Sherstyukov: Flood Risk and Climate Change: Global and Regional Perspectives. *Hydrological Sciences Journal*, 59, 1-28, doi: 10.1080/02626667.2013.857411, 2014.

Lute, A. C., J.T. Abatzoglou, and K.C. Hegewisch: Projected Changes in Snowfall Extremes and Interannual Variability of Snowfall in the Western United States. *Water Resources Research*, 51, 960– 972, doi: 10.1002/2014WR016267, 2015.

Mearns, L.O., Sain, S., Leung, L.R. et al. Climate change projections of the North American Regional Climate Change Assessment Program (NARCCAP). *Climatic Change* 120, 965–975 (2013). <https://doi.org/10.1007/s10584-013-0831-3>.

Najafi, M.R., and H. Moradkhani: Multi-model Ensemble Analysis of Runoff Extremes for Climate Change Impact Assessments. *Journal of Hydrology*, 525, 352-361, doi: 10.1016/j.jhydrol.2015.03.045, 2015.

River Management Joint Operating Committee: Climate and Hydrology Datasets for RMJOC Long-term Planning Studies. Second edition: Part 1—Hydroclimate Projections and Analyses, retrieved from <https://www.bpa.gov/p/Generation/Hydro/Pages/Climate-Change-FCRPS-Hydro.aspx>, 2018.

Rupp, D. E., J.T. Abatzoglou, K.C. Hegewisch, and P.W. Mote: Evaluation of CMIP5 20th Century Climate Simulations for the Pacific Northwest USA. *Journal of Geophysical Research: Atmospheres*, 118, 10,884–10,906, doi: 10.1002/jgrd.50843, 2013.

Rupp, D.E., J.T. Abatzoglou, and P.W. Mote: Projections of 21st Century Climate of the Columbia River Basin. *Clim. Dyn.*, doi: 10.1007/s00382-016-3418-7, 2016.

Salathé, E. P., et al: Estimates of Twenty-First-Century Flood Risk in the Pacific Northwest Based on Regional Climate Model Simulations. *J. Hydrometeorol*, 15, 1881–1899, 2014.

Serinaldi, F., and C.G. Kilsby: Stationarity is Undead: Uncertainty Dominates the Distribution of Extremes. *Advances in Water Resources*, doi: 10.1016/j.advwatres.2014.12.013, 2015.

Sharma, A., C. Wasko, and D.P. Lettenmaier: If Precipitation Extremes Are Increasing, Why Aren't Floods? *Water Resources Research*, doi: 10.1029/2018WR023749, 2018.

Stewart, I. T., D.R. Cayan, and M.D. Dettinger: Changes Toward Earlier Streamflow Timing Across Western North America. *J. Climate*, 18, 1136–1155, 2005.

Surfleet, C. G., and D. Tullos, D.: Variability in Effect of Climate Change on Rain-on-Snow Peak Flow Events in a Temperate Climate. *Journal of Hydrology*, 479, 24-34, doi: 10.1016/j.jhydrol.2012.11.021, 2013.

Thober, S., Kumar, R., Wanders, N., Marx, A., Pan, M., Rakovec, O., Samaniego, L., Sheffield, J., Wood, E.F. and Zink, M., 2018. Multi-model ensemble projections of European river floods and high flows at 1.5, 2, and 3 degrees global warming. *Environmental Research Letters*, 13(1), p.014003.

Tohver, I., A. F. Hamlet, and S.-Y. Lee: Impacts of 21st Century Climate Change on Hydrologic Extremes in the Pacific Northwest Region of North America. *J. Amer. Water Resour. Assoc.*, doi: 10.1111/jawr.12199, 2014.

Vano, J. A., J. B. Kim, D. E. Rupp, and P. W. Mote: Selecting Climate Change Scenarios Using Impact-relevant Sensitivities. *Geophys. Res. Lett.*, 42, 5516–5525, doi: 10.1002/2015GL063208, 2015.

Wood, A., L. Leung, V. Sridhar, and D. Lettenmaier: Hydrologic Implications of Dynamical and Statistical Approaches to Downscaling Climate Model Outputs. *Clim. Change*, 62, 189–216, 2004.

Table 1 Information about locations featured in this paper - location, river, and discharge ratios

River	UW Key	Description	10-year flood discharge ratios				100-year flood discharge ratios			
			Avg.	Coeff. of Var.	Min	Max	Avg.	Coeff. of Var.	Min	Max
Chehalis	CHEGR	Chehalis R nr Grand Mount	1.21	0.09	1.0 ₃	1.42	1.34	0.18	0.87	2.07
Chehalis	CHE	Chehalis R at Porter	1.21	0.08	1.0 ₃	1.40	1.31	0.16	0.91	1.89
Willamette	SVN	T.W. Sullivan	1.33	0.09	1.0 ₇	1.64	1.39	0.22	0.87	2.39
Willamette	WILPO	Portland	1.34	0.09	1.0 ₈	1.69	1.40	0.23	0.86	2.47
Willamette	WILLA	Newberg	1.34	0.09	1.0 ₉	1.66	1.40	0.22	0.88	2.44
Willamette	SLM	Salem	1.37	0.09	1.1 ₀	1.70	1.43	0.22	0.84	2.52
Willamette	ALBO	Albany	1.40	0.09	1.11	1.73	1.47	0.20	0.89	2.40
Willamette	HARO	Harrisburg	1.45	0.10	1.1 ₈	1.86	1.50	0.22	0.88	2.37
Willamette	JASO	Middle fork @ Jasper	1.50	0.14	1.2 ₀	2.13	1.57	0.23	0.93	2.68
Willamette	DEX	Dexter	1.55	0.16	1.1 ₇	2.33	1.61	0.22	1.05	2.67
Willamette	HCR	Hills Creek	1.57	0.18	1.1 ₅	2.46	1.60	0.25	1.10	3.18
Willamette	WILNF	Oakridge	1.57	0.18	1.1 ₆	2.45	1.63	0.24	1.09	2.88

River	UW Key	Description	10-year flood discharge ratios				100-year flood discharge ratios			
			Avg.	Coeff. of Var.	Min	Max	Avg.	Coeff. of Var.	Min	Max
Willamette	EUGO	WR at Eugene (NWP)	1.50	0.12	1.26	2.04	1.54	0.22	0.88	2.57
Willamette	WAV	Walterville	1.54	0.13	1.29	2.13	1.55	0.18	1.04	2.23
Willamette	LEA	Leaburg	1.56	0.14	1.28	2.23	1.56	0.18	1.05	2.34
Willamette	VIDO	McKenzie nr Vida	1.57	0.15	1.28	2.32	1.58	0.19	1.02	2.41
Willamette	COT	Cottage Grove	1.25	0.11	0.97	1.69	1.39	0.29	0.78	2.38
Snake	IHR	Ice Harbor	1.20	0.13	0.92	1.75	1.26	0.28	0.79	2.84
Snake	LMN	Lower Monumental	1.20	0.13	0.92	1.76	1.26	0.28	0.78	2.77
Snake	LGS	Little Goose	1.19	0.13	0.92	1.77	1.26	0.28	0.78	2.83
Snake	LWG	Lower Granite	1.19	0.13	0.92	1.77	1.25	0.29	0.78	2.89
Snake	ANA	Anatone	1.24	0.14	0.95	1.74	1.29	0.29	0.78	2.84
Snake	LIM	Lime Point	1.23	0.14	0.94	1.73	1.28	0.30	0.76	2.81
Snake	HCD	Hells Canyon	1.40	0.18	1.01	2.11	1.55	0.38	0.87	3.62
Snake	OXB	Oxbow	1.41	0.18	1.01	2.11	1.56	0.38	0.86	3.65

River	UW Key	Description	10-year flood discharge ratios				100-year flood discharge ratios			
			Avg.	Coeff. of Var.	Min	Max	Avg.	Coeff. of Var.	Min	Max
Snake	BRN	Brownlee Dam	1.41	0.18	1.0 1	2.12	1.56	0.37	0.86	3.63
Snake	WEII	Weiser, ID	1.39	0.18	1.0 2	2.09	1.53	0.35	0.86	3.28
Snake	SNYI	Nyssa, OR	1.40	0.18	1.0 4	2.16	1.52	0.33	0.89	3.21
Snake	SWAI	Murphy, ID	1.37	0.19	0.9 8	2.09	1.48	0.33	0.84	3.24
Snake	CJSTR	CJ Strike Dam	1.37	0.19	0.9 7	2.08	1.48	0.32	0.86	3.08
Snake	SKHI	King Hill, ID	1.37	0.19	0.9 6	2.08	1.48	0.32	0.85	2.84
Snake	SNKBL WLSAL MON	Hagerman, ID	1.35	0.18	0.9 3	2.05	1.46	0.31	0.83	2.66
Snake	BUHL	Buhl, ID	1.35	0.19	0.9 1	2.05	1.46	0.32	0.73	2.54
Snake	KIMI	Kimberly, ID	1.33	0.19	0.8 9	2.03	1.44	0.33	0.74	2.47
Snake	MILI	Milner, ID	1.33	0.19	0.8 8	2.04	1.44	0.34	0.73	2.52
Snake	MINI	Minidoka, ID	1.33	0.19	0.8 6	2.02	1.45	0.33	0.70	2.53
Snake	AMFI	Neeley American Falls	1.32	0.19	0.8 5	1.99	1.45	0.34	0.67	2.69

River	UW Key	Description	10-year flood discharge ratios				100-year flood discharge ratios			
			Avg.	Coeff. of Var.	Min	Max	Avg.	Coeff. of Var.	Min	Max
Snake	BFTI	nr Blackfoot, ID	1.31	0.19	0.84	1.96	1.43	0.34	0.67	2.72
Snake	SNAI	nr Blackfoot, ID	1.30	0.19	0.84	1.95	1.43	0.34	0.67	2.69
Snake	SHYI	Shelley, ID	1.29	0.18	0.84	1.92	1.40	0.33	0.69	2.62
Snake	LORI	Lorenzo, ID	1.28	0.19	0.86	1.91	1.38	0.34	0.69	2.52
Snake	HEII	Heise, ID	1.28	0.18	0.86	1.91	1.37	0.33	0.70	2.53
Snake	PALI	Irwin Palisades	1.28	0.19	0.87	1.95	1.37	0.34	0.71	2.60
Snake	JKSY	Jackson, WY	1.26	0.15	0.89	1.73	1.35	0.30	0.80	2.46
Snake	SRMO	Moose, WY	1.25	0.13	0.91	1.59	1.35	0.25	0.83	2.34
Grand Ronde	TRY	Troy	1.48	0.19	1.09	2.55	1.68	0.34	1.01	4.38
Salmon	WHB	White Bird	1.07	0.13	0.83	1.57	1.09	0.33	0.72	2.81
Columbia	CRVAN	Vancouver	1.03	0.09	0.90	1.22	1.05	0.13	0.80	1.49
Columbia	BON	Bonneville	1.03	0.09	0.90	1.21	1.05	0.13	0.80	1.49
Columbia	TDA	The Dalles	1.03	0.08	0.90	1.20	1.05	0.13	0.81	1.52

River	UW Key	Description	10-year flood discharge ratios				100-year flood discharge ratios			
			Avg.	Coeff. of Var.	Min	Max	Avg.	Coeff. of Var.	Min	Max
Columbia	JDA	John Day	1.02	0.08	0.9 0	1.19	1.05	0.13	0.80	1.51
Columbia	MCN	McNary Dam	1.02	0.08	0.8 9	1.18	1.05	0.13	0.80	1.45
Columbia	CLKEN	Clover Island @ Kennewick	1.03	0.10	0.8 2	1.22	1.11	0.14	0.84	1.49
Columbia	CHJ	Chief Joseph	1.06	0.11	0.8 3	1.25	1.15	0.15	0.85	1.70
Columbia	GCL	Grand Coulee	1.06	0.11	0.8 3	1.25	1.14	0.14	0.84	1.66
Columbia	PRD	Priest Rapids	1.04	0.10	0.8 2	1.22	1.11	0.13	0.84	1.54
Columbia	WAN	Wanapum	1.04	0.10	0.8 2	1.22	1.11	0.14	0.84	1.58
Columbia	RIS	Rock Island	1.04	0.10	0.8 2	1.23	1.12	0.14	0.84	1.60
Columbia	RRH	Rocky Reach	1.05	0.10	0.8 3	1.23	1.13	0.14	0.84	1.61
Columbia	WEL	Wells Dam	1.05	0.10	0.8 3	1.24	1.14	0.14	0.85	1.63
Columbia	ARD	Hugh Keenleyside (Arrow)	1.13	0.12	0.8 7	1.43	1.24	0.21	0.69	1.83
Columbia	RVC	Revelstoke	1.19	0.12	0.9 1	1.62	1.36	0.23	0.69	2.08

River	UW Key	Description	10-year flood discharge ratios				100-year flood discharge ratios			
			Avg.	Coeff. of Var.	Min	Max	Avg.	Coeff. of Var.	Min	Max
Columbia	MCD	Mica Dam	1.22	0.12	0.9 4	1.66	1.41	0.24	0.72	2.12
Columbia	DONAL	Donald	1.28	0.14	1.0 2	1.79	1.55	0.25	0.94	2.38
Columbia	CRNIC	Nicholson	1.25	0.13	0.9 8	1.61	1.47	0.23	0.94	2.39
Clearwater	SPD	Spalding, ID	1.15	0.15	0.8 5	1.78	1.32	0.30	0.80	2.63
Clearwater	DWR	Dworshak Dam, ID	1.14	0.12	0.8 6	1.55	1.30	0.24	0.89	2.22
Santiam	JFFO	Santiam R nr Jefferson	1.40	0.10	1.1 4	1.81	1.41	0.25	0.81	2.27
Kootenay	COR	Corra Linn Dam, BC	1.08	0.12	0.8 5	1.31	1.15	0.16	0.79	1.67
Kootenai	LIB	Libby Dam, MT	1.17	0.14	0.9 2	1.52	1.32	0.22	0.85	2.01
Kootenay	BFE	Bonner's Ferry, ID	1.13	0.13	0.8 9	1.45	1.26	0.20	0.83	2.02
Pend Oreille	ALF	Albeni Falls, ID	1.26	0.14	0.9 6	1.68	1.65	0.30	1.02	2.97
Flathead	CFM	Columbia Falls, MT	1.24	0.13	0.9 4	1.63	1.65	0.26	1.01	3.19
Flathead	HGH	Hungry Horse Dam, MT	1.30	0.13	1.0 4	1.70	1.78	0.29	1.16	3.56
Yakima	KIOW	Yakima, WA	1.82	0.21	1.3 5	3.11	2.28	0.30	1.57	4.39

