

RE: Submission titled: “ ” by Dziubanski, Franz, and Gutowski

Dear Dr. Toth,

I am submitting the revised version of the above referenced manuscript for publication in the Journal of Hydrology and Earth System Sciences on behalf of all authors.

We have considered all the reviewer comments and have revised the original submission accordingly. We have expanded the references and background discussion to add studies from water resources systems analysis (WRSAs) (reviewer 1 comment 1), which is a precursor to socio-hydrology, and from other ABM applications in hydrology (reviewer 2 comment 1). We were careful to consider the scope and length of this paper when adding the additional references. A focused review paper on the evolution of ABM as applied in WRSAs and hydrology would be a wonderful addition to the literature, and the referencing we provide may give a basis for such a paper. We feel that the discussion we have provided gives sufficient background to place our study within context of prior works.

We strengthened the case for the ABM approach that we selected for our study (reviewer 3 comment 1). We also clarified sections that were not clear to the reviewers. In particular, we carefully reworded the section in the Methods that referred to Bayesian-based ABM (reviewer 1 comment 2). We perhaps were unintentionally giving the wrong impression about how we classify this model. We have added references on ensemble Kalman filter approaches that inspired the methods we used, but point out that this is not a true Bayesian approach.

A couple of the reviewers mentioned that they needed to refer to the supplementary material to better understand the model and results. We moved some of the information from the supplemental material to the main manuscript to provide a better explanation of the model processes in the areas that were deemed confusing by the reviewers (reviewer 2 comment 9, reviewer 3 comment 7). We also moved the calibration section to improve the flow of the paper as suggested by the reviewers.

In some cases, we did not change the manuscript as a result of the reviewer comments. Evaluation of additional model outcomes such as flood damage (reviewer 3 comment 4), sensitivity analysis of the individual modules (reviewer 1 comment 8, reviewer 3 comment 2), or detailed analysis of specific sections of time series data (reviewer 3 comment 9, reviewer 1 comment 10) would add length to the paper without significantly contributing to our overall objective. Because the model has not been published previously, considerable space was needed to adequately present the model. Rather than explore all possible outputs from the model, we have tried to focus the paper on the concepts and legitimacy of the modeling approach. In some cases, such as the questions about the crop model (reviewer 1 comment 8, reviewer 2 comment 7) and curve number (reviewer 3 comment 2), we rely on the efficacy of previously developed methods.

Thank you for your consideration of this revised manuscript.

Sincerely,

David J. Dziubanski, Corresponding author

Lines modified in response to Reviewer 1

Comment 1: lines 71-74

Comment 2: lines 403-404

Comment 3: Calibration section moved and lines 636-663 modified

Comment 4: lines 71-74

Comment 5: line 152

Comment 6: lines 245-246 and 249-250

Comment 7: lines 181-187

Comment 8: No change

Comment 9: lines 428-429

Comment 10: No change

Lines modified in response to Reviewer 2

Comment 1: line 97

Comment 2: lines 115-120

Comment 3: lines 152-153

Comment 4: no change

Comment 5: lines 278-279

Comment 6: lines 457-459

Comment 7: no change

Comment 8: lines 457-459

Comment 9: lines 468-493

Comment 10: lines 819-825

Lines modified in response to Reviewer 3

Comment 1: lines 89-95 and 827-834

Comment 2: no change

Comment 3: lines 760-762

Comment 4: no change

Comment 5: lines 542-544

Comment 6: lines 386-388

Comment 7: lines 468-493

Comment 8: lines 514-515

Comment 9: no change

Comment 10: Figure 10 corrected

Comment 11: lines 136-137 and 773-777

Comment 12: Calibration section moved

Reviewer #1

We thank the reviewer for these helpful suggestions to improve our manuscript.

This paper “Linking economic and social factors to peak flows in an agricultural watershed using socio-hydrologic modeling” develops a coupled agent-based model to evaluate the impact of conversion decision on flood reduction in a watershed. I think the scope fits quite well with the journal and the authors explain their goal and method reasonably well. I do have some comments which I hope can further improve the quality of the manuscript. I would recommend a minor to moderate revision.

1. First, I think the authors can benefit well by enlarging their literature review into the “water resources systems analysis” (WRSA) realm. The study of human-hydrologic cycle interaction started at the Harvard Water Program in the 1960s. A lot of classic issues (including the impact of land use, land cover change) had been addressed extensively in WRSA literature. Compare to “socio-hydrology,” WRSA also has a longer history of incorporating ABM into their modeling framework. I would strongly encourage authors to identify more literature on this aspect.

We thank the reviewer for this helpful comment. This paper was written with the emerging field of “socio-hydrology” strictly in mind. However, we realize that many other areas of water resources research have also utilized ABMs. We feel that the purpose of this paper is not to provide an elaborate and lengthy literature review. Most likely, one can probably write an entire review paper on the subject of incorporating ABMs and humans in water resources/hydrological analysis. We have cited some literature in the introduction (lines 71-74) to make readers aware of the WRSA field. Also, some of the studies that we cite on lines 97-100 do come from the WRSA field (e.g. Schlüter and Pahl-wostl, 2007).

2. Second, following my above comment, studies of ABM become more and more popular in the past decade. Methods used to quantify agents' behavior have been improved a lot as well. Methods proposed by the authors are not entirely new (Section 2.7.2, line 375-385) because it is a Bayesian-based method (the authors even use the terminology: prior and posterior). Authors are encouraged to broaden their literature about ABM that uses Bayesian theory to address behavior uncertainty. The authors should highlight the different settings they used in their ABM compare to other Bayesian-based ABM.

Lines 399-410: The approach that we are using is not a true Bayesian approach. We are not using bayes rule/conditional probabilities to update any sort of probability distributions. The farmer agents are simply using a weighted average formulation that includes a variable taking into account their past land use configuration and several variables taking into account new information such as profits or future crop price projections. This is similar to a data assimilation approach such as the EnKF were the past model state is given a certain weight and new observations are given a certain weight based on a computed Kalman gain. Hence, we point to this field in lines 399-408 to indicate where this idea came from. Many studies in ABM dealing with agriculture and water resource take an optimization approach (e.g. Schreinemachers, P., Berger, T., 2011. An agent-based simulation model of human–environment interactions in agricultural systems.) or a rule-based approach (e.g. van Oel, P.R., Krol, M.S., Hoekstra, A.Y., Taddei, R.R., 2010. Feedback mechanisms between water availability and water use in a semi-arid river basin: A spatially explicit multi-agent simulation approach). We point out the different types of models used in the paragraph on lines 386-398. Some studies do use Bayesian methods, but these methods are usually paired with the main decision model (e.g. optimization). Ng et al 2011, “An agent-based model of farmer decision-making and water quality impacts at the watershed scale under markets for carbon allowances and a second-generation biofuel crop”, is one such study that uses Bayesian updating for updating farmer’s perceptions of variables such as yields or crop prices.

We have updated the sentence on lines 403-404 so as not to confuse readers into thinking that we are using Bayesian methods.

3. Third, I do have a suggestion about paper structure. Currently, the authors put the ABM calibration in Section 5 which reads weird to me. The purpose of calibration and validation of the model is to demonstrate the credibility, therefore, it should be put before the authors use the model for any scenarios. I would suggest move Section 5 before the results. And add more discussion about ABM validation (beyond line 711-720) because this topic is the most popular issue in the ABM community nowadays.

This section has been moved prior to the results. Some further discussion and literature has been added in the paragraphs between lines 636 and 663.

I have some minor comments below:

4. Line 71-73: This kind of argument really needs to incorporate the studies of Water Resources Systems Analysis.


See comment above.

5. Line 137: You mean two “types” of agents?

Line 152: That is correct. The terminology in the manuscript has been changed from “primary” to “types” so as to make this more clear.

6. Line 223-224: This does match with your equation (7) to (9) because I did not see minimize flood damage objective function. Also, why the goal of the city agent is not "minimize the cost = flood damage + contact fee?

Lines 245-246: In the current version of the model, a stronger focus was placed on capturing the various decision variables that farmers may take into account, whereas the decision-making of the city agent was kept rather simple. So the city agent isn't “minimizing” flood damage using an objective function with numerical optimization, but rather the city agent is trying to reduce flooding based on a simplified linear equation, displayed below. Flood damage is computed based on a sigmoid relationship plotted below (left). This is described in detail in section S7 of the supplemental material. The city agent then takes this flood damage and computes a new conservation goal (amount of new land that the city agent would like to convert to conservation as a percentage of the total watershed area) based on the linear relationship plotted below (right).



The city agent is a feature that will be improved in further iterations of the model. Introducing a cost function such as “cost = flood damage – flood reduction + contract_fee” is viable; however, this would require the city agent to have capability to simulate specific flood events in order to estimate flood reduction for a given

amount of conservation implementation. We have updated lines 245-246 and lines 249-250 to make it clear that it's not a minimization via an objective function.

7. "Line 229: I think this is the first time you mention risk-aversion. You need a more detailed description of what does it mean in your model.

Risk-aversion indicates the willingness of a farmer to change his/her land use under uncertainty. Farmers with a high risk aversion will not want to change their previous land use because they are trying to avoid risk (keeping their land use the same represents a more predictable payoff to the farmer, even though their revenue may be smaller). Lines 181-187 were added to clarify this prior to the paragraph containing line 229 (now line 251).

8. Line 337: Since FAO has a physically-based crop model, you might want to test the sensitivity of the current crop model on your results given that this will affect farm agents' decisions.

Line 358: The crop yields in our model are computed using a robust regression model that was formulated using temperature, precipitation, and yield data from 1960-2006. This model gives a reasonable prediction of yields based on environmental conditions. Unlike a physically-based model, there is no feasible way of testing the sensitivity of a regression-based model. We are not changing any of the values associated with specific regression coefficients. We do however take into account differences in yield based on soil types and add stochastic variability based on local differences in environmental conditions.

We thank the reviewer for this good suggestion. One of the goals for the future is to improve the crop model by introducing a physically-based crop model. This will allow us to simulate yields in more detail based on finer level farm management techniques.

Listed below is the reference for the crop regression model that is currently used. Tannura, M. A., Irwin, S. H. and Good, D. L.: Weather, Technology, and Corn and Soybean Yields in the U.S. Corn Belt. [online] Available from: https://farmdoc.illinois.edu/assets/marketing/morr/morr_08-01.pdf, 2008.

9. Line 402: How you define "neighbor?"

Line 427-428: If a farmer is located in subbasin A for example, he/she can make a certain random number of neighboring connections with other farmers in that same subbasin. A sentence was inserted at lines 428-429 to clarify the above. If a subbasin contains 10 farmers, one farmer might form 5 neighboring connections with farmers in that same subbasin while another farmer may form only 2 connections. This process is described in greater detail in section S3 of the supplement.

10. Line 564-Figure 6d: Why is there a jump in all three curves around 2012? The same question for Figure 8d. I hope these comments help the authors for their revision.

In the scenarios plotted in figures 6d and 8d (now figures 8d and 10 d), the farmer is placing an 85% weight on the future price decision variable ($\delta C_{futures:Y}$). Most likely what is happening in this case is that prices are high during 2010-2013, but crop price forecasts are predicting a down turn in crop prices. If farmer agents are considering crop price forecasts several years into the future, conservation land begins increasing while crop prices are still high during 2012 and 2013. Essentially, the farmers are changing their land use in anticipation of lower crop prices. The increase in conservation land will be more dramatic considering that farmer agents are placing such high weight on the future crop price forecasts. In this section of the paper, we are trying to discuss more broadly about what happens under low and high crop prices. This is a very specific question about a small portion of the time series, thus we feel that describing these details does not fit with this section of the paper.

Reviewer #2

We thank the reviewer for comments to help improve this manuscript.

This paper presents a social-hydrological model on the links between farmers' land use decisions and flood consequences at a basin scale, which is a novel effort in this field. The research question, methods and data are well introduced, and the results are valuable. I recommend to consider this paper for publish after minor revisions.

A few detailed comments are listed here, which are hopefully helpful for improvement of the paper.

1. Page 4, line 87-90, the authors listed literature of ABM on water issues, but the references are quite old from the years of 2003-2011 and only one from 2017. In addition, there is a lack of reference about ABM in flood studies. I suppose it would be valuable if the authors read some recent papers:
 - Aerts, J.C.J.H., Botzen, W.J., Clarke, K.C. et al. Integrating human behaviour dynamics into flood disaster risk assessment. *Nature Clim Change* 8,193–199 (2018)
 - Liang E. Yang, Jürgen Scheffran, et al., 2018. Assessment of Flood Losses with Household Responses: Agent-based Simulation in an Urban Catchment Area. *Environmental Modelling and Assessment*, 2018, 23(4):369-388
 - Ahmed Mustafa, et al., 2018. Effects of spatial planning on future flood risks in urban environments. *Journal of Environmental Management*, Volume 225, 1 November 2018, Pages 193-204
 - Omar S. Areu-Rangel, et al., 2019. Impact of Urban Growth and Changes in Land Use on River Flood Hazard in Villahermosa, Tabasco (Mexico). *Water* 2019, 11(2), 304

ABMs have been used in a variety of fields within hydrology. We list studies that are more closely related to the agriculture field, but realize that including some citations that use ABMs for studying floods would be good. Thus, we have modified lines 97 to include a part that mentions ABM use for flood studies.

2. Page 4-5, line 94-102, there are already many research about land use changes impacts on stream flow, what's the new value added by the present study?

Line 103-105: The reviewer is correct in that there is a large body of research examining historical changes in streamflow from land use and climate change. However, most of these studies use more traditional methods to analyze the effects of land use change on hydrology. This study, however, attempts to highlight the use of socio-hydrology and Agent-Based modeling for surface water hydrologic investigations related to land use change. We are trying to show that hydrologic changes in the system can potentially be tied to external economic variables and characteristics of the population (i.e. farmers in this case) residing in that watershed, which is difficult to show with the more traditional methods. Lines 115-120 were modified to highlight the difference between this study and the other studies mentioned.

3. Line 137, it would be more accurate to say “a group of farmer agents and city agents”. Is there only one city agent?

Line 152-153: There is only one city agent, but there are 100 farmer agents in the current simulations. We have corrected this line to say “a group of farmer agents”.

4. ABM proceed with monthly time steps. Hydrological model proceeds with hourly timestep. How are they integrated?

The hydrologic model proceeds in hourly timesteps because it needs to be able to simulate a flood discharge event. Most flood events in the Midwest U.S are associated with intense rainfall, thus it is critical to simulate those events on a fine scale. Every year, the city agent then computes the flood damage based on the maximum discharge event for that year (stated on lines 233-236). The farmer agents may change their land use on an annual basis (in the spring before the growing season). When land use changes occur, CN values are updated within the hydrology module. Those are the two main integrations between the hydrology and the agent modules. The above is described in detail in section 2.2.

5. Section 2.4.2, Line 255-257, why is it assumed that most farmers rent lands for crop production? Do they rent yearly, or how long usually is their land rent contract? I am not sure, but I assume many farmers own lands in the US.

Line 277-280: Majority of the land in the Midwest, particularly in the CornBelt (Eastern Nebraska, Iowa, Illinois, Indiana) is rented, according to the latest Tenure, Ownership, and Transition of Agricultural Land Survey published by the USDA:

**https://www.nass.usda.gov/Publications/Highlights/2015/TOTAL_Highlights.pdf
<https://www.extension.iastate.edu/agdm/wholefarm/html/c2-78.html>**

In fact, 59% of the land is rented in Iowa according to the Iowa Farmland ownership and tenure survey (Table 3.3 in the document – cash rent lease + crop share lease): <https://www.card.iastate.edu/farmland/ownership/FM1893.pdf>

The NASS TOTAL survey puts this number at 80% rented.

The rental contracts are on an annual basis. Average rental rates for Iowa over the last few years have been around \$220-240/acre (\$543-592/hectare).

We modified lines 278-279 to state that a majority of land is rented and have included a reference.

6. Nice to see the many historical data in Figure 3. Are these data used as input in the ABM, and how?

All of the input data in section 2.4 are used as input into the model. Corn prices, crop production costs, land rental values (cash rent), and federal government subsidies are used by the farmers for calculating crop revenue. Section 2.7.2 of the manuscript and particularly section S4 of the supplement describe how these inputs are used. The city agent bases the conservation subsidies on the land rental values. A sentence was added to lines 457-459 stating that costs and revenue of crop production and conservation land are computed based on these inputs.

7. Equation 1, yield is the Arithmetic function of year, precipitation and temperature. I may not understand this correctly. How could you add year, precipitation and temperature together? They don't even have the same unit. How would you explain the equation, e.g. $2200 \text{ MT/ha} = 2005 \text{ year} + 160\text{mm} + 35^\circ\text{C}$?

This equation is a multiple regression between temperature, precipitation, and yield for the years 1960-2006). Thus, it's not like a physical equation (e.g. bernoulli's equation, darcy's law, etc) where the units have to work out.

8. Section 2,7,1, taking conservation option means farmers have to plant native prairie strips. Did the authors consider the costs of planting the prairie strips? How much is it less than planting crops?

Yes, the farmer agents do consider the cost of planting native prairie. Section S4 of the supplement describes how the cost of native prairie strips is incorporated into the farmer agent revenue calculation. Section S1.2 describes in detail the actual cost of native prairie strips, which include establishment costs, continual maintenance costs, and opportunity costs (cost of forgone revenue from the land taken out of crop production). The cost of planting crops is significantly higher than the cost of planting prairie. For example, in 2010, the cost of planting prairie was ~\$216/acre for high quality land (e.g. land that produces the highest crop yields) versus ~\$422/acre from crops (not including land rent). However, with crops, farmers can make significantly more money than they would from conservation subsidies. So in 2010, the farmers had a net revenue of ~ +10\$/acre from crops versus a net revenue of ~ -\$11/acre from prairie. Some years, the net revenue from crops may be -\$40/acre to -\$70/acre (i.e. a loss), whereas other years that net revenue may be +\$150/acre. Of course, all of these numbers vary from farmer to farmer based on quality of land, which influences the cash rental rate, and other stochastic variability. A sentence was added to lines 457-459 briefly stating that costs of

conservation are considered. This sentence also refers the reader to the supplement for further information.

9. How did you quantify the many “ δC ”s in equation 2? There is only one introduction of calculating $\delta C_{\text{profit}:X}$ (equation 4).

At the request of the editor, we moved a significant portion to the supplementary material to make the manuscript more readable. Where needed, we point the reader to the supplement. For instance, line 459 points to supplement section S4 for a more detailed description of how Profit_{diff} is computed. The only difference between $\delta C_{\text{profit}:X}$ and $\delta C_{\text{futures}:Y}$ is the use of realized crop prices versus the use of projected future crop prices. Both variables use the same set of equations. We modified lines 468-493 to provide more detail for the profit function. We also moved figure 4 from the supplement back to the main manuscript to describe the profit function in more detail. Where needed, we point the reader to the supplement for further information.

10. The major finding of the study indicates that peak discharge is most sensitive to changes in crop prices. As we know generally crop price is a market effect and is not controllable by the farmer and city agents in the basin scale. Thus three questions:
- Would you say that crop price in the US or in the globe influences water discharge (flood) in the Squaw Creek basin?
 - Does it mean that local efforts in the basin have little effects regarding flood control?
 - What would you suggest for decision making of flood management in the Squaw Creek basin, based on your research findings?

We thank the reviewer for these insightful questions. We know that local decisions influence the local land use, which along with other physical characteristics of the landscape, ultimately influence discharge outcomes. Our model demonstrates that external factors can also influence local streamflow, albeit in a complex and unpredictable way as the information gets filtered through the complex decision making of local farmers. Social factors (local or external) introduce significant uncertainty in local hydrology outcomes, and by ignoring them, water management plans will be inherently incomplete. Thus, we recommend that multi-scale human factors be explicitly considered when assessing the sustainability of long-term management plans. Some of this description was added to the conclusions lines 819-825.

Reviewer #3

We thank the reviewer for comments to help improve this manuscript.

This is an interesting paper that explores the feedbacks between land-use management changes, driven by economic and social factors, and the hydrological system of an intensively managed agricultural watershed. An agent-based model ABM considers two agent types: a reactive city agent (providing financial incentives for agricultural conservation practices based on prior flooding impacts) and farmer agents that maximize profits, subject to conservation and risk-aversion attitudes. A simple semi-distributed model is used to simulate the hydrological system, perturbed by farmer agent conservation practices through changes to the SCS-CN parameter. The purpose of the modelling is to bring understanding of the impacts of land use decisions on downstream flood impacts. The manuscript reads reasonably well, although some key details regarding the ABM are omitted (description of equation 2) and the scenario outcomes could be made more succinct.

Remarks:

1. Paragraph starting line 71: The authors need to make a stronger case for the use of ABM over, for example, top down approaches using differential equations. This should be followed up in the conclusions.

A section describing the advantage's of ABMs was added in the intro – lines 89-95. This was followed up with a short paragraph in the conclusions – lines 827-834.


2. Comment [Section 2.5]: The ability to reproduce an historical set of flood peaks does not indicate a model has the correct sensitivity for impact assessment (i.e. the selection of CN parameters). However, it is recognised that the method employed is typical of operational practice, and limitations are recognized in the conclusions.

Thank you for this comment. We realize that the current methods employed in the model do have limitations. As is stated, we did try our best to follow typical operational methods. Future work may involve coupling a more advanced hydrologic model that is better able to capture fine-scale processes such as runoff.

3. Paragraph starting line 616: I am unclear how much feedback between there is between the hydrological system and the city agent. The conservational goal of the city (Eq. 7) appears unresponsive to flooding with a value of ~1,100 Ha, after the initial introduction of the conservation scheme.

Paragraph line 754: The city agent calculates a yearly flood damage amount based on the maximum peak discharge for that year (Supplement S7). The flood damage is then used in a linear function to calculate a new conservation goal, which is added to any outstanding conservation goal that was not fulfilled during previous years. Below is plotted the time trend in the conservation goal of the city agent. As you can see, it directly follows the amount of conservation land implemented (see figure 6 in the manuscript). The reason why it may seem like the conservation goal is unresponsive beyond 1100 Ha is because the two

largest flooding events of the entire time series occur in 1975 and 1977. Based on the 1% and maximum flood damage levels set in the model (based on the 10 year and 100 year discharge values, Supplement S7), 1975 flood damage was 42% higher than maximum, while the 1977 flood damage was 72% of the maximum damage. The threshold of minimum flood damage is exceeded 13 times through the entire simulation period, but 11 of those times, damage is less than 30% of maximum, and 7 of those times damage is less than 15% of maximum. Thus, a lot of conservation land isn't being added to the city conservation goal beyond those two flood events. The year 1990 saw a flood event 30% of maximum, which corresponds to a small increase in conservation land (g and h below show this clearly). A similar increase occurs in 2009 in response to a flood event in 2008 that was 24% of maximum damage. Lines 760-762 were added to clarify this. One thing that is clear in this figure is that the city goal is only being fully implemented by the conservation-minded farmers (green) when they place 85% weight on their conservation decision variable (a, b). The amount of land implemented by this group is ~1100 Ha, which corresponds to the approximate goal of the city agent after the 1975 and 1977 floods.



4. Line 113: The results are presented in form of reductions to peak flows [i.e. this represents land use decision making]. However, it may have also been interesting to evaluate reductions in flood damages, given that this is the objective of the city agent [line 223].

While this would be an interesting analysis, it is difficult to compute a reduction to peak flow within a single simulation using a historical time series of precipitation. One would have to have identical precipitation events at different times throughout the simulation period to compute an estimate of reduction in flood damage. However, one interesting analysis that can be done is examining the flood damage reductions for a single event based

on the types of farmers (conservationist or productionist) in the watershed. Just examining the large flood event in 1977 (which follows the large increase in conservation land in response to the 1975 flood), flood damage is 17% lower (~\$8000000 reduction) under conservationists versus productionists when farmers are placing most weight on their past profit variable. For the 2008 flood event, flood damages are reduced by ~\$2400000, or 4.8% of the maximum damage. However, this analysis does not get at the flood reduction within a single simulation. Given the length of the paper and the fact that the focus is on how changes in economic variables may change discharge, we decided not to include this discussion. The figures would also be similar to the figures showing reductions in the 90th percentile discharge, except that the plots would be of flood damage.

5. In Figs 5-8, results are presented as reductions in the “mean 90th percentile discharge”. I am unclear what this represents and why this value was chosen (is this the peak discharge or taken from a flow duration curve).

This value was taken from a flow duration curve. It represents the 0.1 exceedance probability level. This value was simply chosen because we wanted to examine changes to the larger discharge events. Another option, for example, would’ve been to examine changes to the average maximum peak discharge. However, there are some significant flows in the record, so we don’t want results to be skewed based on a few large events. Lines 542-544 were added to clarify this.

6. Line 363: “Rules governing agent decision making need to realistically capture human behaviour without creating an excessively complex model”. Could the authors make some comment on this highly parameterized ABM in this regard? Also I note in the conclusions that there are arguments for introducing further decision processes /state variables.

Line 386: This is a highly complex question that could be extensively discussed beyond a manuscript. Using the ABM approach, you will end up with a larger set of parameters because ultimately, trying to describe human behavior is not simple. This is an issue that cross cuts multiple disciplines such as the natural and social sciences. In hydrology for instance, there is a similar discussion of trying to keep models fairly simple, but still complex enough to realistically simulate hydrologic processes. Do we need to capture every single micro process in detail such as hillslope runoff patterns, or is it sufficient to estimate these processes on a larger scale? So this question does not have a simple answer – it can be extensively debated. To simplify the manuscript, we have decided to eliminate that statement as the main focus of that paragraph is on the different decision models that are used in ABM and not how complex ABMs should or should not be.

7. Section 2.7.1, Farmer agent land use decision process. I found this difficult to follow without referring to the Supplementary Material. The first two paragraphs [p 18] could be moved to a methodology section. The rationale for Eq 4 (the profit function) with the use of a polynomial is unclear in the main manuscript, and I also wonder whether this is a representation of the cognitive process of the farmers (e.g. line 697 states for arise of \$1 in corn prices, 10-15% of land is converted back into production and line 354 “Either 10% or 20% of the total field size is

converted into native prairie vegetation” – these seem to provide a more appropriate basis for forming a profit rule).

At the request of the editor prior to the review process, we moved a significant portion of this section to the supplementary material to make the manuscript more readable. However, we realize that some more detail in the manuscript is necessary. The statement on line 697 (now line 487-490) is precisely why we introduce a profit rule. It’s meant to capture the changes that occur in conservation land with changes in crop price. It also allows for variation between farmer agents since different farmers have difference crop production costs and crop yields. We modified lines 468-493 to provide more detail and rationale for the profit function. We also moved figure 4 from the supplement back to the main manuscript to describe the profit function in more detail. Where needed, we point the reader to the supplement for further information on the polynomial in equation 4. As of the two paragraphs that were on page 18 (now page 19, introduction to section 2.7.2), we feel that leaving this material with this section provides the best flow. Otherwise, transferring it into its own section with further subdivide a paper that already has a lot of sections/subsections.

8. The catchment area is ~56,200 Ha (line 426), but 100 farmers are allocated 121 Ha each (line 485). Eq 7 uses A_{tot} , the total land area of the catchment. This appears inconsistent.

Line 514-515: This has been corrected. A_{tot} actually represents the total agricultural land in the watershed, or as we wrote in the manuscript, “the total land area owned by the farmer agents”. With 100 farmer agents owning 121 Ha each, the total land area operated by these farmers is 12100 Ha. Currently in the model, we scale up the effects of these farmer agents to the size of the watershed. So with 100 farmer agents and 14 subbasins, there are approximately 7 farmer agents controlling the land use in each subbasin. Even though there is currently a mismatch in the land area, we are dealing with percentage of different landcover types. So we are assuming that if each of the 7 farmers has 20% land in native prairie, then 20% of the land in that subbasin is in native prairie. We chose 100 farmer agents because, based on initial modeling testing, we found that 100 farmer agents is a sufficient population to represent variability and interactions in the watershed while reducing computational time. We would actually need ~460 farmer agents to match the land area in the watershed if each farmer agent owns 121 Ha.

9. I am not clear on the large fluctuations in land area for example in Figs 6 c,d and Figs 8 c,d, prices appear more volatile in years after 2000.

In the scenarios plotted in figures 6d and 8d (now figure 8d and 10d), the farmer is placing an 85% weight on the future price decision variable ($\delta C_{futures:y}$). Most likely what is happening in this case is that prices are high during 2010-2013, but crop price forecasts are predicting a down turn in crop prices. If farmer agents are considering crop price forecasts several years into the future, conservation land begins increasing while crop prices are still high during 2012 and 2013. Essentially, the farmers are changing their land use in anticipation of lower crop prices. The increase in conservation land will be more dramatic

considering that farmer agents are placing such high weight on the future crop price forecasts. In this section of the paper, we are discussing more broadly about what happens under low and high crop prices. Since this is a specific question about a small portion of the time series, we feel that describing these details does not fit with this section of the paper.

10. Fig 8: Caption “Yearly crop yields are plotted as bars”; crop prices are displayed on figure.

Thank you for catching this error. The plot has been corrected to show crop yields.

11. The historical scenario needs to be more clearly defined (line 122 and 635) for the interpretation of fig 9 in the Historical Comparison. I interpret this as prices/yield/subsidies use the input time series in the historical scenario (Section 4.2), but land cover can change on an annual basis according to the model.

Under the historical scenario, the farmer agents continue to make annual land use decisions (i.e. land use between crops and conservation can change on an annual basis), but corn prices, conservation subsidies, or yields are not modified like in the other scenarios. This scenario is simply using the historical observed data.

Line 136-137 was modified as such: “Additionally, we simulate land use and hydrologic outcomes for the historical period without any perturbations to these economic data for comparison purposes.”

Lines 773-777 were modified as such: “To gain an understanding of how each of the scenarios differs from the historical 1970-2016 period, the mean peak discharge is compared against the historical scenario (Figure 9). Recall that under the historical scenario, farmer agents make annual land use decisions as in the other scenarios, but corn prices, conservation subsidies, and crop prices are unchanged from historical observed values.”

12. Section 5 Model Calibration and Validation: This should be moved prior to the results section. It is interesting to note that conservation area was more sensitive to crop prices in the mid-1990s than in 2010s, despite the higher price volatility in the latter.

Thank you for this suggestion. The calibration section has been moved.

Linking economic and social factors to peak flows in an agricultural watershed using socio-hydrologic modeling

David Dziubanski¹, Kristie J. Franz¹, William Gutowski¹

¹Department of Geological and Atmospheric Sciences, Iowa State University, Ames, IA

Correspondence to:

David Dziubanski

2027 Agronomy Hall

Iowa State University

Ames, IA 50011

dave.dziubanski@gmail.com

Abstract: Hydrologic modeling studies most often represent humans through predefined actions and fail to account for human responses under changing hydrologic conditions. By treating both human and hydrologic systems as co-evolving, we build a socio-hydrological model that combines an agent-based model (ABM) with a semi-distributed hydrologic model. The curve number method is used to clearly illustrate the impacts of landcover changes resulting from decisions made by two different agent types. Aiming to reduce flooding, a city agent pays farmer agents to convert land into conservation. Farmer agents decide how to allocate land between conservation and production based on factors related to profits, past land use, and willingness. The model is implemented for a watershed representative of the mixed agricultural/small urban area land use found in Iowa, USA. In this preliminary study, we simulate scenarios of crop yields, crop prices, and conservation subsidies along with varied farmer parameters that illustrate the effects of human system variables on peak discharges. High corn prices lead to a decrease in conservation land from historical levels; consequently, mean peak discharge increases by 6%, creating greater potential for downstream flooding within the watershed. However, when corn prices are low and the watershed is characterized by a conservation-minded farmer population, mean peak discharge is reduced by 3%. Overall, changes in mean peak discharge, which is representative of farmer land use decisions, are most sensitive to changes in crop prices as opposed to yields or conservation subsidies.

1. Introduction

Humans change the water cycle through actions that affect physical and chemical aspects of the landscape, and these changes occur from global to local scales and over varying time periods (Vorosmarty and Sahagian, 2000). Despite their significant impacts to the landscape, humans remain the most poorly represented variables in hydrologic models (Sivapalan et al., 2012). Land cover and land use are commonly treated as fixed in time in many hydrologic models through the use of static parameters. When made dynamic, landscape change is often limited to predefined scenarios that are developed without consideration of how economics, local culture, or climate may combine to influence land use decisions. For example, the field of integrated water resources management (IWRM), which attempts to explore the interactions between humans and water, typically uses “scenario-based” approaches (Savenije and Van der Zaag, 2008). While scenario-based studies allow quantification of the impacts of a management decision on the hydrologic system, there are significant limitations (Elshafei et al., 2014; Sivapalan et al., 2012). Human and environmental systems are highly coupled with feedbacks from one system creating stress on the other system, which in turn affects the behavior of the first system. Therefore, representing management decisions as pre-determined will not reproduce the real-world variability that may arise as a result of complex feedbacks between the human system and the physical system.

Arguments have emerged for socio-hydrological in the hydrological sciences and Water Resources Systems Analysis (WRSAs) fields for modeling in which humans and the environment are treated as co-evolving (e.g., Di Baldassarre et al., 2013; Brown et al., 2015; Montanari, 2015; Rosengrant et al., 2002; Sivapalan et al., 2012; Sivapalan and Blöschl, 2015; Wainwright, 2008). In this way, models can account for disturbances to natural systems by humans and

simultaneously assess physical processes and economic and social issues. In the hydrologic literature, two approaches have been used to simulate coupled human and natural systems: a classic top-down approach and a bottom-up approach using agent-based modeling (ABM). In the first approach, all aspects of the human system are represented through a set of parametrized differential equations (e.g., Di Baldassarre et al., 2013; Elshafei et al., 2014; Viglione et al., 2014). For example, Elshafei et al. (2014) characterizes the population dynamics, economics, and sensitivity of the human population to hydrologic change through differential equations to simulate the coupled dynamics of the human and hydrologic systems in an agricultural watershed. In contrast, the ABM approach consists of a set of algorithms that encapsulate the behaviors of agents and their interactions within a defined system, where agents can represent individuals, groups, companies, or countries (Axelrod and Tesfatsion, 2006; Borrill and Tesfatsion, 2011; Parunak et al., 1998). System agents can range from passive members with no cognitive function to individual and group decision-makers with sophisticated learning and communication capabilities. The ABM approach has several advantages over the traditional top down approach (Bonabeau, 2002). Agent-based models are able to capture emergent phenomenon that result from interactions between individual entities. In addition, simulating individual entities through ABM provides for a more natural description of a system in contrast to developing differential equations that capture the behavior of the system as a whole. ABMs also provide for greater modeling flexibility by allowing for different number of agents, various degrees of agent complexity, and behavioral differences among the agents. ABM has been used to study the influence of human decision making on hydrologic topics such as water balance and stream hydrology (Bithell and Brasington, 2009), flooding (Du et al., 2017; Jenkins et al., 2017; Yang et al., 2018), irrigation and water usage (Barreteau et al., 2004; Becu et al., 2003; Berger et

al., 2006; Berglund, 2015; van Oel et al., 2010; Schlüter and Pahl-wostl, 2007), water quality (Ng et al., 2011), and groundwater resources (Noel and Cai, 2017; Reeves and Zellner, 2010).

A dominating topic in the hydrologic sciences that can be studied through use of ABMs is the issue of land use change impacts on hydrologic flows in intensively managed agricultural landscapes (Rogger et al., 2017). A number of studies have attempted to quantify the impact of land use change on streamflow (Ahn and Merwade, 2014; Frans et al., 2013; Naik and Jay, 2011; Schilling et al., 2010; Tomer and Schilling, 2009; Wang and Hejazi, 2011) Ahn and Merwade (2014) is one such study that found that 85% of streamflow stations in Georgia indicated a significant human impact on streamflow. Another study by Schilling et al., (2010) indicated a 32% increase in the runoff ratio in the Upper Mississippi River basin due to land use changes, mainly due to increases in soybean acreage. Results of Wang and Hejazi (2011) are consistent with Schilling et al., (2010). They found a clear spatial pattern of increased human impact on mean annual streamflow over the Midwestern states due to increases in cropland area.

~~Given clear evidence that the human system has a significant effect on streamflow, we use a social hydrologic modeling approach to better understand the effects of land use changes driven by economic and human behavior on hydrologic responses, which would be otherwise difficult to observe with a hydrologic model alone. The above studies use more traditional methods such as hydrologic modeling, trend analysis, or Budyko analysis to determine the impact of land use change on streamflow. We use the social-hydrologic modeling approach to better understand the effects of land use change. Using ABMs may allow for a more in-depth investigation of hydrologic changes and how they may be tied to external economic variables and watershed population characteristics.~~

In this study, we develop a social-hydrologic model that simulates changes in conservation land area over time within an agriculturally-dominated watershed as a function of dynamic human and natural factors. Using a sensitivity analysis approach, we use this model to quantify the impact of economic and human factors on land use changes relating to conservation implementation and subsequently, how these land use changes impact the hydrologic system. We explore the following research questions:

- 1) To what degree do economic and agronomic factors (specifically crop prices, conservation incentives, and crop yields) impact the success of a conservation program designed to reduce peak flows?
- 2) To what degree are hydrologic outcomes sensitive to various factors that commonly influence agricultural land use decisions?

Using simulations of a historical 47 year period, we explore land use and hydrologic outcomes for a typical agricultural watershed in Iowa under the following six scenarios developed from economic data: crop yields 11% above and below historical values, corn prices 19% above and below historical values, and conservation subsidy rates 27% above and below historical cash rent values. Additionally, we simulate [land use and hydrologic outcomes for](#) the historical period without any perturbations to [these](#) economic data for comparison purposes. The following model methodology is described using the ODD (Overview, Design Concepts, and Details) protocol developed by Grimm et al. (2006).

2. Model Purpose

The purpose of the model is to understand the impact of land use decisions by upstream farmers on flooding response in a downstream urban area under perturbations to extrinsic economic and natural factors (e.g. crop prices, land rental values, climate), as well as intrinsic

factors (e.g. internal farmer behavior, local government incentives). System behavior under changes in extrinsic and intrinsic factors is analyzed using a scenario-based ensemble approach.

2.1 State Variables and Scales

The model links an agent-based model of human decision making with a rainfall-runoff model to simulate social and natural processes within highly-managed agricultural watersheds (Figure 1). The agent-based model consists of two types of agents: a group of farmer agents and a city agent.

The primary modeling domain consists of the watershed and the subbasins located within the watershed. The model user must define the subbasins based on external analyses of hydrologic flows and conditions. Each subbasin is populated by one or more farmer agents as specified by the user. A farmer agent modifies the land use of the subbasin in proportion to the subbasin area assigned to that agent. The most downstream subbasin in the watershed is populated by an urban center, which is represented by a city agent. The city agent impacts land use by providing subsidies to upstream farmer agents to change his/her land management.




Figure 1. Flow of information within the agent-based model.

2.1.1 Farmer agent state variables

The primary state variable for a farmer agent is the conservation parameter ($Cons_{max}$), which characterizes the degree to which a farmer agent is “production-minded” versus “conservation-minded”. This concept is based on McGuire et al. (2013) who identified that US cornbelt farmers tend to fall along a spectrum from purely productivist to purely conservationist. $Cons_{max}$ is randomly assigned to each farmer agent upon initialization and provides variation in farmer agent behavior based on how an individual agent may prefer to balance maximizing crop yields versus protecting the environment. $Cons_{max}$ represents the maximum fraction of land a farmer is willing to put into conservation. The minimum value is 0.0, in which case a farmer is purely production-minded and is unwilling to convert any

production land into conservation. We set the maximum value at 10% ($Cons_{max} = 0.10$) based on the conservation practice used in this study (Section 2.7.1). Therefore, a farmer is purely conservation-minded at a parameter value of 0.1, and is willing to convert up to 10% of his/her production land into conservation. This range of values corresponds to the percentage of conservation land implemented over each of the last ten year for the entire state of Iowa (~5-6% conservation land) and the Central Iowa Agricultural District (~3-4% conservation land).

A secondary state variable of importance to the farmer agent is risk aversion attitude (Prokopy et al., 2019). Risk aversion can be defined as the willingness to change land use under uncertainty. Farmers with a high risk aversion are unwilling to change their land use because they are trying to avoid risk. Keeping their land use consistent represents a more predictable payoff, even if the revenue may not be as great as another land use choice. Farmers that are more risk tolerant however, are more likely to adopt new practices such as conservation.

Farmer agents are further characterized by their decision-making preferences, which describe the relative importance that farmer agents place on different decision variables when adjusting their land use. The farmer agent decision characteristics are described in Sect. 2.7.2.

Each farmer agent is assigned state variables characterizing the percent of different soil types associated with the farmer's land. Corn crop productivity and crop production costs (including the land rental value) vary for each soil type. Thus, the soil types associated with a farmer agent's land impact his/her revenue.

2.1.2 City Agent State Variables

The city agent is characterized by a conservation goal that defines the amount of acres of conservation land desired. The purpose of the conservation land is to reduce flooding in the city, and the conservation goal changes from year-to-year depending on prior hydrologic events. The damage that the city agent incurs from a flood event is defined by a flood damage function. A parameter, $ConsGoal_{max}$, in the agent model defines how responsive the city agent is to prior hydrologic outcomes and determines by how much the city agent will change the conservation goal after experiencing a flood event (Section 2.8)

2.2 Model Overview and Scheduling

Each year, the agent-based model proceeds through monthly time steps to simulate the relevant decision making. The hydrologic module proceeds in shorter hourly time steps to capture flood discharge events associated with rainfall events. Figure 2 depicts the decision-scheduling within the agent-based model. In January, the farmer agent calculates his/her preferred land division between production and conservation based on their risk aversion attitude, conservation-mindedness, newly acquired information about the global market (crop prices, crop production costs, and crop insurance), conservation subsidies provided by the city agent, as well as recent farm performance (profits and yields) (Figure 2, purple box).

In February, the city agent contacts farmer agents in random order to establish new conservation contracts if an unmet conservation goal remains or to renew any expiring contracts (Figure 2, yellow box). If the farmer agent wants to add additional conservation acreage, a new contract is established for a 10 year period. The contract length is based on the Conservation Reserve Program (CRP), which is a program administered by the Farm Service Agency that promotes removal of environmentally-sensitive land from agricultural production in exchange for an annual subsidy payment. However, if the farmer agent wants fewer conservation hectares,

expiring contracts are renewed for a smaller number of hectares or are ended. The farmer is obligated to fulfill any contracts that have not yet expired (i.e. contracts less than 10 years old). Any new acreage that has been established in conservation in addition to currently active contracts is subtracted from the city agent's conservation goal that was established in January. The city agent contacts as many farmer agents as needed until the conservation goal is reached. If there are not enough farmer agents willing to enter into conservation contracts and the conservation goal is not reached, the goal rolls into the next year. Because the farmer agents' land use decisions change on a yearly basis, it may be possible for the city agent to establish further contracts in the next year and fulfill the conservation goal.




Figure 2. Timeline of agent decisions and actions within the agent-based model.

Prior to May, the farmer agent establishes any newly contracted conservation land on the historically poorest yielding land. The farmer agent makes no further decisions during May through August (Figure 2). The city agent continuously keeps track of any flooding that occurs during the May-August period (when the maximum discharge is assumed to occur) (Figure 2, orange box). The associated flood damage cost is calculated in September and used to calculate whether any further conservation land should be added (Figure 2, green box). If no flooding occurred, the conservation goal remains unchanged. In October, the farmer agent harvests his/her crop and calculates yields and profits for that year (Figure 2, blue box).

2.3 Design Concepts

Emergence: Patterns in total conservation land and flood magnitude arise over time, depending on a number of variables. Agent decision-making parameters and behavioral characteristics (e.g. conservation-mindedness) influence the total acreage in conservation land, which in turn affects the magnitude of floods through changes in runoff productivity of the landscape.

Objectives and Adaptation: The ~~objective-goal~~ of the city agent is to reduce flood damage in the city. The city agent attempts to meet this ~~objective-goal~~ through an incentive program in which farmer agents are paid to convert production land to a conservation practice that will reduce runoff. If the city agent incurs a large cost from flooding in a given year, the city agent adjusts his/her “conservation goal” upward in order to ~~reduce minimize~~ future flood damage from events of similar magnitude. The objective of the farmer agent is to balance ~~a-maximization~~ ~~of~~ profits with conservation and risk-aversion attitude. The farmer agents incrementally adjust their land use on an annual basis by taking into account profit variables, risk-aversion, and conservation-mindedness.

Stochasticity: Adjustments and stochastic variability are added to key agricultural variables, which include crop yields, production costs, cash rent values, and opportunity costs associated with conservation land in order to account for economic and environmental randomness within the system (Supplement S1.1, S1.2, S2). Random factors for these variables are drawn from uniform continuous distributions that are based on field data of crop yields, empirical survey data, and estimates published by Iowa State University Extension and Outreach. Changes in these distributions are also accounted for, depending on crop price levels.

Learning: As will be outlined further in Sect. 2.7.2, each year, the farmer agents calculate profit differences between crop production and conservation subsidies. Farmer agents save this profit difference information from the beginning of the simulation and use it to adjust their decision-making space on an annual basis. The profit difference information is based on past crop prices, production costs, and conservation subsidies.

2.4 Model Input

2.4.1 Economic Inputs

Inputs to the agent-based models are historical crop prices (\$/MT), production costs (\$/Ha), cash rental rates (\$/Ha), and federal government subsidy estimates (\$/Ha). An example of these model inputs is shown in Fig. 3 in comparison to mean Iowa crop yields.

2.4.2 Production Costs

Production costs are treated as a time series input, with total costs per hectare for each year represented by one lumped value. Production costs used in this model application include land rent, machinery, labor, crop seed, chemicals, and crop insurance (Plastina, 2017). ~~in~~ addition, in ce estimates put the amount of land rented out in Iowa for farming at 60-80%, it is assumed that all farmer agents rent their land (Plastina et al., 2018; Zhang, 2015); ~~which~~ This

significantly increases expenses as land rental costs account for approximately half of total production costs (Plastina, 2017).

2.4.3 Conservation Subsidy and Costs

The conservation subsidy is based on the CRP Contour Grass Strips practice (CP-15A) which includes annual land rental payments and 90% cost share for site preparation and establishment (USDA Conservation Reserve Program Practice CP-15A, 2011). Subsidies are calculated using annual inputs of historical cash rental rates. The cost of establishing and maintaining conservation land is based on analysis conducted by Tyndall et al., (2013). These costs are adjusted based on the land quality of each farmer agent (Supplement S1.2).




Figure 3. Example input time series of corn price, production cost, and cash rent as compared to mean crop yields.

2.4.4 Federal Government Subsidies

Calculation of federal government crop subsidies for individual farmer agents were not included in the agent-based model due to the complexity and variety of commodity programs available to US farmers, each of which focuses on different aspects of revenue protection (e.g., protection against low crop prices, protection against revenue loss). Rather, federal crop subsidies are an input to the model and applied equally to each farmer agent. In this study, crop

subsidy inputs are based on historical estimates produced by Iowa State University Agricultural Extension (Hofstrand, 2018).

2.4.5 Environmental Variables

The hydrology module requires hourly liquid precipitation (mm) as an input to simulate discharge from short-term heavy rainfall events. The crop yield module requires inputs of mean monthly precipitation and temperature to estimate crop yields (Section 2.6). The module calculates mean monthly precipitation based on the hourly precipitation input, however, the user must provide an input of mean monthly temperatures (C).

2.5 Hydrology Module

A model structure that is designed to simulate peak flows was chosen for the hydrology module. Because the city agent in this model is impacted only by the maximum annual peak flow, precisely simulating the full time series of hydrologic flows as well as hydrologic components such as groundwater flow and evapotranspiration were not needed to meet the objectives of the current study. The modeling structure was designed based on a version of the U.S. Army Corps of Engineers' Hydrologic Engineering Center Hydrologic Modeling System (HEC-HMS) (Scharffenberg, 2013) used by the City of Ames, Iowa for flood forecasting in the Squaw Creek watershed in central Iowa. The Squaw Creek watershed represents the type of rural-urban conditions of interest for this study, and is a useful test-bed for this modeling application (Section 3). Further, calibrated parameters were available for the Squaw Creek watershed (Schmieg et al., 2011), providing a realistic baseline for the hydrology module.

Using the configuration and parameters previously defined by Schmieg et al. (2011) for the Squaw Creek watershed, the model on average was within 12.7% of the observed peak discharge for 12 major events simulated. Six of these events were simulated within 3-8% of the

observation, while the least satisfactory simulation overestimated the observed peak discharge by 33%. This error was most likely due to the high spatial variability of precipitation for that event. For the two most recent record flooding events that have occurred, the model underestimated the peak discharge by 6.2% (2008, observed: $356.7 \text{ m}^3\text{s}^{-1}$, simulated: $334.6 \text{ m}^3\text{s}^{-1}$) and 16.6% (2010, observed: $634.3 \text{ m}^3\text{s}^{-1}$, simulated $528.3 \text{ m}^3\text{s}^{-1}$), showing that the model is able to simulate the flooding events needed to run scenarios within the ABM with a fair degree of accuracy. The HEC-HMS model has also been successfully used for simulation of short term rainfall-runoff events and peak flow and flood analysis in other studies (Chu and Steinman, 2009; Cydzik and Hogue, 2009; Gyawali and Watkins, 2013; Halwatura and Najim, 2013; Knebl et al., 2005; Verma et al., 2010; Zhang et al., 2013).

In the module, basin runoff is computed using the Soil Conservation Service (SCS) curve number (CN) method, runoff is converted to basin outflow using the SCS unit hydrograph (SCS-UH) method, and channel flow is routed through reaches in the river network using the Muskingum method (Mays, 2011). A single area-weighted CN parameter is required for each subbasin and is the only hydrology module parameter that changes during the simulation if land cover changes. The SCS-UH method requires specification of subbasin area, time lag, and model timestep. The Muskingum method is based on the continuity equation and a discharge-storage relationship which characterizes the storage in a river reach through a combination of wedge and prism storage (Mays, 2011). The Muskingum method requires specification of three parameters for each reach within the river network: Muskingum X, Muskingum K, and the number of segments over which the method will be applied within the reach (Mays, 2011). Muskingum X describes the shape of the wedge storage within the reach whereas Muskingum K can be approximated as the travel time through the reach.

For the agricultural areas, empirically-derived CN values (Dziubanski et al., 2017) are used for native prairie strips; a CN = 82 is used for 100% row crop production; and a CN = 72 is used for the conservation option implemented by the farmer agents. Urban areas are set to a CN = 90 which is derived from the standard lookup tables for residential areas with lot sizes of 0.051 hectares or less, soil group C (USDA-Natural Resources Conservation Service, 2004). Subbasin delineations and Muskingum parameters previously defined by Schmieg et al. (2011) are used.

The model accepts point-scale rainfall data (e.g., rain gauge data) and calculates mean areal precipitation using the Thiessen Polygon gauge weighting technique (Mays, 2011). The Thiessen weights are entered as parameters to the module. For the initial testing presented in this paper, uniform precipitation over the entire watershed was assumed.

Output from the hydrology module is discharge at the watershed outlet ($\text{m}^3 \text{s}^{-1}$). The hydrology module is run continuously but is designed primarily for simulation of peak flows, which generally occur during the summer in the study region; therefore, for simplicity, a constant baseflow is assumed and snow is ignored. Runoff, river routing processes, and discharge are computed on a timestep identical to the input rainfall data. The model is run at an hourly timestep in this study, but is capable of running at a 30-minute timestep.

2.6 Crop Yield Module

Crop yields are modeled with a multiple regression equation that takes into account monthly precipitation and temperature. The regression equation, which was developed using historical crop yield and meteorological data for Iowa from 1960-2006, can be represented as (Tannura et al., 2008):

$$\begin{aligned}
yield_t = & \beta_0 + \beta_1(year_t) + \beta_2(September\ through\ April\ precipitation) \\
& + \beta_3(May\ precipitation) + \beta_4(June\ precipitation) \\
& + \beta_5(June\ precipitation)^2 + \beta_6(July\ precipitation) \\
& + \beta_7(July\ precipitation)^2 + \beta_8(August\ precipitation) \\
& + \beta_9(August\ precipitation)^2 + \beta_{10}(May\ temperature) \\
& + \beta_{11}(June\ temperature) + \beta_{12}(July\ temperature) \\
& + \beta_{13}(August\ temperature) + \varepsilon_t
\end{aligned} \tag{1}$$

Mean error of the above regression for Iowa over the 1960-2016 period is -0.395 MT/ha, and mean absolute error is +0.542 MT/ha. An error correction factor of +0.395 MT/ha was added to the yield for each year to correct for this error. The above regression model is only appropriate for reproducing mean historical crop yields. Since each farmer's land can be composed of different soil types, adjustments are applied to the crop yield for each soil type to account for differences in soil productivity (Supplement S2).

2.7 Farmer Agent Module

2.7.1 Conservation option

The conservation option implemented by farmer agents is native prairie strips, a practice in which prairie vegetation is planted in multiple strips perpendicular to the primary flow direction upland of and/or at the farm plot outlet (Dziubanski et al., 2017; Helmers et al., 2012; Zhou et al., 2010). Either 10% or 20% of the total field size is converted into native prairie vegetation under this practice. Prairie strips have been shown to reduce runoff by an average of 37% (Hernandez-Santana et al., 2013), and have additional benefits of reducing nutrients (Zhou et al., 2014) and sediments (Helmers et al., 2012) in runoff. The greatest runoff reduction was realized under the 10% native prairie cover; therefore, the most conservation-minded farmers ($Cons_{max} = 0.10$) in the model potentially convert up to 10% of their total land into native prairie.

2.7.2 Farmer agent land use decision process

Agents within an ABM can be modeled using a variety of decision models with varying degrees of complexity. Rules governing agent decision-making need to realistically capture human behavior without creating an excessively complex model (An, 2012; Zenobia et al., 2009). An (2012) compiled a list of nine of the most common decision models used in agent-based modeling studies. Examples of a few of these include micro economic models, space theory based models, cognitive models, and heuristic models. In micro-economic models, agents are typically designed to determine optimal resource allocation or production plans such that profit is maximized and constraints are obeyed (Berger and Troost, 2014). Example studies using optimization include Becu et al. (2003), Ng et al. (2011), Schreinemachers and Berger (2011). In heuristic-based models, agents are set up to use “rules” to determine their final decision (Pahlwostl and Ebenhöf, 2004; Schreinemachers and Berger, 2006). The “rules” are typically implemented using conditional statements (e.g. if-then). Example studies using heuristics include Barreteau et al. (2004), Le et al. (2010), Matthews (2006), van Oel et al. (2010).

We take a different approach from the aforementioned studies by modeling agent decision making using a nudging concept originating in the field of data assimilation (Asch et al., 2017). Agents nudge their decision based on outcomes (i.e. flood damage, farm profitability) from the previous year. Information relevant to an individual agent is mapped into the decision space through a weighting function that updates the previous year's land use prior decision to create a new (posterior) decision for the current year. The approach used for both agents is different from optimization in that the agents are not trying to determine the best decision for each year. These types of agents behave based on the idea of “bounded rationality”. In this case, the rationality of the agents is limited by the complexity of the decision problem and their cognitive ability to process information about their environment (Simon, 1957). These agents try to find a

satisfactory solution for the current year, and are thus termed “satisficers” rather than optimizers (Kulik and Baker, 2008).

At the start of each calendar year, a farmer agent decides how to allocate his/her land between production and conservation based on five variables: risk-aversion, crop price projections, past profits, conservation goal, and neighbor land decisions. These factors were chosen based on numerous studies indicating profits, economic incentives, conservation beliefs, beliefs in traditional practices, neighbor connections, and observable benefits to be the key factors influencing on-farm decision making related to conservation adoption (Arbuckle, 2017; Arbuckle et al., 2013; Burton, 2014; Daloğlu et al., 2014; Davis and Gillespie, 2007; Hoag et al., 2012; Lambert et al., 2007; McGuire et al., 2015; Nowak, 1992; Pfrimmer et al., 2017; Prokopy et al., 2019; Ryan et al., 2003).

A farmer agent’s decision of the total amount of land to be allocated into conservation, C_t , for the current year t is:

$$D_t = W_{risk-averse}[C_{t-1:t-X}] + W_{futures}[D_{t-1} + \delta C_{futures:Y}] + W_{profit}[D_{t-1} + \delta C_{profit:X}] + W_{cons}[D_{t-1} + \delta C_{cons}] + W_{neighbor}[C_{neighbor}] \quad (2)$$

where $C_{t-1:t-X}$ is the mean total amount of land allocated to conservation during the previous X years, D_{t-1} is the prior conservation decision (total amount of land the farmer would have liked to implement in conservation) in year $t - 1$, $\delta C_{futures:Y}$ is the decision based on crop price projections for Y years into the future, $\delta C_{profit:X}$ is the decision based on the mean past profit of the previous X years, δC_{cons} is the decision based on the conservation goal of the farmer, and $C_{neighbor}$ (Supplement S3) is the weighted mean conservation land of the farmer agent’s neighbors (Table 1). A given farmer can make a certain random number of neighboring connections with farmers that are located in the same subbasin (Supplement S3). The variable Y

indicates that Θ one farmer agent might consider his/her history of conservation land implemented over the last year, while another farmer agent might consider his/her conservation land implemented over the last 5 years. Similarly, the variable X indicates that one farmer agent might take into account future crop projections for the next 5 years, while another farmer agent might take into account crop projections for the next 10 years.

Decision weights alter how each of the five components factor into the farmer agent's decision: $W_{risk-averse}$ reflects the unwillingness to change past land use, $W_{futures}$ reflects the consideration of future price projections, W_{profit} reflects the consideration of past profits, W_{cons} is the agent's consideration of his/her conservation goal, and $W_{neighbor}$ reflects the importance that the agent places on his neighbor's decision (Table 2). Upon initializing each farmer agent, values are allocated for each decision weight such that:

$$W_{risk-averse} + W_{futures} + W_{profit} + W_{cons} + W_{neighbor} = 1 \quad (3)$$

The above decision scheme allows for varying decision weights, thus one farmer's decision may be heavily weighted by future crop prices, whereas another farmer's decision may be heavily weighted by past profits. If majority of a farmer's decision is based on $W_{risk-averse}$, then that farmer is less inclined to change his/her previous land use.

The decision components for past profit and future crop prices are based on a partial budgeting approach that compares land use alternatives. Under this budgeting approach, farmer agents take into account added and reduced income, as well as added and reduced costs from changing an acre of land from crop production to conservation (Tigner, 2006). The result from performing this budget indicates the net gain or loss in income that a farmer agent may incur if they make the land conversion.

The past profits decision is solely based on outcomes that have been fully realized for the previous X years. In this decision, the land allocated to conservation is based on the net amount of money that could have been earned per hectare of conservation land versus crop land and is calculated as:

$$\delta C_{profit:X} = [A * Profit_{diff}^2 + B * Profit_{diff} + C] \cdot Cons_{max} \cdot Hectares_{tot} \quad (4)$$

where $Profit_{diff}$ is the difference in profit between a hectare of cropland and a hectare of conservation land (Table 1), $Cons_{max}$ is the farmer agent's maximum conservation parameter, $Hectares_{tot}$ is the area of the agent's land. $Profit_{diff}$ integrates costs and revenue of crop production as well as costs and revenue of conservation land, which are computed based on model input data (Section 2.4, Supplement S4). In the case of $\delta C_{profit:X}$, $Profit_{diff}$ is calculated using realized crop prices from previous years (Supplement S4). The future price decision variable, $\delta C_{futures:Y}$, is also calculated using the same form of Eq. (4). However, $Profit_{diff}$ is calculated using projected crop prices for the Y upcoming growing seasons. These price projections are based on historical crop prices with an added adjustment calculated from historical errors in crop price forecasts produced by the U.S. Department of Agriculture (Supplement S5).




Figure 4. Example of percent conservation change for δC_{profit} and $\delta C_{futures}$. Gray curves indicate negative percent change (decrease conservation land), black curves indicate positive percent change (increase conservation land).

The first term in Eq. (4), ~~the is a~~ second-degree polynomial of form $Ax^2 + Bx + C = y$, is displayed in Fig. 4. At the start of each year, farmers may decide to alter their land use based on observed $Profit_{diff}$ from harvests in previous years ($\delta C_{profit:X}$) or calculated $Profit_{diff}$ based on projected crop prices ($\delta C_{futures:Y}$). If $Profit_{diff}$ is positive (i.e. greater profit is earned from crop production than conservation land), the farmer agent will potentially decrease the amount of land in conservation (gray curve). Likewise, under negative $Profit_{diff}$, conservation land is potentially increased because revenue is lower from crop production (black curve). Half of the maximum allowable percent increase in conservation land is assumed to correspond to the median historical negative $Profit_{diff}$, whereas half of the maximum allowable percent decrease in conservation land corresponds to the median historical positive $Profit_{diff}$ (Figure 4). We assume that farmer agents will not change land use when a very small profit difference between

the two possible options is observed because changing land use requires extra upfront time and resources (Duffy, 2015). Similarly, we assume that farmer agents will fully implement the maximum land conversion possible prior to reaching the most extreme $Profit_{diff}$ values. Three equations need to be simultaneously solved to determine coefficients A, B, C (Supplement S4). The three equations are based on the 25th, median, and 75th percentiles of historical $Profit_{diff}$ information. Thus, farmers are continually utilizing historical observations of $Profit_{diff}$ to formulate their decision space through time.

The use of a profit function (i.e. Eq. (4)) is meant to capture to effects of changes in crop prices on conservation land. In 2008 and 2011, corn prices rose to a record high values, and farmers in the Midwest U.S. (e.g., Iowa, Minnesota) were converting significant portions of CRP land back into crop production (Marcotty, 2011; Secchi and Babcock, 2007). It is estimated that when corn prices rise by \$1.00, 10-15% of CRP land in Iowa is converted back to production (Secchi and Babcock, 2007). Eq. (4) captures this transition between adding and removing conservation land based on crop price change, and it allows for variation in the decision-making between farmer agents since variables such as crop production costs vary from farm to farm.

The total amount of agricultural land that a farmer converts to conservation in any given year based on his/her conservation goal (δC_{cons}) is defined by the Bernoulli distribution:

$$P(n) = p^n(1 - p)^{1-n} \quad n \in \{0,1\} \quad (5)$$

Here, p indicates the probability of fully implementing conservation land and $1 - p$ indicates the probability of not implementing any conservation land. The variable n is simply the support of the distribution that labels a success of full implementation as 1 and a failure of full

implementation as 0. The probability p of fully implementing conservation land is a function of the agent's $Cons_{max}$ parameter and is computed by:

$$p = 10 \cdot Cons_{max} \quad (6)$$

The probability p scales from 0 at a $Cons_{max}$ of 0, to 1 at a $Cons_{max}$ of 0.1. Therefore, farmer agents with a $Cons_{max}$ of 0.05 and 0.1 will have a 50% and 100% probability of fully implementing (10% of total agricultural land) conservation land in any given year based on their conservation decision variable.

2.8 City Agent Module

At the end of each year, the city agent collects discharge data and calculates the damage (Supplement S7) associated with the peak annual discharge at the watershed outlet for that year. In February of the next year, the flood damage for the previous year $t - 1$ is used to compute the conservation goal of the city agent for the current year t .

The conservation goal of the city agent is calculated as:

$$G_t = G_{t-1} + (A_{tot} - C_{tot}) \cdot P \quad (7)$$

$$P = P_{new} \cdot FDam \quad (8)$$

$$P_{new} = \frac{ConsGoal_{max}}{FDmax} \quad (9)$$

where G_t is the conservation goal for the new year t (Table 1), G_{t-1} is the unfulfilled hectares in conservation from the previous conservation goal for year $t - 1$, A_{tot} is the total land area ~~in the~~ catchment owned by the farmer agents, C_{tot} is the total number of hectares currently in conservation, P is the percentage of new production land added into conservation, P_{new} indicates how much land to add into conservation based on the flood damage $FDam$ for year $t - 1$, and $ConsGoal_{max}$ is a parameter that indicates the new percentage of conservation land to be added

if maximum flood damage occurs (Table 2). Currently, $ConsGoal_{max}$ is set to 5% of total land area in the watershed when maximum damage occurs.

3. Scenario Analysis

The study watershed is modeled after the Squaw Creek basin (~56200 Ha) located in central Iowa, USA (Figure 45). This basin is characterized by relatively flat hummocky topography and poorly drained soils with a high silt and clay content (~30-40% silt and clay) (Prior, 1991; USDA-Natural Resources Conservation Service (USDA-NRCS), 2015). The predominant land use is row crop agriculture (~70% of the total watershed area) with one major urban center at the outlet (Ames, Iowa), and several small communities upstream. Average annual precipitation is 32 inches (812 mm), with the heaviest precipitation falling during the months of May and June. The watershed is divided into 14 subbasins.




Figure 45. Squaw Creek watershed and subbasin division used in the hydrology module. Land cover data shown is from the National Land Cover Database (NLCD), 2016.

In this model application, 100 farmer agents are implemented (~7 farmers per subbasin) with 121 hectares total for each farmer. The total acreage per farmer compares reasonably well with average farm size for the state of Iowa in 2017, which was 140 hectares (USDA National Agricultural Statistics Service, 2018). Soil types and the area of land associated with each soil type are randomly assigned to each farmer agent upon model initialization. Assigning different

soil types creates heterogeneous conditions under which farmer agents must operate (Supplement S2) and affects the profitability of each farmer agent differently.

Six scenarios are run: high and low yield ($\pm 11\%$ from historical yield), high and low corn prices ($\pm 19\%$ from historical prices) and high and low conservation subsidies ($\pm 27\%$ from historical cash rent). The watershed was also simulated under historical conditions, in which no economic variables were changed, for comparison purposes. The 90th percentile discharge in analyzed, which represents the 0.1 exceedance probability level, to examine changes in large discharge events. The above percentages were computed using trends and mean absolute deviations of historical economic data. For instance, based on the crop regression model (Section 2.6), crop yields display a relatively linear increase with time. The mean absolute deviation of crop yield was then computed using the linear time trend as a central tendency. The mean absolute deviation was determined to be 11%, thus the yield scenarios are $\pm 11\%$ from the historical yield. The same approach was used for the crop price and conservation subsidy scenarios. A linear and cubic function were found to provide a good estimate of the central tendency of historical cash rent and crop prices, respectively, for those calculations. In addition, four different farmer decision schemes are created in which an 80% weight was assigned to one decision variable, with all other variable weights set to 5% (Table 3). Each scenario is tested with each decision scheme and system outcomes under different farmer behaviors are assessed.

To test the sensitivity of the hydrologic system to farmer types, the conservation parameter ($Cons_{max}$) of the farmer agents is varied using a stratified sampling approach. Each farmer agent is randomly assigned a $Cons_{max}$ value from a predefined normal distribution: $(\overline{Cons_{max}}, \sigma_{Cons_{max}})$. The lowest distribution is defined as $\mathcal{N}(0.01, 0.01)$ and the highest distribution is defined as $\mathcal{N}(0.09, 0.01)$. Any farmer agent that is assigned a parameter value

less than 0 or greater than 0.1 is modified to have a value of 0 or 0.1, respectively. Twelve simulations are performed for each conservation parameter distribution, with a total of 17 conservation parameter distributions. Thus, the first 12 simulations consist of farmer agents with $Cons_{max}$ chosen from $\mathcal{N}(0.01, 0.01)$. For the next 12 simulations, the mean $Cons_{max}$ is shifted up by 0.05, with $Cons_{max}$ chosen from $\mathcal{N}(0.015, 0.01)$. A total of 204 simulations are conducted for each decision scheme under each scenario (Table 3).

Each simulation is run using 47 years of historical climate and market data, with the exception of federal crop subsidies, which are based on 16 years of historical estimates produced by Iowa State University Agricultural Extension (Hofstrand, 2018; Table 4). It is assumed that federal crop subsidy payments from 1970-2000 are similar to levels seen from year 2000-2005 due to relative stability in long-term crop prices and production costs. The hourly 47 year precipitation time series data was obtained from the Des Moines, Iowa airport Automated Surface Observing System. Historical 47 year time series of corn prices, crop production costs, and land rental values are used as economic inputs into the model and were obtained from Iowa State University Agricultural Extension and Illinois FarmDoc (Table 4).

45. Model Calibration and Validation

Calibrating and validating the social part of social-hydrologic models is difficult due to reasons that include lack of sufficiently detailed empirical data or system complexity at various scales (An, 2012; Ormerod and Rosewell, 2009; Troy et al., 2015). Validation of agent-based models is usually performed on what are termed the micro and macro levels. The micro level involves comparing individual agent behaviors to real world empirical data whereas the macro level involves comparing the model's aggregate response to system-wide empirical data (An et al., 2005; Berger, 2001; Troy et al., 2015; Xiang et al., 2005). Troy et al., (2015) suggests that

one or a few model simulations out of an ensemble of simulations should match the real-world observed data.

We conduct an indirect macro-level model calibration for determining an appropriate range of farmer agent decision weights (Windrum et al., 2007). Since the subsidy program offered by the city agent is similar to the federal Conservation Reserve Program (CRP), the model was developed and calibrated to attempt to reproduce the range and variability of conservation land seen in the CRP program. CRP data from 1986-2016 for the Central Iowa Agricultural District was used in the calibration process and two main objectives functions were used:

$$MAE = \frac{\sum_{i=1}^n |y_i - x_i|}{n} \quad (10)$$

$$Pearson's\ r = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2} \sqrt{\sum_{i=1}^n (y_i - \bar{y})^2}} \quad (11)$$

In the first step of calibration, the focus was to determine an appropriate range of mean *ConsMax* of the farmer agent population to match the magnitude of CRP land seen for central Iowa. The model was simulated 360 times using 20 random sets of farmer agent decision weights. Output from the first calibration step was filtered using a criteria of $r > 0.6$ and $MAE < 25\%$, and the optimal *ConsMax* range was reduced to 0.05-0.07. In the second step of calibration, the focus was to determine a singular optimal mean *ConsMax* value and narrow the range for each decision weight. *ConsMax* was incremented by 0.001 within the range derived from step 1, and 20 simulations were performed for each increment using decision weights stochastically drawn from the uniform distribution $\mathcal{U}(0.05, 0.95)$ for a total of 400 simulations. Output was filtered using a stricter criteria of $r > 0.7$ and $MAE < 25\%$. The final calibration

step involved 400 simulations with the optimal mean *ConsMax* value and stochastic sampling from the reduced range of decision weights derived in step 2. Filtering with a criteria of $r > 0.75$ and $MAE < 12.5\%$ was performed to determine the final optimal decision weight ranges.

The optimal mean *ConsMax* value was determined to be 0.06 and the final optimal decision weight ranges were determined to be: $W_{risk-averse} = (0.1, 0.43)$, $W_{futures} = (0.07, 0.24)$, $W_{profit} = (0.07, 0.34)$, $W_{cons} = (0.18, 0.37)$, $W_{neighbor} = (0.05, 0.35)$. The median r and MAE values of the simulations after filtering with the criteria in step three ($r > 0.75$, $MAE < 12.5\%$) were 0.79 and 11% respectively. Sixty-six out of 400 simulations matched this criteria in step three, whereas only seven matched this criteria in step one and 26 matched this criteria in step two.

The model simulated conservation land generally aligns with trends in the observed conservation land (Figure 106). Simulated conservation land is not maintained following a rise in crop prices in the mid-1990s and from 2006-2013, which is similar to the observed data (red). The drop in conservation land during these time periods occurs because the subsidy rate is not modified rapidly enough in comparison to market forces to incentivize the farmer (Newton, 2017). ~~In 2008 and 2011, corn prices rose to a record high values, and farmer in the Midwest U.S. (e.g., Iowa, Minnesota) were converting significant portions of CRP land back into crop production (Marcotty, 2011; Secchi and Babcock, 2007). It is estimated that when corn prices rise by \$1.00, 10-15% of CRP land in Iowa is converted back to production (Secchi and Babcock, 2007).~~ The model does capture the smaller decrease in conservation land between 2007-2014, even though crop prices rose more dramatically than in the mid-1990s.




Figure 106. Simulated conservation land from four model simulations with Pearson's $r > 0.8$ and MAE $< 12.5\%$ in comparison to observed conservation land.

The onset of significant land conversion in the model is offset from the observations. Conservation land is implemented in the mid-1970s, while conservation land in the observation is implemented in the late-1980s. The CRP program did not come into existence until 1985, which partly explains this difference. A large rise in conservation land to roughly 4% occurs from 1975-1978, most likely due to a combination of decreasing crop prices from 1970-1974 and model spin up. This is similar to the rate of rise in conservation land that occurred under the CRP programs from 1985-1987 under a comparable period of decreasing crop prices.

Overall calibration does provide evidence that the model captures changes in CRP land during the appropriate time periods. However, the calibration technique does have limitations. however, it does not provide evidence that any individual agent's decisions are valid. The technique followed here was an indirect calibration approach, whereby the parameters are determined based on the simulations that replicate the empirical data best (Fagiolo et al., 2006). This technique can lead to equifinality since difference parameter sets may reproduce the historical observations with similar degrees on accuracy. Further, this calibration approach does not provide evidence that any individual agent's decisions are valid. The stochastic nature of human behavior coupled with path dependencies makes it difficult to predict individual agent outcomes accurately (Berglund, 2015). A dominating problem with calibrating ABMs is that it may be difficult to find sufficient data sets to support a robust validation at the micro-level. For modeling land use decisions, data is typically available at a larger scale such as county or state level rather than at the individual agent-level (e.g. single farm) (An, 2012; Parker et al., 2008). This introduces difficulty in trying to validate farm-level decisions with respect to farm-level finances (Section 2.7.2). Adding in additional factors, such as Federal Market Loss Assistance and Loan Deficiency Payments, as well as trying to characterize some of the other model parameters that were not a focus of this calibration, may further improve results.

In light of the paper by Windrum et al. (2007), there has been much debate as to the proper methodology and techniques to follow for ABM validation (Bharathy and Silverman, 2013; Hahn, 2013). To fully validate the current model, a more extensive process may be necessary. Macal et al., (2007) introduced a framework for ABM validation that may provide for a more comprehensive evaluation. This framework includes subject matter expert evaluation, participatory simulation, model-to-model comparison, comparison against critical test cases,

invalidation tests, and comprehensive testing of the entire agent strategy and parameter space.
However, following this framework is very time costly, and thus most recent studies have
focused on empirical validation against real world macro level data, with some studies validating
at the individual agent level if data is available (Fagiolo et al., 2019; Guerini and Moneta, 2017;
Langevin et al., 2015; Schwarz and Ernst, 2009).

4.5. Results

4.5.1 Crop Price Scenarios

The 90th percentile peak discharge is 296.4 m³/s when no conservation is occurring in the watershed (~~Figure 5~~Figure 7). The 90th percentile peak discharge decreases for all four decision schemes and under all scenarios as the average conservation-mindedness ($Cons_{max}$) of the population increases (~~Figure 5~~Figure 7). The low crop price scenario produces a larger decline in peak discharge compared to the high crop price scenario, with the exception of the conservation decision scheme (80% weight on conservation) in which both low and high crop price scenarios produce a similar ensemble pattern (~~Figure 5~~Figure 7a).




Figure 7. Mean 90th percentile discharge for high and low crop price scenarios under (a) 80% weight on conservation goal, (b) 80% weight on future price, (c) 80% weight on past profit, and (d) 80% weight on risk aversion. Bars indicate the median (circle) and the 5th and 95th percentiles of discharge for all simulations at a specific $Cons_{max}$.

Under low crop prices, peak discharge reaches an average reduction of 8.18% ($24.27 \text{ m}^3/\text{s}$) when the average $Cons_{max}$ is 0.08-0.09 (conservation-minded population) and 4.67% ($13.85 \text{ m}^3/\text{s}$) when the average $Cons_{max}$ is 0.04-0.06 (mixed population). The decrease in peak discharge corresponds with the 800-1000 hectares and 400-600 hectares converted to conservation by the conservation-minded and mixed farmer populations, respectively (**Figure 8a, c, e, g**). The production-minded populations ($Cons_{max} \sim 0.01-0.02$) implement less than 200 hectares during the entire simulation period. These acreage values represent 6.5-8.2%, 3.3-5.0%, and less than 2.0% of the entire watershed for the conservation-minded, mixed, and production-minded groups, respectively. Given that 10% of the watershed would be in conservation if native prairie strips were fully implemented, about 65-80% of a conservation-

minded population fully implements the practice over the simulation period under low crop prices.

Under the high crop prices, mean peak discharge decreases by 5.6 % (16.6 m³/s) under the future price weighting scheme and 2.9% (8.6 m³/s) under the past profit weighting schemes for the highly conservation-minded population ([Figure 5](#)[Figure 7b](#) and c, respectively), with an even smaller reduction seen for the risk-averse scenario. This represents approximately a 61% smaller decrease in the peak discharge when crop prices are high and the population is conservation-minded as compared to the low crop price scenario. Discharge remains largely unchanged for these decision schemes because generally less than 300 hectares of land is allocated for conservation when corn prices are high ([Figure 6](#)[Figure 8d, f, and h](#)). The small amount of conservation land implemented is due to farmer agents receiving significantly more revenue from crops than conservation subsidies. However, in the case of low crop prices, conservation subsidies allow the farmer agents to approach break even because they are guaranteed a subsidy that covers the cash rent for that land, whereas crop production leads to potential losses due to corn prices being low relative to production costs. Even in these scenarios where farmer agents are heavily considering profit related variables, populations dominated by production-minded farmer agents are still inclined to leave land in production ([Figure 6](#)[Figure 8c](#) and e).




Figure 8. Range of simulated conservation land within the watershed under low (left column) and high (right column) crop prices for conservation-minded populations (green), mixed populations (blue) and production-minded populations (red). Crop prices are plotted as bars for each crop price scenario. Results are for decision schemes of 80% weight on conservation behavior (a, b), 80% weight on future price (c, d), 80% weight on past profit (e, f), and 80% weight on risk aversion (g, h).

4.5.2 Crop Yield Scenarios

Under high and low crop yield scenarios, the 90th percentile peak discharge decreases by an average of 5.9% (17.4 m³/s) and 7.6% (22.7 m³/s), respectively, for the conservation-minded populations (Figure 7, Figure 9). Thus, a smaller decrease in peak discharge occurs with low crop yields relative to low crop prices (Figure 5, Figure 7). In the low crop yield scenario, conservation land was approximately 200 Ha less than in the low crop price scenario, particularly for the past profit and future price decision schemes (Figure 6, Figure 8a, c, e, g and 8a10a, c, e, g).

Conversely, more conservation land is established under the high yield scenario compared to the high crop price scenario (Figure 6, Figure 8b, d, f, h and 108b, d, f, h). As a result, mean peak

discharge decreases in the high yield scenario by 15.6% more compared to the high crop price scenario for the conservation-minded population.




Figure 9. Mean 90th percentile discharge for high and low crop yield scenarios under (a) 80% weight on conservation goal, (b) 80% weight on future price, (c) 80% weight on past profit, and (d) 80% weight on risk aversion. Bars indicate the median (circle) and the 5th and 95th percentiles of discharge for all simulations at a specific Cons_{max} .




Figure 10. Range of simulated conservation land within the watershed under low (left column) and high (right column) crop yields for conservation-minded populations (green), mixed populations (blue) and production-minded populations (red). Yearly crop yields are plotted as bars for crop yield scenario. Results are for decision schemes of 80% weight on conservation behavior (a, b), 80% weight on future price (c, d), 80% weight on past profit (e, f), and 80% weight on risk aversion (g,h).

45.3 Conservation Subsidy Scenarios

Under the low and high subsidies scenarios (not shown), the 90th percentile peak discharge decreases by an average of 5.8% (17.3 m³/s) and 7.6% (22.5 m³/s), respectively, for conservation-minded populations. Similar to the low crop yield scenario, high subsidies do not produce as large of a decrease in mean peak discharge as low crop prices (Figure 7). In the high subsidies scenario, conservation land was approximately 200-300 Ha less than in the low crop price scenario, specifically for the future price and past profit decision scheme. In comparison, low subsidies generate more conservation land than under high crop prices (Figure 8b, d, f, h). As a result, mean peak discharge decreases in the low subsidy scenario by

14.8% more compared to the high crop price scenario for the conservation-minded population. Differences in peak discharge reduction between the high subsidy and low yield scenarios were insignificant, with less than 1% difference between these two scenarios.

4.5.4 Decision Schemes

The future price and past profit decision schemes display the largest spread in discharge outcomes between scenarios (Figure 5 Figure 7, 79). Mean peak discharge decreases on average by 9% (~27.2 m³/s) relative to when no conservation occurs for both decision schemes under all scenarios that encourage more conservation land (i.e. low crop prices, low yields, high subsidies) (Figure 5 Figure 7b, c and 7b9b, c). Under scenarios that encourage less conservation land, mean peak discharge decreases by 5% (~15.4 m³/s). This spread in peak discharge results is not present under the risk-averse and conservation decision schemes.

The spread between the mean peak discharge under the different scenarios is smaller for the future price decision scheme (Figure 5 Figure 7b and 7b9b) compared to the past profit decision schemes (Figure 5 Figure 7c and 7e9c). This smaller spread may be due to uncertainty in future crop price projections. For instance, future crop price projections may underestimate high crop prices, but overestimate low crop prices, as is observed in previous USDA crop price forecasts (Supplement S5). Thus, the farmer agents may be making decisions based on a smaller range of crop prices when under the future price decisions compared to the past profit decision scheme where they use realized crop prices. In addition, the future crop price decision scheme results in greater variability in conservation land over short periods of time under all scenarios (Figure 6 Figure 8c,d and 8e10c,d). This result is evident under the low crop price scenario, with several short periods showing changes in conservation land of 200-400 ha as compared to the

past profit scenario where conservation land remains relatively steady. However, this result does not lead to a larger spread (i.e. red and blue bars) within the mean peak discharge results.

The risk averse decision scheme produces the smallest changes in peak discharge under all scenarios, with an average decrease of less than 2% ($6 \text{ m}^3/\text{s}$) and 3% ($9 \text{ m}^3/\text{s}$) for mixed and conservation-minded populations, respectively (~~Figure 5~~Figure 7d, 7d9d). Because the farmer's past practices are the primary factor in determining land conversion in this scheme, the farmer agents implement a limited number of conservation acres ($\leq 200 \text{ ha}$), regardless of the scenario. Therefore, changes in the economic variables are not having as large of an impact on the farmer agents when they are strongly risk-averse.

Overall, the current city agent conservation goal of 5% new conservation land at maximum flood damage did not have a significant impact on the total amount of land implemented. Following two major flooding events, the conservation goal of the city agent increases from less than 20 ha in 1975 to 620 ha in 1976. A similar event in 1977 increases the conservation goal by another 500 ha for a total goal of approximately 1100 ha. These increases correspond to the large and rapid onset of conservation land seen during those years (~~Figure 6~~Figure 8a, c, e; 8a10a, c, e). After the 1977 flood event, several smaller flood events do occur that are generally less than 15-30% of maximum, which further increases the conservation goal by ~200-300 Ha. When the population has a high average $Cons_{max}$, the conservation goal of the city agent is nearly fulfilled during this period, particularly in the low crop price scenario. In these cases, 900 ha of the conservation goal is implemented, and 200 ha remains unimplemented. This results in the largest reduction in 90th percentile discharge under all scenarios and decision schemes (~~Figure 5~~Figure 7a, 7a9a). When the population has a low average $Cons_{max}$, the majority of the city agent's conservation goal remains unimplemented. Thus, the goal remains at

a constant 1000-1400 ha and discharge remains unchanged. The only case where the city agent conservation goal limits the amount of land implemented is under the conservation weighting scenario since conservation-minded farmers are inclined to add conservation land on a yearly basis.

4.5.5 Historical Comparison

To gain an understanding of how each of the scenarios differs from the historical 1970-2016 period, the mean peak discharge is compared against the historical scenario (Figure 11). Recall that under the historical scenario, farmer agents make annual land use decisions as in the other scenarios, but corn prices, conservation subsidies, and crop prices are unchanged from historical observed values., ~~which does not modify any economic or agricultural variables~~ (Figure 9). Overall, crop prices had the largest impact on mean peak discharge while changes in subsidies had the smallest overall impact. When crop prices were low, mean peak discharge decreased by 1-2% for mixed populations and 2-3% for conservation-minded populations under the future price and past profit schemes compared to the historical scenario (Figure 9 Figure 11a). High crop prices result in an increase in peak discharge from the historical scenario, with an increase of 1-3% for mixed populations, and 3-5% for conservation-minded populations. This indicates that the farmer agents are more likely to convert land back to crop production under high crop prices than convert land to conservation under low crop prices, which is a similar conclusion to Claassen and Tegene, 1999.

The subsidy scenarios produced a similar pattern to the crop price scenarios, where a larger change (increase) in mean peak discharge occurs under low subsidies than under high subsidies (Figure 9 Figure 11b). This pattern was not as clearly evident under the yield scenarios, with similar changes resulting from high and low yields (Figure 9 Figure 11c).




Figure 9 Figure 11. Percent Change in median 90th percentile discharge from the historical scenario for (a) high and low crop prices, (b) high and low subsidies, (c) high and low yields for the conservation, risk, future price, and past profit weighting schemes.

6. Conclusions

Scenarios of historical and low crop yields, as well as high and low corn prices and conservation subsidies, were simulated for an agricultural watershed in the Midwest US corn-

belt using an agent-based model of farmer decision making and a simple rainfall-runoff model. The influence of different farmer agent decision components on model outcomes was also explored. Model results demonstrate causations and correlations between human systems and hydrologic outcomes, uncertainties, and sensitivities (specifically focused on high flows).

The primary findings from this study are:

- Crop prices had the largest impact on mean peak discharge, with a 61% larger reduction in mean peak discharge under low crop prices in comparison to high crop prices.
- Changes in subsidy rates and crop yields produced a smaller impact on mean peak discharge. Only a 25-30% difference in mean peak discharge was realized between high and low subsidies, and high and low yields.
- Farmer agents more often made decisions to eliminate conservation land than to enter into conservation contracts: a 3-5% increase in mean peak discharge occurred under high crop prices, while only a 2-3% decrease in mean peak discharge occurred under low crop prices compared to the historical simulation. Thus, even under low crop prices, the effectiveness of the conservation program is limited either due to economic or behavioral factors.
- Hydrologic outcomes were most sensitive when farmer agents placed more weight on their future price or past profit decision variables and least sensitive when farmer agents were highly risk averse. For instance, under future price and past profit weighting scenarios, a 4% and 7% difference in mean peak discharge is seen between high and low crop prices as opposed to a 0-1% difference under the risk averse weighting scenario.

The ABM modeling approach demonstrated here can be used to advance fundamental understanding of the interactions of water resources systems and human societies, particularly

focusing on human adaptation under future climate change. Our model indicates that external factors can influence local streamflow, albeit in a complex and unpredictable way as the information gets filtered through the complex decision making of local farmers. Social factors, both local and external, introduce significant uncertainty in local hydrology outcomes, and by ignoring them, water management plans will be inherently incomplete. Thus, multi-scale human factors need to be explicitly considered when assessing the sustainability of long-term management plans.

This study additionally demonstrates some of the advantages of the ABM approach. One of the primary advantages of ABMs is the ability to capture emergent phenomenon (Bonabeau, 2002). For instance, in the model, the change in conservation area seen in the mid-1990s is larger than during the period after 2007, despite the much larger volatility in crop prices after 2007. While the primary reason behind this phenomenon may not be clear, the ABM captures this change. The ABM also allows for specifying small scale differences between farmer agents such as variations in conservation-mindedness, production costs, yields, cash rents, etc. Thus, using ABMs allows for a very flexible modelling approach.

The current model design contains limitations in both the hydrologic and agent-based models that should be addressed in future model development. The curve number values that were used to represent the conservation option were derived for small agricultural plots of approximately 0.5-3 Ha in size. The question remains whether these CN values can be scaled up to the size of a several hundred hectare farm plot and still produce reasonable discharge results. In addition, there is no explicit spatial representation of farmer agents within each subbasin, Coupling the agent-based model to a more robust hydrologic model may reduce some of these hydrologic limitations. The Agro-IBIS model, which includes dynamic crop growth and a crop

management module, would be particularly well suited to further investigating various farm-level decisions within an ABM on hydrologic outcomes (Kucharik, 2003).

From the agent-based modeling standpoint, the decision-making of the farmer and city agent could be made more sophisticated by introducing certain state variables, further decision components and longer planning horizons. Studies have identified variables such as farm size, type of farm, age of farmer, off farm income, land tenure agreement, education from local experts, among others, to be significant in determining adoption of conservation practices (Arbuckle, 2017; Daloğlu et al., 2014; Davis and Gillespie, 2007; Lambert et al., 2007; McGuire et al., 2015; Ryan et al., 2003; Salatiel et al., 1994; Schaible et al., 2015). The functionality of the city agent could be expanded by introducing cost-benefit analysis capabilities. Cost-benefit capabilities would allow the city agent to make more advanced decisions such as choosing among a variety of flood reducing investments (Shreve and Kelman, 2014; Tesfatsion et al., 2017). The model is capable of replicating historical trends in observed conservation land in Iowa with a Pearson's $r > 0.75$ and a $MAE < 12.5\%$ for a select number of simulations; however, more work is needed to try to validate the model on a micro-level (farm-level) scale. Finally, future work should more fully explore the feedbacks from the hydrologic system to the human system, which is one of the strengths of the agent-based modeling approach (An, 2012).

Code Availability

Model code can be obtained from the corresponding author.

Author Contribution

David Dziubanski and Kristie Franz were the primary model developers and prepared the manuscript. William Gutowski aided with manuscript preparation and editing.

Competing Interests

The authors declare that they have no conflict of interest.

Acknowledgments

Funding for this project was provided by an Iowa State University College of Liberal Arts and Sciences seed grant. We would like to thank all other seed grant participants, including Jean Goodwin, Chris R. Rehmann, William W. Simpkins, Leigh Tesfatsion, Dara Wald, and Alan Wanamaker.

References

- Ahn, K. H. and Merwade, V.: Quantifying the relative impact of climate and human activities on streamflow, *J. Hydrol.*, 515, 257–266, doi:10.1016/j.jhydrol.2014.04.062, 2014.
- An, L.: Modeling human decisions in coupled human and natural systems : Review of agent-based models, *Ecol. Modell.*, 229, 25–36, doi:10.1016/j.ecolmodel.2011.07.010, 2012.
- An, L., Linderman, M., Qi, J., Shortridge, A. and Liu, J.: Exploring Complexity in a Human–Environment System: An Agent-Based Spatial Model for Multidisciplinary and Multiscale Integration, *Ann. Assoc. Am. Geogr.*, 95(1), 54–79, doi:10.1111/j.1467-8306.2005.00450.x, 2005.
- Arbuckle, J. G.: Iowa Farm and Rural Life Poll 2016 Summary Report, Ames, IA., 2017.
- Arbuckle, J. G., Morton, L. W. and Hobbs, J.: Understanding farmer perspectives on climate change adaptation and mitigation: the roles of trust in sources of climate information, climate change beliefs, and perceived risk, *Environ. Behav.*, 1–30, doi:10.1177/0013916513503832,

2013.

Asch, M., Boquet, M. and Nodet, M.: Nudging Methods, in *Data Assimilation: Methods, Algorithms, and Applications*, pp. 120–123, SIAM., 2017.

Axelrod, R. and Tesfatsion, L.: A Guide for Newcomers to Agent-Based Modeling in the Social Sciences, *Handb. Comput. Econ.*, 2, 1647–1659, doi:10.1016/S1574-0021(05)02044-7, 2006.

Di Baldassarre, G., Viglione, A., Carr, G., Kuil, L., Salinas, J. L. and Blöschl, G.: Socio-hydrology: Conceptualising human-flood interactions, *Hydrol. Earth Syst. Sci.*, 17(8), 3295–3303, doi:10.5194/hess-17-3295-2013, 2013.

Barreteau, O., Bousquet, F., Millier, C. and Weber, J.: Suitability of Multi-Agent Simulations to study irrigated system viability: application to case studies in the Senegal River Valley, *Agric. Syst.*, 80(3), 255–275, doi:10.1016/j.agsy.2003.07.005, 2004.

Becu, N., Perez, P., Walker, a, Barreteau, O. and Page, C. L.: Agent based simulation of a small catchment water management in northern Thailand, *Ecol. Modell.*, 170(2–3), 319–331, doi:10.1016/S0304-3800(03)00236-9, 2003.

Berger, T.: Agent-based spatial models applied to agriculture: A simulation tool for technology diffusion, resource use changes and policy analysis, *Agric. Econ.*, 25(2–3), 245–260, doi:10.1016/S0169-5150(01)00082-2, 2001.

Berger, T. and Troost, C.: Agent-based Modelling of Climate Adaptation and Mitigation Options in Agriculture, *J. Agric. Econ.*, 65(2), 323–348, doi:10.1111/1477-9552.12045, 2014.

Berger, T., Birner, R., Mccarthy, N., DíAz, J. and Wittmer, H.: Capturing the complexity of water uses and water users within a multi-agent framework, *Water Resour. Manag.*, 21(1), 129–148, doi:10.1007/s11269-006-9045-z, 2006.

Berglund, E. Z.: Using agent-based modeling for water resources planning and management, *J.*

Water Resour. Plan. Manag., 141(11), 1–17, doi:10.1061/(ASCE)WR.1943-5452.0000544, 2015.

Bharathy, G. K. and Silverman, B.: Holistically evaluating agent-based social systems models: A case study., 2013.

Bithell, M. and Brasington, J.: Coupling agent-based models of subsistence farming with individual-based forest models and dynamic models of water distribution, Environ. Model. Softw., 24(2), 173–190, doi:10.1016/j.envsoft.2008.06.016, 2009.

Bonabeau, E.: Agent-based modeling: Methods and techniques for simulating human systems, Proc. Natl. Acad. Sci. U. S. A., 99(3), 7280–7287, doi:10.1073/pnas.082080899, 2002.

Borrill, P. and Tesfatsion, L.: Agent-based modeling: the right mathematics for the social sciences?, in The Elgar Companion to Recent Economic Methodology, pp. 228–258, New York, New York., 2011.

Brown, C. M., Lund, J. R., Cai, X., Reed, P. M., Zagana, E. A., Ostfeld, A., Hall, J., Characklis, G. W., Yu, W. and Brekke, L.: Scientific Framework for Sustainable Water Management, Water Resour. Res., 6110–6124, doi:10.1002/2015WR017114.Received, 2015.

Burton, R. J. F.: The influence of farmer demographic characteristics on environmental behaviour: A review, J. Environ. Manage., 135, 19–26, doi:10.1016/j.jenvman.2013.12.005, 2014.

Chu, X. and Steinman, A.: Event and Continuous Hydrologic Modeling with HEC-HMS, J. Irrig. Drain. Eng., 135(1), 119–124, doi:10.1061/(ASCE)0733-9437(2009)135:1(119), 2009.

Claassen, R. and Tegene, A.: Agricultural Land Use Choice: A Discrete Choice Approach, Agric. Resour. Econ. Rev., 28(1), 26–36, doi:10.1017/s1068280500000940, 1999.

Cydzik, K. and Hogue, T. S.: Modeling postfire response and recovery using the hydrologic engineering center hydrologic modeling system (HEC-HMS), J. Am. Water Resour. Assoc.,

45(3), doi:10.1111/j.1752-1688.2009.00317.x, 2009.

Daloğlu, I., Nassauer, J. I., Riolo, R. L. and Scavia, D.: Development of a farmer typology of agricultural conservation behavior in the american corn belt, *Agric. Syst.*, 129, 93–102, doi:10.1016/j.agsy.2014.05.007, 2014.

Davis, C. G. and Gillespie, J. M.: Factors affecting the selection of business arrangements by U.S. hog farmers, *Rev. Agric. Econ.*, 29(2), 331–348, doi:10.1111/j.1467-9353.2007.00346.x, 2007.

Du, E., Cai, X., Sun, Z. and Minsker, B.: Exploring the Role of Social Media and Individual Behaviors in Flood Evacuation Processes: An Agent-Based Modeling Approach, *Water Resour. Res.*, 53(11), 9164–9180, doi:10.1002/2017WR021192, 2017.

Duffy, M.: *Conservation Practices for Landlords*, Ames, IA., 2015.

Dziubanski, D. J., Franz, K. J. and Helmers, M. J.: Effects of Spatial Distribution of Prairie Vegetation in an Agricultural Landscape on Curve Number Values, *J. Am. Water Resour. Assoc.*, 53(2), 365–381, doi:10.1111/1752-1688.12510, 2017.

Elshafei, Y., Sivapalan, M., Tonts, M. and Hipsey, M. R.: A prototype framework for models of socio-hydrology: Identification of key feedback loops and parameterisation approach, *Hydrol. Earth Syst. Sci.*, 18(6), 2141–2166, doi:10.5194/hess-18-2141-2014, 2014.

Fagiolo, G., Windrum, P. and Moneta, A.: Empirical validation of agent-based models: A critical survey, *Econ. Policy*, (May), 1–45 [online] Available from:

<http://www.lem.sssup.it/WPLem/files/2006-14.pdf>, 2006.

Fagiolo, G., Guerini, M., Lamperti, F., Moneta, A. and Roventini, A.: *Validation of Agent-Based Models in Economics and Finance* BT - *Computer Simulation Validation: Fundamental Concepts, Methodological Frameworks, and Philosophical Perspectives*, edited by C. Beisbart

and N. J. Saam, pp. 763–787, Springer International Publishing, Cham., 2019.

Frans, C., Istanbuluoglu, E., Mishra, V., Munoz-Arriola, F. and Lettenmaier, D. P.: Are climatic or land cover changes the dominant cause of runoff trends in the Upper Mississippi River Basin?, *Geophys. Res. Lett.*, 40(6), 1104–1110, doi:10.1002/grl.50262, 2013.

Grimm, V., Berger, U., Bastiansen, F., Eliassen, S., Ginot, V., Giske, J., Goss-Custard, J., Grand, T., Heinz, S. K., Huse, G., Huth, A., Jepsen, J. U., Jørgensen, C., Mooij, W. M., Müller, B., Pe'er, G., Piou, C., Railsback, S. F., Robbins, A. M., Robbins, M. M., Rossmanith, E., Rüger, N., Strand, E., Souissi, S., Stillman, R. a., Vabø, R., Visser, U. and DeAngelis, D. L.: A standard protocol for describing individual-based and agent-based models, *Ecol. Modell.*, 198(1–2), 115–126, doi:10.1016/j.ecolmodel.2006.04.023, 2006.

Guerini, M. and Moneta, A.: A method for agent-based models validation, *J. Econ. Dyn. Control*, 82, 125–141, doi:10.1016/j.jedc.2017.06.001, 2017.

Gyawali, R. and Watkins, D. W.: Continuous Hydrologic Modeling of Snow-Affected Watersheds in the Great Lakes Basin Using HEC-HMS, *J. Hydrol. Eng.*, 18(January), 29–39, doi:10.1061/(ASCE)HE.1943-5584.0000591., 2013.

Hahn, H. A.: The conundrum of verification and validation of social science-based models, *Procedia Comput. Sci.*, 16, 878–887, doi:10.1016/j.procs.2013.01.092, 2013.

Halwatura, D. and Najim, M. M. M.: Environmental Modelling & Software Application of the HEC-HMS model for runoff simulation in a tropical catchment, *Environ. Model. Softw.*, 46, 155–162, doi:10.1016/j.envsoft.2013.03.006, 2013.

Helmets, M. J., Zhou, X., Asbjornsen, H., Kolka, R., Tomer, M. D. and Cruse, R. M.: Sediment Removal by Prairie Filter Strips in Row-Cropped Ephemeral Watersheds, *J. Environ. Qual.*, 41(5), 1531, doi:10.2134/jeq2011.0473, 2012.

Hernandez-Santana, V., Zhou, X., Helmers, M. J., Asbjornsen, H., Kolka, R. and Tomer, M.: Native prairie filter strips reduce runoff from hillslopes under annual row-crop systems in Iowa, USA, *J. Hydrol.*, 477, 94–103, doi:10.1016/j.jhydrol.2012.11.013, 2013.

Hoag, D., Luloff, A. E. and Osmond, D.: *How Farmers and Ranchers Make Decisions on Conservation Practices*, Raleigh, NC., 2012.

Hofstrand, D.: *Tracking the Profitability of Corn Production*, Ames, IA., 2018.

Jenkins, K., Surminski, S., Hall, J. and Crick, F.: Assessing surface water flood risk and management strategies under future climate change: Insights from an Agent-Based Model, *Sci. Total Environ.*, 595, 159–168, doi:10.1016/j.scitotenv.2017.03.242, 2017.

Knebl, M. R., Yang, Z., Hutchison, K. and Maidment, D. R.: Regional scale flood modeling using NEXRAD rainfall , GIS , and HEC-HMS / RAS : a case study for the San Antonio River Basin Summer 2002 storm event, *J. Environ. Manage.*, 75, 325–336, doi:10.1016/j.jenvman.2004.11.024, 2005.

Kucharik, C. J.: Evaluation of a process-based agro-ecosystem model (Agro-IBIS) across the U.S. Corn Belt: Simulations of the interannual variability in maize yield, *Earth Interact.*, 7(14), 1–33, doi:10.1175/1087-3562(2003)007<0001:EOAPAM>2.0.CO;2, 2003.

Kulik, B. and Baker, T.: Putting the organization back into computational organization theory: a complex Perrowian model of organizational action, *Comput. Math. Organ. Theory*, 14, 84–119, doi:10.1007/s10588-008-9022-6, 2008.

Lambert, D. M., Sullivan, P., Claassen, R. and Foreman, L.: Profiles of US farm households adopting conservation-compatible practices, *Land use policy*, 24(1), 72–88, doi:10.1016/j.landusepol.2005.12.002, 2007.

Langevin, J., Wen, J. and Gurian, P. L.: *Simulating the human-building interaction:*

Development and validation of an agent-based model of office occupant behaviors, *Build. Environ.*, 88, 27–45, doi:10.1016/j.buildenv.2014.11.037, 2015.

Le, Q., Park, S. and Vlek, P.: Ecological Informatics Land Use Dynamic Simulator (LUDAS): A multi-agent system model for simulating spatio-temporal dynamics of coupled human – landscape system 2. Scenario-based application for impact assessment of land-use policies, *Ecol. Inform.*, 5(3), 203–221, doi:10.1016/j.ecoinf.2010.02.001, 2010.

Macal, C. M., North, M. J., Cirillo, R., Koratorav, V., Thimmapuram, P. and Veselka, T.: Validation of an Agent-based Model of Deregulated Electric Power Markets Abstract EMCAS (Electricity Market Complex Adaptive System) is an agent-based simulation model of the electric power market designed to investigate market restructuring and deregul. [online]

Available from:

<http://www2.econ.iastate.edu/tesfatsi/EmpValidACE.MacalNorth.ElectricPower.pdf>, 2007.

Marcotty, J.: High crop prices a threat to nature?, *StarTribune*, 11th November, 2011.

Matthews, R.: The People and Landscape Model (PALM): Towards full integration of human decision-making and biophysical simulation models, *Ecol. Model.*, 194, 329–343, doi:10.1016/j.ecolmodel.2005.10.032, 2006.

Mays, L.: *Water Resources Engineering*, 2nd ed., John Wiler & Sons, Inc., Hoboken, NJ., 2011.

McGuire, J., Morton, L. W. and Cast, A. D.: Reconstructing the good farmer identity: Shifts in farmer identities and farm management practices to improve water quality, *Agric. Human Values*, 30(1), 57–69, doi:10.1007/s10460-012-9381-y, 2013.

Mcguire, J. M., Wright, L., Arbuckle, J. G. and Cast, A. D.: Farmer identities and responses to the social-biophysical environment, *J. Rural Stud.*, 39, 145–155, doi:10.1016/j.jrurstud.2015.03.011, 2015.

Montanari, A.: Debates-Perspectives on socio-hydrology: Introduction, *Water Resour. Res.*, 51(6), 4768–4769, doi:10.1002/2015WR017430, 2015.

Naik, P. K. and Jay, D. a.: Distinguishing human and climate influences on the Columbia River: Changes in mean flow and sediment transport, *J. Hydrol.*, 404(3–4), 259–277, doi:10.1016/j.jhydrol.2011.04.035, 2011.

Newton, J.: Change on the Horizon for the Conservation Reserve Program?, [online] Available from: <https://www.fb.org/market-intel/change-on-the-horizon-for-the-conservation-reserve-program> (Accessed 15 January 2018), 2017.

Ng, T. L., Eheart, J. W., Cai, X. and Braden, J. B.: An agent-based model of farmer decision-making and water quality impacts at the watershed scale under markets for carbon allowances and a second-generation biofuel crop, *Water Resour. Res.*, 47(9), doi:10.1029/2011WR010399, 2011.

Noel, P. H. and Cai, X.: On the role of individuals in models of coupled human and natural systems : Lessons from a case study in the Republican River Basin, *Environ. Model. Softw.*, 92, 1–16, doi:10.1016/j.envsoft.2017.02.010, 2017.

Nowak, P.: Why farmers adopt production technology, *Soil Water Conserv.*, 47(1), 14–16, 1992.

van Oel, P. R., Krol, M. S., Hoekstra, A. Y. and Taddei, R. R.: Feedback mechanisms between water availability and water use in a semi-arid river basin: A spatially explicit multi-agent simulation approach, in *Environmental Modelling & Software*, vol. 25, pp. 433–443, Elsevier Ltd., 2010.

Ormerod, P. and Rosewell, B.: Validation and Verification of Agent-Based Models in the Social Sciences, *Epistemol. Asp. Comput. Simul. Soc. Sci.*, 5466, 130–140, doi:10.1007/978-3-642-01109-2_10, 2009.

Pahl-wostl, C. and Ebenhöf, E.: Heuristics to characterise human behaviour in agent based models., 2004.

Parker, D. C., Hessler, A. and Davis, S. C.: Complexity, land-use modeling, and the human dimension: Fundamental challenges for mapping unknown outcome spaces, *Geoforum*, 39(2), 789–804, doi:10.1016/j.geoforum.2007.05.005, 2008.

Parunak, H. V. D., Savit, R. and Riolo, R. L.: Multi-agent systems and agent-based simulation, *Proc. First Int. Work. Multi-Agent Syst. Agent-Based Simul.*, 10–25, doi:10.1007/b71639, 1998.

Pfrimmer, J., Gigliotti, L., Stafford, J. and Schumann, D.: Motivations for Enrollment Into the Conservation Reserve Enhancement Program in the James River Basin of South Dakota, *Hum. Dimens. Wildl.*, 22(4), 1–8, doi:10.1080/10871209.2017.1324069, 2017.

Plastina, A.: *Estimated Costs of Crop Production in Iowa - 2017*, Ames, IA., 2017.

Plastina, A., Zhang, W. and Sawadgo, W.: *Iowa Farmland Ownership and Tenure Survey 1982–2017: A Thirty-five Year Perspective*, Ames, IA. [online] Available from: https://www.researchgate.net/publication/326092108_Iowa_Farmland_Ownership_and_Tenure_Survey_1982-2017_A_Thirty-five_Year_Perspective, 2018.

Prior, J.: *Landforms of Iowa*, 1st ed., University of Iowa Press, Iowa City, Iowa., 1991.

Prokopy, L. S., Floress, K., Arbuckle, J. G., Church, S. P., Eanes, F. R., Gao, Y., Gramig, B. M., Ranjan, P. and Singh, A. S.: Adoption of agricultural conservation practices in the United States: Evidence from 35 years of quantitative literature, *J. Soil Water Conserv.*, 74(5), 520–534, doi:10.2489/jswc.74.5.520, 2019.

Reeves, H. W. and Zellner, M. L.: Linking MODFLOW with an agent-based land-use model to support decision making., *Ground Water*, 48(5), 649–60, doi:10.1111/j.1745-6584.2010.00677.x, 2010.

Rogger, M., Agnoletti, M., Alaoui, A., Bathurst, J. C., Bodner, G., Borga, M., Chaplot, V., Gallart, F., Glatzel, G., Hall, J., Holden, J., Holko, L., Horn, R., Kiss, A., Quinton, J. N., Leitinger, G., Lennartz, B., Parajka, J., Peth, S., Robinson, M., Salinas, J. L., Santoro, A., Szolgay, J., Tron, S. and Viglione, A.: Land use change impacts on floods at the catchment scale: Challenges and opportunities for future research, *Water Resources Res.*, 53(June 2013), 5209–5219, doi:10.1002/2017WR020723. Received, 2017.

Rosengrant, M., Cai, X. and Cline, S.: *World water and food to 2025.*, 2002.

Ryan, R. L., Erickson, D. L. and De Young, R.: Farmers' Motivation for Adopting Conservation Practices along Riparian Zones in a Mid-western Agricultural Watershed, *J. Environ. Plan. Manag.*, 46(1), 19–37, doi:10.1080/713676702, 2003.

Saltiel, J., Bauder, J. W. and Palakovich, S.: Adoption of Sustainable Agricultural Practices: Diffusion, Farm Structure, and Profitability, *Rural Sociol.*, 59(2), 333–349, 1994.

Savenije, H. H. G. and Van der Zaag, P.: Integrated water resources management: Concepts and issues, *Phys. Chem. Earth*, 33(5), 290–297, doi:10.1016/j.pce.2008.02.003, 2008.

Schaible, G. D., Mishra, A. K., Lambert, D. M. and Panterov, G.: Factors influencing environmental stewardship in U.S. agriculture: Conservation program participants vs. non-participants, *Land use policy*, 46, 125–141, doi:10.1016/j.landusepol.2015.01.018, 2015.

Scharffenberg, W. A.: *Hydrologic Modeling System User's Manual*, United State Army Corps Eng. [online] Available from: http://www.hec.usace.army.mil/software/hech-hms/documentation/HEC-HMS_Users_Manual_4.0.pdf, 2013.

Schilling, K. E., Chan, K. S., Liu, H. and Zhang, Y. K.: Quantifying the effect of land use land cover change on increasing discharge in the Upper Mississippi River, *J. Hydrol.*, 387(3–4), 343–345, doi:10.1016/j.jhydrol.2010.04.019, 2010.

Schlüter, M. and Pahl-wostl, C.: Mechanisms of Resilience in Common-pool Resource Management Systems : an Agent-based Model of Water Use in a River Basin, *Ecol. Soc.*, 12(2) [online] Available from: <http://www.ecologyandsociety.org/vol12/iss2/art4/>, 2007.

Schmieg, S., Franz, K., Rehmann, C. and van Leeuwen, J. (Hans): Reparameterization and evaluation of the HEC-HMS modeling application for the City of Ames, Iowa, Ames, IA., 2011.

Schreinemachers, P. and Berger, T.: Land use decisions in developing countries and their representation in multi-agent systems, *L. Use Sci.*, 1(1), 29–44, doi:10.1080/17474230600605202, 2006.

Schreinemachers, P. and Berger, T.: An agent-based simulation model of human–environment interactions in agricultural systems, *Environ. Model. Softw.*, 26(7), 845–859, doi:10.1016/j.envsoft.2011.02.004, 2011.

Schwarz, N. and Ernst, A.: Agent-based modeling of the diffusion of environmental innovations - An empirical approach, *Technol. Forecast. Soc. Change*, 76(4), 497–511, doi:10.1016/j.techfore.2008.03.024, 2009.

Secchi, S. and Babcock, B. A.: Impact of High Corn Prices on Conservation Reserve Program Acreage., *Iowa Ag Rev.*, 13(2), 4–7, 2007.

Shreve, C. M. and Kelman, I.: Does mitigation save? Reviewing cost-benefit analyses of disaster risk reduction, *Int. J. Disaster Risk Reduct.*, 10, 213–235, doi:10.1016/j.ijdr.2014.08.004, 2014.

Simon, H.: *Models of Man*, John Wiley & Sons, New York, New York., 1957.

Sivapalan, M. and Blöschl, G.: Time scale interactions and the coevolution of humans and water, *Water Resour. Res.*, 51(9), 6988–7022, doi:10.1002/2015WR017896, 2015.

Sivapalan, M., Savenije, H. H. G. and Blöschl, G.: Socio-hydrology: A new science of people and water, *Hydrol. Process.*, 26(8), 1270–1276, doi:10.1002/hyp.8426, 2012.

Tannura, M. A., Irwin, S. H. and Good, D. L.: Weather, Technology, and Corn and Soybean Yields in the U.S. Corn Belt. [online] Available from:

<http://www.farmdoc.uiuc.edu/marketing/reports>, 2008.

Tesfatsion, L., Rehmann, C. R., Cardoso, D. S., Jie, Y. and Gutowski, W. J.: An agent-based platform for the study of watersheds as coupled natural and human systems, *Environ. Model. Softw.*, 89, 40–60, doi:10.1016/j.envsoft.2016.11.021, 2017.

Tigner, R.: *Partial Budgeting: A Tool to Analyze Farm Business Changes*, Ames, IA., 2006.

Tomer, M. D. and Schilling, K. E.: A simple approach to distinguish land-use and climate-change effects on watershed hydrology, *J. Hydrol.*, 376(1–2), 24–33,

doi:10.1016/j.jhydrol.2009.07.029, 2009.

Troy, T., Pavao-Zuckerman, M. and Evans, T.: Debates—Perspectives on socio-hydrology: Socio-hydrologic modeling: Tradeoffs, hypothesis testing, and validation, *Water Resour. Res.*, 51, 4806–4814, doi:10.1002/2015WR017046, 2015.

Tyndall, J. C., Schulte, L. A., Liebman, M. and Helmers, M.: Field-level financial assessment of contour prairie strips for enhancement of environmental quality, *Environ. Manage.*, 52(3), 736–747, doi:10.1007/s00267-013-0106-9, 2013.

USDA-Natural Resources Conservation Service (USDA-NRCS): *National Engineering Handbook, Part 630*, Washington, DC., 2004.

USDA-Natural Resources Conservation Service (USDA-NRCS): *Field Office Technical Guide*, [online] Available from: <http://www.nrcs.usda.gov/wps/portal/nrcs/main/national/technical/fotg/> (Accessed 9 April 2016), 2015.

USDA: *Conservation Reserve Program*. [online] Available from: www.nrcs.usda.gov/programs/crp, 2011.

USDA National Agricultural Statistics Service: 2018 Iowa Agricultural Statistics, Des Moines, Iowa., 2018.

Verma, A. K., Jha, M. K. and Mahana, R. K.: Evaluation of HEC-HMS and WEPP for simulating watershed runoff using remote sensing and geographical information system, *Paddy Water Environ.*, 8, 131–144, doi:10.1007/s10333-009-0192-8, 2010.

Viglione, A., Di Baldassarre, G., Brandimarte, L., Kuil, L., Carr, G., Salinas, J. L., Scolobig, A. and Bloßchl, G.: Insights from socio-hydrology modelling on dealing with flood risk - Roles of collective memory, risk-taking attitude and trust, *J. Hydrol.*, 518, 71–82, doi:10.1016/j.jhydrol.2014.01.018, 2014.

Vorosmarty, C. and Sahagian, D.: Anthropogenic Disturbance of the Terrestrial Water Cycle, *Bioscience*, 50(9), 753–765, doi:http://dx.doi.org/10.1641/0006-3568(2000)050[0753:ADOTTW]2.0.CO;2, 2000.

Wainwright, J.: Can modelling enable us to understand the rôle of humans in landscape evolution?, *Geoforum*, 39(2), 659–674, doi:10.1016/j.geoforum.2006.09.011, 2008.

Wang, D. and Hejazi, M.: Quantifying the relative contribution of the climate and direct human impacts on mean annual streamflow in the contiguous United States, *Water Resour. Res.*, 47(9), doi:10.1029/2010WR010283, 2011.

Windrum, P., Fagiolo, G. and Moneta, A.: Empirical Validation of Agent-Based Models: Alternatives and Prospects, *J. Artif. Soc. Soc. Simul.*, 10(2), 2007.

Xiang, X., Kennedy, R. and Madey, G.: Verification and Validation of Agent-based Scientific Simulation Models, *Agent-Directed Simul. Conf.*, 47–55 [online] Available from: http://www.nd.edu/~nom/Papers/ADS019_Xiang.pdf, 2005.

Yang, L. E., Scheffran, J., Süsner, D., Dawson, R. and Chen, Y. D.: Assessment of Flood Losses

with Household Responses: Agent-Based Simulation in an Urban Catchment Area, *Environ. Model. Assess.*, 23(4), 369–388, doi:10.1007/s10666-018-9597-3, 2018.

Zenobia, B., Weber, C. and Daim, T.: Artificial markets : A review and assessment of a new venue for innovation research, *Technovation*, 29, 338–350, doi:10.1016/j.technovation.2008.09.002, 2009.

Zhang, H. L., Wang, Y. J., Wang, Y. Q., Li, D. X. and Wang, X. K.: The effect of watershed scale on HEC-HMS calibrated parameters: A case study in the Clear Creek watershed in Iowa, US, *Hydrol. Earth Syst. Sci.*, 17(7), 2735–2745, doi:10.5194/hess-17-2735-2013, 2013.

Zhang, W.: Who Owns and Rents Iowa's Farmland?, *Ag Decis. Mak.*, C2-78(December), 1–7, 2015.

Zhou, X., Helmers, M. J., Asbjornsen, H., Kolka, R. and Tomer, M. D.: Perennial Filter Strips Reduce Nitrate Levels in Soil and Shallow Groundwater after Grassland-to-Cropland Conversion, *J. Environ. Qual.*, 39(6), 2006, doi:10.2134/jeq2010.0151, 2010.

Zhou, X., Helmers, M. J., Asbjornsen, H., Kolka, R., Tomer, M. D. and Cruse, R. M.: Nutrient removal by prairie filter strips in agricultural landscapes, *J. Soil Water Conserv.*, 69, 54–64, doi:10.2489/jswc.69.1.54, 2014.

Variable	Description	Unit
$C_{t-1:t-X}$	Mean total amount of land allocated to conservation during the previous X years	Hectares
D_{t-1}	Previous year's conservation land decision	Hectares
$\delta C_{futures:Y}$	Conservation decision based on crop price projections for Y years into the future	Hectares
$\delta C_{profit:X}$	Conservation decision based on mean past profit of previous X years	Hectares
δC_{cons}	Conservation decision based on conservation goal	Hectares
$C_{neighbor}$	Weighted mean conservation land of the farmer agent's neighbors	Hectares
$Profit_{diff}$	Differences in profit between an acre of crop and an acre of conservation land	(\$/Hectare)
$Hectares_{tot}$	Total land owned by farmer agent	Hectares
G_t	Government agent conservation goal for the current year t	Hectares
G_{t-1}	Unfulfilled conservation land from the previous year's t-1 conservation goal	Hectares
A_{tot}	Total agricultural land in watershed	Hectares
C_{tot}	Total land currently in conservation	Hectares
P	Total conservation land to be added to the goal as a percentage of production land	Dimensionless
P_{new}	Variable describing change in conservation goal with flood damage	(1/\$)

Table 1. Variables in farmer and city agent equations.

Agent Model Parameters	Description	Range
$W_{risk-averse}$	Weight placed on farmer agent's previous land use	0.0 - 1.0
$W_{futures}$	Weight placed on farmer agent's decision based on future crop price	0.0 - 1.0
W_{profit}	Weight placed on farmer agent's decision based on past profit	0.0 - 1.0
W_{cons}	Weight placed on farmer agent's decision based on his/her conservation goal	0.0 - 1.0
$W_{neighbor}$	Weight placed on farmer agent's decision based on his/her neighbor's decisions	0.0 - 1.0
$Cons_{max}$	Farmer's conservation goal - used to describe the farmer's conservation-mindedness	0.0 - 0.1
X	Number of previous years a farmer agent takes into account for his/her land decision	1 - 5
Y	Number of future years a farmer agent takes into account for his/her land decision	5 - 10
$ConsGoal_{max}$	Conservation goal at maximum flood damage	0.0 - 0.1

Table 2. Primary agent model parameters in decision-making equations.

Decision Scheme	Decision Weight				
	Conservation Goal	Futures	Past Profit	Risk Aversion	Neighbor
Conservation	0.8	0.05	0.05	0.05	0.05
Future price	0.05	0.8	0.05	0.05	0.05
Past profit	0.05	0.05	0.8	0.05	0.05
Risk averse	0.05	0.05	0.05	0.8	0.05

Table 3. Decision weighting scheme tested with each scenario.

Model Inputs	Years	Unit
Historical Cash Rent	1970-2016	(\$/Hectare)
Federal Subsidies	2000-2016	(\$/Hectare)
Historical Production Costs	1970-2016	(\$/Hectare)
Historical Corn Prices	1970-2016	(\$/MT)
Precipitation	1970-2016	(mm/hr)

Table 4. Model Inputs.