

## Response to the referees

The authors gratefully thank to the editor and referees for their critical comments on our manuscript which drives us to improve the manuscript greatly. The comments and questions were addressed point by point.

### Response to Referee #1

#### General Comments

My main comment is there are still some works worth being discussed, though this review is overall complete: 1) The perspective of Lhomme and Guillioni (2006, 2010) which relates potential evaporation to surface resistance. 2) Aminzadeh et al. (2016)'s CR with  $E_p$  defined by a surface temperature. Also, there are a few latest CR studies in 2019 that are highly relevant to the submitted manuscript, e.g., Anayah & Kaluarachchi (2019) and Brutsaert et al. (2019). Could the authors discussed a little bit?

Response: Thanks for the Referee's suggestions. We discussed these publications in the revised manuscript.

In Lhomme and Guillioni (2006, 2010)'s perspective of CR, the surface resistance is related to the equilibrium evaporation and Penman's potential evaporation. We found that it is a linear function relating  $E/E_{Pen}$  to  $E_{rad}/E_{Pen}$  without intercept in the normalized form. Thus, we added following paragraph to Section "Normalized complementary functions". Please refer to Lines 216-221 of the revised manuscript.

Aminzadeh et al. (2016) derived a steady state surface temperature via the surface energy balance at which the sensible heat flux is zero, and calculated  $E_{pa}$  and  $E_{po}$  using a mass-transfer type reference evaporation corresponding to current and saturated surface water content. We discussed it in addition to the works of Morton (1983) and Szilagyi and Jozsa (2008) in the revised manuscript (Lines 193-195)

The latest studies of using CR for global ET estimation is discussed in the revised manuscript as "The polynomial B2015 was applied to estimate global terrestrial evaporation with calibrated  $\alpha$  as a function of aridity index (Brutsaert et al., 2019). The modified Granger's model was also applied for estimating global evaporation with 30 min spatial resolution and monthly time steps (Anayah and Kaluarachchi, 2019)."

#### Specific Comments

Line 19: Is the boundary condition here specified to the wet environment?

Response: Yes. We added "under wet environments" after it in the revised manuscript.

Line 40: Recent publications using GCR in 2019 for estimating evaporation should be

added here.

Response: We added Brutsaert's latest work here.

Brutsaert, W., Cheng, L., and Zhang, L.: Spatial Distribution of Global Landscape Evaporation in the Early Twenty First Century by Means of a Generalized Complementary Approach, J Hydrometeorol, 10.1175/jhm-d-19-0208.1, 2019.

Line 87: "while" should be replaced by "whereas"?

Response: It was revised.

Line 107: the "realistic" is compared to the former model. I think adding "more" here may be better.

Response: "more" was added here.

Line 110: "wss"??

Response: We are sorry for the typo. It should be "was"

Line 167: "The asymmetric CR is widely used", please revised this sentence

Response: We replaced this sentence by "The asymmetric CR is a significant improvement of the symmetric CR, and the opposite changes of  $E/E_{po}$  and  $E_{pa}/E_{po}$  against  $E/E_{pa}$  were treated as an enhanced illustration of the CR (Hu et al., 2018; Zhang et al., 2017; Ma et al., 2015; Brutsaert et al., 2019; Szilagyi, 2007)."

Line 179: More statements on the asymmetric CR should be added, including the negative relationship between  $E/E_{po}$  and  $E_{pa}/E_{po}$  was treated as an extension of the original CR, and the validation in several locations.

Response: More statements on the asymmetric CR were added as "The asymmetric CR is a significant improvement of the symmetric CR, and the opposite changes of  $E/E_{po}$  and  $E_{pa}/E_{po}$  against  $E/E_{pa}$  were treated as an enhanced illustration of the CR (Hu et al., 2018; Zhang et al., 2017; Ma et al., 2015; Brutsaert et al., 2019; Szilagyi, 2007). The performances on evaporation estimation are improved by calibrating the asymmetry parameter  $b$  (Kahler and Brutsaert, 2006; Han et al., 2008; Huntington et al., 2011; Ma et al., 2015). Efforts have also made to calculate  $b$  by using the meteorological variables, which enhance the predict ability of the CR (Szilagyi, 2015; Szilagyi, 2007; Aminzadeh et al., 2016). However, the changes in  $b$  imply a potential nonlinear characteristic of the CR (Han, 2008; Lintner et al., 2015). The observed values of  $E/E_{po}$  and  $E_{pa}/E_{po}$  even exhibit a positive correlation under wet conditions at several flux sites, which challenges the linear CR (Han and Tian, 2018)."

Line 246: "Han and Tian (2018) further validated the sigmoid feature": Please state

the work more detailed because there are still controversies on it.

Response: We revised this sentence as “Han and Tian (2018) further validated the sigmoid feature according to the much larger regression slopes of  $E/E_{pen}$  upon  $E_{rad}/E_{pen}$  in the middle stage than those in the other two stages with smaller or larger values of  $E_{rad}/E_{pen}$  by using 22 eddy covariance towers from the FLUXNET (Baldocchi et al., 2001) dataset which includes representative biomes of grasslands, croplands, shrublands, evergreen needleleaf forests, deciduous broadleaf forests, and wetlands.”

Line 270: What is the essential difference between B15 and H12? Is “B15 inherits all three types of evaporation dated from the original CR”? Please rearrange these sentences.

Response: The two generalized complementary approaches, H12 and B15, are essentially different, with completely different normalized variables (Table 3). B15 inherits the concept of the three types of evaporation dated from the original CR, and its boundary conditions and analytical form are derived for  $x_B = E_{po}/E_{pa}$  and  $y_B = E/E_{pa}$ . By contrast, H12 goes much further from the original CR. The boundary conditions and the analytical form of H12 are derived for  $x_H = E_{rad}/E_{pen}$  and  $y_H = E/E_{pen}$ . We added a new subsection “Comparisons between the two generalized complementary approaches” to discuss the essential difference between B15 and H12. Please refer to subsection “3.4 Comparisons between the two generalized complementary approaches” for details.

Line 304: The varying characteristics of the PT coefficient should be introduced here

Response: We introduced it in the revised manuscript, as “the Priestley-Taylor coefficient varies with several factors, such as the relative transport efficiency of turbulent, or the surface/air temperature (Assouline et al., 2016; Szilagyi, 2014)”

Line 359: Brutsaert’s recent work by using  $c=0$  and varying PT coefficient should be added. Check Brutsaert et al. (2019).

Response: We added it in the revised manuscript

The Conclusion part could be improved. I wonder are there any outlooks for future studies on CR could be summarized using a few sentences here?

Response: We revised the conclusion part and added three points about future studies: 1) “further validation and application of the two types of generalized complementary functions are required with multiple data over the world”; 2) “be

carefully examined for its physical base of the boundary conditions under completely wet environment”, and 3) “be integrated with other approaches to include the information of the land surface properly”. Please refer to the revised manuscript for details.

## **Response to Referee #2**

### **General Comments**

The authors present a very detailed review on the studies and developments of the complementary relationship of evaporation. Although the review is very detailed and scientifically well supported on the existing literature, I think it is too heavy due to the load of parameters introduced and unexplained, the long list of studies mentioned and a weak coherency when enumerating the studies. Can the authors make this easier for the reader to read through?

Response: In the revised manuscript, we tried our best to make it easier for the reader to read through the following aspects: 1) Section “2 Symmetric complementary relationship” and “3 Efforts in maintaining a linear complementary relationship” of the original manuscript was be combined to one section “Linear complementary relationship” to make it more coherent. 2) Subsection “4.5 Improved understanding on the correlation between actual and potential evaporation” of the original manuscript was deleted by considering it is no tightly related to the whole review. 3) The parameters was checked and introduced more clearly in the revised manuscript. 4) We tried to improve the presentation and English.

Furthermore, I am completely missing the incites and perspectives from the authors. It would be very nice to see the opinion from the authors regarding the benefits of the framework. I suggest an extra section discussing 1) the best approach according to the authors criteria, 2) the future of the CR for E estimation, and 3) a comparison highlighting the advantages, disadvantages and opportunities of using the CR principle against other methods of Evaporation estimation that are not mentioned here. After this, the review should be ready for publication.

Response: In the revised manuscript, we tried to give a clear point of view on the different approaches: 1) “The asymmetric CR is a significant improvement of the symmetric CR”; 2) “The generalized complementary principle with earlier linear CRs as special cases has a more rigorous physical base (Brutsaert, 2015; Han and Tian, 2018b), and its methodology based on nonlinear functions is robust and effective.”

We compared the two generalized complementary functions (the sigmoid H2017 and polynomial B2015) in a new subsection “3.4 Comparisons between the two generalized complementary approaches”.

We added a new subsection “4.3 Integrating with other approaches for further development” to discuss the current practice and future development of the CR for E estimation, and to compare it with other methods (the Penman approach, the Budyko approach and others) on the advantages, disadvantages and opportunities of using the

CR principle. At last, we proposed a suggestion of integrating these approaches for a new generation of evaporation estimation method.

### Specific Comments

L. 24 State if it is a positive or negative feedback.

Response: We revised this sentence as “this principle originated from the negative feedback of areal evaporation on evaporation demand (Bouchet, 1963)”.

L. 27 To understand, so the differences between  $E_{pa}$  and  $E_{po}$  is just that  $E_{pa}$  is small and local and  $E_{po}$  large-scale? Can you provide more explanation on what these two variables really mean since they are so important for this discussion? Specially for understanding Figure 1.

Response: The major differences between  $E_{pa}$  and  $E_{po}$  are that they correspond to different atmosphere characteristics.  $E_{pa}$  corresponds to the atmosphere in contact with current non-saturated evaporating surface as the overpassing air is not affected by the small saturated surface, whereas the atmosphere corresponding to  $E_{po}$  is in contact with the large-scale saturated surface. Thus, the surface water availability can be detected from the relative magnitude of  $E_{pa}$  and  $E_{po}$  (as shown in Figure 1), and  $E$  can be estimated without the knowledges of the surface. We provided more explanation in the revised manuscript (Lines 30-34).

L. 32 What complex formulations?

Response: The formulations of  $E_{pa}$  and  $E_{po}$  are introduced in Subection 2.1 and 2.4 of the revised manuscript. We will add a tip (“which will be reviewed in more detail in the following sections”) in the revised manuscript.

L. 35-39 Please rephrase, it is difficult to understand.

L. 38 If there is a Penman calculation variable  $E_{pen}$ , then how do you estimate  $E_{pa}$  and  $E_{po}$ , that is different from Penman. Please specify.

Response: It should be noted that  $E_{pa}$  and  $E_{po}$  are theoretical concepts, and need to be formulated when applying for practical problems. The generalized complementary function comes in two ways. Brutsaert (2015) adopted a polynomial function to describe the relationship between  $E$ ,  $E_{pa}$  and  $E_{po}$ , and suggested to formulate  $E_{pa}$  and  $E_{po}$  by using Penman's potential evaporation ( $E_{pen}$ ) and Priestley-Taylor's minimal advection evaporation to formulate, respectively. By contrast, Han and Tian (2018) abandoned the theoretical concept of  $E_{pa}$  and  $E_{po}$ , yet used a sigmoid function to describe the relationship among  $E$ , Penman's potential evaporation ( $E_{pen}$ ), and its radiation term ( $E_{rad}$ ), which can be directly used for practical problems. We revised this paragraph as:

“Recent studies have adopted the “generalized” complementary principle, which employs nonlinear functions instead of the linear CR (Han et al., 2012; Brutsaert, 2015; Han and Tian, 2018). The generalized complementary function comes in two ways, with the first abandons the theoretical concept of  $E_{pa}$  and  $E_{po}$  yet uses a sigmoid function to describe the relationship among  $E$ , Penman's potential evaporation ( $E_{pen}$ ), and its radiation term ( $E_{rad}$ ) (Han and Tian, 2018; Han et al., 2012). By contrast, the other adopts a polynomial function to describe the relationship between  $E$ ,  $E_{pa}$  and  $E_{po}$ . However,  $E_{pa}$  and  $E_{po}$  need to be formulated before applying the polynomial function to practical problems (Brutsaert, 2015).”

Table 1. Nice table!, but refer to the Appendix for the unexplained parameters.

Response: We added it below the table.

L. 69 why “basin-wide water balance” results? You said before that  $E_{pa}$  is from a “small saturated surface”.

Response:  $E_{pa}$  was derived from a hypothetical theoretical concept of “small saturated surface”, which means that the “small saturated surface” does not affect the atmosphere. Thus, it is determined by the atmosphere corresponding to current unsaturated surface. In application,  $E_{pa}$  is calculated by using the meteorological variables corresponding to current unsaturated surface, and is used to calculate the basin-wide actual evaporation.

L. 70-75 But have these estimates been validated in some way?

Response: Yes. We pointed out it in the revised manuscript.

L. 75 When they found that it is overestimating or underestimating  $E$ , how did these studies obtain the real  $E$  then?

Response: In the revised manuscript, we added the potential causes of the bias “measurement error, imperfect formulations of  $E_{pa}$  and/or  $E_{po}$ , external energy sources, or even the nonlinear nature of the complementary principle were considered as potential causes of this bias (Qualls and Gultekin, 1997; Hobbins et al., 2001; Han et al., 2008, 2012).” There are two approaches to obtain the real  $E$ : 1) maintaining a linear complementary relationship through properly formulating  $E_{pa}$  and/or  $E_{po}$ ; 2) using a nonlinear generalized complementary function. Please refer to section “2.4 Efforts in”, and “3 Generalized complementary principle via nonlinear functions” for the detailed methods.

L. 78 IMPORTANT. Since you are constantly introducing many parameters related to actual or potential evaporation, please include in the appendix a detailed explanation on the difference between each  $E$  parameter. For instance, to know how  $E_{pan}$  differs

from Epa.

Response: Generally speaking,  $E_{pa}$  and  $E_{po}$  are theoretical concepts, whereas Epan, Epen, Ept and others are the specifications of them. They are introduced in the appendix to “Three types of evaporation in CR” and “Specific formulations for  $E_{pa}$  or  $E_{po}$ ” respectively in the revised manuscript.

L. 88 Do you mean that they change in opposite directions with increasing water availability?

Response: Yes. We changed “while” to “whereas” to make the sentence more clear.

L95 “the governing changes”

Response: We changed it in the revised manuscript.

L. 98 Why does Morton say that it is unrealistic and does not have proof, and argue against it, since you are performing a review on the subject.

Response: Morton derived the CR by two assumptions: “(1) the net radiation will not change with the surface, and (2) the heat and vapor eddy transfer characteristics are identical for  $E$  and  $E_{pa}$ . Relaxing the second assumption of Morton (1983), Szilagyi (2001) derived the CR by using the mass conservation equation for water vapor. However, LeDrew (1979) argued that Morton's two assumption do not necessarily hold, and pointed out that the symmetric CR is physically unrealistic by using a diagnostic model of the energy fluxes within a closed system.”

L. 113 You mention an asymmetry, but before you were talking about symmetry?

Response: We revised it to “linear CR”.

L. 136 So when is  $E_{PT}$  different from  $E_{po}$ . In other words, more clarity between these terms.

Response: Please refer Lines 174-180 of the revised manuscript:

“In theory,  $E_{po}$  is the potential evaporation when the land surface is saturated, and should be calculated with a proper formula by using meteorological variables corresponding to the “potential” saturated surface. The Priestley-Taylor equation has been widely accepted to represent evaporation from extensive saturated surfaces by using meteorological variables corresponding to these saturated surfaces (Brutsaert, 1982; Priestley and Taylor, 1972). This way it was suggested to represent  $E_{po}$  (Brutsaert and Stricker, 1979). However, in the AA approach,  $E_{PT}$  is calculated by the Priestley-Taylor equation using the atmospheric variables that correspond to the



current unsaturated surface. But the atmosphere in contact with the land surface will change if the land surface was brought into saturated (Morton, 1983; Brutsaert, 2015). Thus,  $E_{PT}$  is in reality a variable dependent on the meteorological variables at the time of calculation and does not represent the “true”  $E_{po}$ .”

L. 135-156 IMPORTANT I find these paragraphs hard to read and somehow “boring”. As in a review, it would be very good if you can try to articulate all the studies in a more consistent way so that it does not become a list of studies and references each with a brief explanation. Also, many, many terms that have not been previously explained, only in an appendix. As it is, the review paper is now more focus to experts in the CR that common hydrologists.

Response: We rephrased this paragraph to make is more consistent and easy to read. In the first paragraph, the first problem of using EPT to denote  $E_{po}$  is pointed, and is explained as “predicting the air temperature corresponding to the extensive saturated surface is critical for properly formulating  $E_{po}$ ”. Then, three works aiming to settle this problem are introduced one by one: Morton (1983), Szilagyi and Jozsa (2008), and Aminzadeh et al. (2016).

Next, another problem is pointed as “Advection is another factor influencing  $E_{po}$ , which could not adequately considered by  $E_{PT}$  with an assumption of a minimal advection effect (Morton, 1983, 1975; Parlange and Katul, 1992).” The works of Morton (1983) and Parlange and Katul (1992) are introduced.

Section 3. I don’t see the rationale behind the selection of the subtitles 3.1 and 3.2. A brief explanation is needed. Why these subtitles, I assume 3.1 are the symmetric approaches and 3.2 the asymmetric ones? Think on the reader that is reading this review.

Response: We combined section 2 and 3 and change the order of former subsection 3.1 and 3.2. We believe it would be more rationale following this order:

- 2.1 Concept of symmetric complementary relationship;
  - 2.2 Proofs of symmetric CR;
  - 2.3 Asymmetric linear CR as an extension;
  - 2.4 Efforts in maintaining a linear complementary relationship through.
- The new 2.4 is for both the symmetric and asymmetric CR.

L.164 so b=1 means symmetry?

Response: Yes. We will add this in the revised manuscript.

L. 167. “The asymmetric CR is widely used?” Can you make a paragraph saying in your point of view which approach is better and why, symmetric or asymmetric?

Response: We deleted this sentence and add a paragraph as:



“The asymmetric CR is a significant improvement of the symmetric CR, and the opposite changes of  $E/E_{po}$  and  $E_{pa}/E_{po}$  against  $E/E_{pa}$  were treated as an enhanced illustration of the CR (Hu et al., 2018; Zhang et al., 2017; Ma et al., 2015; Brutsaert et al., 2019; Szilagyi, 2007). The performances on evaporation estimation are improved by calibrating the asymmetry parameter  $b$  (Kahler and Brutsaert, 2006; Han et al., 2008; Huntington et al., 2011; Ma et al., 2015). Efforts have also made to calculate  $b$  by using the meteorological variables, which enhance the predict ability of the CR (Szilagyi, 2015; Szilagyi, 2007; Aminzadeh et al., 2016). However, the changes in  $b$  imply a potential nonlinear characteristic of the CR (Han, 2008; Lintner et al., 2015). The observed values of  $E/E_{po}$  and  $E_{pa}/E_{po}$  even exhibit a positive correlation under wet conditions at several flux sites, which challenges the linear CR (Han and Tian, 2018). ”

Can you make a similar Table 1 but for the non-linear relationships? I think that you mention many approaches that are not included in Table 2.

Response: We added a Table for the nonlinear generalized complementary functions as:

**Table 2. Different analytical formulas for generalized complementary functions H12**

Type	Formula <sup>*</sup>	References
Linear	$y = \alpha(1 + \frac{1}{b})x - \frac{1}{b}$	Brutsaert and Stricker (1979)
	$y = (1 + \omega)x$	Lhomme and Guillioni (2010, 2006)
Sigmoid	$y = \frac{1}{1 + c_1 e^{d(1-x)}}$	Granger (1989), Han et al. (2011)
	$y = \frac{1}{1 + m(\frac{1}{x} - 1)^n}$	Han et al. (2012)
	$y = \frac{1}{1 + m(\frac{x_{\max} - x}{x - x_{\min}})^n}$	Han and Tian (2018)
Concave	$y = \frac{1}{1 + k(\frac{1}{x} - 1) + l}$	Katerji and Perrier (1983) , Han et al. (2014b)
	$y = (2 - c)\alpha^2 x^2 - (1 - 2c)\alpha^3 x^3 - c\alpha^4 x^4$	Brutsaert (2015)

<sup>\*</sup>  $x = E_{rad}/E_{Pen}$  and  $y = E/E_{Pen}$  . the other symbols are parameters.

### Response to Referee #3

The authors provided an informative yet in-depth review of the research activities over the past half-century using the complementary principle of evaporation. I enjoyed the reading and feel it is overall a timely and nice contribution to the hydrology community. Nevertheless, I do challenge the authors to elevate the presentation quality, in a spirit to make it valuable for a variety of audiences, from those who are not very familiar with this type of research but would like to have some background information, to those who are really active in the field. In its current form, the writing style is still, more or less, in favor of the latter.

Response: Thanks for the positive comment to our manuscript.

We tried our best to make the review easier for the reader to read through. In the revised manuscript, we added a new subsection “Integrating with other approaches for further development” to compare the complementary approach of evaporation estimation with the approaches which are more popular in the research community, such as the Penman approach, the Budyko approach et., and propose a suggestion of integrating these approaches with the complementary principle for evaporation estimation. We think this new section would help the readers who are not very familiar with the complementary principle, but are interested in evaporation research.

Also, I feel it might be more straightforward to include the word "review" in the title and abstract. Otherwise, some readers may not realize this is a review article until the end of the introduction.

Response: We changed the title to “A review of complementary principle of evaporation: From original linear relationship to generalized nonlinear functions”, and included the word “review” in the abstract and main text.

### Reference:

- Aminzadeh, M., Roderick, M. L., and Or, D.: A generalized complementary relationship between actual and potential evaporation defined by a reference surface temperature, *Water Resour Res*, 52, 385-406, 10.1002/2015wr017969, 2016.
- Anayah, F. M., and Kaluarachchi, J. J.: Estimating Global Distribution of Evapotranspiration and Water Balance Using Complementary Methods, *Atmos Ocean*, 57, 279-294, 10.1080/07055900.2019.1656052, 2019.
- Assouline, S., Li, D., Tyler, S., Tanny, J., Cohen, S., Bou-Zeid, E., Parlange, M., and Katul, G. G.: On the variability of the Priestley-Taylor coefficient over water bodies, *Water Resour Res*, 52, 150-163, 10.1002/2015wr017504, 2016.
- Baldocchi, D., Falge, E., Gu, L., Olson, R., Hollinger, D., Running, S., Anthoni, P., Bernhofer, C., Davis, K., and Evans, R.: FLUXNET: A new tool to study the temporal and spatial variability of ecosystem-scale carbon dioxide, water vapor, and energy flux densities, *B Am Meteorol Soc*, 82, 2415-2434, 2001.
- Bouchet, R.: Evapotranspiration réelle et potentielle, signification climatique, *International Association of Hydrological Sciences Publication*, 62, 134-142, 1963.
- Brutsaert, W., and Stricker, H.: An advection-aridity approach to estimate actual regional

evapotranspiration, *Water Resour Res*, 15, 443-450, 1979.

Brutsaert, W.: *Evaporation into the Atmosphere: Theory, History, and Applications*, D. Reidel-Kluwer, Hingham, 1982.

Brutsaert, W.: A generalized complementary principle with physical constraints for land-surface evaporation, *Water Resour Res*, 51, 8087–8093, doi:8010.1002/2015WR017720., 10.1002/2015wr017720, 2015.

Brutsaert, W., Cheng, L., and Zhang, L.: Spatial Distribution of Global Landscape Evaporation in the Early Twenty First Century by Means of a Generalized Complementary Approach, *J Hydrometeorol*, 10.1175/jhm-d-19-0208.1, 2019.

Han, S.: Study on Complementary Relationship of Evapotranspiration and its Application in Tarim River Basin, Ph.D, Tsinghua University, Beijing, 2008.

Han, S., Hu, H., and Tian, F.: Evaluating the advection-aridity model of evaporation using data from field-sized surfaces of HEIFE, IAHS Publication, 322, 9-14, 2008.

Han, S., Hu, H., and Tian, F.: A nonlinear function approach for the normalized complementary relationship evaporation model, *Hydrol Process*, 26, 3973-3981, 2012.

Han, S., and Tian, F.: Derivation of a sigmoid generalized complementary function for evaporation with physical constraints, *Water Resour Res*, 54, 5050-5068, doi: 10.1029/2017WR021755, 2018.

Hobbins, M. T., Ramí´rez, J. A., Brown, T. C., and Claessens, L. H. J. M.: The complementary relationship in estimation of regional evapotranspiration: the CRAE and Advection-aridity models, *Water Resour Res*, 37, 1367-1388, 2001.

Hu, Z., Wang, G., Sun, X., Zhu, M., Song, C., Huang, K., and Chen, X.: Spatial - Temporal Patterns of Evapotranspiration Along an Elevation Gradient on Mount Gongga, Southwest China, *Water Resour Res*, 2018.

Huntington, J. L., Szilagyi, J., Tyler, S. W., and Pohll, G. M.: Evaluating the complementary relationship for estimating evapotranspiration from arid shrublands, *Water Resour Res*, 47, W05533, 2011.

Kahler, D. M., and Brutsaert, W.: Complementary relationship between daily evaporation in the environment and pan evaporation, *Water Resour Res*, 42, W05413, doi:05410.01029/02005WR004541, 2006.

LeDrew, E. F.: A diagnostic examination of a complementary relationship between actual and potential evapotranspiration, *J Appl Meteorol*, 18, 495-501, 1979.

Lhomme, J. P., and Guilioni, L.: Comments on some articles about the complementary relationship, *J Hydrol*, 323, 1-3, 2006.

Lhomme, J. P., and Guilioni, L.: On the link between potential evaporation and regional evaporation from a CBL perspective, *Theor Appl Climatol*, 101, 143-147, 2010.

Lintner, B., Gentine, P., Findell, K., and Salvucci, G.: The Budyko and complementary relationships in an idealized model of large-scale land-atmosphere coupling, *Hydrol Earth Syst Sc*, 19, 2119-2131, 2015.

Ma, N., Zhang, Y., Szilagyi, J., Guo, Y., Zhai, J., and Gao, H.: Evaluating the complementary relationship of evapotranspiration in the alpine steppe of the Tibetan Plateau, *Water Resour Res*, 51, 1069-1083, 10.1002/2014wr015493, 2015.

Morton, F. I.: Estimating evaporation and transpiration from climatological observations, *J Appl Meteorol*, 14, 488-497, 1975.

Morton, F. I.: Operational estimates of areal evapotranspiration and their significance to the science

and practice of hydrology, *J Hydrol*, 66, 1-76, 1983.

Parlange, M. B., and Katul, G. G.: An advection-aridity evaporation model, *Water Resour Res*, 28, 127-132, 1992.

Priestley, C. H., and Taylor, R. J.: On the assessment of surface heat flux and evaporation using large-scale parameters, *Mon Weather Rev*, 100, 81-92, 1972.

Qualls, R. J., and Gultekin, H.: Influence of components of the advection-aridity approach on evapotranspiration estimation, *J Hydrol*, 199, 3-12, 1997.

Szilagyi, J.: On Bouchet's complementary hypothesis, *J Hydrol*, 246, 155-158, 2001.

Szilagyi, J.: On the inherent asymmetric nature of the complementary relationship of evaporation, *Geophys Res Lett*, 34, L02405, doi:02410.01029/02006GL028708, 2007.

Szilagyi, J.: Temperature corrections in the Priestley–Taylor equation of evaporation, *J Hydrol*, 519, 455-464, 10.1016/j.jhydrol.2014.07.040, 2014.

Szilagyi, J.: Complementary-relationship-based 30 year normals (1981-2010) of monthly latent heat fluxes across the contiguous United States, *Water Resour Res*, 51, 9367-9377, 10.1002/2015wr017693, 2015.

Zhang, L., Cheng, L., and Brutsaert, W.: Estimation of land surface evaporation using a generalized nonlinear complementary relationship, *Journal of Geophysical Research: Atmospheres*, 122, 1475-1487, 10.1002/2016jd025936, 2017.

# A review of complementary principle of evaporation: From original linear relationship to generalized nonlinear functions

Songjun Han<sup>1\*</sup>, Fuqiang Tian<sup>2</sup>

<sup>1</sup>State Key Laboratory of Simulation and Regulation of Water Cycle in River Basin, China Institute of Water Resources and Hydropower Research, Beijing 100038, China, hansj@iwhr.com

<sup>2</sup>State Key Laboratory of Hydro-science and Engineering, Tsinghua University, Beijing 100084, China,

Correspondence to: Songjun Han (hansj@iwhr.com)

**Abstract.** The complementary principle is an important methodology for estimating actual evaporation by using routinely observed meteorological variables. This review summaries its~~Throughout the~~ 56-year development, focusing on how related studies have shifted from adopting a symmetric linear complementary relationship (CR) to employing generalized nonlinear functions. The original CR denotes that the actual evaporation ( $E$ ) and "apparent" potential evaporation ( $E_{pa}$ ) depart from the potential evaporation ( $E_{po}$ ) complementarily when the land surface dries from completely wet environment with constant available energy. The CR was then extended to an asymmetric linear relationship, and the linear nature was retained through properly formulating  $E_{pa}$  and/or  $E_{po}$ .~~Studies based on the linear CR have been maintained for a long time by rationally formulating the potential ( $E_{po}$ ) and "apparent" potential evaporation ( $E_p$ ) and/or employing an asymmetric parameter. Recently, the linear CR was generalized to a sigmoid function and a polynomial function respectively. These works have also advanced two types of generalized nonlinear complementary functions by invoking the boundary conditions. The sigmoid function~~ The first type does not involve the formulations of  $E_{pa}$  and  $E_{po}$ , yet uses the Penman (1948)'s potential evaporation and its radiation component as inputs, whereas the polynomial function inherits  $E_{po}$  and  $E_{pa}$  as inputs and requires proper formulations for application. The generalized complementary principle has a more rigorous physical base and offers a great potential in advancing evaporation estimation. Future studies may cover several topics including the boundary conditions under wet environments, the parameterization and application over different regions of the world, and integrating with other approaches for further development.

## 1 Introduction

The complementary principle provides a framework for estimating terrestrial land surface evaporation by adopting routinely observed meteorological variables, and offers strong potential applications (Brutsaert and Stricker, 1979; Morton, 1983; McMahon et al., 2016). In this review paper, the terms "evaporation" and "evapotranspiration" are considered equivalent. As its underlying physical basis, this principle ~~describes~~ originated from the negative feedback of areal evaporation on evaporation demand (Bouchet, 1963; Brutsaert, 2015) as illustrated by the fact that reducing areal evaporation can make the overpassing air hotter and drier (Morton, 1983). Based on the complementary principle, Bouchet

(1963) first proposed a complementary relationship (CR) among three types of evaporation (Brutsaert, 2015), namely, the actual evaporation ( $E$ ) from an extensive landscape under natural conditions ~~by relating~~, the apparent potential evaporation ( $E_{pa}$ ) of a small saturated surface inside the landscape ~~that does not affect the overpassing air and the natural evaporation process~~, and the potential evaporation ( $E_{po}$ ) that occurs from the same large-size surface of  $E$  when it is saturated. In practice,  $E_{pa}$  corresponds to current atmosphere in contact with the unsaturated evaporating surface as the overpassing air is not affected by the small saturated surface, whereas the atmosphere corresponding to  $E_{po}$  is in contact with the “potential” saturated surface. Thus, the surface water availability can be detected from the relative magnitude of  $E_{pa}$  and  $E_{po}$  because of the land surface-atmosphere interaction, and  $E$  can be estimated without the explicit knowledges of the surface. The original symmetric linear “complementary” relationship (Bouchet, 1963; Brutsaert and Stricker, 1979; Morton, 1983) evolved into an asymmetric linear relationship (Brutsaert and Parlange, 1998; Pettijohn and Salvucci, 2006; Szilagyi, 2007). However, its development and applications are hindered by the use of complex formulations of  $E_{po}$  and  $E_{pa}$  to retain the linear CR, which will be reviewed next in more detail in the following sections.

Recent studies have adopted the “generalized” complementary principle, which employs nonlinear functions instead of the linear CR (Han et al., 2012; Brutsaert, 2015; Han and Tian, 2018a). The generalized complementary function comes in two ~~types~~ways, with the first attempt abandons the theoretical concept of  $E_{pa}$  and  $E_{po}$  yet uses a sigmoid function to describe the relationship among  $E$ , Penman's potential evaporation ( $E_{Pen}$ ), and its radiation term ( $E_{rad}$ ) (Han and Tian, 2018a; Han et al., 2012). ~~adopts a polynomial function to describe the relationship between  $E$ , and inheriting the three types evaporation of linear CR yet adopts a polynomial function to describe their relationship, and and need to be formulated for the function to practical problems (Brutsaert, 2015). By contrast, while the other attempt abandons does not use the theoretical concept of and  $E_{po}$  yet uses a sigmoid function to describe the relationship among  $E$ , Penman's potential evaporation ( $E_{Pen}$ ), and its radiation term ( $E_{rad}$ ) (Han and Tian, 2018b; Han et al., 2012). ~~adopts a polynomial function to describe the relationship between  $E$ ,  $E_{pa}$  and  $E_{po}$ . However,  $E_{pa}$  and  $E_{po}$  still need to be formulated before applying the polynomial function to practical problems (Brutsaert, 2015).~~~~

The generalized complementary principle with earlier linear CRs as special cases has a more rigorous physical base (Brutsaert, 2015; Han and Tian, 2018b), and its methodology based on nonlinear functions is robust and effective. The generalized complementary principle has received much attention for its promising applications in estimating evaporation upon its proposal (Liu et al., 2016; Szilagyi et al., 2016; Ai et al., 2017; Brutsaert et al., 2017; Zhang et al., 2017; Han and Tian, 2018a; Brutsaert et al., 2019). However, the boundary conditions and proper mathematical forms of the generalized complementary functions are still under study (Han and Tian, 2018a; Crago et al., 2016; Ma and Zhang, 2017; Szilagyi et al., 2017). In this review, we summarize the 56-year development of the complementary principle with a specific focus on its

evolution from a symmetric linear CR to generalized nonlinear functions. We also compare the two types of generalized complementary functions, and discuss their future development.

2 **Symmetric-Linear** complementary relationship

2.1 Concept of **symmetric** complementary relationship

The concept of CR is illustrated in Figure 1. When the water availability of the landscape is not limited,  $E$  is assumed to proceed at  $E_{pa}$  and  $E = E_{pa} = E_{po}$ . Given that the surface dries with constant available energy,  $E$  and  $E_{pa}$  depart from  $E_{po}$  with equal yet opposite changes in fluxes and exhibit a **CR-complementary relation** as follows:

$$E_{pa} - E_{po} = E_{po} - E \quad (1)$$

The formulations of  $E_{pa}$  and  $E_{po}$  should be specified in Eq. (1)(+). Bouchet (1963) assumed  $E_{po}$  to be half the input solar radiation. Morton (1976) calculated  $E_{pa}$  by using the modified Penman's (1948) equation proposed by Kohler and Parmele (1967)(  $E_{Pen}^{KP}$  ), in which a constant vapor transfer coefficient was used to replace the wind function, and calculated  $E_{po}$  by using the Priestley–Taylor's (1972) equation (  $E_{PT}$  ) for an extensive saturated surfaced with **vanished-minimal** advection. This method has been used to calculate monthly evaporation in large areas.




Figure 1. Schematic **illustration** of symmetric CR **following Bouchet (1963)**

Table 1. **Different fF**ormulations of  $E_{pa}$  and  $E_{po}$  in **different linear CR formulations**

Types	$E_{pa}$	$E_{po}$	$b$	References
Symmetric	$E_{Pen}^{KP}$	$E_{PT}$	1	Morton (1976)
	$E_{Pen}$	$E_{PT}$		Brutsaert and Stricker (1979)



	$E_{Mor}$	$E_{PT}^{T_p}$		Morton (1983)
	$E_{Pen}^{r_s}$	$E_{Pen}^{r_s=0}$		McNaughton and Spriggs (1989)
	$E_{Pen}$	$E_{PT} +  R_n - G - E_{Pen} $		Parlange and Katul (1992a)
	$E_{PM}^{r_{s,min}}$	$E_{PT}$		Pettijohn and Salvucci (2006)
	$E_{Pen}$	$E_{PT}^{T_{ws}}$		Szilagyi and Jozsa (2008)
	$E_{Pan}$	$E_{PT}$	$b$	Kahler and Brutsaert (2006)
	$ET_0$	$E_{PT}$	$b$	Han et al. (2014c)
Asymmetric	$E_{MT}$	$E_{Pen}$	$\gamma/\Delta(T_a)$	Granger (1989)
	$E_{Pen}$	$E_{PT}$	$\gamma/\Delta(T_s)$	Szilagyi (2007)
	$E_{Pen}$	$E_{PT}^{T_{ws}}$	$f(RH)$	Szilagyi (2015)

\*The symbols can be referred to the main text [and the appendix for details](#).

Brutsaert and Stricker (1979) proposed the advection-aridity (AA) approach at a daily timescale, where  $E_{pa}$  and  $E_{po}$  are directly formulated by  $E_{Pen}$  and  $E_{PT}$ , respectively. Although various combinations of  $E_{pa}$  and  $E_{po}$  exist (Table 1),  $E_{po}$  is widely accepted to reflect the energy input while  $E_{pa}$  includes the drying power of air simultaneously (Bouchet, 1963; Morton, 1983; Lhomme and Guilioni, 2006). Therefore, the AA approach seems logical and convincing (Lhomme and Guilioni, 2006). This approach has been validated based on hourly (Parlange and Katul, 1992a; Crago and Crowley, 2005), daily (Brutsaert and Stricker, 1979; Ali and Mawdsley, 1987; Qualls and Gultekin, 1997), monthly (Xu and Singh, 2005; Lemeur and Zhang, 1990; Hobbins et al., 2001a), and annual (Ramirez et al., 2005; Yu et al., 2009) data from either plot-scale lysimeters and eddy-covariance ~~meas~~[resentsmeasurements](#) or basin-wide water balance-derived results. By calculating  $E_{Pen}$  and  $E_{PT}$  using the standard meteorological data, the AA approach has been applied to estimate evaporation in [regions with](#) various land covers and climatic ~~features~~[regions](#) (Hobbins et al., 2001a; Liu et al., 2006; Wang et al., 2011; Ozdogan and Salvucci, 2004). For instance, this approach has been applied [and validated](#) in China from the Gobi Desert with a mean annual precipitation of less than 150 mm (Liu and Kotoda, 1998; Han et al., 2008; Lemeur and Zhang, 1990) to the humid Eastern China with an annual precipitation of approximately 1,800 mm (Xu and Singh, 2005). Note that however, the AA approach tends to overestimate  $E$  under wet environments but underestimate  $E$  under arid environments. [Measurement error, imperfect formulations of  \$E\_{pa}\$  and/or  \$E\_{po}\$ , external energy sources, or even the nonlinear nature of the complementary principle were considered as potential causes of this bias](#) (Qualls and Gultekin, 1997; Hobbins et al., 2001a; Han et al., 2008, 2012).

## 2.2 Proofs of complementary relationship

Bouchet (1963) and Morton (1965; 1970) approximately validated the CR by using annual and monthly data, respectively. At an annual scale,  $E$  and  $E_{pa}$  (which are represented by  $E_{Pen}$  or pan evaporation ( $E_{Pan}$ )) were plotted against

annual precipitation and their negative relationship was used as an evidence to support the reliable probability of the complementarity (Morton, 1983). Ramirez et al. (2005) tested the CR by using a composite of 192 data pairs from 25 basins across US, and claimed a direct observational evidence ~~for the symmetry~~. Yu et al. (2009) examined the CR at 102 observatories across China and found the CR at low elevations. Su et al. (2015) also showed a negative correlation between  $E$  from atmospheric reanalysis data and  $E_{pan}$  in the non-humid regions of China. The large scale irrigation development in an arid environment provides a large “natural” experimental area for validating the CR by the opposite changes in  $E$  and  $E_{pa}$  (Roderick et al., 2009). A study from Turkey revealed that the warm-season  $E_{pa}$  ~~decreases-decreased~~ progressively along with an increasing ~~irrigation-irrigated~~ area (Ozdogan and Salvucci, 2004). Similar results were obtained from arid irrigation districts in Northwest China, where an increasing irrigation water consumption reduces  $E_{pa}$  (Han et al., 2014c) ~~while whereas~~ a decreasing irrigation water consumption increases  $E_{pa}$  (Han et al., 2017). However, although these studies showed that  $E$  and  $E_{pa}$  move in opposite directions in most cases, there was not solid evidence to support the symmetric nature of CR. ~~A rigorous quantitative assessment of the symmetric CR was not conducted by Ramirez et al. (2005). Yu et al. (2009) found that the CR is asymmetric at high elevations. However, Ma et al. (2015) argued that the asymmetric CR in TP was mainly due to inappropriate parameterizations of the wind function in  $E_{pen}$ , the wet environment air temperature and Priestley Taylor coefficient in  $E_{PT}$ .~~

The plausibility of CR has also been validated on theoretical bases and has been mathematically rationalized by Bouchet (1963), Morton (1971); Morton (1969), and Seguin (1975). The rationalization proposed by Morton (1969, 1971) considers ~~governing~~ the ~~governing~~ changes in the humidity and temperature of the equilibrium sublayer of the atmospheric boundary layer (ABL) ~~by assuming that (1) the net radiation will not change with the surface, and (2) the heat and vapor eddy transfer characteristics are identical for  $E$  and  $E_{pa}$ .~~ Relaxing the ~~second~~ assumption of Morton (1983), Szilagyi (2001) derived the CR by using the mass conservation equation for water vapor. However, ~~by using a diagnostic model of the energy fluxes within a closed system,~~ LeDrew (1979) argued that ~~Morton's two assumption do not necessarily hold, and pointed out that the symmetric CR is physically unrealistic by using a diagnostic model of the energy fluxes within a closed system, the CR proposed by Morton (1969) is physically unrealistic and added that the complementarity of the negative relationship is not supported by any proof.~~

The physical basis of the CR has been further ~~advanced-explored~~ by using climate models. McNaughton and Spriggs (1989) tested the CR by using a simple model of the atmospheric mixed layer with entrainment in which the latent heat of the surface is simulated by using the bulk mass transfer equation with bulk resistance. During the validation,  $E_{pa}$  is calculated via Penman's equation, which uses the temperature and humidity obtained from the results of the mixed-layer model corresponding to certain resistance ( $E_{pen}^{r_s}$ ), while  $E_{po}$  is calculated with the surface resistance set to zero ( $E_{pen}^{r_s=0}$ ). Kim and Entekhabi (1997) added the surface energy balance and atmospheric thermal radiation fluxes into the model to

extend the study of McNaughton and Spriggs (1989). By using the Penman–Monteith equation to govern the areal latent heat flux at the surface, Lhomme (1997a) proposed a closed-box model with an impermeable lid at a fixed height while Lhomme (1997b) used a more realistic open-box model of the ABL with entrainment to assess the CR. Sugita et al. (2001) tested the CR by using a modified version of Lhomme (1997b)'s model, which was calibrated by using a dataset obtained from the Hexi Corridor desert area in Northwest China. But a strict symmetric CR ~~wss-was~~ hardly confirmed by these studies.

### 2.3 Asymmetric linear complementary relationship

With  $E_{pa}$  and  $E_{po}$  denoted by the mass-transfer type potential evaporation  $E_{MT}$  and  $E_{Pen}$ , respectively, Granger (1989) proposed an alternative CR as follows:

$$(E_{MT} - E_{Pen}) = \frac{\Delta(T_a)}{\gamma} (E_{Pen} - E), \quad (2)$$

where  $\gamma$  is the psychrometric constant,  $\Delta(T_a)$  is the slope of the saturation vapor pressure at air temperature  $T_a$ . Despite being identical to the surface energy balance, Eq. (2) has inspired researchers to examine whether the CR should be symmetric or not (Szilagyi, 2007; Pettijohn and Salvucci, 2006). By using pan evaporation to denote  $E_{pa}$ , Brutsaert and Parlange (1998) extended the symmetric CR as follows:


$$(E_{pa} - E_{po}) = b(E_{po} - E), \quad (3)$$

where  $b$  is the coefficient that denotes asymmetry, and the original symmetric CR is characterized with  $b=1$ . Kahler and Brutsaert (2006) clarified and tested the validity of Eq. (3) at a daily timescale and attributed the asymmetry to the nature of the heat transfer between the pan and its surroundings, which made the changes in pan evaporation larger than those in  $E$ . Szilagyi (2007) showed that the asymmetry is not limited only to the evaporation pan but is inherently linked to the definition of  $E_{pa}$ . Brutsaert (2015) stated that asymmetry is an inherent characteristic of the CR.

The asymmetric CR can be illustrated in a dimensionless form (Figure 2) (Kahler and Brutsaert, 2006). Normalized by  $E_{po}$ ,  $E_{pa}$  and  $E$  can be scaled as

$$\frac{E}{E_{po}} = \frac{(1+b)E/E_{pa}}{1+bE/E_{pa}} \text{ and } \frac{E_{pa}}{E_{po}} = \frac{1+b}{1+bE/E_{pa}}. \quad (4)$$

The scaled  $E_{pa}$  and  $E$  are both functions of the dimensionless variable  $E/E_{pa}$ , while  $E/E_{pa}$  serves as the evaporative surface moisture index. Compared with the original form (Eq. (1) and Figure 1), the CR here is illustrated without the presenceappearance of the water availability explicitly. The asymmetric CR has been validated via the opposite changes of  $E/E_{po}$  and  $E_{pa}/E_{po}$  against  $E/E_{pa}$  at several locations over the world. However, the wet conditions were seldom focusedexplored, which may hide the true correlation as the two curves of  $E/E_{po}$  and  $E_{pa}/E_{po}$  approach.



**Figure 2. Scaled  $E_{pa}$  and  $E$ , which serve as functions of the evaporative moisture index  $E/E_{pa}$  and calculated on the basis of the asymmetric CR, according to method of Kahler and Brutsaert (2006).**

The asymmetric CR is a significant improvement of the symmetric CR, and the opposite changes of  $E/E_{po}$  and  $E_{pa}/E_{po}$  against  $E/E_{pa}$  were treated as an enhanced illustration of the CR (Hu et al., 2018; Zhang et al., 2017; Ma et al., 2015a; Brutsaert et al., 2019; Szilagyi, 2007). The performances on evaporation estimation are improved by calibrating the asymmetry parameter  $b$  (Kahler and Brutsaert, 2006; Han et al., 2008; Huntington et al., 2011; Ma et al., 2015a). Efforts have also made to calculate  $b$  by using the meteorological variables, which enhance the predict ability of the CR (Szilagyi, 2015; Szilagyi, 2007; Aminzadeh et al., 2016). However, the changes in  $b$  imply a potential nonlinear characteristic of the CR (Han, 2008; Lintner et al., 2015). The observed values of  $E/E_{po}$  and  $E_{pa}/E_{po}$  can even exhibit a positive correlation under wet conditions at several flux sites, which challenges the linear CR (Han and Tian, 2018a).

### 3 Efforts in maintaining a linear complementary relationship

#### 3.12.4 Efforts in remaintaining thea linear nature of complementary relationship through Rationalproperly formulation formulating of $E_{pa}$ and/or $E_{po}$

The imperfect asymmetriclinear CR has inspired researchers to apply a rational formulations of  $E_{pa}$  and/or  $E_{po}$  for to retaining the symmetry of it CR. One direct method is to improve revise the formulations of  $E_{pen}$  and/or  $E_{PT}$  based on the AA approach through calibration. For  $E_{pen}$ , the empirical wind function was calibrated to improve the CR (Hobbins, 2001). However, Penman's wind function cannot work under the wet and dry conditions simultaneously (Pettijohn and Salvucci, 2006). The wind function derived from Monin–Obukhov's similarity theory was then employed (Crago and Crowley, 2005; Parlange and Katul, 1992b; Pettijohn and Salvucci, 2006; Ma et al., 2015a). The surface roughness and surface albedo were also calibrated to improve the CR (Lemur and Zhang, 1990). Meanwhile, for  $E_{PT}$ , the Priestley–Taylor coefficient ( $\alpha$ ) is regarded varying, thereby leaving a range for calibration (Han et al., 2006; Yang et al., 2012; Xu and Singh, 2005). In

addition to  $E_{pen}$  and  $E_{PT}$ , the mass-transfer type potential evaporation (van Bavel, 1966) ( $E_{MT}$ ) was considered as another formulation of  $E_{pa}$  (Granger, 1989). Different combinations of  $E_{pa}$  and  $E_{po}$ , (i.e.,  $E_{pen}$ ,  $E_{PT}$ , and  $E_{MT}$ ) were tested through the trial-and-error method to retain the linear nature of CR (Anayah and Kaluarachchi, 2014; Crago and Crowley, 2005). ~~The local re-parameterizations and/or calibrations have significantly improved the evaporation estimation (Hobbins et al., 2001b; Xu and Singh, 2005; Ma et al., 2015a). However, the calibration approach and trial and error process are deemed ineffective because of their high computation demand, which is a key stumbling block when applying the CR in large scale (e.g., continental or global) E modelling (Ma et al., 2019).~~

Given the conceptual problems inadequacy in using  $E_{pen}$  and  $E_{PT}$  to denote  $E_{pa}$  and  $E_{po}$  (Morton, 1983; Szilagyi and Jozsa, 2008), a better CR must be obtained by modifying the formulations of  $E_{pen}$  and/or  $E_{PT}$  on physical basis. For  $E_{pa}$ , the net long-wave radiation depends on the land surface temperature; meanwhile, adjusting surface temperature with air temperature to calculate solar radiation in  $E_{pen}$  may be problematic (Morton, 1983). To address these limitations, Morton (1983) combined the energy balance and water vapor transfer equations by using an equilibrium temperature ( $T_p$ ) and derived a Morton-type potential evaporation  $E_{Mor}$  to denote  $E_{pa}$ . By attributing the asymmetry to the assumption that  $E_{pa}$  conceptually includes a transpiration component, Pettijohn and Salvucci (2006) improved the asymmetry by replacing  $E_{pen}$  with the Penman–Monteith equation with a minimum surface resistance ( $E_{PM}^{r_{smin}}$ ). Similarly, the reference evapotranspiration ( $ET_0$ ) was also used to replace  $E_{pen}$  (Han et al., 2014c; Han et al., 2017).

In theory,  $E_{po}$  is the potential evaporation when the land surface is saturated, and should be calculated with a proper formula by using meteorological variables corresponding to the “potential” saturated surface. The Priestley-Taylor equation has been widely accepted to  ~~$E_{PT}$  was proposed by Priestley and Taylor (1972) to represent evaporation from extensive saturated surfaces, and has been widely used by using meteorological variables corresponding to these saturated surfaces~~ (Brutsaert, 1982; Priestley and Taylor, 1972). This way it ~~could be used~~ was suggested to represent  $E_{po}$  (Brutsaert and Stricker, 1979). However, in the The AA approach, ~~calculates~~  $E_{PT}$  is calculated by the Priestley-Taylor equation using the atmospheric variables that correspond to the current unsaturated surface natural landscape. But the atmosphere in contact with the land surface will change if the land surface was brought into saturated (Morton, 1983; Brutsaert, 2015). Thus,  $E_{PT}$  is in reality a variable dependent on the meteorological variables at the time of calculation and does not represent ~~Therefore,  $E_{PT}$  cannot represent~~ the “true”  $E_{po}$ .

Obviously, since calculating ~~the slope of the saturation vapor pressure at the current air temperature ( $\Delta(T_a)$ ) for  $E_{PT}$~~  is imperfect because the temperature corresponding to  $E_{po}$  is different from current  $T_a$  ~~that~~ corresponding to an non-

un-saturated environment (Morton, 1983; Szilagyi and Jozsa, 2008). Thus, predicting the air temperature corresponding to the extensive saturated surface is critical for rational properly formulating  $E_{po}$ . Morton (1983) derived  $E_{po}$  by using a modified Priestley–Taylor equation with net radiation and the slope of the saturation vapor pressure that is calculated at equilibrium temperature  $T_p$  ( $E_{PT}^{T_p}$ ). Szilagyi and Jozsa (2008) argued that  $\Delta$  in  $E_{PT}$  should be calculated at the air temperature corresponding to the wet environment instead of actual air temperature, which is ~~While it is~~ not straightforward to derive  $T_{wa}$ . Thus, Szilagyi and Jozsa (2008) proposed an iterative approach based on the Bowen ratio method ~~for a small wet patch~~ to estimate the surface temperature under wet environments ( $T_{ws}$ ), and replaced  $\Delta(T_a)$  with the slope of the saturation vapor pressure curve at  $T_{ws}$  ( $\Delta(T_{ws})$ ) in the Priestley–Taylor equation ( $E_{PT}^{T_{ws}}$ ) by assuming a negligible temperature gradient over such a small wet area.  $E_{PT}^{T_{ws}}$  was used to improve the symmetry of the CR in arid shrubland environments (Huntington et al., 2011) and in an alpine steppe of the Tibetan Plateau (Ma et al., 2015a). The evaporation estimations across the US were also improved by applying the modified AA approach (Szilagyi and Jozsa, 2008; Szilagyi et al., 2009; Szilagyi, 2015). Aminzadeh et al. (2016) derived a steady state surface temperature via the surface energy balance at which the sensible heat flux is zero, and calculated  $E_{pa}$  and  $E_{po}$  using a mass-transfer type reference evaporation corresponding to current and saturated surface water content.

Advection is another factor influencing  $E_{po}$ , which could not be adequately ~~adequately~~ considered by ~~However,  $E_{PT}$  with an assumption of a minimal advection effect does not fully consider the effects of advection, which are inevitable in reality~~ (Morton, 1983, 1975; Parlange and Katul, 1992a). The effects of advection were considered by an empirical correction factor in  $E_{PT}^{T_p}$  (Morton, 1975, 1983). Parlange and Katul (1992a) attributed the asymmetry to the horizontal advection of dry air, which would make  $E_{pen}$  larger than the available energy ( $R_n - G$ ) (i.e.,  $R_n - G - E_{pen} < 0$ ) and proposed to replace  $E_{PT}$  with  $E_{PT} + |R_n - G - E_{pen}|$  to improve the CR on an hourly basis.

The efforts of reformulating  $E_{pa}$  and/or  $E_{po}$  through the calibration, trial-and-error process and the physical improvement have significantly improved the evaporation estimation (Hobbins et al., 2001b; Xu and Singh, 2005; Ma et al., 2015a; Szilagyi, 2015). However, it is always impossible to find formulations of  $E_{pa}$  and  $E_{po}$  completely rational at present, and these approaches are deemed ineffective because of their high computation demand, which is a key stumbling block when applying the CR at large-scale (e.g., continental or global) (Ma et al., 2019).

### 3.2 Asymmetric linear complementary relationship

~~With  $E_{pa}$  and  $E_{po}$  denoted by  $E_{MT}$  and  $E_{pen}$ , respectively, Granger (1989) proposed an alternative CR as follows:~~

$$(E_{MT} - E_{Pen}) = \frac{\Delta(T_a)}{\gamma} (E_{Pen} - E), \quad (2)$$

where  $\gamma$  is the psychrometric constant. Despite being identical to the surface energy balance, Eq. (2) has inspired researchers to examine whether the CR should be symmetric (Szilagyi, 2007; Pettijohn and Salvucci, 2006). By using pan evaporation to denote  $E_{pa}$ , Brutsaert and Parlange (1998) extended the symmetric CR as follows:

$$(E_{pa} - E_{po}) = b(E_{po} - E), \quad (3)$$


where  $b$  is the coefficient that denotes asymmetry. Kahler and Brutsaert (2006) clarified and tested the validity of Eq. (3) at a daily timescale and attributed the asymmetry to the nature of the heat transfer between the pan and its surroundings, which made the changes in  $E_{pan}$  larger than those in  $E$ . Szilagyi (2007) showed that the asymmetry is not limited only to  $E_{pan}$  but is inherently linked to the definition of  $E_{pa}$ . The asymmetric CR is widely used, and Brutsaert (2015) stated that asymmetry is an inherent characteristic of the CR. The asymmetric CR is widely used, and the parameter  $b$  was considered a calibrated parameter at first (Kahler and Brutsaert, 2006; Han et al., 2008; Huntington et al., 2011; Ma et al., 2015a), but was calculated by using the meteorological variables (Szilagyi, 2015; Szilagyi, 2007; Aminzadeh et al., 2016).

The asymmetric CR can be illustrated in a dimensionless form (Figure 2) (Kahler and Brutsaert, 2006). Normalized by  $E_{po}$ ,  $E_{pa}$  and  $E$  can be scaled as

$$\frac{E}{E_{po}} = \frac{(1+b)E/E_{pa}}{1+bE/E_{pa}} \text{ and } \frac{E_{pa}}{E_{po}} = \frac{1+b}{1+bE/E_{pa}}. \quad (4)$$

The scaled  $E_{pa}$  and  $E$  are both functions of the dimensionless variable  $E/E_{pa}$ , while  $E/E_{pa}$  serves as the evaporative surface moisture index. Compared with the original form (Eq. (1) and Figure 1), the CR here is illustrated without the appearance of the water availability explicitly. The opposite changes of  $E/E_{pa}$  and  $E_{pa}/E_{po}$  were treated as a standard illustration of the CR (Hu et al., 2018; Zhang et al., 2017; Ma et al., 2015a; Brutsaert et al., 2019; Szilagyi, 2007), but the wet conditions where the two curves of  $E/E_{pa}$  and  $E_{pa}/E_{po}$  approach were seldom focused. However, the observed values of  $E/E_{pa}$  and  $E_{pa}/E_{po}$  exhibit a positive correlation at several flux sites, which challenges the CR under wet conditions (Han and Tian, 2018a).





**Figure 2. Scaled  $E/E_{pa}$  and  $E/E_{po}$ , which serve as functions of the evaporative moisture index  $E/E_{pa}$  and calculated on the basis of the asymmetric CR.**

### 4.3 Generalized complementary principle via nonlinear functions

#### 4.3.1 Normalized complementary functions

Unlike the normalization by  $E_{po}$  (Kahler and Brutsaert, 2006), Han (2008) normalized Eq. (3) by using  $E_{pa}$  and found that  $E/E_{pa}$  is expressed as a linear function of  $E_{po}/E_{pa}$ . Normalized by  $E_{pen}$  (Han et al., 2008), the AA approach can be expressed as

$$\frac{E}{E_{pen}} = \alpha \left(1 + \frac{1}{b}\right) \frac{E_{rad}}{E_{pen}} - \frac{1}{b}, \quad (5)$$

where  $E/E_{pen}$  is regarded as a linear function of  $E_{rad}/E_{pen}$ . The bias of the AA function under arid and wet environments can be easily understood in its dimensionless form. Also, but the AA approach with a tuned  $b$  still underestimated evaporation in arid environments (Han et al., 2008). The work of Crago and Brutsaert (1992) in Kansas during FIFE 1987 revealed that the parameter  $b$  are obviously different for days with differing degrees of soil moisture. These studies, which imply implies that the CR may deviate from its linear characteristics.

Based on the examination of the CR using a model of the convective boundary-layer with entrainment (Lhomme, 1997b), Lhomme and Guilioni (2010, 2006) recommended a form of the CR through the effective surface resistance of the region. Integrating this relationship into Penman–Monteith equation and the normalization by  $E_{pen}$  lead to

$$\frac{E}{E_{pen}} = (1 + \omega) \frac{E_{rad}}{E_{pen}}, \quad (6)$$

where  $\omega$  is a coefficient accounting for the entrainment of dry air within the atmospheric boundary layer. Equation (6) is a linear function without intercept, but was not verified and applied using observed data.

The CR model proposed by Granger (1989) based on Eq. (2) has demonstrated promising application across different land covers and regional climate conditions (Carey et al., 2005; Granger, 1999; Granger and Gray, 1989b; Pomeroy et al., 1997; Xu and Singh, 2005). In fact, the relationship between relative evaporation and relative drying power plays a key role in reflecting the dryness of the surface (Granger and Gray, 1989a). This relationship was integrated to a asymmetric CR to improve the performance on evaporation estimation (Anayah and Kaluarachchi, 2014). Normalized by  $E_{pen}$ , Granger's model is similar to the AA function in that  $E/E_{pen}$  is expressed as a function of the relative magnitude of drying power to net radiation (Han et al., 2011). By synthesizing the dimensionless forms of the AA function and the Granger's model, Han et al. (2011) proposed the following ~~logistic~~ function as an alternative:

$$\frac{E}{E_{pen}} = \frac{1}{1 + c_1 e^{d(1 - \frac{E_{rad}}{E_{pen}})}}, \quad (7)$$

where  $c_1$  and  $d$  are the parameters. Eq. (7) approximates the linear AA function under normal conditions neither too wet nor too dry but amends its bias (Han et al., 2011). thus can be regarded an enhanced nonlinear version of the linear CR.

Actual evaporation can be estimated using routinely measured meteorological data by using the climatological resistance to parameterize the bulk surface resistance in the Penman–Monteith equation (Liu et al., 2012; Rana et al., 1997; Katerji and Perrier, 1983; Ma et al., 2015b). A linear relationship between the ratio of surface resistance to aerodynamic resistance and the ratio of climatological resistance to aerodynamic resistance was proposed by Katerji and Perrier (1983). Han et al. (2014d) integrated this linear relationship into the Penman–Monteith equation and derived a dimensionless form via normalization by  $E_{pen}$ :

$$\frac{E}{E_{pen}} = \frac{1}{1 + k(\frac{E_{pen}}{E_{rad}} - 1) + l}, \quad (8)$$

where  $k$  and  $l$  are the empirical calibration parameters. With similar variables yet different mathematical formulations, Eq. (8) can also be considered a complementary function (Han et al., 2014d).

#### 4.2 Sigmoid function relating $E/E_{pen}$ to $E_{rad}/E_{pen}$

By synthesizing the aforementioned ~~three~~ functions (Table 2), Han et al. (2012) generalized the CR as a function that relates  $E/E_{pen}$  to  $E_{rad}/E_{pen}$ :

$$E/E_{pen} = f(E_{rad}/E_{pen}). \quad (9)$$

Eq. (9) shares the same form of Penman's approach for estimating evaporation. The function of surface wetness that denotes the reduction of  $E$  to  $E_{pen}$  is replaced by the function of  $E_{rad}/E_{pen}$ , which is termed “atmospheric wetness” (Han and Tian, 2018b). Despite not explicitly exhibiting a CR, Eq. (9) holds the complementary principle that the land surface wetness is indirectly denoted by the drying power of air with a constant radiation energy input (Brutsaert, 1982).

395

Accordingly, Eq. (9)(9) is considered a "general form" of the CR (Han et al., 2014b) (hereinafter referred to as H12 whereas the other type of generalized complementary function first proposed by Brutsaert (2015) if referred to as B15 for comparison). The existing analytical forms of the function can be classified into linear, concave/convex, or sigmoid (Table 4 in Han and Tian (2018a)2). Studies on the complementary principle can be advanced by formulating a proper analytical form for H12.

**Table 2. Different analytical formulas for normalizedgeneralized complementary functions H12**

Type	Formula *	References
Linear	$y = \alpha(1 + \frac{1}{b})x - \frac{1}{b}$	Brutsaert and Stricker (1979)
	$y = (1 + \omega)x$	<u>Lhomme and Guillioni (2010, 2006)</u>
Sigmoid	$y = \frac{1}{1 + c_1 e^{d(1-x)}}$	Granger (1989), Han et al. (2011)
	$y = \frac{1}{1 + m(\frac{1}{x} - 1)^n}$	Han et al. (2012)
	$y = \frac{1}{1 + m(\frac{x_{\max} - x}{x - x_{\min}})^n}$	Han and Tian (2018)
Concave/ convex	$y = \frac{1}{1 + k(\frac{1}{x} - 1) + l}$	Katerji and Perrier (1983) , Han et al. (2014b)
	$y = (2 - c)\alpha^2 x^2 - (1 - 2c)\alpha^3 x^3 - c\alpha^4 x^4$	Brutsaert (2015)

400

\*  $x = E_{rad} / E_{Pen}$  and  $y = E / E_{Pen}$  . the other symbols are parameters.

The exact analytical form of H12 is inadequately understood at present. However, some of its characteristics can be detected from its boundary conditions under extremely arid and completely wet environments. Han et al. (2012) derived the zero-order and first-order boundary conditions for H12 as

420

$$\begin{cases} y_H = 0, x_H \rightarrow 0 \\ y_H = 1, x_H \rightarrow 1 \\ \frac{dy_H}{dx_H} = 0, x_H \rightarrow 0, \\ \frac{dy_H}{dx_H} = 0, x_H \rightarrow 1 \end{cases} \quad (10)$$

where  $x_H = E_{rad}/E_{Pen}$  and  $y_H = E/E_{Pen}$ . Han et al. (2012) proposed the following sigmoid function (hereinafter [this specific analytical form of H12 is](#) referred to as [sigmoid H2012](#)):

425

$$\frac{E}{E_{Pen}} = \frac{1}{1 + m(\frac{E_{Pen}}{E_{rad}} - 1)^n}, \quad (11)$$

where  $m$  and  $n$  are parameters. ~~The linear AA and nonlinear H2012 have been compared in a 2D space ( $E_{rad}/E_{Pen}$  vs  $E/E_{Pen}$ ) (Han et al., 2012).~~ The results obtained from an extremely dry desert and a wet farmland reveal that the sigmoid H2012 corrects the bias of the linear AA and Equation ~~(7)(7)~~ (Han et al., 2012); the application of this sigmoid function has also been recommended for an alpine meadow region of the Tibetan Plateau (Ma et al., 2015b).

**Table 23. Different forms of the generalized complementary function,  $y = f(x)$**

<a href="#">Approach</a>	Specific function	$E_{po}$	$x$	$y$	Typical type	Reference
H12*	H2017	Not involved	$\frac{E_{rad}}{E_{Pen}}$	$\frac{E}{E_{Pen}}$	Sigmoid	Han et al. (2018)
	B2015	$E_{PT}$	$\frac{\alpha E_{rad}}{E_{Pen}}$	$\frac{E}{E_{Pen}}$	4-order polynomial	Brutsaert (2015)
B15**	C2016	$E_{PT}^{T_{ws}}$	$\frac{E_{PT}^{T_{ws}}/E_{Pen} - E_{PT}^{T_{ws}}/E_{MT}^{max}}{1 - E_{PT}^{T_{ws}}/E_{MT}^{max}}$	$\frac{E}{E_{Pen}}$	Linear	Crago et al. (2016)
	S2017	$E_{PT}^{T_{ws}}$	$\frac{E_{Pen}^{max} - E_{Pen}}{E_{Pen}^{max} - E_{PT}^{T_{ws}}} \frac{E_{PT}^{T_{ws}}}{E_{Pen}}$	$\frac{E}{E_{Pen}}$	3-order Polynomial	Szilagyi et al. (2017)

430

\*H12 denotes the t generalized complementary function  $E/E_{Pen} = f(E_{rad}/E_{Pen})$ , while H2012 and H2017 are the sigmoid analytical formulas for H12.

435

\*\*B15 denotes the generalized complementary function  $E/E_{pa} = f(E_{po}/E_{pa})$ , while B2015, C2016 and S2017 are the analytical formulas for B15 in application.

The zero-order arid boundary condition of H12 adopted in H2012 may be [problematic-imperfect](#) in the sense that the aerodynamic term ( $E_{aero}$ ) of  $E_{Pen}$  may not reach infinity under an arbitrary  $E_{rad}$  (Crago et al., 2016; Szilagyi et al., 2017; Kovács, 1987). Moreover,  $E_{rad}/E_{Pen}$  cannot easily approach unity because of advection (Kovács, 1987; Priestley and Taylor,

460 1972). Therefore, Han and Tian (2018a) brought in the minimum and maximum limits of  $E_{rad}/E_{Pen}$  ( $x_{min}$  and  $x_{max}$ ) under an assumed constant  $E_{rad}$  and re-derived the boundary conditions of H12 by adopting two widely accepted assumptions following Penman's combination theory, namely,  $\partial E/\partial E_{Pen} = 0$  under extremely arid environments and  $E = E_{Pen}$  under completely wet environments. The boundary conditions are set as follows:

$$\begin{cases} y_H = 0, x_H \rightarrow x_{min} \\ y_H = 1, x_H \rightarrow x_{max} \\ \frac{dy_H}{dx_H} = 0, x_H \rightarrow x_{min} \\ \frac{dy_H}{dx_H} = 0, x_H \rightarrow x_{max} \end{cases} \quad (12)$$

465 Based on the boundary conditions, Han and Tian (2018a) ~~found~~ speculated that the growth of  $E/E_{Pen}$  upon  $E_{rad}/E_{Pen}$  exhibits a sigmoid feature, which is a three-stage pattern in which  $E/E_{Pen}$  gradually increases along with  $E_{rad}/E_{Pen}$ , rapidly increases along with  $E_{rad}/E_{Pen}$  in the following stage, and then demonstrates a decelerated growth in the final stage. The sigmoid feature can be detected from the study by Han et al. (2012) in the arid Gobi-HEIFE site and the humid Wudaogou site in China. Han and Tian (2018a) further validated the sigmoid feature according to the much larger regression slopes of  $E/E_{Pen}$  upon  $E_{rad}/E_{Pen}$  in the middle stage than those in the other two stages with smaller or larger values of  $E_{rad}/E_{Pen}$  by

470 using 22 eddy covariance towers from the FLUXNET (Baldocchi et al., 2001) dataset which includes representative biomes of grasslands, croplands, shrublands, evergreen needleleaf forests, deciduous broadleaf forests, and wetlands.

In 2017, Han and Tian (2018a) proposed the following new sigmoid function to accordance with the boundary conditions (hereinafter referred to as sigmoid H2017):


$$475 \quad \frac{E}{E_{Pen}} = \frac{1}{1 + m \left( \frac{x_{max} - E_{rad}/E_{Pen}}{E_{rad}/E_{Pen} - x_{min}} \right)^n}, \quad (13)$$

where  $E_{rad}/E_{Pen}$  adopts the feasible domain ( $x_{min}$ ,  $x_{max}$ ), which is a subdomain of (0, 1). Both the linear AA function and sigmoid H2012 are special cases of sigmoid H2017. Han and Tian (2018a) performed a first-order Taylor expansion of Eq. (13)(13) at the point where  $E/E_{Pen} = 0.5$  and set the linear equation equivalent to the linear AA function. Afterward, the parameters  $m$  and  $n$  of sigmoid H2017 can be transformed from the Priestley-Taylor coefficient  $\alpha$  and parameter  $b$  of the

480 AA function.

Han et al. (2008) was the first to plot the AA function as a linear in the state space ( $E_{rad}/E_{Pen}$  -  $E/E_{Pen}$ ), in which the biases of the AA function under arid and wet environments can be understood easily. The analytical forms of the generalized complementary function H12 listed in Table 2 can be plotted as curves in a 2D space ( $E_{rad}/E_{Pen}$  -  $E/E_{Pen}$ ) (Fig. 3), and the

upper limits of  $E_{pen}$  and  $E_{PT}$  are illustrated as the curve of  $OMN$ . The sigmoid H2012 was compared to the linear AA in the state space ( $E_{rad}/E_{pen}, E/E_{pen}$ ) to demonstrate its improvement (Han et al., 2012). Observed  $E/E_{pen}$  can be plotted against  $E_{rad}/E_{pen}$  and fitted by the analytical functions of H12 in the state space ( $E_{rad}/E_{pen}, E/E_{pen}$ ), which is an obvious improvement compared to the schematic illustrations of the CR in Fig. 1 and 2.



**Figure 3. Generalized complementary functions in the state space ( $E_{rad}/E_{pen}, E/E_{pen}$ ): linear AA, polynomial B2015, and sigmoid H2017, with  $\alpha = 1.26$  and  $b=1$ .  $x_{min}$  and  $x_{max}$  are set to 0 and 0.9, respectively, as revised from Han and Tian (2018).  $OM$  is the edge at which  $E = E_{PT}$ ,  ~~$M$  corresponds to the condition of the minimal advection evaporation where  $E_{PT} = E_{pen}$~~ ,  $MN$  is the edge where  $E = E_{pen}$ ,  $M$  corresponds to the condition of the minimal advection evaporation where  $E_{PT} = E_{pen}$ , and  $N$  corresponds to the condition of the equilibrium evaporation where  $E_{pen} = E_{rad}$ .**

### 43.3 Polynomial function relating $E/E_{pa}$ to $E_{po}/E_{pa}$

Inspired by Han *et al.* (2012), Brutsaert (2015) reformulated another general dimensionless form of the CR,  $E/E_{pa} = f(E_{po}/E_{pa})$ , and proposed its boundary conditions as follows:

$$\begin{cases} y_B = 0, x_B \rightarrow 0 \\ y_B = 1, x_B \rightarrow 1 \\ \frac{dy_B}{dx_B} = 0, x_B \rightarrow 0, \\ \frac{dy_B}{dx_B} = 1, x_B \rightarrow 1 \end{cases} \quad (14)$$

where  $x_B = E_{po}/E_{pa}$  and  $y_B = E/E_{pa}$ . The following fourth-order polynomial function was also derived to satisfy the boundary conditions:

$$\frac{E}{E_{pa}} = (2-c) \left( \frac{E_{po}}{E_{pa}} \right)^2 - (1-2c) \left( \frac{E_{po}}{E_{pa}} \right)^3 - c \left( \frac{E_{po}}{E_{pa}} \right)^4, \quad (15)$$

525 | where  $c$  is a parameter. Brutsaert (2015) regarded Eq. (15)(45) (hereinafter referred to as B15) as a generalization of the linear CR and referred to the corresponding methodology as the “generalized complementary principle.”

530 | ~~B15 is essentially different from H12, with completely different normalized variables. The boundary conditions of H12 are derived for  $x_H = E_{rad}/E_{pen}$  and  $y_H = E/E_{pen}$ , whilst those of B15 are derived for  $x_B = E_{po}/E_{pa}$  and  $y_B = E/E_{pa}$  (Table 2). B15 inherits all three types of evaporation dated from the original CR, while the validity of its boundary conditions depends on the proper definitions of  $E_{pa}$  and  $E_{po}$ . Therefore, B15 still faces the problem of the original CR, that is, formulating  $E_{po}$  and  $E_{pa}$ . By contrast, only the mostly accepted  $E_{pen}$  appears in H12, and the knowledge on  $E_{po}$  is unnecessary in deriving the boundary conditions and the analytical form of H12. By doing so, the corresponding theoretical and practical difficulties can be prevented.~~

535 | The application of Eq. (15)(45) depends on specific formulations of  $E_{pa}$  and  $E_{po}$ . In the manner of the AA approach, Eq. (15)(45) has been applied to estimate evaporation (Brutsaert et al., 2017; Liu et al., 2016; Szilagyi et al., 2016; Zhang et al., 2017; Ai et al., 2017). In this case, we refer to Eq. (15)(45) in the manner of the AA approach as B2015 to avoid confusion. Although estimating  $E_{pa}$  by using  $E_{pen}$  is widely accepted by the research community, prognostically predicting  $E_{po}$  based on  $E_{PT}$  remains a huge challenge considering the theoretical problems of the Priestley-Taylor coefficient. In addition, the lower limit of  $x_B \rightarrow 0$  of B15 may not hold in the manner of the AA approach (Kovács, 1987; Szilagyi et al., 540 | 2017; Crago et al., 2016; Han and Tian, 2018a). To address these challenges, Szilagyi et al. (2017); Crago et al. (2016) used the maximum value of  $E_{pa}$  to rescale  $x_B$  and replaced  $E_{PT}$  with  $E_{PT}^{T_{ws}}$ , the latter of which is based on the air temperature in a wet environment. Crago et al. (2016) applied a mass transfer approach to calculate the maximum value of  $E_{pa}$  ( $E_{MT}^{\max}$ ) and rescaled  $x_B$  as



$$x_C = \frac{E_{PT}^{T_{ws}}/E_{Pen} - E_{PT}^{T_{ws}}/E_{MT}^{\max}}{1 - E_{PT}^{T_{ws}}/E_{MT}^{\max}}. \quad (16)$$

Szilagyi et al. (2017) employed the Penman equation to calculate the maximum value of  $E_{pa}$  ( $E_{Pen}^{\max}$ ) and proposed the following rescaled version:

$$x_S = \frac{E_{Pen}^{\max} - E_{Pen} E_{PT}^{T_{ws}}}{E_{Pen}^{\max} - E_{PT}^{T_{ws}} E_{Pen}}. \quad (17)$$

$x_C$  and  $x_S$  are essentially same (Szilagyi et al., 2017) except for the different formulations for the maximum value of  $E_{pa}$ .

However,  $E_{MT}^{\max}$  in Eq. (16) may become invalid under conditions with relatively strong available energy yet weak winds (Ma and Zhang, 2017), and was replaced with  $E_{Pen}^{\max}$  (Crago and Qualls, 2018). In the latest version of C16,  $E_{MT}^{\max}$  is replaced with  $E_{Pen}^{\max}$  (Crago and Qualls, 2018). After rescaling, Crago et al. (2016) proposed a new linear version of the generalized complementary function (hereinafter referred to as C2016) (i.e.,  $y_B = x_C$ ; Table 2), while Szilagyi et al. (2017) used the third order polynomial function (hereinafter referred to as S2017) by replacing B15 with  $c=0$ . With the same independent variable yet different functions (Table 2), C2016 and S2017 demonstrate improvements in their evaporation estimation performance (Szilagyi et al., 2017; Crago et al., 2016; Crago and Qualls, 2018).

#### 4.3.4 Comparisons between the two generalized complementary approaches between different analytical forms

The two generalized complementary approaches, H12 and B15, are essentially different, with completely different normalized variables (Table 3). The differences in the analytical forms, sigmoid and 4-order polynomial, mainly result from their wet boundary conditions. B15 inherits the concept of the three types of evaporation dated from the original CR, and its boundary conditions and analytical form are derived for  $x_B = E_{po}/E_{pa}$  and  $y_B = E/E_{pa}$ . The original CR adopts the limits of  $E_{pa}$  and  $E_{po}$  on  $E$  in a serial manneres ( $E \leq E_{po} \leq E_{pa}$ ) (Brutsaert, 2015) while considering that the wet regional evaporation must always be smaller than the wet patch evaporation ( $E_{po} \leq E_{pa}$ ). Under wet conditions, B15 adopts  $dy_B/dx_B = 1$  as  $x_B \rightarrow 1$  by considering that any change in  $E$  is the same as the change in  $E_{po}$ , which results in a concave polynomial type function. The limits and boundary conditions of B15 would be appropriate in theory. However,  $E_{po}$  and  $E_{pa}$  should be formulated before B15 is applied to practical problems. Thus, B15 still faces one of the difficulties of that the original CR has, that is, appropriately formulating  $E_{po}$  and  $E_{pa}$ , which determines the validity and application of B15. So, future studies can be conducted towards more proper formulations of  $E_{pa}$  and  $E_{po}$  to satisfy the boundary conditions of B15.

By contrast, H12 goes much further from the original CR. The boundary conditions and the analytical form of H12 are derived for  $x_H = E_{rad}/E_{Pen}$  and  $y_H = E/E_{Pen}$ . The knowledge on  $E_{po}$  is unnecessary, and only the mostly accepted  $E_{Pen}$

and its radiation term appear in H12. By doing so, the corresponding theoretical and practical difficulties of formulating  $E_{po}$  and  $E_{pa}$  are prevented/eliminated. HB12 adopts  $E_{pen}$  as the upper limit  $E \leq E_{pen}$  during the derivation and introduce the limit of  $E \leq E_{PT}$  by considering that  $E_{PT}$  is widely used as an upper limit of  $E$  in practice. Han and Tian (2018a) showed that the upper limits of  $E_{pen}$  and  $E_{PT}$  on evaporation must be in parallel, that is,  $\begin{cases} E \leq E_{pen} \\ E \leq E_{PT} \end{cases}$ , and the complementary curves should be constrained by the limits of OMN as illustrated in (Figure 3). The limits of  $E_{pen}$  and  $E_{PT}$  on  $E$  can be approximately satisfied by the sigmoid function H2017 with the parameters transformed from the linear AA function (Han and Tian, 2018a). Besides, HB12 adopts  $dy_H/dx_H = 0$  as  $y_H \rightarrow 1$  by considering that  $E$  approaches  $E_{pen}$  under wet conditions, which results in a sigmoid type function.

In the manner of the AA approach of formulating  $E_{po}$  and  $E_{pa}$ , B15 evolves to one of its analytical forms, the polynomial B2015. Taking the Priestley-Taylor coefficient as a parameter, B2015 also can be regarded a polynomial analytical forms of H12 (Table 2), and can be compared with the sigmoid H2017 in the state space  $(E_{rad}/E_{pen}, E/E_{pen})$  (Figure 3). In the polynomial B2015, the limits on the  $E$  are specified to  $E \leq \alpha E_{rad} \leq E_{pen}$ . In practice, a constant  $\alpha$  is widely used, and the polynomial curves of B2015 are required to be constrained by the triangle domain OMP (Figure 3), which discards the domain out of OMP. However, the Priestley-Taylor coefficient varies with several factors, such as the relative transport efficiency of turbulent, or the surface/air temperature (Assouline et al., 2016; Szilagyi, 2014). Thus,  $E_{rad}/E_{pen}$  may be larger than  $1/\alpha$ , revealing that the trapezoidal domain adopted by the sigmoid H2017 is more accurate. In the state space  $(E_{rad}/E_{pen}, E/E_{pen})$ , the curve of the sigmoid H2017 exhibits a three-stage pattern, whereas the linear AA and polynomial B2015 have one and two stages respectively. As it is difficult for one site to cover all the three stages with a wide range of wetness, the linear AA can effectively represent the complementary curve under normal conditions falling in the middle stage. The polynomial B2015 is effective if the first two stages exist. Given that the third stage is uncommon, the polynomial B2015 performs well with calibrated parameters (Brutsaert et al., 2017; Liu et al., 2016; Zhang et al., 2017; Han and Tian, 2018a). However, observed points are located in the domain out of OMP at several flux sites, and the sigmoid H2017 shows the best performance in estimating evaporation as validated by using data from FLUXNET (Han and Tian, 2018a; Wang et al., 2019).

#### 4.5 Improved understanding on the correlation between actual and potential evaporation

~~Interpreting the changes in  $E$  based on the trends in  $E_{pen}$  (or pan evaporation) greatly relies on the understanding of whether the correlation between  $E$  and  $E_{pen}$  is positive or negative. The corresponding confusion has resulted in a discrepancy between the Penman hypothesis and the complementary principle (Yang et al., 2006) and encouraged debates on~~

whether the increasing or decreasing trend in  $E$  corresponds to reductions in the observed pan evaporation in the past (Brutsaert and Parlange, 1998; Roderick and Farquhar, 2002; Roderick et al., 2009; Wang et al., 2017). According to the symmetric CR,  $E_{Pen}$  would be negatively correlated with  $E$  when the energy input is constant (Morton, 1983). Based on the asymmetric linear CR, Brutsaert and Parlange (1998) stated that the decreasing  $E_{pan}$  can be used to indicate an increasing  $E$  in water-limited regions. However, the interpretation is not general (Roderick et al., 2009) because of the inherent weakness of the linear CR (Han et al., 2014b).

A proper generalized complementary function offers advantages in assessing the correlation between  $E$  and  $E_{Pen}$  while considering the different impacts of  $E_{rad}$  and  $E_{aero}$  (Hobbins et al., 2004). Han et al. (2014b) proposed a systematic and analytical approach for evaluating the correlation between  $E$  and  $E_{Pen}$  by establishing a linear regression between them via H2012. Han et al. (2014b) also demonstrated that the relative variation of  $E_{rad}$  and  $E_{aero}$  (which significantly vary at different timescales) as well as water availability (which varies across different climate regions) are two factors that affect the correlation between  $E$  and  $E_{Pen}$ . With obvious variations in  $E_{rad}$  and a relative stable  $E_{aero}$ , which are commonly observed on a diurnal or intra-annual basis, the influence of  $E_{rad}$  becomes more significant and  $E$  is always positively correlated with  $E_{Pen}$ . Under conditions where the variations in  $E_{rad}$  and  $E_{aero}$  are comparable or when  $E_{aero}$  obviously varies (which tends to occur on a daily or annual basis), the influence of  $E_{aero}$  comes to force. As a result, the correlation between  $E$  and  $E_{Pen}$  changes from negative to positive along with increasing water availability. The theoretical results were validated in a grassland site in Northeast China, and can rationally interpret the trends in  $E$  over China (Han et al., 2014b).

## 5.4 Current applications and future developments for the generalized complementary principle

### 4.1 Current applications of the generalized complementary functions for evaporation estimation

Morton (1983) thought that the ability of the complementary principle to estimate actual evaporation by using meteorological variables only can significantly influence the science and practice of hydrology. However, the attempts in using the complementary principle for evaporation estimation in hydrological modelling (Oudin et al., 2005; Barr et al., 1997; Nandagiri, 2007) have been suspended, while those attempts in applying such principle in drought assessment (Kim and Rhee, 2016; Hobbins et al., 2016) are still in their infancy. Moreover, the potential applications in agriculture water management are limited in the sense that the irrigation-induced changes in potential evaporation was mainly evaluated at an annual timescale (Ozdogan et al., 2006; Han et al., 2014a; Han et al., 2017). Apparently, the complementary principle did not develop to its full capacity via the linear CR, which leaves a broad space for further applications of applying the generalized complementary functions for evaporation research.

For example, the generalized complementary functions have been validated or applied in evaporation estimation for many sites (Ai et al., 2017; Brutsaert et al., 2017; Zhang et al., 2017; Han and Tian, 2018a; Crago and Qualls, 2018), and several basins in China (Liu et al., 2016; Gao et al., 2018). B2015 was applied to estimate global terrestrial evaporation with calibrated  $\alpha$  as a function of aridity index (Brutsaert et al., 2019). The modified Granger's model was also applied for estimating global evaporation with 0.5° spatial resolution and monthly time steps (Anayah and Kaluarachchi, 2019). It should be noted that most, if not all, above mentioned CR applications need “prior” knowledge in  $E$  (either ground-measured or water-balance-derived) to calibrate the parameters. Recently, Szilagyi et al. (2017)'s model was applied for monthly evaporation estimation without calibration across the conterminous China (Ma et al., 2019) and United States (Ma and Szilagyi, 2019). A wide range of model evaluations against the plot-scale flux measurements and basin-scale water balance results suggested that the generalized complementary functions could serve as a benchmark tool for validating the large-scale  $E$  results simulated by those Land Surface models and Remote Sensing models (Ma and Szilagyi, 2019). However, further applications over the world are still needed to develop more long-term, high-resolution  $E$  datasets for use in hydrological and atmospheric communities.

Boundary conditions are crucial to the derivations of the generalized complementary functions. Under completely wet environment, H12 adopts  $dy_H/dx_H=0$  for the first order wet boundary condition, whereas B15 adopts  $dy_B/dx_B=1$ , which results in the different types of H12 and B15. It should be noted that the boundary conditions of B15 were derived about theoretical wet patch evaporation  $E_{pa}$  and wet regional evaporation  $E_{po}$ , and their validity depends on proper definitions of  $E_{pa}$  and  $E_{po}$ . So, future studies can be conducted towards more proper formulations of  $E_{pa}$  and  $E_{po}$  to satisfy the boundary conditions of B15.

However, the boundary conditions of H12 were derived about  $E_{Pen}$  and its radiation term. The boundary conditions of B15 are only comparable to those of H12 if it is in the specific form of B2015. Han and Tian (2018a) found that the first order wet boundary conditions of H12 and B2015 are two possible solutions of the assumption that actual evaporation proceeds at  $E_{Pen}$ , and they discarded each other. However, there was not perfect instance to demonstrate H12's first order wet boundary condition. Thus, the controversies on the first order wet boundary condition (Szilagyi and Crago, 2019; Han and Tian, 2019) require further studies from both the theoretical and practical aspects. Thus, the flux data of sites over lakes or wetlands need to be well examined.

#### 5.4.2. Parameterizations of generalized complementary functions for future applications

Determining the parameters of the generalized complementary functions is the urgent work for the application of B2015 and H2017 for evaporation estimation, as well as the development of the generalized complementary principle. Given the variations in  $\alpha$ , the linear AA, polynomial B2015 and sigmoid H2012 all have two parameters. The linear AA with a default value of  $b=1$  has achieved a great success was applied at first in evaporation estimation. For the B2015,  $c$  was thought

to be only applied to accommodate unusual situations (Brutsaert, 2015). In practice,  $c = 0$  is adopted and the Priestley-Taylor coefficient is calibrated (Zhang et al., 2017; Brutsaert et al., 2017; Liu et al., 2016; Brutsaert, 2015). But the calibrated  $\alpha$  is smaller than the widely accepted constant 1.26 or even smaller than the unit at several sites, which is physically unrealistic. Han and Tian (2018a) found that  $c$  corresponds to  $b$  in the AA by setting the B2015 approximately equal to the AA in the middle stage. However, the default value of  $c=0$  corresponds to  $b$  with a value around 4.5, not the early default value of  $b=1$ , implying the default value of  $c=0$  may be not suitable. The consistency suggests that  $c$  needs to be calibrated. By calibrating both  $\alpha$  and  $c$ , the B2015 performed well in estimating evaporation for 20 FLUXNET sites, and the value of  $\alpha$  were more rational (Han and Tian, 2018a).

By contrast, two more parameters ( $x_{\min}$  and  $x_{\max}$ ) are added to the sigmoid H2017. Because the sigmoid complementary curve are insensitive to  $x_{\min}$  and  $x_{\max}$ , Han and Tian (2018a) suggested that they could be treated as constant parameters for application convenience.  $x_{\min}$  and  $x_{\max}$  may change along with  $E_{rad}$ , and were thought to vary with the time scales (Han and Tian, 2018a),  $x_{\min} = 0$  and  $x_{\max} = 1$  are appropriate at a daily scale for convenience, as have been evidenced by the well performances when compared to the flux measurements (Han and Tian, 2018a; Han et al., 2012).  $x_{\min}$  and  $x_{\max}$  are expected to be calculated by applying certain approaches to reduce the number of parameters of H2017 to two (Han and Tian, 2019).

Although  $\alpha$  would vary in theory (Assouline et al., 2016), it is widely used with a constant value of 1.26 in practice (Priestley and Taylor, 1972). After calibrating, the variations of  $\alpha$  is much less significant than those of the other parameters. Moreover, the calibrated  $\alpha$  approaches 1.26, especially for the sigmoid H2017. Thus, the constant  $\alpha = 1.26$  was suggested with acceptable weakening of the accuracy of  $E$  estimation (Han and Tian, 2018a; Han et al., 2012). In practice,  $\alpha$  was also determined from the observed  $E$  values in wet condition when  $E$  is close to  $E_{Pen}$  and/or  $E_{PT}$  (Kahler and Brutsaert, 2006; Ma et al., 2015a; Wang et al., 2019). A novel method by using observed air temperature and humidity data under wet environment was proposed by Szilagyi et al. (2017) when measured  $E$  is lacking, and was successfully used for large-scale CR model applications (Ma and Szilagyi, 2019; Ma et al., 2019).

After determining  $\alpha$  in advance, only a single parameter in the generalized complementary functions needs to be calibrated. As the parameters of the B2015 and H2017 can be transferred from the asymmetric parameter  $b$  of the original CR (Han and Tian, 2018a), the former studies on the characteristics of  $b$  could help its parameterization. The  $b$  in the desert was much smaller than those in the oases or irrigated farmlands (Han et al., 2008, 2012).  $b$  was thought to be related to the characteristics of the atmosphere, i.e., the atmospheric humidity (Szilagyi, 2015), the Clausius–Clapeyron relationship between saturation-specific humidity and temperature (Lintner et al., 2015), or the characteristics of the land surface, i.e., the surface temperature (Szilagyi, 2007), the water availability of the land surface (Han and Tian, 2018b; 2010), or the ecosystem types (Wang et al., 2019). Szilagyi (2015) applied a sigmoid function of relative humidity to parameterize  $b^{-1}$ .

Wang et al. (2019) used the ecosystem mean  $b$  values of 217 sites around the world in the B2017 with litter weakening of the evaporation estimation accuracy. However, the characteristics and determination methods of  $b$  need further studies toward a calibration-free evaporation estimation model.

### **5.4.3.2 Applications of generalized complementary principle**

The generalized complementary functions have been validated or applied in evaporation estimation for many sites (Ai et al., 2017; Brutsaert et al., 2017; Zhang et al., 2017; Han and Tian, 2018a; Crago and Qualls, 2018), and several basins in China (Liu et al., 2016; Gao et al., 2018). It should be noted that most, if not all, above mentioned CR applications need “prior” knowledge in  $E$  (either ground measured or water balance derived) to calibrate the parameters. Recently, the calibration free CR model of S2017 was applied for monthly evaporation estimation across the conterminous China (Ma et al., 2019) and United States (Ma and Szilagyi, 2019). (Brutsaert et al., 2019) (Anayah and Kaluarachchi, 2019) A wide range of model evaluations against the plot scale flux measurements and basin scale water balance results suggested that the generalized complementary functions could serve as a benchmark tool for validating the large scale  $E$  results simulated by those Land Surface models and Remote Sensing models (Ma and Szilagyi, 2019). However, further applications over the world are still needed to develop more long term, high resolution  $E$  datasets for use in hydrological and atmospheric communities.

~~Morton (1983) thought that the ability of the complementary principle to estimate actual evaporation by using meteorological variables only can significantly influence the science and practice of hydrology. However, the attempts in using the complementary principle for hydrological modelling (Oudin et al., 2005; Barr et al., 1997; Nandagiri, 2007) have been suspended, while those attempts in applying such principle in drought assessment (Kim and Rhee, 2016; Hobbins et al., 2016) are still in their infancy. Moreover, the potential applications in agriculture water management are limited in the sense that the irrigation induced changes in potential evaporation at an annual timescale (Ozdogan et al., 2006; Han et al., 2014a; Han et al., 2017). Apparently, the complementary principle did not develop to its full capacity via the linear CR, which leaves a broad space for further applications of the generalized complementary functions.~~

### **4.3 Integrating with other approaches for further development**

Actual evaporation is widely estimated as a reduction of the evaporation demand. The reduction factor was first taken as a function of soil moisture (Penman, 1950; Shuttleworth, 1993), or canopy resistance (Monteith, 1965). This Penman approach or Penman-Monteith equation approach has played a great role in in-parameterizing the evaporation process in hydrological models and the land surface models. The canopy or surface temperature has also been widely used as a water stress indicator (Jackson et al., 1981; Jackson et al., 1988), and the approach based on land surface temperature from remote sensing data has generated increasing attention. At the annual or long term time scales, the reduction factor is taken as a function of the humidity index represented by the ratio of precipitation to potential evaporation, and this method is known as Budyko approach (Yang et al., 2006; Zhang et al., 2001; Budyko, 1974). In the above approaches, the evaporation demand is

assumed to be independent of the land surface (Lhomme, 1997c; Morton, 1983). But at a large area where the land surface significantly interacts with the atmosphere, the evaporation demand will be altered by the changes of the land surface, and this the independent assumption does not hold. Although problems may not arise in diagnostic modelling as current evaporation demand can be observed and applied, they should be considered if these approaches are applied to a large area and used for future prediction or management in prognostic modelling (Han and Tian, 2018b).

Compared to the above approaches relied on the land surface properties, the reduction factor is determined from the atmospheric wetness in the generalized complementary functions (Table 3). The changes in evaporation demand due to the land surface properties are conceptually considered in the complementary principle, which is a theoretical improvement and would be helpful in predicting evaporation with land use changes. In addition, under the conditions that the land surface properties are difficult to get, it is an obvious advantage of the complementary principle using the routinely observed meteorological variables in evaporation estimation. However, the complementary principle assumes that the changes in land surface properties can be accurately and timely detected from the changes of the atmospheric conditions. This assumption requires that the effects of regional or large-scale advections are negligible (Morton, 1983). Outside these situations, the generalized complementary functions may not work well because land surface properties are incompletely inadequately involved. Besides, the components of evaporation from different patches of the spatially heterogeneous surfaces, especially the evaporation from bare soil and the transpiration from vegetation, cannot be separated in the complementary principle, which is its disadvantage compared to the other approaches.

Considering the above disadvantages, Han and Tian (2018b) proposed a framework to integrate the complementary principle with other approaches for the advancement of evaporation research, which expresses  $E/E_{pen}$  as a function of both the land surface properties and the atmospheric wetness. Actually, both the land surface characteristics (e.g., soil moisture and vegetation) and atmospheric variables (e.g., radiation, humidity, and temperature) have been used in the Jarvis–Stewart model (Jarvis, 1976; Stewart, 1988) to parameterize the canopy resistance. In fact, several attempts were conducted by integrating the complementary principle with other approaches to derive some of the land surface variables by using the meteorological variables (Mallick et al., 2013; Han et al., 2015; Szilagyi and Jozsa, 2009). A unified formulation of Penman approach and the linear AA function was proposed by Crago and Brutsaert (1992). The integrated approach is a more rational conceptualization of the evaporation process from the unsaturated surface into the unsaturated atmosphere, and is expected to increase the accuracy of evaporation estimation while reducing the burdens of parameterization. The findings of Liu et al. (2018) and Wang et al. (2019) that the parameters of the generalized functions significantly depends on the wetness of the land surface have demonstrated that the integrated approach has a bright prospect. However, proper manners to integrate them need further studies.

## 6.5 Conclusions



The complementary principle conceptualizes the feedbacks of land surface evaporation on atmospheric evaporation demand and offers advantages in evaporation estimation. In this study, the historical development of the complementary principle during the past half century was reviewed and the two types of generalized complementary functions were focused. In addition, future development for the generalized complementary principle was summarized based on the review. The concluding remarks are as follows:

(1) The studies on the complementary principle adopted a symmetric CR at first, and then extended to an asymmetric CR. At present the original CR has evolved to the generalized complementary principle, which employs nonlinear functions as generalizations of the original linear relationship. The generalized complementary principle has a more rigorous physical base and offers potential in advancing actual evaporation estimation by using simple and standardized procedures.

(2) Two types of generalized complementary functions were derived based on different understandings of the boundary conditions under completely wet environments: the sigmoid H12 and polynomial B15. The B15 inherits the concepts of “potential evaporation  $E_{po}$ ” and “apparent potential evaporation  $E_{pa}$ ” from the original CR, and uses a polynomial function relating  $E/E_{pa}$  to  $E_{po}/E_{pa}$ . By contrast, H12 goes further from the original CR without involving the difficulties in formulating  $E_{po}$  and  $E_{pa}$ . Instead, a sigmoid function relating the ratio of actual evaporation to the Penman potential evaporation  $E_{pen}$  and the proportion of the radiation component in  $E_{pen}$  was derived. Nevertheless, further validation and application of the two types of generalized complementary functions are required with multiple data set over from different parts of the world.

(3) Further studies from both the theoretical and practical aspects are still required before the generalized complementary principle achieves its potential. The generalized complementary principle requires a bold attempt for the practice of hydrology through enhancing its ability of evaporation estimation while reducing the burdens of parameterization. Thus, it should be carefully examined for its physical base of the boundary conditions under completely wet environment, and be integrated with other approaches to include the information of the land surface properly.

## Appendix: List of symbols

Abbreviations of complementary functions	AA	Advection-aridity function proposed by Brutsaert and Stricker (1979)
	H12	Generalized complementary function proposed by Han et al., (2012)
	H2012	Sigmoid analytical form of H12 proposed by Han et al., (2012)
	H2017	Sigmoid analytical form of H12 proposed by Han and Tian (2018)
	B15	Generalized complementary function proposed by Brutsaert (2015)
	B2015	Polynomial applicable form of B15 suggested by Brutsaert (2015)
	C2016	Rescaled applicable form of B15 proposed by Crago et al., (2016)
Three types of evaporation in CR	S2017	Rescaled applicable form of B15 proposed by Szilagyi et al., (2017)
	$E$	Actual evaporation
	$E_{pa}$	Apparent potential evaporation in CR
Specific formulations for $E_{pa}$ or $E_{po}$	$E_{po}$	Potential evaporation in CR
	$E_{Pan}$	Pan evaporation
	$E_{Pen}$	Penman's potential evaporation (Penman, 1948)
	$E_{rad}$	Radiation term of $E_{Pen}$
	$E_{aero}$	Aerodynamic term of $E_{Pen}$
	$E_{Pen}^{KP}$	Modified Penman's equation by Kohler and Parmele (1967)
	$E_{Pen}^{r_s}$	Penman's potential evaporation with temperature and humidity calculated from the ABL model corresponding to certain surface resistance ( $r_s$ )
	$E_{Pen}^{r_s=0}$	Penman's potential evaporation with temperature and humidity calculated from the ABL model corresponding to $r_s = 0$
	$E_{PM}^{r_s \min}$	Penman–Monteith (Monteith, 1965) evaporation with a minimum surface resistance
	$ET_0$	Reference <a href="#">crop</a> evapotranspiration (Allen et al., 1998)
	$E_{MT}$	Mass-transfer type potential evaporation (van Bavel, 1966)
	$E_{Mor}$	Morton (1983)'s potential evaporation
	$E_{PT}$	Priestley-Taylor's (Priestley and Taylor, 1972) minimal advection evaporation
	$E_{PT}^{T_p}$	Morton's modified Priestley-Taylor's minimal advection evaporation (Morton, 1983)
	$E_{PT}^{T_{ws}}$	Szilagyi and Jozsa (2008)'s modified Priestley-Taylor's minimal advection evaporation
	$E_{Pen}^{\max}$	Maximum value of $E_{pa}$ calculated by Penman equation (Szilagyi et al., 2017)
	$E_{MT}^{\max}$	Maximum value of $E_{pa}$ calculated by a mass transfer approach (Crago et al., 2016)
Parameters in CR	$\alpha$	Priestley-Taylor coefficient
	$b$	Symmetry parameter of the CR
Meteorological variables used for calculating $E_{pa}$ or $E_{po}$	$T_a$	Air temperature
	$T_s$	Surface temperature
	$T_{ws}$	Surface temperature under wet environment <a href="#">defined by Szilagyi and Jozsa</a> (2008)
	$T_p$	Equilibrium temperature <a href="#">defined by Morton</a> (1983)
	$\Delta$	Slope of the saturation vapor curve
	$\gamma$	Psychrometric constant
	$R_n$	Net radiation
	$G$	Ground heat flux
	$RH$	Relative humidity

## Acknowledgments

805 This research was partially sponsored by the National Natural Science Foundation of China (No. 51579249, 51825902), the National Key Research and Development Program of China (No. 2016YFC0402707), the Research Fund (No. 2016ZY06) of State Key Laboratory of Simulation and Regulation of Water Cycle in River Basin, China Institute of Water Resources and Hydropower Research.

810 **Code and data availability.** There is no code and data used in this review paper.

**Author contributions.** SH and FT jointly developed the review and edited the manuscript. SH drafted the paper.

**Competing interests.** The authors have declared no conflicts of interest for this article.

## References

- 815 Ai, Z., Wang, Q., Yang, Y., Manevski, K., Zhao, X., and Eer, D.: Estimation of land-surface evaporation at four forest sites across Japan with the new nonlinear complementary method, *Scientific reports*, 7, 17793, 10.1038/s41598-017-17473-0, 2017.
- Ali, M. F., and Mawdsley, J. A.: Comparison of two recent models for estimating actual evapotranspiration using only recorded data, *J Hydrol*, 93, 257-276, 1987.
- 820 Allen, R. G., Pereira, L. S., Raes, D., and Smith, M.: Crop evapotranspiration: Guidelines for computing crop water requirements. FAO irrigation and drainage paper No. 56, FAO irrigation and drainage paper No. 56, Food and Agricultural Organization of the U.N. , Rome, Italy, 1998.
- Aminzadeh, M., Roderick, M. L., and Or, D.: A generalized complementary relationship between actual and potential evaporation defined by a reference surface temperature, *Water Resour Res*, 52, 385-406, 10.1002/2015wr017969, 2016.
- 825 Anayah, F., and Kaluarachchi, J.: Improving the complementary methods to estimate evapotranspiration under diverse climatic and physical conditions, *Hydrol Earth Syst Sc*, 18, 2049-2064, 2014.
- Anayah, F. M., and Kaluarachchi, J. J.: Estimating Global Distribution of Evapotranspiration and Water Balance Using Complementary Methods, *Atmos Ocean*, 57, 279-294, 10.1080/07055900.2019.1656052, 2019.
- Assouline, S., Li, D., Tyler, S., Tanny, J., Cohen, S., Bou-Zeid, E., Parlange, M., and Katul, G. G.: On the variability of the Priestley-Taylor coefficient over water bodies, *Water Resour Res*, 52, 150-163, 10.1002/2015wr017504, 2016.
- 830 Baldocchi, D., Falge, E., Gu, L., Olson, R., Hollinger, D., Running, S., Anthoni, P., Bernhofer, C., Davis, K., and Evans, R.: FLUXNET: A new tool to study the temporal and spatial variability of ecosystem-scale carbon dioxide, water vapor, and energy flux densities, *B Am Meteorol Soc*, 82, 2415-2434, 2001.
- Barr, A. G., Kite, G. W., Granger, R., and Smith, C.: Evaluating three evapotranspiration methods in the SLURP macroscale hydrological model, *Hydrological processes*, 11, 1685-1705, 1997.
- 835 Bouchet, R.: Evapotranspiration réelle et potentielle, signification climatique, *International Association of Hydrological Sciences Publication*, 62, 134-142, 1963.
- Brutsaert, W., and Stricker, H.: An advection-aridity approach to estimate actual regional evapotranspiration, *Water Resour Res*, 15, 443-450, 1979.
- Brutsaert, W.: *Evaporation into the Atmosphere: Theory, History, and Applications*, D. Reidel-Kluwer, Hingham, 1982.
- 840 Brutsaert, W., and Parlange, M. B.: Hydrologic Cycle explains the evaporation paradox, *Nature*, 396, 30, 1998.
- Brutsaert, W.: A generalized complementary principle with physical constraints for land-surface evaporation, *Water Resour Res*, 51, 8087–8093, doi:8010.1002/2015WR017720., 10.1002/2015wr017720, 2015.

- Brutsaert, W., Li, W., Takahashi, A., Hiyama, T., Zhang, L., and Liu, W.: Nonlinear advection - aridity method for landscape evaporation and its application during the growing season in the southern Loess Plateau of the Yellow River basin, *Water Resour Res*, 53, 270–282, 2017.
- Brutsaert, W., Cheng, L., and Zhang, L.: Spatial Distribution of Global Landscape Evaporation in the Early Twenty First Century by Means of a Generalized Complementary Approach, *J Hydrometeorol*, 10.1175/jhm-d-19-0208.1, 2019.
- Budyko, M. I.: *Climate and Life*, Academic presss, San Diego, Calif., 508 pp., 1974.
- Carey, S. K., Barbour, S. L., and Hendry, M. J.: Evaporation from a waste rock surface, Key Lake, Saskatchewan, *Canadian Geotechnical Journal*, 42, 1189-1199, 2005.
- Crago, R., and Crowley, R.: Complementary relationships for near-instantaneous evaporation, *Journal of Hydrology*, 300, 199-211, 2005.
- Crago, R., Szilagyi, J., Qualls, R., and Huntington, J.: Rescaling the complementary relationship for land surface evaporation, *Water Resour Res*, 2016.
- Crago, R., and Qualls, R.: Evaluation of the Generalized and Rescaled Complementary Evaporation Relationships, *Water Resour Res*, 54, 8086-8102, 10.1029/2018WR023401, 2018.
- Crago, R. D., and Brutsaert, W.: A comparison of several evaporation equations, *Water Resour Res*, 28, 951-954, 1992.
- Gao, J., Qiao, M., Qiu, X., Zeng, Y., Hua, H., Ye, X., and Adamu, M.: Estimation of Actual Evapotranspiration Distribution in the Huaihe River Upstream Basin Based on the Generalized Complementary Principle, *Advances in Meteorology*, 2018, 1-9, 10.1155/2018/2158168, 2018.
- Granger, R. J.: A complementary relationship approach for evaporation from nonsaturated surfaces, *J Hydrol*, 111, 31-38, 1989.
- Granger, R. J., and Gray, D. M.: Evaporation from natural nonsaturated surfaces, *J Hydrol*, 111, 21-29, 1989a.
- Granger, R. J., and Gray, D. M.: Evaporation from natural nonsaturated surfaces., *Journal of Hydrology*, 111, 21-29, 1989b.
- Granger, R. J.: Partitioning of energy during the snow-free season at the Wolf Creek research basin, *Wolf Creek Research Basin: Hydrology, Ecology, Environment*, Environment Canada: Saskatoon, 1999.
- Han, S., Hu, H., and Tian, F.: Evaluation of three complementary relationship approaches for regional evapotranspiration in hyperarid Akesu alluvial plain, Northwest China, *Proceedings of the International Symposium on Flood Forecast and Water Resources Assessment for IAHS-PUB*, Beijing, 2006, 75-82,
- Han, S.: *Study on Complementary Relationship of Evapotranspiration and its Application in Tarim River Basin*, Ph.D, Tsinghua University, Beijing, 2008.
- Han, S., Hu, H., and Tian, F.: Evaluating the advection-aridity model of evaporation using data from field-sized surfaces of HEIFE, *IAHS Publication*, 322, 9-14, 2008.
- Han, S., Hu, H., and Yang, D.: A complementary relationship evaporation model referring to the Granger model and the advection-aridity model, *Hydrol Process*, 25, 2094-2101, 2011.
- Han, S., Hu, H., and Tian, F.: A nonlinear function approach for the normalized complementary relationship evaporation model, *Hydrol Process*, 26, 3973-3981, 2012.
- Han, S., Tang, Q., Xu, D., and Wang, S.: Irrigation - induced changes in potential evaporation: more attention is needed, *Hydrol Process*, 28, 2717-2720, 2014a.
- Han, S., Tian, F., and Hu, H.: Positive or negative correlation between actual and potential evaporation? Evaluating using a nonlinear complementary relationship model, *Water Resour Res*, 50, 1322–1336, doi: 1310.1002/2013WR014151, 2014b.
- Han, S., Xu, D., Wang, S., and Yang, Z.: Water requirement with irrigation expansion in Jingtai Irrigation District, Northwest China: The need to consider irrigation-induced local changes in evapotranspiration demand, *Journal of Irrigation and Drainage Engineering (ASCE)*, 140, 04013009, 2014c.
- Han, S., Xu, D., Wang, S., and Yang, Z.: Similarities and differences of two evapotranspiration models with routinely measured meteorological variables: application to a cropland and grassland in northeast China, *Theor Appl Climatol*, 117, 501-510, 2014d.
- Han, S., Tian, F., and Shao, W.: An annual evapotranspiration model by combining Budyko curve and complementary relationship, *EGU General Assembly Conference Abstracts*, 2015, EGU2015-8068,
- Han, S., Xu, D., and Yang, Z.: Irrigation-induced changes in evapotranspiration demand of Awati irrigation district, Northwest China: Weakening the effects of water saving?, *Sustainability*, 9, 1531, 2017.

- Han, S., and Tian, F.: Derivation of a sigmoid generalized complementary function for evaporation with physical constraints, *Water Resour Res*, 54, 5050-5068, doi: 10.1029/2017WR021755, 2018a.
- Han, S., and Tian, F.: Integration of Penman approach with complementary principle for evaporation research, *Hydrol*  
895 *Process*, 32, 3051-3058, 2018b.
- Han, S., and Tian, F.: Reply to Comment by J. Szilagyi and R. Crago on “Derivation of a sigmoid generalized complementary function for evaporation with physical constraints”, *Water Resour Res*, 55, 1734-1736, 10.1029/2018WR023844, 2019.
- Hobbins, M. T., Ramí'ez, J. A., Brown, T. C., and Claessens, L. H. J. M.: The complementary relationship in estimation of  
900 regional evapotranspiration: the CRAE and Advection-aridity models, *Water Resour Res*, 37, 1367-1388, 2001a.
- Hobbins, M. T., Ramirez, J. A., and Brown, T. C.: The complementary relationship in estimation of regional evapotranspiration: An enhanced advection - aridity model, *Water Resour Res*, 37, 1389-1403, 2001b.
- Hobbins, M. T., Wood, A., McEvoy, D. J., Huntington, J. L., Morton, C., Anderson, M., and Hain, C.: The Evaporative Demand Drought Index. Part I: Linking Drought Evolution to Variations in Evaporative Demand, *J Hydrometeorol*, 17,  
905 1745-1761, 10.1175/jhm-d-15-0121.1, 2016.
- Hobbins, M. T., J. A. Ramí'ez, and T. C. Brown: The complementary relationship in estimation of regional evapotranspiration: an enhanced Advection-aridity model, *Water Resources Research*, 37, 1389–1404, 2001.
- Hu, Z., Wang, G., Sun, X., Zhu, M., Song, C., Huang, K., and Chen, X.: Spatial - Temporal Patterns of Evapotranspiration Along an Elevation Gradient on Mount Gongga, Southwest China, *Water Resour Res*, 2018.
- 910 Huntington, J. L., Szilagyi, J., Tyler, S. W., and Pohl, G. M.: Evaluating the complementary relationship for estimating evapotranspiration from arid shrublands, *Water Resour Res*, 47, W05533, 2011.
- Jackson, R. D., Idso, S., Reginato, R., and Pinter Jr, P.: Canopy temperature as a crop water stress indicator, *Water Resour Res*, 17, 1133-1138, 1981.
- Jackson, R. D., Kustas, W. P., and Choudhury, B. J.: A reexamination of the crop water stress index, *Irrigation Science*, 9,  
915 309-317, 1988.
- Jarvis, P. G.: The Interpretation of the Variations in Leaf Water Potential and Stomatal Conductance Found in Canopies in the Field, *Philos Trans R Soc Lond Ser B Biol Sci*, 273, 593-610, 1976.
- Kahler, D. M., and Brutsaert, W.: Complementary relationship between daily evaporation in the environment and pan evaporation, *Water Resour Res*, 42, W05413, doi:05410.01029/02005WR004541, 2006.
- 920 Katerji, N., and Perrier, A.: Modelisation de l'evapotranspiration réelle d'une parcelle de luzerne: rôle d'un coefficient cultural, *Agronomie*, 3, 513-521, 1983.
- Kim, C. P., and Entekhabi, D.: Examination of two methods for estimating regional evapotranspiration using a coupled mixed-layer and surface model, *Water Resour Res*, 33, 2109-2116, 1997.
- Kim, D., and Rhee, J.: A drought index based on actual evapotranspiration from the Bouchet hypothesis, *Geophys Res Lett*,  
925 43, 2016.
- Kohler, M. A., and Parmele, L. H.: Generalized estimates of free-water evaporation, *Water Resour Res*, 3, 997–1005, 1967.
- Kovács, G.: Estimation of average areal evapotranspiration — Proposal to modify Morton's model based on the complementary character of actual and potential evapotranspiration, *J Hydrol*, 95, 227-240, 1987.
- LeDrew, E. F.: A diagnostic examination of a complementary relationship between actual and potential evapotranspiration, *J*  
930 *Appl Meteorol*, 18, 495-501, 1979.
- Lemeyre, R., and Zhang, L.: Evaluation of three evapotranspiration models in terms of their applicability for an arid region, *J Hydrol*, 114, 395-411, 1990.
- Lhomme, J. P.: An theoretical basis for the Priestley-Taylor coefficient, *Bound-Lay Meteorol*, 82, 179-191, 1997a.
- Lhomme, J. P.: An examination of the Priestley-Taylor equation using a convective boundary layer model, *Water Resour*  
935 *Res*, 33, 2571-2578, 1997b.
- Lhomme, J. P.: Towards a rational definition of potential evaporation, *Hydrology & Earth System Sciences*, 1, 257-264, 1997c.
- Lhomme, J. P., and Guilion, L.: Comments on some articles about the complementary relationship, *J Hydrol*, 323, 1-3, 2006.
- Lhomme, J. P., and Guilion, L.: On the link between potential evaporation and regional evaporation from a CBL perspective,  
940 *Theor Appl Climatol*, 101, 143-147, 2010.

- Lintner, B., Gentine, P., Findell, K., and Salvucci, G.: The Budyko and complementary relationships in an idealized model of large-scale land-atmosphere coupling, *Hydrol Earth Syst Sc*, 19, 2119-2131, 2015.
- Liu, G., Liu, Y., Hafeez, M., Xu, D., and Vote, C.: Comparison of two methods to derive time series of actual evapotranspiration using eddy covariance measurements in the southeastern Australia, *J Hydrol*, 454-455, 1-6, 2012.
- 945 Liu, J., and Kotoda, K.: Estimation of regional evapotranspiration from arid and semi-arid surfaces, *J Am Water Resour As*, 34, 27-40, 1998.
- Liu, S., Sun, R., Sun, Z., Li, X., and Liu, C.: Evaluation of three complementary relationship approaches for evapotranspiration over the Yellow River basin, *Hydrol Process*, 20, 2347-2361, 2006.
- 950 Liu, X., Liu, C., and Brutsaert, W.: Regional evaporation estimates in the eastern monsoon region of China: Assessment of a nonlinear formulation of the complementary principle, *Water Resour Res*, 52, doi:10.1002/2016WR019340, 2016.
- Liu, X., Liu, C., and Brutsaert, W.: Investigation of a Generalized Nonlinear Form of the Complementary Principle for Evaporation Estimation, *Journal of Geophysical Research: Atmospheres*, 123, 3933-3942, 10.1002/2017jd028035, 2018.
- Ma, N., Zhang, Y., Szilagyi, J., Guo, Y., Zhai, J., and Gao, H.: Evaluating the complementary relationship of evapotranspiration in the alpine steppe of the Tibetan Plateau, *Water Resour Res*, 51, 1069-1083, 10.1002/2014wr015493, 2015a.
- 955 Ma, N., Zhang, Y., Xu, C.-Y., and Szilagyi, J.: Modeling actual evapotranspiration with routine meteorological variables in the data-scarce region of the Tibetan Plateau: Comparisons and implications, *Journal of Geophysical Research: Biogeosciences*, 120, doi:10.1002/2015JG003006, 10.1002/2015jg003006, 2015b.
- Ma, N., and Zhang, Y.: Comment on “Rescaling the complementary relationship for land surface evaporation” by R. Crago et al, *Water Resour Res*, 53, 6340-6342, 10.1002/2017wr020892, 2017.
- 960 Ma, N., and Szilagyi, J.: The CR of Evaporation: A Calibration - Free Diagnostic and Benchmarking Tool for Large - Scale Terrestrial Evapotranspiration Modeling, *Water Resour Res*, 10.1029/2019wr024867, 2019.
- Ma, N., Szilagyi, J., Zhang, Y., and Liu, W.: Complementary - Relationship - Based Modeling of Terrestrial Evapotranspiration Across China During 1982-2012: Validations and Spatiotemporal Analyses, *Journal of Geophysical Research: Atmospheres*, 124, 4326-4351, 10.1029/2018jd029850, 2019.
- 965 Mallick, K., Jarvis, A., Fisher, J. B., Tu, K. P., Boegh, E., and Niyogi, D.: Latent Heat Flux and Canopy Conductance Based on Penman-Monteith, Priestley-Taylor Equation, and Bouchet's Complementary Hypothesis, *J Hydrometeorol*, 14, 419-442, 10.1175/jhm-d-12-0117.1, 2013.
- McMahon, T. A., Finlayson, B. L., and Peel, M. C.: Historical developments of models for estimating evaporation using standard meteorological data, *Wiley Interdisciplinary Reviews: Water*, 3, 788-818, 10.1002/wat2.1172, 2016.
- McNaughton, K. G., and Spriggs, T. W.: An evaluation of the Priestley and Taylor equation and the complementary relationship using results from a mixed layer model of the convective boundary layer, in: *Estimation of Areal Evapotranspiration*, IAHS Press, Wallingford, UK, 1989.
- Monteith, J. L.: *Evaporation and environment*, Symposium of the Society of Experimental Biology, Cambridge, 1965, 205-234,
- 975 Morton, F.: Potential evaporation as a manifestation of regional evaporation, *Water Resour Res*, 5, 1244-1255, 1969.
- Morton, F.: Catchment evaporation as manifested in climatologic observations, 1970.
- Morton, F. I.: Potential evaporation and river basin evaporation, *Journal of the Hydraulics Division*, 1965.
- Morton, F. I.: Catchment evaporation and potential evaporation further development of a climatologic relationship, *J Hydrol*, 12, 81-99, 1971.
- 980 Morton, F. I.: Estimating evaporation and transpiration from climatological observations, *J Appl Meteorol*, 14, 488-497, 1975.
- Morton, F. I.: Climatological Estimates of Evapotranspiration, *Journal of the Hydraulics Division*, 102(HY3), 275-291, 1976.
- Morton, F. I.: Operational estimates of areal evapotranspiration and their significance to the science and practice of hydrology, *J Hydrol*, 66, 1-76, 1983.
- 985 Nandagiri, L.: Calibrating hydrological models in ungauged basins: possible use of areal evapotranspiration instead of streamflows, *Predictions In Ungauged Basins: PUB Kick-off (Proceedings of the Kick-off meeting held in Brasilia, 20-22 November 2002)* IAHS, 2007,
- Oudin, L., Michel, C., Andr, V., Anctil, F., and Loumagne, C.: Should Bouchet's hypothesis be taken into account in rainfall-runoff modelling? An assessment over 308 catchments, *Hydrol Process*, 19, 4093-4106, 2005.
- 990

- Ozdogan, M., and Salvucci, G. D.: Irrigation-induced changes in potential evapotranspiration in southeastern Turkey: Test and application of Bouchet's complementary hypothesis, *Water Resour Res*, 40, W04301, 2004.
- Ozdogan, M., Woodcock, C. E., Salvucci, G. D., and Demir, H.: Changes in summer irrigated crop area and water use in Southeastern Turkey from 1993 to 2002: implications for current and future water resources, *Water Resour Manag*, 20, 467-488, 2006.
- 995 Parlange, M. B., and Katul, G. G.: An advection-aridity evaporation model, *Water Resour Res*, 28, 127-132, 1992a.
- Parlange, M. B., and Katul, G. G.: Estimation of the diurnal variation of potential evaporation from a wet bare soil surface, *J Hydrol*, 132, 71-89, 1992b.
- 1000 Penman, H. L.: Natural Evaporation from open water, bare soil and grass, *Proceedings of the Royal Society of London, Series A, Mathematical and Physical Sciences*, 193, 120-145, 1948.
- Penman, H. L.: The dependence of transpiration on weather and soil conditions, *J Soil Sci*, 1, 74-89, 1950.
- Pettijohn, J. C., and Salvucci, G. D.: Impact of an unstressed canopy conductance on the Bouchet-Morton complementary relationship, *Water Resour Res*, 42, W09418, doi: 09410.01029/02005WR004385, 2006.
- 1005 Pomeroy, J. W., Granger, R. J., Pietroniro, A., Elliot, J. E., Toth, B., and Hedstrom, N.: Hydrological pathways in the Prince Albert Model Forest: Final Report, National Hydrology Research Institute Environment Canada, Saskatoon, SaskatchewanNHRI Contribution Series No. CS-97007, 153, 1997.
- Priestley, C. H., and Taylor, R. J.: On the assessment of surface heat flux and evaporation using large-scale parameters, *Mon Weather Rev*, 100, 81-92, 1972.
- 1010 Qualls, R. J., and Gultekin, H.: Influence of components of the advection-aridity approach on evapotranspiration estimation, *J Hydrol*, 199, 3-12, 1997.
- Ramirez, J. A., Hobbins, M. T., and Brown, T. C.: Observational evidence of the complementary relationship in regional evaporation lends strong support for Bouchet's hypothesis, *Geophys Res Lett*, 32, L15401, doi:10.1029/2005GL023549, 2005.
- 1015 Rana, G., Katerji, N., Mastrorilli, M., and Moujabber, M.: A model for predicting actual evapotranspiration under soil water stress in a Mediterranean region, *Theor Appl Climatol*, 56, 45-55, 1997.
- Roderick, M. L., Hobbins, M. T., and Farquhar, G. D.: Pan Evaporation Trends and the Terrestrial Water Balance. II. Energy Balance and Interpretation, *Geography Compass*, 3, 761-780, 10.1111/j.1749-8198.2008.00214.x, 2009.
- Seguin, B.: Influence de l'évapotranspiration regionale sur la mesure locale d'évapotranspiration potentielle, *Agr Meteorol*, 15, 355-370, 1975.
- 1020 Shuttleworth, W. J.: Evaporation, in: *Handbook of Hydrology*, McGraw-Hill, New York, 1993.
- Stewart, J. B.: Modelling surface conductance of pine forest, *Agricultural & Forest Meteorology*, 43, 19-35, 1988.
- Su, T., Feng, T., and Feng, G.: Evaporation variability under climate warming in five reanalyses and its association with pan evaporation over China, *Journal of Geophysical Research: Atmospheres*, 120, 8080-8098, 10.1002/2014jd023040, 2015.
- 1025 Sugita, M., Usui, J., Tamagawa, I., and Kaihotsu, I.: Complementary Relationship with a Convective Boundary Layer to Estimate Regional Evapotranspiration, *Water Resour Res*, 37, 353-365, 2001.
- Szilagy, J.: On Bouchet's complementary hypothesis, *J Hydrol*, 246, 155-158, 2001.
- Szilagy, J.: On the inherent asymmetric nature of the complementary relationship of evaporation, *Geophys Res Lett*, 34, L02405, doi:02410.01029/02006GL028708, 2007.
- 1030 Szilagy, J., and Jozsa, J.: New findings about the complementary relationship-based evaporation estimation methods, *J Hydrol*, 354, 171-186, 2008.
- Szilagy, J., Hobbins, M. T., and Jozsa, J.: Modified Advection-Aridity Model of Evapotranspiration, *J Hydrol Eng*, 14, 569-574, 2009.
- Szilagy, J., and Jozsa, J.: Complementary relationship of evaporation and the mean annual water-energy balance, *Water Resour Res*, 45, W09201, 2009.
- 1035 Szilagy, J.: Temperature corrections in the Priestley-Taylor equation of evaporation, *J Hydrol*, 519, 455-464, 10.1016/j.jhydrol.2014.07.040, 2014.
- Szilagy, J.: Complementary-relationship-based 30 year normals (1981-2010) of monthly latent heat fluxes across the contiguous United States, *Water Resour Res*, 51, 9367-9377, 10.1002/2015wr017693, 2015.
- 1040 Szilagy, J., Crago, R., and Qualls, R. J.: Testing the generalized complementary relationship of evaporation with continental-scale long-term water-balance data, *J Hydrol*, 540, 914-922, 2016.

- Szilagyi, J., Crago, R., and Qualls, R.: A calibration - free formulation of the complementary relationship of evaporation for continental - scale hydrology, *Journal of Geophysical Research: Atmospheres*, 122, 264–278, 2017.
- van Bavel, C. H. M.: Potential Evaporation: The Combination Concept and Its Experimental Verification, *Water Resour Res*, 2, 455-467, 1966.
- 1045 Wang, L., Tian, F., Han, S., and Wei, Z.: Determinants of the asymmetric parameter in the complementary principle of evaporation, Submitted to *Water Resources Research*, 2019.
- Wang, Y., Liu, B., Su, B., Zhai, J., and Gemmer, M.: Trends of Calculated and Simulated Actual Evaporation in the Yangtze River Basin, *J Climate*, 24, 4494-4507, 2011.
- Xu, C. Y., and Singh, V. P.: Evaluation of three complementary relationship evapotranspiration models by water balance approach to estimate actual regional evapotranspiration in different climatic regions, *J Hydrol*, 308, 105-121, 2005.
- 1050 Yang, D., Fubao, S., Zhiyu, L., and Cong, Z.: Interpreting the complementary relationship in non-humid environments based on the Budyko and Penman hypotheses, *Geophys Res Lett*, 33, L18402, doi:18410.11029/12006GL027657, 2006.
- Yang, H., Yang, D., and Lei, Z.: Seasonal variability of the complementary relationship in the Asian monsoon region, *Hydrol Process*, 27, 2736–2741, 10.1002/hyp.9400, 2012.
- 1055 Yu, J., Zhang, Y., and Liu, C.: Validity of the Bouchet’s complementary relationship at 102 observatories across China, *Science in China Series D: Earth Sciences*, 52, 708-713, 2009.
- Zhang, L., Dawes, W. R., and Walker, G. R.: Response of mean annual evapotranspiration to vegetation changes at catchment scale, *Water Resour Res*, 37, 701-708, 2001.
- Zhang, L., Cheng, L., and Brutsaert, W.: Estimation of land surface evaporation using a generalized nonlinear complementary relationship, *Journal of Geophysical Research: Atmospheres*, 122, 1475-1487, 10.1002/2016jd025936, 2017.
- 1060