

Assessing the impact of seasonal rainfall anomalies on catchment-scale water balance components

Paolo Nasta^{1,*}, Carolina Allocca¹, Roberto Deidda², Nunzio Romano^{1,3}

¹ Department of Agricultural Sciences, AFBE Division, University of Naples Federico II, Portici (Napoli), Italy.

² Department of Civil and Environmental Engineering and Architecture, University of Cagliari, Cagliari, Italy.

³ The Interdepartmental Research Center for Environment (C.I.R.A.M.), University of Naples Federico II, Napoli, Italy.

* Correspondence to: Paolo Nasta (paolo.nasta@unina.it)

Keywords: Mediterranean climate, Budyko curve, drought, Standardized Precipitation Index, SWAT model, Upper Alento River Catchment

Abstract. Although water balance components at the catchment scale are strongly related to annual rainfall, availability of water resources in Mediterranean catchments also depends on rainfall seasonality. Observed seasonal anomalies in historical records are fairly episodic, but an increase in their frequency might exacerbate water deficit or water excess if the rainy season shortens or extends its duration, e.g. due to climate change. This study evaluates the sensitivity of water yield, evapotranspiration, and groundwater recharge to changes in rainfall seasonality by using the Soil Water Assessment Tool (SWAT) model applied to the Upper Alento River Catchment (UARC) in southern Italy where a long time series of daily rainfall is available from 1920 to 2018. We compare two distinct approaches: *i*) a “static” approach, where three seasonal features (namely rainy, dry, and transition fixed-duration 4-month seasons) are identified through the standardized precipitation index (SPI); *ii*) a “dynamic” approach based on a stochastic framework where the duration of two seasons (rainy and dry seasons) varies from year to year according to a probability distribution. Seasonal anomalies occur when the transition season is replaced by the rainy or dry season in the first approach and when season duration occurs in the tails of its normal distribution in the second approach. Results are presented within a probabilistic framework. We also show that the Budyko curve is sensitive to the rainfall seasonality regime in UARC by questioning the implicit assumption of a temporal steady state between annual average dryness and the evaporative index. Although the duration of the rainy season does not exert a major control on water balance, we were able to identify season-dependent regression equations linking water yield to the dryness index in the rainy season.

30 1. Introduction

31 The rainfall regime of the Mediterranean climate is characterized by the alternation of wet and dry periods within the
32 year, with evident out-of-phase seasonal behavior of precipitation and temperature patterns. Most annual rainfall is
33 concentrated in the late fall, winter, and early spring, while late spring, summer, and early fall are usually hot and quite
34 dry. Rainfall seasonality plays a fundamental role in planning and managing water resources in countries subject to a
35 Mediterranean climate. Summer is characterized by water stress due to scarce rainfall supply, combined with high
36 evapotranspiration loss and the seasonal peak in water consumption (comprising agricultural, industrial, and
37 recreational uses, hydroelectric power generation, as well as domestic uses, which are often boosted by tourism
38 pressure). Therefore, it is necessary to store water during the rainy period to cope with the uncertain duration of adverse
39 water deficit conditions during the dry period. Water-supply infrastructures necessitate high investment costs that
40 strongly depend on the expected balance between the amount of water supplied in the rainy period and the amount of
41 water lost and consumed during the dry season. The amount of rainfall in each season can be suitably decomposed and
42 simulated on the basis of the following three main components: *i*) duration of the seasons; *ii*) occurrence probability of a
43 daily rainfall event in each season; *iii*) mean depth of daily rainfall events in each season (Van Loon et al., 2014). A
44 combination of the last two factors determines the rainfall magnitude in each season (Feng et al., 2013).

45 A very small or very large amount of water (exceeding a certain threshold value for a specified return period and
46 duration) supplied in each season can be interpreted as a seasonal precipitation anomaly and is usually observed
47 episodically in a historical multi-decadal time-series of annual rainfall values. Seasonal precipitation anomalies result
48 mainly from a combination of the duration of the wet season and its rainfall magnitude. These two factors should be
49 taken into due account when planning water-supply infrastructures (Apuv et al., 2017). The most recent reports
50 released by the Intergovernmental Panel on Climate Change (IPCC) warn of the projected increase in seasonal
51 anomalies induced by global warming in the Mediterranean region, with a considerable decrease in annual precipitation

52 and warming-enhanced evapotranspiration associated with rather severe and prolonged droughts, as recently observed
53 in southern Europe in 2003, 2015, and 2017 (Mariotti et al., 2008; Laaha et al., 2017; Hanel et al., 2018).

54 Studies underway in the Upper Alento River Catchment (UARC) in southern Italy offer a good chance to understand
55 the effects of seasonal rainfall uncertainty on water supply generation given the presence of a multi-purpose earthen
56 dam (known as Piano della Rocca) constructed to regulate water for irrigation, hydropower generation, flood control,
57 and drinking purposes. The main research question, also raised or prioritized in some way by local stakeholders in their
58 decision-making processes, can be expressed as follows: “*What is the impact of seasonal rainfall anomalies on annual*
59 *average (or seasonal average) water supply in UARC?*”. This question is particularly relevant to hilly catchments
60 similar to UARC within the Mediterranean region such that UARC could become a pilot area for dealing with some
61 specific problems and carrying out paired-catchment analyses.

62 This study therefore aimed to quantify the effects exerted by seasonal rainfall anomalies on water balance components.
63 With a view to coordinating interaction with stakeholders, end-users, and professionals, we performed this task by
64 implementing the well-known and well-validated Soil Water Assessment Tool (SWAT) model (Arnold et al., 1998).
65 Particular attention is devoted to computing water yield supplying the artificial reservoir bounded by the Piano della
66 Rocca earthen dam in ARC (Romano et al., 2018). One of the strengths of our approach lies in the availability of long-
67 term rainfall time-series (about a century of daily data) and detailed soil and land cover maps, enabling reliable
68 catchment-scale model simulations. Reliable scenario-based projections are built to investigate whether the longer-than-
69 average duration of the wet season implies a higher-than-average mean annual rainfall and consequently higher-than-
70 average water yield. To investigate this issue, our research strategy couples the seasonal duration with daily rainfall
71 occurrences and depths by using a Monte Carlo approach to obtain SWAT-simulated water balance components within
72 a general probabilistic framework.

73 Many authors have attempted to quantify rainfall seasonality using different approaches (Ayoade, 1970; Markham
74 1970; Nieuwolt, 1974; Oliver, 1980; Walsh and Lawler, 1981; Zhang and Qian, 2003; Martin-Vide, 2004; Potter et al.,
75 2005; Feng et al., 2013; de Lavenne and Andréassian, 2018). The precipitation concentration index (PCI) proposed by
76 Oliver (1980) is the most popular approach for quantifying the year-round precipitation distribution in a given study
77 area (Raziei, 2018). Sumner et al. (2001) analyzed the spatial and temporal variation of precipitation seasonality over
78 eastern and southern Spain by using the seasonality index (SI). The SI was also utilized to examine the spatial and
79 temporal variability of precipitation seasonality in Greece (Livada and Asimakopoulos 2005), USA (Pryor and Schoof
80 2008), and northern Bangladesh (Bari et al. 2016). Under the typical Mediterranean climate of Sardinia (Italy), Corona
81 et al. (2018) used the SI to evaluate the role of precipitation seasonality on runoff generation. Nonetheless, while PCI
82 and SI are useful indexes to classify rainfall seasonality and the degree of concentration of rainfall within the year, their
83 implementation in a Monte Carlo framework is not straightforward. Therefore, we opted to characterize rainfall
84 seasonality and its anomalies by using the two approaches described as follows. A first approach, which is hereafter
85 referred to as the static approach, is based on the analysis of the standardized precipitation index (SPI) to define the
86 duration of a wet season (4 months), a dry season (4 months) and a transition season (2 months from dry to wet phase
87 plus 2 months from wet to dry phase) in UARC. In this approach, the drought anomaly is rigidly built with the artifact
88 of extending the duration of the dry season to eight months by removing the transition season. The same criterion
89 applies to a prolonged duration of the rainy season. The second approach, instead, exploits the seasonality
90 characterization proposed by Feng et al. (2013) and can be viewed as a dynamic approach since the duration of the rainy
91 season is time-variant (inter-annual variability) and can be stochastically generated with random duration values drawn
92 from their statistical distribution. This second approach investigates what happens to the water budget if the duration of
93 the rainy season becomes shorter-than-normal (i.e. rainfall scarcity) or longer-than-normal (i.e. rainfall excess). As far
94 as we are aware, there is still a lack of knowledge about the effects of possible changes in rainfall seasonality on the
95 water balance of a catchment subject to a Mediterranean climate, and the analyses presented in this paper aim primarily
96 to contribute to fill this gap.

97 **2. Study area and experimental analyses**

98 The Upper Alento River Catchment (UARC) is situated in the Southern Apennines (Province of Salerno, Campania,
99 southern Italy) and has a total drainage area of about 102 km² (Fig.1). The Piano della Rocca dam is an earthen
100 embankment with an impervious core that has been operating since 1995. The area consists mostly of relatively poor-
101 permeable arenaceous-clayey deposits and secondarily of arenaceous-marly-clayey and calcareous-clayey deposits
102 (Romano et al., 2018).

103 *Please insert Fig. 1 here*

104 A weather station managed by the Italian Hydrological Service is located near the village of Gioi Cilento and provides a
105 dataset of daily rainfall values covering the period 1920-2018 (about 90 years), with an interruption of nine years (1942-
106 1950) straddling World War II (Nasta et al., 2017). The data set of annual rainfall sums derived from the daily rainfall
107 time series has a mean of 1,229.3 mm, while other metrics (median, standard deviation and coefficient of variation) are
108 reported in the last row of Table 1. The same statistics are also summarized for rainfall depths in each month of the
109 year. The variability exhibited by the monthly time series of rainfall depths is also depicted in Figure 2, denoting a
110 typical Mediterranean seasonal cycle. A large amount of precipitation occurs in the months from October to March, a
111 period commonly identified as a wet period of the hydrological year, and accounts for about 68% of the annual mean
112 rainfall (i.e. 834.9 mm over 1,229.3 mm) (see Table 1 and Figure 2). November is the wettest month with an average
113 monthly rainfall of 166.9 mm (about 14% of mean annual rainfall). In contrast, lower mean monthly rainfall depths are
114 concentrated from April to September, which commonly identify a dry period of the hydrological year, with a
115 cumulative rainfall over this period of 394.5 mm with respect to the annual mean of 1,229.3 mm, hence representing
116 about 32% of mean annual rainfall. July is the driest month with a monthly mean rainfall of 29.8 mm (i.e. 2% of mean
117 annual rainfall).

118 *Please insert Fig. 2 here*

119 *Please insert Table 1 here*

120 Within the monitoring activities of the MOSAICUS (MONitoring and modeling Soil–vegetation–atmosphere processes
121 in the Alento river basin for Implementing adaptation strategies to Climate and land USE changes) project (Nasta et al.,
122 2013; Romano et al., 2018), an automated weather station was installed in 2004 close to the village of Monteforte
123 Cilento and equipped with sensors for monitoring rainfall, wind speed and direction, air temperature and relative
124 humidity, and solar radiation, to record such meteorological variables at 15 min intervals (Nasta et al., 2019). The
125 statistical distributions of weather data recorded at the weather station of Monteforte Cilento (2004-2018) will be used
126 to calculate potential evapotranspiration as described in Section 3.

127 In this study, we used the most recent available land-use map drawn up in 2015 by using second-level CORINE
128 (Coordination of Information on the Environment) Land-Cover classes (CORINE 2006 land cover dataset;
129 <http://www.eea.europa.eu>): forest, arable land (annual crops), permanent crops (orchards, vineyards, olive groves, and
130 fruit trees), pasture, urban fabric, and water bodies. Forest (evergreen and deciduous trees, and multi-stem evergreen
131 sclerophyllous Mediterranean shrubs) and agricultural (arable land, permanent crops, and orchards) cover about 70%
132 and 20% of the catchment, respectively (Nasta et al., 2017). A five-meter resolution Digital Terrain Model (DTM) was
133 used to generate the hydrographic network and a soil-landscape units map is used to depict soil attributes in UARC
134 (Nasta et al., 2018).

135 **3. Parameterization of the SWAT Model**

136 The Soil Water Assessment Tool (SWAT) is a bucket-type, semi-distributed hydrological model operating on a daily
137 time scale and at a catchment spatial scale (Arnold et al., 1998). The main components of the water balance equation are
138 the daily change in water storage (ΔWS) as affected by rainfall (R), actual evapotranspiration (ET_a), groundwater
139 recharge (GR), and water yield (WY). Water yield is given by the contribution of surface runoff, groundwater
140 circulation, and lateral flow within the soil profile, and is partially depleted by transmission losses from tributary
141 channels and water abstractions. All variables are expressed in units of mm of water height.

142 SWAT requires as input rainfall (R) and potential evapotranspiration (ET_p) time series at a daily scale and is based on
143 the concept of hydrological response units (HRUs), which are areas identified by similarities in soil, land cover, and
144 topographic features, where hydrological processes are represented by a lumped schematization. The five-meter DTM
145 of the study area was used to determine the catchment boundaries, the hydrographic network, and thirteen distinct
146 HRUs. Catchment-lumped parameters are assigned to each HRU through look-up tables. By using the available soil-
147 landscape unit map, the input parameters were assigned according to the model set-up as presented in Nasta et al.
148 (2017). Nine parameters were calibrated to achieve the best model fit between simulated and measured monthly water
149 yield data recorded from 1995 and 2004 (Nasta et al., 2017). Such hydrological parameters include the soil evaporation
150 and compensation factor, plant uptake compensation factor, Manning's value for overland flow, the baseflow recession
151 constant (groundwater flow response to changes in recharge), groundwater delay time, groundwater "revap" coefficient
152 (controlling water that moves from the shallow aquifer into the unsaturated zone), Manning's coefficient for the main
153 channel, effective hydraulic condition in the main channel alluvium, and the bank storage recession curve. Model
154 performance proved to be satisfactory at a monthly time scale. We ran numerical simulations at a daily time step
155 (rainfall was randomly generated at a daily time step) and aggregated the output fluxes at a monthly time resolution.
156 Although there is evidence in the body of scientific literature of a potential misfit between measured and simulated
157 water yield values at a daily time-scale when calibrating a model with data at a monthly time resolution (Adla et al.,
158 2019), we are confident that our results and conclusions will not be affected by this drawback. Our analysis is based on
159 the monthly aggregation of fluxes and is aimed at analyzing seasonal patterns of monthly aggregates.

160 This study is based on modeling scenarios implemented in SWAT through a Monte Carlo approach, where each
161 simulation is three years long. Results from the first two-year warm-up period are discarded, while water balance
162 components simulated for the third year are stored for subsequent analysis. Initial soil water storage is set as field
163 capacity. The model simulations of the first two years are disregarded in order to erase the impact of the initial
164 (unknown) soil moisture values set in the soil domain. We point out that initial soil water content set at field capacity

165 can be considered a realistic situation in winter under the Mediterranean climate. The rainfall data are generated for the
166 static and dynamic approaches (described below) using a probability setting calibrated on daily rainfall values recorded
167 at the Gioi Cilento weather station (1920-2018). Mean and standard deviation of the meteorological data (wind speed,
168 air temperature and relative humidity, and solar radiation) recorded at the second automated weather station (close to
169 the village of Monteforte Cilento) are calculated each month. Daily potential evapotranspiration data were calculated by
170 using random values of weather data drawn from their normal distribution in each month of the year (Allen et al., 1998).
171 Results were provided as input to SWAT to randomly generate daily potential evapotranspiration by using the Penman-
172 Monteith equation (Allen et al., 1998).

173 **4. Determination of rainfall seasonality**

174 **4.1. Static approach based on the SPI drought index**

175 The intra-annual rainfall regime under a Mediterranean climate can be characterized through the distribution of annual
176 rainfall depth among different seasons (Paz and Kutiel, 2003; Kutiel and Trigo, 2013). The seasonal pattern occurring in
177 the study area is here characterized by analyzing the distribution of the standardized precipitation index (SPI) on a long-
178 term monthly rainfall time series. The SPI is a probability index developed to classify rainfall anomalies and often
179 employed as an indicator of potential (meteorological) droughts over many time scales (McKee et al., 1993; Hayes et
180 al., 1999). Computation of the SPI should rely on long-term rainfall datasets (e.g. 30 years, according to climatological
181 standards), and is usually obtained by projecting a Gamma distribution fitted on rainfall depths cumulated on 1, 3, 6, 12,
182 18, or 24 months (referred to as SPI-1, SPI-3, SPI-6, SPI-12, SPI-18, or SPI-24, respectively) into a standardized normal
183 distribution. The short-term SPI (e.g. 3-month time scale) can provide useful information for crop production and soil
184 moisture supply, while the long-term SPI (e.g. 12- or 24-month time scale) can give insights on water availability for
185 groundwater recharge. Negative SPI values indicate drier-than-expected rainfall, whereas positive SPI values refer to
186 wetter-than-expected months. To quantify the degree of departure from median conditions, McKee et al. (1993)
187 proposed a rainfall regime classification. Since the SPI is given in units of standard deviation from the standardized

188 mean, this statistical index enables also the precipitation anomaly to be identified through the magnitude of its value:
189 values ranging from -0.99 to $+0.99$ are considered near normal, from $+1.00$ to $+1.49$ (or from -1.49 to -1.00) indicate
190 moderately wet (or moderately dry) periods, from $+1.50$ to $+1.99$ (or from -1.99 to -1.50) very wet (or very dry)
191 periods, and above $+2.00$ (or below -2.00) extremely wet (or extremely dry) periods. Therefore, the extent of SPI
192 departure from the mean (i.e. from the zero value) gives a probabilistic measure of the severity of a wet (if positive) or
193 dry (if negative) period. By exploiting the properties of the (standard) normal distribution, the probabilities of obtaining
194 SPI values greater than $+1$, $+2$, and $+3$ (or less than -1 , -2 , and -3) are 15.90%, 2.28% and 0.14%, respectively.
195 To emphasize the seasonal cycle of intra-annual rainfall patterns within a probabilistic framework, we used the SPI-1 by
196 fitting the Gamma distribution on all monthly rainfall depths, i.e. pooling observations from all months in each year. In
197 such a way, the months characterized by SPI-1 values below, around or above the zero line can be assumed to belong to
198 the dry, transition or wet seasons, respectively.

199 **4.2. Dynamic approach based on the duration of the wet season proposed by Feng et al. (2013)**

200 According to Feng et al. (2013), the dimensionless seasonality index (DSI) is based on the concept of relative entropy
201 and quantifies the rainfall concentration occurring in the wet season. The DSI is zero when the average annual rainfall is
202 uniformly distributed throughout the year and maximized at 3.585 when maximum average annual rainfall is
203 concentrated in one single month (Pascale et al., 2016); see the Appendix for details. Feng et al. (2013) proposed to
204 describe the rainfall seasonality through the following three components: annual rainfall depth (magnitude), centroid
205 (timing), and spread (duration) of the wet season (see also Pascale et al., 2015; Sahani et al., 2018). As described in
206 Section 5.2 and according to appropriate statistical tests, we found that a normal distribution can reasonably describe the
207 90 wet season durations obtained by applying to the observed rainfall time series the procedure proposed by Feng et al.
208 (2013), and briefly summarized in the Appendix. Thus, each hydrological year will consist of the alternation of only
209 two seasons: the wet season with a duration that is randomly generated by a normal distribution with mean and standard
210 deviation estimated on the Gioi Cilento time series, and a dry season in the subsequent months of the year.

211 **4.3 Set-up of Monte-Carlo rainfall scenarios in SWAT**

212 Seasonal rainfall anomalies, although episodic, can affect the water balance components at the catchment scale. As
213 suggested by Domínguez-Castro et al. (2019), the impact of such anomalies can be quantified within a probabilistic
214 framework. For the Upper Alento River Catchment (UARC), we evaluated the effects of seasonal anomalies by running
215 SWAT simulations with synthetic rainfall time series considering different hypotheses (scenarios) of alternations of
216 seasons, according to the static and the dynamic approaches described above. In each season, we assumed that rainfall
217 evolution in time can be represented by a stochastic Poisson point process of daily rainfall occurrences, with daily
218 rainfall depth following a proper probability distribution (Eagleson, 1972; Rodríguez-Iturbe et al., 1987; Veneziano and
219 Iacobellis, 2002). Synthetic rainfall time series were then generated, keeping constant parameters of the Poisson process
220 and daily rainfall parent distribution in each season.

221 A preliminary analysis was conducted to investigate the best parent distribution for observed rainfall daily depths. With
222 this aim, we used the L-moment ratios diagram proposed by Hosking (1990) (see also Vogel and Fennessey, 1993) as a
223 diagnostic tool. Results are shown in Figure 3 where the L-skewness and L-kurtosis computed on the time series left-
224 censored with a threshold of 3 mm (large filled circle) is compared with the theoretical expectation of the same L-
225 moment ratios for several probability distributions commonly adopted in statistical hydrology. An ideal candidate as
226 parent distribution seems the Generalized Pareto distribution (GPd), although it is also worth noting that sample
227 estimation of L-skewness and L-kurtosis (0.3437, 0.1706) is very close to the expected values for exponential
228 distribution (1/3, 1/6). As visual support for this preliminary analysis, the exponential probability plot in Figure 4
229 compares the empirical cumulative distribution function $F(x)$ of the observed time series (circles) with the fitted GPd
230 (dashed line) and the fitted exponential distribution (continuous line). The two models are very close to each other for
231 the whole body of observation, with only a slight departure of the GPd from the straight line characterizing the
232 exponential distribution due to a very slight right tail. This evidence gave us the confidence to adopt the single-
233 parameter exponential model as parent distribution for series partitioned according to the seasons defined above,
234 thereby reducing the uncertainty related to the additional shape parameter of the GPd. Finally, it is worthwhile

235 mentioning that both distributions shown in Figure 4 were fitted by applying Deidda’s (2010) multiple-threshold-
236 method (MTM) on a range of thresholds from 2.5 to 12.5 mm to prevent biases due to very low records and data
237 discretization (Deidda, 2007). The MTM was then applied to estimate the exponential parameter η (mm) and the
238 probability occurrence of rainy days λ (d^{-1}) for each season considered.

239 For each scenario pertaining to either the static or dynamic approach, we generated 10,000 equi-probable realizations of
240 synthetic daily rainfall time series, each three years long, according to a stochastic Poisson point process model. In each
241 modeling scenario, the synthetic time series was then used as input for the SWAT model to evaluate the effects on the
242 water balance components in UARC. As anticipated in Section 3, the first two years represent warm-up simulations and
243 were thus discarded, while only results for the third year were stored for subsequent analyses presented in the next
244 section.

245 *Please insert Fig. 3 here*

246 *Please insert Fig. 4 here*

247 To further evaluate the hydrologic behavior of the study catchment, an issue deserving more detailed attention is the
248 assessment of the sensitivity of water balance to rainfall seasonality. With this aim, we refer to the Budyko framework
249 (Budyko, 1974), which has been extensively applied to relate water components in different climatic contexts
250 worldwide, including the Mediterranean climate (see e.g. Viola et al., 2017, Caracciolo et al. 2017). Specifically, the
251 Budyko framework relates the evaporative index (ET_a/R) to the dryness index (ET_p/R) computed at an annual time scale
252 in terms of “available water” (i.e., rainfall R). Potential evapotranspiration, ET_p , is limited by either energy supply (for
253 the dryness index less than or equal to one) or water supply (for the dryness index greater than one), and therefore the
254 Budyko space has two physical bounds dictated by either the atmospheric water demand ($ET_a \leq ET_p$) or the atmospheric
255 water supply ($ET_a \leq R$). The first bound is the energy limit (or demand limit, i.e. the 1:1 line corresponding to $ET_a = ET_p$)
256 implying that actual evapotranspiration cannot exceed potential evapotranspiration. The second bound is the water limit

257 (or supply limit, i.e. the horizontal line corresponding to $ET_a=R$) implying that actual evapotranspiration cannot exceed
258 precipitation when the dryness index is greater than one (i.e. $ET_p/R>1$).

259

260 **5. Results and discussion**

261 **5.1. Static approach for assessing rainfall seasonality**

262 The observed temporal evolution of SPI-6 in our 90-year time series (see gray bars in Fig. 5) highlights prolonged
263 droughts amongst the 1980s and 1990s and prolonged wet periods in the last decade when SPI-6 values above the
264 threshold +2 occurred in 2008, 2010, and 2012. Yet, by splitting the SPI-6 values into two 45-year sub-groups, we can
265 observe that the last 45-year period is characterized by a drier climate compared to the first 45-year period. Specifically,
266 in the first sub-group the probabilities of obtaining $SPI-6>+1$ and $SPI-6<-1$ are 17.9% and 7.6%, respectively. In
267 contrast, in the second sub-group there is a general increase in negative SPI-6 values: the probability of obtaining SPI-
268 $6>+1$ becomes 11.9% and that of obtaining $SPI-6<-1$ 19.3%. By analyzing daily rainfall datasets recorded at 55
269 weather stations located in the region of Basilicata near UARC (characterized by similar climatic conditions), Piccarreta
270 et al. (2013) observed a general decreasing trend in the mean annual rainfall over the period 1951–2010 mainly due to
271 the autumn-winter decrease in precipitation.

272 *Please insert Fig. 5 here*

273 We now discuss the results pertaining to the calculation of the seasonal pattern of SPI-1 values. Rainfall seasonality
274 under a Mediterranean climate can be assumed to be roughly represented by the alternation of two six-month seasons,
275 characterized by positive and negative SPI-1 values (wet and dry season, respectively) (Rivoire et al., 2019). The
276 temporal evolution of the SPI-1 values is represented by the gray bars in Fig. 6a and highlights the seasonal cycle within
277 each year, whereas their 12-month moving average (magenta line in Fig. 6a) oscillates around the zero value with
278 prolonged dry periods during the 1980s and 1990s and prolonged wet periods in the 2000s and 2010s. Fig. 6b shows the
279 box and whiskers plots of the SPI-1 values for each month of the year, thus depicting the monthly distribution of this

280 index throughout the available recorded period. The median SPI-1 values (central red line in the blue boxes) are
281 negative only from May to August and positive from September to April, even though the whiskers (identified by the
282 two lines at the 25th and 75th percentile) denote the presence of relatively large variability in almost all months. Closer
283 inspection of this graph enables one to identify three main seasonal features: *i*) a dry period from May till August with
284 median values below zero; *ii*) a rainy period from November till February with median values above zero; *iii*) two
285 transition periods from wet to dry (March and April) and from dry to wet (September and October) with median values
286 near zero. We are aware that the median values in March, April, and October of the transition season are above zero,
287 rather than “near” zero, but we recall that the Mediterranean climate in UARC is sub-humid mainly due to orographic
288 influences. However, this approach is intrinsically a “static” procedure since the subdivision of the twelve months into
289 three groups is rigid even though months in the transition periods have high variability in SPI-1 values. This outcome
290 refines the initial working hypothesis of seasonal alternation of two semesters.

291 *Please insert Fig. 6 here*

292 The frequency distributions of the SPI-1 values computed over the rainy, dry, and transition seasons are illustrated in
293 Fig.6c-6d-6e. The wet season (depicted by the blue histograms) is characterized by probabilities of having SPI-1 values
294 greater than 0, +1, +2, and +3 of 80.60%, 30.50%, 1.90%, and 0.30%, respectively. The dry season (depicted by the red
295 histograms) is associated with SPI-1 values lower than 0, -1, -2, and -3 with probabilities of 78.10%, 31.10%, 0.56%
296 and 0.10%, respectively. Conversely, we warn that probabilities of obtaining positive SPI-1 values in the transition
297 season are 63.30% instead of the expected 50% if the hypothesis were “perfectly true”. Therefore, we considered three
298 different scenarios, each with fixed and recurrent alternation of seasons during the hydrological year: *i*) a “reference
299 scenario” with a four-month wet season (NDJF), a four-month dry season (MJJJA), and a four-month transition season
300 (MA from wet to dry and SO from dry to wet); *ii*) a “dry scenario”, which mimics an extreme drought anomaly,
301 characterized by a prolonged eight-month dry season (from March to October) and abrupt alternations with the four-
302 month wet season (NDJF), without any transition season; *iii*) a “wet scenario”, which mimics an extreme rainy

303 anomaly, characterized by a prolonged eight-month wet season (from September to April) and abrupt alternations with
304 the four-month dry season (MJJA), again with no transition season.

305 In light of the above results, the two Poisson parameters (η and λ) describing daily rainfall values were calculated for
306 each of the three seasons in the “reference scenario” and were then also used to develop synthetic simulations of rainfall
307 time series in the “dry” and “wet” scenarios (see Table 2).

308 *Please insert Table 2 here*

309

310 **5.2. Dynamic approach for assessing rainfall seasonality**

311 The centroid of the monthly rainfall distribution measured at the Gioi Cilento weather station (in the 90 years between
312 1920 and 2018) indicates that the wet season is centered in the second half of December, while its average duration is
313 about 5.44 months (see Fig. 7). Nonetheless, it is worth noting the occurrence of a few extreme situations: the severe
314 drought recorded in 1985 caused a minimum duration of about four months of the rainy period, while the year 1964
315 registered a maximum duration of about 7.0 months. The term “dynamic” used for this approach stems mainly from the
316 fact that the duration of the rainy period is time-variant throughout the years.

317 *Please insert Fig. 7 here*

318 The dimensionless seasonality index (DSI) and the seasonality index (SI) were computed for the Gioi Cilento time
319 series according to procedures proposed by Feng et al. (2013) and by Walsh and Lawler (1981), respectively. The
320 Mann-Kendall nonparametric test (Mann, 1945; Kendall, 1975) was then applied to evaluate possible decreasing,
321 increasing, or absence of temporal trends on these indexes, and revealed that the null hypothesis of absence of trend
322 cannot be neglected at the 0.05 significance level for both indexes. The stationarity in time of the DSI (red line) and SI
323 (green line) is also apparent from a perusal of Fig. 8, where the linear regressions (dashed and dotted for the DSI and SI,
324 respectively) are characterized by very weak downward slopes.

325 *Please insert Fig. 8 here*

326 As described in Section 4.2, the dynamic approach assumes the alternation of only two seasons (wet and dry) with
327 random durations of the rainy period. Figure 9a shows the time series of the 90 durations of the wet season estimated
328 with the procedure proposed by Feng et al. (2013), while their frequency distribution is plotted in Fig. 9b. We then
329 applied the Lilliefors statistical test (Lilliefors, 1967) to the null hypothesis of normality for the estimated wet durations
330 obtaining a p -value of 0.327, meaning that the null hypothesis cannot be rejected with the commonly adopted 5%
331 significance level. For each hydrological year, we thus generate a duration of the wet season from a normal distribution
332 with the same mean and standard deviation of the Gioi Cilento time series (with a mean of 2.71 months and standard
333 deviation of 0.28 months), while the dry seasons were consequently obtained as the complement in the same year to the
334 wet seasons. In this case, the two Poisson parameters (η and λ) for modeling daily rainfall values were computed for the
335 wet and dry seasons (Table 3).

336 *Please insert Fig. 9 here*

337 *Please insert Table 3 here*

338

339 **5.3. Effects of seasonal rainfall anomalies on water balance when using the static approach**

340 The results obtained from the three scenarios pertaining to the static approach are presented using the descriptive
341 statistics of the water balance components at the annual time scale obtained from 10,000 SWAT simulation runs (Table
342 4). The reference scenario represents the normal situation with three seasons (dry, transition, and wet). Even though the
343 range of annual rainfall values is relatively large, the coefficient of variation (CV) is only 14%, implying that very low
344 and very high annual rainfall depths (outliers) occur occasionally. The water balance components, namely water yield
345 (WY), actual evapotranspiration (ET_a), and groundwater recharge (GR), represent on average 35%, 49%, and 16% of the
346 annual mean rainfall depth ($R=1,229$ mm). The annual rainfall depths for the other two scenarios (only two seasons
347 without the transition season) shift down to 988 mm (dry scenario) and up to 1,393 mm (wet scenario), thus affecting

348 the water balance. When the dry season lasts eight months (dry scenario), water yield, actual evapotranspiration, and
349 groundwater recharge decrease by 116 mm, 60 mm, and 66 mm, respectively, when compared to the reference scenario.

350 *Please insert Table 4 here*

351

352 In contrast, when the wet season lasts eight months (wet scenario), the water yield, actual evapotranspiration, and
353 groundwater recharge increase by 93 mm, 21 mm, and 54 mm, respectively, when compared to the reference scenario.

354 Water yield, actual evapotranspiration, and groundwater recharge represent on average 32%, 55%, and 13% of the
355 annual rainfall depth in the extreme dry season (dry scenario), and 38%, 45%, and 18% of annual rainfall depth in the
356 extreme wet season (wet scenario).

357 Decomposition of the annual results into the seasonal components highlights other interesting features that are shown in
358 Fig. 10 (rainfall and potential evapotranspiration forcings) and in Fig. 11 (main water balance components). For the
359 reference scenario the seasonal rainfall depth is 201 mm, 436 mm, and 593 mm for the dry, transition, and wet seasons,
360 respectively, representing 16%, 35%, and 48% of the total annual rainfall (see Fig. 10a). Water yield depths span from
361 44 mm during the dry season to 251 mm during the rainy season (see Fig. 11a). Almost 60% of annual water yield
362 occurs over the wet season, about 30% in the transition season, and about 10% in the dry season. In contrast, the actual
363 evapotranspiration depths are higher than rainfall depths in the dry season (269 mm) and lower than rainfall depths
364 during the transition (226 mm) and rainy (110 mm) seasons (see Fig. 11a).

365 *Please insert Fig. 10 here*

366 *Please insert Fig. 11 here*

367

368 Over the dry scenario (see Figs. 10b and 11b), the months belonging to the transition season become drier-than-normal.
369 The total rainfall depths over the dry and wet seasons are 397 mm and 590 mm, respectively, whereas the extreme
370 drought anomaly induces precipitation loss only in the dry season with a considerable decrease of 239 mm of rainfall

371 depth (Fig. 10b). The consequences of this situation on the average water balance components in the prolonged dry
372 season lead to significant deficits (Fig. 11b). Water yield loss over the dry season is 93 mm, which represents 50% of
373 water yield obtained for the dry and transition seasons in the reference scenario. The wet season (from November to
374 February) provides about 590 mm of water yield per year. The water loss by actual evapotranspiration is limited and
375 represents only 10% of ET_a obtained for the dry and transition seasons in the reference scenario (Fig. 11b).
376 In the wet scenario (see Fig. 10c and Fig. 11c), the months belonging to the transition season become wet (8 wet months
377 and 4 dry months). Total rainfall depths in the dry and wet seasons are 200 mm and 1,193 mm (Fig. 10c). Rainfall depth
378 increases by 164 mm in the wet season (+14% compared with that obtained in the wet and transition seasons in the
379 reference scenario). Water yield gain in the wet season is 89 mm which represents 20% of water yield obtained in the
380 wet and transition seasons in the reference scenario (Fig. 11c). The water lost by actual evapotranspiration is negligible.

381 **5.4. Effects of seasonal rainfall anomalies on water balance when using the dynamic approach**

382 The second approach to assessing the effect of rainfall seasonality extremes on water balance components is based on
383 the stochastic generation of the wet season durations from their normal distribution (see Fig. 9b). This approach helps
384 classify the results within a probabilistic framework according to the following rainy period duration classes: 3-4
385 months, 4-5 months, 5-6 months, 6-7 months, 7-8 months. Seasonal extremes (3-4 months and 7-8 months) have very
386 low probabilities of occurrence (0.60% and 0.30%, respectively). Nonetheless, it is interesting to analyze the effect of
387 rainfall variability on water yield (WY), actual evapotranspiration (ET_a) and groundwater recharge (GR). The most
388 probable (62%) situation occurs when the rainy period lasts 5-6 months. Under these circumstances, the mean annual
389 rainfall depth is 1,275 mm, whereas WY , ET_a , and GR represent 35%, 49%, and 16% of annual average rainfall depth,
390 respectively. These percentages are very close to those observed in the reference scenario of the static approach. If the
391 wet season shortens by one month (23% probability), the mean annual rainfall depth decreases by 62 mm, whereas
392 water yield depth by 33 mm (-7%). In contrast, if the wet season is made up of 6-7 months (14% probability), the
393 annual mean rainfall depth increases by 51 mm and water yield by 27 mm (+6%).

394 Extreme dry and extreme wet situations reflect similar results obtained from the dry and wet scenarios presented above.
395 A prolonged drought (i.e. rainy period only 3-4 months long) leads to an average rainfall loss of 130 mm per year
396 inducing an appreciable annual decrease in both water yield (-68 mm) and groundwater recharge (-30 mm). A
397 prolonged wet season (i.e. lasting 7-8 months), instead, causes average rainfall to gain approximately 108 mm per year,
398 yielding annual increases in both water yield (+59 mm) and groundwater recharge (+12 mm). It is worth noting that the
399 duration of the rainy period does not seem to exert a major control on the water balance. Pearson's linear correlation
400 coefficients between duration and average annual rainfall, water yield, and actual evapotranspiration are 0.22, 0.20, and
401 0.11, respectively.

402 *Please insert Table 5 here*

403 *Please insert Fig. 12 here*

404 Assuming that the long-term mean annual precipitation can be partitioned into the mean annual actual
405 evapotranspiration and mean annual water yield, according to the Budyko framework we assume that larger values of
406 the dryness index (drier climate conditions; $ET_p/R > 1$) induce a greater proportion of rainfall that is partitioned to ET_a .
407 In contrast, data on the left-hand side of the Budyko curve will be characterized by a greater proportion of rainfall that
408 is partitioned to water yield. Fig. 12 shows the Budyko plot of the dryness index (ET_p/R) versus the evaporative index
409 (ET_a/R) together with the Budyko curve (solid garnet line). In this plot we depict the data points (colored dots) for the
410 five different durations of the rainy period in UARC obtained by the dynamic approach. The first comment to be made
411 is that all of these data points gather within the energy-limited region of the Budyko plot, with the longest rainy period
412 (blue dot) favoring conditions of greater discharges (evaporative index $ET_a/R=0.45$) and the shortest rainy period
413 (droughts indicated by the red dot) inducing higher evapotranspiration fluxes (evaporative index $ET_a/R= 0.54$). The
414 latter situation shows that on average the Upper Alento River catchment is characterized by relatively good storage of
415 soil-water made possible by the hydraulic properties of the soils and the large portion of shrub spots and forest areas
416 (mostly deciduous chestnut forests and olive orchards), together with a good amount of annual precipitation in a hilly

417 and mountainous zone in southern Italy. However, it may also be noted that all of these data points cluster below the
418 Budyko curve (Williams et al., 2012). The observed departure below the Budyko curve may be due to several reasons.
419 Allowing for the Budyko assumptions for water balance, the present study refers to a long time scale (90 years), but a
420 relatively small spatial scale since UARC has a drainage area of 102 km². In fact, rainfall seasonality (i.e. intra-annual
421 variability) may be just one of the major factors that could have led to a departure from the Budyko curve. The typical
422 Mediterranean climate, which is characterized by precipitation being out-of-phase with potential evapotranspiration, is
423 also singled out as a cause of the deviations we observed in our case study from the Budyko curve (Milly, 1994).
424 Normal situations, characterized by a wet season lasting 5-6 months (green dot), lead to rainfall being partitioned into
425 49% ET_a , as indicated by the evaporative index value of 0.49. We hereby recall that this study is based on the
426 assumption that the catchment response is not affected by human interferences and their feedbacks (land-use change,
427 change in soil hydraulic properties, enhanced evapotranspiration induced by global warming, etc.), but only by changes
428 in rainfall seasonality which, of course, can undermine Budyko's implicit assumption of temporal steady-state (Feng et
429 al., 2012; Troch et al., 2013).

430 *Please insert Fig. 13 here*

431 *Please insert Table 6 here*

432 The relationships between the seasonal dryness index and water yield to rainfall ratio (WY/R) are affected by the
433 duration of the wet season and are depicted in Fig. 13. The coefficients of the exponential regression models with their
434 corresponding R^2 values pertaining to the wet or dry season are reported for each duration class of the rainy period in
435 Table 6. The exponential curves in the wet season (see plot 13a) are virtually parallel, yielding, for a fixed ET_p/R , more
436 WY/R as the duration of the rainy period increases from 3-4 months to 7-8 months. In contrast, the exponential
437 regression curves belonging to the dry season (see plot 13b) explain only a small amount of the variations of WY/R in
438 response to the dryness index and all seem quite insensitive to rainfall seasonality. Only the exponential model

439 pertaining to the dry season and for the smaller duration of the rainy period (3-4 months) explains slightly less than 50%
440 of the variability of ET_p/R for the study catchment.

441

442 **6. Conclusions**

443 Capturing the relationship between precipitation and catchment-scale water balance components in a Mediterranean
444 context is a scientific challenge in view of expected increasing frequencies in extremes such as droughts and floods
445 induced by climate warming. On the one hand, intense and prolonged droughts induce a steep decline in water
446 availability for irrigation (with a subsequent decrease in crop productivity), domestic use (especially for the tourist
447 sector), clean power generation, to mention just a few. On the other hand, projected increments in runoff and flooding
448 induce higher-than-normal risk of landslides and soil erosion, compromising the local economy and leading to
449 unprecedented hazards for a vulnerable population. Therefore, countries across the Mediterranean region are being
450 forced to pursue drastic adaptive options which in turn depend on modeling scenarios which can be performed by using
451 hydrological models. Indeed, scenarios need to rely on adequate rainfall modeling within the hydrological year by
452 generating multiple data sets of reliable daily rainfall time series drawn from statistical distributions derived from long-
453 term observations. Nonetheless, a key is first to define rainfall seasons, and then optimize parameters featuring in the
454 best statistical distribution describing rainfall data distribution in each season. If this exercise is well posed, one can
455 capture realistic rainfall dynamics occurring in the water balance simulated by a numerical model. Within this
456 framework, the aim of this study is to contribute in understanding the impact of rainfall seasonality and its anomalies on
457 the water balance components by providing reliable and robust scenario-based projections, based on the use of well-
458 posed hydrological models.

459 This study presented a pilot area (UARC in southern Italy) in the Mediterranean region. We applied the SWAT model
460 that was calibrated and validated in a previous paper using a large amount of environmental data and maps (Nasta et al,
461 2017). Moreover, the availability of a long-term time series of daily rainfall data (almost one century) allowed us to

462 detect rainfall seasonality by using a static and a dynamic approach. In both approaches we apply the SWAT model to
463 evaluate the sensitivity of hydrological water balance components to rainfall seasonality, using as input synthetic
464 rainfall time series generated by a Poisson process with two parameters that characterize daily rainfall occurrences and
465 daily rainfall depth in each season. In the static approach, dry or wet anomalies are considered when the transition
466 seasons turn into dry or wet seasons. The advantage of this approach lies in its simplicity and easy reproducibility in
467 other sites. However, it can be considered only an artifact based on criteria to group monthly rainfall amounts that might
468 be subjective. In the dynamic approach, the seasonal anomalies occur on the tails of the normal distribution of the wet
469 season duration. Although this approach seems statistically sound, the main disadvantage is the fact that it requires
470 long-term historical rainfall time-series of daily rainfall data that are unlikely to be available in most weather stations
471 across the Mediterranean region. In this study, both approaches concurred on understanding the impact of seasonal
472 rainfall anomalies on catchment-scale water balance components.

473 Our results show a drought anomaly (i.e. a prolonged duration of the dry season) in just one single year potentially leads
474 to a decrease of even about a fifth of the annual average rainfall and induces a drastic decline in average annual amounts
475 of water yield, actual evapotranspiration, and groundwater recharge. Conversely, an exceptional prolonged wet season
476 is likely to cause a considerable increase in annual average rainfall, hence about a one-third rise in annual average water
477 yield as well as enhanced groundwater recharge. In the dynamic approach, we demonstrated that the implicit
478 assumption of a temporal steady-state in the Budyko relation approach is sensitive to rainfall seasonality. The Budyko
479 evaporative index spans from 0.45 to 0.54 when the wet season lasts from 7-8 months up to 3-4 months. Moreover, it is
480 possible to identify distinct season-dependent regression equations linking seasonal water yield to the dryness index
481 over the wet season.

482 In conclusion this paper provides a framework to analyze the effects of rainfall seasonality changes on the hydrological
483 water budget and partition, while providing some preliminary results that can be representative for Mediterranean
484 catchments. Finer analyses can be performed by considering consecutive years of prolonged drought episodes and/or by

485 adding the effects of temperature trends, which obviously affect potential evapotranspiration forcing and in principle
 486 can produce a further feedback on precipitation cycles. These still unexplored issues will form the subject of future
 487 research investigation and forthcoming communications.

488 **7. Appendix**

489 We set k and m as counters for the hydrological year and the 12 months in each year, respectively. The annual rainfall,
 490 R_k , and associated monthly probability distribution, $p_{k,m}$, are defined as:

$$491 \quad R_k = \sum_{m=1}^{12} r_{k,m} \quad (\text{A1})$$

$$492 \quad p_{k,m} = \frac{r_{k,m}}{R_k} \quad (\text{A2})$$

493 where $r_{k,m}$ represents the rainfall depth recorded in the m -th month in the k -th year.

494 The relative entropy, D_k , is calculated in each hydrological year, k , as:

$$495 \quad D_k = \sum_{m=1}^{12} p_{k,m} \log_2 \left(\frac{p_{k,m}}{q_m} \right) \quad (\text{A3})$$

496 where q_m is equal to $1/12$ (uniform distribution). This statistical index quantifies the distribution of monthly rainfall
 497 within each hydrological year. Finally, the dimensionless seasonality index (DSI_k) in each hydrological year, k , is given
 498 by:

$$499 \quad DSI_k = D_k \frac{R_k}{R_{max}} \quad (\text{A4})$$

500 where \bar{R}_{max} is maximum \bar{R} . This way DSI_k is zero when rainfall is uniformly distributed throughout the year and
 501 reaches its maximum value $\log_2 12$ when rainfall is concentrated in a single month.

502 According to Feng et al. (2013), the magnitude (R_k) represents annual rainfall whereas the centroid (C_k) and the spread
 503 (Z_k) indicate timing and duration of the wet season, respectively, and are calculated in each hydrological year k as:

$$504 \quad C_k = \frac{1}{R_k} \sum_{m=1}^{12} m r_{k,m} \quad (A5)$$

$$505 \quad Z_k = \sqrt{\frac{1}{R_k} \sum_{m=1}^{12} |m - C_k|^2 r_{k,m}} \quad (A6)$$

506

507 **Acknowledgments**

508 The study reported in this paper was partially supported by the MiUR-PRIN Project “Innovative methods for water
 509 resources management under hydro-climatic uncertainty scenarios” (grant 2010JHF437). The Director of the Consorzio
 510 di Bonifica Velia, Marcello Nicodemo, is also acknowledged for his support in providing the datasets recorded at the
 511 Piano della Rocca earth dam. Roberto Deidda acknowledges the financial support received from the Sardinia Regional
 512 Authority under grant L.R. 7/2007, funding call 2017, CUP: F76C18000920002.

513

514 **References**

- 515 Abbott, B.W., Bishop, K.H., Zarnetske, J.P., Minaudo, C., Chapin, F.S., Krause S., Hannah, D.M., Conner, L., Ellison,
 516 D., Godsey, S.E., et al.: Human domination of the global water cycle absent from depictions and perceptions, *Nat.*
 517 *Geosci.*, doi: 10.1038/s41561-019-0374-y, 2019.
- 518 Adla, S., Tripathi, S., and Disse, M.: Can we calibrate a daily time-step hydrological model using monthly time-step
 519 discharge data?, *Water*, 11, 1750, doi:10.3390/w11091750, 2019.
- 520 Allen, R.G., Pereira, L.S., Raes, D., and Smith, M.: *Crop Evapotranspiration: Guidelines for Computing Crop Water*
 521 *Requirements.*, Food and Agriculture Organization of the United Nations, 1998.
- 522 Ayoade, J.O.: The seasonal incidence of rainfall, *Weather* 25, 414-418, [https://doi.org/10.1002/j.1477-](https://doi.org/10.1002/j.1477-8696.1970.tb04132)
 523 [8696.1970.tb04132](https://doi.org/10.1002/j.1477-8696.1970.tb04132), 1970.
- 524 Apurv, T., Sivapalan, M., and Cai, X.: Understanding the role of climate characteristics in drought propagation, *Water*
 525 *Resour. Res.*, 53, 9304–9329. <https://doi.org/10.1002/2017WR021445>, 2017.
- 526 Arnold, J.G., Srinivasan, R., Mutiah, R.S., and Williams, J.R.: Large area hydrologic modeling and assessment part I:
 527 model development, *J. Am. Soc. Water Resour. Assoc.*, 34 (1), 73–89, 1998.
- 528 Bari, S.H., Hussain, M.M., and Husna, N.E.A.: Rainfall variability and seasonality in northern Bangladesh, *Theor.*
 529 *Appl. Climatol.*, <https://doi.org/10.1007/s00704-016-1823-9>, 2016.

530 Budyko, M.I.: *Climate and Life.*, Academic Press, New York, 1974.

531 Caracciolo D., Deidda R., and Viola F.: Analytical estimation of annual runoff distribution in ungauged seasonally dry
532 basins based on a first order Taylor expansion of the Fu's equation, *Adv. Water Resour.*, 109, 320-332,
533 <https://doi.org/10.1016/j.advwatres.2017.09.019>, 2017.

534 Corona, R., Montaldo, N., and Albertson, J.D.: On the Role of NAO-Driven Interannual Variability in Rainfall
535 Seasonality on Water Resources and Hydrologic Design in a Typical Mediterranean Basin, *J. Hydrometeorol.*, 19,
536 485-498, doi: 10.1175/jhm-d-17-0078.1, 2018.

537 de Lavenne, A., and Andréassian, V.: Impact of climate seasonality on catchment yield: A parameterization for
538 commonly-used water balance formulas, *J. Hydrol.*, 558, 266-274, 2018.

539 Deidda, R.: An efficient rounding-off rule estimator: Application to daily rainfall time series, *Water Resour. Res.*, 43,
540 W12405, doi:10.1029/2006WR005409, 2007.

541 Deidda, R.: A multiple threshold method for fitting the generalized Pareto distribution to rainfall time series, *Hydrol.*
542 *Earth Syst. Sci.*, 14, 2559–2575, 2010.

543 Domínguez-Castro, F., Vicente-Serrano, S.M., Tomás-Burguera, M., Peña-Gallardo, M., Beguería, S., El Kenawy, A.,
544 Luna, Y., and Morata, A.: High-spatial-resolution probability maps of drought duration and magnitude across Spain,
545 *Nat. Hazards Earth Syst. Sci.*, 19, 611–628, 2019.

546 Eagleson, P. S.: Dynamics of flood frequency, *Water Resour. Res.*, 8, 878–898, 1972.

547 Feng, X., Porporato, A., and Rodriguez-Iturbe, I.: Changes in rainfall seasonality in the tropics, *Nat. Clim. Change*,
548 <https://doi.org/10.1038/nclimate1907>, 2013.

549 Feng, X., Vico, G., and Porporato, A.: On the effects of seasonality on soil water balance and plant growth, *Water*
550 *Resour. Res.*, 48, W05543, doi:10.1029/2011WR011263, 2012.

551 Hargreaves, G.L., Hargreaves, G.H., and Riley, J.P.: Irrigation water requirements for Senegal river basin, *J. Irrig.*
552 *Drain. Eng.*, 111, 265–275, 1985.

553 Hanel, M., Rakovec, O., Markonis, Y., Máca, P., Samaniego, L., Kyselý, J., and Kumar, R.: Revisiting the recent
554 European droughts from a long-term perspective, *Scientific Reports*, 8:9499, doi:10.1038/s41598-018-27464-4,
555 2018.

556 Hayes, M., Wilhite, D.A., Svoboda, M., and Vanyarkho, O.: Monitoring the 1996 drought using the Standardized
557 Precipitation Index, *Bull. Am. Meteorol. Soc.*, 80, 429-438, 1999.

558 Kendall, M.G.: *Rank Correlation Measures*, Charles Griffin: London, 1975.

559 Kutiel, H., and Trigo, R.M.: The rainfall regime in Lisbon in the last 150 years, *Theor. Appl. Climatol.*, doi
560 10.1007/s00704-013-1066-y, 2013.

561 IPCC, Climate change, 2013: the physical science basis. Contribution of working group I to the fifth assessment report
562 of the intergovernmental panel on climate change. Cambridge, United Kingdom and New York, USA: Cambridge
563 University Press.

564 Laaha, G., Gauster, T., Tallaksen, L. M., Vidal, J.-P., Stahl, K., Prudhomme, C., Heudorfer, B., Vlnas, R., Ionita, M.,
565 Van Lanen, H. A. J., Adler, M.-J., Caillouet, L., Delus, C., Fendekova, M., Gailliez, S., Hannaford, J., Kingston, D.,
566 Van Loon, A. F., Mediero, L., Osuch, M., Romanowicz, R., Sauquet, E., Stagge, J. H., and Wong, W. K.: The
567 European 2015 drought from a hydrological perspective, *Hydrol. Earth Syst. Sci.*, 21, 3001–
568 3024, <https://doi.org/10.5194/hess-21-3001-2017>, 2017.

569 Lilliefors, H.W.: On the Kolmogorov-Smirnov test for normality with mean and variance unknown, *J. Am. Stat. Assoc.*,
570 62, 399–402, 1967.

571 Livada, I., and Asimakopoulos, D.N.: Individual seasonality index of rainfall regimes in Greece, *Clim. Res.*, 28, 155–
572 161, 2005.

573 Mann, H.B.: Non-parametric tests against trend, *Econometrica*, 13, 245-259, 1945.

574 Markham, C.G.: Seasonality of precipitation in the United States, *Ann. Am. Assoc. Geogr.*, 60, 593–597,
575 <https://doi.org/10.1111/j.1467-8306.1970.tb00743.x>, 1970.

576 Mariotti, A., Zeng, N., Yoon, J.-H., Artale, V., Navarra, A., Alpert, P., and Li, L.: Mediterranean water cycle changes:
577 transition to drier 21st century conditions in observations and CMIP3 simulations, *Environ. Res. Lett.*, 3,
578 doi:10.1088/1748-9326/3/4/044001, 2008.

579 Martin-Vide, J.: Spatial distribution of a daily precipitation concentration index in Peninsular Spain, *Int. J. Climatol.*,
580 24, 959-971, 2004.

581 McKee, T.B., Doesken, N.J., and Kleist, J.: The relationship of drought frequency and duration to time scales. In
582 “Eighth conference on applied climatology”, pp. 17–22, Anaheim, California: American Meteorological Society,
583 1993.

584 Miller, S.N., Kepner, W.G., Mehaffey, M.H., Hernandez, M., Miller, R.C., Goodrich, D.C., Devonald, K.K., Heggem,
585 D.T., and Miller, W.P.: Integrating landscape assessment and hydrologic modeling for land cover change analysis, *J.*
586 *Am. Water Resour. Assoc.*, 38, 915-929, 2002.

587 Milly, P.C.D.: Climate, soil water storage, and the average annual water balance, *Water Resour. Res.*, 30, 2143-2156,
588 1994.

589 Nasta, P., Romano, N., and Chirico, G.B.: Functional evaluation of a simplified scaling method for assessing the spatial
590 variability of the soil hydraulic properties at hillslope scale, *Hydrol. Sci. J.*, 58, 1-13, 2013.

591 Nasta, P., Palladino, M., Ursino, N., Saracino, A., Sommella, A., and Romano, N.: Assessing long-term impact of land
592 use change on hydrologic ecosystem functions in a Mediterranean upland agro-forestry catchment, *Sci. Total*
593 *Environm.*, 605-606, 1070-1082, 2017.

594 Nasta, P., Sica, B., Mazzitelli, C., Di Fiore, P., Lazzaro, U., Palladino, M., and Romano, N.: How effective is
595 information on soil-landscape units for determining spatio-temporal variability of near-surface soil moisture?, *J.*
596 *Agric. Eng.*, 49(3), 174-182, doi:10.4081/jae.2018.822, 2018.

597 Nasta, P., Boaga, J., Deiana, R., Cassiani, G., and Romano, N.: Comparing ERT- and scaling-based approaches to
598 parameterize soil hydraulic properties for spatially distributed model applications, *Adv. Water Resour.*, 126, 155-
599 167, 2019.

600 Nieuwolt, S.: Seasonal rainfall distribution in Tanzania and its cartographic representation, *Erdkunde*, 28, 186–194,
601 1974.

602 Oliver, J.E.: Monthly precipitation distribution: A comparative index, *Prof. Geogr.*, 32, 300–309, 1980.

603 Hosking, J.R.M.: L-moments: Analysis and estimation of distributions using linear combinations of order statistics, *J.*
604 *Royal Stat. Soc., Series B (Methodological)*, 52, 105-124, 1990.

605 Pascale, S., Lucarini, V., Feng, X., Porporato, A., and Hasson, S.: Analysis of rainfall seasonality from observations and
606 climate models, *Clim. Dyn.*, 44, 3281–3301, 2015.

607 Pascale, S., Lucarini, V., Feng, X., Porporato, A., and Hasson, S.: Projected changes of rainfall seasonality and dry
608 spells in a high greenhouse gas emissions scenario, *Clim. Dyn.*, 46, 1331-1350, 2016.

609 Paz, S., and Kutiel, H.: Rainfall regime uncertainty (RRU) in an eastern Mediterranean region — a methodological
610 approach, *Isr. J. Earth Sci.*, 52: 47–63, 2003.

611 Piccarreta, M., Pasini, A., Capolongo, D., and Lazzari, M.: Changes in daily precipitation extremes in the
612 Mediterranean from 1951 to 2010: the Basilicata region, southern Italy, *Int. J. Climatol.*, 33, 3229–3248, 2013.

613 Potter, N.J., Zhang, L., Milly, P.C.D., McMahon, T.A., and Jakeman, A.J.: Effects of rainfall seasonality and soil
614 moisture capacity on mean annual water balance for Australian catchments, *Water Resour. Res.*, 41, W06007,
615 doi:10.1029/2004WR003697, 2005.

616 Pryor, S.C., and Schoof, J.T.: Changes in the seasonality of precipitation over the contiguous USA, *J. Geophys. Res.*,
617 113, D21108. <https://doi.org/10.1029/2008JD010251>, 2008.

618 Raziei, T.: An analysis of daily and monthly precipitation seasonality and regimes in Iran and the associated changes in
619 1951–2014, *Theor. Appl. Climatol.*, pp.134:913–934 <https://doi.org/10.1007/s00704-017-2317-0>, 2018.

620 Rivoire, P., Tramblay, Y., Neppel, L., Hertig, E., and Vicente-Serrano, S.M.: Impact of the dry-day definition on
621 Mediterranean extreme dry-spell analysis, *Nat. Hazards Earth Syst. Sci.*, 19, 1629–1638, 2019.

622 Rodríguez-Iturbe, I., Febres de Power, B., and Valdés, J.B.: Rectangular pulses point process models for rainfall:
623 Analysis of empirical data, *J. Geophys. Res.*, <https://doi.org/10.1029/JD092iD08p09645>, 1987.

624 Romano N., Nasta, P., Bogena, H.R., De Vita, P., Stellato, L., and Vereecken, H.: Monitoring hydrological processes
625 for land and water resources management in a Mediterranean ecosystem: the Alento River catchment observatory,
626 *Vadose Zone J.*, 17, 180042. doi:10.2136/vzj2018.03.0042, 2018.

627 Sahany, S., Mishra, S. K., Pathak, R., and Rajagopalan, B.: Spatiotemporal variability of seasonality of rainfall over
628 India, *Geophys. Res. Lett.*, 45, 7140-7147, 2018.

629 SCS, 1972. Hydrology. Section 4 in National Engineering Handbook. Washington, D.C.: USDA Soil Conservation
630 Service.

631 Sumner, G., Homar, V., and Ramis, C.: Precipitation seasonality in eastern and southern coastal Spain, *Int. J. Climatol.*,
632 21, 219–247, <https://doi.org/10.1002/joc.600>, 2001.

633 Troch, P.A, Carrillo, G., Sivapalan, M., Wagener, T., and Sawicz, K.: Climate-vegetation-soil interactions and long-
634 term hydrologic partitioning: signatures of catchment co-evolution, *Hydrol. Earth Syst. Sci.*, 17, 2209–2217, 2013.

635 Van Loon, A.F., Tjeldeman, E., Wanders, N., Van Lanen, H.A.J., Teuling, A.J., and Uijlenhoet, R.: How climate
636 seasonality modifies drought duration and deficit, *J. Geophys. Res. Atmos.*, 119, 4640–4656,
637 doi:10.1002/2013JD020383, 2014.

638 Veneziano, D., and Iacobellis, V.: Multiscaling pulse representation of temporal rainfall, *Water Resour. Res.*, 38, 1138,
639 10.1029/2001WR000522, 2002.

640 Viola, F., Caracciolo, D., Forestieri, A., Pumo, D., and Noto, L.: Annual runoff assess- ment in arid and semi-arid
641 Mediterranean watersheds under the Budyko's framework, *Hydrol. Process.*, 31 (10), 1876–1888,
642 <http://dx.doi.org/10.1002/hyp.11145>, 2017.

643 Vogel, R.M., and Fennessey, N.M.: L moment diagrams should replace product moment diagrams, *Water Resour.*
644 *Res.*, 29, 1745-1752, 1993.

645 Walsh, R.P.D., and Lawler, D.M.: Rainfall seasonality: description, spatial patterns and change through time, *Weather*
646 36, 201–208, <https://doi.org/10.1002/j.1477-8696.1981.tb05400.x>, 1981.

647 Williams, C. A., Reichstein, M., Buchmann, N., Baldocchi, D., Beer, C., Schwalm, C., Wohlfahrt, G., Hasler, N.,
648 Bernhofer, C., Foken, T., Papale, D., Schymansky, S., and Schaefer, K.: Climate and vegetation controls on the
649 surface water balance: Synthesis of evapotranspiration measured across a global network of flux towers, *Water*
650 *Resour. Res.*, 48, W06523, doi:10.1029/2011WR011586, 2012.

651 Zhang, L.J., and Qian, Y.F.: Annual distribution features of precipitation in China and their interannual variations, *Acta*
652 *Meteorol. Sin.*, 17, 146–163, 2003.

653

Figures




Figure 1:Geographical position of the Upper Alento River Catchment (UARC) in Campania (southern Italy) with the locations of the weather stations of Gioi Cilento and Monteforte Cilento. This figure was adapted from © Google Maps




Figure 2: Box plots of monthly rainfall depths recorded at the Gioi Cilento weather station (1920-2018).




Figure 3: Theoretical L-moment ratio of common distribution models, as compared to the sample L-moment ratios of daily rainfall time series at the Gioi Cilento weather station (large filled circle).




Figure 4: Exponential probability plot of empirical and fitted cumulative distribution functions of daily rainfall depths collected at the Gioi Cilento weather station.




Figure 5: Temporal evolution of SPI-6 spanning from 1920 to 2018 (rainfall data were recorded at the Gioi Cilento weather station).




Figure 6: a) Temporal evolution of SPI-1 values (gray bars) and their 12-month moving average (magenta line) spanning from 1920 to 2018 in the static approach; b) Box plots of SPI-1 values and frequency distribution in the c) rainy period (blue histograms corresponding to Nov-Dec-Jan-Feb), d) transition period (yellow histograms corresponding to Mar-Apr-Sep-Oct), e) dry period (red histograms corresponding to May-Jun-Jul-Aug).




Figure 7: Temporal evolution of the centroid (or timing; solid line) and spread (or duration; dashed lines) of the wet seasons estimated as proposed by Feng et al. (2013) within the framework of the dynamic approach (rainfall data were recorded at the Gioi Cilento weather station).




Figure 8: Temporal evolution of a) dimensionless seasonal index, DSI (Feng et al., 2013) represented by a red line with corresponding linear regression (dashed line); b) seasonality index, SI (Walsh and Lawler, 1981) represented by a green line with corresponding linear regression (dotted line).




Figure 9: Time series (a) and frequency distribution (b) of durations of the rainy periods at the Gioi Cilento weather station in the dynamic approach.




Figure 10: Rainfall and potential evapotranspiration forcings in the static approach, namely seasonal rainfall (R) and potential evapotranspiration (ET_p) in the dry (red bars), transition (orange bars), and wet season (blue bars). Three scenarios are presented: a) “reference scenario” with the dry, transition, and wet seasons all lasting 4 months; b) “dry scenario” with the dry and wet seasons lasting 8 and 4 months, respectively; c) “wet scenario” with the dry and wet seasons lasting 4 and 8 months, respectively.




Figure 11: Main water balance components in the static approach, namely seasonal water yield (WY) and actual evapotranspiration (ET_a) in the dry (red bars), transition (orange bars), and wet season (blue bars). Three scenarios are presented: a) “reference scenario” with the dry, transition, and wet seasons all lasting 4 months; b) “dry scenario” with the dry and wet seasons lasting 8 and 4 months, respectively; c) “wet scenario” with the dry and wet seasons lasting 4 and 8 months, respectively.




Figure 12: Budyko diagram relating the dryness index (ET_p/R) with the evaporative (ET_a/R) index classified according to the duration of the rainy period pertaining to the dynamic approach. Circles denote median and vertical colored lines represent the range between 5th and 95th percentiles of evaporative index (red, black, green, cyan and blue colors correspond to duration of the rainy period of 3-4, 4-5, 5-6, 6-7 and 7-8 months, respectively). Solid lines denote energy and water limits, the solid garnet line represents the Budyko curve (Budyko, 1974). The vertical dashed line separates left-hand side from right-hand side of the Budyko curve.




Figure 13: Relationship between dryness index and water yield to rainfall ratio (WY/R) on a seasonal basis and classified according to the duration of the wet season (from shortest to longest denoted by reddish and bluish colors in the color bar) pertaining to the dynamic approach for the wet season (plot 12a) and the dry season (plot 12b). The exponential regression equations are represented in both plots by the dashed black lines according to the duration of the rainy period.

Tables

Table 1: Descriptive statistics of the monthly and annual rainfall distributions recorded at the Gioi Cilento weather station during the period 1920-2018.

	<i>mean</i>	<i>median</i>	<i>min</i>	<i>max</i>	<i>Std. Dev.</i>	<i>CV</i>
	mm	mm	mm	mm	mm	%
Jan	145.6	141.65	0.0	461.2	81.6	56.0
Feb	128.1	120.25	0.8	350.1	76.3	59.6
Mar	112.9	101.1	0.0	302.6	73.4	65.0
Apr	102.5	101	16.2	350.6	59.5	58.0
May	75.2	67.6	1.1	304.8	56.6	75.2
Jun	52.8	45.3	0.0	190.9	38.2	72.3
Jul	29.8	17.6	0.0	140.4	32.8	110.0
Aug	39.7	30.3	0.0	210	42.8	107.7
Sep	94.4	81.9	1.6	296.8	63.0	66.7
Oct	126.8	118.8	0.0	335.5	70.3	55.4
Nov	166.9	152.2	26.0	613.2	94.9	56.9
Dec	154.6	134.55	0.8	411.8	85.1	55.1
Annual	1229.3	1198.3	478.6	2069.6	295.9	24.1

Table 2: Scenario set-up in the “static” approach. Duration and Poisson distribution parameters (η and λ) are reported for each of the considered scenarios.

	Dry season			Transition season			Wet season		
	months	η	λ	months	η	λ	months	η	λ
	-	mm	d ⁻¹	-	mm	d ⁻¹	-	mm	d ⁻¹
Reference scenario (static)	4	8.20	0.196	4	10.53	0.34	4	11.70	0.423
Dry scenario (static)	8	8.20	0.196	0	-	-	4	11.70	0.423
Wet scenario (static)	4	8.20	0.196	0	-	-	8	11.70	0.423

Table 3: Scenario set up in the “dynamic” approach. Duration and Poisson distribution parameters (η and λ) are reported in the dry and wet season.

Dynamic scenario	Dry season			Wet season		
	months	η	λ	months	η	λ
	-	mm	d ⁻¹	-	mm	d ⁻¹
	random	9.34	0.243	random	11.99	0.413

Table 4: Descriptive statistics of annual water balance components obtained in the three scenarios in the “static” approach. Units are mm, except for CV (%).

Scenario	Variable	R	WY	ET_a	GR
		mm	mm	mm	mm
Reference scenario	mean	1229.0	433.3	605.2	194.3
	stand. dev.	176.0	104.2	36.5	48.0
	CV (%)	14.3	24.1	6.0	24.7
	min	586.6	150.8	449.1	44.0
	max	2053.9	1005.9	743.0	389.6
Dry scenario	mean	987.7	317.3	545.1	128.0
	stand. dev.	155.5	88.1	40.8	42.7
	CV (%)	15.7	27.8	7.5	33.4
	min	498.7	96.2	396.0	7.2
	max	1649.9	802.4	691.6	319.3
Wet scenario	mean	1392.8	526.0	625.8	248.1
	stand. dev.	192.4	119.6	34.3	52.6
	CV (%)	13.8	22.7	5.5	21.2
	min	721.9	157.0	481.2	59.0
	max	2179.2	1088.2	748.6	461.6

Table 5: Water balance components associated to occurrence probabilities for each duration of the rainy period.

	Probability	<i>R</i>	<i>WY</i>	<i>ET_a</i>	<i>GR</i>
	%	mm	mm	mm	mm
3-4 months	0.6%	1,145.0	385.3	608.5	169.6
4-5 months	23%	1,213.4	420.0	619.4	188.0
5-6 months	62%	1,275.4	453.0	624.9	199.6
6-7 months	14%	1,326.0	480.2	631.6	210.2
7-8 months	0.3%	1,383.5	511.6	644.2	211.8

Table 6: Exponential regression models, with the corresponding coefficient of determination (*R*²), for the wet and dry seasons as a function of the duration of the rainy period.

Duration	Wet season		Dry season	
	Exp regression function	<i>R</i> ²	Exp regression function	<i>R</i> ²
3-4 months	$WY/R = 0.5914 \times \exp(-1.674 \times ET_p/R)$	0.440	$WY/R = 0.4635 \times \exp(-0.343 \times ET_p/R)$	0.482
4-5 months	$WY/R = 0.6031 \times \exp(-1.536 \times ET_p/R)$	0.579	$WY/R = 0.3675 \times \exp(-0.204 \times ET_p/R)$	0.290
5-6 months	$WY/R = 0.6171 \times \exp(-1.477 \times ET_p/R)$	0.587	$WY/R = 0.3530 \times \exp(-0.174 \times ET_p/R)$	0.279
6-7 months	$WY/R = 0.6313 \times \exp(-1.399 \times ET_p/R)$	0.617	$WY/R = 0.3476 \times \exp(-0.159 \times ET_p/R)$	0.284
7-8 months	$WY/R = 0.6586 \times \exp(-1.389 \times ET_p/R)$	0.585	$WY/R = 0.3137 \times \exp(-0.105 \times ET_p/R)$	0.211

654

655