

Interactive comment on “X Water Worlds and how to investigate them: A review and future perspective on *in situ* measurements of water stable isotopes in soils and plants” by Matthias Beyer and Maren Dubbert

Erik Oerter

erikjoerter@gmail.com

Received and published: 18 December 2019

This is a very nice review of the current state of the art for *in situ* soil and plant water stable isotope analysis techniques, and I applaud the authors for this timely contribution. The focus of the review is on laser-based techniques, and this is the route that most researchers currently take. However, the use of soil gas measurements taken via soil gas wells or probes, specifically CO₂, as a way to measure soil water oxygen stable isotopic composition, has not been included.

C1

The basic principle is that in soils, the molar abundance of oxygen in soil water is orders of magnitude greater than the molar abundance of oxygen in coexisting soil CO₂, so that the CO₂ comes into oxygen isotope equilibrium with that of the soil water. Aliquots of soil gas are sampled via small diameter wells and stored in septum-capped vials for transport back to the lab. In the lab, the $\delta^{18}\text{O}$ of the CO₂ is measured via continuous flow or dual inlet IRMS.

I suggest the inclusion of this technique in the review manuscript because it enables researchers without access to laser instruments, but with access to conventional IRMS instruments to be able to make *in situ* soil water O isotope measurements. Below, I include some notes on potential references that the authors may wish to include in their review.

Stern and colleagues laid the foundation for the O isotope relationship between soil CO₂ and soil water:

Stern, L., Baisden, W. T., & Amundson, R. (1999). Processes controlling the oxygen isotope ratio of soil CO₂: Analytic and numerical modeling. *Geochimica et Cosmochimica Acta*, 63(6), 799-814.

Breecker and Sharp developed a technique for the sampling of *in situ* soil CO₂ and its isotopic analysis: Breecker, D., & Sharp, Z. D. (2008). A field and laboratory method for monitoring the concentration and isotopic composition of soil CO₂. *Rapid Communications in Mass Spectrometry*, 22(4), 449-454.

Breecker and colleagues deployed their *in situ* soil CO₂ system and successfully measured soil water isotope profiles at high depth resolution (~10cm depth increments) at monthly intervals for more than a year at several sites:

Breecker, D. O., Sharp, Z. D., & McFadden, L. D. (2009). Seasonal bias in the formation and stable isotopic composition of pedogenic carbonate in modern soils from central New Mexico, USA. *Geological Society of America Bulletin*, 121(3-4), 630-640.

C2

Oerter and Amundson measured soil water O isotope composition via soil CO₂ at four sites for over a year. They were able to develop estimates of the depth of plant water uptake via comparison with xylem water samples, thus establishing the ecohydrologic applications of the soil CO₂-water technique:

Oerter, E. J., & Amundson, R. (2016). Climate controls on spatial and temporal variations in the formation of pedogenic carbonate in the western Great Basin of North America. *GSA Bulletin*, 128(7-8), 1095-1104.

An added value of the use of soil CO₂ sampling to measure soil water O isotope composition, is that the soil gas samples can also be measured for CO₂, CH₄, NO_x, as well as other trace gases abundance. The ecohydrologic benefits and implications of these types of measurements were demonstrated in a later paper by Oerter and colleagues from the same field sites as the 2016 paper above:

Oerter, E., Mills, J. V., Maurer, G. E., Lammers, L. N., & Amundson, R. (2018). Greenhouse Gas Production and Transport in Desert Soils of the Southwestern United States. *Global Biogeochemical Cycles*, 32(11), 1703-1717.

Interactive comment on Hydrol. Earth Syst. Sci. Discuss., <https://doi.org/10.5194/hess-2019-600>, 2019.