

# ***Interactive comment on “The Influence of a Prolonged Meteorological Drought on the Catchment Water Storage Capacity: A Hydrological Model Perspective” by Zhengke Pan et al.***

**Zhengke Pan et al.**

liupan@whu.edu.cn

Received and published: 27 June 2020

Responses to Referee #1: This paper represents an interesting huge work, the main objective of this paper is to detect the possible changes in catchment water storage capacity induced by a prolonged meteorological drought. It will also help improve our ability to simulate the hydrological system. Meanwhile, the structure is clear and analyses and conclusions in this paper are reasonable and logical. Overall, it is a nice paper with proper methodology and complete discussion on the main findings. A minor modification should be considered before publication.

[Printer-friendly version](#)

[Discussion paper](#)



Reply: We are grateful for the reviewer's appreciation of our work and for the professional comments, which are carefully followed in making revision.

General comments: A1: First, the results indicated that 12 catchments had a significant downward shift in the CWSC. My first principal doubts come from there: are 12 sufficient to be able to analyze regularity about catchments with a significant downward shift. Reply: Thanks. We will make it clearer in the revised version that to ensure the accuracy of the modeling results, strict criteria were adopted to evaluate the model results and to identify the catchments with a significant change in  $\theta_1$  including the minimum NSE requirement, the minimum requirement of significant change, the requirement for maximum performance degradation, and the requirement for the robustness of results (see section 3.2.5). Only these catchments that satisfied all these criteria were identified as catchments undergo significant changes. Actually, only 12 of the 145 catchments, which had experienced a long-term meteorological drought, satisfied all these four criteria and have been identified with a significant downward shift in the CWSC. Thus, the identified 12 catchments that had a significant downward shift should be reasonable results. We will also clarify that this result depends on the criteria used for evaluation. All catchments from southeastern Australia in the dataset have been tested in this study using the above defined criteria, and we did not find more catchments with a significant downward shift in the CWSC. More explanation will be provided in the revision.

A2: Second: Formulating conclusions is similar to drawing up an abstract. I mean that the emphases of the abstract and conclusions are different. Reply: Thank you for your comments, following which, the Abstract and conclusion will be rewritten. The part of Abstract will be modified as follows: Understanding the propagation of prolonged meteorological droughts helps solve the problem of intensified water scarcity around the world. Most of the existing literature studied the propagation of drought from one type to another (e.g., from meteorological to hydrological drought) with statistical approaches, there remains a difficulty in revealing the causality between the meteorological drought

[Printer-friendly version](#)

[Discussion paper](#)



and potential changes in the Catchment Water Storage Capacity (CWSC). This study aims to identify the response of the CWSC to the meteorological drought by examining the changes of hydrological model parameters after the drought events. Firstly, the temporal variation of a model parameter that denotes the CWSC is estimated to reflect the potential changes in real CWSC. Next, the change points of the CWSC parameter were determined based on the Bayesian change point analysis. Finally, the possible association and linkage between the shift in the CWSC and the time-lag of the catchment (i.e., time-lag between the onset of the drought and the change point) with multiple catchment properties and climate characteristics were identified. In total 145 catchments from southeastern Australia were selected as the study areas. Results indicated that (1) significant shifts in the CWSC can be observed in 62.7% of the catchments, which can be divided into two subsets with the opposite response, i.e. 48.2% of catchments had lower runoff generation rates while 14.5% of catchments had higher runoff generation rate; (2) the increase in the CWSC during a chronic drought can be observed in smaller catchments with lower elevation, slope, and forest coverage of Evergreen Broadleaf Forest, while the decrease in the CWSC can be observed in larger catchments with higher elevation and larger coverage of the Evergreen Broadleaf Forest; (3) catchments with a lower proportion of Evergreen Broadleaf Forest usually have longer time-lag and are more resilient. This study improves our understanding of possible changes in the CWSC induced by a prolonged meteorological drought, which will help improve our ability to simulate the hydrological system under climate change. The Conclusion will be modified as follows: This study aims to examine the possible changes in the CWSC as well as the time-lag between the onset of the meteorological drought and the change point of the CWSC. A classical hydrological model, GR4J, was used and its parameter  $\theta_1$  was selected to denote catchment water storage capacity (CWSC). Thus, the temporal variation in parameter  $\theta_1$  was detected to reveal the possible fluctuation in the CWSC, and the causality between the temporal variation in parameter  $\theta_1$  and a persistent meteorological drought was examined. One hundred forty-five catchments in southeastern Australia were selected as the study ar-

eas. Main conclusions can be drawn as follows: (1) Significant changes in the CWSC have been identified in 62.7% (52 in 83) of catchments, which can be divided into two subsets with opposite catchment responses: 48.2% (40 in 83) experienced a significant decrease in the CWSC during the drought period and had lower runoff generation rates, while 14.5% (12 in 83) of catchments experienced a significant decrease in the CWSC during the drought period, and had higher runoff generation rate. (2) Different change directions in the CWSC resulted in the opposite impacts on runoff generation, i.e., catchments with increased CWSC would result in lower runoff generation rates for similar amounts of rainfall than before while those catchments with decreased CWSC would have an opposite response (higher runoff generation rate). Generally, the increase in the CWSC during a chronic drought can be observed in smaller catchments with lower elevation, slope, and forest coverage of Evergreen Broadleaf Forest, while the decrease in the CWSC can be observed in larger catchments with higher elevation and larger coverage of the Evergreen Broadleaf Forest. Among all catchment properties and climate variables considered, our results suggest that two climate variables (i.e., variation in annual rainfall and annual runoff ratio) have the strongest associations with the shift in the CWSC. (3) The responses of different catchments to persistent meteorological drought were not equally susceptible. Catchments with a lower proportion of Evergreen Broadleaf Forest usually have longer time-lag and are more resilient. It is noted that although this study resulted in interesting findings that give new insight and have not been fully outlined before, it is based on the lumped GR4J model and the specific case in Australia, which implies that the main findings/conclusions may not directly extendable to other regions. Thus, to examine the generality of the main conclusions, the response of CWSC to the meteorological drought can be analyzed with the other hydrological models in the other regions.

Detailed comments: A3: Is there a mistake in the inconsistency between the Abstract (L37-L38) and Conclusion (L613-615). Reply: We are sorry for this error. The sentence in the Abstract and Conclusion will be modified as "...the increase in the CWSC during a chronic drought can be observed in smaller catchments with lower elevation, slope,

[Printer-friendly version](#)

[Discussion paper](#)



and forest coverage of Evergreen Broadleaf Forest, while the decrease in the CWSC can be observed in larger catchments with higher elevation and larger coverage of the Evergreen Broadleaf Forest.”.

A4: Previous study (Yan et al., 2015) has indicated that the sensitivity of the parameters in the GR4J model has different performance in different parts of the same basin. Therefore, you should cite more citations or do some researches in the study area to support the sentence in the L224-L226. Reply: Thank you. More citations will be added to the revised manuscript. This sentence will be modified as “Previous studies (Demirel et al., 2013; Pan et al., 2019a; Pan et al., 2019b; Perrin et al., 2003; Westra et al., 2014; Yan et al., 2015) showed that  $\theta_1$ , which denotes the catchment water storage capacity, is the most sensitive parameter in the structure of the GR4J model.”. Added reference: Yan, X. L., Zhang, J. Y., Wang, G. Q., Bao, Z. X., Liu, C. S., and Xuan, Y. Q.: Application of GR4J Rainfall-runoff Model to Typical Catchments in the Yellow River Basin, Proceedings of the 5th International Yellow River Forum on Ensuring Water Right of the River’s Demand and Healthy River Basin Maintenance, Vol V, edited by: Hongqi, S., and Xiangxin, L., Yellow River Conservancy Press, Zhengzhou, 191-198 pp., 2015. Demirel, M. C., Booij, M. J., and Hoekstra, A. Y.: Effect of different uncertainty sources on the skill of 10 day ensemble low flow forecasts for two hydrological models, Water Resour. Res., 49, 4035-4053, 10.1002/wrcr.20294, 2013.

A5: L503-517: You should cut out much of the repetition with the Section 4. Reply: Thanks. Changes will be made as suggested. The paragraph in lines 503-517 will be modified as follows: The results indicate that, under certain circumstances, a long-term meteorological drought would result in a significant change in the CWSC. However, no strong association has been found between the magnitude of the change in the CWSC with any single variable. In addition, the length of dry period was not associated with the shift in the CWSC. Thus, it seems that the catchment response behavior to long-term meteorological drought is controlled by the combination of local catchment properties and climate characteristics rather than a single factor. Thus, fur-

ther studies are still required to confirm which factors played the most important role in the catchment dynamic. A6: In the Section 4.4 You had better discuss the factors for the direction first and then the magnitude of shifts in the CWSC. Reply: Thanks. We will explain better that Section 4.4.1 presents the results that classify the catchments into two groups, one group shows significant shifts in the CWSC, and another does not. While section 4.4.2 continued to discuss the difference between two sub-groups of catchments with significant upward/significant downward shifts in the CWSC. These two sub-groups of catchments were extracted from the group of catchments with significant shifts in the CWSC according to the change direction of the estimated  $\theta_1$ . Thus, we thought it might be better to discuss the difference between two groups of catchments with/without significant shifts at first, and then illustrate the difference between two subsets of catchments with significant upward/significant downward shifts later. To make it clearer, after considering reviewer's advice and reminder, the headings of sections 4.4.1 and 4.4.2 will be modified as "4.4.1 Factors for the significant/non-significant shifts in the CWSC" and "4.4.2 Factors for the significant upward/downward shifts in the CWSC" in the revised manuscript.

A7: Why not analyze the reasons for conclusion 2 in the Section 5.1? (Catchments with larger elevation and slope, lower forest coverage of Evergreen Broad leaf Forest are more likely to have an increase in the CWSC during a chronic drought while smaller catchments with lower elevation, lower coverage of the Evergreen Broad leaf Forest are more likely to have a decrease in the CWSC.) Reply: Thanks. We tried but as shown in Figures 7 and 9, no strong PCC association has been found between the magnitude of the change in the CWSC and the single catchment property or climate variable. It seems that the catchment response behavior to long-term meteorological drought is controlled by the combination of local catchment properties and climate characteristics rather than a single factor. Thus, we prospect that further studies are still required to confirm which factors played the most important role in the catchment dynamic.

A8: The conclusion should be concise and coherent. It is suggested that only the key

[Printer-friendly version](#)

[Discussion paper](#)



and definitive conclusions of this paper should be stated. If you cannot confirm your conclusion, these results may only be used in the Discussion, such as L610-611 and L622-631, etc. Reply: Thanks. Changes will be made as suggested. The sentences in lines 610-611 will be deleted in the conclusion part of the revised manuscript. In addition, the sentences in lines 622-631 will be modified as follows: It is noted that although this study resulted in interesting findings that give new insight and have not been fully outlined before, it is based on the lumped GR4J model and the specific case in Australia, which implies that the main findings/conclusions may not directly extendable to other regions. Thus, to examine the generality of the main conclusions, the response of CWSC to the meteorological drought can be analyzed with the other hydrological models in the other regions.

A9: It's better to add latitude and longitude to the location map of the study catchments (Figure 1). Reply: Thanks. Changes will be made in the revised Figure 1.

A10: Table 3 should add the results about the fourth criteria. Reply: Thanks. Changes will be made in the revised Table 3. The modified Table 3 is presented in the Supplement.pdf.

A11: Figure 4 can be improved. The figure should show the change directions and magnitudes of the CWSC at the same time. It is better for readers to understand the shifts of the CWSC in the adjacent catchments. Reply: Thanks. Changes will be made as suggested in The modified Figure 4 in Figure 2 in the supplement.

A12: Is the supplementary material missing? Reply: We are sorry for this mistake. We have uploaded the supplementary material (Table S1) to the system. Please also refer to the supplement.pdf (Table S1).

Please also note the supplement to this comment:

<https://www.hydrol-earth-syst-sci-discuss.net/hess-2019-692/hess-2019-692-AC1-supplement.zip>

Interactive comment on Hydrol. Earth Syst. Sci. Discuss., <https://doi.org/10.5194/hess-2019-692>, 2020.

**HESSD**


---

Interactive  
comment

[Printer-friendly version](#)

[Discussion paper](#)






**Fig. 1.** Figure 1. Location map of the study catchments in southeastern Australia.

[Printer-friendly version](#)

[Discussion paper](#)





**Fig. 2.** Figure 4. Location map of the catchments with the significant and non-significant shifts in the CWSC.

[Printer-friendly version](#)

[Discussion paper](#)

