

Groundwater fauna in an urban area: natural or affected?

Fabien Koch¹, Kathrin Menberg¹, Svenja Schweikert¹, Cornelia Spengler², Hans Jürgen Hahn², Philipp Blum¹

¹Institute of Applied Geosciences (AGW), Karlsruhe Institute of Technology (KIT), Kaiserstraße 12, 76131 Karlsruhe, Germany

²Faculty of Nature and Environmental Sciences (Working Group: Groundwater Ecology), University Koblenz-Landau, Im Fort 7, 76829 Landau, Germany

Correspondence to: Fabien Koch (fabien.koch@kit.edu)

Abstract. In Germany 70 % of the drinking water demand is met by groundwater, whose quality is the product of ~~manifold~~ multiple physical-chemical and biological ~~cleaning~~ processes. As healthy groundwater ecosystems help to provide clean drinking water, it is necessary to assess the ir ecological conditions ~~of these ecosystems~~. This is particularly true for densely populated, urban areas, where faunistic groundwater investigations are still ~~seare~~ scarce. The aim of this study is therefore to provide a first-tier assessment of the groundwater fauna in an urban area. Thus, we assess the ecological condition of an anthropogenically influenced aquifer by analysing the groundwater fauna in 39 groundwater monitoring wells in Karlsruhe (Germany) and a nearby forest ~~land~~. For classification, we apply the scheme ~~from the Federal Environmental Agency (UBA) of~~ Griebler et al. (2014), in which a threshold of more than 70 % of Crustaceans and of less than 20 % of Oligochaetes serves as an indication for good ecological conditions. In our study it is revealed that only 35 % of the wells in the urban area, and 50 % of wells in the forest fulfil these criteria, ~~and even in the pristine forest land only 50 % of the wells indicate fine ecological conditions~~. While the assessment ~~reveals~~ shows that ecological conditions in the studied urban area are ~~predominantly~~ not in a good ecological state, there is no clear spatial pattern with respect to land use and other anthropogenic impacts, in particular, groundwater temperature and nitrate concentrations. However, there are noticeable differences in the spatial distribution of species and abiotic groundwater characteristics between wells in the forest ~~land~~ and the urban area, which indicates that more comprehensive assessment methods are required to fully ~~capture~~ understand the different effects on groundwater fauna.

In Germany 70 % of the drinking water demand is met by groundwater, whose quality is the product of multiple physical-chemical and biological ~~cleaning~~-processes (Avramov et al., 2010). Groundwater ecosystems are responsible for several services that help to provide clean drinking water, which is a vital resource for humanity (Griebler and Avramov, 2015). Bacteria and ~~also~~-fauna also play an important role in the biological self-purification of groundwater by the retention of organic matter, natural attenuation of pollutants, storing and buffering of nutrients as well as the elimination of pathogens. Organic matter and pollutants can be degraded and converted to ~~valuable~~-biomass or ~~tied-bound~~ by microbial activity. Protozoa and higher ~~animals-organisms~~ can graze resulting biofilms, loosen the substrate and therefore stimulate ~~the~~-biological self-purification (Hancock et al., 2005; Avramov et al., 2010)(~~Avramov et al., 2010~~).

~~Yet, only healthy~~Healthy groundwater ecosystems can provide clean drinking water.~~Groundwater ecosystems however, they~~ are sensitive to external influences, such as chemical and thermal disturbances. The latter drives hydro-geochemical and biological processes in groundwater systems, which are ~~relatively-typically~~ isothermal (Briellmann et al., 2009; 20011). Groundwater fauna mainly consists of stygobion~~nte~~ species, which spend their entire life in groundwater and are adjusted to this habitat (Hahn, 2006). Hence, in Central Europe they are assumed to be cold stenotherm, which means that they prefer cold temperatures. A variability in temperature tolerance among groundwater faunal groups and species is reported in various studies, which explains why the use of individual temperature thresholds is more useful to capture different preferences. According to Spengler (2017) faunal diversity is generally declining at a temperature above 14 °C. Various authors reported species specific temperature preferences between 8 and 16 °C (for individuals of the species *Niphargus inopinatus* and *Proasellus cavaticus* (Briellmann et al., 2009, 2011)) and a specific temperature threshold of up to 19 °C (for *Parastenocaris phyllura* (Glatzel, 1990)). Above these thresholds the mortality of individuals raises until groundwater fauna is almost absent, for example at 22 °C in the study of Foulquier et al. (2011). However, temperature sensitivity is not only an issue at species level but also for the communities as a whole. Spengler (2017) reported 12 °C to be a temperature threshold value indicated by a shift in community structure for faunal communities of groundwater of the Upper Rhine Valley. and can hardly persist water temperatures over 16 °C (Briellmann et al., 2009) or rather 14 °C (Spengler, 2017) for a longer time period.

Nevertheless, in German and European legislation, as in many countries globally, groundwater is not yet recognized as a ~~protected-habitat which is worthy of protection in the German and European legislation, as in many countries globally,~~ and there is no common ~~ground-understanding~~ on the best practice of assessing ~~the groundwater-ecologyecological status of groundwater~~ (Hahn et al., 2018; Spengler and Hahn, 2018). The assessment of surface water is typically based on biological, physical-chemical and supported by hydro-morphological criteria (European Water Framework Directive and German legislation article 5 of the ‘Regulation on the Protection of Surface Water’). While groundwater quality is mostly assessed by physical-chemical and quantitative criteria, very few quantifiable ecological criteria are available for the assessment of ~~ecosystem-the~~ health of groundwater ecosystems. The availability of ecological criteria can only be increased by conducting a large number of studies dealing with the analyses of groundwater ecosystem health by investigating groundwater fauna.

Results from previous faunistic groundwater analyses are contained in a Germany-wide data record (~~data record by~~ Hahn, 2005; Berkhoff, 2010; Stein et al., 2012; Gutjahr, 2013; Spengler, 2017; Spengler and Hahn, 2018). ~~A closer look at the south-western part of Germany, the German federal state Baden-Württemberg, is given~~ The study by Hahn and Fuchs (2009). ~~This large-scaled study~~ focuses on defining stygoregions, ~~which extend over several square kilometres, and are~~ based on different hydrogeological units located in Baden-Württemberg, Germany. They conclude that the observed patterns of groundwater communities reflect a high spatial and temporal heterogeneity of ~~groundwater, with respect to hydrogeological~~ aquifer types ~~(with respect to~~ habitat structure, food ~~and~~ oxygen supply ~~etc.~~ Accordingly, stygobiotic biodiversity is likely to be underestimated ~~Although there are various studies on this topic (e.g. Gibert and Deharveng, 2002; Malard et al., 2002; Deharveng et al., 2009; Dole-Olivier et al., 2009b)-stygobiotic biodiversity is still likely to be underestimated at present.~~

Regional investigations on the spatial variation of groundwater fauna, i.e. stygobiont occurrences, and corresponding environmental parameters, such as geological site characteristics and altitude, are rare (Dole-Olivier et al., 2009; Gibert et al., 2009). An approach to elucidate groundwater biodiversity patterns in six European ~~and seven North American~~ regions was conducted in the PASCALIS project (Protocol for the ~~Assessment and Conservation of Aquatic Life In the Subsurface~~) (Gibert et al., 2009), which aimed at mapping biodiversity and endemism patterns (Deharveng et al., 2009) ~~and~~ The study shows that regional processes, such as hydrological connectivity, ~~in~~ a specific habitat (e.g. river floodplains as in Ward and Tockner, 2001) have a much stronger influence on species composition than local habitat features such as permeability and saturation. Within a region, hydrogeology, altitude, palaeographical factors and human activities can interact in complex ways to produce dissimilar patterns of species compositions and diversity (Gibert et al., 2009). ~~Unfortunately, the~~ The PASCALIS sampling protocol recommends selecting hydro-geographic basins that are not strongly affected by human activities such as groundwater pollutions (Malard et al., 2002), and does not biogeographically classify a groundwater system (Stein et al., 2012). In urban areas, ~~anthropogenic impacts,~~ such as a dense building development, underground car parks, open geothermal systems and injections of thermal wastewater from industry result in local thermal alteration of groundwater up to several degrees (e.g. Taylor and Stefan, 2009; Zhu et al., 2011; Menberg et al., 2013b; Tissen et al., 2019). According to Brielmann et al. (2011) annual temperature fluctuations in aquifers, ~~caused by shallow geothermal energy systems, range between 4 °C in winter and ≤20 °C in summer.~~ In 2000, the European Union (EU) (Water Framework Directive) defined the release of heat in the groundwater as a pollution, whereas the cooling of the groundwater is not ~~particularly~~ mentioned. Until now, there are no scientifically derived threshold values for groundwater temperature in the case of thermal (heat) pollution (Hähnlein et al., 2010; 2013). This results in a tension between conservation, exploitation and thermal use of groundwater. ~~Yet, However, as seen~~ in an aquifer ecosystem downstream of from an industrial facility in Freising (Germany), where groundwater is used for cooling resulting in a warm thermal plume, no relation between faunal abundance and groundwater temperature could be identified (Brielmann et al., 2009). Investigation of hydro-geochemical parameters, microbial activities, bacterial communities and groundwater faunal assemblages indicates that bacterial diversity ~~clearly increaseds~~ with temperature, while faunal diversity ~~usually decreasesdecreased~~ with temperature (Brielmann et al., 2009). Similar results are provided by Griebler et al. (2016), where potential impacts of geothermal energy use and storage of heat on groundwater are investigated. Temperature

changes in groundwater correspond with changes in groundwater chemistry, biodiversity, community composition, microbial processes and function of the ecosystem. How exactly these groundwater communities react to changes in temperature and concentration of nutrients, dissolved organic carbon and oxygen, is not yet fully understood (Briellmann et al., 2009, 2011; Spengler, 2017; Sánchez et al., 2020).

Several approaches exist that allow a local assessment of the ecological state of groundwater based on different faunistic, hydro-chemical and physical parameters. ~~The Commissioned by the~~ Federal Environmental Agency of Germany (Umweltbundesamt, UBA), Griebler et al. (2014), ~~for example,~~ developed a concept for an ecologically based assessment scheme for groundwater ecosystems. This two-step scheme characterizes groundwater on two different levels by using the most important physico-chemical parameters, such as content of dissolved oxygen, as well as microbiological and faunistic characteristics such as amount of Oligochaetes and Crustaceans, and comparing these to reference values for natural, undisturbed and ecologically intact groundwater ecosystems (Griebler et al., 2014). Moreover, Korbel and Hose (2017) introduced the Groundwater Health Index (GHI), which is a tiered framework for assessing the health of groundwater ecosystems. Here, both ~~The GHI uses~~ biotic and abiotic attributes of groundwater ecosystems are used to set ~~as~~ benchmarks, ~~which provide an indication of~~ ecosystem health. ~~In fact,~~ Their study shows that ecosystem health benchmarks are probably more associated ~~with aquifer typology,~~ than being applicable for local areas. This index is applied and tested by di Lorenzo et al. (2020) in unconsolidated aquifers in Italy located in nitrate vulnerable zones. They refined the index (wGHI^N) and demonstrated its applicability on shallow and deep aquifers and also revealed that this new index is limited due to low correlations between the indicators. ~~The common ground of both studies is the assessment of the ecological condition relative to a reference aquifer and the aim of classifying the locations (GHI: impacted or non impacted groundwater).~~

Furthermore, the Groundwater-Fauna-Index (GFI), introduced by Hahn (2006), quantifies the ~~ecological~~-relevant ecological conditions in the groundwater as a result of hydrological exchange between surface and groundwater. It incorporates ecologically important groundwater parameters such as relative amount of detritus, variation of groundwater temperature and concentration of dissolved oxygen (Hahn, 2006). Gutjahr et al. (2014) used the GFI as part of a proposal for a groundwater habitat classification ~~at on a~~ local scale, which introduce five types of faunistic habitats as a result of surface water influence, content of dissolved oxygen and amount of organic matter. Moreover, in the study of Berkhoff (2010) the GFI was used to examine the impact of the surface water influence on groundwater with the aim to develop a faunistic monitoring concept for hydrological exchange processes in the surrounding ~~of waterside river bank~~ filtration plants. Spengler and Hahn (2018) argued for the definition of a regional and ecological temperature threshold and an ecology based assessment of thermal stress in groundwater.

The objective of this study is to investigate specifically the groundwater fauna ~~under beneath~~ an urban area in comparison to a natural forest ~~and to determine whether land use has an impact on groundwater organism communities.~~ Hence, in 39 groundwater monitoring wells in Karlsruhe, Germany, the groundwater fauna is sampled, groundwater temperatures measured and as well as thermal and chemical properties are ~~sampled analysed in 39 groundwater monitoring wells in Karlsruhe, Germany.~~ In our study the classification scheme ~~by the Federal Environmental Agency of Germany (UBA) developed by~~

Griebler et al. (2014) is applied. The wells are characterized regarding the state of ~~their~~ the state of their ecosystem-quality. Hence, we finally aim to distinguish areas with natural state of groundwater ecology from anthropogenically disturbed areas.

2. Material and methods

2.1 Study site

130 The study is performed in Karlsruhe, a city in the Upper Rhine Valley in south-western Germany. The urban region covers an area of 173 km² and has about 310,000 inhabitants (Amt für Stadtentwicklung - Statistikstelle, 2018). The Cenozoic continental rift valley is filled with Tertiary and Quaternary sediments, which are dominated by sands and gravels with minor contents of silt, clay and stones (Geyer et al., 2011). Sporadic layers with lower permeabilities lead to a separation of up to three aquifer levels (Wirsing and Luz, 2007). The upper aquifer is unconfined with a water table between 2 and 10 m below the ground. The
135 flow direction is ~~towards~~ northwest ~~to of~~ the Rhine River with groundwater flow velocities ranging between 0.5 and 1.5 m/d (Technologiezentrum Wasser, 2018).

Based on the land use plan of Karlsruhe, about 20 % of the ~~study~~-area (i.e. urban area, city centre, neighbouring districts, as well as parts of the Hardtwald forest and several outskirts) is covered by buildings. The rest is ~~characterised by~~ vegetation (~ 56 %) and artificial surface covers (~ 24 %), showing the complexity and heterogeneity of the urban environment.
140 According to Benz et al. (2016), the annual mean groundwater temperature (GWT) in Karlsruhe in the years 2011 and 2012 was 13.0 ± 1.0 °C. Distinct temperature hotspots occur mainly below the city centre, where building densities are highest. In the north-western part of Karlsruhe, the increase of GWT ~~with~~ was about 3 K warmer than the annual mean land surface temperature (LST), which is mainly caused by several groundwater reinjections of thermal wastewater (Benz et al., 2016). In general, groundwater in the region of Karlsruhe is of good quality ~~so that and~~ the local drinking water supplier (Stadtwerke
145 Karlsruhe) only needs to ~~removes~~ oxidised iron and manganese from the pumped groundwater. However, two main contaminations, which affect groundwater quality, are known in the urban area (Stadt Karlsruhe, 2006). A contaminant plume, which contains a polycyclic aromatic hydrocarbons concentration of up to 500 µg/l, of 200 m length over the entire aquifer thickness is located at a former gas plant in the east of Karlsruhe (Figure S1b) (Kühlers et al., 2012). ~~in the east of Karlsruhe (Figure S1b). This plume contains a polycyclic aromatic hydrocarbons concentration of up to 500 µg/l caused by the former gas plant in the east of Karlsruhe (Kühlers et al., 2012).~~
150 ~~Moreover, three parallel contamination plumes, of 2.5, 500 km length each, can be found in the southeast of Karlsruhe (Figure S1b), where highly volatile chlorinated hydrocarbons (7 µg/l - 26 µg/l) and their degradation products were detected (Wickert et al., 2006).~~

2.2 Material and sampling

From 2011 to 2014, samplings of groundwater parameters and fauna were performed in 39 groundwater monitoring wells in
155 Karlsruhe, of which eight wells are in the forest and 31 in the urban area. At the beginning of each sampling process, temperature and electrical conductivity were measured with an electric contact gauge (Type 120-LTC, Hydrotechnik) at a

depth interval of 1 m. Using a bailer (Aqua Sampler, Cole-Parmer), water from the bottom of the groundwater monitoring wells was sampled and the pH value (Multiline Type 3430; WTW GmbH, Weilheim Germany) as well as the contents of dissolved oxygen (Multiline Type 3430; WTW GmbH, Weilheim Germany), iron, nitrate (NO_3^-) and phosphate (PO_4^{3-}) (RQflex® plus 10 Reflectoquant®; Merck Millipore KGaG, Darmstadt Germany) were measured.

In accordance with the suggestion made by Hahn and Gutjahr (2014), several integrative samplings (i.e. repeated samples taken over a period of time) ~~(at least three)~~ were conducted to ~~achieve—capture~~ an ecologically ~~representative samplingrepresentation~~ of groundwater fauna; which ~~also~~ reflects the occurring species at a community level. Every well is sampled at least three times. From 2011-2012, 22 measurement wells (mainly in the Hardtwald and the North-West of Karlsruhe) were sampled six times at a minimum interval of two months. In 2014, 17 measurement wells, mainly located in the south/inner city, were sampled three times (see Table S2). As the aim of this study is to provide a first-tier screening of the groundwater ~~ecology~~ecological status, we sampled d the fauna in the monitoring wells in accordance with the sampling manual of the European PASCALIS Project (Malard et al., 2002) and the procedure described by Hahn and Fuchs (2009), using a modified Cvetkov net.

Furthermore, the relative amount of sediment as an indication of the nutrient availability and the cavity system was measured. Before the fauna sample from the net sampler was passed over a sieve with a mesh size of 74 μm , the sediment is separated and classified in different categories (sand, fine sand, ochre, detritus, silt). It should be noted that the detritus content is not recorded quantitatively but on the basis of estimated frequency classes. The estimation of the relative amounts of sediment per sample is based on Table S1 in the supplement.

Mann-Whitney-tests (U-tests) were applied to detect potential impacts of groundwater characteristics (physical-chemical parameters), geology and well design on the groundwater quality as well as on groundwater fauna. Samples were regarded as significantly different if the p -value was $< 5.0 \times 10^{-2}$.









To better understand large-scale relationships as well as fine structures of high-dimensional biological data, the PHATE (potential of heat diffusion for affinity-based transition embedding) analysis introduced by Moon et al. (2019) (<https://github.com/KrishnaswamyLab/PHATE>) was used. This dimensionality-reduction method generates a low-dimensional embedding specific for visualization, which provides an accurate, denoised representation of both local and global structures of a dataset without imposing strong assumptions on the structure of the data. The PHATE algorithm computes the pairwise distances from the data matrix and transforms the distances to affinities to encode local information by applying a kernel function, which is developed to Euclidian distances. By using diffusion processes, global relationships are learned and encoded using the potential distance. Finally, the potential distance information is embedded into low dimensions for visualization by using metric Multi-Dimensional-Scaling (MDS) (Moon et al., 2019). Objects that are close to each other in the final graph therefore have similar characteristics.

~~The organism communities of the groundwater consist of microorganisms and invertebrates (in particular Crustaceans) (Griebler et al., 2014).~~ Crustaceans, especially Amphipods and Copepods, represent the majority of ~~the~~ groundwater fauna.

The identification keys from the following studies were used to identify the different groups in the samples: Einsle (1993),

Janetzka et al. (1996), Meisch (2000), Schellenberg (1942) and Schminke et al. (2007). The sampled fauna for this study can be assigned to the subphylum *Crustacea* and four other subordinate taxa (Table 1).

Table 1: Overview of the sampled fauna, divided into the subphylum *Crustacea* and other subordinate taxa.

Subphylum: <i>Crustacea</i>	Size [mm]	Habitats	Species number
Order: <i>Cyclopoida</i> 	0.4 - 0.7 ¹	Fresh and marine water, groundwater ¹	2982 species and subspecies of the genus <i>Cyclops</i> worldwide ² , <u>8 stygobiotic species in Germany³</u>
Order: <i>Harpacticoida</i> 	<u>< 0.5⁴</u>	<u>Marine, freshwater, semi-terrestrial environments and groundwater⁵</u>	<u>599 (sub-)species worldwide², 20 stygobiotic species in Germany³, 17 stygophile* & stygobiotic species in Baden-Württemberg⁶</u>
Genus: <i>Parastenocaris</i>	<u>0.3 - 0.5¹</u>	<u>Tertiary relic living in cavity rooms of streams, in groundwater and moss¹</u>	<u>206 (sub-)species worldwide² (16 stygophile & stygobiotic species in Baden-Württemberg¹)</u>
Order: <i>Bathynellacea</i> 	0.5 - 5.4 ⁷	Cavity systems ⁷ and in groundwater ⁸ (foreign tropical origin) ⁹	Exclusively 160 real groundwater-stygobiotic <u>stygobiotic</u> species (stygobiotic) worldwide⁹⁵, 8 species in Germany³
Order: <i>Amphipoda</i> 	0.5 – 30 ¹	Sea, fresh water ¹ and in healthy groundwater ecosystems (important ecosystem service providers ¹⁰ & biodiversity indicators in Europe ¹¹)	321 stygophile and & stygobiotic species for in Europe ¹² , <u>24 stygobiotic species in Germany³</u>
Other subordinate taxa	Size [mm]	Habitats	Species number
Subclass: <i>Oligochaeta</i> 	< 1 – 3 ¹³	Colonise every habitat, groundwater ¹³	<u>100 species worldwide¹⁴ and 27 stygobiotic species in Europe¹³ and 100 species worldwide¹⁴</u>
Phylum: <i>Nematoda</i> 	1 – 3 ⁹	Colonise every habitat ⁹ , can live under unfavourable conditions ¹⁵	20,000 species worldwide ¹⁶ , <u>60 stygobiotic species in Europe, 6 species in Germany³</u>
Class: <i>Turbellaria</i> 	0.4 – 5 ¹⁷	Sea, brackish and fresh water and groundwater ¹⁷	The class includes 3,400 species worldwide ¹⁷ , <u>7 stygobiotic species in Germany³</u>
Subclass: <i>Acari</i> 	a few mm ⁹	Colonize every habitat, also groundwater, have high demands on water quality ⁹	Worldwide more than < 5,000 water mite species <u>worldwide¹⁸, 10 stygobiotic species in Germany³</u>
<div> ¹ Fuchs et al. (2006) ² Galassi (2001) ³ Zenker et al. (2020) ⁴ Hahn (1996) ⁵ Galassi et al. (2009) ⁶ Fuchs (2007) </div> <div> ⁷ Sauermost and Freudig (1999a) ⁸ Camacho (2006) ⁹ Hunkeler et al. (2006) ¹⁰ Boulton et al. (2008) ¹¹ Stoch et al. (2009) ¹² Botosaneanu (1986) </div> <div> ¹³ Sauermost and Freudig (1999b) ¹⁴ Batzer and Boix (2016) ¹⁵ Hahn et al. (2013) ¹⁶ Eckert et al. (2008) ¹⁷ Sauermost and Freudig (1999c) ¹⁸ di Sabatino et al. (2000) </div>			

*Stygophile organisms are found primarily in surface water, but they can survive in shallow groundwater for a while (Preuß and Schminke, 2004).

2.3 Classification scheme by ~~the Federal Environmental Agency of Germany~~Griebler et al. (2014)

Commissioned by the Federal Environmental Agency of Germany (UBA), Griebler et al. (2014) developed a two-step ecologically based classification scheme ~~for the Federal Environmental Agency of Germany (UBA)~~ for characterization of groundwater ~~ecology~~ecosystems and. ~~They~~ also defined spatially dependent reference values of ecologically intact groundwater ecosystems. In order to enable a statement about the exposure ~~(organic, chemical, structural)~~ of the groundwater at a specific site, biotic and abiotic parameters, which are determined and compared with reference values, are used to distinguish locations with ~~ecological conditions, which are~~ very good or good (i.e. O.K.), i.e. very good or good ecological conditions or locations which fail these criteria, i.e. affected areas (Figure ~~1~~Figure 4). If an ecological assessment of groundwater ecosystems based on the groundwater fauna takes place, some faunistic criteria must be considered. Invertebrates avoid habitats that are ochred or have a low content of dissolved oxygen. Thus, unstressed or natural habitats are defined as Five criteria, of which at least three biological/ecological indicator parameters are to be selected, are taken as basis for a reliable assessment. The measured indicator parameters of any location have to be compared with the reference values provided by Griebler et al. (2014). Unstressed or natural groundwater habitats are defined as areas with a content of dissolved oxygen > 1.0 mg/l, that are not ochred and have an existing fauna, i.e. an amount of > 50 % of Stygobites, of > 70 % of Crustaceans and of < 20 % of Oligochaetes (Figure 1~~Figure 4~~). This ~~results in~~allows a qualitative interpretation of the ecological condition of the groundwater system. If the results ~~s~~ indicates affected ecological conditions, ~~which means that i.e.~~ one or more biological/ecological indicators are out of the reference range, an assessment according to the Level 2 scheme is necessary. This requires a determination of reference values at local reference locations which are protected and have a weak surface influence as well as a subsequent comparison of these values with measured data. to obtain also a qualitative and quantitative interpretation of the ecological conditions. As our aim is a first-tier screening of an urban area, we only apply Level 1 in our study. ~~In addition, Level 2 requires a determination of reference values at local reference locations, which are well protected and have only a weak surface influence, and a subsequent comparison of these values with measured data.~~




Figure 1: UBA-eClassification scheme by Griebler et al. (2014) according to Level 1 for groundwater ecosystems on the basis of groundwater fauna (modified after Griebler et al. (2014)).

3. Results and discussion

3.1 Physical and chemical parameters

First, the groundwater conditions in the study site are evaluated by their physical-chemical characteristics. The following values are average values of the individual samplings ~~of from~~ each monitoring well. In order to allow a spatially differentiated assessment, the study site is classified in different zones based on land use types provided by the European seamless vector database of the CORINE Land Cover (CLC) inventory (GISAT, 2016). Based on this data the study site is subdivided into: (1) Forest area (local name: Hardtwald) and (2) Urban area containing industrial, commercial and residential areas (Figure 2a). A more detailed subdivision in the urban area did not appear reasonable due to the heterogeneous structure into two separate zones based on land-use types: (1) Forest area (local name: Hardtwald), (2) Urban area containing industrial, commercial and residential areas (Figure 2a).




Figure 2: Overview map city area of Karlsruhe: (a) land use plan (GISAT, 2016) and average groundwater temperature of the multiple measurements [°C] at the bottom of the monitoring wells; (b) detailed groundwater fauna: colours of the circles shows the different taxa in the sample [%], the size indicates the number of individuals; (c) faunistic evaluation according to UBA-classification scheme after Griebler et al. (2014) and groundwater contour map in metres above sea level (modified after the local authority real estate of Karlsruhe).

As expected, measured GWT at the bottom of the wells, in 8.5 to 39.0 m depth, are mainly constant over the repeated measurements. The lowest GWT ranging between 10.5 and 10.9 °C were measured in the eight wells of the forest area (Table S4S2). In contrast, the highest average GWT with 17.5 °C was measured in a well near the city hospital (T113) (Figure 2a). The mean value of all wells is 13.5 ± 2.1 °C, which is similar to the results from Benz et al. (2015) with 13.0 ± 1.0 °C.

According to Benz et al. (2017), annual shallow GWT vary between 6 and 16 °C in the area of Karlsruhe, which is in line with the temperatures measured during fauna sampling (Figure 3Figure 3a). For the urban area in the north-western part of the city, Figure 2Figure 2a shows a clear warming trend, which was also observed by Menberg et al. (2013a,b). The increased GWT in this area can be traced back to effects of urban infrastructures and industries, which use groundwater for cooling purposes.




Figure 3: Boxplots of the physical and chemical parameters for the forest and urban area in the study site and the proportion of wells in which ecological conditions are O.K. in percentage [%] indicated by the blue (forest area) and red (urban area) lines (secondary axis) [%], second axis; (a) average temperature of the repeated measurements [°C] at the bottom of the monitoring wells; (b) average content of dissolved oxygen [mg/l] of the monitoring wells; (c) average nitrate content [mg/l] of each monitoring well. (n = number of wells)

The content of dissolved oxygen acts as a limiting factor for groundwater fauna, since groundwater is usually under-saturated with a varying oxygen content between 0 and 8 mg/l (Griebler et al., 2014; Kunkel et al., 2004). In this study, the average content of dissolved oxygen in all wells is between 1.0 and 12.8 mg/l (Figure 3Figure 3b and Figure S1a). As expected, the monitoring wells, located in the forest area (Hardtwald) show the highest content, while the lowest values are found in urban areas, ~~which and~~ is likely linked to aquifer contamination and other anthropogenic effects (content of dissolved oxygen of forest vs. urban area: U-test: $p\text{-value} = 5.3 \times 10^{-3}$, $n = 8; 31$). Urban water can be polluted in multiple ways, which affects the chemical and biological oxygen consumption in the groundwater. The higher the pollution and/or biological activity, the lower the dissolved oxygen (Kunkel et al., 2004; Griebler et al., 2014). Moreover, it seems that with a greater depth of the measurement wells the content of dissolved oxygen is increasing (U-test: $p\text{-value} = <10^{-13}$, $n = 39$). This can be explained by the fact that shallow wells can have a low water column in which oxygen can rapidly be consumed by groundwater microorganisms, chemical reactions and/or groundwater fauna. In the upper unscreened part of deeper wells, dissolved oxygen can be consumed while in the lower screened part oxygen is continuously refilled by oxidic groundwater from the surrounding (Malard et al., 2002). Furthermore, reducing conditions in the overlaying soil can result in a low content of dissolved oxygen in groundwater.

265 Nitrate is often named as an important pollutant in groundwater. The natural and geogenic concentrations of nitrate in groundwater is usually under 10 mg/l (Griebler et al., 2014). In our study area, the average nitrate content of all wells varies between 1.3 and 14.7 mg/l. In the urban area average nitrate concentrations are generally higher and correlate inversely with the content of the dissolved oxygen (U-test: p -value = 4.0×10^{-3} , $n = 39$) showing the link between pollution-nitrate content and oxygen consumption. Wells with a content of dissolved oxygen below 1.5 mg/l have an average content of nitrate of 1.5 mg/l, caused by nitrate reduction under anoxic conditions. Groundwater with reducing conditions (< 5 mg/l dissolved oxygen) has an average nitrate content of about 7 mg/l in contrast to groundwater with oxidising conditions with 9 mg/l, which is characterised by the oxidation of ammonium to nitrate. The lowest nitrate concentrations are found in the forest area (Figure 3c and Figure S1c), where atmospheric nitrogen is held held back by forest soils (U-test: p -value = 1.7×10^{-3} , $n = 8$) and fertilization is prohibited due to water protection regulations in the forest area. ~~At the same time, anthropogenic impact is minimal as fertilization is forbidden due to the presence of water protection areas in the forest area~~ (Aber et al., 1998; Schönthaler and von Adrian-Werburg, 2008). ~~Within the study, the~~ Moreover, the average ~~contents concentration~~ of iron and phosphate are low and in most cases below the detection limit of the test (Figure S1d, e) and also below the natural and geogenic concentrations (phosphate: 0.05 mg/l (Griebler et al., 2014) and iron: 3.3 mg/l (Kunkel et al., 2004)) within the study site.

270

275

280 Considering these findings, clear differences in the spatial distribution patterns of abiotic groundwater characteristics are noticeable. The rural forest area shows lower average GWT than the urban area (U-test: p -value = 3.3×10^{-5} , $n = 8$; 31), lower nitrate concentrations (U-test: p -value = 4.1×10^{-3} , $n = 8$; 31) and higher dissolved oxygen concentrations (U-test: p -value = 5.3×10^{-3} , $n = 8$; 31), which indicates a correlation between abiotic groundwater characteristics and land use in the study area. Moreover, no impact of groundwater originating from the urban area on the wells in the forest area is observed, as

285 the groundwater flow direction in Karlsruhe is northwest (see Chapter 2.1 and Figure 2c). Further investigations demonstrated that besides one larger and two smaller contamination sites (however, still with concentrations below the threshold values, Figure S1b), only minor groundwater pollution is documented in Karlsruhe (see Supplement). The chemical and physical parameters considered in the long-term monitoring system are within the range of local background and below threshold values of the drinking water ordinance of Germany (see Supplement for more information). Thus, the main documented impacts on


290 groundwater quality in the study area are related to temperature and oxygen. The rural forest area shows a lower average GWT, lower nitrate concentrations and higher dissolved oxygen concentrations, which indicates a correlation between abiotic groundwater characteristics and land use in the study area.

3.2 Groundwater fauna

295 The biotic communities of the groundwater consist of microorganisms and invertebrates (in particular Crustaceans) (Griebler et al., 2014). In the ~~entire pool of~~ samples 3, ~~633-666~~ individuals were detected in 37 of 39 wells (Table ~~S2S3~~). With ~~2,014~~ 2,047 individuals, the group of *Crustacea* was found to be the most abundant ~~group~~ (~~55-56~~ %). 976 individuals (27 %) of the order of *Cyclopoida* dominated this group, followed by the genus *Parastenocaris* with 599 individuals (~~47-16~~ %), by the order of

300 *Bathynellacea* (371), *Amphipoda* (66), *Harpacticoida* (33) and *Nauplia*. The communities of the monitoring wells also frequently contained *Oligochaetes* (1,343 individuals, 37 %). Furthermore, individuals of the phylum *Nematoda* (228 individuals) and *Microturbellaria* (46 individuals) were also often present.

Overall, there is a noticeable difference in the spatial distribution of species within the study area. Individuals of the subphylum *Crustacea* were found in larger numbers, with regard to the number of wells, in the monitoring wells in the forest area (~~660~~ 690 individuals in eight wells) ~~than compared to those~~ in the urban area (~~1,354~~ 1,357 individuals in 31 wells). Furthermore, no individuals of the order *Bathynellacea* and only 135 individuals of the genus *Parastenocaris* were found in the forest area. In contrast, larger numbers of the latter species as well as of *Oligochaetes* are ~~characteristics~~ characteristically found in of the wells in the urban area. However, in contrast to the abiotic characteristics, no clear pattern of faunal diversity and land use was observed; as Crustaceans and individuals of other subordinate taxa were found both in the rural forest and in the urban area.



310 **Figure 4: Boxplots of the amount of fauna [%]: (a) proportion of individuals and of wells in which ecological conditions are O.K. (secondary axis) [%] of the forest area; (b) proportion of individuals and of wells in which ecological conditions are O.K. [%] of the urban area; (c) proportion of individuals and of wells in which ecological conditions are O.K. [%] divided based on the results of the UBA-classification scheme by Griebler et al. (2014); (d) proportion of individuals and of wells with affected ecological conditions**

[%] divided based on the results of the UBA-classification scheme by Griebler et al. (2014). The colour of the boxes shows the different taxa in the samples. (n = number of wells)

Stygobiotic Amphipods, i.e. large-bodied invertebrates that predominantly live within wells which due to their size have a habitat preference for open spaces such as wells (Table 1) (e.g. Hahn and Matzke, 2005; Korbel et al., 2017), were found ~~only~~ in only three wells (~~Figure 2e~~Figure 2b). 46 individuals of this order were detected in the forest and 20 individuals in the urban area (Figure 4a,b). Although statistical analysis showed no clear differences between the abundance of Amphipods and land use (U-test: p -value = 1.5×10^{-1} , $n = 8$; 31), the higher number of individuals in the forest area could support the hypothesis that~~As mentioned above,~~ Amphipods indicate healthy groundwater ecosystems, as they react most sensitive to disturbances such as pollutants (Korbel and Hose, 2011) as well as and groundwater temperature. ~~(11 ± 5 °C (Briellmann et al., 2011) up to 17 °C).~~ In laboratory experiments with a thermal tank, Briellmann et al. (2011) found that 77 % of the individuals of the studied Amphipods (*Niphargus inopinatus*) preferred areas with a temperature between 8 and 16 °C. In addition, Spengler (2017) and
Issartel et al. (2005) observed maximum temperatures up to 17 °C. The lack of a statistically significant correlation might also be related to the low number of wells ($n = 8$ in the forest area) and individuals ($n = 46$). Amphipods ~~They~~ are important ecosystem service providers in terms of bioturbation and organic decomposition (Boulton et al., 2008). As observed in laboratory experiments (Smith et al., 2016), as they show an active movement~~actively move,~~ with ~~possible~~ migration speeds between 1.7 and ~~3.5×10^4 m per year~~ observed in laboratory experiments (Smith et al., 2016). ~~If~~ In most cases when
Amphipods were found, ~~in most cases~~ higher amounts ~~concentrations~~ of individuals of the order *Cyclopoida* were also identified (Abundance *Amphipoda* vs. *Cyclopoida*: U-test: p -value = 9.6×10^{-5} , $n = 39$). Individuals of the latter order were ~~mostly generally~~ be found in larger quantities in the majority of the wells (479 in the forest area and 497 in the urban area), as they are the largest group of Crustaceans in this environment (Fuchs et al., 2006) and can tolerate a wide temperature range (e.g. upper thermal limit of 26.9 ± 0.2 °C in laboratory tests by Sánchez et al. (2020)) (Spengler, 2017).

The order *Harpacticoida*, which includes the genus *Parastenocaris*, have an elongated body shape and a stem-chiselling movement, which is why they are predestined for living in cavities and groundwater (Hahn, 1996; Fuchs, 2007), preferring sand and gravel as a substrate (Galassi et al., 2009). Larger numbers of *Parastenocaris* (464 individuals), which can tolerate GWT from 8 to > 20 °C (Fuchs et al., 2006) (e.g. *Parastenocaris phyllura* up to 22.5 °C in laboratory tests; ~~(Glatzel, 1990)~~), were found in the urban area, especially in the northwest area (Figure 2~~Figure 2b~~). This area is characterised by GWT between
16 and 18 °C, the highest at the study site. This observation is comparable with previous studies (Hahn, 2006; Hahn et al., 2013; Spengler, 2017), which showed that the genus *Parastenocaris* is particularly non-competitive and can often be found isolated in structurally burdened and physico-chemically altered areas. Accordingly, only 135 individuals were detected in the forest area.

In addition, quantities of *Bathynellacea* (371 individuals) were found in five monitoring wells all located in the urban area in a depth of 9 to 13.5 m and by medium GWT (12 – 15 °C)~~at a GWT of 12 – 15 °C~~ all located in the urban area (371 individuals), respectively (Figure 4b). This order typically inhabits the interstitial groundwater, which is characterised by a dominant exchange with the surface water and high variations in GWT, and can tolerate temperatures up to 18 °C (Stein et al., 2012).

Interestingly, one location in the southern city area with 272 individuals is characterised by a high fluctuation in GWT (standard deviation of 3.4 °C) and a rather high nitrate content (8.3 mg/l) compared to wells in the forest area, which are both indications for a disturbed and stressed habitat.

Besides the group of Crustaceans, Oligochaetes, which can tolerate a wide temperature range, were also found in large abundance in the study site. A significant amount of the subclass *Oligochaeta* (996 individuals) was found in the urban area (Figure 4b), compared to an overall number of 1,343 individuals. In general, the number of Oligochaetes is larger in locations with high GWT (12.6--17.3 °C) and high nitrate concentrations (up to 14 mg/l, which is above the geogenic concentration of 10 mg/l and higher compared to wells in the forest area).

Finally, Nematodes and Microturbellarians were found at locations with unfavourable living conditions, such as a low content of dissolved oxygen, or a high amount of fine substrates, as also reported by Hahn et al. (2013). ~~Both~~ can tolerate high temperature ranges (*Turbellaria*: 2 – 20°C (Herrmann, 1985), *Acari*: 9.1 – 18.5 °C (Więcek et al., 2013)). Here, both were found in larger quantities in the urban area of Karlsruhe (Figure 4b). This area has the lowest content of dissolved oxygen, high relative relatively higher amount of detritus (> 2) ~~and the highest nitrate concentrations (> 6 mg/l)~~.

Eventually, correlation analysis between groundwater fauna and the chemical parameters showed that Stygobites are only slightly affected by groundwater quality chemistry (Hahn, 2006; Schmidt et al., 2007; Stein et al., 2010). Only the Spearman's rank correlation coefficient ρ between the number of taxa and the content of dissolved oxygen is significant with a value of $\rho = 0.53$ (p -value = 3.0×10^{-4} , $n = 39$ ~~p -value = 0.0005, $n = 39$~~). Moreover, it is assumed that groundwater fauna can usually cope well with short-term changes of chemical-physical parameters (Griebler et al., 2016). Previous studies showed that some species can even benefit from pollutants (Matzke, 2006; Zuurbier et al., 2013). In case of nitrate, numerous studies underline that nitrate at concentrations below 50 mg/l does not directly affect groundwater fauna (Fakher el Abiari et al., 1998; Mösslacher and Notenboom, 2000; Di Lorenzo and Galassi, 2013; Di Lorenzo et al., 2020). As the highest average nitrate content per well is below 15 mg/l in this study, a direct negative effect of the nitrate concentration on the groundwater fauna is unlikely. Thus, nitrate is only mentioned as one measured parameter and is not discussed as a potential anthropogenic impact in this study.

The natural influence on porosity, groundwater flow and nutrient delivery were also discussed as primary influence on natural Stygobites distribution by previous studies (Hahn, 2006; Korbel and Hose, 2015). One important natural influence is the local geology, as fine sands and silts are typically rather harsh environments, resulting in an impoverishment of specific groundwater fauna such as *Crustacea* (Hahn, 1996). The city of Karlsruhe is located on carbonate ('Würm') gravel and river terrace sands, pervaded by bands of drifting sand and inland dune sands. These sediments are highly water-permeable and show almost exclusively vertical seepage of water movement. Flood sediments (on top of river gravel) and bog formations, are located in the east and west of Karlsruhe (Regierungspräsidium Freiburg, 2019). This local geology limits the cavity size and therefore has impacts on the habitat of the groundwater fauna (Wirsing and Luz, 2007). For example, individuals of the genus *Parastenocaris* typically inhabit small-scale cavity systems (Spengler, 2017). Individuals of this genus can be found both in the wells drilled in gravel (4 wells) and in drifting sand sediments (3 wells) (abundance *Parastenocaris* vs. geological units:

U-test: p -value = 1.4×10^{-9} , $n = 39$). Amphipods are predominantly found in measurement wells located in the 'Würm' gravels (in 5 of 7 wells) (abundance *Amphipoda* vs geological units: U-test: p -value = 9.0×10^{-11} , $n = 39$). Moreover, it seems that differences in the geological units have an influence on the total amount of individuals (U-test: p -value = 1.7×10^{-9} , $n = 39$) and the relative amount of detritus (U-test: p -value = 3.0×10^{-3} , $n = 39$). As these results show, regional geology seems to have an influence on the occurrence of specific groundwater taxa and on the number of individuals as well as on food supply, in terms of available organic matter. However, it is not possible to give a reliable estimate of the strength of the anthropogenic impacts, e.g. if they are strong enough to overrule the regional selective forces. Hence, this should be investigated in more detail in future studies.

Some limitations regarding the sampling method ~~have to~~must be considered when interpreting the faunistic results. In this study, a simple basic screening of well water was conducted, using a net sampler and bailer, to assess conditions in the groundwater monitoring wells (39 wells with an average diameter of 132.5 mm, which corresponds to an area of 0.003 % of the total urban area). According to the sampling manual of the PASCALIS Project 'the use of a phreatobiological net alone is considered as a satisfactory method for sampling groundwater fauna in large diameter wells' (Malard et al., 2002). Yet, several studies (e.g. Scheytt, 2014) report that scooped samples of wells are not representative, and therefore the water remaining in a well has to be purged and discarded before sampling. Nevertheless, pumping can result in the selection of the taxa, especially in the presence of very fine sediments, and can result in changes of the sediment composition in the surrounding of wells and therefore in changes of habitat conditions. Other studies, on the other hand, found no significant differences in hydro-chemical values (temperature, pH, dissolved oxygen, etc.) between the surrounding groundwater and the standing water in a well (Hahn and Matzke, 2005; Korbel et al., 2017). The sampled groundwater fauna of corresponding wells and aquifers were also shown to be similar with respect to the types of faunal communities. However, in terms of total abundance ~~however~~, as well as the numbers of individuals per litre, monitoring wells appear to exhibit larger numbers, caused by filtration effects (Hahn and Matzke, 2005; Hahn and Gutjahr, 2014; Korbel et al., 2017). As the aim of this study is to provide an overview of the groundwater fauna community (assess biodiversity) and to receive a first impression of groundwater ecology, sampling the fauna by using a net sampler is sufficient. In order to achieve a representative sampling of groundwater fauna in the aquifer and to reflect the occurring species at a community level, a more comprehensive sampling method is required, e.g. the use of a defined standard sampling method using a pump to collect animals (Malard et al., 2002). Care should also be taken when interpreting faunistic results of sites that are sampled in different years. To improve comparison of the biotic communities, a consistent sampling period of every well is necessary in the future.

3.3 Classification scheme by ~~the Federal Environmental Agency of Germany~~Griebler et al. (2014)

In three wells, evaluation with the ~~UBA~~-classification scheme by Griebler et al. (2014) was not possible due to ocherous conditions in two monitoring wells and low content of dissolved oxygen (<1 mg/l) in the third well. According to the ~~UBA~~ classification scheme by Griebler et al. (2014), unstressed (meaning no natural or anthropogenic stressors), or natural groundwater habitats have an amount of more than 70 % of Crustaceans and less than 20 % of Oligochaetes. In 36 % of the

415 sampled wells, i.e. 14 out of 39, these criteria ~~are-were~~ fulfilled, indicating O.K. ecological conditions or in other words a natural groundwater habitat (Figure 4c). These natural areas tend to contain more individuals of the orders *Amphipoda*, *Cyclopoida* and *Bathynellacea*. Monitoring wells, which do not fulfil these criteria and are accordingly defined as affected areas not having natural ecological conditions, contain more Oligochaetes and also Nematodes, which is partly explained by the used criteria of this classification scheme (Figure 4d).

420 Surprisingly, only 50 % of the wells in the rural forest, which is also the catchment area of the drinking water supply of Karlsruhe, are described as natural groundwater habitats. An identical number of wells yielded habitats with affected ecological conditions. The main difference between natural and affected wells in the forest area arises from the occurrence of specific species. ~~86 to 100 % of species found in natural wells are Crustaceans. Natural wells have an amount of Crustaceans of 86 to 100 %,~~ in contrast to affected wells with only 33-67 % (Table ~~S1-S2~~ and Table ~~S2S3~~). However, the abiotic parameters scarcely
425 differ between natural and affected wells (average values for GWT: 10.8 and 10.6 °C, dissolved oxygen: 7.1 and 8.8 mg/l, nitrate: 2.5 and 3.0 mg/l), indicating that there are other processes or parameters that influence the groundwater fauna in these wells. One reason could be the varying local geology-, ~~as mentioned above. Moreover, food supply is one of the most limiting parameters for the survival of groundwater fauna (Datry et al., 2005; Hahn, 2006). If the organic carbon supply varies on a small scale, this can influence microbiology and therefore groundwater fauna as well, although, short-term changes in nutrient~~
430 ~~supply can be compensated by groundwater fauna, as fine sands and silts are typical rather harsh environments resulting in an impoverishment of specific groundwater fauna such as Crustacea (Hahn, 1996).~~

In contrast to the forest land, the majority of wells (65 %) in the urban area are categorised as affected habitats. As expected, this indicates anthropogenically influenced groundwater ecosystems beneath the studied urban area. Once more, no significant differences between the abiotic parameters of natural and affected wells are observed (e.g. median of dissolved oxygen: 4.7
435 and 5.8 mg/l, median of nitrate: 7.2 and 7.8 mg/l). On the other hand, the remaining 35 % of the wells in the urban area show natural ecological conditions, even though some of them are located in areas with anthropogenic impacts such as increased groundwater temperatures. Hence, no distinct spatial pattern of the ecological condition with respect to land use could be identified.

~~In future, a further subdivision of a study area in more land use categories could be useful to specifically look at typical anthropogenic impacts. Furthermore, the integration of more biological criteria is useful to improve the results of the assessment according to Griebler et al. (2014) and to use more different assessments, like the similarly structured GHI or wGHI^N (Korbel and Hose, 2017; Di Lorenzo et al., 2020b). Moreover, there are a couple of newly developed indexes, like the D-A-C-Index, which is based on microbiological indicators and shows whether groundwater reserves deviate from natural references (Fillinger et al., 2019), which can be used in the future. As mentioned in the introduction, another way to quantify~~
440 ~~the relevant ecological conditions in the groundwater is the GFI. During the preparation of this study, the GFI was tested on the data (see Supplement), however, it did not provide any additional information or valuable insights and was therefore excluded. The influence of multiple stressors, such as the pollution of the groundwater through industrial plants etc., and their effects on the governing parameters can bias the GFI. In general, the GFI seems to be suitable only for unpolluted and~~

450
455
460
465

anthropogenically undisturbed groundwater with sufficient oxygen concentrations (> 1 mg/l). Moreover, under urban areas changes in GWT are caused by anthropogenic heat inputs (Menberg et al., 2013b, 2013a; Benz et al., 2014; Tissen et al., 2018), rather than being related to surface water influences. Hence, the GFI appears to be unsuitable for the assessment of the groundwater fauna in an urban setting. The same outcome emerges for the Shannon diversity index, which was also tested during the preparation of the study and showed no clear distribution pattern according to faunal diversity and was therefore not considered further.

This observed spatial heterogeneity in ecological conditions and heat anomalies in an urban area therefore also offer the potential to use groundwater for heating and cooling, and even to locally store energy in form of aquifer thermal energy storage (ATES) systems (e.g. Fleuchaus et al., 2018).

3.4 PHATE analysis

A PHATE analysis is conducted using the following 15 input parameters: depth, GWT, nitrate and phosphate content, relative amount of detritus, geological unit, numbers of taxa, number of individuals, Shannon diversity, amount of Crustaceans and Oligochaetes (according to Griebler et al., 2014) and the abundance of Amphipods as well as of individuals of the order *Cyclopoida*, *Bathynellacea* and the genus *Parastenocaris*. The content of dissolved oxygen is not considered in this analysis, since it was always above the limit of 1 mg/l, except for in one case. Thus, dissolved oxygen is not expected to have an influence on the groundwater fauna in our study area.




Figure 5: PHATE visualization showing similarities between measurement wells. Different colours indicate the four clearly separable groups.

Four groups, which can be assigned predominant characteristics, can be distinguished in the PHATE visualization (Figure 5, Figures S3-S4). Three measurement wells (Group IV) contain neither Oligochaetes nor Crustaceans, indicating unfavourable living conditions. In contrast, the nine wells of Group III contain high amounts of Oligochaetes (100 % Oligochaetes according to the scheme of Griebler et al. (2014), and an average GWT of 14.3 °C (Table S4). However, diversity and abundance was found to be low in Group III.

An even higher average GWT of 15.0 °C was found for Group II, which mostly consists of wells drilled in drifting sand sediments. Surprisingly, these wells also show the highest diversity (\geq three Taxa per well), the highest Shannon diversity (see Supplement), highest amount of individuals in total, as well as of individuals of the genus *Parastenocaris*. Individuals of this genus are often found isolated in altered areas (Spengler, 2017). Moreover, in five wells of Group II individuals of the order *Bathynellacea*, which can tolerate temperatures up to 18 °C and typically inhabit interstitial groundwater (Stein et al., 2012), were found. The presence of individuals of the genus *Parastenocaris* and the order *Bathynellacea* in Group II suggests that they may act as type species for urban situations. The observation that Group II shows the highest GWT and the highest Shannon diversity is in contrast to findings of previous studies that noticed decreased diversity at elevated temperatures (Briellmann et al., 2009). These diverging observations suggest that faunal quantities, such as diversity or abundance, are not always suitable indicators for changes within organism communities. For example, if species disappear due to increased temperatures and are substituted by more tolerant species, the difference in diversity may be marginal and the change in the community may not be noticeable.

Wells of Group I (blue) are drilled predominantly in Würm gravel (geological unit of Group I vs. Group II: U-test: p -value = 8.2×10^{-3} , $n = 13; 14$), while having the lowest GWT (GWT of Group I vs. Group II: U-test: p -value = 2.0×10^{-5} , $n = 13; 14$). These wells show a moderate diversity and amount of individuals, yet the highest average amount of Crustaceans as well as the highest amount of Amphipods and individuals of the order *Cyclopoida*. Considering these findings and the U-Test results (see Table S5), the grouping of the measurement wells seems to be influenced by the composition of the groundwater organism communities, the faunal diversity (numbers of taxa and amount of individuals), as well as the geological unit and the GWT (Figure S3-S4).

Considering the spatial distribution of the grouped wells in the study area, it becomes apparent that all wells in the forest area fall within Group I (Figure 5). Those wells which are located outside the forest area are in locations with nearby green areas (parks, recreational areas, etc.). In contrast, the wells of the other three groups are heterogeneously distributed within the urban area. Many of the measurement wells of Group III and IV are associated with suspected or known contaminated sites (Figure S1b). Overall, a spatial pattern of abiotic groundwater characteristics (GWT, nitrate content) and occurrence of particular species (*Parastenocaris*) within the study area is apparent in the PHATE analysis, which confirm the classification according to land use. Yet again, no clear spatial pattern regarding faunal diversity in the study area could be identified. Although, a tendency of clustering of wells from Group III with higher diversity and amount of individuals can be seen in the northwest city area.

4. Conclusion

The aim of this study is to provide a first-tier assessment of the ecological state of groundwater in an urban area and to distinguish areas with a natural state of the groundwater ecology from anthropogenically affected areas. To achieve this, we examine the groundwater fauna, as well as abiotic parameters in 39 groundwater monitoring wells in the urban area (31 wells) of Karlsruhe, Germany, and a nearby forest land (eight wells) using the simple UBA-classification scheme by Griebler et al. (2014) to characterise the sampled monitoring wells.

We found a noticeable difference in the spatial distribution of abiotic groundwater characteristics and special species within the study area. The rural forest area shows lower GWT, lower nitrate concentrations and higher dissolved oxygen concentrations, which indicates a correlation between abiotic groundwater characteristics and land use. Moreover, Amphipods are more abundant in wells in the forest than in urban area. However, both in the rural forest and in the urban area Crustaceans and individuals of other subordinate taxa were widely found and therefore no clear spatial pattern regarding faunal diversity and land use was found, as both in the rural forest and in the urban area Crustaceans and individuals of other subordinate taxa were widely found. In terms of faunal quantity, Crustaceans were found in larger numbers, with respect to the number of wells, in the monitoring wells in the forest area than compared to those in the urban area. Larger amounts of the genus *Parastenocaris* as well as of Nematodes and Oligochaetes were found to be characteristics for wells in the urban area.

Furthermore, no clear spatial pattern of ecological groundwater conditions according to the UBA-classification scheme by Griebler et al. (2014) could be observed. Surprisingly, only 50 % of the sampled wells in the rural forest were described as natural (undisturbed) groundwater habitats, while the other four were characterised as habitats with affected ecological conditions. Yet, the majority of wells (65 %) in the urban area were classified as affected locations, which suggest, suggesting that there are noticeable differences in the groundwater ecosystems between the surrounding rural areas and urban areas. The Level 2 assessment from Griebler et al. (2014) can help to achieve a more reliable and quantitative ecological assessment of urban aquifers as it divides groundwater ecosystems in ecological grades according to the intensity of anthropogenic disturbance. It is based on the use of local reference values and the collaboration with experts, however, is challenging to apply. Therefore, further studies with large-scale and repeated measurement campaigns are needed to verify our findings. This should also include other cities and the determination of undisturbed local reference values which are required for a more reliable but also quantitative ecological assessment of urban aquifers. Moreover, a wider range of indicators should be considered in a classification scheme, such as temperature, porosity of the aquifer, groundwater flow, pollutants and nutrient supply, especially when investigating urban areas. In addition, an important adaptation for an improved evaluation method is the determination of fauna at species level which will provide more information (i.e. about Stygobionts, Stygophiles, Stygoxenes) and also consider the endemism of stygobiotic species. In this context, classification schemes should pay more attention to the different groundwater species and their potential use as indicator species. Thus, further studies with larger scale and repeated measurement campaigns are needed to verify our findings also in other cities, and to provide undisturbed local reference values which are required for a more reliable and also quantitative ecological assessment of urban aquifers.

535 Finally, city and also energy planning should seriously consider urban groundwater ecosystems as they provide valuable information for a sustainable use of the subsurface.

Data availability

Team list

Institute of Applied Geosciences (AGW), Karlsruhe Institute of Technology

540 Prof. Dr. Philipp Blum (philipp.blum@kit.edu)

MSc. Fabien Koch (fabien.koch@kit.edu)

Dr. Kathrin Menberg (menberg@kit.edu)

MSc. Svenja Schweikert (svenja.schweikert@googlemail.com)

545 Faculty of Nature and Environmental Sciences, University Koblenz-Landau

Dr. Hans Jürgen Hahn (hjahn@uni-landau.de)

Dr. Cornelia Spengler (spengler@uni-landau.de)

Author contributions

550 PB and HJH provided the topic and supervised the work, together with KM. SS and CS executed the field work and evaluated the samples. FK evaluated the collected data and interpreted as well as visualised the results and wrote the first draft of the paper. KM, CS, HJH and PB ~~participating-participated~~ in editing the paper.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

555 We would like to thank Annette März (Environmental Service, City of Karlsruhe), Michael Schöenthal (Public Utilities Karlsruhe) and Friedhelm Fischer (Civil Engineering Office of Karlsruhe). Special thanks are also given to Christine Buschhaus and Tanja Liesch for their support with the measurement and sampling (Institute of Applied Geosciences, Karlsruhe Institute of Technology).

Financial support

560 This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Review statement

References

- Aber, J., McDowell, W., Nadelhoffer, K., Magill, A., Berntson, G., McNulty, S., Currie, W., Rustad, L. and Fernandez, I.: Nitrogen saturation in temperate forest ecosystems – Hypotheses revisited, *Bioscience*, 48(11), 921–934, 1998.
- 565 Amt für Stadtentwicklung - Statistikstelle: Statistic Atlas Karlsruhe, [online] Available from: <https://web3.karlsruhe.de/Stadtentwicklung/statistik/atlas/?select=005> (Accessed 27 February 2019), 2018.
- Avramov, M., Schmidt, S. I., München, C. G., Jürgen, H. and Berkhoff, S.: Dienstleistungen der Grundwasserökosysteme, *KW - Korrespondenz Wasserwirtschaft*, 3(2), 74–81, doi:10.3243/kwe2010.02.001, 2010.
- Batzer, D. and Boix, D.: *Invertebrates in Freshwater Wetlands: An International Perspective on their Ecology*, Springer
570 International Publishing, Heidelberg., 2016.
- Benz, S., Bayer, P., Menberg, K. and Blum, P.: Comparison of local and regional heat transport processes into the subsurface urban heat island of Karlsruhe, Germany, *Geophys. Res. Abstr. EGU Gen. Assem.*, 16, 11252, 2014.
- Benz, S., Bayer, S. and Blum, P.: Identifying anthropogenic anomalies in air, surface and groundwater temperatures in Germany, *Sci. Total Environ.*, 584–584, 145–153, 2017.
- 575 Benz, S. A., Bayer, P., Goettsche, F. M., Olesen, F. S. and Blum, P.: Linking Surface Urban Heat Islands with Groundwater Temperatures, *Environ. Sci. Technol.*, 50(1), 70–78, doi:10.1021/acs.est.5b03672, 2016.
- Berkhoff, S.: *Die Meiofauna des Interstitials und Grundwassers als Indikator für Oberflächenwasser-Grundwasser-Interaktionen im Bereich einer Uferfiltrationsanlage*, University Koblenz-Landau., 2010.
- Botosaneanu, L.: *Stygofauna mundi: a faunistic, distributional, and ecological synthesis of the world fauna inhabiting
580 subterranean waters (including the marine interstitial)*, Leiden The Netherlands., 1986.
- Boulton, A. J., Fenwick, G. D., Hancock, P. J. and Harvey, M. S.: Biodiversity, functional roles and ecosystem services of groundwater invertebrates, *Invertebr. Syst.*, 22(2), 103–116, doi:10.1071/IS07024, 2008.
- Briellmann, H., Griebler, C., Schmidt, S. I., Michel, R. and Lueders, T.: Effects of thermal energy discharge on shallow groundwater ecosystems, *FEMS Microbiol. Ecol.*, 68(3), 273–286, doi:10.1111/j.1574-6941.2009.00674.x, 2009.
- 585 Briellmann, H., Lueders, T., Schreglmann, K., Ferraro, F., Avramov, M., Hammerl, V., Blum, P., Bayer, P. and Griebler, C.: Oberflächennahe Geothermie und ihre potenziellen Auswirkungen auf Grundwasserökosysteme, *Grundwasser*, 16(2), 77–91, doi:10.1007/s00767-011-0166-9, 2011.
- Camacho, A. I.: An annotated checklist of the Syncarida (Crustacea, Malacostraca) of the world, *Zootaxa*, 54, 1–54, 2006.
- Datry, T., Malard, F. and Gibert, J.: Response of invertebrate assemblages to increased groundwater recharge rates in a phreatic
590 aquifer, *J. North Am. Benthol. Soc.*, 24(3), 461–477, doi:10.1899/04-140.1, 2005.

- Deharveng, L., Stoch, F., Gibert, J., Bedos, A., Galassi, D., Zagamajster, M., Brancelj, A., Camacho, A., Fiers, F., Martin, P., Giani, N., Magniez, G. and Marmonier, P.: Groundwater biodiversity in Europe, *Freshw. Biol.*, 54(4), 709–726, doi:10.1111/j.1365-2427.2008.01972.x, 2009.
- 595 Dole-Olivier, M. J., Malard, F., Martin, D., Lefébure, T. and Gibert, J.: Relationships between environmental variables and groundwater biodiversity at the regional scale, *Freshw. Biol.*, 54(4), 797–813, doi:10.1111/j.1365-2427.2009.02184.x, 2009a.
- Dole-Olivier, M. J., Castellarini, F., Coineau, N., Galassi, D. M. P., Martin, P., Mori, N., Valdecasas, A. and Gibert, J.: Towards an optimal sampling strategy to assess groundwater biodiversity: Comparison across six European regions, *Freshw. Biol.*, 54(4), 777–796, doi:10.1111/j.1365-2427.2008.02133.x, 2009b.
- 600 Eckert, J., Friedhoff, K. T., Zahner, H. and Deplazes, P.: *Lehrbuch der Parasitologie für die Tiermedizin Teil II Parasiten und Parasitosen: 3 Metazoa*, 2nd ed., Thieme Verlagsgruppe Stuttgart/ Enke Verlag, Stuttgart., 2008.
- Einsle, U.: *Crustacea: Copepoda, Calanoida and Cyclopoida – Süßwasserfauna von Mitteleuropa*, 8/4–1., Gustav Fischer Verlag Stuttgart, Stuttgart., 1993.
- 605 Fakher el Abiari, A., Oulbaz, Z., Yacoubi-Khebiza, M., Coineau, N. and Boutin, C.: Etude expérimentale de la sensibilité comparée de trois crustacés stygobies vis-à-vis de diverses substances toxiques pouvant se rencontrer dans les eaux souterraines, in *Mémoires de Biospéologie*, pp. 167–181., 1998.
- Fillinger, L., Hug, K., Trimbach, A. M., Wang, H., Kellermann, C., Meyer, A., Bendinger, B. and Griebler, C.: The D-A-(C) index: A practical approach towards the microbiological-ecological monitoring of groundwater ecosystems, *Water Res.*, 163, 114902, doi:10.1016/j.watres.2019.114902, 2019.
- 610 Foulquier, A., Malard, F., Mermillod-Blondin, F., Montuelle, B., Dolédec, S., Volat, B. and Gibert, J.: Surface Water Linkages Regulate Trophic Interactions in a Groundwater Food Web, *Ecosystems*, 14(8), 1339–1353, doi:10.1007/s10021-011-9484-0, 2011.
- Fuchs, A.: *Erhebung und Beschreibung der Grundwasserfauna in Baden-Württemberg*, , 1–109 [online] Available from: <https://kola.opus.hbz-nrw.de/frontdoor/index/index/docId/175>, 2007.
- 615 Fuchs, A., Hahn, H. J. and Barufke, K. P.: *Grundwasser-Überwachungsprogramm - Erhebung und Beschreibung der Grundwasserfauna in Baden-Württemberg*, 2006.
- Galassi, D. M. P.: Groundwater copepods: Diversity patterns over ecological and evolutionary scales, *Hydrobiologia*, 453–454(1997), 227–253, doi:10.1023/A:1013100924948, 2001.
- Galassi, D. M. P., Huys, R. and Reid, J. W.: Diversity, ecology and evolution of groundwater copepods, *Freshw. Biol.*, 54(4), 691–708, doi:10.1111/j.1365-2427.2009.02185.x, 2009.
- 620 Geyer, O. F., Gwinner, M. P., Nitsch, E. and Simon, T.: *Geologie von Baden-Württemberg*, Schweizerbart Stuttgart., 2011.

- Gibert, J. and Deharveng, L.: Subterranean Ecosystems: A Truncated Functional Biodiversity, *Bioscience*, 52(6), 473, doi:10.1641/0006-3568(2002)052[0473:seatfb]2.0.co;2, 2002.
- Gibert, J., Culver, D. C., Dole-Olivier, M. J., Malard, F., Christman, M. C. and Deharveng, L.: Assessing and conserving groundwater biodiversity: Synthesis and perspectives, *Freshw. Biol.*, 54(4), 930–941, doi:10.1111/j.1365-2427.2009.02201.x, 625 2009.
- GISAT: Corine Land Cover European seamless vector database, 2016.
- Glatzel, T.: On the biology of *Parastenocaris phyllura* Kiefer 1938 (Copepoda: Harpacticoda, *Stygologia*, 5, 131–136, 1990.
- Griebler, C. and Avramov, M.: Groundwater ecosystem services: A review, *Freshw. Sci.*, 34(1), 355–367, doi:10.1086/679903, 2015.
- 630 Griebler, C., Stein, H., Hahn, H. J., Steube, C., Kellefmann, C., Fuchs, A., Berkhoff, S. and Brielmann, H.: Entwicklung biologischer Bewertungsmethoden und -kriterien für Grundwasserökosysteme, Umweltbundesamt., 2014.
- Griebler, C., Brielmann, H., Haberer, C. M., Kaschuba, S., Kellermann, C., Stumpp, C., Hegler, F., Kuntz, D., Walker-Hertkorn, S. and Lueders, T.: Potential impacts of geothermal energy use and storage of heat on groundwater quality, biodiversity, and ecosystem processes, *Environ. Earth Sci.*, 75(20), 1–18, doi:10.1007/s12665-016-6207-z, 2016.
- 635 Gutjahr, S.: Grundwasserlebensräume in der Landschaft - Untersuchungen zur Bedeutung von Hydrologie und Hydrogeologie für Grundwasserlebensgemeinschaften, Universität Koblenz-Landau., 2013.
- Gutjahr, S., Schmidt, S. I. and Hahn, H. J.: A proposal for a groundwater habitat classification at local scale, *Subterr. Biol.*, 14(1), 25–49, doi:10.3897/subtblol.14.5429, 2014.
- Hahn, H. J.: Die Ökologie der Sedimente eines Buntsandsteinbaches im Pfälzerwald unter besonderer Berücksichtigung der 640 Ostracoden und Harpacticoiden (Crustacea), 62nd ed., Tectum-Verlag, Marburg., 1996.
- Hahn, H. J.: Unbaited phreatic traps: A new method of sampling stygofauna, *Limnologia*, 35(4), 248–261, doi:10.1016/j.limno.2005.04.004, 2005.
- Hahn, H. J.: A first approach to a quantitative ecological assessment of groundwater habitats: The GW-Fauna-Index, *Limnologia*, 36(2), 119–137, 2006.
- 645 Hahn, H. J. and Fuchs, A.: Distribution patterns of groundwater communities across aquifer types in south-western Germany, *Freshw. Biol.*, 54(4), 848–860, doi:10.1111/j.1365-2427.2008.02132.x, 2009.
- Hahn, H. J. and Gutjahr, S.: Bioindikation im Grundwasser funktioniert – Erwiderung zum Kommentar von T. Scheytt zum Beitrag „Grundwasserfauna als Indikator für komplexe hydrogeologische Verhältnisse am westlichen Kaiserstuhl“ von Gutjahr, S., Bork, J. & Hahn, H.J. in *Grundwasser* 18, *Grundwasser*, 19(3), 215–218, doi:10.1007/s00767-014-0266-4, 2014.
- 650 Hahn, H. J. and Matzke, D.: A comparison of stygofauna communities inside and outside groundwater bores, *Limnologia*, 35,

31–44, 2005.

Hahn, H. J., Matzke, D., Kolberg, A. and Limberg, A.: Untersuchung zur Fauna des Berliner Grundwassers – erste Ergebnisse, Berlin., 2013.

655 Hahn, H. J., Schweer, C. and Griebler, C.: Are groundwater ecosystem rights being preserved?: A critical evaluation of the legal background of groundwater ecosystems, *Grundwasser*, 23(3), 209–218, doi:10.1007/s00767-018-0394-3, 2018.

Hähnlein, S., Bayer, P. and Blum, P.: International legal status of the use of shallow geothermal energy, *Renew. Sustain. Energy Rev.*, 14(9), 2611–2625, doi:10.1016/j.rser.2010.07.069, 2010.

Hähnlein, S., Bayer, P., Ferguson, G. and Blum, P.: Sustainability and policy for the thermal use of shallow geothermal energy, *Energy Policy*, 59, 914–925, doi:10.1016/j.enpol.2013.04.040, 2013.

660 Hancock, P. J., Boulton, A. J. and Humphreys, W. F.: Aquifers and hyporheic zones: Towards an ecological understanding of groundwater, *Hydrogeol. J.*, 13(1), 98–111, doi:10.1007/s10040-004-0421-6, 2005.

Herrmann, J.: Dependence of Reproduction in *Dendrocoelum lacteum* (Turbellaria): An Experimental Approach, *Oikos*, 44(2), 268–272, 1985.

665 Hunkeler, D., Goldscheider, N., Rossi, P., Burn, C.: Biozönosen im Grundwasser - Grundlagen und Methoden der Charakterisierung von mikrobiellen Gemeinschaften, Bundesamt für Umwelt (BAFU) Umwelt-Wissen 0603, Bern., 2006.

Issartel, J., Hervant, F., Voituron, Y., Renault, D. and Vernon, P.: Behavioural, ventilatory and respiratory responses of epigean and hypogean crustaceans to different temperatures, *Comp. Biochem. Physiol., Part A: Mol*(1), 1–7, 2005.

Janetzka, W., Enderle, R. and Noodt, W.: Crustacea: Copepoda: Gelyelloida and Harpacticoida – Süßwasserfauna von Mitteleuropa, 8/4–2., Gustav Fischer Verlag Stuttgart, Stuttgart., 1996.

670 Korbel, K., Chariton, A., Stephenson, S., Greenfield, P. and Hose, G. C.: Wells provide a distorted view of life in the aquifer: Implications for sampling, monitoring and assessment of groundwater ecosystems, *Sci. Rep.*, 7(July 2016), 1–14, doi:10.1038/srep40702, 2017.

Korbel, K. L. and Hose, G. C.: A tiered framework for assessing groundwater ecosystem health, *Hydrobiologia*, 661(1), 329–349, doi:10.1007/s10750-010-0541-z, 2011.

675 Korbel, K. L. and Hose, G. C.: Habitat, water quality, seasonality, or site? Identifying environmental correlates of the distribution of groundwater biota, *Freshw. Sci.*, 34(1), 329–342, doi:10.1086/680038, 2015.

Korbel, K. L. and Hose, G. C.: The weighted groundwater health index: Improving the monitoring and management of groundwater resources, *Ecol. Indic.*, 75, 164–181, doi:10.1016/j.ecolind.2016.11.039, 2017.

Kühlers, D., Maier, M. and Roth, K.: Sanierung im Verborgenen, *TerraTech Sanierungspraxis*, 3, 14–16, 2012.

- 680 Kunkel, R., Wendland, F. and Hannappel, S.: Die natürliche, ubiquitär überprägte Grundwasserbeschaffenheit in Deutschland, 47th ed., Schriften des Forschungszentrums Jülich (Forschungszentrum Jülich GmbH), 2004.
- Di Lorenzo, T. and Galassi, D. M. P.: Agricultural impact on Mediterranean alluvial aquifers: Do groundwater communities respond?, *Fundam. Appl. Limnol.*, 182(4), 271–282, doi:10.1127/1863-9135/2013/0398, 2013.
- Di Lorenzo, T., Fiasca, B., Di Cicco, M. and Galassi, D. M. P.: The impact of nitrate on the groundwater assemblages of
 685 European unconsolidated aquifers is likely less severe than expected, *Environ. Sci. Pollut. Res.*, 10, doi:10.1007/s11356-020-11408-5, 2020a.
- Di Lorenzo, T., Fiasca, B., di Camillo Tabilio, A., Murolo, A., di Cicco, M. and Galassi, D. M. P.: The weighted Groundwater Health Index (wGHI) by Korbel and Hose (2017) in European groundwater bodies in nitrate vulnerable zones, *Ecol. Indic.*, 116 [online] Available from: <https://doi.org/10.1016/j.ecolind.2020.106525>, 2020b.
- 690 Malard, F., Dole-Olivier, M.-J., Mathieu, J., Stoch, F., Boutin, C., Brancelj, A., Camacho, A. I., Fiers, F., Galassi, D., Gibert, J., Lefebure, T., Martin, P., Sket, B. and Valdecasas, A. G.: Sampling Manual for the Assessment of Regional Groundwater Biodiversity, Tech. Rep. Eur. Proj. PASCALIS, 2002.
- Matzke, D.: Untersuchungen zum Verhalten von Grundwasserfauna in Altlastflächen mit vorangegangenem Vergleich unterschiedlicher Sammeltechniken., 2006.
- 695 Meisch, C.: Freshwater Ostracoda of Western and Central Europe - Süßwasserfauna von Mitteleuropa, 8/3., Spektrum Akademischer Verlag, Heidelberg., 2000.
- Menberg, K., Blum, P., Schaffitel, A. and Bayer, P.: Long-term evolution of anthropogenic heat fluxes into a subsurface urban heat island, *Environ. Sci. Technol.*, 47(17), 9747–9755, doi:10.1021/es401546u, 2013a.
- Menberg, K., Bayer, P., Zosseder, K., Rumohr, S. and Blum, P.: Subsurface urban heat islands in German cities, *Sci. Total*
 700 *Environ.*, 442, 123–133, doi:10.1016/j.scitotenv.2012.10.043, 2013b.
- Moon, K. R., van Dijk, D., Wang, Z., Gigante, S., Burkhardt, D. B., Chen, W. S., Yim, K., Elzen, A. van den, Hirn, M. J., Coifman, R. R., Ivanova, N. B., Wolf, G. and Krishnaswamy, S.: Visualizing structure and transitions in high-dimensional biological data, *Nat. Biotechnol.*, 37(12), 1482–1492, doi:10.1038/s41587-019-0336-3, 2019.
- Mösslacher, F. and Notenboom, J.: Groundwater biomonitoring, in *Biomonitoring of polluted water*, edited by A. Gerhardt,
 705 pp. 119–139, Trans Tech Publications Ltd, Zurich-Uetikon., 2000.
- Preuß, G. and Schminke, H. K.: Ein globales ökosystem: Grundwasser lebt!, *Chemie Unserer Zeit*, 38(5), 340–347, doi:10.1002/ciuz.200400307, 2004.
- Regierungspräsidium Freiburg: LGRB-Kartenviewer – Layer GK50: Geologische Einheiten (Flächen), [online] Available from: <https://maps.lgrb-bw.de/> (Accessed 6 July 2020), 2019.

- 710 Di Sabatino, A., Gerecke, R. and Martin, P.: The biology and ecology of lotic water mites (Hydrachnidia), *Freshw. Biol.*, 44(1), 47–62, 2000.
- Sánchez, A. C., Hose, G. C. and Reboleira, A. S. P. S.: Salinity and temperature increase impact groundwater crustaceans, *Sci. Rep.*, (0123456789), 1–9, doi:10.1038/s41598-020-69050-7, 2020.
- Sauermost, R. and Freudig, D.: *Bathynellacea*, Spektrum Akad. Verlag. Heidelb. [online] Available from:
715 <https://www.spektrum.de/lexikon/biologie/bathynellacea/7445> (Accessed 27 February 2019a), 1999.
- Sauermost, R. and Freudig, D.: *Oligochaeta*, Spektrum Akad. Verlag. Heidelb. [online] Available from:
<https://www.spektrum.de/lexikon/biologie/oligochaeta/47593> (Accessed 27 February 2019b), 1999.
- Sauermost, R. and Freudig, D.: *Strudelwürmer*, Spektrum Akad. Verlag. Heidelb. [online] Available from:
<https://www.spektrum.de/lexikon/biologie/strudelwuermer/64369> (Accessed 27 February 2019c), 1999.
- 720 Schellenberg, A.: *Krebstiere oder Crustace, IV: Flohkebs oder Amphipoda*, in *Die Tierwelt Deutschlands und der angrenzenden Meeresteile nach ihren Merkmalen und nach ihrer Lebensweise*, p. 252, Gustav Fischer Verlag Jena., 1942.
- Scheytt, T.: Kommentar zur Veröffentlichung von Gutjahr, S., Bork, J. und Hahn, H.J.: Grundwasserfauna als Indikator für komplexe hydrogeologische Verhältnisse am westlichen Kaiserstuhl in *Grundwasser* 18 (3), 173–184 (2013), *Grundwasser*, 19(3), 211–213, doi:10.1007/s00767-014-0267-3, 2014.
- 725 Schmidt, S. I., Hahn, H. J., Hatton, T. J. and Humphreys, W. F.: Do faunal assemblages reflect the exchange intensity in groundwater zones?, *Hydrobiologia*, 583(1), 1–19, doi:10.1007/s10750-006-0405-8, 2007.
- Schminke, H. K., Grad, G., Ahlrichs, W., Bartsch, I., Christl, H., Gerecke, R., Martin, P., Rumm, P. and Wägele, J. W.: *Grundwasserfauna Deutschlands - Ein Bestimmungswerk: DWA-Themen*, 1st ed., Deutsche Vereinigung für Wasserwirtschaft, Abwasser und Abfall., 2007.
- 730 Schönthaler, K. and von Adrian-Werburg, S.: *Erster integrierter Umweltbericht für das länderübergreifende UNESCO-Biosphärenreservat Rhön*, Bayerisches Staatsministerium für Umwelt, Gesundheit und Verbraucherschutz (BayStMUGV) Hessisches Ministerium für Umwelt, ländlichen Raum und Verbraucherschutz (HMULV) Thüringer Ministerium für Landwirtschaft, Naturschutz und Umwelt (TMLNU)., 2008.
- Smith, R. J., Paterson, J. S., Launer, E., Tobe, S. S., Morello, E., Leijs, R., Marri, S. and Mitchell, J. G.: Stygofauna enhance
735 prokaryotic transport in groundwater ecosystems, *Sci. Rep.*, 6(September), 1–7, doi:10.1038/srep32738, 2016.
- Spengler, C.: *Die Auswirkungen von anthropogenen Temperaturerhöhungen auf die Crustaceagemeinschaften im Grundwasser*, Universität Koblenz-Landau., 2017.
- Spengler, C. and Hahn, J.: *Thermostress : Ökologisch begründete , thermische Schwellenwerte und Be- wertungsansätze für das Grundwasser*, *Korrespondenz Wasserwirtschaft Fachbeiträge Gewässer und Böden*, 11(9), 521–525,

740 doi:10.3243/kwe2018.09.001, 2018.

Stadt Karlsruhe: Bodenschutz- und Altlastenkataster der Stadt Karlsruhe, [online] Available from: https://www.karlsruhe.de/b3/natur_und_umwelt/umweltschutz/altlasten.de (Accessed 23 October 2019), 2006.

Stein, H., Kellermann, C., Schmidt, S. I., Brielmann, H., Steube, C., Berkhoff, S. E., Fuchs, A., Hahn, H. J., Thulin, B. and Griebler, C.: The potential use of fauna and bacteria as ecological indicators for the assessment of groundwater quality, J. Environ. Monit., 12(1), 242–254, doi:10.1039/b913484k, 2010.

Stein, H., Griebler, C., Berkhoff, S., Matzke, D., Fuchs, A. and Hahn, H. J.: Stygoregions-a promising approach to a bioregional classification of groundwater systems, Sci. Rep., 2, 1–9, doi:10.1038/srep00673, 2012.

Stoch, F., Artheau, M., Brancelj, A., Galassi, D. M. P. and Malard, F.: Biodiversity indicators in European ground waters: Towards a predictive model of stygobiotic species richness, Freshw. Biol., 54(4), 745–755, doi:10.1111/j.1365-2427.2008.02143.x, 2009.

Taylor, C. A. and Stefan, H. G.: Shallow groundwater temperature response to climate change and urbanization, J. Hydrol., 375(3–4), 601–612, doi:10.1016/j.jhydrol.2009.07.009, 2009.

Technologiezentrum Wasser: Grundwasserdatenbank Wasserversorgung: Regionale Auswertung - Region Mittlerer Oberrhein, [online] Available from: <http://www.grundwasserdatenbank.de/regionmo.htm> (Accessed 27 February 2019), 2018.

755 Tissen, C., Menberg, K., Bayer, P. and Blum, P.: Heat supply by shallow geothermal energy in Karlsruhe, in Groundwater in the surrounding of mining, energy and urban space, Conference of the professional division Hydrogeology in the DGGV, Bochum., 2018.

Tissen, C., Benz, S. A., Menberg, K., Bayer, P. and Blum, P.: Groundwater temperature anomalies in central Europe, Environ. Res. Lett., 14(10), 104012, doi:10.1088/1748-9326/ab4240, 2019.

760 Ward, J. V. and Tockner, K.: Biodiversity: Towards a unifying theme for river ecology, Freshw. Biol., 46(6), 807–819, doi:10.1046/j.1365-2427.2001.00713.x, 2001.

Wickert, F., Müller, A., Schäfer, W. and Tiehm, A.: Vergleich hochauflösender Grundwasserprobennahmeverfahren zur Charakterisierung der vertikalen LCKW-Verteilung im Grundwasserleiter, Altlastenspektrum, 01, 29–35, 2006.

Więcek, M., Martin, P. and Gąbka, M.: Distribution patterns and environmental correlates of water mites (Hydrachnidia, Acari) 765 in peatland microhabitats, Exp. Appl. Acarol., 61(2), 147–160, doi:10.1007/s10493-013-9692-8, 2013.

Wirsing, G. and Luz, A.: Hydrogeologischer Bau und Aquifereigenschaften der Lockergesteine im Oberrheingraben (Baden Württemberg), LGRB-Informationen, 19, 130, 2007.

Zenker, S., Bogon, K. and Weigand, A.: Die Höhlentiere Deutschlands, Quelle & Meyer Verlag, Wiebelsheim., 2020.

Zhu, K., Blum, P., Ferguson, G., Balke, K. D. and Bayer, P.: The geothermal potential of urban heat Islands, Environ. Res.

770 Lett., 6(1), doi:10.1088/1748-9326/6/1/019501, 2010.

Zuurbier, K. G., Hartog, N., Valstar, J., Post, V. E. A. and Van Breukelen, B. M.: The impact of low-temperature seasonal aquifer thermal energy storage (SATES) systems on chlorinated solvent contaminated groundwater: Modeling of spreading and degradation, J. Contam. Hydrol., 147, 1–13, doi:10.1016/j.jconhyd.2013.01.002, 2013.