

Spatio-temporal controls of C-N-P dynamics across headwater catchments of a temperate agricultural region from public data analysis

5 Stella Guillemot^{1,2}, Ophelie Fovet¹, Chantal Gascuel-Oudou¹, Gérard Gruau³, Antoine Casquin¹, Florence Curie², Camille Minaudo⁴, Laurent Strohmenger¹, and Florentina Moatar^{5,2}

¹INRAE, AGROCAMPUS OUEST/INSTITUT AGRO, UMR SAS, 35000 Rennes, France

²Université de Tours, EA 6293 GéHCO, 37200 Tours, France

³OSUR, Geosciences Rennes, CNRS, Université Rennes 1, 35000 Rennes, France

⁴EPFL, Physics of Aquatic Systems Laboratory, 1015 Lausanne, Switzerland

10 ⁵INRAE, RIVERLY, 69625 Villeurbanne, France

Correspondence to: Ophelie Fovet (ophelie.fovot@inrae.fr)

Abstract. Characterizing and understanding spatial variability in water quality for a variety of chemical elements is an issue for present and future water resource management. However, most studies of spatial variability in water quality focus on a single element and rarely consider headwater catchments. Moreover, they assess few catchments and focus on annual means without considering seasonal variations. To overcome these limitations, we studied spatial variability and seasonal variation in dissolved C, N, and P concentrations at the scale of an intensively farmed region of France (Brittany). We analyzed 185 headwater catchments (from 5-179 km²) for which 10-year time series of monthly concentrations and daily stream flow were available from public databases. We calculated interannual loads, concentration percentiles, and seasonal metrics for each element to assess their spatial patterns and correlations. We then performed rank correlation analyses between water quality, human pressures, and soil and climate features. Results show that nitrate (NO₃) concentrations increased with increasing agricultural pressures and base flow contribution; dissolved organic carbon (DOC) concentrations decreased with increasing rainfall, base flow contribution, and topography; and soluble reactive phosphorus (SRP) concentrations showed weaker positive correlations with diffuse and point sources, rainfall and topography. An opposite pattern was found between DOC and NO₃: spatially, between their median concentrations, and temporally, according to their seasonal cycles. In addition, the quality of annual maximum NO₃ concentration was in-phase with maximum flow when the base flow index was low, but this synchrony disappeared when flow flashiness was lower. These DOC-NO₃ seasonal cycle types were related to the mixing of flowpaths combined with the spatial variability of their respective sources and to local biogeochemical processes. The annual maximum SRP concentration occurred during the low-flow period in nearly all catchments. This likely resulted from the dominance of P point sources. The approach shows that despite the relatively low frequency of public water quality data, such databases can provide consistent pictures of the spatio-temporal variability of water quality and of its drivers as soon as they contain a large number of catchments to compare and a sufficient length of concentration time series.

15
20
25
30

1 Introduction

As a condition for human health, food production, and ecosystem functions, water quality is recognized as “one of the main challenges of the 21st century” (FAO and WWC, 2015; UNESCO, 2015), and potential impacts of climate change on water quality are even more challenging (Whitehead et al., 2009). To better estimate and reduce human impact on water quality, water scientists are expected to provide integrated understanding of multiple pollutants (Cosgrove and Loucks, 2015). Eutrophication risks (Dodds and Smith, 2016) are considered the main factors that decrease the quality of surface water, according to objectives set by the European Union Water Framework Directive. Mitigating the problem of eutrophication involves considering at least the three major elements: carbon (C), nitrogen (N), and phosphorus (P) (Le Moal et al., 2019). In addition, the quality of headwater catchments have been studied less than large rivers (Bishop et al., 2008), despite their influence on downstream water quality (Alexander et al., 2007; Barnes and Raymond, 2010; Bol et al., 2018) and higher spatial variability in their concentrations (Abbott et al., 2018a; Temnerud and Bishop, 2005). One reason for this is that most water quality monitoring networks coincide with the location of drinking-water production facilities, which explains why they focus on large rivers. Nonetheless, investigating spatial variability in upstream water quality is relevant for understanding what causes it to degrade, targeting locations with the greatest disturbances, and identifying which remediation policies would be most cost effective.

In non-agricultural headwater catchments, spatial variability in dissolved organic C (DOC) concentrations in streams has been related to topography, wetland coverage, and soil properties such as clay content or pH (Andersson and Nyberg, 2008; Brooks et al., 1999; Creed et al., 2008; Hytteborn et al., 2015; Musolff et al., 2018; Temnerud and Bishop, 2005; Zarnetske et al., 2018). Stream DOC concentrations and composition in agricultural and urbanized areas also generally differ greatly from those in semi-natural or pristine catchments (Graeber et al., 2012; Gücker et al., 2016). Over large gradients of human impact (e.g. from undisturbed to urban catchments), the cover of agricultural and urban land uses often appears as a key factor that explains differences in stream chemistry of C, N, and P species (e.g. Barnes and Raymond, 2010; Edwards et al., 2000; Mutema et al., 2015) and even silica (Onderka et al., 2012). Conversely, in more homogeneous catchments – e.g. mostly undisturbed (Mengistu et al., 2014) or mostly rural (Heppell et al., 2017; Lintern et al., 2018) – “natural” controls such as topography, geology, and flow paths are more frequently highlighted as the main factors that explain spatial variability in C, N and P.

Besides being spatially variable, C, N, and P concentrations also vary temporally. The variability of concentrations with flow has been described in several studies using concentration-flow relationships at event (Fasching et al., 2019) or inter-annual to long-term scales (Basu et al., 2010; 2011; Moatar et al., 2017). Concentrations also vary seasonally in streams and rivers (Aubert et al., 2013; Dawson et al., 2008; Duncan et al., 2015; Exner-Kittridge et al., 2016; Lambert et al., 2013), as does the composition of dissolved organic matter (Griffiths et al., 2011; Gücker et al., 2016). This seasonality can also be spatially structured. Several studies showed that the relative importance of catchment characteristics on water concentrations or loads varied by season because nutrient sources and biological and physico-chemical processes that influence nutrient mobilization and transfer in catchments (e.g. vegetation uptake, in-stream biomass production, denitrification) changed with the

hydrological conditions (Ågren et al., 2007; Fasching et al., 2016; Gardner and McGlynn, 2009). Some variability in seasonal patterns of dissolved C, N, and/or P concentrations among headwater catchments has been reported (e.g. Van Meter et al., 2019; Abbott et al., 2018b; Duncan et al., 2015; Martin et al., 2004). Identifying these patterns is relevant from a management viewpoint as they may indicate changes in the locations of C, N, or P sources or their transfer pathways.

70

Thus, to date, analysis of spatial variability in water quality at the headwater scale:

- 1) is usually restricted to one element, although multi-element approaches are becoming more frequent (Edwards et al., 2000; Heppell et al., 2017; Lintern et al., 2018; Mengistu et al., 2014; Mutema et al., 2015),
- 2) is particularly rare for headwater catchments with similar human pressures (e.g. intensive farming), despite the high variability in water quality sometimes observed among them (e.g. Thomas et al., 2014),
- 3) often uses mean annual values (concentration or load) to describe spatial variability in water quality among catchments, with little or no analysis of seasonal patterns despite their frequent occurrence (Van Meter et al., 2019; Abbott et al., 2018b; Liu et al., 2014; Halliday et al., 2012; Mullholland et al. 1997), and
- 4) is usually restricted to a few catchments: multiple-catchment studies on multiple elements are uncommon, despite their ability to identify dominant controlling factors better.

75

80

We studied the spatial variability and seasonal variation in water quality of 185 headwater catchments (from 5-179 km²) draining Brittany, an intensively farmed region of France. Our analysis focuses on dissolved C, N, and P concentrations as DOC, nitrate (NO₃), and soluble reactive P (SRP), respectively. We hypothesized that:

- 1) Human (i.e. rural and urban) pressures determine spatial variability in NO₃ and SRP concentrations (Preston et al., 2011; Melland et al., 2012; Dupas et al., 2015a; Kaushal et al., 2018), while soil and climate characteristics determine that in DOC and possibly SRP (Lambert et al., 2013; Humbert et al., 2015; Gu et al., 2017).
- 2) Seasonal variations in water quality provide information about spatial variability in biogeochemical sources and/or reactivity in catchments as a function of changes in water pathways and are correlated in part with spatial variability in concentrations and loads.

85

90

We selected headwater catchments for which relevant time series of DOC, NO₃, and SRP concentrations and stream flow were available (10 years of consecutive data measured at least monthly). In addition to estimating interannual loads, we calculated concentration metrics for each element to assess the spatial variability and temporal variation in water quality. Generalized Additive Models (GAMs) were applied to the time series to highlight average patterns of seasonal variation. Correlations between the water quality metrics and the geological, soil, climatic, hydrological, land cover, and human pressure characteristics of the corresponding headwater catchments were evaluated using rank correlation analyses.

95

2 Materials and Methods

2.1 Study area

Brittany is a 27,208 km² region in western France. Its bedrock is composed mainly of a crystalline substratum dominated by granite and schist (Supplement S1b). Its topography is moderate, with elevation ranging from 0-330 m a.s.l. Its climate is temperate oceanic, with precipitation ranging from 531 mm.yr⁻¹ in the east to 1070 mm.yr⁻¹ on the western coasts (regional median of 723.0 mm.yr⁻¹) (S1a), and a mean annual temperature of 12°C. The regional hydrographic network is dense, with a mean density of 1 km.km⁻². Its intensive agriculture has a strong influence on land use and agri-food production. Overall, 56.6% of the region was Utilized Agricultural Area (UAA) in 2017 (data from DREAL Bretagne, Brittany's Agency for Environment, Infrastructure, and Housing), which represented 6% of national UAA in 2016. Of total French production, Brittany produces 17.4% of milk and dairy products, 20% of pork products, and 17% of eggs and poultry (Brittany Chamber of Agriculture, 2016 data). At the canton (administrative district) scale, mean N and P surpluses are high and have high spatial variability (standard deviation (SD)): 50.01 ± 26.59 kg N.ha⁻¹.yr⁻¹ and 22.52 ± 12.66 kg P.ha⁻¹.yr⁻¹ (Supplement S1e,f). The region has a population of ca 3.3 million inhabitants (data 2017), some scattered throughout the region, and some concentrated in a few cities and near the coasts (Supplement S1c,d).

2.2 Stream data selection and headwater characteristics

Water quality data consisted of time series of DOC, NO₃, and SRP concentrations, extracted from two public monitoring networks – OSUR (Loire-Brittany Water Agency, 554 sites) and HYDRE/BEA (DREAL Bretagne, ca. 1964 sites), measured for regulatory monitoring, regional contracts, or specific programs. Concentrations were measured from grab samples. Headwater catchments were selected according to the following two criteria: (i) independence, with no overlap of the drained areas of the water-quality stations selected, and (ii) availability of at least 80 measurements of DOC, NO₃, and SRP concentrations at the same station (after removing outliers based on expert knowledge, i.e. values > 200 mg N.L⁻¹ or 5 g P.L⁻¹) over 10 calendar years (2007-2016). We selected 185 stations (83% and 17% from OSUR and HYDRE/BEA, respectively) (hereafter, “concentration (C) stations”), which had mean frequencies of 12, 14, and 11 analyses per year for DOC, NO₃, and SRP, respectively. We checked that there was no bias in the timing of concentration data: OSUR database has fixed and regular sampling frequencies while we noticed a few time series where summer periods were less sampled in the HYDRE/BEA data for some years only.

Each C station was paired with a hydrometric station (Q). Observed daily streamflow data from the national hydrometric network (<http://hydro.eaufrance.fr/>) were used when draining headwater catchments for C and Q stations shared at least 80% of their areas (25% of cases). When observed Q data were not available, or at a frequency less than 320 measurements per year from 2007-2016 (75% of cases), discharge data were simulated using the GR4J model (Perrin et al., 2003). The headwater catchments selected and their associated C and Q stations were distributed throughout Brittany (Fig. 1).

The 185 headwater catchments selected cover ca. 32% of Brittany's area. Despite having a similar hydrographic context dominated by subsurface flow, the catchments have large differences in topography, geology, hydrology, and diffuse and point-source pressures of N and P. We used a set of catchment descriptors to quantify this variability (Table 1) (see Supplemental S2 for their statistical distribution and S3 for their correlations). The descriptors selected included a set of spatial metrics for element sources (e.g. land use, pressure, soil contents) and for mobilization and retention processes (e.g. hydrology, climate, topography, geology, and soil properties).

The headwater catchments range in area from 5-179 km² (median of 38 km²), and the density of each one's hydrographic network ranges from 0.47-1.49 km.km⁻² (median of 0.90 km.km⁻²). Strahler stream order is 3 for 36% of the catchments, 2 for 18%, 4 for 17%, and 1 for 11%. Substrate composition is dominated by schists/micaschists (44%) or granites/gneisses (31%). In the topsoil horizon (0-30 cm), the soil organic C content varies greatly from 18.6-565.4 g.kg⁻¹ (median of 126.9 g.kg⁻¹), while the total P (Dyer method) content varies from 0.6-1.4 g.kg⁻¹ (median of 0.9 g.kg⁻¹). Land use is largely agricultural, although some catchments have high percentages of forested and urbanized areas. Riparian wetlands cover 12.3-36.3% of catchment area (median of 22.4%), forest covers 1.3-55.7% (median of 13.2%), pasture covers 10.3-46.7% (median of 25.6%), summer crops cover 6.5-50.3% (median of 27.8%), and winter crops cover 7.0-51.0% (median of 22.7%). The N and P surplus (potential diffuse agricultural sources) vary from 12.9-96.0 kg N.ha⁻¹.yr⁻¹ (median of 47.7) and 2.8-63.2 kg P.ha⁻¹.yr⁻¹ (median of 18.9), respectively. Urban areas cover 1.3-31.8% of the headwater catchments (median of 6%), with point-source input estimates ranging from 0-6.2 kg N. ha⁻¹.yr⁻¹ and 0-0.626 kg P. ha⁻¹.yr⁻¹. These data illustrate relative diversity in human pressures among the catchments despite a regional context of intensive agriculture. The daily mean flow (Q_{mean}) varies from 4.8-24.5 l.s⁻¹.km⁻² (median of 10.8 l.s⁻¹.km⁻²), the median of annual minimum of monthly flows (QMNA) varies from 0.2-5.9 l.s⁻¹.km⁻², and the flow flashiness index (W2), defined as the percentage of total discharge that occurs during the highest 2% of flows (Moatar et al., 2020), ranges from 10-28%.

150

2.3 Data analysis

2.3.1 Concentration and load metrics

To analyze spatial variability in DOC, NO₃, and SRP concentrations in streams, we calculated their 10th, 50th, and 90th percentiles of concentration (C₁₀, C₅₀, and C₉₀, respectively) for each headwater catchment from 2007-2016. We also calculated the ratio of the coefficient of variation (CV) of mean concentration (CV_{C_{mean}}) and to that of mean flow (CV_{Q_{mean}}) to compare spatial variabilities in concentrations and stream flow. We estimated interannual loads for a 10-year period (2007-2016), with 8-12 C-Q values per year. However, a 5-year period (2010-2014) was considered to analyze the spatial variability because it minimized data gaps (in C and Q time series) among all stations simultaneously.

To calculate interannual DOC, NO₃, and SRP loads for each headwater catchment, we tested different methods and selected the most suitable, depending on the reactivity of the element with flow. When C-Q relationships were relatively flat or diluted

160

(NO₃) or slowly mobilized (DOC) during high flow (Q>Q₅₀), we used the discharge weighted concentration (DWC) method (Eq. 1), which estimates loads with lower uncertainties (Moatar and Meybeck, 2007; Raymond et al., 2013):

$$DWC = \frac{k}{A} \times \frac{\sum_{i=1}^n C_i Q_i}{\sum_{i=1}^n Q_i} \bar{Q} \quad (1)$$

165 where DWC is the mean of annual loads (kg.y⁻¹.ha⁻¹), C_i is the instantaneous concentration (mg.l⁻¹), Q_i is the corresponding flow rate (m³.s⁻¹), \bar{Q} is the mean annual flow rate calculated from daily data (m³.s⁻¹), A is the area of the headwater catchment (m²), k is a conversion factor (31557.6), and n is the number of C-Q pairs per year.

The loads estimated by the DWC method were corrected for bias. Precisions were calculated from the number of samples (n), number of years, export regime exponent (b_{50high}), and W2 (Moatar et al., 2020).

170 To calculate SRP loads, regression methods were more suitable (because of strong concentration patterns when stream flow increases). We averaged the loads estimated by two regression methods developed by Raymond et al. (2013) – Integral Regression Curve (IRC) and Segmented Regression Curve (SRC) – both based on a regression between concentration and flow:

$$IRC = \frac{k'}{A} \times \sum_{i=1}^n C_i Q_i \quad (2)$$

175
$$SRC = \frac{k'}{A} \times (\sum_{i=1}^n C_{inf} Q_i + \sum_{i=1}^n C_{sup} Q_i) \quad (3)$$

where IRC and SRC are the mean of annual loads (kg.y⁻¹.ha⁻¹); C_i, C_{sup}, and C_{inf} are instantaneous concentrations estimated by the regression curves (mg.l⁻¹); C_{sup} and C_{inf} are concentrations of flows above and below the median flow, respectively; and k' is a conversion factor (86.4).

180

2.3.2 Seasonal signal

Seasonal dynamics of discharge and solute concentrations were modeled using GAMs (Wood, 2017), which can estimate smoothed seasonal dynamics from time series (Musolff et al., 2017). The smoothing function was a cyclic cubic spline fitted to the month of the year (1-12); thus, the ends of the spline were forced to be equal, using the R package mgcv. We did not
 185 consider a long-term trend in the time series over the 10 years, for two reasons. First, significant long-term trends (according to Man-Kendall tests) had low slopes: mean Theil-Sen slopes ranged from -3% to 0% of the median concentration (while mean seasonal relative amplitudes exceeded 50%). Second, performance of the GAMs did not increase significantly when a long-term trend was added: the mean adjusted coefficient of determination (Rsq) increased from 0.16 to 0.18 for DOC and from 0.30 to 0.40 for NO₃. We considered a seasonal dynamic to exist when the GAM adjusted coefficient of determination was
 190 greater than 0.10.


Seasonal dynamics of the concentrations of the three solutes (DOC, NO₃, and SRP) and river discharge were then analyzed using five metrics calculated from the daily simulations of the GAMs. The first three were the annual amplitude (Ampli; i.e. annual maximum minus annual minimum), and the mean time in which annual maximum and minimum concentrations occurred (MaxPhase and MinPhase, respectively; in months from 1 January). The next was Ampli standardized by the corresponding mean concentration to compare the three solutes. The last metric was a seasonality index (SI), which measures the relative importance of summer (1 June to 31 July) concentrations compared to winter (15 January to 15 March) concentrations of an element, as follows (Eq. 4):

$$SI = \frac{C_{\text{winter}} - C_{\text{summer}}}{C_{\text{winter}} + C_{\text{summer}}} \quad (4)$$

where C_{winter} and C_{summer} are the averages of winter and summer concentrations, calculated from daily values from fitted GAM. Positive values of SI (near 1) indicate that $C_{\text{winter}} > C_{\text{summer}}$, while negative values (near -1) indicate that $C_{\text{winter}} < C_{\text{summer}}$. We considered that SI values close to 0 (from -0.1 to 0.1) indicated that C_{winter} equaled C_{summer} . The SI integrates both amplitude and phasing features of the seasonal signal.

2.3.2 Statistical analyses

To compare the concentration metrics of the elements, a multivariate analytical approach, principal component analysis (PCA), was performed for the 9 variables of concentration percentiles (C10, C50, and C90) of DOC, NO₃, and SRP for the dataset of 185 headwater catchments. To identify dominant drivers of spatial variability in concentration percentiles, seasonality, and loads of DOC, NO₃, and SRP, we calculated Spearman's rank correlation (r_s) between these water-quality metrics and the descriptors of the headwater catchments. We considered a rank correlation to be significant if the corresponding p-value was ≤ 0.05 . All analyses were performed using R software (v. 3.6.1) with packages mgcv, hydroGOF, hydrostats, FactoMineR, tidyverse, lubridate, reshape2, plyr, ggcorrplot, and ggplot2 (Grolemund and Wickham, 2011; Le et al., 2008; Wickham, 2016, 2011; Wood, 2017; Zambrano-Bigiarini, 2020).



215

Figure 1. Locations of the 185 study headwater catchments where dissolved organic carbon, nitrate, and soluble reactive phosphorus concentrations were monitored monthly at the outlet from 2007-2016, and paired discharge stations where daily records of stream flow were available from observations or modeling.

220 **Table 1. Headwater catchment descriptors identified as potential explanatory variables of spatial variability and temporal variation in dissolved organic carbon (DOC), nitrate (NO₃), and soluble reactive phosphorus (SRP) in stream and river water.**
 Topo_i = $\log \frac{\alpha}{\tan \beta}$, (Beven and Kirkby, 1979), where α is the drainage area (ha) and β is the downstream slope (%), (Merot et al., 2003).
 225 ^a there are 3 classes of soil thickness: 40-60 cm, 60-80 cm, 80-100 cm and >100 cm. ^b Winter crops have a winter plant cover and a phenological maximum in April (wheat, barley, rapeseed). ^c Summer crops correspond to bare winter soils and a phenological maximum in early summer (corn).

Type	Descriptor name	Unit	Definition	Source
Topography	Area	km ²	Drainage area of the monitoring station	Web Processing Service “Service de Traitement de Modèles Numériques de Terrain” and DEM 50 m by IGN
	Elevation	m	Mean elevation of headwater catchment	DEM 25 m by IGN
	Density_hn	km.km ⁻²	Density of the hydrographic network	
	Topo _i	log(m ³)	Downstream topographic index of the headwater	BD Carthage by IGN
	IDPR	-	Hydrographic Network Development and Persistence Index	http://infoterre.brgm.fr/ BRGM data and geoservices portal (Mardhel and Gravier, 2004)
Geology	Granite_pm	%	Percentage of granite and gneiss area	Web Mapping Service “Carte des Sols de Bretagne” by UMR 1069 SAS
	Schist_pm	%	Percentage of schist and micaschist area	INRAE - Agrocampus Ouest
	Other_pm	%	Percentage of various geological substrata	http://www.sols-de-bretagne.fr/
Soil	Erosion	%	Percentage of area with high to very high erosion risk (derived from land use, topography and soil properties)	Erosion risk map estimated from MESALES by GIS Sol, INRAE from Colmar et al. (2010)
	OC_soil	g.kg ⁻¹	Organic carbon content in the topsoil horizon (0-30 cm)	Web Mapping Service from BDAT database, Saby et al. (2015) by GIS Sol
	Thick_soil	cm	Classes of dominant soil thickness ^a	Web Mapping Service “Carte des Sols de Bretagne” by UMR 1069 SAS
	TP_soil	g.kg ⁻¹	Total phosphorus content in the topsoil horizon (0-30 cm)	INRAE - Agrocampus Ouest Web Mapping Service from BDAT database by GIS Sol
Land use	SummerCrop	%	Percentage of summer crop ^b land	
	WinterCrop	%	Percentage of winter crop ^c land	OSO database, CESBIO, land-cover map 2016 (1 ha) from http://osr-cesbio.ups-tlse.fr/~oso/
	Forest	%	Percentage of forest land	
	Pasture	%	Percentage of pasture land	
	Urban	%	Percentage of urban land	
	Wetland	%	Percentage of potential wetlands	Web Mapping Service “Enveloppe des milieux potentiellement humides de France réalisée par les laboratoires Infosol et UMR SAS” by UMR 1069

Diffuse and point N and P sources	N_surplus	kg.ha ⁻¹ .yr ⁻¹	Nitrogen surplus (= the maximum quantity on a given agricultural area that is likely to be transferred to the stream network)	CASSIS-N estimates by (Poisvert et al., 2017) from https://geosciences.univ-tours.fr/cassis/login
	P_surplus	kg.ha ⁻¹ .yr ⁻¹	Phosphorous surplus	NOPOLU estimates by (SoeS, 2013)
	N_point	kg.ha ⁻¹ .yr ⁻¹	Sum of nitrogen loads from domestic and industrial point sources	Data from Loire-Bretagne Water Agency data (2008-2012)
	P_point	kg.ha ⁻¹ .yr ⁻¹	Sum of phosphorus loads from domestic and industrial point sources	Data from Loire-Bretagne Water Agency (2008-2012)
Hydrology	Qmean	l.s ⁻¹ .km ²	Interannual mean flow	
	QMNA	l.s ⁻¹ .km ²	Median of annual minimum monthly specific discharge	Calculated from flow data observations: HYDRO regional database by DREAL Bretagne & GR4J simulations (Perrin et al., 2003)
	BFI	%	Base flow index (Lyne et Hollick, 1979)	
	W2	%	Percentage of total discharge that occurs during the highest 2% of flows (Moatar <i>et al.</i> , 2013)	
	Rainfall	mm.yr ⁻¹	Mean effective rainfall from 2008-2012	SAFRAN database (8 km ²) by Météo France

3 Results

3.1 Spatial variability in concentrations and loads

The C50 of the 185 headwater catchments ranged from 2-14.6 mg C.l⁻¹ for DOC, 0.9-15.8 mg N.l⁻¹ for NO₃, and 8-241 µg P.l⁻¹ for SRP (with 75% of the SRP C50 < 64 µg P.l⁻¹). The C50 displayed spatial gradients: rivers with DOC concentrations > 5 mg C.l⁻¹ were located in eastern Brittany, while the highest NO₃ concentrations were located on the west coast (Fig. 2). In contrast, the highest concentrations of SRP (C50 > 68 µg P.l⁻¹) were located in northern Brittany.


The two first axes of the PCA (Supplemental S4a) performed on the percentiles of DOC, NO₃, and SRP concentrations of the 185 headwater catchments explained 58% of the variance and revealed three important points. First, percentiles (C10, C50, or C90) were grouped by solute, showing that the spatial organization remained the same regardless of the concentration percentile (Spearman rank correlations between the three indices always greater than 0.56 for all elements). Second, there was a negative correlation between DOC and NO₃ concentrations ($r_s = -0.58$; Supplemental S4b). Third, SRP concentrations had an orthogonal relation compared to DOC and NO₃ concentrations (r_s close to zero).

The ratios of mean concentration ($CV_{c_{mean}}$) to mean flow ($CV_{q_{mean}}$) were < 1 for DOC and NO₃ (Table 2), indicating that concentrations varied less in space than in flow, and vice-versa for SRP.


For DOC and NO₃, Ampli was not correlated significantly with C50, but it was with C90 (Fig. 3). For SRP, correlations between Ampli and the percentiles were high, with $r_s > 0.85$ for C50 and C90 (Fig. 3). The SI and phases were correlated more with C10 for DOC and NO₃ (negatively for SI and positively for the phases), and more with C90 for SRP (negatively, for SI only).

245 Mean (± 1 SD) interannual loads had high spatial variabilities – 20.71 ± 10.52 kg C.ha⁻¹.yr⁻¹ for DOC, 27.48 ± 18.51 kg N.ha⁻¹.yr⁻¹ for NO₃, and 0.315 ± 0.11 kg P.ha⁻¹.yr⁻¹ for SRP – which differed from those observed for concentrations (Fig. 2). Unsurprisingly, interannual loads of the three solutes were significantly ($p < 0.001$) and strongly correlated with annual water fluxes (Pearson $r = 0.88$ for DOC, 0.90 for NO₃, and 0.75 for SRP). There were weak but significant positive correlations between mean interannual loads and seasonality indices (Ampli, SI) or C90 for DOC (Fig. 3). Mean interannual loads of NO₃

250 were significantly and positively correlated with C10 and C50, and negatively with its seasonality indices. The strongest significant correlation was found between mean interannual loads and concentration percentiles for SRP.



255 **Figure 2. Map of median (left) concentrations C50 and (right) loads of dissolved organic carbon (DOC), nitrate N (N-NO₃), and soluble reactive phosphorus (SRP) for the 185 streams. The catchments in gray did not meet the criteria to estimate a mean average interannual load. Classes in the legends have equal numbers of catchments.**



260 **Figure 3. Matrices of Spearman's rank correlations of water quality (load, concentration percentiles (10th (C10), 50th (C50), and 90th (C90)), and seasonality metrics) for (a) dissolved organic carbon (DOC), (b) nitrate N (N-NO₃), and (c) soluble reactive phosphorus (SRP) (c). Only significant (p ≤ 0.05) values are shown.**

Table 2. Coefficients of variation (spatial variability among catchments) of flow-weighted mean concentration (CV_cmean) and mean stream flow (CV_qmean), and the value of their ratio, for dissolved organic carbon (DOC), nitrate (NO₃), and soluble reactive phosphorus (SRP).

Parameter	CV _c mean	CV _q mean	CV _c mean:CV _q mean
DOC	0.2954	0.4614	0.6403
NO ₃	0.3285	0.4709	0.6976
SRP	0.9207	0.4743	1.9412

265

3.2 Characterization of concentrations seasonality

3.2.1 Performance of GAMS

270 Of the 185 catchments, GAMS were fitted for 159 to DOC concentrations time series, 168 to NO₃ concentrations time series, 162 to SRP concentrations time series, and 185 to discharge time series. The cases for which fitting was not possible corresponded to those with no seasonal cyclicality or with excessive interannual variability. The percentage of variance explained by the GAM varied by site and solute. Fitting performed best for NO₃, followed by SRP and then DOC: the means and SDs of the adjusted R_{sq} were 0.30 ± 0.18, 0.16 ± 0.11, and 0.22 ± 0.15 for NO₃, DOC, and SRP, respectively (Supplemental S5 and S6), and the percentages of catchment for which the fitted model had R_{sq} > 0.20 were 67%, 52% and 38%, respectively. Metrics calculated from monthly data differed only moderately from those calculated from sub-monthly data (Supplemental S7), which tended to validate the approach of using monthly data.

275

3.2.2 Types of seasonal cyclicity in DOC, NO₃, and SRP

Most of the catchments had a seasonal concentration cycle: 85%, 71%, 78%, for NO₃, DOC, SRP concentration respectively and 100% of them had a seasonal discharge cycle (Fig. 4). Means and SDs of the standardized Ampli were 0.59 ± 0.46 for
280 NO₃, 0.53 ± 0.30 for DOC, 0.79 ± 0.14 for SRP, and 1.99 ± 0.38 for discharge. The distribution of the calculated seasonality indices is provided in Supplemental S8.

The annual phases for discharge were more stable among all catchments than those for concentrations. The highest discharge period was centered on mid-February (winter) and the lowest discharge period on September. A strong gradient of hydrological dynamics was observed among catchments (Fig. 4d and Supplemental S8). The highest W2 was associated with both severe
285 low-flow discharge and many high discharge events. Values of Q_{mean}, BFI, W2, and QMNA clearly followed an east-west gradient (not shown). Because of similar seasonal discharge dynamics in all catchments, SI can be used to describe the seasonal dynamics of a concentration relative to those of discharge. When SI was positive, the concentration seasonality was in-phase with discharge; when negative, the concentration seasonality was out-of-phase with discharge (Fig. 4).

Most of the catchments had opposite dynamics for DOC and NO₃. For 90% of them, Pearson correlation between the daily
290 GAM estimates of DOC and NO₃ was negative, and for 50% of the catchments, less than -0.79. The remaining 10% of catchments (15) had low Ampli of DOC and NO₃. The DOC and NO₃ concentrations had out-of-phase seasonal cycles, as shown by the negative correlation between SI and DOC or NO₃ for all catchments that had a significant seasonality in these concentrations (Fig. 5; R² = 0.62). We classified two types of catchments according to their seasonality in both DOC (MinPhase) and NO₃ (MaxPhase) concentrations and consistent with the SI (Fig. 5, Supplemental S8). NO₃ MaxPhase and
295 DOC MinPhase that occurred before 1 May were classified as “in-phase” with discharge (Q), while those that occurred after were “out-of-phase” with Q. All catchments experienced high stability of the DOC MaxPhase and NO₃ MinPhase were the same for all catchments as they always occurred between July and December (Fig. 4, Supplemental S8).

The first type, “in-phase” (68% of the catchments with seasonality), had a NO₃ MaxPhase between October and May (Fig. 4, Supplemental S8) (i.e. high-flow period, in-phase with maximum discharge and usually with DOC MinPhase). For these
300 catchments, the mean SI was positive for NO₃ (0.22 ± 0.19) and usually negative or null for DOC (0.00 ± 0.13). They tended to be located toward central Brittany and be associated with mesoscale catchments (mean of 52.6 ± 38.8 km²). They had large Ampli for NO₃ and low Ampli for DOC (mean relative Ampli of 0.83 ± 0.46, and 0.44 ± 0.23 for DOC) and relatively low C50 of NO₃ (means of 5.74 ± 2.46 mg N.l⁻¹ and 5.92 ± 2.00 mg C.l⁻¹).

The second type, “out-of-phase” (32% of the catchments with seasonality), had a DOC MinPhase and NO₃ MaxPhase between
305 May and September (Fig. 4; Supplemental S8) (i.e. low-flow period, out-of-phase with maximum discharge). For most catchments, maximum NO₃ and minimum DOC concentrations occurred a mean of 1.85 months before minimum discharge or 5.5 months after maximum discharge, respectively. For these catchments, the mean SI was negative or null for NO₃ (-0.08 ± 0.06) and weakly positive for DOC (0.21 ± 0.10). These catchments were close to the coast and relatively small (mean of

31.4 ± 21.7 km²). They had smaller Ampli than “in-phase” catchments for NO₃, and higher Ampli for DOC (mean relative
310 Ampli of 0.13 ± 0.13, and 0.74 ± 0.30 for DOC) and relatively high C50 of NO₃ (means of 8.27 ± 2.90 mg N.l⁻¹ and 5.00 ±
1.62 mg C.l⁻¹).

Some catchments had intermediate behavior between these two types (Figs. 4 and 5). Some had a plateau with maximum NO₃
and minimum DOC concentrations from winter to summer, while others showed two maxima for NO₃ or two minima for DOC
315 (one synchronous with maximum discharge and another with minimum discharge). Other catchments also had maximum NO₃
synchronous with discharge, but minimum DOC after maximum discharge.

The seasonal dynamics of SRP were more stable than those of DOC and NO₃, but less stable than those of discharge. Thus,
there was only one type of seasonality for SRP, which was out-of-phase with flow: MaxPhase SRP dominated in summer
(mid-August ± 1.4 months), and MinPhase SRP dominated in late winter (March ± 1.2 months) (Fig. 4, Supplement S7), except
for two catchments with maximum SRP in January-February.

320





Figure 4. Seasonal dynamics of nitrate N (N-NO₃), dissolved organic carbon (DOC), soluble reactive phosphorus (SRP), and daily discharge modeled by Generalized Additive Models for 185 headwater catchments. To compare concentrations, they are standardized by their mean interannual concentration. The color gradient represents the seasonality index of each parameter; thus, a headwater catchment's color can vary among panels.

325

Relationship between the seasonality indices (SI) of NO₃ versus DOC parameter within the headwater catchments where the seasonality is significant for both parameters (n=98).



330 **Figure 5. Relationship between the seasonality indices (SI) of nitrate N (N-NO₃⁻) vs. dissolved organic carbon (DOC) in the headwater catchments for which seasonality was significant for both parameters (n=98). The color and shape of symbols identify the seasonality types based on the NO₃ MaxPhase and DOC MinPhase metrics. The threshold date was 1 May: MaxPhase that occurred before were classified as “in-phase” with discharge (Q), while those that occurred after were “out-of-phase” with Q. The DOC MinPhase metric is shown to highlight the synchrony between minimum DOC and maximum N-NO₃ concentrations.**

3.3 Controlling factors of concentration percentiles and seasonality

The C50 of DOC was correlated significantly with 15 spatial variables and most strongly ($|r_s| \geq 0.4$) with topographic index, 335 QMNA, and the other hydrological indices. The C50 of NO₃ was correlated significantly with 12 spatial variables, in particular diffuse agricultural sources ($r_s = 0.68$ for the percentage of summer crops, $r_s > 0.39$ for N and P surplus, and $r_s = 0.48$ for soil erosion rate) and hydrological indices, through the base flow index (BFI) (positively) and W2 (negatively), (Table 3). The C50 of SRP was correlated significantly with more variables (18), but the correlations were slightly weaker. It correlated most strongly with soil P stock ($r_s = -0.40$), climate and hydrology ($r_s = -0.43$ to -0.34 with effective rainfall, Qmean, QMNA),


340 elevation, and hydrographic network density. It had weaker positive correlations ($r_s < 0.3$) with the soil erosion rate and domestic and agricultural pressures (urban percentage and P surplus).

Ampli and SI for DOC and NO_3 were correlated most with the hydrodynamic properties, followed by agricultural pressures (Fig. 6, Table 3). The catchments “in-phase” with discharge (i.e. positive SI- NO_3 and negative SI-DOC correlations) were associated with high hydrological reactivity (low BFI and high W2) and a low percentage of summer crops (Table 3).

345 Conversely, catchments “out-of-phase” with discharge (i.e. negative SI- NO_3 and positive SI-DOC correlations) were associated with low hydrological reactivity (high BFI and QMNA, low W2) and a high percentage of summer crops.

Correlations of SI with catchment descriptors were weaker ($|r_s| \leq 0.4$) for SRP than for DOC and NO_3 because most catchments had the same seasonal pattern, with maximum SRP concentration during low flow. Catchments with the highest amplitudes of SRP concentration were associated with low QMNA and Q_{mean} , high W2, low effective rainfall, and low soil P stock.

350 Interannual loads were correlated mainly with hydrological descriptors (positively with Q_{mean} and QMNA, and negatively with W2) (Table 3). Interannual NO_3 loads were also correlated with the percentage of summer crops and soil TP content, while interannual SRP loads were correlated weakly with the percentage of summer crops, agricultural surplus, erosion, and point sources.



355

Figure 6. Relationship between the seasonality index (SI) of dissolved organic carbon (DOC) and nitrate (NO_3) and the hydrological reactivity descriptors (A) flow flashiness index (W2) and (B) base-flow index (BFI) for 124 headwater catchments.

360 Table 3. Spearman rank correlations between water quality indices and geographical descriptors for dissolved organic carbon (DOC), nitrate (NO₃), and soluble reactive phosphorus (SRP). Only significant correlations (p≤0.05) are shown, and bold text indicates |r| ≥ 0.40.

		DOC				NO ₃				SRP			
Spatial variable		C50	Ampli	SI	Load	C50	Ampli	SI	Load	C50	Ampli	SI	Load
Topography	Area	-	-0.24	-	-	-	-	-	-	-	-	-	-
	Elevation	-0.46	-0.18	-	-	-	-0.31	-0.20	0.19	-0.20	-	-	-
	Density_hn	-	-	-	-	-	-0.22	-	0.16	-0.30	-0.27	0.19	-
	Topo_i	0.54	-	-	-	-	0.41	0.25	-0.33	0.39	0.25	-	0.18
	IDPR	-	-	-	-	-	-	-	-	-0.21	-0.19	-	-
Geology	Granite_pm	-	-	0.21	0.41	-	-0.43	-0.31	0.27	-0.26	-0.24	-	-
	Schist_pm	-	-0.21	-0.37	-0.29	-0.16	0.25	0.22	-0.23	-	-	-	-0.20
	Other_pm	-	0.32	0.35	-	0.28	-	-	-	0.28	0.16	-	0.35
Soil	Erosion	-0.36	0.24	-	-	0.48	0.16	-0.26	0.39	0.24	0.17	-	0.33
	OC_soil	-0.27	-0.21	-	-	-	-0.29	-	0.18	-0.20	-0.19	-	-
	TP_soil	-0.44	-	-	0.38	-	-0.51	-0.34	0.49	-0.40	-0.32	-	-
Land use	SummerCrop	-0.30	0.28	0.54	-	0.68	-	-0.47	0.54	-	-	0.29	0.36
	WinterCrop	0.19	-	-0.20	-0.29	-	0.48	0.21	-0.23	0.17	-	-0.18	-
	Forest	-	-0.17	-0.30	0.23	-0.37	-0.47	-	-	-0.29	-0.19	-	-0.27
	Pasture	-	-	-	-	-0.30	-	0.26	-0.20	-	-	-	-
	Urban	-	-	-	-	-	-	-	-	0.23	-	-	-
N and P diffuse and point sources	N_surplus	-0.21	0.20	-	-	0.39	-	-	0.38	-	-	0.29	0.29
	P_surplus	-0.24	0.33	-	-0.22	0.49	-	-0.32	0.37	0.20	-0.19	-	0.35
	N_point	-	-0.17	-	-	-	-	-	-	-	-	-	-
	P_point	-	-0.16	-	-	-	-	-	0.21	-	-	-	0.21
Hydrology	Qmean	-0.49	0.19	-	0.53	0.16	-0.58	-0.42	0.67	-0.39	-0.31	0.21	0.18
	QMNA	-0.52	0.25	0.41	0.48	0.42	-0.54	-0.56	0.76	-0.34	-0.32	0.35	0.27
	BFI	-0.41	-0.27	0.64	0.38	0.54	-0.52	-0.69	0.57	-0.20	-0.23	0.32	0.23
	W2	0.43	-	-0.61	-0.46	-0.49	0.54	0.68	-0.59	0.20	0.20	-0.26	-0.24
	Precipitation	-0.50	-	-	0.47	-	-0.60	-0.39	0.60	-0.43	-0.33	0.18	-
	Wetland	0.16	-	0.31	0.38	-	-	-	-	-	-	-	0.35

365 4 Discussion

4.1 Interpretation of the spatial opposition between DOC and NO₃

Spatial opposition between DOC and NO₃ concentrations has been reported for a wide range of ecosystems. Taylor and Townsend (2010) found a non-linear negative relationship between them for soils, groundwater, surface freshwater, and oceans, from global to local scales, and highlighted that this negative correlation prevails in disturbed ecosystems. Goodale et al. (2005) reported a similar negative correlation among 100 streams in the northeastern USA. Heppell et al. (2017) found that DOC and NO₃ concentrations were inversely correlated with the BFI in six reaches of the Hampshire Avon catchment (UK). Our contribution brings an original focus on this relationship in headwater catchments with high domestic and agricultural pressures. Taylor and Townsend (2010) interpreted this spatial opposition as a response of microbial processes (i.e. biomass production, nitrification, and denitrification) to the ratio of ambient DOC:NO₃, which controls NO₃ export/retention in catchments (see also Goodale et al. (2005)). In semi-natural ecosystems, high but poorly labile soil organic C pools were associated with lower N retention capacity and thus higher N leaching (Evans et al., 2006). Similarly, several studies (e.g. Hedin et al., 1998; Hill et al., 2000) suggested that DOC supply limits in- and near-stream denitrification. In contrast, other studies claimed that N can influence loss of DOC from soils by altering substrate availability or/and microbial processing of soil organic matter (Findlay, 2005; Pregitzer et al., 2004). In our study, C50 were correlated with both BFI and QMNA, positively for NO₃ and negatively for DOC, which suggests that catchments strongly sustained by groundwater flow produced higher NO₃ and lower DOC concentrations, as reported in other rural catchments (e.g. Heppell et al., 2017). The C50 of NO₃ increased with agricultural pressures (percentage of summer crop, N surplus), as observed by Lintern et al. (2018), while that of DOC increased in flatter catchments, which is consistent with results of Mengistu et al. (2014) and Musolff et al. (2018). This suggests that this spatial opposition between DOC and NO₃ results from the combination of heterogeneous human inputs, heterogeneous natural pools, and different physical and biogeochemical connections between C and N pools. In surface water, these heterogeneous sources are expressed to differing degrees depending on the catchment's hydrological behavior. When deep or slow flowpaths dominate, they store and release N via groundwater and mobilize little the sources rich in organic matter. When shallower and faster flowpaths dominate, they transport some of the N via compartments rich in organic matter, which causes N depletion and release of more DOC to the streams. The initial amounts of NO₃ along these flowpaths are a function of human pressures.

4.2 Interpretation of the temporal opposition between DOC and NO₃

The seasonal opposition between DOC and NO₃ concentration dynamics could be another manifestation of the spatial opposition between DOC and NO₃ sources, because the strength of the hydrological connection between sources and streams


395 varies seasonally (e.g. Mulholland and Hill (1997), Weigand et al. (2017)). The direct contribution of biogeochemical reactions that connect DOC and NO₃ cycles may also vary seasonally (Mulholland and Hill, 1997; Plont et al., 2020). Indeed, temperature, wetness condition, and light availability influence rates of these organic matter reactions (Davidson et al., 2006; Hénault and Germon, 2000; Luo and Zhou, 2006). In addition, the relative importance of the fluxes produced or consumed via these reactions appears clearer during the low-flow period, when the fluxes exported from the terrestrial ecosystem and delivered to the stream decrease. These reactions consume NO₃ (e.g. denitrification, biological uptake) and release (reductive dissolution) or produce (autotrophic production) DOC. Of the two seasonal NO₃-DOC cycles, the most common in our datasets is thus maximum NO₃ in-phase with maximum discharge and minimum DOC, which has been reported in Brittany (Abbott et al., 2018b; Dupas et al., 2018) and elsewhere (Van Meter et al., 2019; Dupas et al., 2017; Halliday et al., 2012; Minaudo et al., 2015; Weigand et al., 2017). The main control of seasonal DOC-NO₃ cycles appears to be related to hydrological indices (expressed as BFI and W2). Hydrological flashiness reflects the relative importance of subsurface flow compared to deep base flow (Heppell et al., 2017); thus, low BFI (or high W2) would indicate higher connectivity with subsurface riparian sources and shorter transit times. This is consistent with results of Weigand et al. (2017), who observed higher seasonal amplitudes in DOC and NO₃ concentrations and stronger temporal anti-correlation between DOC and NO₃ concentrations in stream water dominated by subsurface runoff.

410 Our results are consistent with these previous results, while the correlations with catchment characteristics can provide some explanation. Catchments with low BFI have larger shallow flows and experience seasonal DOC-NO₃ cycles that are in-phase with flow and have higher NO₃ amplitudes. These cycles can be interpreted as the combination of several mechanisms (Fig. 7):

- 1) Synchronization of NO₃-rich and DOC-poor groundwater contribution with maximum flow.
- 415 2) Large contribution of near-/in-stream biogeochemical processes at reduced low flows that decreases NO₃ concentration (e.g. NO₃ consumption by aquatic microorganisms, biofilms, and macrophytes).
- 3) Large DOC-rich riparian contribution throughout the year, but larger in autumn, when flow starts to increase, as described in detail in previous AgrHys Observatory studies (Aubert et al., 2013; Humbert et al., 2015).

In contrast, catchments with higher BFI have smaller shallow flows and experience mainly DOC and NO₃ cycles that are out-of-phase with flow and have lower amplitudes. These cycles can be attributed to the following:

- 420 1) More continuous groundwater contribution, combined with a decrease in agricultural pressures over time, which could increase NO₃ concentrations more in deeper groundwater than in shallower groundwater (Abbott et al., 2018b; Martin et al., 2004; Martin et al., 2006). This vertical gradient in groundwater supply could explain why NO₃ concentrations peaked during the annual discharge recession, which is sustained mainly by deep groundwater inputs.
- 425 2) Little contribution of near-/in-stream biogeochemical processes at reduced low flows due to larger inputs from groundwater, which maintains a relatively high minimum NO₃ concentration.
- 3) Contribution of DOC-rich riparian sources, mainly in autumn, that are smaller than those in in-phase catchments, again due to a predominantly deeper geometry of water circulation.



430 **Figure 7. Conceptual diagram of seasonal flowpaths involved in the DOC-NO₃ seasonal cycles leading to a) in-phase cycles with discharge or b) out-of-phase cycles with discharge.**

4.3 Interpretation of the spatial and temporal signature of SRP

The correlations between the C50 of SRP and geographic variables highlighted the importance of P sources (soil P stocks, followed by domestic and agricultural pressures) and surface flowpaths (e.g. hydrological indices, elevation, erosion risk). Similarly, analysis of regression models that predicted spatial variability in total P₃ concentration of 102 rural catchments in Australia also indicated positive effects of human-modified land uses, natural land uses prone to soil erosion, mean P content of soils, and to a lesser extent, topography (Lintern et al., 2018). They always included the percentage of urban area, which suggests a considerable effect of sewage discharge, even at low levels of urbanization. The catchments analyzed in the present study have a homogeneous and relatively dense distribution of small villages but no large city, which seems to support this last hypothesis. Sobota et al. (2011) studied spatial relationships among P inputs, land cover and mean annual concentrations of different forms of P in 24 catchments in California, USA. They found that P concentrations were significantly correlated with agricultural inputs and, to a lesser extent, agricultural land cover but not with estimates of sewage discharge. Nonpoint sources of P in agricultural runoff, historical inputs of fertilizer and manure in excess of crop requirements have led to a build-up of soil P levels, particularly in areas of intensive crop and livestock production (Sharpley et al., 1994). This led to correlations between soil P and runoff concentrations in agricultural catchments (Cooper et al., 2015; Sandström et al., 2020), as found here.

The seasonality of SRP was generally the same in the region studied, and C50 and amplitudes were significantly correlated. A peak in seasonal SRP concentrations at low flow has been reported previously (Abbott et al., 2018b; Bowes et al., 2015; Dupas et al., 2018; Melland et al., 2012). It is interpreted as the result of a dominance of point sources diluted during high flow

(Minaudo et al., 2019, 2015; Bowes et al., 2011) or of stream-bed sediment sources for which P release increases with temperature (Duan et al., 2012).

Correlation between spatial patterns of NO₃ and SRP was expected given the dominant agricultural origin of N and substantial agricultural origin of P, but it was not observed in all catchments. The C50 of NO₃ and SRP were high mainly on the northwestern coast, perhaps due to intensive vegetable production associated with a dominance of mineral fertilization (Lemercier et al., 2008). Elsewhere, a high proportion of allochthonous P in the topsoil results from livestock farming and manure application (Delmas et al., 2015). The P-retention capacity of soils (related to their Al, Ca, Fe, and clay contents) is also likely to increase spatial variability in the release of P from catchments (Delmas et al., 2015). Synchronous variations in SRP and DOC, such as those observed in small, completely agricultural headwater catchments without villages (Cooper et al., 2015; Dupas et al., 2015b; Gu et al., 2017), were not observed in the present set of catchments. We assume that synchronicity of SRP and DOC in small catchments depends on soil processes, such as reduction of soil Fe-oxyhydroxides in wetland zones (Gu et al., 2019), which are hidden by in-stream processes (P adsorption on streambed sediments) and downstream point-source inputs (especially P inputs) in the set of larger catchments studied.

Regarding the geographic data used as spatial descriptors, the region studied did not have a few dense urban centers but rather smaller domestic points scattered across the region, which is harder to characterize finely. Moreover, Brittany's coastlines may have higher population densities in spring and summer due to tourism. Refined estimates of domestic point sources and their seasonal variations would be useful in future analyses.

4.4 Hydrological vs. anthropogenic controls of spatial variability in water quality

Among the headwater catchments selected, the human pressures (agriculture for NO₃ and sewage water discharge for SRP) influenced the C50 and loads of NO₃ and SRP. However, the influence of hydrological descriptors on the spatial variability in their loads suggested a transport-limited behavior of these catchments (Basu et al., 2010). Nutrient load estimates had high uncertainties due to i) using modeled flow data when measurements were not available and ii) the frequency of concentration data (monthly), which is low for estimating nutrient loads (especially of P) (Raymond et al., 2013). Thus, these load estimates allowed only their relative spatial variation to be analyzed. Although land-use or agricultural pressure variables, in combination with rainfall and discharge variables, are good predictors of nutrient loads at larger scales (Dupas et al., 2015a; Grizzetti et al., 2005; Preston et al., 2011), the correlations with loads were lower in the set of headwater catchments selected. For NO₃, this can be explained by higher spatial variability (CVs) in water fluxes than in concentrations (Table 2), which can explain the dominance of hydrological fluxes in the spatial organization of nutrient loads. Such dominance was found to increase with the level of human pressure in Thompson et al. (2011) for NO₃. In this study, such relationship was not visible as all the catchments exhibited a transport-limited behavior. It may also suggest that the nutrient-surplus data at the local scale remained uncertain (Poisvert et al., 2017) or that at this scale, data on agricultural practices would be more relevant, and that variability in concentration depends less on the magnitude of nutrient inputs than on their locations.

The catchments studied have clear seasonal dynamics in concentration, which is consistent with previous observations (Minaudo et al., 2019; Abbott et al., 2018a). The seasonal pattern is controlled mainly by hydrological variables. It partly reflects the mixing of contrasting sources that are connected to streams by seasonally varying flowpaths with nutrients that are transferred vs. nutrients that are processed locally in hotspots (e.g. riparian buffer, stream water, stream sediments) or delivered over point sources. The seasonal NO₃-DOC pattern seemed to become somewhat homogenous among catchments larger than 100 km², where seasonal cycles with maximum NO₃ in-phase with flow seemed less common. This may be related to an increase in in-stream biological activity during summer as catchment size increases, enhanced by a lower stream water level and slower discharge (Minaudo et al., 2015). Therefore, the potential relationship between seasonal cycle type and catchment size should be studied over a wider range of catchment sizes and nested catchments to include variations along the hydrographic network.

4.5 Implications for headwater monitoring and management

The high regional and seasonal variations of nutrient concentrations in streams probably drive high variations of nutrient stoichiometry along the water year and over the region, and, consequently, high variations in time and space of eutrophication risks downstream (Westphal et al., 2020). Due to the combination of anthropogenic and hydrological drivers in explaining these stream concentrations, a better estimation on nutrient inputs and discharge in all headwater catchments, as a first step, is important to predict areas at risks. The spatial analysis shows high and poorly structured spatial variations of concentrations over the region. Nevertheless, the opposition between NO₃ and DOC concentrations suggests that the C:N ratios will be even more variable:

- 1) In space: catchments with high DOC C50 and low NO₃ C50 will exhibit very high C:N and vice versa
- 2) Over the seasons: as minimum of DOC and maximum of NO₃ concentrations are in-phase: catchment where DOC-NO₃ variations are in phase with Q will exhibit a low C:N ratio in winter high flow period and higher C:N ratio during low flow period. The N:P ratio in these catchments will be high during the low flow periods (high NO₃ and low SRP concentrations). Catchments where DOC-NO₃ variations are out-of-phase with discharge will exhibit probably less variation in their ratios (because of lower NO₃ amplitude) with relatively higher winter C:N ratio than the previous type of catchments.

510 5 Conclusion

To analyze spatial variability in water quality at a regional scale, we used an original dataset from public databases, little used by the scientific community, for the French region of Brittany with monthly measurements of water quality. The dataset selected covers 185 headwater and agricultural catchments monitored over a period sufficiently long (10 years) to allow the

spatial (regional) variability and temporal (seasonal) variation in DOC, NO₃, and SRP concentrations to be analyzed. We
515 described spatio-temporal variations in concentrations, loads, and seasonal patterns and analyzed their correlations with
geographic variables (related to topography, hydro-climate, geology, soils, land uses, and human pressures). Our study showed
the following:

- 1) Seasonal cycles of DOC and NO₃ concentrations are usually opposite from each other. Catchments with a low base-
flow index exhibit maximum NO₃ in-phase with maximum flow, while those with a higher base-flow index exhibit
520 maximum NO₃ after maximum flow. Both types exhibited maximum DOC in autumn, at the beginning of the annual
increase in flow.
- 2) NO₃ concentrations increased as human pressures and base flow contribution increased. DOC concentrations
decreased as rainfall, base flow contribution, and elevation increased. SRP concentrations showed weaker correlations
with human pressures, rainfall, and hydrological and topographic variables.
- 525 3) Seasonal SRP cycles are synchronized in nearly all catchments that have a clear seasonal amplitude, with maximum
SRP concentrations that occur during the summer low-flow period due to a decreased dilution capacity of point
sources.

The spatial and temporal opposition between DOC and NO₃ concentrations likely results from a combination of heterogeneous
human inputs and biogeochemical connection between these pools. The seasonal cycles in stream concentrations result from
530 the mixing of water parcels that followed contrasting flowpaths, combined with high spatial variability in nutrient sources,
local-scale biogeochemical processes, and point sources. As a perspective, we recommend further studies of multiple elements
that are likely to show contrasting responses to diverse human pressures and to the retention/removal capacities of
hydrosystems.

Acknowledgments

535 The salary of SG was supported by Region Bretagne and Agence de l'Eau Loire Bretagne. We thanks Dr Remi Dupas (INRAE
Rennes) for his valuable contribution for methodological choices and the scientific interpretations and discussions. We thank
Dr Vazken Andreassian (INRAE Anthony) for providing regional simulations of discharge time series with the model GR4J.
We thank also Josette Launay (CRESEB), Elodie Bardon (Observatoire Environnement Bretagne), Yves-Marie Heno and
Olivier Nauleau (DREAL Bretagne) for their contributions to the data selection and to the project. Finally, we thanks all the
540 people who contributed to the collection of public data on surface water quality in French Brittany.

References

- Abbott, B. W., Gruau, G., Zarnetske, J. P., Moatar, F., Barbe, L., Thomas, Z., Fovet, O., Kolbe, T., Gu, S., Pierson-Wickmann, A.-C., Davy, P., and Pinay, G.: Unexpected spatial stability of water chemistry in headwater stream networks, *Ecol. Lett.*, 21, 296-308, <https://doi.org/10.1111/ele.12897>, 2018a.
- 545 Abbott, B. W., Moatar, F., Gauthier, O., Fovet, O., Antoine, V., and Ragueneau, O.: Trends and seasonality of river nutrients in agricultural catchments: 18 years of weekly citizen science in France, *Sci. Total Environ.*, 624, 845-858, <https://doi.org/10.1016/j.scitotenv.2017.12.176>, 2018b.
- Ågren, A., Buffam, I., Jansson, M., and Laudon, H.: Importance of seasonality and small streams for the landscape regulation of dissolved organic carbon export, *J. Geophys. Res-Bioge.*, 112, <https://doi.org/10.1029/2006JG000381>, 2007.
- 550 Alexander, R. B., Boyer, E. W., Smith, R. A., Schwarz, G. E., and Moore, R. B.: The Role of Headwater Streams in Downstream Water Quality, *J Am Water Resour Assoc*, 43, 41-59, [10.1111/j.1752-1688.2007.00005.x](https://doi.org/10.1111/j.1752-1688.2007.00005.x), 2007.
- Andersson, J.-O. and Nyberg, L.: Spatial variation of wetlands and flux of dissolved organic carbon in boreal headwater streams, *Hydrol. Process.*, 22, 1965-1975, <https://doi.org/10.1002/hyp.6779>, 2008.
- Aubert, A. H., Gascuel-Oudou, C., Gruau, G., Akkal, N., Faucheux, M., Fauvel, Y., Grimaldi, C., Hamon, Y., Jaffrézic, A., Lecoq-Boutnik, M., Molénat, J., Petitjean, P., Ruiz, L., and Merot, P.: Solute transport dynamics in small, shallow groundwater-dominated agricultural catchments: insights from a high-frequency, multisolute 10 yr-long monitoring study, *Hydrol. Earth Syst. Sci.*, 17, 1379-1391, <https://doi.org/10.5194/hess-17-1379-2013>, 2013.
- Barnes, R. T. and Raymond, P. A.: Land-use controls on sources and processing of nitrate in small watersheds: insights from dual isotopic analysis, *Ecol. Appl.*, 20, 1961-1978, <https://doi.org/10.1890/08-1328.1>, 2010.
- 560 Basu, N. B., Destouni, G., Jawitz, J. W., Thompson, S. E., Loukinova, N. V., Darracq, A., Zanardo, S., Yaeger, M., Sivapalan, M., Rinaldo, A., and Rao, P. S. C.: Nutrient loads exported from managed catchments reveal emergent biogeochemical stationarity, *Geophys. Res. Lett.*, 37, <https://doi.org/10.1029/2010GL045168>, 2010.
- Basu, N. B., Thompson, S. E., and Rao, P. S. C.: Hydrologic and biogeochemical functioning of intensively managed catchments: A synthesis of top-down analyses, *Water Resources Research*, 47, <https://doi.org/10.1029/2011WR010800>, 2011.
- 565 Beven, K. J. and Kirkby, M. J.: A physically based, variable contributing area model of basin hydrology / Un modèle à base physique de zone d'appel variable de l'hydrologie du bassin versant, *Hydrological Sciences Bulletin*, 24, 43-69, <https://doi.org/10.1080/02626667909491834>, 1979.
- Bishop, K., Buffam, I., Erlandsson, M., Fölster, J., Laudon, H., Seibert, J., and Temnerud, J.: Aqua Incognita: the unknown headwaters, *Hydrol. Process.*, 22, 1239-1242, <https://doi.org/10.1002/hyp.7049>, 2008.
- 570 Bol, R., Gruau, G., Mellander, P.-E., Dupas, R., Bechmann, M., Skarbøvik, E., Bierzoza, M., Djodjic, F., Glendell, M., Jordan, P., Van der Grift, B., Rode, M., Smolders, E., Verbeeck, M., Gu, S., Klumpp, E., Pohle, I., Fresne, M., and Gascuel-Oudou,

- C.: Challenges of reducing phosphorus based water eutrophication in the agricultural landscapes of northwest Europe, *Front. Mar. Sci.*, 5, <https://doi.org/10.3389/fmars.2018.00276>, 2018.
- 575 Bowes, M. J., Jarvie, H. P., Halliday, S. J., Skeffington, R. A., Wade, A. J., Loewenthal, M., Gozzard, E., Newman, J. R., and Palmer-Felgate, E. J.: Characterising phosphorus and nitrate inputs to a rural river using high-frequency concentration–flow relationships, *Sci. Total Environ.*, 511, 608–620, <https://doi.org/10.1016/j.scitotenv.2014.12.086>, 2015.
- Bowes, M. J., Smith, J. T., Neal, C., Leach, D. V., Scarlett, P. M., Wickham, H. D., Harman, S. A., Armstrong, L. K., Davy-Bowker, J., Haft, M., and Davies, C. E.: Changes in water quality of the River Frome (UK) from 1965 to 2009: Is phosphorus mitigation finally working?, *Sci. Total Environ.*, 409, 3418–3430, <https://doi.org/10.1016/j.scitotenv.2011.04.049>, 2011.
- 580 Brooks, P. D., McKnight, D. M., and Bencala, K. E.: The relationship between soil heterotrophic activity, soil dissolved organic carbon (DOC) leachate, and catchment-scale DOC export in headwater catchments, *Water Resour. Res.*, 35, 1895–1902, <https://doi.org/10.1029/1998WR900125>, 1999.
- 585 Colmar, A., Walter, C., Le Bissonnais, Y., and Daroussin, J.: Démarche de validation régionale par avis d'experts du modèle MESALES d'estimation de l'aléa érosif, *Etude et Gestion des Sols*, 17, 19–32, 2010.
- Cooper, R. J., Rawlins, B. G., Krueger, T., Lézé, B., Hiscock, K. M., and Pedentchouk, N.: Contrasting controls on the phosphorus concentration of suspended particulate matter under baseflow and storm event conditions in agricultural headwater streams, *Sci. Total Environ.*, 533, 49–59, <https://doi.org/10.1016/j.scitotenv.2015.06.113>, 2015.
- 590 Cosgrove, W. J. and Loucks, D. P.: Water management: Current and future challenges and research directions, *Water Resour. Res.*, 51, 4823–4839, <https://doi.org/10.1002/2014WR016869>, 2015.
- Creed, I. F., Beall, F. D., Clair, T. A., Dillon, P. J., and Hesslein, R. H.: Predicting export of dissolved organic carbon from forested catchments in glaciated landscapes with shallow soils, *Global Biogeochem. Cy.*, 22, <https://doi.org/10.1029/2008GB003294>, 2008.
- 595 Davidson, E. A., Janssens, I. A., and Luo, Y.: On the variability of respiration in terrestrial ecosystems: moving beyond Q10, *Global Change Biology*, 12, 154–164, <https://doi.org/10.1111/j.1365-2486.2005.01065.x>, 2006.
- Dawson, J. J. C., Soulsby, C., Tetzlaff, D., Hrachowitz, M., Dunn, S. M., and Malcolm, I. A.: Influence of hydrology and seasonality on DOC exports from three contrasting upland catchments, *Biogeochemistry*, 90, 93–113, <https://doi.org/10.1007/s10533-008-9234-3>, 2008.
- 600 Delmas, M., Saby, N., Arrouays, D., Dupas, R., Lemerrier, B., Pellerin, S., and Gascuel-Oudou, C.: Explaining and mapping total phosphorus content in French topsoils, *Soil Use Manage.*, 31, 259–269, <https://doi.org/10.1111/sum.12192>, 2015.
- Dodds, W. K. and Smith, V. H.: Nitrogen, phosphorus, and eutrophication in streams, *Inland Waters*, 6, 155–164, DOI: [10.5268/IW-6.2.909](https://doi.org/10.5268/IW-6.2.909), 2016.
- Duan, S., Kaushal, S. S., Groffman, P. M., Band, L. E., and Belt, K. T.: Phosphorus export across an urban to rural gradient in the Chesapeake Bay watershed, *J. Geophys. Res.-Biogeo.*, 117, <https://doi.org/10.1029/2011JG001782>, 2012.
- 605

- Duncan, J. M., Band, L. E., Groffman, P. M., and Bernhardt, E. S.: Mechanisms driving the seasonality of catchment scale nitrate export: Evidence for riparian ecohydrologic controls, *Water Resour. Res.*, 51, 3982-3997, <https://doi.org/10.1002/2015WR016937>, 2015.
- 610 Dupas, R., Delmas, M., Dorioz, J.-M., Garnier, J., Moatar, F., and Gascuel-Oudou, C.: Assessing the impact of agricultural pressures on N and P loads and eutrophication risk, *Ecol. Indic.*, 48, 396-407, <https://doi.org/10.1016/j.ecolind.2014.08.007>, 2015a.
- Dupas, R., Gruau, G., Gu, S., Humbert, G., Jaffrézic, A., and Gascuel-Oudou, C.: Groundwater control of biogeochemical processes causing phosphorus release from riparian wetlands, *Water Res.*, 84, 307-314, <https://doi.org/10.1016/j.watres.2015.07.048>, 2015b.
- 615 Dupas, R., Minaudo, C., and Abbott, B. W.: Stability of spatial patterns in water chemistry across temperate ecoregions, *Environ. Res. Lett.*, 14, 074015, <https://doi.org/10.1088/1748-9326/ab24f4>, 2019.
- Dupas, R., Minaudo, C., Gruau, G., Ruiz, L., and Gascuel-Oudou, C.: Multidecadal Trajectory of Riverine Nitrogen and Phosphorus Dynamics in Rural Catchments, *Water Resour. Res.*, 54, 5327– 5340, <https://doi.org/10.1029/2018WR022905>, 2018.
- 620 Dupas, R., Musolff, A., Jawitz, J. W., Rao, P. S. C., Jäger, C. G., Fleckenstein, J. H., Rode, M., and Borchardt, D.: Carbon and nutrient export regimes from headwater catchments to downstream reaches, *Biogeosciences*, 14, 4391-4407, <https://doi.org/10.5194/bg-14-4391-2017>, 2017.
- Edwards, A. C., Cook, Y., Smart, R., and Wade, A. J.: Concentrations of nitrogen and phosphorus in streams draining the mixed land-use Dee Catchment, north-east Scotland, *J. Appl. Ecol.*, 37, 159-170, <https://doi.org/10.1046/j.1365-2664.2000.00500.x>, 2000.
- 625 Evans, C. D., Reynolds, B., Jenkins, A., Helliwell, R. C., Curtis, C. J., Goodale, C. L., Ferrier, R. C., Emmett, B. A., Pilkington, M. G., Caporn, S. J. M., Carroll, J. A., Norris, D., Davies, J., and Coull, M. C.: Evidence that Soil Carbon Pool Determines Susceptibility of Semi-Natural Ecosystems to Elevated Nitrogen Leaching, *Ecosystems*, 9, 453-462, <https://doi.org/10.1007/s10021-006-0051-z>, 2006.
- 630 Exner-Kittridge, M., Strauss, P., Blöschl, G., Eder, A., Saracevic, E., and Zessner, M.: The seasonal dynamics of the stream sources and input flow paths of water and nitrogen of an Austrian headwater agricultural catchment, *Sci. Total Environ.*, 542, 935-945, <https://doi.org/10.1016/j.scitotenv.2015.10.151>, 2016.
- FAO and WWC: Towards a water and food secure future: Critical perspectives for policy-makers, Food and Agriculture Organization of the United Nations (FAO) and the World Water Council (WWC) in support to the High Level Panel on Water for Food Security, Seventh World Water Forum in Daegu, South Korea, White paper, 76 pp., <http://www.fao.org/3/a-i4560e.pdf>, 2015.
- 635 Fasching, C., Ulseth, A. J., Schelker, J., Steniczka, G., and Battin, T. J.: Hydrology controls dissolved organic matter export and composition in an Alpine stream and its hyporheic zone, *Limnol. Oceanogr.*, 61, 558-571, <https://doi.org/10.1002/lno.10232>, 2016.

- 640 Fasching, C., Wilson, H. F., D'Amario, S. C., and Xenopoulos, M. A.: Natural Land Cover in Agricultural Catchments Alters Flood Effects on DOM Composition and Decreases Nutrient Levels in Streams, *Ecosystems*, 22, 1530-1545, <https://doi.org/10.1007/s10021-019-00354-0>, 2019.
- Findlay, S. E.: Increased carbon transport in the Hudson River: unexpected consequence of nitrogen deposition?, *Front. Ecol. Environ.*, 3, 133-137, [https://doi.org/10.1890/1540-9295\(2005\)003\[0133:ICTITH\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2005)003[0133:ICTITH]2.0.CO;2), 2005.
- 645 Gardner, K. K. and McGlynn, B. L.: Seasonality in spatial variability and influence of land use/land cover and watershed characteristics on stream water nitrate concentrations in a developing watershed in the Rocky Mountain West: Human impacts on spatial N patterns, *Water Resour. Res.*, 45, <https://doi.org/10.1029/2008WR007029>, 2009.
- Goodale, C. L., Aber, J. D., Vitousek, P. M., and McDowell, W. H.: Long-term decreases in stream nitrate: successional causes unlikely; possible links to DOC?, *Ecosystems*, 8, 334-337, <https://doi.org/10.1007/s10021-003-0162-8>, 2005.
- 650 Graeber, D., Gelbrecht, J., Pusch, M. T., Anlanger, C., and von Schiller, D.: Agriculture has changed the amount and composition of dissolved organic matter in Central European headwater streams, *Sci. Total Environ.*, 438, 435-446, <https://doi.org/10.1016/j.scitotenv.2012.08.087>, 2012.
- Griffiths, N. A., Tank, J. L., Royer, T. V., Warrner, T. J., Frauendorf, T. C., Rosi-Marshall, E. J., and Whiles, M. R.: Temporal variation in organic carbon spiraling in Midwestern agricultural streams, *Biogeochemistry*, 108, 149-169, <https://doi.org/10.1007/s10533-011-9585-z>, 2011.
- 655 Grizzetti, B., Bouraoui, F., de Marsily, G., and Bidoglio, G.: A statistical method for source apportionment of riverine nitrogen loads, *J. Hydrol.*, 304, 302-315, <https://doi.org/10.1016/j.jhydrol.2004.07.036>, 2005.
- Grolemund, G. and Wickham, H.: Dates and Times Made Easy with lubridate, *Journal of Statistical Software*, 40, 1-25, DOI: [10.18637/jss.v040.i03](https://doi.org/10.18637/jss.v040.i03), 2011.
- 660 Gu, S., Gruau, G., Dupas, R., Petitjean, P., Li, Q., and Pinay, G.: Respective roles of Fe-oxyhydroxide dissolution, pH changes and sediment inputs in dissolved phosphorus release from wetland soils under anoxic conditions, *Geoderma*, 338, 365-374, <https://doi.org/10.1016/j.geoderma.2018.12.034>, 2019.
- Gu, S., Gruau, G., Dupas, R., Rumpel, C., Crème, A., Fovet, O., Gascuel-Oudou, C., Jeanneau, L., Humbert, G., and Petitjean, P.: Release of dissolved phosphorus from riparian wetlands: Evidence for complex interactions among hydroclimate variability, topography and soil properties, *Sci. Total Environ.*, 598, 421-431, <https://doi.org/10.1016/j.scitotenv.2017.04.028>, 2017.
- 665 Gücker, B., Silva, R. C. S., Graeber, D., Monteiro, J. A. F., and Boëchat, I. G.: Urbanization and agriculture increase exports and differentially alter elemental stoichiometry of dissolved organic matter (DOM) from tropical catchments, *Sci. Total Environ.*, 550, 785-792, <https://doi.org/10.1016/j.scitotenv.2016.01.158>, 2016.
- 670 Halliday, S. J., Wade, A. J., Skeffington, R. A., Neal, C., Reynolds, B., Rowland, P., Neal, M., and Norris, D.: An analysis of long-term trends, seasonality and short-term dynamics in water quality data from Plynlimon, Wales, *Sci. Total Environ.*, 434, 186-200, <https://doi.org/10.1016/j.scitotenv.2011.10.052>, 2012.

- 675 Hedin, L. O., von Fischer, J. C., Ostrom, N. E., Kennedy, B. P., Brown, M. G., and Robertson, G. P.: Thermodynamic Constraints on Nitrogen Transformations and Other Biogeochemical Processes at Soil-Stream Interfaces, *Ecology*, 79, 684-703, DOI: 10.2307/176963, 1998.
- Hénault, C. and Germon, J. C.: NEMIS, a predictive model of denitrification on the field scale, *European Journal of Soil Science*, 51, 257-270, <https://doi.org/10.1046/j.1365-2389.2000.00314.x>, 2000.
- 680 Heppell, C. M., Binley, A., Trimmer, M., Darch, T., Jones, A., Malone, E., Collins, A. L., Johnes, P. J., Freer, J. E., and Lloyd, C. E. M.: Hydrological controls on DOC : nitrate resource stoichiometry in a lowland, agricultural catchment, southern UK, *Hydrol. Earth Syst. Sc.*, 21, 4785-4802, <https://doi.org/10.5194/hess-21-4785-2017>, 2017.
- Hill, A. R., Devito, K. J., Campagnolo, S., and Sanmugadas, K.: Subsurface denitrification in a forest riparianzone: Interactions between hydrology and supplies of nitrate and organic carbon, *Biogeochemistry*, 51, 193-223, <https://doi.org/10.1023/A:1006476514038>, 2000.
- 685 Humbert, G., Jaffrézic, A., Fovet, O., Gruau, G., and Durand, P.: Dry-season length and runoff control annual variability in stream DOC dynamics in a small, shallowgroundwater-dominated agricultural watershed, *Water Resour. Res.*, 51, 7860-7877, <https://doi.org/10.1002/2015WR017336>, 2015.
- Hytteborn, J. K., Temnerud, J., Alexander, R. B., Boyer, E. W., Futter, M. N., Fröberg, M., Dahné, J., and Bishop, K. H.: Patterns and predictability in the intra-annual organic carbon variability across the boreal and hemiboreal landscape, *Sci. Total Environ.*, 520, 260-269, <https://doi.org/10.1016/j.scitotenv.2015.03.041>, 2015.
- 690 Kaushal, S. S., Gold, A. J., Bernal, S., Johnson, T. A. N., Addy, K., Burgin, A., Burns, D. A., Coble, A. A., Hood, E., Lu, Y., Mayer, P., Minor, E. C., Schroth, A. W., Vidon, P., Wilson, H., Xenopoulos, M. A., Doody, T., Galella, J. G., Goodling, P., Haviland, K., Haq, S., Wessel, B., Wood, K. L., Jaworski, N., and Belt, K. T.: Watershed ‘chemical cocktails’: forming novel elemental combinations in Anthropocene fresh waters, *Biogeochemistry*, 141, 281-305, <https://doi.org/10.1007/s10533-018-0502-6>, 2018.
- 695 Lambert, T., Pierson-Wickmann, A.-C., Gruau, G., Jaffrezic, A., Petitjean, P., Thibault, J.-N., and Jeanneau, L.: Hydrologically driven seasonal changes in the sources and production mechanisms of dissolved organic carbon in a small lowland catchment, *Water Resour. Res.*, 49, 5792-5803, <https://doi.org/10.1002/wrcr.20466>, 2013.
- Le, S., Josse, J., and Husson, F.: FactoMineR: An R Package for Multivariate Analysis, *J. Stat. Softw.*, 25, 1-18, DOI: [10.18637/jss.v025.i01](https://doi.org/10.18637/jss.v025.i01), 2008.
- 700 Lemercier, B., Gaudin, L., Walter, C., Aurousseau, P., Arrouays, D., Schwartz, C., Saby, N. P. A., Follain, S., and Abrassart, J.: Soil phosphorus monitoring at the regional level by means of a soil test database, *Soil Use Manage.*, 24, 131-138, <https://doi.org/10.1111/j.1475-2743.2008.00146.x>, 2008.
- 705 Le Moal, M., Gascuel-Oudou, C., Ménesguen, A., Souchon, Y., Étrillard, C., Levain, A., Moatar, F., Pannard, A., Souchu, P., Lefebvre, A., and Pinay, G.: Eutrophication: A new wine in an old bottle?, *Sci. Total Environ.*, 651, 1-11, <https://doi.org/10.1016/j.scitotenv.2018.09.139>, 2019.

- Lintern, A., Webb, J. A., Ryu, D., Liu, S., Bende-Michl, U., Waters, D., Leahy, P., Wilson, P., and Western, A. W.: Key factors influencing differences in stream water quality across space, *Wiley Interdisciplinary Reviews: Water*, 5, e1260, <https://doi.org/10.1002/wat2.1260>, 2018.
- 710 Liu, W., Xu, X., McGoff, N. M., Eaton, J. M., Leahy, P., Foley, N., and Kiely, G.: Spatial and Seasonal Variation of Dissolved Organic Carbon (DOC) Concentrations in Irish Streams: Importance of Soil and Topography Characteristics, *Environmental Management*, 53, 959-967, <https://doi.org/10.1007/s00267-014-0259-1>, 2014.
- Luo, Y. and Zhou, X.: CHAPTER 5 - Controlling Factors. In: *Soil Respiration and the Environment*, Luo, Y. and Zhou, X. (Eds.), Academic Press, Burlington, <https://doi.org/10.1016/B978-012088782-8/50005-X>, 2006.
- 715 Mardhel, V. and Gravier, A.: Carte de vulnérabilité intrinsèque simplifiée des eaux souterraines du Bassin Seine- Normandie, Report BRGM/RP54148-FR, 92 pp., 2004.
- Martin, C., Aquilina, L., Gascuel-Oudou, C., Molénat, J., Faucheux, M., and Ruiz, L.: Seasonal and interannual variations of nitrate and chloride in stream waters related to spatial and temporal patterns of groundwater concentrations in agricultural catchments, *Hydrol. Process.*, 18, 1237-1254, <https://doi.org/10.1002/hyp.1395>, 2004.
- 720 Martin, C., Molénat, J., Gascuel-Oudou, C., Vouillamoz, J. M., Robain, H., Ruiz, L., Faucheux, M., and Aquilina, L.: Modelling the effect of physical and chemical characteristics of shallow aquifers on water and nitrate transport in small agricultural catchments, *J. Hydrol.*, 326, 25-42, <https://doi.org/10.1016/j.jhydrol.2005.10.040>, 2006.
- Melland, A. R., Mellander, P. E., Murphy, P. N. C., Wall, D. P., Mehan, S., Shine, O., Shortle, G., and Jordan, P.: Stream water quality in intensive cereal cropping catchments with regulated nutrient management, *Environ. Sci. Policy*, 24, 58-70, <https://doi.org/10.1016/j.envsci.2012.06.006>, 2012.
- 725 Mengistu, S. G., Creed, I. F., Webster, K. L., Enanga, E., and Beall, F. D.: Searching for similarity in topographic controls on carbon, nitrogen and phosphorus export from forest headwater catchments, *Hydrol. Process.*, 28, 3201-3216, <https://doi.org/10.1002/hyp.9862>, 2014.
- Merot, P., Squidant, H., Arousseau, P., Hefting, M., Burt, T., Maitre, V., Kruk, M., Butturini, A., Thenail, C., and Viaud, V.: Testing a climato-topographic index for predicting wetlands distribution along an European climate gradient, *Ecological Modelling*, 163, 51-71, [https://doi.org/10.1016/S0304-3800\(02\)00387-3](https://doi.org/10.1016/S0304-3800(02)00387-3), 2003.
- 730 Minaudo, C., Dupas, R., Gascuel-Oudou, C., Roubeix, V., Danis, P.-A., and Moatar, F.: Seasonal and event-based concentration-discharge relationships to identify catchment controls on nutrient export regimes, *Adv. Water Resour.*, 131, 103379, <https://doi.org/10.1016/j.advwatres.2019.103379>, 2019.
- Minaudo, C., Meybeck, M., Moatar, F., Gassama, N., and Curie, F.: Eutrophication mitigation in rivers: 30 years of trends in spatial and seasonal patterns of biogeochemistry of the Loire River (1980–2012), *Biogeosciences*, 12, 2549-2563, <https://doi.org/10.5194/bg-12-2549-2015>, 2015.
- 735 Moatar, F., Abbott, B. W., Minaudo, C., Curie, F., and Pinay, G.: Elemental properties, hydrology, and biology interact to shape concentration-discharge curves for carbon, nutrients, sediment, and major ions, *Water Resources Research*, 53, 1270-1287, <https://doi.org/10.1002/2016WR019635>, 2017.

- 740 Moatar, F., Flourey, M., Gold, A. J., Meybeck, M., Renard, B., Ferréol, M., Chandesris, A., Minaudo, C., Addy, K., Piffady, J., and Pinay, G.: Stream Solutes and Particulates Export Regimes: A New Framework to Optimize Their Monitoring, *Front Ecol Evol*, 7, <https://doi.org/10.3389/fevo.2019.00516>, 2020.
- Moatar, F. and Meybeck, M.: Riverine fluxes of pollutants: Towards predictions of uncertainties by flux duration indicators, *C.R. Geosci.*, 339, 367-382, <https://doi.org/10.1016/j.crte.2007.05.001>, 2007.
- 745 Mulholland, P. J. and Hill, W. R.: Seasonal patterns in streamwater nutrient and dissolved organic carbon concentrations: Separating catchment flow path and in-stream effects, *Water Resour. Res.*, 33, 1297-1306, <https://doi.org/10.1029/97WR00490>, 1997.
- Musolff, A., Fleckenstein, J. H., Opitz, M., Büttner, O., Kumar, R., and Tittel, J.: Spatio-temporal controls of dissolved organic carbon stream water concentrations, *J. of Hydrol.*, 566, 205-215, <https://doi.org/10.1016/j.jhydrol.2018.09.011>, 2018.
- 750 Musolff, A., Selle, B., Büttner, O., Opitz, M., and Tittel, J.: Unexpected release of phosphate and organic carbon to streams linked to declining nitrogen depositions, *Global Change Biol.*, 23, 1891-1901, <https://doi.org/10.1111/gcb.13498>, 2017.
- Mutema, M., Chaplot, V., Jewitt, G., Chivenge, P., and Blöschl, G.: Annual water, sediment, nutrient, and organic carbon fluxes in river basins: A global meta-analysis as a function of scale, *Water Resour. Res.*, 51, 8949-8972, <https://doi.org/10.1002/2014WR016668>, 2015.
- 755 Onderka, M., Wrede, S., Rodný, M., Pfister, L., Hoffmann, L., and Krein, A.: Hydrogeologic and landscape controls of dissolved inorganic nitrogen (DIN) and dissolved silica (DSi) fluxes in heterogeneous catchments, *J. Hydrol.*, 450-451, 36-47, <https://doi.org/10.1016/j.jhydrol.2012.05.035>, 2012.
- Perrin, C., Michel, C., and Andréassian, V.: Improvement of a parsimonious model for streamflow simulation, *J. Hydrol.*, 279, 275-289, [https://doi.org/10.1016/S0022-1694\(03\)00225-7](https://doi.org/10.1016/S0022-1694(03)00225-7), 2003.
- 760 Plont, S., O'Donnell, B. M., Gallagher, M. T., and Hotchkiss, E. R.: Linking carbon and nitrogen spiraling in streams, *Freshw. Sci.*, 39, 126-136, <https://doi.org/10.1086/707810>, 2020.
- Poisvert, C., Curie, F., and Moatar, F.: Annual agricultural N surplus in France over a 70-year period, *Nutr. Cycl. Agroecosyst.*, 107, 63-78, <https://doi.org/10.1007/s10705-016-9814-x>, 2017.
- Pregitzer, K. S., Zak, D. R., Burton, A. J., Ashby, J. A., and MacDonald, N. W.: Chronic nitrate additions dramatically increase the export of carbon and nitrogen from northern hardwood ecosystems, *Biogeochemistry*, 68, 179-197, <https://doi.org/10.1023/B:BIOG.0000025737.29546.fd>, 2004.
- 765 Preston, S. D., Alexander, R. B., Schwarz, G. E., and Crawford, C. G.: Factors affecting stream nutrient loads: a synthesis of regional SPARROW model results for the continental United States, *J. Am. Water Resour. As.*, 47, 891-915, <https://doi.org/10.1111/j.1752-1688.2011.00577.x>, 2011.
- 770 Raymond, S., Moatar, F., Meybeck, M., and Bustillo, V.: Choosing methods for estimating dissolved and particulate riverine fluxes from monthly sampling, *Hydrolog. Sci. J.*, 58, 1326-1339, <https://doi.org/10.1080/02626667.2013.814915>, 2013.

- Saby, N. P. A., Lemercier, B., Arrouays, D., Leménager, S., Louis, B. P., Millet, F., Schellenberger, E., Squidant, H., Swiderski, C., Toutain, B. F. P., Walter, C., and Bardy, M.: Le programme Base de Données des Analyses de Terre (BDAT) : Bilan de 20 ans de collecte de résultats d'analyses, *Etude et Gestion des Sols*, 21, 141-150, 2015.
- 775 Sandström, S., Futter, M. N., Kyllmar, K., Bishop, K., O'Connell, D. W., and Djodjic, F.: Particulate phosphorus and suspended solids losses from small agricultural catchments: Links to stream and catchment characteristics, *Science of The Total Environment*, 711, 134616, <https://doi.org/10.1016/j.scitotenv.2019.134616>, 2020.
- Sharpley, A. N., Chapra, S. C., Wedepohl, R., Sims, J. T., Daniel, T. C., and Reddy, K. R.: Managing Agricultural Phosphorus for Protection of Surface Waters: Issues and Options, *Journal of Environmental Quality*, 23, 437-451, 780 <https://doi.org/10.2134/jeq1994.00472425002300030006x>, 1994.
- Sobota, D. J., Harrison, J. A., and Dahlgren, R. A.: Linking dissolved and particulate phosphorus export in rivers draining California's Central Valley with anthropogenic sources at the regional scale, *J. Environ. Qual.*, 40, 1290-1302, <https://doi.org/10.2134/jeq2011.0010>, 2011.
- SoeS: NOPOLU-Agri. Outil de spatialisation des pressions de l'agriculture. Méthodologie et résultats pour les surplus d'azote et les émissions des gaz à effet de serre. Campagne 2010–2011. Ministère du Développement durable et de l'Énergie, 2013.
- 785 Taylor, P. G. and Townsend, A. R.: Stoichiometric control of organic carbon-nitrate relationships from soils to the sea, *Nature*, 464, 1178-1181, <https://doi.org/10.1038/nature08985>, 2010.
- Temnerud, J. and Bishop, K.: Spatial Variation of Streamwater Chemistry in Two Swedish Boreal Catchments: Implications for Environmental Assessment, *Environ. Sci. Technol.*, 39, 1463-1469, <https://doi.org/10.1021/es040045q>, 2005.
- 790 Thomas, O., Jung, A. V., Causse, J., Louyer, M. V., Piel, S., Baurès, E., and Thomas, M. F.: Revealing organic carbon–nitrate linear relationship from UV spectra of freshwaters in agricultural environment, *Chemosphere*, 107, 115-120, <https://doi.org/10.1016/j.chemosphere.2014.03.034>, 2014.
- Thompson, S. E., Basu, N. B., Lascrain Jr., J., Aubeneau, A., and Rao, P. S. C.: Relative dominance of hydrologic versus biogeochemical factors on solute export across impact gradients, *Water Resources Research*, 47, 795 <https://doi.org/10.1029/2010WR009605>, 2011.
- UNESCO, LIWQ.: International Initiative on Water Quality: promoting scientific research, knowledge sharing, effective technology and policy approaches to improve water quality for sustainable development - UNESCO Bibliothèque Numérique, 23 pp., <https://unesdoc.unesco.org/ark:/48223/pf0000243651>, 2015.
- Van Meter, K. J., Chowdhury, S., Byrnes, D. K., and Basu, N. B.: Biogeochemical asynchrony: Ecosystem drivers of seasonal concentration regimes across the Great Lakes Basin, *Limnol. Oceanogr.*, 9999, <https://doi.org/10.1002/lno.11353>, 2019.
- 800 Weigand, S., Bol, R., Reichert, B., Graf, A., Wiekenkamp, I., Stockinger, M., Luecke, A., Tappe, W., Bogena, H., Puetz, T., Amelung, W., and Vereecken, H.: Spatiotemporal Analysis of Dissolved Organic Carbon and Nitrate in Waters of a Forested Catchment Using Wavelet Analysis, *Vadose Zone J.*, 16, doi:10.2136/vzj2016.09.0077, 2017.

- Westphal, K., Musolff, A., Graeber, D., and Borchardt, D.: Controls of point and diffuse sources lowered riverine nutrient concentrations asynchronously, thereby warping molar N:P ratios, *Environ. Res. Lett.*, 15, 104009, <https://doi.org/10.1088/1748-9326/ab98b6>, 2020.
- Whitehead, P. G., Wilby, R. L., Battarbee, R. W., Kernan, M., and Wade, A. J.: A review of the potential impacts of climate change on surface water quality, *Hydrolog. Sci. J.*, 54, 101-123, <https://doi.org/10.1623/hysj.54.1.101>, 2009.
- Wickham, H.: *ggplot2: Elegant Graphics for Data Analysis*, Springer, 2016.
- 810 Wickham, H.: The Split-Apply-Combine Strategy for Data Analysis, *J. Stat. Softw.*, 40, 1-29, DOI: [10.18637/jss.v040.i01](https://doi.org/10.18637/jss.v040.i01), 2011.
- Wood, S. N.: *Generalized Additive Models : An Introduction with R*, Second Edition, Chapman and Hall/CRC, 2017.
- Zambrano-Bigiarini, M.: hydroGOF: Goodness-of-fit functions for comparison of simulated and observed hydrological time series. R package version 0.4-0. , DOI:10.5281/zenodo.839854., 2020.
- 815 Zarnetske, J. P., Bouda, M., Abbott, B. W., Saiers, J., and Raymond, P. A.: Generality of Hydrologic Transport Limitation of Watershed Organic Carbon Flux Across Ecoregions of the United States, *Geophysical Research Letters*, 45, 11,702-711,711, <https://doi.org/10.1029/2018GL080005>, 2018.