

Final response to referees

We would like to thank referees and the Associate Editor for their valuable comments that helped enhance the quality of the paper.

We took into account all the comments provided by the referees. All the modifications are described in the following text. Line numbers refer to those of the marked manuscript.

- Response to Referee 1

We thank Dr. Holzbecher for his careful and useful review of our manuscript. Below we respond (in bold type) to Dr. Holzbecher's specific comments (in normal type)

RC1: The paper is an excellent study in which different methodologies (hydrogeological, physical and geochemical) and data are combined to obtain a coherent view of the system. I suggest few technical corrections, which I list here:

Line 94: Table 1, do not justify text within table cells

Agree, changes made (line 95)

Line 272: Table 2, use centered format in all cells, including header

Agree, changes made (line 279)

Line 294: Table 3, do not justify text within table cells

Agree, changes made (line 301)

Line 346: leave blank between number and unit, 69 m h⁻¹

Agree, changes made (line 353)

Line 368: Figure 5, the 2x2 sub-figure design extends outside of the page margins; as the reader may want to see the details, I suggest to put sub-figures a-d vertically in a 4x1 design

We agree with the comment and we made the suggested changes. Subfigures in Figure 5 are now presented vertically (line 375)

Line 450: use italics for f_{fresh} and f_{salt} , in order to match with format in the equations

Agree, changes made (line 457)

Line 528: use italics for the f -factors, in order to match with format in equation 3

Agree, changes made (line 535)

- Response to Referee 2

We would like to thank Referee 2 for the constructive feedback and the suggestions to improve the manuscript. Below are our responses (in bold type) to the referee's specific comments (in normal type)

RC2: Authors used multi-tracer approach to understand occasional saltwater intrusions in a karst coastal aquifer in southern France. The topic fits with the journal scopes. In general the paper is well written.

Thank you for this positive feedback

My major concern is the novelty of this work. The methodology and approach are standard. From this point of view, the paper does not present any novelty. However, the novelty relies on the study area. But this point is briefly discussed in the paper. I am suggesting minor revision. I would like to ask the authors to revise the introduction to point out the novelty of this work by discussing more previous studies on understanding occasional saltwater intrusions in the study area and some previous works related to the methodology.

We revised the introduction to better reflect the novelty of our work, as suggested by Referee 2 (lines 48-50, 99-102 and 110-111). As pointed out by Referee 2, the novelty relies in the study area. To the best of our knowledge, the literature on occasional saltwater intrusions through a submarine karst spring is very limited. Our work stands out from the previous studies in the study area in part because our multi-tracer approach includes more tracers and consider multiple inversac events since 1967. Also, previous studies in the Thau basin focused on the hydrochemistry while our work is the first to combine a multi-tracer approach with hydrogeological data to fully describe this phenomenon. Our comprehensive approach was helpful in developing a new conceptual model of the site and provide insights on the management of the groundwater resources.

Other improvements:

- We updated the references list to comply with the HESS standards.
- Line 162, we changed “Paloc et Bonnet 1969” with “Bonnet and Paloc 1969” and updated the associated reference in the references list.
- Line 269, we changed the reference “Hemelsdael et al.” with “Ladouche et al 2019”.
- Line 332, we added the word “that” in the sentence.
- Line 464, we deleted the word “of”.
- Line 481, we deleted “(Fig.7)” as Figure 7 was already cited in the sentence.
- Line 559, we added the word “and” in the sentence for clarity.
- Line 559-600; we changed “Cl/B ratio as a function of B isotopic composition” with “B isotopic composition as a function of Cl/B ratio”.
- Line 602: we changed 52.3‰ with 43.8‰.
- Line 618, we deleted the word “enrichment”.
- Line 766: we changed the word “remainder” with “rest”.
- In the Supplement, we modified the title of Table S2 by changing “2018” with “1996-2000”.

Hydraulic and geochemical impact of occasional saltwater intrusions through a submarine spring in a karst and thermal aquifer (Balaruc peninsula near Montpellier, France)

Marie-Amélie Pétré^{1,2*}, Bernard Ladouche³, Jean-Luc Seidel¹, Romain Hemelsdaël⁴,
Véronique de Montety¹, Christelle Batiot-Guilhe¹, Claudine Lamotte³

¹HydroSciences Montpellier, Montpellier University, CNRS, IRD, Montpellier, France

²Now at North Carolina State University, Marine, Earth, and Atmospheric Sciences Department, Raleigh, NC 27695, United States

³BRGM, Montpellier University, Montpellier, France

⁴Geosciences Montpellier, Montpellier University, CNRS, Montpellier, France

*corresponding author: Marie-Amélie Pétré, marieamelie.petre@gmail.com

Abstract

Submarine springs are a common discharge feature of the karst aquifers along the Mediterranean coast. In some instances, occasional and localized saltwater intrusions can occur through the submarine spring and negatively impact the quality of the groundwater resource. The hydraulic and geochemical behavior of a submarine spring discharging into the Thau lagoon just offshore of the Balaruc peninsula near Montpellier, France has been characterized to determine the impacts of such phenomena to better understand the dynamics of a regional karst aquifer and improve its groundwater management. This work is based on both historical and new hydrogeological and geochemical data, illustrating 6 occasional saltwater intrusion events (from 1967 to 2014) in the Thau lagoon area (southern France).

Hydraulic perturbation of the aquifer is propagated instantly within the Balaruc-les-Bains peninsula and reaches a distance of about 5 km upgradient within 9 days. Comparison of hydraulic heads during seawater intrusion events in 2010 and 2014 indicates an aggravation of the phenomenon with an increase in hydraulic head variations.

In contrast, isotopic tracers ($^{87}\text{Sr}/^{86}\text{Sr}$, D/H, $^{18}\text{O}/^{16}\text{O}$,) and Rare Earth Elements (REE) demonstrate that the geochemical impact of these inversac events is only observed at the local scale, but is still perceptible several years after the event. For example, some of the thermal wells had not recovered their initial geochemical state 20 and 40 months after the last two inversac events (2010 and 2014, respectively), suggesting a geochemical legacy of this phenomenon within the complex karst system. By contrast, an adjacent deep karst compartment

located south of the study area is not affected by the saltwater intrusion and is characterized by distinctly different hydrodynamic behavior.

Overall, this work on occasional and localized saltwater intrusions constitutes a key step in understanding the dynamics of this complex karstic and thermal aquifer and will support the management of the groundwater resource.

1. Introduction

Submarine springs are common discharge features of karst systems along the Mediterranean coast (Bakalowicz, 2014; Fleury, 2005; Fleury et al., 2007; Stieglitz et al., 2013). Water from these springs can support economic development or be exploited for drinking water supply.

In certain instances, especially during periods of low discharge, flow can reverse and instead intrude into the upgradient karst aquifer, causing an adverse impact on the quality of the groundwater resource and the economic activities that depend on it (Arfib and Gilli, 2010; Drogue and Bidaux, 1986). This phenomenon, termed “*inversac*” in French (Gèze, 1987; Pinault et al., 2004), was previously described by Albéric (2004) and Joigneaux et al. (2011) in a related context of river backflooding into karst springs. However, occasional saltwater intrusions through a submarine spring is a largely understudied phenomenon.

The factors responsible for initiating and terminating inversac events in coastal areas are not particularly well understood and the recovery of carbonate aquifers after a seawater intrusion is poorly described (Han et al., 2015). Yet, it is necessary to understand the mechanisms controlling this phenomenon, as well as the long-term impact of occasional and localized saltwater intrusions, to appropriately manage the groundwater resources of such karst systems.

In this context, a hydraulic and geochemical study was undertaken to characterize the occasional and localized saltwater intrusions into the karst aquifer of the Thau basin. More specifically, our study addressed the following questions: (i) What are the hydraulic and geochemical conditions in the karstic aquifer during and after an inversac event? (ii) What are the triggering factors contributing to the occurrence of the inversac event? (iii) To what extent is the aquifer system able to recover from an inversac event and get back to its original geochemical state? (iv) What are the implications in terms of groundwater resources management?

The Thau hydrosystem in the Balaruc-les-Bains area is an example of a karst and thermal aquifer affected by occasional saltwater intrusions through a submarine spring. This complex

hydrologic system has been described by Aquilina et al. (1997, 2002, 2003), Doerfliger et al. (2001) and Ladouche et al. (2001). However, the causes and extent of the impact of the saltwater intrusions here remain poorly understood. The karst and thermal water resources across the Thau hydrosystem are exploited both for drinking water supply and spa activities. Balaruc-les-bains (Fig.1) is the largest spa in France with 55,000 visitors annually. In the past, water from the Vise submarine spring was diverted to the seashore and used for aquaculture activities. Additionally, the Thau lagoon supports an extensive shellfish aquaculture and fishery. Thus, the economic stakes are high in this area and water use conflict has the potential to arise between stakeholders during dry conditions, as a consequence of an increase in water demand, and the occurrence of occasional saltwater intrusions (Chu et al., 2014; La Jeunesse et al., 2015). Indeed, the karst system has been affected by six inversac events through the Vise submarine spring over the past 50 years (Table 1), five of which (1967,1983, 2008, 2010 and 2014) have been documented and described (Ladouche et al., 2011, 2019; Ladouche and Lamotte, 2015). Human activities, such as bauxite mining upstream from the Vise spring, and intervention on the spring itself are thought to have contributed to the saltwater intrusions over the 1967-1993 period by drawing down the hydraulic head of the karst system. Historically, these inversac events produced an increase in the water level and electrical conductivity at the observation points (thermal well and spring) in the Balaruc peninsula. A better understanding of this phenomenon is for local use purposes, but is also of strategic importance for water managers of the Issanka karst spring, a major drinking water supply in the area.

The last two inversac events in 2010 and 2014 lasted seven and five months, respectively. The 2014 inversac event led to the permanent abandonment of the Cauvy spring which provided drinking water to a local population of 30,000.

Date	Duration	Causal Factors	End of Episode
2014 (May-October)	5 months	Very low waters. No storm surge in the lagoon. Groundwater withdrawals	Temporary low water level in the lagoon combined with recharge in the karst aquifer
2010 (June-December)	7 months	Intermediate waters	Heavy rainfall
2008 (January-March)	3 months	Low waters and a pump test on thermal well F14	-
1993 (October)	24 days	Low waters. Human intervention on the submarine spring and pumping test on the Balaruc peninsula	Heavy rainfall
1984-1987 (May-February)	20 months	Low waters. Lagoon with high water level. Human intervention on the griffon of the submarine spring	Heavy rainfall
1967 (October) - 1969 (March)	17/18 months	Dry conditions. Bauxite mining at the Cambelliès site. Large pumping (up to 1000 m ³ h ⁻¹), that lowered the water level to -20 m asl	Cessation of groundwater pumping

96

97 Table 1. Summary of the observed occasional saltwater intrusions (inversac events) in the karst
98 and thermal aquifer of the Thau basin.

99 Previous research in the study area focused primarily on the hydrochemistry and failed to
100 consider the hydraulic impact of the inversac events. To fully describe this phenomenon, the
101 present study combined hydrogeological data and a multi-tracer approach for several inversac
102 events. A larger set of geochemical tracers was used. A multi-tracer approach was followed to
103 better define the origin and contribution of the thermal, karst and saltwater end-members
104 under different conditions. Water stable isotopes (D/H, ¹⁸O/¹⁶O) were used to quantify the
105 contribution of the saltwater end-member, whereas strontium isotopes (⁸⁷Sr/⁸⁶Sr) were used to
106 distinguish the origin of salinity (thermal or marine). Rare Earth Elements (REE) were used
107 to establish the chemical signature of the thermal wells in a reference context and to evaluate
108 the influence of the marine signature over time. Finally, the B- isotope signature (δ¹¹B) was
109 used to trace water-rock interaction initiated during inversac events. Overall, this approach
110 supported the development of a revised new conceptual model of the site and provided new
111 insights on the management of the groundwater resources.

112 2. Study Area and Geological Setting

113

Mis en forme : Gauche, Espace Après : 0 pt, Ne pas ajuster l'espace entre le texte latin et asiatique, Ne pas ajuster l'espace entre le texte et les nombres asiatiques

Mis en forme : Police :(Par défaut) Times New Roman, 12 pt, Non Gras

Mis en forme : Police :(Par défaut) Times New Roman, 12 pt, Non Gras

Mis en forme : Gauche, Espace Après : 0 pt, Interligne : simple, Ne pas ajuster l'espace entre le texte latin et asiatique, Ne pas ajuster l'espace entre le texte et les nombres asiatiques

114 The Balaruc peninsula is located along the Mediterranean coast of southernmost France, west
115 of the city of Montpellier, (Fig. 1a). The karst aquifer of the peninsula lies within the Middle
116 and Upper Jurassic carbonates that outcrop to the north in the Aumelas Causse and the Gardiole
117 Massif. These Jurassic carbonates are buried under Miocene to Plio-Quaternary deposits in the
118 eastern part of the Thau lagoon (Fig 1b). Karstification and burial of the main Jurassic aquifer
119 are the result of the geological history of the region described below.

120 A Mesozoic series of Triassic to Early Cretaceous age was deposited during Tethyan rifting
121 (Baudrimont and Dubois, 1977; Debrand-Passard, 1984). The Mesozoic carbonate platform
122 underwent uplift during the Mid-Cretaceous, leading to exhumation that caused erosion of the
123 Early Cretaceous series and strong karstification of the Jurassic carbonates

124 The Mesozoic series was successively deformed during the Pyrenean collision phase from
125 Upper Cretaceous to Eocene time (Arthaud and Laurent, 1995; Arthaud and Seguret, 1981;
126 Choukroune et al., 1973; Choukroune and Mattauer, 1978) and the later rifting of the Gulf of
127 Lion (Arthaud et al., 1977; Arthaud and Seguret, 1981; Benedicto et al., 1996; Maerten and
128 Séranne, 1995; Séranne, 1999; Thaler, 1962). Then, the sea level rise of the Early Miocene
129 (Burdigalian) deposited transgressive sediments across the region (Fig. 1b). The overall
130 sedimentary record associated with these Upper Cretaceous to Middle Miocene geological
131 events, includes a wide range of detrital and carbonate facies deposited in marine, lacustrine
132 and fluvial environments (Combes, 1990; Marchand, 2019) (Fig. 1a).

133 The sea level fall related to the Messinian Salinity Crisis caused major river incision across
134 southern France which, as a consequence, deepened the karst system (Clauzon, 1982; Hsu,
135 1973; Ryan, 1976). The Messinian paleovalley in the coastal area was then flooded during the
136 early Pliocene when the Mediterranean was connected again to the Atlantic Ocean. Recent sea
137 level fluctuations over the last 5 Ma and associated Plio-Quaternary deposits have generated
138 the present day morphology of the Thau lagoon.

139 Structurally, the Balaruc-les-bains-Bains area is located between both the major NE-trending
140 Cévennes and Nîmes faults, to the south of the Pyrenean Montpellier Thrust (Fig. 1a). The study
141 area is also affected by the presence of the E-W oriented Pyrenean Thau Thrust. The complex
142 fault network at the intersection with the Nîmes Fault allows to bring the Paleozoic basement
143 at shallow depth (less than 2000 m according to borehole data) in the Balaruc-Sète area (Fig.
144 1a)




Figure 1 (a) Simplified geological map showing the main structures of the Montpellier region. The box locates the Balaruc-les-bains study area, between the Montpellier thrust and the Nîmes Fault. (b) Detailed geological map of the eastern Thau lagoon. Sample sites indicated by the yellow dots in both maps correspond to the locations of the thermal wells, karst springs, and other piezometric drilling sites. The location of the ancient bauxite mine (Cambelliès) is indicated by the dashed oval in the northwest of the study area.

3. Hydrogeological conceptual model of the Thau hydrosystem

The Balaruc peninsula is located at the point of natural convergence of different types of waters including seawater from the Thau lagoon and Mediterranean Sea, cold karst groundwater sourced from the Aumelas Causse and Gardiolle Massif, and mineralized hot thermal water rising from depth in the Balaruc peninsula (Aquilina et al, 2002). These different types of water interact together in the Middle to Upper Jurassic carbonate reservoir. The Montpellier Thrust to the north of the study area (Fig. 1a) brings impermeable marly Lias to surface, which delimits the aquifer and acts as barrier to groundwater circulation. Groundwater flow follows a general N-S trend from the Aumelas Causse and Gardiolle Massif to the Balaruc peninsula and the Mediterranean Sea. A clear hydraulic connection from the Vène spring (Aumelas Causse) to Issanka spring (western slope of the Gardiolle Massif) has been demonstrated through artificial tracer testing (Paloc et Bonnet and Paloc, 1969; Ladouche et al., 2001). The main outlets of the Aumelas Causse karst network are the Vise submarine spring, the Issanka spring and the Vène spring which acts as an overflow outlet. The Vise submarine spring is located on the floor of the Thau lagoon at 30 m depth and constitutes the most downgradient discharge feature of the hydrological system. The western part of the Gardiolle Massif is drained by the Vise spring as well as Cauvy and Ambressac springs.

At the regional scale, four levels of groundwater circulations have been recognized by previous studies (Aquilina et al., 2003): 1) a surficial groundwater flow component originating from the recharge areas that has a residence time <20 years; 2) an intermediate groundwater flow system from the Aumelas Causse, that circulates below the Montbazin-Gigean basin and the Gardiole Massif and discharges at points along the continental shelf (residence time of 50 years); 3) a deep regional groundwater flow system directed from the Aumelas Causse to the Villeveyrac Basin where several deep drillings (Pézenas and La Castillonne, depth of 1000 m) present hot karstic waters with low mineralization (residence time of several thousand years); 4) a very deep (>2 km) paleo-karst water circulation mixed with an ancient seawater circulation both recharging the thermal reservoir that has residence times on the order of 100,000 years.

4. Material and methods

4.1 Hydrogeological Monitoring

Water level, specific electrical conductivity and temperature were measured at 10 locations on an hourly basis during the last three inversac events in 2008, 2010 and 2014. The time-series data used in this study come from the "national quantitative groundwater monitoring network of the AFB/BRGM (available on the ADES website: <https://ades.eaufrance.fr> for P4 La Balme (location 10166X0253) and CGE Tennis (location 10166X0212.) Other observation stations (F5, F6, F8, F9 and S12) are managed by Balaruc-les-Bains and the municipality of Sète and the Syndicat Mixte du bassin de Thau (El Cantou, F5-Issanka, Frescaly and Cauvy).

In order to compare the different inversac events, hydraulic head and electrical conductivity were expressed with respect to reference values observed before the inversac events, as discussed in the Results Section below. In Figures 3 and 5) b and 5d), the x-axis corresponds to the number of hours elapsed since the beginning of the inversac event (t_0). The y-axis represents the difference between the parameter (hydraulic head or electrical conductivity) measured during the inversac and the preceding respective reference values. The variable "delta H" is the difference between hydraulic head measured at time = "t" - hydraulic head observed at time = t_0 (before the inversac event). Thus, a positive value corresponds to an increase compared to the baseline condition and vice versa. This relative comparison permits the evaluation of the inversac event perturbations with respect to the initial conditions prevailing in the system and result in a more precise valuation of the transient states of the system.

Code de champ modifié

4.2 Geochemical and Isotopic Analysis

Geochemical data for the 1996-2000 period are from Aquilina et al. (2002, 2003) and Ladouche et al. (2001). They are considered as representative of reference geochemical conditions of the karst system. Indeed, these data were collected three to seven years after the 1993 inversac, which is considered as a sufficient time for the system to have returned to its equilibrium state given the short, 21-day interval of this event.

New geochemical data (major ions and REE compositions) and isotope analyses (D/H, $^{18}\text{O}/^{16}\text{O}$, ^3H , $^{87}\text{Sr}/^{86}\text{Sr}$ and d^{11}B) were determined for samples collected in October 2010, April and September of 2012 and March and August of 2018, i.e. during the 2010 inversac, 17 and 22 months after the end of the 2010 inversac and 3.5 years after the end of the 2014 inversac.

Temperature, pH, Dissolved Oxygen and Electrical Conductivity ($T_{\text{ref}}=25\text{ }^{\circ}\text{C}$) were measured in the field, using a portable pH meter, oxymeter and conductivity meter (WTW 3210).

Water sample for major and trace element analysis were filtered on-site with disposable PP syringe with a $0.22\text{ }\mu\text{m}$ Durapore membrane and stored in acid washed HDPE bottles. Aliquots for cations and trace elements were acidified to pH 2 with ultrapure HNO_3 (1‰ v/v). Samples for H- and O-isotope analysis were collected in 15 mL amber glass vials capped with airtight lids. One liter samples were collected for B- and Sr-isotope analysis in pre-cleaned HDPE bottles and were later filtered in a cleanroom through a $0.22\text{ }\mu\text{m}$ Durapore membrane in a pressurized Nalgene filtration unit, with samples for Sr-isotope analysis acidified with 1 % ultrapure HNO_3 . Samples for tritium measurement were collected in 1 L HDPE bottles. All samples were stored at $4\text{ }^{\circ}\text{C}$ before analysis. Chemical analyses were performed in the HydroSciences Montpellier laboratory at the University of Montpellier. Total alkalinity was measured by acid titration with 0.1N HCl. Major ion (Cl^- , NO_3^- , SO_4^{2-} , Ca^{2+} , Mg^{2+} , Na^+ , and K^+) were analyzed by ion chromatography (ICS 1000 Dionex®). Precision error was $< \pm 5\%$. After acidification with 1% HNO_3 , trace elements (Li, B, Sr, REE and U) were analyzed by inductively-coupled mass spectrometry using a Thermo Scientific® iCAP Q at the AETE-ISO technical platform of the OSU OREME at the University of Montpellier. The use of an in-line Argon Gas Dilution system permits the direct injection of highly mineralized samples without prior dilution. Precision error was typically $<5\%$. Fresh water reference material SLRS-6, and seawater reference materials CASS-6 and NASS-6 for trace metals (National Research Council, Canada) were analyzed every 20 samples to monitor analytical accuracy. Mean results are

within the range of certified uncertainties. Precision error for all analyses was typically <5%. The REE data are represented in profiles, after normalization to a reference geological material, the North American Shale Composite (NASC) for natural waters, which corresponds to an average sample of North American shales (Taylor and McLennan, 1985).

For stable isotopes analysis, samples collected in 2010 were analyzed at BRGM Laboratories using a Finnigan MAT 252 mass spectrometer, whereas those collected in 2012 and 2018 were measured on an Elementar Isoprime stable isotope mass spectrometer at the LAMA laboratory of HydroSciences Montpellier at the University of Montpellier. Calibration was performed by repeated analyses of in-house standards of known isotopic composition in alternation with samples. Oxygen ($^{18}\text{O}/^{16}\text{O}$) and hydrogen (D/H) isotope ratio measurements are expressed in parts per thousand (i.e. ‰) in the familiar δ notation relative to the Vienna Standard Mean Ocean Water (SMOW) standard, where $\delta = ([R_{\text{sample}}/R_{\text{standard}} - 1] \times 1000)$. Samples analyzed by BRGM have a precision of ± 0.8 ‰ for δD values and ± 0.1 ‰ for $\delta^{18}\text{O}$ values, whereas those determined at the University of Montpellier have an overall precision of ± 0.6 ‰ for both δD and $\delta^{18}\text{O}$ values. Tritium analyses were performed at Hydrogeology laboratory at the Avignon and Pays du Vaucluse University.

Sr-isotope analyses were made by thermal ionization mass spectrometry, at the BRGM Laboratories for the 2010 samples and at the Centre de Recherches Pétrographiques et Géochimiques in Nancy for the 2012 samples. Chemical separation of Sr was done using a Sr-Spec ion-exchange column that has a total blank <0.5 ng for the entire chemical separation procedure. Around 150 ng of purified Sr was loaded onto a tungsten filament and analyzed with an average internal precision of $\pm 10 \cdot 10^{-6}$ (2σ) using a Finnigan MAT262 multiple collector thermal ionization mass spectrometer. Measured $^{87}\text{Sr}/^{86}\text{Sr}$ ratios were normalized to a $^{86}\text{Sr}/^{88}\text{Sr}$ ratio of 0.1194. The reproducibility of $^{87}\text{Sr}/^{86}\text{Sr}$ ratio measurements was tested through replicate analyses of the NBS987 standard (0.710240) for which the mean value was $0.710232 \pm 22 \times 10^{-6}$ at the BRGM and $0.710262 \pm 13 \times 10^{-6}$ at the CRPG.

Boron isotopic ratios were measured with a Neptune+, Thermo Electron inductively-coupled mass spectrometer (at the BRGM laboratories for samples collected in 2010 (Guerrot et al., 2011) and at the AETE-ISO technical platform of the OSU OREME at University of Montpellier for the other samples. The average value determined for the NIST SRM 951 standard was 4.67 ± 0.0033 .

5. Results

5.1 Revised Conceptual Model for the Thau hydrosystem

The conceptual hydrogeological model of the Thau system at the local scale of the Balaruc-les-Bains peninsula, is summarized in Figure 2. Thermal waters rise from the deep reservoir along inferred faults in the vicinity of the Balaruc peninsula to reach the top of the Jurassic aquifer. Consistent with the recent data compilation and results acquired in the Balaruc-les-Bains area by the Dem'eaux Thau project (Ladouche et al., 2019), we propose that the Thau Thrust is likely to provide the main pathway for thermal water. The karst freshwater is found below this thermal lens as these cold waters are more dense. The diffuse intrusion of marine waters from the Mediterranean Sea into the Jurassic limestone corresponds to a saltwater wedge and is located below the karst waters which have a lower density. This saltwater wedge can nevertheless have a more complex geometry than that represented in Fig. 2, since this heterogeneous karst system is made up of multiple compartments. These contrasts in water density generate several hydraulic interfaces that change according to the different hydraulic heads resulting from continually- varying lagoon water levels, groundwater withdrawals and recharge from precipitation.

Linear depth (m)	EC (mS cm ⁻¹)	Temperature (°C)
85	16.5	25
145	1	23
223	1	37
237	1	37
249	2	37
297	3.2	37
389	35	37


Table 2 Water temperature and electrical conductivity from the exploratory drilling F13 at different depths. This inclined well (40°NW, N320°) is located near the F14 well (Fig.1b)

This hydrogeological setting illustrated in Figure 2 was confirmed by the F13 exploratory well (Table 2), that displayed contrasting electrical conductivity and temperature over depth.

Under normal hydrologic conditions (Fig. 2a), groundwater from the karst aquifer discharges into the Vise spring, creating a plume of freshwater in the lagoon. During an occasional episode

Mis en forme : Centré

287 of saltwater intrusion (or “inversac” event), the natural flow is reversed and saltwater from the
288 Thau lagoon flows into the Vise spring and enters the karstic aquifer via the Miocene cover
289 (Fig. 2b). The equilibrium between the different water bodies inside the Jurassic reservoir is
290 then strongly disrupted and the outflow of the karstic system through the Vise spring is
291 prevented. The characteristics of the thermal and karstic wells and spring in the Balaruc-les-
292 Bains peninsula are shown in Table 3.



294 Figure 2: Conceptual cross-section oriented NW-SE passing through thermal wells of the Balaruc peninsula, and
295 showing groundwater flows of the karst hydrosystem under (a) normal conditions and (b) inversac conditions.
296 Location of the cross-section is indicated in Figure 1a.

Name and BRGM location reference number	Main water type	Well depth (m)	Depth to the top of the karst aquifer (m)	Distance from the Vise submarine spring (m)
F5 (10165X0185)	Thermal	105	-72	325
F6 (10165X0251)	Karst/thermal	63.5	-58.5	330
F9 (10165X0252)	Thermal	120	-88	375
F3 (10165X0183)	Thermal	175	-65	380
F4 (10165X0184)	Thermal	55	-38	460
F14 (10165X0257)	Thermal	300	-170	585
F8 (10165X0234)	Thermal	407	-136	930
Cauvy spring(10165X0021)	Karst	0	0 (outcrop)	1000
CGE Tennis (10166X0212)	Karst	115	-95	2600
P4 La Balme (10166X0253)	Karst	100	0 (outcrop)	2800
Frescaly (10162X0194)	Karst	54	-22	3400
F7-Issanka (10162X0184)	Karst	58	-12	4000
F5-Issanka (10162X0136)	Karst	27	-14	4450
El Cantou (10162X0197)	Karst			4535

Mis en forme : Gauche

Mis en forme : Gauche

Mis en forme : Gauche

Mis en forme : Gauche

Mis en forme : Gauche

Mis en forme : Gauche

Mis en forme : Gauche

Mis en forme : Gauche

Mis en forme : Gauche

Mis en forme : Gauche

Mis en forme : Gauche

Mis en forme : Gauche

Mis en forme : Gauche

Mis en forme : Gauche

Mis en forme : Gauche

Table 3 Characteristics of the main wells and springs in the study area

5.2 Hydraulic impact of the inversac event

This section provides a detailed hydrodynamic analysis of the most recent inversac event in 2014 and then makes a comparison between the last three inversac events of 2008, 2010, and 2014.

The hydraulic impact of the inversac event of 2014 was observed at all Balaruc peninsula springs and observation wells (Fig.3). From the first hours after the intrusion of lagoon water, a sharp increase in water level is observed for both karst and thermal wells, although the impact is variable for different locations. The hydraulic impact was most significant for thermal well

313 F9, for which a +2.2 m increase in hydraulic head was observed during the first four days of
314 the inversac event. The magnitude of the hydraulic impact is similar for F5 and F6 wells, with
315 experienced respective increases in hydraulic head of +1.8 m and +1.9 m respectively. A lower
316 increase in hydraulic head of just +0.7 m during the first four days of the inversac event was
317 recorded at the Cauvy spring, situated only 1 km from the Vise submarine spring. For this
318 spring, the increase is also partly caused by the interruption of water pumping. The hydraulic
319 disturbance gradually increases over a 3-months period for the CGE Tennis and P4 La Balme
320 wells, reaching a maximum of 2 m at CGE and around 1 m at P4 La Balme piezometers after
321 1,500 h.

322 Two large rainfall events occurred in September and October 2014, some 3,400 h after the
323 beginning of the inversac event. These resulted in recharge to the aquifer that caused a sharp
324 increase in well water levels of +1.3 m at CGE Tennis and +1 m, at P4 La Balme.

325 The end of the inversac event occurred in October during the normal recession period of the
326 groundwater levels across the region. The thermal wells experienced a rapid decrease of
327 hydraulic head, which contrasted with the slow decrease of water levels in the CGE Tennis, P4
328 La Balme and F7-Issanka wells. After the inversac event, the water levels in CGE Tennis, P4
329 La Balme and Cauvy spring were observed in an intermediate position between the maximum
330 inversac and reference levels. In addition, the analysis of the water levels in the Thau lagoon
331 by Ladouche and Lamotte (2015) indicates that the end of the 2014 inversac resulted from
332 strong Tramontane winds that caused an abrupt decrease in the lagoon water level in the vicinity
333 of the Vise submarine spring. Thus, both a temporary condition of a low water level in the Thau
334 lagoon and high waters conditions in the karst aquifer following heavy rainfall events appear to
335 have contributed to the conclusion of the 2014 inversac event.

336 In the Villeveyrac Basin area, the evolution of piezometric levels during 2014 indicates that the
337 hydrology of this region was not affected by the inversac event (Ladouche and Lamotte, 2015).
338 The elevated piezometric level of some 70 m here compared to the Vène area (3 m) strongly
339 suggests a compartmentalization of the karst aquifer.

340

341

Mis en forme : Non Surlignage




Figure 3 Evolution of variations in hydraulic head (m) of wells and springs across the study area during the 2014 inversac event. The rainfall record during this time is shown at the top of the figure.

Figure 4 illustrates the dynamics of the hydraulic disturbance to the karst system caused by the 2014 inversac event across the Balaruc-les-Bains peninsula. This detailed analysis demonstrates the hydrologic connection between the Vise spring and the Issanka area. This perturbation propagates rapidly and reaches a distance of 3 km upgradient in less than 24 h (i.e. a velocity of 125 m h^{-1}). More specifically, the inversac causes an increase in hydraulic head of +1.5 m throughout the peninsula and +0.5 m to the north of the peninsula (Frescaly, CGE Tennis) in the first 24 h. This hydraulic perturbation then reaches the Issanka area in less than 72 h (i.e. a velocity of 69 m h^{-1}) and causes a +0.4 m increase in hydraulic head within 10 days.

In addition, Fig.4 clearly shows that the hydraulic perturbation follows a preferential pathway with a NNE orientation. A second minor flowpath follows a ENE orientation from the Vise spring towards Cauvy and Ambressac springs.




Figure 4 Spatial propagation of the hydraulic perturbation (hydraulic head increase in m) across the Balaruc-les-Bains peninsula during the 2014 inversac event at three times after initiation: T+24 h, T+72 h, and T+216 h.

Comparison between the 2008, 2010, and 2014 inversac events.

Comparison between the last three inversac events of 2008, 2010, and 2014 provides a basis for examining the hydraulic response to the phenomenon under different hydrological states. The 2008 inversac caused an increase in the hydraulic heads of all the wells and piezometers in the study area. For example, karst wells P4 La Balme and CGE Tennis showed respective increases in the hydraulic head of +2 m and +1 m.

The increase in the water levels reached the same wells during the 2014 inversac as in 2008, but was generally lower (more than 1 m for P4 La Balme and Frescaly and about 0.4 m in the Issanka area).

In contrast to the 2008 and 2014 events, the hydraulic impact of the 2010 inversac did not reach the Issanka area. An increase in hydraulic heads ranging from 1.5 to 2 m was only observed north of the Balaruc-les-Bains peninsula for the P4 La Balme and CGE Tennis wells. The explanation for this observation is discussed in the Discussion section.




Figure 5 (a) Hydraulic head (m) and (b) hydraulic head variation (m) of the karst well P4 La Balme during the 2010 and 2014 inversac events; electrical conductivity variation (mS cm^{-1}) for the thermal wells (c) F5 and (d) F9 during the 2008, 2010 and 2014 inversac events.

The hydraulic head of the karst well P4 La Balme was higher during the 2010 inversac (intermediate waters) than during the 2014 inversac (very low waters) reflecting contrasted

hydrologic conditions during these periods (Fig.5 a). However, the hydraulic head difference, with respect to the reference measured before the inversac (Fig. 5b), is higher in 2014 than in 2010. This suggests that the low water conditions magnify the hydraulic impact of inversac events. Similarly, electrical conductivity differences for the thermal wells F5 (Fig. 5c) and F9 (Fig. 5d) show a stronger response to the 2014 inversac than during both the 2008 and 2010 inversacs.

5.3 Geochemical impact of the inversac event

In addition to the hydraulic impact just described, inversac events also have a notable geochemical impact on the water bodies circulating within the karst system. As indicated in the discussion of the “Hydrogeological conceptual model” section, the salty and highly mineralized water of the lagoon mixes with thermal and karst waters to alter their quality. A multi-tracer approach was undertaken to determine the long-term hydrochemical impact of inversac events on the system that employed analysis of major ion contents, REE concentrations, and H-, O-, Sr- and B-isotopic compositions. The geochemical analyses reflect contrasting conditions: the 1996-2000 reference interval, the 2010 inversac event, and the 2012 and 2018 post-inversac period.

Groundwater characteristics- Major ions and water types

Outside of the Balaruc Peninsula, karst groundwater from Issanka spring and Pézenas deep borehole (738 m) is of a Ca-Mg-HCO₃ type, characterized by electrical conductivity (EC) values of about 500 $\mu\text{S cm}^{-1}$, in agreement with their karstic water type. The water from the deep borehole has a higher temperature (37.2 °C) than that of the Issanka spring (about 17.5 °C).

Within the Balaruc peninsula, karst water from the Cauvy spring is of the same water type, but displays elevated Cl contents and EC values of up to 222 mg L⁻¹ and 1200 $\mu\text{S cm}^{-1}$, respectively. In contrast, the thermal waters with temperatures that range up to 49.9°C are of a Na-Cl type with Cl concentrations of up to 7,900 mg L⁻¹. The Ambressac spring, located east of the Balaruc peninsula, is of mixed water type, with higher Cl concentration than at the Cauvy spring (up to 930 mg L⁻¹). Some samples also have high sulfate concentrations that approach 585 mg L⁻¹.

The Cauvy spring and the thermal wells F5, F6, and F9 were the most impacted by the 2010 inversac event, displaying a sharp increase of conductivity as well as high contents of Cl and Na. For example, Cl concentrations in the Cauvy spring and F6 reached levels 8 to 11 times above values typical of normal flow conditions (i.e. 1,240 mg L⁻¹ compared with 157 mg L⁻¹ for

Cauvy and 8611 mg L⁻¹ compare with 759 mg L⁻¹ for F6). For the Cauvy spring, which is used for potable water, this Cl content observed during the inversac event far exceeds the drinking water standard of 250 mg L⁻¹.

In a post-inversac context, Cl and Na concentrations observed for the Cauvy spring and F5, F6 and F9 wells in 2012 decreased significantly, with waters at these sampling points having returned to levels close to those of the 1996-2000 reference period. By contrast, Cl concentrations in wells F8 and F3 increased between 2010 and 2012, suggesting an increase in the relative contribution of the marine water component.

Karst water and saltwater mixing traced by water stable isotopes

As shown in Figure 6a, different waters present in the Balaruc peninsula hydrosystem are characterized by distinct δD and $\delta^{18}O$ values. Normal waters within the karst system lie close to the Local Meteoric Water Line ($\delta D = 8 \delta^{18}O + 14$; Ladouche et al., 1998), illustrating their meteoric origin. Waters sampled from thermal wells are more enriched in D and ^{18}O than the karst waters and are distributed along a karst-saltwater (i.e. lagoon or seawater) mixing line.

This result indicates that thermal waters result from a mixing between karst and marine end-members. However, stable isotopes information alone does not allow a determination of the origin of the marine end-member, which could be either lagoon water or modern or ancient seawater. Samples from the F8 thermal well show a slightly enriched stable isotopes signature compared to the other thermal wells, reflecting a greater contribution of the saltwater end-member in this area, as initially indicated by Aquilina et al. (2002).

F6 samples from the reference period are located within the karst water domain, confirming the strong influence of karst waters in this well (Aquilina et al., 2002) under normal flow conditions.

H- and O-isotopic signatures of groundwater from F5, F9, F8 and F6 thermal wells are shifted towards the saltwater end-member during the 2010 inversac. Then, in 2012 and 2018, their δD and $\delta^{18}O$ values decreased, but still remain in an intermediate position between the inversac maximum and the reference (1996-2000) period minimum. This result raises the question of the persistence of the chemical impact in the aquifer system following occasional saltwater intrusions.

Thermal wells F14, F5 and the thermal spring S12 show similar mixing proportions for the 2012 and 2018 post-inversac times. For F5 and S12, these mixing proportions are intermediate

between that of the 2010 inversac event and the 1996-2000 reference time. Groundwater samples of F3 well do not follow the same evolution, as the shift in its $\delta^{18}\text{O}$ value does not appear inversac related (Fig. 6b).




Figure 6 H- and O-isotopic compositions, expressed as δD and $\delta^{18}\text{O}$ values, of (a) karst waters, thermal waters, F8 well water, seawater, and Thau lagoon water, showing the Global Meteoric Water Line (GMWL) after Craig (1961) and the Local Meteoric Water Line (Ladouche et al., 1998-) for reference; (b) expanded view of the thermal well water array from (a) during reference, inversac, and post-inversac conditions.

Two end-member mixing calculations were made to model the geochemical impact of the 2010 inversac event on waters at the different sampling points across the study area, the first based on stable isotope ratios and the second using Cl concentrations measured in 2010, 2012, and 2018. The proportions of karst water (f_{fresh}) and saltwater (f_{salt}) in a groundwater sample were calculated using the following formula:

$$f_{\text{salt}} = \frac{C_{\text{sample}} - C_{\text{fresh}}}{C_{\text{salt}} - C_{\text{fresh}}} \quad \text{equation 1}$$

$$f_{\text{salt}} + f_{\text{fresh}} = 1 \quad \text{equation 2}$$

Results of the mixing calculations are included in the Supplement (Table S1). This calculation gives an indication of the magnitude of the karst water and saltwater component in each water sample, but does not provide additional insight into the origin of groundwater salinity. The calculated fractions of saltwater for the karst wells and spring are similar for the three tracers ($\delta^{18}\text{O}$, δD and Cl). However, for thermal wells with typical high Cl contents, the fractions obtained are distinct for the different tracers, indicating another source of Cl for these

Mis en forme : Police :Italique

Mis en forme : Police :Italique

469 wells. Indeed, as indicated by Aquilina et al. (2002), Cl in the thermal wells is thought to derive
470 from both seawater and the thermal water end-member, which is itself partly constituted of old
471 seawater.

472 **Inversac persistence revealed by REE**

473 The 14 members of the lanthanide group of elements (from La to Lu), commonly termed rare
474 earth elements (REE), constitute an effective tool to trace water origin, water/rock interaction
475 and mixing processes (Johannesson et al., 1997; Tweed et al., 2006; Zhan et al., 2013; Gil
476 et al., 2018). The REE are naturally present in natural waters at trace concentrations and their
477 normalized abundances, represented as a distribution pattern of individual REE ordered by their
478 atomic number, provide a means of visualizing the behaviour of the entire group of elements.

479 The REE profiles for seawater and groundwater from the thermal wells (S12, F8, F9, F5, F6)
480 and seawater are shown in Figure 7 for the pre-inversac reference and three post-inversac
481 situations (Fig. 7). With the exception of the F6 well, all patterns are generally flat and show a
482 small negative Ce anomaly. The F6 profile exhibits a more pronounced Ce anomaly and an
483 enrichment in heavy REE. These characteristics are those of limestone REE profiles, inherited
484 from the seawater REE profile and confirm the strong karst influence of F6 samples.




Figure 7 REE profiles for thermal wells at different times: the 1998 data are for the pre-inversac reference state, whereas the 2011, 2012 and 2018 profiles are for the three post-inversac situations.

REE contents during the 2011-2012 post-inversac period were generally lower than for the 1998 reference context. This is to say that, REE profiles in a post-inversac context lie in an intermediate position between the seawater profile and the reference profiles. Such an observation indicates that the thermal wells, at the time of sampling 20 months later in 2012, were still under the influence of the perturbations caused by the 2010 inversac event. The post-inversac profiles for 2012 and 2018 are quite similar, except for the F6 well, suggesting a similarity in extent of the 2010 and 2014 inversac events and post-event response of the hydrologic system. The REE profile for the F8 thermal well displays a distinctly different evolution, with an increasing shift away from the seawater profile over time. However, this well


498 is not hydraulically impacted by the occasional saltwater intrusion. The observed evolution does
499 not seem related to an increase in water withdrawals, but may instead reflect a relative decrease
500 in the contribution of thermal component over time.

501 **Characterization of the thermal and marine contributions from the strontium isotope**
502 **ratios $^{87}\text{Sr}/^{86}\text{Sr}$**

503 Consideration of strontium geochemistry allow the characterization of the thermal and marine
504 water contributions to the karst groundwaters. Sr-isotope ratios ($^{87}\text{Sr}/^{86}\text{Sr}$) for the groundwater
505 and lagoon samples are plotted as a function of strontium in Figure 8, where the 2010 and 2012
506 post-inversac data are compared with the 1996-2000 reference situation. It was assumed that
507 the waters from the F5 thermal well in the Balaruc peninsula during the reference interval are
508 representative of the thermal end-member, which is not affected by the karst or marine waters.
509 It is notable of the Sr-isotopic signature and Sr content of the CGE Tennis samples are relatively
510 stable in all contexts and fully representative of karst waters under confined conditions. The
511 Thau lagoon water sample is compositionally similar to that of the Mediterranean Sea and
512 corresponds to the saltwater end-member. The $^{87}\text{Sr}/^{86}\text{Sr}$ isotopic signature of karst waters and
513 thermal end-member are similar, indicative of derivation through water-rock interaction with
514 the Mesozoic carbonate aquifer. The karst end-member is different from the thermal end-
515 member by its lower Sr concentrations ($100\text{ }\mu\text{g L}^{-1}$), suggesting different water residence times.

516 As shown in Fig.8, all water samples plot within a ternary domain, confirming that the
517 geochemistry from the Balaruc peninsula groundwater can be explained by a 3-member mixing.
518 The karst waters from the Issanka spring that drain the Aumelas Massif have a lower $^{87}\text{Sr}/^{86}\text{Sr}$
519 ratios and Sr contents. The Issanka isotopic signature defines an up-gradient end-member for
520 the karst system, unaffected by mixings of the thermal and seawater that characterizes the
521 Balaruc peninsula. Waters from the P4 La Balme piezometer had a very high $^{87}\text{Sr}/^{86}\text{Sr}$ ratio in
522 2012, almost certainly reflecting the influence of rainfall infiltration since this sample site is
523 located at the outcrop of the aquifer and may be representative of surficial recharge water to the
524 karst aquifer.

525



526

527 Figure 8 Plot of $^{87}\text{Sr}/^{86}\text{Sr}$ ratio versus Sr concentration (as $1/\text{Sr}$) for springs and wells of the Balaruc peninsula
528 for different hydrogeological situations: pre-inversac reference (green), 2010 inversac (red) and 2012 post-
529 inversac (yellow).
530

531 The contribution of each end-member (saltwater, thermal water and karst water) to groundwater
532 samples under different hydrogeological situations related to inversac events were estimated
533 through 3-component mass balance equations following Faure (1986):

534
$$f_{\text{salt}} + f_{\text{karst}} + f_{\text{therm}} = 1$$
 equation 3

535 where:

536
$$(^{87}\text{Sr}/^{86}\text{Sr})_{\text{sample}} = f_{\text{salt}} (^{87}\text{Sr}/^{86}\text{Sr})_{\text{salt}} + f_{\text{karst}} (^{87}\text{Sr}/^{86}\text{Sr})_{\text{karst}} + f_{\text{therm}} (^{87}\text{Sr}/^{86}\text{Sr})_{\text{therm}}$$

537 Results of the mixing calculations are in the Supplement (Table S2).

538 The Sr-isotopic composition of the Cauvy spring is quite variable during the reference interval,
539 with samples aligned along the karst-saltwater mixing line in Figure 8. The contribution of the
540 saltwater end-member to the spring discharge is negligible under normal flow conditions, with

Mis en forme : Police :Italique

Mis en forme : Police :Italique

Mis en forme : Police :Italique

541 the Cauvy isotopic signature very close (99%) to the karst end-member. However, the Cauvy
 542 spring is strongly impacted by the inversac phenomenon. Although the karst end-member
 543 remains dominant at $88.1 \pm 0.2 \%$, the contribution of saltwater end-member increases to $8.1 \pm$
 544 0.5% and that of the thermal end-member reaches about $3.8 \pm 0.3 \%$ during the 2010 inversac
 545 event. The contribution of thermal waters to the Cauvy spring is not surprising, as it is located
 546 close to the old thermal spring (S12), but the large contribution of the saltwater end-member
 547 during inversac events implies a preferential groundwater flow axis between the spring and the
 548 Vise submarine spring. Pinault et al. (2004) have shown that variations in the contribution of
 549 thermal waters may be related to the existence of a piston effect on the thermal reservoir,
 550 particularly following an intense recharge event.

551 Thermal well waters of the Balaruc peninsula are all aligned along the thermal-saltwater end-
 552 member mixing line in Figure 8, except for F3, F4 hotel, and F6 waters, each of which
 553 documents a significant contribution from the karst end-member ($87.9 \pm 0.3 \%$, $89.8 \pm 0.2 \%$
 554 and $53.7 \pm 0.6 \%$, respectively).

555 The evolution of the isotopic signature for the F5, F9 and F14 borehole waters shows a similar
 556 pattern. Thermal water $^{87}\text{Sr}/^{86}\text{Sr}$ ratios increase during the 2010 inversac event, but then
 557 decreases in 2012 to values intermediate between the inversac condition and normal state. These
 558 observations suggest that waters from this group of thermal wells were strongly affected by the
 559 2010 inversac and had not returned to their 1996-2000 pre-inversac Sr-isotope signature by the
 560 2012 post-inversac situation.

561 The contribution of the Thau Lagoon end-member increased to $8.3 \pm 1.3 \%$ for the F9 waters
 562 during the 2010 inversac, the thermal end-member was $91.3 \pm 0.6 \%$, the karstic contribution
 563 being negligible ($0.4 \pm 1.9 \%$). The contribution of the Thau lagoon end-member decreased to
 564 $5.7 \pm 1.2 \%$ during the 2012 context.

565 Reporting on water collected from the F8 borehole, Aquilina et al. (2002) noted a seawater
 566 contribution to the thermal waters in this area of the Balaruc peninsula, which we have
 567 estimated to be $8.8 \pm 1.3 \%$ for the 1996-2000 pre-inversac reference period.

568 The strontium concentrations and $^{87}\text{Sr}/^{86}\text{Sr}$ ratios of F8 well waters were very similar in 2010
 569 and in April and September of 2012. Although the contribution from the saltwater end-member
 570 during the 2010 inversac was greater than the reference condition ($18.0 \pm 1.6 \%$ instead of 8.8
 571 $\pm 1.3 \%$), and remained between $17.5 \pm 1.6 \%$ to $21.2 \pm 1.8 \%$ from April to September 2012.
 572 The thermal well F8 does not seem to be impacted by a direct intrusion of lagoon water during
 573 the inversac event but instead influenced by seawater already within the hydrosystem.

Water at the F6 borehole was the most impacted by the 2010 inversac. During this time, the contribution from the saltwater end-member sharply increased from 1.6 ± 0.1 % to 35.4 ± 1.8 % between 1996 to 2010 then decreased to 0.6 ± 0.1 % in 2012, a value below the 1996 to 2000 reference interval. Waters from the F6 borehole are representative of the upper part of the Jurassic aquifer and demonstrate a significant contribution of the karst end-member (94.1 ± 0.1 % to 91.2 ± 0.2 % in 1996 and 2012). This karst contribution remains significant during an inversac context (53.7 ± 0.6 %). Thus, waters from the F6 well illustrate the range of fluctuations of the thermal component.

By contrast to the F6 well, the three end-member contributions are relatively stable for the waters of Ambressac spring. Indeed, water samples display a large variability, with a saltwater end-member contribution of between 1.1 ± 0.1 % and 2.7 ± 0.2 % and a karst contribution between 94.4 ± 0.1 % and 91.5 ± 0.1 % over the reference interval (1996-2000). The thermal contribution ranges between 4.4 ± 0.1 % and 5.8 ± 0.2 % during normal times.

Boron Isotope Tracking of Water-rock interactions

A lack of knowledge of the baseline B-isotopic signature during normal flow conditions does not allow to discuss the origin of boron and its quantification for thermal borehole waters.

However, examination of $\delta^{11}\text{B}$ variations as a function of B/Cl ratio clearly identifies water interactions with clay minerals. Indeed, boron is present in aqueous solution, in the form of boric acid $\text{B}(\text{OH})_3$ and borate ions $\text{B}(\text{OH})_4^-$. The distribution between these two species depends on pH, temperature and salinity (Hershey et al., 1986; Hakihana, 1977). For the boron isotopes, ^{11}B is preferentially incorporated into boric acid whereas ^{10}B has a greater affinity for the borate ion. During the water-rock interaction, $\text{B}(\text{OH})_4^-$ will be preferentially adsorbed onto clay minerals or organic matter, resulting in a decrease of water B concentration and a consequent enrichment of ^{11}B in residual water.

Figure 9 shows the B isotopic composition ~~Cl/B ratio~~ as a function of ~~Cl/B ratio~~ B isotopic composition. $\delta^{11}\text{B}$ of the Thau Lagoon is 39.1 ‰, in agreement with the signature of the modern seawater of 39.5 ‰; (Aggarwal et al., 2004; Aggarwal and Palmer, 1995). By contrast, karst waters display a range of $\delta^{11}\text{B}$ values from 23.1 to ~~52.3~~43.8 ‰, but are strongly ^{10}B enriched relative to seawater signature, whereas, thermal waters have $\delta^{11}\text{B}$ values of up to 44.3 ‰ that are similar to or higher than the seawater signature.


605 There is a good alignment of the data points in Fig. 9, which illustrates the process of water/rock
606 interaction and indicates that the B in the waters across the Balaruc peninsula is derived from
607 clay sediments. That $^{11}\text{B}/^{10}\text{B}$ ratios are well above the seawater values, suggests that water from
608 the Thau lagoon interacts with clay sediments within the karst system during the occasional
609 intrusion into the carbonate aquifer. An exploration dive of the bottom of the Vise spring in
610 April 2017 showed the absence of sediment on the walls and bottom of the spring. The inversac
611 phenomenon therefore seems to be accompanied by a significant intrusion of sediments (clayey
612 marine mud, rich in organic matter). This phenomenon seems to contribute to initiate the
613 process of boron isotope fractionation. This process could also occur within the silty-clay
614 Miocene unit during the transfer of water from the lagoon to the Jurassic aquifer. This Miocene
615 formation overlies the Jurassic aquifer and is about 30 m thick at the Vise spring. The high
616 isotopic enrichment observed for the F3 borehole waters can be explained by their high pH
617 values (8.3) and thus a relatively higher proportion of the $\text{B}(\text{OH})_4^-$ species, leading to a higher
618 adsorption on the solid phase and a higher enrichment of $\delta^{11}\text{B}$ of the residual water.

619 Boron isotopes are used to identify the chemical modification of the water that enters the system
620 and interacts with the clay matrix. There is therefore a modification of the water chemistry
621 before it even enters the karst aquifer, and this modification is initiated during the saltwater
622 intrusion.

Mis en forme : Exposant

Mis en forme : Exposant

623



624

Figure 9: Plot of $\delta^{11}\text{B}$ versus Cl/B ratio for different hydrogeological contexts: 2010 (inversac times in red), 2012 and 2018 (post-inversac times in yellow and blue)

6. Discussion

6.1 Implications for Dynamics of the Thau hydrosystem

Extent of the hydraulic impact and conditions required to reach the upgradient area of Issanka

Analysis of the hydraulic response of the Thau hydrosystem to the last inversac events provides insights into the dynamics of the system and a better understanding of the underlying controlling mechanisms. For example, the 2010 inversac did not impact the upgradient area of Issanka in contrast to the 2008 and 2014 inversac events. This is attributed to the fact that the karst system in 2010 was less in deficit than at the time of the 2008 and 2014 events, when very low flow conditions prevailed. For example, piezometric levels measured for the Issanka area in 2008 and 2014 were lower than in 2010 (i.e. 6.82 m in 2008 and, 6.89 m in 2014 and, 9.33 m in 2010). Considering that the overflow level of the Issanka spring is at ~ 9.5 m, this implies that the hydraulic impact in the Issanka compartment is controlled by the pressure head difference between the upstream (Issanka) and downstream (Balaruc peninsula) compartments. Piezometric monitoring at these different times illustrates that the occurrence of the inversac phenomenon is mainly determined by hydrogeological processes that occur within the downstream compartment of the karst system. Water level of the Thau lagoon is one of the controlling parameters. At inversac initiation in 2008, the water level in the Thau lagoon was higher than in 2010 at 1 m compared to just 0.5 m, but then only of 0.2 m in 2014. For the inversacs of 2008 and 2010, the sudden rise in lagoon waters of + 0.5 m and + 0.4 m, respectively, in a few hours is considered to be the triggering factor. By contrast, no sudden change in the water level of lagoon was observed in 2014. The triggering mechanism of this event is still poorly understood, however it is likely that the withdrawals from the aquifer for drinking water and irrigation purposes as well as the low recharge during the previous winter may be the explanatory factors. Thus, two factors would need to be combined for an inversac to reach the upgradient area of Issanka: (i) first, low water conditions that produces a very low hydraulic head throughout the aquifer. The hydraulic head would need to be lower than 7 m in the Issanka area, i.e. below than the hydraulic head imposed by the saltwater intrusion; (ii) an event-specific triggering

Mis en forme : Espace Après : 0 pt

657 mechanism, such as a sudden rise in the water level of the Thau lagoon or groundwater
658 withdrawals.

659 The intrusion of lagoon waters in the system causes a 2-phase response within the Thau
660 hydrosystem:

661 1) A short initial transient phase of a few days duration during which there is a
662 reorganization or displacement of the different karst, thermal and marine water bodies
663 within the hydrosystem,

664 2) A longer transient phase during which the hydrosystem reaches a new equilibrium (in
665 30 days in 2008 and 2010, 40 days in 2014) with clear evidence of water mixing causing
666 important changes in the physico-chemical parameters of waters from the sampling
667 points F5, F6, F9 and the Cauvy spring.

668

669 **The impact on water quality is local whereas the hydraulic impact is perceived at the**
670 **regional scale**

671 Results in section 5.1 showed that the hydraulic impact of an inversac event is instantly
672 propagated throughout the Balaruc peninsula and the effects of such events can also reach
673 longer distances into the karst system than previously described. The return to the equilibrium
674 situation occurs within days after a significant recharge event and is much more rapid than the
675 dissipation of the chemical impact of the inversac event. Indeed, the 2010 and 2014 inversac
676 events lasted about 6 months and had a persistent chemical impact on the hydrosystem. It stayed
677 perceptible on the quality of the water in certain compartments from 20 months to 42 months
678 after the occasional inversac intrusions ceased.

679 Even though the geochemical conditions within the hydrosystem were not monitored during
680 the 2014 inversac event, the 2018 data are similar to those for 2012, suggesting that the
681 hydrosystem did not reach its geochemical reference conditions and was still under a state of
682 influence by the latest inversac event.

683 Geochemical analysis indicates that the 2010 inversac had the greatest impact on waters from
684 the F6 well and the Cauvy spring, thus confirming the existence of a preferential groundwater
685 flowpath between the Vise spring and those points.

686 The physico-chemical disturbances resulting from inversac event are not observed at sampling
687 points CGE Tennis, P4 La Balme [and](#), Issanka, located in the upgradient area of the Balaruc

688 peninsula. The geochemical impact is, therefore, restricted to within a 1-kilometer radius from
689 the Vise spring (Fig. 10).

690 This slow geochemical recovery is best explained by the characteristic internal heterogeneity
691 of the karst aquifer, which is typically described in terms of a triple porosity model comprising:
692 (i) matrix or primary porosity, (ii) fracture porosity and (iii) conduit porosity (Palmer et al.,
693 1999; Martin and Scream, 2001). Influx of inversac water into the matrix porosity of the karst
694 aquifer, its storage, and then prolonged time of removal and return to fracture and conduit flow
695 explains the persistent fingerprint of the saltwater intrusions in the Balaruc peninsula.

696 Wells F8 and F10 in the southern part of the Balaruc peninsula were not affected by the inversac
697 events. This part of the hydrosystem corresponds to a deeper compartment in the karst system,
698 where the top of the Jurassic aquifer is 136 m deep in the F8 well and 195 m deep in the F10
699 well, which appears to be less hydraulically connected to the rest of the peninsula. Previous
700 studies postulated that an indirect impact of an inversac event might occur (Ladouce et al.,
701 2012). However, the increasing mineralization of F8 waters over time from 20 mS cm⁻¹ in 1996
702 to 26.5 mS cm⁻¹ in 2018 (Figure S1 in the Supplement) suggests that waters from this well are
703 not influenced by the occasional saltwater intrusions into the hydrosystem. This distinct
704 evolution cannot be explained by pumping withdrawals as well withdrawals rates in the F8
705 thermal well were steady over the period of inversac monitoring. However, a decrease in the
706 recharge rate from the karst system could result in an increase in the proportion of the thermal
707 or marine end-member contributions. Recent monitoring of springs (Ambressac and S12) will
708 address this question in the Dem'Eaux Thau project.

709 A comprehensive review of historical inversac events undertaken as part of this study showed
710 that there is a hydraulic connection between the Villeveyrac Basin and the Balaruc peninsula.
711 The relationship between these two regions is clearly demonstrated as dewatering of a bauxite
712 mine in the Cambelliès area (Fig.1) triggered the 1967 inversac event.




Figure 10- Bloc diagram of the study area showing the spatial extent of the hydraulic and geochemical impacts.

6.2 Implications on groundwater management:

Results presented here illustrates the fragile equilibrium that exists between different groundwater bodies in the Thau hydrosystem. During low flow conditions, any modification of the exploitation regime of the system can cause an inversac event and cause deleterious long-term consequences.

Historically, the 1967 and 2010 inversac events led to the complete cessation of withdrawal from the Cauvy spring, followed by its subsequent abandonment because the quality of this major drinking water supply was strongly impacted by the inversac events.

The Issanka spring is another major source of drinking water in the area for a population of 40,000. This study recognized the conditions needed for the hydraulic impact of an inversac event to reach the Issanka well-field. Such new understanding can support water management operations by facilitating the prediction of the arrival of an inversac disturbance, using the criteria highlighted in section 6.1.

While the southern compartment of the hydrosystem has not been impacted by inversac events, the gradual increase in mineralization observed for the F8 thermal well has the potential to affect its quality over time. In order to sustain the water quality of this thermal well, further studies should investigate the specific dynamics of this karstic compartment and the underlying mechanism explaining this observation. More specifically, the hydraulic connection between this deep compartment with the Sète thermal system south of the study area should be investigated.

735 Finally, unlike classical saline intrusion, the inversac events mobilize sediments loaded with
736 organic matter during saltwater intrusions from the Vise spring. This intrusion of sediments
737 may have an additional impact on the water quality within the karstic hydrosystem as suggested
738 by B-isotopic signatures. Future studies could explore the fate and transport of the organic
739 matter in the karst system.

740

741 **7. Conclusion**

742 This study investigated the impact of occasional saltwater intrusions (inversac in French) into
743 the karst hydrosystem of the Balaruc peninsula and provides an improved understanding of the
744 hydrodynamics and recovery of the aquifer from saltwater intrusions. Differences were
745 observed between the hydraulic and geochemical responses, both spatially and temporally. The
746 karst, thermal and marine water bodies within different compartments of the karst system
747 coexist within a very delicate equilibrium that is disturbed by the rapid intrusion of lagoon
748 waters during inversac events. While the hydraulic impact of such saltwater intrusions is
749 immediate and manifest over a distance of some 5 km within the hydrosystem, the geochemical
750 impact focused within a 1 km radius around the Vise spring but temporally persistent.
751 Geochemical tracers showed that the hydrosystem did not reach its pre-inversac reference state
752 20 months and 40 months after the occurrence of inversac of 6-months duration. The slow
753 geochemical recovery of the Balaruc peninsula karst system reflects the triple-porosity
754 character of the karst aquifer. Additionally, preliminary results suggested that the modification
755 of the groundwater chemistry is initiated by an interaction process with clay sediments and
756 organic matter during the saltwater intrusion event and possibly during the subsequent transfer
757 of seawater to the karst aquifer through the Miocene formation. This study demonstrates the
758 potential to couple physico-chemical, hydrogeological and isotopic data to understand the
759 complex inversac phenomenon and recognize preferential flowpaths during such events.

760 The results of this study support groundwater management across the Balaruc peninsula. The
761 triggering factors of the saltwater intrusions were highlighted as well as the conditions needed
762 for the saltwater intrusion to hydraulically impact the upgradient area, which is a major source
763 of drinking water supply. While the water quality of some of wells across the Balaruc peninsula
764 are strongly impacted by the inversac and retain a geochemical signature of the event, the
765 thermal wells located in the southern portion of the hydrosystem are not affected by inversac
766 events, suggesting hydraulic isolation from the ~~remainder-rest~~ of the Balaruc peninsula.

8. Data availability

The geochemical and isotopic data are accessible at <https://doi.org/10.5281/zenodo.3893897>

9. Supplement link

The supplement related to this article is available online at

10. Authors contribution

BL, JLS and CL designed the study and data collection plan. MAP, JLS and BL conducted the field work, collected the hydrogeological and geochemical data, and performed the data analysis. MAP prepared the paper with contributions from RH and JLS. All authors reviewed and edited the manuscript.

11. Competing interests

The authors declare that they have no conflict of interest

12. Acknowledgements

This work was funded by a FEDER-CPER grant (Agence de l'Eau, Balaruc-les-Bains, SMBT, and BRGM) through the Dem'Eaux Thau project. Authors are thankful to the municipalities of Balaruc les Bains and Pézenas for continuous support during the 2018 field work. Thanks are due to Franck Bujaldon (city of Pézenas), Ludovic Sarrou (Thermal spa, Balaruc les Bains), Nathalie Masscheleyn (Balaruc les Bains municipality), David Mimard (Suez) and Vincent Durand (Antéa) for granting access to the thermal boreholes, karst springs and piezometers. We are thankful to Jean-Gilbert Muller from President Electronics for providing access to the Ambressac spring. Gilles Lorente and Kevin Buttaró (Syndicat Mixte du Bassin de Thau) are thanked for their technical support during the March 2018 field campaign. We also acknowledge Nicolas Patris (HydroSciences Montpellier) for his help and useful discussion on stable isotopes. Remi Freydier (HydroSciences Montpellier) and graduate student Hikma Kassime are also thanked for their valuable contribution on the B isotopic analyses. We thank Russell Harmon (North Carolina State University) for a thorough review of an earlier version of the manuscript.

13. References

795 Aggarwal, J. K. and Palmer, M. R.: Boron isotope analysis. A review, Analyst, 120(5), 1301–
796 1307, 1995.

797 Aggarwal, J. K., Mezger, K., Pernicka, E. and Meixner, A.: The effect of instrumental mass
798 bias on $\delta^{11}\text{B}$ measurements: a comparison between thermal ionisation mass spectrometry and
799 multiple-collector ICP-MS, International journal of mass spectrometry, 232(3), 259–263, 2004.

800 Albéric, P.: River backflooding into a karst resurgence (Loiret, France), Journal of Hydrology,
801 286(1–4), 194–202, 2004.

802 Aquilina, L., Deluchat, V., Brach, M., Bakalowicz, M., Le Strat, P. and Giraud, F.: Etude
803 géochimique des eaux souterraines autour du bassin de Thau. Rap. BRGM R39530, 58 p., 15
804 fig., 4 tabl., 2 annexes., 1997.

805 Aquilina, L., Ladouche, B., Dörfliger, N., Seidel, J. L., Bakalowicz, M., Dupuy, C. and Le Strat,
806 P.: Origin, evolution and residence time of saline thermal fluids (Balaruc springs, southern
807 France): implications for fluid transfer across the continental shelf, Chemical Geology, 192(1–
808 2), 1–21, 2002.

809 Aquilina, L., Ladouche, B., Doerfliger, N. and Bakalowicz, M.: Deep water circulation,
810 residence time, and chemistry in a karst complex, Groundwater, 41(6), 790–805, 2003.

811 Arfib, B. and Gilli, E.: Karst côtier et sources sous-marines-fonctionnement et exploitation.
812 Dans Audra P. Grottes et karst de France., 2010.

813 Arthaud, F. and Laurent, P.: Contraintes, déformation et déplacement dans l'avant-pays Nord-
814 pyrénéen du Languedoc méditerranéen, Geodinamica Acta, 8(3), 142–157,
815 doi:10.1080/09853111.1995.11105386, 1995.

816 Arthaud, F. and Seguret, M.: Les structures pyreneennes du Languedoc et du Golfe du Lion
817 (Sud de la France), Bulletin de la Société Géologique de France, S7-XXIII(1), 51–63,
818 doi:10.2113/gssgfbull.S7-XXIII.1.51, 1981.

819 Arthaud, F., Mégard, F. and Séguret, M.: Cadre tectonique de quelques bassins sédimentaires.
820 Bulletin du Centre de Recherche, d'Exploration et de Production, Elf-Aquitaine., 1, 147–188,
821 1977.

822 Bakalowicz, M.: Karst at depth below the sea level around the Mediterranean due to the
823 Messinian crisis of salinity. Hydrogeological consequences and issues, Geologica Belgica,
824 2014.

825 Baudrimont, A. F. and Dubois, P.: Un bassin mésogéen du domaine péri-alpin: le Sud-Est de la
826 France, Bulletin Centre Recherche Exploration Production Elf-Aquitaine, 262–308, 1977.

827 Benedicto, A., Labaume, P., Séguret, M. and Séranne, M.: Low-angle crustal ramp and basin
828 geometry in the Gulf of Lion passive margin: Oligocene-aquitainian Vistrenque graben, SE
829 France, Tectonics, 15(6), 1192–1212, doi:10.1029/96TC01097, 1996.

830 Bonnet, A. and Paloc, H.: Les eaux des calcaires Jurassiques du bassin de Montbazin-Gigean
831 et de ses bordures (pli de Montpellier et massif de la Gardiole Hérault) BRGM 69 SGL 207
832 LRO., 1969.

Mis en forme : Anglais (Canada)

Code de champ modifié

Mis en forme : Anglais (Canada)

Mis en forme : Anglais (Canada)

Mis en forme : Anglais (Canada)

Mis en forme : Anglais (Canada)

Choukroune, P. and Mattauer, M.: Tectonique des plaques et Pyrénées: sur le fonctionnement de la faille transformante nord-pyrénéenne; comparaison avec des modèles actuels, *Bulletin de la Société Géologique de France*, 7(5), 689–700, 1978.

Choukroune, P., Seguret, M. and Galdeano, A.: Caractéristiques et évolution structurale des Pyrénées; un modèle de relations entre zone orogénique et mouvement des plaques, *Bulletin de la Société Géologique de France*, S7-XV(5–6), 600–611, doi:10.2113/gssgfbull.S7-XV.5-6.600, 1973.

Chu, Y., Tournoud, M. G., Salles, C., Got, P., Perrin, J.-L., Rodier, C., Caro, A. and Troussellier, M.: Spatial and temporal dynamics of bacterial contamination in South France coastal rivers: focus on in-stream processes during low flows and floods, *Hydrological processes*, 28(8), 3300–3313, 2014.

Clauzon, G.: Le canyon messinien du Rhône: une preuve décisive du “desiccated deep-basin model” (Hsue, Cita and Ryan, 1973), *Bulletin de la Société Géologique de France*, S7-XXIV(3), 597–610, doi:10.2113/gssgfbull.S7-XXIV.3.597, 1982.

Combes, J.-P.: Typologie, cadre géodynamique des bauxites téthysiennes, *Geodinamica Acta*, 4, 91–109, 1990.

Debrand-Passard, S.: Synthèse géologique du Sud-Est de la France - Stratigraphie et paléogéographie., 1984.

Doerfliger, N., Ladouche, B., Bakalowicz, M., J.-L. Pinault and Chemin, P.: Étude du pourtour est de l'étang de Thau, phase II. Synthèse générale. Volume 4. BRGM/RP-50789-FR, 71 p., 34 fig., 3 tabl., 2 photos., 2001.

Droge, C. and Bidaux, P.: Simultaneous outflow of fresh water and inflow of sea water in a coastal spring, *Nature*, 322(6077), 361–363, doi:10.1038/322361a0, 1986.

Faure, G.: Principles of isotope geology, Second ed Wiley, New-York. 588 p., 1986.

Fleury, P.: Sources sous-marines et aquifères karstiques côtiers méditerranéens : Fonctionnement et caractérisation., 2005.

Fleury, P., Bakalowicz, M. and de Marsily, G.: Submarine springs and coastal karst aquifers: a review, *Journal of Hydrology*, 339(1–2), 79–92, 2007.

Gèse, B.: Les mésaventures des sources de l'Estavelle et de l'Inversac en Languedoc méditerranéen, *Int. J. Speleol*, 16, 101–109, 1987.

Guerrot, C., Millot, R., Robert, M. and Négrel, P.: Accurate and High-Precision Determination of Boron Isotopic Ratios at Low Concentration by MC-ICP-MS (Neptune), *Geostandards and Geoanalytical Research*, 35(2), 275–284, doi:10.1111/j.1751-908X.2010.00073.x, 2011.

Han, D., Post, V. E. and Song, X.: Groundwater salinization processes and reversibility of seawater intrusion in coastal carbonate aquifers, *Journal of Hydrology*, 531, 1067–1080, 2015.

Hershey, J. P., Fernandez, M., Milne, P. J. and Millero, F. J.: The ionization of boric acid in NaCl, NaCaCl and NaMgCl solutions at 25°C, *Geochimica et Cosmochimica Acta*, 50(1), 143–148, doi:10.1016/0016-7037(86)90059-1, 1986.

Mis en forme : Anglais (Canada)

Mis en forme : Anglais (Canada)

Mis en forme : Anglais (Canada)

Mis en forme : Anglais (Canada)

Hsu, K. J.: The origin of the Mediterranean evaporites, edited by W. B. F. Ryan and K. J. Hsu. Initial reports of the deep sea drilling project, 13, 1203–1231, 1973.

Johannesson, K. H., Stetzenbach, K. J. and Hodge, V. F.: Rare earth elements as geochemical tracers of regional groundwater mixing, *Geochimica et Cosmochimica Acta*, 61(17), 3605–3618, doi:10.1016/S0016-7037(97)00177-4, 1997.

Joigneaux, E., Albéric, P., Pauwels, H., Pagé, C., Terray, L. and Bruand, A.: Impact of climate change on groundwater point discharge: backflooding of karstic springs (Loiret, France), , 2459–2470, 2011.

Kakihana, H., Kotaka, M., Satoh, S., Nomura, M. and Okamoto, M.: Fundamental studies on the ion-exchange separation of boron isotopes, *Bulletin of the Chemical Society of Japan*, 50(1), 158–163, 1977.

La Jeunesse, I., Cirelli, C., Sellami, H., Aubin, D., Deidda, R. and Baghdadi, N.: Is the governance of the Thau coastal lagoon ready to face climate change impacts?, *Ocean & Coastal Management*, 118, 234–246, 2015.

Lacombe, O. and Jolivet, L.: Structural and kinematic relationships between Corsica and the Pyrenees-Provence domain at the time of the Pyrenean orogeny, *Tectonics*, 24(1), 1–20, doi:10.1029/2004TC001673, 2005.

Ladouche, B. and Lamotte, C.: Inversac de la source sous-marine de la Vise en 2014 : Recueil et analyse des données disponibles. Rapport final. BRGM/RP - 64812 - FR, 82 p., 56 ill. , 2015.

Ladouche, B., Aquilina, L., Cubizoles, J. and Négrel, P.: Rainfall chemistry in the south of France (Hérault, 1996-1997), in *Mineralogical Magazine*, 8th Annual VM Goldschmidt Conference, vol. 30, pp. 842–843., 1998.

Ladouche, B., Doerfliger, N., Bakalowicz, M. and avec la collaboration de J. Cubizolles: Étude du pourtour est de l'étang de Thau. Phase II. Caractérisation hydrochimique des réservoirs souterrains karstiques et thermaux. Volume 3. BRGM/RP-50788-FR, 79 p., 20 fig., 5 tabl., 5 ann., 2001.

Ladouche, B., Millot, R., Guerrot, C. and Lamotte, C.: Caractérisation géochimique de l'aquifère hydrothermal de Balaruc-les-Bains lors d'un épisode d'inversac, in *Dix-huitièmes journées techniques du Comité Français d'Hydrogéologie de l'Association Internationale des Hydrogéologues*, "Ressources et gestion des aquifères littoraux, Cassis 2012.", pp. 141–149, Cassis, France. [online] Available from: <https://hal-brgm.archives-ouvertes.fr/hal-00680642> (Accessed 16 August 2018), 2012.

Ladouche, B., Lamotte, C., Hemelsdael, R., Pétré, M.-A., Dewandel, B., Leonardi, V., Seidel, J.-L. and Séranne, M.: Dem'Eaux Thau - Synthèse et valorisation préliminaire des données sur l'hydrosystème de Thau (34). Rapport final. BRGM/RP-68483-FR, 313 p., 133 fig., 6 tab., 6 ann., 2019.

Maerten, L. and Séranne, M.: Extensional tectonics of the Oligo-Miocene Hérault Basin (S France), Gulf of Lion margin, *Bulletin de la Société Géologique de France*, 166(6), 739–749, 1995.

Mis en forme : Anglais (Canada)

Mis en forme : Anglais (Canada)

Martin, J. B. and Screaton, E. J.: Exchange of matrix and conduit water with examples from the Floridan aquifer, US Geological Survey Karst Interest Group Proceedings, St. Petersburg, Florida, 38–44, 2001.

Mis en forme : Anglais (Canada)

Palmer, A., Palmer, M. and Sasowsky, I.: Karst modeling, Karst Waters Institute., 1999.

Pinault, J.-L., Dörfliger, N., Ladouche, B. and Bakalowicz, M.: Characterizing a coastal karst aquifer using an inverse modeling approach: The saline springs of Thau, southern France, Water Resources Research, 40(8), 2004.

Ryan, W. B. F.: Quantitative evaluation of the depth of the western Mediterranean before, during and after the Late Miocene salinity crisis, Sedimentology, 23(6), 791–813, doi:10.1111/j.1365-3091.1976.tb00109.x, 1976.

Séranne, M.: The Gulf of Lion continental margin (NW Mediterranean) revisited by IBS: an overview, Geological Society, London, Special Publications, 156(1), 15–36, doi:10.1144/gsl.sp.1999.156.01.03, 1999.

Stieglitz, T. C., van Beek, P., Souhaut, M. and Cook, P. G.: Karstic groundwater discharge and seawater recirculation through sediments in shallow coastal Mediterranean lagoons, determined from water, salt and radon budgets, Marine Chemistry, 156, 73–84, 2013.

Taylor, S. and McLennan, S.: The continental crust: its composition and evolution., Blackwell Scientific Publications, Oxford., 1985.

Thaler, L.: Premiers résultats d'une recherche systématique des dents de rongeurs, par lavage de marnes de l'Oligocène, en Bas-Languedoc, Comptes Rendus Sommaires de la Société Géologique de France, 10, 315–315, 1962.

Tweed, S. O., Weaver, T. R., Cartwright, I. and Schaefer, B.: Behavior of rare earth elements in groundwater during flow and mixing in fractured rock aquifers: An example from the Dandenong Ranges, southeast Australia, Chemical Geology, 234(3), 291–307, doi:10.1016/j.chemgeo.2006.05.006, 2006.

Mis en forme : Anglais (Canada)

Zhan, Y., Guo, H. and Xing, L.: Characteristics of Rare Earth Elements in Groundwaters along the Flow Path in the North China Plain, Procedia Earth and Planetary Science, 7, 940–943, doi:10.1016/j.proeps.2013.03.173, 2013.

Aggarwal and Palmer 1995 Boron isotope analysis. A review. Analyst, 1995, 120, 1301–1307. DOI: 10.1039/AN9952001301

Aggarwal, J.K., Mezger, K., Pernicka, E., Meixner, A., 2004. The effect of instrumental mass bias on d11B measurements: a comparison between thermal ionisation mass spectrometry and multiple collector ICP-MS. Internat. J. Mass Spectrom. 232, 259–263.

Albéric, P.: River backflooding into a karst resurgence (Loiret, France), Journal of Hydrology, 286(1–4), 194–202, 2004.

Aquilina, L., Deluchat, V., Brach, M., Bakalowicz, M., Le Strat, P. and Giraud, F: Etude géochimique des eaux souterraines autour du bassin de Thau. Rap. BRGM R39530, 58 p., 15 fig., 4 tabl., 2 annexes., 1997.

Aquilina, L., Ladouche, B., Dörfliger, N., Seidel, J. L., Bakalowicz, M., Dupuy, C. and Le Strat, P.: Origin, evolution and residence time of saline thermal fluids (Balaruc springs, southern France): implications for fluid transfer across the continental shelf, *Chemical Geology*, 192(1–2), 1–21, 2002.

Aquilina, L., Ladouche, B., Doerfliger, N. and Bakalowicz, M.: Deep water circulation, residence time, and chemistry in a karst complex, *Groundwater*, 41(6), 790–805, 2003.

Arfib, B. and Gilli, E.: Karst côtier et sources sous marines fonctionnement et exploitation., 2010.

Arthaud, F. and Laurent, P.: Contraintes, déformation et déplacement dans l'avant-pays Nord-pyrénéen du Languedoc méditerranéen, *Geodinamica Acta*, 8(3), 142–157, doi:10.1080/09853111.1995.11105386, 1995.

Arthaud, F. and Seguret, M.: Les structures pyrénéennes du Languedoc et du Golfe du Lion (Sud de la France), *Bulletin de la Société Géologique de France*, S7-XXIII(1), 51–63, doi:10.2113/gssgfbull.S7-XXIII.1.51, 1981.

Arthaud, F., Mégard, F. and Séguret, M.: Cadre tectonique de quelques bassins sédimentaires, *Bulletin du Centre de Recherche, d'Exploration et de Production, Elf-Aquitaine*, 1, 147–188, 1977.

Bakalowicz, M.: Karst at depth below the sea level around the Mediterranean due to the Messinian crisis of salinity. Hydrogeological consequences and issues, *Geologica Belgica*, 2014.

Baudrimont, A. F. and Dubois, P.: Un bassin mésogéen du domaine péri-alpin: le Sud-Est de la France, *Bulletin Centre Recherche Exploration Production Elf-Aquitaine*, 262–308, 1977.

Benedieto, A., Labaume, P., Séguret, M. and Séranne, M.: Low-angle crustal ramp and basin geometry in the Gulf of Lion passive margin: Oligocene-aquitainian Vistrenque graben, SE France, *Tectonics*, 15(6), 1192–1212, doi:10.1029/96TC01097, 1996.

Choukroune, P. and Mattauer, M.: Tectonique des plaques et Pyrénées: sur le fonctionnement de la faille transformante nord-pyrénéenne; comparaison avec des modèles actuels, *Bulletin de la Société Géologique de France*, 7(5), 689–700, 1978.

Choukroune, P., Seguret, M. and Galdeano, A.: Caractéristiques et évolution structurale des Pyrénées; un modèle de relations entre zone orogénique et mouvement des plaques, *Bulletin de la Société Géologique de France*, S7-XV(5–6), 600–611, doi:10.2113/gssgfbull.S7-XV.5-6.600, 1973.

Chu, Y., Tournoud, M. G., Salles, C., Got, P., Perrin, J. L., Rodier, C., Caro, A. and Troussellier, M.: Spatial and temporal dynamics of bacterial contamination in South-France coastal rivers: focus on in-stream processes during low-flows and floods, *Hydrological processes*, 28(8), 3300–3313, 2014.

Clauzon, G.: Le canyon messinien du Rhône; une preuve décisive du “desiccated-deep-basin model” (Hsue, Cita and Ryan, 1973), *Bulletin de la Société Géologique de France*, S7-XXIV(3), 597–610, doi:10.2113/gssgfbull.S7-XXIV.3.597, 1982.

988 Combes, J. P.: Typologie, cadre géodynamique des bauxites téthysiennes, *Geodinamica Acta*,
989 4, 91–109, 1990.

990 Debrand-Passard, S.: Synthèse géologique du Sud-Est de la France—Stratigraphie et
991 paléogéographie., 1984.

992 Doerfliger, N., Ladouche, B., Bakalowicz, M., J. L. Pinault and Chemin, P.: Étude du pourtour
993 est de l'étang de Thau, phase II. Synthèse générale. Volume 4. BRGM/RP-50789-
994 FR, 71 p., 34 fig., 3 tabl., 2 photos., 2001.

995 Drogue, C. and Bidaux, P.: Simultaneous outflow of fresh water and inflow of sea water in a
996 coastal spring, *Nature*, 322(6077), 361–363, doi:10.1038/322361a0, 1986.

997 Faure, G.: Principles of isotope geology, Second ed Wiley, New York. 588 p., 1986.

998 Fleury, P.: Sources sous marines et aquifères karstiques côtiers méditerranéens :
999 Fonctionnement et caractérisation., 2005.

1000 Fleury, P., Bakalowicz, M. and de Marsily, G.: Submarine springs and coastal karst aquifers: a
1001 review, *Journal of Hydrology*, 339(1–2), 79–92, 2007.

1002 Gèze, B.: Les mésaventures des sources de l'Estavelle et de l'Inversac en Languedoc
1003 méditerranéen, *Int. J. Speleol*, 16, 101–109, 1987.

1004 Guerrot, C., Millot, R., Robert, M. and Négrel, P.: Accurate and High-Precision Determination
1005 of Boron Isotopic Ratios at Low Concentration by MC-ICP-MS (Neptune), *Geostandards and
1006 Geoanalytical Research*, 35(2), 275–284, doi:10.1111/j.1751-908X.2010.00073.x, 2011.

1007 Han, D., Post, V. E. and Song, X.: Groundwater salinization processes and reversibility of
1008 seawater intrusion in coastal carbonate aquifers, *Journal of Hydrology*, 531, 1067–1080, 2015.

1009 Hemelsdaël, R., Séranne M., Jacob T., Husson E., Courrioux G., Caritg S., Ballas G. (in prep)
1010 Structural style of the Languedoc Pyrenean thrust belt in relation with the polyphased history
1011 of the western Gulf of Lion margin, SE France. To be submitted in *Earth Sciences Bulletin*
1012

1013 Hershey J.P., Fernandez M., Milne P.J., Millero F.J. (1986) The ionization of boric acid in
1014 NaCl, Na-Ca-Cl and Na-Mg-Cl solutions at 25°C. *Geochim. Cosmochim. Acta*, 50: 143–148

1015 Hsu, K. J.: The origin of the Mediterranean evaporites, edited by W. B. F. Ryan and K. J. Hsu,
1016 Initial reports of the deep-sea drilling project, 13, 1203–1231, 1973.

1017 Kakihana H., Kotaka M., Satoh S., Nomura M., Okamoto M. (1977) Fundamental studies on
1018 the ion-exchange separation of boron isotopes. *Bull. Chem. Soc. Japan.*, 50: 158–163.

1019 Johannesson, K. H., Stetzenbach, K. J. and Hodge, V. F.: Rare earth elements as geochemical
1020 tracers of regional groundwater mixing, *Geochimica et Cosmochimica Acta*, 61(17), 3605–
1021 3618, doi:10.1016/S0016-7037(97)00177-4, 1997.

1022 Joigneaux, E., Albéric, P., Pauwels, H., Pagé, C., Terray, L., Bruand, A., 2011. Impact of
1023 climate change on groundwater point discharge: backflooding of karstic springs (Loiret,
1024 France). *Hydrol. Earth Syst. Sci.* 15, 2459–2470. <https://doi.org/10.5194/hess-15-2459-2011>

La Jeunesse, I., Cirelli, C., Sellami, H., Aubin, D., Deidda, R. and Baghdadi, N.: Is the governance of the Thau coastal lagoon ready to face climate change impacts?, *Ocean & Coastal Management*, 118, 234–246, 2015.

Lacombe, O. and Jolivet, L.: Structural and kinematic relationships between Corsica and the Pyrenees-Provence domain at the time of the Pyrenean orogeny, *Tectonics*, 24(1), 1–20, doi:10.1029/2004TC001673, 2005.

Ladouche, B. and Lamotte, C.: Inversac de la source sous-marine de la Vise en 2014: Recueil et analyse des données disponibles. Rapport final. BRGM/RP–64812–FR–82 p., 56 ill., 2015.

Ladouche, B., Aquilina, L., Cubizoles, J. and Négrel, P.: Rainfall chemistry in the south of France (Hérault, 1996–1997), in *Mineralogical Magazine*, 8th Annual VM–Goldschmidt Conference, vol. 30, pp. 842–843., 1998.

Ladouche, B., Doerfliger, N., Bakalowicz, M. and avec la collaboration de J. Cubizoles: Étude du pourtour est de l'étang de Thau. Phase II. Caractérisation hydrochimique des réservoirs souterrains karstiques et thermaux. Volume 3. BRGM/RP–50788–FR, 9 p., 20 fig., 5 tabl., 5 ann., 2001.

Ladouche, B., Millot, R., Guerrot, C. and Lamotte, C.: Caractérisation géochimique de l'aquifère hydrothermal de Balaruc les Bains lors d'un épisode d'inversac, in *Dix huitièmes journées techniques du Comité Français d'Hydrogéologie de l'Association Internationale des Hydrogéologues*. "Ressources et gestion des aquifères littoraux–Cassis 2012.", pp. 141–149, Cassis, France. [online] Available from: <https://hal-brgm.archives-ouvertes.fr/hal-00680642> (Accessed 16 August 2018), 2012.

Ladouche B., Lamotte C., Hemelsdael R., Pétré M.A., Dewandel B., Léonardi V., Seidel J.L., Séranne M. (2019) Dem'Eaux Thau–Synthèse et valorisation préliminaire des données sur l'hydrosystème de Thau (34). Rapport final. BRGM/RP–68483–FR, 313 p., 133 fig., 6 tab., 6 ann.

Maerten, L. and Séranne, M.: Extensional tectonics of the Oligo-Miocene Herault Basin (S-France), Gulf of Lion margin, *Bulletin de la Société Géologique de France*, 166(6), 739–749, 1995.

Martin, J.B. and Sreaton, E.J., 2001, Exchange of matrix and conduit water with examples from the Floridan aquifer, in E.L. Kuniansky (ed.), U.S. Geological Survey Karst Interest Group Proceedings, Water Resources Investigations Report 01–4011, p.38–44.

Palmer, A.N., 1999, Introduction, in A.N. Palmer, M.V. Palmer, and I.D. Sasowsky (eds.), *Karst Modeling*: Charles Town, West Virginia, Karst Waters Institute Special Publication 5, p 1–9.

Paloc and Bonnet 1969, *Définition de la structure géologique profonde du bassin de Montbazin–Gigean*. BRGM/69–SGL–207–LRO

Pinault, J.-L., Dörfliger, N., Ladouche, B. and Bakalowicz, M.: Characterizing a coastal karst aquifer using an inverse modeling approach: The saline springs of Thau, southern France, *Water Resources Research*, 40(8), 2004.

~~Ryan, W. B. F.: Quantitative evaluation of the depth of the western Mediterranean before, during and after the Late Miocene salinity crisis, *Sedimentology*, 23(6), 791–813, doi:10.1111/j.1365-3091.1976.tb00109.x, 1976.~~
~~Séranne, M.: The Gulf of Lion continental margin (NW Mediterranean) revisited by IBS: an overview, *Geological Society, London, Special Publications*, 156(1), 15–36, doi:10.1144/gsl.sp.1999.156.01.03, 1999.~~
~~Stieglitz, T. C., van Beek, P., Souhaut, M. and Cook, P. G.: Karstic groundwater discharge and seawater recirculation through sediments in shallow coastal Mediterranean lagoons, determined from water, salt and radon budgets, *Marine Chemistry*, 156, 73–84, 2013.~~
~~Taylor, S. and McLennan, S.: *The continental crust: its composition and evolution.*, Blackwell Scientific Publications, Oxford., 1985.~~
~~Thaler, L.: Premiers résultats d'une recherche systématique des dents de rongeurs, par lavage de marnes de l'Oligocène, en Bas Languedoc, *Comptes Rendus Sommaires de la Société Géologique de France*, 10, 315–315, 1962.~~
~~Tweed, S. O., Weaver, T. R., Cartwright, I. and Schaefer, B.: Behavior of rare earth elements in groundwater during flow and mixing in fractured rock aquifers: An example from the Dandenong Ranges, southeast Australia, *Chemical Geology*, 234(3), 291–307, doi:10.1016/j.chemgeo.2006.05.006, 2006.~~
~~Zhan, Y., Guo, H. and Xing, L.: Characteristics of Rare Earth Elements in Groundwaters along the Flow Path in the North China Plain, *Procedia Earth and Planetary Science*, 7, 940–943, doi:10.1016/j.proeps.2013.03.173, 2013.~~