

Quantification of Ecohydrological Sensitivities and Their Influencing Factors at the Seasonal Scale

Yiping Hou^{1,2}, Mingfang Zhang^{1,3}, Xiaohua Wei², Shirong Liu⁴, Qiang Li⁵, Tijiu Cai⁶, Wenfei Liu⁷, Runqi Zhao^{1,8}, Xiangzhuo Liu^{1,9}

5 ¹School of Resources and Environment, University of Electronic Science and Technology of China, Chengdu, 611731, China

²Department of Earth, Environmental and Geographic Sciences, University of British Columbia (Okanagan campus), 1177 Research Road, Kelowna, British Columbia, V1V 1V7, Canada

³Center for Information Geoscience, University of Electronic Science and Technology of China, Chengdu, 611731, China

⁴Research Institute of Forest Ecology, Environment and Protection, Chinese Academy of Forestry, Beijing, 100091, China

10 ⁵Center for Ecological Forecasting and Global Change, College of Forestry, Northwest A&F University, Yangling, 712100, China

⁶Department of Forestry, School of Forestry, Northeast Forestry University, Harbin, 150040, China

⁷Jiangxi Provincial Key Laboratory for Restoration of Degraded Ecosystems & Watershed Ecohydrology, Nanchang Institute of Technology, Nanchang, 330099, China

15 ⁸Division of Ocean Science and Technology, Tsinghua Shenzhen International Graduate School, Tsinghua University, Shenzhen, 518055, China

⁹INRA, Centre INRA Bordeaux Aquitaine, URM1391 ISPA, 33140, Villenave d'Ornon, France

Correspondence to: Mingfang Zhang (mingfangzhang@uestc.edu.cn)

20 **Abstract**

Ecohydrological sensitivity, defined as the response intensity of streamflow to per unit vegetation change is an integrated indicator for assessing hydrological sensitivity to vegetation change. Understanding ecohydrological sensitivity and its influencing factors is crucial for managing water supply, reducing water-related hazards, and ensuring aquatic functions by vegetation management. Yet, there still lacks a systematic assessment on ecohydrological sensitivity and associated driving factors especially at a seasonal scale. In this study, 14 large watersheds across various environmental gradients in China were selected to quantify their ecohydrological sensitivities at a seasonal scale and to examine the role of associated influencing factors such as climate, vegetation, topography, soil, and landscape. Based on the variables identified by correlation analysis and factor analysis, prediction models of seasonal ecohydrological sensitivity were constructed to test their utilities for the design of watershed management and protection strategies. Our key findings were: (1) ecohydrological sensitivities were more sensitive under dry conditions than wet conditions, for example, 1% LAI (leaf area index) change averagely induced 5.05% and 1.96% change in the dry and wet season streamflow, respectively; (2) seasonal ecohydrological sensitivities were highly variable across the study watersheds with different climate conditions, dominant soil types, and hydrological regimes; and (3) the dry season ecohydrological sensitivity was mostly determined by topography (slope, slope length, valley depth, downslope distance gradient), soil (topsoil organic carbon, topsoil bulk density), and vegetation (LAI), while the wet season ecohydrological sensitivity was mainly controlled by soil (topsoil available water holding capacity), landscape (edge density),

25
30
35

and vegetation (leaf area index). Our study provided a useful and practical framework to assess and predict ecohydrological sensitivities at the seasonal scale. The established ecohydrological sensitivity prediction models can be applied to ungauged watersheds or watersheds with limited hydrological data to help decision makers and watershed managers effectively manage hydrological impacts through vegetation restoration programs. We conclude that ecohydrological sensitivities at the seasonal scale are varied by climate, vegetation, and watershed property, and their understanding can greatly support the management of hydrological risks and protection of aquatic functions.

1 Introduction

Natural rivers often have a distinctive seasonal pattern of flow, where flow is highly related to precipitation and shows large variations over dry and wet seasons. Seasonal flows determine ecosystem functions (Toledo-Aceves et al., 2011; Bruijnzeel et al., 2011; Salve et al., 2011), and their responses to vegetation change are highly variable and consequently affect watershed ecosystem equilibrium (Maeda et al., 2015). On the one hand, wet season flows and their variability regulate flood magnitudes (Arias et al., 2012), determine the structure of floodplains and channel morphology (Jansen and Nanson, 2010), and provide opportunities for the recruitment of large woody debris (Warfe et al., 2011; de Paula et al., 2011). On the other hand, dry season flows are critical for maintaining a stable water supply and protecting aquatic ecosystem, as well as playing important roles in sustaining aquatic biota and refuging juvenile fishes (Bunn et al., 2006; Palmer and Ruhi, 2019). However, seasonal streamflow can be significantly affected by forest or vegetation change (Dai, 2011; Hirabayashi et al., 2013). Research has shown that vegetation change can influence water retention time (Moore and Wondzell, 2005; Baker and Wiley, 2009; Bisantino et al., 2015), alter snow accumulation and snowmelt processes (Lin and Wei, 2008; Zhang and Wei, 2012; Calder, 2005), and route river flow quickly to downstream (Winkler et al., 2010; Chang, 2012) and consequently increase the frequency and size of floods in wet season. Vegetation change can also affect dry season flows, which may influence baseflow level and the risk of droughts, and degrade or enrich in-channel habitat for aquatic species (Simonit and Perrings, 2013; Sun et al., 2016). Thus, understanding seasonal hydrological variations to vegetation change is critical for maintaining the sustainable water supply, preventing large floods and droughts, and developing the bestwatershed management plans.

Obviously, seasonal streamflow responses to vegetation change are highly variable among watersheds worldwide. To better understand the general pattern of streamflow response to vegetation change, Zhang et al. (2017) has introduced a uniform indicator named ecohydrological sensitivity (defined as the response intensity of streamflow to per unit forest change) to express the hydrological sensitivity to forest change for a given watershed. Ecohydrological sensitivity is believed to be controlled not only by forest or vegetation coverage but also by climate condition, hydrological regime, and forest or vegetation type (Zhang et al., 2017; Li et al., 2017). Assessing ecohydrological sensitivity can provide various benefits. For example, it provides a dimensionless index on the vegetation-water relationship so that any watersheds can be effectively compared. It allows for predicting ecohydrological sensitivities for a landscape or region so that negative hydrological impacts in the areas

with high ecohydrological sensitivities can be minimized through suitable arrangements of vegetation or watershed management strategies.

70 Ecohydrological sensitivity is likely varied with time scales. The hydrological responses to vegetation change at the annual scale are the averaged and cumulative effects from those at shorter time intervals, which are typically associated with total annual magnitudes such as water yield, while those at daily or monthly or seasonal scales affect flow patterns and are normally related to floods and droughts. The seasonal scale is a medium level between daily and annual scales, which can affect both magnitude and pattern in terms of hydrological response and sensitivity. For example, the interactions between vegetation and water are quite different between dry and wet seasons (Donohue et al., 2010; Asbjornsen et al., 2011). Abundant
75 water is available for vegetation growth in wet season, while vegetation in dry season mostly relies on limited soil moisture or groundwater for limited photosynthesis and transpiration. Besides, streamflow generation in wet season is mainly based on precipitation or water input, whereas dry season flow is controlled by soil moisture in the antecedent wet season and groundwater discharge. Thus, the contrasted processes in different seasons suggest that ecohydrological sensitivity must be examined at a seasonal scale.

80 Various factors, including climate, vegetation, and watershed property affect hydrological responses or sensitivities (Zhou et al., 2015; Li et al., 2017; Zhang et al., 2017). For example, hydrological responses to forest change tend to be more sensitive in non-humid regions (Zhang et al., 2017). Evapotranspiration change related to vegetation change is controlled by energy and water (Zhang et al., 2004; Creed et al., 2014; Yang et al., 2007). Topography controls hydrological processes by affecting the distribution and routing of water (Woods, 2007). Soil and landscape conditions are important for erosion,
85 sediment, and flow connectivity (Borselli et al., 2008). Clearly, fully assessing and understanding ecohydrological sensitivity requires a consideration of various influencing variables. Yet, current studies have only focused on the hydrological influences of a single type of variables such as vegetation (Beck et al., 2013; Feng et al., 2016; van Dijk et al., 2012), climate (Creed et al., 2014; Miara et al., 2017), topography (Lyon et al., 2012; Jencso and McGlynn, 2011; Li et al., 2018a), and landscape (Nippgen et al., 2011; Buma and Livneh, 2017; Teutschbein et al., 2018). The inclusion of various types of variables into an
90 integrated assessment framework of hydrological responses remains a challenging subject. Despite the recognition that ecohydrological sensitivity can be a good index that facilitates the understanding of variations in hydrological response to vegetation change, there still lacks a commonly accepted definition or framework for its quantitative assessment and comparisons especially at a seasonal scale. To our best knowledge, there is no study on quantifying seasonal ecohydrological sensitivity.

95 China has experienced substantial and dynamic vegetation change over the past few decades. Deforestation and biomass loss dominated vegetation change from the 1950s to 1980s (Wei et al., 2008), while the large-scale revegetation programs have been implemented since the 1980s (Li et al., 2018b). These large-scale vegetation changes can inevitably affect local and regional water cycles. However, given the large variations in climate, vegetation, soil, topography, and landscapes in China, hydrological responses to vegetation change can be quite different among watersheds. Since assessing the
100 hydrological impact of vegetation change in every single watershed can be very challenging and time-consuming, a general

framework for an efficient evaluation of ecohydrological sensitivity at a watershed scale is in an urgent need for the support of future water and forest resource management. The objectives of this study were: (1) to evaluate seasonal ecohydrological sensitivity in the selected large watersheds across environmental gradients; (2) to examine the role of climate, vegetation, topography, soil, and landscape in seasonal ecohydrological sensitivity; and (3) to simulate and predict seasonal ecohydrological sensitivity based on the selected factors.

2 Study watersheds and data

2.1 Study watersheds

Given that the dominant climate zones in China include subtropical monsoon, alpine, temperate monsoon, and temperate continental climate zones, 2-4 representative study watersheds in each climate zone are identified according to their hydrological data availability, watershed size, climate type, and vegetation type. The selected watersheds in each climatic zone are with the watershed size greater than 500 km² and long-time hydrological data available to meet the data requirements for statistical analysis (≥ 15 yrs). In addition, only vegetative watersheds with vegetation coverage greater than 30% are included since the climate (e.g., precipitation) is a more influencing factor than vegetation on river flows in less vegetative watersheds. With these criteria, fourteen large watersheds across climatic zones with the area ranging from 832 to 19189 km² are selected. They include the Pingjiang and Xiangshui watersheds in Southeast China, the Tangwang River and Xinancha River watersheds in Northeast China, the Upper Zagunao, Zagunao, Upper Heishui River, Heishui River, Gongbujiangda and Gengzhang watersheds in Southwest China and the Dongchuan, Heishuichuan, Jingchuan and Rui River watersheds in Northwest China (Fig. 1). In this study, the dry and wet seasons are defined according to the long-term mean monthly precipitation in a hydrological year. For subtropical monsoon climate dominated watersheds (the Pingjiang and Xiangshui), wet season starts from March to August with its precipitation amount accounting for over 70% of the annual total, while dry season lasts from September to February. For those from the alpine, temperate monsoon, and temperate continental climate zones, wet season is from May to October with dry season from November to April. Table 1 provides a brief summary of seasonality, climate, vegetation, hydrology, and topography in the study watersheds. Detailed descriptions of study watersheds can be found in Sect. S1 in the Supplement. In addition, substantial vegetation restoration programs caused large-scale vegetation change from the 1980s onwards. To evaluate seasonal ecohydrological sensitivity, the study periods start from 1983.

2.2 Data

Daily or monthly discharges for 14 watersheds were obtained from various government agencies. The details about the study periods and hydrometric stations can be found in the Supplement (Table S3). Discharges (m³/s) were converted into the unit of mm according to the drainage area. According to the definitions of seasonality in Table 1, a hydrological year was divided into dry season and wet season, and then seasonal flows were calculated accordingly.

The historical climate data used in this study include three sources: daily climate records from National Meteorological Information Centre of China Meteorological Administration (CMA: <http://data.cma.cn/>), spatial-interpolated gridded climate data by use of the ANUSPLIN model and meteorological data collected at the associated hydrological stations or rain gauges (Sect. S1.2 and Table S3). In this study, daily or monthly climate data including mean temperature (T_{mean}),
135 minimum temperature (T_{min}), maximum temperature (T_{max}), and precipitation (P) were derived and calculated accordingly. Monthly potential evapotranspiration (PET) was calculated based on estimated T_{max} and T_{min} by using Hargreaves' equation (Equation 1) (Hargreaves and Samani, 1985).

$$PET = 0.0023 \times Ra \times [(T_{\text{min}} + T_{\text{max}})/2 + 17.8] \times (T_{\text{max}} - T_{\text{min}})^{0.5} \quad (1)$$

where, Ra is the extraterrestrial radiation; and T_{min} and T_{max} are the minimum and maximum temperatures in °C.

140 Moderate Resolution Imaging Spectroradiometer (MODIS) land cover product MODIS MCD12Q1 with the spatial resolution of 500m was downloaded from Land Process Distributed Active Archive Centre (LPDAAC: <https://lpdaac.usgs.gov/products/mcd12q1v006/>) (Sulla-Menashe et al., 2019). There are 17 types of land covers in MODIS MCD12Q1, including evergreen needleleaf forests, deciduous needleleaf forests, evergreen broadleaf forests, deciduous broadleaf forests, mixed forests, closed shrublands, opened shrublands, woody savannas, savannas, grasslands, permanent
145 wetlands, croplands, urban and built-up land, cropland/natural vegetation mosaics, permanent snow and ice, barren, and water bodies. We reclassified them into forest (evergreen needleleaf forests, deciduous needleleaf forests, evergreen broadleaf forests, deciduous broadleaf forests and mixed forests), shrubland (closed shrublands and opened shrublands), grassland (woody savannas, savannas and grasslands), agricultural (croplands and cropland/natural vegetation mosaics), snow (permanent snow and ice), and other lands (permanent wetlands, urban and built-up land, barren, and water bodies) (Table
150 S2). Vegetation coverage including forest, shrubland and grassland can be then calculated.

Leaf area index (LAI) derived from the Global Land Surface Satellite LAI Product (GLASS LAI) was used as a vegetation index to express vegetation change in this study (GLASS: <http://glass-product.bnu.edu.cn/>). The GLASS LAI product dataset provides continuous global LAI at a high temporal resolution of eight days (Liang et al., 2013; Xiao et al., 2014). There are two types of GLASS LAI products with different spatial resolutions and available periods. The first GLASS
155 LAI product is based on Advanced Very High Resolution Radiometer (AVHRR) reflectance data with the spatial resolution of 0.05°, and this dataset is available from 1982 to 2016. The other one, with a higher spatial resolution of 1 km is retrieved from Moderate Resolution Imaging Spectroradiometer (MODIS) reflectance data, but it only covers a period of 17 years from 2000 to 2016. As the study watersheds are large watersheds (>500 km²) and the study periods are ended before 2006, the former GLASS LAI product was chosen for this study, wherein two data series of LAI, dry season LAI (mean value of the
160 LAIs in the dry season) and wet season LAI (mean value of the LAIs in the wet season) from the entire study period were generated.

Harmonized World Soil Database (HWSD) published by Food and Agriculture Organization (FAO) and International Institute for Applied Systems Analysis (IIASA) with the spatial resolution of 1km was used to collect soil indices (Wieder,

2014). HWSD classifies soil into topsoil from surface to 30 cm below ground, and subsoil between 30 cm and 100 cm below
165 ground.

Digital elevation models (DEMs) with the spatial resolution of 30m derived from GDEM were provided by Geospatial Data Cloud site, Computer Network Information Centre, Chinese Academy of Sciences (<http://www.gscloud.cn>). Topographic information of the study watersheds was derived from DEMs.

3 Methods

170 3.1 Definition and calculation of ecohydrological sensitivity

In this study, an improved single watershed approach was employed to quantify seasonal streamflow variations attributed to climate variability, vegetation change, and other factors (Hou et al., 2018a; Hou et al., 2018b). The modified double mass curve (MDMC) was firstly used to remove the effects of climate variability on seasonal streamflow variation. The multivariate ARIMA (ARIMAX) model was then adopted to quantify seasonal streamflow variation attributed to non-climatic factors
175 (vegetation change and other factors). The 95% confidence intervals (95% CIs) criterion was applied to separate the statistical errors and the seasonal streamflow variation attributed to other factors. The seasonal streamflow variation caused by vegetation change (ΔQ_v) can be quantified eventually and be used to calculate the seasonal ecohydrological sensitivity. A more detailed description of the methodology is provided in the Supplement Sect. 2.

Similar to the concept of ecohydrological sensitivity proposed by Zhang et al. (2017), in this study, we defined
180 seasonal ecohydrological sensitivity (S_f) as the response intensity of seasonal streamflow variations to per unit vegetation change (using the leaf area index (LAI) as a proxy), which can be computed with equations (2)-(3). The value of seasonal ecohydrological sensitivity refers to the percentage of seasonal streamflow changes induced by 1% of LAI change. Given seasonal streamflow response to vegetation change in mm (ΔQ_v) can be influenced by its background value (\bar{Q} , the long-term mean seasonal streamflow during the study period), seasonal streamflow response to vegetation change in percentage ($\Delta Q_v\%$)
185 is used for the calculation of ecohydrological sensitivity. Here, ΔQ_v is divided by \bar{Q} to calculate $\Delta Q_v\%$. Through this normalization, $\Delta Q_v\%$ representing relative change (%) in seasonal streamflow compared to its average state can be a better indicator for hydrological sensitivity analysis than ΔQ_v .

$$\Delta Q_v \% = 100 \times \frac{\Delta Q_v}{\bar{Q}} \quad (2)$$

$$S_f = \left| \frac{\Delta Q_v \%}{\Delta LAI} \right| \quad (3)$$

190 where, \bar{Q} refers to the long-term mean seasonal streamflow during the study period; ΔQ_v is seasonal streamflow response to vegetation change in mm; $\Delta Q_v\%$ is seasonal streamflow response to vegetation change in percentage (%); and ΔLAI is LAI variation compared to average LAI in the reference period in %.

3.2 Comparison of seasonal ecohydrological sensitivities between watershed conditions

195 According to dryness index (*DI*), watersheds were grouped into energy-limited (EL), equitant (EQ) and water-limited (WL) conditions (McVicar et al., 2012). The most widely distributed soil type in a watershed was treated as the dominant soil type. Following our analysis, four dominant soil types (LIXISOLS, LUVISOLS, LEPTOSOLS, and CAMBISOLS) were shown in this study. Additionally, the selected watersheds were categorized into rain-dominated (RD) and rain-snow hybrid (Hybrid) watersheds according to their hydrological regimes. Table 2 showed the detailed classifications for each watershed in terms of climate condition, dominant soil type and hydrological regime.

200 Non-parametric Mann Whitney U test was performed to detect the statistically significant differences between the watershed groups. Mann Whitney U test can test whether there are significant differences in the median values of seasonal ecohydrological sensitivities between two groups (Birnbaum, 1956).

3.3 Prediction of seasonal ecohydrological sensitivity

205 Five types of indices including climate, vegetation, topography, soil, and landscape were adopted in this study. Detailed information on the interpretations and calculations of 40 indices were presented in Table 3. Climate indices, including dryness index and effective precipitation can demonstrate water input and climate condition in a given watershed (van Dijk et al., 2012; Jones et al., 2012; Zhang et al., 2004). Dryness index is calculated at the annual scale to demonstrate dryness condition, while effective precipitation (an integrated index of climatic variability) in dry season and wet season denotes seasonal water inputs. Vegetation growth is highly dependent on temperature, water, soil, and geographical location (Chang, 2012). Vegetation coverage or forest coverage indicates a proportion of vegetation or forest in a watershed, but it cannot express vegetation growth, mortality, and seasonality. LAI is recognized as a better indicator mainly because it is an important biophysical variable relating to photosynthesis, transpiration, and energy balance (Launiainen et al., 2016; Verrelst et al., 2016; González-Sanpedro et al., 2008). Topographic indices can be classified into two groups: primary and secondary (also known as compounded topographic indices) (Li et al., 2018a; Moore et al., 1991). Primary topographic indices can be directly derived from DEM, whilst compounded topographic indices are based on one or more primary indices (Li et al., 2018a). Based on previously published studies, 17 topographic indices including 5 primary indices and 12 compounded indices which are most frequently used in studying the topographic effect on hydrological processes were selected to describe watershed characteristics including visibility, generation process, and morphology (Yokoyama et al., 2002; Park et al., 2001; Jenness, 2004; Li et al., 2018a). Calculations of the topographic indices were made in ArcGIS 10.2 (ERSI) and SAGA GIS 2.1. Soil types were based on the 220 FAO-85 system classification, while soil organic carbon and sandiness were directly derived from HWSD in ArcGIS 10.2 (ERSI), and soil available water holding capacity, saturated hydraulic conductivity, and bulk density were calculated using Soil-Plant-Air-Water (SPAW) hydrology model. We used the weighted average value to represent watershed-scale soil indices. Seven landscape indices including patch number (PN), patch density (PD), largest patch index (LPI), edge density (ED), contagion index (CONTAG), Shannon's diversity index (SHDI), and Simpson's diversity index (SIDI) at the landscape level which are

225 most correlated with hydrological processes were selected in the analysis (Zhou and Li, 2015; Boongaling et al., 2008). The calculations of landscape indices were performed by FRAGSTATS 4.2 software.

Obviously, the prediction with a large number of indices may cause model redundancy. Moreover, some of these indices can be correlated with each other, wherein a multicollinearity problem may arise. To address these issues, we have firstly performed Kendall correlation analysis and linear regression to identify indices that are significantly correlated with seasonal ecohydrological sensitivities, and then have conducted the factor analysis to further reduce the redundancy of indices. 230 Eventually, only a few indices with key influences on seasonal ecohydrological sensitivity are retained for multiple linear regression.

To be specific, kendall correlation analysis and linear regression were used to identify statistically significant correlations between seasonal ecohydrological sensitivities and 40 indices at a significant level of $p=0.10$. The insignificant 235 indices were excluded for prediction described below. Factor analysis (FA) was introduced to further reduce the redundancy of indices. Similar to principal component analysis (PCA), indices after filtering by factor analysis could retain important information, which means that fewer indices can be used to represent most information (Lyon et al., 2012). Three criteria were used to pick highly related indices: the coefficient of Kaiser-Meyer-Olkin (KMO) test, the p -value of Bartlett's test, and the diagonal coefficients of the anti-image correlation matrix (Li et al., 2018a). Indices filtered by factor analysis with the 240 coefficient of KMO being greater than 0.50, the p -value of Bartlett's test being less than 0.05 and the diagonal coefficients of the anti-image correlation matrix being greater than 0.50 were selected for further analysis. After filtering, only a few indices with key influences on seasonal ecohydrological sensitivity were retained for the prediction models. In this way, the correlation between the influencing drivers could be greatly reduced. In addition, the collinearity of inputting variables for the multiple linear regression was assessed by variance inflation factor (VIF). Models with the VIF less than 10 were selected to address 245 collinearity.

Multiple linear regression model modified by stepwise regression was employed to predict seasonal ecohydrological sensitivity. Influencing factors filtered by correlation analysis and factor analysis were regarded as independent variables and ecohydrological sensitivity was considered as a dependent variable in a linear regression model. Independent variables were inputted into a model one by one, and the ANOVA test was conducted accordingly. Once the p -value of the ANOVA test was 250 greater than 0.10, the input independent variable at this stage would be dropped. The optimal linear regression model was reached when no independent variables were inputted and no variables were dropped. The Akaike Information Criterion (AIC) and R^2 were used to find optimal multiple linear regression models for prediction. Except for quantitative indices, climate condition, dominant soil type and hydrological regime might also make contributions to the prediction of ecohydrological sensitivity. As a result, we introduced dummy variables to quantify the influence of climate condition, dominant soil type and 255 hydrological regime on model accuracy (Hardy, 1993). In this study, ecohydrological sensitivity based on the improved single watershed approach was called the observed S_f , while ecohydrological sensitivity from the multiple linear regression model was named as the predicted S_f .

4 Results

4.1 Seasonal ecohydrological sensitivity and its variations

260 Table 4 compared ecohydrological sensitivities between the dry and wet seasons. The ecohydrological sensitivities in the dry season were significantly greater than those in the wet season (Fig. 2 and Fig. S8-S10). As shown in Fig. 2, 1% LAI change averagely induced 5.05% change in dry season streamflow, while in wet season, this value dropped to 1.96%. There were large variations in seasonal ecohydrological sensitivity among the study watersheds. The dry season ecohydrological sensitivity of the Tangwang River watershed was highest, up to 27.75, while the dry season ecohydrological sensitivity of the Upper Heishui
265 River watershed was the lowest (1.01). Similarly, the wet season ecohydrological sensitivity with the value of 4.36 in the Tangwang River watershed was also the highest among all watersheds in the wet season, whereas the lowest wet season ecohydrological sensitivity (0.40) was found in the Xiangshui watershed (Table S4).

Comparisons of seasonal ecohydrological sensitivities were made among the study watersheds grouped by their climate conditions, dominant soil types and hydrological regimes (Fig. 3, Fig. 4 and Fig. 5). As suggested by Fig. 3 and Table
270 5, significant differences in both dry season and wet season ecohydrological sensitivities between energy-limited (EL) and equitant (EQ) watersheds and between energy-limited and water-limited (WL) watersheds were found. Significant differences in the medians of wet season ecohydrological sensitivity in the pair of EQ-WL were also detected. 1% vegetation change caused 2.11%, 5.86% and 5.23% change of dry season streamflow in the energy-limited, equitant and water-limited watersheds, respectively (Fig. 3a), while it only led to 0.58%, 2.82% and 1.64% change of wet season streamflow in the EL,
275 EQ and WL watersheds, respectively (Fig. 3b). These results clearly demonstrated that ecohydrological sensitivity was greater in the EQ and WL conditions, particularly in the dry season.

When seasonal ecohydrological sensitivity in watersheds grouped by dominant soil types was compared (Fig.4 and Table 5), the median of dry season ecohydrological sensitivity in the LIXISOLS-dominated watersheds was significantly different from those of the LUVISOLS- and CAMBISOLS-dominated watersheds at $\alpha=0.05$, and the significant differences
280 in median of dry season ecohydrological sensitivity were also detected in the LUVISOLS-LEPTOSOLS, LIXISOLS-LEPTOSOLS, LUVISOLS-CAMBISOLS and LEPTOSOLS-CAMBISOLS pairs at $\alpha=0.05$ (Table 5). Similarly, the median of dry season ecohydrological sensitivity in the LIXISOLS-dominated watersheds was significantly different from those of the LUVISOLS-, LEPTOSOLS- and CAMBISOLS-dominated watersheds at $\alpha=0.05$. On average 1% change in vegetation led to 2.11%, 3.29%, 5.62% and 13.01% change of dry season streamflow in the LIXISOLS-, LEPTOSOLS-, CAMBISOLS- and
285 LUVISOLS-dominated watersheds, respectively (Fig. 4a), while it caused only 0.58%, 2.20%, 2.11% and 2.24% change of wet season streamflow (Fig. 4b).

Fig. 5 demonstrated the differences of seasonal ecohydrological sensitivity in watersheds grouped by hydrological regime. Mann-Whitney U test showed that there were significant differences between rain-dominated and hybrid watersheds in medians of dry season ecohydrological sensitivity (Table 5). 1% vegetation change can result in 6.51% and 3.29% change

290 of dry season streamflow in rain-dominated and hybrid watersheds, respectively (Fig. 5a), while it only led to 1.75% and 2.20% change of wet season streamflow in rain-dominated and hybrid watersheds, respectively (Fig. 5b).

4.2 Prediction models for seasonal ecohydrological sensitivity

According to correlations between seasonal ecohydrological sensitivity and 40 indices detected by Kendall correlation and linear regression, 17 indices significantly related to dry season ecohydrological sensitivity were identified (Table 6). Dry 295 season ecohydrological sensitivity was significantly and positively correlated with dryness index (DI), topographic wetness index (TWI), downslope distance gradient (DDG), topographic positive openness (PO), topographic negative openness (NO), topsoil salinity (T_{ece}), topsoil bulk density (T_d), while its correlations with all vegetation indices (LAI , vegetation coverage and forest coverage), slope, slope length factor (LS), terrain ruggedness index (TRI), valley depth ($Depth$), topsoil organic carbon (T_{oc}), patch density (PD), and edge density (ED) were significantly negative. In contrast, only 8 indices were significantly 300 correlated with wet season ecohydrological sensitivity. Wet season ecohydrological sensitivity had a significantly positive correlation with convergence (CON), topsoil available water holding capacity (T_w), topsoil saturated hydraulic conductivity (T_{hy}), subsoil saturated hydraulic conductivity (S_{hy}), and subsoil salinity (S_{ece}) whereas a negative relation with effective precipitation (P_e), soil types, and edge density (ED).

8 out of 17 indices significantly related to dry season ecohydrological sensitivity were further identified by factor 305 analysis, which included factors such as DI , slope, LS , TWI , DDG , TRI , $Depth$, and NO . For the factor analysis of dry season ecohydrological sensitivity, the coefficient of KMO was 0.730, the p -value of Bartlett's test was less than 0.05, and diagonal coefficients of the anti-image correlation matrix were greater than 0.53 (Table 7). Meanwhile, factor analysis identified 6 indices (P_e , CON , T_w , T_{hy} , S_{hy} , and ED) associated with wet season ecohydrological sensitivity based on correlation analysis. For wet season subset, the coefficient of KMO with the value of 0.634 was lower than that in dry season subset, but diagonal 310 coefficients of the anti-image correlation matrix were higher than those in wet season subset (≥ 0.57). The p -value of Bartlett's test was 0.00. Given it is an important ecohydrological indicator for vegetation status in a watershed, LAI was also manually added as a predictor in the predicted model. Fig. 6 showed the structure, parameters and statistics of the established prediction models for ecohydrological sensitivity. The dry season model had a better performance with a higher R^2 of 0.966 (Fig. 6a), while the R^2 was only 0.501 for the wet season model (Fig. 6b).

315 5 Discussion

5.1 Seasonal ecohydrological sensitivity and climate conditions

Climate conditions in terms of energy (temperature) and water (precipitation) are the most important drivers for vegetation growth. Ecohydrological processes of vegetative watersheds vary greatly with climate conditions (Donohue et al., 2010). As suggested by our study, both dry season and wet season ecohydrological sensitivities of the water-limited watersheds were 320 higher than those of the energy-limited watersheds (Fig. 3), and the dry season ecohydrological sensitivities were much higher

than the wet season ecohydrological sensitivities (Fig. 2). In addition, the dry season ecohydrological sensitivity significantly increased with rising dryness index while the wet season ecohydrological sensitivity significantly decreased with increasing effective precipitation (Table 6). In other words, under dry conditions (during dry periods or in dry regions), streamflow is more sensitive to vegetation change than under wet conditions (during wet periods or in wet regions). These findings are in accordance with results from previous studies, which indicate streamflow response to vegetation in drier regions might be more pronounced than in wetter regions (Jackson et al., 2005; Vose et al., 2011; Li et al., 2017; Zhang et al., 2017). For example, Farley et al. (2005) demonstrated that afforestation produced 27% water yield reduction in wetter sites, whilst 62% water yield reduction was identified in drier sites based on the analysis of 26 catchments globally. Sun et al. (2006) modelled streamflow responses to large-scale reforestation in China and found increased vegetation cover produced a nearly 30% reduction in streamflow in humid regions, but the streamflow reduction rose to approximately 50% in semi-arid and arid areas. Creed et al. (2014) indicated water use efficiencies in forests were higher in drier years than in wetter years by assessing water yield variations in North America. The different ecohydrological sensitivities between dry and wet seasons might be explained by their various mechanisms of water use by vegetation. Vegetation growth in wet conditions with abundantly available water, sufficient soil moisture and saturated aquifers is more sensitive to energy factors including temperature, radiation and heat input (Newman et al., 2006; Hou et al., 2018a; Zhang et al., 2011; Brooks et al., 2012). Changes in energy input in wet conditions can alter stomatal conductance and transpiration, and consequently affect the photosynthesis, transpiration, and biomass of vegetation (de Sarrau et al., 2012; Van Dover and Lutz, 2004). In contrast, in dry conditions with limited precipitation input, water is more critical for vegetation growth where vegetation mainly relies on its access to soil water through root systems to support photosynthesis and transpiration (Zhou et al., 2015).

340 **5.2 Seasonal ecohydrological sensitivity and soils**

Soils as the interface between streamflow and groundwater play vital roles in water cycle (Bockheim and Gennadiyev, 2010; Schoonover and Crim, 2015). Our study showed that watersheds with different dominant soil types could have contrasting seasonal ecohydrological sensitivity. As shown in Fig. 4, the ecohydrological sensitivities in both dry and wet seasons in the LIXISOLS-dominated watersheds were the lowest compared with those of CAMBISOLS-, LEPTOSOLS- and LUVISOLS-dominated watersheds. This result clearly illustrates the importance of soil types in hydrological responses and sensitivities (Rieu and Sposito, 1991; Srivastava et al., 2010; Chadli, 2016). Soil properties including organic carbon, salinity, available water holding capacity, saturated hydraulic conductivity and bulk density can affect soil water infiltration and lateral movement (Hillel, 1974; Leu et al., 2010). For example, soils with higher available water holding capacity have the ability to store more water for vegetation growth (Mukundan et al., 2010). Saturated hydraulic conductivity is positively correlated to available water holding capacity, suggesting that soils in a watershed with a higher value of saturated hydraulic conductivity might promote interactions between streamflow and groundwater (Sulis et al., 2010). Large differences between topsoil and subsoil bulk densities suggest a frequent moisture movement, leading to more active interactions and feedbacks above and below the soil (Zhao et al., 2010). LIXISOLS is characterized by the lowest saturated hydraulic conductivity and the smallest difference

between topsoil and subsoil bulk densities as compared to other three types of soils (Table S1), indicating its lowest water storage capacity and less frequent water movement between topsoil and subsoil. Therefore, hydrological responses in the LIXISOLS-dominated watersheds were less sensitive to vegetation change, and consequently led to the lowest seasonal ecohydrological sensitivity.

5.3 Seasonal ecohydrological sensitivity and hydrological regimes

Hydrological regime is another determinant for ecohydrological sensitivity (Zhang et al., 2017). Our study found that the dry season ecohydrological sensitivity in the rain-dominated watersheds was significantly higher than that in the hybrid watersheds (Fig. 5), while an insignificant difference in wet season ecohydrological sensitivity between the rain-dominated and hybrid watersheds was estimated (Table 5). The differences in dry season ecohydrological sensitivity between the rain-dominated and hybrid watersheds are associated with their differences in the mechanisms of streamflow generation. In the rain-dominated watersheds, dry season streamflow is mainly maintained by groundwater discharge while both groundwater and snow water might be the sources of dry season streamflow in the hybrid watersheds. Thus, the generation of the dry season streamflow in the hybrid watersheds tend to be more complex and stable, and can be more resilient to vegetation change in comparison with that in rain-dominated watersheds. This is supported by several reviews which found that forest cover change in rain-dominated watersheds can produce greater hydrological impacts than in snow-dominated watersheds (Zhang et al., 2017; Moore and Wondzell, 2005). In hybrid watersheds, forestation or vegetation removal can lead to changes in snowmelt processes by altering snow accumulation, melting timing, energy input and wind speed in dry season (Frank et al., 2015), resulting in hydrological de-synchronization effects. These de-synchronization effects may offset negative impacts of vegetation change on dry season streamflow, and eventually lower dry season ecohydrological sensitivity in the hybrid watersheds.

The lack of a significant difference in the wet season ecohydrological sensitivity between the rain-dominated and hybrid watersheds might be due to the fact that only precipitation form during wet season is rainfall. It is expected that there are similar interactions and feedback mechanisms between vegetation and water in wet season in all watersheds, leading to insignificant differences in wet season ecohydrological sensitivity between the rain-dominated and hybrid watersheds.

5.4 Seasonal ecohydrological sensitivity and topography

Topography as a dominating factor for hydrological processes (Zeng et al., 2016; Jenness, 2004; Scown et al., 2015; Yokoyama et al., 2002; Park et al., 2001; Li et al., 2018a) plays an important role in determining streamflow response to vegetation change (Price, 2011; Smakhtin, 2001). According to the established prediction model of dry season ecohydrological sensitivity (Fig. 6a), topographic factors including slope and downslope distance gradient had positive effects on dry season ecohydrological sensitivity, while slope length factor and valley depth yielded negative effects. The vegetative watersheds with steeper slopes often have faster water movement from slopes to river channel and severe soil erosion in wet season if vegetation is destroyed, which can greatly reduce wet season soil water storage for supply to dry season streamflow, and therefore have greater dry season ecohydrological sensitivity (Desmet and Govers, 1996; Zhang et al., 2012). Similarly, vegetative watersheds with

smaller slope length factor and valley depth can have greater dry season ecohydrological sensitivity. This is probably because these watersheds generally have a generally flatter topography and longer water residence time, and consequently allow for more interactions between vegetation and water, which likely lead to greater ecohydrological sensitivity in dry season.

390 Unlike the dry season ecohydrological sensitivity, no topographic indices were associated with wet season ecohydrological sensitivity (Fig. 6b). As we know, climate and vegetation are two major drivers to hydrological variations in vegetative watersheds (Wei et al., 2018; Li et al., 2017). This indicates that in wet season, climate plays a more dominant role in hydrological responses or variations, which means a decreasing role of vegetation on streamflow and consequently reduction of ecohydrological sensitivity. The decreasing role of vegetation on streamflow in wet season may explain the insignificant impact of topographic indices on wet season ecohydrological sensitivity.

395 **5.5 Seasonal ecohydrological sensitivity and landscape**

Landscape pattern can directly affect hydrological connectivity and indirectly influence hydrological processes by controlling soil activities such as soil erosion and sediment (Buma and Livneh, 2017; Teutschbein et al., 2018; Karlsen et al., 2016). Based on the prediction models (Fig. 6), the landscape pattern played a more important role in wet season ecohydrological sensitivity than in dry season ecohydrological sensitivity. Only edge density was identified as an effective, negative landscape predictor
400 for wet season ecohydrological sensitivity. Watersheds with a higher value of edge density are often featured by landscape fragmentation and segmentation, e.g., scatter distributed vegetation, higher road densities, leading to poor hydrological connectivity and a high risk of soil erosion. The increasing role of watershed property (edge density) means that the relative role of vegetation in hydrological response would be lower, which consequently leads to decreasing of wet season ecohydrological sensitivity.

405 **5.6 Implications**

Ecohydrological studies at the seasonal scale are limited due to the lack of the understanding of complex and variable streamflow responses to climate, vegetation, topography, soil and landscape (McDonnell et al., 2018; Singh et al., 2014; Wei et al., 2018; Li et al., 2018a; Opiel and Schumann, 2020; Guswa et al., 2020). Our findings clearly showed that seasonal ecohydrological sensitivity was not only highly associated with climate and vegetation change, but also significantly related
410 to watershed properties like topography, soil and landscape. As indicated by the constructed prediction models, the dry season ecohydrological sensitivity could be better described by vegetation, topography and soil (Fig. 6a), while the wet season hydrological response was mainly controlled by vegetation (leaf area index), soil (topsoil available water holding capacity) and landscape (edge density) (Fig. 6b). Given complex and variable hydrological responses to vegetation change among the study watersheds due to their differences in watershed properties (Zhou et al., 2015; Wei et al., 2018), our seasonal
415 ecohydrological sensitivity prediction model can provide valuable information for the understanding of the relative role of climate, vegetation and watershed characteristics.

Since many watersheds lack long-term monitoring data on climate, hydrology and vegetation, a quantitative assessment of hydrological response to vegetation change at the watershed scale is very challenging and time-consuming. However, physical watershed data on climate, vegetation, and watershed property can be easily derived from on-line climate datasets, remote sensing-based products, DEMs and soil databases. The development of a seasonal ecohydrological sensitivity prediction model can be an efficient tool for watershed managers to evaluate hydrological impacts of vegetation restoration programs with easily accessible data on climate, vegetation, topography, soil and landscape. Once seasonal ecohydrological sensitivity for different watersheds can be predicted quickly, future forest management can be implemented in a more sustainable way. We expect that the assessment framework from this study can be effectively applied to any watersheds where physical watershed data are available to support sustainable watershed planning and management.

5.7 Uncertainties and limitations

This study may have some uncertainties and limitations regarding the ecohydrological sensitivity quantification and its prediction model development. The accuracy of ecohydrological sensitivity quantification relies on the methodology for quantifying seasonal streamflow variation attributed to vegetation change. In this study, the improved single watershed approach used to separate the effects of vegetation change, climate variability and other factors on seasonal streamflow has several limitations. An important assumption of this approach is that the vegetation-water relationship during the study period must be stationary, which limits its application under nonstationary conditions. In addition, various watershed disturbances such as urbanization, dam regulations, and other human activities are considered as an integrated driver (other factors). Thus, the impact of each watershed disturbance (e.g., urbanization, dam regulation, and irrigation) cannot be quantified separately.

Given the ecohydrological sensitivity prediction models were generated from only 14 large representative watersheds, an uncertainty associated with the sample size may arise. Admittedly, a larger number of study watersheds would yield more robust conclusions. However, the quantification of vegetation impact on seasonal streamflow involves tremendous data processing analyses for each watershed, and there is a trade-off between the number of study watersheds and workload.

The selection of indices and models may also give rise to some uncertainties and limitations of the prediction models. In this study, topographic and landscape indices were identified based on previously published studies, which were most frequently used in studying the topographic and landscape effects on hydrological processes. As is known, every feature can have a certain impact on the watershed hydrological responses. For example, area, perimeter, mean elevation, and elevation differences provide basic topographic conditions for each watershed, showing watershed heterogeneity. Slope, flow path length (Length), and slope length factor (LS) are indices used for assessing erosion hazard. Topographic wetness index (TWI) is a critical topographic index related to soil water content and surface saturation. Shannon's diversity index (SHDI) and Simpson's diversity index (SIDI) could be applied to indicate a patch diversity of landscape. The ideal way is to include all indices in the analysis. Nevertheless, some of these indices are highly linearly related to others, possibly resulting in a multicollinearity problem in a prediction model. In this study, multicollinear relationships between these indices were detected and confirmed first and then to identify the key factors mostly related to seasonal ecohydrological sensitivities by factor analysis and stepwise

450 regression. The whole selection process is a trade-off between the model complexity and model performance. In addition, our linear prediction models fail to capture some non-linear relationships between ecohydrological sensitivity and its influencing factors. Other methodologies such as machine learning or neural network could be applied to explore non-linear relationships between ecohydrological sensitivity and its influencing factors with a sufficient sample size in future studies.

6 Conclusions

455 Ecohydrological sensitivities at the seasonal scale were quantified in 14 large watersheds across various environmental gradients in China. Our main conclusions are: (1) hydrological responses were greater and more sensitive under dry conditions than wet conditions; (2) seasonal ecohydrological sensitivities were highly variable across climate gradient, dominant soil type and hydrological regime; and (3) dry season ecohydrological sensitivity could be better controlled by vegetation, topography and soil while wet season hydrological sensitivity by vegetation, soil and landscape. Our study also demonstrated the
460 usefulness of constructing an ecohydrological sensitivity prediction model for predicting ecohydrological sensitivity in ungauged watersheds or watersheds with insufficient hydrological data to help watershed managers to effectively manage hydrological impacts and risks through vegetation restoration programs.

Data availability. Climate, vegetation, topography, soil and landscape indices of study watersheds are freely available upon
465 request by sending an email to the corresponding author.

Author contributions. YH and MF proposed the analysis, designed the experiment, performed the result analysis and wrote the paper. QL and XW interpreted results and reviewed the manuscript. SL, TC, WL and XL collected the data. RZ calculated landscape indices. All authors participated in the ‘Discussion’ section and manuscript revision.

470
Competing interests. The authors declare that they have no conflict of interest.

Financial support. This paper was supported by National Program on Key Research and Development Project of China (No. 2017YFC0505006) and National Natural Science Foundation of China (No.31770759).

475
Acknowledgements. We would like to thank the editor, Dr. Stacey Archfield, and two anonymous reviewers for their constructive comments and suggestions on this paper.

References

- Ambroise, B., Beven, K., Freer, J.: Toward a generalization of the TOPMODEL concepts: Topographic indices of hydrological similarity. *Water Resour. Res.*, 32(7), 2135-2145, 1996.
- 480 Arias, M. E., Cochrane, T. A., Piman, T., Kumm, M., Caruso, B. S., and Killeen, T. J.: Quantifying changes in flooding and habitats in the Tonle Sap Lake (Cambodia) caused by water infrastructure development and climate change in the Mekong Basin, *J. Environ. Manag.*, 112, 53-66, <https://doi.org/10.1016/j.jenvman.2012.07.003>, 2012.
- Asbjornsen, H., Goldsmith, G. R., Alvarado-Barrientos, M. S., Rebel, K., Van Osch, F. P., Rietkerk, M., Chen, J., Gotsch, S., Tobón, C., Geissert, D. R., Gómez-Tagle, A., Vache, K., and Dawson, T. E.: Ecohydrological advances and applications in plant-water relations research: a review, *J. Plant Ecol.*, 4, 3-22, 10.1093/jpe/rtr005, 2011.
- 485 Baker, M. E., and Wiley, M. J.: Multiscale control of flooding and riparian-forest composition in Lower Michigan, USA, *Ecology*, 90, 145-159, Doi 10.1890/07-1242.1, 2009.
- Beck, H. E., Bruijnzeel, L. A., van Dijk, A. I. J. M., McVicar, T. R., Scatena, F. N., and Schellekens, J.: The impact of forest regeneration on streamflow in 12 mesoscale humid tropical catchments, *Hydrol. Earth Syst. Sci.*, 17, 2613-2635, 10.5194/hess-17-2613-2013, 2013.
- 490 Birnbaum, Z. W.: On a Use of the Mann-Whitney Statistic, in: Third Berkeley Symposium on Mathematical Statistics and Probability, Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, Volume 1: Contributions to the Theory of Statistics, Berkeley, Calif., 13.17, 1956
- 495 Bisantino, T., Bingner, R., Chouaib, W., Gentile, F., and Liuzzi, G. T.: Estimation of runoff, peak discharge and sediment load at the event scale in a medium-size Mediterranean watershed using the Annagnps model, *Land Degrad. Dev.*, 26, 340-355, 10.1002/ldr.2213, 2015.
- Bockheim, J. G., and Gennadiyev, A. N.: Soil-factorial models and earth-system science: A review, *Geoderma*, 159, 243-251, <https://doi.org/10.1016/j.geoderma.2010.09.005>, 2010.
- 500 Boongaling, C. G. K., Faustino-Eslava, D. V., and Lansigan, F. P.: Modeling land use change impacts on hydrology and the use of landscape metrics as tools for watershed management: The case of an ungauged catchment in the Philippines. *Land use policy*, 72, 116-128, 10.1016/j.landusepol.2017.12.042, 2008.
- Borselli, L., Cassi, P., and Torri, D.: Prolegomena to sediment and flow connectivity in the landscape: A GIS and field numerical assessment, *CATENA*, 75, 268-277, <https://doi.org/10.1016/j.catena.2008.07.006>, 2008.
- 505 Brooks, K. N., Ffolliott, P. F., and Magner, J. A.: Integrated Watershed Management, Hydrology and the Management of Watersheds, 329-331, doi:10.1002/9781118459751.part3, 2012.
- Bruijnzeel, L. A., Mulligan, M., and Scatena, F. N.: Hydrometeorology of tropical montane cloud forests: emerging patterns, *Hydrol. Process.*, 25, 465-498, 10.1002/hyp.7974, 2011.
- Buma, B., and Livneh, B.: Key landscape and biotic indicators of watersheds sensitivity to forest disturbance identified using remote sensing and historical hydrography data, *Environ. Res. Lett.*, 12, 074028, 10.1088/1748-9326/aa7091, 2017.
- 510

- Bunn, S. E., Thoms, M. C., Hamilton, S. K., and Capon, S. J.: Flow variability in dryland rivers: boom, bust and the bits in between, *River Res. Appl.*, 22, 179-186, 10.1002/rra.904, 2006.
- Calder, I. R.: *Blue revolution: Integrated land and water resource management*, Routledge, 2005.
- Chadli, K.: Estimation of soil loss using RUSLE model for Sebou watershed (Morocco), *Model. Earth Syst. Environ.*, 2, 51, 10.1007/s40808-016-0105-y, 2016.
- Chang, M.: *Forest hydrology: An introduction to water and forests*, CRC Press, Boca Raton, Florida, 2012.
- Creed, I. F., Spargo, A. T., Jones, J. A., Buttle, J. M., Adams, M. B., Beall, F. D., Booth, E. G., Campbell, J. L., Clow, D., Elder, K., Green, M. B., Grimm, N. B., Miniati, C., Ramlal, P., Saha, A., Sebestyen, S., Spittlehouse, D., Sterling, S., Williams, M. W., Winkler, R., and Yao, H.: Changing forest water yields in response to climate warming: results from long-term experimental watershed sites across North America, *Global Change Biol.*, 20, 3191-3208, 10.1111/gcb.12615, 2014.
- Dai, A.: Characteristics and trends in various forms of the Palmer Drought Severity Index during 1900-2008, *J. Geophys. Res-Atmos.*, 116, 10.1029/2010JD015541, 2011.
- de Paula, F. R., Ferraz, S. F. d. B., Gerhard, P., Vettorazzi, C. A., and Ferreira, A.: Large woody debris input and its influence on channel structure in agricultural lands of Southeast Brazil, *Environ. Manage.*, 48, 750-763, 10.1007/s00267-011-9730-4, 2011.
- de Sarrau, B., Clavel, T., Clerté, C., Carlin, F., Giniès, C., and Nguyen-The, C.: Influence of anaerobiosis and low temperature on *Bacillus cereus* growth, metabolism, and membrane properties, *Appl. Environ. Microb.*, 78, 1715, 10.1128/AEM.06410-11, 2012.
- Desmet, P. J. J., and Govers, G.: A GIS procedure for automatically calculating the USLE LS factor on topographically complex landscape units, *J. Soil Water Conserv.*, 51, 427-433, 1996.
- Donohue, R. J., Roderick, M. L., and McVicar, T. R.: Can dynamic vegetation information improve the accuracy of Budyko's hydrological model?, *J. Hydrol.*, 390, 23-34, <https://doi.org/10.1016/j.jhydrol.2010.06.025>, 2010.
- Farley, K. A., Jobbagy, E. G., and Jackson, R. B.: Effects of afforestation on water yield: a global synthesis with implications for policy, *Global Change Biol.*, 11, 1565-1576, 10.1111/j.1365-2486.2005.01011.x, 2005.
- Feng, X., Fu, B., Piao, S., Wang, S., Ciais, P., Zeng, Z., Lü, Y., Zeng, Y., Li, Y., and Jiang, X.: Revegetation in China's Loess Plateau is approaching sustainable water resource limits, *Nat. Clim. Change*, 6, 1019, 2016.
- Frank, D. C., Poulter, B., Saurer, M., Esper, J., Huntingford, C., Helle, G., Treydte, K., Zimmermann, N. E., Schleser, G. H., Ahlström, A., Ciais, P., Friedlingstein, P., Levis, S., Lomas, M., Sitch, S., Viovy, N., Andreu-Hayles, L., Bednarz, Z., Berninger, F., Boettger, T., D'Alessandro, C. M., Daux, V., Filot, M., Grabner, M., Gutierrez, E., Haupt, M., Hilasvuori, E., Jungner, H., Kalela-Brundin, M., Krapiec, M., Leuenberger, M., Loader, N. J., Marah, H., Masson-Delmotte, V., Pazdur, A., Pawelczyk, S., Pierre, M., Planells, O., Pukiene, R., Reynolds-Henne, C. E., Rinne, K. T., Saracino, A., Sonninen, E., Stievenard, M., Switsur, V. R., Szczepanek, M., Szychowska-Krapiec, E., Todaro, L., Waterhouse, J. S., and Weigl, M.: Water-use efficiency and transpiration across European forests during the Anthropocene, *Nat. Clim. Change*, 5, 579-583, 10.1038/nclimate2614, 2015.

- 545 González-Sanpedro, M. C., Le Toan, T., Moreno, J., Kergoat, L., and Rubio, E.: Seasonal variations of leaf area index of agricultural fields retrieved from Landsat data, *Remote Sens. Environ.*, 112, 810-824, <https://doi.org/10.1016/j.rse.2007.06.018>, 2008.
- Guswa, A. J., Tetzlaff, D., Selker, J. S., Carlyle-Moses, D. E., Boyer, E. W., Bruen, M., Cayuela, C., Creed, I. F., van de Giesen, N., Grasso, D., Hannah, D. M., Hudson, J. E., Hudson, S. A., Iida, S., Jackson, R. B., Katul, G. G., Kumagai, T.,
- 550 Llorens, P., Lopes Ribeiro, F., Michalzik, B., Nanko, K., Oster, C., Pataki, D. E., Peters, C. A., Rinaldo, A., Sanchez Carretero, D., Trifunovic, B., Zalewski, M., Haagsma, M., and Levia, D. F.: Advancing ecohydrology in the 21st century: A convergence of opportunities, *Ecohydrology*, 13, e2208, 10.1002/eco.2208, 2020.
- Hardy, M. A.: *Regression with dummy variables*, Sage Publications, Inc, Thousand Oaks, CA, US, 90, 1993.
- Hargreaves, G., and Samani, Z.: Reference crop evapotranspiration from temperature, *Appl. Eng. Agric.*, 1, 96-99, <https://doi.org/10.13031/2013.26773>, 1985.
- 555 Hernandez, M., Kepner, W. G., Semmens, D. J., Ebert, D. W., Goodrich, D. C., and Miller, S. N.: Integrating a Landscape/Hydrologic Analysis for Watershed Assessment, In proceedings of the First Interagency Conference on Research in the Watersheds, Agricultural Research Service, Benson, AZ, USA, 27-30 October 2003, 2003.
- Hillel, D.: *Soil and water: Physical principles and processes*, 1974.
- 560 Hirabayashi, Y., Mahendran, R., Koiralala, S., Konoshima, L., Yamazaki, D., Watanabe, S., Kim, H., and Kanae, S.: Global flood risk under climate change, *Nat. Clim. Change*, 3, 816-821, 10.1038/nclimate1911, 2013.
- Hou, Y., Zhang, M., Liu, S., Sun, P., Yin, L., Yang, T., Li, Y., Li, Q., and Wei, X.: The hydrological impact of extreme weather-induced forest disturbances in a tropical experimental watershed in South China, *Forests*, 9, 10.3390/f9120734, 2018a.
- 565 Hou, Y., Zhang, M., Meng, Z., Liu, S., Sun, P., and Yang, T.: Assessing the impact of forest change and climate variability on dry season runoff by an improved single watershed approach: A comparative study in two large watersheds, China, *Forests*, 9, 46, 10.3390/f9010046, 2018b.
- Jackson, R. B., Jobbágy, E. G., Avissar, R., Roy, S. B., Barrett, D. J., Cook, C. W., Farley, K. A., le Maitre, D. C., McCarl, B. A., and Murray, B. C.: Trading Water for Carbon with Biological Carbon Sequestration, *Science*, 310, 1944, [10.1126/science.1119282](https://doi.org/10.1126/science.1119282), 2005.
- 570 Jansen, J. D., and Nanson, G. C.: Functional relationships between vegetation, channel morphology, and flow efficiency in an alluvial (anabranching) river, *J. Geophys. Res.*, 115, 10.1029/2010JF001657, 2010.
- Jencso, K. G., and McGlynn, B. L.: Hierarchical controls on runoff generation: Topographically driven hydrologic connectivity, geology, and vegetation, *Water Resour. Res.*, 47, 10.1029/2011WR010666, 2011.
- 575 Jenness, J. S.: Calculating landscape surface area from digital elevation models, *Wildlife Soc. B.*, 32, 829-839, [Doi 10.2193/0091-7648\(2004\)032\[0829:Clsafd\]2.0.Co;2](https://doi.org/10.2193/0091-7648(2004)032[0829:Clsafd]2.0.Co;2), 2004.
- Jones, J. A., Creed, I. F., Hatcher, K. L., Warren, R. J., Adams, M. B., Benson, M. H., Boose, E., Brown, W. A., Campbell, J. L., Covich, A., Clow, D. W., Dahm, C. N., Elder, K., Ford, C. R., Grimm, N. B., Henshaw, D. L., Larson, K. L., Miles, E.

- S., Miles, K. M., Sebestyen, S. D., Spargo, A. T., Stone, A. B., Vose, J. M., and Williams, M. W.: Ecosystem processes and human influences regulate streamflow response to climate change at long-term ecological research sites, *BioScience*, 62, 390-404, 10.1525/bio.2012.62.4.10, 2012.
- Karlsen, R. H., Grabs, T., Bishop, K., Buffam, I., Laudon, H., and Seibert, J.: Landscape controls on spatiotemporal discharge variability in a boreal catchment, *Water Resour. Res.*, 52, 6541-6556, 10.1002/2016WR019186, 2016.
- Kokkonen, T. S., and Jakeman, A. J.: Chapter 14 Structural effects of landscape and land use on streamflow response, In: *Developments in Environmental Modelling*, edited by: Beck, M. B., Elsevier, 303-321, 2002.
- Launiainen, S., Katul, G. G., Kolari, P., Lindroth, A., Lohila, A., Aurela, M., Varlagin, A., Grelle, A., and Vesala, T.: Do the energy fluxes and surface conductance of boreal coniferous forests in Europe scale with leaf area?, *Global Change Biol.*, 22, 4096-4113, 10.1111/gcb.13497, 2016.
- Leu, J., Traore, S., Wang, Y., and Kan, C. E.: The effect of organic matter amendment on soil water holding capacity change for irrigation water saving: case study in Sahelian environment of Africa, *Sci. Res. Essays*, 5, 3564-3571, 2010.
- Li, Q., Wei, X., Zhang, M., Liu, W., Fan, H., Zhou, G., Giles-Hansen, K., Liu, S., and Wang, Y.: Forest cover change and water yield in large forested watersheds: A global synthetic assessment, *Ecohydrology*, 10, 2017.
- Li, Q., Wei, X., Yang, X., Giles-Hansen, K., Zhang, M., and Liu, W.: Topography significantly influencing low flows in snow-dominated watersheds, *Hydrol. Earth Syst. Sci.*, 22, 1947-1956, 10.5194/hess-22-1947-2018, 2018a.
- Li, Y., Piao, S., Li, L., Chen, A., Wang, X., Ciais, P., Huang, L., Lian, X., Peng, S., Zeng, Z., Wang, K., and Zhou, L.: Divergent hydrological response to large-scale afforestation and vegetation greening in China, *Sci. Adv.*, 4, 10.1126/sciadv.aar4182, 2018b.
- Liang, S., Zhao, X., Liu, S., Yuan, W., Cheng, X., Xiao, Z., Zhang, X., Liu, Q., Cheng, J., Tang, H., Qu, Y., Bo, Y., Qu, Y., Ren, H., Yu, K., and Townshend, J.: A long-term Global Land Surface Satellite (GLASS) data-set for environmental studies, *Int. J. Digit. Earth*, 6, 5-33, 10.1080/17538947.2013.805262, 2013.
- Lin, Y., and Wei, X.: The impact of large-scale forest harvesting on hydrology in the Willow watershed of Central British Columbia, *J. Hydrol.*, 359, 141-149, 2008.
- Lyon, S. W., Nathanson, M., Spans, A., Grabs, T., Laudon, H., Temnerud, J., Bishop, K. H., and Seibert, J.: Specific discharge variability in a boreal landscape, *Water Resour. Res.*, 48, 10.1029/2011wr011073, 2012.
- Maeda, E. E., Kim, H., Aragão, L. E. O. C., Famiglietti, J. S., and Oki, T.: Disruption of hydroecological equilibrium in southwest Amazon mediated by drought, *Geophys. Res. Lett.*, 42, 7546-7553, 10.1002/2015GL065252, 2015.
- McDonnell, J. J., Evaristo, J., Bladon, K. D., Buttle, J., Creed, I. F., Dymond, S. F., Grant, G., Iroume, A., Jackson, C. R., Jones, J. A., Maness, T., McGuire, K. J., Scott, D. F., Segura, C., Sidle, R. C., and Tague, C.: Water sustainability and watershed storage, *Nat. Sustain.*, 1, 378-379, 10.1038/s41893-018-0099-8, 2018.
- McVicar, T. R., Roderick, M. L., Donohue, R. J., and Van Niel, T. G.: Less bluster ahead? Ecohydrological implications of global trends of terrestrial near-surface wind speeds, *Ecohydrology*, 5, 381-388, 10.1002/eco.1298, 2012.


- Miara, A., Macknick, J. E., Vörösmarty, C. J., Tidwell, V. C., Newmark, R., and Fekete, B.: Climate and water resource change impacts and adaptation potential for US power supply, *Nat. Clim. Change*, 7, 793-798, 10.1038/nclimate3417, 2017.
- Moore, I. D., Grayson, R. B., and Ladson, A. R.: Digital Terrain Modeling - A Review of Hydrological, Geomorphological, and Biological Applications, *Hydrol. Process.*, 5, 3-30, DOI 10.1002/hyp.3360050103, 1991.
- Moore, R. D., and Wondzell, S. M.: Physical hydrology and the effects of forest harvesting in the Pacific Northwest: A review, *J. Am. Water Resour. As.*, 41, 763-784, 2005.
- Mukundan, R., Radcliffe, D. E., and Risse, L. M.: Spatial resolution of soil data and channel erosion effects on SWAT model predictions of flow and sediment, *J. Soil Water Conserv.*, 65, 92-104, 2010.
- Newman, B. D., Wilcox, B. P., Archer, S. R., Breshears, D. D., Dahm, C. N., Duffy, C. J., McDowell, N. G., Phillips, F. M., Scanlon, B. R., and Vivoni, E. R.: Ecohydrology of water-limited environments: A scientific vision, *Water Resour. Res.*, 42, 10.1029/2005WR004141, 2006.
- Nippgen, F., McGlynn, B. L., Marshall, L. A., and Emanuel, R. E.: Landscape structure and climate influences on hydrologic response, *Water Resour. Res.*, 47, 10.1029/2011WR011161, 2011.
- Oppel, H., Schumann, A. H.: Machine learning based identification of dominant controls on runoff dynamics, *Hydrol. Process.* 34, 2450-2465. <https://doi.org/10.1002/hyp.13740>, 2020
- Palmer, M., and Ruhi, A.: Linkages between flow regime, biota, and ecosystem processes: Implications for river restoration, *Science*, 365, eaaw2087, 10.1126/science.aaw2087, 2019.
- Park, S. J., McSweeney, K., and Lowery, B.: Identification of the spatial distribution of soils using a process-based terrain characterization, *Geoderma*, 103, 249-272, 10.1016/S0016-7061(01)00042-8, 2001.
- Price, K.: Effects of watershed topography, soils, land use, and climate on baseflow hydrology in humid regions: A review, *Prog. Phys. Geog.*, 35, 465-492, 10.1177/0309133311402714, 2011.
- Rieu, M., and Sposito, G.: Fractal fragmentation, soil porosity, and soil water properties: I. Theory, *Soil Sci. Soc. Am. J.*, 55, 1231-1238, 10.2136/sssaj1991.036159950005500050006x, 1991.
- Salve, R., Sudderth, E. A., St. Clair, S. B., and Torn, M. S.: Effect of grassland vegetation type on the responses of hydrological processes to seasonal precipitation patterns, *J. Hydrol.*, 410, 51-61, 10.1016/j.jhydrol.2011.09.003, 2011.
- Schoonover, J. E., and Crim, J. F.: An Introduction to Soil Concepts and the Role of Soils in Watershed Management, *J. Contemp. Water Res. Educ*, 154, 21-47, 10.1111/j.1936-704X.2015.03186.x, 2015.
- Scown, M. W., Thoms, M. C., and De Jager, N. R.: Measuring floodplain spatial patterns using continuous surface metrics at multiple scales, *Geomorphology*, 245, 87-101, 10.1016/j.geomorph.2015.05.026, 2015.
- Simonit, S., and Perrings, C.: Reply to Ogden and Stallard: Phenomenological runoff models in the Panama Canal watershed, *P. Natl. Acad. Sci. USA.*, 110, E5038, 10.1073/pnas.1318590111, 2013.
- Singh, R., Wagener, T., Crane, R., Mann, M. E., and Ning, L.: A vulnerability driven approach to identify adverse climate and land use change combinations for critical hydrologic indicator thresholds: Application to a watershed in Pennsylvania, USA, *Water Resour. Res.*, 50, 3409-3427, 10.1002/2013WR014988, 2014.

- Smakhtin, V. U.: Low flow hydrology: a review, *J. Hydrol.*, 240, 147-186, 10.1016/S0022-1694(00)00340-1, 2001.
- Srivastava, A., Babu, G. L. S., and Haldar, S.: Influence of spatial variability of permeability property on steady state seepage flow and slope stability analysis, *Eng. Geol.*, 110, 93-101, <https://doi.org/10.1016/j.enggeo.2009.11.006>, 2010.
- 650 Sulis, M., Meyerhoff, S. B., Paniconi, C., Maxwell, R. M., Putti, M., and Kollet, S. J.: A comparison of two physics-based numerical models for simulating surface water-groundwater interactions, *Adv. Water Resour.*, 33, 456-467, <https://doi.org/10.1016/j.advwatres.2010.01.010>, 2010.
- Sulla-Menashe, D., Gray, J. M., Abercrombie, S. P., and Friedl, M. A.: Hierarchical mapping of annual global land cover 2001 to present: The MODIS Collection 6 Land Cover product, *Remote Sens. Environ.*, 222, 183-194, <https://doi.org/10.1016/j.rse.2018.12.013>, 2019.
- 655 Sun, G., Zhou, G. Y., Zhang, Z. Q., Wei, X. H., McNulty, S. G., and Vose, J. M.: Potential water yield reduction due to forestation across China, *J. Hydrol.*, 328, 548-558, 10.1016/j.jhydrol.2005.12.013, 2006.
- Sun, G., Amatya, D., and McNulty, S.: Forest hydrology, in, 85-81-85-88, 2016.
- Teutschbein, C., Grabs, T., Laudon, H., Karlsen, R. H., and Bishop, K.: Simulating streamflow in ungauged basins under a changing climate: The importance of landscape characteristics, *J. Hydrol.*, 561, 160-178, 10.1016/j.jhydrol.2018.03.060, 660 2018.
- Toledo-Aceves, T., Meave, J. A., González-Espinosa, M., and Ramírez-Marcial, N.: Tropical montane cloud forests: Current threats and opportunities for their conservation and sustainable management in Mexico, *J. Environ. Manage.*, 92, 974-981, <https://doi.org/10.1016/j.jenvman.2010.11.007>, 2011.
- van Dijk, A. I. J. M., Peña-Arancibia, J. L., and Bruijnzeel, L. A.: Land cover and water yield: inference problems when 665 comparing catchments with mixed land cover, *Hydrol. Earth Syst. Sci.*, 16, 3461-3473, 10.5194/hess-16-3461-2012, 2012.
- Van Dover, C. L., and Lutz, R. A.: Experimental ecology at deep-sea hydrothermal vents: a perspective, *J. Exp. Mar. Biol. Ecol.*, 300, 273-307, <https://doi.org/10.1016/j.jembe.2003.12.024>, 2004.
- Verrelst, J., van der Tol, C., Magnani, F., Sabater, N., Rivera, J. P., Mohammed, G., and Moreno, J.: Evaluating the predictive power of sun-induced chlorophyll fluorescence to estimate net photosynthesis of vegetation canopies: A SCOPE 670 modeling study, *Remote Sens. Environ.*, 176, 139-151, <https://doi.org/10.1016/j.rse.2016.01.018>, 2016.
- Voepel, H., Ruddell, B., Schumer, R., Troch, P. A., Brooks, P. D., Neal, A., Durcik, M., and Sivapalan, M.: Quantifying the role of climate and landscape characteristics on hydrologic partitioning and vegetation response, *Water Resour. Res.*, 47, 10.1029/2010WR009944, 2011.
- Vose, J. M., Sun, G., Ford, C. R., Bredemeier, M., Otsuki, K., Wei, X., Zhang, Z., and Zhang, L.: Forest ecohydrological 675 research in the 21st century: what are the critical needs?, *Ecohydrology*, 4, 146-158, 2011.
- Warfe, D. M., Pettit, N. E., Davies, P. M., Pusey, B. J., Hamilton, S. K., Kennard, M. J., Townsend, S. A., Bayliss, P., Ward, D. P., Douglas, M. M., Burford, M. A., Finn, M., Bunn, S. E., and Halliday, I. A.: The 'wet-dry' in the wet-dry tropics drives river ecosystem structure and processes in northern Australia, *Freshwater Biol.*, 56, 2169-2195, 10.1111/j.1365-2427.2011.02660.x, 2011.

- 680 Wei, X., Sun, G., Liu, S., Jiang, H., Zhou, G., and Dai, L.: The forest-streamflow relationship in China: A 40-year retrospect, *J. Am. Water Resour. As.*, 44, 1076-1085, 10.1111/j.1752-1688.2008.00237.x, 2008.
- Wei, X., Li, Q., Zhang, M., Giles-Hansen, K., Liu, W., Fan, H., Wang, Y., Zhou, G., Piao, S., and Liu, S.: Vegetation cover—another dominant factor in determining global water resources in forested regions, *Global Change Biol.*, 24, 786-795, 10.1111/gcb.13983, 2018.
- 685 Wieder, W.: *Regridded Harmonized World Soil Database v1.2*. ORNL Distributed Active Archive Center. 10.3334/ORNLDAAC/1247, 2014.
- Winkler, R., Boon, S., Zimonick, B., and Baleshta, K.: Assessing the effects of post-pine beetle forest litter on snow albedo, *Hydrol. Process.*, 24, 803-812, 10.1002/hyp.7648, 2010.
- Woods, R.: The relative roles of climate, soil, vegetation and topography in determining seasonal and long-term catchment dynamics, *Adv. Water Resour.*, 30, 1061-1061, 10.1016/j.advwatres.2006.10.010, 2007.
- 690 Xiao, Z., Liang, S., Wang, J., Chen, P., Yin, X., Zhang, L., and Song, J.: Use of General Regression Neural Networks for Generating the GLASS Leaf Area Index Product From Time-Series MODIS Surface Reflectance, *IEEE T. Geosci. Remote Sens.*, 52, 209-223, 10.1109/TGRS.2013.2237780, 2014.
- Yang, D., Sun, F., Liu, Z., Cong, Z., Ni, G., and Lei, Z.: Analyzing spatial and temporal variability of annual water-energy balance in nonhumid regions of China using the Budyko hypothesis, *Water Resour. Res.*, 43, 10.1029/2006WR005224, 2007.
- 695 Yokoyama, R., Shirasawa, M., and Pike, R. J.: Visualizing topography by openness: a new application of image processing to digital elevation models, *Photogramm. Eng. Rem. S.*, 68, 257-266, 2002.
- Zeng, C., Yang, L., Zhu, A., Rossiter, D., Liu, J., Liu, J., Qin, C., and Wang, D.: Mapping soil organic matter concentration at different scales using a mixed geographically weighted regression method, *Geoderma*, 281, 69-82, 10.1016/j.geoderma.2016.06.033, 2016.
- 700 Zhang, L., Dawes, W.R., Walker, G.R.: Response of mean annual evapotranspiration to vegetation changes at catchment scale. *Water Resour. Res.*, 37(3), 701-708. doi:10.1029/2000wr900325, 2001.
- Zhang, L., Hickel, K., Dawes, W. R., Chiew, F. H. S., Western, A. W., and Briggs, P. R.: A rational function approach for estimating mean annual evapotranspiration, *Water Resour. Res.*, 40, 10.1029/2003WR002710, 2004.
- 705 Zhang, M., Ren, Q., Wei, X., Wang, J., Yang, X., and Jiang, Z.: Climate change, glacier melting and streamflow in the Niyang River Basin, Southeast Tibet, China, *Ecohydrology*, 4, 288-298, 10.1002/eco.206, 2011.
- Zhang, M., and Wei, X.: The effects of cumulative forest disturbance on streamflow in a large watershed in the central interior of British Columbia, Canada, *Hydrol. Earth Syst. Sci.*, 16, 2021-2034, 10.5194/hess-16-2021-2012, 2012.
- 710 Zhang, M., Wei, X., Sun, P., and Liu, S.: The effect of forest harvesting and climatic variability on runoff in a large watershed: The case study in the Upper Minjiang River of Yangtze River basin, *J. Hydrol.*, 464, 1-11, 10.1016/j.jhydrol.2012.05.050, 2012.

- Zhang, M., Liu, N., Harper, R., Li, Q., Liu, K., Wei, X., Ning, D., Hou, Y., and Liu, S.: A global review on hydrological responses to forest change across multiple spatial scales: Importance of scale, climate, forest type and hydrological regime, *J. Hydrol.*, 546, 44-59, 10.1016/j.jhydrol.2016.12.040, 2017.
- 715 Zhao, Y., Peth, S., Wang, X., Lin, H., and Horn, R.: Controls of surface soil moisture spatial patterns and their temporal stability in a semi-arid steppe, *Hydrol. Process.*, 24, 2507-2519, 10.1002/hyp.7665, 2010.
- Zhou, G., Wei, X., Chen, X., Zhou, P., Liu, X., Xiao, Y., Sun, G., Scott, D. F., Zhou, S., Han, L., and Su, Y.: Global pattern for the effect of climate and land cover on water yield, *Nat. Commun.*, 6, 10.1038/Ncomms6918, 2015.
- 720 Zhou, Z., and Li, J.: The correlation analysis on the landscape pattern index and hydrological processes in the Yanhe watershed, China. *J. Hydrol.*, 524, 417-426, 10.1016/j.jhydrol.2015.02.028, 2015.

Figures:



725 **Figure 1.** Locations of the the study watersheds.







Figure 2. A comparison of ecohydrological sensitivity in dry season and wet season. (S_d and S_w are dry season ecohydrological sensitivity and wet season ecohydrological sensitivity)



730

Figure 3. Comparisons of ecohydrological sensitivity grouped by energy-limited (EL), equitant (EQ) and water-limited (WL) conditions in (a) dry season and (b) wet season. (S_{fd} and S_{fw} are dry season ecohydrological sensitivity and wet season ecohydrological sensitivity)



735 **Figure 4.** Comparisons of ecohydrological sensitivity grouped by dominant soil type in (a) dry season and (b) wet season.






Figure 5. Comparisons of ecohydrological sensitivity grouped by rain-dominated (RD) and hybrid regimes in (a) dry season and (b) wet season.



740



745

Figure 6. Comparisons of observed and predicted ecohydrological sensitivity in (a) dry season and (b) wet season. (TR, XR, PJ, XS, GBJD, GZ, UZGN, ZGN, UHR, HR, DC, HSC, JC and RR refer to the Tangwang River, Xinancha River, Pingjiang, Xiangshui, Gongbujiangda, Gengzhang, Upper Zagunao, Zagunao, Upper Heishui River, Heishui River, Dongchuan, Heishuichuan, Jingchuan and Rui River watersheds, respectively)

Tables:

Table 1. Watershed characteristics in the study watersheds.

| Watersheds | Area (km ²) | Mean elevation (m) | Slope (°) | Climate zone | Dry season | | | | | Wet season | | | | | | | | |
|---------------------|-------------------------|--------------------|-----------|--------------|--------------------|------------------------|--------|---------|--------|---------------------------------------|--------------|------------------------|-------------|---------|--------|---------------------------------------|-------|------|
| | | | | | Period | T _{mean} (°C) | P (mm) | ET (mm) | Q (mm) | LAI (m ² /m ²) | Period | T _{mean} (°C) | P (mm) | ET (mm) | Q (mm) | LAI (m ² /m ²) | | |
| Pingjiang | 2778 | 314 | 15.1 | SMC | September-February | 13.5 | 501.1 | 254.0 | 236.2 | 1.45 | March-August | 22.3 | 1310.7 | 585.5 | 604.3 | 1.90 | | |
| Xiangshui | 1742 | 440 | 17.6 | | | 14.3 | 472.0 | 274.6 | 242.5 | 2.54 | | 22.0 | 1402.7 | 659.5 | 616.4 | 3.17 | | |
| Tangwang River | 19189 | 447 | 8.7 | TCMC | | -9.6 | 60.3 | 42.4 | 30.1 | 0.73 | | 14.8 | 517.1 | 367.4 | 239.9 | 3.53 | | |
| Xinancha River | 2585 | 507 | 11.3 | | | -11.1 | 81.0 | 47.5 | 37.3 | 0.71 | | 12.8 | 567.5 | 398.9 | 293.6 | 3.59 | | |
| Upper Zagunao | 2442 | 3814 | 31.0 | | | 5.3 | 190.3 | 146.2 | 176.9 | 0.83 | | 17.0 | 848.6 | 500.4 | 672.9 | 1.87 | | |
| Zagunao | 4629 | 3622 | 31.7 | AC | | 5.0 | 164.4 | 139.9 | 144.0 | 0.86 | | 16.6 | 759.9 | 484.2 | 583.2 | 2.14 | | |
| Upper Heishui River | 1710 | 3858 | 27.8 | | | -1.8 | 121.1 | 102.7 | 136.9 | 0.50 | | 9.7 | 599.8 | 408.6 | 630.0 | 1.78 | | |
| Heishui River | 7170 | 3619 | 27.3 | | | November-April | -1.8 | 121.1 | 103.0 | 117.8 | | 0.53 | May-October | 9.7 | 599.8 | 410.5 | 471.2 | 1.93 |
| Gongbujiangda | 6323 | 4946 | 27.2 | | | | 2.5 | 61.0 | 52.3 | 60.8 | | 0.11 | | 12.9 | 611.8 | 352.4 | 530.7 | 0.45 |
| Gengzhang | 16000 | 4752 | 28.3 | | | | 3.8 | 83.1 | 68.2 | 95.8 | | 0.22 | | 13.4 | 783.6 | 404.8 | 880.3 | 0.59 |
| Dongchuan | 3049 | 1415 | 16.3 | TCC | | 0.1 | 68.1 | 59.2 | 6.6 | 0.19 | | 16.6 | 438.5 | 321.0 | 23.1 | 0.59 | | |
| Heshuichuan | 832 | 1340 | 16.8 | | | 0.9 | 83.5 | 71.1 | 4.7 | 0.34 | | 17.0 | 471.9 | 336.8 | 14.6 | 1.51 | | |
| Jingchuan | 3155 | 1678 | 13.5 | | | 2.2 | 81.1 | 66.7 | 16.6 | 0.26 | | 18.1 | 444.4 | 305.3 | 39.5 | 0.98 | | |
| Rui River | 1688 | 1608 | 13.0 | | | 0.1 | 83.6 | 74.0 | 17.6 | 0.28 | | 15.2 | 488.3 | 364.0 | 55.1 | 1.23 | | |

Note: T_{mean}, P, ET, Q, and LAI stand for mean temperature, precipitation, actual evapotranspiration, streamflow, and leaf area index during the study period. SMC,

750 TCMC, AC and TCC refer to subtropical monsoon climate, temperate continental monsoon climate, alpine climate and temperate continental climate, respectively.

Table 2. Classification of watersheds.

| Watersheds | Climate condition | Dominant soil type | Hydrological regime |
|---------------------|-------------------|--------------------|---------------------|
| Pingjiang | Energy-limited | LIXISOLS | Rain-dominated |
| Xiangshui | Energy-limited | LIXISOLS | Rain-dominated |
| Tangwang River | Equitant | LUVISOLS | Rain-dominated |
| Xinancha River | Equitant | LUVISOLS | Rain-dominated |
| Upper Zagunao | Equitant | LEPTOSOLS | Hybrid |
| Zagunao | Equitant | LEPTOSOLS | Hybrid |
| Upper Heishui River | Equitant | LEPTOSOLS | Hybrid |
| Heishui River | Equitant | LEPTOSOLS | Hybrid |
| Gongbujiangda | Water-limited | LEPTOSOLS | Hybrid |
| Gengzhang | Water-limited | LEPTOSOLS | Hybrid |
| Dongchuan | Water-limited | CAMBISOLS | Rain-dominated |
| Heshuichuan | Water-limited | CAMBISOLS | Rain-dominated |
| Jingchuan | Water-limited | CAMBISOLS | Rain-dominated |
| Rui River | Water-limited | CAMBISOLS | Rain-dominated |

Table 3. Definition or description of the selected influencing factors.

| No. | Category | Abbreviation | Metrics | Definition or description |
|-----|------------|------------------------|--|---|
| 1 | Climate | <i>DI</i> | Dryness index | $DI = PET/P$, annual potential evaporation (PET) was calculated by Hargreaves method (Hargreaves and Samani, 1985). It shows interactions between energy and water and indicates the water availability for vegetation growth. |
| 2 | | <i>P_e</i> | Effective precipitation | $P_e = P - E$, actual evapotranspiration was calculated by Zhang's equation (Zhang et al., 2001). |
| 3 | Vegetation | LAI | Leaf area index | One-half of the total green leaf area per unit of horizontal ground surface area. Derived from GLASS Product. |
| 4 | | Forest coverage | Forest coverage | Forest coverage in a watershed. |
| 5 | | Vegetation coverage | Vegetation coverage | Vegetation coverage in a watershed (total coverage of forest, shrubland and grassland). |
| 6 | Soil | Soil types | Number of soil types | Total number of soil types in a watershed. |
| 7 | | <i>T_{oc}</i> | Topsoil organic carbon | Amount of carbon bound in human, animal and plant residues and microorganisms formed by microbial action in soil. |
| 8 | | <i>S_{oc}</i> | Subsoil organic carbon | |
| 9 | | <i>T_{ece}</i> | Topsoil salinity | Soil total salinity. |
| 10 | | <i>S_{ece}</i> | Subsoil salinity | |
| 11 | | <i>T_w</i> | Topsoil available water holding capacity | Soil moisture in a stable level. |
| 12 | | <i>S_w</i> | Subsoil available water holding capacity | |
| 13 | | <i>T_{hy}</i> | Topsoil saturated hydraulic conductivity | Infiltration rate of each hydraulic gradient. |
| 14 | | <i>S_{hy}</i> | Subsoil saturated hydraulic conductivity | |
| 15 | | <i>T_d</i> | Topsoil bulk density | Soil mass of each volume. |
| 16 | | <i>S_d</i> | Subsoil bulk density | |
| 17 | Landscape | PN | Patch number | Total number of patches within a specified land cover class. |
| 18 | | PD | Patch density | The number of patches per unit area. |
| 19 | | LPI | Largest patch index | The ratio of the largest patch area to total area. |
| 20 | | ED | Edge density | The total length of patches per unit area. |
| 21 | | CONTAG | Contagion index | Indicates the aggregation of patches. |
| 22 | | SHDI | Shannon's diversity index | Based on information theory, indicates the patch diversity in landscape. |
| 23 | | SIDI | Simpson's diversity index | Indicates the patch diversity in landscape. |

Table 3. Definition or description of the selected influencing factors (*continued*).

| No. | Category | Abbreviation | Metrics | Definition or description |
|-----|-------------|--------------------|-------------------------------|---|
| 24 | Topographic | Area | Area of a watershed | Area draining to watershed outlet. |
| 25 | | Perimeter | Perimeter of a watershed | Perimeter of a watershed. |
| 26 | | Elevation | Mean elevation | Mean value of all DEM pixels in a watershed. |
| 27 | | Δ Elevation | Elevation difference | Difference between the highest elevation and the lowest elevation in a watershed. |
| 28 | | Slope | Average slope | Slope degree of each DEM pixel, can be used in estimation of energy budgets. |
| 29 | | LS | Slope Length Factor | A combined factor of slope length and slope gradient. |
| 30 | | Length | Flow Path Length | The average flow path length starting from the seeds. |
| 31 | | Max Length | Maximum Flow Path Length | The maximum distance of water flow to a point. |
| 32 | | TWI | Topographic Wetness Index | $TWI = \ln(SCA/\tan(\text{slope}))$, it shows the spatial distribution of zones of surface saturation and soil water content (Ambroise et al., 1996). |
| 33 | | CON | Convergence | Convergence of a cell, which is calculated based on the surrounding eight cells. 100% convergence means all surrounding grid cells flow into the center cell. |
| 34 | | DDG | Downslope distance gradient | An indicator for assessing the impact of the local slope characteristics on a hydraulic gradient. Values are lower on concave slope profiles and higher on convex slope profiles. |
| 35 | | SA | Surface Area | Land area of each DEM. |
| 36 | | TPI | Topographic Position Index | $TPI \approx 0$ indicates flat area. $TPI > 0$ tends towards ridge tops and hilltops. $TPI < 0$ tends towards the valley and canyon bottoms. |
| 37 | | TRI | Terrain Ruggedness Index | The degree of difference in elevation among adjacent cells. |
| 38 | | PO | Topographic Positive Openness | The degree of dominance or enclosure of a location on an irregular surface. Values are high for convex forms. |
| 39 | | NO | Topographic Negative Openness | |
| 40 | | Depth | Valley depth | Difference between the elevation and an interpolated ridge level. |

Table 4. Mann Whitney U test for ecohydrological sensitivity between dry season and wet season.

| Season | Z | p |
|---------------------------|-------------|--------------|
| Dry season vs. Wet season | 5.63 | 0.00* |

765 *Note:* The bolded number with * indicates statistically significant at $p < 0.05$.

Table 5. Mann Whitney U tests for the differences of seasonal ecohydrological sensitivity between climate condition, dominant soil type and hydrological regime.

| Watershed classification | Pairs | Sf_d | | Sf_w | |
|--------------------------|---------------------|--------------|-------------------|--------------|-------------------|
| | | Z | p | Z | p |
| Climate condition | EL-EQ | -2.14 | 0.03* | -3.98 | <0.001* |
| | EL-WL | -3.09 | <0.002* | -3.15 | <0.002* |
| | EQ-WL | -1.41 | 0.16 | 2.20 | 0.03* |
| Dominant soil type | LIXISOLS-LUVISOLS | -3.70 | <0.001* | -2.19 | 0.028* |
| | LIXISOLS-LEPTOSOLS | -1.79 | 0.074* | -2.93 | 0.003* |
| | LIXISOLS-CAMBISOLS | -2.95 | 0.003* | -4.62 | <0.001* |
| | LUVISOLS-LEPTOSOLS | 3.53 | <0.001* | 0.02 | 0.98 |
| | LUVISOLS-CAMBISOLS | 1.88 | 0.059* | -0.80 | 0.42 |
| Hydrological regime | LEPTOSOLS-CAMBISOLS | -2.20 | 0.027* | -1.42 | 0.15 |
| | RD-Hybrid | 1.97 | 0.05* | -0.26 | 0.79 |

770 *Note:* Sf_d and Sf_w are dry season and wet season ecohydrological sensitivity, respectively; EL, EQ and WL refer to energy-limited, equitant and water-limited watersheds, respectively; RD is the rain-dominated watershed.

The bolded number with * indicates statistically significant at $p < 0.10$.

Table 6. Correlation analysis between seasonal ecohydrological sensitivities and contributing factors.

| | Variables | Sfd | | | Sfw | | |
|------------|---------------------------------|---------------|-----------------------|-------------|--------------|-----------------------|-------------|
| | | Kendall | a | R^2 | Kendall | a | R^2 |
| Climate | DI | 0.44* | 2.37* | 0.41 | 0.19 | 0.51 | 0.05 |
| | P_e | -0.23 | -0.01 | 0.09 | -0.32 | -0.03* | 0.23 |
| Vegetation | Vegetation coverage | -0.51* | -0.08* | 0.53 | 0.08 | 0.01 | 0.01 |
| | Forest coverage | -0.36* | -0.03 | 0.13 | 0.21 | 0.01 | 0.06 |
| | LAI | -0.44* | -1.62* | 0.05 | 0.09 | -0.08 | 0.00 |
| Topography | Area ^c | 0.15 | 0.28 | 0.01 | 0.19 | 0.28 | 0.02 |
| | Perimeter ^c | 0.23 | 1.75 | 0.07 | 0.25 | 1.08 | 0.15 |
| | Elevation ^c | 0.00 | -0.10 | 0.00 | 0.12 | 0.23 | 0.03 |
| | Δ Elevation ^c | 0.10 | 0.71 | 0.05 | 0.27 | 0.5 | 0.06 |
| | Slope | -0.39* | -0.15* | 0.28 | -0.03 | -0.01 | 0.00 |
| | LS | -0.40* | -0.20* | 0.24 | 0.04 | 0.48 | 0.01 |
| | Length | -0.18 | -4.3×10^{-3} | 0.10 | 0.21 | -1.2×10^{-2} | 0.03 |
| | Max Length | -0.23 | -1.9×10^{-3} | 0.15 | 0.32 | -1.4×10^{-3} | 0.10 |
| | TWI | 0.62* | 4.30* | 0.51 | 0.19 | 1.05 | 0.15 |
| | CON | 0.12 | 0.04 | 0.04 | 0.20 | 0.05* | 0.20 |
| | DDG | 0.49* | 0.10* | 0.45 | 0.03 | 0.02 | 0.05 |
| | SA | -0.13 | 1.5×10^{-3} | 0.00 | 0.14 | 4.2×10^{-3} | 0.15 |
| | TPI | -0.04 | 3.45 | 0.00 | -0.05 | 8.86 | 0.03 |
| | TRI | -0.33 | -0.32* | 0.23 | 0.01 | 0.02 | 0.00 |
| | Positive Openness | 0.36* | 14.23* | 0.26 | 0.08 | 0.63 | 0.00 |
| | Negative Openness | 0.34 | 14.78* | 0.25 | 0.03 | 0.43 | 0.00 |
| | Depth | -0.31 | -0.01* | 0.32 | -0.10 | -0.01 | 0.01 |
| Soil | T_w | 0.25 | 74.26 | 0.05 | 0.53* | 125.46* | 0.38 |
| | T_{hy} | -0.03 | 0.04 | 0.01 | 0.41* | 0.15* | 0.25 |
| | T_d | 0.28 | 32.32* | 0.28 | 0.10 | 2.49 | 0.01 |
| | T_{oc} | -0.21 | -3.99* | 0.27 | -0.11 | -0.29 | 0.00 |
| | T_{ece} | 0.39* | 10.74* | 0.28 | 0.30 | 3.99 | 0.19 |
| | S_w | -0.09 | -17.10 | 0.03 | 0.06 | -3.80 | 0.01 |
| | S_{hy} | 0.15 | 0.28 | 0.07 | 0.30 | 0.30* | 0.22 |
| | S_d | 0.00 | 13.30 | 0.06 | 0.15 | 8.66 | 0.08 |
| | S_{oc} | 0.17 | 3.80 | 0.03 | -0.09 | 1.76 | 0.02 |
| | S_{ece} | 0.34 | 7.71 | 0.16 | 0.28 | 3.87* | 0.22 |
| | Soil types | -0.30 | -0.11 | 0.14 | 0.37* | 0.06 | 0.13 |
| Landscape | PN | -0.18 | -5.2×10^{-4} | 0.02 | 0.01 | -4.2×10^{-5} | 0.00 |
| | PD | -0.54* | -10.83* | 0.30 | -0.25 | -4.73 | 0.15 |
| | LPI | 0.08 | 0.02 | 0.04 | 0.06 | 0.04 | 0.00 |
| | ED | -0.36* | -0.27 | 0.17 | -0.32 | -0.19* | 0.23 |
| | CONTAG | 0.03 | 0.02 | 0.01 | 0.10 | 0.03 | 0.09 |
| | SHDI | -0.05 | -0.31 | 0.00 | -0.06 | -0.66 | 0.03 |
| | SIDI | -0.08 | -0.74 | 0.00 | -0.03 | -0.69 | 0.01 |

775 Note: Linear regressions are built as $y=ax+b$, where a is the slope of the linear regression; c means parameters are transferred into $\ln()$ format.

The bolded number with * indicates statistically significant at $p < 0.10$.

Table 7. Selected factor analysis models.

| | Influencing factors | MSA | KMO | Bartlett's test |
|------------|---|-------------|-------|-----------------|
| Dry season | DI, slope, LS, TWI, DDG, TRI, Depth, NO | ≥ 0.53 | 0.730 | 0.000 |
| Wet season | P_e , CON, T_w , T_{hy} , S_{hy} , ED | ≥ 0.57 | 0.634 | 0.000 |