

Variations in surface roughness of inhomogeneous underlying surfaces in the Nagqu area of the Tibetan Plateau

Maoshan Li^{*1}, Xiaoran Liu², Lei Shu¹, Lingzhi Wang¹, Wei Fu¹, Yaoming Ma³, Yaoxian Yang⁴, Fanglin Sun⁴

(1.School of Atmospheric Sciences/Plateau Atmosphere and Environment Key Laboratory of Sichuan Province/Joint Laboratory of Climate and Environment Change, Chengdu University of Information Technology, Chengdu 610225, Sichuan China;

2. Climate Center, Meteorological Bureau of Inner Mongolia Autonomous Region, Huhhot 010051, Inner Mongolia Autonomous Region, China)

3.Key Laboratory of Tibetan Environment Changes and Land Surface Processes, Institute of Tibetan Plateau Research, Chinese Academy of Sciences, CAS Center for Excellence in Tibetan Plateau Earth Sciences, Beijing, China;

3. Key Laboratory of Land Surface Process and Climate Change in Cold and Arid Regions, Chinese Academy of Sciences. Lanzhou, China)

Abstract: The temporal and spatial variation characteristics of the surface roughness in the Nagqu area of the northern Tibetan Plateau were analysed in 2008, 2010 and 2012 using MODIS satellite data and station atmospheric turbulence observation data, and the **Massman retrieved model** and measured average wind speed and turbulent flux of a single height ultrasonic anemometer were used to determine the aerodynamic surface roughness. The results showed that the surface roughness length has obvious seasonal variation characteristics. From February to August, Z0m increased constantly with snow ablation and vegetation growth, and the maximum value reached 4-5 cm at the BJ site. From September to February, Z0m gradually decreased with the post-monsoon phase over the plateau, and the values decreased to approximately 1-2 cm. Snowfall in abnormal years was the main reason for the obviously lower Z0m compared with that in normal conditions. The underlying surface can be divided into four categories according to the different

* Corresponding author

Dr. Maoshan Li

Chengdu University of Information Technology

24 Block 1, Xuefu Road, Chengdu 610225, Sichuan, China

E-mail: lims@cuit.edu.cn

26 values of Z0m: snow and ice, sparse grassland, lush grassland and town. Among them, lush
27 grassland and sparse grassland accounted for 62.49% and 33.74% respectively, and their Z0m
28 annual changes are between 2-6 cm and 1-4 cm. The two methods were positively correlated with
29 each other, and **the retrieved data values** were lower than the measured results due to
30 non-uniformity of underlying surface. These results are substituted into Noah-MP **replaces the**
31 **original parameter design** numerical simulation experiment. After replacing the model surface
32 roughness, the sensible heat flux and **latent** heat flux **has a better daily improvement effect.**

33 **Key words:** Northern Tibetan Plateau; Surface roughness; NDVI; MODIS

34

35 **1 Introduction**

36 Known as the "third pole" of Earth (Jane, 2008), the Tibetan Plateau has an average altitude
37 of over 4000 m and accounts for a quarter of China's territory. It is located in southwestern China
38 adjacent to the subtropical tropics in the south, and it reaches the mid-latitudes in the north,
39 making it the highest plateau in the world. Due to its special geographical location and
40 geomorphic characteristics, it plays an important role in the global climate system as well as the
41 formation, outbreak, duration and intensity of the Asian monsoon (Yang et al., 1998; Zhang et al.,
42 1998; Wu et al., 1999, 2004, 2005; Ye et al, 1998; Wu et al, 1998; Tao et al, 1998). Many studies
43 (Wu et al., 2013; Wang, 1999; Ma et al, 2002) have shown that the land-atmosphere interaction on
44 the Tibetan Plateau plays an important role in the regional and global climate. Over the past 47
45 years, the Tibetan Plateau has shown a significant warming trend and increased precipitation (Li et
46 al., 2010). The thermal effects of the Tibetan Plateau not only have an important impact on the
47 Asian monsoon and precipitation variability but also affect the atmospheric circulation and climate
48 in North America, Europe and the South Indian Ocean by **inspiring** large-scale teleconnections
49 similar to the Asia-Pacific Oscillation (Zhou et al., 2009).

50 The various thermal and dynamic effects of the Tibetan Plateau on the atmosphere **affect** the
51 free atmosphere via the atmospheric boundary layer. Therefore, it is particularly important to
52 analyse the micrometeorological characteristics of the atmospheric boundary layer of the Tibetan
53 Plateau, especially the near-surface layer (Li et al., 2000). Affected by the unique underlying

54 surface conditions of the Tibetan Plateau, local heating shows interannual and interdecadal
55 variability (Zhou et al., 2009). Different underlying surfaces have differing diversities, complex
56 compositions and uneven distributions and contribute to diverse and complex **land surfaces**. As the
57 main input ~~factor~~ for atmospheric energy, the surface greatly affects the various interactions
58 between the ground and the atmosphere and even plays a key role in local areas or specific times
59 (Guan et al, 2009). The surface characteristic parameters (dynamic roughness, thermodynamic
60 roughness, etc.) play an important role in the land surface process and are important factors in
61 causing climate change (Jia et al., 2000). The underlying surface of the Tibetan Plateau presents
62 different degrees of fluctuation, which introduces certain obstacles to understanding the
63 land-atmosphere interaction of the Tibetan Plateau. The fluctuating surface may alter the
64 arrangement of roughness elements on the surface and cause changes in surface roughness.
65 Changes in roughness can also affect changes in the characteristics of other surface turbulent
66 transportation, which may also result in changes in surface fluxes. Chen et al. (2015) presented a
67 practical approach for determining the aerodynamic roughness length at fine temporal and spatial
68 resolutions over the landscape by combining remote sensing and ground measurements. Surface
69 roughness is an important parameter in land surface models and climate models. Its size **reflects**
70 the ~~matter-energy~~ exchange, transmission intensity and interactions between the near surface
71 airflow and the underlying surface to some extent (Liu et al., 2007; Irannejad et al, 1998; Shao et
72 al, 2000; Zhang et al., 2003). Zhou et al. (2012) demonstrated that simulated sensible heat flux
73 compared with measurement was significantly improved by using a time-dependent Z0m
74 parameter. Therefore, the primary objective of this study is to calculate the surface roughness and
75 its variation characteristics to further understand the land-atmosphere interactions on the central
76 Tibetan Plateau.

77 Through the study of surface roughness, it is beneficial to obtain the surface feature
78 parameter values in the region, provide the ground truth value for model inputs, improve the
79 simulation level of the model in the Tibetan Plateau, and deepen the understanding of the
80 land-atmosphere interaction process. Research on model simulations of surface flux has achieved
81 good results in many regions (Smirnova et al., 2016). Especially with the continuous development

82 and improvement of numerical models in recent years, research on the applicability of different
83 parameterization schemes in different models to different regions has continued. Luo et al. used
84 the land surface model CoLM to conduct a single-point numerical simulation of the BJ station and
85 successfully simulated the energy exchange process in the Nagqu area (Luo et al., 2009). Zhang et
86 al. evaluated the surface physical process parameterization schemes of the Noah LSM and
87 Noah-MP models in the entire East Asia region and evaluated the simulation of the surface heat
88 flux of the Tibetan Plateau (Zhang et al., 2017). Xie et al. explored the simulation effect of the
89 land surface model CLM4.5 in the alpine meadow area of the Qinghai-Tibet Plateau (Xie et al.,
90 2017). Xu et al. studied the applicability of different parameterization schemes in the WRF model
91 when simulating boundary layer characteristics in the Nagqu area (Xu, et al., 2018). Zhang, et al.
92 Comparative analyses have been performed of the meteorological elements simulated by different
93 land surface process schemes in the WRF model in the Yellow River source region (Zhang et al.,
94 2019). However, the applicability of the model in the Tibetan Plateau needs further study. The
95 terrain of the Tibetan Plateau is complex, the underlying surface is very uneven, and the area has
96 high spatial heterogeneity. Because the condition of the underlying surface has a very significant
97 impact on the surface flux, obtaining information on the surface vegetation status of a certain area
98 is very helpful for analysing the spatial representation of the surface flux.

99 In this study, satellite data are obtained by MODIS and the normalized difference vegetation
100 index (NDVI) in the Nagqu area is used to study the dynamic surface roughness length.
101 Atmospheric turbulence observation data from 2008, 2010, and 2012 and observation data from
102 automatic weather stations were collected for three observation stations in the region, and the
103 measured values of the average wind speed and turbulent flux of a single height ultrasonic
104 anemometer were used to determine the surface dynamic roughness Z_{0m} (Chen et al., 1993). The
105 time-scale dynamics of Z_{0m} and the results of different underlying surfaces were analysed.
106 Through a comparison of the calculation results of the observation data, we studied whether the
107 surface roughness values retrieved by satellites are reliable to provide accurate surface
108 characteristic parameters. Then we used the retrieved surface roughness to replace the surface
109 roughness in the original model for numerical simulation experiments, and evaluated the model

110 simulation results. These researches **will helpful** for the study of land-atmosphere interactions in
111 the plateau area and improve the theoretical research level of the near-surface layer on the Tibetan
112 Plateau. In the following section, we describe the case study area, the MODIS remote sensing data,
113 the ground observations, and the land cover map used to drive the revised Massman model
114 (Massman et al, 1997, 1999). In Section 3, we present the results and then a validation based on
115 flux measurements at Nagqu station. Finally, we provide some concluding remarks on the
116 variation characteristics of aerodynamic roughness lengths and numerical simulation of the
117 surface turbulent flux in the Nagqu area of the central Tibetan Plateau.

118

119 **2 Study area, Data and methods**

120 **2.1 Study area and Data**

121 The area selected in this study is a 200×200 km² area centred on the Nagqu Station of Plateau
122 Climate and Environment of the Northwest Institute of Ecology and Environmental Resources,
123 Chinese Academy of Sciences, Northern Tibet Plateau.

124 In this area, three meteorological observatory stations are located: North Pam (Portable
125 Automated Meso-net) Automatic Meteorological Observatory (NPAM), Nam Co Station for
126 Multisphere Observation and Research, Chinese Academy of Sciences (NAMC), and BJ station
127 (Figure 1). The underlying surface around the observation point is relatively flat on a small spatial
128 scale, and a certain undulation is observed at a large spatial scale. The data used included
129 observations from atmospheric turbulence and automatic meteorological stations.

130 The BJ station is located at coordinates 31.37°N, 91.90°E and an altitude of 4509 m a.s.l. The
131 BJ observation point is located in the seasonal frozen soil area, and the vegetation is alpine
132 grassland. The site measurement equipment includes an ultrasonic anemometer (CAST3,
133 Campbell, Inc.), CO₂/H₂O infrared open **circuit** analyser (LI 7500) and automatic meteorological
134 observation system (Ma et al., 2006). This study uses the BJ station data from 2008 and 2012. The
135 NPAM station is located at 31°56'N, 91°43'E and has an altitude of approximately 4700 m. The
136 ground of the experimental field is flat, and the area is wide. The ground is covered by a plateau
137 meadow that grows 15 cm high in summer. The experimental station observation equipment

138 includes an ultrasonic wind thermometer and humidity probe pulsator and includes data on
139 temperature and humidity, air pressure, average wind speed, average wind direction, surface
140 radiation temperature, soil heat flux, soil moisture and temperature, and radiation (Ma et al., 2006).
141 The NAMC station is located at 30°46.44'N, 90°59.31'E and an altitude of 4730 m. It is located on
142 the southeastern shore of Namco Lake in Namuqin Township, Dangxiong County, Tibet
143 Autonomous Region. It is backed by the Nyainqentanglha Mountain Range, and the underlying
144 surface is an alpine meadow. This study uses NPAM station data for the whole year of 2012 and
145 NAMC station data for the whole year of 2010.

146 **Figure 1 about here**

147 **Table 1 about here**

148 The land cover data used in this study are GLC2009 (Arino et al, 2010) data from the Envisat
149 satellite in 2009, and the spatial resolution is 300 m. The classification standard is the land cover
150 classification system (LCCS), and it divides the global surface into 23 different types, with the
151 study area including 14 of these types. The actual situation in the selected area does not match the
152 data part of GLC2009 because of the lack of an underlying surface, such as farmland, in the
153 selected area. Therefore, according to the actual land cover types obtained by Chu (Chu, 2010),
154 the categories irrigated farmland, dry farmland, mixed farmland vegetation, mixed multivegetation
155 land, closed grassland and open grassland are replaced with 6-grasslands, shrub meadows,
156 mountain meadows, alpine grasslands, alpine meadows and sparse vegetation in the mountains.
157 Since the proportion of the underlying surface of the tree as a whole is only 0.36%, the underlying
158 surface types evergreen coniferous forest, the mixed forest, the multiforest grassland mix and the
159 multigrass forestland mix will no longer be studied.

160 The MODerate-resolution Imaging Spectroradiometer (MODIS) is an important sensor on
161 board the satellites TERRA and AQUA launched by the US Earth Observing System Program. The
162 band of the MODIS sensor covers the full spectrum from visible light to thermal infrared; thus,
163 this sensor can detect surface and atmospheric conditions, such as surface temperature, surface
164 vegetation cover, atmospheric precipitation, cloud top temperature, etc. The maximum spatial
165 resolution is 250 m. The normalized vegetation index obtained by MODIS is the MYD13Q1

166 product, which provides a global resolution of 250 m per 16 days. This study selects 73 materials
 167 for 2008, 2010 and 2012 in Nagqu.

168 2.2 Methodology

169 2.2.1 Method for calculating surface roughness by observation data

170 Using the measured values of the average wind speed and turbulent flux of a single height
 171 ultrasonic anemometer, the calculation scheme of surface roughness proposed by Chen et al.
 172 (Chen et al., 1993) was selected and the dynamic variation in the surface roughness was obtained.

173 According to the Monin-Obukhov similarity theory (Monin et al., 1954), the wind profile
 174 formula with the stratification stability correction function (Panosky et al, 1984) is as follows:

$$175 \quad U(z) = \frac{u_*}{k} \left[\ln \frac{z-d}{Z_{0m}} - \psi_m(\zeta) \right] \quad (1)$$

$$176 \quad \psi_m(\zeta) = 2 \ln\left(\frac{1+x}{2}\right) + \ln\left(\frac{1+x^2}{2}\right) - \tan^{-1}(x) + \frac{\pi}{2} \quad \zeta < 0 \quad (2)$$

$$177 \quad \psi_m(\zeta) = -5\zeta \quad \zeta > 0 \quad (3)$$

178 where $u_* = \sqrt{-\overline{u'w'}}$; Z_{0m} is the dynamic surface roughness length; z is the observed height;
 179 d is zero plane displacement, which is set to 0.03 m and calculated from the average vegetation
 180 height 0.045 m (Stanhill, 1969); U is the average wind speed; k is the Karman constant, which is
 181 set to 0.40 (Högström,1996); $L = -\frac{u_*^3}{(k \frac{g}{\theta}) \overline{\theta'w'}}$ is the Monin-Obukhov length (Monin et al.,
 182 1954); $x = (1-16\zeta)^{1/4}$; and $\zeta = (z-d)/L$ is the atmospheric stability parameter. Available

183 from formula (2.1):

184

$$185 \quad \ln \frac{z-d}{Z_{0m}} = \frac{kU}{u_*} + \psi_m(\zeta) \quad (4)$$

186 Using equations (2)~(4), Z_{0m} can be determined by fitting ζ and observing a single height

$$187 \quad \frac{kU}{u_*}.$$

188 2.2.2 Method for calculating surface roughness by satellite data

189 For a fully covered uniform canopy, Brutsaert suggested that $z_0 m = 0.13 h v$ (Brutsaert, 1982).
190 For a canopy with proportional coverage (partial coverage), Raupach (Raupach, 1994) indicated
191 that $z_0 m/hv$ varies with the leaf area index (LAI). However, Pierce et al. (Pierce et al. 1992)
192 pointed out that for all kinds of biological groups, the leaf area index can be obtained from the
193 NDVI and the **density grade** of vegetation can be related to the NDVI. Asrar et al. (Asrar et al.,
194 1992) pointed out that a mutual relationship occurred among the LAI, NDVI and ground cover
195 through the study of physical models. Moran's study (Moran, et al., 1994) provides another
196 method that uses the function of the relationship between NDVI and Z_{0m} in the growing season of
197 alfalfa.

198 Considering that the main underlying surface of the study area is grassland, this study selects
199 the Massman model (Massman et al, 1997, 1999) to calculate the Z_{0m} in the Nagqu area of the
200 central Tibetan Plateau. The Massman model is calculated as follows:

$$201 \quad \gamma = C_1 - C_2 \cdot \exp(-C_3 \cdot C_d \cdot LAI) \quad (5)$$

$$202 \quad n_{ec} = \frac{C_d \cdot LAI}{2 \cdot \gamma^2} \quad (6)$$

$$203 \quad d_h = 1 - \frac{[1 - \exp(-2 \cdot n_{ec})]}{2 \cdot n_{ec}} \quad (7)$$

$$204 \quad \frac{Z_{0m}}{h} = [1 - d_h] \cdot \exp(-\frac{k}{\gamma}) \quad (8)$$

205 where $C_1=0.32$, $C_2=0.26$, and $C_3=15.1$ are constants in the model and related to the surface
206 drag coefficient; LAI is the leaf area index; $C_d=0.2$ is the drag coefficient of the foliage elements;
207 n_{ec} is the wind speed profile coefficient of fluctuation in the vegetation canopy; and h is the
208 vegetation height. In many earlier studies, the high-altitude environment of the Tibetan Plateau
209 was correlated with a low temperature in the study area and shown to affect the height and
210 sparseness of the vegetation. Based on previous research, this study considers that the vegetation
211 height in northern Tibet is related to the normalized **differential** vegetation index (NDVI) and
212 altitude (Chen et al, 2013) and introduces the altitude correction factor on the original basis. The
213 following is the calculation formula:

214
$$H = h_{\min} + \left(\frac{h_{\max} - h_{\min}}{NDVI_{\max} - NDVI_{\min}} \right) (NDVI - NDVI_{\min}) \quad (10)$$

215
$$h = acf \cdot H \quad (11)$$

216 where h_{\min} and h_{\max} are the minimum and maximum vegetation height observed at the
 217 observation station, respectively, $NDVI_{\max}$ and $NDVI_{\min}$ are the maximum and minimum NDVI of
 218 the observation station, respectively; H is based on the assumption that the vegetation height is
 219 directly proportional to the NDVI; x is the altitude, which is obtained from ASTER's DEM
 220 products; and acf is the altitude correction factor (Chen et al., 2013), which is used to characterize
 221 the effect of elevation on the height of vegetation in northern Tibet. The acf parameter has the
 222 following form:

223
$$acf = \begin{cases} 0.149, x > 4800 \\ 11.809 - 0.0024 \cdot x, 4300 < x < 4800 \\ 1.49, x < 4300 \end{cases} \quad (12)$$

224 The LAI used in this study is calculated by the NDVI of MODIS (Su, 1996). The calculation
 225 formula is as follows:

226
$$LAI = \left(\frac{NDVI * (1 + NDVI)}{1 - NDVI} \right)^{0.5} \quad (13)$$

227 **3 Results analysis**

228 **3.1 Variation characteristics of surface roughness based on measured data**

229 Figure 2 shows the temporal variation characteristics of the surface roughness of sites in
 230 different years in the Nagqu area. The Z0m value has continued to increase since February to
 231 reach a maximum in July and August. The results for the BJ and NPAM stations in 2012 show that
 232 July has slightly larger values than August, and the results for the NAMCO station in 2010 and BJ
 233 station in 2008 show that August has larger than July. ~~The BJ station may have experienced a~~
 234 ~~precipitation process in July 2008, resulting in a July Z0m value less than June.~~ After August, the
 235 Z0m value began to decrease, and in December, the value was approximately the same as the
 236 value in January. In general, the change in the Z0m degree of each station increases from spring to
 237 summer and decreases month by month from summer to winter.

238 **Figure 2 about here**

239

240 **3.2 Spatiotemporal variation characteristics of surface roughness length**
241 **retrieved by MODIS data**

242 Figure 3 shows a plot of the surface roughness distribution of 200×200 km² around the BJ
243 site in 2008. In February, the Z0m decreased from January, which may have been due to snowfall,
244 temperature, etc., resulting in a small Z0m that continued to decrease. Due to the rising
245 temperature and snow melting, Z0m showed a slowly increasing trend from February to May and
246 a rapid increase from June to August. From June onwards, a large number of surface textures were
247 observed, indicating the complexity of the underlying surface. Whether the bulk surface or
248 vegetation had a more important impact on Z0m is not clear. From May to August, obvious
249 changes in humidity, temperature and pressure caused by the plateau summer monsoon led to an
250 increase in the height and coverage of surface vegetation, and Z0m peaked in August. In particular,
251 the change from May to June was very significant, which may have been related to the beginning
252 of the summer monsoon in June, the corresponding increase in precipitation, **the** accelerated the
253 growth of vegetation and the rapid rise of Z0m. In June, July, and August, continuous precipitation
254 and rising temperatures led to vigorous vegetation growth, although changes were not observed
255 after the vegetation reached maturity. The corresponding maximum value of Z0m in the figure
256 remains unchanged, although due to **sufficient** values in these three months, the area with high
257 Z0m values gradually expanded and reached the maximum range in August. From September to
258 December, as the plateau summer monsoon retreated, the temperature and humidity gradually
259 decreased. Compared with the plateau summer monsoon, the conditions were no longer suitable
260 for vegetation growth; thus, the contribution of vegetation to Z0m was weakened, the surface
261 **vegetation height gradually decreased, and Z0m continued to decrease. Moreover, the range of**
262 **large-value regions** also gradually decreased.

263 **Figure 3 about here**

264 **Figure 4 about here**

265 Figures 4 and 5 show the retrieved monthly surface roughness values in the BJ area in 2010
266 and 2012, respectively. Moreover, Z0m also showed a decrease from January to February in the

267 Nagqu area in 2010 and 2012. Starting in February, Z0m increased. Starting in June, Z0m
268 increased rapidly and reached the peak of the whole year in August. Subsequently, Z0m began to
269 decrease.

270 **Figure 5 about here**

271 Figures 3, 4, and 5 show that Z0m changes with the spatial and temporal scale. Z0m shows
272 different trends on different underlying surfaces. In November 2008, the Z0m in the Nagqu area
273 was small overall and generally as low as 1 cm. Historical data show that there is a large-scale
274 snowfall process in the Nagqu area at this time. Snowfall over the meadow causes the underlying
275 surface of the meadow to be homogeneous and flat, and after the snowfall falls, it is easy to form a
276 block with a scattered and discontinuous underlying surface. We subsequently determined that the
277 surface roughness of the area with ice and snow as the underlying surface is not more than 1 cm,
278 which is consistent with historical weather processes. Therefore, we think that snowfall caused the
279 Z0m in November to be very small. From November to December, Z0m showed a growing trend,
280 which may be due to temperature, unfrozen soil or other reasons that resulted in the melting of
281 snow, and then the surface roughness showed a growing trend (Zhou, 2017).

282 **3.3 Evaluation of satellite data retrieved results**

283 **Figure 6 about here**

284 The underlying surfaces of the three sites selected in this study are all alpine meadows. In
285 Figure 6, the NPAM site data calculation results are larger than the satellite data retrieved results
286 throughout the year. Only the values in September and October are very close, and the trends are
287 similar. The maximum value of the site data calculation is 5 cm and the satellite data retrieved
288 result is 4.5 cm. The maximum difference is in May at 1.7 cm. The NAMCO station data
289 calculation results are very close to the satellite data retrieved results from April to November,
290 although the satellite data retrieved results are significantly larger than the site data calculation
291 results in January, March and December. The largest difference occurs in January, and the
292 difference value reaches 1.5 cm. In 2008, the calculation results of the BJ station data were larger
293 than the satellite data retrieved results throughout the year. The calculation results of the site data
294 were very close to the satellite data retrieved results from January to April and July to November,
295 although a large difference was observed in May, June and December, with the largest difference
296 occurring in May at 1.8 cm. In 2012, the BJ site data calculation results were consistent with the

297 satellite data retrieved results for the whole year, although the site data calculation results were
298 larger than the satellite data retrieved results from March to June, and the station data calculation
299 results were smaller than the satellite data retrieved at other times. As a result, the largest
300 difference occurred in June at 1.1 cm. Figure 6 shows that for the overall situation, the seasonal
301 variation trend of the site data calculation results is consistent with the satellite data retrieved
302 results in January, February, March, November and December. However, the site data calculation
303 results from April to October are greater than the satellite data retrieved results. From Figure 6, the
304 Z0m calculated from the site observation data is larger than that of the satellite data, which may be
305 because of the average smoothing effect. From February to July, the single point Z0m value was
306 significantly increased according to the independent method of determining the surface roughness,
307 while the results obtained by using the satellite data did not increase significantly. The satellite
308 results show that the values from January to May, November, and December are basically stable
309 below 2 cm and only change from June to October, which is related to non-uniformity of
310 underlying surface in Massman **mode**. In general, the results calculated by the station are generally
311 larger than those obtained by satellite retrieval.

312 **Figure 7 about here**

313 The Z0m scatter plot is shown in Figure 7. A significant positive correlation is observed
314 between the satellite retrieval and the surface roughness calculated from the site data. The
315 correlation coefficients between the observation result and the retrieved result are large except for
316 at the NAMCO station in 2010 in Figure 7(g). The average result of the underlying surface were
317 consistent with the underlying surface results in different regions, further indicating that the
318 satellite retrieved results are consistent with the site calculation results. However, the results of the
319 NAMCO site are different from those of the other sites. The correlation coefficient with the
320 average results of the underlying surface is 0.83, and the correlation coefficient with the satellite
321 retrieved results is 0.62. Because the Namco Observation Station is closer to the lake (1 km), it is
322 more affected by local microclimates, such as lake and land winds. The results in Figure 7 all
323 passed the F test of $P=0.05$, which indicates that there is no significant difference between the site
324 data calculation results and the satellite data retrieved results.

325 **4 Variation characteristics of the surface roughness of different** 326 **underlying surfaces**

327 According to the vegetation dataset GLC2009 combined with actual local conditions, the
328 200×200 km² area of Nagqu was divided into 10 different underlying surfaces (~~Arino et al., 2010~~):
329 mountain grassland, shrub meadow, mountain meadow, alpine grasslands, alpine meadows, sparse
330 vegetation lap, urban land, bare land, water bodies, ice sheets and snow cover.

331

332

Figure 8 about here

333 The monthly variation in Z0m in different underlying surfaces in the Nagqu area is shown in
334 Figure 8, which indicates that 14 different underlying surfaces in the Nagqu area can be divided
335 into four categories. The first category is urban land, which accounts for 0.07% of the whole study
336 area. The Z0m of this type of underlying surface is greater than that of other types throughout the
337 year, and the change in Z0m is very large, which is probably due to the irregular changes in the
338 underground areas of the selected areas and the irregularities caused by human activities. The
339 second category is lush grassland, including shrub meadows, mountain grasslands, alpine
340 grasslands and mountain meadows, which account for 62.49% of the area. The variation curves of
341 Z0m of the four underlying surfaces are similar, and the Z0m of the urban land is only smaller
342 than that of other underlying surfaces. The third category is sparse grassland, including alpine
343 sparse vegetation, alpine meadows and bare land, and it accounts for 33.74% of the area. The Z0m
344 values of the three underlying surfaces are similar at a medium height. The Z0m of the bare soil is
345 at the lowest point of these underlying surface Z0m, and the Z0m of the alpine meadow is
346 relatively stable and less affected **by the outside**. The fourth category is ice and snow, including ice
347 **sheets** and snow cover, and water bodies are two kinds of underlying surfaces, accounting for
348 3.7% of the area. The Z0m of these three underlying surfaces presents another phenomenon. The
349 variation range of the whole year is relatively small, and the Z0m of these underlying surfaces is
350 also small. It is more than 1 cm in mid-June and less than 1 cm at other times. Figure 8(d) shows
351 the multiyear average seasonal variation in Z0m. The figure clearly shows that the underlying
352 surface can be divided into four categories due to the difference in surface roughness. The change
353 from January to May Z0m is very small, peaking from May to August and then down to the

354 previous January to May level in November and December. The snowfall in November 2008 may
355 have led to the low level of November in Figure 8(d). Table 2 shows that the winter albedo at the
356 BJ station and Namco station is higher than that in other seasons, and the summer is the smallest.
357 Both show that the surface albedo in November 2008 was significantly higher than that in
358 November of the other two years. In fact, the surface roughness on November should be higher
359 than that on December in former years.

360

Table 2 about here

361 Figure 8 also shows that in the Nagqu area, except for the area of the fourth type of
362 underlying surface, the Z0m change in other areas decreases from January to February and begins
363 to increase after February, reaching a peak in August and then starting to decrease. However,
364 Figure 6 clearly shows that there are several stages in which Z0m changes significantly, in early
365 April, mid-May, early July, late August, and late September. The change at the end of August was
366 the most obvious. Each of the underlying surfaces Z0m changes by more than 2 cm on average.
367 The extent of the change in late September was also large, with an average change of more than
368 1.5 cm. Moreover, the change in early July was special because the change resulted in a significant
369 increase in the Z0m of water bodies and ice.

370 Certain factors, such as cloud cover in May, August, and November of 2008; August and
371 September of 2010; and April and July of 2012, caused significant changes in the overall Z0,
372 which resulted in two very significant changes in the three-year average for August and November.
373 In November, the change was caused by snowfall based on other meteorological data. In August
374 2008 and 2010, the changes was caused by precipitation based on an analysis of the sudden
375 increase in the Z0m of the water body and ice and snow surface. Combined with several changes
376 in Z0m, precipitation, snowfall, and snow accumulation will make the underlying surface more
377 uniform and flatter, which will lead to relative reductions in Z0m.

378 5 Simulation and evaluation of Noah-MP model

379 5.1 Model setup

380 According to the surface roughness variation characteristics retrieved from
381 satellite data, the underlying surface of Nagqu area can be divided into four types.

382 They are urban, lush grass, sparse grass, and ice and snow. Among them, urban
383 accounts for 0.07%, its Z0m up to 9 cm, lush grassland accounts for 62.49% of the
384 area, its Z0m can reach up to 6cm, sparse grassland up to 33.74%, its Z0m can reach
385 up to about 4cm, ice and snow accounts for 3.7% of the area, and Z0m does not
386 exceed 1cm. These results are substituted into Noah—MP **replaces** the original
387 parameter design numerical simulation experiment. The model after replacing the
388 surface roughness is set as a sensitivity experiment, and the original model is set as a
389 control experiment. The selection of other parameterization schemes suitable for
390 numerical simulation in Nagqu area is shown in Table 3. The simulation time is from
391 July 1 to 31, 2008, and the spin-up time is 9 days. The forcing field dataset is a
392 Chinese meteorological forcing dataset jointly developed by the Tibetan Plateau Data
393 Assimilation and Modelling Centre and the Institute of Tibetan Plateau Research of
394 the Chinese Academy of Sciences (ITPCAS).

395 **5.2 Evaluation of the simulated single point heat flux**

396 Figure 9(a) shows that the sensible heat flux simulated by the sensitivity
397 experiment is closer to the measured value than the control experiment. In the daytime,
398 the results of sensitivity experiment **were** smaller than those of the control experiment,
399 ~~and some time were larger than those of the control experiment.~~ At night, the results
400 of the two models were close to each other before July 21, and the sensitivity
401 experiment results were significantly improved after July 21. Figure 9(b) shows that
402 the sensitivity experiment results are basically consistent with the control experiment
403 results, which are maintained at about $0\text{W}/\text{m}^2$, and there is no improvement at night.
404 Before July 19, the **two experiments latent heat fluxes** remained at about $200\text{W}/\text{m}^2$,
405 which was less than the observed latent heat flux, and the simulated maximum value
406 of the sensitivity experiment was greater than that of the control experiment. After
407 July 19, the two experiments simulation results began to increase and reached about
408 $400\text{W}/\text{m}^2$ consistent with the observed latent heat flux, indicating that the simulation
409 effect was improved to some extent. Similarly, it can be found from the maximum

410 value that the sensitivity experiment results were slightly greater than the control
411 experiment results. It also shows that improving the accuracy of surface roughness
412 can improve the simulation effect of latent heat flux.

413 **Figure 9 about here**

414 **5.3 Evaluation of regional heat flux simulation results**

415 In order to compare the changes of sensible heat flux and latent heat flux before
416 and after improvement, the sensitivity simulations are used to subtract the control
417 model simulation results. By subtracting the sensible heat flux **from** control **and**
418 sensitivity experiment, the results are shown in the figure 10. It can also be seen from
419 the figure 10 (a) that the difference of sensible heat flux is basically negative in the
420 daytime, indicating that the sensible heat flux after improvement is smaller than that
421 before improvement. The above results show that the modified surface roughness can
422 improve the simulation effect of sensible heat flux in daytime. The results in the
423 figure 10 (c) are basically positive in the daytime, indicating that the latent heat flux
424 after improvement is larger than that before improvement, about $15\text{W}/\text{m}^2$. The above
425 results show that the model simulation results are generally less than the actual
426 observation results, so it can be considered that the improvement of surface roughness
427 in the daytime can improve the simulation of the model. Figure 10 also shows that the
428 improvement of nighttime latent heat flux is not significant, which is basically
429 maintained at $0\text{W}/\text{m}^2$.

430 **Figure 10 about here**

431 **6 Conclusions and discussion**

432 Through the calculation and analysis of the surface roughness of the Nagqu area in the
433 central Tibetan Plateau and comparing the retrieved satellite data with the calculation results of the
434 observational data, the attained main conclusions are as follows.

435 (1) The retrieved results of the satellite data are basically consistent with the calculated
436 results of the measured data. Both results indicate that the surface roughness continued to increase
437 from February to August, began to decrease after reaching a peak in August and reached the

438 lowest value in February of the following year. A strong connection is observed between the
439 monthly variation in surface roughness and the changes in meteorological elements brought by the
440 plateau summer monsoon. Among them, the satellite surface retrieval results in a slow increase in
441 surface roughness from February to May.

442 (2) Through the characteristics of surface roughness variation retrieved by satellite data, the
443 underlying surface can be divided into four categories according to the surface roughness (from
444 large to small: urban, lush grassland, sparse grassland and ice and snow. Among them, lush
445 grassland accounts for 62.49%, and the Z_{0m} can reach 6 cm; sparse grassland accounts for 33.74%,
446 and the Z_{0m} can reach up to 4 cm; and ice and snow account for 3.7%, and the Z_{0m} does not exceed
447 1 cm.

448 (3) Comparing ~~the results of the satellite retrieved calculations~~, satellite retrieved results and
449 measured data shows that the results are positively correlated and the satellite retrieved results are
450 better fit with the measured results. Due to the average sliding effect of the retrieved data, the
451 satellite retrieved data are smaller than the measured results. This method can be used to calculate
452 the surface roughness results for a region and provide a true value for the model for simulation.

453 (4) The accuracy of ground-air flux simulation can be improved after adjusting the surface
454 roughness in Nagqu area. After replacing the model surface roughness, the sensible heat flux has a
455 better daily improvement ~~effect~~, about $20\text{W}/\text{m}^2$. The night improvement ~~effect~~ is poor, about
456 $0.15\text{W}/\text{m}^2$. The improvement of latent heat flux is not obvious, and there is an improvement
457 within $15\text{W}/\text{m}^2$ during the daytime.

458 This study uses remote sensing images and an aerodynamic roughness remote sensing
459 retrieved model to estimate the spatial scale of aerodynamic roughness conditions in northern
460 Tibet, and this method will provide parameter and parameterization scheme improvements for
461 model simulations to study the spatial distribution of the surface flux in the Tibetan Plateau. Air
462 thermodynamics surface roughness is affected by shortwave and longwave radiation (the latter for
463 deriving surface temperature), air temperature, wind speed, precipitation or snowfall. The
464 relationship between air thermodynamics surface roughness and **them** and how to
465 **parameterization** them in Massman **mode** will be studied in the future.

466 **Acknowledgements:**

467 This work was financially supported by the Second Tibetan Plateau Scientific Expedition and
468 Research (STEP) program (Grant No. 2019QZKK0103), the National Natural Science Foundation
469 of China (Grant No. 41675106, 41805009) and National key research and development program
470 of China (2017YFC1505702)

471

472 **References:**

- 473 Arino O, Ramos J, Kalogirou V, et al. "Glob Cover 2009" [R]. Edinburgh, UK: Proceedings of the
474 living planet Symposium, 2010.
- 475 Asrar , G ., Myneni , R.B. and Chaudhury , B.J., 1992, Spatial heterogeneity in vegetation canopies
476 and remote sensing of observed photosynthetically active radiation: a modeling study ,
477 *Ren.Sens.Env .*, 41, 85~103.
- 478 Brutsaert , W.A ., 1982, *Evaporation into the Atmosphere* , Dordrecht in Holland , D .Reidel
479 Publishing Company, 113 ~ 121.
- 480 Chu D., Basabta S., Wang W., et al, 2010. Land Cover Mapping in the Tibet Plateau Using
481 MODIS Imagery[J]. *Resources Science*, 32(11): 2152-2159. (in Chinese with English abstract)
- 482 Chen, J., Wang, J., Mitsuaki, H., 1993. An independent method to determine the surface roughness
483 length. *Chin. J. Atmos. Sci.* 1993, 17, 21–26. (in Chinese with English abstract)
- 484 Chen Q., L. Jia, R. Hutjes, M. Menenti, 2015, Estimation of Aerodynamic Roughness Length over
485 Oasis in the Heihe River Basin by Utilizing Remote Sensing and Ground Data, *Remote Sensing*,
486 2015, 7(4), 3690-3709; doi:10.3390/rs70403690.
- 487 Chen X., et al, 2013. An improvement of roughness height parameterization of the Surface Energy
488 Balance System (SEBS) over the Tibetan Plateau[J]. *Journal of Applied Meteorology and*
489 *Climatology*, 52(3): 607-622.
- 490 Guan X., Huang J., et al, 2009. Variability of soil moisture and its relationship with surface albedo
491 and soil thermal parameters over the Loess Plateau[J]. *Advances in Atmospheric Sciences*, 26(4):
492 692-700, doi: 10.1007/s00376-009-8198-0.
- 493 Högström U., 1996 'Review of Some Characteristics of the Atmospheric Surface Layer'

494 Boundary-Layer Meteorol 78 215–246 10.1007/BF00120937

495 Irannejad P, Shao Y P, 1998. Description and validation of the atmosphere-land-surface interaction
496 scheme(ALSIS)with HAPEX and Cabauw data[J]. Global Planet Change, 19(1): 87-114.

497 Jane Q., 2008. The third pole[J]. Nature, 454(24): 393-396.

498 Jia L., Wang J., Hu Z., 2000. The Characteristics of Roughness Length for Heat and Its Influence
499 on Determination of Sensible Heat Flux in Arid Zone[J]. Plateau Meteorology, 19(04): 495-503.
500 (in Chinese with English abstract)

501 Li F., Li S., Chen T., 2004. A Study on Development of the Tibetan North Grasslands Tourism
502 Stockbreeding in Naqv Tibet[J]. Sichuan Caoyuan, 109(12): 38-42. (in Chinese with English
503 abstract)

504 Li J., Hong Z., Sun S., 2000. An Observational Experiment on the Atmospheric Boundary Layer in
505 Gerze Area of the Tibetan Plateau[J]. Chinese Journal of Atmospheric Sciences, 24(03): 301-312.
506 (in Chinese with English abstract)

507 Li L., Chen X., Wang Z., et al, 2010. Climate Change and Its Regional Differences over the
508 Tibetan Plateau[J]. Advances in Climate Change Research, 6(03): 181-186. (in Chinese with
509 English abstract)

510 Liu J., Zhou M., Hu Y., 2007. Discussion on the Terrain Aerodynamic Roughness[J]. Ecology and
511 Environment, 16(06): 1829-1836. (in Chinese with English abstract)

512 Luo S., Lü S., Yu Z. (2009) Development and validation of the frozen soil parameterization
513 scheme in Common Land Model. Cold Reg Sci Technol 55:130–140

514 Ma Y., Tsukamoto O, Wang J., et al, 2002. Analysis of aerodynamic and thermodynamic
515 parameters on the grassy marshland surface of Tibetan Plateau[J]. Progress in Natural Science,
516 12(1): 36-40.

517 Ma Y., Wang J., et al, 2002. Analysis of Aerodynamic and Thermodynamic Parameters on the
518 Grassy Marshland Surface of Tibetan Plateau[J]. Peogress in Natural Science, 12(01): 36-40. (in
519 Chinese with English abstract)

520 Ma Y., Yao T., Wang J., et al, 2006. The Study on the Land Surface Heat Fluxes over
521 Heterogeneous Landscape of the Tibetan Plateau[J]. Advances in Earth Science, 21(12):

522 1215-1223. (in Chinese with English abstract)

523 Massman W., 1997. An analytical one-dimensional model of momentum transfer by vegetation of
524 arbitrary structure[J]. *Boundary-Layer Meteorology*, 83(3): 407-421.

525 Massman W., 1999. An analytical one-dimensional second-order closure model of turbulence
526 statistics and the Lagrangian time scale within and above plant canopies of arbitrary structure[J].
527 *Boundary-Layer Meteorology*, 91(1): 81-107.

528 Monin A., Obukhov A., 1954. Basic laws of turbulent mixing in the atmosphere near the ground[J].
529 *Tr Akad Nauk SSSR Geofiz Inst*, 24(15): 163-187.

530 Moran, M .S ., Clarke , T .H., Inone, Y.et al., 1994, Estimating crop water deficit using the relation
531 between surface-air temperature and spectral vegetation index, *Rem .Sens. of Env .*, 49 , 246 ~
532 263.

533 Panosky H., Dutton J., 1984. *Atmospheric Turbulence: Models and Methods for Engineering*
534 *Applications*[M]. New York: John Wiley, 1-399.

535 Pierce , L.L., Walker, J., Downling , T .I.et al., 1992, Ecological change in the Murry -Darling
536 Basin-III :A simulation of regional hydrological changes , *Journal of Applied Ecology* , 30, 283~
537 294.

538 Raupach, M .R., 1994, Simplified expressions for vegetation roughness leng h and zero-plane
539 displacement as functions of canopy height and area index , *Boundary -Layer Meteor .* , 71, 211-
540 216.

541 Shao Y., 2000. *Phsics and Modeling of Wind Erosion*[M]. London: Kluwer Academic Publishers,
542 1-452.

543 Smirnova T., Brown J., Benjamin S., Kenyon J. (2016) Modifications to the rapid update cycle
544 land surface model (RUC LSM) available in the weather research and forecasting (WRF) model.
545 *Mon Weather Rev* 144(5):1851–1865

546 Stanhill G., 1969. A simple instrument for the field measurement of turbulent diffusion flux. *J*
547 *Appl Meteorol* 8:509–513

548 Stull B R, 1991. *An Introduction to Boundary Layer Meteorology*[M]. Beijing: China
549 *Meteorological Press*, 1-737.

550 Su, Z., 1996. Remote Sensing Applied to Hydrology: The Sauer River Basin Study.
551 RuhrUniversität Bochum, Lehrstuhl für Hydrologie, Wasserwirtschaft und Umwelttechnik.

552 Tao S., Chen, L., Xu X., et al, 1998. Progresses of the Theoretical Study in the Second Tibetan
553 Plateau Experiment of Atmospheric Sciences (PartI)[M]. Beijing: China Meteorological Press,
554 1-348.

555 Wu G., Mao J., Duan, A., et al., 2004. Recent progress in the study on the impact of Tibetan
556 Plateau on Asian summer climate[J]. *Acta Meteorologica Sinica*, 62(5): 528-540. (in Chinese with
557 English abstract)

558 Wu G., Zhang Y., 1998. Tibetan Plateau forcing and timing of the Monsoon onset over south Asia
559 and the south China sea[J]. *Monthly Weather Review*, 4(126): 913-927.

560 Wu G., Zhang Y., 1999. Thermal and Mechanical Forcing of the Tibetan Plateau and Asian
561 Monsoon Onset.Part II :Timing of the Onset[J]. *Chinese Journal of Atmospheric Sciences*,
562 23(01): 52-62. (in Chinese with English abstract)

563 Wu G., Liu Y., Liu X., et al. 2005. How the heating over the Tibetan Plateau affects the Asian
564 climate in summer[J]. *Chinese Journal of Atmospheric Sciences*, 29(1): 47-56. (in Chinese with
565 English abstract)

566 Wu X., Ma W., Ma Y., 2013. Observation and Simulation Analyses on Characteristics of Land
567 Surface Heat Flux in Northern Tibetan Plateau in Summer[J]. *Plateau Meteorology*, 32(05):
568 1246-1252. (in Chinese with English abstract)

569 Xie Z., Hu Z., Liu H., Sun G., et al., 2017 Evaluation of the Surface Energy Exchange Simulations
570 of Land Surface Model CLM4.5 in Alpine Meadow over the Qinghai-Xizang Plateau. *Plateau
571 Meteorology*, 36(1): 1-12. (in Chinese with English abstract)

572 Xu L., Liu H., Xu X., et al. 2018, Applicability of WRF model to the simulation of atmospheric
573 boundary layer in Nagqu area of Tibetan Plateau[J]. *Acta Meteorologica Sinica*, 2018(6):955-967.
574 (in Chinese with English abstract)

575 Yang M., Yao T., 1998. A Review of the Study on the Impact of Snow Cover in the Tibetan
576 Plateau on Asian Monsoon[J]. *Journal of Glaciology and Geocryology*, 20(2): 90-95. (in Chinese
577 with English abstract)

578 Ye D., Wu G., 1998. The role of heat source of the Tibetan Plateau in the general circulation[J].
579 Meteorological and Atmospheric Physics, 67(1): 181-198.

580 Zhang G., Zhou G., Chen F. 2017, Analysis of Parameter Sensitivity on Surface Heat Exchange in
581 the Noah Land Surface Model at a Temperate Desert Steppe Site in China[J]. Acta Meteorologica
582 Sinica. (6).1167-1182.doi:10.1007/s13351-017-7050-1.

583 Zhang Q., Lv S., 2003. The Determination of Roughness Length over City Surface[J]. Plateau
584 Meteorology, 22(01): 24-32. (in Chinese with English abstract)

585 Zhang Y., Wu G., 1998. Diagnostic Investigations of Mechanism of Onset of Asian Summer
586 Monsoon and Abrupt Seasonal Transitions Over Northern Hemisphere PartI[J]. Acta
587 Meteorologica Sinica, 56(5): 2-17. (in Chinese with English abstract)

588 Zhou X., Zhao P., Chen J., et al, 2009. Impacts of Thermodynamic Processes over the Tibetan
589 Plateau on the Northern Hemispheric Climate[J]. Sci China Ser D-Earth Sci, 39(11): 1473-1486.
590 (in Chinese with English abstract)

591 Zhou, Y., Ju, W., Sun, X., Wen, X., Guan, D. 2012 Significant decrease of uncertainties in sensible
592 heat flux simulation using temporally variable aerodynamic roughness in two typical forest
593 ecosystems of China. J. Appl. Meteorol. Climatol. 2012, 51, 1099–1110.

594 Zhou Y., Xu W., Bai A., et al, 2017. Dynamic Snow-melting Process and its Relationship with Air
595 Temperature in Tuotuohe, TibetanPlateau[J]. Plateau Meteorology, 36(1): 24-32. (in Chinese with
596 English abstract)

597

598 Figure captions

599 Fig. 1 Location of sites and the land cover on the northern Tibetan Plateau. The black solid circle
600 '●' indicates the location of the sites

601 Fig. 2 Surface roughness length of different sites on the northern Tibetan Plateau

602 Fig. 3 Surface roughness length on the northern Tibetan Plateau in 2008

603 Fig. 4 Surface roughness length on the northern Tibetan Plateau in 2010

604 Fig. 5 Surface roughness length on the northern Tibetan Plateau in 2012

605 Fig. 6 Comparison of the surface roughness length by site observations and satellite remote **sense**

606 retrieved data

607 Fig. 7 Scatter plots of the retrieved and calculated surface roughness lengths at four sites

608 Fig. 8 Curve of the surface roughness length for different underlying surfaces








609 Fig.9 Comparison of simulated and observed sensible heat flux (a) and Comparison of simulated
610 and observed latent heat flux (b)

611 Fig. 10 The difference of the control and sensitivity experiments simulated regional sensible heat
612 flux (a) 12:00, (b) 00:00 and latent heat flux (c) 12:00, (d) 00:00







613

614 Tables

615 Table 1. Legend of the land cover map on the northern Tibetan Plateau

Value	Colour	Land Cover Types	Percent (%)
11		Mountain grassland	5.79
14		Shrub meadow	3.25
20		Mountain meadow	8.26
30		Alpine grassland	45.16
70		Needle-leaved evergreen forest	0.23
100		Mixed forest	0.03
110		Mixed forestland and grassland	0.06

616

Value	Colour	Land Cover Types	Percent (%)
120		Mixed grassland and	0.04
140		Alpine meadow	28.28
150		Alpine sparse vegetation	0.29
190		Urban areas	0.07
200		Bare areas	4.90
210		Water bodies	2.57
220		Permanent snow and ice	1.07

617

618

619

620

621

622 Table 2 the observed albedo of the sites (BJ and **Namco**) on the northern Tibetan

623 Plateau

Site	year	Jan.	Feb.	Mar.	Apr.	May.	Jun.	Jul.	Aug.	Sep.	Oct.	Nov.	Dec.
BJ	2008	0.13	0.17	0.14	0.13	0.12	0.10	0.09	0.09	0.09	0.16	0.32	0.16
	2009	0.30	0.26	0.30	0.25	0.26	0.21	0.19	0.19	0.21	0.26	0.24	0.26
	2010	0.26	0.30	0.31	0.30	0.34	0.22	0.19	0.18	0.18	0.35	0.26	0.27
NAMCO	2008	0.28	0.28	0.31	0.28	0.25	0.21	0.17	0.18	0.18	0.30	0.89	0.28
	2009	0.35	0.32	0.28	0.24	0.26	0.22	0.19	0.17	0.22	0.24	0.31	0.27
	2010	0.45	0.23	0.25	0.24	0.29	0.22	0.20	0.17	0.18	0.40	0.35	0.24

624

625

626

627

628

629

630

631

632


633

Table 3 The selected other schemes in Noah-MP

Options for different schemes	Name of the option
dynamic vegetation	Use table LAI; use FVEG=SHDFAC from input
canopy stomatal resistance	Ball-Berry's method
soil moisture factor for stomatal resistance	Noah's method
runoff and groundwater	TOPMODEL with groundwater
surface layer drag coeff	Monin-Obukhov's method
supercooled liquid water	No iteration
frozen soil permeability	Nonlinear effects, less permeable
radiation transfer	Two-stream applied to grid-cell
ground snow surface albedo	Classic method
partitioning precipitation into rainfall & snowfall	Jordan's method
lower boundary condition of soil temperature	TBOT at ZBOT (8m) read from a file
snow/soil temperature time scheme	full implicit (original Noah); temperature top boundary condition

634

635




636

637 Fig. 1 Location of the sites and the land cover on the northern Tibetan Plateau. The black solid

638 circle '●' indicates the location of the sites.


639



640

641 Fig. 2 Surface roughness length of different sites on the northern Tibetan Plateau

642



643


644

645

646 Fig. 3 Surface roughness length on the northern Tibetan Plateau in 2008

647

648



649


650

651

652 Fig. 4 Surface roughness length on the northern Tibetan Plateau in 2010

653

654



655


656

657

658

659

Fig. 5 Surface roughness length on the northern Tibetan Plateau in 2012




660

661 Fig. 6 Comparison of the surface roughness length by site observations and satellite remote sense

662 retrieved data

663



664

665

666

Fig. 7 Scatter plots of the retrieved and calculated surface roughness lengths at four sites

667

(a-d: scatter plot of the observation results and the average result of the underlying surface; e-h:


668

scatter plot of the observation and retrieved results; a, e: BJ station in 2008; b, f: BJ station in


669

2012; c, g: NAMC station in 2010; and d, h: NPAM station in 2012)


670




671



672



673




674

675 Fig. 8 Curve of the surface roughness length for different underlying surfaces

676

677


(a)



678

679

(b)




680

681

682


683

Figure 9 Comparison of simulated and observed sensible heat flux (a) and Comparison of simulated and observed latent heat flux (b)



684

685



686

687 Figure 10 The difference of the control and sensitivity experiments simulated
 688 regional sensible heat flux (a) 12:00, (b) 00:00 and latent heat flux (c) 12:00, (d)
 689 00:00

690

691

692