

# 1 **Satellite soil moisture data assimilation for improved operational** 2 **continental water balance prediction**

3 Siyuan Tian<sup>1</sup>, Luigi J. Renzullo<sup>1</sup>, Robert C. Pipunic<sup>2</sup>, Julien Lerat<sup>2</sup>, Wendy Sharples<sup>2</sup>, Chantal Donnelly<sup>2</sup>

4 <sup>1</sup>Fenner School of Environment & Society, Australian National University, Canberra, 2601, Australia

5 <sup>2</sup>Bureau of Meteorology, Melbourne, 3000, Australia

6 *Correspondence to:* Siyuan Tian (siyuan.tian@anu.edu.au)

7 **Abstract.** A simple and effective two-step data assimilation framework was developed to improve soil moisture representation  
8 in an operational large-scale water balance model. The first step is a *Kalman* filter type sequential state updating process that  
9 exploits temporal covariance statistics between modelled and satellite-derived soil moisture to produce analysed estimates.  
10 The second step is to use analysed surface moisture estimates to impart mass conservation constraints (mass redistribution) on  
11 related states and fluxes of the model using tangent linear modelling theory in a post-analysis adjustment after the state updating  
12 at each time step. In this study, we assimilate satellite soil moisture retrievals from both SMAP and SMOS missions  
13 simultaneously into the Australian Water Resources Assessment Landscape model (AWRA-L) using the proposed framework  
14 and evaluate its impact on the model's accuracy against in-situ observations across water balance components. We show that  
15 the correlation between simulated surface soil moisture and in-situ observation increases from 0.54 (open-loop) to 0.77 (data  
16 assimilation). Furthermore, indirect verification of root-zone soil moisture using remotely sensed Enhanced Vegetation Index  
17 (EVI) time series across cropland areas results in significant improvements from 0.52 to 0.64 in correlation. The improvements  
18 gained from data assimilation can persist for more than one week in surface soil moisture estimates and one month in root-  
19 zone soil moisture estimates, thus demonstrating the efficacy of this data assimilation framework.

## 20 **1 Introduction**

21 Accurate estimation of soil moisture is fundamental to monitoring and forecasting water availability and land surface  
22 conditions under extreme events such as droughts, heatwaves and floods (Ines et al., 2013; Sheffield and Wood, 2007; Tian et  
23 al., 2019b). The spatial pattern of soil moisture can vary significantly due to the heterogeneous spatial distribution of rainfall  
24 and variability in soil properties, land cover type and topography. Due to this large spatial variability, the utility of ground-  
25 based, point-scale measurements is limited in estimating soil water availability at continental scale. Soil moisture estimates  
26 from land surface models are adversely affected by the uncertainties of atmospheric forcing, model dynamics and model  
27 parameterization. Remotely sensed data can provide spatially and temporally varying constraints on the modelling of  
28 biophysical landscape variables that are often superior to that achieved by a single static set of model parameters. Data

29 assimilation merges models and observations in a way that take advantage of their respective strength (e.g. uncertainty,  
30 coverage), resulting in improved accuracy, coverage, and ultimately forecasting capability.

31 The assimilation of satellite soil moisture (SSM) into land surface and hydrology models has been repeatedly demonstrated to  
32 improve model representation of soil water dynamics, evapotranspiration and streamflow (De Lannoy and Reichle,  
33 2016;Draper et al., 2012;Kumar et al., 2009;Li et al., 2012;Pipunic et al., 2008;Reichle and Koster, 2005;Renzullo et al.,  
34 2014;Tian et al., 2019a;Tian et al., 2017;Crow and Yilmaz, 2014;Su et al., 2014a;Alvarez-Garreton et al., 2015;Crow and Ryu,  
35 2009;Baldwin et al., 2017;Patil and Ramsankaran, 2017;Wanders et al., 2014b;Peters-Lidard et al., 2011). Accurate knowledge  
36 of initial soil moisture states gained from data assimilation contributes significantly to the skill of flood forecasting, drought  
37 monitoring and weather forecasts (Bolten et al., 2009;Carrera et al., 2019;Wanders et al., 2014b;Yan et al., 2018;Alvarez-  
38 Garreton et al., 2015). Wanders et al. (2014a) found that the assimilation of remotely sensed soil moisture in combination with  
39 discharge observation can improve the quality of the operational flood alerts, both in terms of timing and in the exact height  
40 of the flood peak.

41 Methods of assimilation are many and varied, however commonalities exist between them. These commonalities are such, that  
42 for any time step, the time integrated first guess (the forecast) of soil moisture states are adjusted by an amount determined by  
43 the difference between observed and modelled soil moisture (the innovation), which is weighted by the respective error  
44 variances of modelled and observed quantities (the gain), to generate revised soil moisture states (the analysis). At the end of  
45 this process, the revised model soil moisture states are out of balance with the other stores and fluxes, until the model integrates  
46 forward to the next time step, whereupon water balance discontinuity is progressively removed through model physics. Soil  
47 moisture is the linchpin between atmospheric fluxes, surface- and ground-water hydrology, thus it is important that any changes  
48 in modelled state variables are not detrimental to other components of the water balance. However, the assimilation of remotely  
49 sensed soil moisture or total water storage data may lead to undesired impacts on groundwater or evapotranspiration  
50 simulations due to the mass imbalance or random error covariances (Giroto et al., 2017;Tangdamrongsub et al., 2020;Tian et  
51 al., 2017). Studies considering mass conservation in data assimilation often require extra data sources such as  
52 evapotranspiration and runoff as constraints or without considering the fluxes (Li et al., 2012;Pan and Wood, 2006).

53 From an operational water balance perspective, is it important that the method of data assimilation be: i) computationally  
54 efficient for routine, automated simulation over the whole model domain; ii) robust to data gaps; and iii) make lasting positive  
55 improvements to future predictions of soil water stores and fluxes. An additional constraint is that if a data assimilation method  
56 is applied to an existing operational system, then it ought to require minimal modification to the system framework, and be as  
57 least disruptive as possible to the model performance. Currently, there are few operational continental water balance modelling  
58 systems that provide near-real time soil moisture estimates that have been constrained through the assimilation of satellite  
59 observations, and mainly at a relatively coarse resolution. Some recent examples include surface soil wetness observations  
60 from Advanced Scatterometer (ASCAT) active radar system, on the meteorological operational satellite (MetOp), being

61 assimilated into Unified Model (Davies et al., 2005) through nudging to provide soil moisture analysis at 40 km globally  
62 (Dharssi et al., 2011). Additionally, ASCAT data are used in the ECMWF (European Centre for Medium-Range Weather  
63 Forecasts) Land Data Assimilation System through a simplified Extended Kalman Filter approach (de Rosnay et al., 2013) to  
64 provide near-real time surface soil moisture and root-zone soil moisture at 25-km resolution globally. SMOS (Soil Moisture  
65 and Ocean Salinity) brightness temperatures have been assimilated in ECMWF's global NWP (Numerical Weather Prediction)  
66 system through the Surface Data Assimilation System, based on the Extended Kalman filter, to produce soil moisture reanalysis  
67 at 40-km resolution (Muñoz-Sabater, 2015). Level-2 Radiometer soil moisture retrievals from SMAP mission (Entekhabi et  
68 al., 2010) have been assimilated into the real-time instance of the NASA Land Information System (LIS) over the  
69 Conterminous United States (CONUS) to produce hourly outputs at 0.03° resolution using ensemble Kalman filter  
70 (Blankenship et al., 2018). However, unlike the aforementioned systems where data assimilation is inherent in the system  
71 design, many operational water balance models, or catchment hydrology models, are calibrated to observations a priori.  
72 Including data assimilation as an afterthought restrains the flexibility of the system, thereby limiting the complexity of the data  
73 assimilation scheme for operational use.

74 In this study, we develop a simple, computationally efficient, and effective data assimilation framework with mass  
75 conservation for incorporating satellite soil moisture products into an existing operational national water balance model. We  
76 demonstrate the application of the method to the Australian Water Resources Assessment Landscape model (AWRA-L), which  
77 provides daily water balance estimates at ~5-km resolution across Australia, with the assimilation of satellite soil moisture  
78 from both SMOS and SMAP. The proposed data assimilation framework is a two-step process that requires minimal  
79 modification of the existing operational system. The first step is the sequential state updating, with weightings between models  
80 and observations derived from the Triple Collocation (TC) approach (Chen et al., 2018; Crow and Van den Berg, 2010; Crow  
81 and Ryu, 2009; Crow and Yilmaz, 2014; Su et al., 2014b; Yilmaz and Crow, 2014). The second step is to impart mass  
82 conservation constraints on related states and fluxes such as root-zone soil water storage, evapotranspiration and streamflow,  
83 thus improving the accuracy of the water balance post assimilation. Accurate initial water balance conditions are of critical  
84 importance in the forecasting of water availability and land surface water dynamics. However, few studies quantify how long  
85 the impacts of data assimilation persist in the model system's memory. To explore the impacts of data assimilation on model  
86 predictions, we quantified the persistence of the correction to key model components with respect to open-loop simulations,  
87 to illustrate the potential gains from data assimilation in improving water balance forecasts.

## 88 **2 Materials**

### 89 **2.1 Australian Water Resources Assessment Modelling system**

90 The Australian Water Resources Assessment Landscape (AWRA-L) model (Van Dijk, 2010) underpins the annual national  
91 water resource assessments and water use accounts for Australia (Frost et al., 2018). The operational implementation of the  
92 AWRA-L by the Australian Bureau of Meteorology provides daily 0.05-degree (approximately 5 km) national gridded water

93 balance estimates. The outputs from the operational AWRA-L has been widely used in various agricultural applications and  
94 natural resources risk assessment and planning, including commodity forecasting, irrigation scheduling, flood and drought risk  
95 analysis, as well as flood forecasting (Frost et al., 2018;Hafeez et al., 2015;Nguyen et al., 2019;Van Dijk et al., 2013;van Dijk  
96 and Renzullo, 2011). The version of the AWRA-L model used in the study was obtained from the Community Modelling  
97 system (AWRA-CMS) and is freely available from [https://github.com/awracms/awra\\_cms](https://github.com/awracms/awra_cms).

98 AWRA-L is a one-dimensional grid-based model that simulates water balance for each grid cell across the modelling domain  
99 by distributing rainfall influx into plant-accessible water, soil moisture and groundwater stores, and computing outflux such  
100 as evapotranspiration, runoff and deep drainage. The soil water column is partitioned into three layers (surface: 0–10 cm,  
101 shallow: 10–100 cm, and deep: 1–6 m) and simulated separately two hydrological response unit, i.e. deep-rooted (trees) and  
102 shallow-rooted (grass) vegetation. The water storage for the surface-layer soil is termed  $S_0$ , while  $S_s$  is used for the shallow-  
103 layer and  $S_d$  for the deeper-layer. In addition to the modelling of soil columns, the model includes a surface water and a  
104 groundwater storage that are simulated at each grid cell and conceptualized as a small unimpaired catchment. In this study, we  
105 used forcing inputs from the AWAP (Australian Water Availability Project) gridded climate data including daily precipitation,  
106 air temperature and solar exposure (Jones et al., 2009), and interpolated site-based wind speed (McVicar et al., 2008). It is  
107 acknowledged that the accuracy of the model estimates is limited in regions with insufficient coverage in the ground-based  
108 observation network (e.g. rain gauges) which is the raw source of AWAP gridded data used to force the model. This is limited  
109 to very remote and mostly uninhabited arid regions in Australia.

## 110 2.2 Satellite soil moisture (SSM)

111 To maximize daily spatial coverage, we used two satellite soil moisture products derived from passive L-band systems: the  
112 Soil Moisture Active-Passive (SMAP) product from NASA (Entekhabi et al., 2010); and the product from the European Space  
113 Agency's (ESA's) Soil Moisture and Ocean Salinity (SMOS) mission (Kerr et al., 2001). The SMAP product is the level-2  
114 enhanced radiometer half-orbit 9-km EASE-grid soil moisture (Chan et al., 2018). The SMOS product is the level-2 soil  
115 moisture product on ~ 25-km grid (Rahmoune et al., 2013). Both SMAP and SMOS produce volumetric soil moisture estimates  
116 (units  $m^3/m^3$ ) of approximately the upper 5 cm of soil. Available swath data for each product covering Australia were collated  
117 for each 24-hour period approximating the AWRA-CMS operational time steps and resampled to a regular 0.05-degree grid  
118 across the modelling domain using linear interpolation from 2015 to 2019. The volumetric soil moisture retrievals from both  
119 SMAP and SMOS were converted into water storage units (mm) to be consistent in units and soil depths with model estimates,  
120 using mean and variance matching to remove the systematic bias. Figure 1 shows an example of daily composites of SMAP  
121 (Fig1.a) and SMOS (Fig1.b) soil moisture retrievals in model units compared to AWRA-L estimates of  $S_0$  (Fig1.c). For regions  
122 with sparse rain-gauge coverage such as central Western Australia (Fig1.c), AWRA-L modeled  $S_0$  persists as zeros or very  
123 low values for the experiment period, reflecting a deficiency in the gauge-based analysis of daily rainfall used to drive model  
124 simulations. The result of mean and variance matching in these gauge-sparse areas will flatten the variability of SSM time

125 series to zero when using values of the modelled  $S_0$  for these areas directly. To resolve this problem, and fully leverage the  
126 information available in the SSM products to fill the gaps in modelled outputs across the continent, we derived a set of  
127 coefficients for the mean and variance matching over the gauge sparse regions by sampling modelled and SSM data from cells  
128 surrounding the gaps. Specifically, we fitted a linear model between the maximum SSM values through time and the  
129 coefficients for mean and variance matching for each cell in neighboring region. We applied the derived linear relationship to  
130 estimate the correspond ‘slope’ and ‘intercept’ from the maximum SSM values in the rainfall gaps. This provided a  
131 transformation of the SSM into water storage unit (mm) and ensures the assimilation can effectively influence the spatial  
132 pattern of soil moisture over the sparsely gauged regions.

## 133 **2.3 Validation data**

### 134 **2.3.1 In-situ measurements**

135 Evaluation of the modelled soil water storages was made against measurements from three soil moisture monitoring networks  
136 in Australia from 2016 to 2018, namely OzNet (Smith et al., 2012), CosmOz (Hawdon et al., 2014) and OzFlux (Fig. 1d).  
137 AWRA-L model estimates of water storage in surface soil layer were compared against in situ measurements from the top 10  
138 cm of soil across all three networks. The depths of in situ measurements of root-zone moisture varied across networks from 0-  
139 30 cm to 0-1 m. As such, AWRA-L soil water storages over the root-zone were constructed by combining surface- and shallow-  
140 layer soil water storage in the appropriate proportions to be consistent with in situ measurement depth. OzFlux sites are also  
141 used for the evaluation of AWRA-L evapotranspiration estimates, which were calculated from accumulated latent heat flux  
142 measurements at each location. In total, there are 45 sites for soil moisture validation and 14 sites for evapotranspiration  
143 validation. Streamflow observations for 110 catchments across Australia have been used in the validation based on the quality  
144 and data availability (Fig. 1d).

### 145 **2.3.2 Vegetation index**

146 In water-limited regions like Australia, shallow-rooted vegetation normally responds quickly to soil water availability,  
147 typically within a month. Consistency between root-zone soil water storage and vegetation greenness may be considered as an  
148 indirect independent verification of the simulation of root-zone soil water dynamics (Tian et al., 2019a; Tian et al., 2019b). The  
149 0.05-degree monthly Enhanced Vegetation Index (EVI) from Moderate Resolution Imaging Spectroradiometer (MODIS,  
150 MYD13C2 v6) was used to evaluate estimates of monthly root-zone soil water storage (the sum of water storage in surface-  
151 layer ( $S_0$ ) and shallow-layer ( $S_s$ ) within the AWRA-L soil column) over cropland regions of the continent. The EVI is used  
152 here to characterize vegetation dynamics since it is not as influenced by atmospheric effects and canopy background noise, and  
153 has a greater dynamic range (i.e., less likely to saturate) in areas of dense vegetation compared to the Normalized Difference  
154 Vegetation Index (NDVI). The choice of root-zone soil water storage at the 0-1 m depth is due to the average rooting depths  
155 varying from 30 - 80 cm over the cropland areas in Australia (Donohue et al., 2012; Figueroa-Bustos et al., 2018; Incerti and

156 O'Leary, 1990). The 250-m land cover classification map from Geoscience Australia (Lymburner, 2015) was resampled to  
157 0.05 degree over the model domain and used in the identification of cropland areas.

### 158 **3 Method**

#### 159 **3.1 Triple collocation-based error characteristics**

160 Triple collocation (TC) was developed as a method of quantifying error characteristics in geophysical variables when the  
161 true error structure is elusive. It was first applied to near-surface wind data (Stoffelen, 1998) and later extensively applied to  
162 soil moisture (Chen et al., 2018; Crow and Yilmaz, 2014; Dorigo et al., 2017; McColl et al., 2014; Scipal et al., 2008; Su et al.,  
163 2014a; Yilmaz and Crow, 2014; Zwieback et al., 2013; Crow and Van den Berg, 2010) and rainfall (Alemohammad et al.,  
164 2015; Massari et al., 2017). The assumption of this approach is that three independent data sets of the same geophysical  
165 variable can be used to infer the error variances in each. Here we use TC as a way of inferring error variances from our three  
166 independent estimates of surface soil moisture, AWRA-L  $S_0$ , SMAP, and SMOS from 2015 to 2019. Those three collocated  
167 measurements were assumed to be linearly related to the true value with additive random errors. To ensure the errors from  
168 the three independent sources were unbiased relative to each other, SMAP and SMOS soil moisture retrievals were rescaled  
169 to the reference model estimates (AWRA-L  $S_0$ ) using temporal mean and variance matching. McColl et al. (2014) shows that  
170 the error variances ( $\sigma^2$ ) of each data set can be calculated from the temporal variance and covariance between data sets  
171 respectively as:

$$172 \sigma_x^2 = \left( Q_{x,x} - \frac{Q_{x,y}Q_{x,z}}{Q_{y,z}} \right), \quad \sigma_y^2 = \left( Q_{y,y} - \frac{Q_{x,y}Q_{y,z}}{Q_{x,z}} \right) \quad \text{and} \quad \sigma_z^2 = \left( Q_{z,z} - \frac{Q_{z,y}Q_{x,z}}{Q_{x,y}} \right) \quad (1)$$

173 where x, y and z denote AWRA-L, SMAP and SMOS soil moisture estimates respectively and Q denotes temporal variance  
174 and covariance between the data sets. These estimates of error variance are used in the determination of the weighting of each  
175 data source in the data assimilation (Section 3.2).

#### 176 **3.2 Sequential state updating**

177 The data assimilation method used here is a time sequential updating of model state(s) given observations of relevant model  
178 variables (Reichle, 2008). There are two key modelling components in data assimilation: the dynamics operator, which  
179 describes the time integration of the system states and fluxes, which in this study is the AWRA-CMS; and the observation  
180 operator, which provides the mathematical mapping from state to observation space. The role of the observation operator is to  
181 perform a mapping between observation and state space, as often observations are not directly comparable to model states.

182 The common state updating equation for sequential data assimilation is written as:

$$183 x_t^a = x_t^f + K_t[y_t - H(x_t^f)] \quad (2)$$

184 which says that the best estimate of model state, known as analysis ( $x_t^a$ ), is equal to the first guess or forecast ( $x_t^f$ ) plus a  
 185 weighted difference between observations,  $y_t$ , and the model equivalent to the observation,  $H(x_t^f)$ , for that time step. In this  
 186 study, the AWRA-L model soil water storage in  $S_0$  for shallow-rooted vegetation and deep-rooted vegetation at surface layer  
 187 are updated directly through the sequential data assimilation. Satellite surface soil moisture (SSM) products from both SMOS  
 188 and SMAP are used as the observations to update the model simulation. The observation operator  $H$  here is the aggregation of  
 189 soil water storage estimates in the top-soil layer for two land cover types, i.e. shallow-rooted vegetation and deep-rooted  
 190 vegetation. The multiplier,  $K$ , is known as the gain factor which contains uncertainty expressed as error variance ( $\sigma^2$ ) for both  
 191 model estimates and observations. For example, the gain factor for the AWRA-L estimates can be written as:

$$192 \quad K_x = \frac{\frac{1}{\sigma_x^2}}{\frac{1}{\sigma_x^2} + \frac{1}{\sigma_y^2} + \frac{1}{\sigma_z^2}}, \quad (3)$$

193 where  $x, y, z$  denotes AWRA-L estimates, SMAP and SMOS soil moisture retrievals. Observation error variance is often  
 194 estimated through field campaigns (Draper et al., 2009; Panciera et al., 2013), but these rarely represent the spatial and temporal  
 195 variability of errors in gridded satellite products. Alternatively, data providers often specify error estimates, but their magnitude  
 196 can be overly optimistic. Here, we applied the triple collocation approach (Section 3.1) to characterise the temporal error  
 197 variances of the model estimates and the two satellite observations for each grid cell across Australia. The analysis receives  
 198 higher contribution from observation with smaller error variance (Eq. 2). Given the relatively short time series (small number)  
 199 of observations, however, a single set of error variances is calculated for all time. This results in spatially varying but  
 200 temporally static error variances (and thus gain weights) for each of the three sources (Fig. 2). We acknowledge the limitations  
 201 of assuming a temporally constant error variances and future refinements to the assimilation method will consider introducing  
 202 seasonally varying error variances.

### 203 3.3 Analysis increment redistribution (AIR)

204 The assimilation of satellite soil moisture temporarily violates mass conservation in the model through the analysis update.  
 205 The difference between the analysis,  $x_t^a$ , and the forecast,  $x_t^f$ , (known as the *analysis increment*) represents an amount of  
 206 water that has been added or subtracted from the system that was not present at the start of model integration for the given  
 207 time step. In this study, we use the concept of tangent linear modelling (Errico, 1997; Giering, 2000) to redistribute the  
 208 analysis increment of surface soil water storage,  $S_0$ , to all the relevant model states and fluxes as a way of maintaining mass  
 209 (i.e. water) balance within each model time step. This adjustment is applied after the sequential state updating as the second-  
 210 step in the assimilation framework, which we refer to as analysis increment redistribution (AIR).

211 The adjoint and tangent linear models were originally used in variational data assimilation (Bouttier and Courtier, 2002) and  
 212 have been used to estimate the sensitivity of model outputs with respect to input (Errico, 1997). We assume the input

213 perturbation here is the analysis increment after the data assimilation (i.e.  $x_t^a - x_t^f$  from Eq. 2), then the change in other model  
214 outputs due to the change in inputs can be determined through tangent linear modelling. Assuming model variable  $b$  is  
215 related to the state variable  $x$ , the relationship between them can be simply described as:

$$216 \quad b = M(x), \tag{4}$$

217 where  $M$  denotes the model operator. The change in output variable  $\Delta b$  at time step  $t$  due to the input change  $\Delta x$  can be  
218 determined by

$$219 \quad \Delta b_t = \frac{\partial M}{\partial x_t} \Delta x_t. \tag{5}$$

220 In this study, we applied the tangent linear modelling approach to correct the model forecast of soil water storage for  
221 shallow-layer ( $S_s$ ), and deep-layer soil water storage ( $S_d$ ), evapotranspiration ( $E_{tot}$ ), and total streamflow ( $Q_{tot}$ ) after the  
222 state updating of surface soil moisture ( $S_0$ ) at each time step. Note that this process ensures that the correction is affecting all  
223 model states in proportion to their sensitivity against changes in the  $S_0$ . All the model equations regarding to the mass  
224 redistribution were derived using model equations (Frost et al., 2018; Van Dijk, 2010) and can be found in the Appendix A.

## 225 **4. Results**

### 226 **4.1 Impact on surface soil water storage estimates**

227 Error variances were derived using TC for AWRA-L model estimates and the SSM products, and showed that for the  
228 majority of the grid cells over the continent SMAP soil moisture had smaller error variance than SMOS and the model  
229 estimates. This is consistent with other studies that have shown SMAP provides the best-performing satellite soil moisture  
230 product over the majority of applicable global land pixels (Chen et al., 2018). Figure 2 shows the relative weightings  
231 (derived from the TC error variances) of model estimates, SMOS and SMAP soil moisture in the data assimilation. The  
232 analysed surface soil water storage estimates ( $S_0$ ) receive a greater contribution from SSM products, in particular SMAP  
233 observations, compared to model simulations (Fig. 2). Figure 3 gives an example of the temporal change in modelled  $S_0$   
234 estimates before and after the assimilation for 2017. The temporal dynamics of  $S_0$  estimates after the assimilation has been  
235 highly adjusted towards SSM retrievals and in consistency with in-situ measurements.

236 AWRA-L model simulations are driven by gauge-based rainfall analyses. As such the model has difficulty in adequately  
237 simulating soil moisture patterns over regions lacking in rain gauge coverage, such as Western Australia and central  
238 Australia (Fig. 1c). Water storage simulations over these regions default to zero, thus very little or no weight was given to  
239 the AWRA-L estimates in these regions (Fig. 2a). Figure 4 shows different spatial patterns of daily average  $S_0$  estimates for  
240 December 2019 from model open-loop (OL) without data assimilation and with data assimilation through TC-derived  
241 weighting (DA-TC). Data assimilation has the effect of adding moisture to AWRA-L  $S_0$  simulations over most of gauge-



242 sparse areas as shown in Figure 4c. Analysed AWRA-L simulations of  $S_0$  are dominated by the satellite SSM data as a result  
243 of TC weighting in the region which largely eliminates the erroneous artefacts associated with deficient rainfall data forcing.  
244 Reduced water storage in the surface layer of the soil column was found over southeast of Australia, particularly within the  
245 Murray-Darling Basin. This suggests that AWRA-L OL simulations underestimated the severity of the drought experienced  
246 in the region in December 2019. The analysis increments of AWRA-L  $S_0$  ( $x^a - x^f$ ) were compared with the difference  
247 between in-situ rainfall observations from OzFlux network,  $P^{OzFlux}$  and AWAP rainfall forcing,  $P^{AWAP}$ , (Fig. 5). The  
248 increasing  $S_0$  simulations align with missing or underestimated rainfall events in the AWAP rainfall forcing  
249 ( $P^{OzFlux} - P^{AWAP} > 0$ ) and vice versa (Fig. 5). This supports the hypothesis that data assimilation correctly distributes  
250 water into the system and mitigates the impact of uncertainty in rainfall forcing.

## 251 4.2 Impact on root-zone soil water storage and fluxes estimates

252 If the analysis increment redistribution (AIR) is not applied, the soil water storage in the surface layer ( $S_0$ ) is the only state  
253 variable directly updated with SSM (DA-TC). Other variables such as root-zone soil water storage, evapotranspiration and  
254 streamflow are adjusted with model integration to the next time step using the analysed  $S_0$  as the surface layer initial condition.  
255 Therefore, the observed changes in those variables following DA-TC (Fig.6, centre column) are relatively small when  
256 compared to model open-loop estimates (Fig.6, left column). For example, the OL soil water storage of shallow-layer ( $S_s$ )  
257 estimates in those gauge-sparse regions of Australia remain zero or very low due to the AWAP rainfall forcing. The predictions  
258 of  $S_s$  receive relatively small contribution from the analysed  $S_0$  since the analysis increment of  $S_0$  is small compared to the  
259 field compacity of  $S_s$ .

260 One known issue of sequential state updating is the temporary break of water balance at each time step until the next model  
261 integration. The proposed AIR approach (Section 3.2) adjusts variables coupled with surface soil moisture after the state  
262 updating at each time step. Significant difference in the spatial patterns of  $S_s$ ,  $E_{tot}$  and  $Q_{tot}$  after the mass redistribution (DA-  
263 TCAIR) can be seen in Fig. 6 (right column) compared to model open-loop or forecasts after only  $S_0$  updating. The changes  
264 in estimates of  $S_s$  and  $E_{tot}$  over coastal regions are relatively small due to more accurate rainfall forcing data with the dense  
265 network of rain-gauges. Finally, the  $Q_{tot}$  estimates after AIR are lower than the DA-TC and OL. This reduction in streamflow  
266 over south-eastern Australia and northern Australia is consistent with the reduced surface soil moisture in those regions (Fig.4c).

## 267 4.3 Quantitative evaluation

268 Estimates of surface soil moisture, root-zone soil moisture, evapotranspiration and streamflow after data assimilation (DA-  
269 TC) and data assimilation with mass redistribution (DA-TCAIR) were compared with time series of in-situ observations. We  
270 compared the model outputs after DA-TC and DA-TCAIR separately to investigate the benefits of maintaining mass balance  
271 in data assimilation. Pearson's correlation coefficients were computed from time series of model estimates and observations  
272 between January 2016 to December 2018 for each site. The distribution of correlation coefficients for OL, DA-TC and DA-

273 TCAIR are displayed as boxplots in Figure 7. Consistent, significant improvement in modelled surface layer soil water storage  
274 estimates ( $S_0$ ) were observed across all sites (Fig. 7a) with the single exception of an OzFlux site located in a tropical rainforest,  
275 where microwave SSM retrievals are known to be typically poor (Njoku and Entekhabi, 1996). TC-based assimilation (DA-  
276 TC) increases the correlation between in-situ surface SM measurements from 0.47 to 0.72 on average for CosmOz sites, 0.54  
277 to 0.69 for OzFlux sites, and 0.56 to 0.77 for OzNet sites compared to OL. This is a significant improvement in AWRA-L  
278 simulations of surface soil moisture dynamics with an increase in correlation of 0.23 on average across all in-situ sites.

279 Overall subtle improvements were observed across the AWRA-L estimates of root-zone soil water storage, evapotranspiration  
280 and streamflow after the assimilation (DA-TC) (Fig. 7b, c, d). The level of improvement is not surprising since those variables  
281 were not directly updated through DA-TC and are only influenced through the integration of the model to the next time step.  
282 Degradation was found in root-zone soil moisture estimation for a few OzFlux and OzNet monitoring sites. This is likely due  
283 to the break of water balance in the assimilation, since the estimates followed by the second step of AIR (DA-TCAIR) slightly  
284 increases the correlation with in-situ observations compared to model open-loop and the estimates after assimilation without  
285 mass redistribution (DA-TC). Moreover, the model estimates of root-zone soil moisture from model OL are in good agreement  
286 with in-situ observations as is with average correlation above 0.8 (Fig. 7b), which leaves little room for improvements.

287 Although the corrections of other water balance estimates from the analysis increments redistribution are relatively small  
288 compared to direct state updating, they are improvements nevertheless. Slight improvements were similarly found in  
289 streamflow estimates after the AIR (Fig. 7d). Figure 8 shows an example of the OL estimates of streamflow, the analysed  
290 streamflow after the application of AIR, and the streamflow observations,  $Q_{tot\ obs}$ . Also displayed is the streamflow analysis  
291 increments, i.e.  $Q_{tot}^a - Q_{tot}^f$  for each time step. The negative streamflow analysis increment (Fig. 8) indicates that water is  
292 removed from the surface water store after the assimilation of SSM and application of AIR, which appears to compensate  
293 for the overall overestimate of OL simulations, in this example. Although the change in streamflow due to the soil moisture  
294 data assimilation is small compared to the disparity between model and observed streamflow, the adjustment in the direction  
295 towards observations highlights the importance of accurate antecedent soil moisture conditions in the simulation of runoff  
296 response. The joint assimilation of gauge-measured streamflow and satellite soil moisture retrievals into AWRA-L is expected  
297 to improve the streamflow simulation.

298 A limited number of root-zone soil moisture monitoring sites as well as the large spatial disparity between the point-scale in-  
299 situ measurements and modelling resolution ( $\sim 5$  km grid cell) represent substantial limitations for wide-area evaluation of  
300 root-zone soil moisture estimates. An indirect verification of AWRA-L simulations of root-zone soil moisture was based on a  
301 comparison against satellite-derived EVI. This provided an independent, albeit indirect, way of evaluating the impact of data  
302 assimilation over larger areas. We calculated the correlation between time series of monthly average AWRA-L root-zone soil  
303 moisture estimates from OL, DA-TC and DA-TCAIR against EVI for cropland across Australia from 2015 to 2018. Cropland

304 cover type was selected based on the rooting depths of the dominant grass species and wheat varieties in the area that have  
305 been shown to have rooting depths spanning at least half the combined soil depths (0-1m) of the surface- and shallow-layer  
306 soil water storage in AWRA-L. Figure 9a shows the relative change in correlation between root-zone soil water storage  
307 simulations from DA-TCAIR and those from model OL against EVI data for cropland areas of Australia. Significant  
308 improvements were found after the data assimilation and mass redistribution for the vast majority of model grid cells (Fig. 9a).  
309 The averaged correlation with EVI is 0.64 from DA-TCAIR compared to 0.52 for model open-loop. The root-zone soil water  
310 storage estimates after the mass redistribution are significantly improved over the cropland in Western Australia and southern  
311 Australia with more than 20% increase in correlation comparing to DA-TC without mass redistribution (Fig. 9b). This  
312 demonstrates that enforcing mass balances as part of the soil moisture data assimilation at each time step is essential to  
313 improving the simulation of root-zone soil water balance. Limited difference between DA-TC and DA-TCAIR were found  
314 over cropland regions over south-eastern Australia, likely due to the overall good performance of AWRA-L OL root-zone soil  
315 moisture estimates in those areas (Fig. 7b). The improved consistency with EVI after data assimilation highlights the potential  
316 of improving agricultural planning with more accurate information of root-zone soil water availability.

#### 317 **4.4 Implications for water balance forecasting**

318 To quantify how long improvements in model state last in AWRA-L simulations, we used OL and DA-TCAIR estimates  
319 between 1 March 2018 and 28 February 2019. The model states for each day over this one-year period served as initial  
320 conditions for 100-day AWRA-L simulations from which we calculated the number of days it took for the simulation from the  
321 analysed DA-TCAIR states to converge to within +/- 5% of those from OL. Results showed that data assimilation can impact  
322 model states and fluxes for weeks and sometimes up to 2-3 months (Fig. 10). The impacts of data assimilation can persist in  
323 simulated  $S_0$  for as long as a week over coastal regions, and longer in central Western Australia and Northern Australia with  
324 up to a month persistence in winter and spring (Fig. 10a). There is less impact on  $S_0$  simulations during wet season (Summer-  
325 Autumn) in Northern Australia since the  $S_0$  can saturate rapidly due to the heavy rainfall. Overall, the longest persistence is  
326 found in winter with a continental average of 13 days; the shortest is 6 days on average in autumn and summer. The memory  
327 of initial conditions in simulations of  $S_s$  can persist even longer due to the slower response to rainfall variability and higher  
328 field capacity (Fig. 10b). Summer persistence for  $S_s$  is the least with a continental average of 30 days; in winter this increased  
329 to 45 days.

330 On average, the impact of antecedent soil moisture conditions on evapotranspiration simulations can persist for 1 week over  
331 coastal areas, but up to months in central Western Australia (Fig. 10c). The continental average varies from 13 to 20 days for  
332 each season. The areas with the longest persistence are those areas with artefacts of zero rainfall in the forcing. This  
333 demonstrates that improvements in AWRA-L estimates after SSM assimilation over regions with sparse rain-gauge coverage  
334 can persist in the system for more than 2 months. The impact on runoff varies from 1 week to 3 months over the continent

335 (Fig. 10d). The majority of areas impacted for more than 2 months are in locations of low rainfall and runoff. However, in  
336 areas of heavy runoff, e.g. north-eastern Australia, there is between 1-2 week of persistence.

## 337 **5. Discussion**

338 In this study, we assimilated SMAP and SMOS data into an operational AWRA-L water balance modelling system through a  
339 simple sequential state updating approach, with weightings derived using triple collocation approach (DA-TC), followed by a  
340 post-adjustment for mass redistribution (DA-TCAIR). Previous data assimilation studies using the AWRA-L model opted for  
341 ensemble-based methods (Renzullo et al., 2014;Shokri et al., 2019;Tian et al., 2019a;Tian et al., 2017;Tian et al., 2019b).  
342 Ensemble based methods rely on *a priori* knowledge of uncertainty in forcing data and model error variances to derive spatially  
343 and temporally varying gain matrices at each time step. However ensembles often require post hoc correction such as state  
344 inflation (Anderson et al., 2009) to achieve optimal performance, and many members (> 10) comprised of multiple model  
345 runs to infer statistically meaningful error variances, which can be computationally costly. In contrast, the proposed DA-TC/  
346 TCAIR framework is simple, effective and computationally efficient and requires minimal modification in the current  
347 operational system. The gain factor in the proposed assimilation framework is temporally constant but spatially varying. It is  
348 derived from the temporal covariances between modelled and satellite-derived soil moisture for each grid cell across the  
349 domain through the widely used triple collocation (TC) method (Chen et al., 2018;Crow and Van den Berg, 2010;Crow and  
350 Yilmaz, 2014;Su et al., 2014b;Yilmaz and Crow, 2014). The significant improvements in AWRA-L model surface soil  
351 moisture estimation demonstrates the efficiency of the proposed assimilation approach (Fig. 7a). Temporally varying gain  
352 factor is considered for future improvement to the approach once a longer time series of SMAP data is available.

353 Pan and Wood (2006) used mass redistribution in a two-step constrained Kalman filter that required error covariances derived  
354 from evapotranspiration and runoff observations. However, these observations are often not available for continental scale of  
355 studies. Li et al. (2012) redistribute the mass imbalance within soil layers during the assimilation but without the updates of  
356 fluxes. Our proposed method based on tangent linear modelling redistributes the mass change across all the states and fluxes  
357 related to surface soil moisture states without the need for extra observations. The analysis increment redistribution (AIR)  
358 method conserves the mass balance thereby improving water balance estimates (Fig. 7), in particular it can improve the root-  
359 zone soil moisture estimates over croplands (Fig. 9). Although the improvements are limited, the streamflow estimates from  
360 the AIR are predominantly a better match to observations (Fig. 8). Model physics limits the strength of coupling between an  
361 analysed state and resulting fluxes (Kumar et al., 2009;Walker et al., 2001). Thus a small level of improvement in performance  
362 in AWRA-L streamflow in response to soil moisture state updating is not unexpected due to a weak coupling between the  
363 states and fluxes. Calibration of model parameters using satellite and in-situ observations may lead to further improvements.

364 Many studies have demonstrated the assimilation of satellite soil moisture can improve model forecasts due to the correction  
365 for initial soil moisture conditions (Crow and Ryu, 2009;Pauwels et al., 2001;Scipal et al., 2008). Getirana et al. (2020a) and

366 Getirana et al. (2020b) found that using initial conditions derived from the assimilation of GRACE (Gravity Recovery and  
367 Climate Experiment) total water storage observations can improve the seasonal streamflow and groundwater forecast due to  
368 the long memory of groundwater and soil moisture. However, few studies quantify how long the impacts of data assimilation  
369 can persist in the model system's memory for different states. In this study, we found that the impact of different initial  
370 conditions of root-zone soil water storage has a long memory in the system, exceeding 2 months (Fig.10b). The constraints on  
371 the simulations of surface soil moisture, evapotranspiration and streamflow can persist 1-2 weeks due to the high temporal  
372 variability. This highlights the potential gains from data assimilation for agricultural planning and flood forecasting, as a result  
373 of improved short-term water balance forecasts.

## 374 **6. Conclusion**

375 In this study, we proposed a simple and robust framework for assimilating SMAP and SMOS soil moisture products into the  
376 operational Australian Water Resources Assessment modelling system. The method involves the sequential (daily) updating  
377 of the model's surface soil water storage with satellite soil moisture observations using weights determined through triple  
378 collocation (DA-TC). Furthermore, we proposed an additional component to the data assimilation whereby the analysis  
379 increment of the upper layer soil water storage is propagated into relevant model states and fluxes as a way of maintaining  
380 mass balance (DA-TCAIR). Evaluation against in-situ measurements showed that simulations of surface soil moisture  
381 dynamics is improved significantly after TC data assimilation with an average increase of 0.23 correlation units compared with  
382 open-loop simulations. An evaluation of the root-zone soil moisture, evapotranspiration and streamflow estimates showed that  
383 the TC-AIR appeared to provide marginal, yet positive, improvement over the TC data assimilation method alone. However,  
384 in an indirect verification of modelled root-zone soil moisture against satellite-derived EVI, DA-TCAIR was seen to provide  
385 significant improvement over the TC method alone. This demonstrates that by enforcing mass balances as part of the SSM  
386 data assimilation each time step, AWRA-L can better represent soil water dynamics such that it has greater consistency with  
387 observed vegetation response.

388

389 The assimilation of satellite soil moisture estimates together with the mass redistribution reduces the uncertainties in model  
390 estimates resulting mainly from uncertain forcing and model physics, and provides temporally and spatially varying constraints  
391 on model water balance estimates. For example, the assimilation resolves the gaps in rainfall forcing over Western Australia  
392 and central Australia. We demonstrate that the impacts of data assimilation can persist in the model system for more than a  
393 week for surface soil water storage and more than a month for root-zone soil water storage. This highlights the importance of  
394 accurate initial hydrological states for improving forecast skill over longer lead times. Hence, an operational water balance  
395 modelling system, with satellite data assimilation, has strong potential to add value for assessing and predicting water  
396 availability for a range of decision makers across industries and sectors.

397

398

399 **Appendix A**

400 For a complete understanding and description of the AWRA-L model equations, please refer to Frost et al (2016). Here we  
401 only present those parts of the model equation related to  $S_0$ .

402

403 The analysis increments after the data assimilation can be calculated as:

404 
$$\Delta S_0 = S_0^a - S_0^f,$$

405 where  $S_0^a$  denotes the analysed upper-layer soil water storage and  $S_0^f$  denotes the forecast, or initial estimate. The change in  $S_0$   
406 affects the drainage to the lower-layer soil water storage ( $D_0$ ) and interflow draining laterally from the top soil layer ( $Q_{I0}$ ). The  
407 corresponding change in drainage to lower-layer soil water storage from the increment  $\Delta S_0$  is calculated as:

408 
$$\Delta D_0 = (1 - \beta_0)k_{0sat} \left[ \left( \frac{S_0^a}{S_{0max}} \right)^2 - \left( \frac{S_0^f}{S_{0max}} \right)^2 \right],$$

409 
$$\Delta Q_{I0} = \beta_0 k_{0sat} \left[ \left( \frac{S_0^a}{S_{0max}} \right)^2 - \left( \frac{S_0^f}{S_{0max}} \right)^2 \right],$$

410 where the  $k_{0sat}$  and  $S_{0max}$  are model parameters representing the saturated hydraulic conductivity and maximum storage of  
411 the upper soil layer, respectively. The proportion of overall top layer drainage that is lateral drainage ( $\beta_0$ ) given as:

412 
$$\beta_0 = \tanh \left( k_\beta \beta \frac{S_0^a}{S_{0max}} \right) \tanh \left( k_\zeta \left( \frac{k_{0sat}}{k_{ssat}} - 1 \right) \frac{S_0^a}{S_{0max}} \right),$$

413 where  $\beta$  and  $k_\beta$  are the slope radians and scaling factor, and  $k_\zeta$  is a scaling factor for the ratio of saturated hydraulic  
414 conductivity. The revised lower-layer soil water storage  $S_s^a$  is then determined as:

415 
$$S_s^a = S_s^f + \Delta D_0.$$

416 The change in  $S_s$  will lead to the change in the shallow soil water storage ( $D_s$ ) and lateral interflow ( $Q_{Is}$ ). The soil water storage  
417 at lower layer is thus updated as:

418 
$$S_d^a = S_s^a + \Delta D_s.$$

419 Similarly, the groundwater storage  $S_g$  will be adjusted with the increment of deep soil layer drainage.

420 The total runoff ( $Q_{tot}^a$ ) should be updated as:

421 
$$Q_{tot}^a = (1 - e^{-k_r})(S_r^f + Q_{tot}^f + \Delta Q_{Is} + \Delta Q_{I0}),$$

422 where  $k_r$  is a routing delay factor.

423 The surface water storage  $S_r$  should be updated accordingly as:

424 
$$S_r^a = S_r^f + \Delta Q_{Is} + \Delta Q_{I0} - \Delta Q_{tot}.$$

425 The total evapotranspiration change ( $\Delta E_{tot}$ ) caused by the changes in  $S_0$  and  $S_s$  can be updated as follow:

426 
$$\Delta E_{tot} = \delta E_s * \Delta S_0 + \delta E_t * \Delta S_s,$$

427 where the  $E_s$  is the evaporation flux from the surface soil store ( $S_0$ ) and  $E_t$  is the total actual plant transpiration. The term  $\delta E_s$   
428 is given as

429  $\delta E_s = (1 - f_{sat})E_{t\_rem}\delta f_{soile},$

430 where  $f_{soile}$  is relative soil evaporation and  $f_{sat}$  is the fraction of the grid cell that is saturated, and

431  $E_{t\_rem} = E_0 - (E_t - \delta E_t),$

432 The term  $\delta E_t$  is from the changes in root-water uptake from shallow and deep soil layers as

433  $\delta E_t = \delta U_s + \delta U_d,$

434 with

435  $\delta U_s = \delta U_{smax} \frac{\max(abs(\delta U_{smax}, \delta U_{dmax}))}{\delta U_{smax} + \delta U_{dmax}}$

436  $\delta U_d = \delta U_{dmax} \frac{\max(abs(\delta U_{smax}, \delta U_{dmax}))}{\delta U_{smax} + \delta U_{dmax}}$

437  $\delta U_{smax} = \frac{U_{s0}}{w_{stim}}\delta w_s, \delta U_{dmax} = \frac{U_{d0}}{w_{dlim}}\delta w_d,$  where  $U_{smax}$  and  $U_{dmax}$  are the maximum root water uptake from the shallow soil

438 store and from deep soil store.  $w_{stim}$  and  $w_{dlim}$  is the water-limiting relative water content from the *shallow and deep* soil

439 layer.

440 Finally,

441  $\delta f_{soile} = \frac{f_{soilmax}}{w_{olim}}\delta w_0,$  where  $f_{soilmax}$  is the scaling factor corresponding to unlimited soil water supply, with

442  $\delta w_0 = \frac{1}{s_{0max}}, \delta w_s = \frac{1}{s_{smax}},$  and  $\delta w_d = \frac{1}{s_{dmax}},$

443 where the  $w_z$  is the relative soil wetness of layer  $z,$  *i.e.* either 0, s or d.

444

## 445 **Data Availability**

446 The AWRA-CMS code is accessible from github ([https://github.com/awracms/awra\\_cms](https://github.com/awracms/awra_cms)). SMAP product used here is the

447 level-2 enhanced radiometer half-orbit 9-km EASE-grid soil moisture from the US National Snow and Ice Data Center

448 (<https://nsidc.org>). SMOS level-2 soil moisture product is available from ESA's SMOS online dissemination service

449 (<https://smos-diss.esa.int/oads/access/>). The MYD13C2 EVI data is accessible through Land Processes Distributed Active

450 Archive Centre (<https://lpdaac.usgs.gov>). The National Dynamic Land Cover Dataset of Australia is available from Geoscience

451 Australia (<https://www.ga.gov.au>).

## 452 **Author contribution**

453 ST developed and led the implementation of the method in AWRA-CMS. ST led the writing of the manuscript and graphics

454 creation. LR co-wrote project plan and co-developed the method. LR guided the application and evaluation. LR contributed to

455 manuscript writing. RP facilitated sharing of data feeds; coordinated the transition of method to operational implementation;

456 editing and review of manuscript. JL guided the selection of streamflow observation and reviewed the manuscript. WS guided

457 the modification to AWRA-CMS and reviewed the manuscript. CD co-wrote project plan and reviewed the manuscript.

## 458 **Competing interests**

459 The authors declare that they have no conflict of interest.

## 460 **Acknowledgements**

461 This project is supported by collaborative research agreement between the Australian Bureau of Meteorology and Australian  
462 National University. We would like to thank Stuart Baron-Hay from the Bureau of Meteorology for his help with  
463 implementation of the in AWRA-CMS. This research was undertaken with the assistance of resources and services from the  
464 National Computational Infrastructure (NCI), which is supported by the Australian Government through the National  
465 Collaborative Research Infrastructure Strategy.

## 466 **References**

- 467 Alemohammad, S. H., McColl, K. A., Konings, A. G., Entekhabi, D., and Stoffelen, A.: Characterization of precipitation  
468 product errors across the United States using multiplicative triple collocation, *Hydrol. Earth Syst. Sci.*, 19, 3489-3503,  
469 10.5194/hess-19-3489-2015, 2015.
- 470 Alvarez-Garreton, C., Ryu, D., Western, A. W., Su, C. H., Crow, W. T., Robertson, D. E., and Leahy, C.: Improving operational  
471 flood ensemble prediction by the assimilation of satellite soil moisture: comparison between lumped and semi-distributed  
472 schemes, *Hydrology and Earth System Sciences*, 19, 1659-1676, 10.5194/hess-19-1659-2015, 2015.
- 473 Anderson, J., Hoar, T., Raeder, K., Liu, H., Collins, N., Torn, R., and Avellano, A.: The data assimilation research testbed: A  
474 community facility, *Bulletin of the American Meteorological Society*, 90, 1283-1296, 2009.
- 475 Baldwin, D., Manfreda, S., Keller, K., and Smithwick, E.: Predicting root zone soil moisture with soil properties and satellite  
476 near-surface moisture data across the conterminous United States, *J Hydrol*, 546, 393-404, 2017.
- 477 Blankenship, C. B., Case, J. L., Crosson, W. L., and Zavodsky, B. T.: Correction of forcing-related spatial artifacts in a land  
478 surface model by satellite soil moisture data assimilation, *IEEE Geoscience and Remote Sensing Letters*, 15, 498-502, 2018.
- 479 Bolten, J. D., Crow, W. T., Zhan, X., Jackson, T. J., and Reynolds, C. A.: Evaluating the utility of remotely sensed soil moisture  
480 retrievals for operational agricultural drought monitoring, *Ieee J-Stars*, 3, 57-66, 2009.
- 481 Bouttier, F., and Courtier, P.: Data assimilation concepts and methods March 1999, Meteorological training course lecture  
482 series. ECMWF, 718, 59, 2002.
- 483 Carrera, M. L., Bilodeau, B., Bélair, S., Abrahamowicz, M., Russell, A., and Wang, X.: Assimilation of passive L-band  
484 microwave brightness temperatures in the Canadian land data assimilation system: Impacts on short-range warm season  
485 numerical weather prediction, *Journal of Hydrometeorology*, 20, 1053-1079, 2019.
- 486 Chan, S. K., Bindlish, R., O'Neill, P., Jackson, T., Njoku, E., Dunbar, S., Chaubell, J., Piepmeier, J., Yueh, S., Entekhabi, D.,  
487 Colliander, A., Chen, F., Cosh, M. H., Caldwell, T., Walker, J., Berg, A., McNairn, H., Thibeault, M., Martinez-Fernandez, J.,  
488 Uldall, F., Seyfried, M., Bosch, D., Starks, P., Collins, C. H., Prueger, J., van der Velde, R., Asanuma, J., Palecki, M., Small,  
489 E. E., Zreda, M., Calvet, J., Crow, W. T., and Kerr, Y.: Development and assessment of the SMAP enhanced passive soil  
490 moisture product, *Remote Sensing of Environment*, 204, 931-941, 10.1016/j.rse.2017.08.025, 2018.
- 491 Chen, F., Crow, W. T., Bindlish, R., Colliander, A., Burgin, M. S., Asanuma, J., and Aida, K.: Global-scale evaluation of  
492 SMAP, SMOS and ASCAT soil moisture products using triple collocation, *Remote Sensing of Environment*, 214, 1-13, 2018.



493 Crow, W., and Van den Berg, M.: An improved approach for estimating observation and model error parameters in soil  
494 moisture data assimilation, *Water Resources Research*, 46, 2010.

495 Crow, W. T., and Ryu, D.: A new data assimilation approach for improving runoff prediction using remotely-sensed soil  
496 moisture retrievals, *Hydrology and Earth System Sciences*, 13, 1-16, DOI 10.5194/hess-13-1-2009, 2009.

497 Crow, W. T., and Yilmaz, M. T.: The auto-tuned land data assimilation system (ATLAS), *Water resources research*, 50, 371-  
498 385, 2014.

499 Davies, T., Cullen, M. J. P., Malcolm, A. J., Mawson, M. H., Staniforth, A., White, A. A., and Wood, N.: A new dynamical  
500 core for the Met Office's global and regional modelling of the atmosphere, *Q J Roy Meteor Soc*, 131, 1759-1782,  
501 10.1256/qj.04.101, 2005.

502 De Lannoy, G. J., and Reichle, R. H.: Global assimilation of multiangle and multipolarization SMOS brightness temperature  
503 observations into the GEOS-5 catchment land surface model for soil moisture estimation, *Journal of Hydrometeorology*, 17,  
504 669-691, 2016.

505 de Rosnay, P., Drusch, M., Vasiljevic, D., Balsamo, G., Albergel, C., and Isaksen, L.: A simplified Extended Kalman Filter  
506 for the global operational soil moisture analysis at ECMWF, *Q J Roy Meteor Soc*, 139, 1199-1213, 10.1002/qj.2023, 2013.

507 Dharssi, I., Bovis, K. J., Macpherson, B., and Jones, C. P.: Operational assimilation of ASCAT surface soil wetness at the Met  
508 Office, *Hydrology and Earth System Sciences*, 15, 2729-2746, 10.5194/hess-15-2729-2011, 2011.

509 Donohue, R. J., Roderick, M. L., and McVicar, T. R.: Roots, storms and soil pores: Incorporating key ecohydrological  
510 processes into Budyko's hydrological model, *J Hydrol*, 436, 35-50, 2012.

511 Dorigo, W., Wagner, W., Albergel, C., Albrecht, F., Balsamo, G., Brocca, L., Chung, D., Ertl, M., Forkel, M., and Gruber, A.:  
512 ESA CCI Soil Moisture for improved Earth system understanding: State-of-the art and future directions, *Remote Sensing of  
513 Environment*, 203, 185-215, 2017.

514 Draper, C. S., Walker, J. P., Steinle, P. J., De Jeu, R. A., and Holmes, T. R.: An evaluation of AMSR-E derived soil moisture  
515 over Australia, *Remote Sensing of Environment*, 113, 703-710, 2009.

516 Draper, C. S., Reichle, R. H., De Lannoy, G. J. M., and Liu, Q.: Assimilation of passive and active microwave soil moisture  
517 retrievals, *Geophysical Research Letters*, 39, Artn L04401  
518 10.1029/2011gl050655, 2012.

519 Entekhabi, D., Njoku, E. G., O'Neill, P. E., Kellogg, K. H., Crow, W. T., Edelstein, W. N., Entin, J. K., Goodman, S. D.,  
520 Jackson, T. J., Johnson, J., Kimball, J., Piepmeier, J. R., Koster, R. D., Martin, N., McDonald, K. C., Moghaddam, M., Moran,  
521 S., Reichle, R., Shi, J. C., Spencer, M. W., Thurman, S. W., Tsang, L., and Van Zyl, J.: The Soil Moisture Active Passive  
522 (SMAP) Mission, *P IEEE*, 98, 704-716, 10.1109/Jproc.2010.2043918, 2010.

523 Errico, R. M.: What is an adjoint model?, *Bulletin of the American Meteorological Society*, 78, 2577-2592, 1997.

524 Figueroa-Bustos, V., Palta, J. A., Chen, Y., and Siddique, K. H.: Characterization of root and shoot traits in wheat cultivars  
525 with putative differences in root system size, *Agronomy*, 8, 109, 2018.

526 Getirana, A., Jung, H. C., Arsenault, K., Shukla, S., Kumar, S., Peters-Lidard, C., Maigari, I., and Mamane, B.: Satellite  
527 Gravimetry Improves Seasonal Streamflow Forecast Initialization in Africa, *Water Resources Research*, 56, ARTN  
528 e2019WR026259  
529 10.1029/2019WR026259, 2020a.

530 Getirana, A., Rodell, M., Kumar, S., Beaudoin, H. K., Arsenault, K., Zaitchik, B., Save, H., and Bettadpur, S.: GRACE  
531 Improves Seasonal Groundwater Forecast Initialization over the United States, *Journal of Hydrometeorology*, 21, 59-71,  
532 10.1175/Jhm-D-19-0096.1, 2020b.

- 533 Giering, R.: Tangent linear and adjoint biogeochemical models, *GEOPHYSICAL MONOGRAPH-AMERICAN*  
534 *GEOPHYSICAL UNION*, 114, 33-48, 2000.
- 535 Giroto, M., De Lannoy, G. J., Reichle, R. H., Rodell, M., Draper, C., Bhanja, S. N., and Mukherjee, A.: Benefits and pitfalls  
536 of GRACE data assimilation: A case study of terrestrial water storage depletion in India, *Geophysical research letters*, 44,  
537 4107-4115, 2017.
- 538 Hafeez, F., Frost, A., Vaze, J., Dutta, D., Smith, A., and Elmahdi, A.: A new integrated continental hydrological simulation  
539 system, *Water J. Aust. Water Assoc*, 42, 75-82, 2015.
- 540 Hawdon, A., McJannet, D., and Wallace, J.: Calibration and correction procedures for cosmic-ray neutron soil moisture probes  
541 located across Australia, *Water Resources Research*, 50, 5029-5043, 10.1002/2013wr015138, 2014.
- 542 Incerti, M., and O'Leary, G.: Rooting depth of wheat in the Victorian Mallee, *Australian Journal of Experimental Agriculture*,  
543 30, 817-824, 1990.
- 544 Ines, A. V. M., Das, N. N., Hansen, J. W., and Njoku, E. G.: Assimilation of remotely sensed soil moisture and vegetation  
545 with a crop simulation model for maize yield prediction, *Remote Sensing of Environment*, 138, 149-164,  
546 10.1016/j.rse.2013.07.018, 2013.
- 547 Jones, D. A., Wang, W., and Fawcett, R.: High-quality spatial climate data-sets for Australia, *Aust Meteorol Ocean*, 58, 233,  
548 2009.
- 549 Kerr, Y. H., Waldteufel, P., Wigneron, J. P., Martinuzzi, J. M., Font, J., and Berger, M.: Soil moisture retrieval from space:  
550 The Soil Moisture and Ocean Salinity (SMOS) mission, *Ieee T Geosci Remote*, 39, 1729-1735, Doi 10.1109/36.942551, 2001.
- 551 Kumar, S. V., Reichle, R. H., Koster, R. D., Crow, W. T., and Peters-Lidard, C. D.: Role of Subsurface Physics in the  
552 Assimilation of Surface Soil Moisture Observations, *Journal of Hydrometeorology*, 10, 1534-1547, 10.1175/2009jhm1134.1,  
553 2009.
- 554 Li, B., Toll, D., Zhan, X., and Cosgrove, B.: Improving estimated soil moisture fields through assimilation of AMSR-E soil  
555 moisture retrievals with an ensemble Kalman filter and a mass conservation constraint, *Hydrology and Earth System Sciences*,  
556 16, 105-119, 10.5194/hess-16-105-2012, 2012.
- 557 Lymburner, L., Tan, P., McIntyre, A., Thankappan, M., Sixsmith, J. : Dynamic Land Cover Dataset Version 2.1, Geoscience  
558 Australia, Canberra, <http://pid.geoscience.gov.au/dataset/ga/83868>, 2015.
- 559 Massari, C., Crow, W., and Brocca, L.: An assessment of the performance of global rainfall estimates without ground-based  
560 observations, *Hydrology and earth system sciences*, 21, 4347, 2017.
- 561 McColl, K. A., Vogelzang, J., Konings, A. G., Entekhabi, D., Piles, M., and Stoffelen, A.: Extended triple collocation:  
562 Estimating errors and correlation coefficients with respect to an unknown target, *Geophysical research letters*, 41, 6229-6236,  
563 2014.
- 564 McVicar, T. R., Van Niel, T. G., Li, L. T., Roderick, M. L., Rayner, D. P., Ricciardulli, L., and Donohue, R. J.: Wind speed  
565 climatology and trends for Australia, 1975–2006: Capturing the stilling phenomenon and comparison with near-surface  
566 reanalysis output, *Geophysical Research Letters*, 35, 2008.
- 567 Muñoz-Sabater, J.: Incorporation of passive microwave brightness temperatures in the ECMWF soil moisture analysis, *Remote*  
568 *Sensing*, 7, 5758-5784, 2015.
- 569 Nguyen, H., Wheeler, M. C., Otkin, J. A., Cowan, T., Frost, A., and Stone, R.: Using the evaporative stress index to monitor  
570 flash drought in Australia, *Environmental Research Letters*, 14, 064016, 2019.
- 571 Njoku, E. G., and Entekhabi, D.: Passive microwave remote sensing of soil moisture, *J Hydrol*, 184, 101-129, Doi  
572 10.1016/0022-1694(95)02970-2, 1996.
- 573 Pan, M., and Wood, E. F.: Data assimilation for estimating the terrestrial water budget using a constrained ensemble Kalman  
574 filter, *Journal of Hydrometeorology*, 7, 534-547, Doi 10.1175/Jhm495.1, 2006.

575 Panciera, R., Walker, J. P., Jackson, T. J., Gray, D. A., Tanase, M. A., Ryu, D., Monerris, A., Yardley, H., Rüdiger, C., and  
576 Wu, X.: The soil moisture active passive experiments (SMAPEX): Toward soil moisture retrieval from the SMAP mission,  
577 *Ieee T Geosci Remote*, 52, 490-507, 2013.

578 Patil, A., and Ramsankaran, R.: Improving streamflow simulations and forecasting performance of SWAT model by  
579 assimilating remotely sensed soil moisture observations, *J Hydrol*, 555, 683-696, 2017.

580 Pauwels, V. R. N., Hoeben, R., Verhoest, N. E. C., and De Troch, F. P.: The importance of the spatial patterns of remotely  
581 sensed soil moisture in the improvement of discharge predictions for small-scale basins through data assimilation, *J Hydrol*,  
582 251, 88-102, Doi 10.1016/S0022-1694(01)00440-1, 2001.

583 Peters-Lidard, C. D., Kumar, S. V., Mocko, D. M., and Tian, Y.: Estimating evapotranspiration with land data assimilation  
584 systems, *Hydrol Process*, 25, 3979-3992, 2011.

585 Pipunic, R. C., Walker, J. P., and Western, A.: Assimilation of remotely sensed data for improved latent and sensible heat flux  
586 prediction: A comparative synthetic study, *Remote Sensing of Environment*, 112, 1295-1305, 10.1016/j.rse.2007.02.038, 2008.

587 Rahmoune, R., Ferrazzoli, P., Kerr, Y. H., and Richaume, P.: SMOS Level 2 Retrieval Algorithm Over Forests: Description  
588 and Generation of Global Maps, *Ieee J-Stars*, 6, 1430-1439, 10.1109/Jstars.2013.2256339, 2013.

589 Reichle, R. H., and Koster, R. D.: Global assimilation of satellite surface soil moisture retrievals into the NASA Catchment  
590 land surface model, *Geophysical Research Letters*, 32, Artn L02404  
591 10.1029/2004gl021700, 2005.

592 Reichle, R. H.: Data assimilation methods in the Earth sciences, *Advances in water resources*, 31, 1411-1418, 2008.

593 Renzullo, L. J., van Dijk, A. I. J. M., Perraud, J. M., Collins, D., Henderson, B., Jin, H., SmiAssimilation of a ERS scatterometer  
594 derived soil moisture index in the ECMWF numerical weather prediction systemth, A. B., and McJannet, D. L.: Continental  
595 satellite soil moisture data assimilation improves root-zone moisture analysis for water resources assessment, *J Hydrol*, 519,  
596 2747-2762, 10.1016/j.jhydrol.2014.08.008, 2014.

597 Scipal, K., Holmes, T., De Jeu, R., Naeimi, V., and Wagner, W.: A possible solution for the problem of estimating the error  
598 structure of global soil moisture data sets, *Geophysical Research Letters*, 35, 2008.

599 Sheffield, J., and Wood, E. F.: Characteristics of global and regional drought, 1950-2000: Analysis of soil moisture data from  
600 off-line simulation of the terrestrial hydrologic cycle, *J Geophys Res-Atmos*, 112, Artn D17115  
601 10.1029/2006jd008288, 2007.

602 Shokri, A., Walker, J. P., van Dijk, A. I., and Pauwels, V. R.: On the use of adaptive ensemble Kalman filtering to mitigate  
603 error misspecifications in GRACE data assimilation, *Water Resources Research*, 55, 7622-7637, 2019.

604 Smith, A. B., Walker, J. P., Western, A. W., Young, R. I., Ellett, K. M., Pipunic, R. C., Grayson, R. B., Siriwardena, L., Chiew,  
605 F. H. S., and Richter, H.: The Murrumbidgee soil moisture monitoring network data set, *Water Resources Research*, 48, Artn  
606 W07701  
607 10.1029/2012wr011976, 2012.

608 Stoffelen, A.: Toward the true near-surface wind speed: Error modeling and calibration using triple collocation, *Journal of*  
609 *geophysical research: oceans*, 103, 7755-7766, 1998.

610 Su, C. H., Ryu, D., Crow, W. T., and Western, A. W.: Beyond triple collocation: Applications to soil moisture monitoring,  
611 *Journal of Geophysical Research: Atmospheres*, 119, 6419-6439, 2014a.

612 Su, C. H., Ryu, D., Crow, W. T., and Western, A. W.: Beyond triple collocation: Applications to soil moisture monitoring, *J*  
613 *Geophys Res-Atmos*, 119, 6419-6439, 10.1002/2013jd021043, 2014b.

614 Tangdamrongsub, N., Han, S.-C., Yeo, I.-Y., Dong, J., Steele-Dunne, S. C., Willgoose, G., and Walker, J. P.: Multivariate data  
615 assimilation of GRACE, SMOS, SMAP measurements for improved regional soil moisture and groundwater storage estimates,  
616 *Advances in Water Resources*, 135, 103477, 2020.

617 Tian, S. Y., Tregoning, P., Renzullo, L. J., van Dijk, A. I. J. M., Walker, J. P., Pauwels, V. R. N., and Allgeyer, S.: Improved  
618 water balance component estimates through joint assimilation of GRACE water storage and SMOS soil moisture retrievals,  
619 *Water Resources Research*, 53, 1820-1840, 10.1002/2016wr019641, 2017.

620 Tian, S. Y., Renzullo, L. J., van Dijk, A. I. J. M., Tregoning, P., and Walker, J. P.: Global joint assimilation of GRACE and  
621 SMOS for improved estimation of root-zone soil moisture and vegetation response, *Hydrology and Earth System Sciences*,  
622 23, 1067-1081, 10.5194/hess-23-1067-2019, 2019a.

623 Tian, S. Y., Van Dijk, A. I. J. M., Tregoning, P., and Renzullo, L. J.: Forecasting dryland vegetation condition months in  
624 advance through satellite data assimilation, *Nat Commun*, 10, ARTN 469  
625 10.1038/s41467-019-08403-x, 2019b.

626 Van Dijk, A. I., Peña-Arancibia, J. L., Wood, E. F., Sheffield, J., and Beck, H. E.: Global analysis of seasonal streamflow  
627 predictability using an ensemble prediction system and observations from 6192 small catchments worldwide, *Water Resources*  
628 *Research*, 49, 2729-2746, 2013.

629 Van Dijk, A. I. J. M.: AWRA Technical Report 3, Landscape Model (version 0.5) Technical Description, WIRADA, CSIRO  
630 Water for a Healthy Country Flagship, Canberra, 2010.

631 van Dijk, A. I. J. M., and Renzullo, L. J.: Water resource monitoring systems and the role of satellite observations, *Hydrol.*  
632 *Earth Syst. Sci.*, 15, 39-55, 10.5194/hess-15-39-2011, 2011.

633 Walker, J. P., Willgoose, G. R., and Kalma, J. D.: One-dimensional soil moisture profile retrieval by assimilation of near-  
634 surface measurements: A simplified soil moisture model and field application, *Journal of Hydrometeorology*, 2, 356-373, 2001.

635 Wanders, N., Karssenber, D., de Roo, A., de Jong, S. M., and Bierkens, M. F. P.: The suitability of remotely sensed soil  
636 moisture for improving operational flood forecasting, *Hydrology and Earth System Sciences*, 18, 2343-2357, 10.5194/hess-  
637 18-2343-2014, 2014a.


638 Wanders, N., Karssenber, D., Roo, A. d., De Jong, S., and Bierkens, M.: The suitability of remotely sensed soil moisture for  
639 improving operational flood forecasting, *Hydrology and Earth System Sciences*, 18, 2343-2357, 2014b.

640 Yan, H., Zarekarizi, M., and Moradkhani, H.: Toward improving drought monitoring using the remotely sensed soil moisture  
641 assimilation: A parallel particle filtering framework, *Remote sensing of environment*, 216, 456-471, 2018.

642 Yilmaz, M. T., and Crow, W. T.: Evaluation of assumptions in soil moisture triple collocation analysis, *Journal of*  
643 *hydrometeorology*, 15, 1293-1302, 2014.


644 Zwieback, S., Dorigo, W., and Wagner, W.: Estimation of the temporal autocorrelation structure by the collocation technique  
645 with an emphasis on soil moisture studies, *Hydrological sciences journal*, 58, 1729-1747, 2013.

646  
647



649 **Figure 1:** Satellite soil moisture retrievals in model unit (mm) for (a) SMAP and (b) SMOS compared to (c) AWRA-L estimates of  
 650 surface soil water storage for 1 Jan 2019. (d) Locations of in-situ soil moisture monitoring networks (CosmOz, OzNet and OzFlux),  
 651 catchments for streamflow validation and grid cells classified as cropland. The rectangular inset map provides a zoomed view into  
 652 the OzNet network region in south eastern Australia.


653



654


655 **Figure 2: Gain weights for sequential data assimilation derived from Triple Collocation (TC) showing the relative contribution of**  
 656 **the respective estimate in (a) AWRA-simulated surface soil water storage  $S_0$ , (b) SMOS soil moisture, and (c) SMAP soil moisture.**

657



658

659 **Figure 3: Time series of AWRA-L surface soil water storage estimates from open-loop (OL) compared to estimates after data**  
 660 **assimilation (DA-TC) of SMAP and SMOS soil moisture retrievals at CosmOz monitoring site: Bennets (35.826°E, 143.004°S). Note**  
 661 **that the in-situ soil moisture values are in volumetric unit.**




662

663

664

665 **Figure 4: Comparison of daily average surface soil water storage estimates ( $S_0$ ) for December 2019 from (a) model open-loop (OL),**


666 **(b) joint assimilation of SMAP and SMOS with Triple Collocation (DA-TC) and (c) difference between estimates DA-TC and OL.**



667


668 **Figure 5: Analysis increments of AWRA-L surface soil water storage ( $x_{50}^a - x_{50}^f$ ) in comparison with difference between in-situ rainfall**  
 669 **observations and rainfall forcing from AWAP used in AWRA-L modelling ( $p^{OzFlux} - p^{AWAP}$ ) for (a) Yanco site (34.989°E,**  
 670 **146.291°S) and (b) Wombat Forest (37.422°E, 144.094°S).**





671  
 672 **Figure 6: Averaged estimates of (a) shallow layer (10-100cm) soil water storage ( $S_s$ ), (b) evapotranspiration ( $E_{tot}$ ), and (c) total**  
 673 **streamflow ( $Q_{tot}$ ) for December 2019 from model open-loop, data assimilation (DA-TC), and after the analysis increments**  
 674 **redistribution (DA-TCAIR).**

675  
 676



677

678 **Figure 7: Distribution of correlation statistics of AWRA-L water balance estimates against in-situ measurements of (a) surface soil**  
 679 **moisture, (b) root-zone soil moisture, (c) evapotranspiration and (d) streamflow.**

680

681

682

683

684

685


686

687

688

689


690



691

692 **Figure 8:** Changes in streamflow  $Q_{tot}$  estimates after the analysis increments redistribution (DA-TCAIR) for a catchment in south-  
 693 eastern Australia (centre coordinates: 36.63°E, 147.43°S) compared to in-situ streamflow observations ( $Q_{tot}$  obs) and model open-  
 694 loop.


695



696

697 **Figure 9:** Comparison of vegetation index, EVI, with modelled root-zone soil moisture over cropland: (a) changes in correlations  
 698 after data assimilation (DA-TCAIR,  $r^{air}$ ) compared to model OL ( $r^o$ ); (b) changes in correlations between DA-TCAIR and DA-TC.

699



700  
 701 **Figure 10: Quantified impacts of data assimilation on forecasting AWRA-L state variables using the initial condition from DA-**  
 702 **TCAIR: average time period that the impact of data assimilation can persist in autumn (2018.03-2018.05), Winter (2018.06-2018.08),**  
 703 **Spring (2018.09-2018.11) and Summer (2018.12-2019.02) on (a) upper-layer soil water storage  $S_0$ , (b) lower-layer soil water storage**  
 704  **$S_s$ , (c) total evapotranspiration  $E_{tot}$  and (d) total runoff  $Q_{tot}$ .**