

Aquifer recharge in the Piedmont Alpine zone: Historical trends and future scenarios

Elisa Brussolo¹, Elisa Palazzi^{2,3}, Jost von Hardenberg^{4,2}, Giulio Masetti⁵, Gianna Vivaldo⁵, Maurizio Previati⁶, Davide Canone⁶, Davide Gisolo⁶, Ivan Bevilacqua⁶, Antonello Provenzale⁵, and Stefano Ferraris⁶

¹Research Center, Società Metropolitana Acque Torino S.p.A., Torino, Italy

²Institute of Atmospheric Sciences and Climate, National Research Council of Italy (CNR), Torino, Italy

³Department of Physics, Università di Torino, Torino, Italy

⁴Department of Environment, Land and Infrastructure Engineering (DIATI), Politecnico di Torino, Torino, Italy

⁵Institute of Geosciences and Earth Resources, National Research Council of Italy (CNR), Pisa, Italy

⁶Interuniversity Department of Regional and Urban Studies and Planning (DIST), Politecnico di Torino and Università di Torino, Torino, Italy

Correspondence: Elisa Palazzi (e.palazzi@isac.cnr.it)

Abstract. The spatial and temporal variability of air temperature, precipitation, actual evapotranspiration (*AET*) and of the related water balance components, as well as their responses to anthropogenic climate change, provide fundamental information for an effective management of water resources and for a proactive involvement of users and stakeholders, in order to develop and apply adaptation and mitigation strategies at the local level.

- 5 In this study, using an interdisciplinary research approach tailored to water management needs, we evaluate the past, present and future quantity of water potentially available for drinking supply in the water catchments feeding the about 2.3 million inhabitants of the Turin Metropolitan Area (the former Province of Turin, North-Western Italy), considering climatologies at the quarterly and yearly timescales. Observed daily maximum surface air temperature and precipitation data from 1959 to 2017 were analyzed to assess historical trends, their significance and the possible cross-correlations between the water
- 10 balance components. Regional climate model (RCM) simulations from a small ensemble were analysed to provide mid-century projections of the difference between precipitation and *AET* for the area of interest in the future CMIP5 scenarios RCP4.5 (stabilization) and RCP8.5 (business as usual). Temporal and spatial variations in recharge were approximated with variations of drainage. The impact of irrigation, and of snowpack variability, on the latter were also assessed. The other terms of water balance were disregarded because they are affected by higher uncertainty.
- 15 The analysis over the historical period indicated that the driest area of the study region displayed significant negative annual (and spring) trends of both precipitation and drainage. Results from field experiments were used to model irrigation and we found that relatively wetter watersheds in the northern and in the southern parts behave differently, with significant increase of *AET* due to irrigation. The analysis of future projections suggested almost stationary conditions for annual data. Regarding quarterly data, a slight decrease in summer drainage was found in 3 out of 5 models in both emission scenarios. The RCM
- 20 ensemble exhibits a large spread in the representation of the future drainage trends. The large interannual variability of precip-

itation was **also quantified, and identified** as a relevant risk factor for water management, expected to play a major role also in future decades.

Copyright statement. TEXT

1 Introduction

25 Water is a crucial resource, intrinsically linked to society and culture development, food and energy security, well-being, environmental sustainability and poverty reduction. However, several factors, including urbanization, population growth, land use and soil consumption, industrial and agricultural development, endanger water resource sustainability in terms of availability, quality, management and demand (IPCC, 2014; WWAP, 2015). Groundwater resources represent about 97% of liquid freshwater resources on Earth (WHO, 2006; Healy, 2010) and play a key role in water supply and proper ecosystems preservation (WWAP, 2015). Groundwater resources help to maintain river discharges and, together with surface freshwaters, are accounted for in water budget considerations at the river basin scale (Rumsey et al., 2015). **The hydrological connection between groundwater and surface water is primarily controlled by: (1) the driving force generated by the hydraulic gradient between groundwater and surface water, (2) the permeability degree of the aquifer in comparison to a streambed (i.e. different hydraulic conductivity) due to the geological context (Lasagna et al., 2016; Epting et al., 2018). Groundwater and surface water interaction is influenced by both local and regional regimes (Epting et al., 2018). Local interaction could be very complex and different methods were developed to quantify this interaction in different locations (Bertrand et al., 2014; Kalbus et al., 2006).** Groundwater resources are of utmost importance for their mitigation effects during dry periods and their reduction can impact the whole hydrological cycle. Groundwater is a fundamental natural resource that acts as a reservoir from which good quality water can be collected for drinking purposes, requiring few purifying treatments compared to surface water. Climate change influences several components of the water cycle, including groundwater resources, causing a lowering of piezometric levels due to discharge modifications as a result of snow retention reduction, changes in precipitation regimes and potential evapotranspiration **that increases with increasing temperatures. In Alpine regions, shorter and thinner snow pack will decrease late spring flows, while the air temperature rise will increase stream flows in fall and winter due to trading of snowfall for rainfall (Confortola et al., 2013), leading to a shift of groundwater recharge from summer to winter, as evaluated by CH2018-Project-Team (2018) and Epting et al. (2021) in Switzerland.**

Surface water and pollutants infiltration together with over-exploitation of wells can further deplete groundwater resources, triggering the competition between irrigation and potable uses. While the degradation of water quality mostly depends on land use and saltwater intrusions into coastal groundwater (Jiménez Cisneros et al., 2014), also **climate change may affect, either directly or indirectly, the quality of groundwater resources. Temperature impacts on biological, chemical and physical properties of groundwater resources (Epting et al., 2021), even if the increase of air temperature is not necessary directly correlated with groundwater temperature increase. In fact, this correlation depends on the intrinsic properties of aquifers,**

on local and regional spatiotemporal scales and on different anthropic inputs (Epting et al., 2021; Bastiancich et al., 2021). Moreover, the interaction between surface water and groundwater flow systems influences the water chemistry (Lasagna et al., 2016), since in areas where rainfall intensity is expected to increase pollutants will be increasingly washed from soils to water bodies (Parry et al., 2007). Finally, water-level changes are a key indicator that flow patterns are changing and that low-quality water may be mobilized (Moench et al., 2003).

Climate change also exacerbates the risks associated with changes in the distribution and availability of water resources (Jiménez Cisneros et al., 2014), with consequences on water-demand management and infrastructural system planning (van der Gun, 2012). In this framework, assessing climate change impacts on Integrated Urban Water Management (IUWM), considering a worsening of preexisting conditions and/or an occurrence of new hazards or risk factors, and planning climate change adaptation strategies are fundamental challenges that IUWM is expected to face in the near future, using an integrated approach based on prevention, preparedness and risk assessment. To this end, an appropriate proactive engagement of stakeholders is needed, in order to develop and apply adaptation and mitigation strategies quantified and driven by state-of-the art climate information and projections.

The Alps and the Mediterranean area are recognised as two climate hot-spot regions (IPCC, 2014), showing amplified climate change signals and associated environmental, social and economical impacts. Future projections for the Italian territory, in particular, show an increase of high precipitation intensity typically distributed in more intermittent events, together with an increase of the duration of dry periods (Desiato et al., 2015). However, in northwestern Italy these trends are not clear and they are very site-dependent (Baiamonte et al., 2019). In a recent paper analyzing European floods, in Piedmont almost no precipitation trend emerged, with the exception of the Dora Riparia Valley, where precipitation is declining (Blöschl et al., 2019). Another recent work shows the differences in freshwater trends between Central and Southern Europe (Gudmundsson et al., 2017). In fact Italy is a climatic bridge between the Mediterranean and the inland European climate (Libertino et al., 2019) and in Piedmont there are 4000-meter-high mountains at just 160 km distance from the Mediterranean sea.

Northern Italy is also a bridge between areas where actual evapotranspiration is mainly soil-moisture-limited (Mediterranean) and areas where it is energy-limited (Central Europe).

For precipitation, the transient development of the impingement of cold fronts on the Alps induces a wide range of mesoscale phenomena. On one hand, when the Alpine chain is subject to a southerly flow of moist and relatively warm air from the Mediterranean Sea, very intense precipitation episodes can take place, such as the Piedmont flood in November 1994. The Piedmont lowlands act as a natural trap for moist airflows from the southeast, particularly in Autumn. On the other hand, northerly flows lead to dry weather, particularly in Winter (Pradier et al., 2002). The main part of relevant rainfall and snowfall episodes, including extreme events, are thus due to southerly flow.

For water management, the IUWM in Italy is geographically organized into local districts (called “ATO”– Ambiti Territoriali Ottimali, which translates into “Optimal Territorial Divisions”) whose domains were defined based on various criteria, either administrative boundaries or physical ones, including river basin boundaries (Legislative Decree No 152/2006, as further amended). The boundaries of these districts mostly coincide with administrative borders; in the Piedmont region, the Turin Metropolitan Area represents local district ATO3, where the IUWM is provided by Società Metropolitana Acque Torino

(SMAT). This is a wide and geographically complex area and SMAT exploits many and diverse supply sources, with ground-water resources representing about 80% of the whole water supply in terms of volume available to SMAT.

It is therefore important to evaluate the balance between precipitation and actual evapotranspiration (AET), and their related spatial and temporal variabilities. Several studies can be found in the literature which evaluate the impacts of climate change on groundwater resources, e.g.: Jiménez Cisneros et al. (2014) and Taylor et al. (2013).

Recharge varies in space and time and it is difficult to measure directly, therefore a comprehensive understanding is lacking. In this paper we define recharge as the difference between precipitation and actual evapotranspiration. This simplification follows a recent review paper, where mesoscale is defined as the ideal scale for simulating the effect of climate change on recharge and precipitation is mentioned as the largest source of recharge variability (Smerdon, 2017). Moreover, in a recent study, considering several wells located in nine regions of the central and northeastern U.S., the recharge (evaluated as drainage from the lowest model soil layer) was shown to be compatible with observed monthly groundwater storage anomalies and month-to-month changes in groundwater storage (Li et al., 2015). In the selected study area, a recent study commissioned by the Water Department of the Piedmont Region shows that the recharge areas of the deep aquifers occur in the high plain sectors, close to the Alps (Regione Piemonte, 2018b; De Luca et al., 2020), justifying our choice to avoid river-fed recharge. A new global dataset encompassing more than 5000 locations has shown that precipitation amounts and seasonality of temperature and precipitation are the most important variables (Moeck et al., 2020; Condon et al., 2020). Epting et al. (2021), in a recent paper on Swiss alluvial aquifers, underline the importance of both spatial and temporal variability in recharge related studies carried out in a region close to our study area. Regarding spatial variability, Pangle et al. (2014) identified drainage as a proxy of recharge in a controlled mesocosm. Their results highlighted the potential for local interactions between temperature, vegetation, and soils to moderate the hydrological response to climate warming. The importance of soil and vegetation was also underlined by Condon et al. (2020). The outputs of Pangle et al. (2014) study show that AET decreased in summer because of soil moisture shortage. This result is due to the fact that precipitation is out of phase with the growing season cycle, and irrigation is operated only for reproducing natural rainfall events. At the yearly timescale, they did not find a reduction of AET in their evaluation of trends.

Regarding temporal variability, interannual variability also plays a major role in groundwater recharge, and it is critical for water managers. Masbruch et al. (2016) have shown how quasi-decadal large groundwater recharge events can be important for replenishing the aquifers. These events are characterised by large precipitation (both rainfall and snow water equivalent) and by below-average seasonal temperatures.

Also, the effects of climate change are not all in the same direction. In a study using 16 GCMs, a considerable uncertainty in both the magnitude and direction of recharge changes was shown for 2050 year projections in different parts of the High Plains (Crosbie et al., 2013). A more recent study has revealed variability in both direction and magnitude of hydrological changes for the Great Lake basin of North America, with a combination of different RCMs (Persaud et al., 2020).

Konapala et al. (2020) reported at the global scale an increase in annual mean evaporation over the land surface, attributed to the increase in temperature. They deal with water availability as the difference between precipitation and actual evapotranspiration. In this paper we call this quantity drainage, as a proxy of recharge, computed by the soil model in each pixel, without

modeling both runoff and the underlying aquifer flow. The other terms of water balance have more uncertainty and probably lower impact: soil water storage is small, river runoff has scattered measurements and complex process modelling, subsurface flow always shows large uncertainties associated with its estimation (Healy, 2010).

125 In this paper we develop a stakeholder-driven interdisciplinary research study, where scientists in atmospheric/climate research and hydrologists work together with agricultural and soil scientists and experts from a water utility to quantify the role of groundwater focusing on an area providing water to about 2.3 million people. This area is characterized by a very large spatial precipitation and air temperature variability, owing to the proximity of high mountains and of the sea.

The main research questions, relevant also for other regions worldwide, are: a) Can the water balance show a significant trend only because of a significant trend in *AET*? Is also a significant trend in precipitation necessary? b) How different are water balance trends in three different parts of the area, namely a drier west-east oriented mountain area, a wetter mountain area, a mostly irrigated agricultural area? In fact in about 7000 square kilometers there are quite different situations, the former two with an impact of snow versus rain and anticipated snowmelt trends, and the latter with a *AET* temperature increase compensation with irrigation; c) To what extent are the spatial variability and trends observed in the past 60 years expected to undergo changes during the next 30 years?

130 We evaluate the temporal variability and the trends of the water balance terms, estimating the quantity of groundwater resource available for drinking water supply in the water catchments of the area managed by SMAT. The analyses are performed at both the quarterly and the hydrological year timescale, for both past and future conditions, analyzing a historical climate dataset for the period 1959–2017 and future projections from regional climate model RCA4 RCM simulations up to 2050, in order to be compliant with relatively short-term water management objectives.

140 The expected result will form a knowledge basis for operational indications and will be useful to characterize future groundwater resources availability in other border areas between Mediterranean and continental climate, especially where the water resource is subject to multiple anthropogenic pressures. This paper represents a scientific contribution to the management and the governance of water resources and water supply that could be applied worldwide, through e.g. the implementation of scientifically-driven guidelines and strategic agendas on water supply and water policies.

2 Study area and methodology

2.1 Study area

Many foothill zones in the Alps and Apennines contain aquifer systems of strategic interest for water supply, especially for drinking purposes (Doveri et al., 2016). In this framework, the aquifer system extending in the foothill plain located between the western Alps and the Turin hills represents one of the most significant and studied groundwater bodies in the Piedmont region (Raco et al., 2021).

This study area, within the administrative borders of the Turin Metropolitan Area (encompassing the entire territory of the former Province of Turin), has a complex orography and is surrounded on the western and northern sides by the Alps (with elevation peaks higher than 4000 meters above sea level at the border with the Valle d'Aosta region) and on the eastern




Figure 1. The study area including river catchments and sub-catchments. **Illustration of the water balance terms at the catchment scale is shown in a bottom right box.** Topographic shading is based on DEM data from the “Progetto Risknat - Base topografica transfrontaliera, ARPA Piemonte” (http://webgis.arpa.piemonte.it/ags101free/rest/services/topografia_dati_di_base/Sfumo_Europa_WM/MapServer).

155 and southern sides by hills and plains. Precipitation in the study area is characterized by relatively high spatial and temporal variability (yearly total ranging from about 500 mm in the plain to 2000 mm for high elevation gauges). The study area, in fact, is prone to topographically-induced precipitation and is exposed to the inflow of moisture-rich air from the Mediterranean sea (Cicarelli et al., 2008). It is also an area characterized by the occurrence of relatively long dry periods, up to 6.4 days on average during winter (Agnese et al., 2012; Baiamonte et al., 2019).

160 Owing to its hydrogeological features (see also the rivers system shown in Fig. 1), the foothill aquifer systems generally sustain the infiltration of both local rainfall and streamwater originating from mountain catchments. These systems are highly sensitive to variations in meteorological variables, first of all in precipitation regimes.

2.2 Water balance terms

In this paper we refer to water balance as the balance between water inputs and outputs at the catchment scale, as schematically shown in Figure 1.

165 Given the timescales of interest and the uncertainties inherent in some terms of the water balance equation, in this study the standard formulation of this equation has been simplified, including precipitation, actual evapotranspiration (AET), and drainage (obtained by subtracting AET from the sum of rainfall and snowmelt) which is used as a proxy of the groundwater recharge. This approach is common to other studies and many of them (Healy, 2010) do not make a distinction between drainage and groundwater recharge. This choice has also been discussed in the introduction. Also, in the study area measured riverflow data cover a temporal interval shorter than two decades (2000-2017), not allowing to build robust regressive models able to estimate deep percolation outside the time interval of data availability. Regarding the time step of calculation, the daily scale is recommended because of precipitation intermittency, in order to avoid the underestimation of drainage and recharge (e.g. Healy, 2010). Therefore in this work the computation is done at the daily scale in the soil column, and results are then aggregated at the yearly and quarterly time scale and, spatially, at the catchment scale.

175 The daily soil model will be described later in the text. The yearly catchment water balance can be simplified as follows (Healy, 2010):

$$P_q = AET + Q_{out} \quad (1)$$

where P_q represents the sum of liquid precipitation (rainfall) and snowmelt, AET the Actual Evapotranspiration and Q_{out} the drainage. Surface and subsurface catchments are assumed to be coincident as they are bounded by the border mountain divide (De Luca et al., 2020). Storage variations in time are disregarded, because the control volume is composed of vadose zone soil columns, with an aggregation of results at the quarterly and yearly time scale. The soil model presented in the following is used to calculate AET and Q_{out} in each pixel at the daily time scale.

To account for the different ground characteristics and variations, together with the physical description of the hydrological processes, all the water balance variables (except for output discharge) must be first evaluated at a fine spatial resolution and then aggregated (i.e., upscaled) at watershed level. To this end, a horizontal spatial resolution of 250 m was considered as a good compromise between the necessity of high-resolution and the computational resources required: the study area was discretised into a grid of 652 x 521 pixels.

185 The model takes into account the effects of irrigation on the value of AET . However, input and output irrigation terms in surface and groundwater balance are assumed to balance out at the catchment level. Also, industrial water withdrawals are not

190 changing the water balance at the watershed scale, because they give back the captured water at short distance, **being performed mostly for hydroelectric energy production.**

The contribution of glacier and permanent snow melting can be disregarded **because it is very small, in comparison** with the catchment **areas.** However the evaluation of snowmelt from the snowpack is quite important as it represents the amount of daily snow equivalent contributing to the water balance. **In fact, as outlined in the review by Taylor et al. (2013), at high altitudes,**
195 **increasing temperatures lead to less snow accumulation and earlier snowmelt, and to more winter precipitation in form of rain.**

To this end, as a first step, air temperature was used to distinguish between snowfall (P_s) and rainfall (P_r) (DeWalle and Rango, 2008): for temperatures lower or equal to -2°C all the precipitation amount can be considered as snowfall, while for temperature higher or equal to 5°C all the precipitation amount can be considered as rainfall. For temperatures between -2°C and 5°C , rainfall and snowfall amounts **were linearly interpolated, in order** to breakdown the different percentages of
200 precipitation type. The resulting snowfall P_s was used as an input to a simple bucket model in each pixel, used to integrate the storage of snow on ground, S . The daily snowmelt, N_f , was subtracted from the same bucket and it was evaluated using the method by Zeinivand and Smedt (2009), where the melted snow is a function of the difference between the daily mean temperature T_{mean} and melting temperature T_0 (0°C), and of the rainfall amount (P_r evaluated in mm/day), as follows:

$$N_f = \min \{S/(1 \text{ day}), \max [0, (k_{snow} + k_{rain} \cdot P_r) \cdot (T_{mean} - T_0)]\} \quad (2)$$

were k_{rain} (rainfall melt-rate factor) and k_{snow} (melt-rate factor) are fixed parameters. The same values as in Zeinivand and
205 Smedt (2009) **were used:** $k_{rain} = 0.0757^\circ\text{C}^{-1}$ and $k_{snow} = 3 \text{ mm day}^{-1} \cdot \text{C}^{-1}$.

A correct evaluation of the water balance allows a reliable estimation of water availability. To this end the following chain **was applied** in the study area:

1. identification and retrieval of the daily meteorological (i.e. temperature and precipitation) data;
2. regridding of the meteorological and hydrological data at the spatial resolution required by the hydrological model,
210 namely a square 250 m grid;
3. setting-up of a mathematical model that accounts for soil-vegetation-atmosphere interactions and providing actual evapotranspiration estimates (see Section 2.3).

Observational and reanalysis datasets have been used to obtain meteorological data for the period 1959-2017, as described in Section 3, while for the assessment of future water balance scenarios climate projections of temperature and precipitation **were**
215 **used** (see Section 4).

In order to be compliant with management objectives for strategic long-term planning, the input datasets, available at daily resolution (see Section 3), were aggregated to quarterly and yearly timescales using the definition of hydrological year (the first quarter includes January, February and March (*JFM*), the second April, May and June (*AMJ*), the third July, August and September (*JAS*) and the fourth October, November and December (*OND*)). The hydrological year N , starts on October 1st
220 of year $N - 1$ and ends on September 30th of year N . During the hydrological year a complete snowfall melting occurs, making

the breakdown of precipitation between rainfall and melted snow negligible. At high altitudes this melting is not complete and the annual variability entails the fluctuation of snow **on the ground** from one (hydrological) year to the following one. This phenomenon is limited to the highest peaks and is relevant only for limited areas. For this reason, for evaluations at the yearly timescale, P_q was assumed **equal to the** total average yearly precipitation.

225 2.3 Soil-water model

A simple bucket soil-water model was developed in order to estimate AET values at the daily scale for each $250\text{ m} \times 250\text{ m}$ pixel of the study area. The soil is schematised with seven different layers of increasing thickness with depth, similar to the FAO56 model (Allen et al., 1998). The water inputs for the model are precipitation (sum of rainfall and melted snow) and irrigation, if any. The outputs are AET and drainage (sum of runoff and deep percolation). The distinction between runoff and
230 deep percolation is not included in the model, as well as the modelling of capillary rise.

For each layer, the bucket equation is as follows:

$$P_q + I^* - Q_{out} - AET - \Delta S = 0 \quad (3)$$

where P_q is rainfall plus snowmelt; I^* is irrigation; Q_{out} is drainage; AET is actual evapotranspiration; ΔS is the variation of water storage in the soil per day. A bucket model (the one by Baudena et al. (2012), which was applied in the same area in northwestern Italy) water can flow upwards within each layer via evapotranspiration or evaporation, and downwards via deep
235 percolation, as a function of soil water content of each layer: if it exceeds field capacity, the water surplus percolates into the layer immediately below.

2.4 Actual Evapotranspiration computation

As reported by the Fifth Assessment Report of the IPCC (IPCC, 2014), a rise in GHG concentrations is associated with reduced soil moisture in Northern Hemisphere mid-latitude summers. This is the result of higher winter and spring evaporation, caused
240 by higher temperatures and reduced snow cover, and of lower rainfall inputs during summer. Regarding the effects on recharge of managed agrosystems, Taylor et al. (2013) state that changes in surface energy budgets are associated with enhanced soil moisture from irrigation. The widespread use of irrigation in most parts of the Po Valley plain cancels the dampening role of AET soil moisture limitation. At the same time, roughly half of the surface of the study area is covered by mountain grasslands and not-irrigated areas, where moisture limitation can play an important role.

245 The Actual Evapotranspiration AET was calculated starting from potential evapotranspiration PET , then reduced by considering the actual soil water content, obtained from the bucket soil model for each layer, with AET of each day being the sum of the depletions from the single layers. PET was calculated with the model of Hargreaves and Samani as in Allen et al. (1998), using daily maximum and minimum air temperature data from the Regional dataset described in Section 3.1 and extraterrestrial radiation modeled as in Aguilar et al. (2010). Also, the reduction of evapotranspiration (from PET to AET) due
250 to actual soil water content was modeled by using the coefficients Kc and Ks related to crop and soil respectively according to Allen et al. (1998), comparing the modeling results with real world measurements in different sites (Raffelli et al., 2017).

The soil water model reproduces very well the data in the case of no irrigation. However, a relevant contribution of irrigation, especially for highly water-demanding crops such as maize, can increase the *AET* term of the water balance. These water-demanding crops are quite widespread in the study area (about 750 square km) and therefore a novel procedure reproducing agricultural irrigation techniques was implemented in the soil-water model. This module is based on previous research in three farms (Canone et al., 2015, 2016), reproducing the farmers' decision rules and tuned in order to obtain irrigation events similar to the observed ones.

3 Observed data

3.1 Temperature and precipitation

The past and present precipitation and temperature data necessary to force the hydrological model employed in this study were derived from the Regional Environmental Protection Agency of Piedmont (ARPA Piemonte) databases. In particular they have been extracted from the OI dataset (in Italian, ARPA Piemonte, 2010a; Ciccarelli et al., 2008). This dataset provides cumulative daily precipitation, and maximum and minimum daily temperature at a spatial resolution of 0.125° longitude-latitude (corresponding to ~ 12 km) over the entire Piedmont region, for a time period extending from 1959 to 2017. The OI dataset was obtained by interpolation of in-situ observations collected by the Hydrographic Office network and by the network of the ARPA telemetry stations through the technique of the Optimal Interpolation, which allows to obtain data on a regular grid homogenizing observational data from different measurement networks and sources (ARPA Piemonte, 2010b). A preliminary quality check of the OI data revealed the existence of days (all referring to years before 1990) where the minimum temperature showed a higher value than the maximum temperature, probably owing to issues in the data acquisition. These data were excluded from the analysis and replaced by new values obtained through linearly interpolation in time, rather than in space, in order not to smooth the orographic information inherent in the original OI dataset.

The OI dataset provides a set of meteorological variables at a coarser spatial resolution than that required to describe the small scale hydrological processes and to accurately estimate the water balance terms, calling for the application of interpolation and downscaling techniques. After re-projecting the OI temperature and precipitation data into a WGS84/UTM 32N coordinate system useful for the subsequent analyses, they were further interpolated at the finer resolution of 250 m over the domain of interest ($x_{min} = 304250W$, $x_{max} = 434500W$, $y_{min} = 4951750N$, $y_{max} = 5094750N$). The conversion was preceded by a preliminary bilinear interpolation of the OI dataset, in order to produce an intermediate higher resolution dataset at a resolution of 0.001° latitude-longitude, and to avoid artifacts when re-projecting into UTM coordinates. Post-processing was performed using the GDAL (Geospatial Data Abstraction Library; v. 2.1) and CDO (Climate Data Operators; v. 1.7.0) softwares.

A further adjustment to account for orographic effects was applied to maximum and minimum temperature data, using the environmental lapse-rate correction coefficients derived for the Alpine region by Rolland (2003), shown in Table 1. Figure 2 shows an example of orographic correction (middle panel) applied to a maximum temperature field (left panel) for a selected day (January 1st, 2009). The orography of the study area is shown in the right panel, based on a 90 m resolution Digital

Table 1. Average climatological monthly adiabatic lapse-rate from Rolland (2003) for the Alpine region (expressed in K/km)

Jan	Feb	Mar	Apr	May	June	July	Aug	Sep	Oct	Nov	Dec
4.5	5.0	5.8	6.2	6.5	6.5	6.5	6.5	6.0	5.5	5.0	4.5




Figure 2. Maximum temperature field at (left) 0.125° lon-lat spatial resolution from the original OI dataset and at (middle) 250 m resolution after interpolation with orographic correction, for January 1st, 2009. The right panel shows the orography of the study area from a Digital Elevation Model derived from the SRTM project, interpolated at 250 m of spatial resolution.

Elevation Model derived from the SRTM (Shuttle Radar Topography Mission) project (Jarvis et al., 2008), interpolated at 250
 285 m.

3.2 Input data of the soil water model

The soil hydraulic properties have been estimated via *pedotransfer* (PTF) following Schaap et al. (2001) from the sand, clay and silt percentages taken from the Soil map of Regione Piemonte (scale 1:250000, IPLA, 2007). Computing via PTF the wilting point (*WP*) and field capacity (*FC*), the Total Available Water (*TAW*) is calculated for each layer as follows:

$$TAW = (FC - WP) \cdot (1 - r) \cdot L_d \quad (4)$$

290 where r is the fraction of volume occupied by stones, and L_d is the layer depth, which is obtained by dividing the root zone
 depth z by the number of layers (7, see Section 2.3). The root zone depth has been obtained using the land cover classes from
 the BDTRE database (Regione Piemonte, 2018a), listed in Table 2, converted into root zone depth using different root zone
 depths for each class, as listed in Table 3. The resulting root zone depth was compared with the soil depth provided by the
 Regione Piemonte Soil Map (IPLA, 2007), choosing the minimum between the two depths. For trees the water from the whole
 295 soil depth can be depleted and so soil depth has been used. Because of the high spatial resolution of BDTRE, the raster has

Table 2. Land cover classes

Class	ID Code
Winter crops	1
Summer crops-irrigated	2
Plain meadows (< 1000 m a.s.l.)	3
Orchards	4
Horticultural	5
Plain broadleaves (< 800 m a.s.l.)	6
Mountain broadleaves (> 800 m a.s.l.)	7
Coniferous	8
Vineyards	9
Mountain grassland (> 1000 m a.s.l.)	10
Bare soil	11
Bare rocks	12
Impervious surfaces	13
Water	14
Others	15
Glaciers and permanent snow	16

Table 3. Soil bucket depth as a function of land cover class. For trees (codes 4,6,7,8,9) the full soil depth from the Soil Map of Regione Piemonte is used.

ID Code	Soil use	Soil bucket depth [mm]
1 2	Irrigated crops	1000
3 5	Plain grassland and horticultures	500
10	Mountain grassland	250
11	Bare soil	150
12 13 14 16	Impermeable surfaces, water, glaciers	0

been produced with a 5 m × 5 m grid, then resampled at 250 m by the mode criterium. Homogenising data as a function of the evapotranspiration behavior of each class, the different land cover classes have been aggregated (Table 2).

300 Finally, considering that irrigation (here considered as water input) significantly modifies actual evapotranspiration, for quantifying the actually irrigated fields, we used the Regional Irrigation Information system (Regione Piemonte, 2016), the agricultural crop survey and the historical maps within the actually utilised agricultural areas survey database (Regione Piemonte, 2006), for the years 2013, 2014 and 2015. The actual quantity of irrigation water, and the number of irrigations, have been

evaluated using the measured data collected in experiments performed in real world farms both for surface irrigation (Canone et al., 2015) and for sprinklers (Canone et al., 2016).

305 **Even if runoff** was not considered in this study, following Kumar et al. (2016) it could be useful to quantify its variability with climate change. **As suggested by Epting et al. (2021), this should be taken into account where river-fed aquifers are considered.**

4 Future projections of precipitation and temperature

Reliable estimates of the hydrological response and of water availability in the coming decades generally require the implementation of a modelling chain consisting of global climate models (GCMs), which provide climate scenarios for the entire planet, regional climate models (RCMs) nested into global models providing lateral and boundary conditions for the regional simulation and, depending on the resolution which needs to be achieved, further downscaling procedures. At the end of the modelling chain, a hydrological model is thus forced with a high-resolution climatic input to simulate the hydrological response at the scale of interest. In this study, we used a small multi-member ensemble of the RCA4 RCM, (Strandberg et al., 2014) forced by five GCM simulations, able to provide the climatic variables of interest – surface air temperature and precipitation – at a spatial resolution of ~ 12.5 km. More detailed analyses of the interplay between several GCMs and RCMs (Sorland et al., 2018) are out of the operational water management scope of this paper. In this work we proceeded similarly to another recharge study (Allen et al., 2010) which used four GCMs and one RCM. They concluded that a range of GCMs should be considered for water management planning. The simulation outputs of this RCM from 1970 to 2050 have been analysed, considering two different emission scenarios for future projections among those defined by the IPCC (Intergovernmental Panel on Climate Change, IPCC (2013); Moss et al. (2010)), as described in section 4.1. The model data were subsequently interpolated applying the same procedure employed for the OI observational dataset (see section 3) and adjusted to correct the systematic bias which the model displays with respect to the OI reference climatology in a common time period (1986–2015, see section 4.3 for a description of the post-processing procedure).

4.1 RCA4 Regional Climate Model

325 **In the following, we used model simulations of the** RCA4 RCM (Strandberg et al., 2014) driven by five different GCMs, namely EC–Earth, CNRM–CM5, IPSL–CM5A–MR, HadGEM2–ES, MPI–ESM–LR, which provide lateral and boundary conditions for the regional simulation. Using one single RCM allowed us to obtain an ensemble of reasonably homogeneous simulations at the regional level, but representing at the same time model uncertainties in future projections captured by the different large-scale GCMs. RCA4 is one state-of-the-art RCM which participated to CORDEX, the Coordinated Regional Climate Downscaling Experiment (<http://wcrp-cordex.ipsl.jussieu.fr/>, Giorgi et al., 2009) sponsored by WRCP (World Climate Research Program) aimed at providing a global coordination of regional climate downscaling activities useful to support climate change adaptation policies. The simulations used in this study, in particular, are part of the EURO-CORDEX initiative (<http://www.euro-cordex.net/>) which provides regional climate projections for Europe at two different spatial resolutions, ~ 50

km (EUR-44, 0.44 degrees resolution) and ~12 km (EUR-11, 0.11 degrees resolution). EURO-CORDEX includes a total of
335 seven RCMs nested into several GCMs whose simulations belong to the most recent Climate Model Intercomparison Project
phase 5 (CMIP5 Taylor et al., 2012). **The RCA4 model was chosen** because, at the time **of downloading** the data from the
CORDEX archive, it was the only RCM providing data at the finest available spatial resolution (~12 km) and with sub-daily
(3hr) temporal resolution (WCRP, 2009). For a detailed description of the RCA4 model and its validation, please refer to the
technical report by Strandberg et al. (2014).

340 **4.2 Emission scenarios**

Future projections provided by climate models are based on a set of assumptions about the future evolution of the society in
terms of energetic and technological choices, population growth, land use changes, and others, which correspond to possible
greenhouse gas emission and concentration pathways in the atmosphere. The fifth IPCC Assessment Report (IPCC, 2013) uses
four different Representative Concentration Pathway (RCP) scenarios (Moss et al., 2010; van Vuuren et al., 2011) to evaluate
345 how climate is likely to change by the end of the 21st century. For this study two of these scenarios **were considered**, referred
to as RCP4.5 and RCP8.5, as they were the only ones available for the model under consideration. RCP4.5 is a stabilisation
scenario in which emissions will be stabilized by 2070 and carbon dioxide concentrations in the atmosphere are expected to
stabilise at about twice the pre-industrial level by the end of the century. RCP8.5 is an extreme business-as-usual scenario in
which greenhouse gas emissions are not expected to stabilise and carbon dioxide concentrations will be more than tripled at
350 the end of the century compared to preindustrial levels.

4.3 RCM data post-processing

RCA4 simulation outputs were linearly interpolated on the UTM grid at 1 km resolution in the study area using the same
method applied to the OI data and illustrated in section 3. To avoid distortions and artifacts, the data were first mapped with
the tool CDO (Climate Data Operators; v 1.7) on a lon-lat grid at 0.001° resolution and then projected with the GDALwarp
355 (Geospatial Data Abstraction Library; v. 2.1) on the final UTM grid (WGS 84 / UTM zone 32N) at 1 km resolution. The intrinsic
imperfections of climate model parameterizations and the errors in the model initialization are often reflected in an imperfect
representation of the observed climate, which can give rise to biases. Model biases must be taken into account when the climate
model outputs are used in impact studies, as impacts and feedbacks can be sensitive to the absolute values and the statistical
properties of the climatic input. Bias correction methods are usually applied to correct the differences between climate model
360 output and observed climatologies and are different depending on the variable and specific application which is considered
(Hempel et al., 2013; Maraun, 2013; Maraun et al., 2010). In this study we used an additive correction factor for adjusting the
temperature and a multiplicative correction factor for precipitation (a standard procedure for positive-defined fields) applied
pixel by pixel and constant in time, in order to correct the differences in the long-term climatology calculated over a common
time period between the simulated and observed fields. To this end, we calculated the long term mean of the historical Euro-
365 CORDEX simulations and of the OI dataset, already interpolated on the UTM grid, in the 30-year long period from 1986 to
2015. In order to maintain the physical consistency between the minimum temperature and the maximum temperature, and thus

avoid possible inversions, the same correction factor was used for daily minimum and maximum temperature data, calculated from the average between the daily minimum temperature and the maximum temperatures. The correction factor calculated to correct the model bias in the historical reference period was then applied to the future simulations. In addition to the bias adjustment, temperatures have been further corrected for the lapse rate, as already done for the OI data, based on Rolland (2003).

5 Results and discussion

5.1 Trends in observed data

Historical data from 1959 to 2017 were analyzed in each river catchment and sub-catchment, aggregating all data previously evaluated at the spatial resolution of 250 m, to find out significant trends both of the water balance terms and of the meteorological variables, for both the hydrological year and the quarterly analysis.

A linear regression was performed to calculate the temporal trends. The goodness of line fit was estimated with the coefficient of determination R^2 and its statistical significance was evaluated considering the p -value at 5% level (i.e., 95% significance) (Wilks, 2011).

5.1.1 Hydrological year analysis

During the hydrological year, an almost complete snow melting usually occurs in the considered area. Thus, in Eq. 1 P_q refers to the whole precipitation amount and $P_q - AET$ to drainage. Figure 3 shows the time series from 1959 to 2017 of the daily maximum temperature (in Kelvin, top left), P_q (mm/year, top right), AET (mm/year, bottom left), and drainage (mm/year, bottom right), for the Dora Riparia station in Turin as an example. It represents the driest watershed in this study, located in the middle of the study area with a west-east orientation (Figure 1). In the Alps this kind of valley is often characterised by foehn wind, corresponding to dry weather. Other examples in the Alps are Valais, Valle d'Aosta, Valtellina and Val Venosta.

Figure 3 reveals that, for this catchment, the daily maximum temperature has increased during the study period, with a statistically significant trend, as in all other catchments. The total precipitation, P_q , has decreased ($p=0.059$) while AET has increased ($p=0.054$), and drainage has significantly decreased ($p=0.032$). The catchment considered here is in the driest part of the study region, which is also the part where most of the significant trends of precipitation and AET were detected in the data analysis. The same approach is adopted for all the catchments in the study area. The results are summarised in Table S1 of the Supplementary Information. For all catchments, hydrological year trends in maximum temperature are positive and statistically significant, ranging between 0.032 and 0.078 K per year. AET trends are also positive for all catchments and statistically significant in 14 out of 23 cases. In 13 cases out of 14, they were either in the western mountain Dora Riparia area or in the southern irrigated area. The finding of an increase in AET is not obvious. Pangle et al. (2014) found a decreasing trend of AET looking at data from a mesocosm experiment. Their results highlight that the hydrological response to climate warming can be attenuated where precipitation is out of phase with the vegetation growing season. Our results refer instead to

an area where precipitation and growing season are not markedly out of phase. This can be due to the presence of the irrigated fields in the southern plains, where *AET* is mostly water limited and to the forested areas in the western mountain part of the region where *AET* is mostly energy limited. Blyth et al. (2018) also found an increase in evapotranspiration under similar conditions, using a land surface model in Great Britain from 1961 to 2015. In a more theoretical work Fatichi and Ivanov (2014) found *AET* to be quite unaffected by the imposed climate fluctuations, using the input data from four very different sites. This confirms the large role of the spatial variability in the response of water balance to climate change.

The map in Figure 4 represents the spatial distribution of annual actual evapotranspiration. It clearly shows the higher values of *AET* in the southern part of the region, where the irrigated fields play a major role, highlighting the importance of correctly modeling irrigation, as in the irrigation module implemented in our soil-water model. The surface energy budget is heavily influenced by agricultural water management and, together with air temperature increase, this leads to higher evapotranspiration average fluxes in comparison to non irrigated crops.

Most catchments exhibit negative precipitation (P_q) trends but only 5 cases display statistically significant trends. Precipitation at all catchments exhibits a high interannual variability (a quantification of interannual variability can be provided by the standard deviation, evaluated considering the detrended timeseries, shown in Table S1 of the Supplementary Information), in accordance with the results of previous studies in the same area (Pavan et al., 2019; Baiamonte et al., 2019; Ciccarelli et al., 2008). A recent study has shown a low frequency variability in the same historical years of this study for the Alpine region (Haslinger et al., 2021).

Drainage shows negative but mostly non-significant trends in all catchments. The seven catchments with significant decreasing trends all belong to the western and driest part of the region (Dora Riparia catchment). Four of them have significant trends both in P_q and *AET*, two of them have significant trends only in *AET*, and one has significant trends only in P_q .

This area is characterised by much lower yearly total precipitation values than the northern one. Long-term climatological values of cumulative yearly precipitation in the western Dora Riparia area are: Oulx 552 mm, Susa 710 mm, Beaulard 680 mm, Bardonecchia 724 mm, Pragelato 818 mm, while in the northern and wetter part they are: Pont Canavese 1228 mm, Viu 1338 mm, Germagnano 1342 mm. The southern part is instead characterized by the following values: Cumiana 924 mm, Moncalieri 882 mm.

To summarise, a slight negative trend is observed for precipitation and drainage over the last 60 years, showing a high spatial and interannual variability, as highlighted by the existing literature (Blöschl et al., 2019; Gudmundsson et al., 2017; Libertino et al., 2019). In Figure 3 the values for the whole Dora Riparia catchment down to Turin are shown. This western part of the study area can be identified as the driest one, as already observed in a recent work by Blöschl et al. (2019), where decreasing precipitation trends combine with increasing evapotranspiration trends. This can be attributed to non water-limited mountain vegetation in combination with increasing temperatures. Finally, the increasing evapotranspiration trends that characterize the southern part of the study area (due to farmers' irrigation), are not associated with significant decreasing precipitation trends, leading to a non significant drainage variation over time, even if *AET* has relatively high values in the southern area (Figure 4). Thus, in regions such as the one considered in this paper, when focusing on trend analysis, precipitation plays a major role in affecting drainage tendencies.




Figure 3. Hydrological year time series of T_{max} (top left), P_q (top right), AET (bottom left) and drainage (bottom right) along with their trends, R^2 , p -values and standard deviations, for the time period from 1959 to 2017, at the Turin cross-section 66 (the location is shown in Figure 1) of Dora Riparia river basin (average at the catchment level). The standard deviation, evaluated considering the detrended timeseries, provides a quantification of interannual variability.




Figure 4. Spatial distribution of annual actual evapotranspiration.

5.1.2 Analysis at the quarterly scale

The meteorological and hydrological variables at the quarterly time scale provide information on intra-annual variations. Figure 435 5 shows the trend results (color code) for all catchments (identified by their ID number in the y-axis of each panel) and for T_{max} (top left), rainfall plus snowmelt (top right), AET (bottom left) and drainage (bottom right) from 1959 to 2017. The numerical value of the trend is displayed in a cell when it is statistically significant ($p < 0.05$). When quarters are considered for the analysis, total precipitation (water input) is defined as the sum of liquid precipitation P_r and snowmelt N_f (see Section 2.3).

T_{max} shows positive and statistically significant trends in all catchments and quarters in the period 1959–2017, with values 440 between 0.022 and 0.086 K/year. The increase in the daily maximum temperature is more evident in winter (first quarter) and autumn (fourth quarter). The time series of actual evapotranspiration show positive and significant trends in the majority of catchments and quarters, indicating an overall increase of AET in the entire study area. AET has the greater increase in the summer quarter (JAS). Positive trends in the first quarter are consistent with those observed for T_{max} , both suggesting a possible anticipation of the growing season.




Figure 5. Trend slope of all quarters and catchments considered. Top: T_{max} (left) and precipitation (right), bottom: AET (left) and drainage (right). Trend slope is specified if significant at 5% level. When quarters are considered, the precipitation is represented by the sum of liquid precipitation P_r and melted snow N_f .

445 While historical yearly precipitation trends are overall negative, the quarterly analysis shows a significant decrease of precipitation in the first and second quarters, confirming that precipitation trends in NW Italy depend on the considered temporal aggregation (Brunetti et al., 2006). The drainage trend analysis shows an overall reduction in the first and second quarters, with larger decreases in *AMJ*, but significant only for six watersheds, again in the western driest part of the region. The third and fourth quarters show slight positive non-significant trends.

450 A visual comparison between yearly and quarterly analysis is shown in Figure 6, where precipitation (top panel) and drainage (bottom panel) trends are displayed over the whole study area. This figure clearly shows again the spatial pattern of hydrological changes, with negative trends in both precipitation plus snowmelt and discharge in spring. The dry western area is mainly represented by the Dora Riparia valley. Beside the spatial issue, the quarterly results show the importance of the time variability of precipitation. It confirm the importance of the computation at the daily scale (Healy, 2010). Also Masbruch et al. (2016) has
455 shown the sensitivity of drainage to rainfall time variability, and Fatichi and Ivanov (2014) stressed the importance of short periods (order of hours or days) with high *AET*.

5.2 Mid-century projections of drainage

This section reports the results for the future projections of drainage. As already mentioned, this variable allows to quantify the groundwater resource availability by using only meteorological variables provided by the climate models. The simulations
460 have been performed up to 2050 using the ensemble of simulations obtained with the RCM described in Section 4.1, at both hydrological year and quarterly scales.

5.2.1 Hydrological year analysis

Table S2 of the Supplementary Information shows the drainage trend slopes from 2018 up to 2050 for each catchment in the study area using climate projections under the RCP4.5 and the RCP8.5 scenarios. Drainage trends in the RCP4.5 scenario are
465 often negative but not statistically significant. These negative trends become stronger and occasionally statistically significant in the more extreme RCP8.5 scenario.

As a general statement, taking into account also the variability within the projection ensemble, the rather small values of drainage trends and their limited significance do not suggest a strong variation from a steady-state yearly drainage condition until 2050, with total precipitation that shows a slight decrease. We find a slight drainage increase with higher precipitation
470 amounts, decreasing trends with higher daily maximum temperature and, above all, a strong interannual variability (see, as an example, the standard deviation evaluated for the detrended timeseries shown in Figure 7).

The projections of $P_q - AET$ (drainage) for two different cross-sections of the study area, the Dora Riparia station in Turin (catchment ID 66) and the Orco station in S. Benigno (catchment ID 109), are shown in Figure 7. These two cases where shown
475 because the two catchments are among the largest ones (as shown in Table S1 of the Supplementary Information, 1243 km² and 829 km², respectively) and they represent, respectively, the drier (western) and the wetter (northern) parts of the region.

The overall results at yearly time scale can be summarized as follows:




Figure 6. Trend slope at the catchment level for all **studied** catchments. Top: **total precipitation**, bottom: drainage. Hydrological year (full dots) and quarterly trend values (four-segments circles) are jointly represented, and trend significance is also indicated. When quarters are considered, the precipitation is the sum of liquid precipitation P_r and melted snow N_f . Topographic shading is based on DEM data from the Risknat **Project** - “Base topografica transfrontaliera, ARPA Piemonte” (http://webgis.arpa.piemonte.it/ags101free/rest/services/topografia_dati_di_base/Sfumo_Europa_WM/MapServer).




Figure 7. Yearly drainage projections (RCP4.5 scenario in green and RCP8.5 scenario in red) for two different cross-sections (Dora Riparia in Torino, cross-section 66, and Orco in San Benigno, cross-section 109) of the studied area. Each row corresponds to the results obtained with one GCM driving the RCA4 RCM. The slope of the regression line, the coefficient of determination R^2 , the p-value and the standard deviation evaluated for the detrended timeseries are indicated.

- T_{max} shows positive trends, that are almost always significant for all model realizations and both scenarios;
- P_q has either positive or negative trends (according to the different GCMs and scenarios), rarely significant. In particular, RCA4 driven by the CNRM-CM5 and MPI-ESM-LR models shows positive trends for all catchments and scenarios. RCA4 driven by EC-Earth shows negative trends in all catchments when the RCP4.5 scenario is considered, and positive trends for the RCP8.5 scenario. Finally, RCA4 driven by IPSL-CM5A-MR and HadGEM2-ES shows negative trends in the whole study area for both the RCP4.5 and RCP8.5 scenarios;
- AET has either positive and negative trends, rarely significant. In detail, considering the RCP4.5 scenario, RCA4 driven by the HadGEM2-ES and MPI-ESM-LR models shows positive trends in the whole study area; when driven by EC-Earth and IPSL-CM5A-MR it shows negative trends in almost all catchments and when driven by CNRM-CM5 it shows spatial heterogeneity (only Malone, Sangone, Chisola and Orco primary catchments have positive trends). Considering the RCP8.5 scenario, RCA4 driven by CNRM-CM5, EC-Earth and MPI-ESM-LR shows positive trends in the whole study area, that are almost always significant for the case of the CNRM-CM5 model. RCA4 driven by IPSL-CM5A-MR has negative trends for all catchments, while RCA4 driven by HadGEM2-ES shows negative trends only in Banna, Chisola, Malone and Sangone primary catchments;
- slight positive drainage trends with higher precipitation amounts, negative trends with higher T_{max} and high interannual variability.

To summarize, we observe a large variability between the different projections, as also found in Crosbie et al. (2013) and in Persaud et al. (2020), with a much less clear pattern in spatial variability if compared with the historical data trend evaluations.

5.2.2 Quarterly analysis

Projections of drainage evaluated at the seasonal timescale show again a large variability within the ensemble of projections. However, a general tendency to drainage increase in the first quarter and to drainage decrease in the third and fourth quarters, *JAS* and *OND*, emerges. Table S3–S6 of the Supplementary Information show the drainage trend (expressed in mm/quarter/year) in the four quarters.

In the first quarter (*JFM*) the models mostly agree to indicate a drainage increase up to 2050 over the whole study area, with a wider agreement for RCP4.5 (4 out of 5 models concur in all catchments). In the second quarter (*AMJ*) in all river basins at least 3 out of 5 models estimate a drainage decrease for the RCP8.5 scenario, not for RCP4.5. In the third (3 out of 5 models) and fourth (4 out of 5 models) quarters an overall tendency of drainage decrease in the whole study area for RCP4.5 is evident.

For the other meteorological and water balance variables, at quarterly timescale we can report that:

- T_{max} shows positive trends in all quarters, almost always significant for all models and scenarios;

- in all quarters both rainfall plus snowmelt P_q and rainfall (P_r) have both positive and negative trends, rarely significant. Rainfall has almost **everywhere** a weak positive trend in the first quarter, rarely significant. Snowfall shows broad negative trends, almost always significant;
- 510 – *AET* shows broad positive trends in the first and second quarters, despite some differences between the models. In the third quarter an overall decreasing tendency can be observed, while in the fourth quarter trend slopes have values close to zero.

Also Stoll et al. (2011) found a major role for the time variability of precipitation, in recharge projections for a catchment in northern Switzerland. Konapala et al. (2020) indicate changes in long-term drainage, but they recall the limitations in the
515 ability of current generation coupled climate models to capture the key drivers of persistent weather extreme.

6 Conclusions

Assessing the impacts of climate change on groundwater resources represents a priority in water management, besides being an important scientific challenge. **In this framework, a proactive engagement of stakeholders is still lacking to a large part, and stakeholders are mainly considered as final users who download pre-computed decision-relevant scientific information in order**
520 **to develop and apply adaptation or mitigation strategies.** In this study a stakeholder-driven research study was carried out to **quantify the role of groundwater** on an area providing water to about 2.3 million people. **This area is** characterized by a very large spatial variability of precipitation, due to the proximity of high mountains and of the sea.

This work quantifies the trends of precipitation, temperature and actual evapotranspiration in order to estimate the trend of drainage as a proxy of the water available for drinking purposes. The analyses have been performed both at the hydrological
525 year and at the quarterly timescales. We analysed past and future conditions, using a historical climate dataset providing **minimum and maximum daily air** temperature and precipitation data for the period 1959–2017 and future projections from a multi-member ensemble of a regional climate model up to 2050, in order to be compliant with the time frame of water management objectives.

As suggested by Taylor et al. (2013), the role of irrigated agriculture is considered. In such a context, irrigation, that signifi-
530 **cantly contributes to increasing actual evapotranspiration as a consequence of the air temperature increase, was simulated in a novel way. More specifically, the results of field studies with both surface and sprinkler irrigation methods were used, combining them with crop and irrigation regional databases. *AET* has its greater increase in summer, while in the first three months of the year there is a consistent increase of both T_{max} and *AET*, suggesting a possible anticipation of the growing season. Regarding drainage, our analysis revealed a very strong interannual variability in the historical period, as well as remarkable**
535 **spatial differences, i.e., across the different catchments. It was found that the driest part of the region (the Dora Riparia Valley) shows significant negative precipitation and drainage trends both yearly and from January to June, confirming the seasonal variations found by Epting et al. (2021) and giving interesting hints for other dry climate valleys in the Alps. The increasing trend of yearly actual evapotranspiration is positive for all catchments and statistically significant in 14 out of 23 cases, namely both**

in the Dora Riparia Valley and in the southern irrigated area. This quantity, together with the relevant interannual variability of precipitation, has to be monitored in the future for its potential effects on drainage. Interestingly, only the Dora Riparia Valley and the most western part of Pellice catchment (cross-section 39), show significant negative trends for drainage, combining significant decreasing of precipitation and increasing of *AET* due to mountain non water-limited vegetation. The significant increasing trend of *AET* due to irrigation of the southern irrigated plain area lead to a non significant drainage because it was not combined with a significant decreasing of precipitation.

As for the future projections, there is a large inter-model variability, as in Crosbie et al. (2013) and in Persaud et al. (2020), of all hydrological variables, which is evident for all catchments, and almost no significant trends are present. This last result is in line with Gudmundsson et al. (2017) for this part of Europe. Also interestingly, the future scenarios are not reproducing the spatial variability of the historical analyses, although the soil and irrigation model used was the same.

This study constitutes a knowledge basis which helps for a better informed management, infrastructural and supply decisions in the considered study area, with a methodology that could be applied also in other areas in the world. In the Dora Riparia area a new drinking water aqueduct (70 km long, 800 liter per second discharge and 130 million euro cost) has been built to provide water from an existing hydroelectric reservoir at 1600 m asl, close to the border with France, down to Turin (Società Metropolitana Acque Torino, 2019). It will allow to avoid to continue pumping from the Dora Riparia aquifer, which have a negative trend of drainage, and has so far provided drinking water to 27 municipalities. A complementary hydraulic research has been performed by the Polytechnic of Torino (Fellini et al., 2018). The characteristics of spatial and temporal variability of aquifer recharge found in this paper will be of interest for other European areas, particularly in the Mediterranean area and in the Alps, where multiple anthropogenic pressures act on groundwater resources and where climate change will exacerbate competition between different users. Finally, climate change related spatial variability of precipitation has been similarly shown in a recent work about European floods for the period 1960-2010 (Blöschl et al., 2019). Even if that is a larger scale study, it is possible to see a decreasing trend for Dora Riparia, to be compared to a slightly increasing trend for the rest of Northern Italy, a behavior similar to the one found in this work. The outcomes of this paper reinforce the findings of two different precipitation regimes over Europe, the Mediterranean and the continental one. Both can be found in our study area and the Dora Riparia valley seems to be the bridge between them.

In this research, climate scientists, hydrologists, agricultural and soil scientists worked together with the experts from the local water utility to assess the water potential of the study area and better understand the role of groundwater in the future provision of water to the community. The approach followed is tailored to stakeholder needs, and the outcomes of the study are intended to drive or support local policies, in a general context which stimulated and supported a participatory planning, driven by state-of-the art climate information and projections. Our results will be integrated in the definition and refinement of the study-area long term guidelines and strategic development for groundwater resources protection and infrastructural provision and planning . It will be a part of a wider effort to strengthen good water resource management and governance.

Data availability. The derived datasets presented in this study can be obtained upon request to the corresponding author. Post-processed data are reproducible following the detailed descriptions of Section 2 and 3 and available upon request to the authors. All other datasets cited in the text can be retrieved following the instructions in the relevant citations.

Author contributions. All the authors conceived the study. EP and JvH provided and pre-processed the meteorological data. GM, GV, MP, DC, DG, IB and SF performed and analysed the hydrological simulations and prepared some of the figures. EB and SF analysed the results. EB and IB prepared the figures and EB all the tables. EB, EP, JvH, AP and SF wrote the paper with support from all the authors.

Competing interests. The authors declare that they have no conflict of interest

Acknowledgements. This work has been funded by Società Metropolitana Acque Torino S.p.A. Research has been supported in part by “Dipartimento di Eccellenza” DIST department funds and “PRIN MIUR 2017SL7ABC_005 WATZON Project” and by the Project of National Interest NextData of the MIUR (Italian Ministry for Education, University and Research). The authors would like to thank the Risk Responsible Resilience Interdepartmental Centre (R3C) DIST- PoliT0 for valuable collaboration and ARPA Piemonte for kind support.

References

- Agnese, C., Baiamonte, G., Cammalleri, C., Cat-Berro, D., Ferraris, S., and Mercalli, L.: Statistical analysis of inter-arrival times of rainfall events for Italian Sub-Alpine and Mediterranean areas, *Adv. Sci. Res.*, 8, 171–177, <https://doi.org/10.5194/asr-8-171-2012>, 2012.
- 585 Aguilar, C., Herrero, J., and Polo, M.: Topographic effects on solar radiation distribution in mountainous watersheds and their influence on reference evapotranspiration estimates at watershed scale, *Hydrology and Earth System Sciences*, 14, 2479–2494, <https://doi.org/10.5194/hess-14-2479-2010>, 2010.
- Allen, D., Cannon, A., Toewa, M., and Scibek, J.: Variability in simulated recharge using different GCMs, Tech. rep., <https://doi.org/10.1029/2009WR008932>, 2010.
- 590 Allen, R., Pereira, L., Raes, D., and Smith, M.: FAO irrigation and drainage paper No. 56, Tech. Rep. 56, Food and Agriculture Organization of the United Nations, 1998.
- ARPA Piemonte: NWIOI daily data, Version 2.1, data updated daily. Retrieved online from Rischì Naturali Archive Center, <http://www.arpa.piemonte.it/rischinaturali/tematismi/clima/confronti-storici/dati/dati.html>, accessed 2020-14-08., 2010a.
- ARPA Piemonte: Metodologia dell’Optimal Interpolation, Tech. rep., Arpa Piemonte, Dipartimento Sistemi Previsionali, retrieved online
595 from <http://rsaonline.arpa.piemonte.it/meteoclima50/pdf/metodologia.pdf>, accessed 2020-14-08, 2010b.
- Baiamonte, G., Mercalli, L., Cat-Berro, D., Agnese, C., and Ferraris, S.: Modelling the frequency distribution of interarrival times from daily precipitation time-series in North-West Italy, *Hydrology Research*, <https://doi.org/10.2166/nh.2018.042>, 2019.
- Bastiancich, L., Lasagna, M., Mancini, S., Falco, M., and Luca, D. A. D.: Temperature and discharge variations in natural mineral water springs due to climate variability: a case study in the Piedmont Alps (NW Italy), *Environ. Geochem. Health*,
600 <https://doi.org/10.1007/s10653-021-00864-8>, 2021.
- Baudena, M., Bevilacqua, I., Canone, D., Ferraris, S., Previati, M., and Provenzale, A.: Soil water dynamics at a midlatitude test site: Field measurements and box modeling approaches, *Journal of Hydrology*, 414–415, 329–340, <https://doi.org/10.1016/j.jhydrol.2011.11.009>, 2012.
- Bertrand, G., Siergieiev, D., Ala-Aho, P., and Rossi, P.: Environmental tracers and indicators bringing together groundwater, surface
605 water and groundwater-dependent ecosystems: importance of scale in choosing relevant tools, *Environ. Earth Sci.*, 72, 813–827, <https://doi.org/10.1007/s12665-013-3005-8>, 2014.
- Blöschl, G., Hall, J., Viglione, A., Perdigão, R. A. P., Parajka, J., Merz, B., Lun, D., Arheimer, B., Aronica, G. T., Bilibashi, A., Boháč, M., Bonacci, O., Borga, M., Čanjevac, I., Castellarin, A., Chirico, G. B., Claps, P., Frolova, N., Ganora, D., Gorbachova, L., Gül, A., Hannaford, J., Harrigan, S., Kireeva, M., Kiss, A., Kjeldsen, T. R., Kohnová, S., Koskela, J. J., Ledvinka, O., Macdonald, N., Mavrova-
610 Guirguinova, M., Mediero, L., Merz, R., Molnar, P., Montanari, A., Murphy, C., Osuch, M., Ovcharuk, V., Radevski, I., Salinas, J. L., Sauquet, E., Šraj, M., Szolgay, J., Volpi, E., Wilson, D., Zaimi, K., and Živković, N.: Changing climate both increases and decreases European river floods, *Nature*, 573, 108–111, <https://doi.org/10.1038/s41586-019-1495-6>, <https://doi.org/10.1038/s41586-019-1495-6>, 2019.
- Blyth, M., de la Torre, M. M., and E.L.Robinson: Trends in evapotranspiration and its drivers in Great Britain: 1961 to 2015, *Hydrol. Earth Syst. Sci. Discuss.*, <https://doi.org/10.5194/hess-2018-153>, 2018.
- 615 Brunetti, M., Maugeri, M., Nanni, T., Auer, I., Bohm, R., and Schoner, W.: Precipitation variability and changes in the greater Alpine region over the 1800–2003 period, *J. Geophys. Res (Atmospheres)*, 111, <https://doi.org/10.1029/2005JD006674>, 2006.

- Canone, D., Previati, M., Bevilacqua, I., Salvai, L., and Ferraris, S.: Field measurements based model for surface irrigation efficiency assessment, *Agricultural Water Management*, 156, 30–42, 2015.
- 620 Canone, D., Previati, M., and Ferraris, S.: Evaluation of stem-flow effects on the spatial distribution of soil moisture using TDR monitoring and an infiltration model, *Asce Journal of Irrigation and Drainage Engineering*, 143, 04016075–1–04016075–14, [https://doi.org/10.1061/\(ASCE\)IR.1943-4774.0001120](https://doi.org/10.1061/(ASCE)IR.1943-4774.0001120), 2016.
- CH2018-Project-Team: CH2018 – Climate Scenarios for Switzerland, Tech. rep., National Centre for Climate Services, Zurich, Switzerland, <https://doi.org/10.18751/Climate/Scenarios/CH2018/1.0>, 2018.
- 625 Ciccarelli, N., von Hardenberg, J., Provenzale, A., Ronchi, C., Vargiu, A., and Pelosini, R.: Climate variability in north-western Italy during the second half of the 20th century, *Global Planet. Change*, 63, 185–195, <https://doi.org/10.1016/j.gloplacha.2008.03.006>, 2008.
- Condon, L., Atchley, A., and Maxwell, R.: Evapotranspiration depletes groundwater under warming over the contiguous United States, *Nature Communications*, 11:873, 1–8, <https://doi.org/10.1038/s41467-020-14688-0>, 2020.
- Confortola, G., Soncini, ., and Bocchiola, D.: Climate change will affect hydrological regimes in the Alps, *Journal of Alpine Research*, 101-3, 630 <https://doi.org/10.4000/rga.2176>, 2013.
- Crosbie, R., Scanlon, B., Mpelasoka, F., Reedy, R., and Gates, J.: Potential climate change effects on groundwater recharge in the High Plains Aquifer, USA, *Water Resources Research*, 49, 3936–3951, <https://doi.org/10.1002/wrcr.20292>, 2013.
- De Luca, D., Lasagna, M., and Debernardi, L.: Hydrogeology of the western Po plain (Piedmont, NW Italy), *Journal of Maps*, 16, 265–273, <https://doi.org/10.1080/17445647.2020.1738280>, 2020.
- 635 Desiato, F., Fioravanti, G., Frascchetti, P., Perconti, W., and Piervitali, E.: Il clima futuro in Italia: analisi delle proiezioni dei modelli regionali. Stato dell' Ambiente 58/2015, Tech. rep., ISPRA, 2015.
- DeWalle, D. and Rango, A.: Principles of snow hydrology, Cambridge Univ.Press, Cambridge, UK, 2008.
- Doveri, M., Menichini, M., and Scozzari, A.: Protection of groundwater resources: worldwide regulations, scientific approaches and case study, vol. 40, pp. 13–30, Springer, Berlin, DEU, https://doi.org/10.1007/698_2015_421, 2016.
- 640 Epting, J., Huggenberger, P., Radny, D., Hammes, F., Hollender, J., Page, R. M., Weber, S., Bänninger, D., and Auckenthaler, A.: Spatiotemporal scales of river-groundwater interaction - The role of local interaction processes and regional groundwater regimes, *Sci Total Environ.*, 618, 1224–1243, <https://doi.org/10.1016/j.scitotenv.2017.09.219>, 2018.
- Epting, J., Michel, A., Affolter, A., and Huggenberger, P.: Climate change effects on groundwater recharge and temperatures in Swiss alluvial aquifers, *Journal of Hydrology X*, 11, 100071, <https://doi.org/10.1016/j.hydroa.2020.100071>, 2021.
- 645 Fatichi, S. and Ivanov, V.: Interannual variability of evapotranspiration and vegetation productivity, *Water Resources Research*, 50, 3275–3294, <https://doi.org/10.1002/2013WR015044>, 2014.
- Fellini, S., Vesipa, R., Boano, F., and Ridolfi, L.: Multipurpose Design of the Flow-Control System of a Steep Water Main, *Journal of Water Resources Planning and Management ASCE*, 144, [https://doi.org/10.1061/\(ASCE\)WR.1943-5452.0000867](https://doi.org/10.1061/(ASCE)WR.1943-5452.0000867), 2018.
- Giorgi, F., Jones, C., and Asrar, G.: Addressing climate information needs at the regional level: the CORDEX framework, *World Meteorological Organization (WMO) Bulletin*, 58, 175, 2009.
- 650 Gudmundsson, L., Seneviratne, S., and Zhang, X.: Anthropogenic climate change detected in European renewable freshwater resources, *Nat. Climate Change*, 7, 813–817, <https://doi.org/10.1038/NCLIMATE3416>, 2017.
- Haslinger, K., Hofstatter, M., Schoener, W., and G.Bloeschl: Changing summer precipitation variability in the Alpine region: on the role of scale dependent atmospheric drivers, *Climate Dynamics*, <https://doi.org/10.1007/s00382-021-05753-5>, 2021.
- 655 Healy, R. W.: Estimating groundwater recharge, Cambridge, Cambridge, UK, 2010.

- Hempel, S., Frieler, K., Warszawski, L., Schewe, J., and Piontek, F.: A trend-preserving bias correction - the ISI-MIP approach, *Earth System Dynamics*, 4, 219–236, <https://doi.org/10.5194/esd-4-219-2013>, 2013.
- IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/CBO9781107415324>, www.climatechange2013.org, 2013.
- IPCC: Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II, and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, IPCC, Geneva, Switzerland, 2014.
- IPLA: Carta dei suoli del Piemonte a scala 1:250.000, Tech. rep., IPLA, Regione Piemonte, Firenze, Italy, 2007.
- Jarvis, A., Guevara, E., Reuter, H., and Nelson, A.: Hole-filled SRTM for the globe: version 4: data grid, published by CGIAR-CSI on 19 August 2008, available from the CGIAR-CSI SRTM 90m Database ([//srtm.csi.cgiar.org](http://srtm.csi.cgiar.org)), 2008.
- Jiménez Cisneros, B., Oki, T., Arnell, N., Benito, G., Cogley, J., Doll, P., Jiang, T., and Mwakalila, S.: *Freshwater resources*, Cambridge University Press, 2014.
- Kalbus, E., Reinstorf, F., and Schirmer, M.: Measuring methods for groundwater and surface water interactions: a review, *Hydrology and Earth System Sciences*, 10, 873–887, <https://doi.org/10.5194/hess-10-873-2006>, 2006.
- Konapala, G., Mishra, A., and et al., Y. W.: Climate change will affect global water availability through compounding changes in seasonal precipitation and evaporation, *Nat. Commun.*, 11, 3044, <https://doi.org/10.1038/s41467-020-16757-w>, 2020.
- Kumar, S., Zwiers, F., Dirmeyer, P., Lawrence, D., Shresta, R., and A.T.Werner: Terrestrial contribution to the heterogeneity in hydrological changes under global warming, *Water Resources Research*, 52, 3127–3142, 2016.
- Lasagna, M., Luca, D. D., and Franchino, E.: Nitrate contamination of groundwater in the western Po Plain (Italy): the effects of groundwater and surface water interactions, *Environ. Earth Sci.*, 75, 240, <https://doi.org/10.1007/s12665-015-5039-6>, 2016.
- Li, B., Rodell, M., and Famiglietti, J. S.: Groundwater variability across temporal and spatial scales in the central and northeastern U.S., *Journal of Hydrology*, 525, 769–780, <https://doi.org/10.1016/j.jhydrol.2015.04.033>, <https://www.sciencedirect.com/science/article/pii/S0022169415002929>, 2015.
- Libertino, A., Ganora, D., and Claps, P.: Evidence for increasing rainfall extremes remains elusive at large spatial scales: the case of Italy, *Geophysical Research Letters*, 46, <https://doi.org/10.1029/2019GL083371>, 2019.
- Maraun, D.: Bias correction, quantile mapping, and downscaling: revisiting the inflation issue, *Journal of Climate*, 26, 2137–2143, 2013.
- Maraun, D. et al.: Precipitation downscaling under climate change: Recent developments to bridge the gap between dynamical models and the end user, *Rev. Geophys.*, 48, RG3003, <https://doi.org/10.1029/2009RG000314>, 2010.
- Masbruch, M., Rumsey, C., Gangopadhyay, S., Susong, D., and Pruitt, T.: Analyses of infrequent (quasi-decadal) large groundwater recharge events in the northern Great basin: their importance for groundwater availability, use, and management, *Water Resources Research*, 52, 7819–7836, 2016.
- Moeck, C., Grech-Cumbo, N., Podgorski, J., Bretzler, A., and Gurdak, J.: A global-scale dataset of direct natural groundwater recharge rates: a review of variables, processes and relationships, *Science of the total environment*, 717, 1–19, <https://doi.org/10.1016/j.scitotenv.2020.137042>, 2020.
- Moench, M., Burke, J., and Moench, Y.: *Rethinking the Approach to Groundwater and Food Security*, Tech. Rep. Water Reports 24, Food and Agriculture Organization of the United Nations, Rome, Italy, 2003.
- Moss, R., Edmonds, J., Hibbard, K., et al.: The next generation of scenarios for climate change research and assessment, *Nature*, 463, 747–756, <https://doi.org/10.1038/nature08823>, 2010.

- Pangle, L., Gregg, J., and McDonnell, J.: Rainfall seasonality and an ecohydrological feedback offset the potential impact of climate warming on evapotranspiration and groundwater recharge, *Water Resources Research*, 50, 1308—1321, <https://doi.org/10.1002/2012WR013253>, 2014.
- 695
- Parry, M., Parry, M., Canziani, O., Palutikof, J., P. Van der Linden, P., and Hanson, C.: *Climate Change 2007-Impacts, Adaptation and Vulnerability*, Cambridge University Press, 2007.
- Pavan, V., Antolini, G., Barbiero, R., et al.: High resolution climate precipitation analysis for north-central Italy, 1961–2015, *Climate Dynamics*, 52, 3435—3453, <https://doi.org/10.1007/s00382-018-4337-6>, 2019.
- 700
- Persaud, E., Levison, J., MacRitchie, S., Berg, S., Parker, B., and Sudicky, E.: Integrated modelling to assess climate change impacts on groundwater and surface water in the Great lakes Basin using diverse climate forcing, *Journal of Hydrology*, 584, 1–15, <https://doi.org/10.1016/j.jhydrol.2020.124682>, 2020.
- Pradier, S., Chong, M., and Roux, F.: Radar observations and numerical modeling of a precipitation line during MAP IOP 5, *Monthly Weather Review*, 130, 2533–2553, 2002.
- 705
- Raco, B., Vivaldo, G., Doveri, M., Menichini, M., Masetti, G., Battaglini, R., Irace, A., Fioraso, G., Marcelli, I., and Brussolo, E.: Geochemical, geostatistical and time series analysis techniques as a tool to achieve the Water Framework Directive goals: An example from Piedmont region (NW Italy), *Journal of Geochemical Exploration*, 229, 106 832, <https://doi.org/10.1016/j.gexplo.2021.106832>, 2021.
- Raffelli, G., Previati, M., Canone, D., Gisolo, D., Bevilacqua, I., Capello, G., Biddoccu, M., Cavallo, E., Deiana, R., Cassiani, G., and Ferraris, S.: Local- and Plot-Scale Measurements of Soil Moisture: Time and Spatially Resolved Field Techniques in Plain, Hill and Mountain Sites, *Water*, 9, <https://www.mdpi.com/2073-4441/9/9/706>, 2017.
- 710
- Regione Piemonte: Anagrafe agricola del Piemonte, retrieved online from <http://www.sistemapiemonte.it/cms/privati/agricoltura/servizi/367-anagrafe-agricola-unica-data-warehouse>, accessed 2020-14-08., 2006.
- Regione Piemonte: SIBI Sistema Informativo della Bonifica e Irrigazione, retrieved online from Regional Irrigation Information System Archive Center, <https://www.regione.piemonte.it/web/temi/agricoltura/agroambiente-meteo-suoli/bonifica-irrigazione-sibi>, accessed 2020-14-08., 2016.
- 715
- Regione Piemonte: BDTRE, Base Dati Territoriale di Riferimento degli Enti, cartographic reference material, 1:250.000, retrieved online from Sistema informativo territoriale e ambientale Archive Center, <http://www.geoportale.piemonte.it/geocatalogorp/index.jsp>, accessed 2020-14-08., 2018a.
- 720
- Regione Piemonte: Piano di Tutela delle Acque – Revisione 2018, Tech. rep., Regione Piemonte, Direzione Ambiente, Governo e Tutela del territorio, Settore Tutela delle Acque, retrieved online from https://www.regione.piemonte.it/web/sites/default/files/media/documenti/2019-01/pta2018_tavole_di_piano.pdf, accessed 2021-07-05, 2018b.
- Rolland, C.: Spatial and seasonal variations of air temperature lapse rates in Alpine regions, *J. Climate*, 16, 1032–1046, [https://doi.org/10.1175/1520-0442\(2003\)016<1032:SASVOA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<1032:SASVOA>2.0.CO;2), 2003.
- 725
- Rumsey, C. A., Miller, M., Susong, D., Tillman, F., and Anning, D.: Regional scale estimates of baseflow and factors influencing baseflow in the Upper Colorado River Basin, *Journal of Hydrology: Regional Studies*, 4, 91–107, <https://doi.org/10.1016/j.ejrh.2015.04.008>, 2015.
- Schaap, M., Leij, F., and van Genuchten, M.: Rosetta: a computer program for estimating soil hydraulic parameters with hierarchical pedo-transfer functions, *Journal of Hydrology*, 251, 163–176, [https://doi.org/10.1016/S0022-1694\(01\)00466-8](https://doi.org/10.1016/S0022-1694(01)00466-8), 2001.
- Smerdon, B. D.: A synopsis of climate change effects on groundwater recharge, *Journal of Hydrology*, 555, 125–128, <https://doi.org/10.1016/j.jhydrol.2017.09.047>, <https://www.sciencedirect.com/science/article/pii/S0022169417306510>, 2017.
- 730

- Società Metropolitana Acque Torino: Consolidated financial statement and fiscal year financial statement, Tech. rep., retrieved online from https://www.smatorino.it/wp-content/uploads/2020/07/EN-BILANCIO_SMAT_31_12_2019.pdf, accessed 2021-09-08, 2019.
- Sorland, S., Schar, C., D.Luthi, and Kjellstrom, E.: Bias patterns and climate change signals in GCM-RCM model chains, *Environ. Res. Lett.*, 13, <https://doi.org/10.1088/1748-9326/aacc77>, 2018.
- 735 Stoll, M., Franssen, H. H., M.Butts, and W.Kinzelbach: Analysis of the impact of climate change on groundwater related hydrological fluxes: a multi-model approach including different downscaling methods, *Hydrol. Earth Syst. Sci.*, <https://doi.org/10.5194/hess-15-21-2011>, 2011.
- Strandberg, G., Barring, L., Hansson, U., Jansson, C., Jones, C., Kjellström, E., Kolax, M., M. Kupiainen, M., Nikulin, G., Samuelsson, P., Ullerstig, A., and Wang, S.: CORDEX scenarios for Europe from the Rossby Centre regional climate model RCA4, <http://urn.kb.se/resolve?urn=urn:nbn:se:smhi:diva-2839>, 2014.
- 740 Taylor, K., Stouffer, R., and Meehl, G.: An overview of CMIP5 and the experiment design, *Bulletin of the American Meteorological Society*, 93, 485–498, <https://doi.org/10.1175/BAMS-D-11-00094.1>, 2012.
- Taylor, R., B. Scanlon, B., and et al., P. D.: Ground water and climate change, *Nature Clim Change*, 3, 322–329, <https://doi.org/10.1038/nclimate1744>, 2013.
- 745 van der Gun, J.: Groundwater and Global Change: Trends, Opportunities and Challenges, Tech. rep., United Nations Educational, Paris, France, 2012.
- van Vuuren, D., Stehfest, E., Elzen, M. D., Deetman, S., et al.: RCP2.6: Exploring the possibility to keep global mean temperature change below 2°C, *Climatic Change*, 109, 95–116, <https://doi.org/10.1007/s10584-011-0152-3>, 2011.
- WCRP: CORDEX data access, retrieved online from ESGF Archive Center, <https://cordex.org/data-access/esgf/>, accessed 2020-14-08.,
- 750 2009.
- WHO: Protecting groundwater for health: Managing the quality of drinking-water sources, IWA Publishing for World Health Organization, 2006.
- Wilks, D. S.: *Statistical Methods in the Atmospheric Sciences*, vol. 100, Academic Press, third edn., <http://www.sciencedirect.com/science/bookseries/00746142/100/supp/C>, 2011.
- 755 WWAP: The United Nations World Water Development Report 2015: Water for a Sustainable World, Tech. rep., UNESCO, Paris, 2015.
- Zeinivand, H. and Smedt, F. D.: Hydrological modeling of snow accumulation and melting on river basin scale, *Water Resources Management*, 23, 2271–2287., 2009.