

1 River runoff in Switzerland in a changing climate - runoff regime 2 changes and their time of emergence

3 Regula Muelchi¹, Ole Rössler^{1,2}, Jan Schwanbeck¹, Rolf Weingartner^{1,3}, Olivia Martius^{1,3}

4 ¹Institute of Geography and Oeschger Centre for Climate Change Research, University of Bern, Switzerland

5 ²now at German Federal Institute of Hydrology (BfG), Germany

6 ³Mobilair Lab for Natural Risks, University of Bern, Switzerland

7
8 *Correspondence to* Regula Muelchi (regula.muelchi@giub.unibe.ch)

9
10 **Abstract.** Assessments of climate change impacts on runoff regimes are essential to climate change adaptation and mitigation
11 planning. Changing runoff regimes and thus changing seasonal patterns of water availability strongly influence various
12 economic sectors such as agriculture, energy production, and fishery and also affect river ecology. In this study, we use new
13 transient hydrological scenarios driven by the most up-to-date local climate projections for Switzerland, the Swiss Climate
14 Change Scenarios. These provide detailed information on changes in runoff regimes and their time of emergence for 93 rivers
15 in Switzerland under three Representative Concentration Pathways (RCPs), RCP2.6, RCP4.5, and RCP8.5. These transient
16 scenarios also allow changes to be framed as a function of global mean temperature.
17 The new projections for seasonal runoff changes largely confirm the sign of changes in runoff from previous hydrological
18 scenarios with increasing winter runoff and decreasing summer and autumn runoff. Spring runoff is projected to increase in
19 high-elevation catchments and to decrease in lower-lying catchments. Despite the increases in winter and some increases in
20 spring, the annual mean runoff is projected to decrease in most catchments. Compared to lower-lying catchments, runoff
21 changes in high-elevation catchments (above 1500 m a.s.l.) are larger in winter, spring, and summer due to the large influence
22 of reduced snow accumulation and earlier snow melt and glacier melt. The changes in runoff and the agreement between
23 climate models on the sign of change both increase with increasing global mean temperatures and stronger emission scenarios.
24 This amplification highlights the importance of climate change mitigation.
25 The time of emergence is the time when the climate signal emerges significantly from natural variability. Under RCP8.5, times
26 of emergence were found early, before the period 2036–2065, in winter and summer for catchments with mean altitudes above
27 1500 m a.s.l. Significant changes in catchments below 1500 m a.s.l. emerge later in the century. Not all catchments show
28 significant changes in the distribution of seasonal means, and thus no time of emergence could be determined in these

29 catchments. Furthermore, the significant changes of seasonal mean runoff are not persistent over time in some catchments due
30 to nonlinear changes in runoff.

31 **1 Introduction**

32 Anthropogenic climate change is certain to affect regional and local hydrology (IPCC, 2013; IPCC, 2014a; IPCC, 2014b).
33 Major research efforts have enabled more precise and more reliable projections of regional and local temperature and
34 precipitation changes (CH2018, 2018). The new Swiss Climate Change Scenarios (CH2018) are the result of a large modelling
35 effort to downscale regional climate projections for Switzerland to local scales (CH2018, 2018). Warming in Switzerland is
36 projected to be higher than the global mean warming. Without major mitigation efforts, mean temperatures are projected to
37 increase by up to 6.8°C by end of the 21st century in Switzerland (CH2018, 2018). This large increase in temperatures is
38 accompanied by changes in many other hydrologically relevant variables such as precipitation amounts, precipitation type,
39 and glacier volumes. The projected combination of increasing temperatures, changing precipitation patterns, retreating
40 glaciers, and changes in snowpack will have a potentially strong impact on runoff regimes. Runoff regimes reflect the integral
41 response in time and space of the hydrological conditions within a catchment and hence its water supply. Understanding and
42 assessing changes in runoff regimes is crucial both for many economic sectors, such as agriculture, fishery, hydropower
43 generation, and tourism, and for the ecology within and around the rivers. Assessments of climate change impacts on river
44 runoff are therefore particularly important for decision makers in adaptation planning and can also provide a basis for
45 mitigation policies.

46 Several studies on the impacts of climate change on hydrology in Switzerland have been conducted in recent years. These have
47 focused on changes in runoff regimes (Horton et al., 2006; Koeplin et al., 2012; Koeplin et al., 2014; Addor et al., 2014, Milano
48 et al., 2015), on changes in low flows (Jenicek et al., 2018; Brunner et al., 2019a,b), on changes in high flows (Keller et al., 2018;
49 Brunner et al., 2019a), and on the effects of retreating glaciers for rivers (Huss et al., 2008; Farinotti et al., 2012; Finger et al.,
50 2012; Huss et al., 2014, Fatichi et al., 2015; Etter et al., 2017). Studies on climate change impacts on various aspects of the
51 hydrology in the Alpine area have also been carried out in Austria (e.g., Weber et al., 2010; Blöschl et al., 2011; Prash et al.,
52 2011; Tecklenburg et al., 2012; Goler et al., 2016; Wijngaard et al., 2016; Hanzer et al., 2018), in Italy (e.g., Gropelli et al.,
53 2011), in Germany (e.g., Hattermann et al., 2015; Nilson et al., 2014), and in France (e.g., Ruiz-Villanueva et al., 2015, Vidal
54 et al., 2016). Most of these studies are case studies or focus on specific aspects of hydrology. Previous studies have focused
55 on changes in runoff regimes and included many catchments with diverse properties in Switzerland (Horton et al., 2006;
56 Koeplin et al., 2012; Koeplin et al., 2014; Addor et al., 2014). Koeplin et al. (2012) found a shift in seasonality with increasing
57 winter runoff, decreasing summer runoff, unchanging annual mean runoff for lower-lying catchments and increasing annual
58 mean runoff for high Alpine catchments. However, these studies are based either on older climate model generations or on
59 climate simulations downscaled with a delta change approach. The delta change method is based on climate simulations
60 downscaled for 30-year periods both in the reference period and in future periods and does not provide continuous transient

61 simulations for the whole 21st century. Therefore, this approach does not simulate daily to interannual variability (Bosshard et
62 al., 2011). More sophisticated downscaling approaches have been developed, such as quantile mapping (Teutschbein and
63 Seibert, 2012; Gudmundsson et al., 2012). They correct not only for the mean bias but for the full distribution, are applicable
64 to long-term climate simulations, and allow transient scenarios to be established. Using quantile mapping as a downscaling
65 approach can intensify and increase the number of projected extremes (Roessler et al., 2019). Therefore, the present study uses
66 the Hydro-CH2018-Runoff ensemble (dataset: Muelchi et al., 2020; Muelchi et al., 2021) run with the most up-to-date local
67 climate change scenarios for Switzerland, CH2018. These scenarios used the quantile mapping approach to downscale the
68 coarse climate model output. The Hydro-CH2018-Runoff ensemble includes transient daily simulations from 1981 to 2099 for
69 93 catchments in Switzerland and three Representative Concentration Pathways (RCPs): RCP2.6, RCP4.5, and RCP8.5. The
70 new ensemble enables detailed quantification of changes in runoff regimes. In this study, we investigate changes in runoff
71 regimes under a range of RCP scenarios. These 119-year continuous daily runoff simulations enable the estimation of the time
72 of emergence (Giorgi and Bi, 2009; Leng et al., 2016) of those changes. The time of emergence is the time when the climate
73 signal emerges significantly from natural variability. It is of particular importance to the question of how much time is left for
74 adaptation planning. In this study, we analyze not only the time of emergence but also the evolution of time of emergence over
75 time. Moreover, the ensemble allows the quantification of changes at different global warming levels; these warming levels
76 are particularly relevant for policy makers (e.g., James et al., 2017). The framing of the results as a function of global mean
77 temperature change rather than time enables a direct link to the Paris agreement target and indicates the changes to be expected
78 in a +2°C world and the consequences of missing this target. In this study, we assess changes for global warming levels of
79 +1.5°C, +2°C, and +3°C.

80

81 **2 Data**

82 For this study, we use the Hydro-CH2018-Runoff ensemble of daily discharge simulations for 93 medium-sized (catchment
83 size between 14–1700 km²) catchments in Switzerland (Fig. 1; Muelchi et al., 2020; Muelchi et al., 2021). The Precipitation-
84 Runoff-Evapotranspiration HRU-related Model (PREVAH) hydrological modelling system (Viviroli et al., 2009) was used
85 for simulating the hydrological response to the climate change scenarios. PREVAH is a semidistributed model based on
86 hydrological response units and includes submodules to account for important hydrological processes related to snow, glacier,
87 and soil moisture dynamics and evapotranspiration. PREVAH was calibrated and validated for each catchment (for more
88 details, see Muelchi et al., 2021). The calibrated parameters were kept constant for the simulation of runoff under climate
89 change, and land use was held unchanged for nonglaciated catchments. The minor impact of land use changes on changes in
90 the runoff regime for Switzerland was assessed by Koeplin et al. (2014). For glaciated catchments, glacier extents were updated
91 every 5 years according to the glacier projections by Zekollari et al. (2019), which are based on the same climatic data set. The

92 land use of areas from which glaciers disappear was replaced by rock for areas above 3000 m a.s.l. and by bare soil for areas
93 below 3000 m a.s.l.

94 The meteorological input used for the simulations comes from the new Swiss climate change scenarios CH2018 (CH2018,
95 2018). The CH2018 scenarios used EURO-CORDEX simulations (Jacob et al., 2014; Kotlarski et al., 2014) and applied
96 statistical downscaling (quantile mapping; Teutschbein and Seibert, 2012; Gudmundsson et al., 2012) on chains of Global
97 Circulation Models and Regional Climate Models (GCM-RCM chains). The results consist of gridded high-resolution (2x2
98 km) daily temperature and precipitation data for Switzerland for the period 1981–2099. This dataset was then used to drive the
99 hydrological model, resulting in the Hydro-CH2018-Runoff ensemble, which comprises daily runoff time series for 1981–
100 2099 for each of the GCM-RCM chains. The Hydro-CH2018-Runoff ensemble contains simulations for three RCPs: 8
101 simulations under RCP2.6, 16 simulations under RCP4.5, and 20 simulations under RCP8.5. The GCM-RCM chains and their
102 underlying emission scenarios used for this study are listed in Table 1. The Hydro-CH2018-Runoff ensemble comprises
103 transient long-term daily runoff projections for Switzerland for the period 1981–2099. These transient simulations incorporate
104 the daily to interannual climate variability. Changes in runoff constrained by the different warming levels were analyzed using
105 the global mean temperatures of the driving GCMs (CMIP5, Taylor et al., 2012).

106

107 **3 Methods**

108 **3.1 Study area**

109 The study area consists of 93 catchments distributed over Switzerland (Fig. 1) and covering a wide range of catchment
110 characteristics with an average catchment size of 314 km² (from 14 km² to 1700 km²) and a mean altitude of 1344 m a.s.l.
111 (range in mean altitude: 476–2700 m a.s.l.). Of the 93 catchments, 22 are glaciated with a modelled degree of glaciation
112 varying between 1% and 29% (Fig. S1). The present runoff regimes range from glacier-fed catchments in high Alpine areas,
113 mainly snow-driven catchments (mean altitude above 1550 m a.s.l.) in the Alps and pre-Alps to rain-fed catchments
114 predominant in the Swiss Plateau and at lower elevations in the southern part of Switzerland. Six catchments (highlighted in
115 Fig. 1) are used as example catchments representing typical runoff regimes: Rosegbach, highly glaciated (29%); Kander,
116 slightly glaciated (5%); Plessur, high-Alpine snow influenced; Emme, pre-Alpine rain and snow influenced; Venoge,
117 lowland rain dominated; and Verzasca, southern Alpine rain and snow dominated. An overview of the catchment
118 characteristics (Table S1), the degree of glaciation (Fig. S1), and the fraction of precipitation falling as snow (Fig. S2) can be
119 found in the Supplement.

120 **3.2 Determining changes in seasonal and annual mean runoff**

121 The simulations were analyzed using annual and seasonal changes of runoff under the three RCPs. The seasons were defined
122 as winter (December, January, February), spring (March, April, May), summer (June, July, August), and autumn (September,
123 October, November). All changes were specified for 30-year periods to remove interannual variability. The reference period
124 covers the years 1981–2010 and was compared with the far future period 2070–2099. The median among all simulations within
125 an RCP pathway was considered the best estimate. To obtain an indication of the robustness of the estimation, the changes
126 were highlighted when at least 90% of the simulations showed the same direction of change, whether positive or negative.
127 This criterion corresponds to “very likely” in the terminology of the Intergovernmental Panel on Climate Change (IPCC) that
128 the runoff changes are either positive or negative (Mastrandrea et al., 2010). The changes in seasonal and annual runoff were
129 also analyzed as a function of the mean elevation of the catchments to show the elevation dependence of runoff responses.
130 Other elevation-related characteristics of a catchment might have been used. However, a study by Koeplin et al. (2012) showed
131 that catchment responses to climate change in Switzerland can be directly linked to mean altitude.
132 For the analysis of changes in the runoff regime, monthly means for 30-year periods were calculated with the median
133 representing the best estimate and the uncertainty band showing the full range among all models within an RCP. To simplify
134 the interpretation of the results, this article focuses only on changes up to the period 2070–2099 under RCP8.5 and RCP2.6.
135 These two emission scenarios provide the broadest range of changes available from the full Hydro-CH2018-Runoff ensemble.
136 Results for the near-future period of 2045–2074 and for RCP4.5 can be found in the Supplement.

137 **3.3 Determination of the time of emergence of seasonal and annual runoff changes**

138 The time of emergence (Giorgi and Bi, 2009; Leng et al., 2016) indicates the time when significant changes in the distribution
139 of seasonal and annual means emerge from natural variability. The Kolmogorov-Smirnov test was used to test whether two
140 30-year samples of seasonal or annual means are drawn from the same distribution. This test was conducted on the distributions
141 of moving 30-year windows and the distribution of the reference period. The Kolmogorov-Smirnov test procedure was also
142 used in previous studies (e.g., for precipitation Mahlstein et al., 2011; Gaetani et al., 2020), but other definitions of time of
143 emergence also exist. Although Mahlstein et al. (2011) did not find significant differences from other definitions, Gaetani et
144 al. (2020) found that the Kolmogorov-Smirnov testing procedure results in a more robust and earlier time of emergence. The
145 testing was performed for each simulation under RCP8.5 and each catchment separately. Constraining the analysis to the
146 RCP8.5 ensemble was motivated by the sufficiently large number of simulations, 20, within the ensemble. The time of
147 emergence was then defined following the procedure used in Mahlstein et al. (2011) and refers to the last year of the 30-year
148 moving window in which the Kolmogorov-Smirnov test was rejected for the first time at 95% significance. We then considered
149 the significance of changes in the seasonal and annual mean when at least 66% of the models detect a significant change in the
150 same 30-year window. Sixty-six percent corresponds to the threshold referred to as “likely” in the IPCC terminology
151 (Mastrandrea et al., 2010). Because changes in runoff may not be linear over time, the time of emergence may not be stable

152 after the first detection. Even though changes in seasonal and annual runoff are tested as significant in one period, they may
153 not be significant in all subsequent periods (e.g., due to nonlinear effects in snow melt or glacier melt contributions). Therefore,
154 we also analyzed the temporal evolution of rejections of the null hypothesis for the Kolmogorov-Smirnov test (p -values smaller
155 than 0.05).

156 **3.4 Stratification of seasonal and annual runoff changes by increases in global mean temperature**

157 For the analysis of runoff changes as a function of global mean temperature change, temperature targets were defined of
158 +1.5°C, +2°C, and +3°C with respect to the pre-industrial state. Because the temperature targets are defined with respect to
159 the pre-industrial state, the warming observed between the pre-industrial state (1864–1900) and the reference period (1981–
160 2010) was subtracted from the temperature targets. The observed warming is estimated to be 0.6°C, and thus the remaining
161 global warming for the 1.5°C, 2°C, and 3°C temperature targets is 0.9°C, 1.4°C, and 2.4°C, respectively (Morice et al., 2012;
162 for technical details: CH2018, 2018). For each of the driving GCMs used in the Hydro-CH2018-Runoff ensemble for RCP8.5,
163 we computed differences in moving 30-year averages of global mean temperatures compared to the reference period. The 30-
164 year windows when global mean temperature change exceeds +0.9°C, +1.4°C, and +2.4°C were selected for each GCM.
165 Subsequently, the seasonal and annual changes in runoff were extracted for each of the time periods and the driving GCM in
166 the GCM-RCM combination. Again, catchments with robust signs were highlighted if at least 90% of simulations agreed on
167 the direction of change.
168

169 **4 Results**

170 **4.1 Changes in seasonal and annual mean runoff for Switzerland**

171 Changes in the multimodel median of seasonal and annual mean runoff by end of the century (2070–2099) are shown in Fig.
172 2 for RCP8.5 and in Fig. 3 for RCP2.6 (see Fig. S3 for RCP4.5 and Figs. S4-S6 for the period 2045–2074). Highlighted
173 catchments show changes where at least 90% of the models agree on the sign of change.

174 **4.1.1 Changes in winter runoff for Switzerland**

175 In winter, all catchments show positive mean runoff changes under RCP8.5 by the end of the century (Fig. 2a) compared to
176 the reference period. The mean runoff changes range from +2% to +221% with larger changes in higher-elevation catchments.
177 The mean change among all catchments is +48%. In all, 84 out of 93 catchments show good agreement ($\geq 90\%$) on the sign
178 of change. Under RCP2.6, 87 out of 93 catchments show positive changes, with mean runoff changes across all catchments of
179 +13% (Fig. 3a). However, only 41 catchments show robust model agreement on the sign of change. The range among the
180 catchments is between -3% and +58%, again with stronger changes in the mountainous areas.

181 **4.1.2 Changes in spring runoff for Switzerland**

182 In spring, both positive and negative changes are found in mean runoff (Figs. 2b and 3b). The mean change across all
183 catchments is +9% under RCP8.5 (Fig. 2b). Although most of the lower catchments show a decrease in runoff (up to -21%),
184 the higher-elevation catchments exhibit an increase (up to +166%) under RCP8.5. The strong increase in spring runoff is
185 mainly found in the highest elevation catchments (Fig. 2b), where snowmelt is increased due to higher temperatures that shift
186 the snowmelt season from early summer to spring. However, only 34 out of 93 catchments exhibit robust changes across the
187 climate models. Compared to the high emission scenario, the changes in the low emission scenario (RCP2.6) tend to be more
188 moderate (Fig. 3b). Under RCP2.6, the changes range from -15% to +70% with a slightly positive mean (+3%) across all
189 catchments. 58 catchments show robust changes in spring mean runoff under RCP2.6. Some catchments show negative
190 changes under RCP2.6, but positive changes under RCP8.5. This transition from negative to positive changes is also found in
191 RCP4.5 (Fig. S3).

192

193 **4.1.3 Changes in summer runoff for Switzerland**

194 In summer and under RCP8.5, all catchments show a decrease in mean runoff, ranging from -16% to -59% with mean change
195 of -35% across all catchments (Fig. 2c). Again, more pronounced negative changes are found in higher-elevation catchments,
196 where summer runoff is projected to decrease by up to half of the runoff of the reference period. All catchments except one
197 agree on the sign of change among the climate models. Under RCP2.6, the signs and magnitude of the changes are less clear
198 (Fig. 3c). The average change across all catchments is negative (-6%), but mean runoff change ranges from -26% to +4%. Just
199 9 catchments show positive but non-robust mean runoff changes in summer whereas 34 catchments yield robust negative signs.

200 **4.1.4 Changes in autumn runoff for Switzerland**

201 In autumn, all catchments but one show decreasing runoff under RCP8.5 (Fig. 2d). Changes range from -36% to +4% with an
202 average change of -19% across all catchments. More than 50% of the catchments reveal robust changes. In contrast to the
203 changes in summer, the autumn runoff tends to decrease more strongly in the lower-lying catchments than in the higher-
204 elevation catchments. Under RCP2.6, the changes are much smaller than under RCP8.5, with an average change of -5%
205 (ranging from -20% to +10%; Fig. 3d). The changes are also less robust, with only 17 catchments showing good agreement on
206 the sign of change.

207

208 **4.1.5 Changes in annual runoff for Switzerland**

209 Despite the increases in winter and partial increases in spring, the annual runoff is projected to decrease by 8% (range: -23%
210 to +4%) under RCP8.5 (Fig. 2e) and by 2% (range: -13% to +11%) under RCP2.6 (Fig. 3e) on average. Whereas 82 out of 93
211 catchments show negative changes under RCP8.5, only 65 catchments exhibit negative changes under RCP2.6. However, the
212 robustness of the sign of change in the annual mean runoff is weaker than that of the seasonal mean changes.

213

214 **4.1.6 Elevation dependence of mean seasonal and annual changes in runoff**

215 Considering the results above, changes in seasonal and annual mean runoff are heavily dependent on the mean elevation of the
216 catchment. This dependence is highlighted in Fig. 4, where changes in runoff are plotted against mean altitude of the
217 catchments. The higher-elevation catchments generally show larger changes in winter, spring, and summer than the lower
218 elevation catchments. For autumn and annual runoff, no distinct pattern can be seen. Under RCP8.5 and for catchments with
219 mean altitude below 1500 m a.s.l., the average change is +27% in winter, -5% in spring, -31% in summer, -21% in autumn,
220 and -8% in the annual mean (Fig. 4a). For catchments with mean elevation above 1500 m a.s.l., runoff changes on average by
221 +77% in winter, by +28% in spring, by -41% in summer, by -15% in autumn, and by -9% in the annual mean. However, the
222 changes in the higher-elevation catchments are less pronounced under RCP2.6, with an average change in catchments below
223 1500 m a.s.l. of +5% in winter, -6% in spring, -4% in summer, -6% in autumn, and -4% in annual mean runoff (Fig. 4b). In
224 higher-elevation catchments (>1500 m a.s.l.), the mean changes under RCP2.6 amount to +24% in winter, +16% in spring, -
225 9% in summer, -4% in autumn, and -0.6% across the year.

226 **4.2 Changes in the runoff regime of six representative catchments**

227 Changes in runoff regime for six example catchments representing typical runoff regime types in Switzerland are depicted in
228 Fig. 5. These are the catchments highlighted in Fig. 1: Rosegbach–Pontresina, Kander–Hondrich, Plessur–Chur, Emme–
229 Emmenmatt, Venoge–Ecublens, and Verzasca–Lavertezzo. The runoff regimes with absolute monthly mean runoff presented
230 in this chapter help to interpret the relative changes of seasonal and annual runoff means discussed in Section 4.1. Figures with
231 Pardé coefficients and runoff regimes for each RCP separately can be found in the Supplement (Figs. S7-S11).

232 **4.2.1 Highly glaciated catchment: Rosegbach–Pontresina**

233 The glaciated catchment, Rosegbach–Pontresina, exhibits a pronounced decrease in July, August, and September (-3.1 to -6.7
234 mm/day) and a small increase in winter runoff (+0.5 to +1 mm/day) under RCP8.5 (Fig. 5a). Monthly mean runoff does not
235 change significantly in October. The typical glacier runoff regime with low flows in winter and peak flows in summer in the
236 reference period changes to a nival regime with a peak runoff shifted from July or August to June under RCP8.5 (Fig. 5a).
237 Although there are large relative (+200%) increases in the winter months, the contribution of winter runoff to the total runoff
238 remains small. The mean runoff between June and September drops dramatically due to the absence of snow and glacier melt
239 contributions. However, runoff in late spring and summer remains the major contributor to the annual volume. Under RCP2.6,
240 the changes for winter, spring, and autumn runoff are small (+0.1 to +0.7 mm/day). For July, August, and September, the
241 runoff decreases significantly (-1.7 to -3.5 mm/day), and the summer peak shifts from July or August to June or July. The
242 change in summer runoff under RCP2.6 is approximately halved compared to the changes under RCP8.5.

243 **4.2.2 Slightly glaciated catchment: Kander–Hondrich**

244 A similar behavior with a shift of the peak in the runoff regime from summer to late spring/early summer is also found for the
245 runoff regime in the slightly glaciated catchment, Kander–Hondrich, under RCP8.5 (Fig. 5b). The regime is characterized by
246 a peak in early summer runoff in the reference period. Under RCP8.5, the summer and early autumn runoff (July, August,
247 September) decreases significantly (-1.8 to -3.2 mm/day) whereas winter runoff increases (+0.8 to +1.3 mm/day). This leads
248 to a flattening of the runoff regime curve, resulting in similar contributions of winter and summer runoff to the annual volume.
249 Under RCP2.6, there is also a decrease in July, August, and September (-0.7 to -1.1 mm/day), but less pronounced than the
250 decrease under RCP8.5.

251 **4.2.3 Nival catchment: Plessur–Chur**

252 The nival regime of the river Plessur–Chur shows a shift in peak flow from June to May under both emission scenarios (Figure
253 5c). Due to the increase in winter months (+0.4 to +0.9 mm/day) and decrease in summer runoff (-0.9 to -2.1 mm/day), the
254 regime curve flattens under RCP8.5. The results also show increasing winter runoff and decreasing summer runoff under
255 RCP2.6 but far less pronounced than under RCP8.5.

256 **4.2.4 Nival-to-pluvial catchment: Emme–Emmenmatt**

257 In the reference period, the runoff regime in the nival-to-pluvial catchment, Emme–Emmenmatt, shows a peak in spring and
258 early summer due to snowmelt and stable mean runoff from August to February (Figure 5d). By end of the century under
259 RCP8.5, the peak runoff in spring almost disappears. The runoff decreases strongly in the summer months (-0.7 to -1.3
260 mm/day) and less in the autumn months (-0.1 to -1.2 mm/day).

261 **4.2.5 Pluvial catchment: Venoge–Ecublens**

262 The shape of the runoff regime curve for the pluvial river Venoge–Ecublens remains the same for the reference period and the
263 two emission scenarios with higher runoff in winter and lower runoff in summer (Figure 5e). Although the regime changes
264 only marginally under RCP2.6, the amplitudes under RCP8.5 become more distinct, with higher winter runoff (+0.3 to +0.9
265 mm/day) and lower summer runoff (-0.2 to -0.3 mm/day) than in the reference period. However, in comparison to other
266 catchments, this change is smaller.

267 **4.2.6 Southern Alpine catchment: Verzasca–Lavertezzo**

268 The southern Alpine catchment, Verzasca–Lavertezzo, shows a two peaked runoff regime, with a first runoff peak in late
269 spring and a second runoff peak in autumn in the reference period (Figure 5f). This pattern is still present at the end of the
270 century under both scenarios. However, the amplitudes of the peaks are less pronounced under RCP8.5 because of increasing
271 winter runoff and decreasing spring (-0.2 to -1.9 mm/day) and summer (-1.4 to -1.7 mm/day) runoff.

272

273 Summarizing the differences under RCP8.5 and RCP2.6 shows that the sign of change under RCP2.6 is equal to the sign under
274 RCP8.5 in almost all months and catchments. Comparisons with RCP4.5 (see Fig. S5 in the Supplement) show that the
275 magnitude of changes increases with the strength of the emission scenario and the distance in time. Also, the model agreement
276 on the direction of change among the climate models is weaker under the low emission scenario, RCP2.6, than under RCP8.5.

277 **4.3 Time of emergence of changes in the seasonal and annual means**

278 The time of emergence, when at least 66% of the models under RCP8.5 agree on significant changes in the distribution of
279 seasonal and annual means, is depicted in Fig. 6. In winter, 45 out of 93 catchments show a time of emergence in the 21st
280 century (Fig. 6a). An elevation dependence can be identified: times of emergence are earlier in higher elevated catchments and
281 later in lower-lying catchments. The high Alpine catchments show particularly early times of emergence, with the end of the
282 period 2017-2046 the earliest. The mean elevation of catchments with times of emergence earlier than 2065 is higher than
283 1500 m a.s.l., with one exception. Among the 48 catchments that do not show a time of emergence, 46 catchments have mean
284 elevations lower than 1200 m a.s.l.

285 In summer, 73 catchments exhibit a time of emergence and again generally an earlier time of emergence for higher-elevation
286 catchments (Fig. 6c). Catchments showing a time of emergence earlier than 2065 are all located in mountainous areas with
287 mean elevations higher than 1500 m a.s.l., again with one exception. The earliest time of emergence in summer is found for
288 the year 2043 (period 2014-2043). Catchments without time of emergence show a mean elevation lower than 1000 m a.s.l.,
289 with one exception.

290 In spring, only 20 catchments exhibit a time of emergence (Fig. 6b). Again, significant changes in the distribution are mainly
291 found in the higher Alpine catchments (above 1500 m a.s.l.). Only 14 catchments out of 93 exhibit a time of emergence in
292 autumn (Fig. 6d). In contrast to the other seasons, there is no clear elevation pattern distinguishable in autumn.

293 In all seasons, some of the catchments do not show a time of emergence, meaning that there is no statistically significant
294 change in the distribution of the seasonal means. Clear patterns of significant changes in the distribution of seasonal means are
295 mainly found in winter and summer.

296 For the annual means, 11 catchments reveal a time of emergence in the 21st century (Fig. 6e). Time of emergence in the annual
297 mean is not restricted to high Alpine catchments, but only two catchments below 1500 m a.s.l. show significant changes, and
298 that after 2095.

299 Due to the definition of time of emergence as the last year of a moving window in which the Kolmogorov-Smirnov test is
300 rejected for the first time, time of emergence is not necessarily given for all periods after the first detection. Fig. 7 shows the
301 temporal evolution of the time of emergence for the seasons under RCP8.5. Most of the catchments show persistent significant
302 changes after the first detection of a time of emergence. However, few catchments do not reveal persistent significant changes
303 after the first detection due to non-linear changes in the runoff response to climate change. The problem of non-constant

304 rejections affects 17 catchments in winter, 3 catchments in spring, 25 catchments in summer, and 6 catchments in autumn.
305 Most of these catchments show a persistent time of emergence for the rest of the century a few years after the first detection.

306 **4.4 Changes in seasonal means with increasing global mean temperatures**

307 Changes in the multimodel median of seasonal and annual mean runoff for different levels of global warming are shown in
308 Fig. 8 for warming targets +1.5°C, +2°C, and +3°C. Generally, the patterns of change with levels of global warming are similar
309 to the patterns for the two emission scenarios. Both, the range of change between the catchments and the climate model
310 agreement increase with higher global warming levels.

311 In winter, the mean runoff change across all catchments is +17% for a global warming of +1.5°C, +23% for +2°C, and +35%
312 for +3°C (Fig. 8 a-c). Stronger global warming increases not only the mean but also the range of change across the catchments.
313 At +1.5°C global warming, the range across all catchments is between -1% and +53%, but at +3°C global warming, the range
314 is +5% to +127%. Whereas two catchments show slightly negative changes at +1.5°C warming, all catchments show positive
315 changes for higher warming levels. Moreover, the agreement across the climate models per catchment increases with increasing
316 warming.

317 The mean change in spring runoff among all catchments is +4% for +1.5°C global warming, +6% for 2°C, and +10% for 3°C
318 (Fig. 8 d-f). The changes in spring vary with elevation (see results in section 4.1), with positive changes mainly for the higher
319 elevated catchments and negative changes for lower-lying catchments. Even though the average across all catchments changes
320 little between warming levels, the range of change across the catchments and the model agreement per catchment increases
321 with higher warming levels, and thus the regional, elevation-dependent patterns become more pronounced.

322 In summer, the average among all catchments is -7% for 1.5°C global warming, -13% for 2°C, and -23% for 3°C (Fig. 8g-j).
323 Again, the ranges across catchments and the model agreement increase with higher warming levels. Compared to summer, the
324 decrease in autumn runoff is smaller, with an average across all catchments of -9% for 1.5°C global warming, -7% for 2°C,
325 and -13% for 3°C. Most catchments, 89 out of 93, show a negative multimodel median for both 1.5°C and 3°C warming levels.
326 Although only three catchments show a model agreement of more than 90% for 1.5°C global warming, 43 catchments show
327 robust model agreement on the direction of change for 3°C global warming.

328 For annual mean runoff, the mean among all catchments is +0.2% for 1.5°C, -0.9% for 2°C, and -3.7% for 3°C. Despite the
329 slightly positive sign for 1.5°C warming, 53 catchments out of 93 show negative changes. This number increases to 66
330 catchments for 2°C and to 77 catchments for 3°C. Moreover, model agreement increases from 3 catchments with robust model
331 agreement for 1.5°C to 17 catchments for 3°C.

332

333 **5 Discussion**

334 Winter runoff is projected to increase in Switzerland due to enhanced winter precipitation and increasing temperatures with
335 climate change (CH2018, 2018). The higher temperatures result in more liquid precipitation and less solid precipitation in
336 winter. This leads to less snow accumulation and thus to more direct runoff in winter. The relative changes in winter mean
337 runoff in glaciated catchments are very high but still negligible as a contribution to the annual runoff volume. In nival and
338 pluvial catchments, the contribution of winter runoff to the annual volume increases.

339 In spring, runoff in glaciated and nival catchments is projected to increase due to increased snowmelt, particularly in early
340 spring. The combination of reduced snow accumulation in winter and earlier snowmelt shifts the peak in runoff regime to a
341 month earlier. Lower-lying catchments show decreasing runoff in spring due to reduced snow accumulation in winter, which
342 reduces the snow available for snowmelt.

343 The summer runoff in Switzerland generally decreases with climate change. The processes governing this decrease differ with
344 location and elevation of the catchments. In lower-lying catchments, reduced summer precipitation and enhanced
345 evapotranspiration result in decreasing runoff in Switzerland. In high Alpine regions, where summer snowmelt and glacier
346 melt dominate the runoff generation in the reference period, reduced snowpack and glacier retreat amplify the decrease in
347 summer runoff. The large model spread in glaciated catchments in summer stems from the sensitivity of the runoff response
348 to the glaciation of the catchment. In catchments where not all climate projections result in a complete disappearance of
349 glaciers, the model spread increases, and so future glacier retreat is a major source of uncertainty.

350 Autumn runoff is also projected to decrease due to increased evapotranspiration and slightly reduced precipitation in most
351 catchments (CH2018, 2018). In a few of the eastern catchments, autumn precipitation slightly increases, but the effect of
352 enhanced evapotranspiration and the reduced contribution from glacier and snow melt are more dominant, and autumn runoff
353 also decreases in those catchments. Again, this decrease is amplified in catchments where glacier melt is important in early
354 autumn under current climate conditions. However, the decrease in autumn runoff is most noticeable in early autumn, whereas
355 in late autumn, at the end of October and November, the changes are less significant and can even change direction to positive
356 values. This pattern is mainly found in very high Alpine catchments, where late autumn precipitation can fall more often as
357 rain instead of snow and thus cannot be stored as snowpack. For the annual mean, a decrease in runoff was found in most
358 catchments, but with less robust signs among the climate models. This leads to the conclusion that on the annual average, less
359 water will be available in Swiss rivers. The shift in seasonality and thus a shift in the seasonal availability of water will impact
360 many economic sectors. For example, increasing winter runoff may be beneficial for energy production, but decreasing
361 summer runoff may lead to limitations in irrigation, particularly in lower-lying catchments where agricultural irrigation plays
362 a crucial role.

363 In most catchments and seasons, the sign of change is the same for the high-emission scenario, RCP8.5, and the low-emission
364 scenario, RCP2.6. Changes in seasonality or in the runoff regime are amplified by higher emissions and thus by increasing
365 global mean temperatures. This amplification due to increased emissions and with intensified global warming shows the large

366 benefits of mitigation. By mitigating climate change and following the RCP2.6 pathway, the magnitude of change can be
367 reduced or even avoided, depending on the season and the runoff regime. However, some of the seasonal changes can no
368 longer be avoided by mitigation actions, particularly in high Alpine catchments with glacier influence. Responsible planners
369 and policy makers need to adapt to this shift in the seasonal availability of water in our rivers.

370 Previous studies on climate change impacts on the runoff regime in Switzerland (e.g., Koeplin et al., 2012, 2014; Horton et
371 al., 2006) were driven by other emission scenarios, other and fewer climate model chains, different methods of postprocessing
372 the climate model output, and/or different hydrological models and calibration. Despite these differences, the sign of change
373 in those studies agrees in most seasons with the signs in this study. A pronounced dependence of runoff changes on elevation
374 was also found by Koeplin et al. (2012), with lower-lying catchments being less affected by climate change. The largest
375 difference concerns the annual mean runoff. Koeplin et al. (2012) found an increase (up to +50%) in annual runoff for high-
376 elevation catchments and no change for lower-lying catchments in the annual volume. In contrast, our study projects a decrease
377 in the annual mean runoff not only for high-elevation catchments but also for most of the lower-lying catchments. However,
378 not all catchments show a robust decrease among the climate model chains. This difference between our results and previous
379 studies may arise from the use of the most recent generation of Swiss climate change scenarios (CH2018), which project
380 slightly different precipitation changes with less summer drying and wetter winters (CH2018, 2018) than the previous
381 generation of scenarios, CH2011, which was used by Koeplin et al. (2012). The different handling of glacier melt processes
382 and the new projections of glacier extents (Zekollari et al., 2019) may also result in slightly different projections in glaciated
383 catchments. The Hydro-CH2018-Runoff ensemble uses transient glacier projections that are updated every 5 years in the
384 hydrological model. Koeplin et al. (2012) used static glacier projections for 30 years. The largest uncertainties in glaciated
385 catchments were also found by Addor et al. (2014) due to the different handling of glacier extents and resulting glacier melt.
386 Furthermore, the difference in the input data, transient projections versus delta change projections with same baseline time
387 series for the reference period, may add to the different signs.

388 Even though not all catchments show a time of emergence in the 21st century, significant changes in the distribution of seasonal
389 means emerge particularly early in high-elevation catchments. This is due to the importance of snow and glacier melt for
390 Alpine runoff regimes. With climate change, the influence of snow and glacier melt decreases due to higher temperatures and
391 its subsequent glacier retreat. Lower-lying catchments generally show a later time of emergence. This may arise from the large
392 interannual variability in pluvial catchments in the reference period and in future periods. Koeplin et al. (2014) also assessed
393 a time of emergence for Swiss catchments but only based on significant changes for two scenario periods and 10 climate
394 models. Since their climate models were postprocessed with a delta change approach, only the natural variability of the
395 reference period is reflected in their simulations. Despite these differences in methodology and data, they also found earlier
396 time of emergence in winter and summer for high-elevation catchments. Although the definition of time of emergence applied
397 here is commonly used in other climate change studies (e.g., Mahlstein et al., 2011), this definition also has disadvantages. For
398 example, if the rejection of the null hypothesis is unstable in its temporal evolution, the time of emergence may be determined
399 too early. This has been shown for some of the catchments in the present study. However, in most catchments a persistent

400 detection of time of emergence (p -value < 0.05) in the distribution of the seasonal means was found shortly after the first
401 detection of a time of emergence.

402 The findings of this study concern runoff regime changes for Switzerland. The pronounced changes in high-elevation
403 catchments highlight the important influence of temperature changes on snow- and glacier-melt-driven catchments, and thus
404 they also indicate that similar runoff responses may be found in other snow- and glacier-dominated regions. The results for
405 lower-lying catchments, which are mainly driven by evapotranspiration and precipitation changes, may not be directly
406 transferable to other regions. The runoff response in such catchments depends on local precipitation patterns and their changes
407 under climate change.

408 Various sources of uncertainty affect our results. A detailed discussion of the uncertainties for the simulations is provided in
409 Muelchi et al. (2021). Uncertainties arise from all the steps in the modelling chain: the scenarios of greenhouse gas
410 concentrations, the climate models and their boundary and initial conditions, the postprocessing method (Gutiérrez et al.,
411 2018), the hydrological model (Addor et al., 2014) and its calibration, and the underlying glacier projections; all these need to
412 be considered in adaptation planning (e.g., Wilby and Dessai, 2010). Working with three emission scenarios and an ensemble
413 of climate models partly addresses some of the uncertainty issues. Muelchi (2021) compared the results for three catchments
414 to simulations from three different versions of hydrological models, and the key findings are robust among both the models
415 and the catchments.

416

417 **6 Conclusions**

418 Changes in runoff regimes and their time of emergence were assessed with the new hydrological scenarios for 93 catchments
419 in Switzerland. This study is based on the most recent generation of climate change scenarios, postprocessed with quantile
420 mapping. Compared to previous studies on runoff regime changes, the results show similar signs of change for most seasons.
421 The largest differences were found for the high-elevation catchments, which is likely due to the transient characteristic of the
422 simulations and the implementation of transient glacier projections driven by the same climate model chains.

423 In general, winter runoff is projected to increase and summer runoff to decrease in Switzerland. The sign of change is robust
424 across catchments, but the magnitude of change is more pronounced for high-elevation catchments. Particularly in summer,
425 when snow and glacier melt play an important role in runoff generation, glaciated catchments will face a strong decrease in
426 runoff due to the retreating glaciers. In rainfall-dominated catchments, the changes are also often robust, but at a lower
427 magnitude. Whereas the higher elevated catchments show increasing spring runoff due to earlier snowmelt, the pluvial
428 catchments in the lowlands will face decreasing spring runoff. A decrease in runoff is also found for autumn and annual mean
429 runoff in most catchments. These seasonal patterns amplify with global warming and with higher emission scenarios.
430 Furthermore, the model agreement among the climate models increases with the strength of the emission scenario and the
431 distance in time. A time of emergence of seasonal mean runoff was mainly found in summer and winter and for only a few

432 catchments in spring and autumn. Early times of emergence, before 2060 in winter and before 2065 in summer, were found
433 for catchments above 1500 m a.s.l.. Significant changes in catchments below 1500 m a.s.l. emerge later in the century.
434 However, not all catchments show a time of emergence in all seasons.

435 The amplification of changes by enhanced global warming highlights the importance of climate change mitigation. By
436 mitigating climate change and following the RCP2.6 pathway to keep global warming below 2 °C, the magnitude of change
437 can be reduced substantially. The large decrease in summer runoff in glaciated catchments can be strongly dampened but not
438 avoided entirely because glacier retreat is projected even for the low emission scenario. The present study can help to support
439 adaptation planning in various sectors by presenting detailed information on changes in mean runoff.

440
441

442 **Data availability**

443 The data used in this study is available under <https://doi.org/10.5281/zenodo.3937485> (Muelchi et al., 2020).

444

445 **Author contributions**

446 RM performed the analysis of the results and drafted the manuscript. JS, OR, RW, and OM helped in interpreting the results.

447 All authors reviewed the resulting data and assisted with paper writing.

448 **Competing interests**

449 The authors declare that they have no conflict of interest.

450 **Acknowledgements**

451 Authors would like to thank Harry Zekollari for processing and providing the glacier projections used in this study. We thank
452 MeteoSwiss and FOEN for providing the data necessary for this study and acknowledge the funding of the Swiss Federal
453 Office for the Environment under the project titled Hydro-CH2018.

454

455

456 **References**

- 457 Addor, N., Rössler, O., Köplin, N., Huss, M., Weingartner, R., and Seibert, J.: Robust changes and sources of uncertainty in
458 the projected hydrological regimes of Swiss catchments, *Water Resour. Res.*, 50, 7541-7562,
459 <https://doi.org/10.1002/2014WR015549>, 2014.
- 460 Blöschl, G., Schöner, W., Kroiß, H., Blaschke, A. P., Böhm, R., Haslinger, K., Kreuzinger, N., Merz, R., Parajka, J., Salinas,
461 J.L. and Viglione, A.: Anpassungsstrategien an den Klimawandel für Österreichs Wasserwirtschaft-Ziele und
462 Schlussfolgerungen der Studie für Bund und Länder, *Österr. Wasser- und Abfallw.*, 63, 1-10, <https://doi.org/10.1007/s00506-010-0274-2>, 2011.
- 464 Bosshard, T., Kotlarski, S., Ewen, T., and Schär, C.: Spectral representation of the annual cycle in the climate change signal,
465 *Hydrol. Earth Syst. Sci.*, 15, 2777–2788, <https://doi.org/10.5194/hess-15-2777-2011>, 2011.
- 466 Brunner, M. I., Gurung Björnsen, A., Zappa, M., Zekollari, H., Farinotti, D., and Stähli, M.: Present and future water scarcity
467 in Switzerland: Potential for alleviation through reservoirs and lakes, *Sci. Total Environ.*, 666, 1033-1047,
468 <https://doi.org/10.1016/j.scitotenv.2019.02.169>, 2019a.
- 469 Brunner, M. I., Farinotti, D., Zekollari, H., Huss, M., and Zappa, M.: Future shifts in extreme flow regimes in Alpine regions,
470 *Hydrol. Earth Syst. Sci.*, 23, 11, 4471-4489, <https://doi.org/10.5194/hess-23-4471-2019>, 2019b.
- 471 CH2018: CH2018 – Climate Scenarios for Switzerland, Technical Report, National Centre for Climate Services, Zurich,
472 Switzerland, 271 pp., 2018.
- 473 Etter, S., Addor, N., Huss, M., and Finger, D.: Climate change impacts on future snow, ice and rain runoff in a Swiss mountain
474 catchment using multi-dataset calibration, *J. Hydrol. Regional Studies*, 13, 222-239,
475 <https://doi.org/10.1016/j.ejrh.2017.08.005>, 2017.
- 476 Farinotti, D., Usselman, S., Huss, M., Bauder, A., and Funk, M.: Runoff evolution in the Swiss Alps: Projections for selected
477 high-alpine catchments based on ENSEMBLES scenarios, *Hydrol. Process*, 26, 13, 1909-1924,
478 <https://doi.org/10.1002/hyp.8276>, 2012.
- 479 Fatichi, S., Rimkus, S., Burlando, P., Bordoy, R., and Molnar, P.: High-resolution distributed analysis of climate and
480 anthropogenic changes on the hydrology of an Alpine catchment, *J. Hydrol.*, 525, 362-382,
481 <https://doi.org/10.1016/j.jhydrol.2015.03.036>, 2015.
- 482 Finger, D., Heinrich, G., Gobiet, A., and Bauder, A.: Projections of future water resources and their uncertainty in a glacierized
483 catchment in the Swiss Alps and the subsequent effects on hydropower production during the 21st century, *Water Resour.*
484 *Res.*, 48, <https://doi.org/10.1029/2011WR010733>, 2012.
- 485 Gaetani, M., Janicot, S., Vrac, M., Famién, A. M., and Sultan, B.: Robust assessment of the time of emergence of precipitation
486 change in West Africa, *Sci. Rep.*, 10, 1-10, <https://doi.org/10.1038/s41598-020-63782-2>, 2020.
- 487 Giorgi, F., and Bi, X.: Time of emergence (TOE) of GHG-forced precipitation change hot-spots, *Geophys. Res. Lett.*, 36,
488 <https://doi.org/10.1029/2009GL037593>, 2009.

489 Goler, R. A., Frey, S., Formayer, H., and Holzmann, H.: Influence of climate change on river discharge in Austria, *Meteorol. Z.*, 25, 621-626, <https://doi.org/10.1127/metz/2016/0562>, 2016.

491 Groppelli, B., Soncini, A., Bocchiola, D., and Rosso, R.: Evaluation of future hydrological cycle under climate change
492 scenarios in a mesoscale Alpine watershed of Italy, *Nat.l Hazards Earth Syst. Sci.*, 11, 6, 1769, [https://doi.org/10.5194/nhess-](https://doi.org/10.5194/nhess-11-1769-2011)
493 11-1769-2011, 2011.

494 Gudmundsson, L., Bremnes, J. B., Haugen, J. E., and Engen-Skaugen, T.: Technical Note: Downscaling RCM precipitation to
495 the station scale using statistical transformations – A comparison of methods, *Hydrol. Earth Syst. Sci.*, 16, 3383–3390,
496 <https://doi.org/10.5194/hess-16-3383-2012>, 2012.

497 Gutiérrez, J. M., Maraun, D., Widmann, M., Huth, R., Hertig, E., Benestad, R., Rössler, O., Wibig, J., Wilcke, R., Kotlarski,
498 S., San Martín, D., Herrera, S., Bedia, J., Casanueva, A., Manzananas, R., Iturbide, M., Vrac, M., Dubrovsky, M., Ribalaygua,
499 J., Pórtoles, J., Rätty, O.E., Räisänen, J.A., Hingray, B., Raynaud, D., Casado, M., Ramos, P., Zerenner, T., Turco, M.,
500 Bosshard, T., Stepanek, P., Bartholy, J., Pongracz, R., Keller, D., Fischer, A., Cardoso, R., Soares, P., Czernecki, B., and Pagé,
501 C.: An intercomparison of a large ensemble of statistical downscaling methods over Europe: Results from the VALUE perfect
502 predictor cross-validation experiment, *Int. J. Climatol.*, 39, 3750–3785, <https://doi.org/10.1002/joc.5462>, 2019.

503 Hanzer, F., Förster, K., Nemeč, J., and Strasser, U.: Projected cryospheric and hydrological impacts of 21st century climate
504 change in the Ötztal Alps (Austria) simulated using a physically based approach, *Hydrol. Earth Syst. Sci.*, 22, 2, 1593-1614,
505 <https://doi.org/10.5194/hess-22-1593-2018>, 2018.

506 Hattermann, F. F., Huang, S., & Koch, H.: Climate change impacts on hydrology and water resources, *Meteorol. Z.*, 24, 2,
507 201-211, [10.1127/metz/2014/0575](https://doi.org/10.1127/metz/2014/0575), 2015.

508 Horton, P., Schaeffli, B., Mezghani, A., Hingray, B., and Musy, A.: Assessment of climate-change impacts on alpine discharge
509 regimes with climate model uncertainty, *Hydrol. Process.*, 20, 2091–2109, <https://doi.org/10.1002/hyp.6197>, 2006.

510 Huss, M., Farinotti, D., Bauder, A., and Funk, M.: Modelling runoff from highly glacierized alpine drainage basins in a
511 changing climate, *Hydrol. Process.*, 22, 19, 3888-3902, <https://doi.org/10.1002/hyp.7055>, 2008.

512 Huss, M., Zemp, M., Joerg, P. C., and Salzmann, N.: High uncertainty in 21st century runoff projections from glacierized
513 basins, *J. Hydrol.*, 510, 35-48, <https://doi.org/10.1016/j.jhydrol.2013.12.017>, 2014.

514 IPCC: Climate Change 2013: The Physical Science Basis, in: Contribution of Working Group I to the Fifth Assessment Report
515 of the Intergovernmental Panel on Climate Change, edited by Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J.
516 Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley, Cambridge University Press, Cambridge, United Kingdom and New
517 York, NY, USA, 1535 pp., 2013.

518 IPCC: Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of
519 Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by Field, C.B.,
520 V.R. Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B.
521 Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R. Mastrandrea, and L.L. White, Cambridge University Press, Cambridge,
522 United Kingdom and New York, NY, USA, 1132 pp., 2014a.

523 IPCC: Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working
524 Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by Barros, V.R., C.B.
525 Field, D.J. Dokken, M.D. Mastrandrea, K.J. Mach, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma,
526 E.S. Kissel, A.N. Levy, S. MacCracken, P.R. Mastrandrea, and L.L. White, Cambridge University Press, Cambridge, United
527 Kingdom and New York, NY, USA, 688 pp., 2014b.

528 Jacob, D., Petersen, J., Eggert, B., Alias, A., Christensen, O. B., Bouwer, L. M., Braun, A., Colette, A., Déqué, M., Georgievski,
529 G., Georgopoulou, E., Gobiet, A., Menut, L., Nikulin, G., Haensler, A., Hempelmann, N., Jones, C., Keuler, K., Kovats, S.,
530 Kröner, N., Kotlarski, S., Kriegsmann, A., Martin, E., van Meijgaard, E., Moseley, C., Pfeifer, S., Preuschmann, S.,
531 Radermacher, C., Radtke, K., Rechid, D., Rounsevell, M., Samuelsson, P., Somot, S., Soussana, J.-F., Teichmann, C.,
532 Valentini, R., Vautard, R., Weber, B., and Yiou, P.: EUROCORDEX: new high-resolution climate change projections for
533 European impact research, *Reg. Environ. Change*, 14, 563–578, <https://doi.org/10.1007/s10113-013-0499-2>, 2014.

534 James, R., Washington, R., Schleussner, C. F., Rogelj, J., and Conway, D.: Characterizing half-a-degree difference: a review
535 of methods for identifying regional climate responses to global warming targets, *Wires. Clim. Change*, 8, e457,
536 <https://doi.org/10.1002/wcc.457>, 2017.

537 Jenicek, M., Seibert, J., and Staudinger, M.: Modeling of future changes in seasonal snowpack and impacts on summer low
538 flows in alpine catchments, *Water Resour. Res.*, 54, 1, 538-556, <https://doi.org/10.1002/2017WR021648>, 2018.

539 Keller, L., Rössler, O., Martius, O., and Weingartner, R.: Delineation of flood generating processes and their hydrological
540 response, *Hydrol. Process*, 32, 2, 228-240, <https://doi.org/10.1002/hyp.11407>, 2018.

541 Köplin, N., Schädler, B., Viviroli, D., and Weingartner, R.: Relating climate change signals and physiographic catchment
542 properties to clustered hydrological response types, *Hydrol. Earth Syst. Sci.*, 16, 7, 2267-2283, <https://doi.org/10.5194/hess-16-2267-2012>, 2012.

544 Köplin, N., Rössler, O., Schädler, B., and Weingartner, R.: Robust estimates of climate-induced hydrological change in a
545 temperate mountainous region, *Climatic Change*, 122, 171-184, <https://doi.org/10.1007/s10584-013-1015-x>, 2014.

546 Kotlarski, S., Keuler, K., Christensen, O. B., Colette, A., Déqué, M., Gobiet, A., Goergen, K., Jacob, D., Lüthi, D., Van
547 Meijgaard, E., Nikulin, G., Schär, C., Teichmann, C., Vautard, R., Warrach-Sagi, K., and Wulfmeyer, V.: Regional climate
548 modeling on European scales: A joint standard evaluation of the EURO-CORDEX RCM ensemble, *Geosci. Model Dev.*, 7,
549 1297–1333, <https://doi.org/10.5194/gmd-7-1297-2014>, 2014.

550 Leng, G., Huang, M., Voisin, N., Zhang, X., Asrar, G. R., and Leung, L. R.: Emergence of new hydrologic regimes of surface
551 water resources in the conterminous United States under future warming, *Environ. Res. Lett.*, 11, 11, 114003,
552 <https://doi.org/10.1088/1748-9326/11/11/114003>, 2016.

553 Mahlstein, I., Knutti, R., Solomon, S., and Portmann, R. W.: Early onset of significant local warming in low latitude countries,
554 *Environ. Res. Lett.*, 6, 3, 034009, [10.1088/1748-9326/6/3/034009](https://doi.org/10.1088/1748-9326/6/3/034009), 2011.

555 Mastrandrea, M.D., Field, C.B., Stocker, T.F., Edenhofer, O., Ebi, K.L., Frame, D.J., Held, H., Kriegler, E., Mach, K.J.,
556 Matschoss, P.R., Plattner, G.-K., Yohe, G.W., and Zwiers, F.W.: Guidance Note for Lead Authors of the IPCC Fifth
557 Assessment Report on Consistent Treatment of Uncertainties, Intergovernmental Panel on Climate Change (IPCC), 2010.

558 Milano, M., Reynard, E., Köplin, N., and Weingartner, R.: Climatic and anthropogenic changes in Western Switzerland:
559 Impacts on water stress, *Sci. Total Environ.*, 536, 12-24, <https://doi.org/10.1016/j.scitotenv.2015.07.049>, 2015.

560 Morice, C. P., Kennedy, J. J., Rayner, N. A., and Jones, P. D.: Quantifying uncertainties in global and regional temperature
561 change using an ensemble of observational estimates: The HadCRUT4 data set, *J. Geophys. Res-Atmos*, 117, D8,
562 <https://doi.org/10.1029/2011JD017187>, 2012.

563 Muelchi, R., Rössler, O., Schwanbeck, J., Weingartner, R., and Martius, O.: Hydro-CH2018-Runoff ensemble (version v1),
564 Zenodo, <http://doi.org/10.5281/zenodo.3937485>, 2020.

565 Muelchi, R.: Future changes of Swiss river runoff and extreme vertically integrated moisture transport, Ph.D thesis, University
566 of Bern, Switzerland, 198 pp., 2021.

567 Muelchi, R., Rössler, O., Schwanbeck, J., Weingartner, R., and Martius, O.: An ensemble of daily simulated runoff data (1981–
568 2099) under climate change conditions for 93 catchments in Switzerland (Hydro-CH2018-Runoff ensemble), *Geosci. Data J.*,
569 2021.

570 Nilson, E., Krahe, P., Klein, B., Lingemann, I., Horsten, T., Carambia, M., Larina, M., and Maurer, T.: Auswirkungen des
571 Klimawandels auf das Abflussgeschehen und die Binnenschifffahrt in Deutschland. Schlussbericht KLIWAS-Projekt 4.01,
572 BfG, Koblenz, 10.5675/Kliwas_43/2014_4.01, 2014.

573 Prash, M., Marke, T., Strasser, U., and Mauser, W.: Large scale integrated hydrological modelling of the impact of climate
574 change on the water balance with DANUBIA, *Adv. Sci. Res.*, 7, 61, <https://doi.org/10.5194/asr-7-61-2011>, 2011.

575 Rössler, O., Kotlarski, S., Fischer, A. M., Keller, D., Liniger, M., and Weingartner, R.: Evaluating the added value of the new
576 Swiss climate scenarios for hydrology: An example from the Thur catchment, *Climate Services*, 13, 1-13,
577 <https://doi.org/10.1016/j.cliser.2019.01.001>, 2019.

578 Ruiz-Villanueva, V., Stoffel, M., Bussi, G., Francés, F., and Bréthaut, C.: Climate change impacts on discharges of the Rhone
579 River in Lyon by the end of the twenty-first century: model results and implications, *Reg. Environ. Change*, 15, 3, 505-515,
580 <https://doi.org/10.1007/s10113-014-0707-8>, 2015.

581 Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An overview of CMIP5 and the experiment design, *B. Am. Meteorol. Soc.*,
582 93, 4, 485-498, <https://doi.org/10.1175/BAMS-D-11-00094.1>, 2012.

583 Tecklenburg, C., Francke, T., Kormann, C., and Bronstert, A.: Modeling of water balance response to an extreme future
584 scenario in the Ötztal catchment, Austria, *Adv. Geosci.*, 32, 63-68, <https://doi.org/10.5194/adgeo-32-63-2012>, 2012

585 Teutschbein, C., and Seibert, J.: Bias correction of regional climate model simulations for hydrological climate-change impact
586 studies: Review and evaluation of different methods. *Journal of Hydrology*, 456, 12-29,
587 <https://doi.org/10.1016/j.jhydrol.2012.05.052>, 2012.

588 Vidal, J.-P., Hingray, B., Magand, C., Sauquet, E., and Ducharne, A.: Hierarchy of climate and hydrological uncertainties in
589 transient low-flow projections, *Hydrol. Earth Syst. Sci. Disc.*, 20, 3651-3672, <https://doi.org/10.5194/hess-20-3651-2016>,
590 2016.

591 Viviroli, D., Zappa, M., Gurtz, J., and Weingartner, R.: An introduction to the hydrological modelling system PREVAH and
592 its pre and post-processing-tools, *Environ. Model. Softw.*, 24, 1209–1222, <https://doi.org/10.1016/j.envsoft.2009.04.001>,
593 2009.

594 Weber, M., Braun, L., Mauser, W., and Prasch, M.: Contribution of rain, snow-and icemelt in the Upper Danube discharge
595 today and in the future, *Geogr. Fis. Dinam. Quat*, 33, 2, 221-230, 2010.

596 Wijngaard, R. R., Helfricht, K., Schneeberger, K., Huttenlau, M., Schneider, K., and Bierkens, M. F.: Hydrological response
597 of the Ötztal glacierized catchments to climate change, *Hydrol. Res.*, 47, 5, 979-995, <https://doi.org/10.2166/nh.2015.093>,
598 2016.


599 Wilby, R. L., and Dessai, S.: Robust adaptation to climate change, *Weather*, 65, 180-185, <https://doi.org/10.1002/wea.543>,
600 2010.

601 Zekollari, H., Huss, M., and Farinotti, D.: Modelling the future evolution of glaciers in the European Alps under the
602 EUROCORDEX RCM ensemble, *The Cryosphere*, 13, 1125–1146, <https://doi.org/10.5194/tc-13-1125-2019>, 2019.

603

Table 1: List of GCM-RCM chains used, their initial resolution, and the RCP available.

GCM	RCM	RCP8.5		RCP4.5		RCP2.6	
		EUR -11	EUR -44	EUR -11	EUR -44	EUR -11	EUR -44
ICHEC-EC-EARTH	KNMI-RACMO22E		X		X		
	DMI-HIRMAM5	X		X		X	
	CLMcom-CCLM4-8-17	X		X			
	CLMcom-CCLM5-0-6		X				
	SMHI-RCA4	X		X		X	
MOHC-HadGEM2-ES	CLMcom-CCLM4-8-17	X		X			
	CLMcom-CCLM5-0-6		X				
	KNMI-RACMO22E		X		X		X
	SMHI-RCA4	X		X			X
MPI-M-MPI-ESM-LR	CLMcom-CCLM4-8-17	X		X			
	CLMcom-CCLM5-0-6		X				
	SMHI-RCA4	X		X			X
	MPI-CSC-REMO2009-2	X		X		X	
MIROC-MIROC5	CLMcom-CCLM5-0-6		X				
	SMHI-RCA4		X		X		X
CCCma-CanESM2	SMHI-RCA4		X		X		
CSIRO-QCCCE-CSIRO-Mk3-6-0	SMHI-RCA4		X		X		
IPSL-IPSL-CM5A-MR	SMHI-RCA4	X		X			
NCC-NorESM1-M	SMHI-RCA4		X		X		X
NOAA-GFDL-GFDL-ESM2M	SMHI-RCA4		X		X		




606

607 **Figure 1: Overview of the study region and the location of the gauging stations in Switzerland. Shadings indicate the mean elevation**
 608 **of the respective catchment. Red contours indicate the six example catchments: Rosegbach–Pontresina (1), Kander–Hondrich (2),**
 609 **Plessur–Chur (3), Emme–Emmenmatt (4), Venoge–Ecublens (5), Verzasca–Lavertezzo (6).**


610

611

612




614
 615 **Figure 2: Multimodel median of seasonal and annual mean changes of runoff under RCP8.5 by 2070–2099 for winter (a), spring (b),**
 616 **summer (c), autumn (d), and annual means (e). Black circles indicate catchments whose direction of change agrees across at least**
 617 **90% the models.**



619


620 **Figure 3: Equivalent to Figure 2 but with projections for RCP2.6**

621




622
 623 **Figure 4: Elevation dependence of the multimodel median (dots) of seasonal and annual mean changes of runoff by 2070–2099 under**
 624 **RCP8.5 (a) and RCP2.6 (b). Colors indicate the mean elevation of the 93 catchments.**

625
 626
 627




628
 629
 630
 631
 632
 633

Figure 5: Runoff regimes for the six representative catchments: Rosegbach (a), Kander (b), Plessur (c), Emme (d), Venoge (e), and Verzasca (f). Thick lines represent the multimodel median for the reference period (grey), for 2070–2099 under RCP2.6 (turquoise), and for 2070–2099 under RCP8.5 (orange). Shadings show the full model range for each RCP.




634 **Figure 6: Time of emergence for winter (a), spring (b), summer (c), autumn (d), year (e) when at least 66% of the models agree on**
 635 **significant changes in the distribution of seasonal and annual means.**
 636

634
 635
 636
 637



638
639
640
641

Figure 7: Temporal evolution of time of emergence for the seasons. Periods where p -value of the Kolmogorov-Smirnov test is lower than 0.05 are highlighted. Only catchments with at least one detection of time of emergence in one of the seasons are shown. Catchments are ordered by decreasing mean elevation of catchment.



642

643 **Figure 8: Multimodel median of seasonal and annual mean changes of runoff due to +1.5°C (left panels), +2°C (middle**
 644 **panels), and +3°C global warming for winter (a-c), spring (d-f), summer (g-i), autumn (j-l), and year (m-o). Black circles**
 645 **indicate catchments whose direction of change agrees across at least 90% the models.**