

Editorial comments:

Comments to the author:

Dear Authors,

Based on the second round of review by one of the original reviewers, I am happy to accept this manuscript after minor revision. Please see the comment from the reviewer regarding the LAI-SPEI and LAI-PRECIPITATION correlation and testing for the significance of the differences in correlation.

Thanks,
Shrad

Response:

Thank you, Editor. We have now tested and reported the significance of the differences in correlation coefficients. These are shown in the newly added Table 5. The text has also been updated.

Please see our response below.

Suggestions for revision or reasons for rejection (will be published if the paper is accepted for final publication)

The authors have done a great job in revising section 3.9. I only have one minor suggestion regarding the statement of "Generally, the pattern correlation coefficient values between the LAI and SPEI are higher than those between the LAI and precipitation in extreme dry and wet years.". The reported coefficient between LAI-SPEI are small and may not be significant. I would recommend the authors also report the significance of the coefficients for both LAI-SPEI and LAI-PRECIPITATION. If the coefficients are not even significant, then the conclusion of LAI-SPEI correlation are in general higher than that of LAI-PRECIPITATION needs to be modified.

Response:

Once again, we thank the reviewer for his time and efforts, whose comments have been used to improve the quality of the manuscript.

As suggested, we have computed the significance of the coefficients for both LAI-SPEI and LAI-PRECIPITATION. We found the coefficients to be significant, except for 1983, an extremely dry year.

The revised figure 10, Table 5 and text are copied below.

3.9 Impacts of extreme events on LAI

The impacts of extreme events on LAI are shown in Fig. 10. The objective was to discuss the impacts and compound influences of extreme events on LAI during extreme hot/dry and wet years. Here, extreme events are the wet (2000, 2010, 2011) years - i.e. the periods with precipitation higher than normal; and the dry (1983, 1984, 1991) years which include the periods of very high dry spells. To achieve this, we used the anomaly of precipitation, SPEI and LAI relative to the long-term mean. The anomaly was computed as a difference between a

particular extreme dry or wet year and 30-year mean representing dry and wet conditions. The anomaly is the magnitude of impacts added by the extreme event in a particular year. The spatial pattern of the changes in LAI, SPEI and precipitation were then plotted. Our analyses follow Pan et al. (2015). Furthermore, we computed the pattern correlation coefficients between the LAI and climate variables for each extreme dry and wet year. The statistical significance of the coefficients was also calculated. The goal of this is to ascertain whether the sign of anomaly of the variables correspond in the same locations on two different maps. Note, although the SPEI is a drought index, it was also considered in a wet year because the impact of drought usually lasts beyond a dry year especially in semi-arid regions of southern Africa. In addition, hot temperature has an influence on the worsening of drought by causing water to evaporate from the soil. SPEI, as a drought index, considers temperature effects on moisture availability.

We considered only observation i.e. CRU (precipitation and drought) and satellite-calculated LAI. The observed climate (CRU) data are not sub-monthly. Only the CRUJRA (reanalysis), which was used for model correlation, is sub-monthly. Model was not analyzed in this section because our goal was simply to examine the observed impacts (or influence) of an extreme event.

The spatial pattern of change of LAI and SPEI are mostly similar during extreme dry and wet years (Fig. 10). For example, in 1983 (a dry year), the negative anomaly of LAI in some parts of the region largely follows the negative anomaly of the SPEI, except in western and central parts. (Figs. 10A & 10B). In 1984, both variables show a strong positive anomaly over Madagascar, Swaziland and Kwazulu Natal Province of South Africa (Figs. 10D & 10E). The pattern of change of both SPEI and LAI are also comparable during extreme wet year. In the wet year of 2000, the positive anomaly of SPEI that is observed in Namibia and South Africa is also evident for the LAI (Figs. 10J & 10K). In a like manner, both variables show negative anomaly over Malawi and Zambia. The strongest pattern (magnitudes) of change of the SPEI in the region is observed in the dry year of 1991 (Fig. 10H). However, the pattern of change of the LAI and SPEI are not similar over some regions in some periods. For instance, in 1991, while a negative anomaly of the SPEI is observed in northern Madagascar and central parts of southern Africa, LAI shows a positive anomaly (Figs. 10G and 10H). The opposite and decreasing relationship between the two variables in 1991 is also evident in the pattern correlation coefficient value of -0.16 (Fig. 10H). The variation in anomaly in these parts and period may be due to the exertion of stronger influence by other factors such as residual soil moisture and precipitation (see Fig. 10I), with temperature having negligible impacts.

The influence of precipitation (as a standalone meteorological factor) on LAI during extreme events is limited. This is observed from the disparity in the spatial pattern of the LAI and precipitation over some regions and periods. For example, in 1984, the wide negative anomaly of precipitation that is shown over Zimbabwe, Mozambique and southern Madagascar is opposite to the LAI, which shows positive anomaly (Figs. 10D and 10F). The LAI anomaly is more similar to that of the SPEI (Fig. 10E). Also, in wet year of 2000, while precipitation shows preponderant increase over northeastern parts of southern Africa, there is a decrease in the LAI as is the case with SPEI (Figs. 10J – 10L). Nevertheless, precipitation plays a primary/major role in the pattern of change of LAI, as is observed over the most parts of the region during the years considered.

Generally, the pattern correlation coefficient values between the LAI and SPEI are higher than those between the LAI and precipitation in extreme dry and wet years (Table 5). Although the coefficients are small, they are mostly significant across the different extreme periods, except for 1983, an extremely dry year, where the p-value is 0.58, for LAI-SPEI (Table 5). The extremely low statistical significance indicate low standard errors and large number of gridsizes for the region.




Figure 10. Spatial pattern changes in satellite-calculated LAI, observed SPEI and precipitation during extreme dry (1983, 1984, 1991) and wet (2000, 2010, 2011) years. For (A) – (I), the changes in LAI, SPEI and precipitation were calculated as a difference between the dry year and the 30-year mean, and for (J) – (R), changes in LAI, SPEI and precipitation were calculated as the difference between the wet year and the 30-year mean. White areas indicate no correlation. The pattern correlation coefficients and the statistical significance of the coefficients between the variables are given in Table 5.

Table 5. Pattern correlation coefficients (*r*-values) and significance of the coefficients calculated at significance level of 5% (*p*-values, 0.05) between the LAI and SPEI, for extreme dry and wet years; as well as between the LAI and precipitation during extreme dry and wet years.

Year	LAI – SPEI		LAI – PRECIPITATION	
	r-values	p-values (0.05)	r-values	p-values (0.05)
1983	0.1	0.58	-0.02	2.43E-06
1984	0.17	7.09E-64	0.04	8.33E-05
1991	-0.16	1.1E-41	0.16	9.83E-60
2000	0.23	1.5E-121	0.14	2.81E-27
2010	0.14	6.3E-17	0.1	0.008
2010	0.14	2.2E-48	0.1	1.47E-06