

# **Investigating the response of LAI to droughts in southern African vegetation using observations and model-simulations.**

Shakirudeen Lawal<sup>1</sup> Stephen Sitch<sup>2</sup> Danica Lombardozzi<sup>3</sup> Julia E.M.S. Nabel<sup>4</sup> Hao-Wei Wey<sup>4</sup> Pierre Friedlingstein<sup>5</sup> Hanqin Tian<sup>6</sup> Bruce Hewitson<sup>1</sup>

5           <sup>1</sup>Climate System Analysis Group, Department of Environmental and Geographical Science, University of Cape Town, Cape Town, 7700, South Africa

<sup>2</sup>College of Life and Environmental Sciences, University of Exeter, Exeter EX4 4QE, UK

10           <sup>3</sup>National Center for Atmospheric Research, Climate and Global Dynamics, Terrestrial Sciences Section, Boulder, CO 80305, USA

<sup>4</sup>Max Planck Institute for Meteorology, Hamburg, Germany

<sup>5</sup>College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter EX4 4QF, UK

15           <sup>6</sup>School of Forestry and Wildlife Sciences, Auburn University, 602 Duncan Drive, Auburn, AL 36849, USA

\*Corresponding author: [lasd\\_dr@yahoo.com](mailto:lasd_dr@yahoo.com)

20

25

## 30 Abstract

In many regions of the world, frequent and continual dry spells are exacerbating drought conditions, which have severe impacts on vegetation biomes. Vegetation in southern Africa is among the most affected by drought. Here, we assessed the spatiotemporal characteristics of meteorological drought in southern Africa using the Standardized Precipitation Evapotranspiration Index (SPEI) over a 30-year period (1982 – 2011). The severity and the effects of droughts on vegetation productiveness were examined at different drought time-scales (1- to 24-month time-scales). In this study, we characterized vegetation using the Leaf Area Index, after evaluating its relationship with the Normalized Difference Vegetation Index. Correlating the LAI with the SPEI, we found that the LAI responds strongly ( $r = 0.6$ ) to drought over the central and south eastern parts of the region, with weaker impacts ( $r < 0.4$ ) over parts of Madagascar, Angola and western parts of South Africa. Furthermore, the latitudinal distribution of LAI responses to drought indicates a similar temporal pattern but different magnitudes across timescales. The results of the study also showed that the seasonal response across different southern African biomes varies in magnitude and occurs mostly at shorter to intermediate timescales. The semi-desert biome strongly correlates ( $r = 0.95$ ) to drought as characterized by the SPEI at 6-month timescale in the MAM (summer) season, while the tropical forest biome shows the weakest response ( $r = 0.35$ ) at 6-month timescale in the DJF (hot and rainy) season. In addition, we found a stronger response (in the year 1983,  $r = 0.84$  over Namibia and eastern parts of South African) of the LAI to drought during dry years as compared to wet years; and we found different temporal variability in global and regional responses across different biomes.

We also examined how well an ensemble of state-of-the-art dynamic global vegetation models (DGVMs) simulate the LAI and its response to drought. The spatial and seasonal response of the LAI to drought is mostly overestimated in the DGVM multi-model ensemble compared to the response calculated for the observation-based data. The correlation coefficient values for the multi-model ensemble are as high as 0.76 (annual) over South Africa, and 0.98 in MAM season over the temperate grassland biome. Furthermore, the DGVM model ensemble shows positive biases (3-month or longer) in the simulation of spatial distribution of drought timescales and overestimates the seasonal distribution timescales. The results of this study highlight the areas to target for further development of DGVMs and can be used to improve the models' capability in simulating the drought-vegetation relationship.

**Keywords:** Drought intensity; Drought indices; Standardized precipitation evapotranspiration index; DGVMs; southern Africa; Drylands

# 1 Introduction

70 Drought can be described as a natural occurrence whereby the natural accessibility of water for a region is beneath the normal state over a long period of time (Xu *et al.*, 2015). Globally, it is considered one of the world's most important climate risks, with significant environmental, social and ecological impacts on different sectors (e.g. agriculture, forestry, hydrology) and human lives (Naumann *et al.*, 2018). Increasing trends in the occurrence and severity of drought in West Africa and Mediterranean have huge impacts on water resources and agriculture (Sultan and Gaetani, 75 2016). In southern Africa, a region regarded as a climate hotspot because of the projected impacts of climate change on its numerous endemic vegetation, an understanding of these impacts is important for mitigation options in managing future drought events. Therefore, it is important to examine drought impacts on vegetation and evaluate how this is simulated in models.

80 Drought is a frequent occurrence in southern Africa, and has enormous impacts on vegetation in the region. For instance, drought has resulted in a significant loss of biomes and death of plants (Masih *et al.*, 2014, Hoffman *et al.*, 2009). It is reported that there has been a significant loss of vegetation cover over the region over the last 30 years (Driver *et al.*, 2012; DEA, 2015). Drought has also impacted speciation of vegetation thereby causing significant changes to the region's rich biomes, through the lack of formation of new species or even growth of species with 85 underdeveloped morphological and physiological characteristics (Hoffman *et al.*, 2009). Drought-induced vegetation loss has both ecological and socio-economic consequences on human lives. For instance, studies have shown that food security in the region is threatened due to the continual mortality of vegetation (FAO, 2000b; Müller *et al.*, 2011). Other studies (e.g. Wang, 2010; Khosravi, 2017) have also reported that southern Africa could lose more than \$200 billion of its 90 GDP from the effects of drought on vegetation. The enormous impacts on vegetation have thus made it imperative to investigate how vegetation might respond to different drought intensities at varying timescales.

In order to monitor and quantify drought characteristics, drought indices are used (Wilhite and Glantz, 1985). Drought indices including the SPI (i.e. the standardized precipitation index), 95 standardized water-level index and standardized anomaly index are derived from a single hydrological variable, which is rainfall (Kwon *et al.*, 2019). Other indices such as the Palmer drought severity index, multivariate standardized drought index and standardized precipitation evapotranspiration index (SPEI) combine two or more variables related to other atmospheric or soil and environmental conditions that may predispose a plant to water stress (Palmer, 1965; 100 Vicente-Serrano *et al.*, 2010; Hao and AghaKouchak, 2013). Among the drought indices, the SPI is the most widely used because of the adjustable timescale and its relatively simple calculation (McKee *et al.*, 1993). It is also recognized as appropriate for use in southern Africa (Hoffman *et al.*, 2009). However, the SPI has a significant shortcoming, which is that its computation uses only rainfall without considering the effect of other meteorological variables in the development of 105 drought occurrence (Teuling *et al.*, 2013). In order to address this shortcoming, SPEI was developed for drought monitoring and it is regarded as a more suitable drought index in the region to investigate the spatiotemporal scale of drought (Ujeneza and Abiodun, 2014). SPEI is computed from the difference between potential evapotranspiration (PET) and rainfall (Vicente-Serrano *et al.*, 2010). PET can be computed using different methods such as Hargreaves (HG) and Penman-Monteith (PM) method. Although studies (e.g. Vicente-Serrano *et al.*, 2010) have found that PM 110

method captures drought better than HG, other studies (e.g. Lawal *et al.*, 2019a) showed this difference is negligible over southern Africa.

115 Many studies have used different indices to quantify observed drought, characterize vegetation, and study drought effects on the productiveness of vegetation across different timescales. Several studies (e.g. Vicente-Serrano *et al.*, 2015; Zhang *et al.*, 2012; Lawal *et al.*, 2019a,b) have shown that the satellite-derived normalized difference vegetation index (NDVI) is one the most important indicators of vegetation health and greenness. These studies applied NDVI in examining drought impacts on global vegetation biomes. However, other studies (Gitelson, 2004; Santin-Janin *et al.*, 2009) have argued that while the NDVI is a true proxy for vegetation trends, its potential saturation makes it difficult to fully estimate biomass. In addition, because the NDVI parameters are not well calibrated and often missing in models, simulated NDVI can be biased. Due to its high correlation with NDVI, the leaf area index (LAI) is instead used to characterize vegetation conditions (Fan *et al.*, 2008; Zhao *et al.*, 2013). Although the LAI is an important vegetation proxy, it is rarely considered in the estimation of drought impacts on vegetation. Thus, quantifying the response of the LAI to drought over southern Africa is important for understanding the processes that modulate ecosystem services produced by vegetation which are crucial for human survival. (Melillo 2015).

130 Previous studies have also evaluated the performance of coupled climate models in simulating the response of vegetation to drought. For instance, Lawal *et al.* (2019a) reported that an ensemble of the Community Earth System Model (CESM) showed biases in response simulation of vegetation to drought. This was attributed to the parameterizations of the land component (i.e. Community Land Model, CLM) which poorly simulated observed NDVI. Given the poor replication of vegetation response to drought by a coupled climate model, there is a need to examine land-only models and whether they might better capture drought-vegetation relationship when the atmospheric forcings are derived from observations. The present study used Dynamic Global Vegetation models (DGVMs) to study vegetation response to drought, as little is known on how the LAI response to drought is simulated by DGVMs. The choice of DGVMs is because of their capability in simulating mostly accurate carbon exchange between the atmosphere and vegetation ecosystems (Lu *et al.*, 2011).

140 The aim of this study is to investigate the response of LAI to droughts in southern African vegetation using observations. We also examined how well the responses are represented in model simulations. We used satellite-derived and simulated LAI to quantify vegetation responses to drought. We characterized the spatiotemporal extent of drought and its severity using the SPEI and then assessed the influence of drought using the LAI from satellite data and model simulations.

## 145 **2 Data and Methodology**

### **2.1 Data**

In this study, we used satellite-calculated (hereafter, observed LAI/observation-based LAI) and simulated LAI, and satellite-derived NDVI; gridded observation and reanalysis climate datasets. The gridded observation climate datasets include precipitation, maximum, mean and minimum temperature. These data were gotten from CRU (i.e. the Climate Research Unit; Mitchell & Jones, 2005; Harris *et al.*, 2014). These are global monthly data which have 0.5° x 0.5° as spatial

155 resolution and spans 1901 – 2019 period. Here, we used the CRU data for the period 1982 – 2011  
to compute observed drought indices (i.e. SPEI) to characterize the spatiotemporal severity of  
drought. CRU is a gridded observed dataset, which was used because of its suitable spatial and  
temporal resolutions. Previous studies (e.g. New 1999; New 2000; Wolski *et al.*, 2018; Otto *et al.*,  
2018; Harris *et al.*, 2020) have shown that there is a good and robust agreement between  
observation network and CRU over most parts of southern Africa. We should note that sparseness  
and missing data generally affect the correlation between CRU and station data in the region.  
160 Furthermore, with respect to inter-annual variability, CRU robustly captures the climate factors in  
southern Africa. The major exception is with the long-term trend of precipitation particularly over  
Western Cape province of South Africa as well as wetter than normal condition over the same  
province. These limitations do not affect the validity of our results because we are looking at  
below-normal precipitation, and temperature.

165 The reanalysis climate data we used are the CRUJRA, which is a combination of CRU and the  
Japanese Reanalysis data (JRA) (University of East Anglia Climatic Research Unit; Harris, I.C.,  
2019). It is a 6-hourly, land surface, gridded data with a spatial resolution  $0.5^\circ \times 0.5^\circ$ . CRUJRA  
was used to compute reanalysis drought indices and used for model simulations. Here we  
aggregated CRUJRA to monthly samples and used the data at the same spatial and temporal  
170 resolution as CRU.

For the satellite vegetation indices, first, we used the third generation of NDVI (hereafter,  
NDVI3g) from the Global Inventory Modelling and Mapping Studies (GIMMS), and spans period  
175 from 1981 – 2015, with a temporal resolution of biweekly and a spatial resolution of about 8km  
(Pinzon and Tucker, 2014; National Center for Atmospheric Research Climate Data Guide,  
accessed 2019). Here, we used the data for the period 1982 – 2011. Furthermore, we used the third  
generation of the GIMMS LAI (LAI3g) which also spans the period 1981 – 2015 and has a  
temporal resolution of biweekly and spatial resolution as GIMMS3g. The LAI data had been  
180 processed (at source) using a set of neural networks which were first trained on highest-quality  
and post-processed MODIS LAI and FPAR products and AVHRR GIMMS NDVI3g data for the  
overlapping period (2000 to 2009). The trained neural networks were then used to produce the  
LAI3g and FPAR3g data sets (Mao and Yan, 2019). For the study, LAI3g was also used for the  
period 1982 – 2011. We note that GIMMS LAI and the NDVI, used in this study, are two different  
185 indices. The LAI was post-processed using different data (MODIS LAI, fPAR, AVHRR NDVI)  
for the period of 2000-2009. GIMMS LAI product is superior here over the GIMMS NDVI, which  
is due to the information derived from the MODIS LAI. The additional properties on GIMMS LAI  
by MODIS differentiate the index from the NDVI. Thus, it was necessary to investigate how the  
two indices differ. In addition, other studies (Forkel *et al.*, 2013., Schaefer *et al.*, 2012., Rezaei *et*  
*al.*, 2016., Lawal *et al.*, 2019a) have investigated how well satellite derived LAI estimate actual  
190 and ground-measured LAI.

The simulated monthly LAI data were obtained from eleven Dynamic Global Vegetation Models  
(DGVMs) which are part of the Trendy-version 7 (Sitch *et al.*, 2008; Le Quéré *et al.*, 2014). These  
DGVMs are CABLE-POP (Haverd *et al.*, 2018), CLM (Oleson *et al.*, 2013), CLASS-CTEM  
195 (Melton and Arora, 2016), DLEM (Tian *et al.*; 2015), JSBACH (Mauritsen *et al.*, 2018), LPX  
(Lienert and Joos; 2018), OCN (Zaehle *et al.*; 2011), ORCHIDEE (Goll *et al.*, 2017), SURFEX  
(Joetzer *et al.*, 2015), JULES (Clark *et al.*; 2011) and VISIT (Kato *et al.*, 2013). LAI from the

models have a monthly temporal resolution spanning period from 1901 – 2017. We selected these DGVMs because they have been run with similar protocol (S3 simulations) and forcing datasets (i.e. CRUJRA).

## 2.2 Methods overview

### 2.2.1 Evaluation of DGVMs and the relationship between NDVI and LAI

The relationship between the NDVI and LAI was evaluated by computing the grid cell spatiotemporal correlation between GIMMS NDVI and GIMMS LAI. The spatiotemporal correlation between GIMMS LAI and simulated LAI from individual DGVMs was also calculated. This was necessary to show whether LAI is an appropriate estimator of NDVI, and how well the models simulate the LAI in the region. Furthermore, we made comparisons of seasonal and interannual variations (IAV) of observed and modelled LAI. We note that the lack of available data makes it difficult to compare GIMMS LAI and actual LAI. Nevertheless, the GIMMS LAI has been evaluated and agrees well with observations in other regions (Fan *et al.*, 2019).

The climatology of observed and simulated climatic variables as well as LAI over six major biomes in southern Africa for the period 1982- 2011 were computed. These biomes are semi desert, Mediterranean, dry savanna, moist savanna, temperate grassland and tropical forest (Fig. 1; Sinclair & Beyers, 2015; Lawal *et al.*, 2019a, b).

### 2.2.2 Description of Drought

For the present study, we adopted the definition of meteorological drought, “which is described as a period (e.g. a season) during which there is a deficit in the magnitude of precipitation in a particular area compared to the long-term normal (Palmer, 1965; Wilhite & Glantz, 1985)”. The deficit in magnitude of precipitation compared to long-term normal is mostly accounted for by temperature, and less by humidity, wind or other variables. Here, we used meteorological drought because it does not make any presumptions about soil characteristics or run-off. In addition, it is acknowledged to be a primary component in the depletion of vegetation productiveness and reduction of biomass (Vicente-Serrano, *et al.*, 2010). Previous studies (Vicente-Serrano *et al.*, 2006; Vicente-Serrano *et al.*, 2013) have also used meteorological drought in the investigation of drought impacts on biomass and vegetation productiveness.

### 2.2.3 Drought computation and correlation with LAI

The analyses include calculating drought (i.e. SPEI) using CRU data over a 30-year (1982 – 2011) period for different drought timescales. The drought time-scale can be described as the aggregation of temporal duration (Vicente-Serrano *et al.*, 2010). SPEI is an index that is used to quantify drought (see Table 1). Therefore, the quantified values of the index give the state of drought in a space. Our definition and approaches follow numerous previous studies (Vicente-Serrano *et al.*, 2013; Khosravi *et al.*, 2017; Zhao *et al.*, 2013; Hao *et al.*, 2013).

A time series of the evolution of drought for the 30-year period was plotted. The present study extends the timeframe for understanding drought impacts from 1982 to 2011 mainly because there were frequent droughts in the 2005 – 2011 window (Masih *et al.*, 2014). The timeframe was then extended back to cover a 30 year period to be long enough to cover impacts of climate change, which is particularly important considering that southern Africa experiences more frequent droughts with impacts exacerbated by climate change. This information is important for considering adaptation measures and understanding the role of climate change.

The drought index, SPEI, is calculated from the deduction between precipitation (P) and potential evapotranspiration (PET) as shown below:

$$D = P - PET \dots \dots \dots (1)$$

where D-values represent a measurement of water deficit or surplus aggregated at different time scales. D values are obtained through aggregation over individual time-scales which span 1- to 24-months (i.e. 1-, 3-, 6-, 9-, 12-, 15-, 18-, 21- and 24-month). The time-scales were calculated by including the past values of the variable. For example, a time-scale of 15-month suggests that input from the preceding 15 months, which includes the present month, was used for calculating SPEI (Begueria *et al.*, 2014). “For the 1-month timescale, only the current month data is used for the calculation. The D values were standardized by assuming a suitable statistical distribution (e.g. gamma, log-logistic). The log-logistic distribution was used to standardize the D values in this study” (see also Lawal *et al.*, 2019a). For more details on the timescale computation, please see Vicente-Serrano *et al.* (2010); <https://rdr.io/cran/SPEI/man/spei.html>. PET is computed from maximum temperature, minimum temperature and mean temperature, using Hargreaves technique (e.g. Vicente-Serrano *et al.*, 2012; Begueria *et al.*, 2014; Stagge *et al.*, 2014).

We note that the PET in the SPEI was computed using Hargreaves (HG) method rather than Penman-Monteith (PM) because the data (e.g. vapour pressure, maximum and minimum humidity) required for computing PM over southern Africa are sometimes missing or not available at the needed gridded spatial resolutions and timespan. Although PM is considered better in most regions, Lawal *et al.* (2019a) showed that the variation between the PM and HG is negligible for southern African region. The study only considered observed SPEI\_PM which was obtained from <https://spei.csic.es/database.html> and not modelled SPEI\_PM due to the unavailability of simulated data required for its computation. Other studies (e.g. Begueria *et al.*, 2014) have also found that there is an insignificant contrast in the strength of PM and HG for reproducing their divergence on measured variables such as vegetation indices. We should also note that the SPEI (unlike SPI or PDSI) is the most appropriate index for measuring drought in southern Africa, as it accounts for the effect of evaporative demand from the atmosphere in drought monitoring (Vicente-Serrano *et al.*, 2010; Ujeneza *et al.*, 2014). In addition, the SPEI is reported to be able to identify the geographical and temporal coverage of droughts (Vicente-Serrano *et al.*, 2010; Ujeneza *et al.*, 2014).

285

**Table 1.** Definition of drought thresholds based on the SPEI scale

<b>SPEI</b>	<b>Drought thresholds</b>
2 or more	Extreme wet
1.5 to 1.99	Severe wet
1 to 1.49	Moderate wet
0 to 0.99	Mild wet
0 to -0.99	Mild drought
-1 to -1.49	Moderate drought
-1.5 to -1.99	Severe drought
-2 or less	Extreme drought

Source: Wang *et al.* (2014), modified as shown in Lawal, 2018.

290 We deseasonalized GIMMS LAI by transforming monthly LAI series per pixel to symbolize the standardized deviations from extended mean. This was to make the sequence of LAI commensurate to SPEI (Vicente-Serrano *et al.*, 2013) and eliminate the impact of periodicity on vegetation response. We note that the SPEI is intrinsically deseasonalized.

295 In order to reconcile the difference in spatial resolutions of CRU and GIMMS, we regridded the data to the same spatial resolution using the bilinear interpolation method. We then computed the correlation per grid-cell between SPEI (based on CRU) and deseasonalized LAI over the 30-year period at the different drought timescales using Pearson correlations. We then compared the spatial distribution of maximum (peak) correlation and the comparable time-scales of drought for  
300 observed SPEI. Our analyses take into account the periods which LAI responds to the presence/absence as well as the severity of drought. This is referred to as “drought time scale” in the study.

Next, we investigated correlations at each grid cell between the drought (from reanalysis -  
305 CRUJRA) and an ensemble median of deseasonalized modelled LAI from individual DGVMs. The peak (maximum) correlations and equivalent time-scales from the complete 1- to 24-month time-scales were mapped for the ensemble median over the 30-year period. We used ensemble median because of its less sensitivity to independent outliers (Reuter *et al.*, 2013). In summary, we calculated model ensemble drought from the median from individual members’ drought indices.  
310 The inter-annual variation of drought impacts on LAI by individual DGVM was also calculated for different timescales. We also examined the observed and an ensemble mean of simulated LAI response to drought across latitudes.

In summary, CRU was used to compute observed drought while CRUJRA was used to calculate  
315 modeled drought because it is what was used to force the models. This will allow for easy observation and model comparisons.

Similar to Lawal *et al.*, 2019a, we calculated the seasonal mean for four seasons i.e. a) December-  
320 January-February (DJF); b) March-April-May (MAM); c) June-July-August (JJA); and d) September-October-November (SON) from the correlations of monthly series of drought and LAI.



These were computed from correlating monthly series (twelve series per year) per pixel of GIMMS-LAI and each monthly series of 1- to 24-months drought (SPEI) series over the 30-year period with Pearson correlation. The same technique was used for the model ensemble. In simpler terms, we calculated the correlations (twelve sequences in a year) of monthly LAI to monthly sequences of 1- to 24-months SPEI using 30-years of data. Subsequently, the seasonal mean of these correlations was calculated. The peak correlations and drought timescales of the models were calculated over six major biomes in southern Africa, namely (Fig. 1) – Temperate grassland, Tropical forest, Moist Savanna, Dry Savanna, Semi-desert and Mediterranean vegetation. These regions were selected because of their relative importance and are most affected by drought.

Finally, the impacts of extreme events (wet and dry years) at different time periods were compared; and the comparison of global and regional responses to drought across biomes for the period 1982 – 2011 was investigated.




Figure 1. Major vegetation biomes in southern Africa (adapted from UNEP, 2008, Sinclair & Beyers, 2015 and Lawal *et al.*, 2019a,b). The black lines indicate political boundaries.

### 3 Results


#### 3.1 Grid cell correlations between NDVI and LAI, seasonal and interannual variation of observed and modelled LAI

Figure 2 illustrates the relationships between the NDVI and LAI for observations, as well as the comparison between observed and model LAI. There is a strong linear relationship between

observed NDVI and LAI (Fig. 2a). The correlation (0.94) is high between both variables and the standard deviation is low (0.005). The standard deviation being referred to is for GIMMS LAI and individual DGVMs, as well as GIMMS LAI and GIMMS NDVI. From the figure, there is a log-like shape, where NDVI grows faster when LAI is low ( $<0.2$ ) and becomes saturated when LAI goes higher ( $>0.3$ ). A linear regression of the data shows a slope of 2.15. The low standard deviation indicates that the values from the two indices are close. Although there is a good agreement between observed NDVI and LAI, the 1:1 line shows that the datasets are not exactly equal.

Furthermore, there is good agreement between observed and the simulated LAI (Fig. 2). JULES has the highest correlation (0.97) with observation (Fig. 2l). CLM has the weakest (0.73) correlation with the observations (Fig. 2c). DLEM and LPX have the same correlation coefficient value of 0.87 with observation (Fig. 2e, 2g). The positive relationships between simulated and observed LAI indicate a general applicability in investigating the model's performance of vegetation response to drought. It also shows that the correlation is strong enough to compare how the LAI reacts to drought in the ensemble. An aggregation of observation along the gradient of simulated LAI shows that most of the models have similar slopes with observation.


The comparison of seasonal and interannual variation of observed and modelled LAI is given in Fig. 3. The model shows a stronger positive bias in JJA and SON in comparison to summer and winter months; and a negative bias over the tropical forest region of Madagascar (Fig. 3A – 3D). In addition, models mostly overestimate the seasonal patterns of LAI in regions during DJF and JJA, and underestimate LAI in MAM and SON (Figs 3E – 3L). Generally, there is a good agreement between observed and modelled LAI in DJF, although it is weaker in SON (Figs. 3M – 3P). Furthermore, models largely simulate the interannual variability of LAI, with the observed LAI magnitudes falling within the modelled spread (Fig. 3Q).



375

Figure 2. Scatterplots of correlations between vegetation indices (observation and model) for the period 1982 – 2011 over southern Africa. Inset values indicate the correlation coefficient (r) and standard deviation (Stdev) between GIMMS LAI and GIMMS NDVI, as well as GIMMS LAI and modelled LAI. The colour represents each grid cell. The pink solid line is the linear regression, while the dashed black line shows 1:1 line. The unequal x-axes are to visualize the detailed data for the models.

380



390

Figure 3. Spatial seasonal distribution and inter-annual variability (IAV) of satellite-calculated and modelled LAI (multi-model mean) over southern Africa. (A) – (D) show the difference (bias); (E) – (H) and (I) – (L) show their standard deviation (Stdev); (M) – (P) show their spatiotemporal correlations; and (Q) shows their inter-annual variability for the period 1982 – 2011.

### 3.2 Climatology of observed and, simulated climate variables and LAI

395

This section compares the seasonal cycle of observational (CRU and CRUJRA) climate variables; as well as observed and simulated LAI from GIMMS LAI and TRENDY models, respectively.

Precipitation and temperature are seasonally variable and their climatologies are mostly similar. For example, precipitation is higher in MAM and DJF over many of the biomes except in Mediterranean vegetation where precipitation is higher in JJA (Fig. 4a, 4d, 4g, 4m, 4p). The wettest month occurs over TF (i.e. tropical forest biome) where the precipitation is about 350mm. Conversely, during the dry season (JJA), there is little rainfall in the biome, although, it experiences some precipitation June and July. Over the Mediterranean vegetation (Fig. 4j), a winter (JJA) rainfall region, rainfall variability is lower and is mostly dry in the DJF and SON. Similarly, the highest minimum and maximum temperature in the region is observed in the DJF season, where the highest temperature value exceeds 30°C. Over the tropical forest biome, although the distribution pattern of precipitation and temperature are similar for most months, they differ during June and July months. The pattern of precipitation and temperature distribution generally differ over the Mediterranean vegetation. The pattern of temperature and precipitation from CRUJRA follows CRU, although the ensemble spread is much narrower (Fig. 4c, 4f, 4i, 4l, 4o, 4r). The spatial patterns of the climate variables from CRU and CRUJRA are shown in Fig. 5. CRUJRA well simulates the pattern of precipitation over southern Africa (Figs. 5a and 5b) as CRU although it shows some biases in magnitudes, as well as for minimum temperature (Figs. 5g and 5h).

There are not strong seasonality for LAI, with maximum observed LAI values less than 4 in all biomes. The models well reproduce the climatology of LAI over the southern African biomes with a few exceptions (Fig. 4b, 4e, 4h, 4k, 4n, 4q). For instance, the models simulate the drop in LAI over the semi desert, temperate grassland, tropical forest, dry savanna and moist savanna biomes in JJA. The highest increase in observed LAI occurs over the tropical forest in April, although the models simulate a decrease in LAI over tropical forest during this time. On the other hand, lowest amount (less than 0.1) of LAI is observed in September and this occurs over the semi-desert biome. Observations typically fall within the range of the model ensemble. In addition, the distribution pattern of simulated LAI is similar to observation in most biomes except in the Mediterranean and tropical forest biomes. The LAI pattern also follows that of the climatic variables although the former lag behind. The lag effect is accounted for in this study, and is known as drought time scale.

430




Figure 4. Annual cycle of observed climate variables (precipitation, mm/month; maximum, minimum and mean temperature, °C) and LAI for observation and multi-model mean (TRENDY) across six southern African biomes over for the period 1982 – 2011. The annual cycle of the LAI for individual models are shown in Figure S5.

435




Figure 5. Spatial distribution of precipitation, mean temperature, maximum temperature and minimum temperature over southern Africa in CRU and CRUJRA; for the periods 1982 – 2011.

### 3.3 The evolution of drought in southern Africa


445 Figure 6 shows the evolution of observed SPEI in southern Africa between 1982 and 2011. Here, drought indices from CRUJRA are not included because preliminary investigation showed close magnitudes for drought indices computed from CRU and CRUJRA. We note that CRU was used to calculate SPEI for observation while simulated SPEI was computed with CRUJRA.

450 There is inter-annual, seasonal and decadal variability of the drought indices during dry and wet conditions over southern Africa. Although 1-, 3- and 6-month SPEI indicate no trend in wet or dry spells, they show the intensity of drought event for the 30-year period (Fig. 6a, 6b, 6c). The highest magnitude of drought is captured by 1-month SPEI while the lowest is shown in 21-month SPEI. The severity of drought intensity is similar for all SPEI (i.e. 1-month to 24-month SPEI). The magnitude of the severity is on the y-axis of 1-month to 24-month SPEI.

We note an increasing trend in SPEI (9- to 24-month). Tables 2 and 3 illustrate when droughts occurred, as well as the severity of drought within the 30-year period. They also show, however, that droughts were most frequent and intense in the second decade, thus, indicating how climate change is expected to increase the frequency and severity of droughts.

460





465

Figure 6. Evolution of SPEI in southern Africa for the period 1982 – 2011. The trend is significant at 90% confidence interval for all timescales.

470

**Table 2. Characteristics of drought occurrence for 1- to 24-month drought timescale for 1<sup>st</sup> decade (1982-1991), 2<sup>nd</sup> decade (1992-2001) and 3<sup>rd</sup> decade (2002- 2011).**

Drought Timescale	Number of drought events			Year of moderate drought events		
	1 <sup>st</sup> Decade	2 <sup>nd</sup> Decade	3 <sup>rd</sup> Decade	1 <sup>st</sup> Decade	2 <sup>nd</sup> Decade	3 <sup>rd</sup> Decade
1-month	53	62	53	1987	1992, 1990	2004, 2007, 2008, 2011
3-month	55	64	58	1988	1991, 1992	2004, 2008, 2011
6-month	44	62	60	1982	1992, 1993	2004
9-month	55	66	62	-	1992, 1994	-
12-month	53	68	56	-	1992, 1995	-
15-month	58	63	56	-	1992	-
18-month	54	69	51	-	1992, 1994	-
21-month	51	69	56	-	1992, 1995	-
24-month	41	71	54	-	1992, 1994	-

**Table 3. Statistics of the severity of drought for 1- to 24-month drought timescale for 1<sup>st</sup> decade (1982-1991), 2<sup>nd</sup> decade (1992-2001) and 3<sup>rd</sup> decade (2002- 2011). SD is the standard deviation and Max is the highest magnitude of drought occurrence.**

Drought Timescale	Mean			SD			Max		
	1 <sup>st</sup> Decade	2 <sup>nd</sup> Decade	3 <sup>rd</sup> Decade	1 <sup>st</sup> Decade	2 <sup>nd</sup> Decade	3 <sup>rd</sup> Decade	1 <sup>st</sup> Decade	2 <sup>nd</sup> Decade	3 <sup>rd</sup> Decade
1-month	0.30	0.39	0.31	0.26	0.26	0.27	1.1	1.02	1.15
3-month	0.32	0.36	0.29	0.24	0.25	0.25	1.02	1.01	1.12
6-month	0.38	0.38	0.26	0.31	0.28	0.24	1.04	1.3	1.33
9-month	0.31	0.41	0.21	0.26	0.35	0.18	0.86	1.32	0.83
12-month	0.30	0.43	0.21	0.21	0.35	0.18	0.81	1.27	0.7
15-month	0.28	0.47	0.20	0.21	0.31	0.18	0.77	1.13	0.7
18-month	0.27	0.48	0.20	0.17	0.28	0.12	1.02	1.02	0.77
21-month	0.18	0.50	0.17	0.2	0.28	0.14	0.62	1.00	0.61
24-month	0.28	0.51	0.15	0.19	0.24	0.11	0.55	0.93	0.46

475 **3.4 Spatial distribution of LAI response to drought and the timescales**

Figure 7 presents the spatial distribution of the peak correlation between the SPEI and the LAI, and the timescales at which the correlation occurs. This is to show the magnitude of response of LAI to drought in southern Africa, and the length of the period for the response.

480 Observations show that southern African LAI can respond fairly strongly to droughts (peak correlation magnitudes of between 0.4 and 0.6), though the response is much weaker ( $r < 0.4$ ) in eastern Madagascar, Angola and parts of South Africa (Fig. 7a). The TRENDY multi model median generally overestimate the observed magnitude of the LAI to drought response (Fig. 7c).  
 485 Peak correlations for the models seem to be much stronger – in the 0.6-0.8 range for most of the regions. In addition, over the arid areas of Namibia, models simulate a LAI while observations depict no measurable LAI, indicating that models simulate the LAI in areas where observations show no measurable LAI.

The multi model median have a drought timescale that is mostly longer than the observations (Fig. 7b, d). For instance, drought response of simulated LAI occurs mostly over a longer time period  
 490 (6-, 9-month timescale) than in the observation over eastern Madagascar. Over southern areas of Madagascar and central Zambia, the multi model median overestimates the drought timescale. Over central areas of South Africa and Mozambique, simulated LAI responds at intermediate (9-month) timescales. In comparison, similar drought timescales for the observation and the model ensemble median are shown in parts of Angola.

495




Figure 7. Spatial distribution of peak correlation between drought (SPEI) and LAI over the region of southern Africa in the observation and in the model ensemble median; for the period 1982 – 2011. Panels (a) and (c) show the peak correlation per pixel, which is independent of the timescale and the month of the year. Panels (b) and (d) indicate the timescales at which the peak correlation between SPEI and LAI is found. Areas with no significant correlation are white.

### 3.5 Latitudinal distributions of LAI response to drought and the timescales

The present study investigated and discussed the implications of drought on different vegetation/biome types across latitudes in the region. Here, we stratified the LAI response to drought based on latitude, because we intend to investigate and identify the shift in response based on the vegetation types across the latitudinal belt.

While the pattern of the latitudinal distribution of the LAI response to drought is identical across different timescales, the magnitudes generally differ (Fig. 8). The response is much weaker (less than 0.15) for the 1-month timescale than for other timescales. The strongest response is observed at 6-month timescale between latitudes 25 and 30°. The model ensemble mean generally agrees with the pattern of the observed LAI-SPEI correlations across all the timescales. However, the

515 magnitudes differ from the observation. Modeled correlation is stronger than observations for the longer (6-month or more) timescales. This means that the models are oversimplifying how LAI responds to drought by neglecting other climate factors, such that in models, LAI only correlates to water deficit (SPEI). Furthermore, there is an offset between observation and model mean which is consistent across most of the timescales, perhaps due to strong memory of the some of the models.

520




Figure 8. Mean correlation (observed and multi-model ensemble) of annual LAI and SPEI for 1982 – 2011 across latitudes over southern Africa for 1- to 24-month timescales.

525 **3.6 Response of LAI to droughts across seasons**

Observations show similar correlations between LAI and drought across all seasons in the biomes (Fig. 9). For the dry savanna, which is one of the most climate-impacted biomes in the region, LAI

530 response to drought is strong and the correlation is as high as 0.8 in MAM season and it occurs at  
12-month timescale. The correlations between drought and LAI are also very strong in other  
seasons over the same biome and occur at 6- and 12-month drought timescale, except over the  
Mediterranean vegetation where the response occurs at 18-month in DJF season. Similarly, the  
peak correlations between drought and LAI are strong across the other biomes. With an exception  
of the tropical forest biome, the drought timescale is at longer time periods (> 6-months).

535 The model ensemble generally overestimates correlations across the biomes in different seasons.  
Whilst the correlation magnitude remains mostly larger than observation, nonetheless, models  
simulate closer correlation with observation in some biomes and seasons. For instance, over  
Mediterranean vegetation, models simulate fairly good response of LAI to drought in all seasons.  
540 Furthermore, in nearly all other biomes, the ensemble spread overlaps with the observations. In  
addition, simulations mostly overestimate drought timescale, except over dry savanna. A possible  
reason for the difference why the time scale for Dry savanna was underestimated may be because  
phenological triggers for dry savannah vegetation types respond differently to environmental  
variables, which the models do not capture. The African Dry savanna region is characterized by  
545 rapid vegetation changes due to fire, land-use among others, as well as senescence for prolonged  
dry periods (Rahimzadeh-Bajgiran *et al.*, 2012; Zhu and Liu, 2015), which may have contributed  
to the underestimation in the response of the models. Similarly, models have different  
representations of fire, which could also indirectly contribute to the underestimated model  
responses to drought.


550

555

560

565

570



575

Figure 9. Seasonal correlations of drought (SPEI) and LAI across six southern African biomes. The values on the left axis show the peak correlation in observation and TRENDY models. The values on the right axis indicate the corresponding drought timescale.

### 3.7 Inter-annual variation of model simulation of drought impacts on LAI

580

Table 4 shows the correlations between observed mean SPEI and LAI for the period 1982 and 2011 as well simulations by individual DGVM across different timescales. Unlike Fig. 7 which shows the peak correlations, the table shows the mean correlation for the 30-year period.

585

There is variation in the inter-annual simulation of LAI response to drought across different timescales by individual models. For instance, on the 1-month timescale, JULES simulates the least correlation value while JSBACH shows the highest correlation value. Furthermore, JSBACH simulates the highest correlation value for most of the timescales while CLM simulates the least correlation values for most (3-, 9-, 12-, 15- and 18-month) of the timescales. A possible reason for the weak performance of CLM may be its representation of the canopy construction of the PFTs and of its foliage clumping representation. In addition, CLM is limited in its simulations of vegetation with regards to transpiration, due to rooting depth among others (Dahlin *et al.*, 2020). Furthermore, CLM does not well simulate savanna ecosystems, but instead uses a combination of grasses, shrubs, and trees. There are also some problems (such as an unusual green-up in dry season) identified with stress deciduous responses (Dahlin *et al.*, 2015).

590


595 Table 4. Model simulation of mean SPEI and LAI correlations between 1982 and 2011. \* indicates the model with the lowest mean correlation.

Correl.	GIMMS LAI	CABLE- POP	CLM	CLASS- CTEM	DLEM	JSBACH	LPX	OCN	ORCHIDEE	SURFEX	JULES	VISIT
1-month	0.0066	0.1017	0.1188	0.1266	0.085	0.1388	0.0842	0.0579	0.0884	0.1094	0.027*	0.086
3-month	0.0775	0.3084	0.0187*	0.29	0.2499	0.4165	0.2403	0.1644	0.266	0.2791	0.1434	0.1108
6-month	0.091	0.3474	0.1806	0.3562	0.3305	0.5404	0.2386	0.3377	0.3733	0.3759	0.2429	0.1703*
9-month	0.0832	0.333	0.2034*	0.3496	0.3437	0.5734	0.2505	0.4055	0.39	0.3994	0.3189	0.23
12-month	0.0813	0.3053	0.2155*	0.3231	0.3229	0.5398	0.304	0.4109	0.4054	0.3911	0.3663	0.2892
15-month	0.0642	0.2773	0.2161*	0.2912	0.2846	0.4925	0.3208	0.4106	0.4132	0.3547	0.3623	0.3198
18-month	0.0452	0.2534	0.2349*	0.276	0.2381	0.4599	0.2405	0.3998	0.4109	0.3289	0.3586	0.3472
21-month	0.0406	0.239	0.2402	0.27	0.2119	0.4334	0.1569*	0.3739	0.3962	0.2955	0.3621	0.3501
24-month	0.0409	0.2302	0.2355	0.2668	0.2186	0.4696	0.2064*	0.3528	0.3777	0.2682	0.3617	0.327

### 3.8 Impacts of extreme events on LAI

600 The impact of extreme events on LAI is shown in Fig. 10. The objective was to discuss the impacts of extreme drought events, and compound influences of drought on LAI during extreme hot and dry years. Here, extreme events are the wet years - i.e. the periods with precipitation higher than normal; and the dry years which include the periods of very high dry spells.

605 The magnitudes of response of LAI are generally stronger during the dry years than in wet years (Fig. 10). The distribution pattern of the magnitude of response however, varies across the region. With the exception of the year 2010, the strongest drought response is observed in parts of Namibia, Botswana, Tanzania and South Africa. During the 2010 wet year period, the response is weaker in most parts except for the northern part of Madagascar and southwestern part of South Africa (Figure 10F).



610 Figure 10. Spatial pattern changes of observed LAI response during extreme wet and dry years. For (A) – (C) changes were calculated as a difference between the wet year and the 30-year mean and for (D) – (F) between the dry year and the 30-year mean. White areas indicate no correlation.

### 615 3.9 Comparison of global and regional distribution of LAI response to droughts (1982 – 2011)

There is variability in the global and regional temporal distribution of LAI response to drought (at 12-month timescale) when global vegetation biomes are split into regional biomes (Fig. 11). The map of the global biomes is shown in Fig. S1 in the supplementary material. The observed global response, indicates a decreasing trend of LAI response to drought while the model mean shows an increasing estimate.


625 The semi-desert biome dominates the LAI response as higher drought-vegetation correlations are observed (Figures 11G – I). Over the biome, there is more marked interannual variability which



makes the biome an important player in global carbon cycling (Poulter *et al.*, 2014). The response over the semi-desert in southern Africa is however weaker in comparison to the other semi-desert biomes.

630 The response over the Mediterranean vegetation in Australia is stronger than the Mediterranean vegetation over the rest of inland southern Africa. Over the biome, model simulates closer magnitudes in the latter than over Australia (Figures 11B and 9C).

635 Over the tropical forest biomes, there is a weaker response in Central Africa compared to Southern Africa and South America; the model simulates a closest response magnitude in South America (Figures 11D – F).



640 Figure 11. Correlations between SPEI (12-month timescale) and ensemble mean of LAI from TRENDY (blue line), GIMMS LAI (black line) for (A) Global (B) Mediterranean vegetation over Australia (C) Mediterranean vegetation over southern Africa (D) Tropical Forest over southern Africa (E) Tropical Forest over Central Africa (F) Tropical Forest over South America (G) Semi-desert biome over southern Africa (H) Semi-desert over Asia and (I) Semi-desert over South America.

645

## 4 Discussion

### 4.1 Relationship of LAI to phenological changes

LAI is a variable that is needed for global modelling of biogeochemistry, climate, ecology and hydrology and different primary production models (e.g., Running & Coughlan, 1988; Sellers *et al.*, 1996; Bonan, 2002). In view of the need to run biogeochemical models at regional and global scales, accurate LAI data at moderate – high resolutions is crucial (Wang *et al.*, 2004). The relationship between NDVI and LAI is applied as a support algorithm in MODIS LAI. Thus, from the viewpoint of availability of data, retrieving LAI from analyzing NDVI-LAI relationship remain the main perspective for high temporal resolution in regional and global-wide studies (Wang *et al.*, 2004).

LAI showed a linear relationship with NDVI. This suggests that the NDVI is associated with the phenological changes of plants, the parts of surface cover class which contribute to the general reflectance as well as the variations in the angle of solar zenith (Wang *et al.*, 2004). Studies (e.g. Myoung *et al.*, 2013) have however, found that the relationship between NDVI and LAI varies intra- and inter-annually; and both vegetation indices differ temporally and seasonally over deciduous forests, which are sometimes not accounted for in models that test their relationship (Wang *et al.*, 2004). For instance, while the relationship is strong during periods of leaf production and senescence, no relationship is observed during the period of leaf constant due to NDVI saturation above certain LAI values (Xue and Su, 2017).

Although the present study found a strong linear relationship between the NDVI and LAI in southern Africa, other studies (Potitthep *et al.*, 2010; Towers *et al.*, 2019) have shown the two indices are not always directly proportional. For example, both indices do not exhibit the same relationships over different eco-regions such as the Evergreen Broadleaf Forest, Deciduous Needleleaf Forest. Furthermore, other studies (Fan *et al.*, 2008; Tian *et al.*, 2017) found that the LAI may be better indicator of plant biomass and health because of the saturation associated with the NDVI, particularly in drylands. This makes the LAI more applicable in monitoring vegetation response to drought. Evaluating how the LAI differs from the NDVI over different biomes (such as dry savanna, tropical forest, etc), with regards to temporal difference is shown in Fig. S7. Both the LAI and NDVI show similar annual cycles over southern Africa, except for the Tropical forest and Mediterranean vegetation.

### 4.2 The importance of sub-monthly data in drought computation and monitoring

The data used to evaluate drought indices is CRUJRA. JRA is a reanalysis and has 6-hourly temporal resolution. Additionally, CRUJRA is the data used to force the DGVMs, so the drought indices are being calculated based on the same data the models use for their simulations. JRA is a reanalysis but the combined CRUJRA product uses the sub-monthly information from JRA, and is constrained to the monthly CRU observations. The comparisons of the data are shown in Figs. 5 and S6. It is useful to use data with shorter times because the study focuses on an evaluation of drought impact, which is sensitive to timescale. In drylands, for instance, the uncertainties

associated with monthly data in drought monitoring are reduced when sub-monthly data are used (Mukherjee *et al.*, 20117). With regards to the precipitation and temperature fields, the difference is negligible.

690 We note that, over different parts of the world, CRU has been widely validated against station data (Harris *et al.*, 2020) and there is a high accuracy of the validation. Therefore, observed SPEI gives a high accuracy of measured drought. Another major advantage of using the observational-based CRU data is its spatial and temporal coverage. Station data are available for very few points and for limited times in the region of interest. The few data that are available are fraught with missing  
695 data, rendering them an unreliable data source (Harris *et al.*, 2020).

### 4.3 Annual cycle of climate and vegetation in southern Africa

Climatologies of meteorological variables show that precipitation drops in JJA and SON seasons over the biomes except over Mediterranean vegetation. The dry condition that is experienced during these seasons could be attributed to the subtropical high pressure system which suppresses  
700 rainfall by shifting the ITCZ (Inter-Tropical Convergence Zone) away from these regions (Naik and Abiodun, 2016).

Observations also show low LAI over some parts of southern Africa (please see Figure S2). The weak gradient in some parts of the region may be due to low winter rains produced by the frontal system which is not sufficient for growth of expanse vegetation (Lange *et al.*, 1999). The  
705 aridification of the western part of southern Africa may be attributed to the influence of cold sea surface temperature (SSTs) of the Namibian Upwelling system along the Namibian coasts (Ward *et al.*, 1983). The aridification does not only result in cessation of river discharge but also sediments that would have favored the growth of drier vegetation (Dupont, 2006).

For Fig. S5, some of the models do not capture this peak around September over the Mediterranean and Tropical biomes. One possible explanation is that the models do not well reproduce the  
710 changes in the biomass and leaf area cover around that period (due to phenological responses to environmental variables). For both biomes, spring rainfall contribute to vegetation growth in the region, which may not be well reproduced by the models. For Fig. 4, the magnitudes ( $< 0.5$ ) of the observation-based LAI over (southern Africa) semi-desert biome is quite “high” because the  
715 region is a pseudo-desert, which experiences very high summer temperatures but does receives some rainfall, and the Okavango river is flowing through it permanently. This region is rich in biodiversity such as *Acacia spp* (trees) and *Aristida and Schmidita spp* (savanna) (see WWF 2001; Street and Prinsloo, 2013; Lawal *et al.*, 2018.) Thus, a 0.5 LAI in the biome, which may be higher than other desert biomes, is reasonable in this region.

### 720 4.4 LAI response to drought in observation

Drought is becoming frequent and more intense in southern Africa (Masih *et al.*, 2014). The frequent and stronger dry spells is observed in Fig. 6. Climate change is expected to increase the frequency and severity of droughts (Tables 2 and 3). The severity and longer durations of drought have enormous impacts on the already endangered vegetation biomes in the region (Hoffman *et al.*, 2009). The results show that drought impacts on vegetation occur across the different seasons  
725

in the region. The seasonal difference in the response of vegetation to drought across biomes is influenced by numerous factors such as vegetation adaptive capacity and resilience, reproduction process and growth stage among others (Zeppel *et al.*, 2014; Corlett, 2016). For instance, over the tropical forest biome, drought has the least impact on vegetation in the region, which could be because of the deeper rooting system of the vegetation which allows them access to soil at the deeper water table (El-Vilaly *et al.*, 2017). It is reported that major drivers of vegetation resilience and productivity are precipitation and temperature which control the evapo-transpirative rate (Allen *et al.*, 2010). It is worth noting that vegetation in southern Africa will be severely impacted if the trends continue in the same trajectory. For instance, the regions where there is a strong vegetation response to drought are experiencing wood encroachment and thus, will likely worsen based on the current trajectory of drought occurrence.

We note that the performance of drought indices is not only limited by the variables used in their computation but also by biomes and location where they are used (Xu *et al.*, 2015). For example, SPI which is calculated using simple methods and has adaptable timescales performs better than SPEI in arid regions (Begueria *et al.*, 2014). However, SPEI which requires more variables for its computation captures drought better in relatively humid zones (Begueria *et al.*, 2014). SPEI is however limited by the potential evapotranspiration (PET) because of its sensitivity to the variable (Xu *et al.*, 2015).

The vegetation situated at the borders of Botswana-Namibia and Mozambique-Zambia respond to droughts at an intermediate timescale (i.e. 9-month). The types of vegetation inhabiting these regions, which are well adapted to water shortage because of their physiological and morphological characteristics, takes prolonged period to respond to drought and thus, do not easily shows symptoms of water strain (Vicente-Serrano *et al.*, 2013). The activities through which vegetation minimize water loss include reduction in photosynthesis, reduced canopy cover (Schwinning & Sala, 2004). The capacity for vegetation to store water is one adaptation for low water ecosystems, as is reduced daytime stomatal conductance and CAM photosynthesis. The disparity in the timescales spatial distribution in models from observation might be because the parameters are represented and estimated in the models (Murray *et al.*, 2011). In addition, the models are not similar in their drought timescales simulations.

The varying response of the tropical forests in different regions may be because of the interplay of precipitation and temperature at different longitudes. Ahlstrom *et al* (2015) showed that temperature is particularly a strong factor in the response of this vegetation. Wang *et al* (2008) also reported that soil moisture variations play a key role in the magnitude of vegetation response. The weak response of LAI to drought over Madagascar and Angola may be attributed to the fact that the vegetation in these regions is able to store water for a long time which it uses during deficit (Chapotin *et al.*, 2006). It may also be because rainfall is not the main regulatory component in the growth of vegetation in these regions (Fuller and Prince, 1996). In regions such as Botswana and Namibia, vegetation is highly dependent on water availability for their ecosystem functions (Anyamba *et al.*, 2003). Please see Fig. S3 in the supplementary material for observed correlations at different timescales.

770 The seasonal response of LAI to drought varies and this could be attributed to many factors. For instance, the sensitivity of the semi-desert to water shortage makes them show quick response to drought (New, 2015). The response of vegetation to drought is particularly stronger in the MAM season, because it is during this period that fruit, leaves and biomass are produced by vegetation (Zeppel *et al.*, 2014). Critical water requirements by vegetation for these developmental activities in the MAM season is the reason for the vegetation respond to droughts at a short timescale (Zeppel *et al.*, 2014). The strongest vegetation response (a correlation of about 0.92) is observed over the semi desert biome (which is in the semi-arid environment) in the JJA season. This is a region with  
775 vegetation which heavily depend on water for all their ecosystems functioning without which they would not survive (New, 2015). The drought response of tropical forest is weaker compared to all other biomes. Over the tropical forest biome, the fairly subtle drought response may be because the biome can be tolerant to drought, have stronger robust capacity and is therefore, not extremely impacted by droughts as are the other biomes (Gilgen *et al.*, 2005; Corlett, 2016).

780 The different response by the global biomes in different geographical locations may be because of climate variations and the sensitivity of LAI to climate variations (Ahlstrom *et al.*, 2015). The dominance of LAI response to drought over the semi-desert biome could be attributed to the global bush encroachment and is in consonance with the increasing greenness (Donohue *et al.*, 2009; Fensholt *et al.*, 2012; Andela *et al.*, 2013). Studies (Cai *et al.*, 2014; Trenberth *et al.*, 2014; Dai *et al.*, 2013; Wang *et al.*, 2014; Ahlstrom *et al.*, 2015) have also found that increased and frequent ENSO events due to climate change have not only led to the expansion of LAI but could increase the water demand by semi-desert vegetation. This comparison is particularly important because until now, there has been little or no study on this.

#### 790 **4.5 How well drought and LAI response is represented in DGVM simulations**

The observed LAI is simulated within the models and calculated by GIMMS based on Mao and Yan, 2019. Lu *et al.*, 2011 found that DGVMs perform better against observations than Earth  
795 system models (ESM) because they use observational-derived climate and can include more complex representations of vegetation processes. The ESM is a coupled model simulating its own climate, while the individual DGVMs models used in the present study are standalone, i.e. are applied with observational based meteorological forcing, and thus we remove one uncertainty. Since offline studies target the DGVM itself, removing one possible issue (incorrect climate  
800 drivers), it became imperative to use DGVM to study drought impacts.

DGVMs simulate the vegetation characteristics and impacts of climate on them. The validation of DGVM simulations of variables such as LAI is quite difficult. This is because of the unavailability of data on large spatiotemporal scale for the different vegetation classes (Potter and Klooster,  
805 1998). Studies (e.g. Potter and Klooster, 1998) have also shown that errors present in the prediction of plant functional types (PFTs) tend to spread to biomass prediction in the model, thus possibly biasing estimates of carbon stored in terrestrial ecosystems. Nevertheless, the DGVMs used in this study simulate the spatial patterns of vegetation distribution though with a magnitude bias as shown in Fig. S2 in the supplementary material.

810 TRENDY models mostly simulate the temporal patterns of global and regional distributions of LAI response to drought. The biases shown by the models have been attributed to the fact that the

models do not factor land use changes (Ahlstrom *et al.*, 2015). This is evident in the simulation of LAI (please see Fig. S2).

815 The models' weaker simulation might also be because some of the DGVMs do not well reproduce the LAI magnitude. The negligible difference in the spatial distributions of SPEI of the models could be due to fact that the model PET does not play a strong role in drought occurrence in the southern Africa and that precipitation is the main driver of drought in the region. The variations in the characterization of hydrological processes in the models are also a source of uncertainty  
820 because they reinforce the bifurcation in runoff outputs which has cascading effects on biospheric changes and evapotranspiration (Murray *et al.*, 2011; Stewart *et al.*, 2004). Also see Fig. S4 for correlations of the model ensemble median at different timescales. Another reason for the biases in the simulations may be to the design of the DGVM experimental set-up, which include the flux deviation between simulations without and with (Murray *et al.*, 2011).

#### 825 **4.6 Variations in observed and simulated vegetation response to drought, and implications on model development**

The biases shown by models could be attributed to the different limitations of individual DGVMs, and addressing these shortcomings would improve models' performances. For example, the sub-  
830 optimal performance of CLM may be partly due to the inability of the model to capture foliage production and root system of vegetation for transpiration. The model is also unable to produce savanna ecosystems, which it simulates by approximating vegetation of forest and grassland ecoregions (Dahlin *et al.*, 2020). In addition, the ineffective simulation of deciduousness would have contributed to the model biases in response simulations. Therefore, targeting these limitations  
835 is important for improving model's performance in simulating morphology and physiological functioning of vegetation biomes. Furthermore, the DGVMs (e.g. JULES, DLEM) used in the study poorly replicate important ecological and physiological processes that are critical to capture the dynamics of savanna systems. Other DGVMs (e.g. JSBACH) poorly simulate significant environmental variables such as fire, which is very crucial for the vegetation growth cycle,  
840 particularly in the savanna biome (Thonicke *et al.*, 2001; Romps *et al.*, 2014; Kim *et al.*, 2018; D'Onofrio *et al.*, 2020). Also, over southern Africa, land use change (LUC) is a common and frequent occurrence, and is an important factor for vegetation turnover. However, most models do not well capture land management, which is an important driver of land cover change in the region. Thus, there is a need for future model development to account for rapid LUC over different regions.  
845 However, the disparity in observed and simulated response of vegetation to drought cannot be fully accounted by the DGVMs alone. The reanalysis (CRUJRA) has also shown some limitations in the simulation of climate variables. Compared to observation-based CRU, CRUJRA has closer magnitudes of maximum and minimum temperature, and addressing this would improve simulated response.

850

## 5 Summary and conclusions

Southern African vegetation is continually affected by drought. In this study, we estimated the spatiotemporal characteristics of meteorological drought in southern Africa using the Standardized Precipitation Evapotranspiration Index (SPEI) over a 30-year period (1982 – 2011). The severity of drought and its impacts on vegetation production were examined at various drought time-scales (1- to 24-month timescales) by correlating the drought index (SPEI) with GIMMS LAI at different timescales. We found that the LAI responds strongly ( $r = 0.6$ ) to drought over the central and south eastern parts of the region, with weaker impacts ( $r < 0.4$ ) over parts of Madagascar and Angola, mostly at a shorter time period (3-, 6-month timescale). We note that, comparing CRU to flux tower precipitation in Skukuza (Kwazulu Natal region) illustrates that CRU captures the timing of precipitation relatively well, though it underestimates the magnitude of dry season precipitation (Fig. S8). For seasonal responses, the semi-desert biome showed the strongest response ( $r = 0.95$ ) to drought at 6-month timescale in the autumn season while the tropical forest biome shows the weakest response ( $r = 0.35$ ) at 6-month timescale in the DJF season.

We assessed the relationship between the NDVI and LAI by computing a grid cell correlation of NDVI and LAI, and examined how well state-of-the-art dynamic global vegetation models (DGVMs) simulate LAI and its response to drought. The DGVM multi-model ensemble mostly overestimated the spatial and seasonal distribution of LAI response to drought in most parts of the region. The results also show that:

- The relationship between the NDVI and LAI is linear, implying that the vegetation index is connected to the changes in phenology of plant, reflectance and the angle of zenith variations from the surface cover class.
- The model ensemble simulates the correlation and timescale of LAI response to drought with biases.
- The model ensemble overestimates observed responses on the seasonal distribution of vegetation – drought correlations across different biomes.
- There is a stronger LAI response to rainfall deficit in dry years than in wet years (for instance, in the year 1983 which was a dry year,  $r > 0.84$  over Namibia and northeastern parts of South Africa and  $r < 0.70$  in 2000, a wet year in the same region); and a variability in the distribution patterns of global and regional response across biomes.

The present study has shown how the LAI responds to drought across the different southern African biomes. Given the present spatial coverage of space monitoring of vegetation in the region, the methods used in the study may be extended towards monitoring and characterizing the impacts of droughts on land cover change, as this may permit real-time monitoring of extreme events on terrestrial vegetation (Yin *et al.*, 2020; Moore *et al.*, 2018). The findings of this study (e.g. timescales of LAI response to drought) could also be used for the development of drought early warning systems in agriculture and forestry sectors. This could assist in the mitigation of direct and indirect costs associated with vegetation production.

Furthermore, this study has applied eleven DGVMs to study how well DGVMs can reproduce the response of LAI in southern African vegetation to drought . While this study may have provided an insight into the capability of DGVMs to simulate vegetation response to drought, the results of the study can however, be improved in some ways. For example, we applied eleven models from the TRENDY DGVMs. For future studies, the number of models should be increased, perhaps from other model intercomparison experiments, because using more models, as well as better quantification, might capture uncertainty better in their simulation of drought response by vegetation. The limitation of the DGVMs can be addressed by optimizing the models so that their capability in reproducing vegetation indices is enhanced and better quantified. In addition, there is a need to improve mechanistic relationships in the models, which could be achieved by enhancing the model approximations which had been done to achieve computational efficiencies (Transtrum *et al.*, 2016). Furthermore, simple phenomenological model could be developed from the complex model. These simple models would use correlations among observations, unlike mechanistic relationships which exploit causative individual constituent and suffer over-fitting problems (Transtrum *et al.*, 2016). Lastly, there may be the need for hybridizing machine learning and mechanistic models (Fayyad *et al.*, 1996; Mitchell, 1997) to simulate vegetation parameters. This is because machine learning models have shown certain advantage in the prediction of outcomes of complex mechanisms by using databases of inputs and outputs for a given task (Fayyad *et al.*, 1996; Mitchell, 1997).

*Data availability.* Sources of data used in this work are provided in Section 2.1 and comprise CRU, CRUJRA, NDVI3g and Trendy DGVMs.

*Author contributions.* S.L. was responsible for conceptualization, developing the initial content of the manuscript, including literature search, data analysis and writing of the manuscript. S.S. and D.L. provided model outputs, and guided on data analysis. J.N., H.W.W., P.F, H.T. and B.H provided model output and guidance in terms of the article structure and finalization of the manuscript.

*Competing interests.* Authors declare no competing of interest.

*Acknowledgements.* The work was supported by the South African National Research Foundation (NRF).



## 940 **References**

- Allen, C.D., Macalady, A.K., Chenchouni, H., Bachelet, D., McDowell, N.; Vennetier, M., and Gonzalez, P.: A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests, *For. Ecol. Manag.*, 259, 660–684, <https://doi.org/10.1016/j.foreco.2009.09.001>, 2010.
- 945 Ahlström, A., Raupach, MR., Schurgers, G., Smith, B., Arneth, A., Jung, M., Reichstein, M., Canadell, JG., Friedlingstein, P., Jain, AK., Kato, E., Poulter, B., Sitch, S., Stocker, BD., Viovy, N., Wang, YP., Wiltshire, A., Zaehle, S., and Zeng, N.: The dominant role of semi-arid ecosystems in the trend and variability of the land CO<sub>2</sub> sink, *Science*, 348, 895– 899, <https://doi.org/10.1126/science.aaa1668>, 2015.
- 950 Andela, N., Liu, YY., van Dijk, A. I. J. M., de Jeu, R. A. M., and McVicar, T. R.: Global changes in dryland vegetation dynamics (1988–2008) assessed by satellite remote sensing: Comparing a new passive microwave vegetation density record with reflective greenness data, *Biogeosciences* 10, 6657–6676, doi:10.5194/bg-10-6657-2013, 2013.
- 955 Anyamba, A., Justice, CO., Tucker, CJ., and Mahoney, R.: Seasonal to interannual variability of Vegetation and fires at SAFARI 2000 sites inferred from advanced very high resolution Radiometer time series data, *Journal of Geophysical Research*, vol 10 no D13, <https://doi.org/10.1029/2002JD002464>, 2003.
- Beguéría, S., Vicente-Serrano, SM., Reig, F., and Latorre, B.: Standardized precipitation evapotranspiration index (SPEI) revisited: parameter fitting, evapotranspiration models, tools, datasets and drought monitoring. *International Journal of Climatology* 34(10), 3001–3023, <https://doi.org/10.1002/joc.3887>, 2014.
- 960 Bonan, G. B., Oleson, KW., Vertenstein, M., Lewis, S., Zeng, X., Dai, Y., Dickinson, DE., and Yang, Z.: The land surface climatology of the NCAR land surface model coupled to the NCAR community climate model, *Journal of Climate*, 11, 1307 – 1327, <https://doi.org/10.1175/1520-0442>, 2002.
- 965 Chapotin, SM., Razanameharizaka, JH., and Holbrook, NM.: Water relations of baobab trees (*Adansonia spp L.*) during the rainy season. Does stem water buffer daily water deficit. *Plant and Cell Environment* 29, 1021–1032. doi. 10.1111-3040.2005.01456.x, 2006.
- 970 Cai, W., Borlace, S., Lengaigne, M., van Rensch, P., Collins, M., Vecchi, G., Timmermann, A., Santoso, A., McPhaden, MJ., Wu, M., England, L., H., Wang, G., Guilyardi, E., Jin, F.-F: Increasing frequency of extreme El Nino events due to greenhouse warming. *Nat. Clim. Change* 4, 111–116, doi:10.1038/nclimate2100, 2014.
- 975 Clark, D. B., Mercado, L. M., Sitch, S., Jones, C. D., Gedney, N., Best, M. J., Pryor, M., Rooney, G. G., Essery, R. L. H., Blyth, E., Boucher, O., Harding, R. J., Huntingford, C., and Cox, P. M.: The Joint UK Land Environment Simulator (JULES), model description – Part 2: Carbon fluxes and vegetation dynamics, *Geosci. Model Dev.*, 4, 701–722, <https://doi.org/10.5194/gmd-4-701-2011>, 2011.
- Corlett, RT.: The Impacts of Droughts in Tropical Forests. *Trends Plant Sci* 21(7) 584–593, <https://doi.org/10.1016/j.tplants.2016.02.003>, 2016.
- 980 Dahlin, K. M. et al. 2015. Environmental drivers of drought deciduous phenology in the Community Land Model. – *Biogeosciences* 12: 5061– 5074.
- Dahlin KM, Akanga D, Lombardozzi DL, Reed DE, Shirkey G, Lei C, Abrahama M & J Chen. (2020) Challenging a global land surface model in a local socio-environmental system. *Land*. 9(398): 1- 21. DOI: 10.3390/land9100398

- 985 Dai, A.: Increasing drought under global warming in observations and models, *Nat. Clim. Change* **3**, 52–58, doi:10.1038/nclimate1633, 2013.
- Department of Environmental Affairs (DEA). 2015. Climate Change Adaptation Plans for South African Biomes (ed. Kharika, J.R.M., Mkhize, N.C.S., Munyai, T., Khavhagali, V.P., Davis, C., Dziba, D., Scholes, R., van Garderen, E., von Maltitz, G., Le Maitre, D., 990 Archibald, S., Lotter, D., van Deventer, H., Midgely, G. and Hoffman, T). Pretoria.
- Donohue, R. J., McVicar, T. R., and Roderick, M. L.: Climate-related trends in Australian vegetation cover as inferred from satellite observations, 1981–2006. *Glob. Change Biol.* **15**, 1025–1039, doi:10.1111/j.1365-2486.2008.01746.x, 2009.
- D’Onofrio, D., Baudena, M., Lasslop, G., Nieradzick, L. P., Wärlind, D., and von Hardenberg, 995 J. (2020). Linking vegetation-climate-fire relationships in subsaharan africa to key ecological processes in two dynamic global vegetation models *front. Environ. Sci.* **8**, 136. 10.3389/fenvs.2020.00136.
- Driver A, Sink, KJ, Nel, JN, Holness, S, Van Niekerk, L, Daniels, F, Jonas, Z, Majiedt, PA, Harris, L and Maze, K 2012. National Biodiversity Assessment 2011: An assessment of South 1000 Africa’s biodiversity and ecosystems. Synthesis Report. South African National Biodiversity Institute and Department of Environmental Affairs, Pretoria.
- Dupont, LM.: Late Pliocene vegetation and Climate and Namibia (Southern Africa) derived From Palynology of ODP site 1082. *Geochemistry, Geophysics and Geosystems, An AGU Journal.* Vol 7 issue 5. doi: 10.1029.2005/gc001208, 2006.
- 1005 Earth Summit, United Nation Conference on Environment and Development.: <https://sustainabledevelopment.un.org/outcomedocuments/agenda21>, 1992.
- El-Vilaly, M.A.S., Didan, K., Marsh, S.E., van Leeuwen, W.J., Crimmins, M.A., and Munoz, A.B.: Vegetation productivity responses to drought on tribal lands in the four corners region of the Southwest USA, *Front. Earth Sci.*, **12**, 37–51, doi. 1010 <https://doi.org/10.1007/s11707-017-0646-z>, 2017.
- Fan, L., Gao, Y., Brück, H., and Bernhofer, Ch.: Investigating the relationship between NDVI and LAI in semi-arid grassland in Inner Mongolia using in-situ measurements *Theoretical and Applied Climatology*, 10.1007/s00704-007-0369-2, 2008.
- Fan, L. et al. Satellite observed pantropical carbon dynamics (2019). *Nat. Plants* **5**, 944–951
- 1015 Fayyad, U., Piatetsky-Shapiro, G., Smyth, P. (Eds): From data mining to knowledge discovery Databases. *AI Magazine* **17**, 37. 1996.
- Fensholt, R., Langanke, T., Rasmussen, K., Reenberg, A., Prince, S. D., Tucker, C., Scholes, R. J., Le, Q. B., Bondeau, A., Eastman, R., Epstein, H., Gaughan, A. E., Hellden, U., Mbow, C., Olsson, L., Paruelo, J., Schweitzer, C., Seaquist, J., and 1020 Wessels. K.: Greenness in semi-arid areas across the globe 1981–2007—an Earth Observing Satellite based analysis of trends and drivers, *Remote Sens. Environ.* **121**, 144–158. doi:10.1016/j.rse.2012.01.017, 2012.
- FAO. 2000b (Eds). *FAO/ESAF Handbook for Defining and Setting up a Food Security Information and Early Warning System (FSIEWS)*. Rome
- 1025 Forkel, M. et al. Trend change detection in NDVI time series: Effects of inter-annual variability and methodology. *Remote Sens.* **5**, 2113–2144 (2013)
- Friederike E L Otto *et al* 2018 *Environ. Res. Lett.* **13** 124010
- Fuller, DO., and Prince, SD.: Rainfall and foliar dynamics in tropical southern Africa: Potential impacts of global climatic change on savanna vegetation, *Clim Change*

- 1030 33,69-96. [10.1007/BF00140514](https://doi.org/10.1007/BF00140514), 1996.
- Gitelson, A. A.: Wide dynamic range vegetation index for remote quantification of biophysical characteristics of vegetation, *Journal of Plant Physiology*, vol. 161, no. 2, pp. 165–173, <https://doi.org/10.1078/0176-1617-01176>, 2004.
- 1035 Gielen B, De Boeck H, Lemmens CMHM, Valcke R, Nijs I, Ceulemans R (2005) Grassland species will not necessarily benefit from future elevated air temperatures: a chlorophyll fluorescence approach to study autumn physiology. *Physiol Plant* 125:52–63
- Glantz, M.H., Betsill, M. and Crandall, K.: Food security in southern Africa. Assessing the use and value of ENSO information. National Center for Atmospheric Research, Boulder, CO, USA, 1997.
- 1040 Goll, D. S., Vuichard, N., Maignan, F., Jornet-Puig, A., Sardans, J., Violette, A., Peng, S., Sun, Y., Kvakic, M., Guimberteau, M., Guenet, B., Zaehle, S., Penuelas, J., Janssens, I., and Ciais, P.: A representation of the phosphorus cycle for ORCHIDEE, *Geosci. Model Dev.*, 10, 3745–3770, <https://doi.org/10.5194/gmd-10-3745-2017>, 2017.
- 1045 Hao, Z., and AghaKouchak, A.: Multivariate standardized drought index: a parametric multi-index model, *Advances in Water Resources*, 57, 12– 18., <https://doi.org/10.1016/j.advwatres.2013.03.009> 2013
- Haverd, V., Smith, B., Nieradzic, L., Briggs, P. R., Woodgate, W., Trudinger, C. M., Canadell, J. G., and Cuntz, M.: A new version of the CABLE land surface model (Subversion revision r4601) incorporating land use and land cover change, woody vegetation demography, and a novel optimisation-based approach to plant coordination of photosynthesis, *Geosci. Model Dev.*, 11, 2995– 3026, <https://doi.org/10.5194/gmd-11-2995-2018>, 2018.
- 1050 Harris, I., Jones, PD., Osborn, TJ., and Lister, DH.: Updated high-resolution grids of monthly climatic observations – the CRU TS3.10 Dataset. *Int. J. Climatol.* 34(3), 623–642, doi: 10.1002/joc.3711, 2014.
- 1055 Harris, I., Osborn, T. J., Jones, P., & Lister, D. Version 4 of the CRU TS monthly high-resolution gridded multivariate climate dataset. *Scientific Data*7, 109 (2020)
- Hoffman, MT., Carrick, PJ., and West, AG.: Drought, climate change and vegetation response in the succulent karoo, South Africa. *South African Journal of Science S.Afr. j. sci.* vol05:1-2, S. Afr. j. sci. vol.105 n.1-2 Pretoria Jan./Feb., 2009
- 1060 Homdee, T., Pongput, K., and Kanae, S.: A comparative performance analysis of three standardized climatic drought indices in the Chi River basin, Thailand. *Agriculture and Natural Resources*, 50(3), 211– 219. <https://doi.org/10.1016/j.anres.2016.02.002>, 2016.
- 1065 Joetzjer E., Delire C., Douville H., Ciais P., Decharme B., D. Carrer, H. Verbeeck, M. De Weirtd, D. Bonal (2015), Improving the ISBACC land surface model simulation of water and carbon fluxes and stocks over the Amazon forest, *Geosci. Model Dev.* 8, 1709-1727, doi:10.5194/gmd-8-1709-2015
- Kato, E., Kinoshita, T., Ito, A., Kawamiya, M., and Yamagata, Y.: Evaluation of spatially explicit emission scenario of landuse change and biomass burning using a process-based biogeochemical model, *Journal of Land Use Science*, 8, 104–122, <https://doi.org/10.1080/1747423X.2011.628705>, 2013.
- 1070 Kim, K.; Wang, M.C.; Ranjitkar, S.; Liu, S.H.; Xu, J.C.; Zomer, R.J. Using leaf area index (LAI) to assess vegetation response to drought in Yunnan province of China. *J. Mt. Sci.* 2017, 14, 1863–1872.
- 1075 Khosravi, H., Haydari, E., Shekoohizadegan, S., and Zareie, S.: Assessment the effect of

- drought on vegetation in desert area using landsat data Egypt. *J. Remote Sens. Space Sci.*, 2007 (20), pp. S3-S12 ISSN 1110-9823, <https://doi.org/10.1016/j.ejrs.2016.11.007>, 2017.
- 1080 Kim, J.B., Kerns, B.K., Drapek, R.J., Pitts, G.S., Halofsky J.E. (2018). Simulating vegetation response to climate change in the Blue Mountains with MC2 dynamic global vegetation model *Clim. Serv.*, 10 , pp. 20-32
- Kwon, M., Kwon, H.H., Han, D.: Spatio-temporal drought patterns of multiple drought indices based on precipitation and soil moisture: a case study in South Korea *Int. J. Climatol.*, 1–19 (2019), [10.1002/joc.6094](https://doi.org/10.1002/joc.6094), 2019.
- 1085 Lange, C. B., Berger, W. H. , Lin, H.-L., Wefer, G. and Shipboard Scientific Party Leg 175.: The early Matuyama diatom maximum off SW Africa, Benguela current system (ODP Leg 175), *Mar. Geol.*, 161, 93–114, [https://doi.org/10.1016/S0025-3227\(99\)00081-X](https://doi.org/10.1016/S0025-3227(99)00081-X), 1999
- 1090 Lawal S.A. The response of southern African vegetation to drought in past and future climate. PhD Thesis University of Cape Town, South Africa (2018)
- Lawal S., Lennard C., jack C. Wolski P., Hewitsin B., and Abiodun B.: The observed and model-simulated response of southern African vegetation to drought. <https://doi.org/10.1016/j.agrformet.2019.107698>, 2019.
- 1095 Lawal S., Lennard C and Hewitson B.: Response of southern African vegetation to climate change at 1.5 and 2.0 degrees global warming above the pre-industrial level. *Climate Services* 16C 100134, 2019.
- 1100 Le Quéré, C., Peters, G. P., Andres, R. J., Andrew, R. M., Boden, T. A., Ciais, P., Friedlingstein, P., Houghton, R. A., Marland, G., Moriarty, R., Sitch, S., Tans, P., Arneeth, A., Arvanitis, A., Bakker, D. C. E., Bopp, L., Canadell, J. G., Chini, L. P., Doney, S. C., Harper, A., Harris, I., House, J. I., Jain, A. K., Jones, S. D., Kato, E., Keeling, R. F., Klein Goldewijk, K., Körtzinger, A., Koven, C., Lefèvre, N., Maignan, F., Omar, A., Ono, T., Park, G.-H., Pfeil, B., Poulter, B., Raupach, M. R., Regnier, P., Rödenbeck, C., Saito, S., Schwinger, J., Segschneider, J., Stocker, B. D., Takahashi, T., Tilbrook, B., van Heuven, S., Viovy, N., Wanninkhof, R., Wiltshire, A., and Zaehle, S.: Global carbon budget 2013, *Earth Syst. Sci. Data*, 6, 235–263, <https://doi.org/10.5194/essd-6-235-2014>, 2014.
- 1105 Lienert, S., and Joos, F.: A Bayesian ensemble data assimilation to constrain model parameters and land-use carbon emissions, *Biogeosciences*, 15, 2909–2930, <https://doi.org/10.5194/bg-15-2909-2018>, 2018.
- 1110 Lu, E., Luo, Y., Zhang, R., Wu, Q., and Liu, L.: Regional atmospheric anomalies responsible for the 2009–2010 severe drought in China, *J. Geophys. Res.*, 116 (D21), <https://doi.org/10.1029/2011JD015706>, 2011.
- Lu, E., Cai, W., Jiang, Z., Zhang, Q., Zhang, C., Higgins, R.W., and Halpert, M.S.: The day-to-day monitoring of the 2011 severe drought in China, *Clim. Dynam.*, <https://doi.org/10.1007/s00382-013-1987-2>, 2014.
- 1115 Luo, L.; Tang, W.; Lin, Z.; Wood, E.F. Evaluation of summer temperature and precipitation predictions from NCEP CFSV2 retrospective forecast over China. *Clim. Dyn.* 2013, 41, 2213–2230
- 1120 Mao, J., and Yan, B.: Global Monthly Mean Leaf Area Index Climatology, 1981-2015. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORN LDAAC/1653>, 2019.
- Masih, I., Uhlenbrook, S., Maskey, S., and Smakhtin, V.: Stream-flow trends and climate

- linkages in the Zagros Mountain, Iran. *Clim. Change*, 104, 317–338, doi:10.1007/s10584-009-9793-x, 201, 2014.
- 1125 Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., Fast, I., Fiedler, S., Popke, D., Gayler, V., Giorgetta, M., Goll, D., Haak, H., Hagemann, S., Hedemann, C., Hohenegger, C., Ilyina, T., Jahns, T., Jimenez Cuesta de la Otero, D., Jungclaus, J., Kleinen, T., Kloster, S., Kracher, D., Kinne, S., Kleberg, D., Lasslop, G., Kornbluh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Möbis, B., Müller, W., Nabel, J. E. M. S., Nam, C., Notz, D., Nyawira, S., Paulsen, H., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Popp, M., 1130 Raddatz, T., Rast, S., Redler, R., Reick, C., Rohrschneider, T., Schemann, V., Schmidt, H., Schnur, R., Schulzweida, U., Six, K., Stein, L., Stemmler, I., Stevens, B., von Storch, J., Tian, F., Voigt, A., de Vrese, P., Wieners, K.-H., Wilkenskjeld, S., Roeckner, E., and Winkler, A Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1.2) and its response to increasing CO<sub>2</sub>, *J. Adv. Model. Earth Sy.*, <https://doi.org/10.1029/2018MS001400>, 2018.
- 1135 McKee, T. B., Doesken, N. J. and Kleist, J.: The relationship of drought frequency and duration to time scales. *Proc. Eight Conf. on Applied Climatology*. Anaheim, CA, Amer. Meteor. Soc. 179–184, 1993
- 1140 Melillo, J.: Climate change, risky business, and a call to action for ecologists. *Ecosystem Health and Sustainability* 1(1), <https://doi.org/10.1890/EHS14-0016.1>, 2015.
- Melton, J. R., and Arora, V. K.: Competition between plant functional types in the Canadian Terrestrial Ecosystem Model (CTEM) v. 2.0, *Geosci. Model Dev.*, 9, 323–361, <https://doi.org/10.5194/gmd-9-323-2016>, 2016.
- 1145 Mitchell TM. 1997*Machine learning*. Boston, MA: McGraw-Hill Series in Computer Science. Mitchell, TD., and Jones, PD.: An improved method of constructing a database of monthly climate observations and associated high- resolution grids. *Int. J. Climatol.* 25: 693–712, <https://doi.org/10.1002/joc.1181>, 2005.
- 1150 Moore, B., Crowell, S. M. R., Rayner, P. J., Kumer, J., O'Dell, C. W., O'Brien, D., Utembe, S., Polonsky, I., Schimel, D., and Lemen, J.: The potential of the geostationary carbon cycle observatory (GeoCarb) to provide multi-scale constraints on the carbon cycle in the Americas. *Frontiers in Environmental Science*, 6, 109, <https://doi.org/10.3389/fenvs.2018.00109> , 2018.
- 1155 Mukherjee, N., Zabala, A., Hugel, J., Nyumba, T. O., Esmail, B. A., and Sutherland, W. J.: Comparison of techniques for eliciting views and judgements in decision-making. *Methods in Ecology and Evolution*, 9, 54– 63, <https://doi.org/10.1111/2041-210X.12940>, 2017.
- Müller C, Cramer W, Hare WL, Lotze-Campen H (2011) Climate change risks for African agriculture. *Proc Natl Acad Sci USA* 108(11):4313–4315. doi:[10.1073/pnas.1015078108](https://doi.org/10.1073/pnas.1015078108)
- 1160 Murray, SJ., Foster, PN., and Prentice, IC.: Evaluation of global continental hydrology as simulated by the Land-surface Processes and eXchanges Dynamic Global Vegetation Model. *Hydrology and Earth System Sciences* 15: 91–105, <https://doi.org/10.5194/hess-15-91-2011>, 2011.
- 1165 Myoung, B., Choi, Y.-S., Hong, S., and Park S. K.: Inter- and intra-annual variability of vegetation in the northern hemisphere and its association with precursory meteorological factors, *Global Biogeochem. Cycles*, 27, 31– 42, doi:10.1002/gbc.20017, 2013.
- Naik, M., and Abiodun, BJ.: Potential impacts of forestation on future climate change in

- Southern Africa. *International Journal of Climatology*. doi:10.1002/joc.4652, 2016.
- 1170 Naumann, G., Alfieri, L., Wyser, K., Mentaschi, L., Betts, R.A., Carrao, H., Spinoni, J., Vogt, J., Feyen, L.: Global changes in drought conditions under different levels of warming. *Geophys. Res. Lett.*, *45*, 3285–3296., <https://doi.org/10.1002/2017GL076521>, 2018.
- 1175 New, M. G., M. Hulme, and P. D. Jones, 1999: Representing twentieth-century space–time climate variability. Part II: Development of 1901–1996 monthly terrestrial climate fields. *J. Climate*
- New M, Hulme M, Jones PD. 2000. Representing twentieth century space-time climate variability. II: development of 1901–1996 monthly grids of terrestrial surface climate. *Journal of Climate* **13**: 2217– 2238.
- 1180 New, M.: Are semi-arid regions climate change hot-spots? Evidence from Southern Africa, African Climate and Development Initiative (ACDI) blog, 2015.
- Oleson, K. W., Dai, Y., Bonan, G., Bosilovich, M., Dickinson, R., Dirmeyer, P., Hoffman, F., Houser, P., Levis, S., Niu, G.-Y., Thornton, P., Vertenstein, M., Yang, Z.-L. and Zeng, X.: Technical description of the Community Land Model (CLM). NCAR Tech. Note NCAR/TN-461+STR, 174 pp, 2004.
- 1185 Oleson, K., Lawrence, D., Bonan, G., Drewniak, B., Huang, M., Koven, C., Levis, S., Li, F., Riley, W., Subin, Z., Swenson, S., Thornton, P., Bozbiyik, A., Fisher, R., Heald, C., Kluzek, E., Lamarque, J., Lawrence, P., Leung, L., Lipscomb, W., Muszala, S., Ricciuto, D., Sacks, W., Tang, J., and Yang, Z.: Technical Description of version 4.5 of the Community Land Model (CLM), NCAR, available at:
- 1190 [http://www.cesm.ucar.edu/models/cesm1.2/clm/CLM45\\_Tech\\_Note.pdf](http://www.cesm.ucar.edu/models/cesm1.2/clm/CLM45_Tech_Note.pdf), 2013.
- Palmer WC.: Keeping track of crop moisture conditions, nationwide: the new Crop Moisture Index. *Weatherwise* 21:156–161, <https://doi.org/10.1080/00431672.1968.9932814>, 1965.
- Pinzon, J. E. and Tucker, C. J.: A Non-Stationary 1981-2012 AVHRR NDVI3g Time Series, *Remote Sensing*, *6*, 6929–6960, <https://doi.org/10.3390/rs6086929>, 2014.
- 1195 Potitsep, S., Nasahara, N., Muraoka, H., Nagai, S. and Suzuki, R.: What is the actual relationship between LAI and VI in a deciduous broadleaf forest. *International Archives of the Photogrammetry, Remote Sensing and Spatial Information Science* *38*, 609–614, 2010.
- Poulter, B., Frank, D., Ciais, P., Myneni, R. B., Andela, N., Broquet, G. J. B, Canadell, J. G., Chevallier, F., Liu, Y. Y., Running, S. W., Sitch, S., and van der Werf, G. R.:
- 1200 Contribution of semi-arid ecosystems to interannual variability of the global carbon cycle. *Nature* *509*, 600–603, doi:10.1038/nature13376pmid:2484788, 2014.
- Potter, C.S., and Klooster, S.A.: Global model estimates of carbon and nitrogen storage in litter and soil pools: response to change in vegetation quality and biomass allocation. *Tellus*, *49B*, 1, <https://doi.org/10.3402/tellusb.v49i1.15947>, 1997.
- 1205 Potter, CS, and Klooster, SA.: Dynamic global vegetation modelling for prediction of plant types and biogenic trace gas fluxes. *Global Ecol Biogeogr* *8*:473–488, <https://doi.org/10.1046/j.1365-2699.1999.00152.x>, 1998
- Rahimzadeh-Bajgiran, P., Omasa, K., & Shimizu, Y. (2012). Comparative evaluation of the Vegetation Dryness Index (VDI), the Temperature Vegetation Dryness Index (TVDI) and the improved TVDI (iTVDI) for water stress detection in semi-arid regions of Iran. *ISPRS Journal of Photogrammetry and Remote Sensing*, *68*, 1–12. doi:10.1016/j.isprsjprs.2011.10.009
- 1210 Rezaei M, Sameni A, Shamsi SRF, Bartholomeus H (2016) Remote sensing of land use/cover

- changes and its effect on wind erosion potential in southern Iran. PeerJ.  
doi:10.7717/peerj.1948
- 1215 Reuter, M., Bösch, H., Bovensmann, H., Bril, A., Buchwitz, M., Butz, A., Burrows, J. P., O'Dell, C. W., Guerlet, S., Hasekamp, O., Heymann, J., Kikuchi, N., Oshchepkov, S., Parker, R., Pfeifer, S., Schneising, O., Yokota, T., and Yoshida, Y.: A joint effort to deliver satellite  
1220 retrieved atmospheric CO<sub>2</sub> concentrations for surface flux inversions: the ensemble median algorithm EMMA, *Atmos. Chem. Phys.*, 13, 1771–1780, doi:10.5194/acp-13-1771-2013, 2013.
- Romps, D.M., Seeley, J.T., Vollaro, D., Molinari J. (2014). Projected increase in lightning strikes in the United States due to global warming *Science*, 346 (6211) , pp. 851-854.
- Running, S. W., & Coughlan, J. C.: A general model of forest ecosystem processes for  
1225 regional applications. *Ecological Modelling*, 42, 124 – 154.,  
[https://doi.org/10.1016/0304-3800\(88\)90112-3](https://doi.org/10.1016/0304-3800(88)90112-3), 1988.
- Santin-Janin, H., Garel, M., Chapuis, J.-L., and Pontier, D.: Assessing the performance of NDVI as a proxy for plant biomass using non-linear models: a case study on the Kerguelen archipelago *Polar Biology*, 32 (2009), pp. 861-871, <https://doi.org/10.1007/s00300-009-0586-5>, 2009.
- 1230 Schaefer, K., Schwalm, C. R., Williams, C., Altaf Arain, M., Barr, A., Chen, J. M., et al. (2012). A model-data comparison of gross primary productivity: Results from the North American Carbon Program site synthesis. *Journal of Geophysical Research*, 117, G03010. <http://dx.doi.org/10.1029/2012JG001960>.
- 1235 Schwinning, S., Starr, B. I., and Ehleringer, J. R., Summer and winter drought in a cold desert ecosystem (Colorado Plateau) part II: effects on plant carbon assimilation and growth. *Journal of Arid Environments* 61:61–78.,  
<https://doi.org/10.1016/j.jaridenv.2004.07.013>, 2005.
- Sinclair, REA., and Beyers, RL.: African Biomes. *Ecology*. doi.10.10/OBO978199830060.,  
1240 2015.
- Singh, S.K., A. Chamorro, M.S. Srinivasan and L. Breuer. 2017. A copula-based analysis of severity-duration-frequency of droughts in six climatic regions of New Zealand. *Journal of Hydrology (New Zealand)*, 56(1):13
- Sellers, P. J., Randall, D. A., Collatz, G. J., Berry, J. A., Field, C. B., Dazlich, D. A., Zhang, C.,  
1245 and Collelo, GD., and Bounuo, L.: A revised land surface parameterization (SIB2) for Atmospheric GCMs: Part I. Model formulation. *Journal of Climate*, 9, 676 – 705.,  
<https://doi.org/10.1175/1520-0442, 1996>.
- Sitch, S., Huntingford, C., Gedney, N., Levy, P.E., Lomas, M., Piao, S.L., Betts, R., Ciais, P., Cox, P., Friedlingstein, P., Jones, CD., Prentice, IC., and Woodward, FI.: Evaluation of the terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using five Dynamic Global Vegetation Models (DGVMs). *Global Change Biol.*, 14, 2015– 2039, <https://doi.org/10.1111/j.1365-2486.2008.01626.x>, .2008.
- 1250 Stagge, J. H., Tallaksen, L.M., Gudmundsson, L., van Loon, A.F. and Stahl, K.: Pan-European comparison of candidate distributions for climatological drought indices (SPI and SPEI) *Hydrology in a Changing World: Environmental and Human Dimensions*  
1255 *Proceedings of FRIEND -Water 2014, Montpellier, France, October 2014 (IAHS Publ. 363, 2014.*
- Stewart, IT., Cayan, DR., Dettinger, MD.: Changes in snowmelt runoff timing in western

- 1260 North America under a ‘business as usual’ climate change scenario. *Climatic Change* 62(1–3): 217–232., <https://doi.org/10.1023/B:CLIM.0000013702.22656.e8>, 2004.
- Street RA & Prinsloo G. (2013). Commercially Important Medicinal Plants of South Africa: A Review. *Journal of Chemistry Vol 2013. Article ID 205048 16pages*.
- 1265 Sultan, B., and Gaetani, M.: Agriculture in West Africa in the twenty-first century: Climate change and impacts scenarios, and potential for adaptation. *Frontiers in Plant Science*, **7**, 1– 20, doi: 10.3389/fpls.2016.01262, 2016
- Thenkabail, P.S., Gamage, M.S.D.N., Smakhtin, V.U., 2004. The use of remote sensing data for drought assessment and monitoring in Southwest Asia. Research report. 85, Int. Water Manage. Inst. Colombo, Sri Lanka
- 1270 Teuling, AJ., vanLoon A., Seneviratne S., Lehner, M., Aubinet M., Heinesch B., Bernhofer, C., Grunwald, T., Prasse, H., and Spank, U.: Spank Evapotranspiration amplifies European summer drought *Geophys. Res. Lett.*, **40** (10), pp. 2071-2075, <https://doi.org/10.1002/grl.50495>, 2013.
- 1275 Tian, H. Q., Chen, G. S., Lu, C. Q., Xu, X. F., Hayes, D. J., Ren, W., Pan, S. F., Huntzinger, D. N., and Wofsy, S. C.: North American terrestrial CO<sub>2</sub> uptake largely offset by CH<sub>4</sub> and N<sub>2</sub>O emissions: toward a full accounting of the greenhouse gas budget, *Climatic Change*, **129**, 413–426, <https://doi.org/10.1007/s10584-014-1072-9>, 2015.
- Tian, F., Brandt, M., Liu, Y. Y., Rasmussen, K., and Fensholt, R.: Mapping gains and losses in woody vegetation across global tropical drylands. *Global Change Biology*, **23**(4), 1748– 1760. <https://doi.org/10.1111/gcb.13464>, 2017.
- 1280 Towers, Pedro C.; Strever, Albert; Poblete-Echeverría, Carlos. 2019. "Comparison of Vegetation Indices for Leaf Area Index Estimation in Vertical Shoot Positioned Vine Canopies with and without Grenbiule Hail-Protection Netting" *Remote Sens.* **11**, no. 9: 1073. <https://doi.org/10.3390/rs11091073>
- 1285 Transtrum, MK., Qiu, P.: Bridging mechanistic and phenomenological models of complex biological systems. *PLoS Comput Biol* **12**(5):e1004, 915, <https://doi.org/10.1371/journal.pcbi.1004915>, 2016.
- Trenberth, K. E., A. Dai, G. van der Schrier, P. D. Jones, J. Barichivich, K. R. Briffa, and J. Sheffield.: Global warming and changes in drought, *Nat. Clim. Change*, **4**(1), 17– 22., <https://doi.org/10.1038/nclimate2067>, 2014.
- 1290 Thonicke, K., Venevsky, S., Sitch, S., and Cramer, W.: The role of fire disturbance for global vegetation dynamics: coupling fire into a Dynamic Global Vegetation Model, *Global Ecol. Biogeogr.*, **10**, 661–677, 2001.
- Ujeneza, E., and Abiodun, BJ., Drought regimes in Southern Africa and how well GCMs simulate them? *Clim Dyn*, **44** (5), 1595-1609, 10.1007/s00382-014-2325-z, 2014.
- 1295 Ujeneza E.L. Simulating the Characteristics of Droughts in Southern Africa M.Sc. Thesis University of Cape Town, South Africa (2014)
- United Nations Environmental Protection – UNEP (2008). Atlas of changing environment. Biomes of Africa. *Nairobi, Kenya: UNEP*.
- 1300 Vicente-Serrano, S.M. and López-Moreno, J.I. (2006) The influence of atmospheric circulation at different spatial scales on winter drought variability through a semi-arid climatic gradient in northeast Spain. *International Journal of Climatology*, **26**(11), 1427– 1453.
- Vicente-Serrano, SM., Beguería, S., López-Moreno, JI., Angulo, M. El Kenawy, A.: A new



- global 0.5° gridded dataset (1901-2006) of a multiscalar drought index: comparison with current drought index datasets based on the Palmer Drought Severity Index. *Journal of Hydrometeorology* 11: 1033–1043, <https://doi.org/10.1175/2010JHM1224.1>, 2010.
- 1305 Vicente-Serrano, S. M, Bengueria, S., and Lopez-Moreno, J.: Comment on “Characteristics and trends in various forms of the Palmer Drought Severity Index (PDSI) during 1900–2008” by Aiguo Dai. *J. Geophys. Res.*, 116, D19112, doi:10.1029/2011JD016410, 2011.
- 1310 Vicente-Serrano, S.M., Gouvelia, C., Camarero, J.J, Begueria, S., Trigo, R., Lopez-Moreno, JI., Azorin-Molina, A., Pasho, E., Lorenzo-Lacruz, J, Revuelto, J., Moran-Tejeda, E., and Sanchez-Lorezo, A.: Response of vegetation to drought time-scales across global land biomes. *PNAS Vol 110 1*. [www.pnas.org/cgi/doi/10.1073/pnas.1207068110](http://www.pnas.org/cgi/doi/10.1073/pnas.1207068110), 2012.
- Vicente-Serrano, S.M. (2013) Spatial and temporal evolution of precipitation droughts in Spain in the last century. In: C.C.-L. Martínez and F.V. Rodríguez (Eds.) *Adverse Weather in Spain*, pp. 283–296. Madrid, Spain: Consorcio de Compensación de Seguros.
- 1315 Vicente-Serrano, Sergio M. and National Center for Atmospheric Research Staff (Eds.): "The Climate Data Guide: Standardized Precipitation Evapotranspiration Index (SPEI)." Retrieved from <https://climatedataguide.ucar.edu/climate-data/standardized-precipitation-evapotranspiration-index-spei>, 2015.
- 1320 Wang, Q., Tenhunen, J., Dinh, NQ., Reichstein, M., Vesala, T., and Keronen, P.: Similarities in ground- and satellite-based ndvi time series and their relationship to physiological activity of a Scots pine forest in Finland. *Remote Sensing of Environment* 93, 225–237., 10.1016/j.rse.2004.07.006, 2004.
- 1325 Wang A, Li, KY., and Lettenmaier DP.: Integration of the Variable Infiltration Capacity Model Soil hydrology scheme into the Community Land Model. *Climate and Dynamics*, <https://doi.org/10.1029/2007JD009246>, 2008.
- Wang S 2010 Three characteristics of meteorological drought in southwest china *China Meteorol. News* 3 3–4 (in Chinese)
- 1330 Wang, X., Piao, S., Ciais, P., Friedlingstein, P., Myneni, R.B., Cox, P., Heimann, M., Miller, J., Peng, S., Wang, T., Yang, H., and Chen, A.: A two-fold increase of carbon cycle sensitivity to tropical temperature variations. *Nature* 506, 212–215 (2014). doi:10.1038/nature12915 pmid:24463514, 2014.
- Ward, J. D., Seely, M. K., and Lancaster, N.: On the antiquity of the Namib, *S. Afr. J. Sci.*, 79, 175–183, 1983.
- 1335 Wilhite, D.A., and Glantz, M.H.: Understanding the drought phenomenon: the role of definitions, *Water Int.*, 10 (3), pp. 111-120, <https://doi.org/10.1080/02508068508686328>, 1985.
- 1340 Woodward, FI., and Lomas, MR.: Vegetation-dynamics – simulating responses to climate Change. *Biological Reviews* 79, 643 – 670, <https://doi.org/10.1017/S1464793103006419>, 2004.
- Wolski P, Otto F E L, Odoulami R and New M 2018 Attribution of the three-year drought that triggered the 2017 ‘Day Zero’ water crisis in Cape Town, South Africa. In explaining extreme events of 2017 from a climate perspective *Spec. Suppl. Bull. Am. Meteorol. Soc.*
- 1345 Wolski P 2018 How severe is Cape Town’s ‘Day Zero’ drought? *Significance* 15 24–7
- WRI (World Resources Institute). 2000. World resources 2000–2001. People and ecosystems: WWF(2001). Assessed on May 17, 2017.  
[http://www.wwf.org.za/our\\_work/initiatives/grasslands.cfm](http://www.wwf.org.za/our_work/initiatives/grasslands.cfm)
- Xu, K., Yang, D., Yang, H., Li, Z., Qin, Y., Shen, Y.: Spatio-temporal variation of drought

- in China during 1961–2012: a climatic perspective *J. Hydrol.*, 526, pp. 253-264, 10.1016/j.jhydrol.2014.09.047, 2015.
- 1350 Xue, J., and Su, B.: Significant remote sensing vegetation indices: a review of developments and applications, *J. Sens.*, p. 17, <https://doi.org/10.1155/2017/1353691>, 2017.
- Yang, Y. J., Fang, J., W, Ma and Wang, W.: Relationship between variability in aboveground Net primary production in global grasslands. *Geophy Res Lett* Vol 35, 123710, 1355 [doi:10.1029/2008GLO35408](https://doi.org/10.1029/2008GLO35408), 2008.
- Yin, Y., Byrne, B., Liu, J., Wennberg, P., Davis, K. J., Magney, T., et al.: Cropland carbon uptake delayed and reduced by 2019 Midwest floods. *AGU Advances*, 1, e2019AV000140. <https://doi.org/10.1029/2019AV000140>, 2020.
- 1360 Zaehle, S., Ciais, P., Friend, A. D., and Prieur, V.: Carbon benefits of anthropogenic reactive nitrogen offset by nitrous oxide emissions, *Na. Geosci.*, 4, 601–605, <https://doi.org/10.1038/NCEO1207>, 2011.
- Zeppel MJB, Wilk JV and Lewis JD.: Impacts of extreme precipitation on plants. *Biogeosciences*, 11, 3083 – 3093. [doi. 10.5194/bg-11 3083-2014](https://doi.org/10.5194/bg-11-3083-2014), 2014.
- 1365 Zhang, L., Xiao, J., Li, J., Wang, K., Lei, L., and Guo, H.: The 2010 spring drought reduced primary productivity in southwestern China, *Environ. Res. Lett.*, 7 (4), p. 045706, [10.1088/1748-9326/7/4/045706](https://doi.org/10.1088/1748-9326/7/4/045706), 2012
- Zhao, C., Deng, X., Yuan, Y., Yan, H., and Liang, H.: Prediction of drought risk based on the WRF model in yunnan province of China *Adv. Meteorol.*, pp. 1-9, <https://doi.org/10.1155/2013/295856>, 2013.
- 1370 Zhu, X., Liu, D., 2015. Improving forest aboveground biomass estimation using seasonal Landsat NDVI time-series. *ISPRS J. Photogr. Rem. Sens.* 102, 222–231.