



Impact of frozen soil processes on soil thermal characteristics at seasonal to decadal scales over the Tibetan Plateau and North China

Qian Li¹, Yongkang Xue^{2,3}, Ye Liu²

5 ¹Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, 100029, China.

²Department of Geography, University of California Los Angeles (UCLA), CA, 90095-1524, USA.

³Department of Atmospheric and Oceanic Sciences, UCLA, CA, 90095-1524, USA.

Correspondence to: Qian Li (qian@mail.iap.ac.cn)

10 **Abstract.** Frozen soil processes are of great importance in controlling surface water and energy balances during
the cold season and in cold regions. Over recent decades, considerable frozen soil degradation and surface soil
warming have been reported over the Tibetan Plateau and North China, but most land surface models have
difficulty in capturing the freeze-thaw cycle and few validations focus on the effects of frozen soil processes on
15 soil thermal characteristics in these regions. This paper addresses these issues by introducing a physically more
realistic and computationally more stable and efficient frozen soil module (FSM) into a land surface model—the
third-generation Simplified Simple Biosphere model (SSiB3-FSM). To overcome the difficulties in achieving
stable numerical solutions for frozen soil, a new semi-implicit scheme and a physics-based freezing-thawing
scheme were applied to solve the governing equations. The performance of this model, as well as the effects of
20 frozen soil process on the soil temperature profile and soil thermal characteristics, were investigated over the
Tibetan Plateau and North China using observation and models. Results show that the SSiB3 model with the FSM
produces more realistic soil temperature profile and its seasonal variation than that without FSM during the
freezing and thawing periods. The freezing process in soil delays the winter cooling, while the thawing process
delays the summer warming. The time lag and amplitude damping of temperature become more pronounced with
25 increasing depth. These processes are well simulated in SSiB3-FSM. The freeze-thaw processes could increase
the simulated phase lag days and land memory at different soil depths, as well as the soil memory change with the
soil thickness. Furthermore, compared with observations, SSiB3-FSM produces a realistic change of maximum
frozen soil depth at decadal scales. This study shows the soil thermal characteristics at seasonal to decadal scales
over frozen ground can be greatly improved in SSiB3-FSM and SSiB3-FSM can be used as an effective model for
TP and NC simulation during cold seasons.



30 **1. Introduction**

The freeze-thaw process affects the surface thermal characteristics of frozen soil. At short timescales, the freeze-thaw process could delay the winter cooling/spring warming in the frozen soil because of the latent heat received/released through liquid-ice phase change, which affects surface hydrology (Poutou et al., 2004; Li et al., 2010; Bao et al., 2016). At longer timescales, the change of frozen soil and the variations of the freeze-thaw process affect the shrink or expansion of seasonally frozen ground or permafrost, which can affect the active layer or maximum frozen soil depth, water resources (Cuo et al., 2015; Liljedahl et al., 2016; Guo and Wang, 2017) and ecosystems (Yang et al., 2010; Qin et al., 2014; Yi et al., 2014), and is also a crucial response to climate change (Collins et al., 2013; Zhao and Wu, 2019).

Studies have shown that the soil thermal conditions in the frozen ground area of the Tibetan Plateau (TP) and North China (NC) have been experiencing widespread changes since the 1980s, such as a distinct rise in soil temperature at different soil depths (Wu and Zhang, 2008; Zhang et al., 2002) and changes in the soil freeze-thaw processes (Li et al., 2012; Guo and Wang, 2014; Jin et al., 2015; Yang et al., 2018; Li et al., 2020). In recent years, surface water and energy budget modeling on the frozen ground has advanced, especially over the TP (Yang et al., 2009; Zheng et al., 2016, 2017), and current land surface models (LSMs) exhibit improved simulation of soil temperature profiles as soil thaws during the warm monsoon season (Chen et al., 2010; Zeng et al., 2012; Zheng et al., 2014; Cuo et al., 2015). However, more severe warming rates are observed in winter (Zhang et al., 2019), when most LSMs have difficulty in simulating the deep soil temperature and capturing freezing processes over the TP (Su et al., 2013; Zheng et al., 2017). In addition, large discrepancies have been found in the simulation of surface water and energy budgets by different models driven with the same forcing data (Luo et al., 2003; Slater et al., 2007; Zheng et al., 2017) and the most common problem is the systematic under-estimation of soil temperature (Yang et al., 2009; Bi et al., 2016). Unstable simulations are considered to be one of the key obstacles of frozen soil models in frozen ground (Sun, 2005; Bao et al., 2016), and are considered to come from the numerical schemes because the relationship among soil temperature, soil moisture and ice content are highly nonlinear. To date, shortening the time step duration (Flerchinger and Saxton, 1989) and pre-estimating the ice content during numerical iteration are commonly used in frozen soil numerical schemes, but they may make it difficult for the models to reach convergence. Moreover, a heavy computation cost is unavoidable with those approaches. An enthalpy-based soil algorithm was recently applied to solve the nonlinear governing equations in the frozen soil model (Li et al., 2009; Bao et al., 2016). However, it produced a stable solution only at limited sites and has not been tested in regional or global domains.

Moreover, few studies have focused on the model performance based on observed soil temperature anomalies over frozen ground. The ability to preserve of soil temperature anomalies is known as “land memory”, which is characterized by exponential decay in amplitude and linear lag in phase of soil temperature with depth. Characteristics of land memory have been documented through observation analysis and modeling studies (Hu and Feng, 2004; Xue et al., 2018; Liu et al., 2020). Hu and Feng (2004) found that the anomaly of soil enthalpy, which can represent integration of soil temperature through the soil column, could persist for 2–3 months in the top 1 m of soil over the eastern United States, and affect



70 the surface temperature via soil heat flows then affect the variations of summer monsoon rainfall in the southwest. Another study found the soil enthalpy anomaly in soil column of below 40 cm could persist for 3–4 months at three sites over the TP (Xue et al., 2018). Over frozen ground, the effects of frozen soil processes in the land memory are not yet well understood.

Another important issue is the maximum frozen depth (MFD), which occurs during the freezing period in seasonally frozen ground and can be used to quantify long-term changes in seasonally frozen ground regions (Zhang et al., 2001). The MFD decreased at 5.58 cm/decade during 1960–2014 over the TP (Fang et al., 2019). Although the active layer depth (ALD) for permafrost has been investigated over the TP by different models and compared against field measurements (Oelke and Zhang, 2007; Guo and Wang, 2013; Li et al., 2020), the MFD has rarely been investigated by models.

Therefore, comprehensive assessments and improvement of the performance of land surface models for the frozen ground are imperative. In this paper, the third-generation simplified simple biosphere model (SSiB3) was further improved by coupling with a comprehensive multi-layer frozen soil model (FSM) (Zhang et al., 2007; Li et al., 2010). By using one host-biophysical model (SSiB3) with freeze-thaw processes in multi-layer soil (SSiB3-FSM), and comparing its simulated results with observation data as well as the SSiB3 results, this study focused on the soil temperature profile during freezing and thawing periods, the change of annual freeze soil depth and its memory capability during the past decades, to investigate the effects of frozen soil process on these soil thermal characteristics.

This paper is structured as follows. Section 2 describes the models used in this study and the coupling schemes. Section 3 presents the used data and experimental designs and the calculation methods of soil temperature memory. The major results obtained in this study, including the characteristics of the soil temperature profile, variation of MFD and the soil temperature memory, are given in Section 4. A summary is presented in Section 5.

2. Description of the Models and the Coupling Scheme

2.1 Background

The SSiB3 model (Xue et al., 1991; Sun and Xue, 2001) substantially enhances the previous model's ability to simulate cold season temperature dynamics (Xue et al., 2003). However, it only predicts temperatures of near-surface soil layer (T_{gs}) and deep-soil layer (T_d) based on the force-restore method (Deardorff, 1977; Xue et al., 1996). As for the soil water, soil wetness for three soil layers is predicted and the deepest soil depth is 3.5 m under forests or trees. There are some rough estimations on the soil freezing and thawing but realistic physical processes in cold season/regions are absent. It is necessary to introduce a multi-layer frozen soil module based on physical process into SSiB3 for more realistic cold season/region research under the climate change scenarios.

A comprehensive multi-layer FSM (Zhang et al., 2007; Li et al., 2010), which takes into account of the interactions between mass and heat transport including ice and liquid water phase exchange, was coupled with SSiB3 (referred to SSiB3-FSM) for this study.


In the FSM, the freezing-thawing scheme derived from the freezing point depression equation and the soil matric potential equation is based on thermodynamic equilibrium theory and both liquid water and ice content have been taken into account in the frozen soil hydrological and thermal property



parameterization. In addition, a variable transformation approach introducing enthalpy and total water
 mass in the prognostic equations as substitutes for temperature and liquid water content was used so that
 110 the phase change between liquid and ice can be calculated more efficiently. FSM has previously been
 evaluated using observational data from the field station at Rosemount, Minnesota, and many TP sites
 with satisfactory results (Li et al., 2009; 2010).

2.2. Model coupling scheme

The FSM was implemented into the SSiB3 model to describe multi-layer soil heat transfer and water
 115 flow in SSiB3 affected by freeze-thaw processes in soil. A schematic of the coupled model is shown in
 Figure 1. The definition of the symbols in the figure and following equations can be found in Appendix
 A. In the soil part, the soil thermal diffusion scheme, soil moisture transport scheme and the freeze-thaw
 scheme are designed to solve the soil thermal diffusion, soil water diffusion and ice-liquid phase change,
 respectively.



120

Fig. 1 Schematic diagram of SSiB3-FSM. Soil temperature, soil volumetric water content and soil
 volumetric ice content are T , θ and θ_i , respectively. The heat and water flux between soil layers are
 represented by H and Q . The soil layer number is k , which ranges from 1 to N .



125 The soil column is discretized into eight, eleven and twelve layers for desert, grassland and trees,
 respectively. The thickness of each soil layer increases with the soil depth and the depths of the soil
 column vary with vegetation types in SSiB3. The surface soil layer was assigned as 2 cm since the
 variables in the surface are sensitive to the atmospheric diurnal forcing.

2.2.1 Energy balance equations

130 The energy balance equation for canopy indicates that the canopy energy storage change with time
 is affected by the net radiation at the canopy layer and can be written as:

$$C_c \frac{\partial T_c}{\partial t} = R_{nc} - H_c - \lambda E_c \quad (1)$$

The heat budget of the uppermost soil layer is affected by the net radiation at soil surface (R_{ngs} ; W
 m^{-2}), sensible heat (H_{gs} ; $W m^{-2}$), latent heat fluxes (λE_{gs} ; $W m^{-2}$), energy exchange with lower soil layer
 and the phase change between ice and liquid, and can be written as:

135
$$\frac{\partial(C_s T_{gs})}{\partial t} - L_{il} \rho_i \frac{\partial \theta_i}{\partial t} = \frac{\partial}{\partial z} \left(\lambda_{eff} \frac{\partial T_{gs}}{\partial z} \right) + R_{ngs} - H_{gs} - \lambda E_{gs} \quad (2)$$

The energy distribution inside the soil column is controlled by the heat conduction between layers and
 the phase change inside each individual layer, so it can be written as:

$$\frac{\partial(C_s T_s)}{\partial t} - L_{il} \rho_i \frac{\partial \theta_i}{\partial t} = \frac{\partial}{\partial z} \left(\lambda_{eff} \frac{\partial T_s}{\partial z} \right) \quad (3)$$

140 The first term of Eq. (3) on the left is the heat storage change with time in each soil layer. The
 second term is the latent heat due to freezing/thawing. The first term on the right is the convective heat
 transferred between the soil layers. At the bottom boundary layer, it is assumed that there is no heat flux
 from the deeper soil. The differences of energy balance equations for soil between SSiB3 and SSiB3-
 FSM is the phase change between ice and liquid ($L_{il} \rho_i \frac{\partial \theta_i}{\partial t}$) in the SSiB3-FSM and directly use the heat
 conduction equation rather than the force-restore method.

145 2.2.2 Water balance equations in soil layers

The water distribution within the soil is driven by the liquid water movement and liquid-ice phase
 change. This scheme treats the freeze-thaw process as continuous, without a fixed freezing point, and
 allows the coexistence of water and ice to modify the hydraulic and thermal properties of the soil. The
 conservation of liquid flow is expressed as a one-dimensional Richards' equation:

150
$$\frac{\partial \theta_l}{\partial t} = - \frac{\rho_i}{\rho_l} \frac{\partial \theta_i}{\partial t} - \frac{\partial Q_l}{\partial z} - E \quad (4)$$

The liquid water flow rate of Q_l ($m s^{-1}$) is described by Darcy's Law (see (A5) in Appendix B).
 In the SSiB3-FSM, a freeze-thaw process scheme is used, which is derived from the freezing point
 depression and soil water potential curve in frozen soil:

$$\theta_l = \theta_s \left[\frac{L_{il} T}{\rho \psi_0 T_f} (1 + C_k \theta_i)^{-2} \right]^{-\frac{1}{b}} \quad (5)$$

155 This equation has been employed to describe the relationships among soil temperature, soil liquid water
 content and ice content (Li et al., 2010).



2.2.3 Numerical scheme for the thermal and hydrological equations

Equations (1–5) are highly non-linear systems because the ice content and liquid water change rapidly with little soil temperature change during soil freezing or thawing. We previously substituted soil enthalpy and total water mass for soil temperature and volumetric liquid water content in governing equations (Li et al., 2010) to solve highly nonlinear differential equations. This method also retains energy and water conservation and represents the continuous and slow energy change of the frozen soil system during the freezing/thawing process. However, this approach was only tested for limited field sites. While the method was used in the coupled SSiB3-FSM and tested over a global domain, the numerical solutions become unstable during the long-term integrations for some grid points because the global soil properties and meteorological forcing vary widely. Therefore, a semi-implicit solution procedure for the soil energy and water prognostic equations was developed with SSiB3-FSM for this study.

Figure 2 presents a flow chart of the semi-implicit solution procedure for SSiB3-FSM. A semi-implicit backward finite difference approximation was used for the thermal diffusion equations for canopy and soil (Eq. (1–3)). The numerical Eqs. (A1–A3) are shown in Appendix B. Meanwhile Eq. (5) was transformed to a numerical form (A4) so that it can represent relationship between the change of soil temperature (ΔT_s) and the change of soil ice content ($\Delta \theta_i$) assuming the total water mass is conserved during one time step. Then a tridiagonal linear equation system (A5) for the change of soil temperature was derived based on Eqs. (A1–A3, A4). After solving the tridiagonal matrix at different soil layers, the phase change between liquid water and ice ($\Delta \theta_i$) in soil was decided using the change of soil temperature (ΔT_s) during one time step (Eq. A4). Because the phase change has been included while solving the temperature tridiagonal matrix, here obtained the soil water content (θ_i) and soil temperature (T_s) for each soil layer. After the change of ice content has been decided, the water balance equations do not involve the prognostic variable of the ice content. Subsequently, we can solve the tridiagonal matrix for water fluxes at the interface of the soil layers (see Appendix B (A6–A10)). The liquid water content at the current time step can be easily obtained from Eq. (A11).

This semi-implicit scheme for soil temperature, liquid water and ice content in SSiB3-FSM has been tested over the global domain. It can effectively produce stable solutions for long-term integrations with the heat and mass balances.




Fig. 2 Flowchart of the semi-implicit solution procedure for SSiB3-FSM.

3. Data Sets and Experimental Design

3.1 Data sets

190 From 1948 to 2007, the SSiB3 model and coupled offline SSiB3-FSM model have been driven using
 the meteorological forcing from the Princeton global meteorological data set (Sheffield et al., 2006),
 which is developed by combining a suite of global observation-based data sets with the National Centers
 for Environmental Prediction/National Center for Atmospheric Research reanalysis data. The data set
 includes surface air temperature, pressure, specific humidity, wind speed, downward short-wave
 195 radiation flux, downward long-wave radiation flux and precipitation. The spatial resolution is $1^\circ \times 1^\circ$,
 and the temporal resolution is 3 h.

Several observation data sets have been used to evaluate the performance of SSiB3-FSM and SSiB3
 in cold regions. For the near surface temperature (T_{2m}), we used the Global Historical Climatology
 Network version 2 and the Climate Anomaly Monitoring System (GHCN-CAMS) gauge-based 2-m
 200 temperature over land for 1979–2007, which provides global coverage of monthly means on a regular
 resolution of 0.5° latitude \times 0.5° longitude grids (Fan and van den Dool, 2008).

For the soil temperature profile and the MFD over the TP and NC, the monthly mean soil temperature



of 626 stations over China for 1981–2005 has been used (Yang and Zhang, 2016), provided by the China
Meteorological Administration. The data set has nine soil layers at 0, 5, 10, 15, 20, 40, 80, 160 and 320
205 cm.

Since the calculation of land soil temperature memory requires a long time series of soil temperature
data, only the stations with complete records for 1981–2005 and nine soil layers over the TP region
(elevation > 2500 m) and NC (110°E–120°E, 34.5°N–42°N) were selected. There are 14 stations over
the TP and 16 sites over NC used for this study. Fig. 3 shows the spatial distribution of the stations with
210 available soil temperature data for all nine soil layers and for all 12 months of all 25 years over the TP
and NC.




Fig. 3 Geographical distribution of stations with complete soil temperature records for all nine soil
layers for 1981–2005. The red boxed region is North China (110°E–120°E, 34.5°N–42°N) and the
215 locations of 16 sites used in this study are marked by solid circles. The grey line represents the elevation
is above 2500 m and the grey empty circles denote the locations of the 14 sites on the TP.

3.2 Experimental Design and Methods

3.2.1 Control run

A global simulation by SSiB3-FSM and SSiB3 was carried out forced by the Princeton global
220 meteorological data set from 1948–2007. The initial soil temperature and liquid water content profiles
were derived by interpolating the NCEP-DOE Reanalysis II (R2) (NCEP-R2, Kanamitsu et al., 2002)
soil temperature and soil moisture data linearly to the model's soil layers. Because the soil ice content
measurements are unavailable, and the initial soil ice content is essential for the soil hydrological and
thermal properties, we set ice content to zero at the beginning. The first 10 years (1948–1957) were used
225 for model spin up and the simulation for the last 50 years (1958–2007) was analyzed. The observational



data were used to evaluate the model performance and the results from this run were used to analyze the cold regions' thermal characteristics and MFD, as well as their variations under global warming.

3.2.2 Sensitivity run

230 To investigate the sensitivity of the soil temperature profile and other thermal characteristics to the freeze-thaw process, we conducted a sensitive simulation using the SSiB3-FSM under the same initial land surface conditions but without freeze-thaw process in soil. This sensitivity run is referred to as SSiB3-FSMnoICE run hereafter. Both the SSiB3-FSM run and the SSiB3-FSMnoICE run produce multi-layer's soil temperature and soil moisture, MFD, net radiation, latent heat flux and sensible heat flux, as well as the canopy temperature, canopy water and interception.

235 3.2.3 Methodology to determine MFD and soil memory

Based on the classification of the permafrost by Frauenfeld et al. (2004), a site was deemed to be seasonally frozen ground while the soil temperature at 3.2 m is above 0°C. Based on this criterion, the 14 stations over the TP and the 16 sites over NC in this study were all classified as seasonally frozen ground. According to the seasonal characteristic of soil temperature over seasonally frozen ground, the MFD for each year can be defined as an index for the study of seasonal frozen soil variability and change. This paper gives a preliminary estimation of MFD variations based on monthly soil temperature. Following Frauenfeld et al. (2004), the maximum depth of zero isothermal line for some year is defined as the MFD for this year. Frauenfeld et al. (2004) validated this robustness of this approach. It should be noted that the MFD is different from active layer thickness (ALT), because the active layer is defined as "the top layer of ground subject to annual thawing and freezing in areas underlain by permafrost" (van Everdingen, 1998). ALT is suitable for the permafrost but MFD is more suitable for the seasonally frozen ground. As the ALT increases, the permafrost thaws deeper, whereas as the MFD increases, the frozen soil freezes deeper.

240 The persistence of soil temperature can be quantified by temporal scale analysis. Hu and Feng (2004) assumed that the temporal variation of the soil enthalpy in North America followed the first-order Markov process. Instead of analyzing soil temperature only, the variations of soil enthalpy, which represents integration of soil temperature through the soil column, was used to examine the land memory (Hu and Feng, 2004). This study uses the observed and simulated soil temperature from ground stations in NC and the TP, and the method presented in Entin et al. (2000) and Hu and Feng (2004) to analyze the persistence. Land memory is characterized by the variable's autocorrelation, r , satisfying the following:

$$r(\delta t) = \exp\left(-\frac{\delta t}{S}\right) \quad (6)$$

in which δt is the time lag, S is the decay time scale that can characterize a red noise process and $r(\delta t)$ is the autocorrelation coefficient at the lag time (e.g., 1 month, 2 months, 3 months, ...)

4. Results and Discussions

260 4.1 Assessment of simulated surface 2 m temperature and temperature profile

4.1.1 Surface 2 m temperature



Before investigating the frozen soil thermal characteristics and MFD, as well as their variability, the SSiB-FSM is first evaluated using the observational data. The root mean square error (RMSE) and absolute bias (BIAS) between CAMS and simulated surface temperature globally as well as TP and NC from the SSiB3-FSM and SSiB3 are assessed (Table 1). Table 1 shows the annual RMSE and BIAS of SSiB3-FSM are less than those of SSiB3. In addition, in different seasons the SSiB3-FSM shows less bias than SSiB3. Overall, the SSiB3-FSM produces more realistic estimates of surface temperature than SSiB3 and it can predict the heat transfer processes globally and locally with a reliable accuracy, which provides a basis for further discussions.

Table 1. Error statistics of the simulated surface temperature by SSiB3-FSM and SSiB3.

Global	BIAS (°C)		RMSE (°C)	
	SSiB3	SSiB3-FSM	SSiB3	SSiB3-FSM
Annual	1.27	0.97	2.25	1.93
MAM	1.79	1.66	2.71	2.47
JJA	1.81	1.37	2.87	2.42
SON	0.80	0.29	2.34	2.16
DJF	0.64	0.51	2.47	2.24

(2) Soil temperature profile in the TP

The seasonal vertical soil temperature profile strongly mirrors the influence of the air temperature forcing in contrast to the almost isothermal average annual temperature profile (Oelke and Zhang, 2004). The averaged seasonal profiles for observed soil temperature at 14 sites over the TP are shown in Fig. 4a. For easy comparison, only soil temperature profiles in Jan., Apr., July and Oct. are shown, since the four curves represent the characteristics of soil temperature profile in winter, spring, summer and autumn, respectively. At the seasonal scale, the surface layer and the subsurface layers (here referred to surface to ~1 m) are frozen during winter (Jan.), whereas the temperature of deep layers (below 1 m) is above 0°C. The surface and subsurface soil begins to thaw from Mar.. In Apr. the soil is almost unfrozen. Deep soil temperature always stays above 0°C.

There is a generally rising trend in monthly temperature from winter to summer (from Jan. to July) above 2 m soil depth. Between the 2 m and 2.5 m depth there are no apparent changes in soil temperature from Jan. to Apr. Below 2.5 m, there is an inverse trend compared with upper soil temperature. For instance, the temperature in Jan. is higher than that in Apr. at 3 m soil depth. From Apr. to July, the temperature of all soil layers (surface to ~3.2 m) increases with the air temperature due to the increasing solar radiation. As the autumn approaches, the soil temperature above 1.5 m begins to decrease. Below



1.5 m there is a time lag behind the rising trend, leading to higher temperature at 3.2 m in Oct. than that in July. From Oct. to Jan., soil temperature on all layers shows a decreasing trend. Generally, the soil is characterized as seasonally frozen ground. Deep soil temperature shows a time lag compared with the surface layer, and the upper soil temperature (<1.5 m) shows larger seasonal variability than the deep soil temperature.





Fig. 4 The seasonal soil temperature profile over the TP (14 sites) for 1981–2005. (a) observation; (b) simulated by SSiB3-FSM; (c) comparison between the observation and the SSiB3-FSM shifts to the observation climatology.

The simulated soil temperature profile by SSiB3-FSM over the TP in different seasons is shown in Fig. 4b. There is a general consistency between the simulated temperature profile and the observed profile in both vertical distribution and the seasonal variations. Compared against observations, however, the simulated soil temperatures underestimate the temperature in whole soil column throughout all seasons. The air temperature at 2 m in Apr., July and Oct. in SSiB3-FSM is lower than that observed. This systematic bias arises from the forcing data. For example, the observed air temperature in Apr. at 2 m is about 10°C but the forcing for the models is only 6°C. The greatest difference (about 6°C) is in July. Considering the forcing data's bias, we can parallelly move the SSiB3-FSM soil temperature profile to make the simulated and the observed 2m temperature climatology at the same position. Subsequently, the observed and simulated soil temperature profiles almost coincided (Fig. 4c).

(3) Soil temperature profile in NC

The simulated and observed seasonal soil temperature profiles over NC are displayed in Fig. 5a and Fig. 5b, respectively. They show a similar seasonal frozen soil temperature variability to those over the TP but its thawing season is earlier than that of TP. As the observations show, in winter (Jan.), the soil freezes above 40 cm and it begins to thaw in Feb. until Mar. The soil under 40 cm stays above 0°C throughout the year. The simulated temperature profiles and their seasonal variations are adequately consistent with the observations. However, the frozen depth is deeper in SSiB3-FSM than that of the observations in Jan. (Fig. 5c). These differences may be attributed to the parameterization of soil thermal and hydrological process in SSiB3-FSM.



315


Fig. 5 The seasonal soil temperature profile over NC (16 sites) for 1981–2005. (a) Observation; (b) simulated by SSiB3-FSM; (c) observation and SSiB3-FSM.

(4) Comparison with the force-restore method

In SSiB3 with the force-restore method, only surface temperature and deep soil temperature are considered. For the seasonal change of soil temperature, both the seasonal variation of surface soil temperature (T_{gs}) and deep soil temperature (T_d) (Fig. 6a and 6c) are on the same phase, only with a weak lag in T_d . It is difficult to define its precise position of the deep soil temperature layer, which is dependent on the vegetation types and soil conditions. By introducing a multi-layer frozen soil model, SSiB3-FSM not only presents more precise soil temperature profile but also clearly shows the seasonal change of soil temperature at different soil depths (Fig. 6b and 6d). The time lag and amplitude damping of temperature become more pronounced with increasing depth and they are well described in SSiB3-FSM (Fig. 6b and 6d). This improved performance of SSiB3-FSM lays a foundation for further investigation of the characteristics of MFD changes and soil memory.

320

325



330 **Fig. 6** The seasonal soil temperature simulated by SSiB3 and SSiB-FSM over the TP (14 sites) and
 NC (16 sites) for 1981–2005. (a) The seasonal climatology of T_{gs} (0.02 m) and T_d (2.5 m) by SSiB3 over
 the TP; (b) the seasonal temperature climatology by SSiB3-FSM over the TP; (c) the seasonal
 climatology of T_{gs} (0.02 m) and T_d (2.5 m) by SSiB3 over NC, and (d) the seasonal temperature
 climatology by SSiB3-FSM over NC.


335 **4.2. Characteristics of the soil temperature profile over the TP and NC**

4.2.1 Temporal variability of the soil temperature profile over the TP and NC


The temporal variability with depth of soil temperature was further explored by analyzing the phase
 variations with depth. Here, we used the cross-correlation statistical method to analyze how the seasonal
 variability in soil temperature decreases with depth. Because the SSiB3-FSM produced reasonable
 surface and subsurface temperature profiles (as discussed in Section 4.1) and the observational data are
 only at monthly resolution, the 50-year simulated daily soil temperature were used to represent the time-
 space variability of soil temperature over a wide area. Fig. 7a and Fig. 8a show the lag cross-correlations
 between soil temperature of the first layer with other layers over the TP and NC, respectively. The time
 lag at which the maximal correlations occur increases with soil depth. For instance, the soil layer at 59
 cm reaches maximum at about 10 days while the layer at 312 cm needs about 90 days (a season). The
 cross-correlation values decrease after they reach the maximum value and reach zero at about 110 and
 180 days for the soil layer at 59 and 312 cm, respectively. To more clearly display these relationships,
 the soil temperature phase lag time, defined as the point at which the cross-correlation with the first soil
 layer reaches its maximum, is shown in Fig. 7b and Fig. 8b.



layer equals 1, with depth is shown in Fig. 7b and Fig. 8b. The phase lag time increases linearly with the
 350 soil depth, the details of which are presented in Table 2. For the soil depth at 1.5 m over the TP, the phase
 lag for soil temperature is about 43 days (~1.5 months). For the soil depth at 3 m, the phase lag could be
 87 days (~3 months). Over NC, the phase lag for soil temperature is 11 and 32 days at 59 cm and 1.5 m,
 respectively.



355 **Fig. 7** The time–space variability of soil temperature over the TP for 1978–2007. (a) Simulated cross-correlation of first layer soil temperature with other soil layers (red line: 59 cm; purple line: 96 cm; green line: 147 cm; black line: 312 cm) by SSiB3-FSM; (b) the phase lag (days) of simulated temperature over the TP by SSiB3-FSM.



360 **Fig. 8** The time–space variability of soil temperature over NC for 1978–2007. (a) Simulated cross-correlation of the first layer soil temperature with other soil layers (red line: 59 cm; purple line: 96 cm; green line: 147 cm; black line: 312 cm) by SSiB3-FSM; (b) the phase lag (days) of simulated temperature over NC by SSiB3-FSM.



365

Table 2. Phase lag (days) of simulated soil temperature at different soil depths by SSiB3-FSM and SSiB-FSMnoICE.

Depth (m)	Phase lag (days)			
	TP		NC	
	SSiB3-FSM	SSiB3-FSMnoICE	SSiB3-FSM	SSiB3-FSMnoICE
0.59	15	12	11	9
0.96	27	22	20	19
1.47	43	35	32	31
3.12	87	72	71	71
4.47	117	103	100	100

4.2.2 Land memory

370 The land surface temperature anomaly over the TP and North America has been recognized as an
 indicator for extreme hydroclimate events (Xue et al., 2018) because of its preservation of the snow and
 other climate signatures in previous months. Evaluating the soil persistence of SSiB3 and SSiB3-FSM
 and comparison with observed soil memory are crucial for its application in climate studies. The above
 analyses of temporal variability in soil temperature with depth shows the soil temperature simulated by
 SSiB3-FSM is characterized by increasing persistence with depth. This suggests SSiB3-FSM can be used
 375 to study the land persistence of soil enthalpy, which represents integration of soil temperature through
 the soil column. The land persistence of soil enthalpy over the TP has been preliminary investigated by
 Xue et al. (2018) and another paper (Liu et al., 2020). The current paper therefore focuses on analyzing
 the land memory over NC.

380 Taking the natural log on both sides of Eq. (6) and rearranging, we can obtain $\delta t = -\ln[r(\delta t)]$, which
 describes a straight line in the two-dimensional domain of δt and the natural log of autocorrelation, r .
 Following this procedure, we calculated autocorrelations of observed monthly soil enthalpy anomalies
 between 5 and 320 cm at time lags from 1 to 4 months at the 16 stations over NC and plotted their average
 autocorrelations in the $\delta t - \ln[r(\delta t)]$ domain (Fig. 9a). The lagged autocorrelations of the simulated
 monthly soil enthalpy anomalies at different soil layers between 5 and 312 cm were also calculated and
 385 are shown in Fig. 9b. The persistence of soil enthalpy anomalies is determined by the negative inverse
 of the slope of the straight line for each case in Fig. 9. The slope of these lines varies, indicating a different
 persistence time of the soil enthalpy anomaly at different depths.




Fig. 9 The natural log for the auto-correlation of over NC for 1981–2005. (a) Observed; (b) SSiB3-FSM.

390

The persistence values over NC are given in Table 3. For the observations, the persistence of soil enthalpy anomalies is about 1.34 months in the top 40 cm column and increases to longer than 2 months in the top 160 cm column under the soil surface. In the 320 cm soil column the persistence of soil enthalpy anomalies reaches 4.4 months. Similarly, the persistence of simulated soil enthalpy anomalies is 1.37 months in the 43 cm below the surface. It increases to longer than 2 months about 167 cm soil column. The observed persistence change with the soil thickness is reasonably simulated with the SSiB3-FSM.

395

Table 3. The persistence of soil temperature at different soil depths by SSiB3-FSM and the observations.

Depth(m) (obs./model)	Persistence (month)	
	Obs.	SSiB3-FSM
0.40/0.43	1.34	1.37
0.80/0.75	1.60	1.50
1.60/1.67	2.25	2.13
3.20/3.5	4.4	4.75

400

The persistence by SSiB3 with the force-restore method is only 1.2 and 1.23 months for the surface temperature and deep temperature, respectively. SSiB3-FSM shows better performance in simulating the land persistence for the anomalies in soil than SSiB3, which only considers two layers of temperature data.

4.3 Sensitivity of the soil temperature profile to the freeze-thaw process


405

A sensitivity experiment, test-SSiB3-FSMnoICE, in which the freeze-thaw process in soil is not included, was detailed in Section 3.2.2. A comparison of soil temperature profiles was made between SSiB3-FSM and SSiB3-FSMnoICE and the results are shown in Fig. 10. With freeze-thaw parameterization, the latent heat released while freezing, e.g., in Oct., could offset the decreasing soil temperature. The soil temperature, therefore, would be higher than that of SSiB3-FSMnoICE. Over the TP, the largest difference is found for Jan., especially between 50 cm and 1.5 m. The simulated soil temperature by SSiB3-FSM is about 1–1.7°C higher than that by SSiB3-FSMnoICE. Over NC, the large

410



differences are also shown in winter, especially in Jan. The difference in Jan. is about 1–1.2°C from 1.5 cm to 1.5 m, which means the freeze process in soil delays the winter cooling in freezing seasons (from Oct. to Jan.), and delays the summer warming in thawing seasons (from Apr. to July).



415

Fig. 10 Differences in the seasonal soil temperature profile between the SSiB3-FSMnoICE and SSiB3-FSM for 1981–2005 over (a) the TP (14 sites) and (b) NC (16 sites).

The freeze-thaw process has a major impact on soil temperature profile simulation, especially when freezing or thawing occurs. This effect would exert an impact on the spatial and temporal variability of soil temperature and play an essential role in the soil temperature time lag. Table 2 shows that the time lag of SSiB3-FSMnoICE is less than that of SSiB3-FSM in almost every soil layer. In particular, over the TP, the difference of lag days between the SSiB3-FSM scheme and SSiB3-FSMnoICE scheme increases with depth. For instance, at 59 cm, the time lag of SSiB3-FSM scheme is 3 days longer than that of no ice scheme; however, at about 3 m depth, the difference is about 15 days. For NC, the freeze-thaw processes also increase the phase lag days even though the number of phase lag days between the SSiB3-FSM and SSiB3-FSMnoICE show less difference than those over the TP in the upper soil depths. This is because the maximum freezing depth over NC is about 30 cm, much shallower than that over the TP. Correspondingly, the effects of the freeze-thaw process are only exerted at shallower soil depths.

420

425

4.4 MFD over the TP


430

The long-term temperature profiles at 14 stations over the TP and 16 sites over NC exhibit characteristics of seasonally frozen ground, which freezes in winter and thaws in spring at the surface soil and remains unfrozen at 3.2 m depth throughout the entire year. The simulated annual soil temporal variation with soil depth over the TP and NC stations are shown in Fig. 11, which displays the seasonal freezing and thawing processes during 1981–2005 for the TP and NC. Over the TP, the surface soil starts to freeze in the middle of Oct. and the MFD occurs around Apr. at 1.8 m. For NC, the surface soil starts

435



to freeze at the beginning of Dec. and the MFD occurs in Feb. at around 60 cm soil depth, which is much shallower than that of the TP.



440 **Fig. 11** The climatology of simulated daily soil temperature (Left) over the TP (14 sites averaged) and (Right) NC (16 sites averaged) for 1981–2005 (Unit: °C).

The MFD at 14 sites over the TP and 16 sites over NC simulated by SSiB3-FSM was averaged to analyze the changes of the MFD for 1981–2005. Zero-score normalization, which is a commonly used normalization method, was employed to normalize the observed and simulated MFD:

$$y_i = \frac{x_i - \bar{x}}{s} \quad (7)$$

445
$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i, s = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2} \quad (8)$$

where x_i is the original MFD value for 1981–2005 and y_i is the normalized MFD corresponding to x_i .

Both the observed MFD over the TP and NC showed a significant decreasing trend from 1981 to 2005 (Fig. 12). These decreasing trends indicate that, in areas of seasonally frozen ground, the freezing ground became increasingly shallower during this time period. Over the TP, the observed net change is a 23 cm decrease in MFD in 2005 compared with 1981 and the rate of decrease is about 0.92 cm year⁻¹. However, from 1983 to 1990 the rate is about 4.5 cm year⁻¹, which is about four times as much as that in 1981–2005. In the 1990s, the decreasing rate is about 3 cm year⁻¹. After the 2000s, the decreasing trend reduced. A similar decreasing in the simulated normalized deviation of MFD at TP 14 sites for 1981–2005 is shown in Fig. 12a. The rate of decrease intensified during 1983–1990, which was also shown in the observations.

Over NC, the observed decrease in MFD is 13 cm from 1981 to 2005. The highest decreasing rate is about 3.1 cm year⁻¹ from 1981 to 1990, about six times higher than in 1981–2005. The simulated results by SSiB3-FSM also show the consistency with the observations, especially during the 1980s, when the MFD decreasing trend is 1.8 cm year⁻¹.

460 The MFD decreasing trend during the 1980s over the TP may be related to the significant increasing trend in the winter and spring air temperature. Wei et al. (2003) analyzed the inter-decadal



variations of air temperature over the TP and found a global climatic jump in the 1980s on the TP and the air temperature increased more strongly in the winter and spring from the 1980s to 2000s. They showed that the rates of increasing air temperature at most stations were $0.02\text{--}0.04^\circ\text{C year}^{-1}$ in the winter and spring, which leads to an increasing trend in the 10–20 cm soil temperature over the TP (Zhang et al., 2008). For the NC, an increasing winter temperature trend has been detected since 1985 (Zhang et al., 2002; Shen et al., 2010), which may lead to the decreasing MFD over NC since 1980s.

465



470 **Fig. 12** Normalization of MFD over (a) the TP and (b) NC for 1981–2005 for SSiB3-FSM and the observations.

A comparison of MFD between SSiB3-FSM and SSiB3-FSMnoICE was conducted to evaluate the effects of freeze-thaw processes on the MFD. Although the heat and water mass balance due to ice–liquid phase change is not included in SSiB3-FSMnoICE, the soil temperature still experience a large range of variation. Both MFDs show almost the same variations but the MFD in SSiB3-FSM is shallower than that in SSiB3-FSMnoICE (figure not presented). This can be explained by the phase change energy released while freezing, which could offset the decreasing temperature during the freezing period and lead to a higher soil temperature at the same soil depth than that simulated by the SSiB3-FSMnoICE.

475

5. Conclusions

To improve the accuracy of soil temperature simulation in frozen ground, a multi-layer FSM was incorporated to the SSiB3 to represent the freezing-thawing process and the heat and water transfer in a multi-layer frozen soil. By introducing a semi-implicit backward finite difference approximation and a freezing-thawing scheme based on the freezing depression equation, the highly nonlinear equations in multi-layer frozen soil can be efficiently and stably solved by two tridiagonal matrixes in SSiB3-FSM. The simulated results show that with the frozen soil component, the SSiB3-FSM produces realistic soil thermal characteristics than that of SSiB3, especially soil vertical temperature profiles in different seasons.

485

Furthermore, our results confirm the important role of frozen soil processes on soil thermal characteristics at different time scales over NC and the TP. The results show that the phase-change latent heat released while freezing can offset the decreasing soil temperature, therefore the soil temperature could be higher than that of the experiment without the frozen soil process as soil freezes. Further analysis into the spatial and temporal variability of soil temperature showed that the seasonal variability of soil

490



temperature decreases with soil depths and the phase lag damps linearly. The frozen soil process could increase the phase lag of soil temperature from several days in the surface layer to 15 days in deep layers.

495 The investigation of SSiB3-FSM's ability to simulate the variability of maximum frozen depth at decadal scales showed that simulated normalized of MFD over the TP and NC by SSiB3-FSM are in good agreement with the observations for 1981–2005, including the substantial decreasing trends and the variabilities at decadal scales. The frozen soil processes affect the magnitudes but do not change the decreasing trends of MFD. SSiB3-FSM shows shallower MFD than SSiB3-FSMnoICE because the simulated soil temperature in SSiB3-FSM is higher than that in SSiB3-FSMnoICE. In addition, the
500 SSiB3-FSM also can reproduce the reliable soil memory at different soil depths.

The changes of soil properties and their parameterizations have great effects on the surface energy balance. In particular, the soil thermal conductivity shows large spatial variabilities, and the soil thermal properties are heterogeneous in the vertical direction. A disparity of the soil properties between models and observations may result in the difference between the observations and the simulations.

505 Although the SSiB3-FSM is capable of capturing the basic soil thermal characteristics at seasonal and decadal scales over regions of seasonally frozen ground, further analyses into soil hydrological characteristics in the freezing and thawing phases remain to be conducted. In addition, the better performance of SSiB3-FSM than that by the SSiB3 or SSiB3-FSMnoICE is not only attributed to the frozen soil process but also to the multi-layer heat and water transfer scheme. The effects of the soil
510 stratification and the soil column depth on the model's performance over seasonal frozen ground require further study.

515

520

525



Appendix A. List of symbols with units and definition

Symbol	Units	Definition
b	–	Exponent in the Clapp and Hornberger (1978) parameterization
C_c	$\text{J m}^{-3} \text{K}^{-1}$	Canopy volumetric heat capacity
C_k	–	An adjustable constant parameter in Eq. (5)
C_s	$\text{J m}^{-3} \text{K}^{-1}$	Soil volumetric heat capacity
E	$\text{m}^3 \text{m}^{-3} \text{s}^{-1}$	Evaporation for the surface soil or transpiration for the soil root zone layers
E_{ct}	$\text{m}^3 \text{m}^{-3} \text{s}^{-1}$	Canopy transpiration
E_{gs}	$\text{m}^3 \text{m}^{-3} \text{s}^{-1}$	Soil surface evaporation
λE_c	W m^{-2}	Latent heat flux at the canopy layer
λE_{gs}	W m^{-2}	Latent heat flux at the soil surface
g	m s^{-2}	Gravity
H_c	W m^{-2}	Sensible heat flux at the canopy layer
H_{gs}	W m^{-2}	Sensible heat flux at the soil surface
L_{il}	J kg^{-1}	Specific latent heat of fusion
Q_l	m s^{-1}	Liquid water flow rate
q_G	m s^{-1}	Gravitational drainage at the bottom soil layer
R_{nc}	W m^{-2}	Net solar radiation flux at the canopy layer
R_{ngs}	W m^{-2}	Net solar radiation flux at the soil surface
t	s	Time
T_c	K	Canopy temperature
T_d	K	Deep-soil layer temperature in SSiB3
T_f	K	Freezing temperature (273.15)
T_{gs}	K	Near-surface soil layer temperature in SSiB3
T_s	K	Soil temperature for inner soil layers
ρ_l	kg m^{-3}	Density of liquid water
ρ_i	kg m^{-3}	Density of ice
θ_l	$\text{m}^3 \text{m}^{-3}$	Soil volumetric liquid water content
θ_i	$\text{m}^3 \text{m}^{-3}$	Soil volumetric ice content
θ_s	$\text{m}^3 \text{m}^{-3}$	Soil porosity
θ_T	$\text{m}^3 \text{m}^{-3}$	Total soil water during one time step
λ_{eff}	$\text{W m}^{-1} \text{K}^{-1}$	Effective soil thermal conductivity
ψ_0	m	Soil saturated water potential
ψ	m	Soil matric potential

530 Appendix B. Numerical scheme for solving governing equations in SSiB3-FSM

In SSiB3-FSM, a semi-implicit backward finite difference approximation was used for the thermal diffusion in the soil.

For energy balance equation at canopy layer, Eq. (1) can be written as:



$$\begin{aligned}
 C_c \frac{\Delta T_c}{\Delta t} &= R_{nc} + \frac{\partial R_{nc}}{\partial T_c} \cdot \Delta T_c + \frac{\partial R_{nc}}{\partial T_{gs}} \cdot \Delta T_{gs} \\
 &- H_c - \frac{\partial H_c}{\partial T_c} \cdot \Delta T_c - \frac{\partial H_c}{\partial T_{gs}} \cdot \Delta T_{gs} \\
 &- \lambda E_c - \lambda \frac{\partial E_c}{\partial T_c} \cdot \Delta T_c - \lambda \frac{\partial E_c}{\partial T_{gs}} \cdot \Delta T_{gs}
 \end{aligned} \quad (A1)$$

535 where ΔT_c and ΔT_{gs} denote the change of T_c and T_{gs} during a time step.

For groundcover and soil, Eq. (2) can be written as:

$$\begin{aligned}
 C_s \frac{\Delta T_{gs}}{\Delta t} &= (R_{ngs} - H_{gs} - \lambda E_{gs}) + \left(\frac{\partial R_{ngs}}{\partial T_c} \cdot \Delta T_c + \frac{\partial R_{ngs}}{\partial T_{gs}} \cdot \Delta T_{gs} \right) \\
 &- \left(\frac{\partial H_{gs}}{\partial T_c} \cdot \Delta T_c + \frac{\partial H_{gs}}{\partial T_{gs}} \cdot \Delta T_{gs} \right) - \left(\lambda \frac{\partial E_{gs}}{\partial T_c} \cdot \Delta T_c - \lambda \frac{\partial E_{gs}}{\partial T_{gs}} \cdot \Delta T_{gs} \right) \\
 &+ L_{il} \rho_i \frac{\Delta \theta_i}{\Delta t} + 2\lambda_{eff} \frac{T_2 - T_{gs}}{\Delta z_1 + \Delta z_2}
 \end{aligned} \quad (A2)$$

For the inner layer, Eq. (3) can be written as:

$$C_s \cdot \Delta z_j \cdot \frac{\Delta T_{s,j}}{\Delta t} = L_{il} \rho_i \frac{\Delta \theta_{i,j}}{\Delta t} \cdot \Delta z_j + 2\lambda_{eff,j} \frac{T_{s,j+1} - T_{s,j}}{\Delta z_j + \Delta z_{j+1}} - 2\lambda_{eff,j-1} \frac{T_{s,j} - T_{s,j-1}}{\Delta z_{j-1} + \Delta z_j} \quad (A3)$$

540 Assuming the total water mass is conserved during one time step, the change of soil temperature (ΔT_s) and the change of soil ice content ($\Delta \theta_i$) can be derived based on Eq. (5):

$$\Delta \theta_i = \frac{(\theta_s)^{-b} \left(\frac{L_{il}}{g \psi_0 T_f} \right)}{(1+c_k \theta_i)(\theta_T - \theta_i)^{-b} [b(\theta_T - \theta_i)^{-1} (1+c_k \theta_i) + 2c_k]} \Delta T_{gs} \quad (A4)$$

Combining Eq. (5) and Eq. (A1)–(A3), the energy balance equation system can be reorganized as a tri-diagonal linear equation system with soil temperature of all the soil layers:

$$545 \quad A_j \cdot \Delta T_{s,j-1} + B_j \cdot \Delta T_{s,j} + C_j \cdot \Delta T_{s,j+1} = D_j \quad (A5)$$

where A_j , B_j and C_j are the known coefficients and functions of $T_{s,j-1}$, $T_{s,j}$ and $T_{s,j+1}$ at the previous time step. D_j also represents the known values at the previous time.

550 After solving the tri-diagonal matrix for the soil temperature change at different soil layers, we can obtain the soil temperature at the current time step ($T_{s,j}$). In addition, the phase change between liquid water and ice in soil can be decided using the change of soil temperature during one time step. Because the phase change has been included while solving the temperature tri-diagonal matrix, here we can obtain $\Delta \theta_i$ using Eq. (A4). Then we can solve the water fluxes at the interface of the soil layers:

$$Q_{l,j} = K_{l,j} \left[2 \frac{\psi_j - \psi_{j+1}}{z_j + z_{j+1}} + 1 \right] \quad (A6)$$

Combining Eq. (A6) with Eq. (4), we can obtain water balance equations for the surface layer:

$$555 \quad \psi_1^{k+1} = \psi_1^k - \frac{\Delta t}{z_1} \left(\frac{\partial \psi_1}{\partial \theta_{l,1}} Q_{l,1} \right) - \frac{\Delta t}{z_1} E_{gs} \quad (A7)$$

for the root zone layer:

$$\psi_j^{k+1} = \psi_j^k + \frac{\Delta t}{z_j} \frac{\partial \psi_j}{\partial \theta_{l,j}} (Q_{l,j-1} - Q_{l,j} - E_{ct}) \quad (A8)$$



and for the bottom soil layer:

$$\psi_N^{k+1} = \psi_N^k + \frac{\Delta t}{z_N} \frac{\partial \psi_N}{\partial \theta_{l,N}} (Q_{l,N-1} - q_G) \quad (\text{A9})$$

560 Inserting Eqs. (A7)–(A9) into Eq. (A6) and then regrouping Eq. (A6), then we obtain a tridiagonal linear system for liquid water flow as follows:

$$a_j Q_{j-1} + b_j Q_j + c_j Q_{j+1} = d_j \quad (\text{A10})$$

where a_j , b_j and d_j are the known coefficients and functions of ψ_j , ψ_{j+1} and $\theta_{l,j}$, $\theta_{l,j+1}$ at the previous time step. Therefore, the water fluxes at the current time step can be solved using the above
565 tridiagonal matrix. Then the liquid water content at the current time step can be easily obtained from the following equation:

$$\theta_{l,j}^{k+1} = \theta_{l,j}^k + \frac{\Delta t}{z_j} (Q_j - Q_{j+1}) \quad (\text{A11})$$

Code and data availability

570 The SSIB3-FSM code is available upon request from the first author. The analyses are developed within the Grads and Python software environment. The scripts are also available upon request from the first author. The CAMS gridded 2-m temperature is available at <https://www.esrl.noaa.gov/psd/data/gridded/data.ghencams.html>. The Princeton global meteorological forcing data set is available at <http://hydrology.princeton.edu/data.php>. The China Ground 2-m
575 Temperature Grid Dataset is available at <http://cdc.nmic.cn>.

Author contributions

QL conceived the study and developed the model. QL and YKX developed the framework of the study. QL, YKX analyzed all of the simulation results and wrote the paper. QL and YL drafted the figures. All
580 authors discussed the results and contributed to finalizing the paper.

Competing interests.

The authors declare that they have no conflict of interest.

585 Acknowledgements

This research is financially supported by the National Key Research and Development Program of China (2018YFC1505701) and U.S. National Science Foundation AGS-1849654.

References

- 590 Bao, H., Koike, T., Yang, K., Wang, L., Shrestha, M., and Lawford, P.: Development of an enthalpy-based frozen soil model and its validation in a cold region in China, *J. Geophys. Res. Atmos.*, 121(10), 5259–5280, doi:10.1002/2015jd024451, 2016.
- Bi, H., Ma, J., Zheng, W., and Zeng, J.: Comparison of soil moisture in GLDAS model simulations and in situ observations over the Tibetan Plateau, *J. Geophys. Res. Atmos.*, 121(6), 2658–2678, 2016.



- 595 Chen, Y., Yang, K., Zhou, D., Qin, J., and Guo, X.: Improving the Noah Land Surface Model in Arid Regions with an Appropriate Parameterization of the Thermal Roughness Length, *J. Hydrometeorol.*, 11(4), 995-1006, doi:10.1175/2010jhm1185.1, 2010.
- Collins, M., Knutti, R., Arblaster J., Dufresne, J.L., Fichet, T., Friedlingstein, P., Gao, X., Gutowski, W.J., Johns, T., Krinner, G., Shongwe, M., Tebaldi, C., Weaver A.J., and Wehner, M.: Long-term
600 Climate Change: Projections, Commitments and Irreversibility, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex V., and Midgley, P.M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA., 1029-1136, 2013.
- 605 Cuo, L., Zhang, Y., Bohn, T. J., Zhao, L., Li, J., Liu, Q., and Zhou, B.: Frozen soil degradation and its effects on surface hydrology in the northern Tibetan Plateau, *J. Geophys. Res. Atmos.*, 120, 8276-8298, doi:10.1002/2015jd023193, 2015.
- Deardorff, J. W.: Parameterization of the Planetary Boundary layer for Use in General Circulation Models, *Mon. Weather Rev.*, 100, 93-106, 1972.
- 610 Entin, J. K., Robock, A., Vinnikov, K. Y., Hollinger, S. E., Liu, S. X., and Namkhai, A.: Temporal and spatial scales of observed soil moisture variations in the extratropics, *J. Geophys. Res. Atmos.*, 105, 11865-11877, doi:10.1029/2000jd900051, 2000.
- Fan, Y., and van den Dool, H.: A global monthly land surface air temperature analysis for 1948-present, *J. Geophys. Res. Atmos.*, 113, doi:10.1029/2007jd008470, 2008.
- 615 Fang, X., Luo, S., and Lyu, S.: Observed soil temperature trends associated with climate change in the Tibetan Plateau, 1960-2014, *Theor. Appl. Climatol.*, 135, 169-181, doi:10.1007/s00704-017-2337-9, 2019.
- Frauenfeld, O. W., Zhang, T. J., Barry, R. G., and Gilichinsky, D.: Interdecadal changes in seasonal freeze and thaw depths in Russia, *J. Geophys. Res. Atmos.*, 109, doi:10.1029/2003jd004245, 2004.
- 620 Flerchinger, G. N., and Saxton, K. E.: Simultaneous heat and water model of a freezing snow-residue-soil system: 1. Theory and development, *Trans. ASAE*, 32, 565-571, 1989.
- Guo, D. L., and Wang, H. J.: Simulation of permafrost and seasonally frozen ground conditions on the Tibetan Plateau, 1981-2010, *J. Geophys. Res. Atmos.*, 118, 5216-5230, 2013.
- Guo, D., and Wang, H.: Simulated change in the near-surface soil freeze/thaw cycle on the Tibetan
625 Plateau from 1981 to 2010, *Chinese Sci. Bull.*, 59, 2439-2448, doi:10.1007/s11434-014-0347-x, 2014.
- Guo, D., and Wang, H.: Simulated Historical (1901-2010) Changes in the Permafrost Extent and Active Layer Thickness in the Northern Hemisphere, *J. Geophys. Res. Atmos.*, 122, 12285-12295, 2017.
- Jin, R., Zhang, T., Li, X., Yang, X., and Ran, Y.: Mapping surface soil freeze-thaw cycles in China based
630 on SMMR and SSM/I brightness temperatures from 1978 to 2008, *Arct. Antarctic Alp Res.*, 47(2), 213-229, doi: 10.1657/aaar00c-13-304, 2015.
- Kanamitsu, M., Ebisuzaki, W., Woollen, J., Yang, S.-K., Hnilo, J.J., Fiorino, M., Potter, G.L.: NCEP-DOE AMIP-II Reanalysis (R-2), *Bull. Am. Meteor. Soc.*, 83, 1631-1643, 2002.
- Li, H., Wang, J., and Hao, X.: Influence of Blowing Snow on Snow Mass and Energy Exchanges in the



- 635 Qilian Mountainous, *Journal of Glaciology and Geocryology*, 34, 1084-1090, 2012.
- Li, Q., Sun, S., and Dai, Q.: The Numerical Scheme Development of a Simplified Frozen Soil Model, *Adv. Atmos. Sci.*, 26, 940-950, 2009.
- Li, Q., Sun, S., and Xue, Y.: Analyses and development of a hierarchy of frozen soil models for cold region study, *J. Geophys. Res. Atmos.*, 115, doi:10.1029/2009jd012530, 2010.
- 640 Li, X., Wu, T., Zhu, X., Jiang, Y., and Ying, X.: Improving the Noah-MP Model for Simulating Hydrothermal Regime of the Active Layer in the Permafrost Regions of the Qinghai-Tibet Plateau, *J. Geophys. Res. Atmos.*, 125, doi:10.1029/2020jd032588, 2020.
- Liljedahl, A. K., Boike, J., Daanen, R. P., Fedorov, A. N., Frost, G. V., Grosse, G., Hinzman, L. D., Iijima, Y., Jorgenson, J. C., Matveyeva, N., Necsoiu, M., Reynolds, M. K., Romanovsky, V. E., Schulla, J., Tape, K. D., Walker, D. A., Wilson, C. J., Yabuki, H., and Zona, D.: Pan-Arctic ice-wedge degradation in warming permafrost and its influence on tundra hydrology, *Nat. Geosci.*, 9, 312–318, 2016.
- 645 Liu, Y., Xue, Y., Li, Q., Lettenmaier, D., and Zhao, P.: Investigation of the Variability of Near-Surface Temperature Anomaly and Its Causes Over the Tibetan Plateau, *J. Geophys. Res. Atmos.*, 125, e2020JD032800, doi:10.1029/2020jd032800, 2020.
- 650 Luo, L. F., Robock, A., Vinnikov, K. Y., Schlosser, C. A., Slater, A. G., Boone, A., Braden, H., Cox, P., de Rosnay, P., Dickinson, R. E., Dai, Y. J., Duan, Q. Y., Etchevers, P., Henderson-Sellers, A., Gedney, N., Gusev, Y. M., Habets, F., Kim, J. W., Kowalczyk, E., Mitchell, K., Nasonova, O. N., Noilhan, J., Pitman, A. J., Schaake, J., Shmakin, A. B., Smirnova, T. G., Wetzel, P., Xue, Y. K., Yang, Z. L., and Zeng, Q. C.: Effects of frozen soil on soil temperature, spring infiltration, and runoff: Results from the PILPS 2(d) experiment at Valdai, Russia, *J. Hydrometeorol.*, 4, 334-351, 2003.
- Oelke, C., and Zhang, T.: A model study of circum-arctic soil temperatures, *Permafrost Periglac.*, 15, 103-121, doi:10.1002/ppp485, 2004.
- 660 Oelke, C., and Zhang, T.: Modeling the active-layer depth over the Tibetan plateau, *Arctic Antarc. Alp. Res.*, 39, 714-722, 2007.
- Poutou, E., Krinner, G., Genthon, C., and de Noblet-Ducoudré, N.: Role of soil freezing in future boreal climate change, *Climate Dyn.*, 23, 621-639, doi:10.1007/s00382-004-0459-0, 2004.
- Qin, D. H., Zhou, B. T., and Xiao, C. D.: Progress in studies of cryospheric changes and their impacts on climate of China, *J. Meteorol. Res.*, 28, 732-746, 2014.
- 665 Sheffield, J., Goteti, G., and Wood, E. F.: Development of a 50-year high-resolution global dataset of meteorological forcings for land surface modeling, *J. Clim.*, 19, 3088-3111, 2006.
- Slater, A. G., Bohn, T. J., Mc Creight, J. L., Serreze, M. C., and Lettenmaier, D. P.: A multimodel simulation of pan-Arctic hydrology, *J. Geophys. Res. Biogeo.*, 112, doi:10.1029/2006jg000303, 2007.
- 670 Su, Z., de Rosnay, P., Wen, J., Wang, L., and Zeng, Y.: Evaluation of ECMWF's soil moisture analyses using observations on the Tibetan Plateau, *J. Geophys. Res. Atmos.*, 118, 5304-5318, doi:10.1002/jgrd.50468, 2013.
- Sun, S.: Physical, Biochemical Mechanism of Land Surface Process and Its Parameterization [in



- 675 Chinese], Meteorol. Press, Beijing, China, 2005.
- Sun, S., and Xue Y.: Implementing a new snow scheme in simplified simple biosphere model, *Adv. Atmos. Sci.*, 18(3), 335-354, 2001.
- Van Everdingen R. (Ed.): Multi-language glossary of permafrost and related ground-ice terms (revised May 2005), National Snow and Ice Data Center, Boulder, USA., 1998.
- 680 Wei, Z. G., Huang, R., and Dong, W. J.: Interannual and Interdecadal Variations of Air Temperature and Precipitation over the Tibetan Plateau, *Chinese Journal of Atmospheric Sciences*, 27(2), 157-170, 2003.
- Wu, Q., and Zhang, T.: Changes in active layer thickness over the Qinghai-Tibetan Plateau from 1995 to 2007, *J. Geophys. Res. Atmos.*, 115, doi:10.1029/2009jd012974, 2010.
- 685 Xue, Y., Sellers, P. J., Kinter, J. L., and Shukla, J.: A Simplified Biosphere Model for Global Climate Studies, *J. Clim.*, 4(3), 345-364, 1991.
- Xue, Y. K., Zeng, F. J., and Schlosser, C. A.: SSiB and its sensitivity to soil properties - A case study using HAPEX-Mobilhy data, *Global Planet. Change*, 13, 183-194, 1996.
- Xue, Y. K., Sun, S. F., Kahan, D. S., and Jiao, Y. J.: Impact of parameterizations in snow physics and interface processes on the simulation of snow cover and runoff at several cold region sites, *J. Geophys. Res. Atmos.*, 108, doi:10.1029/2002jd003174, 2003.
- 690 Xue, Y., Diallo, I., Li, W., David Neelin, J., Chu, P. C., Vasic, R., Guo, W.D., Li, Q., Robinson, D.A., Zhu, Y.J., Fu, C.B., and Oaida, C.M.: Spring Land Surface and Subsurface Temperature Anomalies and Subsequent Downstream Late Spring-Summer Droughts/Floods in North America and East Asia, *J. Geophys. Res. Atmos.*, 123, 5001-5019, doi:10.1029/2017jd028246, 2018.
- 695 Yang, K., Chen, Y. Y., and Qin, J.: Some practical notes on the land surface modeling in the Tibetan Plateau, *Hydrol. Earth Syst. Sc.*, 13, 687-701, doi:10.5194/hess-13-687-2009, 2009.
- Yang, K., and Zhang, J.: Spatiotemporal characteristics of soil temperature memory in China from observation, *Theor. Appl. Climatol.*, 126, 739-749, 2016.
- 700 Yang, M. X., Nelson, F., Shiklomanov, N., Guo, D. L., and Wan, G.: Permafrost degradation and its environmental effects on the Tibetan Plateau: A review of recent research, *Earth-Sci. Rev.*, 103, 31-44, 2010.
- Yang, S., Wu, T., Li, R., Zhu, X., Wang, W., Yu, W., Qin, Y., and Hao, J.: Spatial-temporal Changes of the Near-surface Soil Freeze-thaw Status over the Qinghai-Tibetan Plateau, *Plateau Meteorology*, 37, 43-53, 2018.
- 705 Yang, Z. P., Hua, O.Y., Song, M.H., Zhou, C.P., Yang, W.B., and Liu, X.P.: Species diversity and above-ground biomass of alpine vegetation in permafrost region of Qinghai-Tibetan Plateau, *Chinese Journal of Ecology*, 29, 617-623, 2010.
- Yi, S., Wang, X., Qin, Y., Xiang, B., and Ding, Y.: Responses of alpine grassland on Qinghai-Tibetan plateau to climate warming and permafrost degradation: a modeling perspective, *Environ. Res. Lett.*, 9, 074014, 2014.
- 710 Zeng, X., Wang, Z., and Wang, A.: Surface Skin Temperature and the Interplay between Sensible and Ground Heat Fluxes over Arid Regions, *J. Hydrometeorol.*, 13, 1359-1370, doi:10.1175/jhm-d-11-0117.1, 2012.



- 715 Zhang, G., Nan, Z., Wu, X., Ji, H., and Zhao, S.: The Role of Winter Warming in Permafrost Change Over the Qinghai-Tibet Plateau, *Geophys. Res. Lett.*, 46, 11261-11269, doi:10.1029/2019gl084292, 2019.
- Zhang, T., Barry, R. G., Gilichinsky, D., Bykhovets, S. S., Sorokovikov, V. A., and Ye, J. P.: An amplified signal of climatic change in soil temperatures during the last century at Irkutsk, Russia, *Climatic Change*, 49, 41-76, doi:10.1023/a:1010790203146, 2001.
- 720 Zhang, W.G., Li, S. X., and Pang, Q. Q.: Variation Characteristics of Soil Temperature over Qinghai-Xizang Plateau in the Past 45 Years, *Acta Geographica Sinica*, 63, 1151-1159, 2008.
- Zhang, X., Sun, S. F., and Xue, Y. K.: Development and testing of a frozen soil parameterization for cold region studies, *J. Hydrometeorol.*, 8, 690-701, 2007.
- 725 Zhang, Y., Wang, Q., Qian, Y., and Sun, Y.: Spatial/Temporal Variations of Winter Air Temperature in North China in recent 50 Years, *Journal of Nanjing Institute of Meteorology*, 25(5), 633-639, 2002.
- Zhao, D., and Wu, S.: Projected Changes in Permafrost Active Layer Thickness Over the Qinghai-Tibet Plateau Under Climate Change, *Water Resour. Res.*, 55, 7860-7875, doi:10.1029/2019wr024969, 2019.
- 730 Zheng, D., van der Velde, R., Su, Z., Booi, M. J., Hoekstra, A. Y., and Wen, J.: Assessment of roughness length schemes implemented within the Noah land surface model for high-altitude regions, *J. of Hydrometeorol.*, 15, 921-937, doi:10.1175/jhm-d-14-0140.1, 2014.
- Zheng, D., Van der Velde, R., Su, Z., Wen, J., and Wang, X.: Assessment of Noah land surface model with various runoff parameterizations over a Tibetan river, *J. Geophys. Res. Atmos.*, 122, 1488-1504, doi:10.1002/2016jd025572, 2017.
- 735 Zheng, D., Van der Velde R., Su, Z., Wen, J., Wang, X., Booi, M. J., Hoekstra, A. Y., Lv, S., Zhang, Y., and Ek, M. B.: Impacts of Noah model physics on catchment-scale runoff simulations, *J. Geophys. Res. Atmos.*, 121, 807-832, doi:10.1002/2015jd023695, 2016.