

# Causal effects of dams and land cover changes on flood changes in mainland China

Wencong Yang<sup>1,2</sup>, Hanbo Yang<sup>1,2</sup>, Dawen Yang<sup>1,2</sup>, and Aizhong Hou<sup>3</sup>

<sup>1</sup> Department of Hydraulic Engineering, Tsinghua University, Beijing 100084, China.

5 <sup>2</sup> State Key Laboratory of Hydro-Science and Engineering, Tsinghua University, Beijing, 100084, China.

<sup>3</sup> Hydrological Forecast Center, Ministry of Water Resources of the People's Republic of China, Beijing, China

*Correspondence to:* Hanbo Yang (yanghanbo@tsinghua.edu.cn)

**Abstract.** Quantifying the effects of human activities on floods is challenging because of limited knowledge and observations~~because the knowledge and observations toward the effects are limited~~. Many previous methods fail to isolate  
10 different effects and reduce the uncertainty caused by small samples. We use panel regressions to derive the sensitivity of annual maximum discharges ( $Q$ ) to the changing values of three human factors: urban areas, cropland areas, and reservoir indexes for large and medium middle~~middle~~ dams. We also test whether the effects increase or decrease with increasing initial values of human factors. This method is applied in 2739 hydrological stations~~757 non-nested catchments~~ in China. Results show that a 1% point increase of urban areas causes around a 2.93.9%~~2.93.9%~~ increase of  $Q$  with a confidence interval  $CI=[1.9%,$   
15  $5.7%]$ . Cropland areas have no significant effect on  $Q$ . Reservoir index has a decreasing effect: a 1 unit increase of reservoir index causes a decrease in  $Q$  from 23.121.4%~~23.121.4%~~ (with  $CI=[11.4%, 29.9%]$ ) to 5.46.2%~~5.46.2%~~ (with  $CI=[3.2%, 9.1%]$ ) for catchments with initial reservoir indexes from 0 to 53~~53~~. Among 61 catchments with significant increases in observed  $Q$  From 1992 to 2017~~in 1992-2017~~, increasing urban areas cause more than 10% increases in  $Q$  in 10.4%~~only 5 (8.2% of 61) catchment~~  
20 of the 2739 catchments, most of which are located in the North Plain of China. From 1960 to 2017~~Among 234 catchments with at least one dam and significant decreases in observed  $Q$  in 1960-2017~~, increasing reservoir indexes cause a more than 10% decreases of in  $Q$  in 53.4% of 777138 (59.8% of 234) catchments~~with at least one dam~~. Among 1249 1074~~1249 1074~~ catchments with limited impacts from urban areas and reservoir indexes, 210 403 (19.632.3%)~~210 403 (19.632.3%)~~ catchments have significant decreases in  $Q$  during 1960-2017~~more than 10% unexplained decreasing rates in  $Q$  per decade during 1960-2017~~, and 62.446.7%~~62.446.7%~~ of the 210 403~~210 403~~ catchments are located in the middle and down streams of the Yellow River Basin and the upper  
25 streams of the Haihe River Basin. This study extends the panel regression method in hydrology and sheds light on the attribution of flood changes on a national scale.

## 1 Introduction

River flooding is one of the most severe disasters in the world. China experiences tremendous damages from floods in the past decades with expanding urban areas, a booming economy, and increasing populations (Du et al., 2019; Kundzewicz et  
30 al., 2019). Sharply changing flood characteristics make flood risk management more difficult. According to a national

investigation of flood peak changes in China conducted by Yang et al. (2019), abrupt changes due to human activities are the predominant mode of flood changes. Understanding how floods change in a changing environment helps flood risk management in the future. Therefore, a quantitative attribution of flood changes is urgent on a national scale for policy decisions.

35 To detect flood changes and pinpoint the underlying reasons, scientists need to answer the following questions: 1. does a factor affect floods~~whether a factor affects floods~~? 2. If the effect presents, how strong is the effect? The drivers of flood changes can be classified into three categories: atmospheric factors, catchment factors, and river factors (Merz et al., 2012; Blöschl et al., 2015). Atmospheric factors refer to the meteorological forcing of water fluxes such as natural climate variability and anthropogenic climate change; catchment factors refer to the alternating physiographic characteristics of catchments, such as land cover changes; river factors refer to hydraulic infrastructures that change river morphology and flood routing, such as dams and channelization (Merz et al., 2012; Blöschl et al., 2015). Catchment and river factors are mainly attributed to human activities, which attract increasing attention in hydrological systems in the era of “socio-hydrology” (Di Baldassarre et al., 2019; Müller and Levy, 2019). However, quantifying human impacts on floods is challenging for the following reasons. Firstly, due to the highly unpredictable human behaviors, we have limited knowledge to reproduce the process of how human activities affect floods (Pande and Sivapalan, 2017). For example, the expansion of cropland and urban areas not only casts deterministic effects on floods through changing soil physics and surface roughness but also brings uncertain effects through irrigations and water diversions. We consider these effects “uncertain” because they are related to unknown human decisions. Secondly, the observations of human activities are limited (Pande and Sivapalan, 2017). In the example above, many regions lack long-term and large-scale data of soil physics, roughness, irrigations, and water diversions, which are highly dependent on a high-cost network of in-site measurements.

Previous studies have used three methods to quantify human impacts on floods. ~~The~~A first method is a physical model simulation~~model-based approach~~. This method regards the impacts of human activities as either the difference between actual observations and the model simulations of floods (Viglione et al., 2016; Lu et al., 2018) or flood changes with time-varying model parameters (Peña et al., 2016; Umer et al., 2019). However, this method suffers from limited model accuracy.

55 The second method is a paired-catchment experiment. This method either compares the floods before and after human impacts in one catchment or compares floods in two groups of catchments with and without human impacts (Prosdocimi et al., 2015; Hodgkins et al., 2019). However, the comparisons above cannot rigorously isolate multiple impacts on floods since we cannot actually control everything except one targeted human factor (Runge et al., 2019). The third method is empirical variable dependence, i.e., using regressions or non-stationary probability distributions to link human factors to flood characteristics (FitzHugh et al., 2011; Prosdocimi et al., 2015; Bertola et al., 2019; De Niel and Willems, 2019). The third method is cost-efficient for large-scale studies, but it has two problems. Firstly, to derive the causal effects of human factors, all confounders — which correlates with human factors and floods at the same time — should be explicitly accounted for in the empirical relationships. However, defining numerous variables to represent confounders may be an endless task. For example, climatic confounders are ambiguous because floods are caused by different climatic factors (e.g., long rainfall,

65 short rainfall, snowmelt, and rain on snow) in different regions (Stein et al., 2020; Yang et al., 2020a; Merz et al., 2020). Therefore, in a large-sample study, we do not have a unified regression form to control all possible variables for all catchments. Secondly, empirical methods require sufficient data for robust statistical inference, while flood samples are rare for each catchment.

Panel regression (Steinschneider et al., 2013; Wooldridge, 2016) solves the problems of the empirical method in two ways. 70 Firstly, panel regression adds virtual variables to the regression to represent a fixed individual or regional effect (Steinschneider et al., 2013). In such a way, the regression can account for the effects of ambiguous confounders that are constant in time or region. Secondly, panel regression pools all samples into one model and trades space for time (Steinschneider et al., 2013). Therefore, the regression result is more reliable even with short flood records for each catchment. Although panel regression is a tool in economics, it has been introduced in hydrology to estimate the effects of 75 forests on floods (Ferreira and Ghimire, 2012), urbanization on runoff coefficients (Steinschneider et al., 2013), dams on streamflow (McManamay et al., 2014), rainfall on streamflow (Bassiouni et al., 2016), deforestation on streamflow (Levy et al., 2018), urbanization on floods (Blum et al., 2020), and rain/snow fraction on floods (Davenport et al., 2020). However, these studies only focused on one factor at one time. Considering more human factors can provide a more comprehensive picture of the human impacts on floods. In addition, only Blum et al. (2020) and Davenport et al. (2020) tested the nonlinear 80 effects of factors. Previous studies rarely examined whether the effects increase or decrease with increasing initial values of human factors.

In this study, supported by ~~an unprecedentedly~~ a large dataset of Chinese floods from ~~2739-757~~ streamflow gauge stations, we quantify the national average sensitivities of annual flood peaks to changing urban areas, cropland areas, and reservoir indexes for large and ~~middle-medium~~ dams using panel regression. We also ~~design a workflow to~~ test whether effects 85 increase or decrease for catchments with increasing initial values of the targeted factor. The causal effects of factors distinguish the flood changes explained and unexplained by the three human factors in recent decades. This study is organized as follows. Section 2 introduces methods. Section 3 describes the data. Section 4 presents the results. Section 5 discusses the methods and the insights gained by this study. Section 6 gives conclusions.

## 2 Methods

### 90 2.1 Causal map of flooding

Causal maps depict the dependency relationship between variables, and they help discover confounders and focus on the causal effects of different factors when fitting a regression model (Pearl and Mackenzie, 2020). A confounder is a variable that influences both human factors and floods. A causal effect is defined as the sensitivity of floods to a factor when all possible confounding variables are controlled. Similar to Blum et al. (2020), we draw a causal map of flooding in Fig. 1. 95 This study estimates the causal effects of the changes in dams, urban areas, and cropland areas on floods, as the three dashed lines show in the figure. Variables lying above the dashed lines are unknown or unobserved mediators. Urban areas and

cropland areas are interrelated because they may change into each other during the process of land cover change. We consider two major confounders. The first confounder is the ~~regional~~-time-varying confounder, which ~~means that it changes over time but keeps~~can be unique for a catchment or spatially constant in a region. ~~Climate is such a variable since it varies temporally and has spatial homogeneity. The trends of climate may correlate with human activities.~~ For example, increasing event precipitation during floods, which varies by individual catchments, may promote dam constructions; decreasing annual precipitation, which happens at a regional scale, exacerbates water shortage and may therefore promote the reservoir constructions or the implementation of the Grain for Green Project. We delineate regions by climate since the climate is the first-order driver of catchment similarity (Jehn et al., 2020). In this way, we can control the effect of many omitted variables that have spatial homogeneity. The second confounder is the individual time-invariant confounder, ~~which means that it keeps constant in time but varies by different catchments.~~ This confounder is mainly represented by the characteristics of catchment landscapes, e.g., topography, soil types, geology. ~~They may also correlate with human activities.~~ For example, urban areas are likely built on flat and plain catchments.




Figure 1. Causal map illustrating the relationships between human factors and floods.

## 2.2 Design of panel regression

115 Panel regression is a statistical technique for panel data (Steinschneider et al., 2013; Wooldridge, 2016). Panel data are observations on several subjects in different periods. Panel regression controls the constant effects of each subject or each period to mitigate regression bias due to omitted variables. Panel regressions in this study are extended from the equation in Blum et al. (2020) and are presented in Eq. (1) ~~and (2)~~ as follows.

$$120 \quad \log(Q_{i,t}) = \alpha_i + g_1(Urban_{i,t}) + g_2(Crop_{i,t}) + g_3(RI_{i,t}) + \pi_{r,t}D_rD_t + D_r(\varphi_rP_{i,t}^{(3)} + \lambda_rP_{i,t}^{(30)}) + \varepsilon_{i,t} \log(Q_{i,t}) = \alpha_i^{(1)} + g_1(Urban_{i,t}) + g_2(Crop_{i,t}) + \pi_{r,t}^{(1)}D_rD_t + \varepsilon_{i,t}^{(1)} \quad (1)$$

$$\log(Q_{i,t}) = \alpha_i^{(2)} + g_3(RI_{i,t}) + \pi_{r,t}^{(2)}D_rD_t + \varepsilon_{i,t}^{(2)} \quad (2)$$

$Q_{i,t}$  is the annual flood peak of catchment  $i$  in year  $t$  ( $m^3 \cdot s^{-1}$ ).  $Urban_{i,t}$  is the urban percentage of catchment  $i$  in year  $t$  (%).  $Crop_{i,t}$  is the cropland percentage of catchment  $i$  in year  $t$  (%).  $RI_{i,t} = \sum_j (A_i^{(j)}/A_i) \cdot DOR_i^{(j)}$ , is the reservoir index of catchment  $i$  in year  $t$ ;  $DOR_i^{(j)}$  is the degree of regulation of reservoir  $j$  in catchment  $i$ , which is the ratio between the storage capacity and total annual flow of the reservoir;  $A_i^{(j)}$  is the upstream area of reservoir  $j$ ;  $A_i$  is the area of catchment  $i$ .  $D_r$  is a region dummy which equals 1 or 0.  $D_t$  is a year dummy which equals 1 or 0.  $P_{i,t}^{(3)}$  is the 3-day total precipitation before the flood peak in year  $t$  of catchment  $i$ , which accounts for the rainfall that causes the flood;  $P_{i,t}^{(30)}$  is the 30-day total precipitation before the flood peak in year  $t$  of catchment  $i$ , which accounts for the soil moisture and snowmelt that cause the flood. The coefficients of  $P_{i,t}^{(3)}$  and  $P_{i,t}^{(30)}$ , namely  $\varphi_r$  and  $\lambda_r$ , are assumed to be constant within a climatic region  $r$ .  $\alpha_i^{(1)}$  and  $\alpha_i^{(2)}$  are the time-invariant constant effects of catchment  $i$  in Eq. (1) and (2) respectively.  $\pi_{r,t}^{(1)}$  and  $\pi_{r,t}^{(2)}$  are the time-varying constant effects of region  $r$  in year  $t$  in Eq. (1) and (2) respectively.  $\varepsilon_{i,t}^{(1)}$  and  $\varepsilon_{i,t}^{(2)}$  are the model residuals in Eq. (1) and (2) respectively. The response functions  $g_1(\cdot)$ ,  $g_2(\cdot)$ , and  $g_3(\cdot)$  represent various response types of  $Q$  to different factors. We assigned  $Urban$ ,  $Crop$ , and  $RI$  into two regressions because their data sources had different temporal lengths. Although  $RI$  may correlate with  $Urban$  and  $Crop$ , the effects of dams and land cover on floods can be derived independently since we have controlled their common drivers (Pearl and Mackenzie, 2020) in each equation, i.e., the regional time-varying term and the individual time-invariant term.

135 A region consists of a group of spatially coherent catchments with a similar climate. Unlike Blum et al. (2020) who used predefined physiographic regions, we delineated regions by using the partitioning around medoids (PAM) algorithm (Reynolds et al., 2006) based on the distance matrix of all catchments defined as follow:

$$140 \quad dist(i, j) = dist_{KG}(i, j) + dist_{cen}(i, j) \quad (32)$$

$$dist_{KG}(i, j) = \frac{1}{2} \sum_{l=1}^{30} |k_i^{(l)} - k_j^{(l)}| \quad (43)$$


$$dist_{cen}(i, j) = \left( \frac{earth.dist(i, j)}{\max\{earth.dist(i, j) | \forall i, j\}} \right)^{1/2} \quad (54)$$

where  $dist_{KG}(i, j)$  is the distance of Köppen–Geiger class (Beck et al., 2018) ratios between catchment  $i$  and  $j$ ;  $k_i^{(l)}$  is the area percentage of Köppen–Geiger class  $l$  in catchment  $i$ ;  $dist_{cen}(i, j)$  is the standardized distance between the geometric centers of catchment  $i$  and  $j$ ;  $earth.dist(i, j)$  is the spherical distance on the earth between the geometric centers of catchment  $i$  and  $j$ .

The effect of a factor  $X$  on  $Q$ , i.e., the percentage change in  $Q$  given a fixed change in  $X$ , is expressed as:

$$\Delta Q(\%) = \frac{\Delta Q_{i,t}}{Q_{i,t}} = \frac{e^{g(X_{i,t}+\Delta X)} - e^{g(X_{i,t})}}{e^{g(X_{i,t})}} \exp(g(X_{i,t} + \Delta X) - g(X_{i,t})) - 1 \quad (65)$$

We considered three types of response functions  $g(\cdot)$ .  $g(X_{i,t}) = \beta X_{i,t}$  indicated a stable effect where the percentage change in  $Q$  only depended on  $\Delta X$ ;  $g(X_{i,t}) = \gamma X_{i,t}^2$  indicated an increasing effect where the percentage change in  $Q$  increased with increasing  $X_{i,t}$ ;  $g(X_{i,t}) = \theta X_{i,t}^{1/2}$  indicated a decreasing effect where the percentage change in  $Q$  decreased with increasing  $X_{i,t}$ . To determine the specific effect type, we fitted regressions with 27 possible combinations of  $[g_1(\cdot), g_2(\cdot), g_3(\cdot)]$  types and tested the p values of the coefficients related to different  $g(\cdot)$  functions in regressions selected the one with the lowest AIC value, as shown in Fig. 2. We used bootstrapping to test the significance and derive the confidence intervals of coefficients  $\beta, \gamma$ , and  $\theta$  so that the model residuals were allowed to be non-Gaussian and the sampling uncertainty could be accounted for. Since the pooling samples were sufficient for statistical inference, we regarded p values smaller than 0.01 as significant.



**Figure 2. Workflow for testing the formulation of response functions  $g(\cdot)$  for factors *Urban*, *Crop* and *RI*. All tests are conducted using Eq. (1) and (2).**

The mathematical assumptions of the panel regressions in this study are as follows: 1. ~~There are n~~No other important time-varying sub-regional variables that significantly affect both human factors and floods; 2. ~~There are n~~No interactions between human factors and regional or individual characteristics that produce significant spatially heterogeneous effects. The regressions and statistical tests were performed in R (R Core Team, 2019) using packages *lfe* (Gaure, 2019), ~~lmtest (Hothorn et al., 2020), and sandwich (Zeileis et al., 2020).~~

## 2.3 Flood change quantification

~~To examine whether the changes in observed floods can be explained by the changes in human factors, we first detected catchments with significant changes in  $Q$  using Mann-Kendall test (Mann, 1945) and Pettitt's test (Pettitt, 1979), and then After obtaining the coefficients in the response function  $g(\cdot)$ , we can derived the ~~actual-accumulated~~ flood changes attributed to ~~factor the~~ change in factor  $X_s$  in a long period. The regressions in this study calculate a common percentage change in all flood peaks rather than heterogeneous changes of different flood events, according to Eq. (6). Therefore, the percentage change in annual maximum discharges for catchment  $i$  from year  $t_1$  to  $t_2$  can be written as:~~

$$\Delta Q(\%) = \exp(g(X_{i,t_2}) - g(X_{i,t_1})) - 1 \quad (76)$$

To examine ~~how floods changed in catchments that were free from the impacts of the changes in flood peaks unexplained by changing~~ urban areas, cropland areas, and dams, we selected catchments with less than 10% changes in flood peaks due to those factors respectively. Specifically, for a factor  $X$ , we selected catchments with  $|\exp(g(X_{i,t_2}) - 0) - 1| < 10\%$  where  $t_2$  was the most recent year of the data. ~~For the annual maximum discharges of each selected catchment Then, we used applied the Theil-Sen slope estimator (Theil, 1992) to derive the trend and Mann-Kendall test (Mann, 1945) and Pettitt's test (Pettitt, 1979) in those catchments to derive the statistical significance and detect the ones with significant changes in  $Q$ .~~


## 3 Data

### 3.1 Streamflow and precipitation data

Annual maximum instantaneous discharge data in 2739 streamflow gauge stations were obtained from the Ministry of Water Resources in China (<http://www.mwr.gov.cn/english/>). Figure 3-2 shows the outlet locations of all stations. The catchment areas are from 1 km<sup>2</sup> to 1,705,383 km<sup>2</sup> with a median of 1,660 km<sup>2</sup>. Catchment boundaries were extracted using MERIT-Hydro hydrography data (Yamazaki et al., 2019). Differences between extracted catchment areas and reported areas were less than 20% for all catchments. We only used data from 1960 to 2019 because less than 1,000 stations had available data before 1960. Notice that a few stations in the northeast lie outside mainland China. They were not excluded from this study



because all other data were globally available. The 1-km resolution data of Köppen-Geiger climate classes were obtained from Beck et al. (2018). [A 3-hourly and 0.1° precipitation dataset in 1979-2017, the Multi-Source Weighted-Ensemble Precipitation Version 2.2 \(MSWEP V2.2; Beck et al., 2019\), was used.](#)



195

**Figure 32.** National 2739 streamflow stations and the number of stations with available annual maximum discharges each year. The Köppen-Geiger climate classes are obtained from Beck et al. (2018). The boundary lines delineate nine major river basins of China: 1. the Pearl River Basin, 2. the Southeast Basin, 3. the Yangtze River Basin, 4. the Southwest Basin, 5. the Huaihe River Basin, 6. the Yellow River Basin, 7. the Haihe River Basin, 8. the Songliao River Basin, and 9. the Continental Basin.

200

### 3.2 Land cover and dams data

Land cover maps were obtained from the CCI-LC product produced by the European Space Agency (ESA) Climate Change Initiative (CCI). This product provides global yearly 300m-resolution land cover data in 1992-2015 in version 2.0.7 and 2016-2018 in version 2.1.1 (<http://maps.elie.ucl.ac.be/CCI/viewer/download.php>). Urban areas of catchments can be extracted from the maps directly. Cropland areas consist of rain-fed cropland, irrigated or post-flooding cropland, and mosaic cropland.

205

Dam data were available in the Global Reservoir and Dam (GRanD) v1.3 database (Lehner et al., 2011). GRanD collected information about 7,320 global dams in 1948-2017 and recorded 923 dams with storage capacities larger than 10 million m<sup>3</sup> in China. These 923 dams were categorized as large and ~~medium middle~~-dams, according to the Bulletin of the first National Water Conservancy Survey (<http://www.chinawater.com.cn/ztgz/xwzt/2013slpczt/1/>). The total storage capacity of all 4,694 large and ~~medium middle~~-dams is 861,961 million m<sup>3</sup> in China, according to the bulletin. The total storage capacity of all the 923 GRanD dams in China is 670,158 million m<sup>3</sup>, approximately 78% of that recorded in the bulletin. It suggests that the GRanD database is reliable for quantifying the effects of large and ~~medium middle~~-dams on floods while it is unsuitable for considering small dams. For simplicity, we use “dams” to represent large and ~~medium middle~~-dams in the rest of the paper. We obtained the locations, upstream areas, storage capacities, and total annual flows of each dam from the database. The reservoir index can be calculated with the information above.

### 3.3 Catchment selection for regression setup


We selected catchments with ~~suffieient numbers~~ at least 20 years of annual maximum discharges ( $Q$ ) to fit Eq. (1) ~~and (2)~~ in the common period of CCI-LC and GRanD data, i.e., 1992-2017. To avoid inaccurate statistical inference on the regression coefficients due to the correlated model residuals caused by nested catchments, we selected the most upstream catchments with large or medium dams (if possible) among overlapping catchments. We got 757 independent (non-nested) catchments, among which 207 catchments had as least one dam. The statistics of catchment characteristics in those 757 catchments are presented in Table 1 and the spatial distribution of catchments is presented in Fig. 3, as shown in Table 1. For Eq. (1), the CCI-LC data were available in 1992-2018, and thus, we selected 1644 catchments with at least 20 years of  $Q$  data in this period. For Eq. (2), the discharge data were limited before 1960 and the GRanD data were available until 2017, and thus, we selected 1744 catchments with at least 30 years of  $Q$  data in 1960-2017. The number of catchment groups  $k$  in Section 2.2 had no optimal value. In light of the number of selected catchments in the regression models, we set  $k$  to be 10, 20, 30, 40, 50, ~~and 60, 70, and 80~~ to test the robustness of the models. ~~We premised that groups were too small when  $k \geq 70$ , since the average number of catchments in a group was smaller than 25 for Eq. (1) and (2).~~

**Table 1. Summary of catchment characteristics for 757 independent catchments. For each catchment, among its all years with available flood data in 1992-2017, we choose the last year to calculate *Urban*, *Crop*, and *RI*, and choose the first and last year to calculate  $\Delta Urban$ ,  $\Delta Crop$ , and  $\Delta RI$ . The summaries of *RI* and  $\Delta RI$  are calculated based on 207 catchments with at least one large and medium dam. Setup of panel regression models for different factors.**

Factor to be tested	Data period	Required $Q$ length (years)	Number of stations	Preset number $k$ of regions		
<i>Urban &amp; Crop</i>	1992-2018	$\geq 20$	1644	10, 20, 30, 40, 50, 60		
<i>RI</i>	1960-2017	$\geq 30$	1744	10, 20, 30, 40, 50, 60		
Variables	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
<i>Area</i> ( km <sup>2</sup> )	29	499	1096	3341	2763	142372

<u>Urban (%)</u>	<u>0</u>	<u>0.06</u>	<u>0.30</u>	<u>1.52</u>	<u>1.10</u>	<u>65.07</u>
<u>ΔUrban (%)</u>	<u>0</u>	<u>0.05</u>	<u>0.23</u>	<u>1.14</u>	<u>0.85</u>	<u>24.66</u>
<u>Crop (%)</u>	<u>0</u>	<u>10.63</u>	<u>24.71</u>	<u>32.75</u>	<u>48.99</u>	<u>99.58</u>
<u>ΔCrop (%)</u>	<u>-21.58</u>	<u>-0.81</u>	<u>-0.02</u>	<u>0.38</u>	<u>0.87</u>	<u>32.04</u>
<u>RI</u>	<u>0.01</u>	<u>0.09</u>	<u>0.21</u>	<u>0.51</u>	<u>0.61</u>	<u>7.45</u>
<u>ΔRI</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0.17</u>	<u>0.07</u>	<u>7.44</u>

235



**Figure 3. Urban percentages (*Urban*), cropland percentages (*Crop*), and reservoir indexes (*RI*) of 757 independent catchments in their last years with available flood data.**

## 240 4 Results

### 4.1 The sensitivity of floods to human factors

Table 2 shows the optimal regression forms according to AIC and the corresponding coefficient estimations of factor-effect types of factors *Urban*, *Crop* and *RI* on *Q* according to in Eq. (1) and (2). A positive or negative effect means *Q* increases or decreases with the increasing values of the factor respectively. For *Urban* and *RI*, the effect types are consistent for varying values of *k*, with only one exceptional type of effect for *Urban* and *Crop*. *Urban* exhibits a positive and stable effect ( $g(X_{i,t}) = \beta X_{i,t}$  and  $\beta > 0$ ), which means a fixed percentage point increase of *Urban* brings a fixed percentage increase of *Q* no matter how large the initial *Urban* is. *Crop* exhibits no effect. *RI* exhibits a negative and decreasing effect ( $g(X_{i,t}) = \theta X_{i,t}^{1/2}$  and  $\theta < 0$ ), which means a fixed increase of *RI* brings a lower percentage decrease of *Q* with a larger value of initial *RI*. *Crop* exhibits no significant effect. The details about the regression coefficients (e.g., values, standard deviations, and p values) in the effect testing process (Fig. 2) can be seen in Table S1 and Table S2. The maps of catchment groups for all *k* values can also be seen in Fig. AS1 of Appendix A and Fig. S2.

250

255

**Table 2. Effect types on annual maximum discharges ( $Q$ ) of factor urban percentage ( $Urban$ ), cropland percentage ( $Crop$ ) and reservoir index ( $RI$ ).  $k$  is the preset number of catchment groups.**

$k$	$Urban$ effect	$Crop$ effect	$RI$ effect
10	No	No	Negative, decreasing
20	Positive, stable	No	Negative, decreasing
30	Positive, stable	No	Negative, decreasing
40	Positive, stable	Negative, stable	Negative, decreasing
50	Positive, stable	No	Negative, decreasing
60	Positive, stable	No	Negative, decreasing


260

265


Figure 4 shows the percentage change in  $Q$  caused by a 1% point increase of urban area according to Eq. (65). The values of  $\Delta Q(\%)$  are relatively consistent with varying values of  $k$  when the number of catchment groups  $k \geq 30$ . We regard the average value  $\Delta Q = 2.9\%$  for  $k \in \{30, 40, 50, 60\}$  as the sensitivity of  $Q$  to  $Urban$ .  $Crop$  has no significant effect on  $Q$ , therefore we do not calculate the corresponding sensitivity. Figure 5 shows the percentage change in  $Q$  caused by a 1 unit increase of  $RI$  according to Eq. (65). The relationship between  $\Delta Q$  and  $RI$  has little change when  $k \geq 20$  with varying values of  $k$ . Therefore In summary, the the average values of coefficient  $\theta$  for all six  $k$  values are considered to calculate the sensitivity of  $Q$  to  $RI$  method is robust to  $k$ , and thus, we regard the model with  $k = 50$  as the main model in the remaining part of the study. The sensitivity of  $Q$  to  $Urban$  is  $\Delta Q = 3.9\%$  with the 95% confidence interval  $CI = [1.9\%, 5.7\%]$  when  $\Delta Urban = 1\%$ . In addition, the absolute values of  $\Delta Q$  decrease with increasing initial values of  $RI$ . For initial  $RI = 0$ ,  $\Delta Q = -213.21\%$  with  $CI = [-29.9\%, -11.4\%]$  when  $RI$  increases by 1; for initial  $RI = 35$ ,  $\Delta Q = -6.25.4\%$  with  $CI = [-9.1\%, -3.2\%]$  when  $RI$  increases by 1.

**Table 2. Optimal regression forms (with the lowest AIC) and the corresponding coefficient estimations of factor urban percentage (*Urban*), cropland percentage (*Crop*) and reservoir index (*RI*) in Eq. (1). \* denotes that the coefficient does not equal to 0 at a 0.05 significance level using bootstrapping inference. *k* is the preset number of catchment groups.**

	<i>k</i>	Optimal $g(\cdot)$	2.5% Qu.	Mean	97.5% Qu.
<i>Urban effect</i>	<u>10</u>	$\beta Urban_{i,t}^*$	<u>1.20E-02</u>	<u>2.72E-02</u>	<u>4.21E-02</u>
	<u>20</u>	$\beta Urban_{i,t}^*$	<u>2.31E-02</u>	<u>3.89E-02</u>	<u>5.59E-02</u>
	<u>30</u>	$\beta Urban_{i,t}^*$	<u>9.35E-03</u>	<u>2.65E-02</u>	<u>4.40E-02</u>
	<u>40</u>	$\beta Urban_{i,t}^*$	<u>1.59E-02</u>	<u>3.64E-02</u>	<u>5.68E-02</u>
	<u>50</u>	$\beta Urban_{i,t}^*$	<u>1.88E-02</u>	<u>3.71E-02</u>	<u>5.57E-02</u>
	<u>60</u>	$\beta Urban_{i,t}^*$	<u>1.08E-02</u>	<u>3.10E-02</u>	<u>5.03E-02</u>
	<u>70</u>	$\beta Urban_{i,t}^*$	<u>1.32E-02</u>	<u>3.18E-02</u>	<u>5.05E-02</u>
	<u>80</u>	$\beta Urban_{i,t}^*$	<u>1.22E-02</u>	<u>3.23E-02</u>	<u>5.27E-02</u>
<i>Crop effect</i>	<u>10</u>	$\theta Crop_{i,t}^{1/2}$	<u>-6.14E-02</u>	<u>1.43E-02</u>	<u>8.87E-02</u>
	<u>20</u>	$\theta Crop_{i,t}^{1/2}$	<u>-4.59E-02</u>	<u>2.95E-02</u>	<u>1.01E-01</u>
	<u>30</u>	$\gamma Crop_{i,t}^2$	<u>-1.41E-04</u>	<u>-5.36E-05</u>	<u>3.42E-05</u>
	<u>40</u>	$\gamma Crop_{i,t}^2$	<u>-1.36E-04</u>	<u>-3.81E-05</u>	<u>5.61E-05</u>
	<u>50</u>	$\theta Crop_{i,t}^{1/2}$	<u>-5.21E-02</u>	<u>2.49E-02</u>	<u>1.09E-01</u>
	<u>60</u>	$\gamma Crop_{i,t}^2$	<u>-1.66E-04</u>	<u>-7.54E-05</u>	<u>2.77E-05</u>
	<u>70</u>	$\gamma Crop_{i,t}^2$	<u>-1.62E-04</u>	<u>-6.71E-05</u>	<u>2.85E-05</u>
	<u>80</u>	$\gamma Crop_{i,t}^2$	<u>-1.71E-04</u>	<u>-7.21E-05</u>	<u>2.74E-05</u>
<i>RI effect</i>	<u>10</u>	$\theta RI_{i,t}^{1/2} *$	<u>-2.90E-01</u>	<u>-1.74E-01</u>	<u>-5.84E-02</u>
	<u>20</u>	$\theta RI_{i,t}^{1/2} *$	<u>-3.43E-01</u>	<u>-2.11E-01</u>	<u>-1.01E-01</u>
	<u>30</u>	$\theta RI_{i,t}^{1/2} *$	<u>-3.48E-01</u>	<u>-2.20E-01</u>	<u>-1.08E-01</u>
	<u>40</u>	$\theta RI_{i,t}^{1/2} *$	<u>-3.64E-01</u>	<u>-2.45E-01</u>	<u>-1.25E-01</u>
	<u>50</u>	$\theta RI_{i,t}^{1/2} *$	<u>-3.55E-01</u>	<u>-2.39E-01</u>	<u>-1.21E-01</u>
	<u>60</u>	$\theta RI_{i,t}^{1/2} *$	<u>-3.85E-01</u>	<u>-2.55E-01</u>	<u>-1.30E-01</u>
	<u>70</u>	$\theta RI_{i,t}^{1/2} *$	<u>-3.80E-01</u>	<u>-2.38E-01</u>	<u>-1.14E-01</u>
	<u>80</u>	$\theta RI_{i,t}^{1/2} *$	<u>-3.88E-01</u>	<u>-2.36E-01</u>	<u>-9.03E-02</u>



275 Figure 4. Percentage change in annual maximum discharge ( $Q$ ) caused by a 1% point increase of urban area (*Urban*) based on different numbers of catchment groups ( $k$ ). The error bars are 95% confidence intervals.



280


Figure 5. (a) Percentage change in annual maximum discharge ( $Q$ ) caused by a 1 unit increase of reservoir index ( $RI$ ) from different initial  $RI$  values based on different numbers of catchment groups ( $k$ ). (b) The same as (a) but with the shaded areas are 95% confidence intervals shown by shaded areas.

## 285 4.2 National flood changes and attributions

Figure 6 shows the changes in  $Q$  due to the changes in Urban from 1992 to 2017 in for 2739-1625 catchments with at least 20 years of data in 1992-2017, according to Eq. (76). Increasing Urban causes increases in  $Q$  of more than 10% of increases in  $Q$  attributed to increasing Urban occur in 284-184 (11.310.4% of 16252739) catchments, which are mainly located in the North Plain of China, especially in the Huaihe River Basin and the middle and down streams of the Haihe

290 River Basin. Among these ~~284-184~~ catchments, increasing *Urban* causes increases in *Q* of more than 25% in 71 (4.4% of  
1625) catchments-65 (2.4% of 2739) have more than 25% increases of *Q* attributed to increasing *Urban*. In 61 catchments  
with significant increases in observed *Q*, increasing *Urban* causes increases in *Q* of more than 10% in only 5 catchments,  
which means urbanization is not a predominant driver of flood changes in 1992-2017 in China. The changes in *Q* due to the  
changes in *Crop* cannot be effectively quantified because *Crop* has no statistically significant effects on *Q* according to the  
295 results in Section 4.1.










Figure 6. Accumulated Changes-increases in annual maximum discharges ( $Q$ ) due to the change-increases in urban areas (*Urban*) from 1992 to 2017 infor 2739-1625 catchments with at least 20 years of flood data in 1992-2017, according to Eq. (6). Large dots represent catchments with significant increases in observed  $Q$  if  $p < 0.05$  for any one of the Mann-Kendall test and the Pettitt's test. The boundary lines delineate nine major river basins of China.

Figure 7 shows the changes in  $Q$  due to the changes in  $RI$  from 1960 to 2017for in 777-536out of 2739 catchments with at least one dam and at least 30 years of data in 1960-2017, according to Eq. (7). In 415-196 (53-436.6% of 777536) catchments, increasing  $RI$  leads to more than 10% decreases in  $Q$ . It indicates that flood peaks are likely to decrease severely if dams are built in the catchment. Among these 415-196 catchments, increasing  $RI$  leads to more than 25% decreases in  $Q$  in 68-28 (8-85.2% of 777536) catchments have more than 25% decreases of  $Q$  attributed to increasing  $RI$ . Spatially, the impacts of dams on floods are larger in northern basins (the Huaihe River Basin, the Haihe River Basin, the Yellow River Basin, and the Songhua and Liaohe River Basin) than that in southern basins (the Yangtze River Basin, the Southeast River

Basin, the Southwest River Basin, and the Pearl River Basin). In the northern basins, increasing  $RI$  leads to more than 10% and 25% decreases  $\phi$ -in  $Q$  in ~~66.147.8%~~ 66.147.8% and ~~20.010.0%~~ 20.010.0% catchments, respectively. By comparison, in southern basins, increasing  $RI$  leads to more than 10% and 25% decreases  $\phi$ -in  $Q$  in only ~~45.426.3%~~ 45.426.3% and ~~2.00.7%~~ 2.00.7% catchments, respectively. In 234 catchments with significant decreases in observed  $Q$ , increasing  $RI$  leads to more than 10% decreases in  $Q$  in 138 (59.0% of 234) catchments, which means dam construction is a predominant driver of the decreases in flood magnitudes for catchments with dams in 1960-2017.

315






320 **Figure 7. Accumulated Changes-decreases in annual maximum discharges ( $Q$ ) due to the change-increases in reservoir index ( $RI$ ) for 536 catchments with at least 30 years of flood data and at least one dam in 1960-2017, according to Eq. (6). Large dots represent catchments with significant decreases in observed  $Q$  if  $p < 0.05$  for any one of the Mann-Kendall test and the Pettitt's test, from 1960 to 2017 in 777 out of 2739 catchments with at least one dam. The boundary lines delineate nine major river basins of China.**

325

Figure 8 shows the trends-change directions of  $Q$  during 1960-2017 in 1074-1249 catchments with at least 40-30 years of data, Urban  $< 3.42.6\%$  and RI  $< 0.196$  in 2017. These catchments were selected based on the thresholds that ensure each factor leads to no more than 10% changes in  $Q$ , as stated in Section 2.3. In these catchments, the changes in  $Q$  are unexplained-by-free from the changes-inimpacts of Urban, Crop, and RI. Significant increases in  $Q$  occur in 85 (6.9% of 1249) catchments. Significant decreases in  $Q$  occur in 403 (32.3% of 1249) More than 10% decreasing rates per decade are detected in 210 (19.6% of 1074) catchments, among which 131-188 (46.762.4% of 210/403) are located in the middle and

330

~~down streams of~~ the Yellow River Basin (mainly in the middle and down streams) and ~~the upper streams of~~ the Haihe River Basin (mainly in the upper streams). Such regional coherence of similar trends cannot be found in other regions.






Figure 8. Trends Change directions of annual maximum discharges ( $Q$ ) during 1960-2017 in the for 1074-1249 catchments with at least 40-30 years of flood data,  $Urban < 3.42. 6\%$  and  $RI < 0.196$  in 2017. These catchments are considered to be free from the impacts of urbanization and dam constructions. The change is significant if  $p < 0.05$  for any one of the Mann-Kendall test and Pettitt's test. have less than 10% changes in  $Q$  attributed to Urban or  $RI$ . Catchments with statistically insignificant trends are labeled as "no trend". The boundary lines delineate nine major river basins of China.

## 5 Discussion

### 5.1 Strengths and limitations of panel regressions

345 We use panel regressions to derive the causal effects of urban areas, cropland areas, and dams on annual maximum discharges across mainland China. In this study, the panel regressions exhibit the following strengths. 1. We obtain a nationally generalizable sensitivity of floods to each human factor. This sensitivity helps understand the overall added risks



of specific human activity on floods on a national scale. In addition, with quantitative sensitivity, scientists are able to select catchments with limited impacts of dams and land cover changes for studying the effects of climate change, e.g., Blöschl et al. (2019). 2. Compared with previous studies using panel regressions in hydrology (Ferreira and Ghimire, 2012; Steinschneider et al., 2013; McManamay et al., 2014; Bassiouni et al., 2016; Levy et al., 2018; Blum et al., 2020; Davenport et al., 2020), we take a further step by considering multiple types of human impacts simultaneously and distinguishing their increasing or decreasing effects. Blum et al. (2020) and Davenport et al. (2020) considered non-linear forms of response functions for the targeted factors, but they did not distinguish increasing and decreasing effects. These improvements provide a more comprehensive understanding of human impacts on floods.

The limitations are as follows. 1. The assumptions in the regressions are difficult to test. As stated in Section 2, we assume (i) no more important time-varying sub-regional confounders and (ii) no interaction terms between human factors and regional or individual characteristics that produce significant spatially heterogeneous effects. These assumptions may be violated in some cases. For example, the effect of urbanization on floods may be larger in regions with higher soil permeability, which means spatially heterogeneous effects may be nonnegligible. Testing these assumptions requires detailed information about catchment characteristics such as topography and geology. Moreover, adding too many variables into the regressions will decrease model interpretability. 2. The method cannot distinguish the heterogeneous effects of human factors on different floods. As stated in Section 2.3, the method derives a common percentage change in all flood peaks given changing human factors, which means no changes in coefficients of variation. However, practically, the variability of floods may change by human activities. For example, reservoirs tend to regulate extreme floods but omit small floods. 3. This study does not comprehensively assess the effects of total human impacts on floods. We omit many other human factors due to the lack of data. For example, the data about water diversion, irrigation, channelization, and afforestation on a national scale are currently not available to the public.

## 5.2 Consistency with knowledge and other large-sample studies

We detect a stable positive effect of urban areas on floods. In theory, expanding urban areas magnify floods in two major ways. Firstly, natural soil grounds are replaced by impervious surfaces, which lead to more rainfall water appearing on the surface rather than infiltrating into the soil (Villarini and Slater, 2017). Second, urban areas have smooth surfaces, where floods propagate faster and become more flashy (Mogollón et al., 2016). This study finds a 2-93.9% (with CI = [1.9%, 5.7%]) increase in annual maximum discharges given a 1% point increase in urban areas. This finding accords with the result from a national investigation in the US (Blum et al., 2020), which reported a 3.3% (with CI = [1.9%, 4.7%]) increase in annual maximum discharges based on panel regressions.

The cropland areas impose no significant impacts on floods according to our results. Theoretically, expanding cropland areas affect floods in many ways. For example, during agricultural practices, soil depths may decrease due to erosion while increase due to soil compaction (Rogger et al., 2017). Cropland may also bring artificial drainages that lower groundwater tables (Rogger et al., 2017). Some effects may be offset by others, which masks the relationship between cropland areas and

floods. Similar to our result, Bertola et al. (2019) found that agricultural land-use intensification rarely caused flood changes in 95 catchments of Austria using covariate-based non-stationary flood probability distributions. To our knowledge, large-sample studies are limited on the relationship between cropland and floods. Therefore, more detailed in-site investigations are required to uncover the causal chain from cropland changes to flood changes.

385 This study suggests that dams have a negative decreasing effect on floods. Generally, dams buffer water during floods and thus decrease flood peaks. More dams may not necessarily decrease floods at a constant rate because existing dams with sufficient storage capacities are already capable to control floods. This effect was confirmed by Wang et al. (2017), who used detailed conceptual models of reservoir regulation and found that the mean annual floods had a slowing decrease with increasing reservoirs. In a large-sample study on 4859 catchments in the US (FitzHugh et al., 2011), median annual 1-day  
390 maximum flows were estimated to decrease by more than 20% when the storage ratios, i.e., the total storage capacity of upstream dams divided by average annual runoff, were larger than 1. If a dam with storage capacity equaling the annual runoff is established at the outlet of the catchment without any dam before, both the reservoir index defined in this study and the storage ratio defined by FitzHugh et al. (2011) increase from 0 to 1. In this special case, the annual maximum discharges  
| ~~decrease-change~~ by -21.2% (with  $CI = [-29.9\%, -11.4\%]$ ) ~~23.1%~~ in this study, comparable to the 20% decrease from  
395 FitzHugh et al. (2011). It is noteworthy that this study only focuses on the effects of human factors on annual maximum discharges. Generally, the effects are larger for less frequent floods. Zhao et al. (2020) investigated floods in 1403 catchments in the US and found a decrease of 100-year floods by more than 60% in 47% of catchments with a dam upstream.

### 5.3 Insights toward a national investigation of flood changes

This study takes the first step to explain flood changes quantitatively on a national scale in China. In this study, urbanization  
400 and dam constructions significantly change annual maximum discharges in the middle and down streams of the Yellow River Basin and the Haihe River Basin, where step changes were detected by Yang et al. (2019). As a major human residence with a high density of population, the North Plain of China experiences fast urbanization in recent years (Du et al., 2018), which brings larger flood risks on lives and properties. In addition, the degree of dam regulation is larger in northern China because the annual runoff is much smaller than that in wet southern China. In this study, after removing the  
405 catchments with nonnegligible impacts of urbanization and dams, ~~the-unexplained decreasing~~ annual maximum discharges  
~~trends occur-show spatial coherence~~ in the middle and down streams of the Yellow River Basin and the upper streams of the Haihe River Basin, where decreasing trends were also derived by Yang et al. (2019). Yang et al. (2019) interpreted these trends as the results of soil conservation practices (Bai et al., 2016) and decreasing extreme rainfall (Yang et al., 2013; Wu et al., 2016). Besides, other reasons include decreasing soil moisture (Cheng et al., 2015; Yang et al., 2020a) and the impacts of  
410 cascade small soil-retaining dams (Yang et al., 2020b). It indicates that the impact factors of floods are complex in this region and further studies are required.

## 6 Conclusions

We conducted a data-based analysis on the causal effects of human impacts on floods using a panel regression on a national scale, based on annual maximum discharges ( $Q$ ) from ~~2739-757 non-nested stations-catchments~~ in China, CCI-LC land cover data, and GRanD dam data. ~~Furthermore~~Specifically, we derived nationally generalizable information about the sensitivity of  $Q$  to human factors, namely the changes in urban areas, cropland areas, and reservoir indexes for large and medium middle-dams. ~~Furthermore, using a dataset of 2739 streamflow stations, and then we~~ determined the explained and unexplained changes in floods by the human factors on a national scale based on the sensitivity of  $Q$  to human factors. The major findings are as follows.

— Floods are sensitive to the changes in urban areas and dams.

- Urban areas have a positive and stable effect on floods, i.e., a 1% point increase in urban areas causes a 2.93.9% increase in annual maximum discharges with a confidence interval  $CI = [1.9\%, 5.7\%]$ . Cropland areas have no significant effect on  $Q$ . Reservoir index has a negative and decreasing effect on  $Q$ , i.e., the decrease of  $Q$  caused by a 1 unit increase of reservoir indexes ranges from 21.4% (with  $CI = [11.4\%, 29.9\%]$ )~~23.1%~~ to 5.4%6.2% (with  $CI = [3.2\%, 9.1\%]$ ) corresponding to initial reservoir indexes from 0 to ~~53~~.

— Urbanization is not a predominant driver of the increases in flood magnitudes on a national scale. I

- From 1992 to 2017n 1992-2017, increasing urban areas cause increases in  $Q$  of more than 10% in 184 (11.3%) of 1625 catchments~~more than 10% of increases in  $Q$  were caused by increasing urban areas in 10.4% of the 2739 catchments.~~


These catchments are mainly located in the North Plain of China, especially in the Huaihe River Basin and the middle and down streams of the Haihe River Basin. However, among these 184 catchments, only 5 of them have significant increases in observed  $Q$ .

- Dam construction is a predominant driver of the decreases in flood magnitudes on a national scale. A~~From 1960 to 2017,~~ among the ~~777-536~~ catchments with at least one dam in 1960-2017, 53.4% have more than 10% decreases in  $Q$  caused by increasing reservoir indexes cause decreases in  $Q$  of more than 10% in 196 (36.6%) catchments. - Spatially, the impacts of dams on floods are larger in northern basins, including the Huaihe River Basin, the Haihe River Basin, the Yellow River Basin, and the Songhua and Liaohe River Basin, ~~where 66.1% catchments have more than 10% decreases of  $Q$  attributed to increasing reservoir indexes.~~ There are 138 of those 196 catchments having significant decreases in observed  $Q$ , accounting for 59.0% of the total 234 catchments with significant decreases in observed  $Q$ .

- Unexplained decreases in flood magnitudes show spatial coherence in the middle and down streams of the Yellow River Basin and the upper streams of the Haihe River Basin. Among ~~4074-1249~~ catchments with less than 10% changes in  $Q$  caused by urban areas or dams, ~~210-403 (49.632.3%)~~ catchments have significant decreases in  $Q$ ~~a more than 10% decrease per decade~~ during 1960-2017, and ~~62.446.7%~~ of the ~~210-403~~ catchments are located in ~~the middle and down streams of the Yellow River Basin~~ and the upper streams of the Haihe River Basin.

445 This study extends the panel regression method to quantify the effects of multiple human factors on floods, which helps understand the causes of flood changes on a national scale in China. Future studies may collect more data to consider more human factors and quantify the effects on different return periods of floods.

### Appendix A



**Figure A1. Catchment groups for all  $k$  values in Eq. (1). The points are the geometric centers of catchments.**

### 450 **Data availability**

Annual maximum discharge data are obtained from the Water Resources Information Center of the Ministry of Water Resources in China (<http://www.mwr.gov.cn/english/>). CCI-LC data are obtained from the ESA Climate Change Initiative -

Land Cover project 2017 (<http://maps.elie.ucl.ac.be/CCI/viewer/download.php>). GRanD data are obtained from Global Dam Watch (<http://globaldamwatch.org>). [Köppen-Geiger climate class data and MSWEP V2.2 precipitation data are obtained from GLOH2O \(http://www.gloh2o.org/\).](#)

### Author contribution

Wencong Yang: Conceptualization, Methodology, Software, Visualization, Writing – original draft preparation.

Hanbo Yang: Conceptualization, Data curation, Funding acquisition, Supervision, Writing – review & editing.

Dawen Yang: Data curation, Funding acquisition.

460 Aizhong Hou: Data curation.

### Competing interests

The authors declare that they have no conflict of interest.

### Financial support

This research was partially supported by funding from the National Natural Science Foundation of China (Grant Nos. 465 51979140 and 41661144031), the Scientific Research Project from the China Three Gorges Corporation (Grant No. 202003098), and the National Program for Support of Top-notch Young Professionals.

### References

- Bai, P., Liu, X. M., Liang, K., and Liu, C. M.: Investigation of changes in the annual maximum flood in the Yellow River basin, China, *Quatern Int*, 392, 168-177, 10.1016/j.quaint.2015.04.053, 2016.
- 470 Bassiouni, M., Vogel, R. M., and Archfield, S. A.: Panel regressions to estimate low-flow response to rainfall variability in ungaged basins, *Water Resour Res*, 52, 9470-9494, 10.1002/2016wr018718, 2016.
- [Beck, H. E., Wood, E. F., Pan, M., Fisher, C. K., Miralles, D. G., van Dijk, A. I. J. M., McVicar, T. R., and Adler, R. F.: MSWEP V2 Global 3-Hourly 0.1 degrees Precipitation: Methodology and Quantitative Assessment, \*B Am Meteorol Soc\*, 100, 473-502, 10.1175/Bams-D-17-0138.1, 2019.](#)
- 475 Beck, H. E., Zimmermann, N. E., McVicar, T. R., Vergopolan, N., Berg, A., and Wood, E. F.: Present and future Köppen-Geiger climate classification maps at 1-km resolution, *Sci Data*, 5, ARTN 180214, 10.1038/sdata.2018.214, 2018.
- Bertola, M., Viglione, A., and Blöschl, G.: Informed attribution of flood changes to decadal variation of atmospheric, catchment and river drivers in Upper Austria, *J Hydrol*, 577, ARTN 123919, 10.1016/j.jhydrol.2019.123919, 2019.

- 480 Bloeschl, G., Hall, J., Viglione, A., Perdigao, R. A. P., Parajka, J., Merz, B., Lun, D., Arheimer, B., Aronica, G. T., Bilibashi,  
A., Bohac, M., Bonacci, O., Borga, M., Canjevac, I., Castellarin, A., Chirico, G. B., Claps, P., Frolova, N., Ganora, D.,  
Gorbachova, L., Gul, A., Hannaford, J., Harrigan, S., Kireeva, M., Kiss, A., Kjeldsen, T. R., Kohnova, S., Koskela, J. J.,  
Ledvinka, O., Macdonald, N., Mavrova-Guirguinova, M., Mediero, L., Merz, R., Molnar, P., Montanari, A., Murphy, C.,  
Osuch, M., Ovcharuk, V., Radevski, I., Salinas, J. L., Sauquet, E., Sraj, M., Szolgay, J., Volpi, E., Wilson, D., Zaimi, K.,  
485 and Zivkovic, N.: Changing climate both increases and decreases European river floods, *Nature*, 573, 108-+,  
10.1038/s41586-019-1495-6, 2019.
- Bloschl, G., Gaal, L., Hall, J., Kiss, A., Komma, J., Nester, T., Parajka, J., Perdigao, R. A. P., Plavcova, L., Rogger, M.,  
Salinas, J. L., and Viglione, A.: Increasing river floods: fiction or reality?, *Wires Water*, 2, 329-344, 10.1002/wat2.1079,  
2015.
- Blum, A. G., Ferraro, P. J., Archfield, S. A., and Ryberg, K. R.: Causal Effect of Impervious Cover on Annual Flood  
490 Magnitude for the United States, *Geophys Res Lett*, 47, ARTN e2019GL086480, 10.1029/2019GL086480, 2020.
- Cheng, S. J., Guan, X. D., Huang, J. P., Ji, F., and Guo, R. X.: Long-term trend and variability of soil moisture over East  
Asia, *J Geophys Res-Atmos*, 120, 8658-8670, 10.1002/2015jd023206, 2015.
- Davenport, F. V., Herrera-Estrada, J. E., Burke, M., and Diffenbaugh, N. S.: Flood Size Increases Nonlinearly Across the  
Western United States in Response to Lower Snow-Precipitation Ratios, *Water Resour Res*, 56, ARTN  
495 e2019WR025571, 10.1029/2019WR025571, 2020.
- De Niel, J., and Willems, P.: Climate or land cover variations: what is driving observed changes in river peak flows? A data-  
based attribution study, *Hydrol Earth Syst Sc*, 23, 871-882, 10.5194/hess-23-871-2019, 2019.
- Di Baldassarre, G., Sivapalan, M., Rusca, M., Cudenneq, C., Garcia, M., Kreibich, H., Konar, M., Mondino, E., Mard, J.,  
Pande, S., Sanderson, M. R., Tian, F. Q., Viglione, A., Wei, J., Wei, Y. P., Yu, D. J., Srinivasan, V., and Bloschl, G.:  
500 Sociohydrology: Scientific Challenges in Addressing the Sustainable Development Goals, *Water Resour Res*, 55, 6327-  
6355, 10.1029/2018wr023901, 2019.
- Du, S. Q., He, C. Y., Huang, Q. X., and Shi, P. J.: How did the urban land in floodplains distribute and expand in China from  
1992-2015?, *Environ Res Lett*, 13, ARTN 034018, 10.1088/1748-9326/aaac07, 2018.
- Du, S. Q., Cheng, X. T., Huang, Q. X., Chen, R. S., Ward, P. J., and Aerts, J. C. J. H.: Brief communication: Rethinking the  
505 1998 China floods to prepare for a nonstationary future, *Nat Hazard Earth Sys*, 19, 715-719, 10.5194/nhess-19-715-  
2019, 2019.
- Ferreira, S., and Ghimire, R.: Forest cover, socioeconomics, and reported flood frequency in developing countries, *Water  
Resour Res*, 48, Artn W08529, 10.1029/2011wr011701, 2012.
- FitzHugh, T. W., and Vogel, R. M.: The impact of dams on flood flows in the United States, *River Res Appl*, 27, 1192-1215,  
510 10.1002/rra.1417, 2011.
- Gaure, S.: lfe: Linear group fixed effects, available at: <https://cran.r-project.org/web/packages/lfe/index.html>, R package  
version 2.8-5.1, 2019.

- Hodgkins, G. A., Dudley, R. W., Archfield, S. A., and Renard, B.: Effects of climate, regulation, and urbanization on historical flood trends in the United States, *J Hydrol*, 573, 697-709, 10.1016/j.jhydrol.2019.03.102, 2019.
- 515 ~~Hothorn, T., Zeileis, A., Farebrother, R., Cummins, C., Millo, G., and Mitchell, D.: *lmtest: Testing Linear Regression Models*, available at: <https://cran.r-project.org/web/packages/lmtest/index.html>, R package version 0.9-38, 2020.~~ Jehn, F. U., Bestian, K., Breuer, L., Kraft, P., and Houska, T.: *Using hydrological and climatic catchment clusters to explore drivers of catchment behavior*, *Hydrol Earth Syst Sc*, 24, 1081-1100, 10.5194/hess-24-1081-2020, 2020.
- Kundzewicz, Z. W., Su, B. D., Wang, Y. J., Xia, J., Huang, J. L., and Jiang, T.: Flood risk and its reduction in China, *Adv Water Resour*, 130, 37-45, 10.1016/j.advwatres.2019.05.020, 2019.
- 520 Lehner, B., Liermann, C. R., Revenga, C., Vorosmarty, C., Fekete, B., Crouzet, P., Doll, P., Endejan, M., Frenken, K., Magome, J., Nilsson, C., Robertson, J. C., Rodel, R., Sindorf, N., and Wisser, D.: High-resolution mapping of the world's reservoirs and dams for sustainable river-flow management, *Front Ecol Environ*, 9, 494-502, 10.1890/100125, 2011.
- 525 Levy, M. C., Lopes, A. V., Cohn, A., Larsen, L. G., and Thompson, S. E.: Land Use Change Increases Streamflow Across the Arc of Deforestation in Brazil, *Geophys Res Lett*, 45, 3520-3530, 10.1002/2017gl076526, 2018.
- Lu, W. W., Lei, H. M., Yang, D. W., Tang, L. H., and Miao, Q. H.: Quantifying the impacts of small dam construction on hydrological alterations in the Jiulong River basin of Southeast China, *J Hydrol*, 567, 382-392, 10.1016/j.jhydrol.2018.10.034, 2018.
- 530 Mann, H. B.: Nonparametric Tests Against Trend, *Econometrica*, 13, 10.2307/1907187, 1945.
- McManamay, R. A.: Quantifying and generalizing hydrologic responses to dam regulation using a statistical modeling approach, *J Hydrol*, 519, 1278-1296, 10.1016/j.jhydrol.2014.08.053, 2014.
- 535 Merz, B., Vorogushyn, S., Uhlemann, S., Delgado, J., and Hundscha, Y.: HESS Opinions 'More efforts and scientific rigour are needed to attribute trends in flood time series', *Hydrol Earth Syst Sc*, 16, 1379-1387, 10.5194/hess-16-1379-2012, 2012.
- Merz, R., Tarasova, L., and Basso, S.: The flood cooking book: ingredients and regional flavors of floods across Germany, *Environ Res Lett*, 15, ARTN 114024, 10.1088/1748-9326/abb9dd, 2020.
- Mogollon, B., Frimpong, E. A., Hoegh, A. B., and Angermeier, P. L.: Recent Changes in Stream Flashiness and Flooding, and Effects of Flood Management in North Carolina and Virginia, *J Am Water Resour As*, 52, 561-577, 10.1111/1752-1688.12408, 2016.
- 540 Muller, M. F., and Levy, M. C.: Complementary Vantage Points: Integrating Hydrology and Economics for Sociohydrologic Knowledge Generation, *Water Resour Res*, 55, 2549-2571, 10.1029/2019wr024786, 2019.
- Pande, S., and Sivapalan, M.: Progress in socio-hydrology: a meta-analysis of challenges and opportunities, *Wires Water*, 4, ARTN e1193, 10.1002/wat2.1193, 2017.
- 545 Pearl, J., and Mackenzie, D.: *The book of why: the new science of cause and effect*, Basic Books, New York, 2020.

- Pena, L. E., Barrios, M., and Frances, F.: Flood quantiles scaling with upper soil hydraulic properties for different land uses at catchment scale, *J Hydrol*, 541, 1258-1272, 10.1016/j.jhydrol.2016.08.031, 2016.
- ~~Pettitt, A. N.: A Non-Parametric Approach to the Change-Point Problem, *Applied Statistics*, 28, 10.2307/2346729, 1979.~~
- 550 Prosdocimi, I., Kjeldsen, T. R., and Miller, J. D.: Detection and attribution of urbanization effect on flood extremes using nonstationary flood-frequency models, *Water Resour Res*, 51, 4244-4262, 10.1002/2015wr017065, 2015.
- Reynolds, A. P., Richards, G., de la Iglesia, B., and Rayward-Smith, V. J.: Clustering Rules: A Comparison of Partitioning and Hierarchical Clustering Algorithms, *Journal of Mathematical Modelling and Algorithms*, 5, 475-504, 10.1007/s10852-005-9022-1, 2006.
- R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, 555 Austria, available at: <http://www.R-project.org>, 2019.
- Rogger, M., Agnoletti, M., Alaoui, A., Bathurst, J. C., Bodner, G., Borga, M., Chaplot, V., Gallart, F., Glatzel, G., Hall, J., Holden, J., Holko, L., Horn, R., Kiss, A., Kohnova, S., Leitinger, G., Lennartz, B., Parajka, J., Perdigao, R., Peth, S., Plavcova, L., Quinton, J. N., Robinson, M., Salinas, J. L., Santoro, A., Szolgay, J., Tron, S., van den Akker, J. J. H., Viglione, A., and Bloschl, G.: Land use change impacts on floods at the catchment scale: Challenges and opportunities 560 for future research, *Water Resour Res*, 53, 5209-5219, 10.1002/2017wr020723, 2017.
- Runge, J., Bathiany, S., Bollt, E., Camps-Valls, G., Coumou, D., Deyle, E., Glymour, C., Kretschmer, M., Mahecha, M. D., Munoz-Mari, J., van Nes, E. H., Peters, J., Quax, R., Reichstein, M., Scheffer, M., Scholkopf, B., Spirtes, P., Sugihara, G., Sun, J., Zhang, K., and Zscheischler, J.: Inferring causation from time series in Earth system sciences, *Nat Commun*, 10, ARTN 2553, 10.1038/s41467-019-10105-3, 2019.
- 565 Stein, L., Pianosi, F., and Woods, R.: Event-based classification for global study of river flood generating processes, *Hydrol Process*, 34, 1514-1529, 10.1002/hyp.13678, 2020.
- Steinschneider, S., Yang, Y. C. E., and Brown, C.: Panel regression techniques for identifying impacts of anthropogenic landscape change on hydrologic response, *Water Resour Res*, 49, 7874-7886, 10.1002/2013wr013818, 2013.
- ~~Theil, H.: A Rank Invariant Method of Linear and Polynomial Regression Analysis, in: *Henri Theil's Contributions to Economics and Econometrics, Advanced Studies in Theoretical and Applied Econometrics*, 345-381, 1992.~~
- 570 Umer, Y. M., Jetten, V. G., and Ettema, J.: Sensitivity of flood dynamics to different soil information sources in urbanized areas, *J Hydrol*, 577, ARTN 123945, 10.1016/j.jhydrol.2019.123945, 2019.
- Viglione, A., Merz, B., Dung, N. V., Parajka, J., Nester, T., and Bloschl, G.: Attribution of regional flood changes based on scaling fingerprints, *Water Resour Res*, 52, 5322-5340, 10.1002/2016wr019036, 2016.
- 575 Villarini, G., and Slater, L.: Climatology of Flooding in the United States, in: *Oxford Research Encyclopedia of Natural Hazard Science*, 2017.
- Wang, W., Li, H. Y., Leung, L. R., Yigzaw, W., Zhao, J. S., Lu, H., Deng, Z. Q., Demisie, Y., and Bloschl, G.: Nonlinear Filtering Effects of Reservoirs on Flood Frequency Curves at the Regional Scale, *Water Resour Res*, 53, 8277-8292, 10.1002/2017wr020871, 2017.



- 580 Wooldridge, J. M.: Introductory econometrics a modern approach [Hauptband] [Hauptband], 2016.
- Wu, Y. J., Wu, S. Y., Wen, J. H., Xu, M., and Tan, J. G.: Changing characteristics of precipitation in China during 1960-2012, *Int J Climatol*, 36, 1387-1402, 10.1002/joc.4432, 2016.
- Yamazaki, D., Ikeshima, D., Sosa, J., Bates, P. D., Allen, G. H., and Pavelsky, T. M.: MERIT Hydro: A High-Resolution Global Hydrography Map Based on Latest Topography Dataset, *Water Resour Res*, 55, 5053-5073, 10.1029/2019wr024873, 2019.
- 585 Yang, L., Villarini, G., Smith, J. A., Tian, F. Q., and Hu, H. P.: Changes in seasonal maximum daily precipitation in China over the period 1961-2006, *Int J Climatol*, 33, 1646-1657, 10.1002/joc.3539, 2013.
- Yang, L., Wang, L. C., Li, X., and Gao, J.: On the flood peak distributions over China, *Hydrol Earth Syst Sc*, 23, 5133-5149, 10.5194/hess-23-5133-2019, 2019.
- 590 Yang, W. C., Yang, H. B., and Yang, D. W.: Classifying floods by quantifying driver contributions in the Eastern Monsoon Region of China, *J Hydrol*, 585, ARTN 124767, 10.1016/j.jhydrol.2020.124767, 2020a.
- Yang, X. N., Sun, W. Y., Mu, X. M., Gao, P., and Zhao, G. J.: Run-off affected by climate and anthropogenic changes in a large semi-arid river basin, *Hydrol Process*, 34, 1906-1919, 10.1002/hyp.13702, 2020b.
- ~~Zeileis, A., Lumley, T., Graham, N., and Koell, S.: sandwich: Robust Covariance Matrix Estimators, available at: <https://cran.r-project.org/web/packages/sandwich/index.html>, R package version 3.0-0, 2020.~~
- 595 Zhao, G., Bates, P., and Neal, J.: The Impact of Dams on Design Floods in the Conterminous US, *Water Resour Res*, 56, ARTN e2019WR025380, 10.1029/2019WR025380, 2020.