



Rainbow colors distort and mislead research in hydrology – guidance for better visualizations and science communication

Michael Stoelzle¹ and Lina Stein²

¹Faculty of Environment and Natural Resources, University of Freiburg, Freiburg, Germany

5 ²Department of Civil Engineering, University of Bristol, Bristol, UK

Correspondence to: Michael Stoelzle (michael.stoelzle@hydro.uni-freiburg.de)

Abstract.

Nowadays color in scientific visualizations is standard and extensively used to group, highlight or delineate different parts of data in visualizations. The rainbow color map (also known as jet color map) is famous for its appealing use of the full visual
10 spectrum with impressive changes in chroma and luminance. Beside attracting attention, science has for decades criticized the rainbow color map for its non-linear and erratic change of hue and luminance along the data variation. The missed uniformity causes a misrepresentation of data values and flaws in science communication. The rainbow color map is scientifically incorrect and hardly decodable for a considerable number of people due to color-vision deficiency (CVD) or other vision
15 impairments. Here we aim to raise awareness how widely used the rainbow color maps still is in hydrology. To this end we perform a paper survey scanning for color issues in around 1000 scientific publications in three different journals including papers published between 2005 and 2020. In this survey, depending on the journal, 16-24% of the publications have a rainbow color map and around the same ratio of papers (18-29%) use red-green elements often in a way that color is the only possibility to decode the visualized groups of data. Given these shares, there is a 99.6% chance to pick at least one visual problematic
20 publication in 10 randomly chosen papers from our survey. To overcome the use of the rainbow color maps in science, we propose some tools and techniques focusing on improvement of typical visualization types in hydrological science. Consequently, color should be used with more care to highlight most important aspects of a visualization and the identification of correct data types such as categorical or sequential data is essential to pick appropriate color maps. We give guidance how to avoid, improve and trust and color in a proper and scientific way. Finally, we sketch a way to improve the communication of rainbow flaws between different status groups in science, publishers, and the media.

25 1 Why does the rainbow color map distort and mislead scientific visualizations?

Colorful visualizations are deeply integrated in science communication. In hydrology, visualization of water fluxes like precipitation, evapotranspiration, discharge or percolation and terms like green and blue water, humidity and aridity or flood and drought are subjects of the daily hydrologists work. Our presentation of patterns, relationships, compositions, distributions and comparisons of multivariate datasets is often multifaceted. And they are most often encoded with color. This is first of all



30 reasonable as human perception is dominated by visual perception (70% compared to 30% by the other senses). The human eyes can recognize around 10 millions of unique colors but only 30 shades of grey (Kreit et al., 2013). Today computer software and freely available programming tools like R or Python simplify the use of color and color gradients in color maps. Also, the rise of online only journals reduced the necessity for a good perception of black-white printed graphs or papers. Although colorful graphs and maps can be created with a few clicks, the development of a compelling visualization is a complex task.

35 In terms of correct encoding visual mappings such as position, length, angle, direction, area and volume rank higher in efficiency and accuracy than color (e.g. Wong, 2010). Roughly said, we are stronger in encoding data that is mapped in a bar- or scatterplot than in a colorful heatmap. The encoding accuracy of color maps has primarily been criticized when it comes to the rainbow color map. This color map uses all wavelengths of the visible spectrum between 380 and 750 nanometers, impresses with high lightness and chroma, encapsulates the most saturated colors and hence looks at first sight very appealing

40 and eye-catching (Figure 1). In the past, the rainbow or jet color map or red and green colors for sequential data were often the software standard causing a wide use in the scientific communities and hence also in publications. For example, in former versions of the statistics software R (version 3.x) the pre-set color map uses black, red, green and dark blue as the first four colors and failed numerous colorblind checks.

In general, there are two main reasons why the rainbow color map in scientific visualization is “(still) considered harmful”

45 (Borland and Taylor, 2007). First, color vision deficiency (CVD) affects the perception of up to 8-10% of the male and up to 0.4-0.5% of the female population, depending on earth regions and, thus up to 4% of world’s population (Geissbuehler and Lasser, 2013; Nuñez et al., 2018; Pramanik et al., 2012). CVD shares are given for Caucasian people and might be lower among other ethnic groups. The simultaneous and side by side use of red and green as in the rainbow color map obstruct an unbiased access to the visualization for these people. Putting an 8% CVD ratio into perspective, a Caucasian male team of one


50 editor and two reviewers during a paper review has a chance up to 22.1% that at least one person has a CVD (Wong, 2011a). Secondly, the rainbow color map attracts attention but is weak in representing data in a scientifically correct way (Figure 1). This affects all people, even those with normal color vision. The same Euclidean distances in mapping or the same data ranges in continuous or binned variables are not equally represented by a rainbow color map (Cramer et al., 2020; Sharma and Trussell, 1997). Especially for data comparisons over a wider distance in the color map the distorted colors of the rainbow

55 impedes reliable judgements (Liu and Heer, 2018). Abrupt changes of lightness and saturation often leads to an unintended focus on some sections of the data range (Thyng et al., 2016). The high lightness of the yellow, cyan or magenta segments in the rainbow color map make it difficult to perceive a consistent color and data value ordering (Kovesi, 2015). Also, high and low values could be confused if both are represented by reddish colors at the edges of the rainbow color map. The color map distorts the data representation if the change in value is not visually commensurate with the change in color (Wong, 2010,

60 2011b). Discordant false coloring may lead to visual errors up to 7.5% of the total displayed data variation (Cramer et al., 2020). For example, research has also shown that replacing a rainbow color map with a perceptual uniform color map could identify hidden structures in mapping (Rogowitz et al., 1996). Comparisons of rainbow color maps and perceptual uniform color maps in cartographic mapping have demonstrated that rainbow colors can emphasize strong gradients where actually



smooth data variation is apparent (Figure 3 in Thyng et al., 2016). Empirical judgement of different quantitative color maps
 65 has hence identified the rainbow color map as perceptually much slower and more error-prone compared to single-hue color
 maps or perceptual uniform designed multi-hue color maps (Liu and Heer, 2018).



70 **Figure 1: a) Comparison of different color maps along an arbitrary scale. The same delta changes (+1) in values are not represented uniformly in the rainbow or heated body color map due to unordered luminance. Alternatives are monochromatic, single-hue color maps (greys, blues) or perceptual uniformly designed multi-hue color maps like viridis or plasma. Visualization inspired by literature (Crameri et al., 2020; Wong, 2011a).**

A thoughtful and scientifically correct color map should allow for all types of dichromatic views (i.e., color vision deficiency) and unambiguous perception of the displayed data. As it is now scientific standard and best practice to avoid any rainbow or rainbow-derivates color map (Crameri et al., 2020) we want to challenge the use of rainbow color maps in hydrological science
 75 by analyzing the status quo of rainbow visualizations in hydrological and environmental publications. In this study we evaluate the use of rainbow and red-green color use in the journal Hydrology and Earth System Sciences (HESS) in preprints and publications in 2020 and their use over time (2005-2020). We then compare the results to two other journals which cover



different disciplines. Finally, we discuss alternatives for using color overall in scientific publications and how to improve and trust the use of color.

80 2 Meta analysis with paper survey

2.1 How often is the rainbow color map used in scientific visualizations?

There are discrepancies between theoretically known scientific standards and the de facto use of the rainbow color map. A non-representative survey of presentations and posters at EGU 2018 found 60% included at least one rainbow scale figure (McNeall, 2018). Compared to publications, visualizations are even more essential for poster presentations and conference talks as less time and text is available to present the research results. The appealing effect of rainbow color maps is often used as eye-catcher along the poster walls. Due to the peer-review process we hypothesized that the ratio of rainbow color maps in publications should be notable lower than 60%. If there is a considerable number of scientific publications with rainbow color maps is there at least a decreasing tendency towards less rainbow color maps in recent years?

First, we examined all preprints that were published in October 2020 (n=36) in the journal Hydrology and Earth System Sciences (HESS). We counted the occurrence of rainbow color maps in each preprint. We found 9 (25%) of these 36 preprints having at least one graph or map with a rainbow color map (called *rainbow preprints* here). In these 9 rainbows preprints the rainbow color map was used to encode variables such as precipitation, temperature, humidity, discharge, soil water content and resistivity or to visualize ratios between variables or to color specific hydrological events. Three of the nine rainbow preprints colored 70-80% of all figures with a rainbow color map. Interestingly, none of the rainbow preprints were single-author papers. The median authors number was five suggesting that rainbow-colored visualizations are not necessarily seen as a critical issue during manuscript preparation and internal submission processes. We then consulted the author guidelines from the journal (<https://www.hydrology-and-earth-system-sciences.net/submission.html>) and found in total two occurrences of the term “color” on the webpage. In the section “Figures and tables” recommendations for high quality graphics are given with: “For maps and charts, please keep color blindness in mind and avoid the parallel usage of green and red. For a list of color scales that are illegible to a significant number of readers, please visit “ColorBrewer 2.0.”.

In a second step after the preprint analysis, we evaluated if the review process reduces the use of rainbow-colored visualizations. We did that by screening in total 263 peer-reviewed papers published in HESS in the year 2020. To our knowledge no systematic review of rainbow color maps in environmental journals exists so far. The journal guidelines of HESS also recommend to avoid green and red colors side by side in visualizations. We therefore classify the papers into four groups also considering pure black-white papers:



- A) Black-white paper without use of any color,
- B) paper has no rainbow-colored visualization or supports distinction with additional elements,
- C) paper has at least one visualization with rainbow-related coloring or use of green and red elements without a
110 good chance to separate these elements.
- D) paper has at least one rainbow-colored visualization (graph or map).


That means a graph with a red and green boxplot could be classified as acceptable (Cat. B) as often axis labels explain the
boxplot groups. A graph with two lines (red and green) encoding continuous variables over time without any annotations more
115 than the legend is classified as rainbow-related (Cat. C). If a paper has a rainbow color map visualization then potential misuse
of red/green is not further counted in our statistics.

The majority of 168 (64%) papers in 2020 have not included any rainbow-colored visualization. For 58 papers (22%) we found
at least one graph or map that uses explicitly the rainbow color map. In 37 papers (14%) we classified at least one graph as
“rainbow-related” (e.g., use of the spectral color map) or identified red-green data encodings without a good chance to
120 distinguish different lines or points. Summarizing these color issues, our survey shows that around 36% of the publications in
HESS in 2020 had visualizations that are not scientifically correct, not perceptual uniform and not or hard to access for around
4% of the readership due to color vision deficiency. This indicates that the awareness of misleading color choice is rather low
during publication process in both authors and reviewers. This was further confirmed by evaluating reviewer comments of
articles published in 2020. We searched for keywords “blind”, “color”, “colour”, “green”, “deficiency”. Of 263 articles
125 published, 9 reviewer comments (3.4 %) mention necessary improvements regarding color or problems with readability of the
graph. Only two comments specifically address the issue a red-green color scale will have for some readers. In reaction one
published article changed the color scale to orange-blue instead of red-green. The other article changed most graphs to a vision
deficiency scale, but also did change one color scale to rainbow. For a further three articles the reviewer criticized readability
of the graph. In reaction, one article changed a red-green coloring to red-black. The other two did not change their figures and
130 continued the use of red-green or rainbow colors. This demonstrates that reviewers and editors are not sufficiently aware of
this problem.

We then extended the survey for all papers published in 2015, 2010 and 2005 to better understand if there is or was a tendency
towards more or less rainbow-colored visualizations in scientific publications (Table A1). Nearly half of the examined 800
papers (from 2005, 2010, 2015 and 2020) have been classified with critical use of ambiguous, not color blind-friendly color
135 maps (Fig. 2). Survey results indicated that the ratio of papers with rainbow or rainbow-related color maps has been stable
between 2010 and 2020 but markedly increased between 2005 and 2010. From 2005 to 2010 there was a clear increase in color
use (from 56% to 82%) and black and white papers dropped by 26%. The survey of 797 papers results in 9% pure black and
white papers, 47% of chromatic papers showed no color issues, and 44 % of all papers have either used a rainbow color map
in at least one visualization and/or have embedded a visualization with red-green-issues. Two cross checks with 30 and 50
140 randomly chosen papers lead to minor deviations in color classification due to the personal judgement of our reviewer team.



However, a high fraction of rainbow classified papers from the main survey were also classified as rainbow papers in the cross checks, with 86% (6 out of 7 papers) and 92% (12 out of 13 papers) agreement.



145

Figure 2: Proportion of surveyed papers without and with color issues. In total 797 peer-reviewed papers from Hydrology and Earth System Sciences (HESS) with different shares for the years 2005, 2010, 2015, and 2020 were analyzed.

2.2 Is the use of rainbow color maps a journal- or discipline-specific artefact?

To answer this question, we screened in total 200 additional publications in the journals Nature Scientific Reports and Nature Communications. For Nature Scientific Reports we looked at the Top 100 of most downloaded papers in the section Earth Science in 2019 (accessible via <https://www.nature.com/collections/agegihhehi>). On the corresponding website graphical thumbnails are given to preview the research findings. Here we found 10 out of 100 thumbnails have rainbow color maps or rainbow-related coloring. Going into more detail we scanned the 100 papers and found that 26 % of the papers used rainbow color maps in at least one figure (Cat. D) and 18% have figure(s) with potential red-green issues in color perception (Cat. C). We also counted 6% of the surveyed papers in this journal as black and white papers (Cat. A). However, more than half of the

155



investigated papers in Nature Scientific Reports have been classified to be color-issue free (Figure 3). In Nature Communications (survey of 100 papers published between 4 and 6 November 2020, mostly biology and genetics topics) we found no papers in Cat. A (black-white), 55% of papers without color issues (Cat. B), 29% of papers that used red and green (Cat. C), and 16% with a rainbow color map (Cat. D) in at least one visualization (Figure 3).

160 All together we surveyed 997 scientific papers from three journals (published between 2005 and 2020, with 46% of papers from 2019 and 2020) and found 23.7 % of all papers have at least one visualization colored with a rainbow color map. The ratio of misused red-green color combinations is most likely even higher than reported here as red-green issues in rainbow papers are not separately counted in the statistics. However, our results revealed a considerable lower ratio of rainbow color maps compared to Borland and Taylor (2007). Their survey from the 2001 through 2005 IEEE Visualization Conference

165 proceedings found around 50-60% of papers having at least one rainbow visualization. Putting our survey results into perspective, when picking randomly 5 (10) papers one has still a 75.0% (93.7%) chance to encounter at least one paper with a rainbow visualization. If red-green issues are also considered then the chance of at least one visual problematic paper in a selection of 5 (10) papers is 94.4% (99.6%). Our survey suggests a current redistribution of disappearing black and white papers into papers with and without color issues. In 2005 we found 20% of the papers published in HESS having color issues,

170 in 2020 there were 47% of papers with color issues. In the same time the share of black and white papers dropped from 44% to less than 1% in 2020.




Figure 3: Share of papers with at least one visualization with red-green issues or a rainbow color map across different journals.

When analyzing the effect of color issue awareness among author teams we clearly see that a higher number of authors did not necessarily lead to a lower share of color issue-free papers (Figure 4). Chromatic papers without color issues had a 40 – 50% share regardless how many authors were included in these publications. Black and white papers are mostly published in single-author papers or by teams of two authors (27% and 14%, respectively). However, only 4% of all papers are published with visualizations in black and white and the share is decreasing since 2005 (Figure 2). The share of papers with rainbow color

175



180 issue for single-author papers is around 9%. In contrast, considering multiple-author publications, around 20-30% of the papers have rainbow color issues.




Figure 4: Composition of color issues with focus on number of authors. Labels of x-axes show minimum and maximum value among the categories. Each row adds up to 100%.

185 We speculate that single-author publications are more often composed by senior authors preferring simpler, but often clearer visualizations where color is not (primarily) needed to encode the data. Another feasible explanation is that in former years papers were written by less authors than today. Average author number per paper in our HESS paper survey was 3.72 in 2005 and 5.06 in 2020.

190 Text mining analysis of paper title terms (see Appendix, Fig. A1) suggests that often studies with spatial analyses or cartographic maps have an above-average chance that a visualization with color issues is embedded. Here papers with title terms such as “terrestrial”, “map(s)” or “mapping”, “radar”, “satellite”, and “region” or “regional” have a 73-92% chance that a rainbow-colored figure or a figure with red-green issues was included in the paper.

3 Four steps to go beyond the rainbow color map

195 To overcome the need for rainbow color map we present four possibilities to avoid, improve, trust and communicate color in scientific visualizations. The central questions for these four steps are given in Figure 5 as techniques to improve the use of color and to communicate misuse of color.




Figure 5: Circular flow of central questions to improve and communicate color in scientific visualizations.

3.1 Avoid color - learning from black and white visualizations

200 Taking inspiration from older papers with black and white visualization is a valuable approach to identify potential improvement of colored visualizations. In former times technology and/or computer software did not allow for the same use of color in visualizations as today. Years ago, colored pages in visualization have also been additionally charged by the publishers.

205 Today color is often the first choice of data encoding in visualizations, but colors are also often used without any reason. Despite the fact that the human eye can differentiate millions of colors, Stauffer et al. (2015) stated that only a small number of different hues can be processed for important classification tasks (search and distinguishing). Healey (1996) showed that only around seven different hues can be found accurately and rapidly on a map or cartographic application. On maps, and also heatmaps, the neighboring colors and the distance between two colored elements bias the perception of data variability (Brychtová and Çöltekin, 2017). If so, visualizations with extensive color use should be revised to reduce the number of colors or redesigned using other graphical encodings. Structure, hierarchy, clarity and completeness can instead be used to create an appealing look on a figure instead of color.

210 Some examples in the surveyed and other literatures illustrate potential ways of doing this: Response surfaces of model bias with a grey color gradient (Schaeffli et al., 2005), black and white map shadings (Milly, 1994), response time distributions of different catchments with lines with various greys, thickness and line types (Roa-García and Weiler, 2010), monthly regime curves of different climate models by lines with different line types and additional point symbols to highlight a specific baseline (Kingston and Taylor, 2010), monochromatic mappings and cumulative fluxes (Campbell et al., 2015), stacked bar charts with a sequential grey color scale (Sunyer et al., 2015) and various point shapes to represent groups (Burden et al., 2019; Sunyer et al., 2015) or visualizations with direct labelling instead of legends (Hoellein et al., 2019).

220 Hydrology as a science particularly uses line charts to illustrate change over time, for example in streamflow analysis. We thus suggest to find visual inspiration from black and white papers to demonstrate how color could be avoided or reduced. At the

same time specific aspects of a visualization can be explicitly highlighted for the reader (Figure 6). If a single technique is not sufficient to improve the visual statement then a combination of techniques could be also feasible, e.g., lines with various width and types, additional overlaid points on the lines or direct labels to highlight specific lines such as the baseline or the mean (Figure 6). Especially direct labelling could improve the clarity of the (line) graphs leading to less cluttered graphs due to additional white space when the legend box is removed. Text elements, rich in contrast, give guidance for the reader to comprehend the story and support people with low vision to easily identify the major elements of the visualization.




Figure 6: Recommendations to improve colorful line graphs or line graphs with red and green lines next to each other. Improvements can be achieved by adjustment of line width, type or brightness (a-c), adding additional points or labels to lines (d-f), focusing on one specific line (g), using small multiples to allow for easier comparison (h) and ensuring that the figure is fully self-explanatory with a precise and complete figure caption and an informative graph title (i). Colorblind-friendly color map, increased contrast and dark(er) backgrounds (e.g., increasing contrast to highlight lines during a presentation) could also improve line charts.

Scientists have the possibility to use other visual encodings than color such as position (e.g. scatterplots), length (e.g. bar charts) or different points shapes or line width (Fig. 6a-d) to increase the perception of data variation in the visualization (Kelleher and Wagener, 2011; Wong, 2010). Furthermore, variation in transparency could also improve the clarity of visualization especially when data points are plotted above each other (e.g., in large sample hydrology or scatterplots). If data encoding is shifted from color to other mappings such as position or line width, a pointedly use of one or two colors is a strong technique



to highlight specific parts of the visualization such as baselines, extremes, averages or specific periods or regions (Fig. 6g). Improved visualization should be accompanied by elaborative figure captions and all authors are asked to create self-explanatory figures that are fully understandable solely with the information in the caption (Rougier et al., 2014). Informative, story-telling graph titles, may enhance data visualizations (Wanzer et al., 2021), although they are unpopular in the scientific community yet.

Beside improvements in single visualizations, splitting figures and maps in different subplots (i.e., facets in Fig. 6h) allows for multiple views on the story of a visualization (Shoresh and Wong, 2012). Here the facets or sparklines (Streit and Gehlenborg, 2015) replace different colors in a single plot. A common technique is to present all data points in all facets as a background data variation and then use color or luminance to highlight specific data groups in the single facets (Gnann et al., 2019, 2020). This multi-faceted view might be valuable for Budyko curve analyses, visualization of different model runs, catchment comparisons or to highlight different distributions of data groups along one axis (histograms, density plots, area charts), on two axes (scatterplots) or if grouping in stacked visualizations (bar or area charts) is encoded by color. For example, dense scatterplots with a lot of overplotting like storage-discharge-plots in hydrological recession analysis (Stoelzle et al., 2013) or in large sample hydrology could profit from faceting as then data encoding is shifted from color to position.

Another important issue in hydrological science are heatmaps as they allow to visualize a third variable in a two-dimensional coordinate system. Heatmaps notably rely on color encoding, thus need an appropriate color map and a meaningful order of the data (categories) on rows and columns. Rethinking the order of rows and columns in heatmaps often reveals that an alphabetical or a chronological order is not the best choice. Similarity and clustering of categories may help out here (Gehlenborg and Wong, 2012a), and clustering can be done by splitting a single heatmap in multiple heatmaps. As color perception in heatmaps depends on colors of neighboring cells, varying stroke color or line width around the tiles will improve the data perception. For example, white, black or grey strokes and light or dark background color around the heatmap tiles could improve color perception. Adding numeric values as text to heatmap's tiles reduces the ambiguity of color perception. Parallel coordinate plots might be a valuable alternative to heatmaps by shifting the data encoding from color to position (Gehlenborg and Wong, 2012a). Multivariate data can be split into several two-dimensional visualizations to improve the clarity of the visualization (Gehlenborg and Wong, 2012c).

But there is another very important aspect in advancing data visualizations. Extensive use of color and perceptual ineffective color maps like the rainbow color map impede not only people with CVD but also affect people with low or reduced vision. The World Health Organization (WHO) states that “at least 2.2 billion people have a vision impairment” (World Health Organization, 2019). Although a lot of those people have received professional eye care, a reduction of visual acuity could impede access to overloaded and cluttered visualizations with less contrast. It is important to recognize that the group of people with low or limited vision or visibility is much larger (up to 28% of world's population) than the group of people with CVD. To visualize for people with low vision, high contrast and supportive text elements or pointers are most important. Visualization should hence be improved with high text and element contrasts, annotations, the ease of horizontal labeling without line breaks, clear figure structure with elaborated hierarchy and focus within the visualization (Tufte, 1983). Gestalt



principles such as similarity, closure, proximity and common regions help to achieve grouping and partitioning in graphs. The idea of data-ink-ratio (Tufte, 1983) could help to remove cluttered, non-data elements and allow for more white space as this helps to focus on important parts of the visualization. Less, thinner or removed grid lines also increase the data-ink-ratio and sharpen the view on the data.

In a second step, changing perspective from authors to publishers, the community needs more advanced tools than static PDF or printed papers (Vandemeulebroecke et al., 2019). If articles are more and more published in HTML format, interactive visualizations could allow for multiple perspectives (e.g., zoom, selection, layering). Interactive visualizations give the possibility for people with low vision or CVD to select supportive elements like tooltips during mouse hovering or by highlighting selected elements and give direct labeling and annotations. Using alt attributes in HTML can be used to specify alternative text if a visualization cannot be rendered or perceived. Interactive elements on basis of HTML and R packages are, for example, `dygraphs` (Vanderkam et al., 2018) with sliders to select specific time periods in time series analysis or `leaflet` maps (Cheng et al., 2021) giving the possibility to zoom into maps and to select different background layers (i.e., leading to higher contrast for people with vision impairment). Authors can also accompany their data analysis in a paper with an online available data dashboard such as `shiny` apps (Chang et al., 2020) further data exploration.

3.2 Improve color - what are alternatives to the rainbow color map?

Although black and white visualizations could inspire a thoughtful revision of colored figures or maps, removing all color is not always the best choice. Depending on data dimensions the use of color is sometimes unavoidable. Typically, we want to use color to convey data and also to create a figure or map that looks appealing. When that is the case, there are several techniques how color can be used in an accessible way.

Crameri et al. (2020) presented a thoughtful decision tree explaining how color depending on data types should be used in visualizations. They differentiated between the direction of color gradients to encode higher or lower values according to the chosen background color (light or dark). Dark(er) backgrounds have recently gained more and more attention in the visualization community as possibility to increase the contrast of visualizations (Crameri et al., 2020). Although dark figure backgrounds make for an unusual sight in articles, the increase in contrast might be appropriate for presentations helping people with vision impairments. If a single hue-color map such as the blue color map in Figure 1 is not sufficient to encode data by color then blended hue-color maps are a feasible solution. For example, to visualize depth below and elevation above sea level a combination of two monochromatic color maps with a reasonable mid- or breakpoint can be used (Gehlenborg and Wong, 2012b). If small data variations in a continuous color map are not needed, then a discrete or binned color map with less but more distinguishable colors support faster decoding of color and data variation. Thoughtful breaks for color binning and better boundaries among important data ranges or regions give guidance for the reader. Here the proximity and orientation of the legend could also help to gain undistorted allocation of the presented data range. Depending on the degree of data variability on x- and y axis the legend could be horizontally or vertically aligned to the graph.




In summary, finding an alternative for the rainbow color map can be seen as a relatively straightforward process when two
305 main aspects of color maps are considered. Firstly, there are many perceptually uniform color maps available that are well
documented and professionally designed (e.g., to consider people with CVD). Such color maps are often available for all
visualization software or programming languages (e.g., R, Matlab, Python) for example the *viridis* (Garnier, 2018) or
scico (Pedersen and Crameri, 2020) packages in R. A thoughtful color map should be perceptually uniform (i.e., delta change
in color is equal to delta change in data), color-blind friendly, strong in greyscale conversion and - if possible - pretty and
310 appealing to attract the reader (Figure 7). Famous examples are the Okabe color map (Okabe and Ito, 2008), the *viridis*
color map (Garnier, 2018) or the recently published scientific color maps by Crameri (2020).

Secondly, a simple guidance on the necessity for color should be considered. The main uses of color are within these four
categories:

- A) Distinguish categories (i.e., each color is a category),
- 315 B) visualize sequential data values (i.e., each color is a numeric value),
- C) visualize diverging data values (i.e., each color is a numeric value and data has a direction and a meaningful mid-
point or centric value), and
- D) highlight as special for of categorical scale (i.e., a few categories have a color, the remaining have a grey gradient).

We recommend as a first step to assign used data to one of the four cases A) to D). If this is not possible, color might not be
320 the best way to convey the data. Our survey revealed that especially the cases B) and C) were often mixed up or used color in
the wrong way (Figure 7). Examples how to find appropriate color maps based on data types can be found in literature (Coalter,
2020; Light and Bartlein, 2004; Zeileis et al., 2019). Empirical assessment of quantitative color maps has also shown that
diverging color maps (e.g. blue-white-orange) are slower and more error-prone during data encoding compared to single-hue
color maps (Liu and Heer, 2018). Here comparisons of data values across the white mid-point are critical. For example, there
325 is no need to have diverging color maps to visualize correlation coefficient between 0 and 1, but they may be useful for ranges
between -1 and 1.

Another important point is to know typical color maps of specific types of visualization or in specific research areas. For
hydrology or environmental science, the use of specific diverging color scales such as red/blue for temperature or green/brown
for aridity can also function as a first and easy decodable signal for the reader to give advice what is actually shown in the
330 presented figure or map (Figure 7). Text elements or pointers will support colored data decoding for people with low vision.
Such thematic and discipline-specific color choices aligned with peer discussions should be part of a thoughtful visualization
development within each author team.



335 **Figure 7: Comparison of mapping with rainbow and perceptually uniform color maps for an arbitrary variable across the counties**
of the US state Texas. For each map also the vision for people with deuteranopia (red-green blindness) is given. With the rainbow
color map values around 60 and 20 stick out (a, c) and greenish or yellowish colors dominate a wide range of data variation.
Compared to that uniform and colorblind-friendly color maps (b, d) support a more exact perception of data variation and the
extreme values. White strokes between the map elements (b, d) increase the data-ink-ratio. Maps with emulation of color vision
deficiency were created with the R package `colorblindr`.

340 3.3 Trust color – tools to check for colorblind-friendly visualizations

“In perceptual-uniformity we trust!” (Shephard et al., 2017). There are mainly two possibilities to increase the trustworthiness
of color choice in scientific visualizations. Firstly, by learning from surveys testing the trustworthiness of colors and different
color maps. Examples are the Marie Curie (Cramer et al., 2020) or the Which Blair project (Rogowitz and Kalvin, 2001)
where different color palettes mapped to well-known photos to highlight the effect of different color choice. These studies
345 showed with experiments that color maps not based on a monotonically increasing luminance component produced no positive
rating scores (Liu and Heer, 2018). Considering these findings, tools like Colorbrewer 2.0 (<https://colorbrewer2.org>) give color
advice what kind of single- or multi-hue color maps can be used for sequential, diverging or qualitative data and let the user
explicitly filter for colorblind safe and printer friendly color maps (Coalter, 2020; Harrower and Brewer, 2003). With the R
package `colorspace` (Zeileis et al., 2019) a set of over 80 color palettes can be visualized and compared among each other.
350 In the R package `scico` (Pedersen and Cramer, 2020) users will easily find palettes that embed common colors for
visualization in their disciplines (see Sect. 3.2). The R package `RColorBrewer` (Neuwirth, 2012) provides color maps for



visualizations and offers the possibility to use only colorblind friendly color maps with the option `colorblindFriendly=TRUE`.

355 Secondly, when a visualization is ready for publication authors should aim for testing the specific figure or map how trustworthy the used color map actually is. With the R package `colorblindr` (McWhite and Wilke, 2021) various types of colorblindness are simulated for production-ready R visualizations (Figure 7). Open-source software application for smartphones and computers offer a livestream of color-blindness emulation via camera or screen capture (e.g., <https://github.com/michelf/sim-daltonism/>). Such emulations are also available on the internet without using a specific software, e.g., <https://www.color-blindness.com> or <http://hclwizard.org>. Figure files can be uploaded to compare different
360 sights such as normal, deuteranopia, tritanopia or monochromacy. To improve visualizations for people with other vision impairments online tools like <https://contrastchecker.com> help to test color contrast compliance or to perform greyscale emulation. As a last step the R package `paletteer` (Hvitfeldt, 2021) offer the possibility to produce some descriptive statistics on a color map or publication-ready figure and to run routines that optimize the color map, e.g. avoid colors that are appearing too similar.

365 **3.4 Communicate (rainbow) color - what should scientists and publishers do?**

Literature review and our paper survey of color maps in scientific publications suggest that there is a considerable discrepancy between what science knows about the rainbow color map and what scientists do about it. The presented paper survey shows this discrepancy. Around 47% of published HESS papers in 2020 have color issues but at the same time only less than 4% of all reviewer comments see the choice of color as an issue before publication. From a knowledge perspective the rainbow color
370 map distorts a correct representation of the data variation and exclude people with CVD or other vision impairments to understand the presented scientific work correctly. With that a reliable scientific communication is not possible. This knowledge is not reflected in the submission process of many journals, although the journals have been requested to raise author's awareness about CVD accessibility issues (Albrecht, 2010). If the extensive use of rainbow color maps in science continues then also journalists and (social) media will most likely continue to circulate those rainbow figures and maps. Today
375 rainbow thumbnails appear in graphical abstracts, as thumbnails on journal websites or as screenshots in paper announcements on Twitter. This suggests for a broader audience that the rainbow color map is state of the art and scientifically correct as the visualizations have been produced by scientists with a high reputation in the public opinion. For young(er) scientists the high reputation of scientific journals justifies rainbow color maps as appropriate for their own scientific work.

To leave this vicious circle a major effort in science communication is needed. Publishers and editorial teams should review
380 graphical abstracts and summary thumbnails for rainbow color maps as authors tend to use rainbow-colored figures from the paper to attract readers' attention on websites with paper previews. Journal's author guidelines should specifically advice against the use of rainbow and red-green scales. Editors and reviewers should ask for revisions of rainbow figures and should be more relentless here. At scientific conferences, short courses for improved data analysis or environmental visualizations



385 should rise awareness for the rainbow color map topic, especially for young(er) scientists. This all is not about blaming the
authors of rainbow visualizations, but to clearly criticize those figures and mappings in a fair and constructive way proposing
methods to improve or avoid the (rainbow) coloring. Although the rainbow color map has more or less a tradition in various
hydrological subdisciplines (e.g., in visualizations of water velocities, heat or solute transport and cartographic maps in
general), we encourage especially the networks of young scientists to take responsibility for visualizations with valid color
maps and a clear undistorted message. Communication of rainbow flaws should take place in all parts of science: during
390 lectures, with colleagues, in network meetings, as feedback for presentations, as a conference attendee or paper reviewer, but
also as a journal editor, senior scientist or professor. The anti-rainbow Marie Curie poster from Crameri et al. (2020) is freely
available and could be a communication starter at the wall near the coffee machine of your institute.

Conclusions

The rainbow color map attracts attention but distorts and misleads scientific visualizations. Major rainbow pitfalls are the non-
395 linear data encoding, steps and disorder in luminance and minor perceptual accessibility for people with CVD or other vision
impairments. Here we investigated the use of rainbow color maps in around 1000 papers in different environmental journals
and found that the misleading rainbow color map or red-green color issues are present in around 44% of all papers. Compared
to the knowledge about the flaws of the rainbow color map this share is alarming high. Moreover, our hypothesis that rainbow
color maps are on the decline could not be confirmed. Color issues in papers stay constant or even increase between 2005 and
400 2020. Multi-author papers are not less prone to (rainbow) color issues, even though more people could weigh in against
inaccessible visualizations. Analysis of reviewer comments highlights that the awareness of those issues is alarmingly low
during the review process.


Our survey indicates that past campaigns to banish the use of rainbow color maps were not sufficient. We strongly recommend
that this issue should be raised across the hydrologic community. It will take students, researchers, lectures, professors, editors,
405 reviewers and publishers to banish the rainbow color map to make visualizations accessible for all and to insist on correct data
representation. As a guide we presented manifold visual techniques on how to avoid, improve, trust and communicate color in
data visualizations. Such guidance is given with focus on important graph types in hydrology to attenuate the role and risks of
color use in data encoding. Visualizations could not only be improved for people with CVD but should be drafted with more
care in terms of less exclusive data encoding by color. Such efforts could also bring advantages for a much larger group of
410 people with low vision or vision impairments if more focus is given to visualizations with less clutter, higher contrast and
supportive graphical elements like annotations.



Appendix

415 **Table A1: Survey of papers in the Hydrology and Earth System Sciences journal. Classification is done based on expert judgement. Missing % to 100 is due to rounding. In 2020 only papers are considered that were published before 1st November.**

Category	Description	Year	2005	2010	2015	2020	total
		Number of papers	54	191	289	263	797
A	Pure black-white		44%	18%	4%	<1%	9%
B	Chromatic without color issues		35%	41%	48%	53%	47%
C	Red-green color issues		9%	16%	22%	23%	20%
D	Rainbow color map issues		11%	25%	26%	24%	24%



420 **Figure A1: Text mining analysis finding 27 groups of papers sharing the same title term and having an above-average share of color issues. A paper title is only considered for analysis if the group of papers with the specific term in the title has more than 44% of papers with color issues (above the average of 797 papers). Asterisk (*) acts as a wildcard in the regular expressions during text mining, e.g., with *map** also the terms *maps* and *mapping* are searched. Listed terms have a minimum of 5 letters (exception *map** and *snow**) and occur in at least 10 individual papers (exception *bias correct** with 7 papers). A specific paper could be part of multiple paper groups. More generic words like effect, approach, change, based, water or model and country names like China or France were excluded by expert judgement. Analysis is based on all 797 surveyed papers from the journal Hydrology and Earth System Sciences (2005-2020). The median author number across all groups of papers with the same title term ranges between 3.0 (map* and seasonal*) and 5.0 (evapotranspiration, Mediterranean, region*, and soil moist*).**

425



Data availability

Data of the paper survey and sample code snippets are online available at https://github.com/modche/rainbow_hydrology and as Kaggle notebook for exploration at <https://www.kaggle.com/modche/rainbow-papersurvey-hydrology>.

Author contribution

430 MS developed the research idea and designed the paper survey. MS and LS carried out the paper survey and analyzed the results. MS wrote the manuscript and designed the figures with contributions from LS.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

435 Robin Schwemmler evaluated the color issue classification and Angelika Kuebert suggested R packages for colorblind-friendly color mapping. Thanks are due to Robin Schwemmler and Markus Weiler whose comments greatly helped to improve the visualizations and the paper.

References

- 440 Borland, D. and Taylor, R.: Rainbow Color Map (Still) Considered Harmful, *IEEE Comput. Grap. Appl.*, 27(2), 14–17, <https://doi.org/10.1109/MCG.2007.323435>, 2007.
- Brychtová, A. and Çöltekin, A.: The effect of spatial distance on the discriminability of colors in maps, *Cartography and Geographic Information Science*, 44(3), 229–245, <https://doi.org/10.1080/15230406.2016.1140074>, 2017.
- Burden, C. M., Morgan, M. O., Hladun, K. R., Amdam, G. V., Trumble, J. J. and Smith, B. H.: Acute sublethal exposure to toxic heavy metals alters honey bee (*Apis mellifera*) feeding behavior, *Sci Rep*, 9(1), 4253, <https://doi.org/10.1038/s41598-019-40396-x>, 2019.
- 445 Campbell, J. M., Jordan, P. and Arnscheidt, J.: Using high-resolution phosphorus data to investigate mitigation measures in headwater river catchments, *Hydrol. Earth Syst. Sci.*, 12, 2015.
- Chang, W., Cheng, J., Allaire, J., Xie, Y. and McPherson, J.: shiny: Web Application Framework for R. <https://CRAN.R-project.org/package=shiny>, 2020.
- 450 Cheng, J., Karambelkar, B. and Xie, Y.: leaflet: Create Interactive Web Maps with the JavaScript “Leaflet” Library. <https://CRAN.R-project.org/package=leaflet>, 2021.



- Coalter, J.: ColorBrewer 2.0 and the Rainbow: Using Color Tools to Choose Appropriate Color Schema for your Data Visualization, *ISTL*, (94), <https://doi.org/10.29173/istl63>, 2020.
- Crameri, F.: Scientific Colour Maps, <http://www.fabiocrameri.ch/colourmaps>, last access: 12 February 2020, 2020.
- 455 Crameri, F., Shephard, G. E. and Heron, P. J.: The misuse of colour in science communication, *Nat Commun*, 11(1), 5444, <https://doi.org/10.1038/s41467-020-19160-7>, 2020.
- Garnier, S.: viridis: Default Color Maps from “matplotlib.” <https://CRAN.R-project.org/package=viridis>, 2018.
- Gehlenborg, N. and Wong, B.: Heat maps, *Nat Methods*, 9(3), 213–213, <https://doi.org/10.1038/nmeth.1902>, 2012a.
- Gehlenborg, N. and Wong, B.: Mapping quantitative data to color, *Nat Methods*, 9(8), 769–769, 460 <https://doi.org/10.1038/nmeth.2134>, 2012b.
- Gehlenborg, N. and Wong, B.: Power of the plane, *Nat Methods*, 9(10), 935–935, <https://doi.org/10.1038/nmeth.2186>, 2012c.
- Geissbuehler, M. and Lasser, T.: How to display data by color schemes compatible with red-green color perception deficiencies, *Opt. Express*, 21(8), 9862, <https://doi.org/10.1364/OE.21.009862>, 2013.
- Gnann, S. J., Woods, R. A. and Howden, N. J. K.: Is There a Baseflow Budyko Curve?, *Water Resour. Res.*, 55(4), 2838–465 2855, <https://doi.org/10.1029/2018WR024464>, 2019.
- Gnann, S. J., McMillan, H., Woods, R. A. and Howden, N. J. K.: Including Regional Knowledge Improves Baseflow Signature Predictions in Large Sample Hydrology, *Water Resour. Res.*, <https://doi.org/10.1029/2020WR028354>, 2020.
- Harrower, M. and Brewer, C. A.: ColorBrewer.org: An Online Tool for Selecting Colour Schemes for Maps, *The Cartographic Journal*, 40(1), 27–37, <https://doi.org/10.1179/000870403235002042>, 2003.
- 470 Healey, C. G.: Choosing effective colours for data visualization, in *Proceedings of Seventh Annual IEEE Visualization '96*, pp. 263-270, ACM, San Francisco, CA, USA, <https://doi.org/10.1109/VISUAL.1996.568118>, , 1996.
- Hoellein, T. J., Shogren, A. J., Tank, J. L., Risteca, P. and Kelly, J. J.: Microplastic deposition velocity in streams follows patterns for naturally occurring allochthonous particles, *Sci Rep*, 9(1), 3740, <https://doi.org/10.1038/s41598-019-40126-3>, 2019.
- 475 Hvitfeldt, E.: paletteer: Comprehensive Collection of Color Palettes. <https://CRAN.R-project.org/package=paletteer>, 2021.
- Kelleher, C. and Wagener, T.: Ten guidelines for effective data visualization in scientific publications, *Environmental Modelling & Software*, 26(6), 822–827, <https://doi.org/10.1016/j.envsoft.2010.12.006>, 2011.
- Kingston, D. G. and Taylor, R. G.: Sources of uncertainty in climate change impacts on river discharge and groundwater in a headwater catchment of the Upper Nile Basin, Uganda, *Hydrol. Earth Syst. Sci.*, 12, 2010.
- 480 Kovesi, P.: Bad Colour Maps Hide Big Features and Create False Anomalies, *ASEG Extended Abstracts*, 2015(1), 1–4, <https://doi.org/10.1071/ASEG2015ab107>, 2015.
- Kreit, E., Mäthger, L. M., Hanlon, R. T., Dennis, P. B., Naik, R. R., Forsythe, E. and Heikenfeld, J.: Biological versus electronic adaptive coloration: how can one inform the other?, *J. R. Soc. Interface.*, 10(78), 20120601, <https://doi.org/10.1098/rsif.2012.0601>, 2013.



- 485 Light, A. and Bartlein, P. J.: The end of the rainbow? Color schemes for improved data graphics, *Eos Trans. AGU*, 85(40), 385, <https://doi.org/10.1029/2004EO400002>, 2004.
- Liu, Y. and Heer, J.: Somewhere Over the Rainbow: An Empirical Assessment of Quantitative Colormaps, in *Proceedings of the 2018 CHI Conference on Human Factors in Computing Systems - CHI '18*, pp. 1–12, ACM Press, Montreal QC, Canada, <https://doi.org/10.1145/3173574.3174172>, , 2018.
- 490 McNeill, D.: How many rainbows at EGU 2018?, *Better Figures* <https://betterfigures.org/2018/04/16/how-many-rainbows-at-egu-2018/>, last access: 12 April 2020, 2018.
- McWhite, C. D. and Wilke, C. O.: colorblindr: Simulate colorblindness in R figures. <https://github.com/clauswilke/colorblindr>, 2021.
- Milly, P. C. D.: Climate, soil water storage, and the average annual water balance, *Water Resources Research*, 30(7), 2143–
495 2156, <https://doi.org/10.1029/94WR00586>, 1994.
- Neuwirth, E.: RColorBrewer: ColorBrewer Palettes. <https://CRAN.R-project.org/package=RColorBrewer>, 2012.
- Nuñez, J. R., Anderton, C. R. and Renslow, R. S.: Optimizing colormaps with consideration for color vision deficiency to enable accurate interpretation of scientific data, edited by J. Malo, *PLoS ONE*, 13(7), e0199239, <https://doi.org/10.1371/journal.pone.0199239>, 2018.
- 500 Okabe, M. and Ito, K.: Color universal design (CUD) - how to make figures and presentations that are friendly to colorblind people, *J* Fly: Data Depository for Drosophila Researchers* <http://jfly.iam.u-tokyo.ac.jp/color/>, last access: 12 February 2020, 2008.
- Pedersen, T. L. and Cramer, F.: scico: Colour Palettes. Based on the Scientific Colour-Maps. <https://CRAN.R-project.org/package=scico>, 2020.
- 505 Pramanik, T., Khatiwada, B. and Pandit, R.: Color vision deficiency among a group of students of health sciences, *Nepal Med Coll J*, 4, 2012.
- Roa-García, M. C. and Weiler, M.: Integrated response and transit time distributions of watersheds by combining hydrograph separation and long-term transit time modeling, *Hydrol. Earth Syst. Sci.*, 14(8), 1537–1549, <https://doi.org/10.5194/hess-14-1537-2010>, 2010.
- 510 Rogowitz, B. E. and Kalvin, A. D.: The “Which Blair project”: a quick visual method for evaluating perceptual color maps, in *Proceedings Visualization, 2001. VIS '01.*, pp. 183–556, IEEE, San Diego, CA, USA, <https://doi.org/10.1109/VISUAL.2001.964510>, , 2001.
- Rogowitz, B. E., Treinish, L. A. and Bryson, S.: How Not to Lie with Visualization, *Comput. Phys.*, 10(3), 268, <https://doi.org/10.1063/1.4822401>, 1996.
- 515 Rougier, N. P., Droettboom, M. and Bourne, P. E.: Ten Simple Rules for Better Figures, *PLoS Comput Biol*, 10(9), e1003833, <https://doi.org/10.1371/journal.pcbi.1003833>, 2014.
- Schaeffli, B., Hingray, B., Niggli, M. and Musy, A.: A conceptual glacio-hydrological model for high mountainous catchments, *Hydrology and Earth System Sciences*, 15, 2005.



- Sharma, G. and Trussell, H. J.: Digital color imaging, *IEEE Trans. on Image Process.*, 6(7), 901–932,
520 <https://doi.org/10.1109/83.597268>, 1997.
- Shephard, G., Meier, T. and Paul, J.: The Rainbow Colour Map (repeatedly) considered harmful, *EGU Blogs*
<https://blogs.egu.eu/divisions/gd/2017/08/23/the-rainbow-colour-map/>, last access: 13 January 2021, 2017.
- Shoresh, N. and Wong, B.: Data exploration, *Nat Methods*, 9(1), 5–5, <https://doi.org/10.1038/nmeth.1829>, 2012.
- Stauffer, R., Mayr, G. J., Dabernig, M. and Zeileis, A.: Somewhere Over the Rainbow: How to Make Effective Use of Colors
525 in Meteorological Visualizations, *Bulletin of the American Meteorological Society*, 96(2), 203–216,
<https://doi.org/10.1175/BAMS-D-13-00155.1>, 2015.
- Stoelzle, M., Stahl, K. and Weiler, M.: Are streamflow recession characteristics really characteristic?, *Hydrol. Earth Syst. Sci.*,
17(2), 817–828, <https://doi.org/10.5194/hess-17-817-2013>, 2013.
- Streit, M. and Gehlenborg, N.: Temporal data, *Nat Methods*, 12(2), 97–97, <https://doi.org/10.1038/nmeth.3262>, 2015.
- 530 Sunyer, M. A., Hundecha, Y., Lawrence, D., Madsen, H., Willems, P., Martinkova, M., Vormoor, K., Bürger, G., Hanel, M.,
Kriaučiūnienė, J., Loukas, A., Osuch, M. and Yücel, I.: Inter-comparison of statistical downscaling methods for projection
of extreme precipitation in Europe, *Hydrol. Earth Syst. Sci.*, 19(4), 1827–1847, <https://doi.org/10.5194/hess-19-1827-2015>,
2015.
- Thyng, K., Greene, C., Hetland, R., Zimmerle, H. and DiMarco, S.: True Colors of Oceanography: Guidelines for Effective
535 and Accurate Colormap Selection, *Oceanog.*, 29(3), 9–13, <https://doi.org/10.5670/oceanog.2016.66>, 2016.
- Tufte, E. R.: *The visual display of quantitative information*, CT Graphics, Cheshire., 1983.
- Vandemeulebroecke, M., Baillie, M., Margolskee, A. and Magnusson, B.: Effective Visual Communication for the
Quantitative Scientist, *CPT Pharmacometrics Syst. Pharmacol.*, 8(10), 705–719, <https://doi.org/10.1002/psp4.12455>, 2019.
- Vanderkam, D., Allaire, J., Owen, J., Gromer, D. and Thieurmel, B.: dygraphs: Interface to “Dygraphs” Interactive Time Series
540 Charting Library. <https://CRAN.R-project.org/package=dygraphs>, last access: 23 February 2021, 2018.
- Wanzer, D. L., Azzam, T., Jones, N. D. and Skousen, D.: The role of titles in enhancing data visualization, *Evaluation and
Program Planning*, 84, 101896, <https://doi.org/10.1016/j.evalprogplan.2020.101896>, 2021.
- Wong, B.: Design of data figures, *Nat Methods*, 7(9), 665–665, <https://doi.org/10.1038/nmeth0910-665>, 2010.
- Wong, B.: Points of view: Color blindness, *Nat Methods*, 8(6), 441–441, <https://doi.org/10.1038/nmeth.1618>, 2011a.
- 545 Wong, B.: Points of view: Points of review (part 2), *Nat Methods*, 8(3), 189–189, <https://doi.org/10.1038/nmeth0311-189>,
2011b.
- Zeileis, A., Fisher, J. C., Hornik, K., Ihaka, R., McWhite, C. D., Murrell, P., Stauffer, R. and Wilke, C. O.: colorspace: A
Toolbox for Manipulating and Assessing Colors and Palettes, arXiv:1903.06490 [cs, stat] <http://arxiv.org/abs/1903.06490>,
last access: 13 November 2020, 2019.