



# The International Soil Moisture Network: serving Earth system science for over a decade

Wouter Dorigo<sup>1</sup>, Irene Himmelbauer<sup>1</sup>, Daniel Aberer<sup>1</sup>, Lukas Schremmer<sup>1</sup>, Ivana Petrakovic<sup>1</sup>, Luca Zappa<sup>1</sup>, Wolfgang Preimesberger<sup>1</sup>, Angelika Xaver<sup>1</sup>, Frank Annor<sup>2,3</sup>, Jonas Ardö<sup>4</sup>, Dennis Baldocchi<sup>5</sup>, Günter Blöschl<sup>6</sup>, Heye Bogena<sup>7</sup>, Luca Brocca<sup>8</sup>, Jean-Christophe Calvet<sup>9</sup>, Julio J. Camarero<sup>10</sup>, Giorgio Capello<sup>11</sup>, Minha Choi<sup>12</sup>, Michael C. Cosh<sup>13</sup>, Jerome Demarty<sup>14</sup>, Nick van de Giesen<sup>3</sup>, Istvan Hajdu<sup>15</sup>, Karsten H. Jensen<sup>16</sup>, Kasturi Devi Kanniah<sup>17</sup>, Ileen de Kat<sup>18</sup>, Gottfried Kirchengast<sup>19</sup>, Pankaj Kumar Rai<sup>20</sup>, Jenni Kyrouac<sup>21</sup>, Kristine Larson<sup>22</sup>, Suxia Liu<sup>23</sup>, Alexander Loew<sup>24,†</sup>, Mahta Moghaddam<sup>25</sup>, José Martínez Fernández<sup>26</sup>, Cristian Mattar Bader<sup>27</sup>, Renato Morbidelli<sup>28</sup>, Jan P. Musial<sup>29</sup>, Elise Osenga<sup>30</sup>, Michael A. Palecki<sup>31</sup>, Isabella Pfeil<sup>1</sup>, Jarrett Powers<sup>32</sup>, Jaakko Ikonen<sup>33</sup>, Alan Robock<sup>34</sup>, Christoph Rüdiger<sup>35</sup>, Udo Rummel<sup>36</sup>, Michael Strobel<sup>37</sup>, Zhongbo Su<sup>38</sup>, Ryan Sullivan<sup>21</sup>, Torbern Tagesson<sup>4,16</sup>, Mariette Vreugdenhil<sup>1</sup>, Jeffrey Walker<sup>35</sup>, Jean Pierre Wigneron<sup>39</sup>, Mel Woods<sup>40</sup>, Kun Yang<sup>41</sup>, Xiang Zhang<sup>42</sup>, Marek Zreda<sup>43</sup>, Stephan Dietrich<sup>44</sup>, Alexander Gruber<sup>45</sup>, Peter van Oevelen<sup>46</sup>, Wolfgang Wagner<sup>1</sup>, Klaus Scipal<sup>47</sup>, Matthias Drusch<sup>48</sup>, and Roberto Sabia<sup>47</sup>

<sup>1</sup>Department of Geodesy and Geoinformation, TU Wien, Vienna, Austria

<sup>2</sup>Trans-African Hydro-Meteorological Observatory, Delft, The Netherlands

<sup>3</sup>Department of Water Resources, Delft University of Technology, Delft, Netherlands

<sup>4</sup>Department of Physical Geography and Ecosystem Science, Lund University, Lund, Sweden

<sup>5</sup>Department of Environmental Science, Policy and Management, University of California, Berkeley, CA, United States

<sup>6</sup>Institute of Hydraulic Engineering and Water Resources Management, TU Wien, Vienna, Austria

<sup>7</sup>Forschungszentrum Juelich GmbH, Juelich, Germany

<sup>8</sup>Research Institute for Geo-Hydrological Protection, CNR, Perugia, Italy

<sup>9</sup>CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France

<sup>10</sup>Instituto Pirenaico de Ecología, IPE-CSIC, Zaragoza, Spain

<sup>11</sup>Institute of Sciences and Technologies for Sustainable Energy and Mobility, National Research Council of Italy, Torino, Italy

<sup>12</sup>Department of Water Resources, Graduate School of Water Resources, Sungkyunkwan University, Suwon, Republic of Korea

<sup>13</sup>USDA-Agricultural Research Service, Hydrology and Remote Sensing Laboratory, Beltsville, MD, United States

<sup>14</sup>Hydrosciences Montpellier, IRD, CNRS, Université Montpellier, France

<sup>15</sup>PlantTech Research Institute, Tauranga, New Zealand

<sup>16</sup>Department of Geosciences and Natural Resource Management, University of Copenhagen, Denmark

<sup>17</sup>Research Institute for Sustainable Environment, Universiti Teknologi Malaysia, Johor Bahru, Malaysia

<sup>18</sup>VanderSat B.V., Haarlem, The Netherlands

<sup>19</sup>Wegener Center for Climate and Global Change and Institute of Physics, University of Graz, Austria

<sup>20</sup>Indian Institute of Technology Kanpur, India

<sup>21</sup>Environmental Science Division, Argonne National Laboratory, Lemont, IL, United States

<sup>22</sup>University of Colorado, Boulder, CO, United States

<sup>23</sup>Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing, China; College of Resources and Environment/Sino-Danish Center, University of Chinese Academy of Sciences, Beijing, China

<sup>24</sup>Department of Geography, Ludwig-Maximilian-Universität, Munich, Germany



<sup>25</sup>Ming Hsieh Department of Electrical and Computer Engineering, University of Southern California, Los Angeles, CA, United States

<sup>26</sup>Instituto Hispano Luso de Investigaciones Agrarias, CIALE, Universidad de Salamanca, Villamayor, Spain

<sup>27</sup>Laboratory of Analysis of the Biosphere, University of Chile, Santiago, Chile

<sup>28</sup>Department of Civil and Environmental Engineering, Perugia University, Italy

<sup>29</sup>Institute of Geodesy and Cartography, Warsaw, Poland

<sup>30</sup>Aspen Global Change Institute, Basalt, CO, United States

<sup>31</sup>NOAA National Centers for Environmental Information, Asheville, NC, United States

<sup>32</sup>Agriculture and Agri-Food Canada - Science and Technology Branch, Winnipeg, MB, Canada

<sup>33</sup>Finnish Meteorological Institute, Space and Earth Observation Centre, Helsinki, Finland

<sup>34</sup>Department of Environmental Sciences, Rutgers University, New Brunswick, NJ, United States

<sup>35</sup>Department of Civil Engineering, Monash University, Clayton, Australia

<sup>36</sup>DWD, Meteorologisches Observatorium Lindenberg – Richard-Aßmann-Observatorium, Lindenberg, Germany

<sup>37</sup>USDA-NRCS National Water and Climate Center, Portland, OR, United States

<sup>38</sup>University of Twente, Faculty of Geo-Information Science and Earth Observation, Enschede, the Netherlands

<sup>39</sup>MOST, ISPA, INRAE Bordeaux Aquitaine, France

<sup>40</sup>University of Dundee, UK

<sup>41</sup>Department of Earth System Science, Tsinghua University, Beijing, China

<sup>42</sup>State Key Laboratory of Information Engineering in Surveying, Wuhan University, Wuhan, China

<sup>43</sup>Department of Hydrology and Atmospheric Sciences, University of Arizona, Tucson, AZ, United States

<sup>44</sup>Federal Institute of Hydrology, Koblenz, Germany

<sup>45</sup>Department of Earth and Environmental Sciences, KU Leuven, Heverlee, Belgium

<sup>46</sup>International GEWEX Project Office, Washington, DC, United States

<sup>47</sup>European Space Agency, ESRIN, Frascati, Italy

<sup>48</sup>European Space Agency, ESTEC, Noordwijk, The Netherlands

<sup>†</sup>Deceased July 02, 2017

**Correspondence:** Wouter Dorigo (wouter.dorigo@tuwien.ac.at)

## Abstract.

In 2009, the International Soil Moisture Network (ISMN) was initiated as a community effort, funded by the European Space Agency, to serve as a centralised data hosting facility for globally available in situ soil moisture measurements (Dorigo et al., 2011b, a). The ISMN brings together in situ soil moisture measurements collected and freely shared by a multitude of organisations, harmonizes them in terms of units and sampling rates, applies advanced quality control, and stores them in a database. Users can freely retrieve the data from this database through an online web portal (ismn.earth). Meanwhile, the ISMN has evolved into the primary in situ soil moisture reference database worldwide, as evidenced by more than 3000 active users and over 1000 scientific publications referencing the data sets provided by the network. As of December 2020, the ISMN now contains data of 65 networks and 2678 stations located all over the globe, with a time period spanning from 1952 to present. The number of networks and stations covered by the ISMN is still growing and many of the data sets contained in the database continue to be updated. The main scope of this paper is to inform readers about the evolution of the ISMN over the past decade, including a description of network and data set updates and quality control procedures. A comprehensive review of existing literature making use of ISMN data is also provided in order to identify current limitations in functionality and data usage, and to shape priorities for the next decade of operations of this unique community-based data repository.



## 15 1 Introduction

Ground-based soil moisture measurements of land surface variables are indispensable in the process of developing, validating, and advancing spatially contiguous data sets derived from satellites or models (Loew et al., 2017; Gruber et al., 2020). Although the first systematic measurements of soil moisture started well before the satellite era in the former Soviet Union to support agricultural decision making (Robock et al., 2000), it was not until the early 2000s that soil moisture monitoring networks started being widely established as part of hydrological and meteorological observing capacities. Particularly, the launch of the Soil Moisture Ocean Salinity (SMOS) mission of the European Space Agency (ESA) in 2009 (Kerr et al., 2016), and the launch of the Soil Moisture Active Passive (SMAP) mission of the National Aeronautics and Space Administration (NASA) in 2015 (Entekhabi et al., 2010), boosted the establishment of new research networks (Colliander et al., 2017).

While all networks are a valuable asset for assessing the skill of soil moisture products under various conditions and scales, their usage is hampered by the diversity of sensors, data formats, quality control, and accessibility mechanisms. The need of bringing together and harmonising available soil moisture data was recognised by the international soil moisture community and expedited by the Global Energy and Water cycle Exchanges (GEWEX) project of the World Climate Research Programme (WCRP) with support of the Commission on Earth Observation Satellites (CEOS), the Global Climate Observing System (GCOS), and the Group on Earth Observation (GEO). In the advent of the SMOS mission, ESA decided to provide the financial impetus to establish a global reference database of in situ soil moisture measurements for the purpose of satellite product development and validation. As a result, the International Soil Moisture Network (ISMN) went online in 2010 (Dorigo et al., 2011b, a).

The primary objective of the ISMN is to collect in situ soil moisture data sets shared by various data organisations on a voluntary basis and make them available in a harmonized format through a centralised free and open web portal (ismn.earth). While 10 years after its launch the core objective of the ISMN remains valid, its functionality has expanded since then. This new functionality includes the integration of advanced quality control methods (Dorigo et al., 2013), the provision of additional metadata and ancillary variables (e.g., precipitation, soil and air temperature), ongoing automation, the provision of software code to users, and the implementation of various tools to promote the information exchange between users, the ISMN, and the data providers. Moreover, the ISMN has substantially grown in terms of networks, stations, and data sets.

Data from the ISMN has supported hundreds of scientific papers on soil moisture, satellite product and land surface model validation in particular (e.g., Al-Yaari et al. (2019b); Brocca et al. (2014a); Beck et al. (2020)). Several operational data producing services routinely access the ISMN data for repeated quality assurance, including ESA's Climate Change Initiative (Dorigo et al., 2017), the Copernicus Global Land Service (Bauer-Marschallinger et al., 2018), and the Copernicus Climate Change Service (C3S; Dorigo et al. (2017)). Other domains have also exploited the ISMN data, e.g. in meteorology, drought monitoring, or land-atmosphere coupling (Section 4).

Despite the valuable contribution of the ISMN to satellite and climate communities, multiple challenges have yet to be mastered, including the heterogeneous availability in space and time (Dorigo et al., 2015), scale differences between in situ measurements and satellite or model sampling (Gruber et al., 2013), full characterisation and traceability of uncertainties,



and differences in spatial-temporal support of the observations caused by different measurement techniques and landscape heterogeneity (Ochsner et al., 2013). New scientific avenues to improve the spatial coverage could be the inclusion of soil moisture data sets from low cost sensors collected by citizens. For climate applications, stable long-term reference data are required, calling for the coordinated establishment and maintenance of Fiducial Reference Measurement (FRM) stations, as outlined by the GEO/CEOS Quality Assurance framework for Earth Observation (QA4EO) (GCOS, 2016; Montzka et al., 2020; Gruber et al., 2020).

The scope of this paper is to inform readers about the evolution of the ISMN over the past decade, including a description of network and data set updates, new quality control procedures, and new functionality of the data portal. We also review scientific literature making use of ISMN data to assess the achievements facilitated by the ISMN and to identify current limitations in data availability and functionality, and challenges in data provision and use. Based on this review, prerequisites and priorities needed to ensure another decade of this unique community-based data repository are defined.


## 2 The ISMN data hosting facility

Although the ISMN may be considered a mere data repository, there is much more to it: its core functionality includes collecting data from participating data providing networks, harmonizing them in terms of units, sampling rates, naming, and metadata, performing automated quality control, storing the data and metadata in a queryable database, and making them available through a web interface. And from a system perspective, it entails even more, e.g., communication with (potential) data providers and users (Appendix C1).

### 2.1 Data and metadata summary

Currently, the ISMN contains 65 networks providing access to a total of 2678 stations, approximately 10,000 soil moisture data sets, and an additional 10,000 data sets of other meteorological variables (collocated with the soil moisture measurements). Although the ISMN is a global network, most networks and stations are located in North America, Europe, Australia and Asia (Figure 1). New networks continuously being added, while many existing networks are still being upgraded with additional stations, soil moisture sensors, and meteorological variables. The diversity of networks is large, ranging from networks with a single station to networks comprising more than 400 stations covering different landscape types as well as periods.


Most of the networks originate from scientific initiatives in various disciplines (e.g., remote sensing, soil sciences, agriculture, meteorology), and only a few are run by operational entities like national weather or environmental services. Consequently, a lack of sustainable project funding has forced several scientific networks to close after some time. 17 out of the 65 networks contained in the ISMN have become inactive and will no longer provide data set updates (Figure 2). Data go back as far as 1952 but none of the data sets spans the entire 70 year period. The longest available time series (40 years) is provided by the RUSWET-AGRO network in the former Soviet Union, while the longest operating network still active is SNOTEL in the United States.



**Figure 1.** Locations of ISMN networks and sites plotted with the ISMN package described in the *Code and data availability* section [Status December 2020].



80 Most networks provide data set updates at yearly or irregular intervals. data sets from six networks (COSMOS, FMI, SCAN,  
 SNOTEL, USCRN, WEGENERNET), comprising approximately 900 stations, are updated in near-real-time (NRT; status  
 December 2020), which is currently defined as once per day. While the earliest networks were sampled manually at weekly,  
 fortnightly, or even monthly intervals, most current networks take their measurements using electronic devices at daily, hourly  
 or even more frequent sampling rates. For more details on the networks see Appendix A, Dorigo et al. (2011b), Dorigo et al.  
 85 (2013) or the individual references herein.



**Figure 2.** Overview of all available networks, the individual time span of the data availability within the ISMN, and their operational status (Status December 2020).





The variables contained in the ISMN (Table 1 and 2) originate from networks that were built for various purposes, which consequently do not all contribute the same information. Since the ISMN was initially founded as a validation database for satellite (surface) soil moisture data, each station in the database provides soil moisture for the upper soil layer ( $\leq 0.10$  meter depth). Soil moisture data sets in the ISMN can go as deep as two metres, but generally with a decreasing number of measurements locations with depth (Table D1). Some stations deploy more than one sensor at a certain depth, either as replacement in case of failure of one of the sensors or to characterise local soil moisture variability. The availability of meteorological data like precipitation, soil and air temperature, is even more heterogeneous, depending on the scope of the network or the data sharing policy of the data providing organisation. It is also quite common that single time series in the database are composed of the consecutive measurements of two or more different sensors, when a sensor is replaced after failure.

**Table 1.** Overview of all available temporally dynamic variables stored in the ISMN database. \*Note that for precipitation and air temperature the measurement height above the ground surface is indicated.

variable name	abbreviation	units	measurement depth* [m]	variable with depth?
soil moisture	sm	$\text{m}^3\text{m}^{-3}$	0.00 - 2.10	Y
soil suction	su	kPa	0.04 - 0.75	Y
soil temperature	ts	$^{\circ}\text{C}$	0.00 - 2.03	Y
air temperature	ta	$^{\circ}\text{C}$	2.00 - 12.00	Y
surface temperature	tsf	$^{\circ}\text{C}$	0.00 - 0.00	N
precipitation	p	mm	0.00 - 2.00	N
snow depth	sd	mm	0.00	N
snow water equivalent	sweq	mm	0.00	N

**Table 2.** Overview of all available temporally static variables stored in the ISMN database.

variable name	abbreviation	units	measurement depth [m]	sensors dependency
climate classification	clcl	none	none	N
land cover classification	lccl	none	on surface and above	N
soil classification	socl	none	none	N
bulk density	bd	$\text{g cm}^{-3}$	0.00 - 1.50	Y
sand fraction	sa	% weight	0.00 - 1.50	Y
silt fraction	si	% weight	0.00 - 1.50	Y
clay fraction	cl	% weight	0.00 - 1.50	Y
organic carbon	oc	% weight	0.00 - 1.00	Y
saturation	sat	% volume	0.00 - 1.00	Y

Table – continued on next page



**Table 2.** Overview of all available temporally static variables stored in the ISMN database.

variable name	abbreviation	units	measurement depth [m]	sensors dependency
field capacity	fc	% volume	0.00 - 2.00	Y
potential plant available water	ppaw	% volume	0.00 - 2.00	Y
permanent wilting point	wp	% volume	0.00 - 2.00	Y

95 Metadata information can be divided into two categories, i.e., mandatory metadata, which allow for an unambiguous identification of each network, station, and measurement in the ISMN database (Figure E1), and optional metadata, shared by data providers to allow more in-depth analysis of their data sets. To be consistent between sites, the mandatory variables climate, land cover, and soil characteristics are taken from external databases: Climate classification is taken from the Köppen-Geiger database with a resolution of 0.1° (Peel et al., 2007); Dynamically evolving land cover for 2000, 2005, and 2010 is obtained  
 100 from ESA's Climate Change Initiative (CCI) land cover v1.6.1 with 300 meter resolution; Soil information is retrieved from the Harmonized World Soil Database (HWSD v1.1; FAO/IIASA/ISRIC/ISS-CAS/JRC, 2009) with a 30" sampling, although the actual resolution may strongly vary from location to location.

If available, data providers can optionally share their own, more detailed, characterisations of land cover, soil, and quality flags with the ISMN. These are stored in addition to the same variables from external sources. All static variables per  
 105 measurement site and depth are listed in Table 2.

## 2.2 Data collection and harmonisation

Data collection is done either manually (mostly by email) or automated, depending on the degree of automation at the data provider side. Although standards for in situ soil moisture data collection are available (Colliander et al., 2017; Montzka et al., 2020), there is no general agreement within the community, neither is it prescriptive for participation in the ISMN. Thus, the  
 110 data being contributed to the ISMN are heterogeneous with regard to units (e.g., volumetric soil moisture, water depth, mass, soil saturation, plant available water), installation depth, integration length, and positioning of the sensors (vertical, horizontal), the metrical system, the sampling interval, and the time zones used for the measurement time stamps.

The first harmonisation step for all data and metadata involves the conversion of units to internationally agreed scientific units (e.g., m and °C). Then, following the recommendations of the World Meteorological Organization for weather observation  
 115 and forecasting (Williams, 2010), all data are resampled to hourly UTC reference time steps. Data that are available at higher sampling rates are thinned by selecting the individual measurements at the hourly UTC reference time step or the ones that are closest in time within a window of +/- 0.5 h. If there is no measurement available within this interval, the respective time step is not stored in the database but can be recreated and filled with a no-data value upon download. The temporal resampling scheme is valid for all included dynamic variables, except precipitation. Since precipitation is a flux and not a state variable,  
 120 all measurements within the hourly interval are summed to represent the full amount of precipitation within the preceding hour (Dorigo et al., 2011b).





All soil moisture measurements provided to the ISMN are converted to volumetric soil moisture in  $\text{m}^3\text{m}^{-3}$ . Since the majority of networks already shares their measurements in volumetric soil moisture units often no unit conversion is needed. For the other (mostly historical) networks, measurements are converted to volumetric units using metadata on soil properties (in case measurements are provided as saturation level or plant available water) and/or the thickness of the soil layer represented by the measurement (in case measurements are provided as water height or mass) (Dorigo et al., 2011b).

The harmonisation of measurement depths is particularly challenging, as different networks adopt different measurement depths, similar sensors are positioned differently (horizontally vs. vertically), or their measurements represent different observation volumes, which may even differ according to the soil wetness (as for cosmic ray probes). Thus, harmonising soil moisture measurements to one or several reference depths would require either assumptions on the measurements and soil properties or supplemental soil modelling. Additionally, since there are lots of potential uses for the data, there is no common agreement on the optimum sampling depths. For example, satellite calibration-validation generally requires observations of the 0-5 cm layer while land surface model evaluation requires reference measurements that are representative for the layers defined in the model (Dorigo et al., 2011b). Consequently, the ISMN does not harmonise measurement depths.

After data harmonisation, the data sets are submitted to extensive quality control procedures (Section 3. After quality control, all data sets of soil moisture and other variables, metadata information on networks, responsible organisations, sites, sensors as and static soil attributes for each station are stored in a relational database.

### 2.3 Data portal

The ISMN can be accessed at [ismn.earth](http://ismn.earth) and consists of a project website containing, e.g., information about networks, data, quality control, and partners, and a data interface where users can view, query, and download the data contained in the database.

The data interface displays the location and information of networks and single stations, and allows for plotting the available data to gain an impression of data availability and quality. Data can be selected for time period, area, single networks, or entire continents. Alternatively, the data download can be selected via an advanced SQL query, which allows users to make more specific selections (e.g., for sensor brand, or a certain depth range).

The selected data are directly extracted from the database and downloads are organised per network. For each network, the download contains (i) the measurements and their quality flags, (ii) information about the file data organisation, (iii) a description of the ISMN quality flags, (iv) a metadata file compliant with ISO 19115 and INSPIRE (Infrastructure for Spatial Information in the European Community) metadata standards, and (v) information about static site characteristics (e.g., land cover, climate class, soil characteristics).

The data set files are formatted according to the CEOP (Coordinated Energy and water cycle Observations Project) (Williams, 2010) standard, or a slightly modified version of it, in order to improve data usability (Dorigo et al., 2011b). These standards use the ascii format but a NetCDF format is foreseen for the near future.



## 2.4 Data provider and user involvement

The ISMN is entirely built on the voluntary, free-of-charge contributions from scientific and operational providers. This prevents the ISMN from being too prescriptive towards the data providers in terms of delivery intervals, automation and formatting. Hence, a careful balance is needed between inclusiveness on the one hand and data quality standards on the other. The ISMN facilitates between users and data providers by reporting data quality issues and user feedback to the providers very six months. This is done by means of a report on data usage statistics for each individual network, e.g., on the number of downloads, the usage of their data in scientific publications, and flagging statistics. Together with obtaining visibility and citations, obtaining feedback on data usage and data quality is one of the primary motivations for data providers to join the ISMN.

More than 3200 active users have registered since 2009 (status December 2020). Data download is free of charge but user registration (compliant with the latest GDPR privacy standards) is required to prevent misuse of the data and to track (undisclosed) statistics on data usage for the half-yearly provider reports.

News feeds on the ISMN web page and a bi-annual newsletter informs the users about new networks, new data sets, data quality issues, important publications, workshops, and more. In addition, a dedicated forum and classical email exchange allow users to raise and get response to issues. Moreover, open source software packages are available for reading and plotting the data (Section 6).


## 3 Quality control

### 3.1 Quality flagging methodology

The wide variety of sensors types and installations, measurement protocols, calibration methods, preprocessing, and quality control procedures adopted by the data providers result in data sets with large differences in quality and filtering. In an attempt to harmonise the reliability of the data from different networks and sensors and to allow for marking spurious observations in near-real-time, the ISMN has adopted automated quality procedures, which are applied to all observations feeding into the ISMN (Dorigo et al., 2013). It uses several approaches to detect dubious soil moisture measurements, which can be subdivided into geophysical dynamic range verification, geophysical consistency methods, and spectrum-based methods (Table 3). While the first category of methods applies simple threshold checks directly to the measurements, the geophysical consistency methods make use of ancillary data, either observed in situ at the same site or derived from model data (i.e., GLDAS-Noah). The spectrum-based flags are the result of a series of conditions applied to the soil moisture measurement time series and their first and second derivatives. The geophysical consistency and spectrum-based methods are only applied to the soil moisture observations, while geophysical dynamic range verification is applied to all dynamic variables in the database.

Recently, a few refinements of the original flagging procedures as described and assessed in Dorigo et al. (2013) were implemented to increase the flagging accuracy:

- The outlier detection method (Flag D06) now allows spikes to last two consecutive time steps instead of the initial one hour. This occurs when all conditions of Eq.[4-6] in Dorigo et al. (2013) are met, but the peak value remains unchanged



**Figure 3.** Additionally flagged observations by network shown as fraction of total number of observations. Note that, in principle, the cumulative sum of fractions can be >1 as the occurrence of each flag ranges between 0 and 1.

185 for an additional hour. The overall impact is small (flagging numbers increase from 0.30% to 0.33%) but its impact on extreme values can be significant.

- Flag D07 ("negative breaks (drops)") was extended with an extra possibility, which detects drops from values greater than 0.05 to exactly 0 m<sup>3</sup>m<sup>-3</sup> soil moisture:

$$x_t \geq 0.05 \wedge x_{t+1} = 0 \quad (1)$$

190 Since a spurious soil moisture drop is precondition for a low plateau (D09; "constant low values following negative break"), also the latter is affected. Adding the extra drop detection increased flagging numbers for D07 from 0.01 to 0.03% and for D09 from 0.9 to 1.1%.

- In case more than one soil temperature, air temperature, or precipitation sensor is available at a site, a flag is raised when at least one of them indicates a spurious observation, which leads to a small overall increase in flags D01, D02 and D04 (< 0.7%).

195



All quality control procedures adopted by the ISMN have been made available under the open source license agreement (see Section 6, <https://github.com/TUW-GEO/flagit>).

We assessed the refined flagging procedures by applying them to 10 networks with hourly data that include stations with multiple soil temperature, air temperature, or precipitation sensors. Despite the very low overall impact of the refined flags, for some networks they are substantial (Figure 3).

Measurements that are detected as erroneous by the quality control procedures are not deleted from the database, but tagged as such (Table 3). The flag is provided as additional attribute ("ISMN Quality Flag") to the observation upon download and can take one of the main categories C (exceeding plausible geophysical range), D (questionable/dubious), M (missing) or G (good). The D flag is raised when either a geophysical-consistency or a spectrum-based check is positive. An additional number indicates the actual cause for flagging (Table 3). A soil moisture observation may receive multiple C- and D-type flags but the good and missing flags are exclusive. Seven networks provide their own soil moisture quality flags, which are added to the ISMN database in addition to the ISMN flags common to all time series. Examples are flags for data quality (without further methodological description) or simple thresholds. For instance, RISMA tags soil moisture observations for frozen soils when the average temperature of two adjacent soil layers is below 0°C (Pacheco et al., 2014). By contrast, the ISMN flag D01 "Soil temperature <0°C" only considers the corresponding depth. For RISMA, the frozen soil flags provided by the network and those computed by the ISMN (D01 "soil temperature <0°C") agree for 87.8%.

### 3.2 Flagging occurrence

The most commonly raised flag is when one of the ancillary temperature observations, i.e., in situ soil temperature (D01), in situ air temperature (D02), or GLDAS soil temperature (D03) is <0°C (Table 3; Figure 4). Since in situ temperature measurements are not available for all networks and to keep consistency between networks, flags D01 and D02 are not shown in Figure 4. The number of observations flagged as frozen soil are not an indicator of the site in general but show which networks are located in areas with a pronounced cold season, e.g., stations from the FMI, RISMA, MAQU, SCAN, and SNOTEL networks.

The second-most common flag is C03 "soil moisture above the site-specific saturation point", which is computed from the HWSD soil properties. The site-specific saturation point is usually lower than the globally adopted less conservative threshold of 0.6 m<sup>3</sup>/m<sup>3</sup> (flag C02) and thus raised more often (Figure 4). However, the HWSD soil properties are uncertain and consequently the C03 flag should be considered as indicative rather than as absolute quality indicator. Values exceeding the saturation point are often an indication of calibration biases or atypical site conditions. For example, the large number of C03 flags obtained for BIEBRZA-S-1 network is because it is installed in a temporarily flooded marshlands with peat porosity exceeding 80%.


Constant values as a result of saturation plateaus (D10) or after a negative break (D09) are the most common spectrum-based flags (Figure 5). The latter are often a sign of sensor drop outs and mostly limited to the GROW, ARM, HYDROL-NET-PERUGIA, and USDA-ARS networks. For example, Xaver et al. (2020) evaluated the sensors used in the GROW network and found occasional drops in soil moisture to zero, which may be the result of corroding contacts. Spikes and breaks are by nature




**Table 3.** Occurrence of flags for all variables and measurements contained in the ISMN (Status December 2020). The soil moisture flags are not exclusive, i.e., an observation can be tagged with multiple flags.

Variable	Flag	Type	Description	Occurrence [%]
Soil moisture	C01	range verification	soil moisture $< 0 \text{ m}^3/\text{m}^3$	0.05
	C02	range verification	soil moisture $> 0.60 \text{ m}^3/\text{m}^3$	1.17
	C03	range verification	soil moisture $>$ saturation point (based on HWSD)	3.99
	D01	geophysical consistency	negative soil temperature (in situ)	6.39
	D02	geophysical consistency	negative air temperature (in situ)	16.57
	D03	geophysical consistency	negative soil temperature (GLDAS)	4.53
	D04	geophysical consistency	rise in soil moisture without precipitation (in situ)	0.28
	D05	geophysical consistency	rise in soil moisture without precipitation (GLDAS)	0.26
	D06	spectrum based	spikes	0.20
	D07	spectrum based	negative breaks (drops)	0.02
Soil temperature	D08	spectrum based	positive breaks (jumps)	0.01
	D09	spectrum based	constant low values following negative break	0.40
	D10	spectrum based	saturated plateaus	2.01
	C01	range verification	soil temperature $< -60^\circ\text{C}$	0.07
	C02	range verification	soil temperature $> 60^\circ\text{C}$	0.17
Soil surface temperature	C01	range verification	soil surface temperature $< -60^\circ\text{C}$	0.01
	C02	range verification	soil surface temperature $> 60^\circ\text{C}$	0.09
Air temperature	C01	range verification	air temperature $< -60^\circ\text{C}$	0.03
	C02	range verification	air temperature $> 60^\circ\text{C}$	0.04
Precipitation	C01	range verification	precipitation $< 0 \text{ mm h}^{-1}$	0.08
	C02	range verification	precipitation $> 100 \text{ mm h}^{-1}$	0.24
Soil suction	C01	range verification	soil suction $< 0 \text{ kPa}$	0.26
	C02	range verification	soil suction $> 2500 \text{ kPa}$	0.00
Snow water equivalent	C01	range verification	snow water equivalent $< 0 \text{ mm}$	20.37
	C02	range verification	snow water equivalent $> 10000 \text{ mm}$	0.01
Snow depth	C01	range verification	snow depth $< 0 \text{ mm}$	19.23
	C02	range verification	snow depth $> 10000 \text{ mm}$	0.00

isolated events that do not occur over an extended period of time and thus appear less frequent in the flagging statistics. The relatively large number of spikes for the network SNOTEL of 0.4% is due to some extremely noisy time series.



**Figure 4.** Fractions of geophysical dynamic range and consistency quality flags per network. Note that, in principle, the cumulative sum of fractions can be >1 as occurrence of each flag ranges between 0 and 1.



**Figure 5.** Fractions of spectrum-based quality flags per network. Notice that, in principle, the cumulative sum of fractions can be >1 as occurrence of each flag ranges between 0 and 1.





**Table 4.** Global validation results of surface soil moisture of ERA and ESA CCI Soil moisture against ISMN (masked and unmasked for quality flags) for the period 2001-2019; the results for ESA CCI were produced with the QA4SM validation service (<https://qa4sm.eu/> Results accessible at: [doi.org/10.5281/zenodo.4288919](https://doi.org/10.5281/zenodo.4288919) (absolute values - no flags); [doi.org/10.5281/zenodo.4288921](https://doi.org/10.5281/zenodo.4288921) (absolute values - with flags); [doi.org/10.5281/zenodo.4288915](https://doi.org/10.5281/zenodo.4288915) (anomalies - no flags); [doi.org/10.5281/zenodo.4288913](https://doi.org/10.5281/zenodo.4288913) (anomalies - with flags). ).

Type	Mask	# time series [ $\bar{x}$ ]	R Pearson [ $\bar{x}$ ]	R Spearman [ $\bar{x}$ ]	ubRMSD [ $\bar{x}$ ]
ERA5 absolute values	None	7822	0.53	0.55	2.06
	Flagged values	5878	0.59	0.60	0.08
ERA5 anomalies	None	7822	0.44	0.46	1.21
	Flagged values	5878	0.48	0.49	0.05
CCI absolute values	None	1178	0.46	0.47	2.98
	Flagged values	1115	0.46	0.46	0.09
CCI anomalies	None	1178	0.34	0.34	1.60
	Flagged values	1115	0.36	0.35	0.06

### 3.3 Effect of flagging on applications

For a selected ISMN site (SCAN, Mayday station), Dorigo et al. (2013) showed that masking flagged values has a small but positive impact on the validation of GLDAS-Noah v1 modelled surface soil moisture and the remotely sensed VUA-NASA AMSR-E soil moisture product. Here, we performed a more extensive assessment of the impact of excluding automatically  
 235 detected spurious observations by the revised flagging methods (Section 3.1) by using ISMN observations available in the period 2001-2019 to validate both ERA5 top layer (0-0.07 m) water content (Hersbach et al., 2020) and ESA CCI Soil Moisture (v5.2; Gruber et al. (2019b), Dorigo et al. (2017), Gruber et al. (2017)). While the impact of flagging is positive for temporal agreement (R Pearson and R Spearman) between the ISMN and ERA5, the effect is negligible for ESA CCI (Table 4). On the other hand, the ISMN flagging reduces the ubRSMD for both comparisons. The benefit of excluding spurious values is also  
 240 obvious in Figure G1a where points are located below the 1:1 line. Again, the benefit is less clear for ESA CCI (Figure G1c). Although validations of a satellite and a model-based are not directly comparable, one reason for the different impact is that the ESA CCI retrievals were already flagged in the production process for values outside valid geophysical range, inconsistencies, dense vegetation, freezing and snow-cover, while this is not the case for the ERA5 model data. Consequently, a positive effect of the ISMN flags is more effective for data sets that were not *a priori* masked than for data sets that were already filtered for  
 245 spurious observations.




### 3.4 Other quality indicators

The automated quality control algorithms offer insight into the quality of the respective measurement time series but not necessarily of the usability of the data sets for specific applications. Gruber et al. (2013) adopted a triple collocation approach using soil moisture data from the ISMN, ESA CCI SM v0.1 (Liu et al., 2012b), and ERA-interim land surface model (Dee et al., 2011) to characterise the representativeness errors of ISMN data for coarse-scale ( $\sim 25$  km) use. Here, we apply a similar approach to estimate the representativeness errors of the ISMN data of all networks with sufficient sampling in the period 2001-2019 for application at the coarse scale. ESA CCI SM passive soil moisture (v5.2, Gruber et al. (2019b), Dorigo et al. (2017)) and top-layer ERA5 volumetric soil water content (Hersbach et al., 2020) are used to complement the triplets. Spatial collocation is carried out using a nearest neighbor method with a maximum distance of 30 km, while the temporal collocation uses a maximum time difference of 1:20 h between the triplets. An exception was solely made for the PBO-H<sub>2</sub>O network because its observations are provided daily at 12:00 UTC, while ESA CCI SM is given at 00:00 UTC daily. All measurements that cover at least partly the 0.00-0.07 m depth interval are used.

The results for different networks are quite diverse (Figure 6). For example, the spread of errors is relatively large for ORACLE but small for others (e.g., AMMA-CATCH or SKKU). The median errors per network vary between 0.03 and 0.05  $m^3 m^{-3}$  with some outliers in both directions. Note that the triple collocation analysis estimates the combined representativeness and sensor errors, although the latter are assumed small compared to the natural spatial-temporal variability Gruber et al. (2013).

There is a clear trend of decreasing mean errors with increasing sensor depths (Figure 7a), which is likely due to a reduction of high-frequency signals and sensor perturbations with depth. Note, that this decreasing trend is observed despite the increasing discrepancy in depth support between the ISMN data and the two surface soil moisture data sets (ERA5 and ESA CCI SM). Thus, theoretically, in situ representativeness errors are expected to be even lower than computed. A potential explanation for the slight increase in the median error for the deepest layer (100-255 cm) may be small sample size and the poor soil parameterisation of the land surface model at this depth. Similar patterns were observed by Gruber et al. (2013).

Concerning the sensors used, there is a large spread in computed representativeness errors for TDR and capacitance sensors. While the costly hygrometric sensors have the lowest mean error, the mean error of resistance probes is the highest (Figure 7b). Since cosmic ray and GNSS/GPS reflectometry sensors integrate over larger horizontal and vertical domains, one would expect lower representativeness errors for these sensors compared to the point observations. However, this is not confirmed by our triple collocation results. Possibly, the advantage of the larger spatial support of these systems is counteracted by their lower signal-to-noise ratio. Error information at the site-, sensor-, or data set level is currently not routinely available for the stations in the ISMN, but would be required for a proper weighting of individual stations in large-scale validations (Gruber et al., 2018, 2019b).




**Figure 6.** Representativeness errors of ISMN networks calculated with triple collocation analysis using top-layer ERA5 volumetric soil water and ESA CCI SM v05.2 passive product. Values in brackets show the average number of triplets per time series and the total number of sensors, respectively, for each network. Following Dorigo et al. (2010) we only used triplets with a Spearman correlation  $>0.2$  between each respective data set pair in the calculation.


## 4 Impact of the ISMN on Earth system sciences

### 4.1 User uptake

As mentioned earlier, over 3200 users have registered to the ISMN while 20-30 new users register each month. Most users are based in the US, China, India and Europe (Figure 8). When asked for the intended use of the data, the four main GEO benefit areas are water, disaster, agriculture, and climate sectors, all with a similar share between 15-20%. Most users come from non-profit organisations (32%), research organisations (30%), and higher or secondary education (28%). Only few users come from public bodies or private companies (both 5%).



**Figure 7.** Representativeness errors for different sensor depths (a) and sensor types (b) derived with triple collocation. Summary of triple collocation characterization of random errors per sensor type. Values in brackets show the total number of sensors for each depth interval and type respectively.



**Figure 8.** Number of users per country (status December 2020).



**Table 5.** Overview of purposes ISMN data are used for in scientific studies (status September 2020; n=288).

Purpose	%
Satellite Validation	53.1
Model development and validation	14.8
Meteorological applications	8.9
Drought monitoring	4.2
Other applications	19.0

The large uptake of the ISMN for soil moisture studies is particularly due to the simplicity of accessing and using multiple data sets from a wide variety of networks. Initially, the ISMN was established to facilitate the calibration and validation of SMOS-based soil moisture products (Dorigo et al., 2011b) and still, satellite soil moisture calibration, validation, and algorithm improvement are the primary applications served by the ISMN (Table 5). In the following, we discuss the major purposes the ISMN has been used for, based on a comprehensive review of all studies that used and cited the ISMN in peer-reviewed scientific publications.

## 4.2 Satellite product evaluation and development

### 4.2.1 Scientific studies

Soil moisture measurements from the ISMN have been widely used as reference data set for the development and evaluation of satellite soil moisture products, mostly global coarse-scale surface soil moisture products (Table F1; n=190). Although initially the goal was to support algorithm development and validation of the SMOS satellite, which was indeed the case for 79 studies, many other satellite missions profited from the ISMN data, most notably ASCAT (n=45), AMSR-E (n=44), SMAP (n=39), and ESA CCI (n=22). Fortuitously, the data have also been discovered for the evaluation of soil moisture products from less used sensors, including the Chinese Feng-Yun 3B, HY-2 and Gaofen-1 satellites (Parinussa et al., 2014a, 2018; Zhao et al., 2014; Xing et al., 2017), MSG SIVIRI (Leng et al., 2015, 2017), MODIS (Gumbricht et al., 2017; Gumbricht, 2018), Aquarius (González-Zamora et al., 2016), and Landsat (Zhao et al., 2017; Pradhan, 2019).

Recently, the ISMN was recognised as validation source for testing algorithms to derive soil moisture from Global Navigation Satellite Systems (GNSS; e.g., Kim and Lakshmi (2018); Chew and Small (2020)). Similarly, there is an increasing trend in the use of ISMN data for the validation of novel high-resolution satellite soil moisture products, which either downscale coarse-resolution products through the use of other finer resolution satellite or ancillary data (e.g., Sheng et al. (2019); Helgert and Khodayar (2020)), or directly derive soil moisture from high-resolution Synthetic Aperture Aperture satellites like Sentinel-1 (e.g., Rodionova (2019b); Foucras et al. (2020)). It should be noted that the ISMN and its contributing networks are mostly designed for analysing time series, thus lacking reference data to assess spatial patterns in the data, particularly in high-resolution products (de Jeu and Dorigo, 2016).



Soil moisture measurements from the ISMN have been used as training set for various data-driven approaches. In situ observations were ingested into machine learning frameworks together with several ancillary predictor variables, either to simulate soil moisture at very high spatial resolution (Zappa et al., 2019) or to create long-term records (O and Orth, 2020). Large-scale monitoring networks are necessary to build reliable models for spatially-wide analysis, while dense networks are ideal for accurate localized models (Senanayake et al., 2019; Abbaszadeh et al., 2019).

Usage-oriented evaluation studies have focussed on the intercomparison of multiple coarse-scale satellite products, using the ISMN data as a reference, either to select the best performing sensor for a specific application or geographic region (e.g., Beck et al. (2020); Karthikeyan et al. (2017)) or to combine them in an optimal way to build a superior product (e.g., Liu et al. (2011); Hagan et al. (2020)).

Since direct satellite observations of soil moisture are only possible for the surface layer, most studies concentrate on the evaluation of surface soil moisture. Yet, various studies also focus on products derived from surface soil moisture that represent moisture in deeper (root-zone) layers, either through exponential filtering (e.g., Paulik et al. (2014); Tobin et al. (2017)), empirical models (Pradhan, 2019), or by assimilating them in land surface models (e.g., Pablos et al. (2018); Blyverket et al. (2019)). Occasionally, the ISMN is used for the evaluation of other satellite data sets, e.g. soil temperature data to validate freeze/thaw state (Rautiainen et al., 2016; Hu et al., 2019).

#### 4.2.2 ISMN as part of operational services

Because of its operational nature and advanced quality control procedures, the ISMN has been identified as the primary reference data source for future operational validation systems for global satellite-based soil moisture products (?). But already today, the ISMN is integrative part of several operational satellite soil moisture production chains:

- In 2005, the **Satellite Application Facility on support to operational Hydrology and water management (H SAF)** started to operationally produce and validate precipitation, soil moisture, and snow products from satellites operated by EUMETSAT (Rinollo et al., 2013). ISMN data are used for calibration and validation of various soil moisture products produced in H SAF from the Metop Advanced Scatterometers, including global climate data records and near-real-time products of surface soil moisture, and a root-zone soil moisture product over Europe.
- The **Climate Change Initiative of the European Space Agency (ESA CCI)** uses data from the ISMN to assess each year the quality of new soil moisture climate data record releases and their improvements with respect to forerunner versions (Gruber et al., 2019b; Dorigo et al., 2015). Within ESA CCI, ISMN data were also used to quantify the spatial representativeness of ESA CCI satellite and climate model data sets (Nicolai-Shaw et al., 2015b).
- **C3S** produces authoritative, quality-assured climate data records of soil moisture and other essential climate variables (ECVs). The satellite soil moisture products produced within C3S are routinely updated every 10 days with the latest available satellite observations. C3S uses the ISMN soil moisture data in combination with the metadata provided to categorise product performance per land cover and climate type (Scanlon et al., 2019).





340 – **Copernicus Global Land Service (CGLS)** produces within 1-2 days after satellite overpass soil moisture data sets from Sentinel-1 and from a combination of Sentinel-1 and ASCAT (SCATSAR) (Bauer-Marschallinger et al., 2018, 2019). The latter propagates the surface observations to deeper layers by means of the so-called soil water index (SWI). Both products are provided at 1 km resolution for Europe. ISMN data are used for validating moisture in surface and deeper soil layers.

345 Recently, the **Quality Assurance for Soil Moisture (QA4SM)** service (<https://qa4sm.eu>) was initiated to bring together validation methodologies, community protocols, reference data, and satellite observations to evaluate and intercompare soil moisture data products in a coherent, standardised, and transparent way. In situ data sourced from the ISMN are integral part of this validation system (Scanlon et al., 2019; Gruber et al., 2020).

### 4.3 Model development and validation

350 In situ measurements are the most important reference source when assessing the performance of land surface models, reanalyses products, and hydrological models. Although also satellite observations are a valuable validation source (e.g., Szczypta et al. (2014)), these measure only the upper 5 cm of the soil, and hence do not allow for the validation of deeper layers. Besides, state-of-the-art reanalysis products like ERA5 (Hersbach et al., 2020)) assimilate satellite soil moisture observations, so that for these products in situ data remain the only truly independent validation reference. Since the sampling rate (hourly) of the  
 355 ISMN observations is generally higher, and their spatial support lower than that of most models (Reichle et al., 2004), model evaluation is in principle not limited by the spatial and temporal resolution of the in situ data, although representativeness issues often remain.

In particular for global assessments, the availability of harmonised data over multiple networks makes the ISMN a preferred reference source over data from individual networks (Xia et al., 2019). Table F2 shows that many well-established state-of-  
 360 the-art land surface model and reanalysis products have been evaluated against data from the ISMN, not only by users of these products (e.g. (Li et al., 2020b)), but also by the developer teams themselves, e.g., Reichle et al. (2017) for MERRA2, Albergel et al. (2020) for LDAS-Monde, and Balsamo et al. (2015); Fairbairn et al. (2019) for various ECMWF products.

Also a suite of new products have been assessed against the ISMN, including multi-model ensembles (Schellekens et al., 2017; Cammalleri et al., 2015), data-driven evaporation models (Martens et al., 2017; Lievens et al., 2017) and statistical  
 365 infiltration models Pal and Maity (2019). As for remote sensing products, a trend towards higher-resolution regional to global model-based products can be observed. Among others, ISMN data have been used to validate new high-resolution reanalysis products over the United States (McDonough et al., 2018) and Europe (Naz et al., 2020). Apart from soil moisture also soil temperature data have been drawn from the ISMN for model validation purposes (e.g., Wang et al. (2016); Albergel et al. (2015)).

370 Besides the evaluation of hydrological or land surface model improvements, the ISMN has also frequently served model development in a more fundamental way. For example, Hartmann et al. (2015) developed a large-scale karstic groundwater recharge model over Europe and the Mediterranean and calibrated and evaluated this model with observations of actual evap-



otranspiration from FLUXNET (Baldocchi et al., 2001) and soil water content data from the ISMN. Calvet et al. (2016) used soil moisture and temperature data from the SMOSMANIA network contained in the ISMN to derive pedotransfer functions for soil quartz fraction in southern France. Pal et al. (2016) developed a statistical model to estimate vertical soil moisture profile using SCAN data of the ISMN as source, while on a similar note, Shin et al. (2018) developed a non-parametric evolutionary algorithm to predict soil moisture dynamics using ISMN data over Oklahoma and Illinois. Similarly, Jalilvand et al. (2018) estimated the drainage rate from surface soil moisture drydowns from ISMN data. A bridge between soil moisture and vegetation dynamics was made by Sawada (2018), who used ISMN data to validate a new ecohydrological land reanalysis to better simulate the link between sub-surface soil moisture and vegetation dynamics. Finally, Brocca et al. (2015) developed a water-balance approach to estimate rainfall from soil moisture observations based on reverse modelling and evaluated this at several ISMN sites in Europe. Later refinements of this approach were also evaluated against data from the ISMN (Hoang and Lu, 2019).

#### 4.4 Meteorological applications

Soil moisture from the ISMN has often been used to validate the land surface representations of meteorological forecasting models. However, as meteorological forecasts often rely on the latest generation of land surface models, in practice there is often no strict distinction between meteorological and land surface model development as described in the previous section. Notable examples are the various generations of TESSEL models used both in the Integrated Forecasting Systems and reanalysis products of ECMWF, the development of which greatly profited from soil moisture and temperature data from the ISMN (Albergel et al., 2012a, 2015). Dirmeyer et al. (2016) assessed the skill and soil moisture memory effects of various weather and climate models with ISMN data over the United States. Similarly, Angevine et al. (2014) assessed the soil moisture skill of the Weather Research and Forecasting Model (WRF) (Table F2).

Several studies have used the ISMN data to assess the forecast skill or new implementations of numerical weather prediction models. For example, de Rosnay et al. (2019) and Rodriguez-Fernandez et al. (2019) used in situ soil moisture observations from several ISMN networks to validate the impact of assimilating SMOS brightness temperatures and soil moisture, respectively, to predict soil moisture up to 5 days ahead. Similarly, Lin and Pu (2019, 2020) used ISMN data to examine the impact of MAP soil moisture assimilation in WRF for near-surface short-range weather forecasts. Boussetta et al. (2015) used over 500 ISMN sites to assess the impact of assimilating surface albedo and vegetation states from satellite observations on numerical weather prediction.

From a more methodological, land-atmosphere perspective, Lee (2018) used the ISMN data to study the role of soil moisture in triggering rainfall over West Africa. Vice versa, Zhang et al. (2019c) studied the role of rainfall on soil cooling using data from the SMOSMANIA network in southern France.

#### 4.5 Drought monitoring

In a drought monitoring context, ISMN data have frequently been used in a "convergence of evidence" approach in combination with other drought-related variables or indicators. For example, Scaini et al. (2015) compared the variability of in situ soil



moisture measurements from the ISMN, SMOS surface soil moisture, and two drought indices based on climatic information to study droughts in Spain. Mu et al. (2019) used VNIR satellite data and soil moisture distributed by the ISMN to monitor drought in the southern United States. On a more technical level, Gruber et al. (2018), assessed the use of spatially sparse ISMN in combination with a continuous model for operational agricultural drought monitoring over the United States.

410 ISMN data have also been used to classify new and more "traditional" drought indices, such as the Standardized Precipitation (Evaporation) Index and the Palmer Drought Severity Index (Vicente-Serrano et al., 2012; Krueger et al., 2019). Sadri et al. (2020) used the ISMN data from the RISMA network to assess a global near-real-time soil moisture index monitor for food security using SMOS and SMAP data. Chen et al. (2019) used precipitation distributed through the ISMN to evaluate a drought index derived from Sentinel-2 in Spain. Due to relatively good coverage over Europe, Cammalleri et al.  
 415 (2015) used ISMN data to assess various model soil moisture products as input to the European Drought Observatory (EDO, <http://edo.jrc.ec.europa.eu>). Also Mishra et al. (2018) used data from the ISMN to assess a suite of land surface models used to reconstruct drought events in India since the mid 1900s.

#### 4.6 Other applications

ISMN data have been used for various other purposes, going beyond what the ISMN was originally developed for. In addition  
 420 to supporting satellite and land surface model soil moisture product development, the ISMN has played a fundamental role in the validation of a wide range of hybrid observation-based products, including a Soil Moisture Saturation Index (Campo et al., 2011), apparent thermal inertia surface estimates (Notarnicola et al., 2012), data-driven surface and root-zone soil moisture predictions (Kornelsen and Coulbaly, 2014; Manfreda et al., 2014; Wang et al., 2020a), improved satellite albedo products (Liu et al., 2014), and estimates of effective permittivity and brightness temperature of organic soils Park et al. (2019).

425 The ISMN has frequently been used to assess the impact of assimilating satellite observations into hydrological models (Khaki et al., 2019; Nair et al., 2020a; Gruber et al., 2015; Shin et al., 2016; Li et al., 2020b; Wang et al., 2020b), land surface models (Nair and Indu, 2016; Zhao and Yang, 2018; Nair et al., 2020b) and carbon models (Scholze et al., 2016). On a more methodological level, Zhang et al. (2019a) assessed a new data assimilation scheme against ISMN observations. Gruber et al. (2018) even assimilated spatially sparse ISMN observations directly into a spatially continuous land surface model over the  
 430 Continental US, and tested the preconditions (e.g., requirements on signal-to-noise ratio and number of sites) for having a significant positive impact.

The wealth of data covering a wide range of surface and climate conditions has been frequently exploited to study soil water dynamics at various spatial and temporal scales and their (climatic) drivers (Brocca et al., 2014b; Hirschi et al., 2014; Ojha et al., 2014; Qin et al., 2018; Kumar et al., 2019a; Tian et al., 2019a; Nicolai-Shaw et al., 2015b; Dong et al., 2020; Wang et al.,  
 435 2017; Verrier, 2020; Deng et al., 2020a), flood dynamics (Esposito et al., 2018), or the impact of soil moisture on plant growth (Hottenstein et al., 2015).

The ISMN has been used to develop and test new statistical validation approaches (Zwieback et al. (2013), Gruber et al. (2016), Afshar et al. (2019)). Finally, the standards and quality control procedures adopted by the ISMN have served as a guideline for establishing and benchmarking new networks ((Kim et al., 2019; Skierucha et al., 2012b, a; Petropoulos and



440 McCalmont, 2017)), soil moisture metadatabases (Liao et al., 2019; Xia et al., 2015)), or validation services (Kumar et al., 2012), and to assess or improve alternative sensing techniques and constellations (Xaver et al., 2020; Kapilaratne and Lu, 2017; Nguyen et al., 2017; Mahecha et al., 2017).

## 5 Challenges and opportunities

### 5.1 Scientific challenges

#### 445 5.1.1 Diversity of measurements

The in situ measurements of soil moisture data in the ISMN have been collected by a large variety of sampling techniques. The early networks contained in the ISMN (e.g., RUSWET, CHINA and MONGOLIA) are based on gravimetric sampling, which is still considered the most accurate approach (Romano, 2014). However, it is labor-intensive and invasive and thus measurements are infrequent (weekly or even coarser sampling), every time being taken from a slightly different location, may  
 450 contain systematic errors between sampling dates.

Nowadays, the most commonly used techniques for systematic in situ sampling are based on the contrasting dielectric properties of soil and water and comprise time domain reflectometry (TDR) and frequency domain reflectometry (FDR) (Robinson et al., 2008), e.g., AMMA-CATCH, BNZ-LTER, CARBOAFRICA, FMI, GTK, HYDROL-NET-PERUGIA, LAB-net, OR-  
 455 ACLE, OZNET, SASMAS, SWEX-POLAND, UDC-SMOS, WSMN, UMSOL networks). In particular, capacitance sensors based on FDR are becoming more and more widespread because of their lower cost compared to TDR sensors, despite their lower accuracy (Romano, 2014; Brocca et al., 2017). However, great technical improvements are being achieved with capacitance sensors, thus improving their reliability. Yet, TDR and FDR techniques only provide point measurements, i.e. they are representative of small volumes of soil.

Slightly larger soil volumes (diameter of 15 to 60 cm) are observed with the neutron scattering method, where the density of  
 460 thermal neutrons produced by scattering of fast neutrons on soil hydrogen can be related can be related to soil moisture through a calibration curve (Gardner and Kirkham, 1952; Romano, 2014) (e.g., IOWA network). The cosmic-ray method has been developed based on similar principles of neutron scattering, (Zreda et al., 2012, 2008). Cosmic-ray neutron sensors (CRNS) are located on or above the soil surface and measure and count the number of cosmogenic neutrons in air above the land surface and that are in equilibrium with the soil. CRNS measurements are representative of significantly larger volumes compared to  
 465 neutron probes (i.e., radius of a few 100 metres and depths between 0.12-0.70 m, depending on the soil wetness), and thus can be used to bridge the scales between point observations and coarse resolutions of satellite observations and models. CRNS measurements, which are used in the COSMOS network, are sensitive to vegetation and have relatively high noise levels. To reduce this noise, the data are resampled to daily mean values.

Also the measurements of the PBO\_H2O network, which uses Global Positioning System (GPS) receivers, can be used to  
 470 bridge the scales between point and satellite sensors. GPS sensors, initially used for geophysical and geodetic applications, have been found well suited to measure soil moisture (Larson et al., 2008). The GPS signals are representative for an area



of approximately 300 m<sup>2</sup> and are L-band, hence making them ideal for comparison against satellite missions like SMAP and SMOS (Larson et al., 2008).

As shown, all measurement techniques have their strengths and limitations Bogaen et al. (2015); Dorigo et al. (2011b), which complicates their combined, bulk use. Depending on the application and process scales, the user needs to carefully consider which networks to use or exclude and how to interpret the results obtained. A task of the ISMN could be to translate community-based guidelines (Gruber et al., 2020; Montzka et al., 2020) to recommendations for the use of individual datasets.

### 5.1.2 Spatial representativeness and scaling

Soil moisture is highly variable in space as a result of complex interactions between soil characteristics, topography, vegetation, and meteorological conditions. Depending on the spatial scale considered, the dominant controlling factor(s) can be different (Crow et al., 2012; Western et al., 2002). Hence, validation of satellite-derived products is hindered by the spatial mismatch between ground observations and satellite footprints (Gruber et al., 2020; Molero et al., 2018; Gruber et al., 2013). Ideally, to reduce the representativeness error of in situ references, enough stations should be deployed within a satellite footprint to develop robust areal soil moisture estimates (Brocca et al., 2007; Famiglietti et al., 2008; Colliander et al., 2017). However, this is a costly solution and therefore only a limited number of sites provide such setup (Crow et al., 2012; Colliander et al., 2017). In the ISMN the networks VDS, BIEBRZA\_S-1, RSMN, UDC\_SMOS, LAB-net, FMI, USDA-ARS have been set up in this way. When available, in situ stations within the same satellite pixel should be averaged, either through arithmetic mean or weighted average. Higher weights should be given to stations expected to be more representative of the satellite grid average, e.g., by using Voronoi diagrams (Colliander et al., 2017), the inverse footprint method (Nicolai-Shaw et al., 2015a), the time stability concept (Vachaud et al., 1985), or landscape properties such as land cover and/or soil texture (Bircher et al., 2012). Alternatively, the triple collocation (TC) can be used to quantify and correct for spatial sampling errors of in situ stations (Miralles et al., 2010; Gruber et al., 2013). The resulting pixel-scale soil moisture ground reference, i.e., the averaged value from dense networks with several stations per satellite grid or the original time series in the case of sparse networks with a single station per pixel, should undergo a statistical rescaling (Gruber et al., 2020). Indeed, a direct comparison of in situ and satellite products would be subject to representativeness errors, which may dominate the total soil moisture retrieval errors (Chen et al., 2017; Gruber et al., 2013; Molero et al., 2018). Rescaling accounts for systematic representativeness errors arising from different spatial resolution and different vertical measurement support, i.e., penetration depths of microwave sensors and in situ sensor placement depths (Gruber et al., 2013, 2020) but does not correct for random representativeness errors. One way to address this is the triple collocation described above. Also, systematic representation errors may have a time-varying (e.g., seasonal) component, which, unless explicitly accounted for, may lead to temporally aliased results.

### 5.1.3 Integration of low-cost sensors

The development and use of low-cost sensing technologies, especially in the environmental sciences, has seen a pronounced increase during the last decade. Such a rise is driven by several factors, e.g., the reduced cost of micro-controllers, electric components, and sensors (Mao et al., 2019). Even though a rich variety of low-cost soil moisture sensors based on different



505 measurement principles has been developed (Chawla et al., 2019; González-Teruel et al., 2019; Kumar et al., 2016), capacitance sensors gained most popularity because they are relatively inexpensive, easy to operate, and provide reliable observations (Kojima et al., 2016).

The considerably lower cost of these sensors compared to traditional probes makes them suitable for high-density and/or large-scale monitoring of soil moisture. The possibility to map soil moisture (as well as other environmental variables) with an unprecedented spatial coverage can generate new insights into its dynamics and create new opportunities. For instance, high-density networks of low-cost sensors can be used to reduce spatial representativeness errors by providing numerous observations within a satellite footprint (Teuling et al., 2006). Similarly, one can deploy a temporary low-cost sensors network to identify the most suitable location(s) for long-term monitoring. Such locations could then be equipped with professional sensors, while moving the low-cost network to other sites (Zappa et al., 2019). Another exciting opportunity offered by low-cost sensors is the deployment of networks in low-income countries.

However, a number of practical challenges arise when integrating low-cost sensors in the ISMN. A key aspect to consider is the lifetime: depending on the robustness of the sensor (both of electronic components, such as micro controllers, and sensor housing), reliable measurements can be recorded for a period ranging from a few months to years (Xaver et al., 2020) but even be shorter in extreme environments. Another aspect, particularly affecting automation, is data storage and transmission. Some low-cost sensors allow for wireless communication with a main server ((Bogena et al., 2007), (Majone et al., 2013)), while other sensors have a limited internal storage and data should be collected persistently over time (e.g., every 80 days, Xaver et al. (2020)). Furthermore, it is necessary to assess their accuracy and robustness (Castell et al., 2017). Therefore, low-cost sensors should always undergo thorough evaluation to i) quantify the agreement of low-cost sensor measurements with gravimetric samples and/or professional probes, ideally considering a wide range of soil and climatic conditions, ii) assess the inter-sensor variability, and iii) test the suitability for usage in field conditions (Domínguez-Niño et al., 2019; Kizito et al., 2008; Mittelbach et al., 2012; Adla et al., 2020).

#### 5.1.4 Integration of citizen observations

Citizen science is defined as the involvement of non-experts in collecting data (Bonney et al., 2009). Crowd-sourced measurements of soil moisture are now possible because of the development of low-cost sensors (Section 5.1.3). Crowd-sourcing has the potential to overcome some of the most challenging issues of soil moisture monitoring, such as the use of many sensors to address scaling issues, and the fact that the observations can be carried out anywhere, as long as there are citizens willing to collect the data.

An outstanding example of a citizen science project focusing on soil moisture is the GROW Observatory (growobservatory.org). Within GROW, thousands of low-cost sensors have been distributed to farmers, gardeners, and growers across Europe (Kovács et al., 2019). Focus areas have been identified based on a number of scientific criteria and the presence of active and engaged communities. Within each focus area, covering spatial scales from 20 to 200 km, hundreds of sensors have been distributed. Some sensors served as back-up for potential failures, so that malfunctioning sensors could be promptly replaced enabling long-term continuity of observations. In order to ensure high standards of the measurements, citizens have been





trained in the selection of the correct locations where to install sensors, how to properly install and maintain them through  
 540 field manuals, online courses, meet-ups and remote support (Kovács et al., 2019). Overall, more than 6000 sensors were de-  
 ployed and provided soil moisture measurements across 13 European countries, demonstrating that citizen observatories can  
 be integrated in EO activities and contribute to validation of remotely sensed products (Zappa et al., 2020).

The integration of citizen observations in the ISMN is challenging for multiple reasons. Crucial is the long-term engagement  
 of citizens, which needs to be thoroughly addressed from the early stages of designing a citizen observatory. It is necessary to  
 545 create long-lasting communities that go beyond the duration of the contributory projects (Grainger, 2017). Successful citizen  
 observatories have been those where citizens were able to benefit directly from the data collected by others (Fritz et al., 2017).  
 Additionally, education and resources are essential to boost individual motivation to continuous participation and to sustain re-  
 newal of membership as natural participation cycles change (Zappa et al., 2020). Even if proper training is provided to citizens,  
 uncertainty in the quality of the data is the main limitation for the use of crowd-sourced observations in science (Lukyanenko  
 550 et al., 2020). The reliability of measurements from individual sensors is unknown, because of, e.g., the selection of unsuitable  
 locations, incorrect sensor installation, and existence of defective sensors. However, increasing the number of sensors within the  
 same satellite pixel reduces this uncertainty. Additionally, visual inspection of time series, as well as application of automated  
 quality flagging controls, such as those developed within the ISMN, could be used to mask suspicious observations (Xaver  
 et al., 2020). Overall, it has been shown that well-organized citizen science projects can provide trustworthy contributions to  
 555 the scientific community (Kosmala et al., 2016; Palmer et al., 2017).

## 5.2 Operational challenges

### 5.2.1 Automation

Part of the contributions to the ISMN (6 networks with over 900 stations) are inserted into the ISMN in NRT. This process  
 is fully automated, including data downloaded from data providers, harmonisation, flagging, insertion into the database, and  
 560 updating metadata tables. However, because of different data recording and handling mechanisms at the provider side and  
 differences in data sharing mechanisms and policies the ISMN is also partially manually operated and will probably also have  
 to continue in this way in the future. The differences between the fully automated and the manual approach is confined to data  
 download and data ingestion into the processing chain. Data harmonisation and quality control is automated for all data sets  
 (see Figure C1).

565 A major challenge in the automation process is the enormous heterogeneity of input data formats. Moreover, these change  
 over time for individual networks, stations, and even sensors, as sensors may fail or the method for data logging is changed.  
 Thus, error detection and handling is of utmost importance and frequent adaptation of the system required to cover changes in  
 input data.

A potential way to promote the automated insertion of new data is to allow only data that comply with a strict, yet to be  
 570 defined, data standard. At the same time, this may bear the risk of raising the barrier to contribute to the ISMN too high for  
 several scientists.



### 5.2.2 Including new networks

Since data sharing with the ISMN is entirely built upon a voluntary basis and data usage is open and free, new networks may be reluctant to join. The ISMN is in contact with several network providers who are happy to collaborate but are restrained by data sharing policies, which does not allow data sharing at all or only after a certain time. Furthermore, for governmentally operated networks it is often not allowed to share data as open-access or it is unclear who can make these decisions.

Solving such issues can be supported by collaborative data hosting facilities like the Global Terrestrial Network - Hydrology (GTN-H; [gtn-h.info](http://gtn-h.info)), who have a strong connection to governmental bodies like the International Centre for Water Resources and Global Change under the auspices of UNESCO, the United Nations Environment Programme, the International Science Council, and WMO.

### 5.2.3 Appropriate recognition of data providers

Not all users correctly cite the ISMN and the involved networks, as stated in the terms and conditions for ISMN data use (ISMN). Proper citation of single networks is important for giving data providers the recognition that is required to convince funding agencies to continue supporting the maintenance and operation of these networks. In the end this does not only affect the networks themselves but also the open-access availability of data through the ISMN as a whole. Some network providers have raised their concerns in this matter and, hence, continuous efforts are needed to maintain a strong visibility of network data providers towards users.

### 5.2.4 Transparency and traceability

The ISMN data collection is constantly evolving: New data records are added and existing ones are extended or, if necessary, reprocessed and corrected. Not only the data, but also the underlying code changes. These updates and any retroactive changes made to the data archive are tracked within the ISMN but not yet readily passed on to the users. The updates can lead to differences in analyses on the user side, e.g., when considering obvious changes such as temporal extensions or new stations, and lead to non-reproducible results. Therefore, any update must be traceable and clearly communicated, requiring a system to track and store these changes. Version tracking and digital object identifiers (DOIs) are ways to identify each database access and therefore allow tracing back to past states of the ISMN. Such a mechanism is required to make the ISMN compliant with the FAIR Data Principles (Wilkinson et al., 2016).

## 6 Conclusions

In this study we reviewed the first decade of operations of the ISMN. Besides satisfactorily fulfilling its initial target, i.e., supporting satellite soil moisture product validation and calibration, many additional more or less foreseen uses have emerged. In addition, an increasing number of services and product development chains have routinely included the use of ISMN data in their operational structure. While the development and operations of the ISMN itself have received continuous financial support



from ESA, all network data sets have been freely contributed by dedicated researchers. To guarantee the availability of these resources for climate and environmental monitoring also for the next decade, we plead with governments and international bodies for systematic funding of both the ISMN and its participating data-providing networks.

605 To maximise geographic coverage and data usage, the policy of the ISMN has always been to integrate datasets without strict requirements on sensors, sampling protocol, or data quality. The emerging strongly heterogeneous data set characteristics call for far-going quantification and traceability of errors, from sensor calibration to data download and from the point measurement to the spatial-temporal support of the application. Thus, besides expanding its coverage to data-poor regions and landscapes, the ISMN shall focus the next 10 years of operation on developing methodologies that allow observations to qualify as Fiducial  
610 Reference Measurements for soil moisture applications.

#### *Code and data availability.*

Upon registration, all data and metadata described in this paper can be downloaded for free from <https://ismn.earth>. A python package to read and plot the data and metadata ((TUW-GEO/ismn: v0.4) can be accessed on [doi.org/10.5281/zenodo.3966399](https://doi.org/10.5281/zenodo.3966399).  
615 An example of output of this package is shown in Figure 1. A python package containing the ISMN quality control procedures for in situ soil moisture is available on GitHub (<https://github.com/TUW-GEO/flagit>).

*Author contributions.* Wouter Dorigo conceived the manuscript and wrote it together with Irene Himmelbauer, Daniel Aberer, Lukas Schremmer, Ivana Petrakovic, Luca Zappa, and Wolfgang Preimesberger. All others contributed data to the ISMN and provided feedback on the manuscript

620 *Competing interests.* No competing interests

*Acknowledgements.* The authors greatly acknowledge the financial support provided by ESA through various projects: SMOSnet International Soil Moisture Network (contract No. 4000102722/10/NL/FF/fk and operations contract No. 3-13185/NL/FF/fk); IDEAS+ (contract No. TVUK/AG/18/02082); QA4EO (contract No. TPZV/UK/AG/19/02321). Additional funding for methodological advances has been received from the EU FP7 Earth2Observe project (Contract No. 603608), the EU H2020 GROW project (Contract No. 690199), and the  
625 QA4SM project funded by the Austrian Space Applications Programme 14 (Contract No. 866004). We acknowledge the endorsement of various international bodies, including CEOS, WCRP GEWEX, GCOS, GTN-H, and GEO. We greatly thank all staff members from the participating networks for their continued technical support.



## Appendix A: Network overview

A summary of each network is given in Table A1 while more details are given in the subsequent paragraphs.

**Table A1.** Main properties of networks and data contained in the ISMN.

Network abbreviation	Location	Number of stations	Start date	End date	Status
AACES	Australia	49	2005-05-09	2010-09-24	inactive
AMMA-CATCH	Benin, Niger, Mali	7	2006-01-01	2018-12-31	running
ARM	USA	35	1993-06-29	2020-12-10	running
AWDN	USA	50	1997-12-31	2010-12-30	running
BIEBRZA_S-1	Poland	30	2015-04-23	2018-12-01	running
BNZ-LTER	Alaska	12	1988-06-01	2013-01-01	running
CALABRIA	Italy	5	2001-01-01	2012-12-31	running
CAMPANIA	Italy	2	2000-07-27	2012-12-31	inactive
CARBOAFRICA	Sudan	1	2002-02-08	2010-01-20	running
CHINA	China	40	1981-01-08	1999-12-28	inactive
COSMOS	Worldwide	108	2008-04-28	2020-03-29	running
CTP_SMTMN	China	57	2010-08-01	2016-09-19	running
DAHRA	Senegal	1	2002-07-04	2016-01-01	running
FLUXNET-AMERIFLUX	USA	2	2000-01-01	2020-02-12	running
FMI	Finland	27	2007-01-25	2020-07-04	running
FR_Aqui	France	5	2012-01-01	2020-01-01	running
GROW	Europe	150	2017-02-08	2019-10-08	closed
GTK	Finland	7	2001-05-16	2012-05-29	running
HiWATER_EHWSN	China	174	2012-05-04	2012-09-20	inactive
HOBE	Denmark	32	2009-09-08	2019-03-13	running
HSC_SEOLMACHEON	Korea	1	2008-01-01	2009-01-01	running
HYDROL-NET_PERUGIA	Italy	2	2010-01-01	2013-12-31	running
ICN	USA	19	1983-01-03	2010-11-21	inactive
IIT_KANPUR	India	1	2011-06-16	2012-11-22	running
IMA_CAN1	Italy	12	2011-08-23	2015-12-03	closed
IOWA	USA	6	1972-04-04	1994-11-15	inactive
IPE	Spain	2	2008-04-03	2020-03-25	running
iRON	USA	9	2012-08-21	2019-12-31	running
KHOREZM	Uzbekistan	7	2010-04-17	2011-09-10	inactive



**Table A1 continued from previous page**

Network abbreviation	Location	Number of stations	Start date	End date	Status
KIHS_CMC	South Korea	18	2019-03-20	2019-12-10	running
KIHS_SMC	South Korea	19	2019-03-21	2019-12-05	running
LAB-net	Chile	3	2014-07-18	2017-11-21	running
MAQU	China	20	2008-06-30	2010-07-31	running
METEROBS	Italy	1	2011-10-23	2012-05-09	inactive
MOL-RAO	Germany	2	2003-01-01	2014-01-01	running
MONGOLIA	Mongolia	43	1964-04-08	2002-10-18	inactive
MySMNet	Malaysia	7	2014-05-31	2015-12-31	running
ORACLE	France	6	1985-10-18	2013-09-09	running
OZNET	Australia	38	2001-09-12	2018-08-27	running
PBO_H2O	America, Africa, Asia	158	2004-09-27	2017-12-16	inactive
PTSMN	New Zealand	20	2016-10-30	2018-11-15	running
REMEDHUS	Spain	24	2005-03-15	2020-01-01	running
RISMA	Canada	23	2013-04-24	2020-03-25	running
RSMN	Romania	20	2014-04-09	2020-06-30	running
RUSWET-AGRO	Former Soviet Union	212	1958-04-08	2002-06-28	inactive
RUSWET-GRASS	Former Soviet Union	121	1952-06-08	1985-12-28	inactive
RUSWET-VALDAI	Former Soviet Union	3	1960-01-15	1990-12-15	inactive
SASMAS	Australia	14	2005-12-31	2007-12-31	running
SCAN	USA	239	1996-01-01	2020-07-12	running
SKKU	South Korea	5	2014-05-08	2014-11-19	running
SMOSMANIA	France	22	2007-01-01	2019-01-01	running
SNOTEL	USA	438	1980-10-01	2020-07-12	running
SOILSCAPE	USA	169	2011-08-03	2017-03-29	running
SWEX_POLAND	Poland	6	2000-01-01	2013-05-06	inactive
SW-WHU	China	7	2014-01-12	2015-06-03	running
TAHMO	Africa (Sahel Zone)	70	2015-01-01	2020-10-31	running
TERENO	Germany	5	2009-12-31	2020-07-09	running
UDC_SMOS	Germany	11	2007-11-08	2011-11-18	inactive
UMBRIA	Italy	13	2002-10-09	2017-12-31	partially running
UMSUOL	Italy	1	2009-06-12	2010-09-30	running



**Table A1 continued from previous page**

Network abbreviation	Location	Number of stations	Start date	End date	Status
USCRN	USA	115	2000-11-15	2020-07-11	running
USDA-ARS	USA	4	2002-06-01	2009-07-31	inactive
VAS	Spain	3	2010-01-01	2012-01-01	running
VDS	Myanmar	4	2017-06-01	2020-04-06	running
WEGENERNET	Austria	12	2007-01-01	2020-07-11	running
WSMN	UK	8	2011-09-02	2016-02-29	running

## 630 A1 AACES

AACES stands for the Australian Airborne Cal/val Experiments for SMOS. This campaign network covers a 500 × 100 km<sup>2</sup> study area located in South-East Australia, covering a variety of topography, vegetation- and climate classes (Rüdiger et al., 2007; Peischl et al., 2012). Measurements of soil moisture, soil temperature, and precipitation were taken between May 2009 and September 2010. The AACES calibration and validation campaigns were a temporary project therefore no further data sets  
 635 will be available.

## A2 AMMA-CATCH

The AMMA-CATCH observatory gathered data from densely instrumented mesoscale sites in West Africa (Benin, Niger, Mali). The network is devoted to long-term regional monitoring of global change impacts on the critical zone. Height stations in Benin and four stations in Niger of the network are included in the ISMN from 2006 to present, including surface soil  
 640 moisture and root-zone soil moisture until one-meter depth. For more information, see Galle et al. (2018).

## A3 ARM

The Atmospheric Radiation Measurement (ARM) Program has three soil sensor networks across north-central Oklahoma and southern Kansas in the U.S.: The Soil Water and Temperature System (SWATS) through 2016, and presently the Soil Temperature and Moisture Profile (STAMP) and Surface Energy Balance System (SEBS). The SWATS and STAMP have  
 645 profiles at 5-8 depths up to 175 cm, while the SEBS measure at 2.5 cm. All sites are co-located with a suite of meteorological and radiative measurements available from ARM.gov (Cook, 2016a, b; Cook and Sullivan, 2018).





#### A4 AWDN

The AWDN network is located in Nebraska, United States, and consists of 50 stations. The data sets were collected by the High Plains Regional Climate Center", Data availability is from 1998 to 2010 but varies per station.

#### 650 A5 BIEBRZA S-1

Two dense soil moisture sites suited for the validation of high-resolution Sentinel-1 soil moisture products were established in the Biebrza Wetlands in Northeastern Poland in May 2015 (Musial et al., 2016). One site is located across drained grassland and the second one across natural temporarily flooded marshland. They are located within 7 km distance, therefore weather conditions are similar but soil moisture regimes differ. Both sites are equipped with 9 soil moisture stations with 5 soil probes  
 655 each and a weather station. The sites are homogeneous regarding vegetation cover. The organic peat soils feature porosity values up to 82%. The sites are maintained by the Remote Sensing Centre of the Institute of Geodesy and Cartography (IGiK).

#### A6 BNZ-LTER

The Bonanza Creek Long Term Ecological Research (BNZ-LTER) network consists of 12 stations located in the boreal forest near Fairbanks, Alaska (Van Cleve, 2015). Soil moisture measurements start around the year 2000 for each station. In addition  
 660 to soil moisture in several depths, observations are available of soil temperature, air temperature, precipitation, snow depth and snow water equivalent.

#### A7 CALABRIA

The Calabria network operates 5 TDR stations measuring volumetric soil moisture at 30, 60 and 90 cm depth. The stations were installed in 2000 by the "Centro Funzionale Decentrato" of the Calabria Region for hydrometeorological monitoring for  
 665 flood and landslide risk mitigation. For more information on the performance of the network see (Brocca et al., 2011).

#### A8 CAMPANIA

The CAMPANIA network consisted of two stations located near the city of Naples, southern Italy. It was managed by the "Centro Funzionale per la Previsione Meteorologica e il Monitoraggio Meteo-Pluvio-Idrometrico e delle Frane". The ISMN contains data from the operational start in 2000 until the end of 2008. The data sets include soil moisture measured at a depth of  
 670 0.30 m, precipitation, and air temperature. For more information on the performance of the network, see (Brocca et al., 2011).

#### A9 CARBOAFRICA

CARBOAFRICA is located outside El Obeid in Kordofan in Sudan and has been active since February 2002. It is operated by the Department of Physical Geography and Ecosystem Science of the Lund University, Sweden, in cooperation with the Agricultural Research Corporation (ARC) in El Obeid, Sudan. From 2007-2009, eddy covariance measurements were undertaken,  
 675 Which are available from FLUXNET. For more information, see Ardö (2013).



## A10 CHINA

This agricultural monitoring campaign from 1981 to 1999 includes 40 stations with soil moisture measurements until one meter depth and was hosted by the Institute of Geographical Sciences and Natural Resources Research of the Chinese Academy of Sciences in Beijing (Liu et al., 2001). The data set were transferred to the ISMN from the Global Soil Moisture Data Bank  
680 (Robock et al., 2000) and contain measurements amde at the 8th, 18th and 28th of each month.

## A11 COSMOS

The Cosmic-ray Soil Moisture Observing System (COSMOS; Zreda et al., 2012) started in 2009 with a grant from the US National Science Foundation as a four-year project for demonstration of the then new technology of sensing soil moisture with cosmogenic neutrons (Zreda et al., 2008). On the completion of the project the network had 60 sites, most of them in the USA,  
685 and a few in South America, Europe and Africa. The network produces hourly soil moisture data available in real time to all without restrictions. After the project funding ended in 2013, the network operations continued with the support by Quaesta Instruments, a private company. The current status is active, but the sensors are being relocated and repurposed.

## A12 CTP\_SMTMN

CTP-SMTMN is a multiscale Soil Moisture and Temperature Monitoring Network on the central Tibetan Plateau (Yang et al.,  
690 2013). The network, with an average elevation of 4650 m a.s.l., consists of 56 stations that measure soil moisture and soil freeze/thaw status at three spatial scales (100, 25, and 9 km). The terrain is relatively flat and covered by sparse and short grasses; the annual precipitation is about 400-500 mm. The network has been in operation since 2010.

## A13 DAHRA

The Dahra field site is located in a typical low tree and shrub savanna environment in Senegal. To limit the uncertainty in the  
695 comparison of remote sensing products and models, the site was selected to be flat with homogeneous vegetation cover within a radius of at least three km. The site is equipped with two towers: a meteorological tower with meteorological, hydrological and radiation sensors, and an eddy covariance flux tower. More info can be found at (?).

## A14 FLUXNET-AMERIFLUX

AMERIFLUX is the North-American contribution to the global FLUXNET. At this moment two sites close to Sacramento,  
700 CA, i.e., Tonzi and Vaira Ranch, are distributed through the ISMN. Both stations provide soil moisture measurements at eight different depths down to 0.60 m. Additionally, soil temperature, air temperature, and precipitation are provided.

## A15 FMI

This distributed network of in situ measurement stations gathering information on soil moisture and soil temperature has been set up in recent years at the Finnish Meteorological Institute's (FMI) Sodankylä Arctic research station in Northern Finland.



705 Between 2010 and 2017, 16 stations were installed around Sodankylä and 3 further north at Saariselkä. Each station covers a  
vertical measuring profile and two additional horizontal measuring points. The vertical profiles have five sensors placed close  
to the station at the following depths: 5, 10, 20, 40, and 80 cm in mineral and semi-organic soils, and at 5, 10, 20, 30, 40 cm in  
organic soils. The two additional horizontal measuring points, at depths of 5 and 10 cm, have been installed approximately ten  
meters from the station in opposing directions to catch small-scale variations in top soil moisture. A more detailed description  
710 is provided in (Ikonen et al., 2015, 2018).

#### **A16 FR\_Aqui**

The FR-AQUI network has been set up by INRAE in the Bordeaux–Aquitaine region (southwestern France) in 2012 (Wigneron  
et al., 2018; Al-Yaari et al., 2018a). Measurements taken at this five-station network (plus the nearby ICOS Bilos site), include  
soil moisture and temperature at various depths, and the height of the groundwater table. Four sites are installed in the Les  
715 Landes forest, which is one of the largest coniferous forests in Europe, and one site is installed close to vineyards of the  
Bordeaux Graves region.

#### **A17 GROW**

GROW gathered crowd sourced observations to assess the temporal and spatial consistency of various satellite-derived soil  
moisture products. 6,500 low cost sensors were deployed in 24 GROW places, in 13 Countries across Europe. A subset of 150  
720 sensors were transferred to the ISMN (Zappa et al., 2019, 2020)) and contain measurements of Soil Moisture (%), Temperature  
(C) and metadata: Latitude and Longitude of the Sensor, Name of the Sensor, Date of Observation. Complete data set is licensed  
and available at <https://doi.org/10.15132/10000156>. More information on the quality of the data can be found in Xaver et al.  
(2020).

#### **A18 GTK**

725 The GTK network is operated by the Geological Survey of Finland and contains 7 stations. The data are available from the  
years 2001 to 2012 but the availability varies per station.

#### **A19 HiWATER\_EHWSN**

The HiWATER EHWSN network is located in an irrigated farmland in the middle stream of the Heihe River Basin close to  
Gobi desert, China. It consists of short time series between April and September 2012 collected at 174 stations by the Cold  
730 and Arid Regions Environmental and Engineering Research Institute (CAREERI) of the Chinese Academy of Science (Kang  
et al., 2014; Jin et al., 2014).



## A20 HOAL-SoilNet

The Hydrological Open Air Laboratory - Soil Network (HOAL-Soilnet) has been set up in Petzenkirchen, Austria, as part of a concerted effort to advance the understanding of water-related flow and transport processes in a 66 ha agricultural catchment (Blöschl et al., 2016). Soil moisture has been monitored since 2013 at about 30 locations at four depths at time intervals of 30 minutes using TDR sensors. Measurements are taken from permanent stations, located in grassland, forest or at field boundaries, as well as from stations that are temporarily installed in cropland (Vreugdenhil et al., 2013).

## A21 HOBE

HOBE is a hydrological observatory established in the western part of Denmark in the Skjern River catchment Jensen and Refsgaard (2018). Within the sub-catchment Ahlergaarde data have been collected from a network 30 soil moisture stations distributed within the sub-catchment according to respective fractions of classes representing mainly land cover and soil type (Bircher et al., 2012). At each station Decagon 5TE capacitance sensors have been installed at 2.5, 22.5 and 52.5 cm depths. The sensors are logged every 30 minutes.

## A22 HSC\_SEOLMACHEON

HSC\_SELMACHEON was a single station located in South Korea. Data were collected by the Hydrological Survey Center (HSC) and WRRSL and are available for the period August - September 2011.

## A23 HYDROL-Net Perugia

The HYDROL-Net Perugia network Morbidelli et al. (2014, 2017) consists of 1 station (called WEEF: Water Engineering Experimental Field) located near the city of Perugia, Central Italy. Measurements of soil moisture at 4 different depths (0.05, 0.15, 0.25 and 0.35 m), soil temperature at 2 depths (0.05 and 0.30 m), precipitation and air temperature from January 2010 to December 2013 are included in the ISMN. HYDROL-Net Perugia is operated by the Department of Civil and Environmental Engineering of the Perugia University.

## A24 ICN

The former Illinois Climate Network was operated by the Water and Atmospheric Resources Program of the Illinois State Water Survey and formerly integrated in the Global Soil Moisture Data Bank (Hollinger and Isard, 1994). Between 1983 and 2010, ICN covered 19 stations measuring soil moisture and soil temperature down to two metres depth, as well as precipitation.

## A25 IIT\_KANPUR

The network IIT\_KANPUR network consisted of a single station and was managed by Hydraulics and Water Resources Laboratory, Indian Institute of Technology Kanpur, India. Soil moisture measurements were made at four depths (10, 25, 50 and 80



cm) between from June 2011 to November 2012. The station was situated in the Ganga river basin, which is the largest river basin in India and the soil type at the observation site is clayey silt.

## A26 IMA\_CAN1

The IMA\_CAN1 network is operated by the Institute for Agricultural and Earthmoving Machines (IMAMOTER) of the Italian National Research Council (CNR), now part of STEMS-CNR. It is located in an experimental vineyard in Carpeneto, in the “Alto Monferrato” hilly region, which is a valuable vine-growing and wine production area in the Piedmont Region in northwestern Italy. The monitored vineyards are part of the “Experimental Vine and Wine Centre of Agrion Foundation”. The stations in the network provide measurements of soil moisture, precipitation, air temperature and humidity, hourly runoff and event soil losses. Hourly volumetric soil moisture was measured by 12 5TM sensors in the period 2011 - 2015, both in grassed and tilled vineyards, in correspondence of the track and no-track position, down and up the hill (Biddoccu et al., 2016; Capello et al., 2019; Raffelli et al., 2018).

## A27 IOWA

The IOWA network was located in two catchments in the southwestern part of Iowa, USA. Soil moisture observations from 6 stations, until 2.6 meter depth, with an interval of twice a month during 1972 to 1994 (April to October) are included in the ISMN (Robock et al., 2000). This network was formerly included in the Global Soil Moisture Data Bank.

## 775 A28 IPE

The Instituto Pirenaico de Ecologia (IPE) network runs two stations located in Aragón, north-eastern Spain. The stations have been collecting meteorological data with at least hourly time resolution (air and soil temperatures, soil moisture, relative humidity, radiation, wind speed) since 2008 in a Mediterranean oak forest (Agüero) and a semi-arid pine forest (Peñaflor). These measurements are related to dendrometer hourly records of changes in root and stem (Agüero; see (Alday et al., 2020)) or stem (Peñaflor) increment of the main tree species.

## A29 iRON

The interactive Roaring Fork Observation Network (iRON) is a series of 10 stations operated by the Aspen Global Change Institute spread across the elevations of the Roaring Fork Watershed, located in the Southern Rocky Mountains of the United States. This data set includes soil moisture at 5, 20, and 50cm, soil temperature at 20cm, and additional weather measurements variable by station. Further information can be found in Osenga et al. (2019) or on [agci.org/iron/about](https://agci.org/iron/about).

## A30 KHOREZM

The KHOREZM network in Uzbekistan is located between the Amudarya river and the border to Turkmenistan and was part of a project conducted by Uni Würzburg, Germany (Patrick Knöfel). Seven stations with soil moisture, soil temperature, air tem-



perature and surface temperature measurements from 2010 April to 2011 September are included in the ISMN. Although soil  
790 moisture was not observed continuously, the measurements are still a valuable contribution since no other recent observations  
are available in this region.

### **A31 KIHS\_CMC**

The Korea Institute of Hydrological Survey has been running the Cheongmicheon (CMC) network since 2009 with annually  
returning measurements from March till December. It comprises 56 TDR Buriable Waveguide soil moisture sensors at 18  
795 stations located on an area the size of a football field. All stations have a sensor installed at 0.1 m and additional sensors at  
varying depths, i.e., at 0.3, 0.4, 0.6 and 0.9 m.

### **A32 KIHS\_SMC**

KIHS\_SMC is operated by the Environmental and Remote Sensing Lab of the Korea Institute of Hydrological Survey. The 51  
soil moisture sensors (depths from 0.1 to 0.6 m) are located on a mountain slope distributed over 19 stations in close proximity  
800 to each other.

### **A33 LAB-net**

LAB-net was created as the first soil moisture network in Chile to support remote sensing research on drought and water  
use conflicts (Mattar et al., 2016). Three stations measuring soil moisture, soil temperature, precipitation, and air temperature  
between 2014 until 2017 over various land cover types have been included in the ISMN.

### **805 A34 MAQU**

The Maqu monitoring network ((Su et al., 2011; Dente et al., 2012) is situated on the north-eastern fringe of the Tibetan  
Plateau, covering an area of approximately 40 km×80 km with the elevation varying from 3200 m to 4200 m above sea level.  
The network have SMST profiles (5 cm, 10 cm, 20 cm, 40 cm, and 80 cm) measured with a 15-min interval. Soil moisture and  
temperature data from 2008 to 2018 are included in the ISMN.

### **810 A35 METEROBS**

The METEROBS (MET European Research OBServations) network measured soil moisture at a single site between October  
2011 October and May 2012. It was located in the Apennines mountains in the rural area of Benevento, Southern Italy. Soil  
Moisture measurements from five layers until 0.5 meter depth have been included in the ISMN.

### **A36 MOL-RAO**

815 The MOL-RAO network is operated by the German Meteorological Service (DWD) and part of the operational measuring  
program of the MOL-RAO (Meteorological Observatory Lindenberg – Richard-Aßmann-Observatory). The network is situated



in the north-east of Germany and consists of two stations. While the Station Falkenberg has a grass type vegetation, Kehrigk is situated in a pine forest (Beyrich and Adam, 2007). Volumetric soil moisture, soil temperature, air temperature and precipitation are provided in half hour resolution since 2003.

## 820 **A37 MONGOLIA**

Soil moisture data sets for 44 stations were collected by the National Agency of Meteorology, Hydrology and Environment Monitoring in Ulaanbaatar. All observations were taken using the gravimetric technique and initially provided as volumetric plant-available water [%]. Volumetric soil moisture [ $\text{m}^3/\text{m}^3$ ] was calculated by first extracting texture properties of all sites from the Harmonized World Soil Database and subsequently calculating the wilting levels for all stations using the equations of Saxton and Rawls (2006). Soil moisture measurements are provided three times a month (on the 8<sup>th</sup>, 18<sup>th</sup> and 28<sup>th</sup>) from 1964 to 2002 during the warm period of the year, which runs from April until the end of October (e.g., Robock et al. (2000)).

## **A38 MySMNet**

The Malaysian Soil Moisture Network (MySMNet) is operational since 2014. It deploys seven stations (four on oil palm plantation, two on shrubland and one on orchard) that collect soil moisture at 5, 50, and 100 cm depth, soil temperature at 5 cm, air temperature, and relative humidity, all on an hourly basis. The soil moisture sensors used are the WaterScout SM100 (Kang et al., 2019).

## **A39 ORACLE**

The network ORACLE network includes 6 stations. The datasets reach back to the year 1985 and are available until 2013. ORACLE is a research observatory east of Paris to study of the "Grand Morin" and "Petit Morin" river catchments, particularly floods, low water periods, water quality, and the impact of human activities on the environment.

## **A40 OZNET**

OzNet was established in 2001 with 8 sites across the Murrumbidgee catchment and a cluster of a further 5 sites in the Adelong and Kyeamba catchments. This was further extended in 2003 to include a total of 11 sites at Kyeamba for a GRACE validation experiment and 13 sites at Yanco for SMOS pre-launch algorithm development and post-launch calibration-validation. The Yanco site was further extended to have clusters of stations across 9 km and 3 km grids in crop and grasslands areas for SMAP algorithm development, calibration, and validation (Smith et al., 2012).

## **A41 PBO\_H2O**

The PBO\_H2O network is a former near real time network of the ISMN hosted by the University of Colorado, USA. It consisted of 159 stations distributed in the west of the USA, Bahamas, Dominican Republic, Puerto Rico, Colombia, South Africa and Saudi Arabia. Soil moisture (measured using GPS reflections), precipitation, air temperature and snow depth from 2004 to





2017 is stored in the ISMN database (Larson et al., 2008). The network was discontinued in 2018 because of lacking financial support.

#### **A42 PTSMN**

850 The Patitapu Soil Moisture Network (PTSMN) was established in 2016 on the hill country landscapes of the East Coast of New Zealand's North Island. PTSMN was deployed to capture spatiotemporal soil moisture trends on various topographical positions distributed over a 13.8 km<sup>2</sup> area. The network is composed of twenty multi-sensor probes that were calibrated to the site-specific soils (Hajdu et al., 2019). The sensors provide readings at four consecutive depths down to 0.43 m.

#### **A43 REMEDHUS**

855 REMEDHUS is the University of Salamanca Soil Moisture Measurement Stations Network and was installed in March 1999, with a set of old TDR stations and manual measurements. It is one of the first soil moisture networks in Europe. The network was automated and upgraded with capacitance probes in 2005. The REMEDHUS data available in the ISMN cover the period since the automation. REMEDHUS is located in an agricultural area in the central part of the Duero basin (Spain). The network currently has 20 stations that hourly measure soil moisture and soil temperature in the 0-5 cm layer (González-Zamora et al., 2018).

#### **860 A44 RISMA**

The Real-time in situ Soil Monitoring for Agriculture (RISMA) network was established in 2011 by Agriculture and Agri-Food Canada at agricultural locations in Ontario, Manitoba, and Saskatchewan (Ojo et al., 2015; J., 2011; Canisius, 2011). There are currently 23 RISMA stations collecting hourly soil moisture/temperature data at depths to 1 – 1.5 metres in combination with meteorological data. Calibrated, quality controlled soil moisture and weather data are provided to the ISMN on an annual basis.

#### **865 A45 RSMN**

The Romanian Soil Moisture Network (RSMN) consists of 19 stations homogeneously distributed over Romania. The network is managed by the Romanian National Meteorological Administration and is part of the ASSIMO project, which aims to create a framework for the evaluation of current and future satellite microwave-derived soil moisture products.

#### **A46 RUSWET-AGRO, RUSWET-GRASS, RUSWET-VALDAI**

870 The three historical RUSWET networks were agricultural prediction campaigns conducted by the State Hydrological Institute of the Former Soviet Union within the area of present Russia. Measurements were taken from 1952 until 2002 and initially distributed through the Global Soil Moisture Data Bank. Altogether, the networks operated 337 sites at which soil moisture was measured three times per month via gravimetric sampling. At RUSWET-VALDAI also soil temperature, precipitation, and



air temperature were collected. RUSWET contributes both the northernmost (on Mc-Clintock Island) and the earliest (June 8th 1952) observations to the ISMN (Robock et al., 2000).

#### **A47 SASMAS**

The Scaling and Assimilation of Soil Moisture and Streamflow (SASMAS) monitoring network commenced commissioning in late 2002 with a total of 26 stations in operation by 2003 across a 6,500 km<sup>2</sup> catchment (Rüdiger et al., 2007). Soil moisture is observed up to a depth of 90 cm (where possible) at depth intervals of 0-30, 30-60, and 60-90 cm. The network was developed as a nested catchment with three different types of scales (low-resolution across the entire catchment, medium density across to smaller subcatchment, and very high density across a 175 ha single reach. This site also hosted the second NAFE campaign for which additional data was collected.

#### **A48 SCAN**

The Natural Resources Conservation Service (NRCS) operates the comprehensive, US-wide Soil Climate Analysis Network. SCAN supports natural resource assessments and conservation activities through its network of automated climate monitoring and data collection sites. SCAN focuses primarily on agricultural areas of the U.S., Puerto Rico and the Virgin Islands. The network consists of 216 stations located across the United States and reports soil moisture, soil temperature, precipitation, temperature and other climatic variables hourly. Soil sensors are situated at 5, 10, 20, 50 and 100 cm depth (Schaefer et al., 2007).

#### **A49 SKKU**

The SKKU network was located at the evenly and moderately vegetated botanical garden in South Korea. It was operated by Sungkyunkwan University from 2014 to 2016 as part of a project for evaluating Cosmic-Ray Neutron Probe (CRNP) soil moisture (Nguyen et al., 2017). The network consisted of 10 stations, and soil moisture measurements were taken from 4 different depths (i.e., 10, 20, 30, and 40 cm).

#### **A50 SMOSMANIA**

The SMOSMANIA network was installed in southern France by Météo-France, the French national meteorological service, in order to monitor in situ soil moisture and soil temperature in contrasting soil and climatic conditions at operational automatic weather stations (Calvet et al., 2007). The SMOSMANIA network is composed of 21 stations forming an Atlantic–Mediterranean transect over a large variety of mineral soils ranging from sand to clay and silt loam (Calvet et al., 2016).

#### **A51 SNOTEL**

NRCS installs, operates, and maintains an extensive, automated data collection network called SNOTEL (short for Snow Telemetry) (Leavesley et al., 2008; Leavesley, 2010). SNOTEL is part of the Snow Survey and Water Supply Forecasting



(SSWSF) Program and is designed to collect snowpack and related climatic data in the western U.S. and Alaska. The Program operates under technical guidance from the NRCS National Water and Climate Center (NWCC). With the majority of the water supply in the West arriving in the form of snow, data on snowpack provide critical information to decision-makers and water managers. SNOTEL currently consists of a network of over 860 automated SNOTEL stations of which 463 stations have soil moisture and soil temperature sensors.

## A52 SOILSCAPE

The Soil moisture Sensing Controller And oPtimal Estimator (SoilSCAPE) is a wireless soil moisture sensor network for measurements of surface-to-root-zone profiles of soil moisture (Moghaddam et al., 2011; MOGHADDAM et al., 2016; Shuman et al., 2010). It is designed for long-term, ultra-efficient, unattended field operations. Several networks are deployed in the US, including in the states of California, Arizona, Colorado, Alaska, and New York. Additional deployments are planned in New Mexico, US, and in New Zealand. The SoilSCAPE architecture includes a local coordinator (LC) and multiple end devices (EDs). The LC is the central command center of the network, receiving data from the several soil moisture and temperature sensors connected to each ED. The sensors include 5TM and Teros-12, depending on how recently they were deployed.

## A53 SWEX\_POLAND

The oSil Water and Energy exchange - Poland (SWEX\_POLAND) network was operated between 2000 and 2013 by the Institutes of Agrophysics PAS in Lublin. The network consisted of 6 stations located in the wetlands of Poleski Park Krajobrazowy to support SMOS product calibration and validation. Soil moisture and temperature measurements were taken down to 1 meter depth, along with precipitation observations (Marczewski et al., 2010).

## A54 SW-WHU

The SW-WHU network was hosted by Wuhan University (Chen et al., 2015a, b). SW-WHU is a high-density network, with nearly 100 soil moisture and temperature sensors within 1 km<sup>2</sup>. It adopts an NB-IOT technique for data transmission at low power consumption. Therefore, the data of SW-WHU is particularly valuable for soil moisture validation, monitoring, and application at a very high spatio-temporal resolution (Zhang et al., 2018b).

## A55 TAHMO

The Trans-African Hydro-Meteorological Observatory (TAHMO; tahmo.org) presently runs a network of over 600 meteorological weather stations in more than twenty African countries. The stations measure all standard weather parameters, such as barometric pressure, wind speed, rainfall, and radiation. Each station also has five open ports that can be used for additional sensors, such as soil moisture sensors. Presently, 70 stations have soil moisture sensors, some of them at several depths. Ideally, many stations will be upgraded by soil moisture sensors in the near future.



## A56 TERENO

TERENO consists of four terrestrial observatories that represent typical landscapes in Germany and Central Europe and are considered to be highly vulnerable to the effects of global and climate change (Zacharias et al., 2011; Bogen et al., 2012).  
 935 TERENO combines observations with comprehensive large-scale experiments, integrated modeling, remote sensing and novel measurement technologies to increase our understanding of the functioning and feedbacks of terrestrial ecosystems (Bogen, 2016). The long-term observation platform of TERENO is composed of various measurement systems, including networks of climate and lysimeter stations, eddy-covariance towers, and networks of soil, surface water and groundwater sensors. Almost all online measurements are freely accessible via the TERENO data portal (<http://www.tereno.net/ddp/>).

## 940 A57 UDC\_SMOS

The former German UDC\_SMOS network was hosted by the University of Munich, Department of Geography in cooperation with the Bavarian State Research Center for Agriculture and funded by the German Aerospace Centre (DLR). It was located in grassland in the Bavarian region around Munich city, as an official European SMOS calibration/ validation test site. Eleven stations provided soil moisture data from 2007 until 2011 until 40 cm depth measured by several types of sensors (Loew et al., 2009; Schlenz  
 945 et al., 2012).

## A58 UMBRIA

This soil moisture monitoring network in the north of the Umbria region, in the upper Tiber River basin operates three stations in real-time: Torre dell'Olmo, Petrelle, Cerbara. Each station measures at 10 cm, 20 cm, and 40 cm depth. Additional stations of the network have not been operational since 2015 due to a lack of resources for their maintenance. Provided financial resources  
 950 become available, the stations that are no longer functioning will be restored.

## A59 UMSUOL

UMSUOL ("Umidità del Suolo") is a one-station network located close to Bologna, northern Italy. Soil moisture measurements at seven different depths are provided by the Agenzia Regionale Prevenzione Ambiente (ARPA). The ISMN contains data from the years 2009 and 2010.

## 955 A60 USCRN

The U.S. Climate Reference Network (USCRN) contains 114 stations sparsely distributed across the contiguous United States. Each station has three sets of soil moisture/temperature probes at 5 depths: 5, 10, 20, 50, and 100 cm in addition to air and surface temperature, precipitation, relative humidity, and solar radiation (Bell et al., 2013). The stations were installed between 2009 and 2011 and are still in operation in 2021. The stations are maintained by the National Oceanic and Atmospheric  
 960 Administration Atmospheric Turbulence and Diffusion Division (NOAA ATDD) and their data are maintained by NOAA National Centers for Environmental Information (NOAA NCEI).



#### A61 USDA-ARS

The United States Department of Agriculture-Agricultural Research Service (USDA-ARS) operates a number of Long Term Agroecosystem Research (LTAR) sites, some of which have spatial coverage of soil moisture and soil temperature. As experimental sites, the locations and configuration of the stations can change depending on the current scientific questions being addressed. A description of the sites can be found in Jackson et al. (2011).

#### A62 VDS

The VDS network is run by VanderSat, a dutch company specialised in providing global satellite-observed data and services over land. The VDS network consists of four stations located near the city of Bago in Myanmar. The network was installed to validate satellite soil moisture products in the tropics. The network has two measurement periods, one from June 2017 to July 2018 and one from March 2020 onwards.

#### A63 VAS

The "Valencia Anchor Station" network is operated by the Climatology from Satellites Group and Júcar River Basin Authority of the University of Valencia, Spain. The network is located in Spain and consists 3 stations. The datasets are available for the years 2010 and 2011.

#### A64 WEGENERNET

The WegenerNet, located in the foreland of the southeastern Austrian Alps, is a long-term weather and climate monitoring facility comprising 155 hydrometeorological stations in a dense grid with one station every about 2 km<sup>2</sup> Kirchengast et al. (2014); Fuchsberger et al. (2020). Together with a range of meteorological variables such as temperature, humidity, precipitation and wind, it also measures soil moisture and temperature at 12 stations. These variables are measured at 0.2-0.3 m depth in diverse soil types representative for the region.

#### A65 WSMN

The Wales Soil Moisture Network was founded in July 2011. It consists of a total of 9 monitoring sites located in Mid-Wales representing a range of conditions typical of the Welsh environment, with climate ranging from oceanic to temperate and the most typical land use/cover types. The data set acquired in the network are composed of 0-5 (or 0-10) cm soil moisture, soil temperature, precipitation as well as other ancillary data (Petropoulos and McCalmont, 2017).



## Appendix B: Network abbreviations

Acronym/ definition	Full name
AACES	Australian Airborne Cal/val Experiment for SMOS
AMMA-CATCH	African Monsoon Multidisciplinary Analysis – Coupling the Tropical Atmosphere and the Hydrological Cycle
ARM	Atmospheric Radiation Measurement Climate Research Facility
AWDN	Automated Weather Data Network (name of in situ network in the USA)
BIEBRZA_S-1	Biebrza Sentinel-1 Supersite
BNZ-LTER	Bonanza Creek Long-Term Ecological Research
CALABRIA	Name of in situ network in Italy
CAMPANIA	Name of in situ network in Italy
CARBOAFRICA	Name of in situ network in Sudan
CHINA	Name of in situ network in China
COSMOS	Cosmic-ray Soil Moisture Observing System
CTP_SMTMN	Central Tibetan Plateau Soil Moisture and Temperature Monitoring Network
DAHRA	Name of in situ network in Senegal
ECMWF	European Centre for Medium-Range Weather Forecasts
FLUXNET_AMERIFLUX	Name of in situ network in USA
FMI	Finnish Meteorological Institute (name of in situ network in Finland)
FR_Aqui	Name of in situ network in France
GDPR	General Data Protection Regulation
GROW	GROW citizen science project (name of in situ network based in the UK but distributed through out Europe)
GTK	Geological Survey of Finland (name of in situ network in Finland)
HiWATER_EHWSN	Heihe Watershed Allied Telemetry Research Wireless Sensor Network (in China)
HOBE	Hydrological Observatory in Denmark
HSC_SEOLMACHEON	Hydrological Survey Center (HSC) in –situ network in South Korea
HWSD	Harmonized World Soil Database
HYDROL-NET_PERUGIA	Name of in situ network in Italy
ICN	Illinois Climate Network (name of in situ network in USA)
IIT_KANPUR	Indian Institute of Technology Kanpur (name of in situ network in India)
IGiK	Institute of Geodesy and Cartography

*Table continued on next page*



Acronym/ definition	Full name
IMA_CAN1	STEMS-CNR in situ soil monitoring at the Tenuta Cannona Experimental Vine and Wine Centre of Agrion Foundation
IOWA	Name of in situ network in USA
IPE	Instituto Pirenaico de Ecología (Organisation name and the name of in situ network in Spain)
iRON	Roaring Fork Observation Network
ISMN	International Soil Moisture Network
KHOREZM	Name of in situ network in Uzbekistan
KIHS_CMC	Name of in situ network in South Korea
KIHS_SMC	Name of in situ network in South Korea
LAB-net	Laboratory for Analysis of the Biosphere Network (name of in situ network in Chile)
MAQU	Name of in situ network in China
METEROBS	Met European Research Observatory (name of in situ network in Italy)
MOL-RAO	Lindenberg Meteorological Observatory, Richard-Aßmann Observatory
MONGOLIA	Name of in situ network in Mongolia
MySMNet	Name of in situ network in Malaysia
NRT	Near Real Time
ORACLE	Name of in situ network in France
OZNET	Name of in situ network in Australia
PBO_H2O	Plate Boundary Observatory (Name of in situ network in USA)
REMEDHUS	University of Salamanca Soil Moisture Measurement Stations Network
RISMA	Real-time in situ Soil Monitoring for Agriculture Network in Canada
RSMN	Romanian Soil Moisture Network
RUSWET-AGRO	Name of in situ network in Former Soviet Union
RUSWET-GRASS	Name of in situ network in Former Soviet Union
RUSWET-VALDAI	Name of in situ network in Former Soviet Union
SASMAS	Scaling and Assimilation of Soil Moisture and Streamflow
SCAN	Soil Climate Analysis Network
SWATS	Surface Energy Balance System, sensor network of ARM
SKKU	Name of in situ network in South Korea
SMOS	Soil Moisture and Ocean Salinity
SMOSMANIA	Soil Moisture Observing System – Meteorological Automatic Network Integrated Application
SNOTEL	SNOWpack TELelemetry


*Table continued on next page*





Acronym/ definition	Full name
SOILSCAPE	Soil Moisture Sensing Controller and Optimal Estimator Network in the USA
STAMP	Soil Temperature and Moisture Profile, sensor network of ARM
SWATS	Soil Water and Temperature System, sensor network of ARM
SWEX_POLAND	Soil Water Experiment Poland (name of in situ network in Poland)
SW-WHU	Name of in situ network in Poland
TAHMO	Trans-African Hydro-Meteorological Observatory (name of in situ network in Sahel Zone in Africa)
TERENO	Terrestrial Environmental Observatories (name of in situ network in Germany)
UDC_SMOS	Upper Danube Catchment SMOS (name of in situ network in Germany)
UMBRIA	Name of in situ network in Italy
UMSUOL	Umidita del Suolo (name of in situ network in Italy)
USCRN	US Climate Reference Network
USDA_ARS	Research Watersheds Supporting Soil Moisture Remote Sensing Network in USA
UTC	Universal Time Coordinated
VAS	Valencia Anchor Station
VDS	Vandersat, in situ network in Myanmar
WEGENERNET	Name of in situ network in Austria
WSMN	Wales Soil Moisture Network in the UK

## Appendix C: Detailed system overview



**Figure C1.** Detailed system overview of the ISMN (Status January 2021).




## Appendix D: Soil moisture sensor availability per soil depth interval

**Table D1.** Overview of data availability per depth layers showing the total number of individual sensor brands, data sets and stations integrated in the ISMN database. The calculations have been made in respect to horizontally and vertically placed sensors for each depth layer.

depth from - to [m]	sensors	data sets	stations
0.00 - 0.05	131	2116	1696
0.05 - 0.10	123	2482	1459
0.10 - 0.20	138	2690	1523
0.20 - 0.30	104	1899	1463
0.30 - 0.40	43	562	483
0.40 - 0.50	73	1463	1207
0.50 - 0.60	65	1389	1125
0.60 - 0.70	25	210	185
0.70 - 0.80	20	113	106
0.80 - 0.90	24	183	159
0.90 - 1.00	25	346	302
1.00 - 1.10	42	571	450
1.10 - 1.20	5	7	5
1.20 - 1.30	7	44	24
1.30 - 1.40	4	8	6
1.40 - 1.50	8	13	9
1.50 - 1.60	12	21	15
1.60 - 1.70	0	0	0
1.70 - 1.80	3	35	18
1.80 - 1.90	1	1	1
1.90 - 2.00	8	20	20
2.00 - 2.10	13	10	8



## 990 Appendix E: Metadata structure



**Figure E1.** Overview of all mandatory metadata information available for each station within the ISMN database.



## Appendix F: Scientific studies using ISMN data for satellite and model product evaluation and development.

**Table F1.** Overview of scientific studies using ISMN data for satellite product evaluation and development.

Sensor	Publications
SMOS	Albergel et al. (2012c), Mecklenburg et al. (2012), Parrens et al. (2012), Sanchez et al. (2012), Schlenz et al. (2012), Wanders et al. (2012), Pellarin et al. (2013), Fascetti et al. (2014), Petropoulos et al. (2014), Pierdicca et al. (2014), Zeng et al. (2014), Rodriguez-Fernandez et al. (2015), Gruber et al. (2015), Hottenstein et al. (2015), Kornelsen and Coulibaly (2015), Lee and Im (2015), Parinussa et al. (2015), Pierdicca et al. (2015a), Pierdicca et al. (2015b), Scaini et al. (2015), González-Zamora et al. (2015), Fascetti et al. (2016), Fascetti et al. (2016), Kędzior and Zawadzki (2016), Kerr et al. (2016), Pablos et al. (2016), Piles et al. (2016), Rautiainen et al. (2016), Scholze et al. (2016), van der Schalie et al. (2016), González-Zamora et al. (2016), Ángel González-Zamora et al. (2016), Al-Yaari et al. (2017), Cui et al. (2017), Fascetti et al. (2017), Fernandez-Moran et al. (2017b), Fernandez-Moran et al. (2017a), Karthikeyan et al. (2017), Lievens et al. (2017), Martens et al. (2017), Mohanty et al. (2017), Montzka et al. (2017), Muñoz Sabater et al. (2017), Pan et al. (2017), Peng et al. (2017), Pierdicca et al. (2017), Rains et al. (2017), van der Schalie et al. (2017), Yao et al. (2017), Al-Yaari et al. (2018b), Ebrahimi et al. (2018), Ebrahimi-Khusfi et al. (2018), Lee (2018), Liu et al. (2018), Pablos et al. (2018), Tagesson et al. (2018), Tian (2018), van der Schalie et al. (2018), Wu et al. (2018), Al-Yaari et al. (2019b), Al-Yaari et al. (2019a), Blyverket et al. (2019), Blyverket (2019), de Rosnay et al. (2019), ?, Ma et al. (2019), Piles et al. (2019), Rodriguez-Fernandez et al. (2019), Tian et al. (2019b), González-Zamora et al. (2019), Beck et al. (2020), Chen et al. (2020), Helgert and Khodayar (2020), Herbert et al. (2020), Link et al. (2020), Moreno-Martínez et al. (2020), Portal et al. (2020), Sadri et al. (2020), Dong et al. (2020), Li et al. (2020c), Wigneron et al. (2021)
ASCAT	Brocca et al. (2011), Liu et al. (2011), Albergel et al. (2012c), Liu et al. (2012b), Parrens et al. (2012), Wanders et al. (2012), Albergel et al. (2013b), Pellarin et al. (2013), Zwieback et al. (2013), Fascetti et al. (2014), Gruber et al. (2014), Parinussa et al. (2014b), Paulik et al. (2014), Pierdicca et al. (2014), Gruber et al. (2015), Parinussa et al. (2015), Pierdicca et al. (2015a), Pierdicca et al. (2015b), Su et al. (2015), Zwieback et al. (2015), Fascetti et al. (2016), Fascetti et al. (2016), Zwieback et al. (2016), Fascetti et al. (2017), Gruber et al. (2017), Karthikeyan et al. (2017), Kolassa et al. (2017), Lievens et al. (2017), Massari et al. (2017), Mohanty et al. (2017), Montzka et al. (2017), Pierdicca et al. (2017), Sure et al. (2017), Al-Yaari et al. (2018b), Bauer-Marschallinger et al. (2018), Kim et al. (2018), Afshar et al. (2019), Al-Yaari et al. (2019b), Fairbairn et al. (2019), Gruber et al. (2019a), Beck et al. (2020), Chen et al. (2020), Link et al. (2020), Moreno-Martínez et al. (2020), Zappa et al. (2020)
AMSR-E	Brocca et al. (2011), Liu et al. (2011), Liu et al. (2012b), Parinussa et al. (2012), Wanders et al. (2012), Albergel et al. (2013b), Dente et al. (2013), Dorigo et al. (2013), Zhao and Li (2013), de Jeu et al. (2014), Parinussa et al. (2014b), Zeng et al. (2014), Coopersmith et al. (2015), Parinussa et al. (2015), Su et al. (2015), Zhao and Li (2015), Al-Yaari et al. (2016), Du et al. (2016), Han et al. (2016), Parinussa et al. (2016), Santi et al. (2016), Zhao et al. (2016), Gruber et al. (2017), Han et al. (2017), Karthikeyan et al. (2017), Kolassa et al. (2017), Massari et al. (2017), Mohanty et al. (2017), Su et al. (2017), Sure et al. (2017), Tobin et al. (2017), van der Schalie et al. (2017), WANG Rui (2017), Yao et al. (2017), Lei et al. (2018), Liu et al. (2018), van der Schalie et al. (2018), Afshar et al. (2019), Hu et al. (2019), Xie et al. (2019), Chen et al. (2020), Deng et al. (2020c), Hagan et al. (2020), Dong et al. (2020)
SMAP	Fascetti et al. (2016), Zeng et al. (2016), Al-Yaari et al. (2017), Cui et al. (2017), Fascetti et al. (2017), Karthikeyan et al. (2017), Mohanty et al. (2017), Montzka et al. (2017), Pierdicca et al. (2017), Alemohammad et al. (2018), Ebrahimi et al. (2018), Ebrahimi-Khusfi et al. (2018), Kim et al. (2018), Kim and Lakshmi (2018), Kolassa et al. (2018), Pablos et al. (2018), Santi et al. (2018), Xu et al. (2018), Zhao et al. (2018), Al-Yaari et al. (2019b), Bai et al. (2019), Blyverket et al. (2019), Blyverket (2019), Ebtehaj and Bras (2019), Fang et al. (2019), Ma et al. (2019), Park et al. (2019), Zhang et al. (2019b), Beck et al. (2020), Chen et al. (2020), Fang et al. (2020), Gao et al. (2020), Gao et al. (2020), Link et al. (2020), Park et al. (2020), Portal et al. (2020), Sadri et al. (2020), Suman et al. (2020), Sun et al. (2020), Zappa et al. (2020)



**Table F1 continued from previous page**

Sensor	Publications
ESA CCI	Gruber et al. (2013), Enenkel et al. (2016), Su et al. (2016), Cui et al. (2017), Dorigo et al. (2017), Karthikeyan et al. (2017), Martens et al. (2017), Tobin et al. (2017), Al-Yaari et al. (2018b), Al-Yaari et al. (2019b), Al-Yaari et al. (2019a), Blyverket et al. (2019), Blyverket (2019), Gruber et al. (2019b), Guevara and Vargas (2019), Ma et al. (2019), González-Zamora et al. (2019), Zhu et al. (2019), Beck et al. (2020), Deng et al. (2020c), Kovačević et al. (2020), Zappa et al. (2020), Dorigo et al. (2015), Pratola et al. (2015), Nicolai-Shaw et al. (2015b), An et al. (2016), Liu et al. (2018), Liu et al. (2020), Chen et al. (2016)
AMSR2	Kim et al. (2015a), Kim et al. (2015b), Parinussa et al. (2015), Kim et al. (2016), Wu et al. (2016), Anoop et al. (2017), Cui et al. (2017), Karthikeyan et al. (2017), Montzka et al. (2017), Yao et al. (2017), Fang et al. (2018), Kim et al. (2018), Liu et al. (2018), Santi et al. (2018), Hu et al. (2019), Ma et al. (2019), Beck et al. (2020), Chen et al. (2020), Hagan et al. (2020), Link et al. (2020), Moreno-Martínez et al. (2020), Anoop et al. (2017)
MODIS	Liu et al. (2014), Boussetta et al. (2015), Zhao and Li (2015), Pablos et al. (2016), Przeździecki et al. (2017), Alemohammad et al. (2018), Gumbrecht (2018), Pablos et al. (2018), Kumar et al. (2019b), Park et al. (2019), Han et al. (2020)
Sentinel-1	Paloscia et al. (2013), Liu et al. (2017), Bao et al. (2018), Bauer-Marschallinger et al. (2018), Dabrowska-Zielinska et al. (2018), Bauer-Marschallinger et al. (2019), Greifeneder et al. (2019), Rodionova (2019b), Rodionova (2019a), Wang et al. (2019a), Foucras et al. (2020), Han et al. (2020), Ma et al. (2020), Zappa et al. (2020)
SEVIRI	Zhao and Li (2013), Liu et al. (2014), Leng et al. (2015), Leng et al. (2016), Piles et al. (2016), Leng et al. (2017), Tagesson et al. (2018), Ghilain et al. (2019)
FY-3B	Parinussa et al. (2014a), Cui et al. (2017), Parinussa et al. (2018), Sheng et al. (2019), Chen et al. (2020), Hagan et al. (2020)
TMI	Liu et al. (2012b), Albergel et al. (2013b), Gruber et al. (2015), Karthikeyan et al. (2017), Chen et al. (2020), Hagan et al. (2020)
ERS	Liu et al. (2012b), Albergel et al. (2013b), Dente et al. (2013), Kolassa et al. (2013), Karthikeyan et al. (2017)
GNSS-IR (CYGNSS)	Zhang et al. (2018a), Kim and Lakshmi (2018), Xu et al. (2018), Eroglu et al. (2019), Chew and Small (2020), Senyurek et al. (2020)
SSM/I	Liu et al. (2012b), Albergel et al. (2013b), Kolassa et al. (2013), Karthikeyan et al. (2017), Hagan et al. (2020)
WindSat	Parinussa et al. (2012), Karthikeyan et al. (2017), Chen et al. (2020), Hagan et al. (2020)
ERS-2	Reimer et al. (2012), Pierdicca et al. (2014)
Landsat	Zhao et al. (2017), Bao et al. (2018), Pradhan (2019)
GRACE	Khaki et al. (2019), Sadeghi et al. (2020)
Aquarius/SAC-D	González-Zamora et al. (2016)
GF-1 WFV	Xing et al. (2017)
HY-2	Zhao et al. (2014)
MeMo	Beck et al. (2020)
SMmodel	Mimeau et al. (2020)




**Table F2.** Overview of scientific studies using ISMN data for model evaluation and development.

Model	Publications
ERA-Interim, ERA-Interim\Land	Albergel et al. (2012b), Albergel et al. (2012a), Albergel et al. (2013a), Albergel et al. (2013b), Gruber et al. (2013), Pierdicca et al. (2015a), Zwieback et al. (2013), Angevine et al. (2014), Balsamo et al. (2015), Cammalleri et al. (2015), Zwieback et al. (2015), Kim et al. (2016), Deng et al. (2020c), Deng et al. (2020b), Li et al. (2020a), Pierdicca et al. (2015b), Nicolai-Shaw et al. (2015b), Beck et al. (2020)
ECMWF IFS	Albergel et al. (2012c), Mecklenburg et al. (2012), Pellarin et al. (2013), Albergel et al. (2015), Rodriguez-Fernandez et al. (2015)
ERA-40	Liu et al. (2012a)
ERA5	Li et al. (2020a)
H-TESSEL	Albergel et al. (2010), Albergel et al. (2015), Dirmeyer et al. (2016)
ISBA	Albergel et al. (2010), Parrens et al. (2012), Barbu et al. (2014)
GLDAS Noah	Liu et al. (2011), Dorigo et al. (2013), Angevine et al. (2014), Bi et al. (2016), Dirmeyer et al. (2016), Zawadzki and Kędzior (2016), Pan et al. (2017), Lee (2018), McDonough et al. (2018), Mishra et al. (2018), Afshar et al. (2019), Kędzior and Zawadzki (2016), Beck et al. (2020), Deng et al. (2020c), Solander et al. (2020)
GLDAS Mosaic	Bi et al. (2016), Solander et al. (2020)
MERRA, MERRA\Land, MERRA-2	Albergel et al. (2013b), Pellarin et al. (2013), Kim et al. (2016), Reichle et al. (2017), Draper and Reichle (2019), Wang et al. (2019c), Li et al. (2020a)
CLM	Cammalleri et al. (2015), Bi et al. (2016), Dirmeyer et al. (2016), Mishra et al. (2018), Zhao and Yang (2018)
VIC	Mishra et al. (2014), Bi et al. (2016), Mishra et al. (2018), Beck et al. (2020), Solander et al. (2020)
GLEAM	Lievens et al. (2017), Martens et al. (2017), Martens et al. (2018)
LDAS	Fairbairn et al. (2015), Albergel et al. (2018), Xia et al. (2019), Albergel et al. (2020)
BEACH	Van doninck et al. (2012)
CAS-LSM	Wang et al. (2019b)
CFSR	Li et al. (2020a)
CLSM	Dong et al. (2020)
CPC	Liu et al. (2012a)
CSSP	Ji et al. (2017)
ESSMRA	Naz et al. (2020)
GSFC	Dirmeyer et al. (2016)
HBV	Beck et al. (2020)
JRA-5	Li et al. (2020a)
LISFLOOD	Cammalleri et al. (2015)
NLDAS Noah	Dong et al. (2020)
NLDAS Mosaic	Dong et al. (2020)
NCEP/NCAR	Pierdicca et al. (2015b)
Noah-MP	Dong et al. (2020)
SSA	Fairbairn et al. (2019)
SSMP	Pal and Maity (2019)
WCDA, SCDA, OPNL	Lin and Pu (2019)
WRF, WRF3.3-CLM4crop	Angevine et al. (2014), Lu et al. (2015)
ORCHIDEE-MICT	Yin et al. (2018)







## Appendix G: Effect of quality flagging on validation of ERA5 and ESA CCI soil moisture




(a) ERA5 absolute values



(b) ERA5 anomalies



(c) ESA CCI absolute values



(d) ESA CCI anomalies

**Figure G1.** Scatterplots of R Spearman between ISMN and ERA5 and ISMN and ESA CCI v05.2 combined for absolute values and anomalies. A 35-day moving average window was used to calculate anomalies. Each dot represents a single time series.



## References

- Abbaszadeh, P., Moradkhani, H., and Zhan, X.: Downscaling SMAP Radiometer Soil Moisture Over the CONUS Using an Ensemble Learning Method, *Water Resources Research*, 55, 324–344, <https://doi.org/10.1029/2018WR023354>, 2019.
- Adla, S., Rai, N. K., Karumanchi, S. H., Tripathi, S., Disse, M., and Pande, S.: Laboratory Calibration and Performance Evaluation of Low-Cost Capacitive and Very Low-Cost Resistive Soil Moisture Sensors, *Sensors*, 20, 363, <https://doi.org/10.3390/s20020363>, 2020.
- Afshar, M., Yilmaz, M., and Crow, W.: Impact of rescaling approaches in simple fusion of soil moisture products, *Water Resources Research*, 55, 7804–7825, 2019.
- Al-Yaari, A., Wigneron, J., Kerr, Y., de Jeu, R., Rodriguez-Fernandez, N., van der Schalie, R., Al Bitar, A., Mialon, A., Richaume, P., Dolman, A., and Ducharne, A.: Testing regression equations to derive long-term global soil moisture datasets from passive microwave observations, *Remote Sensing of Environment*, 180, 453–464, <https://doi.org/10.1016/j.rse.2015.11.022>, special Issue: ESA's Soil Moisture and Ocean Salinity Mission - Achievements and Applications, 2016.
- Al-Yaari, A., Wigneron, J.-P., Kerr, Y., Rodriguez-Fernandez, N., O'Neill, P., Jackson, T., De Lannoy, G., Al Bitar, A., Mialon, A., Richaume, P., Walker, J., Mahmoodi, A., and Yueh, S.: Evaluating soil moisture retrievals from ESA's SMOS and NASA's SMAP brightness temperature datasets, *Remote Sensing of Environment*, 193, 257–273, <https://doi.org/10.1016/j.rse.2017.03.010>, 2017.
- Al-Yaari, A., Dayau, S., Chipeaux, C., Aluome, C., Kruszewski, A., Loustau, D., and Wigneron, J.-P.: The AQUIC Soil Moisture Network for Satellite Microwave Remote Sensing Validation in South-Western France, *Remote Sensing*, 10, <https://doi.org/10.3390/rs10111839>, 2018a.
- Al-Yaari, A., Dayau, S., Chipeaux, C., Aluome, C., Kruszewski, A., Loustau, D., and Wigneron, J.-P.: The AQUIC Soil Moisture Network for Satellite Microwave Remote Sensing Validation in South-Western France, *Remote Sensing*, 10, 1839, <https://doi.org/10.3390/rs10111839>, 2018b.
- Al-Yaari, A., Ducharne, A., Cheruy, F., Crow, W. T., and Wigneron, J.-P.: Satellite-based soil moisture provides missing link between summertime precipitation and surface temperature biases in CMIP5 simulations over conterminous United States, *Scientific Reports*, 9, 1657, <https://doi.org/10.1038/s41598-018-38309-5>, 2019a.
- Al-Yaari, A., Wigneron, J.-P., Dorigo, W., Colliander, A., Pellarin, T., Hahn, S., Mialon, A., Richaume, P., Fernandez-Moran, R., Fan, L., Kerr, Y., and Lannoy, G. D.: Assessment and inter-comparison of recently developed/reprocessed microwave satellite soil moisture products using ISMN ground-based measurements, *Remote Sensing of Environment*, 224, 289–303, <https://doi.org/10.1016/j.rse.2019.02.008>, 2019b.
- Albergel, C., Calvet, J.-C., Rosnay, P., Balsamo, G., W, W., Hasenauer, S., V, N., Martin, E., Bazile, E., F, B., and Mahfouf, J.-F.: Cross-evaluation of modelled and remotely sensed surface soil moisture with in situ data in Southwestern France, *Hydrology and Earth System Sciences Discussions*, 14, <https://doi.org/10.5194/hess-14-2177-2010>, 2010.
- Albergel, C., Balsamo, G., de Rosnay, P., Muñoz Sabater, J., and Boussetta, S.: A bare ground evaporation revision in the ECMWF land-surface scheme: evaluation of its impact using ground soil moisture and satellite microwave data, *Hydrology and Earth System Sciences*, 16, 3607–3620, <https://doi.org/10.5194/hess-16-3607-2012>, 2012a.
- Albergel, C., de Rosnay, P., Balsamo, G., Isaksen, L., and Muñoz-Sabater, J.: Soil Moisture Analyses at ECMWF: Evaluation Using Global Ground-Based In Situ Observations, *Journal of Hydrometeorology*, 13, 1442–1460, <https://doi.org/10.1175/JHM-D-11-0107.1>, 2012b.



- Albergel, C., de Rosnay, P., Gruhier, C., Muñoz-Sabater, J., Hasenauer, S., Isaksen, L., Kerr, Y., and Wagner, W.: Evaluation of remotely sensed and modelled soil moisture products using global ground-based in situ observations, *Remote Sensing of Environment*, 118, 215–226, <https://doi.org/10.1016/j.rse.2011.11.017>, 2012c.
- Albergel, C., Dorigo, W., Balsamo, G., Muñoz-Sabater, J., de Rosnay, P., Isaksen, L., Brocca, L., de Jeu, R., and Wagner, W.: Monitoring multi-decadal satellite earth observation of soil moisture products through land surface reanalyses, *Remote Sensing of Environment*, 138, 77–89, <https://doi.org/10.1016/j.rse.2013.07.009>, 2013a.
- Albergel, C., Dorigo, W., Reichle, R. H., Balsamo, G., de Rosnay, P., Muñoz-Sabater, J., Isaksen, L., de Jeu, R., and Wagner, W.: Skill and Global Trend Analysis of Soil Moisture from Reanalyses and Microwave Remote Sensing, *Journal of Hydrometeorology*, 14, 1259–1277, <https://doi.org/10.1175/JHM-D-12-0161.1>, 2013b.
- Albergel, C., Dutra, E., Muñoz-Sabater, J., Haiden, T., Balsamo, G., Beljaars, A., Isaksen, L., de Rosnay, P., Sandu, I., and Wedi, N.: Soil temperature at ECMWF: An assessment using ground-based observations, *Journal of Geophysical Research: Atmospheres*, 120, 1361–1373, <https://doi.org/10.1002/2014JD022505>, 2015.
- Albergel, C., Munier, S., Bocher, A., Bonan, B., Zheng, Y., Draper, C., Leroux, D., and Calvet, J.-C.: LDAS-Monde Sequential Assimilation of Satellite Derived Observations Applied to the Contiguous US: An ERA-5 Driven Reanalysis of the Land Surface Variables, *Remote Sensing*, 10, 1627, <https://doi.org/10.3390/rs10101627>, 2018.
- Albergel, C., Zheng, Y., Bonan, B., Dutra, E., Rodríguez-Fernández, N., Munier, S., Draper, C., de Rosnay, P., Muñoz Sabater, J., Balsamo, G., Fairbairn, D., Meurey, C., and Calvet, J.-C.: Data assimilation for continuous global assessment of severe conditions over terrestrial surfaces, *Hydrology and Earth System Sciences*, 24, 4291–4316, <https://doi.org/10.5194/hess-24-4291-2020>, 2020.
- Alday, J. G., Camarero, J. J., Revilla, J., and Dios, V. R.: Similar diurnal, seasonal and annual rhythms in radial root expansion across two coexisting Mediterranean oak species, *Tree Physiology*, 2020.
- Alemohammad, S. H., Kolassa, J., Prigent, C., Aires, F., and Gentile, P.: Global downscaling of remotely sensed soil moisture using neural networks, *Hydrology and Earth System Sciences*, 22, 5341–5356, <https://doi.org/10.5194/hess-22-5341-2018>, 2018.
- An, R., Zhang, L., Wang, Z., Quaye-Ballard, J. A., You, J., Shen, X., Gao, W., Huang, L., Zhao, Y., and Ke, Z.: Validation of the ESA CCI soil moisture product in China, *International Journal of Applied Earth Observation and Geoinformation*, 48, 28–36, <https://doi.org/10.1016/j.jag.2015.09.009>, advances in the Validation and Application of Remotely Sensed Soil Moisture - Part 2, 2016.
- Ángel González-Zamora, Sánchez, N., Martínez-Fernández, J., and Wagner, W.: Root-zone plant available water estimation using the SMOS-derived soil water index, *Advances in Water Resources*, 96, 339–353, <https://doi.org/10.1016/j.advwatres.2016.08.001>, 2016.
- Angevine, W. M., Bazile, E., Legain, D., and Pino, D.: Land surface spinup for episodic modeling, *Atmospheric Chemistry and Physics*, 14, 8165–8172, <https://doi.org/10.5194/acp-14-8165-2014>, 2014.
- Anoop, S., Maurya, D. K., Rao, P. V. N., and Sekhar, M.: Validation and Comparison of LPRM Retrieved Soil Moisture Using AMSR2 Brightness Temperature at Two Spatial Resolutions in the Indian Region, *IEEE Geoscience and Remote Sensing Letters*, 14, 1561–1564, <https://doi.org/10.1109/LGRS.2017.2722542>, 2017.
- Ardö, J.: A 10-Year Dataset of Basic Meteorology and Soil Properties in Central Sudan, *Dataset Papers in Geosciences*, 2013, <https://doi.org/10.7167/2013/297973/dataset>, 2013.
- Bai, L., Lv, X., and Li, X.: Evaluation of Two SMAP Soil Moisture Retrievals Using Modeled- and Ground-Based Measurements, *Remote Sensing*, 11, 2891, <https://doi.org/10.3390/rs11242891>, 2019.
- Baldocchi, D., Falge, E., Gu, L., Olson, R., Hollinger, D., Running, S., Anthoni, P., Bernhofer, C., Davis, K., Evans, R., Fuentes, J., Goldstein, A., Katul, G., Law, B., Lee, X., Malhi, Y., Meyers, T., Munger, J., Oechel, W., and Richardson, F.: FLUXNET: A New Tool to Study



- the Temporal and Spatial Variability of Ecosystem-Scale Carbon Dioxide, Water Vapor, and Energy Flux Densities, ©2001 American Meteorological Society, 82, [https://doi.org/10.1175/1520-0477\(2001\)082<2415:FANTTS>2.3.CO;2](https://doi.org/10.1175/1520-0477(2001)082<2415:FANTTS>2.3.CO;2), 2001.
- Balsamo, G., Albergel, C., Beljaars, A., Boussetta, S., Brun, E., Cloke, H., Dee, D., Dutra, E., Muñoz Sabater, J., Pappenberger, F., de Rosnay, P., Stockdale, T., and Vitart, F.: ERA-Interim/Land: a global land surface reanalysis data set, *Hydrology and Earth System Sciences*, 19, 389–407, <https://doi.org/10.5194/hess-19-389-2015>, 2015.
- Bao, Y., Lin, L., Wu, S., Deng, K. A. K., and Petropoulos, G. P.: Surface soil moisture retrievals over partially vegetated areas from the synergy of Sentinel-1 and Landsat 8 data using a modified water-cloud model, *International Journal of Applied Earth Observation and Geoinformation*, 72, 76–85, <https://doi.org/10.1016/j.jag.2018.05.026>, 2018.
- Barbu, A. L., Calvet, J.-C., Mahfouf, J.-F., and Lafont, S.: Integrating ASCAT surface soil moisture and GEOV1 leaf area index into the SURFEX modelling platform: a land data assimilation application over France, *Hydrology and Earth System Sciences*, 18, 173–192, <https://doi.org/10.5194/hess-18-173-2014>, 2014.
- Bauer-Marschallinger, B., Paulik, C., Hochstöger, S., Mistelbauer, T., Modanesi, S., Ciabatta, L., Massari, C., Brocca, L., and Wagner, W.: Soil Moisture from Fusion of Scatterometer and SAR: Closing the Scale Gap with Temporal Filtering, *Remote Sensing*, 10, 1030, <https://doi.org/10.3390/rs10071030>, 2018.
- Bauer-Marschallinger, B., Freeman, V., Cao, S., Paulik, C., Schaufler, S., Stachl, T., Modanesi, S., Massari, C., Ciabatta, L., Brocca, L., and Wagner, W.: Toward Global Soil Moisture Monitoring With Sentinel-1: Harnessing Assets and Overcoming Obstacles, *IEEE Transactions on Geoscience and Remote Sensing*, 57, 520–539, <https://doi.org/10.1109/TGRS.2018.2858004>, 2019.
- Beck, H. E., Pan, M., Miralles, D. G., Reichle, R. H., Dorigo, W. A., Hahn, S., Sheffield, J., Karthikeyan, L., Balsamo, G., Parinussa, R. M., van Dijk, A. I. J. M., Du, J., Kimball, J. S., Vergopolan, N., and Wood, E. F.: Evaluation of 18 satellite- and model-based soil moisture products using in situ measurements from 826 sensors, *Hydrology and Earth System Sciences Discussions*, 2020, 1–35, <https://doi.org/10.5194/hess-2020-184>, 2020.
- Bell, J., Palecki, M., Baker, B., Collins, W., Lawrimore, J., Leeper, R., Hall, M., Kochendorfer, J., Meyers, T., Wilson, T., and Diamond, H.: U.S. Climate Reference Network Soil Moisture and Temperature Observations, *Journal of Hydrometeorology*, 14, 977–988, <https://doi.org/10.1175/JHM-D-12-0146.1>, 2013.
- Beyrich, F. and Adam, W.: Site and Data Report for the Lindenberg Reference Site in CEOP - Phase 1, *Berichte des Deutschen Wetterdienstes*, 230, Offenbach am Main, 2007, 2007.
- Bi, H., Ma, J., Zheng, W., and Zeng, J.: Comparison of soil moisture in GLDAS model simulations and in situ observations over the Tibetan Plateau, *Journal of Geophysical Research: Atmospheres*, 121, 2658–2678, <https://doi.org/10.1002/2015JD024131>, 2016.
- Biddoccu, M., Ferraris, S., Opsi, F., and Cavallo, E.: Long-term monitoring of soil management effects on runoff and soil erosion in sloping vineyards in Alto Monferrato (North West Italy), *Soil and Tillage Research*, 155, 176–189, <https://doi.org/10.1016/j.still.2015.07.005>, 2016.
- Bircher, S., Skou, N., Jensen, K., Walker, J., and Rasmussen, L.: A soil moisture and temperature network for SMOS validation in Western Denmark, *Hydrology and Earth System Sciences*, 16, <https://doi.org/10.5194/hess-16-1445-2012>, 2012.
- Blöschl, G., Blaschke, A. P., Broer, M., Bucher, C., Carr, G., Chen, X., Eder, A., Exner-Kittridge, M., Farnleitner, A., Flores-Orozco, A., Haas, P., Hogan, P., Kazemi Amiri, A., Oismüller, M., Parajka, J., Silasari, R., Stadler, P., Strauss, P., Vreugdenhil, M., Wagner, W., and Zessner, M.: The Hydrological Open Air Laboratory (HOAL) in Petzenkirchen: a hypothesis-driven observatory, *Hydrology and Earth System Sciences*, 20, 227–255, <https://doi.org/10.5194/hess-20-227-2016>, 2016.



- Blyverket, J., Hamer, P., Bertino, L., Albergel, C., Fairbairn, D., and Lahoz, W.: An Evaluation of the EnKF vs. EnOI and the Assimilation of SMAP, SMOS and ESA CCI Soil Moisture Data over the Contiguous US, *Remote Sensing*, 11, 478, <https://doi.org/10.3390/rs11050478>, 2019.
- Blyverket, J.; Hamer, P. B. L. A. C. F. D. L. W.: An Evaluation of the EnKF vs. EnOI and the Assimilation of SMAP, SMOS and ESA CCI Soil Moisture Data over the Contiguous US., *Remote Sens.*, 11, 012 046, <https://doi.org/10.3390/rs11050478>, 2019.
- Bogena, H., Huisman, J., Oberdörster, C., and Vereecken, H.: Evaluation of a low-cost soil water content sensor for wireless network applications, *Journal of Hydrology*, 344, 32–42, <https://doi.org/10.1016/j.jhydrol.2007.06.032>, 2007.
- Bogena, H., Kunkel, R., Pütz, T., Vereecken, H., Kruger, E., Zacharias, S., Dietrich, P., Wollschläger, U., Kunstmann, H., Papen, H., Schmid, H., Munch, J., Priesack, E., Schwank, M., Bens, O., Brauer, A., Borg, E., and Hajnsek, I.: TERENO - Long-term monitoring network for terrestrial environmental research, *Hydrologie und Wasserbewirtschaftung*, 56, 138–143, 2012.
- Bogena, H. R.: TERENO: German network of terrestrial environmental observatories, *Journal of large-scale research facilities JLSRF*, 2, 52, 2016.
- Bogena, H. R., Huisman, J. A., Güntner, A., Hübner, C., Kusche, J., Jonard, F., Vey, S., and Vereecken, H.: Emerging methods for noninvasive sensing of soil moisture dynamics from field to catchment scale: a review: Emerging methods for noninvasive sensing of soil moisture dynamics, *Wiley Interdisciplinary Reviews: Water*, 2, 635–647, <https://doi.org/10.1002/wat2.1097>, 2015.
- Bonney, R., Cooper, C. B., Dickinson, J., Kelling, S., Phillips, T., Rosenberg, K. V., and Shirk, J.: Citizen Science: A Developing Tool for Expanding Science Knowledge and Scientific Literacy, *BioScience*, 59, 977–984, <https://doi.org/10.1525/bio.2009.59.11.9>, 2009.
- Boussetta, S., Balsamo, G., Dutra, E., Beljaars, A., and Albergel, C.: Assimilation of surface albedo and vegetation states from satellite observations and their impact on numerical weather prediction, *Remote Sensing of Environment*, 163, 111–126, <https://doi.org/10.1016/j.rse.2015.03.009>, 2015.
- Brocca, L., Morbidelli, R., Melone, F., and Moramarco, T.: Soil moisture spatial variability in experimental areas of central Italy, *Journal of Hydrology*, 333, 356–373, <https://doi.org/10.1016/j.jhydrol.2006.09.004>, 2007.
- Brocca, L., Hasenauer, S., Lacava, T., Melone, F., Moramarco, T., Wagner, W., A. D., Matgen, P., Martínez-Fernández, J., Llorens, P., Latron, J., Martin, C., and Bittelli, M.: Soil moisture estimation through ASCAT and AMSR-E sensors: An intercomparison and validation study across Europe, *Remote Sensing of Environment*, 115, 3390–3408, <https://doi.org/10.1016/j.rse.2011.08.003>, 2011.
- Brocca, L., Ciabatta, L., Massari, C., Moramarco, T., Hahn, S., Hasenauer, S., Kidd, R., Dorigo, W., Wagner, W., and Levizzani, V.: Soil as a natural rain gauge: Estimating global rainfall from satellite soil moisture data, *Journal of Geophysical Research: Atmospheres*, 119, 5128–5141, <https://doi.org/10.1002/2014jd021489>, 2014a.
- Brocca, L., Zucco, G., Mittelbach, H., Moramarco, T., and Seneviratne, S. I.: Absolute versus temporal anomaly and percent of saturation soil moisture spatial variability for six networks worldwide, *Water Resources Research*, 50, 5560–5576, <https://doi.org/10.1002/2014WR015684>, 2014b.
- Brocca, L., Massari, C., Ciabatta, L., Moramarco, T., Penna, D., Zuecco, G., Pianezzola, L., Borga, M., Matgen, P., and Martínez-Fernández, J.: Rainfall estimation from in situ soil moisture observations at several sites in Europe: an evaluation of the SM2RAIN algorithm, *Journal of Hydrology and Hydromechanics*, 63, 201–209, <https://doi.org/10.1515/johh-2015-0016>, 2015.
- Brocca, L., Ciabatta, L., Massari, C., Camici, S., and Tarpanelli, A.: Soil Moisture for Hydrological Applications: Open Questions and New Opportunities, *Water*, 9, 140, <https://doi.org/10.3390/w9020140>, 2017.
- Calvet, J.-C., Fritz, N., Froissard, F., Suquia, D., Petitpa, A., and Pignat, B.: In situ soil moisture observations for the CAL/VAL of SMOS: the SMOSMANIA network, pp. 1196–1199, <https://doi.org/10.1109/IGARSS.2007.4423019>, 2007.



- Calvet, J.-C., Fritz, N., Berne, C., Piguet, B., Maurel, W., and Meurey, C.: Deriving pedotransfer functions for soil quartz fraction in southern France from reverse modeling, *SOIL*, 2, 615–629, <https://doi.org/10.5194/soil-2-615-2016>, 2016.
- Cammalleri, C., Micale, F., and Vogt, J.: On the value of combining different modelled soil moisture products for European drought monitoring, *Journal of Hydrology*, 525, 547–558, <https://doi.org/10.1016/j.jhydrol.2015.04.021>, 2015.
- 1145 Campo, L., Castelli, F., Caparrini, F., and Entekhabi, D.: Production of a 5-years Dataset of Soil Moisture Maps on Italian Territory with an Operational Algorithm, *European Water*, 35, 45–56, 2011.
- Canisius, F.: Calibration of Casselman, Ontario Soil Moisture Monitoring Network, Agriculture and Agri-Food Canada, Ottawa, ON, 37pp, 2011.
- Capello, G., Biddoccu, M., Ferraris, S., and Cavallo, E.: Effects of Tractor Passes on Hydrological and Soil Erosion Processes in Tilled and  
 1150 Grassed Vineyards, *Water*, 11, 2118, <https://doi.org/10.3390/w11102118>, 2019.
- Castell, N., Dauge, F. R., Schneider, P., Vogt, M., Lerner, U., Fishbain, B., Broday, D., and Bartonova, A.: Can commercial low-cost sensor platforms contribute to air quality monitoring and exposure estimates?, *Environment International*, 99, 293–302, <https://doi.org/10.1016/j.envint.2016.12.007>, 2017.
- Chawla, S., Bachhtey, S., Gupta, V., Sharma, S., Seth, S., Gandhi, T., Varshney, S., Mehta, S., and Jha, R.: Low Cost Soil Moisture Sensors  
 1155 and their Application in Automatic Irrigation System, p. 4, 2019.
- Chen, F., Crow, W. T., Colliander, A., Cosh, M. H., Jackson, T. J., Bindlish, R., Reichle, R. H., Chan, S. K., Bosch, D. D., Starks, P. J., Goodrich, D. C., and Seyfried, M. S.: Application of Triple Collocation in Ground-Based Validation of Soil Moisture Active/Passive (SMAP) Level 2 Data Products, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 10, 489–502, <https://doi.org/10.1109/JSTARS.2016.2569998>, 2017.
- 1160 Chen, N., Xiang, Z., and Wang, C.: Integrated open geospatial web service enabled cyber-physical information infrastructure for precision agriculture monitoring, *Computers and Electronics in Agriculture*, 111, <https://doi.org/10.1016/j.compag.2014.12.009>, 2015a.
- Chen, N., Xiao, C., Pu, F., Wang, X., Wang, C., Wang, Z., and Gong, J.: Cyber-Physical Geographical Information Service-Enabled Control of Diverse In-Situ Sensors, *Sensors (Basel, Switzerland)*, 15, 2565–92, <https://doi.org/10.3390/s150202565>, 2015b.
- Chen, X., Su, Y., Liao, J., Shang, J., Dong, T., Wang, C., Liu, W., Zhou, G., and Liu, L.: Detecting significant decreasing trends of land  
 1165 surface soil moisture in eastern China during the past three decades (1979–2010), *Journal of Geophysical Research: Atmospheres*, 121, 5177–5192, <https://doi.org/10.1002/2015JD024676>, 2016.
- Chen, Y., Sun, L., Wang, W., and Pei, Z.: Application of Sentinel 2 data for drought monitoring in Texas, America, in: 2019 8th International Conference on Agro-Geoinformatics (Agro-Geoinformatics), pp. 1–4, <https://doi.org/10.1109/Agro-Geoinformatics.2019.8820491>, 2019.
- Chen, Y., Feng, X., and Fu, B.: A new dataset of satellite observation-based global surface soil moisture covering 2003–2018, *Earth System  
 1170 Science Data Discussions*, 2020, 1–46, <https://doi.org/10.5194/essd-2020-59>, 2020.
- Chew, C. and Small, E.: Estimating inundation extent using CYGNSS data: A conceptual modeling study, *Remote Sensing of Environment*, 246, 111 869, <https://doi.org/10.1016/j.rse.2020.111869>, 2020.
- Colliander, A., Jackson, T., Bindlish, R., Chan, S., Das, N., Kim, S., Cosh, M., Dunbar, R., Dang, L., Pashaian, L., Asanuma, J., Aida, K., Berg, A., Rowlandson, T., Bosch, D., Caldwell, T., Caylor, K., Goodrich, D., al Jassar, H., Lopez-Baeza, E., Martínez-Fernández, J.,  
 1175 González-Zamora, A., Livingston, S., McNairn, H., Pacheco, A., Moghaddam, M., Montzka, C., Notarnicola, C., Niedrist, G., Pellarin, T., Prueger, J., Pulliainen, J., Rautiainen, K., Ramos, J., Seyfried, M., Starks, P., Su, Z., Zeng, Y., van der Velde, R., Thibeault, M., Dorigo, W., Vreugdenhil, M., Walker, J., Wu, X., Monerris, A., O'Neill, P., Entekhabi, D., Njoku, E., and Yueh, S.: Validation of SMAP surface soil





- moisture products with core validation sites, *Remote Sensing of Environment*, 191, 215–231, <https://doi.org/10.1016/j.rse.2017.01.021>, 2017.
- 1180 Cook, D. R.: Soil temperature and moisture profile (STAMP) system handbook, Tech. rep., DOE Office of Science Atmospheric Radiation Measurement (ARM) Program ..., 2016a.
- Cook, D. R.: Soil Water and Temperature System (SWATS) Instrument Handbook, Tech. rep., DOE Office of Science Atmospheric Radiation Measurement (ARM) Program ..., 2016b.
- Cook, D. R. and Sullivan, R. C.: Surface Energy Balance System (SEBS) Instrument Handbook, Tech. rep., DOE Office of Science Atmospheric Radiation Measurement (ARM) Program ..., 2018.
- 1185 Coopersmith, E. J., Cosh, M. H., Bindlish, R., and Bell, J.: Comparing AMSR-E soil moisture estimates to the extended record of the U.S. Climate Reference Network (USCRN), *Advances in Water Resources*, 85, 79–85, <https://doi.org/10.1016/j.advwatres.2015.09.003>, 2015.
- Crow, W. T., Berg, A. A., Cosh, M. H., Loew, A., Mohanty, B. P., Panciera, R., de Rosnay, P., Ryu, D., and Walker, J. P.: Upscaling sparse ground-based soil moisture observations for the validation of coarse-resolution satellite soil moisture products: UPSCALING SOIL MOISTURE, *Reviews of Geophysics*, 50, <https://doi.org/10.1029/2011RG000372>, [Online; accessed 2020-07-07], 2012.
- 1190 Cui, C., Xu, J., Zeng, J., Chen, K.-S., Bai, X., Lu, H., Chen, Q., and Zhao, T.: Soil Moisture Mapping from Satellites: An Intercomparison of SMAP, SMOS, FY3B, AMSR2, and ESA CCI over Two Dense Network Regions at Different Spatial Scales, *Remote Sensing*, 10, 33, <https://doi.org/10.3390/rs10010033>, 2017.
- Dabrowska-Zielinska, K., Musial, J., Malinska, A., Budzynska, M., Gurdak, R., Kiryla, W., Bartold, M., and Grzybowski, P.: Soil Moisture in the Biebrza Wetlands Retrieved from Sentinel-1 Imagery, *Remote Sensing*, 10, 1979, <https://doi.org/10.3390/rs10121979>, 2018.
- 1195 de Jeu, R. and Dorigo, W.: On the importance of satellite observed soil moisture, *International Journal of Applied Earth Observation and Geoinformation*, 45, 107–109, <https://doi.org/10.1016/j.jag.2015.10.007>, advances in the Validation and Application of Remotely Sensed Soil Moisture - Part 1, 2016.
- de Jeu, R. A., Holmes, T. R., Parinussa, R. M., and Owe, M.: A spatially coherent global soil moisture product with improved temporal resolution, *Journal of Hydrology*, 516, 284–296, <https://doi.org/10.1016/j.jhydrol.2014.02.015>, determination of soil moisture: Measurements and theoretical approaches, 2014.
- 1200 de Rosnay, P., Muñoz, J., Sabater, C. A., Lawrence, H., Isaksen, L., and English, S.: ECMWF Final Report on SMOS brightness temperature activities over land: Monitoring and Data Assimilation, 2019.
- Dee, D. P., Uppala, S. M., Simmons, A., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M., Balsamo, G., Bauer, d. P., et al.: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, *Quarterly Journal of the royal meteorological society*, 137, 553–597, 2011.
- 1205 Deng, Y., Wang, S., Bai, X., Luo, G., Wu, L., Chen, F., Wang, J., Li, C., Yang, Y., Hu, Z., Tian, S., and Lu, Q.: Vegetation greening intensified soil drying in some semi-arid and arid areas of the world, *Agricultural and Forest Meteorology*, 292–293, 108 103, <https://doi.org/10.1016/j.agrformet.2020.108103>, 2020a.
- 1210 Deng, Y., Wang, S., Bai, X., Luo, G., Wu, L., Chen, F., Wang, J., Li, Q., Li, C., Yang, Y., Hu, Z., and Tian, S.: Spatiotemporal dynamics of soil moisture in the karst areas of China based on reanalysis and observations data, *Journal of Hydrology*, 585, 124 744, <https://doi.org/10.1016/j.jhydrol.2020.124744>, 2020b.
- Deng, Y., Wang, S., Bai, X., Wu, L., Cao, Y., Li, H., Wang, M., Li, C., Yang, Y., Hu, Z., Tian, S., and Lu, Q.: Comparison of soil moisture products from microwave remote sensing, land model, and reanalysis using global ground observations, *Hydrological Processes*, 34, 836–851, <https://doi.org/10.1002/hyp.13636>, 2020c.
- 1215





- Dente, L., Su, Z., and Wen, J.: Validation of SMOS soil moisture products over the Maqu and Twente regions, *Sensors*, 12, 9965–9986, 2012.
- Dente, L., Vekerdy, Z., de Jeu, R., and Su, Z.: Seasonality and autocorrelation of satellite-derived soil moisture products, *International Journal of Remote Sensing*, 34, 3231–3247, <https://doi.org/10.1080/01431161.2012.716923>, 2013.
- Dirmeyer, P. A., Wu, J., Norton, H. E., Dorigo, W. A., Quiring, S. M., Ford, T. W., Santanello, Joseph A., J., Bosilovich, M. G., Ek, M. B.,  
 1220 Koster, R. D., Balsamo, G., and Lawrence, D. M.: Confronting Weather and Climate Models with Observational Data from Soil Moisture Networks over the United States, *Journal of Hydrometeorology*, 17, 1049–1067, <https://doi.org/10.1175/JHM-D-15-0196.1>, 2016.
- Domínguez-Niño, J. M., Bogen, H. R., Huisman, J. A., Schilling, B., and Casadesús, J.: On the Accuracy of Factory-Calibrated Low-Cost Soil Water Content Sensors, *Sensors*, 19, 3101, <https://doi.org/10.3390/s19143101>, 2019.
- Dong, J., Crow, W. T., Tobin, K. J., Cosh, M. H., Bosch, D. D., Starks, P. J., Seyfried, M., and Collins, C. H.: Comparison of microwave  
 1225 remote sensing and land surface modeling for surface soil moisture climatology estimation, *Remote Sensing of Environment*, 242, 111 756, <https://doi.org/10.1016/j.rse.2020.111756>, 2020.
- Dorigo, W., van Oevelen, P., Wagner, W., Drusch, M., Mecklenburg, S., Robock, A., and Jackson, T.: A New International Network for in Situ Soil Moisture Data, *EOS*, 92, 141–142, <https://doi.org/10.1029/2011EO170001>, 2011a.
- Dorigo, W., Xaver, A., Vreugdenhil, M., Gruber, A., Dostálová, A., Sanchis-Dufau, A. D., Zamojski, D., Cordes, C., Wagner, W., and Drusch,  
 1230 M.: Global Automated Quality Control of In Situ Soil Moisture Data from the International Soil Moisture Network, *Vadose Zone Journal*, 12, <https://doi.org/10.2136/vzj2012.0097>, 2013.
- Dorigo, W., Gruber, A., Jeu, R. D., Wagner, W., Stacke, T., Loew, A., Albergel, C., Brocca, L., Chung, D., Parinussa, R., and Kidd, R.: Evaluation of the ESA CCI soil moisture product using ground-based observations, *Remote Sensing of Environment*, 162, 380–395, <https://doi.org/10.1016/j.rse.2014.07.023>, 2015.
- 1235 Dorigo, W., Wagner, W., Albergel, C., Albrecht, F., Balsamo, G., Brocca, L., Chung, D., Ertl, M., Forkel, M., Gruber, A., Haas, E., Hamer, P. D., Hirschi, M., Ikonen, J., de Jeu, R., Kidd, R., Lahoz, W., Liu, Y. Y., Miralles, D., Mistelbauer, T., Nicolai-Shaw, N., Parinussa, R., Pratola, C., Reimer, C., van der Schalie, R., Seneviratne, S. I., Smolander, T., and Lecomte, P.: ESA CCI Soil Moisture for improved Earth system understanding: State-of-the art and future directions, *Remote Sensing of Environment*, 203, 185–215, <https://doi.org/10.1016/j.rse.2017.07.001>, 2017.
- 1240 Dorigo, W. A., Scipal, K., Parinussa, R. M., Liu, Y. Y., Wagner, W., de Jeu, R. A. M., and Naeimi, V.: Error characterisation of global active and passive microwave soil moisture datasets, *Hydrology and Earth System Sciences*, 14, 2605–2616, <https://doi.org/10.5194/hess-14-2605-2010>, 2010.
- Dorigo, W. A., Wagner, W., Hohensinn, R., Hahn, S., Paulik, C., Xaver, A., Gruber, A., Drusch, M., Mecklenburg, S., van Oevelen, P., Robock, A., and Jackson, T.: The International Soil Moisture Network: a data hosting facility for global in situ soil moisture measurements,  
 1245 *Hydrology and Earth System Sciences*, 15, 1675–1698, <https://doi.org/10.5194/hess-15-1675-2011>, 2011b.
- Draper, C. and Reichle, R. H.: Assimilation of Satellite Soil Moisture for Improved Atmospheric Reanalyses, *Monthly Weather Review*, 147, 2163–2188, <https://doi.org/10.1175/MWR-D-18-0393.1>, 2019.
- Du, J., Kimball, J. S., and Jones, L. A.: Passive Microwave Remote Sensing of Soil Moisture Based on Dynamic Vegetation Scattering Properties for AMSR-E, *IEEE Transactions on Geoscience and Remote Sensing*, 54, 597–608, 2016.
- 1250 Ebrahimi, M., Alavipanah, S. K., Hamzeh, S., Amiraslani, F., Neysani Samany, N., and Wigneron, J.-P.: Exploiting the synergy between SMAP and SMOS to improve brightness temperature simulations and soil moisture retrievals in arid regions, *Journal of Hydrology*, 557, 740–752, <https://doi.org/10.1016/j.jhydrol.2017.12.051>, 2018.



- Ebrahimi-Khusfi, M., Alavipanah, S. K., Hamzeh, S., Amiraslani, F., Neysani Samany, N., and Wigneron, J.-P.: Comparison of soil moisture retrieval algorithms based on the synergy between SMAP and SMOS-IC, *International Journal of Applied Earth Observation and Geoinformation*, 67, 148–160, <https://doi.org/10.1016/j.jag.2017.12.005>, 2018.
- Ebtehaj, A. and Bras, R. L.: A physically constrained inversion for high-resolution passive microwave retrieval of soil moisture and vegetation water content in L-band, *Remote Sensing of Environment*, 233, 111 346, 2019.
- Enenkel, M., Reimer, C., Dorigo, W., Wagner, W., Pfeil, I., Parinussa, R., and De Jeu, R.: Combining satellite observations to develop a global soil moisture product for near-real-time applications, *Hydrology and Earth System Sciences*, 20, 4191–4208, <https://doi.org/10.5194/hess-20-4191-2016>, 2016.
- Entekhabi, D., Njoku, E. G., O'Neill, P. E., Kellogg, K. H., Crow, W. T., Edelstein, W. N., Entin, J. K., Goodman, S. D., Jackson, T. J., Johnson, J., Kimball, J., Piepmeier, J. R., Koster, R. D., Martin, N., McDonald, K. C., Moghaddam, M., Moran, S., Reichle, R., Shi, J. C., Spencer, M. W., Thurman, S. W., Tsang, L., and Van Zyl, J.: The Soil Moisture Active Passive (SMAP) Mission, *Proceedings of the IEEE*, 98, 704–716, <https://doi.org/10.1109/JPROC.2010.2043918>, 2010.
- Eroglu, O., Kurum, M., Boyd, D., and Gurbuz, A. C.: High spatio-temporal resolution CYGNSS soil moisture estimates using artificial neural networks, *Remote Sensing*, 11, 2272, 2019.
- Esposito, G., Matano, F., and Scepi, G.: Analysis of Increasing Flash Flood Frequency in the Densely Urbanized Coastline of the Campi Flegrei Volcanic Area, Italy, *Frontiers in Earth Science*, 6, 63, <https://doi.org/10.3389/feart.2018.00063>, 2018.
- Fairbairn, D., Barbu, A. L., Mahfouf, J.-F., Calvet, J.-C., and Gelati, E.: Comparing the ensemble and extended Kalman filters for in situ soil moisture assimilation with contrasting conditions, *Hydrology and Earth System Sciences*, 19, 4811–4830, <https://doi.org/10.5194/hess-19-4811-2015>, 2015.
- Fairbairn, D., de Rosnay, P., and Browne, P. A.: The New Stand-Alone Surface Analysis at ECMWF: Implications for Land–Atmosphere DA Coupling, *Journal of Hydrometeorology*, 20, 2023–2042, <https://doi.org/10.1175/JHM-D-19-0074.1>, 2019.
- Famiglietti, J. S., Ryu, D., Berg, A. A., Rodell, M., and Jackson, T. J.: Field observations of soil moisture variability across scales: SOIL MOISTURE VARIABILITY ACROSS SCALES, *Water Resources Research*, 44, <https://doi.org/10.1029/2006WR005804>, [Online; accessed 2020-06-18], 2008.
- Fang, B., Lakshmi, V., Bindlish, R., and Jackson, T.: AMSR2 Soil Moisture Downscaling Using Temperature and Vegetation Data, *Remote Sensing*, 10, 1575, <https://doi.org/10.3390/rs10101575>, 2018.
- Fang, B., Lakshmi, V., Bindlish, R., Jackson, T. J., and Liu, P.-W.: Downscaling and Validation of SMAP Radiometer Soil Moisture in CONUS, in: *IGARSS 2019-2019 IEEE International Geoscience and Remote Sensing Symposium*, pp. 6194–6197, IEEE, 2019.
- Fang, B., Lakshmi, V., Bindlish, R., Jackson, T. J., and Liu, P.-W.: Evaluation and validation of a high spatial resolution satellite soil moisture product over the Continental United States, *Journal of Hydrology*, 588, 125 043, <https://doi.org/10.1016/j.jhydrol.2020.125043>, 2020.
- FAO/IIASA/ISRIC/ISS-CAS/JRC: Harmonized World Soil Database (version 1.1). FAO, Rome, Italy and IIASA, Laxenburg, Austria., <http://www.fao.org/3/a-aq361e.pdf>, 2009.
- Fascetti, F., Pierdicca, N., Pulvirenti, L., Crapolicchio, R., and Sabater, J. M.: Soil moisture comparison through triple and quadruple collocation between: Metop, ERA, SMOS and in-situ data, in: *2014 IEEE Geoscience and Remote Sensing Symposium*, pp. 2451–2454, 2014.
- Fascetti, F., Pierdicca, N., Crapolicchio, R., Pulvirenti, L., and Muoz-Sabater, J.: An assessment of SMOS version 6.20 products through Triple and Quadruple Collocation techniques considering ASCAT, ERA/Interim LAND, ISMN and SMAP soil moisture data, in: *2016 14th Specialist Meeting on Microwave Radiometry and Remote Sensing of the Environment (MicroRad)*, pp. 91–94, 2016.



- Fascetti, F., Pierdicca, N., Pulvirenti, L., Crapolicchio, R., and Muñoz-Sabater, J.: A comparison of ASCAT and SMOS soil moisture retrievals over Europe and Northern Africa from 2010 to 2013, *International Journal of Applied Earth Observation and Geoinformation*, 45, 135–142, <https://doi.org/10.1016/j.jag.2015.09.008>, advances in the Validation and Application of Remotely Sensed Soil Moisture - Part 1, 2016.
- 1295 Fascetti, F., Pierdicca, N., Pulvirenti, L., and Crapolicchio, R.: Error characterization of SMOS, ASCAT, SMAP, ERA and ISMN soil moisture products: Automatic detection of cross-correlation error through extended quadruple collocation, in: 2017 IEEE International Geoscience and Remote Sensing Symposium (IGARSS), pp. 4154–4157, <https://doi.org/10.1109/IGARSS.2017.8127916>, 2017.
- Fernandez-Moran, R., Al-Yaari, A., Mialon, A., Mahmoodi, A., Al Bitar, A., De Lannoy, G., Rodriguez-Fernandez, N., Lopez-Baeza, E., Kerr, Y., and Wigneron, J.-P.: SMOS-IC: An Alternative SMOS Soil Moisture and Vegetation Optical Depth Product, *Remote Sensing*, 9, <https://doi.org/10.3390/rs9050457>, 2017a.
- 1300 Fernandez-Moran, R., Wigneron, J.-P., De Lannoy, G., Lopez-Baeza, E., Parrens, M., Mialon, A., Mahmoodi, A., Al-Yaari, A., Bircher, S., Al Bitar, A., Richaume, P., and Kerr, Y.: A new calibration of the effective scattering albedo and soil roughness parameters in the SMOS SM retrieval algorithm, *International Journal of Applied Earth Observation and Geoinformation*, 62, 27–38, <https://doi.org/10.1016/j.jag.2017.05.013>, 2017b.
- 1305 Foucras, M., Zribi, M., Albergel, C., Baghdadi, N., Calvet, J.-C., and Pellarin, T.: Estimating 500-m Resolution Soil Moisture Using Sentinel-1 and Optical Data Synergy, *Water*, 12, 866, <https://doi.org/10.3390/w12030866>, 2020.
- Fritz, S., Fonte, C., and See, L.: The Role of Citizen Science in Earth Observation, *Remote Sensing*, 9, 357, <https://doi.org/10.3390/rs9040357>, 2017.
- Fuchsberger, J., Kirchengast, G., and Kabas, T.: WegenerNet high-resolution weather and climate data 2007 to 2019, *Earth System Science Data Discussions*, 2020, 1–49, <https://doi.org/10.5194/essd-2020-302>, 2020.
- 1310 Galle, S., Manuela, G., Peugeot, C., Bouzou-Moussa, I., Cappelaere, B., Demarty, J., Mougin, E., Panthou, G., Adjomayi, P., Agbossou, E., Abdramane, B., Boucher, M., Cohard, J.-M., Descloitres, M., Descroix, L., Diawara, M., Do, M., Favreau, G., Fabrice, G., and Wilcox, C.: AMMA-CATCH, a critical zone observatory in West Africa monitoring a region in transition, *Vadose Zone Journal*, 17, 1–24, <https://doi.org/10.2136/vzj2018.03.0062>, 2018.
- 1315 Gao, L., Sadeghi, M., and Ebtehaj, A.: Microwave retrievals of soil moisture and vegetation optical depth with improved resolution using a combined constrained inversion algorithm: Application for SMAP satellite, *Remote Sensing of Environment*, 239, 111 662, <https://doi.org/10.1016/j.rse.2020.111662>, 2020.
- Gao, L., Sadeghi, M., Feldman, A. F., and Ebtehaj, A.: A Spatially Constrained Multichannel Algorithm for Inversion of a First-Order Microwave Emission Model at L-Band, *IEEE Transactions on Geoscience and Remote Sensing*, pp. 1–13, <https://doi.org/10.1109/TGRS.2020.2987490>, 2020.
- 1320 Gardner, W. and Kirkham, D.: Determination of soil moisture by neutron scattering, *Soil Science*, 73, 391–401, 1952.
- GCOS: The Global Climate Observing System, <https://gcoss.wmo.int/en/gcos-implementation-plan>, 2016.
- Ghilain, N., Arboleda, A., Batelaan, O., Ardö, J., Trigo, I., Barrios, J.-M., and Gellens-Meulenberghs, F.: A new retrieval algorithm for soil moisture index from thermal infrared sensor on-board geostationary satellites over Europe and Africa and its validation, *Remote Sensing*, 11, 1968, 2019.
- 1325 González-Teruel, J., Torres-Sánchez, R., Blaya-Ros, P., Toledo-Moreo, A., Jiménez-Buendía, M., and Soto-Valles, F.: Design and Calibration of a Low-Cost SDI-12 Soil Moisture Sensor, *Sensors*, 19, 491, <https://doi.org/10.3390/s19030491>, 2019.



- González-Zamora, A., Sánchez, N., Gumuzzio, A., Piles, M., Olmedo, E., and Martínez-Fernández, J.: Validation of SMOS L2 and L3 soil moisture products over the Duero Basin at different spatial scales, *ISPRS - International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, XL-7/W3, 1183–1188, <https://doi.org/10.5194/isprsarchives-XL-7-W3-1183-2015>, 2015.
- González-Zamora, A., Sánchez, N., and Martínez-Fernández, J.: Validation of Aquarius Soil Moisture Products Over the Northwest of Spain: A Comparison With SMOS, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 9, 2763–2769, 2016.
- González-Zamora, Á., Sánchez, N., Pablos, M., and Martínez-Fernández, J.: CCI soil moisture assessment with SMOS soil moisture and in situ data under different environmental conditions and spatial scales in Spain, *Remote Sensing of Environment*, 225, 469–482, <https://doi.org/10.1016/j.rse.2018.02.010>, 2019.
- González-Zamora, , Sanchez, N., Pablos, M., and Martínez-Fernández, J.: CCI soil moisture assessment with SMOS soil moisture and in situ data under different environmental conditions and spatial scales in Spain, *Remote Sensing of Environment*, 225, <https://doi.org/10.1016/j.rse.2018.02.010>, 2018.
- Grainger, A.: Citizen Observatories and the New Earth Observation Science, *Remote Sensing*, 9, 153, <https://doi.org/10.3390/rs9020153>, 2017.
- Greifeneder, F., Khamala, E., Sendabo, D., Wagner, W., Zebisch, M., Farah, H., and Notarnicola, C.: Detection of soil moisture anomalies based on Sentinel-1, *Physics and Chemistry of the Earth, Parts A/B/C*, 112, 75–82, <https://doi.org/https://doi.org/10.1016/j.pce.2018.11.009>, 18th WaterNet/WARFSA/GWPSA Symposium on Integrated Water Resources Development and Management: Innovative Technological Advances for Water Security in Eastern and Southern Africa., 2019.
- Gruber, A., Dorigo, W., Zwieback, S., Xaver, A., and Wagner, W.: Characterizing Coarse-Scale Representativeness of in situ Soil Moisture Measurements from the International Soil Moisture Network, *Vadose Zone Journal*, 12, vzj2012.0170, <https://doi.org/10.2136/vzj2012.0170>, 2013.
- Gruber, A., Paloscia, S., Santi, E., Notarnicola, C., Pasolli, L., Smolander, T., Pulliainen, J., Mittelbach, H., Dorigo, W., and Wagner, W.: Performance inter-comparison of soil moisture retrieval models for the MetOp-A ASCAT instrument, in: 2014 IEEE Geoscience and Remote Sensing Symposium, pp. 2455–2458, <https://doi.org/10.1109/IGARSS.2014.6946969>, 2014.
- Gruber, A., Crow, W., Dorigo, W., and Wagner, W.: The potential of 2D Kalman filtering for soil moisture data assimilation, *Remote Sensing of Environment*, 171, 137–148, <https://doi.org/10.1016/j.rse.2015.10.019>, 2015.
- Gruber, A., Su, C.-H., Crow, W. T., Zwieback, S., Dorigo, W. A., and Wagner, W.: Estimating error cross-correlations in soil moisture data sets using extended collocation analysis, *Journal of Geophysical Research: Atmospheres*, 121, 1208–1219, <https://doi.org/10.1002/2015JD024027>, 2016.
- Gruber, A., Dorigo, W. A., Crow, W., and Wagner, W.: Triple Collocation-Based Merging of Satellite Soil Moisture Retrievals, *IEEE Transactions on Geoscience and Remote Sensing*, 55, 6780–6792, <https://doi.org/10.1109/TGRS.2017.2734070>, 2017.
- Gruber, A., Crow, W. T., and Dorigo, W. A.: Assimilation of Spatially Sparse In Situ Soil Moisture Networks into a Continuous Model Domain, *Water Resources Research*, 54, 1353–1367, <https://doi.org/10.1002/2017WR021277>, 2018.
- Gruber, A., De Lannoy, G., and Crow, W.: A Monte Carlo based adaptive Kalman filtering framework for soil moisture data assimilation, *Remote Sensing of Environment*, 228, 105–114, 2019a.
- Gruber, A., Scanlon, T., van der Schalie, R., Wagner, W., and Dorigo, W.: Evolution of the ESA CCI Soil Moisture climate data records and their underlying merging methodology, *Earth System Science Data*, 11, 717–739, <https://doi.org/10.5194/essd-11-717-2019>, 2019b.



- 1365 Gruber, A., De Lannoy, G., Albergel, C., Al-Yaari, A., Brocca, L., Calvet, J.-C., Colliander, A., Cosh, M., Crow, W., Dorigo, W., Draper, C., Hirschi, M., Kerr, Y., Konings, A., Lahoz, W., McColl, K., Montzka, C., Muñoz-Sabater, J., Peng, J., Reichle, R., Richaume, P., Rüdiger, C., Scanlon, T., van der Schalie, R., Wigneron, J.-P., and Wagner, W.: Validation practices for satellite soil moisture retrievals: What are (the) errors?, *Remote Sensing of Environment*, 244, 111 806, <https://doi.org/10.1016/j.rse.2020.111806>, 2020.
- Guevara, M. and Vargas, R.: Downscaling satellite soil moisture using geomorphometry and machine learning, *PloS one*, 14, e0219 639, 1370 2019.
- Gumbricht, T.: Detecting Trends in Wetland Extent from MODIS Derived Soil Moisture Estimates, *Remote Sensing*, 10, 611, <https://doi.org/10.3390/rs10040611>, 2018.
- Gumbricht, T., Roman-Cuesta, R. M., Verchot, L., Herold, M., Wittmann, F., Householder, E., Herold, N., and Murdiyarso, D.: An expert system model for mapping tropical wetlands and peatlands reveals South America as the largest contributor, *Global Change Biology*, 23, 3581–3599, <https://doi.org/10.1111/gcb.13689>, 2017.
- 1375 Hagan, D. F. T., Wang, G., Kim, S., Parinussa, R. M., Liu, Y., Ullah, W., Bhatti, A. S., Ma, X., Jiang, T., and Su, B.: Maximizing Temporal Correlations in Long-Term Global Satellite Soil Moisture Data-Merging, *Remote Sensing*, 12, 2164, <https://doi.org/10.3390/rs12132164>, 2020.
- Hajdu, I., Yule, I., Bretherton, M., Singh, R., and Hedley, C.: Field performance assessment and calibration of multi-depth AquaCheck capacitance-based soil moisture probes under permanent pasture for hill country soils, *Agricultural Water Management*, 217, 332–345, <https://doi.org/10.1016/j.agwat.2019.03.002>, 2019.
- 1380 Han, M., Lu, H., and Yang, K.: Development of passive microwave retrieval algorithm for estimation of surface soil temperature from AMSR-E data, in: 2016 IEEE International Geoscience and Remote Sensing Symposium (IGARSS), pp. 1671–1674, 2016.
- Han, M., Lu, H., Yang, K., Qin, J., Chen, Y., Zhao, L., and Lazhu: A surface soil temperature retrieval algorithm based on AMSR-E multi-frequency brightness temperatures, *International Journal of Remote Sensing*, 38, 6735–6754, 1385 <https://doi.org/10.1080/01431161.2017.1363438>, 2017.
- Han, Y., Bai, X., Shao, W., and Wang, J.: Retrieval of Soil Moisture by Integrating Sentinel-1A and MODIS Data over Agricultural Fields, *Water*, 12, <https://doi.org/10.3390/w12061726>, 2020.
- Hartmann, A., Gleeson, T., Rosolem, R., Pianosi, F., Wada, Y., and Wagener, T.: A large-scale simulation model to assess karstic groundwater recharge over Europe and the Mediterranean, *Geoscientific Model Development*, 8, 1729–1746, <https://doi.org/10.5194/gmd-8-1729-2015>, 1390 2015.
- Helgert, S. and Khodayar, S.: Improvement of the soil-atmosphere interactions and subsequent heavy precipitation modelling by enhanced initialization using remotely sensed 1 km soil moisture information, *Remote Sensing of Environment*, 246, 111 812, <https://doi.org/10.1016/j.rse.2020.111812>, 2020.
- 1395 Herbert, C., Pablos, M., Vall-llossera, M., Camps, A., and Martínez-Fernández, J.: Analyzing Spatio-Temporal Factors to Estimate the Response Time between SMOS and In-Situ Soil Moisture at Different Depths, *Remote Sensing*, 12, 2614, <https://doi.org/10.3390/rs12162614>, 2020.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 Global Reanalysis, *Quarterly Journal of the Royal Meteorological Society*, n/a, <https://doi.org/10.1002/qj.3803>, 2020.
- 1400



- Hirschi, M., Mueller, B., Dorigo, W., and Seneviratne, S.: Using remotely sensed soil moisture for land–atmosphere coupling diagnostics: The role of surface vs. root-zone soil moisture variability, *Remote Sensing of Environment*, 154, 246–252, <https://doi.org/doi.org/10.1016/j.rse.2014.08.030>, 2014.
- 1405 Hoang, K. O. and Lu, M.: Impacts of temperature effect removal on rainfall estimation from soil water content by using SM2RAIN algorithm, *IOP Conference Series: Earth and Environmental Science*, 344, 012 046, <https://doi.org/10.1088/1755-1315/344/1/012046>, 2019.
- Hollinger, S. and Isard, S.: A Soil Moisture Climatology of Illinois, *Journal of Climate*, 7, 822–833, [https://doi.org/10.1175/1520-0442\(1994\)007<0822:ASMCOT>2.0.CO;2](https://doi.org/10.1175/1520-0442(1994)007<0822:ASMCOT>2.0.CO;2), 1994.
- 1410 Hottenstein, J. D., Ponce-Campos, G. E., Moguel-Yanes, J., and Moran, M. S.: Impact of Varying Storm Intensity and Consecutive Dry Days on Grassland Soil Moisture, *Journal of Hydrometeorology*, 16, 106–117, <https://doi.org/10.1175/JHM-D-14-0057.1>, 2015.
- Hu, T., Zhao, T., Zhao, K., and Shi, J.: A continuous global record of near-surface soil freeze/thaw status from AMSR-E and AMSR2 data, *International Journal of Remote Sensing*, 40, 6993–7016, <https://doi.org/10.1080/01431161.2019.1597307>, 2019.
- Ikonen, J., Vehviläinen, J., Rautiainen, K., Smolander, T., Lemmetyinen, J., Bircher, S., and Pulliainen, J.: The Sodankylä in-situ soil moisture observation network: an example application to Earth Observation data product evaluation, *GID*, 5, 599–629, 2015.
- 1415 Ikonen, J., Smolander, T., Rautiainen, K., Cohen, J., Lemmetyinen, J., Salminen, M., and Pulliainen, J.: Spatially distributed evaluation of ESA CCI Soil Moisture products in a northern boreal forest environment, *Geosciences*, 8, 51, 2018.
- ISMN: ISMN stated terms and conditions for data use, <https://ismn.geo.tuwien.ac.at/en/terms-and-conditions/>, last accessed 20 August 2020.
- J., L.: 2011 Installation Report for AAFC- SAGES Soil Moisture Stations in Kenaston, SK. *Agriculture*, 2011.
- 1420 Jackson, T., Cosh, M., Bindlish, R., Starks, P., Bosch, D., Seyfried, M., Goodrich, D., Moran, M., and Du, J.: Validation of Advanced Microwave Scanning Radiometer Soil Moisture Products, *Geoscience and Remote Sensing, IEEE Transactions on*, 48, 4256–4272, <https://doi.org/10.1109/TGRS.2010.2051035>, 2011.
- Jalilvand, E., Tajrishy, M., Brocca, L., Massari, C., Ghazi Zadeh Hashemi, S., and Ciabatta, L.: Estimating the drainage rate from surface soil moisture drydowns: Application of DfD model to in situ soil moisture data, *Journal of Hydrology*, 565, 489–501, <https://doi.org/10.1016/j.jhydrol.2018.08.035>, 2018.
- 1425 Jensen, K. and Refsgaard, J.: HOBE: The Danish Hydrological Observatory, *Vadose Zone Journal*, 17, <https://doi.org/10.2136/vzj2018.03.0059>, 2018.
- Ji, P., Yuan, X., and Liang, X.-Z.: Do Lateral Flows Matter for the Hyperresolution Land Surface Modeling?, *Journal of Geophysical Research: Atmospheres*, 122, 12,077–12,092, <https://doi.org/10.1002/2017JD027366>, 2017.
- 1430 Jin, R., Li, X., Yan, B., Li, X., Luo, W., Ma, M., Guo, J., Kang, J., Zhu, Z., and Zhao, S.: A Nested Ecohydrological Wireless Sensor Network for Capturing the Surface Heterogeneity in the Midstream Areas of the Heihe River Basin, China, *IEEE Geoscience and Remote Sensing Letters*, 11, 2015–2019, <https://doi.org/10.1109/LGRS.2014.2319085>, 2014.
- Kang, C. S., Kanniah, K. D., and Kerr, Y. H.: Calibration of SMOS Soil Moisture Retrieval Algorithm: A Case of Tropical Site in Malaysia, *IEEE Transactions on Geoscience and Remote Sensing*, 57, 3827–3839, 2019.
- 1435 Kang, J., Li, X., Jin, R., Ge, Y., Wang, J., and Wang, J.: Hybrid Optimal Design of the Eco-Hydrological Wireless Sensor Network in the Middle Reach of the Heihe River Basin, China, *Sensors*, <https://doi.org/10.3390/s141019095>, 2014.
- Kapilaratne, R. J. and Lu, M.: Automated general temperature correction method for dielectric soil moisture sensors, *Journal of Hydrology*, 551, 203–216, <https://doi.org/10.1016/j.jhydrol.2017.05.050>, investigation of Coastal Aquifers, 2017.





- Karthikeyan, L., Pan, M., Wanders, N., Kumar, D. N., and Wood, E. F.: Four decades of microwave satellite soil moisture observations: Part 2. Product validation and inter-satellite comparisons, *Advances in Water Resources*, 109, 236–252, <https://doi.org/10.1016/j.advwatres.2017.09.010>, 2017.
- Kędzior, M. and Zawadzki, J.: Comparative study of soil moisture estimations from SMOS satellite mission, GLDAS database, and cosmic-ray neutrons measurements at COSMOS station in Eastern Poland, *Geoderma*, 283, 21–31, <https://doi.org/10.1016/j.geoderma.2016.07.023>, 2016.
- 1445 Kerr, Y., Al-Yaari, A., Rodriguez-Fernandez, N., Parrens, M., Molero, B., Leroux, D., Bircher, S., Mahmoodi, A., Mialon, A., Richaume, P., Delwart, S., Al Bitar, A., Pellarin, T., Bindlish, R., Jackson, T., Rüdiger, C., Waldteufel, P., Mecklenburg, S., and Wigneron, J.-P.: Overview of SMOS performance in terms of global soil moisture monitoring after six years in operation, *Remote Sensing of Environment*, 180, 40–63, <https://doi.org/10.1016/j.rse.2016.02.042>, special Issue: ESA's Soil Moisture and Ocean Salinity Mission - Achievements and Applications, 2016.
- 1450 Khaki, M., Hoteit, I., Kuhn, M., Forootan, E., and Awange, J.: Assessing data assimilation frameworks for using multi-mission satellite products in a hydrological context, *Science of The Total Environment*, 647, 1031–1043, <https://doi.org/10.1016/j.scitotenv.2018.08.032>, 2019.
- Kim, H. and Lakshmi, V.: Use of Cyclone Global Navigation Satellite System (CyGNSS) Observations for Estimation of Soil Moisture, *Geophysical Research Letters*, 45, 8272–8282, <https://doi.org/10.1029/2018GL078923>, 2018.
- 1455 Kim, H., Parinussa, R., Konings, A. G., Wagner, W., Cosh, M. H., Lakshmi, V., Zohaib, M., and Choi, M.: Global-scale assessment and combination of SMAP with ASCAT (active) and AMSR2 (passive) soil moisture products, *Remote Sensing of Environment*, 204, 260–275, <https://doi.org/10.1016/j.rse.2017.10.026>, 2018.
- Kim, K., Jung, S., and Lee, Y.: A Study for establishment of soil moisture station in mountain terrain (1): the representative analysis of soil moisture for construction of Cosmic-ray verification system, *Journal of Korea Water Resources Association*, 52, 51–60, 2019.
- 1460 Kim, S., Liu, Y., Johnson, F. M., Parinussa, R. M., and Sharma, A.: A global comparison of alternate AMSR2 soil moisture products: Why do they differ?, *Remote Sensing of Environment*, 161, 43–62, <https://doi.org/10.1016/j.rse.2015.02.002>, 2015a.
- Kim, S., Parinussa, R. M., Liu, Y. Y., Johnson, F. M., and Sharma, A.: A framework for combining multiple soil moisture retrievals based on maximizing temporal correlation, *Geophysical Research Letters*, 42, 6662–6670, <https://doi.org/10.1002/2015GL064981>, 2015b.
- Kim, S., Parinussa, R., Liu, Y., Johnson, F., and Sharma, A.: Merging Alternate Remotely-Sensed Soil Moisture Retrievals Using a Non-Static Model Combination Approach, *Remote Sensing*, 8, 518, <https://doi.org/10.3390/rs8060518>, 2016.
- 1465 Kirchengast, G., Kabas, T., Leuprecht, A., Bichler, C., and Truhetz, H.: WegenerNet: A Pioneering High-Resolution Network for Monitoring Weather and Climate, *Bulletin of the American Meteorological Society*, 95, <https://doi.org/10.1175/BAMS-D-11-00161.1>, 2014.
- Kizito, F., Campbell, C., Campbell, G., Cobos, D., Teare, B., Carter, B., and Hopmans, J.: Frequency, electrical conductivity and temperature analysis of a low-cost capacitance soil moisture sensor, *Journal of Hydrology*, 352, 367–378, <https://doi.org/10.1016/j.jhydrol.2008.01.021>, 2008.
- 1470 Kojima, Y., Shigeta, R., Miyamoto, N., Shirahama, Y., Nishioka, K., Mizoguchi, M., and Kawahara, Y.: Low-Cost Soil Moisture Profile Probe Using Thin-Film Capacitors and a Capacitive Touch Sensor, *Sensors*, 16, 1292, <https://doi.org/10.3390/s16081292>, 2016.
- Kolassa, J., Aires, F., Polcher, J., Prigent, C., Jimenez, C., and Pereira, J. M.: Soil moisture retrieval from multi-instrument observations: Information content analysis and retrieval methodology, *Journal of Geophysical Research: Atmospheres*, 118, 4847–4859, <https://doi.org/10.1029/2012JD018150>, 2013.





- Kolassa, J., Gentine, P., Prigent, C., Aires, F., and Alemohammad, S.: Soil moisture retrieval from AMSR-E and ASCAT microwave observation synergy. Part 2: Product evaluation, *Remote Sensing of Environment*, 195, 202–217, <https://doi.org/10.1016/j.rse.2017.04.020>, 2017.
- Kolassa, J., Reichle, R., Liu, Q., Alemohammad, S., Gentine, P., Aida, K., Asanuma, J., Bircher, S., Caldwell, T., Colliander, A., et al.:  
 1480 Estimating surface soil moisture from SMAP observations using a Neural Network technique, *Remote sensing of environment*, 204, 43–59, <https://doi.org/10.1016/j.rse.2017.10.045>, 2018.
- Kornelsen, K. C. and Coulibaly, P.: Root-zone soil moisture estimation using data-driven methods, *Water Resources Research*, 50, 2946–2962, <https://doi.org/10.1002/2013WR014127>, 2014.
- Kornelsen, K. C. and Coulibaly, P.: Reducing multiplicative bias of satellite soil moisture retrievals, *Remote Sensing of Environment*, 165,  
 1485 109–122, <https://doi.org/10.1016/j.rse.2015.04.031>, 2015.
- Kosmala, M., Wiggins, A., Swanson, A., and Simmons, B.: Assessing data quality in citizen science, *Frontiers in Ecology and the Environment*, 14, 551–560, <https://doi.org/10.1002/fee.1436>, 2016.
- Kovačević, J., Cvijetinović, Z., Stančić, N., Brodić, N., and Mihajlović, D.: New Downscaling Approach Using ESA CCI SM Products for Obtaining High Resolution Surface Soil Moisture, *Remote Sensing*, 12, 1119, <https://doi.org/10.3390/rs12071119>, 2020.
- Kovács, K. Z., Hemment, D., Woods, M., van der Velden, N. K., Xaver, A., Giesen, R. H., Burton, V. J., Garrett, N. L., Zappa, L., Long, D.,  
 1490 Dobos, E., and Skalsky, R.: Citizen observatory based soil moisture monitoring – the GROW example, *Hungarian Geographical Bulletin*, 68, 119–139, <https://doi.org/10.15201/hungeobull.68.2.2>, 2019.
- Krueger, E. S., Ochsner, T. E., and Quiring, S. M.: Development and Evaluation of Soil Moisture-Based Indices for Agricultural Drought Monitoring, *Agronomy Journal*, 111, 1392–1406, <https://doi.org/10.2134/agronj2018.09.0558>, 2019.
- 1495 Kumar, M. S., Chandra, T. R., Kumar, D. P., and Manikandan, M. S.: Monitoring moisture of soil using low cost homemade Soil moisture sensor and Arduino UNO, pp. 1–4, 2016 3rd International Conference on Advanced Computing and Communication Systems (ICACCS), IEEE, Coimbatore, India, <https://doi.org/10.1109/ICACCS.2016.7586312>, [Online; accessed 2020-06-18], 2016.
- Kumar, S., Newman, M., Wang, Y., and Livneh, B.: Potential Reemergence of Seasonal Soil Moisture Anomalies in North America, *Journal of Climate*, 32, 2707–2734, <https://doi.org/10.1175/JCLI-D-18-0540.1>, 2019a.
- 1500 Kumar, S. V., Peters-Lidard, C. D., Santanello, J., Harrison, K., Liu, Y., and Shaw, M.: Land surface Verification Toolkit (LVT) – a generalized framework for land surface model evaluation, *Geoscientific Model Development*, 5, 869–886, <https://doi.org/10.5194/gmd-5-869-2012>, 2012.
- Kumar, S. V., Mocko, D., Wang, S., Peters-Lidard, C. D., and Borak, J.: Assimilation of Remotely Sensed Leaf Area Index into the Noah-MP Land Surface Model: Impacts on Water and Carbon Fluxes and States over the Continental United States, *Journal of Hydrometeorology*, 20, 1359–1377, <https://doi.org/10.1175/JHM-D-18-0237.1>, 2019b.
- 1505 Larson, K., Small, E., Gutmann, E., Bilich, A., Braun, J., Zavorotny, V., and Larson, C.: Use of GPS receivers as a soil moisture network for water cycle studies, *Geophysical Research Letters - GEOPHYS RES LETT*, 35, <https://doi.org/10.1029/2008GL036013>, 2008.
- Leavesley: A Modelling Framework for Improved Agricultural Water-Supply Forecasting, 2010.
- Leavesley, G., David, O., Garen, D., Lea, J., Marron, J., Pagano, T., Perkins, T., and Strobel, M.: A Modeling Framework for Improved  
 1510 Agricultural Water Supply Forecasting, AGU Fall Meeting Abstracts, 2008.
- Lee, J.: The consecutive dry days to trigger rainfall over West Africa, *Journal of Hydrology*, 556, 934–943, <https://doi.org/10.1016/j.jhydrol.2016.06.003>, 2018.



- Lee, J. and Im, J.: A Novel Bias Correction Method for Soil Moisture and Ocean Salinity (SMOS) Soil Moisture: Retrieval Ensembles, *Remote Sensing*, 7, 16 045–16 061, <https://doi.org/10.3390/rs71215824>, 2015.
- 1515 Lei, F., Crow, W. T., Shen, H., Su, C.-H., Holmes, T. R., Parinussa, R. M., and Wang, G.: Assessment of the impact of spatial heterogeneity on microwave satellite soil moisture periodic error, *Remote Sensing of Environment*, 205, 85–99, <https://doi.org/10.1016/j.rse.2017.11.002>, 2018.
- Leng, P., Song, X., Li, Z.-L., Wang, Y., and Wang, R.: Toward the Estimation of Surface Soil Moisture Content Using Geostationary Satellite Data over Sparsely Vegetated Area, *Remote Sensing*, 7, 4112–4138, <https://doi.org/10.3390/rs70404112>, 2015.
- 1520 Leng, P., Song, X., Duan, S.-B., and Li, Z.-L.: Preliminary validation of two temporal parameter-based soil moisture retrieval models using a satellite product and in situ soil moisture measurements over the REMEDHUS network, *International Journal of Remote Sensing*, 37, 5902–5917, <https://doi.org/10.1080/01431161.2016.1253896>, 2016.
- Leng, P., Song, X., Duan, S.-B., and Li, Z.-L.: Generation of continuous surface soil moisture dataset using combined optical and thermal infrared images, *Hydrological Processes*, 31, 1398–1407, <https://doi.org/10.1002/hyp.11113>, 2017.
- 1525 Li, M., Wu, P., and Ma, Z.: A comprehensive evaluation of soil moisture and soil temperature from third-generation atmospheric and land reanalysis data sets, *International Journal of Climatology*, n/a, <https://doi.org/10.1002/joc.6549>, 2020a.
- Li, P., Zha, Y., Tso, C.-H. M., Shi, L., Yu, D., Zhang, Y., and Zeng, W.: Data assimilation of uncalibrated soil moisture measurements from frequency-domain reflectometry, *Geoderma*, 374, 114 432, <https://doi.org/10.1016/j.geoderma.2020.114432>, 2020b.
- Li, X., Al-Yaari, A., Schwank, M., Fan, L., Frappart, F., Swenson, J., and Wigneron, J.-P.: Compared performances of SMOS-IC soil moisture and vegetation optical depth retrievals based on Tau-Omega and Two-Stream microwave emission models, *Remote Sensing of Environ-*  
1530 *ment*, 236, 111 502, <https://doi.org/10.1016/j.rse.2019.111502>, 2020c.
- Liao, W., Wang, D., Wang, G., Xia, Y., and Liu, X.: Quality Control and Evaluation of the Observed Daily Data in the North American Soil Moisture Database, *Journal of Meteorological Research*, 33, 501–518, <https://doi.org/10.1007/s13351-019-8121-2>, 2019.
- Lievens, H., Martens, B., Verhoest, N., Hahn, S., Reichle, R., and Miralles, D.: Assimilation of global radar backscatter and radiometer  
1535 brightness temperature observations to improve soil moisture and land evaporation estimates, *Remote Sensing of Environment*, 189, 194–210, <https://doi.org/10.1016/j.rse.2016.11.022>, 2017.
- Lin, L.-F. and Pu, Z.: Examining the Impact of SMAP Soil Moisture Retrievals on Short-Range Weather Prediction under Weakly and Strongly Coupled Data Assimilation with WRF-Noah, *Monthly Weather Review*, 147, 4345–4366, <https://doi.org/10.1175/MWR-D-19-0017.1>, 2019.
- 1540 Lin, L.-F. and Pu, Z.: Improving Near-Surface Short-Range Weather Forecasts Using Strongly Coupled Land–Atmosphere Data Assimilation with GSI-EnKF, *Monthly Weather Review*, 148, 2863–2888, <https://doi.org/10.1175/MWR-D-19-0370.1>, 2020.
- Link, M., Drusch, M., and Scipal, K.: Soil Moisture Information Content in SMOS, SMAP, AMSR2, and ASCAT Level-1 Data Over Selected In Situ Sites, *IEEE Geoscience and Remote Sensing Letters*, 17, 1213–1217, 2020.
- Liu, G., Chen, J.-M., Ji, L.-R., and Sun, S.-Q.: Relationship of summer soil moisture with early winter monsoon and air temperature over  
1545 eastern China, *International Journal of Climatology*, 32, 1513–1519, <https://doi.org/10.1002/joc.2352>, 2012a.
- Liu, S., Mo, X., Li, H., Peng, G., and Robock, A.: Spatial Variation of Soil Moisture in China: Geostatistical Characterization, *Journal of The Meteorological Society of Japan - J METEOROL SOC JPN*, 79, 555–574, <https://doi.org/10.2151/jmsj.79.555>, 2001.
- Liu, S., Roujean, J.-L., Kaptue Tchuenté, A. T., Ceamanos, X., and Calvet, J.-C.: A parameterization of SEVIRI and MODIS daily surface albedo with soil moisture: Calibration and validation over southwestern France, *Remote Sensing of Environment*, 144, 137–151,  
1550 <https://doi.org/10.1016/j.rse.2014.01.016>, 2014.



- Liu, Y., Dorigo, W., Parinussa, R., de Jeu, R., Wagner, W., McCabe, M., Evans, J., and van Dijk, A.: Trend-preserving blending of passive and active microwave soil moisture retrievals, *Remote Sensing of Environment*, 123, 280–297, <https://doi.org/10.1016/j.rse.2012.03.014>, 2012b.
- Liu, Y., Yang, Y., and Yue, X.: Evaluation of Satellite-Based Soil Moisture Products over Four Different Continental In-Situ Measurements, *Remote Sensing*, 10, 1161, <https://doi.org/10.3390/rs10071161>, 2018.
- Liu, Y., Jing, W., Wang, Q., and Xia, X.: Generating high-resolution daily soil moisture by using spatial downscaling techniques: a comparison of six machine learning algorithms, *Advances in Water Resources*, 141, 103 601, <https://doi.org/10.1016/j.advwatres.2020.103601>, 2020.
- Liu, Y. Y., Parinussa, R. M., Dorigo, W. A., De Jeu, R. A. M., Wagner, W., van Dijk, A. I. J. M., McCabe, M. F., and Evans, J. P.: Developing an improved soil moisture dataset by blending passive and active microwave satellite-based retrievals, *Hydrology and Earth System Sciences*, 15, 425–436, <https://doi.org/10.5194/hess-15-425-2011>, 2011.
- Liu, Z., Li, P., and Yang, J.: Soil Moisture Retrieval and Spatiotemporal Pattern Analysis Using Sentinel-1 Data of Dahra, Senegal, *Remote Sensing*, 9, <https://doi.org/10.3390/rs9111197>, 2017.
- Loew, A., dall’Amico, J., Schlenz, F., and Mauser, W.: The Upper Danube soil moisture validation site: measurements and activities, 2009.
- Loew, A., Bell, W., Brocca, L., Bulgin, C. E., Burdanowitz, J., Calbet, X., Donner, R. V., Ghent, D., Gruber, A., Kaminski, T., Kinzel, J., Klepp, C., Lambert, J.-C., Schaepman-Strub, G., Schröder, M., and Verhoelst, T.: Validation practices for satellite-based Earth observation data across communities, *Reviews of Geophysics*, 55, 779–817, <https://doi.org/10.1002/2017RG000562>, 2017.
- Lu, Y., Jin, J., and Kueppers, L.: Crop growth and irrigation interact to influence surface fluxes in a regional climate-cropland model (WRF3.3-CLM4crop), *Climate Dynamics*, 45, <https://doi.org/10.1007/s00382-015-2543-z>, 2015.
- Lukyanenko, R., Wiggins, A., and Rosser, H. K.: Citizen Science: An Information Quality Research Frontier, *Information Systems Frontiers*, 22, 961–983, <https://doi.org/10.1007/s10796-019-09915-z>, 2020.
- Ma, C., Li, X., and McCabe, M. F.: Retrieval of High-Resolution Soil Moisture through Combination of Sentinel-1 and Sentinel-2 Data, *Remote Sensing*, 12, 2303, <https://doi.org/10.3390/rs12142303>, 2020.
- Ma, H., Zeng, J., Chen, N., Zhang, X., Cosh, M. H., and Wang, W.: Satellite surface soil moisture from SMAP, SMOS, AMSR2 and ESA CCI: A comprehensive assessment using global ground-based observations, *Remote Sensing of Environment*, 231, 111 215, 2019.
- Mahecha, M., Gans, F., Sippel, S., Donges, J., Kaminski, T., Metzger, S., Migliavacca, M., Papale, D., Rammig, A., and Zscheischler, J.: Detecting impacts of extreme events with ecological in-situ monitoring networks, *Biogeosciences*, 14, 4255–4277, <https://doi.org/10.5194/bg-14-4255-2017>, 2017.
- Majone, B., Viani, F., Filippi, E., Bellin, A., Massa, A., Toller, G., Robol, F., and Salucci, M.: Wireless Sensor Network Deployment for Monitoring Soil Moisture Dynamics at the Field Scale, *Procedia Environmental Sciences*, 19, 426–435, <https://doi.org/10.1016/j.proenv.2013.06.049>, 2013.
- Manfreda, S., Brocca, L., Moramarco, T., Melone, F., and Sheffield, J.: A physically based approach for the estimation of root-zone soil moisture from surface measurements, *Hydrology and Earth System Sciences*, 18, 1199–1212, <https://doi.org/10.5194/hess-18-1199-2014>, 2014.
- Mao, F., Khamis, K., Krause, S., Clark, J., and Hannah, D. M.: Low-Cost Environmental Sensor Networks: Recent Advances and Future Directions, *Frontiers in Earth Science*, 7, 221, <https://doi.org/10.3389/feart.2019.00221>, 2019.
- Marczewski, W., Slominski, J., Slominska, E., Usowicz, B., Usowicz, J., S, R., O, M., Nastula, J., and Zawadzki, J.: Strategies for validating and directions for employing SMOS data, in the Cal-Val project SWEX (3275) for wetlands, *Hydrology and Earth System Sciences Discussions*, 7, <https://doi.org/10.5194/hessd-7-7007-2010>, 2010.



- Martens, B., Miralles, D., Lievens, H., van der Schalie, R., de Jeu, R. A., Fernández-Prieto, D., Beck, H. E., Dorigo, W., and Verhoest, N.: GLEAM v3 : satellite-based land evaporation and root-zone soil moisture, *Geoscientific Model Development*, 10, 1903–1925, <https://doi.org/10.5194/gmd-10-1903-2017>, 2017.
- Martens, B., de Jeu, R., Verhoest, N., Schuurmans, H., Kleijer, J., and Miralles, D.: Towards Estimating Land Evaporation at Field Scales Using GLEAM, *Remote Sensing*, 10, 1720, <https://doi.org/10.3390/rs10111720>, 2018.
- Massari, C., Su, C.-H., Brocca, L., Sang, Y.-F., Ciabatta, L., Ryu, D., and Wagner, W.: Near real time de-noising of satellite-based soil moisture retrievals: An intercomparison among three different techniques, *Remote Sensing of Environment*, 198, 17–29, <https://doi.org/10.1016/j.rse.2017.05.037>, 2017.
- Mattar, C., Santamaría-Artigas, A., Durán-Alarcón, C., Olivera-Guerra, L., Fuster, R., and Borvarán, D.: The LAB-Net Soil Moisture Network: Application to Thermal Remote Sensing and Surface Energy Balance, *Data*, 1, <https://doi.org/10.3390/data1010006>, 2016.
- McDonough, K. R., Hutchinson, S. L., Hutchinson, J. S., Case, J. L., and Rahmani, V.: Validation and assessment of SPoRT-LIS surface soil moisture estimates for water resources management applications, *Journal of Hydrology*, 566, 43–54, <https://doi.org/10.1016/j.jhydrol.2018.09.007>, 2018.
- Mecklenburg, S., Drusch, M., Kerr, Y. H., Font, J., Martin-Neira, M., Delwart, S., Buenadicha, G., Reul, N., Daganzo-Eusebio, E., Oliva, R., and Crapolicchio, R.: ESA's Soil Moisture and Ocean Salinity Mission: Mission Performance and Operations, *IEEE Transactions on Geoscience and Remote Sensing*, 50, 1354–1366, <https://doi.org/10.1109/TGRS.2012.2187666>, 2012.
- Mimeau, L., Trambly, Y., Brocca, L., Massari, C., Camici, S., and Finaud-Guyot, P.: Modeling the response of soil moisture to climate variability in the Mediterranean region, *Hydrology and Earth System Sciences Discussions*, 2020, 1–29, <https://doi.org/10.5194/hess-2020-302>, 2020.
- Miralles, D. G., Crow, W. T., and Cosh, M. H.: Estimating Spatial Sampling Errors in Coarse-Scale Soil Moisture Estimates Derived from Point-Scale Observations, *Journal of Hydrometeorology*, 11, 1423–1429, <https://doi.org/10.1175/2010JHM1285.1>, 2010.
- Mishra, V., Shah, R., and Thrasher, B.: Soil Moisture Droughts under the Retrospective and Projected Climate in India, *Journal of Hydrometeorology*, 15, 2267–2292, <https://doi.org/10.1175/JHM-D-13-0177.1>, 2014.
- Mishra, V., Shah, R., Azhar, S., Shah, H., Modi, P., and Kumar, R.: Reconstruction of droughts in India using multiple land-surface models (1951–2015), *Hydrology and Earth System Sciences*, 22, 2269–2284, <https://doi.org/10.5194/hess-22-2269-2018>, 2018.
- Mittelbach, H., Lehner, I., and Seneviratne, S. I.: Comparison of four soil moisture sensor types under field conditions in Switzerland, *Journal of Hydrology*, 430–431, 39–49, <https://doi.org/10.1016/j.jhydrol.2012.01.041>, 2012.
- Moghaddam, M., Entekhabi, D., Goykhman, Y., Li, K., Liu, M., Mahajan, A., Nayyar, A., Shuman, D., and Teneketzis, D.: A Wireless Soil Moisture Smart Sensor Web Using Physics-Based Optimal Control: Concept and Initial Demonstrations, *Selected Topics in Applied Earth Observations and Remote Sensing*, *IEEE Journal of*, 3, 522–535, <https://doi.org/10.1109/JSTARS.2010.2052918>, 2011.
- MOGHADDAM, M., SILVA, A., CLEWLEY, D., AKBAR, R., HUSSAINI, S., Whitcomb, J., DEVARAKONDA, R., Shrestha, R., COOK, R., PRAKASH, G., SANTHANA VANNAN, S., and BOYER, A.: Soil Moisture Profiles and Temperature Data from SoilSCAPE Sites, USA, <https://doi.org/10.3334/ORNLDAAAC/1339>, 2016.
- Mohanty, B. P., Cosh, M. H., Lakshmi, V., and Montzka, C.: Soil Moisture Remote Sensing: State-of-the-Science, *Vadose Zone Journal*, 16, vzj2016.10.0105, <https://doi.org/10.2136/vzj2016.10.0105>, 2017.
- Molero, B., Leroux, D. J., Richaume, P., Kerr, Y. H., Merlin, O., Cosh, M. H., and Bindlish, R.: Multi-Timescale Analysis of the Spatial Representativeness of In Situ Soil Moisture Data within Satellite Footprints: Soil Moisture Time and Spatial Scales, *Journal of Geophysical Research: Atmospheres*, 123, 3–21, <https://doi.org/10.1002/2017JD027478>, 2018.



- Montzka, C., Bogaen, H., Zreda, M., Monerris, A., Morrison, R., Muddu, S., and Vereecken, H.: Validation of Spaceborne and Modelled Surface Soil Moisture Products with Cosmic-Ray Neutron Probes, *Remote Sensing*, 9, 103, <https://doi.org/10.3390/rs9020103>, 2017.
- 1630 Montzka, C., Cosh, M., Bayat, B., Bitar, A. A., Berg, A., Bindlish, R., Bogaen, H. R., Bolton, J. D., Cabot, F., Caldwell, T., Chan, S., Colliander, A., Crow, W., Das, N., Lannoy, G. D., Dorigo, W., Evett, S. R., Gruber, A., Hahn, S., Jagdhuber, T., Jones, S., Kerr, Y., Kim, S., Koyama, C., Kurum, M., Lopez-Baeza, E., Mattia, F., McColl, K., Mecklenburg, S., Mohanty, B., O'Neill, P., Or, D., Pellarin, T., Petropoulos, G. P., Piles, M., Reichle, R. H., Rodriguez-Fernandez, N., Rüdiger, C., Scanlon, T., Schwartz, R. C., Spengler, D., Srivastava, P., Suman, S., van der Schalie, R., Wagner, W., Wegmüller, U., Camacho, F., and Nickeson, J.: Soil Moisture Product Validation Best Practice Protocol, Version 1.0, 2020.
- 1635 Morbidelli, R., Saltalippi, C., Flammini, A., Rossi, E., and Corradini, C.: Soil water content vertical profiles under natural conditions: matching of experiments and simulations by a conceptual model, *Hydrological Processes*, 28, 4732–4742, <https://doi.org/10.1002/hyp.9973>, 2014.
- Morbidelli, R., Saltalippi, C., Flammini, A., Cifrodelli, M., Picciafuoco, T., Corradini, C., and Govindaraju, R. S.: In situ measurements of soil saturated hydraulic conductivity: Assessment of reliability through rainfall–runoff experiments, *Hydrological Processes*, 31, 3084–3094, <https://doi.org/10.1002/hyp.11247>, 2017.
- 1640 Moreno-Martínez, Á., Piles, M., Muñoz-Marí, J., Campos-Taberner, M., Adsua, J. E., Mateo, A., Perez-Suay, A., Javier García-Haro, F., and Camps-Valls, G.: Machine Learning Methods for Spatial and Temporal Parameter Estimation, pp. 5–35, Springer International Publishing, Cham, [https://doi.org/10.1007/978-3-030-38617-7\\_2](https://doi.org/10.1007/978-3-030-38617-7_2), 2020.
- Mu, L., Li, S., Qin, B., and Liu, K.: Drought Monitoring Using Tiangong-2 Wide-Band Spectrometer Data, in: Proceedings of the Tiangong-2 Remote Sensing Application Conference, edited by Gu, Y., Gao, M., and Zhao, G., pp. 277–285, Springer Singapore, Singapore, [https://doi.org/10.1007/978-981-13-3501-3\\_26](https://doi.org/10.1007/978-981-13-3501-3_26), 2019.
- 1645 Muñoz Sabater, J., Rodriguez-Fernandez, N., Sabater, J. M., Richaume, P., De Rosnay, P., Kerr, Y., Albergel, C., Drusch, M., and Mecklenburg, S.: SMOS near-real-time soil moisture product: processor overview and first validation results, *Hydrology and Earth System Sciences*, 21, 5201–5216, <https://doi.org/10.5194/hess-21-5201-2017>, 2017.
- 1650 Musial, J. P., Dabrowska-Zielinska, K., Kiryla, W., Oleszczuk, R., Gnatowski, T., and Jaszczynski, J.: Derivation and validation of the high resolution satellite soil moisture products: a case study of the Biebrza Sentinel-1 validation sites, *Geoinformation Issues*, 8, 37–53, 2016.
- Nair, A. and Indu, J.: Enhancing Noah Land Surface Model Prediction Skill over Indian Subcontinent by Assimilating SMOPS Blended Soil Moisture, *Remote Sensing*, 8, 976, <https://doi.org/10.3390/rs8120976>, 2016.
- Nair, A. S., Mangla, R., P, T., and Indu, J.: Remote sensing data assimilation, *Hydrological Sciences Journal*, 0, 1–33, <https://doi.org/10.1080/02626667.2020.1761021>, 2020a.
- 1655 Nair, A. S., Mangla, R., P, T., and Indu, J.: Remote sensing data assimilation, *Hydrological Sciences Journal*, 0, 1–33, <https://doi.org/10.1080/02626667.2020.1761021>, 2020b.
- Naz, B. S., Kollet, S., Franssen, H.-J. H., Montzka, C., and Kurtz, W.: A 3 km spatially and temporally consistent European daily soil moisture reanalysis from 2000 to 2015, *Scientific Data*, 7, 111, <https://doi.org/10.1038/s41597-020-0450-6>, 2020.
- 1660 Nguyen, H. H., Kim, H., and Choi, M.: Evaluation of the soil water content using cosmic-ray neutron probe in a heterogeneous monsoon climate-dominated region, *Advances in Water Resources*, 108, 125–138, <https://doi.org/10.1016/j.advwatres.2017.07.020>, 2017.
- Nicolai-Shaw, N., Hirschi, M., Mittelbach, H., and Seneviratne, S. I.: Spatial representativeness of soil moisture using in situ, remote sensing, and land reanalysis data, *Journal of Geophysical Research: Atmospheres*, 120, 9955–9964, <https://doi.org/10.1002/2015JD023305>, 2015a.



- Nicolai-Shaw, N., Hirschi, M., Mittelbach, H., and Seneviratne, S. I.: Spatial representativeness of soil moisture using in situ, remote sensing, and land reanalysis data, *Journal of Geophysical Research: Atmospheres*, 120, 9955–9964, <https://doi.org/10.1002/2015JD023305>, 2015b.
- Notarnicola, C., Caporaso, L., Giuseppe, F. D., Temimi, M., Ventura, B., and Zebisch, M.: Inferring soil moisture variability in the Mediterranean Sea area using infrared and passive microwave observations, *Canadian Journal of Remote Sensing*, 38, 46–59, <https://doi.org/10.5589/m12-011>, 2012.
- O, S. and Orth, R.: Global soil moisture from in-situ measurements using machine learning – SoMo.ml, arXiv:2010.02374 [physics], <http://arxiv.org/abs/2010.02374>, arXiv: 2010.02374, 2020.
- Ochsner, T., Cosh, M., Cuenca, R., Dorigo, W., Draper, C., Hagimoto, Y., Kerr, Y., Larson, K., Njoku, E., Small, E., and Zreda, M.: State of the Art in Large-Scale Soil Moisture Monitoring, *Soil Science Society of America Journal*, 77, 1888–1919, <https://doi.org/10.2136/sssaj2013.03.0093>, 2013.
- Ojha, R., Morbidelli, R., Saltalippi, C., Flammini, A., and Govindaraju, R. S.: Scaling of surface soil moisture over heterogeneous fields subjected to a single rainfall event, *Journal of Hydrology*, 516, 21–36, <https://doi.org/10.1016/j.jhydrol.2014.01.057>, determination of soil moisture: Measurements and theoretical approaches, 2014.
- Ojo, E. R., Bullock, P., L. Heures, J., Powers, J., McNairn, H., and Pacheco, A.: Calibration and Evaluation of a Frequency Domain Reflectometry Sensor for Real-Time Soil Moisture Monitoring, *Vadose Zone Journal*, 14, <https://doi.org/10.2136/vzj2014.08.0114>, 2015.
- Osenga, E., Arnott, J., Endsley, K., and Katzenberger, J.: Bioclimatic and Soil Moisture Monitoring Across Elevation in a Mountain Watershed: Opportunities for Research and Resource Management, *Water Resources Research*, <https://doi.org/10.1029/2018WR023653>, 2019.
- Pablos, M., Martínez-Fernández, J., Piles, M., Sánchez, N., Vall-Ilossera, M., and Camps, A.: Multi-Temporal Evaluation of Soil Moisture and Land Surface Temperature Dynamics Using in Situ and Satellite Observations, *Remote Sensing*, 8, 587, <https://doi.org/10.3390/rs8070587>, 2016.
- Pablos, M., González-Zamora, Á., Sánchez, N., and Martínez-Fernández, J.: Assessment of Root Zone Soil Moisture Estimations from SMAP, SMOS and MODIS Observations, *Remote Sensing*, 10, 981, <https://doi.org/10.3390/rs10070981>, 2018.
- Pacheco, A., L'Heures, J., McNairn, H., Powers, J., Howard, A., Geng, X., Rollin, P., Gottfried, K., Freeman, J., Ojo, R., et al.: Real-time in-situ soil monitoring for agriculture (RISMA) network metadata, *Science and Technology*, 2014.
- Pal, M. and Maity, R.: Development of a spatially-varying Statistical Soil Moisture Profile model by coupling memory and forcing using hydrologic soil groups, *Journal of Hydrology*, 570, 141–155, 2019.
- Pal, M., Maity, R., and Dey, S.: Statistical Modelling of Vertical Soil Moisture Profile: Coupling of Memory and Forcing, *Water Resources Management*, 30, <https://doi.org/10.1007/s11269-016-1263-4>, 2016.
- Palmer, J. R. B., Oltra, A., Collantes, F., Delgado, J. A., Lucientes, J., Delacour, S., Bengoa, M., Eritja, R., and Bartumeus, F.: Citizen science provides a reliable and scalable tool to track disease-carrying mosquitoes, *Nature Communications*, 8, 916, <https://doi.org/10.1038/s41467-017-00914-9>, 2017.
- Paloscia, S., Pettinato, S., Santi, E., Notarnicola, C., Pasolli, L., and Reppucci, A.: Soil moisture mapping using Sentinel-1 images: Algorithm and preliminary validation, *Remote Sensing of Environment*, 134, 234–248, <https://doi.org/10.1016/j.rse.2013.02.027>, 2013.
- Pan, X., Kornelsen, K. C., and Coulilaly, P.: Estimating Root Zone Soil Moisture at Continental Scale Using Neural Networks, *JAWRA Journal of the American Water Resources Association*, 53, 220–237, <https://doi.org/10.1111/1752-1688.12491>, 2017.
- Parinussa, R., Wang, G., Holmes, T., Liu, Y., Dolman, A., de Jeu, R., Jiang, T., Zhang, P., and Shi, J.: Global surface soil moisture from the Microwave Radiation Imager onboard the Fengyun-3B satellite, *International Journal of Remote Sensing*, 35, 7007–7029, <https://doi.org/10.1080/01431161.2014.960622>, 2014a.





- Parinussa, R., de Jeu, R., van der Schalie, R., Crow, W., Lei, F., and Holmes, T.: A Quasi-Global Approach to Improve Day-Time Satellite Surface Soil Moisture Anomalies through the Land Surface Temperature Input, *Climate*, 4, 50, <https://doi.org/10.3390/cli4040050>, 2016.
- 1705 Parinussa, R. M., Holmes, T. R. H., and de Jeu, R. A. M.: Soil Moisture Retrievals From the WindSat Spaceborne Polarimetric Microwave Radiometer, *IEEE Transactions on Geoscience and Remote Sensing*, 50, 2683–2694, <https://doi.org/10.1109/TGRS.2011.2174643>, 2012.
- Parinussa, R. M., Yilmaz, M. T., Anderson, M. C., Hain, C. R., and de Jeu, R. A. M.: An intercomparison of remotely sensed soil moisture products at various spatial scales over the Iberian Peninsula, *Hydrological Processes*, 28, 4865–4876, <https://doi.org/10.1002/hyp.9975>, 2014b.
- 1710 Parinussa, R. M., Holmes, T. R. H., Wanders, N., Dorigo, W. A., and de Jeu, R. A. M.: A Preliminary Study toward Consistent Soil Moisture from AMSR2, *Journal of Hydrometeorology*, 16, 932–947, <https://doi.org/10.1175/JHM-D-13-0200.1>, 2015.
- Parinussa, R. M., Wang, G., Liu, Y., Lou, D., Hagan, D. F. T., Zhan, M., Su, B., and Jiang, T.: Improved surface soil moisture anomalies from Fengyun-3B over the Jiangxi province of the People’s Republic of China, *International Journal of Remote Sensing*, 39, 8950–8962, <https://doi.org/10.1080/01431161.2018.1500729>, 2018.
- 1715 Park, C.-H., Montzka, C., Jagdhuber, T., Jonard, F., De Lannoy, G., Hong, J., Jackson, T. J., and Wulfmeyer, V.: A Dielectric Mixing Model Accounting for Soil Organic Matter, *Vadose Zone Journal*, 18, 190 036, 2019.
- Park, C.-H., Jagdhuber, T., Colliander, A., Lee, J., Berg, A., Cosh, M., Kim, S.-b., Kim, Y., and Wulfmeyer, V.: Parameterization of Vegetation Scattering Albedo in the Tau-Omega Model for Soil Moisture Retrieval on Croplands, <https://doi.org/10.20944/preprints202007.0717.v1>, 2020.
- Parrens, M., Zakharova, E., Lafont, S., Calvet, J.-C., Kerr, Y., Wagner, W., and Wigneron, J.-P.: Comparing soil moisture retrievals from SMOS and ASCAT over France, *Hydrology and Earth System Sciences*, 16, 423–440, <https://doi.org/10.5194/hess-16-423-2012>, 2012.
- 1720 Paulik, C., Dorigo, W., Wagner, W., and Kidd, R.: Validation of the ASCAT Soil Water Index using in situ data from the International Soil Moisture Network, *International Journal of Applied Earth Observation and Geoinformation*, 30, 1–8, <https://doi.org/10.1016/j.jag.2014.01.007>, 2014.
- Peel, M. C., Finlayson, B. L., and McMahon, T. A.: Updated world map of the Köppen-Geiger climate classification, *Hydrology and Earth System Sciences Discussions*, 4, 439–473, <https://hal.archives-ouvertes.fr/hal-00298818>, 2007.
- 1725 Peischl, S., Walker, J., Rüdiger, C., Ye, N., Kerr, Y., Kim, E., Bandara, R., and Allahmoradi, M.: The AACES field experiments: SMOS calibration and validation across the Murrumbidgee River catchment, *Hydrology and Earth System Sciences Discussions*, 9, 2763–2795, <https://doi.org/10.5194/hessd-9-2763-2012>, 2012.
- Pellarin, T., de Rosnay, P., Albergel, C., Abdalla, S., and al Bitar, A.: Root-zone soil moisture index complementary validation at global scale based on triple collocation method . Comparison with State-OfThe-Art global scale root-zone soil moisture products, in: H-SAF Visiting Scientist Program HSAF\_CDOP2\_VS12\_02, 2013.
- 1730 Peng, J., Loew, A., Merlin, O., and Verhoest, N. E. C.: A review of spatial downscaling of satellite remotely sensed soil moisture: Downscale Satellite-Based Soil Moisture, *Reviews of Geophysics*, 55, 341–366, <https://doi.org/10.1002/2016RG000543>, 2017.
- Petropoulos, G. P. and McCalmont, J. P.: An operational in situ soil moisture & soil temperature monitoring network for West Wales, UK: The WSMN network, *Sensors*, 17, 1481, 2017.
- 1735 Petropoulos, G. P., Ireland, G., Srivastava, P. K., and Ioannou-Katidis, P.: An appraisal of the accuracy of operational soil moisture estimates from SMOS MIRAS using validated in situ observations acquired in a Mediterranean environment, *International Journal of Remote Sensing*, 35, 5239–5250, <https://doi.org/10.1080/2150704X.2014.933277>, 2014.



- Pierdicca, N., Pulvirenti, L., and Fascetti, F.: Multitemporal soil moisture retrieval from 3-days ERS-2 data: Comparison with  
 1740 ASCAT, SMOS and in situ measurements, in: 2014 IEEE Geoscience and Remote Sensing Symposium, pp. 2459–2462,  
<https://doi.org/10.1109/IGARSS.2014.6946970>, 2014.
- Pierdicca, N., Fascetti, F., Pulvirenti, L., Crapolicchio, R., and Muñoz Sabater, J.: Quadruple Collocation Analysis for Soil Moisture Product  
 Assessment, IEEE Geoscience and Remote Sensing Letters, 12, 1595–1599, <https://doi.org/10.1109/LGRS.2015.2414654>, 2015a.
- Pierdicca, N., Fascetti, F., Pulvirenti, L., Crapolicchio, R., and Muñoz-Sabater, J.: Analysis of ASCAT, SMOS, in-situ and land  
 1745 model soil moisture as a regionalized variable over Europe and North Africa, Remote Sensing of Environment, 170, 280–289,  
<https://doi.org/10.1016/j.rse.2015.09.005>, 2015b.
- Pierdicca, N., Fascetti, F., Pulvirenti, L., and Crapolicchio, R.: Error Characterization of Soil Moisture Satellite Products: Retrieving Error  
 Cross-Correlation Through Extended Quadruple Collocation, IEEE Journal of Selected Topics in Applied Earth Observations and Remote  
 Sensing, 10, 4522–4530, <https://doi.org/10.1109/JSTARS.2017.2714025>, 2017.
- 1750 Piles, M., Petropoulos, G. P., Sánchez, N., Ángel González-Zamora, and Ireland, G.: Towards improved spatio-temporal resolution soil mois-  
 ture retrievals from the synergy of SMOS and MSG SEVIRI spaceborne observations, Remote Sensing of Environment, 180, 403–417,  
<https://doi.org/10.1016/j.rse.2016.02.048>, special Issue: ESA's Soil Moisture and Ocean Salinity Mission - Achievements and Applica-  
 tions, 2016.
- Piles, M., Ballabrera-Poy, J., and Muñoz-Sabater, J.: Dominant features of global surface soil moisture variability observed by the SMOS  
 1755 satellite, Remote Sensing, 11, 95, 2019.
- Portal, G., Jagdhuber, T., Vall-llossera, M., Camps, A., Pablos, M., Entekhabi, D., and Piles, M.: Assessment of Multi-Scale SMOS and  
 SMAP Soil Moisture Products across the Iberian Peninsula, Remote Sensing, 12, 570, <https://doi.org/10.3390/rs12030570>, 2020.
- Pradhan, N. R.: Estimating growing-season root zone soil moisture from vegetation index-based evapotranspiration fraction and soil proper-  
 ties in the Northwest Mountain region, USA, Hydrological Sciences Journal, 64, 771–788, 2019.
- 1760 Pratola, C., Barrett, B., Gruber, A., and Dwyer, E.: Quality Assessment of the CCI ECV Soil Moisture Product Using ENVISAT ASAR Wide  
 Swath Data over Spain, Ireland and Finland, Remote Sensing, 7, 15 388–15 423, <https://doi.org/10.3390/rs71115388>, 2015.
- Przeździecki, K., Zawadzki, J., Cieszewski, C., and Bettinger, P.: Estimation of soil moisture across broad landscapes of Georgia and South  
 Carolina using the triangle method applied to MODIS satellite imagery, Silva Fennica, 51, article id 1683, <https://doi.org/10.14214/sf.1683>,  
 2017.
- 1765 Qin, M., Giménez, D., and Miskewitz, R.: Temporal dynamics of subsurface soil water content estimated from surface measurement using  
 wavelet transform, Journal of Hydrology, 563, 834–850, <https://doi.org/10.1016/j.jhydrol.2018.06.023>, 2018.
- Raffelli, G., Id, M., Previati, M., Id, D., Canone, D., Gisolo, D., Bevilacqua, I., Capello, G., Biddoccu, M., Cavallo, E., Id, R., Deiana, R., Id,  
 G., Cassiani, G., and Ferraris, S.: Local-and Plot-Scale Measurements of Soil Moisture: Time and Spatially Resolved Field Techniques in  
 Plain, Hill and Mountain Sites, Water, 9, <https://doi.org/10.3390/w9090706>, 2018.
- 1770 Rains, D., Han, X., Lievens, H., Montzka, C., and Verhoest, N.: SMOS brightness temperature assimilation into the Community Land Model,  
 HYDROLOGY AND EARTH SYSTEM SCIENCES, 21, 5929–5951, <https://doi.org/10.5194/hess-21-5929-2017>, 2017.
- Rautiainen, K., Parkkinen, T., Lemmetyinen, J., Schwank, M., Wiesmann, A., Ikonen, J., Derksen, C., Davydov, S., Davydova, A., Boike, J.,  
 Langer, M., Drusch, M., and Pulliainen, J.: SMOS prototype algorithm for detecting autumn soil freezing, Remote Sensing of Environment,  
 180, 346–360, <https://doi.org/10.1016/j.rse.2016.01.012>, special Issue: ESA's Soil Moisture and Ocean Salinity Mission - Achievements  
 1775 and Applications, 2016.





- Reichle, R. H., Koster, R. D., Dong, J., and Berg, A. A.: Global Soil Moisture from Satellite Observations, Land Surface Models, and Ground Data: Implications for Data Assimilation, *Journal of Hydrometeorology*, 5, 430–442, [https://doi.org/10.1175/1525-7541\(2004\)005<0430:GSMFSO>2.0.CO;2](https://doi.org/10.1175/1525-7541(2004)005<0430:GSMFSO>2.0.CO;2), 2004.
- Reichle, R. H., Draper, C. S., Liu, Q., Girotto, M., Mahanama, S. P. P., Koster, R. D., and De Lannoy, G. J. M.: Assessment of MERRA-2 Land Surface Hydrology Estimates, *Journal of Climate*, 30, 2937–2960, <https://doi.org/10.1175/JCLI-D-16-0720.1>, 2017.
- Reimer, A. C., Melzer, B. T., Kidd, C. R., and Wagner, D. W.: Validation of the enhanced resolution ERS-2 scatterometer soil moisture product, in: 2012 IEEE International Geoscience and Remote Sensing Symposium, pp. 1208–1211, <https://doi.org/10.1109/IGARSS.2012.6351330>, 2012.
- Rinollo, A., Vulpiani, G., Puca, S., Pagliara, P., Kaňák, J., Labo, E., Okon, L., Roulin, E., Baguis, P., Cattani, E., Laviola, S., and Levizzani, V.: Definition and impact of a quality index for radar-based reference measurements in the H-SAF precipitation product validation, *Natural hazards and earth system sciences*, 13, 2695–2705, <https://doi.org/10.5194/nhess-13-2695-2013>, 2013.
- Robinson, D. A., Campbell, C. S., Hopmans, J. W., Hornbuckle, B. K., Jones, S. B., Knight, R., Ogden, F., Selker, J., and Wendroth, O.: Soil Moisture Measurement for Ecological and Hydrological Watershed-Scale Observatories: A Review, *Vadose Zone Journal*, 7, 358–389, <https://doi.org/10.2136/vzj2007.0143>, 2008.
- Robock, A., Vinnikov, K., Srinivasan, G., Entin, J., Hollinger, S., Speranskaya, N., Liu, S., and Namkhai, A.: The Global Soil Moisture Data Bank, *Bulletin of the American Meteorological Society*, 81, [https://doi.org/10.1175/1520-0477\(2000\)081<1281:TGSMDDB>2.3.CO;2](https://doi.org/10.1175/1520-0477(2000)081<1281:TGSMDDB>2.3.CO;2), 2000.
- Rodionova, N.: Correlation of the Sentinel 1 Radar Data with Ground-Based Measurements of the Soil Temperature and Moisture, *Izvestiya, Atmospheric and Oceanic Physics*, 55, 939–948, 2019a.
- Rodionova, N. V.: Backscattering from the near-surface layer of Thawed/Frozen Soils of Alaska from Sentinel 1 Radar Data, *RENSIT*, 11(1), 21–30, <https://doi.org/10.17725/rensit.2019.11.021>, 2019b.
- Rodriguez-Fernandez, N., Aires, F., Richaume, P., Kerr, Y., Prigent, C., Kolassa, J., Cabot, F., Jimenez, C., Mahmoodi, A., and Drusch, M.: Soil Moisture Retrieval Using Neural Networks: Application to SMOS, *IEEE Transactions on Geoscience and Remote Sensing*, 53, 5991–6007, <https://doi.org/10.1109/TGRS.2015.2430845>, 2015.
- Rodriguez-Fernandez, N., de Rosnay, P., Albergel, C., Richaume, P., Aires, F., Prigent, C., and Kerr, Y.: SMOS Neural Network Soil Moisture Data Assimilation in a Land Surface Model and Atmospheric Impact, *Remote Sensing*, 11, 1334, 2019.
- Romano, N.: Soil moisture at local scale: Measurements and simulations, *Journal of Hydrology*, 516, 6–20, <https://doi.org/10.1016/j.jhydrol.2014.01.026>, 2014.
- Rüdiger, C., Hancock, G., Hemakumara, H., Jacobs, B., Kalma, J., Martinez, C., Thyer, M., Walker, J., Wells, T., and Willgoose, G.: Goulburn River experimental catchment data set, *Water Resources Research*, 43, <https://doi.org/10.1029/2006WR005837>, 2007.
- Sadeghi, M., Gao, L., Ebtehaj, A., Wigneron, J.-P., Crow, W. T., Reager, J. T., and Warrick, A. W.: Retrieving global surface soil moisture from GRACE satellite gravity data, *Journal of Hydrology*, 584, 124 717, <https://doi.org/10.1016/j.jhydrol.2020.124717>, 2020.
- Sadri, S., Pan, M., Wada, Y., Vergopolan, N., Sheffield, J., Famiglietti, J. S., Kerr, Y., and Wood, E.: A global near-real-time soil moisture index monitor for food security using integrated SMOS and SMAP, *Remote Sensing of Environment*, 246, 111 864, <https://doi.org/10.1016/j.rse.2020.111864>, 2020.
- Sanchez, N., Martinez-Fernandez, J., Scaini, A., and Perez-Gutierrez, C.: Validation of the SMOS L2 Soil Moisture Data in the REMEDHUS Network (Spain), *IEEE Transactions on Geoscience and Remote Sensing*, 50, 1602–1611, <https://doi.org/10.1109/TGRS.2012.2186971>, 2012.



- Santi, E., Paloscia, S., Pettinato, S., Brocca, L., and Ciabatta, L.: Robust Assessment of an Operational Algorithm for the Retrieval of Soil  
 1815 Moisture From AMSR-E Data in Central Italy, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 9,  
 2478–2492, 2016.
- Santi, E., Paloscia, S., Pettinato, S., Brocca, L., Ciabatta, L., and Entekhabi, D.: Integration of microwave data from SMAP and AMSR2 for  
 soil moisture monitoring in Italy, *Remote Sensing of Environment*, 212, 21–30, <https://doi.org/10.1016/j.rse.2018.04.039>, 2018.
- Sawada, Y.: Quantifying Drought Propagation from Soil Moisture to Vegetation Dynamics Using a Newly Developed Ecohydrological Land  
 1820 Reanalysis, *Remote Sensing*, 10, 1197, <https://doi.org/10.3390/rs10081197>, 2018.
- Scaini, A., Sánchez, N., Vicente-Serrano, S. M., and Martínez-Fernández, J.: SMOS-derived soil moisture anomalies and drought indices: a  
 comparative analysis using in situ measurements, *Hydrological Processes*, 29, 373–383, <https://doi.org/10.1002/hyp.10150>, 2015.
- Scanlon, T., Dorigo, W., Baum, D., Preimesberger, W., Buttinger, P., Bakcsa, Z., van der Schalie, R., Reimer, C., Hirschi, M., Kidd, R., et al.:  
 ESA CCI and C3S Soil Moisture-Recent advances and quality assurance based on the online validation platform QA4SM., in: *Geophysical*  
 1825 *Research Abstracts*, vol. 21, 2019.
- Schaefer, G., Cosh, M., and Jackson, T.: The USDA natural resources conservation service soil climate analysis network (SCAN), *Journal of*  
*Atmospheric and Oceanic Technology - J ATMOS OCEAN TECHNOL.*, 24, <https://doi.org/10.1175/2007JTECHA930.1>, 2007.
- Schellekens, J., Dutra, E., Martínez-de la Torre, A., Balsamo, G., van Dijk, A., Sperna Weiland, F., Minvielle, M., Calvet, J.-C., Decharme,  
 B., Eisner, S., Fink, G., Flörke, M., Peßenteiner, S., van Beek, R., Polcher, J., Beck, H., Orth, R., Calton, B., Burke, S., Dorigo, W., and  
 1830 Weedon, G. P.: A global water resources ensemble of hydrological models: the earthH2Observe Tier-1 dataset, *Earth System Science Data*,  
 9, 389–413, <https://doi.org/10.5194/essd-9-389-2017>, 2017.
- Schlenz, F., dall’Amico, J. T., Loew, A., and Mauser, W.: Uncertainty Assessment of the SMOS Validation in the Upper Danube Catchment,  
*IEEE Transactions on Geoscience and Remote Sensing*, 50, 1517–1529, 2012.
- Schlenz, F., dall’Amico, J. T., Mauser, W., and Loew, A.: Analysis of SMOS brightness temperature and vegetation optical depth data  
 1835 with coupled land surface and radiative transfer models in Southern Germany, *Hydrology and Earth System Sciences*, 16, 3517–3533,  
<https://doi.org/10.5194/hess-16-3517-2012>, 2012.
- Scholze, M., Kaminski, T., Knorr, W., Blessing, S., Vossbeck, M., Grant, J., and Scipal, K.: Simultaneous assimilation of SMOS soil mois-  
 ture and atmospheric CO<sub>2</sub> in-situ observations to constrain the global terrestrial carbon cycle, *Remote Sensing of Environment*, 180,  
 334–345, <https://doi.org/10.1016/j.rse.2016.02.058>, special Issue: ESA’s Soil Moisture and Ocean Salinity Mission - Achievements and  
 1840 Applications, 2016.
- Senanayake, I., Yeo, I.-Y., Tangdamrongsub, N., Willgoose, G., Hancock, G., Wells, T., Fang, B., Lakshmi, V., and Walker, J.: An in-situ  
 data based model to downscale radiometric satellite soil moisture products in the Upper Hunter Region of NSW, Australia, *Journal of*  
*Hydrology*, 572, 820–838, <https://doi.org/10.1016/j.jhydrol.2019.03.014>, 2019.
- Senyurek, V., Lei, F., Boyd, D., Kurum, M., Gurbuz, A. C., and Moorhead, R.: Machine Learning-Based CYGNSS Soil Moisture Estimates  
 1845 over ISMN sites in CONUS, *Remote Sensing*, 12, 1168, <https://doi.org/10.3390/rs12071168>, 2020.
- Sheng, J., Rao, P., and Ma, H.: Spatial Downscaling of the FY3B Soil Moisture Using Random Forest Regression, in: *2019 8th International*  
*Conference on Agro-Geoinformatics (Agro-Geoinformatics)*, pp. 1–6, 2019.
- Shin, Y., Lim, K., Park, K., and Jung, Y.: Development of Dynamic Ground Water Data Assimilation for Quantifying Soil Hydraulic Proper-  
 ties from Remotely Sensed Soil Moisture, *Water*, 8, 311, <https://doi.org/10.3390/w8070311>, 2016.
- 1850 Shin, Y., Mohanty, B. P., and Ines, A. V.: Development of non-parametric evolutionary algorithm for predicting soil moisture dynamics,  
*Journal of Hydrology*, 564, 208–221, <https://doi.org/10.1016/j.jhydrol.2018.07.003>, 2018.



- Shuman, D. I., Nayyar, A., Mahajan, A., Goykhman, Y., Li, K., Liu, M., Teneketzis, D., Moghaddam, M., and Entekhabi, D.: Measurement Scheduling for Soil Moisture Sensing: From Physical Models to Optimal Control, *Proceedings of the IEEE*, 98, 1918–1933, <https://doi.org/10.1109/JPROC.2010.2052532>, 2010.
- 1855 Skierucha, W., Sławiński, C., Wilczek, A., Żyromski, A., and Biniak-Pieróg, M.: Telemetric system for the measurement of soil moisture based on the TDR technique, *Woda Środowisko Obszary Wiejskie*, 12, 257–267, 2012a.
- Skierucha, W., Wilczek, A., Szyplowska, A., Sławiński, C., and Lamorski, K.: A TDR-Based Soil Moisture Monitoring System with Simultaneous Measurement of Soil Temperature and Electrical Conductivity, *Sensors*, 12, 13 545–13 566, <https://doi.org/10.3390/s121013545>, 2012b.
- 1860 Smith, A., Walker, J., Western, A., Young, R., Ellett, K., Pipunic, R., Grayson, R., Siriwardena, L., Chiew, F., and Richter, H.: The Murrumbidgee Soil Moisture Monitoring Network data set, *Water Resources Research*, 48, 7701–, <https://doi.org/10.1029/2012WR011976>, 2012.
- Solander, K. C., Newman, B. D., Carioca de Araujo, A., Barnard, H. R., Berry, Z. C., Bonal, D., Bretfeld, M., Burban, B., Antonio Candido, L., Céleri, R., Chambers, J. Q., Christoffersen, B. O., Detto, M., Dorigo, W. A., Ewers, B. E., José Filgueiras Ferreira, S., Knohl, A.,  
 1865 Leung, L. R., McDowell, N. G., Miller, G. R., Terezinha Ferreira Monteiro, M., Moore, G. W., Negron-Juarez, R., Saleska, S. R., Stiegler, C., Tomasella, J., and Xu, C.: The pantropical response of soil moisture to El Niño, *Hydrology and Earth System Sciences*, 24, 2303–2322, <https://doi.org/10.5194/hess-24-2303-2020>, 2020.
- Su, B., Wen, J., Dente, L., Velde, R., Wang, L., Ma, Y., Yang, K., and Hu, Z.: The Tibetan Plateau observatory of plateau scale soil moisture and soil temperature (Tibet-Obs) for quantifying uncertainties in coarse resolution satellite and model products, *Hydrology and Earth  
 1870 System Sciences*, 15, 2303, <https://doi.org/10.5194/hess-15-2303-2011>, 2011.
- Su, C.-H., Narsey, S. Y., Gruber, A., Xaver, A., Chung, D., Ryu, D., and Wagner, W.: Evaluation of post-retrieval de-noising of active and passive microwave satellite soil moisture, *Remote Sensing of Environment*, 163, 127–139, <https://doi.org/10.1016/j.rse.2015.03.010>, 2015.
- Su, C.-H., Ryu, D., Dorigo, W., Zwieback, S., Gruber, A., Albergel, C., Reichle, R. H., and Wagner, W.: Homogeneity of a global multisatellite  
 1875 soil moisture climate data record, *Geophysical Research Letters*, 43, 11,245–11,252, <https://doi.org/10.1002/2016GL070458>, 2016.
- Su, Y., Chen, X., Su, H., Liu, L., and Liao, J.: Digitizing the thermal and hydrological parameters of land surface in subtropical China using AMSR-E brightness temperatures, *International Journal of Digital Earth*, 10, 687–700, <https://doi.org/10.1080/17538947.2016.1247472>, 2017.
- Suman, S., Srivastava, P. K., Petropoulos, G. P., Pandey, D. K., and O'Neill, P. E.: Appraisal of SMAP Operational Soil Moisture Product  
 1880 from a Global Perspective, *Remote Sensing*, 12, 1977, <https://doi.org/10.3390/rs12121977>, 2020.
- Sun, H., Zhou, B., Zhang, C., Liu, H., and Yang, B.: DSCALE\_mod16: A Model for Disaggregating Microwave Satellite Soil Moisture with Land Surface Evapotranspiration Products and Gridded Meteorological Data, *Remote Sensing*, 12, 980, <https://doi.org/10.3390/rs12060980>, 2020.
- Sure, A., Varade, D., and Dikshit, O.: Factors determining spatio-temporal variations of soil moisture using microwave data,  
 1885 in: 2017 International Conference on Emerging Trends in Computing and Communication Technologies (ICETCCT), pp. 1–5, <https://doi.org/10.1109/ICETCCT.2017.8280301>, 2017.
- Szczypta, C., Calvet, J.-C., Maignan, F., Dorigo, W., Baret, F., and Ciais, P.: Suitability of modelled and remotely sensed essential climate variables for monitoring Euro-Mediterranean droughts, *Geoscientific Model Development*, 7, 931–946, <https://doi.org/10.5194/gmd-7-931-2014>, 2014.



- 1890 Tagesson, T., Horion, S., Nieto, H., Zaldo Fornies, V., Mendiguren González, G., Bulgin, C., Ghent, D., and Fensholt, R.: Disaggregation of SMOS soil moisture over West Africa using the Temperature and Vegetation Dryness Index based on SEVIRI land surface parameters, *Remote Sensing of Environment*, 206, 424–441, <https://doi.org/10.1016/j.rse.2017.12.036>, 2018.
- Teuling, A. J., Uijlenhoet, R., Hupet, F., van Loon, E. E., and Troch, P. A.: Estimating spatial mean root-zone soil moisture from point-scale observations, *Hydrol. Earth Syst. Sci.*, p. 13, 2006.
- 1895 Tian, J., Zhang, B., He, C., Han, Z., Bogena, H. R., and Huisman, J. A.: Dynamic response patterns of profile soil moisture wetting events under different land covers in the Mountainous area of the Heihe River Watershed, Northwest China, *Agricultural and forest meteorology*, 271, 225–239, 2019a.
- Tian, S.: Monitoring and forecasting drought through the assimilation of satellite water observations, Ph.D. thesis, Research School of Earth Sciences, The Australian National University, <https://doi.org/doi.org/10.25911/5c94abe15c9ec>, 2018.
- 1900 Tian, S., Renzullo, L. J., van Dijk, A. I. J. M., Tregoning, P., and Walker, J. P.: Global joint assimilation of GRACE and SMOS for improved estimation of root-zone soil moisture and vegetation response, *Hydrology and Earth System Sciences*, 23, 1067–1081, <https://doi.org/10.5194/hess-23-1067-2019>, 2019b.
- Tobin, K. J., Torres, R., Crow, W. T., and Bennett, M. E.: Multi-decadal analysis of root-zone soil moisture applying the exponential filter across CONUS, *Hydrology and Earth System Sciences*, 21, 4403–4417, <https://doi.org/10.5194/hess-21-4403-2017>, 2017.
- 1905 Vachaud, G., Passerat De Silans, A., Balabanis, P., and Vauclin, M.: Temporal Stability of Spatially Measured Soil Water Probability Density Function, *Soil Science Society of America Journal*, 49, 822–828, <https://doi.org/10.2136/sssaj1985.03615995004900040006x>, 1985.
- Van Cleve, Keith; Chapin, F. S. R. R. W.: Bonanza Creek Long Term Ecological Research Project Climate Database - University of Alaska Fairbanks. <http://www.lter.uaf.edu/>, 2015.
- van der Schalie, R., Kerr, Y., Wigneron, J., Rodríguez-Fernández, N., Al-Yaari, A., and Jeu, R.: Global SMOS Soil Moisture Retrievals from The Land Parameter Retrieval Model, *International Journal of Applied Earth Observation and Geoinformation*, 45, 125–134, <https://doi.org/10.1016/j.jag.2015.08.005>, advances in the Validation and Application of Remotely Sensed Soil Moisture - Part 1, 2016.
- 1910 van der Schalie, R., de Jeu, R., Kerr, Y., Wigneron, J., Rodríguez-Fernández, N., Al-Yaari, A., Parinussa, R., Mecklenburg, S., and Drusch, M.: The merging of radiative transfer based surface soil moisture data from SMOS and AMSR-E, *Remote Sensing of Environment*, 189, 180–193, <https://doi.org/10.1016/j.rse.2016.11.026>, 2017.
- 1915 van der Schalie, R., de Jeu, R., Parinussa, R., Rodríguez-Fernández, N., Kerr, Y., Al-Yaari, A., Wigneron, J.-P., and Drusch, M.: The Effect of Three Different Data Fusion Approaches on the Quality of Soil Moisture Retrievals from Multiple Passive Microwave Sensors, *Remote Sensing*, 10, 107, <https://doi.org/10.3390/rs10010107>, 10.3390/rs10010107, 2018.
- Van doninck, J., Peters, J., Lievens, H., De Baets, B., and Verhoest, N.: Accounting for seasonality in a soil moisture change detection algorithm for ASAR Wide Swath time series, *HYDROLOGY AND EARTH SYSTEM SCIENCES*, 16, 773–786, <https://doi.org/10.5194/hess-16-773-2012>, 2012.
- 1920 Verrier, S.: Multifractal and multiscale entropy scaling of in-situ soil moisture time series: Study of SMOSMANIA network data, southwestern France, *Journal of Hydrology*, 585, 124–131, <https://doi.org/10.1016/j.jhydrol.2020.124821>, 2020.
- Vicente-Serrano, S. M., Beguería, S., Lorenzo-Lacruz, J., Camarero, J. J., López-Moreno, J. I., Azorin-Molina, C., Revuelto, J., Morán-Tejeda, E., and Sanchez-Lorenzo, A.: Performance of Drought Indices for Ecological, Agricultural, and Hydrological Applications, *Earth Interactions*, 16, 1–27, <https://doi.org/10.1175/2012EI000434.1>, 2012.



- Vreugdenhil, M., Dorigo, W., Broer, M., Haas, P., Eder, A., Hogan, P., Bloeschl, G., and Wagner, W.: Towards a high-density soil moisture network for the validation of SMAP in Petzenkirchen, Austria, in: 2013 IEEE International Geoscience and Remote Sensing Symposium - IGARSS, pp. 1865–1868, <https://doi.org/10.1109/IGARSS.2013.6723166>, 2013.
- Wanders, N., Karssenbergh, D., Bierkens, M., Parinussa, R., de Jeu, R., van Dam, J., and de Jong, S.: Observation uncertainty of satellite soil moisture products determined with physically-based modeling, *Remote Sensing of Environment*, 127, 341–356, <https://doi.org/10.1016/j.rse.2012.09.004>, 2012.
- Wang, L., Li, X., Chen, Y., Yang, K., Chen, D., Zhou, J., Liu, W., Qi, J., and Huang, J.: Validation of the global land data assimilation system based on measurements of soil temperature profiles, *Agricultural and Forest Meteorology*, 218–219, 288–297, <https://doi.org/10.1016/j.agrformet.2016.01.003>, 2016.
- Wang, L., He, B., Bai, X., and Xing, M.: Assessment of different vegetation parameters for parameterizing the coupled water cloud model and advanced integral equation model for soil moisture retrieval using time series Sentinel-1A data, *Photogrammetric Engineering & Remote Sensing*, 85, 43–54, 2019a.
- Wang, L., Xie, Z., Jia, B., Xie, J., Wang, Y., Liu, B., Li, R., and Chen, S.: Contributions of climate change and groundwater extraction to soil moisture trends, *Earth System Dynamics*, 10, 599–615, <https://doi.org/10.5194/esd-10-599-2019>, 2019b.
- Wang, L., Fang, S., Pei, Z., Zhu, Y., Khoi, D. N., and Han, W.: Using FengYun-3C VSM Data and Multivariate Models to Estimate Land Surface Soil Moisture, *Remote Sensing*, 12, 1038, <https://doi.org/10.3390/rs12061038>, 2020a.
- Wang, T., Franz, T. E., Li, R., You, J., Shulski, M. D., and Ray, C.: Evaluating climate and soil effects on regional soil moisture spatial variability using EOFs, *Water Resources Research*, 53, 4022–4035, <https://doi.org/10.1002/2017WR020642>, 2017.
- Wang, Y., Yang, J., Chen, Y., Fang, G., Duan, W., Li, Y., and De Maeyer, P.: Quantifying the Effects of Climate and Vegetation on Soil Moisture in an Arid Area, China, *Water*, 11, <https://doi.org/10.3390/w11040767>, 2019c.
- Wang, Y., Shi, L., Lin, L., Holzman, M., Carmona, F., and Zhang, Q.: A robust data-worth analysis framework for soil moisture flow by hybridizing sequential data assimilation and machine learning, *Vadose Zone Journal*, 19, e20026, <https://doi.org/10.1002/vzj2.20026>, 2020b.
- WANG Rui, ZHU Qingke, M. H. e. a.: Spatial and temporal variations of the surface soil moisture in the source region of the Yellow River from 2003 to 2010 based on AMSR-E[J]., *SSWC*, 15(1), 22–32, 2017.
- Western, A. W., Grayson, R. B., and Blöschl, G.: Scaling of Soil Moisture: A Hydrologic Perspective, *Annual Review of Earth and Planetary Sciences*, 30, 149–180, <https://doi.org/10.1146/annurev.earth.30.091201.140434>, 2002.
- Wigneron, J.-P., Dayan, S., Kruszwski, A., Aluome, C., Al-Yaari, A., Fan, L., Guven, S., Chipeaux, C., Moisy, C., Guyon, D., and Loustau, D.: The Aqui Network: Soil Moisture Sites in the “Les Landes” Forest and Graves Vineyards (Bordeaux Aquitaine Region, France), pp. 3739–3742, <https://doi.org/10.1109/IGARSS.2018.8517392>, 2018.
- Wigneron, J.-P., Li, X., Frappart, F., Fan, L., Al-Yaari, A., De Lannoy, G., Liu, X., Wang, M., Le Masson, E., and Moisy, C.: SMOS-IC data record of soil moisture and L-VOD: Historical development, applications and perspectives, *Remote Sensing of Environment*, 254, 112 238, <https://doi.org/10.1016/j.rse.2020.112238>, 2021.
- Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., Gonzalez-Beltran, A., Gray, A. J., Groth, P., Goble, C., Grethe, J. S., Heringa, J., ’t Hoen, P. A., Hooft, R., Kuhn, T., Kok, R., Kok, J., Lusher, S. J., Martone, M. E., Mons, A., Packer, A. L., Persson, B., Rocca-Serra, P., Roos, M., van Schaik, R., Sansone, S.-A., Schultes, E., Sengstag, T., Slater, T., Strawn, G., Swertz, M. A., Thompson, M., van der Lei, J., van Mulligen, E., Velterop, J., Waagmeester, A., Wittenburg, P.,



- Wolstencroft, K., Zhao, J., and Mons, B.: The FAIR Guiding Principles for scientific data management and stewardship, *Scientific Data*, 3, 160018, <https://doi.org/10.1038/sdata.2016.18>, 2016.
- Williams, S.: The Coordinated Energy and Water cycle Observations Project (CEOP) Data Integration Approach, EMS, p. 798, 2010.
- Wu, M., Scholze, M., Voßbeck, M., Kaminski, T., and Hoffmann, G.: Simultaneous Assimilation of Remotely Sensed Soil Moisture and FAPAR for Improving Terrestrial Carbon Fluxes at Multiple Sites Using CCDAS, *Remote Sensing*, 11, 27, <https://doi.org/10.3390/rs11010027>, 2018.
- Wu, Q., Liu, H., Wang, L., and Deng, C.: Evaluation of AMSR2 soil moisture products over the contiguous United States using in situ data from the International Soil Moisture Network, *International Journal of Applied Earth Observation and Geoinformation*, 45, 187–199, <https://doi.org/10.1016/j.jag.2015.10.011>, advances in the Validation and Application of Remotely Sensed Soil Moisture - Part 1, 2016.
- Xaver, A., Zappa, L., Rab, G., Pfeil, I., Vreugdenhil, M., Hemment, D., and Dorigo, W. A.: Evaluating the suitability of the consumer low-cost Parrot Flower Power soil moisture sensor for scientific environmental applications, *Geoscientific Instrumentation, Methods and Data Systems*, 9, 117–139, <https://doi.org/10.5194/gi-9-117-2020>, 2020.
- Xia, Y., Ford, T. W., Wu, Y., Quiring, S. M., and Ek, M. B.: Automated Quality Control of In Situ Soil Moisture from the North American Soil Moisture Database Using NLDAS-2 Products, *Journal of Applied Meteorology and Climatology*, 54, 1267–1282, <https://doi.org/10.1175/JAMC-D-14-0275.1>, 2015.
- Xia, Y., Hao, Z., Shi, C., Li, Y., Meng, J., Xu, T., Wu, X., and Zhang, B.: Regional and Global Land Data Assimilation Systems: Innovations, Challenges, and Prospects, *Journal of Meteorological Research*, 33, 159–189, <https://doi.org/10.1007/s13351-019-8172-4>, 2019.
- Xie, Q., Menenti, M., and Jia, L.: Improving the AMSR-E/NASA Soil Moisture Data Product Using In-Situ Measurements from the Tibetan Plateau, *Remote Sensing*, 11, <https://doi.org/10.3390/rs11232748>, 2019.
- Xing, C., Chen, N., Zhang, X., and Gong, J.: A Machine Learning Based Reconstruction Method for Satellite Remote Sensing of Soil Moisture Images with In Situ Observations, *Remote Sensing*, 9, <https://doi.org/10.3390/rs9050484>, 2017.
- Xu, H., Yuan, Q., Li, T., Shen, H., Zhang, L., and Jiang, H.: Quality Improvement of Satellite Soil Moisture Products by Fusing with In-Situ Measurements and GNSS-R Estimates in the Western Continental U.S., *Remote Sensing*, 10, 1351, <https://doi.org/10.3390/rs10091351>, 2018.
- Yang, K., Qin, J., Zhao, L., Chen, Y., Tang, W., Han, M., Zhu, L., Chen, Z., Lv, N., Ding, B., Wu, H., and Lin, C.: A Multi-Scale Soil Moisture and Freeze-Thaw Monitoring Network on the Third Pole, *Bulletin of the American Meteorological Society*, 94, 1907–1916, <https://doi.org/10.1175/BAMS-D-12-00203.1>, 2013.
- Yao, P., Shi, J., Zhao, T., Lu, H., and Al-Yaari, A.: Rebuilding Long Time Series Global Soil Moisture Products Using the Neural Network Adopting the Microwave Vegetation Index, *Remote Sensing*, 9, <https://doi.org/10.3390/rs9010035>, 2017.
- Yin, Z., Ottlé, C., Ciais, P., Guimberteau, M., Wang, X., Zhu, D., Maignan, F., Peng, S., Piao, S., Polcher, J., Zhou, F., Kim, H., and other China-Trend-Stream project members: Evaluation of ORCHIDEE-MICT-simulated soil moisture over China and impacts of different atmospheric forcing data, *Hydrology and Earth System Sciences*, 22, 5463–5484, <https://doi.org/10.5194/hess-22-5463-2018>, <https://hess.copernicus.org/articles/22/5463/2018/>, 2018.
- Zacharias, S., Bogen, H., Samaniego, L., Mauder, M., Fuß, R., Pütz, T., Frenzel, M., Schwank, M., Baessler, C., Butterbach-Bahl, K., Bens, O., Borg, E., Brauer, A., Dietrich, P., Hajnsek, I., Helle, G., Kiese, R., Kunstmann, H., Klotz, S., and Vereecken, H.: A Network of Terrestrial Environmental Observatories in Germany, *Vadose Zone Journal*, 10, 955–973, <https://doi.org/10.2136/vzj2010.0139>, 2011.
- Zappa, L., Forkel, M., Xaver, A., and Dorigo, W.: Deriving Field Scale Soil Moisture from Satellite Observations and Ground Measurements in a Hilly Agricultural Region, *Remote Sensing*, 11, 2596, <https://doi.org/10.3390/rs11222596>, 2019.





- Zappa, L., Woods, M., Hemment, D., Xaver, A., and Dorigo, W.: Evaluation of Remotely Sensed Soil Moisture Products using Crowdsourced Measurements, Eighth International Conference on Remote Sensing and Geoinformation of Environment, SPIE, Cyprus, 2020.
- Zawadzki, J. and Kędzior, M.: Soil moisture variability over Odra watershed: Comparison between SMOS and GLDAS data, International Journal of Applied Earth Observation and Geoinformation, 45, 110–124, <https://doi.org/10.1016/j.jag.2015.03.005>, advances in the Validation and Application of Remotely Sensed Soil Moisture - Part 1, 2016.
- Zeng, J., Li, Z., Quan, C., and Bi, H.: A simplified physically-based algorithm for surface soil moisture retrieval using AMSR-E data, Frontiers of Earth Science, 8, <https://doi.org/10.1007/s11707-014-0412-4>, 2014.
- Zeng, J., Chen, K., Bi, H., and Chen, Q.: A Preliminary Evaluation of the SMAP Radiometer Soil Moisture Product Over United States and Europe Using Ground-Based Measurements, IEEE Transactions on Geoscience and Remote Sensing, 54, 4929–4940, 2016.
- Zhang, Q., Shi, L., Holzman, M., Ye, M., Wang, Y., Carmona, F., and Zha, Y.: A dynamic data-driven method for dealing with model structural error in soil moisture data assimilation, Advances in Water Resources, 132, 103 407, <https://doi.org/10.1016/j.advwatres.2019.103407>, 2019a.
- Zhang, R., Kim, S., and Sharma, A.: A comprehensive validation of the SMAP Enhanced Level-3 Soil Moisture product using ground measurements over varied climates and landscapes, Remote Sensing of Environment, 223, 82–94, 2019b.
- Zhang, S., Calvet, J.-C., Darrozes, J., Roussel, N., Frappart, F., and Bouhours, G.: Deriving surface soil moisture from reflected GNSS signal observations from a grassland site in southwestern France, Hydrology and Earth System Sciences, 22, 1931–1946, <https://doi.org/10.5194/hess-22-1931-2018>, 2018a.
- Zhang, S., Meurey, C., and Calvet, J.-C.: Identification of soil-cooling rains in southern France from soil temperature and soil moisture observations, Atmospheric Chemistry and Physics, 19, 5005–5020, <https://doi.org/10.5194/acp-19-5005-2019>, 2019c.
- Zhang, X., Chen, N., Chen, Z., Wu, L., Li, X., Zhang, L., Di, L., Gong, J., and Li, D.: Geospatial sensor web: A cyber-physical infrastructure for geoscience research and application, Earth-science reviews, 185, 684–703, 2018b.
- Zhao, L. and Yang, Z.-L.: Multi-sensor land data assimilation: Toward a robust global soil moisture and snow estimation, Remote Sensing of Environment, 216, 13–27, <https://doi.org/10.1016/j.rse.2018.06.033>, 2018.
- Zhao, L., Yang, Z.-L., and Hoar, T. J.: Global Soil Moisture Estimation by Assimilating AMSR-E Brightness Temperatures in a Coupled CLM4–RTM–DART System, Journal of Hydrometeorology, 17, 2431–2454, <https://doi.org/10.1175/JHM-D-15-0218.1>, 2016.
- Zhao, T., Shi, J., Lin, M., Yin, X., Liu, Y., Lan, H., and Xiong, C.: Potential soil moisture product from the Chinese HY-2 scanning microwave radiometer and its initial assessment, Journal of Applied Remote Sensing, 8, 1–12, <https://doi.org/10.1117/1.JRS.8.083560>, 2014.
- Zhao, W. and Li, A.: A Downscaling Method for Improving the Spatial Resolution of AMSR-E Derived Soil Moisture Product Based on MSG-SEVIRI Data, Remote Sensing, 5, 6790–6811, <https://doi.org/10.3390/rs5126790>, 2013.
- Zhao, W. and Li, A.: A comparison study on empirical microwave soil moisture downscaling methods based on the integration of microwave-optical/IR data on the Tibetan Plateau, International Journal of Remote Sensing, 36, 4986–5002, <https://doi.org/10.1080/01431161.2015.1041178>, 2015.
- Zhao, W., Li, A., Jin, H., Zhang, Z., Bian, J., and Yin, G.: Performance Evaluation of the Triangle-Based Empirical Soil Moisture Relationship Models Based on Landsat-5 TM Data and In Situ Measurements, IEEE Transactions on Geoscience and Remote Sensing, 55, 2632–2645, <https://doi.org/10.1109/TGRS.2017.2649522>, 2017.
- Zhao, W., Sánchez, N., Lu, H., and Li, A.: A spatial downscaling approach for the SMAP passive surface soil moisture product using random forest regression, Journal of Hydrology, 563, 1009–1024, <https://doi.org/10.1016/j.jhydrol.2018.06.081>, 2018.



- Zhu, L., Wang, H., Tong, C., Liu, W., and Du, B.: Evaluation of ESA Active, Passive and Combined Soil Moisture Products Using Upscaled  
 2040 Ground Measurements, *Sensors*, 19, 2718, 2019.
- Zreda, M., Desilets, D., Ferré, T., and Scott, R.: Measuring soil moisture content non-invasively at intermediate spatial scale using cosmic-ray  
 neutrons, *Geophysical Research Letters*, 35, <https://doi.org/10.1029/2008GL035655>, 2008.
- Zreda, M., Shuttleworth, W. J., Zeng, X., Zweck, C., Desilets, D., Franz, T., and Rosolem, R.: COSMOS: the COsmic-ray Soil Moisture  
 Observing System, *Hydrology and Earth System Sciences*, 16, 4079–4099, <https://doi.org/10.5194/hess-16-4079-2012>, 2012.
- 2045 Zwieback, S., Dorigo, W., and Wagner, W.: Estimation of the temporal autocorrelation structure by the collocation technique with an emphasis  
 on soil moisture studies, *Hydrological Sciences Journal*, 58, 1729–1747, <https://doi.org/10.1080/02626667.2013.839876>, 2013.
- Zwieback, S., Paulik, C., and Wagner, W.: Frozen Soil Detection Based on Advanced Scatterometer Observations and Air Temperature Data  
 as Part of Soil Moisture Retrieval, *Remote Sensing*, 7, 3206–3231, <https://doi.org/10.3390/rs70303206>, 2015.
- Zwieback, S., Su, C.-H., Gruber, A., Dorigo, W. A., and Wagner, W.: The Impact of Quadratic Nonlinear Relations between Soil Moisture  
 2050 Products on Uncertainty Estimates from Triple Collocation Analysis and Two Quadratic Extensions, *Journal of Hydrometeorology*, 17,  
 1725–1743, <https://doi.org/10.1175/JHM-D-15-0213.1>, 2016.