

Dear Editor, dear Erwin Zehe,

We thank you for the promising words and are very glad to submit the revised manuscript after incorporating the minor revisions made by the referees, which further contribute to improving the manuscript.

In the revision, we mainly adapted Figure 2 as suggested by Referee #1 and we have revised the discussion based on the comments made by Referee #2.

We look forward to hearing from you.

Best wishes,

Laurène Bouaziz and co-authors.

[Reply to Anonymous Referee # 1](#)

Overall assessment by Referee #1:

I want to thank the authors for their detailed reply to my comments. I think that they have been well addressed in the revised version of the manuscript which further improved the manuscript.

Reply:

We thank Referee #1 for the positive words on our revised manuscript. We have addressed the remaining minor suggestions and reply to them below.

Comment 1:

L. 19-21: Revise this sentence for better readability (“with in the non-stationary scenarios upto...” is hard to follow)

Reply 1:

Agreed, we have split this sentence in two to increase readability.

Comment 2:

L. 740-743: “This root-accessible water volume is independent of soil type ...”, is there a reference for these two sentences?

Reply 2:

We have added relevant references for these sentences.

Comment 3:

- L. 793: “increasing its root-zone storage capacity **by** 34%” -> replace “with” with “by”

Reply 3:

Agreed, we have corrected with “by”.

Comment 4:

Figure 2: This graph is a nice addition to the method section, as it nicely supports understanding of the methods workflow. However, I think that it should be revised a bit for more structure, readability and consistency. The workflow/concept presented in the figure should also be clear to someone that has not read the full method section. I have a few suggestions that might help revising the figure:

- Text:

- Could be a shorter and more consistent across the columns to make clear how climate and input parameters differ between columns.
 - E.g. “fixed $S_{R,max,A}$ parameter” (col 3) vs. “ $S_{R,max,A}$ ” (col 4) is inconsistent
 - Maybe instead of text, you could put two bullet points *Climate: xxx* and *Rootzone storage capacity: xxx* in each modeling column. This would make it easy to compare at one glance in which ways columns differ from each other?
- Headings in the sub columns might help for structure of workflow
 - E.g. col (3) Calibration/Model calibration, col (4) Evaluation/Validation
- 4 Scenarios:
 - It is a bit confusing to have the historical scenario from (4) also in each of the scenario description of (5). At a first glance, it looks like the reader has to understand 8 instead of 4 scenarios until they figure out that 4 of them are the same. I think the concept of 4 climate and land-use change scenarios that are all compared to the same historical scenario could be made clearer. You could, for example, have the scenario description of the historical scenario in col (4) and the four 2K scenarios in column (5). Then, in column (6) you explain that you compared all scenarios from (5) to the scenario from (4)
- Currently, there is only one arrow linking the output $S_{R,max,A}$ from col (2) with column (3). But shouldn't there also be arrows linking the 4 Q/P Outputs from col (1) with col (2) and the remaining 3 $S_{R,max}$ B-D from col (2) with the scenarios in col (5)? I realize that in the current structure of the plot this might be difficult. Maybe it's easier if you flipped the plot structure to a top-bottom instead of left-right structure?
- Shading and boxes could be improved to support the workflow better:
 - Too many shades of gray (e.g. headings are different from columns) make it difficult to see the clear main workflow from “Estimate $S_{R,max}$ ” -> Modeling -> Change evaluation.

Reply 4:

We thank the referee for the good suggestions to improve the consistency and readability of the Figure, we have been playing around with the design and have made the following changes:

- *We think it is indeed a very good idea to flip the layout, this also reduces the shades of grey and makes it easier to add titles for each step of the procedure*
- *We rearranged the 4 scenarios and only show the historical schema once.*
- *We removed the arrow, as it should already be clear from the text that we are using results from a previous step.*

[Reply to Anonymous Referee # 2](#)

Overall assessment by Referee #2:

I have already reviewed the previous version of this manuscript. In the revision the authors have really improved the accessibility of the manuscript and included much more interpretation of the results in an extended discussion.

The Methods section is much improved, the collection of all the model information in Appendix 3 works well. Thanks for including some details, e.g. a short note on how transpiration decreases when root zone storage dries. Those pointers will really help readers (like me) to focus on the main message and not be distracted by searching. The discussion has much increased in size and enhances the quality of the manuscript. Most of my comments have been satisfactorily addressed or ruled out. Mainly on the new discussion, I have some more comments below. I propose to include a couple more aspects. Also I propose to bring some of the sections, especially in the „limitations“-part, to a resolution, e.g. how can those be tackled in the future?

I believe only minor revisions are required.

[Reply:](#)

We thank Referee #2 for positively reassessing our revised manuscript. We have addressed the remaining constructive comments and reply to them below.

Comment 1:

L 308 „We repeat this random sampling several times .. several model parameters“

Ok, I did not understand this in the previous version. It remains unclear at this point: ^[1]_{SEP} How much is several, and what happens to the sample? Do you proceed with each individual realization or some aggregated (average) runoff coefficient?

[Reply 1:](#)

We have clarified that we repeated the random sampling seven times and that we indeed work with the ensemble of each individual realization in the subsequent analyses, e.g. when plotting the uncertainty bands in Fig 4c, 7 and 8.

Comment 2:

L 279-281: It is really mostly the rooting depth that will be adaptable. I propose to include a short comment here, to help the reader make the association between rooting storage and rooting depth, which now change with vegetation type.

[Reply 2:](#)

This is a good suggestion, we have included a short sentence in the following paragraph (line 288 in tracked change manuscript) to clarify that the adaptation of the root-zone storage capacity mainly relates to changes in the rooting system (rooting depth and/or rooting density).

Comment 3:

Discussion: The role of the investigation of land use change is still a bit opaque in the manuscript. The Methods section says on lines 224-226: „These scenarios are meant as a sensitivity analysis in the spirit of trading space-for-time (Singh et al., 2011) to evaluate the effect of potential future land-use management on the overall water balance.“

I have interpret this to say that climate change scenarios will be evaluated in the light of the additional affect of land use change? This sounds very reasonable to me. But I wondered, why this idea was not taken up in the discussion? The „implications“ section of the discussion revolves mostly around climate change and I missed the reflection on the effects of the vegetation transformation. Can you more specifically return to the comment made earlier in the manuscript?

Reply 3:

Indeed, we evaluate the effect of the additional effect of potential land use change and the referee is completely right that it would be good to come back to it in the Implication section of the discussion. Now, the relatively limited additional effects of land-use change on the hydrological response are mentioned in section 6.2.2 “On the potential land-use change scenarios”, but this is indeed part of the Limitation section. We have added a sentence (lines 537-539 of the tracked change manuscript) on this also at the end of the first paragraph of the Implication section of the discussion.

Comment 4:

L 553-562 & L 565-579: Here you discuss one the one hand that present management practices affect the future and second the limitation that we do not know whether plants will be readily adapt to the root zone storages required in the altered climate. It would have been nice to have this connected a bit more, and given some idea on why the investigation is nevertheless useful. For example, it proposes an envelope of the hydrological response, which is based on the best of our knowledge. Furthermore, do you agree that the method can also be used to estimate the degree of adaptation in existing catchments? This would be one research outlook to improving the predictions.

Reply 4:

Thank you, we agree with this very good suggestion. We have added a sentence (lines 569-570 of the tracked change manuscript) to make it more explicit that indeed, despite the uncertainty, the study is useful as it quantifies an envelope of the hydrological response when we consider ecosystem adaptation in response to climate change and that the method can be used to estimate the potential adaptation of ecosystems in catchments.

Comment 5:

L 609-612: I believe this has been added in response to one of my comments. It now comes a bit out of context here, and I propose to erase it. Mentioning the constant potential evaporation in the methods section works well. Adding a comment in the Methods section that swapping omega does not affect the HRUs would be helpful for readers like me.

Reply 5:

Agreed, we have moved these sentences of the discussion to method section 4.4 (Hydrological change

evaluation).

Comment 6:

L 639-640: „very interesting“ is a bit vague. Can you be a bit more specific? Why would it be and do you have an intuition how the water fluxes may change with adaptive roots compared to the original HTESSSEL runs? Maybe also have a look at this publication in this context:

Stevens, D., Miranda, P. M. A., Orth, R., Boussetta, S., Balsamo, G., & Dutra, E. (2020). Sensitivity of surface fluxes in the ECMWF land surface model to the remotely sensed leaf area index and root distribution: Evaluation with tower flux data. *Atmosphere*, 11(12), 1–19.
<https://doi.org/10.3390/atmos11121362>

Reply 6:

Thank you for this suggestion and the provided reference. We have clarified that the implementation of a time-dynamic root-zone storage capacity in land surface model could improve the estimation of evaporative fluxes (and by extension also streamflow). However, quantifying how the water fluxes may change with adaptative root-zone storage capacity in the HTESSSEL runs would really require a dedicated and thorough analysis for the study area, which is outside the scope of the current work.

Comment 7:

L 650-651: „may allow us to estimate future ω or root-zone storage capacity values in a region where the future climate may resemble today's climate elsewhere“

I am admittedly not sure where this is going. Can you be more specific on the expected insights? Do you propose to test how omega relates to root storage in different climates? This would be very nice and eventually help to validate one of the major assumptions of the manuscript, which is that omega changes due to vegetation and not climate.

Finally, I would really like to read a short discussion on how changes in climate dynamics would affect omega independently of vegetation in a future climate?

Reply 7:

What we mean here is to develop data-based models/relations which relate either omega and/or the root-zone storage capacity in a large sample of catchments to relevant climatic, topographic, soil or land-use indicators. This would enable us to predict how omega or the root-zone storage capacity may change in response to changes in the combination of these various indicators. This could be a more robust, data-based method to estimate potential changes in root-zone storage capacity or omega values. We have clarified this in the revised manuscript.

In our study, we indeed assume that changes in omega are mainly related to changes in vegetation in a future climate. While we already mention in Section 6.2.2 “On the potential land use change scenarios” that omega values are likely a manifestation of multiple climatic, landscape and vegetation characteristics, we have now more strongly emphasized that climate variability, CO₂-vegetation feedbacks and/or anthropogenic modifications may also influence omega values (Berghuijs et al., 2017).

Comment 8:

Appendix S3: It is good to have all the model related details collected in this section. Very clear now.

It seems that some of the equations in Table S5 are potentially not correct? Should it not read dS_w/dt instead of S_w/dt ? This is repeated in several lines. Also in the line relating to E_R : I am not sure I saw a definition of S_u .

Reply 8:

Great to hear that it is now very clear to have all the information related to the model in the Appendix!

Thank you for pointing out the S_U which should indeed be replaced by S_R . However, the presence of S_w/dt is not incorrect, as explicitly mentioning the time step of the model in the equations ensures that the units of the stores S [mm] divided by the timestep [d] match the units of the fluxes [mm/d].

References:

Berghuijs, W.R., Larsen, J.R., Van Emmerik, T.H. and Woods, R.A., 2017. A global assessment of runoff sensitivity to changes in precipitation, potential evaporation, and other factors. Water Resources Research, 53(10), pp.8475-8486.

Reply to Patricia Saco (Referee #3)

Overall assessment by Patricia Saco:

I believe that the authors have carefully and thoroughly addressed all the reviewers comments. The revised manuscript has been greatly improved in the revision process.

Reply:

We would like to thank Patricia Saco for her very positive assessment of our revised manuscript.

Ecosystem adaptation to climate change: the sensitivity of hydrological predictions to time-dynamic model parameters

Laurène J. E. Bouaziz^{1,2}, Emma E. Aalbers^{3,4}, Albrecht H. Weerts^{2,5}, Mark Hegnauer², Hendrik Buiteveld⁶, Rita Lammersen⁶, Jasper Stam⁶, Eric Sprokkereef⁶, Hubert H. G. Savenije¹, and Markus Hrachowitz¹

¹Department of Water Management, Faculty of Civil Engineering and Geosciences, Delft University of Technology, P.O. Box 5048, NL-2600 GA Delft, The Netherlands

²Department Catchment and Urban Hydrology, Deltares, Boussinesqweg 1, 2629 HV Delft, The Netherlands

³Royal Netherlands Meteorological Institute (KNMI), P.O. Box 201, 3730 AE De Bilt, the Netherlands

⁴Institute for Environmental Studies (IVM), Vrije Universiteit, Amsterdam, 1081 HV, the Netherlands

⁵Hydrology and Quantitative Water Management Group, Wageningen University and Research, P.O. Box 47, 6700 AA Wageningen, The Netherlands

⁶Rijkswaterstaat, P.O. Box 2232, 3500 GE Utrecht, the Netherlands

Correspondence: Laurène Bouaziz (laurene.bouaziz@deltares.nl)

Abstract. Future hydrological behavior in a changing world is typically predicted based on models that are calibrated on past observations, disregarding that hydrological systems, and therefore model parameters, may change as well. In reality, hydrological systems experience almost continuous change over a wide spectrum of temporal and spatial scales. In particular, there is growing evidence that vegetation adapts to changing climatic conditions by adjusting its root-zone storage capacity, which is the key parameter of any terrestrial hydrological system. In addition, other species may become dominant, both under natural and anthropogenic influence. In this study, we test the sensitivity of hydrological model predictions to changes in vegetation parameters that reflect ecosystem adaptation to climate and potential land-use changes. We propose a top-down approach, which directly uses projected climate data to estimate how vegetation adapts its root-zone storage capacity at the catchment scale in response to changes in magnitude and seasonality of hydro-climatic variables. Additionally, long-term water balance characteristics of different dominant ecosystems are used to predict the hydrological behavior of potential future land-use change, in a space-for-time exchange. We hypothesize that changes in the predicted hydrological response as a result of 2K global warming are more pronounced when explicitly considering changes in the sub-surface system properties induced by vegetation adaptation to changing environmental conditions. We test our hypothesis in the Meuse basin in four scenarios designed to predict the hydrological response to 2K global warming in comparison to current-day conditions using a process-based hydrological model with (a) a stationary system, i.e. no assumed changes in the root-zone storage capacity of vegetation and historical land use, (b) an adapted root-zone storage capacity in response to a changing climate but with historical land use, and (c,d) an adapted root-zone storage capacity considering two hypothetical changes in land use. We found that the larger root-zone storage capacities (+34 %) in response to a more pronounced climatic seasonality with warmer summers under 2K global warming result in strong seasonal changes of the hydrological response, ~~with in the non-stationary scenarios~~. More specifically, streamflow and groundwater storage are up to -15 % and -10 % lower ~~streamflow and groundwater storage respectively in~~

~~autumn and an increase of summer evaporation of~~ in autumn respectively due to an up to +14 % higher summer evaporation in the non-stationary scenarios compared to the stationary benchmark scenario. By integrating a time-dynamic representation of changing vegetation properties in hydrological models, we make a potential step towards more reliable hydrological predictions under change.

25 1 Introduction

Hydrological models are required to provide robust short-term hydrological forecasts and long-term predictions of the impact of natural and human-induced change on the hydrological response. Common practice is to predict the future using a hydrological model calibrated to the past (Vaze et al., 2010; Blöschl and Montanari, 2010; Peel and Blöschl, 2011; Coron, 2013; Seibert and van Meerveld, 2016). For the near future, it seems acceptable to assume no fundamental change in the hydrological system, although we know that vegetation has the capacity to adapt to changing climate conditions at scales reaching from individual plants to the composition of entire plant communities at larger ecosystem scales (Guswa, 2008; Schymanski et al., 2008; Gentine et al., 2012; Harman and Troch, 2014; Savenije and Hrachowitz, 2017; Hrachowitz et al., 2021). For longer term predictions, it is therefore problematic to assume an unchanged system within a changing world. This raises the question on the robustness of hydrological predictions, especially in the context of climate change (Coron et al., 2012; Stephens et al., 2019).

For example, Merz et al. (2011) clearly shows the non-stationarity of hydrological model parameters when calibrating 273 Austrian catchments in subsequent 5-years periods between 1976 and 2006. Being the core parameter of any hydrological system, Merz et al. (2011) report almost a doubling of the root-zone storage capacity and this gradual increase is assumed to be related to changing climatic conditions, such as increased evaporation and drier conditions in the more recent years. The temporal variability of model parameters could also be attributed to uncertainties in input and model structure or inadequate calibration strategies. However, the observed trends in model parameters are also likely to reflect transient catchment conditions over the historical period.

Under continued global warming, precipitation and temperature extremes are expected to further increase and the hydrological cycle is likely to further accelerate (Allen et al., 2010; Kovats et al., 2014; Stephens et al., 2021). In addition, natural land cover change and anthropogenic activities of land-cover change and land-use management can substantially alter a catchment's water balance (Brown et al., 2005; Wagener, 2007; Fenicia et al., 2009; Jaramillo and Destouni, 2014; Nijzink et al., 2016a; Hrachowitz et al., 2021; Levia et al., 2020; Stephens et al., 2020, 2021). Considering the unprecedented speed of change, Milly et al. (2008) postulated that "stationarity is dead" and no longer should serve as a default assumption in water management. They advocate the development of methods that quantify the non-stationarity of relevant hydrological variables.

However, describing and quantifying non-stationarity is challenging due to the complex interactions and associated feedback between climate, vegetation, soils, ecosystems and humans at multiple spatial and temporal scales (Seibert and van Meerveld, 2016; Stephens et al., 2020). The main approaches to understand how changes in hydrological functioning relate to changes in catchment characteristics rely on paired watershed studies and hydrological modeling (Andréassian et al., 2003). In many

modeling studies, a selection of one or more parameters are changed using values from literature in combination with adapted
55 land-cover maps to (partly) reflect the characteristics of the altered system (Mao and Cherkauer, 2009; Buytaert and Beven,
2009; Pomeroy et al., 2012; Gao et al., 2015). Alternatively, Duethmann et al. (2020) uses satellite observations of vegetation
indices to improve the representation of the surface resistance dynamics to calculate reference evaporation used in conceptual
hydrological models over a historical record. A similar approach is applied by Fenicia et al. (2009) to account for changes in
evaporation as a result of land-use management changes in the Meuse basin.

60 While these approaches are valuable to test the sensitivity of the hydrological response to changes in catchments charac-
teristics (Seibert and van Meerveld, 2016), they require an understanding of how catchment characteristics (e.g. land use, soil
properties) relate to model parameters. Yet, there is considerable uncertainty in the a priori estimation of parameter values
through the use of look-up tables relating physical catchments properties to parameter values and the use of regionalization
approaches to transfer parameter values from one location to another (Wagener, 2007). Besides, the required data (e.g. future
65 land-use maps or vegetation indices) may not be available in the context of climate change impact assessment (Duethmann
et al., 2020). Instead, a way forward may be to develop robust top-down modeling approaches based on optimality principles of
vegetation growth by considering the co-evolution of soils, vegetation and climate in a holistic way (Schymanski et al., 2009;
Blöschl and Montanari, 2010).

As complex and heterogeneous as landscapes may be across a diversity of climates, the long-term hydrological partitioning
70 of a catchment is governed by a surprisingly simple and predictable relation, which relies on the available water and energy
for evaporation (Turc, 1954; Mezentsev, 1955; Budyko, 1961; Fu, 1981; Zhang et al., 2004). This is described in the Budyko
hypothesis, which suggests that the ratio of mean annual evaporation over precipitation (E_A/P) is mainly controlled by the
aridity index, defined as the ratio of mean annual potential evaporation over precipitation (E_P/P). However, Troch et al. (2013)
found catchments to deviate from the Budyko hypothesis when exchanging climates across different catchments in a modeling
75 experiment. Their results suggest that long-term hydrological partitioning results from the co-evolution of catchment properties
and climate characteristics, including not only the aridity index but also climate seasonality, topography, vegetation and soils.

The combination of these other factors influencing the water balance partitioning besides the aridity index are explicitly
considered in the ω parameter of the parametric description of the Budyko hypothesis (Fu, 1981; Zhang et al., 2004). Deviations
from the Budyko curve suggest that different vegetation develops in different climates, along a different ω curve. If climate
80 changes, catchments are likely not only to shift in the Budyko space as a result of a changing aridity index, but also as a result
of a changing vegetation cover (Jaramillo and Destouni, 2014). Other factors affect the positions and trajectories of catchments
in the Budyko space, including the complex and counteracting responses of ecosystems to elevated CO₂ levels (Jasechko,
2018). More specifically, vegetation density may increase from CO₂ fertilization, leading to increased transpiration, implying
an upward shift in the Budyko space. While higher water-use efficiencies may lead to declining transpiration rates, leading to a
85 downward shift in the Budyko space (Keenan et al., 2013; van der Velde et al., 2014; van Der Sleen et al., 2015; Ukkola et al.,
2016; Jaramillo et al., 2018; Stephens et al., 2020).

The fact that most catchments worldwide scatter closely around the analytical Budyko curve is evidence for the co-evolution
of vegetation and soils with climate at the catchment scale (Gentine et al., 2012; Donohue et al., 2012). Vegetation tends to

efficiently adapt its root-zone storage capacity to satisfy canopy water demand (Milly, 1994; Schymanski et al., 2008; Gerrits et al., 2009; Gentine et al., 2012; Gao et al., 2014; Yang et al., 2016). This implies that vegetation creates a larger buffer to survive dry spells when seasonal water supply and demand are out of phase, than in a climate where demand and supply are in phase. The root-zone storage capacity is, therefore, the key element regulating the partitioning of water fluxes in many terrestrial hydrological systems. In addition, not only natural changes to the environment, but also human interference with vegetation affect transpiration water demand and hence the root-zone storage capacity (Nijzink et al., 2016a; Hrachowitz et al., 2021).

Detailed observations of rooting-systems are very scarce in time and space and difficult to integrate to the catchment scale due to heterogeneity of landscapes (de Boer-Euser et al., 2016; Hrachowitz et al., 2021). Instead, the catchment-scale root-zone storage capacity is often estimated through calibration of a hydrological model. Other methods rely on optimality principles that maximize net primary production, carbon gain or transpiration (van Wijk and Bouten, 2001; Kleidon, 2004; Collins and Bras, 2007; Guswa, 2008; Speich et al., 2018). Alternatively, there is increasing evidence that the catchment-scale root-zone storage capacity can be robustly and directly estimated from annual water deficits using water-balance data (Gao et al., 2014; de Boer-Euser et al., 2016; Wang-Erlandsson et al., 2016; Nijzink et al., 2016a; Bouaziz et al., 2020; Hrachowitz et al., 2021). However, it remains unclear how vegetation may adapt its root-zone storage capacity to climate change and how these changes affect future hydrological behavior.

In contrast to current studies assuming stationary model parameters in a changing system, the objective of this study in the Meuse basin (Western Europe) is to test how sensitive hydrological predictions are when changing vegetation related parameters, thereby accounting for the adaption of vegetation to future climate conditions. We test the hypothesis that changes in the predicted hydrological response to a 2K global warming are more pronounced when explicitly considering time-dynamic values for the root-zone storage capacity parameter, which reflect vegetation adaptation to changes in hydro-climatic variables and potential land-use changes.

Using the Budyko framework, we first estimate changes in the long-term hydrological partitioning. To evaluate the effect of land-use change under potential future conditions, we exchange space-for-time by connecting the spatially variable ω parameter of the Budyko curve to different dominant land uses. We then use water-balance data to estimate how the root-zone storage capacity may adapt to increasing seasonal water deficits under climate change.

2 Study area

2.1 Climate and landscape

The Meuse basin upstream of Borgharen, at the border between Belgium and the Netherlands, covers an area of approximately 21300 km² with an elevation ranging between 50 and 700 m and can be divided into three main zones (Fig. 1). The French Southern part of the basin in the Grand Est region is characterized by relatively thick soils, broad valleys bottoms and gentle slopes underlain by sedimentary consolidated rocks. Metamorphic rocks and relatively thin impermeable soils dominate the

steeper Ardennes Massif in Belgium. On the West bank of the Meuse in Wallonia, the lithology is characterized by porous chalk layers with deep groundwater systems.

The Meuse is a rain-fed river with relatively short response times of several hours up to few days. Streamflow has a strong seasonality with low summer and high winter streamflow reflecting the seasonality of potential evaporation, while precipitation is relatively uniformly distributed throughout the year. The large storage capacity due to relatively thick soils in the French part of the basin increases the hydrological memory of the system, implying a strong influence of winter precipitation on streamflow deficits in the subsequent summer (de Wit et al., 2007). Snow is not a major component of the water balance, but snow melt can have a significant influence during some events (de Boer-Euser, 2017; Bouaziz et al., 2021). Mean annual precipitation, potential evaporation and streamflow is approximately 950 mm yr^{-1} , 590 mm yr^{-1} and 407 mm yr^{-1} , as derived from the historical observed climate and streamflow data (Sect. 3.1 and 3.3).

2.2 Land use

Land use in the basin consists of 35 % forest, 32 % agriculture, 21 % pasture and 9 % urban areas (Fig. 1c, European Environment Agency, 2018). The large majority of forests in the French part of the basin is characterized as "old growth", here defined as forested area which has been continuously wooded since at least the middle of the 19th century (Cateau et al., 2015). These broadleaved forests consist primarily of European Oaks, Sessile Oak and Beech (Institut National de l'Information Géographique et Forestière, 2019). In contrast, only 44 % of the 18th century Walloon forests of Belgium has remained from the original broadleaved forest, the rest being cleared for agriculture on high fertility soils in the North West (30 %) or converted to coniferous plantations (Scots pine, Norway spruce and Douglas-fir) on the poor soils of the Ardennes (26 %, Kervyn et al., 2018). The status of "old growth" forest does not exclude human disturbances, but assumes a relatively limited impact. Soils are less disturbed and their structure and biochemical composition have been preserved for several centuries. This favors a high degree of biodiversity, which is a key element for the resilience of forest ecosystems to perturbations (Kervyn et al., 2018). In contrast, recent short-rotation plantations lack many of these characteristics. Particularly thick canopy plantations, such as the spruce and Douglas-fir, significantly alter the typical biodiversity of forests. Additionally, relatively higher evaporation water use is expected in these recent, short-rotation exotic plantations in comparison to older, more natural forests (Fenicia et al., 2009; Stephens et al., 2021).

3 Data

3.1 Observed historical climate data

We use the European daily high-resolution gridded dataset, termed E-OBS (v20.0e), as observed historical climate data, which includes daily precipitation, temperature and radiation fields for the period 1980-2018 at a 25 km^2 resolution (Table 1, Cornes et al., 2018). The data are based on station data collated by the European Climate Assessment & Dataset (ECA&D) initiative. Temperature is downscaled using the digital elevation model and a fixed lapse rate of $0.0065^\circ\text{C m}^{-1}$. Potential evaporation is

estimated using the Makkink formula (Hooghart and Lablans, 1988). There is a relatively large underestimation of precipitation (> 20 %) in the E-OBS dataset in the center of the basin when compared to an operational dataset, which is based on local precipitation data provided by the Service Public de Wallonie for the period 2005-2017 and used in Bouaziz et al. (2020). A monthly bias-correction factor is applied to correct the E-OBS precipitation in the center of the basin to better represent the local precipitation data provided by the Service Public de Wallonie (Sect. S1 of the Supplement).

3.2 Simulated historical and 2K climate data

To study the impact of 2 Kelvin (2K) global warming on the hydrological response of the Meuse basin, we use climate simulations of precipitation, temperature and potential evaporation for the historical period 1980-2018 and a 2K global warming simulation (Table 1), provided by the Royal Netherlands Meteorological Institute (KNMI). The simulations are generated with the KNMI Regional Atmospheric Climate Model (KNMI-RACMO2) (van Meijgaard et al., 2008) at 12 km x 12 km resolution. RACMO2 uses as land surface scheme the Hydrology Tiled European Centre for Medium-Range Weather Forecasts (ECMWF) Scheme for Surface Exchanges over Land (HTESSEL) (Balsamo et al., 2009), which employs four soil layers with a total depth of 2.9 m. Each land-grid cell includes separate tiles for high and low vegetation (16 vegetation types), bare soil, snow and intercepted water, for which the energy and water balances are solved individually.

The historical simulation uses ECMWF Re-Analysis 5th Generation (ERA5) data (Hersbach et al., 2020) as initial- and lateral boundary conditions. The 2K simulation is a so-called pseudo-global warming simulation (e.g. Schär et al., 1996; Attema et al., 2014; Prein et al., 2017; Brogli et al., 2019), which is an alternative method to generate high-resolution climate change information. Instead of downscaling global climate model (GCM) projections, the historical period is re-simulated, but set against a warmer climate background by adding perturbations to the ERA5 initial- and boundary conditions. The perturbations represent the change in the mean climate state in a globally 2K warmer world, derived from a large initial condition GCM ensemble (Aalbers et al., 2018). The method minimizes biases in the mean climate state of the historical simulation, guaranties a realistic atmospheric circulation under both historical and ‘future’ conditions and increases the signal-to-noise ratio of the climate change response. A full description of the dataset is provided in Aalbers et al. (2021).

3.3 Streamflow

Streamflow data is available for 35 catchments nested within the Meuse basin upstream of Borgharen for the period 2005-2017 (Fig. 1, Service Public de Wallonie, 2018; Banque Hydro, 2018). The streamflow at Borgharen is a constructed time series which sums the observed streamflow of the Meuse at St Pieter and of the Albert Canal at Kanne to represent the total flow from the tributaries before part of it is extracted in the Albert Canal (de Wit et al., 2007).

4 Methods

In four scenarios, we evaluate the sensitivity of hydrological model predictions to ecosystem adaptation in response to climate and potential land use change by changing a key vegetation parameter, i.e. the root-zone storage capacity, in a process-based

model. Our methodology to estimate how vegetation adapts its root-zone storage capacity to changing climate and land-use conditions requires to estimate future transpiration. The following stepwise approach is designed: (1) we first estimate the long-term runoff coefficient (Q/P) in a 2K warmer world from movements in the Budyko space as a result of a shift in aridity index and a potential shift in dominant land-use; (2) we then use these estimates of projected future runoff coefficients to estimate future evaporative demand E_A by closing the water balance ($E_A/P = 1 - Q/P$). Based on these values of E_A and future projections of P we then apply a water balance approach to estimate how the root-zone storage capacity adapts to a changing climate. (3) Next, we calibrate a hydrological model using the observed historical E-OBS climate data to represent current-day hydrological conditions and (4) test if the use of the historical climate data simulated by the regional climate model leads to a plausible representation of current-day hydrological conditions. (5) In a next step, we run the hydrological model with the 2K climate data in four scenarios describing (a) a stationary system with historical root-zone storage capacity and historical land-use, (b) an adapted root-zone storage capacity in response to a changing climate but a historical land-use, (c,d) an adapted root-zone storage capacity and a shift in dominant land-use. (6) Finally, we compare the change in hydrological response between the 2K and historical conditions for these four scenarios. An overview of the procedure is summarized in Fig. 2.

4.1 Changing climate, vegetation and land use

4.1.1 Estimating future runoff coefficients

The long-term partitioning of precipitation (P) into evaporation (E_A) and streamflow (Q) is mainly controlled by the long-term aridity index (ratio of potential evaporation over precipitation, E_P/P), according to the Budyko hypothesis. To account for additional factors that influence the relation between aridity and the evaporative index E_A/P (and runoff coefficient $Q/P = 1 - E_A/P$), Fu (1981) introduces a parameter ω to encapsulate the combined influences of climate, soils, vegetation and topography (Eq. 1). It should be noted that throughout this manuscript, we use the term evaporation to represent all different evaporative fluxes (interception, soil evaporation and transpiration), following the terminology proposed by Savenije (2004) and Miralles et al. (2020).

$$\frac{E_A}{P} = 1 - \frac{Q}{P} = 1 + \frac{E_P}{P} - \left(1 + \left(\frac{E_P}{P}\right)^\omega\right)^{1/\omega} \quad (1)$$

We solve Eq. 1 to determine the value of ω for each of the 35 catchments of the Meuse basin for observed historical conditions for the period 2005 to 2017 (ω_{obs}), using the meteorological E-OBS data (P_{obs} and $E_{P,\text{obs}}$) and observed streamflow (Q_{obs}). Assuming only a change in long-term mean climate conditions, i.e. aridity index, a catchment will move along its ω_{obs} -parameterized curve from its original position p_1 to a new position p_2 due to the horizontal shift in aridity index $(E_P/P)_\Delta$ (Fig. 3a). Here, we use the simulated historical and 2K precipitation and potential evaporation climate data to determine how the change in potential evaporation ($\Delta E_P = E_{P,2K} - E_{P,\text{hist}}$) and precipitation ($\Delta P = P_{2K} - P_{\text{hist}}$) lead to a change in aridity

index (Eq. 2) and therefore in actual evaporation ($\Delta E_A = E_{A,2K} - E_{A,hist}$) and streamflow ($\Delta Q = Q_{2K} - Q_{obs}$), using the Budyko hypothesis in Eq. 1.

$$215 \quad \left(\frac{E_P}{P}\right)_{2K} = \frac{E_{P,obs} + \Delta E_P}{P_{obs} + \Delta P} \quad (2)$$

However, land cover and vegetation are likely to also change in response to a changing climate, introducing an additional shift toward a position p_3 on a different curve with ω_{change} (Fig. 3a). A downward shift from ω_{obs} to ω_{change} indicates less water use for evaporation, as opposed to an upward shift for higher evaporative water use. These shifts in ω values represent changes in drivers other than aridity index, including e.g. land cover, tree species, forest age, biomass growth and water use efficiency (Jaramillo et al., 2018).

To test the sensitivity of the hydrological response to a change in ω in addition to a change in aridity index, we consider two scenarios. The catchments with relatively high percentages of broadleaved forests (25-38% as in the French part of the basin) receive the ω values of catchments with relatively low percentages of broadleaved forests (1-12% as mainly in the Belgian Ardennes) and vice-versa (Fig. 1b). We denote $\omega_{broadleaved}$ for the catchments with relatively high percentages of broadleaved forests and $\omega_{coniferous}$ for the catchments with relatively low percentages of broadleaved forest, i.e. where broadleaved forests were largely converted to coniferous plantations or agriculture. These scenarios are meant as a sensitivity analysis in the spirit of trading space-for-time (Singh et al., 2011) to evaluate the effect of potential future land-use management on the overall water balance.

When converting broadleaved forest to coniferous plantations, we expect an increase in water use for evaporation and therefore an upward shift in ω values, as opposed to a downward shift when converting coniferous plantations to more natural broadleaved forests (Fenicia et al., 2009; Teuling et al., 2019; Stephens et al., 2021). The described movements in the Budyko space are used to estimate the projected long-term runoff coefficients ($(Q/P)_{2K}$, Eq. 3) as a result of, both, climate change but no changes in vegetation cover (ω_{obs}), and climate change in combination with changes in vegetation cover (by swapping $\omega_{broadleaved}$ values to $\omega_{coniferous}$ for a selection of catchments and vice-versa). The projected runoff coefficients are subsequently used to estimate the long-term projected 2K evaporation E_A by closing the water balance, which in turn allows us to estimate projected transpiration E_R and changes in the root-zone storage capacity parameter, as described in detail in Sect. 4.1.2.

$$\left(\frac{Q}{P}\right)_{2K,\omega} = \frac{Q_{obs} + \Delta Q}{P_{obs} + \Delta P} = - \left(\left(\frac{E_P}{P}\right)_{2K} - \left(1 + \left(\frac{E_P}{P}\right)_{2K}^\omega\right)^{1/\omega} \right) = 1 - \left(\frac{E_A}{P}\right)_{2K,\omega} \quad (3)$$

240

4.1.2 Estimating the root-zone storage capacity $S_{R,max}$

The root-zone storage capacity represents the maximum volume of water which can be held against gravity in pores of unsaturated soil and which is accessible to roots of vegetation for transpiration. It is a key element controlling the hydrological response of hydrological systems. The long-term partitioning of precipitation into streamflow and evaporation of catchments worldwide plots closely around the Budyko curve suggesting that vegetation has adapted its root-zone storage capacity to offset hydro-climatic seasonality, by creating a buffer large enough to overcome dry spells (Gentine et al., 2012; Donohue et al., 2012; Gao et al., 2014). This is the main assumption underlying the water-balance method to estimate the root-zone storage capacity at the catchment scale (Gao et al., 2014; Nijzink et al., 2016a; de Boer-Euser et al., 2016; Wang-Erlandsson et al., 2016; Bouaziz et al., 2020; Hrachowitz et al., 2021).

The water-balance method requires daily time series of precipitation, potential evaporation and a long-term runoff coefficient to estimate transpiration, which depletes the root-zone storage during seasonal dry periods. Annual water deficits ($S_{R,def}$) stored in the root-zone of vegetation to fulfill canopy water demand for transpiration are estimated on a daily time step as the cumulative sum of daily effective precipitation (P_E) minus transpiration (E_R).

First, effective precipitation, i.e. the amount of precipitation that reaches the soil after interception evaporation (E_I), is estimated by solving the water balance of a canopy storage (S_I) with maximum interception storage capacity (I_{max} , here taken as 2.0 mm), according to Eq. 4. A value of 2.0 mm for I_{max} was previously also used by de Boer-Euser et al. (2016) and Bouaziz et al. (2020), who also show a low sensitivity of the root-zone storage capacity to the value of I_{max} .

$$P_E(t) = P(t) - E_I(t) - \frac{dS_I(t)}{dt} \quad (4)$$

With daily interception evaporation $E_I(t) = \min(E_P(t), S_I(t)/dt)$ and daily effective precipitation $P_E(t) = \max(0, (S_I(t) - I_{max})/dt)$, both in mm d^{-1} .

Next, the long-term historical and projected “future” transpiration \bar{E}_R is estimated from the long-term water balance, using mean annual historical and projected “future” streamflow, estimated by movements in the Budyko space, and effective precipitation (\bar{Q} and \bar{P}_E , all in mm yr^{-1} , Eq. 5), assuming negligible changes in storage and intercatchment groundwater flows (catchments where $\bar{E}_A = \bar{E}_I + \bar{E}_R = \bar{P} - \bar{Q} < \bar{E}_P$).

$$\bar{E}_R \approx \bar{P}_E - \bar{Q} \quad (5)$$

The long-term transpiration \bar{E}_R is subsequently scaled to daily transpiration estimates E_R , using the daily signal of potential evaporation minus interception evaporation, according to Eq. 6 (Nijzink et al., 2016a; Bouaziz et al., 2020).

$$E_R(t) = (E_P(t) - E_I(t)) \cdot \frac{\bar{E}_R}{(\bar{E}_P - \bar{E}_I)} \quad (6)$$

The maximum annual storage deficits can then be derived from the cumulative difference of effective precipitation (P_E) and transpiration (E_R), assuming an "infinite" storage, according to Eq. 7 and illustrated in Fig. 3b. For each year, $S_{R,def}$ represents the amount of water accessible to the roots of vegetation for transpiration during a dry period. Storage deficits are assumed to be zero at the end of the wet period (T_0 , here April) and increase when transpiration exceeds effective precipitation during dry periods, until they become zero again (T_1) in the fall and winter, when excess precipitation is assumed to drain away as direct runoff or recharge.

$$S_{R,def}(t) = \min \int_{T_0}^{T_1} (P_E(t) - E_R(t)) dt \quad (7)$$

By fitting the extreme value distribution of Gumbel to the series of annual maximum storage deficits, the root-zone storage capacity at the catchment scale $S_{R,max}$ can be derived for various return periods. The Gumbel distribution was previously shown to be a suitable choice for the estimation of the root-zone storage capacity through the water-balance approach by several other studies (Gao et al., 2014; Nijzink et al., 2016a; de Boer-Euser et al., 2016; Bouaziz et al., 2020; Hrachowitz et al., 2021). These studies used a return period of 20 years for forested areas, suggesting that forests develop root systems to survive droughts with a return period of ~ 20 years. The root-zone storage capacity of cropland and grasslands is assumed to correspond to deficits with lower return periods of ~ 2 years (Wang-Erlandsson et al., 2016). It should be noted that the methodology assumes that vegetation taps its water from the unsaturated zone and not from the groundwater.

Using the above described methodology, we determine several sets of $S_{R,max}$ values for each of the 35 catchments of the Meuse basin to represent the historical and adapted root-zone storage capacity in response to a changing climate and changing/historical land-use, using historical climate observations (E-OBS) and the historical and 2K climate simulations. [The estimated adaptation of the root-zone storage capacity mainly relates to changes in the rooting system, i.e. rooting density and/or rooting depth.](#) Technical details for the calculation of the various $S_{R,max}$ are given in Sect. 2 of the Supplement.

290 – **$S_{R,max,A}$: Historical root-zone storage capacity**

The first set is $S_{R,max,A}$, which represents the historical meteorological and land-use conditions, derived from observed historical E-OBS data and observed streamflow data (P_{obs} , $E_{P,obs}$ and Q_{obs} for the period 2005-2017). $S_{R,max,A}$ is used as parameter for three model runs, each forced with a different dataset: historical E-OBS observations, simulated historical and 2K climate data (Sect. 4.4).

295 – **$S_{R,max,B}$: Adapted root-zone storage capacity in response to a changing climate**

We then estimate the root-zone storage capacity $S_{R,max,B}$ based on the 2K climate and historical land use to reflect adaptation of existing vegetation to changing climatic conditions such as differences in seasonality, aridity index (Eq. 2) and the resulting runoff coefficient (Eq. 3), but under the assumption that the vegetation cover remains unchanged.

– $S_{R,max,C}$ and $S_{R,max,D}$: **Adapted root-zone storage capacity in response to a changing climate and land use**

300 Subsequently, the root-zone storage capacity is estimated for the 2K climate under two land-use change scenarios, considering that if climate changes, a different vegetation cover might become dominant under natural and anthropogenic influence. In the first scenario, land use in the catchments with mainly coniferous plantations and agriculture (as mainly in the Belgian Ardennes, Sect. 2.2 and Fig. 1) is assumed to be converted to broadleaved forest. While in the second scenario, the broadleaved forests in the French part of the basin are converted to coniferous plantations and agriculture.

305 To simulate these land use changes, we first group the ω_{obs} parameter of the Budyko curves of the 35 catchments in three categories according to their areal fraction of broadleaved forest (Fig. 1b). Then, making use of a space-for-time exchange, we randomly sample from ω_{obs} values of one category to represent potential future changes in the other category. In other words, we use ω_{obs} values of catchments with high percentage of broadleaved forest to represent a potential conversion of land use to broadleaved forest in catchments with current-day low percentage of broadleaved forest ($S_{R,max,C}$) and vice-versa ($S_{R,max,D}$). These exchanged ω values are used to estimate the 2K runoff coefficient with Eq. 3. We repeat this random sampling ~~several~~ seven times, which results in ~~several parameter realizations of both~~ seven parameter realizations for each of $S_{R,max,C}$ and $S_{R,max,D}$. The sampling is performed because the variability in ω_{obs} values in each category is also influenced by other factors besides the dominant presence of broadleaved forest. In the subsequent analyses, we consider the ensemble of individual realizations to account for this uncertainty.

310

315 **4.2 Hydrological model**

The wflow_FLEX-Topo model (de Boer-Euser, 2017; Schellekens et al., 2020) is a fully distributed process-based model, which uses different flexible model structures for a selection of Hydrological Response Units (HRUs), delineated from the topography and the land use, to represent the spatial variability of hydrological processes. Here, we develop a model with three HRUs for wetlands, hillslopes and plateaus connected through their groundwater storage (model schematization and equations

320 in Sect. S3 of the Supplement; Savenije, 2010; de Boer-Euser, 2017). Thresholds of 5.9 m for the Height Above the Nearest Drainage (HAND Rennó et al., 2008) and 0.129 for slope are used to delineate the three HRUs, as suggested by Gharari et al. (2011), using the MERIT hydro dataset at ~ 60 m x 90 m resolution (Yamazaki et al., 2019). As additional step, given the high proportion of forest on hillslope and of agriculture on plateau, we here associated hillslope with forest and agriculture with plateau, using the CORINE land cover data (European Environment Agency, 2018). The areal fraction of each HRU are then

325 derived for each cell at the model resolution of 0.00833° (or ~ 600 m x 900 m).

The model is forced with precipitation, potential evaporation and temperature. Snow, interception, root-zone, fast and slow storage components are included in the model. Actual evaporation from the root-zone is linearly reduced when storage is below a certain threshold parameter. This standard formulation to express vegetation water stress (Sect. S3 of the Supplement) is used in many process-based models including NAM, HBV and VHM (Bouaziz et al., 2021). Streamflow is routed through the up-

330 scaled river network at the model resolution (Eilander et al., 2021) with the kinematic wave approach. Similar implementations of that model were previously successfully used in a wide variety of environments (e.g. Gharari et al., 2013; Gao et al., 2014; Nijzink et al.,

[\(e.g. Gharari et al., 2013; Gao et al., 2014; Nijzink et al., 2016b; de Boer-Euser, 2017; Hulsman et al., 2021; Bouaziz et al., 2021\)](#)

4.3 Model calibration and evaluation

335 4.3.1 Calibration and evaluation using the observed historical E-OBS climate data

The wflow_FLEX-Topo model is calibrated using streamflow at Borgharen and the observed historical E-OBS meteorological forcing data for the period 2007-2011, using 2005-2006 as warm-up years. The observed historical E-OBS dataset is used for calibration of the model as it is assumed to most closely represent current-day conditions. The parameter space is explored with a Monte Carlo strategy, sampling 10000 realizations from uniform prior parameter distributions (Sect. S4 of the Supplement).
340 The limited number of samples is due to the high computational resources required to run the distributed model. However, our aim is not to find the "optimal" parameter set, but rather to retain an ensemble of plausible parameter sets based on a multiobjective calibration strategy (Hulsman et al., 2019). To best reflect different aspects of the hydrograph, including high flows, low flows and medium-term partitioning of precipitation into drainage and evaporation, parameter sets are selected based on their ability to simultaneously and adequately represent four objective functions, including the Nash-Sutcliffe efficiencies of
345 streamflow, the logarithm of streamflow and, monthly runoff coefficients as well as the Kling-Gupta efficiency of streamflow. Only parameter sets that exceed a performance threshold of 0.9 for each metric are retained as feasible. The root-zone storage capacity parameter $S_{R,max,A}$ is a fixed parameter for each subcatchment, which is derived from annual maximum storage deficits with a return period of 2 years for the wetland and plateaus HRUs and 20 years for the hillslopes HRU (Sect. 4.1.2). Next, model performance is evaluated in the 2012-2017 post-calibration period using the same performance metrics, a visual
350 inspection of the hydrographs and the mean monthly streamflow regime. The performance metrics are also evaluated for the 34 remaining nested sub-catchments.

4.3.2 Evaluation using the simulated historical climate data

The performance of the calibrated model for the ensemble of retained parameter sets is also evaluated when the model is forced with the simulated historical climate data, using $S_{R,max,A}$ for the root-zone storage capacity parameter. This is the
355 reference historical run against which the relative effect of 2K global warming is evaluated for different scenarios to enable a fair comparison (Fig. 2 and Sect. 4.4). In addition, in Sect. S5 of the Supplement, we evaluate the performance of the calibrated model forced with the simulated historical climate data but with a root-zone storage capacity parameter derived directly from this data.

4.4 Hydrological change evaluation

360 We then force the calibrated wflow_FLEX-Topo model for the ensemble of retained parameter sets with the same 2K climate forcing data in four scenarios, each using a different root-zone storage capacity parameter to represent either stationary or adapted conditions in response to a changing climate and land use (Fig. 2 and Sect. 4.1.2). [Using the same climate forcing](#)

for each of the 2K scenarios implies that we do not change potential evaporation as a result of potential land-use changes. In addition, the percentages of each HRU in our model structure are not changed for the potential land-use change scenarios. This is because the top-down approach to estimate the effect of land-use change does not provide the level of detail required to specifically change land-use type at the pixel level.

The difference between the modelled historical hydrological response (1980-2018) and the hydrological responses predicted by each of the four model scenarios based on the 2K climate is evaluated in terms of runoff coefficient, evaporative index, annual statistics (runoff coefficient, evaporative index, mean, maximum, minimum 7-days streamflow and median annual sum of streamflow below the 90th percentile historical streamflow), and monthly patterns of flux and state variables (streamflow, evaporation, root-zone storage, groundwater storage) for a hypothetical 38-year period. The four scenarios, which each use $S_{R,max,A}$ as root-zone storage capacity in the historical run, as shown in Fig. 2, are listed below.

– **Scenario 2K_A: stationarity**

In scenario 2K_A, we assume an unchanged land use and that vegetation has not adapted its root-zone storage capacity to the aridity and seasonality of the 2K climate. This scenario implies stationarity of model parameters by using $S_{R,max,A}$ in both the historical and 2K runs, a common assumption of many climate change impact assessment studies (Booij, 2005; de Wit et al., 2007; Prudhomme et al., 2014; Hakala et al., 2019; Brunner et al., 2019; Gao et al., 2020; Rottler et al., 2020; Hanus et al., 2021). This is the benchmark scenario against which we compare the hydrological response considering non-stationarity of the system, as in the following three scenarios.

– **Scenario 2K_B: non-stationarity of the root-zone storage capacity in response to a changing climate**

In scenario 2K_B, we again assume an unchanged land use (ω_{obs}). However, we assume that vegetation has adapted its root-zone storage capacity to the aridity and seasonality of the 2K climate conditions by selecting $S_{R,max,B}$ as parameter for the 2K model run.

– **Scenario 2K_C: non-stationarity of the root-zone storage capacity in response to a changing climate and land-use conversion to broadleaved forest**

In scenario 2K_C, we use the root-zone storage capacity $S_{R,max,C}$ to reflect adaptation in response to the changing aridity index and seasonality of the 2K climate and changes in vegetation cover for the catchments located mainly in the Belgian Ardennes and dominated by coniferous plantation and agriculture to a land use of broadleaved forest as in the French part of the basin.

– **Scenario 2K_D: non-stationarity of the root-zone storage capacity in response to a changing climate and land-use conversion to coniferous plantation and agriculture**

In scenario 2K_D, the approach of scenario 2K_C is repeated. However, now the broadleaved forest in the French catchments are assumed to be converted to coniferous plantations or agriculture as in the Belgian Ardennes. The parameter $S_{R,max,D}$ is used in the model run forced with the 2K climate.

5.1 Adapted root-zone storage capacity $S_{R,max}$

5.1.1 Long-term water balance characteristics across catchments

In solving the parametric Budyko curve (Eq. 1) for the 35 catchments of the Meuse basin using historical E-OBS data and observed streamflow (Fig. 4a), we found that ω_{obs} values tend to be lower, with a median and standard deviation (subsequent values in next sections represent the same variables) of 2.43 ± 0.48 , for catchments with relatively high percentages of broadleaved forests (25-38 % as in the French part of the basin) as compared to catchments with relatively low percentages of broadleaved forests (1-12 % as in the Belgian part of the catchment) with ω_{obs} values of 3.04 ± 0.54 , as shown in Fig. 4b. Higher values of ω for a same aridity index indicate more water use for evaporation, which is likely related to the increased water use of relatively young coniferous plantations and agriculture as opposed to older broadleaved forests (Fenicia et al., 2009; Teuling et al., 2019).

5.1.2 $S_{R,max,A}$ Historical root-zone storage capacity

The root-zone storage capacity $S_{R,max,A}$ derived with observed historical E-OBS climate data and observed streamflow is estimated at values of 100 ± 17 mm and 169 ± 24 mm across all study catchments for 2 and 20 years return periods, respectively (Fig. 4c).

5.1.3 $S_{R,max,B}$ Adapted root-zone storage capacity in response to a changing climate

The adapted root-zone storage capacity $S_{R,max,B}$, in response to changing climate conditions and an unchanged land use, strongly increases with respect to historical conditions ($S_{R,max,A}$) with estimated values of 127 ± 18 mm (+27 %) and 226 ± 27 mm (+34 %) for return periods of 2 and 20 years, respectively (Fig. 4c). This strong increase is explained by larger storage deficits during summer due to an increase of about +10 % in summer potential evaporation in the 2K climate and, therefore, a more pronounced seasonality (Fig. 3b). In contrast, the change in aridity index between the historical and 2K climate simulations is relatively small with a median of +0.01 across all study catchments. This can be explained by a simultaneous increase in mean annual precipitation (+4 %) and potential evaporation (+7 %) on average over the basin area in the 2K climate compared to the simulated historical climate data. This increase in precipitation mostly occurs during the winter half year (Nov-Apr). In contrast, there is a relatively large variability in precipitation change in summer, characterized by years with wetter and drier summers.

5.1.4 $S_{R,max,C}$ Adapted root-zone storage capacity in response to a changing climate and land use conversion to broadleaved forest

The adapted root-zone storage capacity $S_{R,max,C}$, in response to changing climate conditions and a land use conversion from coniferous plantation and agriculture to broadleaved forest, results in estimated values of 123 ± 17 mm and 216 ± 27 mm for

425 return periods of 2 and 20 years, respectively (Fig. 4c). These values are similar to $S_{R,max,B}$, with a difference of about -4 %. This small decrease is in line with the expected reduced water use of broadleaved forests compared to coniferous plantations.

5.1.5 $S_{R,max,D}$ Adapted root-zone storage capacity in response to a changing climate and land use conversion to coniferous plantation and agriculture

In contrast, the root-zone storage capacity $S_{R,max,D}$, in response to changing climate conditions and a conversion of broadleaved
430 forest to coniferous plantation, result in estimated values of 137 ± 22 mm and 235 ± 35 mm for return periods of 2 and 20 years, respectively (Fig. 4c). This corresponds to an additional increase of +8 % and +4 % for both return periods in comparison with $S_{R,max,B}$, which does not consider additional land-use changes.

Therefore, the increase in root-zone storage capacity between the 2K and historical climate simulations can, here with +57 mm or +34 % for a return period of 20 years, be mostly attributed to a changing climate (aridity and seasonality) than to
435 changes in land use (-10 mm or -4 % for $S_{R,max,C}$ and +9 mm or +4 % for $S_{R,max,D}$). This indicates that with the assumed land-use change in scenarios 2K_C and 2K_D, the strong increase in water demand during summer as a result of a more pronounced seasonality has greater impact on the estimation of the root-zone storage capacity than a change in ω values. However, note that land use is changed in only part of the catchment for both land use change scenarios and that it is plausible to assume that more pronounced changes in land use will reinforce the observed effects.

440 5.2 Model evaluation (historical period)

5.2.1 Model forced with observed historical climate data

The modeled streamflow obtained using the ensemble of parameter sets retained as feasible after calibration mimics the observed hydrograph at Borgharen reasonably well for the evaluation period (Fig. 5a). Also the seasonal streamflow regime is well reproduced by the model, except for an underestimation of -9 % in the first half year (Fig. 5b). The four objective functions
445 show a relatively similar performance during calibration and evaluation with median values of 0.93 and 0.78 at Borgharen and for the ensemble of 34 additional not individually calibrated nested subcatchments of the Meuse, respectively (Fig. 6a,c).

5.2.2 Model forced with simulated historical climate data

When the calibrated model is instead forced with the simulated historical climate data, peaks are slightly overestimated in comparison to the model run forced with the observed historical E-OBS data (Fig. 5c). This is due to the on average +9 %
450 overestimation of precipitation in the simulated historical climate data compared to the observed historical E-OBS climate data. This precipitation overestimation results in an overestimation of about +12 % of modeled mean monthly streamflow during the wet December and January months (Fig. 5d). However, flows in spring and early summer are better captured when using the simulated historical climate data instead of historical E-OBS data (Fig. 5d). Overall, the streamflow model performance at Borgharen slightly decreases when the simulated historical climate data is used instead of E-OBS, but median values across
455 the ensemble of feasible parameter sets are still above 0.77 for each of the objective functions (Fig. 6b). Although a decrease in

model performance is found in a few nested catchments, the performance in the ensemble of nested catchments of the Meuse remains at median values of around 0.67 (Fig. 6d). The results of the model run forced with the simulated historical data and with the root-zone storage capacity parameter derived directly from this data show a relatively similar behavior, as further detailed in Sect. S5 of the Supplement.

460 The calibrated model forced with the simulated historical climate data shows a plausible behavior with respect to observed streamflow and is also reasonably close to the performance achieved with the observed historical E-OBS climate data. This is important because the effect of the 2K climate on the hydrological response is evaluated with respect to the model run forced with the simulated historical climate data, as they are both generated with the regional climate model. Therefore, the relatively high model performance in the evaluation period enables us to use the retained parameter sets from the calibration with E-OBS
465 data for the subsequent analyses with the simulated historical and 2K climate data.

5.3 Hydrological change evaluation (2K warmer climate)

5.3.1 Scenario 2K_A: stationarity

In the 2K_A scenario, representing a stationary system with identical parameters in the historical and 2K climate, runoff coefficients are projected to increase with a median of +3 %, the evaporative index (E_A/P) decreases with a median of -2 % and
470 mean annual streamflow increases with a median of +7 %. Maximum annual streamflow is also projected to increase with a median of about +5 %, while the median change in annual minimum of 7-days mean streamflow remains close to zero. The median annual sum of streamflow below the 90th percentile historical streamflow increases with +10 %, as shown in Fig. 7.

From the intra-annual perspective, streamflow is projected to increase from December until August with +8 % on average and decreases between September and November with -7 % (Fig. 8a). In the months where evaporation demand exceeds
475 precipitation, the root-zone soil moisture decreases, with a maximum of -22 % in September (Fig. 8b). Actual evaporation increases throughout the year with around +3 % on average except in July and August (-4 %) when the availability of water in the root-zone of vegetation is not sufficient to supply canopy water demand (Fig. 8c). The groundwater storage increases with approximately +5 % in all months except November, as shown in Fig. 8d.

5.3.2 Scenario 2K_B: non-stationarity of the root-zone storage capacity in response to a changing climate

480 Changes are substantially different in the 2K_B scenario which considers that the root-zone storage capacity of vegetation has adapted to the change in aridity and seasonality of the 2K climate. In contrast to scenario 2K_A, runoff coefficients are projected to decrease with a median of -2 %, while the evaporative index increases with a median of +2 % and the median change of mean annual streamflow is close to zero (Fig. 7). Similarly, the median change of, both, annual maximum streamflow and minimum
485 7-days mean streamflow remain close to zero. However, there is a substantial increase of +38 % in the median annual sum of streamflow below the 90th percentile historical streamflow. This result suggests that while the minimum streamflow remains relatively similar, the length of the low flow period strongly increases when considering an adaptation of the root-zone storage capacity to the 2K climate (Fig. 7).

Seasonal changes indicate a decrease in streamflow of -19% between September and November, which is longer and considerably more pronounced than in the $2K_A$ scenario (Fig. 8a). The root-zone soil moisture increases throughout the year with an average of +34 % due to the larger root-zone storage capacities (Fig. 8b). Actual evaporation is no longer reduced as a result of moisture stress in the root-zone that has adapted to a changing climate and strongly increases with on average approximately +7 % from May to October to supply canopy water demand (Fig. 8c). However, the considerable increase in spring and summer evaporation strongly reduces the groundwater storage with -5 % from October to February (Fig. 8d).

495 **5.3.3 Scenario $2K_C$: non-stationarity of the root-zone storage capacity in response to a changing climate and land-use conversion to broadleaved forest**

The predicted hydrological response in the $2K_C$ scenario is very similar to the response of the $2K_B$ scenario, despite considering additional changes in the root-zone storage capacity as a result of a land-use conversion from coniferous plantations and agriculture to broadleaved forest (Figs. 7 and 8). This is in line with the limited differences in root-zone storage capacities of around only +4 % between both scenarios (Sect. 5.1).

500 **5.3.4 Scenario $2K_D$: non-stationarity of the root-zone storage capacity in response to a changing climate and land-use conversion to coniferous plantation and agriculture**

In contrast, the change in hydrological response is most pronounced for the scenario $S_{R,max,D}$, which considers land use conversion of the broadleaved forests in the French part of the basin to coniferous plantations and agriculture (Figs. 7 and 8). Runoff coefficients decrease with a median of -4 %, while the evaporative index increases with a median value of +4 % and mean annual streamflow decreases with a median of -2 %, as opposed to a +5 % increase in the stationary $2K_A$ scenario (Fig. 7a,b,c). If the median change in streamflow extremes remains relatively close to zero, there is a strong increase of +54 % in the median annual sum of streamflow below the 90th percentile historical streamflow, suggesting a strong increase in the length of the low flow period (Fig. 7d,e,f).

Streamflow decreases from August to January with an average of -23 % and evaporation strongly increases from May to October with an average of +9 %. This increased evaporation during summer further reduces the groundwater storage from October to February with -7 % (Fig. 8). In comparison with the hydrological response of scenario $2K_B$, the additional land-use conversion in scenario $2K_D$ results in relatively similar patterns of change but with an additional +2 % increase in evaporation, -2 % decrease in streamflow and -2 % decrease in groundwater storage on average throughout the year.

5.3.5 Stationary versus adaptive ecosystems

515 There is a difference of up to -7 % in the change of mean annual streamflow between the stationary scenario $2K_A$ and the scenarios $2K_B$, $2K_C$, $2K_D$ with adaptive ecosystems (Fig. 7c). Additionally, the scenarios with adaptive ecosystems show a substantially more pronounced decrease in streamflow from September to January and a delay in the occurrence of the lowest streamflow from September to October (Fig. 8a). Associated to these seasonal differences in streamflow, the scenarios with

520 adaptive ecosystems are characterized by summer evaporation that is up to +14 % higher and a mean annual evaporation that is around +4 % higher than in the stationary scenario (Fig. 8c). Instead of a year-round increase in groundwater storage in the 2K_A scenario, there is a decrease in winter groundwater storage in the three other scenarios, resulting in a mean annual difference of -6 % between the scenarios with ecosystem adaption and the stationary scenario 2K_A, expressed as constant model parameters (Fig. 8d). Overall these results suggest that the hydrological response in the 2K climate of the stationary scenario 2K_A is substantially different from the responses of the three scenarios 2K_B, 2K_C, 2K_D, which consider a change in
525 the root-zone storage capacity to reflect ecosystem adaptation in response to climate change.

6 Discussion

6.1 Implications

Current practice in many climate change assessment studies is to assume unchanged system properties in the future (benchmark scenario 2K_A in our analysis), thereby neglecting potential changes in the system, such as adaption of vegetation to local
530 climate conditions. In addition to this scenario, we suggest a possible approach to consider ecosystem adaptation in response to climate change and test the sensitivity in the resulting hydrological response. Our analysis is, therefore, a first step in evaluating what may happen if we consider ecosystem adaptation in response to climate change in hydrological model predictions. The hydrological response under 2K global warming compared to historical conditions shows distinct patterns of change if we explicitly consider the non-stationarity of climate-vegetation interactions in a process-based hydrological model. More
535 specifically, in the non-stationary scenarios, there is an up to -15 % and -10 % stronger decrease in streamflow and groundwater storage respectively after summer due to an up to +14 % stronger increase in actual evaporation during the warmer summers, compared to the stationary scenario. These results suggest a substantially different hydrological response between the stationary and the non-stationary scenarios. However, the additional land-use conversion has relatively limited effect within the non-stationary scenarios, which could be attributed to the limited areal fraction of land-use conversion.

540 Our method is based on readily available data and is therefore widely applicable. The choice of hydrological model is open as root-zone storage capacity estimates derived from the water-balance approach are applicable in various hydrological and land surface models, provided that they include a root-zone parameterization, which is the case for most models (Clark et al., 2008; Nijzink et al., 2016a; Bouaziz et al., 2021; van Oorschot et al., 2021). The water-balance approach to estimate the root-zone storage capacity has previously successfully been applied in a variety of climate zones and across various ecosystems
545 (Donohue et al., 2012; Gentine et al., 2012; Gao et al., 2014; Wang-Erlandsson et al., 2016; de Boer-Euser et al., 2016; Singh et al., 2020; Hrachowitz et al., 2021; Dralle et al., 2021; McCormick et al., 2021). Here, the estimated values of the root-zone storage capacity for the different scenarios have median values below 250 mm for a return period of 20 years, which is within the range of global root-zone storage capacity values estimated by Wang-Erlandsson et al. (2016).

A time-dynamic parameterization of the root-zone storage capacity in the context of deforestation was previously introduced
550 by Nijzink et al. (2016a) and Hrachowitz et al. (2021), who demonstrated that an observed 20 % increase of the runoff coefficient in a catchment can to a large part be attributed to the > 20 % reduction of root zone storage capacity after deforestation.

Speich et al. (2020) implemented a time-dynamic root-zone storage capacity in the context of climate change. In their study, forest growth in response to climate change leads to a six times higher reduction of streamflow if a dynamic representation of, both, the Leaf Area Index and the root-zone storage capacity is implemented as opposed to a study in which only the Leaf Area Index varies (Schattan et al., 2013). Although these results are more pronounced than our findings, they point into the same direction of change. As the future is unknown, we cannot test our results against observations, but given the changes in temperature and precipitation, the future predicted hydrological response does not seem implausible. While Speich et al. (2020) combine a forest landscape model with a hydrological model to simultaneously represent the spatio-temporal forest and water balance dynamics, we rely on a simpler approach of movements in the Budyko framework to include potential land-use change.

The impact of climate change on low flows in the Meuse basin has been previously studied by de Wit et al. (2007). Using simulations from regional climate models which project wetter winters and drier summers, they question if the increase in winter precipitation reduces the occurrence of summer low flows due to an increase in groundwater recharge. However, they were unable to address this question with their model due to its poor low-flow performance. Our results indicate an increase in groundwater storage during winter in the stationary scenario, as opposed to a decrease for the models with a time dynamic root-zone storage capacity, as a result of an increased water demand for evaporation during summer. In comparing several scenarios for the root-zone storage capacity parameter, we include some form of system representation uncertainty, which improves our understanding in the modeled changes by placing them in a broader context (Blöschl and Montanari, 2010). Despite considerable uncertainty on the capacity of vegetation to adapt (Sect. 6.2.1), we quantify with the best of our knowledge an envelope of the hydrological response by considering the potential degree of ecosystem adaptation to climate change.

The conversion to native species may increase biodiversity and the resilience of ecosystems to climate change (Schelhaas et al., 2003; Klingen, 2017; Levia et al., 2020). However, these processes are slow, implying that current management practices shape the forests of decades and centuries to come in an uncertain future. Increasing our understanding on how to include vegetation changes in hydrological models to reliably quantify their impact is a way forward in the development of strategies to mitigate the adverse effects of climate change.

6.2 Limitations and knowledge gaps

6.2.1 On the estimation of adapted root-zone storage capacities

The proposed methodology to estimate future root-zone storage capacities relies on the underlying assumption that past empirical relations between aridity index and evaporative index (i.e. the Budyko framework) still apply in the future. While the Budyko framework is a well-established concept, recent studies by Berghuijs et al. (2020) and Reaver et al. (2020) show that it should be cautiously applied in changing systems. The Budyko framework reflects the long-term hydrological partitioning under dynamic equilibrium conditions. Therefore, when using the Budyko framework to estimate the future rate of transpiration, we assume that the future vegetation has adapted to the future climatic conditions and that it is in a state of dynamic equilibrium. The uncertainty of this assumption is considerable because it implies that vegetation has had the time to adapt

585 to the rapidly changing environmental conditions. There is increasing evidence that vegetation will eventually adapt to ensure sufficient access to water (Guswa, 2008; Schymanski et al., 2008; Gentine et al., 2012); otherwise, we would not see the hydrological partitioning of catchments around the world broadly plotting along the Budyko curve (Troch et al., 2013). Yet, considering the unprecedented scale and rate of change (Gleeson et al., 2020), it is unclear how ecosystems will cope with climate change, also considering the impact of potentially increasing storm intensities and frequencies, heatwaves, fires and
590 biotic infestations as a result of water stress on forest ecosystems (Lebourgeois and Mérian, 2011; Allen et al., 2010; Latte et al., 2017; Stephens et al., 2021). In particular, there is considerable uncertainty on how long it will take for vegetation to adapt and how it will adapt, assuming thresholds of adaptability are not exceeded.

In this analysis, we did not explicitly consider that vegetation can adapt to drier conditions by regulating its stomata and hence reduce transpiration. Furthermore, there is evidence that increasing atmospheric CO₂ concentrations may increase both,
595 water use efficiency but also increase green foliage due to fertilization effects (Keenan et al., 2013; Donohue et al., 2013; van der Velde et al., 2014; Frank et al., 2015; van Der Sleen et al., 2015; Ukkola et al., 2016; Jaramillo et al., 2018; Yang et al., 2019; Stephens et al., 2020). While the adaptation strategies of individual plants, e.g. root biomass adjustments, anatomical alterations and physiological acclimatisation, depend on vegetation type and species (Brunner et al., 2015; Zhang et al., 2020), here, we determine effective values of the root-zone storage capacity at the catchment scale to reflect the adaptation of the whole
600 ecosystem. Yet, our ability to predict what will happen is limited, but we can at least test the sensitivity of the hydrological response to changes in the system representation.

Once future transpiration rates are estimated, also the water-balance approach to estimate the root-zone storage capacity is subject to limitations. The method is not suitable in areas where the water table is very close to the surface and where vegetation directly can tap from the available groundwater instead of creating a buffer capacity (e.g Fan et al., 2017). Another limitation
605 of the water-balance approach relates to Eq. 6, in which we scale the daily transpiration estimates with a constant factor to the patterns of potential evaporation minus interception evaporation, implying that vegetation can extract water for transpiration from dry soils as easily as from wet soils.

6.2.2 On the potential land-use change scenarios

We quantify the changes in the hydrological response as a result of a changing climate in combination with several land use
610 scenarios (historical, conversion of broadleaved forests to coniferous plantations and agriculture and vice-versa). The relatively limited additional effects of land-use change on the hydrological response should be understood in the context of the relatively limited areal fraction under potential land-use conversion.

The land-use changes are integrated in the root-zone storage capacity as single parameter. However, climate and land use changes likely affect other aspects of catchment functioning, e.g. interception, infiltration and runoff generation (Seibert and
615 van Meerveld, 2016). Changes in the maximum interception storage capacity (Calder et al., 2003) are not explicitly considered in the estimation of the adapted root-zone storage capacities in the land-use change scenarios, as the impact was shown to be relatively minor in Bouaziz et al. (2020). Additional effects of soil compaction and artificial drainage on peak flows as a result of potential future land conversion (Buytaert and Beven, 2009; Seibert and van Meerveld, 2016) are difficult to quantify but

may partly be captured in the changed ω values. As mutual interactions between parameters remain problematic to quantify,
620 we use an ensemble of parameter sets to somehow account for the uncertainty in model parameters and the possibility that
parameters compensate for each other due to simplistic process representation.

~~In the potential land-use change scenarios, we did not change the percentages of each HRU in our model structure. The
top-down approach to estimate the effect of land-use change does not provide the level of detail required to specifically change
land-use type at the pixel level. In addition, the same climate forcing was used in each of the 2K scenarios, implying that we
625 did not change potential evaporation as a result of potential land-use changes.~~

While ω -values are likely a manifestation of multiple climatic, landscape and vegetation characteristics, and may change
due to climatic variability, CO₂-vegetation feedbacks and/or anthropogenic modifications (Berghuijs et al., 2017), we assumed
that vegetation type is the dominant control to explain differences in ω -values. As this parameter describes the hydrological
partitioning and because transpiration is a major continental flux (Jasechko, 2018; Teuling et al., 2019), it is reasonable to
630 assume that the variability in ω -values is largely controlled by the water volume accessible to the roots of vegetation for
transpiration (i.e. $S_{R,max}$). This root-accessible water volume is independent from the soil type, as root systems will develop
in a way to ensure sufficient access to water (Kleidon and Heimann, 1998; Guswa, 2008). In clayey soils, the rooting depth
might be shallower than in sandy soils for an identical root-zone storage capacity (de Boer-Euser et al., 2016). Topography,
geology and soil type are likely implicitly integrated in other model parameters, e.g. the various recession time-scales of the
635 linear reservoirs, which represent subsurface flow resistance throughout the system.

We also did not consider how the relatively high ω values may be related to intercatchment groundwater losses (Bouaziz
et al., 2018). Note that as our analyses should be understood in the context of a sensitivity analysis of the impact of potential
additional vertical shifts in the Budyko space as a result of a changing land use (Fig. 3), the potential effects on groundwater
losses on the results are likely to be minor.

640 **6.2.3 On the calibration**

Lastly, we performed a limited calibration of the hydrological model to retain an ensemble of plausible solutions and only used
a single climate simulation despite the uncertainty in initial and boundary conditions of regional climate models. Our analyses
are intended as a sensitivity analysis to quantify the effect of non-stationarity of the root-zone storage capacity parameter on
hydrological model predictions through optimal use of projected climate data, rather than a comprehensive climate change
645 impact assessment of the Meuse basin.

6.3 Outlook

6.3.1 Applications in land surface and ecohydrological models

The land surface scheme HTESSEL, that is used in the regional climate model RACMO2 to generate the historical and 2K cli-
mate simulations, assumes, as most land surface models, a fixed root-zone storage capacity over time. Ideally, this discrepancy
650 between the land surface model and the non-stationary hydrological models could be reduced by updating the adapted root-zone

storage capacity from one model to the other in several iteration steps, thereby including soil moisture - atmosphere feedback mechanisms. The implementation of the root-zone storage capacity parameter estimated from water-balance data in land surface models is, however, not trivial and requires further research (van Oorschot et al., 2021). Accounting for this climate control on root development and root-zone parameterization in ecohydrological ~~model could potentially also be very interesting~~ models could potentially improve the estimation of evaporative fluxes (Stevens et al., 2020; van Oorschot et al., 2021), as rooting depth and root distributions estimates are often estimated from fixed look-up tables (Tietjen et al., 2017; Schaphoff et al., 2018).

6.3.2 Climate analogues or trading space-for-time

The concept of trading space-for-time, which uses space as a proxy for time (Singh et al., 2011) could be further explored by selecting regions outside the Meuse basin with a current climate similar to the projected climate. This approach is also commonly referred to as climate-analogue mapping, i.e. statistical techniques to quantify the similarity between the future climate of a given location and the current climate of another location (Rohat et al., 2018; Bastin et al., 2019; Fitzpatrick and Dunn, 2019). This climate matching could be applied over distant regions, using datasets which combine landscape and climatological data over large samples of catchments (e.g. the various Catchment Attributes and Meteorology for Large-sample Studies (CAMELS) datasets, Addor et al., 2017; Klingler et al., 2021). Despite considerable uncertainties, finding a climate analogue for future projections in present conditions, may allow us to estimate future ω or root-zone storage capacity values in a region where the future climate may resemble today's climate elsewhere. This data-driven approach could potentially lead to more robust estimates of future ω or root-zone storage capacity values in response to changes in various climate or landscape indicators. These methods are intuitive but not straightforward, as they rely on the selection and combination of relevant climate variables and their relation with vegetation, despite non-linear vegetation responses to climate change (Reu et al., 2014).

670 7 Conclusions

Understanding non-stationarity of hydrological systems under climate and environmental changes has been recognized as a major challenge in hydrology (Blöschl et al., 2019). Despite our strong awareness of non-stationarity of hydrological parameters, we often lack knowledge to implement system changes in hydrological models. In this study in the Meuse basin, we introduce a method to estimate future changes in vegetation model parameters and we evaluate the sensitivity of hydrological model predictions to these changes in vegetation parameters as a results of ecosystem adaptation in response to climate and land-use changes. Our top-down approach uses readily available regional climate model simulations to allow a time-dynamic representation of the root-zone storage capacity, which is a key parameter in process-based hydrological models. Our approach relies, on the one hand, on a space-for-time exchange of Budyko characteristics of dominant land-use types to estimate the hydrological behavior of potential land-use changes and, on the other hand, on the interplay between the long-term and seasonal water budgets to represent climate-vegetation interactions under climate and land-use change. Despite knowledge gaps on future ecosystem water use, we implement potential system changes in a hydrological model based on our current understanding of hydrological systems. The predicted hydrological response to 2K warming is strongly altered if we consider that

vegetation has adapted by increasing its root-zone storage capacity ~~with~~by 34 % to offset the more pronounced hydro-climatic seasonality under 2K global warming compared to a stationary system. The increased vegetation water demand under global
685 warming results on average annually in -7 % less streamflow, +4 % more evaporation and -6 % less groundwater storage for the scenarios assuming non-stationary conditions compared to a stationary system. These differences even imply decreasing instead of increasing median annual streamflow under 2K global warming under non-stationary conditions. More importantly, the seasonal changes are considerable with up to +14 % higher evaporation in summer and up to -15 % and -10 % lower stream-
690 flow and groundwater storage respectively in autumn, for the non-stationary scenarios. Our study contributes to the quest for more plausible representations of catchment properties under change and, therefore, more reliable long-term hydrological predictions.

Table 1. Mean annual precipitation P , potential evaporation E_P , temperature T and aridity index E_P/P for the observed historical E-OBS data and the simulated historical and 2K climate data for the Meuse basin upstream of Borgharen.

	Observed historical climate	Simulated historical climate	Simulated 2K climate
P (mm yr ⁻¹)	950	1035	1070
E_P (mm yr ⁻¹)	590	670	720
T (°C)	9.6	8.9	10.9
E_P/P (-)	0.62	0.64	0.67





Figure 1. (a) Location of the Meuse basin in North-West Europe. (b) Elevation in the basin and categorization of catchments according to their areal percentage of broadleaved forest. Black dots indicate the locations where streamflow observations are available. (c) Main land-use types according to CORINE Land Cover (European Environment Agency, 2018). The three main zones are the French Southern part of the basin in the Grand Est region, the Ardennes and the area on the West bank of the Meuse.

Estimate historical and adapted root-zone storage capacity ($S_{R,max}$)


(1) Long-term runoff coefficient Q/P :
Use the Budyko framework to estimate Q/P under climate and potential land-use change



Output:

$$\begin{aligned} & Q_{obs}/P_{obs} \\ & (Q/P)_{2K,B} \\ & (Q/P)_{2K,C} \\ & (Q/P)_{2K,D} \end{aligned}$$

(2) Root-zone storage capacity $S_{R,max}$:
Use $Q/P = 1 - E_A/P$ to quantify E_A . Next, use E_A to estimate historical and adapted $S_{R,max}$ from the water-balance method



Output:

$$\begin{aligned} & S_{R,max,A} \text{ (hist.)} \\ & S_{R,max,B} \\ & S_{R,max,C} \\ & S_{R,max,D} \end{aligned}$$

Hydrological modeling

(3) Calibration:
Calibrate the hydrological model with observed historical data (E-OBS) and $S_{R,max,A}$ (as fixed calibration parameter)

(4) Evaluation:
Evaluate the calibrated hydrological model with simulated historical climate data, and $S_{R,max,A}$

(5) Runs:
Run the calibrated model with simulated 2K climate data and each of $S_{R,max,A}$, $S_{R,max,B}$, $S_{R,max,C}$ and $S_{R,max,D}$ for the 4 scenarios.

Hydrological change evaluation

(6) Compare sensitivity of hydrological response:
Compare change in hydrological response between the 2K and historical conditions for 4 scenarios.

Four scenarios




Figure 2. Overview of the methodological procedure and schematic representation of the four model scenarios.




Figure 3. (a) Representation of the Budyko space, which shows the evaporative index ($E_A/P = 1 - Q/P$) as a function of the aridity index (E_P/P) and the water and energy limit. A catchment with aridity index $(E_P/P)_{obs}$ and evaporative index $(E_A/P)_{obs}$, which is derived from observed historical data assuming $(E_A/P)_{obs} = 1 - (Q/P)_{obs}$, plots at location p_1 on the parametric Budyko curve with ω_{obs} . A movement in the Budyko space towards p_2 along the ω_{obs} curve is shown as a result of a change in aridity index $(E_P/P)_{\Delta}$ towards a projected $(E_A/P)_{2K, \omega_{obs}}$ associated with aridity $(E_P/P)_{2K}$. An additional shift from the ω_{obs} curve towards a location p_3 on a ω_{change} curve is assumed if additional factors (e.g. land use) are projected to change besides aridity index. Here, the represented downward shift in ω reduces the change in evaporative index to $(E_A/P)_{2K, \omega_{change}}$. (b) Cumulative storage deficits ($S_{R, def}$) derived from effective precipitation (P_E) and transpiration (E_R) using the simulated historical and 2K climate data. Estimates of transpiration (E_R) are derived from long-term water balance projections as a result of movements within the Budyko framework in response to climate and potential land use changes.




Figure 4. (a) Budyko space with parametric ω_{obs} curves for each of the 35 catchments of the Meuse basin. The dashed curves represent the median ω_{obs} curves for each category of areal broadleaved forest percentage. The change in median aridity index across catchments of the three categories from historical to 2K climate conditions is shown by the circles and triangles. (b) Parameterized ω_{obs} values for each of the 35 catchments of the Meuse basin, categorized according to their percentage of broadleaved forest. (c) Range of root-zone storage capacities across the 35 catchments of the Meuse basin for the four scenarios. $S_{R,max,A}$ represents the root-zone storage capacity for historical conditions. $S_{R,max,B}$ represents an adapted root-zone storage capacity in response to the 2K climate but no land use change. In the estimation of $S_{R,max,C}$, catchments with a low percentage of broadleaved forest (1-12%) receive ω values sampled from catchments with a high percentage of broadleaved forest (25-38%), to represent changes in land use towards a conversion to broadleaved forest. The same but reversed approach is applied for the estimation of $S_{R,max,D}$.




Figure 5. Observed and modeled hydrographs and mean monthly streamflow at Borgharen for the ensemble of parameter sets retained as feasible after calibration when the model is: **(a,b)** forced with E-OBS historical data and using $S_{R,max,A}$ as model parameter, and **(c,d)** forced with the simulated historical climate data using $S_{R,max,A}$ as model parameter.




Figure 6. Streamflow model performance during calibration and evaluation for the four objective functions when the model is forced with (a,c) observed historical E-OBS data and (b,d) simulated historical climate data at (a,b) Borgharen and (c,d) for the ensemble of nested catchments in the Meuse basin. The four objective functions are the Nash-Sutcliffe efficiencies of streamflow, logarithm of streamflow and monthly runoff coefficient ($E_{NS,Q}$, $E_{NS,\log Q}$, $E_{NS,RC}$) as well as the Kling-Gupta efficiency of streamflow ($E_{KG,Q}$). Note the different y-axis scales between rows.




Figure 7. Percentage change in annual hydrological response indicators between the 2K and historical model runs for the four scenarios, each based on different assumptions for the root-zone storage capacity parameter $S_{R,max}$. Percentage change in **(a)** runoff coefficient Q/P , **(b)** evaporative index E_A/P , **(c)** mean annual streamflow, **(d)** mean annual maximum streamflow, **(e)** minimum annual 7-days mean streamflow, **(f)** median annual deficit volume below the reference 90th percentile streamflow. Note the different y-axis scales.




Figure 8. Percentage change in mean monthly hydrological response of several flux and state variables between the 2K and historical model runs for the four scenarios, each based on different assumptions for the root-zone storage capacity parameter $S_{R,max}$. Percentage change in mean monthly (a) streamflow Q , (b) root-zone storage S_R , (c) actual evaporation E_A , (d) groundwater storage S_S . The lines and shaded areas show the median and 5-95th percentiles from the ensemble of parameter sets retained as feasible.

8 Data availability

Streamflow data for the Belgian stations are provided by the Service Public de Wallonie in Belgium (Direction générale opérationnelle de la Mobilité et des Voies hydrauliques, Département des Etudes et de l'Appui à la Gestion, Direction de la Gestion hydrologique intégrée (Bld du Nord 8-5000 Namur, Belgium)). Streamflow data for the French stations are retrieved from the Banque Hydro portal (<http://www.hydro.eaufrance.fr/>). The E-OBS dataset (v20.0e) for daily precipitation, temperature and radiation fields for the historical period is used (Cornes et al., 2018) and can be downloaded from <https://www.ecad.eu/download/ensembles/download.php>. The simulated historical and 2K climate data are provided by the Royal Netherlands Meteorological Institute (KNMI).

700 *Author contributions.* HHG, MH, EEA and LJEB designed the study. EEA provided the simulated historical and 2K climate data. LJEB conducted all the analyses and wrote the manuscript. All authors discussed the results and contributed to the final manuscript.

Competing interests. The authors declare that they have no competing interests.

Acknowledgements. We thank Deltares and Rijkswaterstaat for the financial support to conduct this analysis. The authors would like to thank the Service Public de Wallonie, Direction générale opérationnelle de la Mobilité et des Voies hydrauliques, Département des Etudes et de l'Appui à la Gestion, Direction de la Gestion hydrologique intégrée (Bld du Nord 8-5000 Namur, Belgium) for providing the streamflow data. We thank Erik van Meijgaard for performing the historical and 2K climate data simulations. We also thank Wouter Berghuijs for his valuable advice on the Budyko framework. [We are grateful for the constructive comments of Patricia Saco and two anonymous referees which helped us to considerably improve the manuscript and would like to thank the editor Erwin Zehe.](#)

References

- 710 Aalbers, E., van Meijgaard, E., Lenderink, G., de Vries H., and van den Hurk, B.: The 2018 European drought under future climate conditions, in prep., 2021.
- Aalbers, E. E., Lenderink, G., van Meijgaard, E., and van den Hurk, B. J.: Local-scale changes in mean and heavy precipitation in Western Europe, climate change or internal variability?, *Climate Dynamics*, 50, 4745–4766, <https://doi.org/10.1007/s00382-017-3901-9>, <http://dx.doi.org/10.1007/s00382-017-3901-9>, 2018.
- 715 Addor, N., Newman, A. J., Mizukami, N., and Clark, M. P.: The CAMELS data set: catchment attributes and meteorology for large-sample studies, *Hydrology and Earth System Sciences*, 21, 5293–5313, <https://doi.org/10.5194/hess-21-5293-2017>, <https://www.hydrol-earth-syst-sci.net/21/5293/2017/>, 2017.
- Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., Kitzeberger, T., Rigling, A., Breshears, D. D., Hogg, E. H., Gonzalez, P., Fensham, R., Zhang, Z., Castro, J., Demidova, N., Lim, J. H., Allard, G., Running, S. W., Semerci, A., and
- 720 Cobb, N.: A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests, *Forest Ecology and Management*, 259, 660–684, <https://doi.org/10.1016/j.foreco.2009.09.001>, 2010.
- Andréassian, V., Parent, E., and Michel, C.: A distribution-free test to detect gradual changes in watershed behavior, *Water Resources Research*, 39, 1–11, <https://doi.org/10.1029/2003WR002081>, 2003.
- Attema, J. J., Loriaux, J. M., and Lenderink, G.: Extreme precipitation response to climate perturbations in an atmospheric mesoscale model,
- 725 *Environmental Research Letters*, 9, <https://doi.org/10.1088/1748-9326/9/1/014003>, 2014.
- Balsamo, G., Viterbo, P., Beijaars, A., van den Hurk, B., Hirschi, M., Betts, A. K., and Scipal, K.: A revised hydrology for the ECMWF model: Verification from field site to terrestrial water storage and impact in the integrated forecast system, *Journal of Hydrometeorology*, 10, 623–643, <https://doi.org/10.1175/2008JHM1068.1>, 2009.
- Banque Hydro: Banque Hydro, <http://hydro.eaufrance.fr/>, 2018.
- 730 Bastin, J. F., Clark, E., Elliott, T., Hart, S., Van Den Hoogen, J., Hordijk, I., Ma, H., Majumder, S., Manoli, G., Maschler, J., Mo, L., Routh, D., Yu, K., Zohner, C. M., and Crowther, T. W.: Correction: Understanding climate change from a global analysis of city analogues (*PLoS ONE* (2019) 14:7 (e0217592) DOI: 10.1371/journal.pone.0217592), *PLoS ONE*, 14, 1–13, <https://doi.org/10.1371/journal.pone.0224120>, 2019.
- Berghuijs, W. R., Larsen, J. R., van Emmerik, T. H., and Woods, R. A.: A Global Assessment of Runoff Sensitivity to Changes in Precip-
- 735 itation, Potential Evaporation, and Other Factors, *Water Resources Research*, 53, 8475–8486, <https://doi.org/10.1002/2017WR021593>, 2017.
- Berghuijs, W. R., Gnann, S. J., and Woods, R. A.: Unanswered questions on the Budyko framework, *Hydrological Processes*, pp. 1–5, <https://doi.org/10.1002/hyp.13958>, 2020.
- Blöschl, G. and Montanari, A.: Climate change impacts-throwing the dice?, *Hydrological Processes*, 24, 374–381, <https://doi.org/10.1002/hyp.7574>, <http://jamsb.austms.org.au/courses/CSC2408/semester3/resources/ldp/abs-guide.pdf>, 2010.
- 740 Blöschl, G., Bierkens, M. F. P., Chambel, A., Cudennec, C., Destouni, G., Fiori, A., Kirchner, J. W., McDonnell, J. J., Savenije, H. H. G., Sivapalan, M., Stumpff, C., Toth, E., Volpi, E., Carr, G., Lupton, C., Salinas, J., Széles, B., Viglione, A., Aksoy, H., Allen, S. T., Amin, A., Andréassian, V., Arheimer, B., Aryal, S. K., Baker, V., Bardsley, E., Barendrecht, M. H., Bartosova, A., Batelaan, O., Berghuijs, W. R., Beven, K., Blume, T., Bogaard, T., de Amorim, P. B., Böttcher, M. E., Boulet, G., Breinl, K., Brilly, M., Brocca, L., Buytaert, W.,
- 745 Castellarin, A., Castelletti, A., Chen, X., Chen, Y., Chen, Y., Chiffard, P., Claps, P., Clark, M. P., Collins, A. L., Croke, B., Dathe, A.,

- David, P. C., de Barros, F. P. J., de Rooij, G., Baldassarre, G. D., Driscoll, J. M., Duethmann, D., Dwivedi, R., Eris, E., Farmer, W. H., Feiccabrino, J., Ferguson, G., Ferrari, E., Ferraris, S., Fersch, B., Finger, D., Foglia, L., Fowler, K., Gartsman, B., Gascoïn, S., Gaume, E., Gelfan, A., Geris, J., Gharari, S., Gleeson, T., Glendell, M., Bevacqua, A. G., González-Dugo, M. P., Grimaldi, S., Gupta, A. B., Guse, B., Han, D., Hannah, D., Harpold, A., Haun, S., Heal, K., Helfricht, K., Herrnegger, M., Hipsey, M., Hlaváčiková, H., Hohmann, C., Holko, L., Hopkinson, C., Hrachowitz, M., Illangasekare, T. H., Inam, A., Innocente, C., Istanbuluoglu, E., Jarihani, B., Kalantari, Z., Kalvans, A., Khanal, S., Khatami, S., Kiesel, J., Kirkby, M., Knoben, W., Kochanek, K., Kohnová, S., Kolechkina, A., Krause, S., Kreamer, D., Kreibich, H., Kunstmann, H., Lange, H., Liberato, M. L. R., Lindquist, E., Link, T., Liu, J., Loucks, D. P., Luce, C., Mahé, G., Makarieva, O., Malard, J., Mashtayeva, S., Maskey, S., Mas-Pla, J., Mavrova-Guirguinova, M., Mazzoleni, M., Mernild, S., Misstear, B. D., Montanari, A., Müller-Thomy, H., Nabizadeh, A., Nardi, F., Neale, C., Nesterova, N., Nurtaev, B., Odongo, V. O., Panda, S., Pande, S., Pang, Z., Papacharalampous, G., Perrin, C., Pfister, L., Pimentel, R., Polo, M. J., Post, D., Sierra, C. P., Ramos, M.-H., Renner, M., Reynolds, J. E., Ridolfi, E., Rigon, R., Riva, M., Robertson, D. E., Rosso, R., Roy, T., Sá, J. H. M., Salvadori, G., Sandells, M., Schaeffli, B., Schumann, A., Scolobig, A., Seibert, J., Servat, E., Shafiei, M., Sharma, A., Sidibe, M., Sidle, R. C., Skaugen, T., Smith, H., Spiessl, S. M., Stein, L., Steinsland, I., Strasser, U., Su, B., Szolgay, J., Tarboton, D., Tauro, F., Thirel, G., Tian, F., Tong, R., Tussupova, K., Tyralis, H., Uijlenhoet, R., van Beek, R., van der Ent, R. J., van der Ploeg, M., Loon, A. F. V., van Meerveld, I., van Nooijen, R., van Oel, P. R., Vidal, J.-P., von Freyberg, J., Vorogushyn, S., Wachniew, P., Wade, A. J., Ward, P., Westerberg, I. K., White, C., Wood, E. F., Woods, R., Xu, Z., Yilmaz, K. K., and Zhang, Y.: Twenty-three unsolved problems in hydrology (UPH) – a community perspective, *Hydrological Sciences Journal*, 64, 1141–1158, <https://doi.org/10.1080/02626667.2019.1620507>, <https://doi.org/10.1080/02626667.2019.1620507>, 2019.
- Booij, M. J.: Impact of climate change on river flooding assessed with different spatial model resolutions, *Journal of Hydrology*, 303, 176–198, <https://doi.org/10.1016/j.jhydrol.2004.07.013>, 2005.
- Bouaziz, L., Weerts, A., Schellekens, J., Sprokkereef, E., Stam, J., Savenije, H., and Hrachowitz, M.: Redressing the balance: Quantifying net intercatchment groundwater flows, *Hydrology and Earth System Sciences*, 22, 6415–6434, <https://doi.org/10.5194/hess-22-6415-2018>, 2018.
- Bouaziz, L. J., Steele-Dunne, S. C., Schellekens, J., Weerts, A. H., Stam, J., Sprokkereef, E., Winsemius, H. H., Savenije, H. H., and Hrachowitz, M.: Improved understanding of the link between catchment-scale vegetation accessible storage and satellite-derived Soil Water Index, *Water Resources Research*, <https://doi.org/10.1029/2019WR026365>, 2020.
- Bouaziz, L. J., Fenicia, F., Thirel, G., De Boer-Euser, T., Buitink, J., Brauer, C. C., De Niel, J., Dewals, B. J., Drogue, G., Grelier, B., Melsen, L. A., Moustakas, S., Nossent, J., Pereira, F., Sprokkereef, E., Stam, J., Weerts, A. H., Willems, P., Savenije, H. H., and Hrachowitz, M.: Behind the scenes of streamflow model performance, *Hydrology and Earth System Sciences*, 25, 1069–1095, <https://doi.org/10.5194/hess-25-1069-2021>, 2021.
- Brogli, R., Kröner, N., Sørland, S. L., Lüthi, D., and Schär, C.: The role of hadley circulation and lapse-rate changes for the future European summer climate, *Journal of Climate*, 32, 385–404, <https://doi.org/10.1175/JCLI-D-18-0431.1>, 2019.
- Brown, A. E., Zhang, L., McMahon, T. A., Western, A. W., and Vertessy, R. A.: A review of paired catchment studies for determining changes in water yield resulting from alterations in vegetation, *Journal of Hydrology*, 310, 28–61, <https://doi.org/10.1016/j.jhydrol.2004.12.010>, 2005.
- Brunner, I., Herzog, C., Dawes, M. A., Arend, M., and Sperisen, C.: How tree roots respond to drought, *Frontiers in Plant Science*, 6, 1–16, <https://doi.org/10.3389/fpls.2015.00547>, 2015.
- Brunner, M. I., Farinotti, D., Zekollari, H., Huss, M., and Zappa, M.: Future shifts in extreme flow regimes in Alpine regions, *Hydrology and Earth System Sciences Discussions*, pp. 1–26, <https://doi.org/10.5194/hess-2019-144>, 2019.

- Budyko, M. I.: The heat balance of the earth's surface, *Soviet Geography*, 2, 3–13, 1961.
- 785 Buytaert, W. and Beven, K.: Regionalization as a learning process, *Water Resources Research*, 45, 1–13, <https://doi.org/10.1029/2008WR007359>, 2009.
- Calder, I. R., Reid, I., Nisbet, T. R., and Green, J. C.: Impact of lowland forests in England on water resources: Application of the Hydrological Land Use Change (HYLUC) model, *Water Resources Research*, 39, 1–10, <https://doi.org/10.1029/2003WR002042>, 2003.
- Cateau, E., Larrieu, L., Vallauri, D., Savoie, J. M., Touroult, J., and Brustel, H.: Ancienneté et maturité: Deux qualités complémentaires d'un
790 écosystème forestier, *Comptes Rendus - Biologies*, 338, 58–73, <https://doi.org/10.1016/j.crvi.2014.10.004>, 2015.
- Clark, M. P., Slater, A. G., Rupp, D. E., Woods, R. A., Vrugt, J. A., Gupta, H. V., Wagener, T., and Hay, L. E.: Framework for Understanding Structural Errors (FUSE): A modular framework to diagnose differences between hydrological models, *Water Resources Research*, 44, 1–14, <https://doi.org/10.1029/2007wr006735>, 2008.
- Collins, D. B. and Bras, R. L.: Plant rooting strategies in water-limited ecosystems, *Water Resources Research*, 43, 1–10,
795 <https://doi.org/10.1029/2006WR005541>, 2007.
- Cornes, R. C., van der Schrier, G., van den Besselaar, E. J., and Jones, P. D.: An Ensemble Version of the E-OBS Temperature and Precipitation Data Sets, *Journal of Geophysical Research: Atmospheres*, 123, 9391–9409, <https://doi.org/10.1029/2017JD028200>, 2018.
- Coron, L.: Les modèles hydrologiques conceptuels sont-ils robustes face à un climat en évolution ?, Ph.D. thesis, AgroParisTech, France, https://webgr.irstea.fr/wp-content/uploads/2012/11/these_Coron.pdf, 2013.
- 800 Coron, L., Andréassian, V., Perrin, C., Lerat, J., Vaze, J., Bourqui, M., and Hendrickx, F.: Crash testing hydrological models in contrasted climate conditions: An experiment on 216 Australian catchments, *Water Resources Research*, 48, 1–17, <https://doi.org/10.1029/2011WR011721>, 2012.
- de Boer-Euser, T.: Added value of distribution in rainfall-runoff models for the Meuse basin, Ph.D. thesis, Delft University of Technology, The Netherlands, 2017.
- 805 de Boer-Euser, T., McMillan, H. K., Hrachowitz, M., Winsemius, H. C., and Savenije, H. H.: Influence of soil and climate on root zone storage capacity, *Water Resources Research*, <https://doi.org/10.1002/2015WR018115>, 2016.
- de Wit, M. J., van den Hurk, B., Warmerdam, P. M., Torfs, P. J., Roulin, E., and Van Deursen, W. P.: Impact of climate change on low-flows in the river Meuse, *Climatic Change*, 82, 351–372, <https://doi.org/10.1007/s10584-006-9195-2>, 2007.
- Donohue, R. J., Roderick, M. L., and McVicar, T. R.: Roots, storms and soil pores: Incorporating key ecohydrological processes into
810 Budyko's hydrological model, *Journal of Hydrology*, 436–437, 35–50, <https://doi.org/10.1016/j.jhydrol.2012.02.033>, <http://dx.doi.org/10.1016/j.jhydrol.2012.02.033>, 2012.
- Donohue, R. J., Roderick, M. L., McVicar, T. R., and Farquhar, G. D.: Impact of CO₂ fertilization on maximum foliage cover across the globe's warm, arid environments, *Geophysical Research Letters*, 40, 3031–3035, <https://doi.org/10.1002/grl.50563>, 2013.
- Dralle, D. N., Hahm, W. J., Chadwick, K. D., McCormick, E., and Rempe, D. M.: Technical note: Accounting for snow in the estimation of
815 root zone water storage capacity from precipitation and evapotranspiration fluxes, *Hydrology and Earth System Sciences*, 25, 2861–2867, <https://doi.org/10.5194/hess-25-2861-2021>, <https://hess.copernicus.org/articles/25/2861/2021/>, 2021.
- Duethmann, D., Blöschl, G., and Parajka, J.: Why does a conceptual hydrological model fail to correctly predict discharge changes in response to climate change?, *Hydrology and Earth System Sciences*, 24, 3493–3511, <https://doi.org/10.5194/hess-24-3493-2020>, 2020.
- Eilander, D., van Verseveld, W., Yamazaki, D., Weerts, A., Winsemius, H. C., and Ward, P. J.: A hydrography upscaling method
820 for scale-invariant parametrization of distributed hydrological models, *Hydrology and Earth System Sciences*, 25, 5287–5313, <https://doi.org/10.5194/hess-25-5287-2021>, 2021.

- European Environment Agency: Corine Land Cover (CLC) 2018, Version 2020 20u1, <https://land.copernicus.eu/pan-european/corine-land-cover/>, 2018.
- 825 Fan, Y., Miguez-Macho, G., Jobbágy, E. G., Jackson, R. B., and Otero-Casal, C.: Hydrologic regulation of plant rooting depth, *Proceedings of the National Academy of Sciences*, p. 201712381, <https://doi.org/10.1073/pnas.1712381114>, <http://www.pnas.org/lookup/doi/10.1073/pnas.1712381114>, 2017.
- Fenicia, F., Savenije, H. H., and Avdeeva, Y.: Anomaly in the rainfall-runoff behaviour of the Meuse catchment. Climate, land-use, or land-use management?, *Hydrology and Earth System Sciences*, 13, 1727–1737, <https://doi.org/10.5194/hess-13-1727-2009>, 2009.
- 830 Fitzpatrick, M. C. and Dunn, R. R.: Contemporary climatic analogs for 540 North American urban areas in the late 21st century, *Nature Communications*, 10, 1–7, <https://doi.org/10.1038/s41467-019-08540-3>, <http://dx.doi.org/10.1038/s41467-019-08540-3>, 2019.
- Frank, D. C., Poulter, B., Saurer, M., Esper, J., Huntingford, C., Helle, G., Treydte, K., Zimmermann, N. E., Schleser, G. H., Ahlström, A., Ciais, P., Friedlingstein, P., Levis, S., Lomas, M., Sitch, S., Viovy, N., Andreu-Hayles, L., Bednarz, Z., Berninger, F., Boettger, T., D'alessandro, C. M., Daux, V., Filot, M., Grabner, M., Gutierrez, E., Haupt, M., Hiltunen, E., Jungner, H., Kalela-Brundin, M., Krapiec, M., Leuenberger, M., Loader, N. J., Marah, H., Masson-Delmotte, V., Pazdur, A., Pawelczyk, S., Pierre, M., Planells, O., Pukiene, R., 835 Reynolds-Henne, C. E., Rinne, K. T., Saracino, A., Sonninen, E., Stievenard, M., Switsur, V. R., Szczepanek, M., Szychowska-Krapiec, E., Todaro, L., Waterhouse, J. S., and Weigl, M.: Water-use efficiency and transpiration across European forests during the Anthropocene, *Nature Climate Change*, 5, 579–583, <https://doi.org/10.1038/nclimate2614>, 2015.
- Fu, B.: On the calculation of the evaporation from land surface [in Chinese], *Scientia Atmospherica Sinica*, 5, 23–31, 1981.
- 840 Gao, C., Booij, M. J., and Xu, Y.-P.: Assessment of extreme flows and uncertainty under climate change: Disentangling the uncertainty contribution of representative concentration pathways, global climate models and internal climate variability, *Hydrology and Earth System Sciences*, 24, 3251–3269, <https://doi.org/10.5194/hess-24-3251-2020>, 2020.
- Gao, H., Hrachowitz, M., Schymanski, S. J., Fenicia, F., Sriwongsitanon, N., and Savenije, H. H. G.: Climate controls how ecosystems size the root zone storage capacity at catchment scale, *Geophysical Research Letters*, 41, 7916–7923, <https://doi.org/10.1002/2014GL061668>, 2014.
- 845 Gao, J., Holden, J., and Kirkby, M.: A distributed TOPMODEL for modelling impacts of land-cover change on river flow in upland peatland catchments, *Hydrological Processes*, 29, 2867–2879, <https://doi.org/10.1002/hyp.10408>, 2015.
- Gentine, P., D'Odorico, P., Lintner, B. R., Sivandran, G., and Salvucci, G.: Interdependence of climate, soil, and vegetation as constrained by the Budyko curve, *Geophysical Research Letters*, 39, 2–7, <https://doi.org/10.1029/2012GL053492>, 2012.
- 850 Gerrits, A. M., Savenije, H. H., Veling, E. J., and Pfister, L.: Analytical derivation of the Budyko curve based on rainfall characteristics and a simple evaporation model, *Water Resources Research*, 45, 1–15, <https://doi.org/10.1029/2008WR007308>, 2009.
- Gharari, S., Hrachowitz, M., Fenicia, F., and Savenije, H. H. G.: Hydrological landscape classification: Investigating the performance of HAND based landscape classifications in a central European meso-scale catchment, *Hydrology and Earth System Sciences*, 15, 3275–3291, <https://doi.org/10.5194/hess-15-3275-2011>, 2011.
- 855 Gharari, S., Hrachowitz, M., Fenicia, F., and Savenije, H. H.: An approach to identify time consistent model parameters: Sub-period calibration, *Hydrology and Earth System Sciences*, 17, 149–161, <https://doi.org/10.5194/hess-17-149-2013>, 2013.
- Gleeson, T., Wang-Erlandsson, L., Porkka, M., Zipper, S. C., Jaramillo, F., Gerten, D., Fetzer, I., Cornell, S. E., Piemontese, L., Gordon, L. J., Rockström, J., Oki, T., Sivapalan, M., Wada, Y., Brauman, K. A., Flörke, M., Bierkens, M. F., Lehner, B., Keys, P., Kummu, M., Wagener, T., Dadson, S., Troy, T. J., Steffen, W., Falkenmark, M., and Famiglietti, J. S.: Illuminating water cycle modifications and Earth system resilience in the Anthropocene, *Water Resources Research*, 56, 1–24, <https://doi.org/10.1029/2019WR024957>, 2020.

- 860 Guswa, A. J.: The influence of climate on root depth: A carbon cost-benefit analysis, *Water Resources Research*, 44, 1–11, <https://doi.org/10.1029/2007WR006384>, 2008.
- Hakala, K., Addor, N., Gobbe, T., Ruffieux, J., and Seibert, J.: Risks and opportunities for a Swiss hydropower company in a changing climate, *Hydrology and Earth System Sciences Discussions*, pp. 1–35, <https://doi.org/10.5194/hess-2019-475>, 2019.
- Hanus, S., Hrachowitz, M., Zekollari, H., Schoups, G., Vizcaino, M., and Kaitna, R.: Future changes in annual, seasonal and
865 monthly runoff signatures in contrasting Alpine catchments in Austria, *Hydrology and Earth System Sciences*, 25, 3429–3453, <https://doi.org/10.5194/hess-25-3429-2021>, <https://hess.copernicus.org/articles/25/3429/2021/>, 2021.
- Harman, C. and Troch, P. A.: What makes Darwinian hydrology "darwinian"? Asking a different kind of question about landscapes, *Hydrology and Earth System Sciences*, 18, 417–433, <https://doi.org/10.5194/hess-18-417-2014>, 2014.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Sim-
870 mons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 875 Hooghart, J. C. and Lablans, W. N.: Van Penman naar Makkink: een nieuwe berekeningswijze voor de klimatologische verdampingsgetallen, De Bilt, Royal Netherlands Meteorological Institute (KNMI), De Bilt, the Netherlands, 1988.
- Hrachowitz, M., Stockinger, M., Coenders-gerrits, M., Ent, R. V. D., Lücke, A., and Stumpp, C.: Reduction of vegetation-accessible water storage capacity after deforestation affects catchment travel time distributions and increases young water fractions in a headwater catchment, *Hydrol. Earth Syst. Sci.*, 25, 4887–4915, 2021.
- 880 Hulsman, P., Winsemius, H., Michailovsky, C., Savenije, H., and Hrachowitz, M.: Using altimetry observations combined with GRACE to select parameter sets of a hydrological model in data scarce regions, *Hydrology and Earth System Sciences*, pp. 1–35, <https://doi.org/10.5194/hess-2019-346>, 2019.
- Hulsman, P., Hrachowitz, M., and Savenije, H. H.: Improving the representation of long-term storage variations with conceptual hydrological models in data-scarce regions, *Water Resources Research*, 57, e2020WR028 837, 2021.
- 885 Institut National de l'Information Géographique et Forestière: La base de données Forêt version 2.0, <https://inventaire-forestier.ign.fr/spip.php?rubrique227>, 2019.
- Jaramillo, F. and Destouni, G.: Developing water change spectra and distinguishing change drivers worldwide, *Geophysical Research Letters*, 41, 8377–8386, <https://doi.org/10.1002/2014GL061848>, 2014.
- Jaramillo, F., Cory, N., Arheimer, B., Laudon, H., Van Der Velde, Y., Hasper, T. B., Teutschbein, C., and Uddling, J.: Dominant effect of
890 increasing forest biomass on evapotranspiration: Interpretations of movement in Budyko space, *Hydrology and Earth System Sciences*, 22, 567–580, <https://doi.org/10.5194/hess-22-567-2018>, 2018.
- Jasechko, S.: Plants turn on the tap, *Nature Climate Change*, 8, 560–563, 2018.
- Keenan, T. F., Hollinger, D. Y., Bohrer, G., Dragoni, D., Munger, J. W., Schmid, H. P., and Richardson, A. D.: Increase in forest water-use efficiency as atmospheric carbon dioxide concentrations rise, *Nature*, 499, 324–327, <https://doi.org/10.1038/nature12291>, 2013.
- 895 Kervyn, T., Scohy, J.-P., Marchal, D., Collette, O., Hardy, B., Delahaye, L., Wibail, L., Jacquemin, F., Dufrêne, M., and Claessens, H.: La gestion patrimoniale des forêts anciennes de Wallonie (Belgique), *Forêt Nature*, 148, <https://doi.org/10.4267/2042/67878>, 2018.

- Kleidon, A.: Global datasets and rooting zone depth inferred from inverse methods, *Journal of Climate*, 17, 2714–2722, [https://doi.org/10.1175/1520-0442\(2004\)017<2714:GDORZD>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<2714:GDORZD>2.0.CO;2), 2004.
- 900 Kleidon, A. and Heimann, M.: A method of determining rooting depth from a terrestrial biosphere model and its impacts on the global water and carbon cycle, *Global Change Biology*, 4, 275–286, <https://doi.org/10.1046/j.1365-2486.1998.00152.x>, 1998.
- Klingen, S.: *Twaalf boslessen, Klingen Bomen*, Doorn, 2017.
- Klingler, C., Schulz, K., and Herrnegger, M.: LamaH-CE: LARge-SaMple DATA for Hydrology and Environmental Sciences for Central Europe, *Earth System Science Data*, 13, 4529–4565, <https://doi.org/10.5194/essd-13-4529-2021>, <https://essd.copernicus.org/articles/13/4529/2021/>, 2021.
- 905 Kovats, R. S., Valentini, R., Bouwer, L. M., Georgopoulou, E., Jacob, D., Martin, E., Rounsevell, M., and Soussana, J. F.: Europe, Climate Change 2014: Impacts, Adaptation and Vulnerability: Part B: Regional Aspects: Working Group II Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, pp. 1267–1326, <https://doi.org/10.1017/CBO9781107415386.003>, 2014.
- Latte, N., Lebourgeois, F., Kint, V., Drouet, T., and Claessens, H.: Le hêtre face au changement climatique: Le cas de la Belgique, *Revue Forestière Française*, 69, 205–218, <https://doi.org/10.4267/2042/65336>, 2017.
- 910 Lebourgeois, F. and Mérian, P.: La sensibilité au climat des arbres forestiers a-t-elle changé au cours du XXe siècle?, *Revue Forestière Française*, 63, 17–32, <https://doi.org/10.4267/2042/43091>, 2011.
- Levia, D. F., Creed, I. F., Hannah, D. M., Nanko, K., Boyer, E. W., Carlyle-moses, D. E., Giesen, N. V. D., Grasso, D., Guswa, A. J., Hudson, J. E., Hudson, S. A., Iida, S., Jackson, R. B., Katul, G. G., Kumagai, T., Llorens, P., Ribeiro, F. L., Pataki, D. E., Peters, C. A., Carretero, D. S., and Selker, J. S.: Homogenization of the terrestrial water cycle, *Nature Geoscience*, 13, 656–658, <https://doi.org/10.1038/s41561-020-0641-y>, 2020.
- 915 Mao, D. and Cherkauer, K. A.: Impacts of land-use change on hydrologic responses in the Great Lakes region, *Journal of Hydrology*, 374, 71–82, <https://doi.org/10.1016/j.jhydrol.2009.06.016>, <http://dx.doi.org/10.1016/j.jhydrol.2009.06.016>, 2009.
- McCormick, E. L., Dralle, D. N., Hahm, W. J., Tune, A. K., Schmidt, L. M., Chadwick, K. D., and Rempe, D. M.: Widespread woody plant use of water stored in bedrock, *Nature*, 597, 225–229, 2021.
- 920 Merz, R., Parajka, J., and Blöschl, G.: Time stability of catchment model parameters: Implications for climate impact analyses, *Water Resources Research*, 47, 1–17, <https://doi.org/10.1029/2010WR009505>, 2011.
- Mezentsev, V.: Back to the computation of total evaporation, *Meteorologia i Hidrologia*, 5, 24–26, 1955.
- Milly, P. C.: Climate, interseasonal storage of soil water, and the annual water balance, *Advances in Water Resources*, 17, 19–24, [https://doi.org/10.1016/0309-1708\(94\)90020-5](https://doi.org/10.1016/0309-1708(94)90020-5), 1994.
- 925 Milly, P. C., Betancourt, J., Falkenmark, M., Hirsch, R. M., Kundzewicz, Z. W., Lettenmaier, D. P., and Stouffer, R. J.: Climate change: Stationarity is dead: Whither water management?, *Science*, 319, 573–574, <https://doi.org/10.1126/science.1151915>, 2008.
- Miralles, D. G., Brutsaert, W., Dolman, A. J., and Gash, J. H.: On the use of the term “Evapotranspiration”, *Earth and Space Science Open Archive*, p. 8, <https://doi.org/10.1002/essoar.10503229.1>, <https://doi.org/10.1002/essoar.10503229.1>, 2020.
- Nijzink, R., Hutton, C., Pechlivanidis, I., Capell, R., Arheimer, B., Freer, J., Han, D., Wagener, T., McGuire, K., Savenije, H., and Hrachowitz, M.: The evolution of root-zone moisture capacities after deforestation: A step towards hydrological predictions under change?, *Hydrology and Earth System Sciences*, 20, 4775–4799, <https://doi.org/10.5194/hess-20-4775-2016>, 2016a.
- 930 Nijzink, R. C., Samaniego, L., Mai, J., Kumar, R., Thober, S., Zink, M., Schäfer, D., Savenije, H. H., and Hrachowitz, M.: The importance of topography-controlled sub-grid process heterogeneity and semi-quantitative prior constraints in distributed hydrological models, *Hydrology and Earth System Sciences*, 20, 1151–1176, <https://doi.org/10.5194/hess-20-1151-2016>, 2016b.

- 935 Peel, M. C. and Blöschl, G.: Hydrological modelling in a changing world, *Progress in Physical Geography*, 35, 249–261, <https://doi.org/10.1177/0309133311402550>, 2011.
- Pomeroy, J., Fang, X., and Ellis, C.: Sensitivity of snowmelt hydrology in Marmot Creek, Alberta, to forest cover disturbance, *Hydrological Processes*, 26, 1891–1904, <https://doi.org/10.1002/hyp.9248>, 2012.
- Prein, A. F., Rasmussen, R. M., Ikeda, K., Liu, C., Clark, M. P., and Holland, G. J.: The future intensification of hourly precipitation extremes, *Nature Climate Change*, 7, 48–52, <https://doi.org/10.1038/nclimate3168>, 2017.
- 940 Prudhomme, C., Giuntoli, I., Robinson, E. L., Clark, D. B., Arnell, N. W., Dankers, R., Fekete, B. M., Franssen, W., Gerten, D., Gosling, S. N., Hagemann, S., Hannah, D. M., Kim, H., Masaki, Y., Satoh, Y., Stacke, T., Wada, Y., and Wisser, D.: Hydrological droughts in the 21st century, hotspots and uncertainties from a global multimodel ensemble experiment, *Proceedings of the National Academy of Sciences of the United States of America*, 111, 3262–3267, <https://doi.org/10.1073/pnas.1222473110>, 2014.
- 945 Reaver, N., Kaplan, D., Klammler, H., and Jawitz, J.: Reinterpreting the Budyko Framework, *Hydrology and Earth System Sciences Discussions*, pp. 1–31, <https://doi.org/10.5194/hess-2020-584>, 2020.
- Rennó, C. D., Nobre, A. D., Cuartas, L. A., Soares, J. V., Hodnett, M. G., Tomasella, J., and Waterloo, M. J.: HAND, a new terrain descriptor using SRTM-DEM: Mapping terra-firme rainforest environments in Amazonia, *Remote Sensing of Environment*, 112, 3469–3481, <https://doi.org/http://dx.doi.org/10.1016/j.rse.2008.03.018>, <http://www.sciencedirect.com/science/article/pii/S003442570800120X>, 2008.
- 950 Reu, B., Zaehle, S., Bohn, K., Pavlick, R., Schmidlein, S., Williams, J. W., and Kleidon, A.: Future no-analogue vegetation produced by no-analogue combinations of temperature and insolation, *Global Ecology and Biogeography*, 23, 156–167, <https://doi.org/10.1111/geb.12110>, 2014.
- Rohat, G., Goyette, S., and Flacke, J.: Characterization of European cities' climate shift – an exploratory study based on climate analogues, *International Journal of Climate Change Strategies and Management*, 10, 428–452, <https://doi.org/10.1108/IJCCSM-05-2017-0108>, 2018.
- 955 Rottler, E., Bronstert, A., Bürger, G., and Rakovec, O.: Projected changes in Rhine River flood seasonality under global warming, *Hydrol. Earth Syst. Sci. Discuss.*, pp. 1–25, 2020.
- Savenije, H. H.: The importance of interception and why we should delete the term evapotranspiration from our vocabulary, *Hydrological Processes*, 18, 1507–1511, <https://doi.org/10.1002/hyp.5563>, 2004.
- Savenije, H. H.: HESS opinions "topography driven conceptual modelling (FLEX-Topo)", *Hydrology and Earth System Sciences*, 14, 2681–2692, <https://doi.org/10.5194/hess-14-2681-2010>, 2010.
- 960 Savenije, H. H. and Hrachowitz, M.: HESS Opinions "catchments as meta-organisms - A new blueprint for hydrological modelling", *Hydrology and Earth System Sciences*, 21, 1107–1116, <https://doi.org/10.5194/hess-21-1107-2017>, 2017.
- Schaphoff, S., Von Bloh, W., Rammig, A., Thonicke, K., Biemans, H., Forkel, M., Gerten, D., Heinke, J., Jägermeyr, J., Knauer, J., Langerwisch, F., Lucht, W., Müller, C., Rolinski, S., and Waha, K.: LPJmL4 - A dynamic global vegetation model with managed land - Part 1: *Model description*, *Geoscientific Model Development*, 11, 1343–1375, <https://doi.org/10.5194/gmd-11-1343-2018>, 2018.
- 965 Schär, C., Frei, C., Lüthi, D., and Davies, H. C.: Surrogate climate-change scenarios for regional climate models, *Geophysical Research Letters*, 23, 669–672, <https://doi.org/10.1029/96GL00265>, 1996.
- Schattan, P., Zappa, M., Lischke, H., Bernhard, L., Thürig, E., and Dieckkrüger, B.: An approach for transient consideration of forest change in hydrological impact studies, *IAHS-AISH Proceedings and Reports*, 359, 311–319, 2013.
- 970 Schelhaas, M. J., Nabuurs, G. J., and Schuck, A.: Natural disturbances in the European forests in the 19th and 20th centuries, *Global Change Biology*, 9, 1620–1633, <https://doi.org/10.1046/j.1365-2486.2003.00684.x>, 2003.

- Schellekens, J., Verseveld, W. V., Visser, M., Winsemius, H. H., de Boer-Euser, T., Bouaziz, L. J., Thiange, C., de Vries, S., Boisgontier, H., Eilander, D., Tollenaar, D., Weerts, A. H., Baart, F., Hazenberg, P., Lutz, A., ten Velden, C., Jansen, M., and Benedict, I.: openstreams/wflow, <https://github.com/openstreams/wflow>, 2020.
- 975 Schymanski, S. J., Sivapalan, M., Roderick, M. L., Beringer, J., and Hutley, L. B.: An optimality-based model of the coupled soil moisture and root dynamics, *Hydrology and Earth System Sciences*, 12, 913–932, <https://doi.org/10.5194/hess-12-913-2008>, <https://www.hydrol-earth-syst-sci.net/12/913/2008/>, 2008.
- Schymanski, S. J., Sivapalan, M., Roderick, M. L., Hutley, L. B., and Beringer, J.: An optimality-based model of the dynamic feedbacks between natural vegetation and the water balance, *Water Resources Research*, 45, 1–18, <https://doi.org/10.1029/2008WR006841>, 2009.
- 980 Seibert, J. and van Meerveld, H. I.: Hydrological change modeling: Challenges and opportunities, *Hydrological Processes*, 30, 4966–4971, <https://doi.org/10.1002/hyp.10999>, 2016.
- Service Public de Wallonie: Service Public de Wallonie, 2018.
- Singh, C., Wang-Erlandsson, L., Fetzer, I., Rockström, J., and van der Ent, R.: Rootzone storage capacity reveals drought coping strategies along rainforest-savanna transitions, *Environmental Research Letters*, <https://doi.org/10.1088/1748-9326/abc377>, 2020.
- 985 Singh, R., Wagener, T., Van Werkhoven, K., Mann, M. E., and Crane, R.: A trading-space-for-time approach to probabilistic continuous streamflow predictions in a changing climate-accounting for changing watershed behavior, *Hydrology and Earth System Sciences*, 15, 3591–3603, <https://doi.org/10.5194/hess-15-3591-2011>, 2011.
- Speich, M. J., Lischke, H., and Zappa, M.: Testing an optimality-based model of rooting zone water storage capacity in temperate forests, *Hydrology and Earth System Sciences*, 22, 4097–4124, <https://doi.org/10.5194/hess-22-4097-2018>, 2018.
- 990 Speich, M. J., Zappa, M., Scherstjanoi, M., and Lischke, H.: FORests and HYdrology under Climate Change in Switzerland v1.0: A spatially distributed model combining hydrology and forest dynamics, *Geoscientific Model Development*, 13, 537–564, <https://doi.org/10.5194/gmd-13-537-2020>, 2020.
- Stephens, C. M., Marshall, L. A., and Johnson, F. M.: Investigating strategies to improve hydrologic model performance in a changing climate, *Journal of Hydrology*, 579, 124 219, <https://doi.org/10.1016/j.jhydrol.2019.124219>, <https://doi.org/10.1016/j.jhydrol.2019.124219>, 2019.
- 995 Stephens, C. M., Marshall, L. A., Johnson, F. M., Lin, L., Band, L. E., and Ajami, H.: Is Past Variability a Suitable Proxy for Future Change? A Virtual Catchment Experiment, *Water Resources Research*, 56, 1–25, <https://doi.org/10.1029/2019WR026275>, 2020.
- Stephens, C. M., Lall, U., Johnson, F. M., and Marshall, L. A.: Landscape changes and their hydrologic effects: Interactions and feedbacks across scales, *Earth-Science Reviews*, 212, 103 466, <https://doi.org/10.1016/j.earscirev.2020.103466>, <https://doi.org/10.1016/j.earscirev.2020.103466>, 2021.
- 1000 Stevens, D., Miranda, P. M. A., Orth, R., Boussetta, S., Balsamo, G., and Dutra, E.: Sensitivity of Surface Fluxes in the ECMWF Land Surface Model to the Remotely Sensed Leaf Area Index and Root Distribution: Evaluation with Tower Flux Data, *Atmosphere*, 11, <https://doi.org/10.3390/atmos11121362>, <https://www.mdpi.com/2073-4433/11/12/1362>, 2020.
- Teuling, A. J., De Badts, E. A., Jansen, F. A., Fuchs, R., Buitink, J., Van Dijke, A. J., and Sterling, S. M.: Climate change, reforestation/afforestation, and urbanization impacts on evapotranspiration and streamflow in Europe, *Hydrology and Earth System Sciences*, 23, 3631–3652, <https://doi.org/10.5194/hess-23-3631-2019>, 2019.
- 1005 Tietjen, B., Schlaepfer, D. R., Bradford, J. B., Lauenroth, W. K., Hall, S. A., Duniway, M. C., Hochstrasser, T., Jia, G., Munson, S. M., Pyke, D. A., and Wilson, S. D.: Climate change-induced vegetation shifts lead to more ecological droughts despite projected rainfall increases in many global temperate drylands, *Global Change Biology*, 23, 2743–2754, <https://doi.org/10.1111/gcb.13598>, 2017.

- Troch, P. A., Carrillo, G., Sivapalan, M., Wagener, T., and Sawicz, K.: Climate-vegetation-soil interactions and long-term hydrologic partitioning: Signatures of catchment co-evolution, *Hydrology and Earth System Sciences*, 17, 2209–2217, <https://doi.org/10.5194/hess-17-2209-2013>, 2013.
- Turc, L.: Le Bilan d'eau des sols: relations entre les précipitations, l'évaporation et l'écoulement, *Annales Agronomiques*, 1954.
- Ukkola, A. M., Prentice, I. C., Keenan, T. F., Van Dijk, A. I., Viney, N. R., Myneni, R. B., and Bi, J.: Reduced streamflow in water-stressed climates consistent with CO₂ effects on vegetation, *Nature Climate Change*, 6, 75–78, <https://doi.org/10.1038/nclimate2831>, 2016.
- 1015 van Der Sleen, P., Groenendijk, P., Vlam, M., Anten, N. P., Boom, A., Bongers, F., Pons, T. L., Terburg, G., and Zuidema, P. A.: No growth stimulation of tropical trees by 150 years of CO₂ fertilization but water-use efficiency increased, *Nature Geoscience*, 8, 24–28, <https://doi.org/10.1038/ngeo2313>, 2015.
- van der Velde, Y., Vercauteren, N., Jaramillo, F., Dekker, S. C., Destouni, G., and Lyon, S. W.: Exploring hydroclimatic change disparity via the Budyko framework, *Hydrological Processes*, 28, 4110–4118, <https://doi.org/10.1002/hyp.9949>, 2014.
- 1020 van Meijgaard, E., Ulft, L. H. V., Bosveld, F. C., Lenderink, G., and Siebesma, a. P.: The KNMI regional atmospheric climate model RACMO version 2.1, Technical report; TR - 302, p. 43, 2008.
- van Oorschot, F., van der Ent, R. J., Hrachowitz, M., and Alessandri, A.: Climate-controlled root zone parameters show potential to improve water flux simulations by land surface models, *Earth System Dynamics*, 12, 725–743, 2021.
- van Wijk, M. T. and Bouten, W.: Towards understanding tree root profiles: Simulating hydrologically optimal strategies for root distribution, *Hydrology and Earth System Sciences*, 5, 629–644, <https://doi.org/10.5194/hess-5-629-2001>, 2001.
- 1025 Vaze, J., Post, D. A., Chiew, F. H., Perraud, J. M., Viney, N. R., and Teng, J.: Climate non-stationarity - Validity of calibrated rainfall-runoff models for use in climate change studies, *Journal of Hydrology*, 394, 447–457, <https://doi.org/10.1016/j.jhydrol.2010.09.018>, <http://dx.doi.org/10.1016/j.jhydrol.2010.09.018>, 2010.
- Wagener, T.: Can we model the hydrological impacts of environmental change?, *Hydrological Processes*, 21, 3233–3236, <https://doi.org/10.1002/hyp.6873>, <http://jamsb.austms.org.au/courses/CSC2408/semester3/resources/ldp/abs-guide.pdf>, 2007.
- 1030 Wang-Erlandsson, L., Bastiaanssen, W. G., Gao, H., Jägermeyr, J., Senay, G. B., Van Dijk, A. I., Guerschman, J. P., Keys, P. W., Gordon, L. J., and Savenije, H. H.: Global root zone storage capacity from satellite-based evaporation, *Hydrology and Earth System Sciences*, 20, 1459–1481, <https://doi.org/10.5194/hess-20-1459-2016>, 2016.
- Yamazaki, D., Ikeshima, D., Sosa, J., Bates, P. D., Allen, G. H., and Pavelsky, T. M.: MERIT Hydro: A High-Resolution Global Hydrography Map Based on Latest Topography Dataset, *Water Resources Research*, 55, 5053–5073, <https://doi.org/10.1029/2019WR024873>, 2019.
- 1035 Yang, Y., Donohue, R. J., and McVicar, T. R.: Global estimation of effective plant rooting depth: Implications for hydrological modeling, *Water Resources Research*, 52, 8260–8276, <https://doi.org/10.1002/2016WR019392>, https://www.cambridge.org/core/product/identifier/CBO9781107415324A009/type/book_parthttp://doi.wiley.com/10.1002/2016WR019392, 2016.
- Yang, Y., Roderick, M. L., Zhang, S., McVicar, T. R., and Donohue, R. J.: Hydrologic implications of vegetation response to elevated CO₂ in climate projections, *Nature Climate Change*, 9, 44–48, <https://doi.org/10.1038/s41558-018-0361-0>, <http://dx.doi.org/10.1038/s41558-018-0361-0>, 2019.
- 1040 Zhang, B., Hautier, Y., Tan, X., You, C., Cadotte, M. W., Chu, C., Jiang, L., Sui, X., Ren, T., Han, X., and Chen, S.: Species responses to changing precipitation depend on trait plasticity rather than trait means and intraspecific variation, *Functional Ecology*, pp. 2622–2633, <https://doi.org/10.1111/1365-2435.13675>, 2020.
- 1045 Zhang, L., Hickel, K., Dawes, W. R., Chiew, F. H., Western, A. W., and Briggs, P. R.: A rational function approach for estimating mean annual evapotranspiration, *Water Resources Research*, 40, 1–14, <https://doi.org/10.1029/2003WR002710>, 2004.