

Daily Temporally resolved coastal hypoxia forecasting and uncertainty assessment via Bayesian mechanistic modeling for the Northern Gulf of Mexico modelling

Alexey Katin¹, Dario Del Giudice¹, Daniel R. Obenour¹

¹Department of Civil, Construction, & Environmental Engineering, North Carolina State University, Raleigh, 27606, USA

Correspondence to: Alexey Katin (akatin@ncsu.edu)

Abstract. Low bottom water dissolved oxygen conditions (hypoxia) occur almost every summer in the northern Gulf of Mexico due to a combination of nutrient loadings and water column stratification. Several [statistical and mechanistic](#) models have been used to forecast the midsummer hypoxic area based on spring nitrogen loading from major rivers. However, sub-seasonal forecasts are needed to fully characterize the dynamics of hypoxia over the summer season, which is important for informing fisheries and ecosystem management. Here, we present an approach to forecast hypoxic conditions at daily resolution through Bayesian mechanistic [modeling/modelling](#) that allows for rigorous uncertainty quantification. Within this framework, we develop and test different representations and projections of [hydro-meteorological/hydrometeorological](#) model inputs. We find that May precipitation over the Mississippi River Basin is a key predictor of summer discharge and loading that substantially improves forecast performance. Accounting for spring wind conditions also improves forecast performance, though to a lesser extent. The proposed approach generates forecasts for two different sections of the Louisiana–Texas shelf (east and west), and it explains about 50% of the variability in total hypoxic area when tested against historical observations (1985–2016). Results also show how forecast uncertainties build over the summer season, with longer lead times from the nominal forecast release date of [31-May/1-June](#), due to increasing stochasticity in riverine and meteorological inputs. Consequently, the portion of overall forecast variance associated with uncertainties in data inputs increases from 26% to 41% from June–July to August–September, respectively. Overall, the study demonstrates a unique approach to assessing and reducing uncertainties in [dynamic/temporally resolved](#) hypoxia forecasting.

Introduction

The Northern Gulf of Mexico ([NGoM](#)) has one of the largest hypoxic zones in the world, forming virtually every summer over the last three decades (Rabalais and Turner, 2019). Hypoxic or “dead” zones occur when dissolved oxygen concentrations fall below critical thresholds (e.g., 2 mg/L) threatening aquatic ecosystems (Craig, 2012; Craig and Crowder, 2005; Thronson and Quigg, 2008), fisheries (Purcell et al., 2017; Smith et al., 2017), and coastal economies (Díaz and Rosenberg, 2011). Two major causes of hypoxia in the Gulf are water column stratification and nutrient loadings (Krug, 2007; Obenour et al., 2012; Rabalais et al., 2002), which are both influenced by Mississippi and Atchafalaya River ([MAR](#)) discharges. Additionally, wind controls both the structure of the river plume (Hetland, 2005) and the rates of oxygen supply to the water column (Fennel et al., 2013; Justić et al., 1996). Overall, a complex combination of biophysical factors including long-term accumulation

of organic matter (Del Giudice et al., 2020; Turner et al., 2008) and short-term events like storms and droughts (Bianchi et al., 2010) control hypoxia dynamics in the [NGoMnorthern Gulf](#).

Mathematical models are useful for elucidating important relationships between hypoxia and environmental drivers, and for evaluating the consequences of possible actions to improve water quality (Justić and Rose, 2017).

40 The approaches developed to predict hypoxia in the Gulf of Mexico included statistical regressions (Forrest et al., 2011; Greene et al., 2009; Turner et al., 2012), as well as both parsimonious (Obenour et al., 2015; Scavia et al., 2013) and complex (Justić and Wang, 2014; Yu et al., 2015) mechanistic models. Among these alternatives, parsimonious process-based models attempt to balance biophysical detail with computational efficiency and resilience to overfitting. When embedded in a Bayesian framework, these models describe eutrophication
45 processes and hypoxia formation while enabling data-driven parameter estimation and rigorous uncertainty analysis (Ménesguen and Lacroix, 2018). The latter is especially important for assessing our confidence in the potential outcomes of environmental change and management decisions (Reichert and Borsuk, 2005; Schuwirth et al., 2019).

Currently, a probabilistic ensemble of four models is used to inform stakeholders and fishery managers about the
50 expected extent of the [NGoMnorthern Gulf](#) hypoxic zone (Scavia et al., 2017). This ensemble provides predictions with estimates of uncertainty of the midsummer hypoxic area (HA) ~~in the NGoM~~. However, the forecast lacks dynamic oxygen predictions over the summer season. The lack of subseasonal information on dissolved oxygen variability has been identified as an important limitation in understanding how hypoxia affects fisheries in the region, which occur primarily during the summer season but are highly dynamic in space and time (Langseth et al., 2016; Purcell et al., 2017; Smith et al., 2014). Laurent and Fennel (2019) used a weighted aggregation of
55 seasonal hindcasts generated by a three-dimensional model to produce spatially and ~~temporally~~temporally resolved seasonal hypoxia forecasts, but without accounting for uncertainties related to the model parameterization. (Mattern et al., 2013). Further, the aforementioned forecasting approaches are informed only by observed spring nutrient loading, without considering variability in spring wind conditions (Obenour et al., 2015)
60 or projected summer [MARriver](#) discharge and loading.

Here, we use ~~a Bayesianan existing~~ mechanistic model, [calibrated within a Bayesian inference framework](#), to forecast the temporal dynamics of hypoxia, ~~with a focus on how different model inputs influence forecast accuracy and uncertainty in the northern Gulf over the summer season~~. The model was initially developed by Obenour et al. (2015) and later enhanced by Del Giudice et al. (2020) (hereafter referred to as DMO20). While the model
65 performed well in hindcasting, its ability to forecast hypoxia forward in time has not been explored. In order to provide sufficient lead for environmental planning and fisheries management, we produce a June–September hypoxia forecast based on data available at the end of May. The main objectives of this study are to: a) develop daily forecasted spatial-mean bottom water dissolved oxygen (BWDO) concentrations and HA estimates for targeted portions of the Louisiana–Texas Shelf with accompanying measures of uncertainty; b) understand the
70 major sources of forecast uncertainty; ~~and~~ c) characterize how forecast accuracy degrades over time; and d) explore how different applications of spring-summer riverine and meteorological data influence forecast performance.

2 Methods


We first outline the underlying model and required data inputs. Next, we describe the proposed forecasting procedure, along with regression models to project discharge and nitrogen loading over the summer. Third, we describe the approach to evaluate BWDO and HA forecast performance and analyse how the forecast varies in relation to alternative combinations of data inputs.

2.1 Bayesian mechanistic model and bias adjustment

The [BWDO hypoxia](#) forecast is based on the model described in DMO20, which has a parsimonious mechanistic formulation and coarse spatial resolution. Specifically, the model represents the Louisiana–Texas shelf from Galveston Bay to the Mississippi River Delta, divided into [west four compartments](#). West and east sections are separated at the Atchafalaya River mouth (Fig. S1). The water column of each section is divided by the pycnocline into two layers, assuming that discharge and nutrient loadings are transported within the top layer. Additionally, wind speed and direction control the distribution of flow and loadings between the east and west sections as well as the rate of reoxygenation across the pycnocline. The biogeochemistry is based on the transformation of bioavailable nitrogen (sum of nitrate, nitrite, ammonia, and 12% organic nitrogen (Obenour et al., 2015)) into organic matter, which settles to the bottom layer and is subject to aerobic decomposition. BWDO is depleted due to both near- and long-term oxygen demands, reflecting the effects of nitrogen loadings over different time scales. Therefore, the model uses both recent inputs of daily discharge, loading, and wind (up to 90 d before the date of prediction) and long-term November–March loading. The model predicts BWDO, which is then transformed into HA via regressions. In addition, the Bayesian calibration framework provides systematic estimation of model parameters and their uncertainties (DMO20 Table S1). All major equations of DMO20, including regressions to convert BWDO to HA, are presented in Section S1. Predictions of BWDO and HA generated by DMO20 using known nutrient and hydrometeorological inputs throughout the summer are hereinafter referred to as “hindcasts” (Fig. 2A).

Prior to developing the daily forecast, we examined DMO20 predictions for systematic biases during specific months and found that hindcasted predicted BWDO was (0–20%) lower than observations for the west section of the shelf in June. However, this discrepancy diminished toward the end of the month (Section S2). This apparent bias, which could be due to an overestimation of June oxygen demands or other structural limitation within DMO20 model, was corrected using a linear regression, with the day number (June 1 to June 30) as a predictor and the BWDO adjustment factor as the response (Fig. S2.1). This adjustment factor was applied to all June model predictions (hindcasts and forecasts) unless otherwise indicated.

Formatted: English (United States)



105 [Figure 1: Map of the study area located in the Northern Gulf of Mexico. Mississippi and Atchafalaya River Basin \(top left, light blue filled\) is used in estimation of monthly precipitation and temperature.](#)

2.2 Data

The forecast utilized the same observational data inputs described in DMO20, including monthly discharge and nitrogen loading from the U.S. Geological Survey (USGS, 2019), daily discharge from the U.S. Army Corps of Engineers at Simmesport and Tarbert Landing (USACE, 2019), and wind speed and zonal-velocities from the National Data Buoy Center (NDBC, 2019). Additionally, monthly precipitation and temperature were obtained from gridded data for the MARMississippi River Basin (Hart and Bell, 2015; Schwartz, 2012). Section-specific observations of For both shelf sections, estimates of mean BWDO and HA with associated uncertainties used for DMO20 calibration were obtained via averaging using the space-time geostatistical estimates from model of Matli et al. (2018). Similar This model relies on BWDO samples from ship-based monitoring cruises, and similar to DMO20, we only used the geostatistical space-time estimates corresponding to times of monitoring cruises, as these estimates generally have lower uncertainties, cruises (when uncertainty is typically lowest). At least one monitoring cruise was conducted every year of the study period (1985–2016) and there were a total of 149 cruises (34 in June, 63 in July, 35 in August, and 17 in September).

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Italic, Underline, English (United States)


120 2.3 Forecast procedure

To capture the uncertainty in hydrometeorology, nutrient loading, model parameters, and residual error, the forecast for a given year was determined through 1000 Monte Carlo simulations (Fig. 4-2B) implemented in R (R Core Team, 2019). Each simulation included a random draw from the Bayesian joint posterior parameter and error distribution of the mechanistic model (DMO20), and the uncertainty in the regression to convert regressions for bias correction and for converting BWDO to HA. The simulations used actual November–May riverine and meteorological inputs for the forecast year, since these inputs would be known by the nominal forecast release date of 31 May 1 June. However, summer (June–September) inputs were sampled from historical records of multiple years (always omitting the forecast year). To retain the temporal correlation in these inputs, data were sampled in blocks (i.e., the complete daily time series for each summer). As wind conditions cannot be accurately


130 forecasted beyond 10 days (Zhang et al., 2019), summer wind velocity data were sampled from the complete historical record (1985–2016).

135 Summer riverine inputs can potentially be projected in advance through regression (Section 2.4). Thus, riverine time series were sampled from only the 10 most “relevant” historical years. For each historical year, relevancy to the forecast year was determined by computing the differences between the summer historical records and the regression-projected discharge and bioavailable nitrogen loading. ~~Both monthly~~ (Section 2.4). Monthly projections and observations were standardized based on the mean and standard deviation of the historical data for each summer month (1980–2016), so that the differences in loading and flow could be combined on the same unitless scale. The 10 years with the smallest aggregated differences were selected as the relevant years for use in the Monte Carlo simulations.

140



(A) Hindcast



(B) Pseudo-forecast




Figure 12: Flowchart summarizing processes and data inputs required to generate the hypoxia hindcast (A) and pseudo-forecast (B) for a given year. Parallelograms represent data inputs and Bayesian posterior parameter distributions. Rectangles represent forecasting computations. Squares with bell-shaped curves indicate steps that propagate uncertainty (including stochastic hydrometeorology).

145

2.4 Regressions for June-September discharge and loading

Regression modeling was used to project riverine inputs, which were employed to constrain the historical records used in the Monte Carlo simulations to relevant years (Section 2.3, Fig. 42B). June, July, August, and September MARiver (Atchafalaya and Mississippi) discharge (Q_A and Q_M , m^3/s) and bioavailable nitrogen loading (L_A and L_M , T/mo) were estimated through multiple linear regression. The candidate predictor variables (predictors) included monthly (January–May) and 4-month average (January–April) discharge, loading, total river basin precipitation (P , in), and river basin temperature (T , °C) (Fig. S3.1). Response variables were square-root transformed to account for the skewness of their distributions and comply with error normality assumption for linear regression (Faraway, 2015). Predictors for each model were selected using the Bayesian Information Criterion (BIC) through an exhaustive search (Lumley, 2017). BIC prioritizes models based on log likelihood while penalizing for the number of parameters to prevent overfitting (Faraway, 2015). The performance of the

150

155

regression was measured by the coefficient of determination, R^2 , in the square-root transformed space. If any of the sixteen regressions had $R^2 \leq 30\%$, the associated month and variable was excluded from determining relevant years for hypoxia forecasting (Section 2.3). [To further check the validity of these regressions, we performed leave-one year-out-cross validation \(LOOCV\), where we excluded years by one, calibrated the models to the reduced dataset, and predicted for the excluded year.](#)

2.5 Forecast assessment

The forecasting approach was applied to the [complete](#) historical record (1985–2016), ~~while~~ excluding the summer input data of the forecast year from the Monte Carlo simulation. This retrospective forecast (i.e., pseudo-forecast) performance was evaluated through comparison of the daily forecasted values with both hindcasted (generated by DMO20) and geostatistically estimated (referred to as “observed” for brevity) BWDO and HA for the two shelf sections. The approach also allowed for determining 95% inter-quantile ranges (IQR) of the pseudo-forecasts, [encompassing variability in summer riverine and meteorological inputs, accounting for parameter uncertainty, and residual error \(mechanistic model residual and transformation of BWDO to HA error, and bias adjustment uncertainties, as well as the stochasticity in riverine and meteorological inputs \(Section 2.3\).](#)


We also assessed how inclusion of various hydrometeorological inputs affected pseudo-forecast accuracy and uncertainty. Specifically, we compared four cases with different types of spring–summer wind and summer riverine data. Case 1 included summer riverine and spring–summer wind records randomly sampled from the complete historical data (thus they are treated as unknown, consistent with conventional Gulf forecasting approaches). Case 2 was similar to Case 1, except it included actual spring wind data (to 31 May). Case 3 was also similar to Case 1, except it used summer riverine records sampled from only the 10 most relevant historical years, as determined from the regression projections (Section 2.4). Finally, Case 4 (our proposed approach, [Section 2.3, Fig. 4B](#)) used both actual spring wind data and riverine records from the 10 most relevant years.

3 Results and Discussion

3.1 Monthly discharge and loading projections

~~Sixteen multiple~~Multiple linear regressions predict average monthly (June to September) summer river discharge and bioavailable nitrogen loading [for the MAR at each river outlet](#). The performance of ~~these 16~~ these 16 regressions generally decreases from the beginning to the end of summer (Table 1), reflecting the increasing temporal gap (i.e., lead time) between the available spring predictors and the forecast response. For instance, the regressions explain 78% and 9% of the variability in (square-root transformed) Mississippi River bioavailable nitrogen loading in June and September, respectively. The residuals for all selected models appear evenly distributed with minimal heteroscedasticity (Fig. S3.21–S3.54) and mostly weak serial correlation of residuals (Pearson lag-1 ~~correlation~~ correlations ranging from -0.02 to 0.35). The predictive variables chosen via exhaustive BIC selection include May discharge [from both rivers](#) (Q_{A5} and/or Q_{M5}) or bioavailable nitrogen loading (L_{A5} and/or L_{M5}) in 13 of the 16 models. In other words, high flow and nutrient loading in May is indicative of high flow and nutrient loading in summer. However, the most consistent individual predictor (present in 12 out of 16 regressions) is [MAR Mississippi River](#) Basin precipitation in May (P_5), likely due to the hydrologic lag between rainfall and basin discharge. Note that the correlation between P_5 and May discharge is relatively weak ($r = 0.36$ for both

220 Pseudo-forecasts can also be compared to observed (geostatistically estimated) BWDO and HA at the times of
 monitoring cruises (Fig. 23, bottom). The forecasted BWDO fits moderately well to the observations with an R^2
 of 0.39 and 0.50 for the west and east sections, respectively. When BWDO is transformed to HA, the pseudo-
 forecast explains 41% and 48% of variability in observed HA in the west and east sections, respectively
 (Fig. S4.1), which is similar to the hindcast explanatory power of 46% and 58% (as in DMO20). In comparison,
 225 hindcasting studies using three-dimensional models have generally explained a lower 27–37% of the variability
 in Gulf BWDO, but at finer spatial resolution (Fennel et al., 2016). To our knowledge, this is the first time Gulf
 hypoxia forecasts have been rigorously compared to observations across the entire summer season (June–
 September). Previous studies have generally focused on assessing forecast performance relative to the Louisiana
 Universities Marine Consortium midsummer shelf-wide hypoxia cruises, which typically take place
 230 place within a two-week window beginning in late July (Laurent and Fennel, 2019; Scavia et al., 2017).
[A tighter selection of relevant years for forecast generation may produce more accurate forecasting results, but
 may capture less of the true stochasticity in the hydrometeorology. If the forecasting approach is revised to include
 only the 5 most relevant years, the predictive accuracy slightly improves based on comparisons with hindcasted
 values \(\$R^2\$ increases by 2.8%\), while there is virtually no improvement based on comparisons with observed HA.](#)
 235 [At the same time, the uncertainty \(standard error\) for the population variance increases by a factor of 1.5 when
 using 5 years instead of 10 years \(Benhamou, 2018\). Therefore, using 10 relevant years appears to provide a more
 reasonable balance between predictive accuracy and uncertainty characterization, though this could be explored
 further in future research.](#)




240

Figure 23: Daily hindcasted (available BWDO (predicted for every summer day from DMO20_1985-2016, assuming hydrometeorology and loading is known throughout the summer) and observed BWDO (geostatistically estimated from monitoring cruises) BWDO versus pseudo-forecasted BWDO for west and east shelf sections. Diagonal line represents perfect prediction.

245

Forecasting performance gradually degrades with longer lead times. The ability of pseudo-forecasts to match DMO20 hindcasts of shelfwide HA declines by about 50% (comparing R^2 values, Fig. 34, top) from June to September due to increasing uncertainty in riverine and meteorological inputs toward the end of the summer season. In comparison, the ability of forecasts to match actual observations declines by nearly 70% (Fig. 3, 4, bottom). Forecasts and hindcasts benefit from the same seasonal patterns inherent to the DMO20 model structure, while observations may deviate from these patterns due to additional drivers of variability not captured in the mechanistic formulation.

250



255 Figure 34: Month-by-month comparison of the daily hindcasted (top) and observed (geostatistically estimated, bottom) shelfwide HA versus pseudo-forecasted HA. Diagonal line represents perfect prediction.





260


Our forecast quantifies predictive uncertainty associated with the Bayesian model-parameter estimates, variability in riverine and meteorological inputs, and residual errors, summer model inputs, bias adjustment, and transformation of BWDO to HA (Fig. 45, Figs. S5.1–S5.11). The results indicate that the 95% IQR for the west section is on average 2.6 times higher than for the east section, due to greater overall size of the west section (and greater HA) and the complex effect of both river outfalls (Atchafalaya and Mississippi) on BWDO in this section (DMO20). Although the forecasts generally follow the shape of the hindcasts over time, some dissimilarities exist due to hydrometeorological variability (Fig. 45). The pseudo-forecast captures the large HA during summer 1993 that was caused by extremely high May–September MARriver flow and nutrient loadings (Larson, 1997).

265

Interestingly, however, the pseudo-forecast in 2009 overpredicts HA in the west section (two observations are outside of the 95% IQR), due to unusually strong westerly summer winds in this year (Turner et al., 2012). Generally, high wind stress increases water column reaeration and disrupts stratification (Justić and Wang, 2014; Obenour et al., 2015), while upwelling westerly winds disperse the river plume offshore, reducing the consequent oxygen demand (Feng et al., 2012; Le et al., 2016). Overall, only 6.7% (10% (9 of 149) of the observations of total HA are outside of the 95% IQR (Figs. S6.1–S6.8). Also, the geostatistically estimated 95% confidence intervals of observed HA always overlap the forecasted 95% IQR except for one observational cruise in 1988. This discrepancy is caused by anomalously strong summer winds combined with low discharge and nutrient loading in 1988 (Tables S3.1–S3.2). In general, these results suggest the forecasts realistically characterize predictive uncertainties.

Our forecasting approach also allows for the contribution of uncertainty in parameters, data inputs, disentangling and quantifying various sources of forecast variance (Fig. S4.2), residual error to be disentangled. Note that the relative magnitudes of the variance components are somewhat different from the relative magnitudes of the 95% IQR components (e.g., Fig. 4), as the IQR is related to standard deviation (i.e., square root of variance), which is not readily separable (Fig. 5) because variance has squared units. For total HA, the variance associated with stochasticity in riverine and meteorological inputs is 40 times greater than variance associated with parameter uncertainty (on average). On the other hand, the variance associated with residual error is 1.7 times greater than the variance related to the data inputs. Also, the variance associated with these summer data inputs is more influential in later months, as with its portion of the overall forecast contribution to total variance increases from 26% in June–July to 41% in August–September. Furthermore, residual uncertainty is heavily influenced by the remaining sources of forecast variance (dominated by the residual error, but also including the June bias adjustment and transformation of BWDO to HA (Section 2.1), and exhibits higher variance with larger HA (Fig. 5) are 1.9 times greater than the variance related to the stochastic data inputs, suggesting limitations in the model structure or available modelling data (e.g., accuracy or resolution) (S4.2). The relatively low parameter uncertainty reflects the long calibration record available (currently 1985–2016) and is consistent with the underlying model’s robust performance in cross validation (Obenour et al., 2015).





295 **Figure 45:** Daily pseudo-forecasts of HA for the west and east sections in 1993 (top) and 2009 (bottom), including 95%
 IQR of the predictive distribution, distinguishing between i) parameter, ii) riverine and meteorological inputs,
 iii) mechanistic model residual error and iv) regression regressions related to transformation of BWDO to HA and bias
 adjustment uncertainties (shades of gray from darkest to lightest to darkest). Yellow dashed line is hindcasted estimate,
 black dashed line is the 32 year average hindcast, orange points and error bars represent the mean and associated 95%
 300 confidence interval of the (geostatistically estimated) hypoxia observations.

The forecasting uncertainties vary throughout the summer, mostly

Interestingly, the predictive intervals shown in Fig. 5 do not clearly increase over the course of the summer. This
 is largely because predictive intervals tend to increase with increasing predicted hypoxic area, and hypoxic area
 tends to decline after midsummer. If predictive intervals are normalized (i.e., IQR divided by predicted HA), there

305 is a clearer increase in uncertainty over the summer (Fig. 6). The normalized IQRs increase due to the transition
 from observed to randomly sampled model inputs. The riverine and meteorological inputs to DMO20 are lagged
 rolling window averages that include mostly actual observed data (i.e., data prior to 1 June) at the beginning of
 the summer, and an increasing proportion of randomly sampled historical data thereafter (Section 2.1). As a result,
 the normalized IQR (i.e., IQR divided by predicted HA) for pseudo-forecasted HA for June–July is 30% lower
 310 than for August–September on average (Fig. 5A6A, boxplots). Interestingly, there is a rapid increase in
 normalized IQR during the second week of June. In the model, mean water column reaeration is determined by
 wind speeds over the preceding two weeks (Obenour et al., 2015). Consequently, the large and highly variable

wind speeds of June (Fig. 5B6B) (de Velasco and Winant, 1996) quickly increase predictive uncertainties, as these stochastic inputs replace the known wind speed inputs from May. Finally, increases in IQR are also noted in August and September (Fig. 56A), which is consistent with limited ability of the regression analysis (Section 3.1) to accurately project flow and load for such long lead times.




Figure 56: (A) Boxplots represent daily pseudo-forecasted normalized IQR (IQR/HA) including uncertainty due to variability in parameters, riverine inputs and meteorology. Red dashed and solid lines show the daily mean HA and (non-normalized) IQR, respectively; (B) Boxplots show 14 day weighted average squared wind speed near the west shelf section. Boxplots represent interannual variability in the results. The center of each box is the median, while whiskers extend to the extreme value or 1.5 times the IQR of the corresponding variable (whichever is less).

3.3 Sensitivity to riverine and meteorological inputs

The results presented in the previous section are for the proposed forecasting approach with known spring loadings, discharge and winds, and with summer riverine inputs constrained through the regression projections (i.e., Case 4 of Section 2.5). In comparison, the more conventional Gulf forecasting approach, using known spring riverine inputs but with unknown wind and summer riverine inputs (i.e., randomly sampled from the entire historical record, Case 1) explains only around half of variability in hindcasted and observed HA (i.e., 56% and 44%, respectively, Table 2). Also, the performance of Case_1 declines greatly from the beginning (June) to the end (September) of the summer. The inclusion of summer riverine records constrained through regression projections substantially increases the variance explained in both hindcasted and observed HA (Table 2, Cases 3 and 4). This improvement in performance is the most notable for July–September, indicating that the [constraining constrained summer riverine historical records provides inputs provide](#) a more accurate determination of water column stratification and oxygen demand within the biophysical model. Additionally, the summer-wide average of the normalized IQR for Case 4 is on average 22% lower than that of the conventional forecasting approach (Case 1).

340 Addition of actual spring wind data to the conventional approach (Table 2, Case 2) slightly increases the explained
variance in hindcasted and observed HA by 4% (i.e., from 56% to 60%) and 1%, respectively. This forecast
improvement is most notable in June and July because zonal wind velocities up to three months in advance
regulate the transport of water and nutrients over the shelf (Obenour et al., 2015; Walker et al., 2005).
Interestingly, about a quarter of the variability in hindcasted June HA remains unexplained even when actual
spring wind data [isare](#) included (Table 2, Case 2). This is consistent with the importance of near-term wind and
345 discharge inputs in controlling reaeration in the model. It is also consistent with the uncertainties presented in
Fig. [56](#) and other [modelingmodelling](#) studies exploring the influence of wind on hypoxia formation (Forrest et al.,
2011; Yu et al., 2015). Overall, the forecast is only moderately sensitive to the inclusion of actual spring wind
velocities (compare Cases 1 and 2 to Cases 3 and 4). However, we anticipate that inclusion of actual spring wind
data may still be important, especially for years with anomalous wind patterns.

350 Finally, for the preferred forecasting approach (Case 4) we examine an alternative way of determining the relevant
years that constrain the distribution of riverine inputs for the forecast year. If only nitrogen loading regressions
are used to constrain summer inputs (instead of both flow and loading regressions), the explained variability in
the hindcasted total HA drops from 71% (Table 2, Case 4) to 69%. This relatively small drop in predictive
performance is not too surprising, as monthly nitrogen loading, which is the primary driver of many hypoxia
355 models (Turner et al., 2006), is highly correlated with monthly discharge ($r = 0.90$). However, employing the
discharge regressions (in addition to the loading regressions) better accounts for the influence of river flow on
stratification and hypoxia formation (Obenour et al., 2012).

360

Table 2: Explained variance (R^2) in hindcasted and geostatistically observed HA by pseudo-forecasts based on the different data input cases. Here, *act* indicates actual data for the specific year being forecasted are used, *ran-all* indicates input data are randomly sampled from the complete historical record, and *ran-sel* indicates that riverine input data are randomly sampled from a subset of relevant historical years based on the regression projections for flow and loading. Spring includes March–May while summer includes June–September records. The highest R^2 values for each month are highlighted with bold.

Case	Data input				R^2 , Forecasted vs Hindcasted					R^2 , Forecasted vs Observed				
	Riverine		Wind		Month				Overall	Month				Overall
	Spring	Summer	Spring	Summer	Jun	Jul	Aug	Sep		Jun	Jul	Aug	Sep	
1	<i>act</i>	<i>ran-all</i>	<i>ran-all</i>	<i>ran-all</i>	0.69	0.38	0.32	0.21	0.56	0.57	0.28	0.28	0.15	0.44
2	<i>act</i>	<i>ran-all</i>	<i>act</i>	<i>ran-all</i>	0.76	0.56	0.33	0.19	0.60	0.53	0.32	0.28	0.14	0.45
3	<i>act</i>	<i>ran-sel</i>	<i>ran-all</i>	<i>ran-all</i>	0.73	0.68	0.52	0.36	0.68	0.58	0.39	0.32	0.19	0.49
4	<i>act</i>	<i>ran-sel</i>	<i>act</i>	<i>ran-all</i>	0.80	0.73	0.54	0.38	0.71	0.54	0.42	0.34	0.17	0.50

365

3.4 Implications for hypoxia forecasting and fisheries management

To our knowledge, there is only one hypoxia forecasting study (i.e. Laurent and Fennel, 2019), with a similar temporal scope to the current study. That study applied three-dimensional hydrodynamic–biogeochemical model hindcasts, weighted based on comparisons with historical May nitrogen loading only. Other predictive hypoxia studies have employed both discharge and wind (Forrest et al., 2011; Testa et al., 2017), however they lacked the desired temporal resolution of this study. Here, we demonstrate how projections of summer riverine inputs based on spring discharge, loading, and watershed precipitation (Section 3.1) can be used to constrain model inputs, substantially improving hypoxia forecasting skill (Case 4, Table 2). We suggest that other hypoxia forecasting efforts could also benefit from such expanded and projected model inputs.

370

Our approach allows for daily forecasts of BWDO and HA for two shelf sections of the NGoM throughout the entire summer season. Generally, results indicate that HA can be forecasted up to four months ahead, but predictions for later months should be treated with increased caution given their higher uncertainties (Fig. 6). Note that the MAR river input regressions only explain 25–30% of the variability in August nitrogen loadings (Table 1), which is a major factor underlying the decrease in forecast skill in late summer. Additional sources of uncertainty in model inputs may arise from extreme climatic events (e.g. tropical storms and hurricanes), which are unpredictable at the forecasting time scales considered here. These storm events have a multifaceted effect on BWDO, potentially reerating the bottom water column but also providing additional terrestrially derived or sediment-resuspended nutrients that may exacerbate hypoxia (Bianucci et al., 2018; Yu et al., 2015). Therefore, it is not surprising that the pseudo-forecast can explain only about a third of the variability in hindcasted September total HA (Fig. 3). Future4. Overall, future hypoxia forecasting efforts would benefit from improvements in weather and riverine forecasting systems that provide more reliable projections for longer time periods.

375

380

385

The forecasts also explicitly distinguish between different sources of uncertainty in BWDO and HA (Fig. 45). Most previous forecasting studies for the NGoM northern Gulf (Forrest et al., 2011; Scavia et al., 2013; Turner et al., 2012) and other systems like Chesapeake Bay (Testa et al., 2017) implicitly represent uncertainty associated with unknown summer data inputs, sometimes accounting for it in the residual error. On the other hand, the recent

390

Formatted: English (United States)

study by Laurent and Fennel (2019) explicitly considers uncertainty due to stochastic summer riverine inputs, but does not include parameter and residual error uncertainties in the generated forecasts. Thus, this study provides a more comprehensive uncertainty assessment, allowing for management decisions that are robust to potential extremes (Keeney, 1982; Schuwirth et al., 2019).

395 Finally, the proposed approach has potential benefits for short- and long-term environmental planning and fisheries management. In the [NGoMnorthern Gulf](#), the largest volume (Atlantic Menhaden) and highest valued (Penaeid shrimp) fisheries occur during the summer months concurrent with seasonal hypoxia. These fisheries are highly mobile and hypoxia is known to affect the dynamics of both targeted species (Craig and Crowder, 2005; Craig and Bosman, 2013) and fishing fleets (Langseth et al., 2014; Purcell et al., 2017), with potential implications
400 for catch (Craig, 2012), economic condition (Smith et al., 2017), and management (Langseth et al., 2016). However, previous attempts to correlate fishery performance (e.g. catch) with annual measures of hypoxic severity (e.g., area of hypoxia in late July) have had limited success (O'Connor and Whittall, 2007; Zimmerman and Nance, 2001) because neither the spatio-temporal dynamics of hypoxia or of the fishery have been considered. Thus, the proposed daily forecasts can potentially be linked to fisheries and ecosystem models (e.g. de Mutsert et al., 2016),
405 to provide more actionable management guidance. [In addition, while this study focuses on a 1 June forecast release date, consistent with current Gulf forecasting practices, future modelling enhancements might focus on updating the forecast over the summer, as additional hydrometeorological data become available. Such updating could potentially benefit real-time, adaptive management of the fishery.](#)

4 Conclusion

410 In this study, we demonstrate a novel approach for forecasting intra-seasonal variability in BWDO and HA in the [NGoMnorthern Gulf of Mexico](#) by leveraging a Bayesian mechanistic model. This study generates the first daily hypoxia forecasts across the summer season (up to four months ahead) with comprehensive uncertainty assessment. We show that the major sources of uncertainty include variability in data inputs and residual error, while model parameter uncertainty is relatively small. This study also compares how different methods for
415 specifying riverine and meteorological model inputs influence forecast accuracy. In particular, we show how constraining summer riverine inputs based on spring conditions, including precipitation over the [MARMississippi River](#) Basin, can be used to improve hypoxia forecasting skill. We also show that inclusion of monitored spring wind data further improves hypoxia forecasts. Together, these enhancements increase retrospective pseudo-forecast accuracy from 44% to 50% (R^2) while reducing forecast uncertainty by 22% across summers, relative to
420 the conventional approach using spring loadings and flows only (with randomly sampled summer riverine and spring-summer wind records). Thus, the forecasting system developed here provides an enhanced capacity to inform natural resources management in hypoxic coastal systems.

Acknowledgements

425 We thank Dr. Kevin Craig [and anonymous reviewers](#) for [their detailed review of the comments on this manuscript](#). This work was funded by the NOAA grant NA16NOS4780203. [This is NGOMEX contribution #999.](#)

Code/Data Availability

Hydrometeorological and water quality data can be obtained from the public sources described in the methods. A complete set of compiled input data can be provided upon request (akatin@ncsu.edu).

Author Contribution

430 All authors designed the research, analysed the results and prepared the manuscript. Alexey Katin developed the codes and created tables and figures.

Competing interest

The authors declare that they have no conflict of interest.

References

- 435 [Benhamou, E.: A few properties of sample variance, \[online\] Available from: <http://arxiv.org/abs/1809.03774>, 2018.](https://arxiv.org/abs/1809.03774)
- Bianchi, T. S., DiMarco, S. F., Cowan, J. H., Hetland, R. D., Chapman, P., Day, J. W. and Allison, M. A.: The science of hypoxia in the Northern Gulf of Mexico: A review, *Sci. Total Environ.*, 408(7), 1471–1484, doi:10.1016/j.scitotenv.2009.11.047, 2010.
- 440 Bianucci, L., Balaguru, K., Smith, R. W., Leung, L. R. and Moriarty, J. M.: Contribution of hurricane-induced sediment resuspension to coastal oxygen dynamics, *Sci. Rep.*, 8(1), 15740, doi:10.1038/s41598-018-33640-3, 2018.
- Craig, J.: Aggregation on the edge: effects of hypoxia avoidance on the spatial distribution of brown shrimp and demersal fishes in the Northern Gulf of Mexico, *Mar. Ecol. Prog. Ser.*, 445, 75–95, doi:10.3354/meps09437, 2012.
- 445 Craig, J. and Crowder, L.: Hypoxia-induced habitat shifts and energetic consequences in Atlantic croaker and brown shrimp on the Gulf of Mexico shelf, *Mar. Ecol. Prog. Ser.*, 294, 79–94, doi:10.3354/meps294079, 2005.
- Craig, J. K. and Bosman, S. H.: Small Spatial Scale Variation in Fish Assemblage Structure in the Vicinity of the Northwestern Gulf of Mexico Hypoxic Zone, *Estuaries and Coasts*, 36(2), 268–285, doi:10.1007/s12237-012-9577-9, 2013.
- 450 Díaz, R. J. and Rosenberg, R.: Introduction to environmental and economic consequences of hypoxia, *Int. J. Water Resour. Dev.*, 27(1), 71–82, doi:10.1080/07900627.2010.531379, 2011.
- Donner, S. D. and Scavia, D.: How climate controls the flux of nitrogen by the Mississippi River and the development of hypoxia in the Gulf of Mexico, *Limnol. Oceanogr.*, 52(2), 856–861, doi:10.4319/lo.2007.52.2.0856, 2007.
- 455 Faraway, J. J.: *Linear Models with R*, 2nd ed., Taylor & Francis Group., 2015.
- Feng, Y., DiMarco, S. F. and Jackson, G. A.: Relative role of wind forcing and riverine nutrient input on the extent of hypoxia in the northern Gulf of Mexico, *Geophys. Res. Lett.*, 39(9), n/a–n/a, doi:10.1029/2012GL051192, 2012.
- Fennel, K., Hu, J., Laurent, A., Marta-Almeida, M. and Hetland, R.: Sensitivity of hypoxia predictions for the northern Gulf of Mexico to sediment oxygen consumption and model nesting, *J. Geophys. Res. Ocean.*, 118(2), 460 990–1002, doi:10.1002/jgrc.20077, 2013.

Fennel, K., Laurent, A., Hetland, R., Justić, D., Ko, D. S., Lehrter, J., Murrell, M., Wang, L., Yu, L. and Zhang, W.: Effects of model physics on hypoxia simulations for the northern Gulf of Mexico: A model intercomparison, *J. Geophys. Res. Ocean.*, 121(8), 5731–5750, doi:10.1002/2015JC011577, 2016.

465 Forrest, D. R., Hetland, R. D. and DiMarco, S. F.: Multivariable statistical regression models of the areal extent of hypoxia over the Texas–Louisiana continental shelf, *Environ. Res. Lett.*, 6(4), 045002, doi:10.1088/1748-9326/6/4/045002, 2011.

Gentry, L. E., David, M. B. and McIsaac, G. F.: Variation in Riverine Nitrate Flux and Fall Nitrogen Fertilizer Application in East-Central Illinois, *J. Environ. Qual.*, 43(4), 1467–1474, doi:10.2134/jeq2013.12.0499, 2014.

470 Del Giudice, D., Matli, V. R. R. and Obenour, D. R.: Bayesian mechanistic modeling characterizes Gulf of Mexico hypoxia: 1968–2016 and future scenarios, *Ecol. Appl.*, 30(2), eap.2032, doi:10.1002/eap.2032, 2020.

Greene, R. M., Lehrter, J. C. and III, J. D. H.: Multiple regression models for hindcasting and forecasting midsummer hypoxia in the Gulf of Mexico, *Ecol. Appl.*, 19(5), 1161–1175, doi:10.1890/08-0035.1, 2009.

Hart, E. M. and Bell, K.: prism: Download data from the Oregon prism project, , doi:10.5281/zenodo.33663, 2015.

475 Hetland, R. D.: Relating river plume structure to vertical mixing, *J. Phys. Oceanogr.*, 35(9), 1667–1688, doi:10.1175/JPO2774.1, 2005.

Hinsby, K., Markager, S., Kronvang, B., Windolf, J., Sonnenborg, T. O. and Thorling, L.: Threshold values and management options for nutrients in a catchment of a temperate estuary with poor ecological status, *Hydrol. Earth Syst. Sci.*, 16(8), 2663–2683, doi:10.5194/hess-16-2663-2012, 2012.

480 Justić, D. and Rose, K. A.: Modeling Coastal Hypoxia, edited by D. Justic, K. A. Rose, R. D. Hetland, and K. Fennel, Springer International Publishing, Cham., 2017.

Justić, D. and Wang, L.: Assessing temporal and spatial variability of hypoxia over the inner Louisiana–upper Texas shelf: Application of an unstructured-grid three-dimensional coupled hydrodynamic–water quality model, *Cont. Shelf Res.*, 72, 163–179, doi:10.1016/j.csr.2013.08.006, 2014.

485 Justić, D., Rabalais, N. N. and Turner, R. E.: Effects of climate change on hypoxia in coastal waters: A doubled CO₂ scenario for the northern Gulf of Mexico, *Limnol. Oceanogr.*, 41(5), 992–1003, doi:10.4319/lo.1996.41.5.0992, 1996.

Keeney, R. L.: Decision Analysis: An Overview, *OPER RES*, V 30(N 5), 803–838, doi:10.1287/opre.30.5.803, 1982.

490 Krug, E. C.: Coastal change and hypoxia in the northern Gulf of Mexico: Part I, *Hydrol. Earth Syst. Sci.*, 11(1), 180–190, doi:10.5194/hess-11-180-2007, 2007.

Langseth, B. J., Purcell, K. M., Craig, J. K., Schueller, A. M., Smith, J. W., Shertzer, K. W., Creekmore, S., Rose, K. A. and Fennel, K.: Effect of Changes in Dissolved Oxygen Concentrations on the Spatial Dynamics of the Gulf Menhaden Fishery in the Northern Gulf of Mexico, *Mar. Coast. Fish.*, 6(1), 223–234, doi:10.1080/19425120.2014.949017, 2014.

495 Langseth, B. J., Schueller, A. M., Shertzer, K. W., Craig, J. K. and Smith, J. W.: Management implications of temporally and spatially varying catchability for the Gulf of Mexico menhaden fishery, *Fish. Res.*, 181, 186–197, doi:10.1016/j.fishres.2016.04.013, 2016.

Larson, L. W.: The great USA flood of 1993, *IAHS Publ. Proc. Reports-Intern Assoc Hydrol. Sci.*, 239, 13–29 [online] Available from: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.595.3792&rep=rep1&type=pdf>, 1997.

500

- Laurent, A. and Fennel, K.: Time-Evolving, Spatially Explicit Forecasts of the Northern Gulf of Mexico Hypoxic Zone, *Environ. Sci. Technol.*, 53(24), 14449–14458, doi:10.1021/acs.est.9b05790, 2019.
- Le, C., Lehrter, J. C., Hu, C. and Obenour, D. R.: Satellite-based empirical models linking river plume dynamics with hypoxic area and volume, *Geophys. Res. Lett.*, 43(6), 2693–2699, doi:10.1002/2015GL067521, 2016.
- 505 Lumley, T.: leaps: Regression Subset Selection. Based on Fortran code by Alan Miller, [online] Available from: <https://cran.r-project.org/package=leaps>, 2017.
- Matli, V. R. R., Fang, S., Guinness, J., Rabalais, N. N., Craig, J. K. and Obenour, D. R.: Space-Time Geostatistical Assessment of Hypoxia in the Northern Gulf of Mexico, *Environ. Sci. Technol.*, 52(21), 12484–12493, doi:10.1021/acs.est.8b03474, 2018.
- 510 [Mattern, J. P., Fennel, K. and Dowd, M.: Sensitivity and uncertainty analysis of model hypoxia estimates for the Texas-Louisiana shelf, *J. Geophys. Res. Ocean.*, 118\(3\), 1316–1332, doi:10.1002/jgrc.20130, 2013.](#)
- Ménesguen, A. and Lacroix, G.: Modelling the marine eutrophication: A review, *Sci. Total Environ.*, 636, 339–354, doi:10.1016/j.scitotenv.2018.04.183, 2018.
- de Mutsert, K., Steenbeek, J., Lewis, K., Buszowski, J., Cowan, J. H. and Christensen, V.: Exploring effects of hypoxia on fish and fisheries in the northern Gulf of Mexico using a dynamic spatially explicit ecosystem model, *Ecol. Modell.*, 331, 142–150, doi:10.1016/j.ecolmodel.2015.10.013, 2016.
- 515 NDBC: National Data Buoy Center, [online] Available from: <https://www.ndbc.noaa.gov/> (Accessed 5 May 2019), 2019.
- O'Connor, T. and Whittall, D.: Linking hypoxia to shrimp catch in the northern Gulf of Mexico, *Mar. Pollut. Bull.*, 54(4), 460–463, doi:10.1016/j.marpolbul.2007.01.017, 2007.
- Obenour, D. R., Michalak, A. M., Zhou, Y. and Scavia, D.: Quantifying the Impacts of Stratification and Nutrient Loading on Hypoxia in the Northern Gulf of Mexico, *Environ. Sci. Technol.*, 46(10), 5489–5496, doi:10.1021/es204481a, 2012.
- Obenour, D. R., Michalak, A. M. and Scavia, D.: Assessing biophysical controls on Gulf of Mexico hypoxia through probabilistic modeling, *Ecol. Appl.*, 25(2), 492–505, doi:10.1890/13-2257.1, 2015.
- 525 Purcell, K. M., Craig, J. K., Nance, J. M., Smith, M. D. and Benneer, L. S.: Fleet behavior is responsive to a large-scale environmental disturbance: Hypoxia effects on the spatial dynamics of the northern Gulf of Mexico shrimp fishery, edited by J. Xu, *PLoS One*, 12(8), e0183032, doi:10.1371/journal.pone.0183032, 2017.
- [R Core Team: R: A Language and Environment for Statistical Computing, \[online\] Available from: <https://www.r-project.org/>, 2019.](#)
- 530 Rabalais, N. N. and Turner, R. E.: Gulf of Mexico Hypoxia: Past, Present, and Future, *Limnol. Oceanogr. Bull.*, 28(4), 117–124, doi:10.1002/lob.10351, 2019.
- Rabalais, N. N., Turner, R. E. and Scavia, D.: Beyond science into policy: Gulf of Mexico hypoxia and the Mississippi River, *Bioscience*, 52(2), 129–142, doi:10.1641/0006-3568(2002)052[0129:BSIPGO]2.0.CO;2, 535 2002.
- Reichert, P. and Borsuk, M. E.: Does high forecast uncertainty preclude effective decision support?, in *Environmental Modelling and Software*, vol. 20, pp. 991–1001, Elsevier., 2005.
- Scavia, D., Evans, M. A. and Obenour, D. R.: A Scenario and Forecast Model for Gulf of Mexico Hypoxic Area and Volume, *Environ. Sci. Technol.*, 47(18), 10423–10428, doi:10.1021/es4025035, 2013.
- 540 Scavia, D., Bertani, I., Obenour, D. R., Turner, R. E., Forrest, D. R. and Katin, A.: Ensemble modeling informs

- hypoxia management in the northern Gulf of Mexico, *Proc. Natl. Acad. Sci.*, 114(33), 8823–8828, doi:10.1073/pnas.1705293114, 2017.
- Schuwirth, N., Borgwardt, F., Domisch, S., Friedrichs, M., Kattwinkel, M., Kneis, D., Kuemmerlen, M., Langhans, S. D., Martínez-López, J. and Vermeiren, P.: How to make ecological models useful for environmental management, *Ecol. Modell.*, 411, 108784, doi:10.1016/j.ecolmodel.2019.108784, 2019.
- 545 Schwartz, M.: Boundary of Mississippi River Basin derived by dissolving WBD HU-4, [online] Available from: <https://www.sciencebase.gov/catalog/item/55de04d5e4b0518e354dfcf8>, 2012.
- Sinha, E. and Michalak, A. M.: Precipitation dominates interannual variability of riverine nitrogen loading across the continental United States, *Environ. Sci. Technol.*, 50(23), 12874–12884, doi:10.1021/acs.est.6b04455, 2016.
- 550 Smith, M. D., Asche, F., Benneer, L. S. and Oglend, A.: Spatial-dynamics of Hypoxia and Fisheries: The Case of Gulf of Mexico Brown Shrimp, *Mar. Resour. Econ.*, 29(2), 111–131, doi:10.1086/676826, 2014.
- Smith, M. D., Oglend, A., Kirkpatrick, A. J., Asche, F., Benneer, L. S., Craig, J. K. and Nance, J. M.: Seafood prices reveal impacts of a major ecological disturbance, *Proc. Natl. Acad. Sci.*, 114(7), 1512–1517, doi:10.1073/pnas.1617948114, 2017.
- 555 Testa, J. M., Clark, J. B., Dennison, W. C., Donovan, E. C., Fisher, A. W., Ni, W., Parker, M., Scavia, D., Spitzer, S. E., Waldrop, A. M., Vargas, V. M. D. and Ziegler, G.: Ecological Forecasting and the Science of Hypoxia in Chesapeake Bay, *Bioscience*, 67(7), 614–626, doi:10.1093/biosci/bix048, 2017.
- Thronson, A. and Quigg, A.: Fifty-Five Years of Fish Kills in Coastal Texas, *Estuaries and Coasts*, 31(4), 802–813, doi:10.1007/s12237-008-9056-5, 2008.
- 560 Turner, R. E., Rabalais, N. N. and Justic, D.: Predicting summer hypoxia in the northern Gulf of Mexico: Riverine N, P, and Si loading, *Mar. Pollut. Bull.*, 52(2), 139–148, doi:10.1016/j.marpolbul.2005.08.012, 2006.
- Turner, R. E., Rabalais, N. N. and Justic, D.: Gulf of Mexico Hypoxia: Alternate States and a Legacy, *Environ. Sci. Technol.*, 42(7), 2323–2327, doi:10.1021/es071617k, 2008.
- Turner, R. E., Rabalais, N. N. and Justic, D.: Predicting summer hypoxia in the northern Gulf of Mexico: Redux, *Mar. Pollut. Bull.*, 64(2), 319–324, doi:10.1016/j.marpolbul.2011.11.008, 2012.
- 565 USACE: Atchafalaya and Mississippi River daily discharge, New Orleans Dist. Website [online] Available from: <https://www.mvn.usace.army.mil/Missions/Engineering/Stage-and-Hydrologic-Data/Historical-Discharges/> (Accessed 5 May 2019), 2019.
- USGS: Streamflow and Nutrient Delivery to the Gulf of Mexico, [online] Available from: https://toxics.usgs.gov/hypoxia/mississippi/flux_estimates/delivery/index.html (Accessed 5 May 2019), 2019.
- de Velasco, G. G. and Winant, C. D.: Seasonal patterns of wind stress and wind stress curl over the Gulf of Mexico, *J. Geophys. Res. Ocean.*, 101(C8), 18127–18140, doi:10.1029/96JC01442, 1996.
- Walker, N. D., Wiseman, W. J., Rouse, L. J. and Babin, A.: Effects of River Discharge, Wind Stress, and Slope Eddies on Circulation and the Satellite-Observed Structure of the Mississippi River Plume, *J. Coast. Res.*, 216(6), 1228–1244, doi:10.2112/04-0347.1, 2005.
- 575 Yu, L., Fennel, K. and Laurent, A.: A modeling study of physical controls on hypoxia generation in the northern Gulf of Mexico, *J. Geophys. Res. Ocean.*, 120(7), 5019–5039, doi:10.1002/2014JC010634, 2015.
- Zhang, F., Sun, Y. Q., Magnusson, L., Buizza, R., Lin, S.-J., Chen, J.-H. and Emanuel, K.: What Is the Predictability Limit of Midlatitude Weather?, *J. Atmos. Sci.*, 76(4), 1077–1091, doi:10.1175/JAS-D-18-0269.1, 2019.
- 580

Zimmerman, R. J. and Nance, J. M.: Effects of hypoxia on the shrimp fishery of Louisiana and Texas, pp. 293–310., 2001.

Formatted: Normal