

Spatial extrapolation of stream thermal peaks using heterogeneous time series at a national scale

Aurelien Beaufort^{1,2}, Jacob S. Diamond^{1,2}, Eric Sauquet¹, Florentina Moatar^{1*}

¹RiverLy, INRAE, Centre de Lyon-Grenoble Auvergne-Rhône-Alpes, France 69100

²Université de Tours, GéoHydrosystèmes COntinentaux, Tours, France 38000

Correspondence to: Florentina Moatar (florentina.moatar@inrae.fr)

Abstract. Spatial reconstruction of stream temperature is relevant to water quality standards and fisheries management, yet large, regional scale datasets are rare because data are decentralized and inharmonious. This data discordance is a major limitation for understanding thermal regimes of riverine ecosystems. To overcome this barrier, we first aggregated one of the largest stream temperature databases on record with data from 1700 individual stations over nine years from 2009–2017 (n=45,000,000 hourly measurements) across France (area = 552,000 km²). For each station, we calculated a simple, ecologically relevant metric—the thermal peak—that captures the magnitude of summer thermal extremes. We then used three statistical models to extrapolate the thermal peak to nearly every stream reach in France and Corsica (n=105,800) and compared relative model performances ~~among each other and~~ with an air temperature metric. In general, the hottest thermal peaks were found along major rivers, whereas the coldest thermal peaks were found along small rivers with forested riparian zones, strong groundwater inputs, and which were located in mountainous regions. Several key predictors of the thermal peak emerged, including drainage area, mean summer air temperature, minimum monthly specific discharge, and vegetation cover in the riparian zone. Despite differing predictor importance across model structures, we observed strong concordance among models in their spatial distributions of the thermal peak, suggesting its robustness as a useful metric at the regional scale. Finally, air temperature was found ~~asto be~~ a poor proxy for the stream temperature thermal peak across nearly all stations and reaches, highlighting the growing need to measure and account for stream temperature in regional ecological studies.

1 Introduction

Stream water temperature ~~is a master variable affecting ecosystem processes in lotic systems. It controls the solubility rates of gases and related biogeochemical reactions, regulates metabolism (processes (Nimick et al., 2011; Song et al. 2018) and the distribution and phenology of aquatic communities (Hawkins et al. 1997; Wolter, 2007), nutrient cycling (Malard et al., 2002) and decomposition rates, and dictates animal ingestion and digestion rates (Elliott, 1976), reproduction cycles (Daufresne et al., 2004) and mobility (Ojanguren and Braña, 2000). Stream).~~ Water temperature ~~can also be a source of stress and mortality for aquatic organisms, especially when coupled to additional stressors~~ extremes, like low water levels (Miller) summer maxima, are particularly strong drivers of aquatic biota behavior and habitat selection (Carlson et al., 2007). ~~Consequently, the dynamics of populations and communities, their relative composition (Kishi et al., 2005) and their size structure (Daufresne; Xu et al., 2009) are intimately related to stream thermal regimes.~~

35 ~~The five main components of stream thermal regimes comprise temperature magnitude, frequency, duration, rate of change, and timing, with different metrics to quantify the biological or ecological (2010). Despite their importance of each component (Steel et al., 2017; Olden and Naiman, 2010; Tsang et al., 2016). Importantly, these metrics can be accurately determined only if continuous time series of, we lack an understanding of the spatial distributions of thermal extremes, largely due to a paucity of consistent stream temperature data.~~

40 ~~There is thus a need to develop and improve spatially distributed stream temperature are available (Jones and Schmidt, 2018). Moreover, while each of these metrics can provide complementary information about stream thermal regimes, many of them are strongly correlated (Ashley Steel et al., 2016; Rivers Moore et al., datasets with a focus on identifying areas most affected by thermal extremes (2013; Isaak et al., 2020) and their complete description may be unnecessary to understand critical temperature effects on ecosystems. Indeed, the collinearity of these metrics suggests the utility of a single metric to understand extreme stream temperature regimes, especially a metric that does not require continuous datasets for its calculation. Such a simple metric would allow for rapid spatial comparison that may help managers understand which rivers currently exceed thermal tolerances of important biota.~~

45 ~~Although recent National- and even regional-scale stream temperature data are a relatively recent phenomenon (Jackson et al., 2018), in part due to advancements in *in-situ* sensor technology have greatly expanded stream temperature data availability (Isaak et al., 2017) there is still a lack of long-term data for the vast majority of stream reaches, limiting understanding of thermal regimes at river network and regional scales (sensors (Isaak et al., 2017), but these Arismendi et al., 2014). In addition, stream temperature data collection is oftentend to be spatiotemporally uncoordinated across regions and river networks (Arismendi et al., 2014).~~

50 ~~resultingThis results in snapshot datasets that are regularly supported or replaced with, necessitating the use of air temperature proxies in aquatic ecology studies to supplement or even replace stream temperature time-series (Conti et al., 2015; Logez et al., 2012; Tisseuil et al., 2012). Indeed, ecologists regularly use air temperature as a proxy or predictor of stream temperature (Buisson and Grenouillet, 2009; Durance and Ormerod, 2009; Mayer 2012) despite the fact that water temperature better explains the spatial organization of aquatic biota (Kirk and Rahel, 2022). While air temperature can be used to fill temporal data gaps (Buisson and Grenouillet, 2009; Durance and Ormerod, 2009Morrill et al. 2005), air temperature is a poor surrogate for stream temperature in several cases. In particular, air temperature correlates poorly with stream temperature in headwater reaches (Caissie, 2006), in reaches with strong local controls (e.g., riparian vegetation, groundwater inflows, bed form, impoundments), and in reaches with large environmental heterogeneity (Moatar and Gailhard, 2006; Hill and Hawkins, 2014; Loicq et al., 2018; Chandesris et al., 2019; Seyedhashemi et al., 2021). Hence, usingwhile air temperature metrics to study climate change impacts onhas clear utility for understanding stream temperature, the aquatic species distributions may result in misguided inference and consequent management decisionsspatial distribution of thermal extremes will be modulated by other variables.~~

55 ~~The spatial variation of summer stream temperature is controlled by several factors, each with varying spatiotemporal influence (Fullerton et al., 2015). The most commonly observed factor is stream size-~~

60

65

70

often evaluated with a drainage area or distance-from-source proxy—which tends to have a positive relationship with stream thermal maxima (Hrachowitz et al., 2010; Imholt et al., 2013; Isaak et al., 2017). Instead of using air temperature proxies, many studies have turned to regionalized deterministic and statistical models to fill stream temperature data gaps (e.g. Mohseni et al., 1998; Segura et al., 2015; Chang and Pсарis, 2013; Beaufort et al., 2016; Westhoff et al., 2007; Yearsley, 2012). Deterministic models rely on a physically based formulation of stream energy conservation to compute water temperature (Yearsley, 2012; van Vliet et al., 2012; Beaufort et al., 2016). However, such models suffer from large data requirements leading to a preference for statistical approaches, especially in ungauged catchments. Some of these approaches empirically relate stream temperature to climatic and environmental variables, such as air temperature, discharge, altitude or channel width (Benyahya et al., 2007; Moore et al., 2013). Other statistical methods include lumped regression-based models (Daigle et al., 2010; Ducharme, 2008; Hrachowitz et al., 2010; Bustillo et al., 2014), distributed stream network models (Detenbeck et al., 2016; Isaak et al., 2017), and machine learning methods (Chenard and Caissie, 2008; DeWeber and Wagner, 2014). Still, these approaches often have dense spatial data requirements and their estimates are usually temporally limited (Isaak et al., 2010; Pratt and Chang, 2012; Hill et al., 2013). Indeed, few studies estimate stream temperature over a full year, likely because of non-linear relationships and seasonal hysteresis between air and stream temperature, missing data, and autocorrelation (Jackson et al., 2018; Letcher et al., 2016; Sohrabi et al., 2017). However, for management purposes in the context of climate change, annual stream temperature patterns may be unnecessary. Indeed, stream temperature metrics that focus on extreme periods (e.g., summer) are likely adequate to understand trends of increasing pressures on aquatic ecosystems.

To that end (Isaak et al., 2017). As streams increase in size, their surface area increases leading to rapid heat transfer and subsequent equilibration with air temperatures. Intuitively, discharge is also a strong control on stream temperature distributions, with increasing groundwater influence (e.g., measured with monthly flow minima or the base flow index [BFI]) often cited as a mitigating effect on thermal extremes (Chang and Pсарis, 2013; Hare et al. 2021). An obvious candidate factor is summer air temperature, which exhibits clear positive relationships with stream thermal maxima, albeit at regional scales (Moore et al., 2013; Isaak et al., 2017). In addition to these three dominant factors, an array of additional controls like local geology, slope, altitude, and riparian cover can modulate these relationships, leading to unexpected spatial distributions of stream water temperature (Fullerton et al., 2015).

To account for the variety of controls on stream temperature spatial distributions in a data scarce landscape, researchers turn to predictive models. Still, despite a recent growth in modeling techniques, including statistical (Segura et al. 2015, Benyahya et al. 2007) and physical (Beaufort et al. 2016a,b) approaches, models that can capture regional or national-scale stream temperature distributions remain scarce (but see Hare et al. 2021). At these larger spatial scales, statistical models, which relate stream temperature to climatic and environmental variables, are likely the most pertinent and parsimonious for management. Indeed, geostatistical river network models that can incorporate network covariance structure are likely the most appropriate and accurate models to estimate spatial distributions of stream temperature (Isaak et al., 2017), but their use is limited to regions with high data density. As most countries lack such the required data for such approaches, researchers are more prone to employ

110 empirical regressions or machine learning approaches, but the comparative accuracy and utility of these approaches is still unknown.

In light of these gaps in data and understanding, our objectives are twofold: 1) to ~~create of a harmonious~~aggregate and develop a homogeneous national-scale stream temperature database for France, and 2) to develop empirical statistical models to predict a simple, ecologically relevant stream thermal metric that captures the magnitude of the stream temperature extremes at the regional scale. To do so, we first define an interannual thermal metric, which we term the “thermal peak”, using heterogeneous and non-concomitant time series of stream temperature and estimate it at 1700 stations for 2009–2017. We then test three statistical models and one multi-model approach to predict this metric at the regional scale and compare their predictive capacity with that of air temperature. Specifically, we used these models to address the following question: what are the spatial patterns of stream temperature extremes in France and their drivers, and are these patterns consistent across modeling approaches? We hypothesized that spatial patterns would be consistent, whereas the drivers would depend on the modeling approach used. We also hypothesized that stream size, air temperature, and groundwater contributions would emerge as important regardless of approach.

125 2. Methods

2.1 Study area and monitoring network

The study area is continental France and Corsica (550,000 km²). France is located in a temperate zone characterized by a variety of climates due to the influences of the Atlantic Ocean, the Mediterranean Sea, and mountain areas. We assembled ~~the most exhaustive a large~~ stream temperature dataset for ~~our study this area to date~~ by combining data from both public (national, regional; approximately 600 stations available from <http://www.naiades.eaufrance.fr/>), fishing associations and other private and public sources sub-national agencies. Due to the diversity of station ownership data origins, the assembled times series ~~did~~ not have a consistent spatial and temporal structure. Some regions are densely monitored while others ~~have had~~ few instrumented streams and much of the data do not exhibit temporal overlap, challenging our ability to define comparable metrics among streams. Hence, our main challenge was to coalesce and harmonize/homogenize all the disparate data sources.

The stream temperature data used here comprise approximately 45,000,000 hourly measurements from 1,700 unique measurement stations (n=2,107,623 site-days) collected between 2009 and 2017, primarily during summer. All the stations under strong human influence (i.e., dam releases and nuclear power thermal effluent) ~~and stations without seasonal dynamics have been were~~ excluded from this data set. All data were recorded by automatic data loggers managed by professional biologists, hydrologists, and fishermen. Outliers from each station’s time series were removed with automatic outlier detection filters and the resulting hourly data were screened visually before being averaged into daily mean stream temperature data (hereafter referred to as T_w). The automatic outlier detection consists of three steps with eight unique filters that remove, in order, 1) stream temperature anomalies based on hourly data, 2) anomalies between T_w and daily mean air temperature (T_{air}) at monthly scales, and 3) anomalies between T_w and T_{air} at daily scales (Table 1; Moatar et al., 2001; Beaufort et al., 2020b).

Table 1. Outlier detection and data filtering process of stream temperature dataset

Filter type	Definition	Threshold (°C)	Data removed (%)
Stream temperature anomalies from hourly data	Maximum temperature > threshold (by month)	14, 15, 20, 24, 28, 30, 32, 33, 29, 28, 18, 17 (Jan.–Dec.)	
	Minimum temperature < threshold	-0.5	
	Difference in consecutive data > threshold	2	2
	Daily diel range > threshold	7	
	Difference in daily max. or min. in consecutive data > threshold	3	
Monthly $T_w - T_{air}$ anomalies	R^2 of daily regressions by month < threshold	0.1 [unitless]	
	Deviation of the monthly difference ($T_w - T_{air}$) to its interannual mean > threshold	4	
Daily $T_w - T_{air}$ anomalies	Daily difference ($T_w - T_{air}$) > the monthly mean of daily differences by threshold	2.5	5

To understand spatial patterns in ecologically meaningful temperature metrics, we focused on the two hottest stream temperature months, July and August (hereafter referred to as summer), a time period essential for the growth and survival of many aquatic species. This focus also ~~hashad~~ the benefit of maximizing the number of observation stations for analysis. Still, out of the 1700 stations, 490 stations ~~havehad~~ just one year of data, 88 stations ~~havehad~~ observations covering summer over all nine years, and only 30 ~~havehad~~ year-round observations for all nine years (Figure 1a,b). To obtain hydraulic and hydrologic characteristics for each station, stations were projected onto the Theoretical Hydrographic Network for France (RHT; Pella et al., 2012), an oriented hydrographic network with defined flow directions that ~~comprised~~ 114,600 reaches of median length 1,961 m ($2,475 \pm 1,512$ m, mean \pm sd). A majority of stations were located on RHT river reaches with drainage areas 20–500 km², whereas most reaches are small streams with a drainage area of less than 20 km² (Figure 1c).




Figure 1. Data availability for each temperature station used in this study. a) Map of stream temperature stations in France with RHT network shown for all reaches of Strahler order >4, b) Heatmap of data availability by year (x-axis) and station (y-axis) with the total stations per year listed at the top of each column. Sites are colored by the number of years with observations, and c) Distributions of drainage area for RHT reaches (blue) and of thermal stations (black).

2.2 Defining the thermal peak metric

Due to the limited concordance among stream temperature time series (Figure 1b) and to focus our analysis towards ecologically relevant ends, we summarized stream temperature data with a simple metric, the thermal peak. This metric has precedent in regional species distribution models that instead used air temperature (hereafter referred to as T_{air}) as a proxy (Buisson and Grenouillet, 2009). We refer to this metric as the thermal peak (T_p), and define it as the interannual average of the mean temperature of the 30 hottest consecutive days of each year ($\overline{T_{w,30}}_i$):

$$T_p = \frac{\sum_{i=1}^N (\overline{T_{w,30}})_i}{N} \quad (1)$$

170 where:

i = year index; and

N = the number of years of observation available from 2009–2017 (N = 1–9)

We did not know *a priori* the 30 hottest consecutive days of each year, but a sensitivity analysis on the sites with annual data suggests that July and August were regularly the hottest months (Fig: A1). Indeed, across sites with annual data, the hottest day of the year always occurred within the approximate 30 day period between July 28 and August 30 (mean±sd: August 12±16 days), and only 3% of site-years had their hottest 30-days outside of this period. This fact further allowed us to take advantage of many of the sub-annual time series, particularly those generated by fishing agencies, which only contain July and August data.

2.3 Climate correction of the thermal peak

180 The thermal peak can be biased depending on the climatic variability of the years of observation for each station. Indeed, only 30 of our stations ~~have had~~ T_p calculated using all nine years of data, and therefore ~~have had~~ the highest level of confidence in their estimate. We refer to the T_p from these 30 stations as $T_{p,\text{ref}}$, indicating that these are reference, or true estimates of T_p (Table 2). To account for the bias associated with missing data at the remaining 1670 stations, we gap-filled missing data at these stations using site-specific stream-air temperature regressions. 185 This method accounts for interannual variation in climatic forcing on stream temperature, and we therefore refer to it as a climate correction.

The climate correction is achieved by first calculating station-specific regressions during summer between daily T_w and a right-aligned moving average of T_{air} at lags ranging from 2–10 days. The moving average lag whose regression produced the highest coefficient of determination was then used to fill gaps in the time series of T_w for each station. Stations with watershed areas greater than 1000 km² tended to have the best R² at the longest lags, but there were no clear trends at smaller watershed areas (Fig. A2) Next, we reconstructed summer stream temperature using this regression and subsequently recalculated T_p based on this reconstructed data. We refer to T_p from these climate-corrected data as $T_{p,\text{clim}}$ to indicate that missing data were gap-filled with the climate correction procedure (Table 2).

195 To validate this approach, we conducted a permutation test on the 30 stations with full annual monitoring from 2009–2017 (i.e., sites where a $T_{p,ref}$ is known). At each site, we introduced randomly placed, artificial annual gaps into observed data ranging from 1–9 years to simulate missing data. We then backfilled these introduced gaps according to the climate correction method. Following gap-filling, we calculated two metrics: 1) T_p using gap-induced data without gap-filling ($T_{p,gap}$), and 2) the thermal peak using the gap-filled data ($T_{p,fill}$; Table 2). We then
 200 compare these two metrics to the reference thermal peak ($T_{p,ref}$) using absolute biases at each tested permutation (i.e., the number of introduced gap years). This approach allowed us to assess whether the climate-corrected reconstruction of the gaps in time series is 1) a useful approach, and 2) lower in bias and uncertainty compared to using observed data alone.

Table 2. List of thermal peak terminology with the count of [dayssites](#) (n) used in their calculation

Notation	Definition	n
$T_{p,obs}$	thermal peak, or the interannual average of the mean temperature of the 30 hottest consecutive days of each year for station with observations	1700
$T_{p,ref}$	thermal peak for reference stations with all nine years of data	30
$T_{p,clim}$	thermal peak for stations with less than nine years of data to which climate correction was applied	1670
$T_{p,gap}$	thermal peak for reference stations with introduced data gaps	30
$T_{p,fill}$	thermal peak for reference stations whose introduced data gaps were filled with climate correction	30
$T_{p,m}$	modeled thermal peak using statistical extrapolation to the RHT network using statistical method m	114,600
$T_{p,air}$	thermal peak estimated using the SAFRAN reanalysis T_{air} data	114,600

2.4 Extrapolating the thermal peak to national scale with statistical modeling

205 We estimated T_p throughout the entire RHT network using four distinct statistical models. For modeling ($T_{p,m}$) it was not clear how to choose *a priori* a particular model structure due to the complexity of the processes involved in determining local stream temperature. Therefore, we tested four different structures: 1) a multiple linear regression model (REG), 2) an artificial neural network model (ANN) that is potentially non-linear, but encompasses a linear model as a special case, 3) a random forest model (RF) that can also be non-linear, but with
 210 a different strategy than ANN, and 4) a multi-model combination (MM), which combines the three prior structures ANN, RF, and REG. All these models are based on a function to estimate T_p at all stations (i):

$$T_{p,m,i} = f(g_{1,i}, \dots, g_{n,i}) \quad (5)$$

Where $g_{n,i}$ = the nth explanatory variable defined for each ith station.

To measure how stream thermal regime estimations might be different if using a T_{air} proxy, we also calculated
 215 $T_{p,air}$ using T_{air} from the Système d'Analyse Fournissant des Renseignements Atmosphériques à la Neige dataset (SAFRAN). SAFRAN is a mesoscale atmospheric analysis system for surface variables with reanalysis gridded data at hourly time steps with a resolution of 8 km using ground data observations ([Durand et al. 1993](#); Vidal et al., 2010). Climatic variables including T_{air} were extracted from the SAFRAN meshes overlapping the station location. The $T_{p,air}$ is calculated identically to T_p , but using daily T_{air} instead of daily T_w , as would be done in
 220 ecological studies using T_{air} as a proxy for T_w . Finally, to determine the effect of climate corrections on the full RHT extrapolation, we compared the distribution of $T_{p,m}$ values when models were fit using either $T_{p,obs}$ or $T_{p,clim}$.

2.4.1 Explanatory variables

We selected sixteen variables (Table 3) to explain the spatial distribution of $T_{p,m}$ based both on results from a prior analysis (Beaufort et al., 2019, 2020a), review of the literature and an effort to minimize variable collinearity (Fig. A3). We further considered the ability to calculate or estimate each variable at the scale of the entire RHT network. The variables fall into three categories: climate, hydrology, and catchment watershed characteristics.

The four climatic variables were determined from SAFRAN reanalysis data for the years 2009–2017: 1) mean annual precipitation, 2) mean summer precipitation, 3) mean annual snowfall, and 4) mean summer air temperature.

The four hydrological variables were determined by extrapolation based on prior datasets. In general, we wanted to quantify the influence of groundwater on T_p , but lacked the local hydrometric data necessary to calculate the typically used BFI (Hare et al. 2021) at a majority of sites. Hence, we turned to the following variables that could potentially quantify groundwater influence across all of our sites. The first two variables, monthly minimum discharge (Sauquet et al. 2008) and the annual minimum monthly discharge with a return period of five years (Catalogne, 2012), describe the low-flow regime of each site. The remaining two variables, the hydrologic regime (HR; Sauquet et al., 2008) and the concavity index (CI; Sauquet and Catalogne, 2011), are dimensionless and characterize the general hydrology of each site. More specifically, the HR groups sites into one of 12 classes ranging from rainfall-dominated, to transitional, to glacial and snow melt dominated. These classes generally fall into buffered (i.e., high baseflow) or highly variable (i.e., low baseflow) hydrologic regimes. The CI describes the concavity of the flow duration curve, where values close to 1 indicate low flow variability (e.g., large high storage capacity in aquifer or snow) and values close to 0 indicate high flow variability (e.g., low storage capacity exemplary of Mediterranean systems).

The eight variables relating to catchment watershed characteristics were extracted from either the SYRAH-CE database (Valette et al., 2012) or the RHT database (Pella et al., 2012). The three variables from RHT comprise: 1) mean altitude, 2) catchment watershed drainage area, and 3) mean slope. The five variables from SYRAH-CE comprise: 1) riparian vegetation cover in a 10 m buffer, 2) linear upstream weir density along the stream, 3) areal upstream weir density for the catchment watershed, 4) upstream pond cover as a fraction of stream area, and 5) incision class describing the rate of incision of the valley.

Table 3. List of explanatory variables used in models and hypothesized effects on thermal peaks

Category	Variable [units]	Notation	Source	Hypothesized effect	Reference
Climate ^a	Mean annual precipitation [mm]	P_{annual}	SAFRAN	Reduced T_p via increased baseflow	Strauch et al. 2017
	Mean summer ^b precipitation [mm]	P_{summer}	SAFRAN	Increased summer T_p via warm runoff	Nelson and Palmer, 2008, 2007
	Mean annual snow accumulation [mm]	S_{annual}	SAFRAN	Reduced T_p via colder snowmelt water	Caissie, 2006; Webb et al., 2008
	Mean summer ^b air temperature [°C]	T_{summer}	SAFRAN	Increased summer T_p	Moore et al. 2013; Isaak et al., 2017
Hydrology	Mean annual specific discharges [$L s^{-1} km^{-2}$]	Q_{mean}	RHT	Reduced T_p via greater thermal capacity	Caissie, 2006

	Mean monthly minimum specific discharge* [L s ⁻¹ km ⁻²]	q _{min}	RHT	Increased T _p via lower baseflow	Chang and Psaris, 2013
	Concavity index† [-]	CI	RHT	Proxy of water storage in the catchment) Reduced T _p at higher CI	This paper
	Hydrological regime‡ [-]	HR	RHT	Contrast between hydrological regimes under high baseflow and vice versa	This paper
Catchment Watershed characteristics	Mean watershed elevation [m]	elev	RHT	Reduced T _p via low T _{air}	Isaak and Hubert, 2001
	Drainage area [km ²]	area	RHT	Increased T _p due to greater thermal exchange	Hrachowitz et al., 2010; Imholt et al., 2013 ; Isaak et al., 2017
	Mean slope of the watershed [m km ⁻¹]	slope	RHT	Reduced T _p via reduced insolation exposure time	Daigle et al., 2010
	Riparian vegetation cover ratio in 10 meters buffer (%)**	veg	SYRAH	Reduced T _p via shading	Moore et al., 2005
	Linear weir density upstream of stations (# km ⁻¹)**	weirs	SYRAH	Increased T _p via warming	Chandesris et al., 2019
	Areal weir density upstream of stations (# km ⁻²)**	weir area	SYRAH	Increased T _p via warming	Chandesris et al., 2019
	Pond cover ratio upstream of stations (%)**	ponds	SYRAH	Increased T _p via warming	Seyedhashemi et al., 2021
	Stream incision class **	SI	SYRAH	Reduced T _p for greater incision	Webb et al., 2008

*all climate variables are calculated on data from 2009–2017

^bsummer refers to July–August

*determined by geostatistical interpolation on the RHT network (Sauquet et al., 2000) with return period of five years

†ratio of flow duration quantiles Q₁₀-Q₉₉/Q₁-Q₉₉, as defined by Sauquet and Catalogne (2011)

‡classes from 1–12 with pluvial regimes from 1–6, transition regimes from 7–8; and glacial and snow melting regimes from 9–12 (Sauquet et al., 2008)

**Detailed description of these variables in Valette et al. (2012).

250 2.4.2 Multiple regression

We first fit a multiple linear regression model between T_p and explanatory variables using all possible variables characterized in Table 3. Prior to fitting, we scaled the explanatory variables so that their fitted coefficients could be compared in terms of relative importance. We did not use any variable selection techniques in the multiple regression approach because our goal was to compare across the four modeling approaches (regression, ANN, random forest, and multi-model) that use the same independent variables. In other words, we did not seek to have the most parsimonious multiple regression model, but instead used all 16 variables (Table 3) to create the model. Each of these variables was previously assessed for multicollinearity (Fig. A3), and variance inflation factors were all less than 3.

2.4.3 Artificial neural network

260 We then used an ANN—specifically a feed-forward neural network with one hidden layer (R package *nnet*; Venables and Ripley 2002)—to estimate T_p as a potentially non-linear function of covariates. We included a direct connection between covariate inputs and outputs so that the case with zero hidden units corresponded to a linear relationship. We used weight decay regularization, also known as ridge regression, to control overfitting by decreasing less relevant coefficients. Both the number of hidden units and the amount of weight decay were

265 selected with a first cross-validation procedure (Bishop, 2006). To quantify the importance of the different
covariates, we used a connection weight approach (Olden and Jackson, 2002; Olden et al., 2004):

$$W_V = \sum_{h=1}^{nhu} A_{V,h} B_h \quad (6)$$

where

270 $W_V (-)$ = the relevance of covariate V ,
 $A_{V,h} (-)$ = the ANN coefficients connecting hidden unit h to covariate V ,
 $B_h (-)$ = the ANN coefficients connecting hidden unit h to the output, and
 nhu = the number of hidden units

2.4.4 Random forest

275 We used a random forest for the third statistical model structure, using Breiman's algorithm (Breiman, 2001) with
the R package *randomForest* (Liaw and Wiener, 2002) allowing 500 trees. The importance of each predictor
variable was provided as standard output by the *randomForest* package, which determines how much the mean
square errors in prediction increases when that covariate is randomly permuted within the tree.

2.4.5 Multi-model combination

280 Finally, we used a multi-model combination approach to obtain a consensus estimation map of $T_{p,m}$. The
temperature predictions from each previously described model were linearly combined to reduce the associated
uncertainties through multiple linear regression.

$$T_{p,mm,i} = aT_{p,ANN,i} + bT_{p,RF,i} + cT_{p,REG,i} + d \quad (7)$$

where:

285 $T_{p,mm,i}$ =multi-model T_p based on observed and reconstructed $T_{w,i}$ for each observation station, i ;
 $T_{p,ANN,i}$ = T_p estimated by ANN for each observation station i ;
 $T_{p,RF,i}$ = T_p estimated by RF at each observation station i ;
 $T_{p,REG,i}$ = T_p estimated by multiple regression at each observation station i ; and
 a, b, c, d =fitted regression coefficients.

290 Temperature predictions made by the three models are used as multi-model independent variables and each model
prediction is weighted with a coefficient to match the observations as closely as possible. Hence, the multi-model
coefficients are calculated only in relation to observations at the 1700 stations. Then, using equation (7) with
calculated coefficients, we extrapolate T_p along the river network.

2.4.4 Model cross-validation and comparison with air temperature

295 To assess each model's performance prior to spatial extrapolation to the entire RHT network, we conducted a
cross-validation on observed data from our 1700 stations. For each model, including $T_{p,air}$, we used 80% of stations
($n = 1360$) as training data to estimate model parameters. Using those model parameters, we estimated validation
data T_p at the remaining 20% of stations ($n=340$) and cross-validated those estimates with $T_{p,obs}$. We conducted
this cross-validation 100 times allowing for random selection of stations used in the training and validation data
sets. We evaluated the results of the cross-validation with the Nash-Sutcliffe efficiency criterion (NSE, Nash and
300 Sutcliffe, 1970), the RMSE (Root Mean Square Error) and biases between observed $T_{p,obs}$ and $T_{p,m}$. We also
conducted regressions of $T_{p,obs}$ and $T_{p,m}$.

We also compared all T_p model results with a thermal peak calculated using only T_{air} from SAFRAN reanalysis data, $T_{p,air}$. The goal was to evaluate model performance with the simplest and most widely available stream temperature proxy.


305 2.4.4 Hypothesis testing

We evaluated the hypothesis that spatial patterns across models would be consistent, but that drivers would depend on the modeling approach in two ways. First, we compared relative importance of the most important variables across models to see if the same variables emerged. Second, we visually compared maps of $T_{p,m}$ and compared their distributions across watershed area and stream order. We evaluated our second hypothesis that stream size, 310 air temperature, and groundwater contributions would emerge as important regardless of approach by comparing their relationships with $T_{p,m}$ across modeling approaches using Kendall tau correlation.

3. Results

3.1 Validation of the climate correction approach

Gap-filling T_w with our climate correction approach resulted in consistently lower absolute biases for T_p (i.e., $T_{p,fill}$) compared to only using available data (i.e., $T_{p,gap}$), regardless of the length of introduced annual gaps (Figure 2). This discrepancy in bias between uncorrected and corrected data increased exponentially with the number of introduced gaps, growing from a median of 0.03°C with one year of gaps to 0.25°C with eight years of gaps. Climate corrected biases remained below 0.25°C for 75% of the stations with up to six years of introduced gaps. There was no systematic bias regardless of the years taken into account in the regressions.



320


Figure 2. Improvement in absolute bias of thermal peaks across all reference sites ($T_{p,ref}$; $n=30$) when introduced annual gaps were filled with the climate correction procedure ($|T_{p,fill}-T_{p,ref}|$, red boxplots) compared to when gaps were unfilled ($|T_{p,gap}-T_{p,ref}|$, blue boxplots) regardless of the number of introduced gaps.

3.2 Model cross-validation

325 In cross validation, all four statistical models performed substantially better than air temperature at accurately predicting T_p (Figure 3). Indeed, $T_{p,air}$ overestimates the observed $T_{p,ref}$ by 2.5°C on average (Figure 3a), and the negative NSE for $T_{p,air}$ indicates that using a simple mean of T_w observations is a better predictor of T_p than using T_{air} (Figure 3b). Overall, models tend to slightly underestimate T_p (Figure 3a), but mean biases are close to 0°C . The REG and ANN models had similar performance (median RMSE $> 1.5^\circ\text{C}$; median NSE = 0.6), whereas the RF and MM models obtained the best performances (median RMSE $< 1.5^\circ\text{C}$; median NSE = 0.7) with the MM slightly superior for NSE (Figure 3b,c). The models explain in cross-validation between 70% and 78% of the variation in $T_{p,obs}$ with the best performances for RF and MM (Figure 3d-f). Sites with larger watershed areas had consistent positive bias (lighter colors in Figure 3d-g), but bias was more evenly distributed for sites with smaller watershed areas, though the smallest watersheds tended to exhibit negative bias. We did not observe any strong spatial patterning in performance metrics, although sites with larger watershed areas tended to have positive bias (Figure 3d-h).

330

335






Figure 3. Cross validation performance metrics for modeled thermal peaks ($T_{p,m}$), where 80% of stations ($n = 1360$) were training data and 20% of stations ($n=340$) were validation data. Violin plots show distributions of 100 replicates of training-validation data for each model's performance metrics: a) bias relative to $T_{p,ref}$; b) NSE; c) RMSE of $T_{p,m}$ relative to observed T_p . Horizontal lines indicate sample medians and red points indicate sample means. d-f) $T_{p,m}$ vs. $T_{p,obs}$ for each model colored by watershed area at the site. Simple linear regression results ($n = 1700$) are shown in blue with comparison to a 1:1 line (dashed).

3.3 Explanatory variables importance and effects in models

Variable importance differed among modeling approaches, and the maximum importance value for any one variable was between 25–30% (Figure 4). The two most important variables were **catchment watershed area** (area) and mean summer air temperature (T_{summer}) for the RF and multiple regression models (Figure 4b,c). For the ANN model, minimum monthly specific discharge (q_{min}) and riparian vegetation cover (veg) were most important (Figure 4a; note that q_{min} is not correlated with area, Figure A3). Surprisingly, area and T_{summer} obtained relative importances of less than 5% in ANN whereas they were the most important variables in the RF and the REG models. Across models, none of the three precipitation variables (Table 3) emerged as important, nor did the stream incision class.

The cumulative importance of the four most relevant variables of the REG and ANN models is respectively 92% and 81%, which means that the other variables **have had** very little weight in the estimates. This sum is only 69% for RF, which indicates that the relative importance of the explanatory variables are more distributed and the other explanatory variables **have had** a significant weight in the estimates which could explain the best performances in cross validation of RF. In particular, linear and areal weir density upstream of stations occupied a cumulative 7% relative importance. These variables also occupied the two next greatest importance rankings (at 1.7% and 0.9% relative importance, respectively) for the multiple regression model after those shown in Figure 4.






Figure 4. Relative importance of top four explanatory variables calculated in cross validation for the estimation of T_p with the models: a) ANN; b) REG and c) RF. To compare the relative importance of each variable for each model, all variables were centered and scaled. Text in the upper right of each panel refers to the sum of the relative importances of the first four explanatory variables in each model; colors indicate the explanatory variable.

The direction and magnitude of effects from the hypothesized variables (area, groundwater contributions, and summer air temperature) did not systematically differ across modelling approaches (Figure 5). However, there were minor differences in that the RF model tended towards higher $T_{p,m}$ than ANN and REG models at small watershed areas and tended towards lower $T_{p,m}$ than ANN and REG models at larger watershed areas (compare Fig. 5g with 5a and 5d). The RF model was also less sensitive to T_{summer} , with RF producing lower $T_{p,m}$ than ANN and REG models at sites with cooler summer temperatures and lower $T_{p,m}$ than ANN and REG models at sites with warmer summer temperatures (compare Fig. 5i with 5c and 5f).





375

Figure 5. Modeled T_p at each of the 1700 observation stations as a function of the variables hypothesized to be equally important across approaches. Rows indicate the model used (points also colored accordingly) and columns are separated by area, q_{min} , and T_{summer} ; area and q_{min} are natural-log transformed. Lines indicate best-fit smoothers using a generalized additive model and text indicates Kendall tau correlation values (all $p < 0.001$). There was no systematic difference in variable effects on $T_{p,m}$.

380

3.4 Spatial extrapolation of thermal peaks and comparison with air temperature

The statistical models could extrapolate T_p to 92% (105,800 reaches) of the RHT network, and the resulting spatial structure of the extrapolations was consistent across models (Figure 6). On average, T_p was 18.2°C and ranged between 6.3°C and 27.0°C. The highest $T_{p,m}$ (i.e., $T_p > 22^\circ\text{C}$) were generally found on the largest rivers located in the southeast and in the sedimentary plains. The lowest $T_{p,m}$ are found in the mountain streams of the Alps, Pyrenees and Massif Central, and in the northwest. Although the distribution $T_{p,m}$ is consistent among models (Figure 6), there are some clear disparities, particularly at the extremes. The ANN model simulates/predicts lower $T_{p,m}$ (20% of reaches $T_{p,m} < 15^\circ\text{C}$) compared to the RF and multi-multi-model (< 10%; Fig A4a). Application of the climate correction prior to model fitting and subsequent extrapolation showed that differences at the RHT are less

385

390 than 0.5°C at 89% of the reaches (Fig A4b). Still, 10% of the reaches exhibited $T_{p,m}$ differences greater than 0.5°C, and the vast majority of these differences are negative (9.4% vs. 1.6%), suggesting that climate correction more often than not reduces overestimates in T_p .

Estimating T_p with air temperature (i.e., $T_{p,air}$) led to consistently higher values than were obtained with statistical models, with $T_{p,air}$ greater than 20°C for more than 70% of the reaches (Figure 6a, Fig A4a). Indeed, depending on
395 the model, 94–97% of reaches ~~have had~~ $T_{p,m}$ lower than $T_{p,air}$ (Figure 6) and more than 95% of these were in reaches with watershed areas less than 1000 km². $T_{p,air}$ exhibited its greatest overestimations in the smallest watersheds (Figure 7a) and switched from overestimation to underestimation relative to $T_{p,m}$ at reaches with watershed areas between 2000–5000 km². While this behavior was similar across models, the REG and ANN models tended to be produce lower $T_{p,m}$ for small areas and higher $T_{p,m}$ for large areas compared to RF and MM models, suggesting their increased sensitivity to this variable. $T_{p,air}$ also increased its overestimations relative to
400 $T_{p,m}$ in reaches with the most groundwater contributions (Figure 7b). Similar to watershed area, the REG and ANN models tended to be produce higher $T_{p,m}$ for low q_{min} and lower $T_{p,m}$ for high q_{min} than the RF and MM models, suggesting higher sensitivity to this variable. In other words, in small ~~watersheds~~ watersheds with low baseflow, $T_{p,m}$ from REG and ANN models were 2–3°C less than from RF and MM models, but in large watersheds with high baseflow, $T_{p,m}$ from REG and ANN models were 3–5°C greater than from RF and MM models.
405




Figure 6. Thermal peaks (T_p) of stream water extrapolated to all reaches of the French hydrographical network RHT with the different predictive model structures: a) air temperature (T_{air}), b) multiple regression (REG), c) artificial neural network (ANN), d) random forest (RF), and e) multi-model combination of all previous models
410 (MM). All reaches are colored by their modeled range of T_p and colors are chosen to improve visualization.




Figure 7. Differences between stream thermal peaks estimated by air temperature and models depending on a) watershed area, and b) minimum monthly specific discharge. To simplify visualization across the 114,600 reaches, points were summarized across 5% bins in the x-axis; points indicate mean values at each bin, with vertical error bars indicating standard error. a) As watershed area increases, T_{air} shifts from overestimation to underestimation relative to $T_{p,m}$, with a shift in sign between 2,000 and 5,000 km^2 . b) As minimum monthly specific discharge increases, T_{air} tends to increase its overestimation relative to $T_{p,m}$, especially for ~~catchment watershed~~ size greater than 100 km^2 (colors). Only results for RF model are shown- for clarity of presentation and because it had the best performance (each model had similar patterns).

420 4. Discussion

We compiled one of the largest regional summertime stream temperature datasets to address the growing need to understand the spatial distributions of stream thermal ~~regimes in the context of climate change maxima~~. This database is available for use at the following website: <https://thermie-rivieres.inrae.fr>. Using this database, we demonstrate that a simple, ecologically meaningful metric, which we term the thermal peak (T_p), can be reliably estimated at the regional scale using a few easily accessible explanatory variables.

4.1 Horizons and limitations in estimating large-scale stream thermal metrics

Spatiotemporally comprehensive stream temperature datasets are rare because interest in these data is relatively recent and there is little money to support instrumentation at regional or national scales. This lack of centralized data has been recognized as a major limitation for understanding thermal regimes of riverine ecosystems (Arismendi et al., 2012; Ouellet et al., 2020). ~~Existing data typically come from different entities~~ Here, we alleviate these limitations by homogenizing existing disparate datasets and are not managed according to a predefined regional strategy, precluding broad scale synthesis and demonstrate the aggregated utility of such datasets on improving our understanding of ~~controls on~~ stream temperature ~~and its subsequent effects on ecosystems and~~

435 ~~society. To overcome these barriers, we employed a combined empirical approach that~~distributions. Indeed, our
~~homogenized dataset~~ allowed us to identify, at ~~the regional~~a national scale, a map of summer stream temperature
maxima with important implications for aquatic species distributions under climate change. We note here that the
more detailed computation of T_p (i.e., rolling window of 30-days) could be simplified with a mean of August
stream temperature data, as we observed this to regularly be the hottest month across all observable data.

440 We observed the hottest T_p along major rivers and the coldest T_p along small rivers and in mountainous regions
(Fig. 6). The downside of the current approach is that ~~it remains~~is based on interannual metrics. Indeed, the non-
concomitance of the time series ~~does not allow~~precluded us ~~to compare~~from comparing extreme years (hot vs.
cold). However, the stream temperature dataset used here contains ~~benchmark~~enough interannual information,
~~which that it~~ could be used to calibrate models and ~~asses~~assess the impact of climate change on thermal regimes
445 (cf. Isaak et al., 2017, 2020). ~~This map can presently be used to predict and manage future cold-water habitat~~
~~streams~~stream habitats, with potential for regular multi-annual updates.

To enable unbiased comparison among stations from time series with gaps, we used a climate correction of T_w
based on regression with T_{air} . The efficacy of regressions between T_w and T_{air} is well understood (Ducharne, 2008;
Segura et al., 2015, Moatar and Gailhard, 2006), and the approach can be easily transferred to other stations and
450 regions. On the other hand, a cautious approach is required for such regressions, because they assume seasonal
correlation between T_{air} and T_w , and can therefore only apply to rivers having natural seasonal dynamics, without
dam release, thermal peaking, or major weirs regulating the flow of streams (Bruno et al., 2013; Chandesris et al.,
2019; Seyedhashemi et al., 2021; Casado et al., 2013). Moreover, our rolling window approach to T_w - T_{air}
regression suggests that the watershed size may be an important factor in developing these relationships, with
455 regressions for larger rivers benefitting from longer lags in T_{air} (Fig. A2). However, where possible, climate
correction makes it possible to significantly reduce the biases in T_p estimates when compared ~~to~~ those made using
observed data without climate corrections (Figs. 2 and 3). The biases are reduced when the number of years of
observation available is less than four. Beyond this limit, the meteorological variability specific to each year is
sufficient to estimate T_p , which confirms results obtained by (Jones and Schmidt, 2018). Moreover, climate
460 correction reduces overestimation of T_p , reinforcing the importance of taking into account these climatic
corrections of temperature metrics even if it only slightly affects the majority of rivers (Isaak et al., 2017)

4.2 Spatial extrapolation of T_p is consistent and best predicted with a random forest model

All four statistical models achieved similar predictive performance, but the RF model exhibited marginal
improvements over the others (Fig. 3). The MM approach only slightly improves performance in cross-validation
465 in comparison with RF (vis-à-vis the NSE criterion), so it is unlikely to be a useful approach in future applications
due to its complexity. Moreover, whereas the multi-model has the best performance, it lacks the explanatory power
and relative simplicity of the other approaches. In contrast, a potential benefit of the multi-model approach is that
by leveraging multiple approaches, it can compensate for errors particular to individual models.

Overall, our model performances (NSE = 0.75, RMSE <2) were of the same order as other large scale stream
470 temperature studies found in the literature (cf. Segura et al., 2015; Daigle et al., 2010; Wehrly et al., 2009),
suggesting that our approach is reasonable and broadly applicable. Indeed, literature ranges of RMSE for monthly
stream thermal maxima are consistent with values we obtained here (e.g., 0.9–2.1°C for 16 sites across Canada
(Daigle et al. 2010) and 2.0–2.3°C for 1131 sites across USA (Wehrly et al., 2009)), suggesting a general accuracy

limit to current modeling approaches. This is likely because RF models can be prone to overfitting such that they can accurately predict a set of observations, but their performance may decline when predictions are made at unsampled locations. They also have less robust means of model selection and significance testing than the much simpler multiple linear regression approach.

In spatial extrapolation, the T_p estimates are globally consistent between the models and the same spatial structures are found regardless of the approach used (Figure 6). We observed divergences between the models in particular for T_p less than 14°C where ANN tends to ~~simulate~~ predict coldest T_p for more reaches compared to other models (Fig. A4). Still, because our analysis is limited to 2009–2017, these model differences may diminish as T_w measurements grow in time and space. We further note that the performance of the modeling techniques used here was less than that of spatial stream-network models (SSNs) applied to similar temperature datasets, which typically have $R^2 \sim 0.90$ and $\text{RMSE} \sim 1.0^\circ\text{C}$ (Issak ~~Issak~~ et al. 2020). ~~A2017). We tested a~~ SSN model was tested on a small region well covered by data (9000 km², 92 stations) for a robust estimation of parameters with the R package SSN (see Fig. A5 and Table A1) ~~), and indeed; SSN had reduced bias relative to than random forest model (absolute bias decreased by 0.2°C). By comparing the observed and estimated values, we can see that compared to the SSN, the RF model tends to underestimate the high values (on major rivers) and to overestimate the low values (on smaller reaches). Still, the spatial patterns are very consistent among the two approaches, though there are important differences between the SSN and RF model estimates which can be $\pm 2^\circ\text{C}$. However, SSNs are labor intensive to apply in comparison to non-geospatial techniques and require specialized geospatial algorithms for fitting. Hence, despite lower accuracy, the approaches used here may be useful in more generic use cases when geospatial data and computing time are limiting. Overall, while the presented models may not be optimal, we are confident in the predicted spatial patterns.~~

4.3 Drivers of thermal peak depend on model structure

We observed clear divergence of variable importance for the estimation of T_p among the ANN, RF, and REG models. The two most relevant variables in RF and REG were ~~catchment watershed~~ area- and mean summer air temperature, consistent with other studies (Laanaya et al., 2017; McGarvey et al., 2018). The importance of mean summer air temperature on T_p is consistent with other studies (Moore et al, 2010, 2013). Drainage area also emerged as one of the most important variables driving thermal peaks, behind ~~catchment watershed~~ elevation and slope, in a recent regional study (Johnson et al, 2020). Likewise, distance from source and Strahler order, which directly correlate with drainage area, are known to be important drivers of thermal peaks (Hrachowitz et al, 2010; Imholt et al, 2013; Ducharne, 2008, Mohseni et Stephan, 1999). The clear rationale for this common effect is that longer water travel times in large rivers allow more time for temperature equilibration with the atmosphere compared to small rivers (Beaufort et al., ~~2019~~2020a; Mohseni et al., 1998, Fullerton et al., 2015). Large rivers are also less shaded by riparian vegetation and are less influenced by groundwater inflows, which compounds residence time effects.

Importantly, RF had a much more even distribution of variable importances relative to the other models structures, which is likely due to its non-linear structure. In contrast, for ANN, the most important variables were minimum monthly specific discharge (43%) followed by riparian vegetation cover (23%). Higher minimum flows imply a consistent groundwater supply, leading to cool surface waters in summer (Hannah et al., 2004; Kelleher et al., 2012; Lalot et al., 2015). Similarly, more riparian vegetation implies greater shading and reduced temperature

increases from solar radiation (Dugdale et al., 2018; Loicq et al., 2018; Moore et al., 2013). These differences in relevance between the variables for each model underlines the importance of using several approaches and shows that the multi-model approach makes it possible to take into account all these across-model divergences. We caution however, that while predictability may increase, interpretation of variable importance in multi-models is complex, and their utility may therefore decrease when project goals are focused on critical variables for stream temperature habitat restoration.

4.4 Air temperature is not an appropriate proxy for stream temperature

Estimates of T_p produced by stream temperature were clearly more accurate than those produced by air temperature (Figs. 3 and 5). In cross-validation, $T_{p,air}$ overestimated observed T_p by more than 2°C , could not differentiate stream temperature among regions (Figs. 6 and 7), and could not differentiate large rivers from small rivers. These results highlight the need to account for stream-specific temperature estimates, especially when such $T_{p,air}$ overestimates inflate potential ecological change (cf. Kirk and Rahel 2022). Our results further demonstrated that this overestimation of $T_{p,air}$ was greatest in smaller rivers, and that these biases were amplified in streams with significant groundwater contribution (Fig. 7), implying that groundwater buffers effects of increasing T_{air} . This aligns with recent results that found sites with deep groundwater contributions ~~can buffer~~ are much less likely to exhibit increasing summer stream temperatures compared to sites with shallow groundwater contributions (Hare et al. 2021). However, we note that the influence of q_{min} on $T_{p,air}-T_{p,w}$ is weak up to values of approximately $5 \text{ L s}^{-1} \text{ km}^{-2}$ (Fig. 7b), which is in accordance with recent work in the same region (Beaufort et al., 2019). This small effect may in part be because q_{min} is not an effective proxy for groundwater contributions; the base flow index is likely more appropriate (Kelleher et al, 2012; O’Driscoll & DeWalle, 2004, Hare et al, 2021; Johnson et al, 2020), but we were unable to obtain this parameter in this work at a national scale.

Overall, this work clearly demonstrates that T_{air} is an inappropriate proxy for T_w , with important implications for ecological studies, especially those that consider temperature tolerance thresholds of aquatic species. Species distribution models may need to use T_{air} instead of T_w because data from T_w are not sufficient in the regions studied (McGarvey et al., 2018). It is therefore important to introduce T_w in input of these models rather than T_{air} in order to limit the biases linked to the poor spatial representation of T_{air} .


This research further emphasizes the importance of spatial scale and heterogeneity for water temperature studies. As streams increase in size, they become more coupled to air temperature dynamics. Hence, smaller reaches more influenced by groundwater inputs and vegetation may serve as “climate refugia” for ectotherms species especially in the context of climate change. Therefore, choosing the best predictors at a spatial scale for thermal peaks and other thermal regime metrics is essential to accurately predict and manage future stream cold-water habitat. Groundwater and shading proxies could be good candidates.

4. Conclusion


Stream thermal ~~regimes~~ regimes maxima are essential controls on aquatic ecosystems, but our understanding of ~~these regimes~~ their spatial distribution and thus our ability to adequately manage them accordingly ~~in the context of climate change~~ is limited by data availability and simple metrics applicable at large scales. To address these gaps, we created a publicly available, harmonized dataset of stream temperature that can be used by ecologists in France and scientists more broadly. We then developed a simple, ecologically relevant metric—the thermal peak, T_p —that

can be extrapolated at large scales, even when data are sparse. We developed an innovative climate correction method to reduce biases related to such data sparseness, when applied to summer data. The T_p provides an important perspective on the magnitude of thermal extremes during summer, but development of additional metrics such as threshold exceedance frequency, duration, and timing will continue to grow our understanding of stream temperature behavior under climate change. However, development of these metrics will require longer and more spatially explicit time series. Hence, we argue that to improve our capacity to manage and benefit from aquatic ecosystems, it is critical to continue and expand our stream temperature measurement networks.

5. Appendix A



560 **Figure A1.** Histogram of the median date for the hottest 30-day periods across all sites that had annual data. Thirty day rolling windows of mean daily stream temperature were calculated across the entire time series and for each site, and the 30-day windows with the maximum values were selected here. The median date here refers to day 15 in the 30 day period.



565 **Figure A2.** Daily lags for T_{air} that produce the highest R^2 (colors) for a regression between T_w and a right-
aligned moving average of T_{air} at each site as a function of watershed area. The moving average lag whose
regression produced the highest R^2 was used to fill gaps in the time series of T_w for each station. Each point
represents a station, with the x-axis being the watershed area at that station, the y-axis being the lag in T_{air} that
produced the best regression. Points are jittered for visual clarity; only integer lag values were possible. Black
570 points indicate the mean and standard error (horizontal bars) of watershed areas within each possible lag (2–10
days).






Figure A3. Correlation matrix of all 16 considered environmental variables prior to selection in the thermal peak analysis. All variables had collinearity values less than 0.6 or greater than -0.6.



575

Figure A4. Distributions of $T_{p,m}$ extrapolated to all RHT network reaches. a) Histogram of models (10 bins equally spaced between 0°C and 30°C) according their extrapolated T_p value, colored by model, and b) differences between $T_{p,m}$ calculated with and without climate corrections on the 1,630 stations with gaps in their data.



580 **Figure A5.** Figure comparing modeled thermal peak ($T_{p,m}$) estimates from (a) SSN, (b) RF, and (c) the
 difference between RF and SSN. The SSN model here is from a benchmark test on a small region well covered
 by data (9000 km², 92 stations) for a robust estimation of parameters with the R package *SSN*. SSN had reduced
 bias relative to than random forest model (absolute bias decreased by 0.2°C). Additionally, by comparing the
 observed and estimated values, we can see that RF tends to underestimate the high values and to overestimate the
 585 low values. Unfortunately, due to the lack of an RHT with upstream-downstream information, we could not
 apply at the scale of the whole catchmentwatershed. Still, the spatial patterns are very consistent among the two
 approaches, though there are important differences between the SSN and RF model estimates which can be +/-
 2°C. The estimates of the SSN model are generally warmer than those of RF on the main major river axes and
 colder on the small tributaries. This is consistent also with observations. So, while the presented models may not
 590 be optimal, we are confident the spatial patterns are correct.

Table A1. Comparison of model performance metrics for the region tested in Figure A5

Model	RMSE (-)	NSE (-)	Bias (°C)	Absolute bias (°C)
RF	1.24	0.44	0.01	0.95
SSN	0.99	0.81	0.05	0.76

References

595 Arismendi, I., Johnson, S. L., Dunham, J. B., Haggerty, R., and Hockman-Wert, D.: The paradox of cooling streams in a warming world: regional climate trends do not parallel variable local trends in stream temperature in the Pacific continental United States, *Geophysical Research Letters*, 39, 2012.

Arismendi, I., Safeeq, M., Dunham, J. B., and Johnson, S. L.: Can air temperature be used to project influences of climate change on stream temperature?, *Environmental Research Letters*, 9, 084015, 2014.

- 600 ~~Ashley Steel, E., Sowder, C. and Peterson, E.E.: Spatial and temporal variation of water temperature regimes on the Snoqualmie River network, JAWRA Journal of the American Water Resources Association, 52(3), 769–787, 2016.~~
- Beaufort, A., Moatar, F., Curie, F., Ducharme, A., Bustillo, V., and Thiéry, D.: River Temperature Modelling by Strahler Order at the Regional Scale in the Loire River Basin, France, River Research and Applications, 32, 597-609, 10.1002/rra.2888, ~~2016~~2016a.
- 605 ~~Beaufort, A., Curie, F., Moatar, F., Ducharme, A., Melin, E. and Thiery, D.: T-NET, a dynamic model for simulating daily stream temperature at the regional scale based on a network topology. Hydrological Processes, 30(13), pp.2196-2210, 2016b.~~
- Beaufort, A., Moatar, F., Sauquet, E., Loicq, P., and Hannah, D. M.: Influence of landscape and hydrological factors on stream–air temperature relationships at regional scale, Hydrological Processes, 34, 583-597, 2020a.
- 610 Beaufort, A., Moatar, F., Sauquet, E., and Magand, C.: Thermie en rivière : Analyse géostatistique et description de régime : Application à l'échelle de la France, INRAE, 106 pages, 2020b.
- Benyahya, L., Caissie, D., St-Hilaire, A., Ouarda, T. B., and Bobée, B.: A review of statistical water temperature models, Canadian Water Resources Journal, 32, 179-192, 2007.
- Bishop, C. M.: Pattern recognition and machine learning, springer, 2006.
- 615 Breiman, L.: Random forests, Machine learning, 45, 5-32, 2001.
- Bruno, M. C., Siviglia, A., Carolli, M., and Maiolini, B.: Multiple drift responses of benthic invertebrates to interacting hydropeaking and thermopeaking waves, Ecohydrology, 6, 511-522, 2013.
- Buisson, L., and Grenouillet, G.: Contrasted impacts of climate change on stream fish assemblages along an environmental gradient, Divers. Distrib., 15, 613-626, 2009.
- 620 ~~Bustillo, V., Moatar, F., Ducharme, A., Thiery, D., and Poirel, A.: A multimodel comparison for assessing water temperatures under changing climate conditions via the equilibrium temperature concept: case study of the Middle Loire River, France, Hydrological Processes, 28, 1507–1524, 10.1002/hyp.9683, 2014.~~
- Caissie, D.: The thermal regime of rivers: a review, Freshwater biology, 51, 1389-1406, 2006.
- 625 ~~Carlson, S.M., Hendry, A.P. and Letcher, B.H.: Growth rate differences between resident native brook trout and non-native brown trout. Journal of Fish Biology, 71(5), pp.1430-1447, 2007~~
- Casado, A., Hannah, D. M., Peiry, J. L., and Campo, A. M.: Influence of dam-induced hydrological regulation on summer water temperature: Sauce Grande River, Argentina, Ecohydrology, 6, 523-535, 2013.
- Catalogne, C.: Amélioration des méthodes de prédétermination des débits de référence d'étiage en sites peu ou pas jaugés, Doctorat Ocean Atmosphere Hydrologie, Université Joseph Fourier, Grenoble, 2012.
- 630 Chandesris, A., Looy, K. V., Diamond, J. S., and Souchon, Y.: Small dams alter thermal regimes of downstream water, Hydrology and Earth System Sciences, 23, 4509-4525, 2019.
- Chang, H., and Parris, M.: Local landscape predictors of maximum stream temperature and thermal sensitivity in the Columbia River Basin, USA, Science of the Total Environment, 461, 587-600, 2013.
- 635 ~~Chenard, J. F., and Caissie, D.: Stream temperature modelling using artificial neural networks: application on Catamaran Brook, New Brunswick, Canada, Hydrological Processes: An International Journal, 22, 3361–3372, 2008.~~
- Conti, L., Comte, L., Hugueny, B., and Grenouillet, G.: Drivers of freshwater fish colonisations and extirpations under climate change, Ecography, 38, 510-519, 2015.

- 640 Daigle, A., St-Hilaire, A., Peters, D., and Baird, D.: Multivariate modelling of water temperature in the Okanagan watershed, *Canadian Water Resources Journal*, 35, 237-258, 2010.
- ~~Daufresne, M., Roger, M., Capra, H., and Lamouroux, N.: Long term changes within the invertebrate and fish communities of the Upper Rhône River: effects of climatic factors, *Global Change Biology*, 10, 124-140, 2004.~~
- ~~Daufresne, M., Lengfellner, K., and Sommer, U.: Global warming benefits the small in aquatic ecosystems, *Proceedings of the National Academy of Sciences*, 106, 12788-12793, 2009.~~
- 645 ~~Detenbeck, N. E., Morrison, A. C., Abele, R. W., and Kopp, D. A.: Spatial statistical network models for stream and river temperature in New England, USA, *Water Resources Research*, 52, 6018-6040, 2016.~~
- ~~DeWeber, J. T., and Wagner, T.: A regional neural network ensemble for predicting mean daily river water temperature, *Journal of Hydrology*, 517, 187-200, 2014.~~
- Ducharme, A.: Importance of stream temperature to climate change impact on water quality, *Hydrology and Earth System Sciences*, 12, 797-810, 2008.
- 650 Dugdale, S. J., Malcolm, I. A., Kantola, K., and Hannah, D. M.: Stream temperature under contrasting riparian forest cover: Understanding thermal dynamics and heat exchange processes, *Science of The Total Environment*, 610, 1375-1389, 2018.
- Durance, I., and Ormerod, S. J.: Trends in water quality and discharge confound long-term warming effects on river macroinvertebrates, *Freshwater Biology*, 54, 388-405, 2009.
- 655 ~~Durand, Y., Brun, E., Merindol, L., Guyomarc'h, G., Lesaffre, B. and Martin, E.: A meteorological estimation of relevant parameters for snow models, *Annals of glaciology*, 18, 65-71, 1993.~~
- ~~Elliott, J.: The energetics of feeding, metabolism and growth of brown trout (*Salmo trutta* L.) in relation to body weight, water temperature and ration size, *The Journal of Animal Ecology*, 923-948, 1976.~~
- 660 ~~Fullerton, A.H., Torgersen, C.E., Lawler, J.J., Faux, R.N., Steel, E.A., Beechie, T.J., Ebersole, J.L. and Leibowitz, S.G.: Rethinking the longitudinal stream temperature paradigm: region-wide comparison of thermal infrared imagery reveals unexpected complexity of river temperatures. *Hydrological Processes*, 29(22), pp.4719-4737, 2015.~~
- 665 Hannah, D. M., Malcolm, I. A., Soulsby, C., and Youngson, A. F.: Heat exchanges and temperatures within a salmon spawning stream in the Cairngorms, Scotland: seasonal and sub-seasonal dynamics, *River Research and Applications*, 20, 635-652, 2004.
- Hare, D.K., Helton, A.M., Johnson, Z.C., Lane, J.W. and Briggs, M.A.: Continental-scale analysis of shallow and deep groundwater contributions to streams, *Nature communications*, 12(1), 1-10, 2021.
- 670 ~~Heino, J., Virkkala, R., and Toivonen, H.: Climate change and freshwater biodiversity: detected patterns, future trends and adaptations in northern regions, *Biological Reviews*, 84, 39-54, 2009.~~
- ~~Hill, R. A., Hawkins, C. P., and Carlisle, D. M.: Predicting thermal reference conditions for USA streams and rivers, *Freshwater Science*, 32, 39-55, 2013.~~
- ~~Hawkins, C.P., Hogue, J.N., Decker, L.M. and Feminella, J.W.: Channel morphology, water temperature, and assemblage structure of stream insects, *Journal of the North American benthological society*, 16(4), pp.728-749, 1997.~~
- 675 Hill, R. A., and Hawkins, C. P.: Using modelled stream temperatures to predict macro-spatial patterns of stream invertebrate biodiversity, *Freshwater Biology*, 59, 2632-2644, 2014.

- 680 Hrachowitz, M., Soulsby, C., Imholt, C., Malcolm, I., and Tetzlaff, D.: Thermal regimes in a large upland salmon river: a simple model to identify the influence of landscape controls and climate change on maximum temperatures, *Hydrological Processes*, 24, 3374-3391, 2010.
- Imholt, C., Soulsby, C., Malcolm, I.A., Hrachowitz, M., Gibbins, C.N., Langan, S. and Tetzlaff, D.: Influence of scale on thermal characteristics in a large montane river basin, *River Research and Applications*, 29(4), 403-419, 2013.
- 685 Isaak, D.J., and Hubert, W. A.: A hypothesis about factors that affect maximum summer stream temperatures across montane landscapes 1. *Journal of the American Water Resources Association*, 37(2), 351-366, 2001.
- Isaak, D. J., ~~Luce, C. H., Rieman, B. E., Nagel, D. E., Peterson, E. E., Horan, D. L., Parkes, S., and Chandler, G. L.: Effects of climate change and wildfire on stream temperatures and salmonid thermal habitat in a mountain river network, *Ecological Applications*, 20, 1350-1371, 2010.~~
- 690 ~~Isaak, D. J.,~~ Wenger, S. J., Peterson, E. E., Ver Hoef, J. M., Nagel, D. E., Luce, C. H., Hostetler, S. W., Dunham, J. B., Roper, B. B., and Wollrab, S. P.: The NorWeST summer stream temperature model and scenarios for the western US: A crowd-sourced database and new geospatial tools foster a user community and predict broad climate warming of rivers and streams, *Water Resources Research*, 53, 9181-9205, 2017.
- Isaak, D.J., Luce, C.H., Horan, D.L., Chandler, G.L., Wollrab, S.P., Dubois, W.B. and Nagel, D.E.: Thermal regimes of perennial rivers and streams in the Western United States, *JAWRA Journal of the American Water Resources Association*, 56(5), 842-867, 2020.
- 695 Jackson, F. L., Fryer, R. J., Hannah, D. M., Millar, C. P., and Malcolm, I. A.: A spatio-temporal statistical model of maximum daily river temperatures to inform the management of Scotland's Atlantic salmon rivers under climate change, *Science of the Total Environment*, 612, 1543-1558, 2018.
- 700 Johnson, Z.C., Johnson, B.G., Briggs, M.A., Devine, W.D., Snyder, C.D., Hitt, N.P., Hare, D.K. and Minkova, T.V.: Paired air-water annual temperature patterns reveal hydrogeological controls on stream thermal regimes at watershed to continental scales, *Journal of Hydrology*, 587, p.124929, 2020.
- Jones, N., and Schmidt, B.: Thermal regime metrics and quantifying their uncertainty for North American streams, *River Research and Applications*, 34, 382-393, 2018.
- 705 Kelleher, C., Wagener, T., Gooseff, M., McGlynn, B., McGuire, K., and Marshall, L.: Investigating controls on the thermal sensitivity of Pennsylvania streams, *Hydrological Processes*, 26, 771-785, 2012.
- ~~Kishi, D., Murakami, M., Nakano, S., and Mackawa, K.: Water temperature determines strength of top-down control in a stream food web, *Freshwater biology*, 50, 1315-1322, 2005.~~
- ~~Kirk, M.A. and Rahel, F.J.: Air temperatures over-predict changes to stream fish assemblages with climate warming compared with water temperatures, *Ecological Applications*, 32(1), p.e02465, 2022.~~
- 710 Laanaya, F., St-Hilaire, A. and Gloaguen, E.: Water temperature modelling: comparison between the generalized additive model, logistic, residuals regression and linear regression models, *Hydrological Sciences Journal*, 62(7), 1078-1093, 2017
- Lalot, E., Curie, F., Wawrzyniak, V., Baratelli, F., Schomburgk, S., Flipo, N., Piegay, H., and Moatar, F.: Quantification of the contribution of the Beauce groundwater aquifer to the discharge of the Loire River using thermal infrared satellite imaging, *Hydrology and Earth System Sciences*, 19, 4479-4492, 2015.
- 715

- ~~Letcher, B. H., Hocking, D. J., O'Neil, K., Whiteley, A. R., Nislow, K. H., and O'Donnell, M. J.: A hierarchical model of daily stream temperature using air-water temperature synchronization, autocorrelation, and time lags, *PeerJ*, 4, e1727, 2016.~~
- 720 Liaw, A., and Wiener, M.: Classification and regression by randomForest, *R news*, 2, 18-22, 2002.
- Logez, M., Bady, P., and Pont, D.: Modelling the habitat requirement of riverine fish species at the European scale: sensitivity to temperature and precipitation and associated uncertainty, *Ecology of Freshwater Fish*, 21, 266-282, 2012.
- Loicq, P., Moatar, F., Jullian, Y., Dugdale, S. J., and Hannah, D. M.: Improving representation of riparian
- 725 vegetation shading in a regional stream temperature model using LiDAR data, *Science of the total environment*, 624, 480-490, 2018.
- ~~Malard, F., Toekner, K., DOLE OLIVIER, M. J., and Ward, J.: A landscape perspective of surface-subsurface hydrological exchanges in river corridors, *Freshwater Biology*, 47, 621-640, 2002.~~
- ~~Mayer, T.D.: Controls of summer stream temperature in the Pacific Northwest. *Journal of Hydrology*, 475, pp.323-335, 2012.~~
- 730 McGarvey, D. J., Menon, M., Woods, T., Tassone, S., Reese, J., Vergamini, M., and Kellogg, E.: On the use of climate covariates in aquatic species distribution models: are we at risk of throwing out the baby with the bath water?, *Ecography*, 41, 695-712, 2018.
- ~~Miller, S. W., Wooster, D., and Li, J.: Resistance and resilience of macroinvertebrates to irrigation water withdrawals, *Freshwater Biology*, 52, 2494-2510, 2007.~~
- 735 Moatar, F., Miquel, J., and Poirel, A.: A quality-control method for physical and chemical monitoring data. Application to dissolved oxygen levels in the river Loire (France), *Journal of Hydrology*, 252, 25-36, 10.1016/s0022-1694(01)00439-5, 2001.
- Moatar, F., and Gailhard, J.: Water temperature behaviour in the River Loire since 1976 and 1881, *Comptes*
- 740 *Rendus Geoscience*, 338, 319-328, 10.1016/j.crte.2006.02.011, 2006.
- ~~Morrill, J.C., Bales, R.C. and Conklin, M.H.: Estimating stream temperature from air temperature: implications for future water quality. *Journal of Environmental Engineering*, 131(1), pp.139-146, 2005.~~
- Mohseni, O., Stefan, H. G., and Erickson, T. R.: A nonlinear regression model for weekly stream temperatures, *Water Resources Research*, 34, 2685-2692, 1998.
- 745 Mohseni, O. and Stefan, H.G.: Stream temperature/air temperature relationship: a physical interpretation, *Journal of hydrology*, 218(3-4), 128-141, 1999.
- Moore, R. D., Spittlehouse, D. L., & Story, A.: Riparian microclimate and stream temperature response to forest harvesting: a review1. *JAWRA Journal of the American Water Resources Association*, 41(4), 813-834, 2005.
- Moore, R., Nelitz, M., and Parkinson, E.: Empirical modelling of maximum weekly average stream temperature
- 750 in British Columbia, Canada, to support assessment of fish habitat suitability, *Canadian Water Resources Journal*, 38, 135-147, 2013.
- Nash, J. E., and Sutcliffe, J. V.: River flow forecasting through conceptual models part I—A discussion of principles, *Journal of hydrology*, 10, 282-290, 1970.
- Nelson, K.C. and Palmer, M.A.: Stream temperature surges under urbanization and climate change: data, models,
- 755 and responses 1, *JAWRA journal of the American water resources association*, 43(2), 440–452, 2007.

~~Nimick, D.A., Gammons, C.H. and Parker, S.R.: Diel biogeochemical processes and their effect on the aqueous chemistry of streams: A review. *Chemical Geology*, 283(1-2), pp.3-17, 2011.~~

O'Driscoll, M.A. and DeWalle, D.R.: Stream-air temperature relationships as indicators of groundwater inputs, Watershed Update (AWRA Hydrology and Watershed Management Technical Committee), 2, 2004.

760 ~~Ojanguren, A., and Braña, F.: Thermal dependence of swimming endurance in juvenile brown trout, *Journal of Fish Biology*, 56, 1342-1347, 2000.~~

Olden, J. D., and Jackson, D. A.: Illuminating the “black box”: a randomization approach for understanding variable contributions in artificial neural networks, *Ecological modelling*, 154, 135-150, 2002.

765 Olden, J. D., Joy, M. K., and Death, R. G.: An accurate comparison of methods for quantifying variable importance in artificial neural networks using simulated data, *Ecological modelling*, 178, 389-397, 2004.

~~Olden, J. D., and Naiman, R. J.: Incorporating thermal regimes into environmental flows assessments: modifying dam operations to restore freshwater ecosystem integrity, *Freshwater Biology*, 55, 86-107, 2010.~~

770 Ouellet, V., St-Hilaire, A., Dugdale, S. J., Hannah, D. M., Krause, S., and Proulx-Ouellet, S.: River temperature research and practice: Recent challenges and emerging opportunities for managing thermal habitat conditions in stream ecosystems, *Science of The Total Environment*, 139679, 2020.

Pella, H., Lejot, J., Lamouroux, N., and Snelder, T.: Le réseau hydrographique théorique (RHT) français et ses attributs environnementaux, *Géomorphologie: relief, processus, environnement*, 18, 317-336, 2012.

~~Pratt, B., and Chang, H.: Effects of land cover, topography, and built structure on seasonal water quality at multiple spatial scales, *Journal of hazardous materials*, 209, 48-58, 2012.~~

775 ~~Quintana-Segui, P., Le Moigne, P., Durand, Y., Martin, E., Habets, F., Baillon, M., Canellas, C., Franchisteguy, L., and Morel, S.: Analysis of near surface atmospheric variables: Validation of the SAFRAN analysis over France, *Journal of applied meteorology and climatology*, 47, 92-107, 2008.~~

~~Rivers Moore, N.A., Dallas, H.F. and Morris, C., 2013: Towards setting environmental water temperature guidelines: A South African example, *Journal of environmental management*, 128, 380-392, 2013.~~

780 Sauquet, E., Gottschalk, L., and Leblois, E.: Mapping average annual runoff: a hierarchical approach applying a stochastic interpolation scheme, *Hydrological sciences journal*, 45, 799-815, 2000.

~~Sauquet, E.: Mapping mean annual river discharges: geostatistical developments for incorporating river network dependencies, *Journal of Hydrology*, 331, 300-314, 2006.~~

785 ~~Sauquet E., Gottschalk L., and Krasovskaia I. Estimating mean monthly runoff at ungauged locations: an application to France. *Hydrology Research*, 39, 403-423, 2008.~~

Sauquet, E., and Catalogne, C.: Comparison of catchment grouping methods for flow duration curve estimation at ungauged sites in France, *Hydrology and Earth System Sciences Discussions*, 15, p. 2421-p. 2435, 2011.

Segura, C., Caldwell, P., Sun, G., McNulty, S., and Zhang, Y.: A model to predict stream water temperature across the conterminous USA, *Hydrological Processes*, 29, 2178-2195, 2015.

790 Seyedhashemi, H., Moatar, F., Vidal, J.-P., Diamond, J. S., Beaufort, A., Chandesaris, A., and Valette, L.: Thermal signatures identify the influence of dams and ponds on stream temperature at the regional scale, *Science of The Total Environment*, 766, 142667, <https://doi.org/10.1016/j.scitotenv.2020.142667>, 2021.

~~Sohrabi, M. M., Benjankar, R., Tonina, D., Wenger, S. J., and Isaak, D. J.: Estimation of daily stream water temperatures with a Bayesian regression approach, *Hydrological Processes*, 31, 1719-1733, 2017.~~

- 795 [Steel, E. A., Beechie, T. J., Torgersen, C. E., and Fullerton, A. H.: Envisioning, quantifying, and managing thermal regimes on river networks, *BioScience*, 67, 506-522, 2017.](#)
- [Song, C., Dodds, W.K., Rüegg, J., Argerich, A., Baker, C.L., Bowden, W.B., Douglas, M.M., Farrell, K.J., Flinn, M.B., Garcia, E.A. and Helton, A.M.: Continental-scale decrease in net primary productivity in streams due to climate warming. *Nature Geoscience*, 11\(6\), pp.415-420, 2018.](#)
- 800 Strauch, A.M., MacKenzie, R.A. and Tingley III, R.W.: Base flow-driven shifts in tropical stream temperature regimes across a mean annual rainfall gradient, *Hydrological Processes*, 31(9), pp.1678-1689, 2017.
- Tisseuil, C., Vrac, M., Grenouillet, G., Wade, A. J., Gevrey, M., Oberdorff, T., Grodwohl, J.-B., and Lek, S.: Strengthening the link between climate, hydrological and species distribution modeling to assess the impacts of climate change on freshwater biodiversity, *Science of the total environment*, 424, 193-201, 2012.
- 805 ~~[Tsang, Y. P., Infante, D. M., Stewart, J., Wang, L., Tingley III, R. W., Thornbrugh, D., Cooper, A. R., and Daniel, W. M.: StreamThermal: A software package for calculating thermal metrics from stream temperature data, *Fisheries*, 41, 548-554, 2016.](#)~~
- Valette, L., Piffady, J., Chandesris, A., and Souchon, Y.: SYRAH-CE: description des données et modélisation du risque d'altération hydromorphologique des cours d'eau pour l'état des lieux DCE, Rapport Technique Onema-Irstea, 2012.
- 810 ~~[van Vliet, M. T. H., Yearsley, J. R., Franssen, W. H. P., Ludwig, F., Haddeland, I., Lettenmaier, D. P., and Kabat, P.: Coupled daily streamflow and water temperature modelling in large river basins, *Hydrology and Earth System Sciences*, 16, 4303-4321, 10.5194/hess-16-4303-2012, 2012.](#)~~
- Venables, W. N. and Ripley, B. D.: *Modern Applied Statistics with S*. Fourth Edition. Springer, New York. ISBN 0-387-95457-0, 2002.
- 815 Vidal, J. P., Martin, E., Franchistéguy, L., Baillon, M., and Soubeyroux, J. M.: A 50-year high-resolution atmospheric reanalysis over France with the Safran system, *International Journal of Climatology*, 30, 1627-1644, 2010.
- Webb, B.W., Hannah, D.M., Moore, R.D., Brown, L.E. and Nobilis, F.: Recent advances in stream and river temperature research, *Hydrological Processes: An International Journal*, 22(7), 902-918, 2008.
- 820 Wehrly, K. E., Brenden, T. O., and Wang, L.: A comparison of statistical approaches for predicting stream temperatures across heterogeneous landscapes 1, *JAWRA Journal of the American Water Resources Association*, 45, 986-997, 2009.
- ~~[Westhoff, M., Savenije, H., Luxemburg, W., Stelling, G., Van de Giesen, N., Selker, J., Pfister, L., and Uhlenbrook, S.: A distributed stream temperature model using high resolution temperature observations, *Hydrology and Earth System Sciences*, 11, 1469-1480, 2007.](#)~~
- 825 Wolter, C.: Temperature influence on the fish assemblage structure in a large lowland river, the lower Oder River, Germany, *Ecology of Freshwater Fish*, 16, 493-503, 2007.
- ~~[Yearsley, J.: A grid-based approach for simulating stream temperature, *Water Resources Research*, 48, 2012.](#)~~
- 830 [Xu, C.L., Letcher, B.H. and Nislow, K.H.: Size-dependent survival of brook trout *Salvelinus fontinalis* in summer: effects of water temperature and stream flow. *Journal of Fish Biology*, 76\(10\), pp.2342-2369, 2010.](#)