

Recent hydrological response of glaciers in the Canadian Rockies to changing climate and glacier configuration

Dhiraj Pradhananga^{1,2,3}, John W. Pomeroy¹

5 ¹Centre for Hydrology, University of Saskatchewan, Canmore, 1151 Sidney Street, Alberta, Canada T1W 3G1

²Department of Meteorology, Tri-Chandra Multiple Campus, Tribhuvan University, Kathmandu, Nepal

³The Small Earth Nepal, P.O. Box 20533, Kathmandu, Nepal

Correspondence to: Dhiraj Pradhananga (dhiraj.pradhananga@usask.ca)

10

Abstract. Mountain snow and ice greatly influence the hydrological cycle of alpine regions by regulating both the quantity and seasonal variations of water availability downstream. This study considers the combined impacts of climate and glacier changes due to recession on the hydrology and water balance of two high-elevation basins in the Canadian Rockies. A distributed, physically based, uncalibrated glacier hydrology model developed in the Cold Regions Hydrological Modelling platform (CRHM) was used to simulate the glacier mass balance and basin hydrology of Peyto and Athabasca Glacier basins in Alberta. Bias-corrected reanalysis data were used to drive the model. The model calculates the water balance of glacierized basins, influenced by the surface energy and mass balance, and considering redistribution of snow by wind and avalanches. It was set up using hydrological response units based on elevation bands, surface slope and aspect, as well as changing land cover. Aerial photos, satellite images and Digital Elevation Models (DEM) were assimilated to represent the changing configurations of glacier area and the exposure of ice and firn. Observations of glacier mass balance, snow and glacier ice surface elevation changes at glacier and alpine tundra meteorological stations and streamflow discharge at the glacier outlets were used to evaluate the model performance. Basin hydrology was simulated over two periods, 1965-1975 and 2008-2018, using observed glacier configurations for those time periods. Both basins have undergone continuous glacier loss over the last three to five decades, leading to a 6-31% reduction in glacierized area, a 78-109% increase in ice exposure, and changes to the elevation and slope of the glacier surfaces. Air temperatures are increasing, mainly due to increasing winter maximum and summer minimum daily temperatures. Annual precipitation has increased by less than 11%, but rainfall ratios have increased by 29-44%. The results show that changes in both climate and glacier configuration have influenced changes in melt rates and runoff, and a shift of peak flows in Peyto Glacier basin from August to July. Glacier melt contributions increased/decreased from 27-61% to 43-59% of annual discharges. Recent discharges were 3-19% higher than in the 1960s and 1970s. The results suggest that increased exposure of glacier ice and lower surface elevation due to glacier thinning were less influential than climate warming in increasing streamflow. Streamflow from these glaciers continues to increase.

1 Introduction

Mountain streamflow profoundly affects the quantity, quality and seasonal variation of water in major downstream river systems, particularly in the arid and semiarid regions of western North America (Marks et al., 2008). Glaciers contribute significantly to this streamflow during warm and dry periods and in doing so moderate inter-annual variability and contribute to flow water during extreme warm and dry periods (Comeau et al., 2009; Fountain and Tangborn, 1985; Hopkinson and Young, 1998). North American mountain glaciers began to retreat after the Little Ice Age ended in the early 1800s (Barry, 2006; Riedel *et al.*, 2015). In addition to warming after the Little Ice Age, Western Canada has experienced a warming climate in recent decades, with changes in precipitation regimes and decline in snow cover (DeBeer et al., 2016). As a result, glaciers in the region are retreating more rapidly than in the past (DeBeer et al., 2016; Schiefer et al., 2007; Tennant et al., 2012) and additional glacier meltwater is contributing to headwater streamflow discharge and groundwater recharge (Castellazzi et al., 2019).

Marshall *et al.* (2011) projected glacier volumes of the Canadian Rockies (eastern slopes) for the next century. Their projected values are alarming, as they indicate a further ~85% loss of glacier volume by 2100 and an order of magnitude decrease in glacier contribution to streamflow in Alberta from 1.1 km³ per year at present to 0.1 km³ per year at the end of this century. Similarly, Clarke *et al.* (2015) projected a 75% loss of glacier mass in western Canada by the end of the 21st Century compared to 2005.

It has been proposed that as the climate warms, flow originating from glaciers will increase for a certain time due to increased melt rates, then decline as the glacier-covered area decreases (Moore *et al.*, 2009). The duration and timing of this change from increasing to decreasing flow, however, will be regionally dependent on basin elevation and/or glacier coverage of the basin (Casassa *et al.*, 2009). Stahl and Moore (2006) observed that late summer streamflow from many British Columbia glacierized mountain basins has been declining, which suggests that most source glaciers have already completed the phase of increased flow due to rising temperatures and increasing melt rates and now contribute less streamflow as their areas decline. Chernos et al. (2020) projected a rise in glacier discharge in Athabasca River Basin in Alberta until the mid-21st century and then reduced discharge. Similarly, Neupane et al., (2017) modelled the upper Athabasca River basin and assessed the effects of changes in temperature and precipitation on simulated future discharge for the 2080-2099 period. They projected a reduction in water availability in the basin during the summer months.

Glacier mass loss is associated with a reduction in glacier-covered area, an increase in ice exposure compared to that of snow and firn, and changes to the elevation and slope of the glacier surface. However, it is yet to be fully understood how the changes in glacier configuration and climate impact the streamflow jointly and individually. The integrated impacts of climate change on mountain streamflow are complex and sometimes changes in hydrological processes can have compensating effects on streamflow generation under a changing climate, e.g., a reduction in one contribution is compensated for by increase in another contribution (Fang and Pomeroy, 2020; Harder et al., 2015). Therefore, the impacts from climate change and glacier change on hydrology need to be diagnosed both separately and together. This study investigates the individual and combined impacts of changing climate and receding glaciers on the headwater hydrology of two well instrumented glacierized basins on the

eastern slopes of the Canadian Rockies using a cold regions glacier hydrological model, CRHM-glacier (Cold Regions Hydrological Model – Glacier), forced by bias-corrected reanalysis data.

2 Study basins, data, methods

5

2.1 Study basins

Two alpine glacier basins in the Canadian Rockies, Alberta, Canada (Figure 1) were chosen for this research, Peyto Glacier Research Basin (PGRB, 22.43 km²) in Banff National Park and Athabasca Glacier Research Basin (AGRB, 29.3 km²) in Jasper National Park. The details of these basins are provided by Pradhananga and Pomeroy (2022) and Pradhananga et al. (2021). Both glaciers are instrumented with on-ice meteorological stations on the lower ice tongues of the glacier and off-ice meteorological stations on moraine below the tongue and their outlet streams are gauged at the outlets of their current proglacial lakes. Both glaciers have been losing mass continuously since the mid-1970s (Demuth and Keller, 2006; Intsiful and Ambinakudige, 2021; Kehrl et al., 2014; Reynolds and Young, 1997; Tennant and Menounos, 2013). Clarke et al. (2015) projected that AGRB will lose half of its glacier coverage by 2050. Kehrl *et al.* (2014) estimated that Peyto Glacier may lose about 85% of its present-day mass by 2100.

15

2.2 Meteorological forcing datasets

Bias-corrected ERA-40 (Uppala et al., 2005) and ERA-Interim reanalysis data (Dee et al., 2011) representing surface levels for air temperature, vapour pressure, wind speed, precipitation, incoming short- and longwave radiation were used to force the CRHM-glacier model. The ERA-40 was available for the period of 1957-2002 and ERA-Interim for the period of 1979-2019 with an overlapping period of 1979-2002. These ERA global reanalyses were first bias corrected to *in situ* observational datasets from meteorological stations near to the glaciers (Athabasca Moraine Station for AGRB and Peyto Main Station for PGRB, Figure 1). A monthly quantile mapping approach (Gudmundsson, 2016) with monthly bias correction factors were used for the bias correction of ERA-40 and ERA-Interim.

25

For PGRB, ERA-Interim data (2008-2018) were bias-corrected to Peyto Main Station observations from 2013-2018 and ERA-40 data (1965-1975) were bias-corrected to the archived observations from the station for the common overlap period of 1992-2001 (Munro, 2011), as described by Pradhananga et al. (2021). For AGRB, ERA-Interim data were bias-corrected to 2014-2018 observations at the Athabasca Moraine Station. There were not any *in-situ* meteorological observations available at AGRB prior to 2014. Therefore, the biases in ERA-40 for AGRB were identified by comparing the overlapping period of ERA-40 and bias corrected ERA-Interim, similar to the procedure outlined by Krogh and Pomeroy (2018).

35

2.3 CRHM-glacier model

The CRHM-glacier model (Pradhananga and Pomeroy, 2022), developed in the Cold Regions Hydrological Modelling Platform (Pomeroy *et al.*, 2007) was applied in this study to evaluate the impacts of changes in climate and in glacier configuration on the hydrology of glacierized basins. CRHM-glacier is a physically based, flexible, multi-physics hydrological model (Pradhananga and Pomeroy, 2022). It downscales and distributes

40

meteorological variables (shortwave and longwave radiation, air temperature, relative humidity, wind speed, precipitation and its phase) to differing slopes, aspects, elevations and groundcovers defined by hydrological response units (HRU), using in-built algorithms and macros (Ellis et al., 2010; Harder and Pomeroy, 2013). The HRUs used to discretize these basins are presented in Pradhananga and Pomeroy (2022). CRHM-glacier simulates the hydrology of both glacier and non-glacier areas in a basin. It redistributes snow by coupling the blowing snow transport and sublimation processes with snow avalanching. The implementation used here does not calculate ice flow. Melt energies for snow and icemelt are calculated separately, based on *Snobal* and energy budget glacier melt modules respectively, to calculate snow, firn and ice mass balances (Pradhananga and Pomeroy, 2022). Meltwater routing is through three glacier reservoirs (snow, firn, and ice) modified to the de Woul *et al.* (2006) approach. Once water leaves the glacier, rain and meltwater are routed further into the soil surface, subsurface and groundwater using well developed alpine routing routines (Fang *et al.*, 2013). The model includes calculation of actual evapotranspiration using the Penman-Monteith method, and soil moisture and groundwater dynamics based on infiltration to frozen and unfrozen soils, and use of saturated and unsaturated hydraulic conductivities to calculate flow velocities in porous media. The model parameters are not calibrated from its fit to streamflow. Parameters are selected primarily from local and regional observations (e.g., temperature and precipitation lapse rates) and its operation has been verified against observed albedo, mass balance, melt rate and streamflow at PGRB and AGRB (Pradhananga and Pomeroy, 2022).

2.4 Modelling scenarios

CRHM-glacier was run to simulate the hydrological responses of the two glacier research basins to four experimental scenarios. Observed glacier configurations, “past” corresponding to 1966 (PGRB) and 1983-1984 (AGRB) and “present” to 2014 (PGRB) and 2011-2014 (AGRB), were set as initial conditions for two modelling scenarios in each basin. Initializing the model with observed glacier configurations, and restricting simulation periods to one decade, compensated for the lack of ice flow dynamics in the model scenarios.

Glacier configuration maps for the two periods were prepared from available DEM and landcover information. These maps were used to delineate the configuration, namely the accumulation (snow or firn-covered) and ablation (ice-exposed) areas and to determine ice elevations, slopes and aspects. The glacier area was taken as the sum of the snow, firn and the ice exposed areas. The accumulation area ratio (AAR) of the glaciers was calculated as the ratio of snow/firn-covered area to the total glacier area.

A topographic map of Peyto Glacier from 1966 (Sedgwick and Henoeh, 1975) was used to prepare the past glacier configuration scenario. Both the 1966 DEM (10 m resolution) and the 1966 landcover map were developed from this topographic map, which was produced from aerial photographs taken in August 1966. The 2014 DEM was prepared at 10 m resolution from aerial photographs taken during July and September 2014 by Parks Canada over Banff National Park and made available for this study. The landcover map for the present basin was prepared based upon the Landsat image taken on 18 August 2014. Bolch *et al.* (2010) found only 1.7% deviation between aerial photograph and satellite imagery approaches for delineating glacier area and snow/firn/ice coverage. For AGRB, DEM and landcover map were not available from the 1960s. Therefore, DEM at 20 m horizontal resolution from 1983 (Canadian Digital Elevation Model, CDEM) and the satellite image taken on 28 July 1984

(Landsat 5) was used to prepare the past glacier configuration. The 2011 DEM (20 m resolution from JAXA) and the 2014 satellite image (Landsat 8) taken on 18 August were used to prepare the present glacier configuration.

The models were then run for two climate periods, past (1965-1975) and present (2008-2018). A novel approach was used in that past and present climate forced both past and present glacier configurations to diagnose how past glaciers would respond to the present-day climate and present-day glaciers to the past climate. Therefore, there was a combination of four model simulation scenarios, A-D using two separate decades of climate forcing from past and present periods, with past and present glacier configurations:

A: Past Climate, Past Glacier

B: Past Climate, Present Glacier

C: Present Climate, Past Glacier

D: Present Climate, Present Glacier

Based on these four experimental scenarios (A-D), five comparisons (C1-C5, Table 1) were employed to diagnose the impacts of climate and glacier changes on streamflow. C1 represents realistic conditions of both climate and glacier configuration; it compares A and D, i.e., past climate – past glacier with present climate – present glacier. The other comparisons are falsified modeling experiments to segregate the impacts of changing climate and glacier configurations. C2 and C5 consider only changes in the glacier configuration, while keeping the climate fixed in either the past or present. C3 and C4 compare the impacts from changing the climate while keeping the glacier configuration constant, as either the past or present glacier. Simulated runoff from these model outputs was examined to diagnose the hydrological response to both glacier change and climate change.

Statistical tests, the Student's t-test and the Wilcoxon Signed-Rank test (Wilcoxon, 1945) in the R environment (R Core Team, 2017) were applied to test the significance of the changes between the results obtained from the model scenarios. These tests were used to assess whether one population of model output metrics is statistically distinguishable from the other. All tests were conducted at the 5% level of significance.

3 Results

3.1 Change in climate

Air temperature and precipitation over PGRB and AGRB were analyzed for the two decades – 1965-1975 (past) and 2008-2018 (present). Daily mean (T_{mean}), maximum (T_{max}), and minimum (T_{min}) temperature (Figure 2 and Table ST2-ST4) and monthly precipitation (Figure 3 and Table ST1), averaged and aggregated over the two climatic periods, were compared using C1.

annual and seasonal average temperatures generally increased in the present decade compared to the past except for the summer maximum at AGRB. Annual T_{max} increased by 1 °C at AGRB and by 1.5 °C at PGRB. Analysis for monthly time periods also shows that air temperatures have increased at both glaciers. The greatest increases in the monthly T_{max} were in January, when they rose by 5.1 °C at AGRB and 4.8 °C at PGRB. Monthly T_{min} at AGRB increased by 3.6 °C in January and by 1 °C in July, and that at PGRB increased by 3.5 °C in January and

1.9 °C in July. Annual T_{mean} at AGRB increased by 0.5 °C, and that at PGRB increased by 1 °C. Temperature differences over time that were found to be significantly different from zero are more consistently evident at PGRB than at AGRB.

5 Annual precipitation increased slightly in the present decade compared to the past (Figure 3 and Table ST1), 15.2 mm (2.5%) at AGRB and 71.7 mm (10.3%) at PGRB. but these differences are not statistically significant. Winter (Dec-Feb) precipitation decreased at both basins, by 49.5 mm at AGRB and by 35.2 mm at PGRB. But precipitation in the other seasons (spring, summer and fall) increased by 13.1 mm, 38.5 mm and 13.1 mm respectively at AGRB and by 29.7 mm, 50.2 mm and 27.0 mm respectively at PGRB. The increases in summer
10 precipitation at both basins and decrease in winter precipitation at AGRB were statistically significant (Table ST1).

There was a statistically significant increase in rainfall ratio (rainfall divided by total precipitation), for D compared A (C1) in both research basins (Figure 4), from 0.161 to 0.232 (44%) at AGRB and from 0.220 to 0.283
15 (29%) at PGRB (Table ST5). The average annual rainfall ratios increased with reductions in glacier area and surface elevation; at AGRB from 0.161 to 0.163 for the past climate (C2) and from 0.163 to 0.232 for the present climate (C5); and at PGRB from 0.220 to 0.283 for the past climate (C2) and from 0.268 to 0.283 for the present climate (C5). However, the increase in rainfall ratios was mainly due to changes in climate (temperature increases) over time. The average annual rainfall ratio for the past glacier configuration (C3) increased from 0.161 to 0.229
20 at AGRB (statistically significant) and from 0.220 to 0.268 at PGRB; and for the present glacier configuration (C4), the ratio increased from 0.163 to 0.232 at AGRB (statistically significant) and from 0.234 to 0.283 at PGRB.

3.2 Change in glacier configuration

Figure 5 compares the configuration of both glaciers between two periods and shows changes in accumulation
25 and ablation areas. During the period 1966-2014, the area of Peyto Glacier shrank from 14.4 km² to 9.9 km² (31%) and its AAR dropped from 0.75 to 0.35, exposing more ice in 2014, more than double the area exposed in 1966. The exposed ice area increased from 3.6 km² to 6.4 km² (78%), whereas the snow/firn area decreased from 10.8 km² to 3.5 km². Though to a lesser degree than Peyto Glacier, the area of Athabasca Glacier also decreased from 18 km² to 16.9 km² (6%), and its AAR decreased from 0.76 to 0.47. The exposed ice area of Athabasca
30 increased from 4.3 km² to 9.0 km² (109%), and the snow/firn area decreased from 13.6 km² to 7.9 km². The firn line moved to a higher elevation in both glaciers, and glacier surfaces have become steeper from the past to present period, slope of glacier increased from 14.2 to 16.8 in AGRB and that increased in PGRB from 19.9 to 22.0. The other change in the two glacier configurations was in elevation; the mean glacier surface elevation Peyto Glacier decreased from 2628 m to 2615 m, whilst that of Athabasca Glacier increased from 2799 m to 2826 m.

35

3.3 Change in glacier mass balance

Seasonal and annual mass balance for AGRB and PGRB from the four model scenarios, A-D, are presented in Figure 6. The results from the statistical analysis are presented in Table ST7. Except for the change in winter mass balance between A and D (C1), the mass balance changes are not statistically significant at AGRB. There were
40 significant changes in winter and annual mass balances between past and present climates and glaciers at PGRB

(C1). Mean annual winter accumulation decreased from an average of 586 mm [1965-1975] to 324 mm [2008-2018], resulting in negative mean annual mass balances, from -271 mm in the past climate to -733 mm in the present climate (Table ST7). These changes are more due to the change in climate than the change in glacier configuration. Summer ablation increased significantly from past to present climate for both past and present glacier configurations (C3 and C4). The changes are not statistically significant in model runs for C2 and C5. However, the past glacier configurations resulted in greater winter snow accumulations in both basins for both past and present climates.

3.4 Change in runoff and runoff generation processes

Figure 7 shows runoff and volumetric melt components of runoff from AGRB and PGRB for A and D scenarios (C1). Snowmelt runoff dominated both the basins, in comparison to rainfall runoff, icemelt runoff, and firmmelt runoff. The present climate and present glacier configurations (D) produced more runoff than the past climate and past glacier configurations (A). There was a 19% significant increase ($p=0.005$) in mean annual runoff, from 1581 mm to 1888 mm at PGRB (Table ST5). This was mainly due to an increased contribution from icemelt, from 265 mm to 667 mm. There was a decrease in mean annual snowmelt and firmmelt, but the changes in these and the other fluxes were not statistically significant. For AGRB, the 3% increase in runoff from 1320 mm to 1365 mm was not statistically significant ($p=0.578$, Table ST5), though there was a significant increase in rainfall from 175 mm to 262 mm. Monthly statistical values are shown in Table ST6. AGRB experienced increased snowmelt and firmmelt but decreased icemelt.

Snowmelt dominated basin runoff, providing 52-70% and 52-53% of all runoff for PGRB and AGRB, respectively. At PGRB, rainfall-runoff provided 19-20% of basin runoff, which exceeded icemelt and firmmelt contributions, except for icemelt in the current climate and glacier configuration. The sum of firmmelt and icemelt contributions at PGRB increased from 27% in the past to 43% in the present (C1). At AGRB, rainfall-runoff provided 13-19% of basin runoff and was smaller than either firmmelt or icemelt contributions. Firmmelt was smaller than both snowmelt and icemelt. The sum of firmmelt and icemelt contributions at AGRB decreased slightly from 61% in the past to 59% in the present. Basin runoff increased 19% at PGRB but only 3% at AGRB from the past to present climate for the present glacier configuration (C1). This increase was due to increases in rainfall (statistically significant), snowmelt and firmmelt at AGRB, whereas it was due to increase in rainfall and a remarkable 152% increase in icemelt (statistically significant) at PGRB (Table ST5). Snowmelt declined 12% at PGRB. Basin runoff increased due to the warming climate for both past and present glaciers (C3 and C4), whereas it decreased with decreasing glacier configurations for both the past and present climates (C2 and C5).

Monthly averaged runoff from the four model scenarios is presented in Figure 8. The glaciers in AGRB and PGRB produced more runoff in the recent period (D) than in the past (A). Both past and present glaciers produced more runoff in the present than in the past climate. However, the past glaciers generated more streamflow compared to the present glaciers with both present and past climates. As the climate shifted there was a shift in peak from August in the past (A) to July in the present (D) at PGRB.

Significant changes in runoff, firn melt and snowmelt occurred for PGRB, suggesting that the increase in runoff over time (C3 and C4) was due to an increase in firn melt and icemelt (Table ST5). The large loss of firn coverage from past to present at PGRB resulted in a decrease of firn melt by 65% (from 414 mm to 146 mm), when the past glacier configuration was replaced by the present one, with the climate for both glacier configurations held constant at the present climate (C5). In the case of AGRB, only increases in rainfall in the C3 and C4 and in snowmelt in C4 were significant. More rainfall occurred with both past and present glacier configurations.

4 Discussion

The glacio-hydrological model, CRHM, was used to simulate headwater hydrology of two glacierized mountain basins with four scenarios by combining the climate and glacier configurations of two periods, mid-20th century and early 21st century. These scenarios were used to diagnose the hydrological responses to changes in glacier and climate, individually and jointly presenting how glaciers in the past would respond to present day climate and glaciers today to past climate.

Changes in glacier configurations included changes in glacier extent, glacier surface elevation and slope, and change in AAR, which also changed the albedo feedback of glacier surfaces. Compared to the past, present glacier areas have declined, elevations have decreased/increased, surface slopes have become steeper, AARs have become smaller, exposing more ice, and thus reducing the glacier albedos.

Air temperatures increased from past to present; the changes were more often significant at PGRB than at AGRB. Though there was not much change in the total precipitation, rainfall ratios increased in the present compared to past due to changes in both climate and glacier configuration and this caused reduced snowfall at both basins. The differences for the same climate were due to the change in glacier configurations, for instance the present glacier configurations feature a smaller glacier surface at a lower surface height than do past configurations; both factors contribute to increased rainfall ratios. Increases in the rainfall ratio in these basins are consistent with other studies, for example, in western Canada by DeBeer *et al.*, (2016) and in Europe by Hynčica and Huth (2019). In the case of glacier mass balance, the present climate and present glacier were responsible for making the winter mass balance less positive. Compared to the present glacier configuration, the past glacier gained more mass during winter.

There was a reduction in peak monthly flows from both basins as the glacier area declined over time with the climate held constant (Figure 8). However, with the climate changing and the glacier configuration held constant, peak monthly flows increased over time. The combination of moving from past to present climate and changing glacier configuration shifted peak monthly flows forward by a month at PGRB, but the impact of changing climate was greater than that of the changing glacier configuration on this shift. The shift in PGRB's peak monthly flow from August to July is consistent with the future prediction by Kienzle *et al.* (2012) for the Cline River watershed that spring runoff and peak streamflow would shift 18–26 days advance in the 21st century (2020–2080) compared to the baseline period (1961-1989).

In general, there was increased runoff in the present compared with the past for both basins, particularly from PGRB. This shows that these glaciers in Alberta, in the cold, high elevation headwaters on the northeastern slopes of the Canadian Rockies, are still in the initial phase of warming-induced increased runoff, in contrast to most
5 glaciers in British Columbia, including the more temperate southwestern slopes of the Canadian Rockies (Stahl and Moore, 2006) and western Canada, in general (DeBeer et al., 2016). However, this is similar to the results of Chernos et al. (2020), who projected glacier contributions to streamflow to increase till the middle of the twenty-first century (2040) and to decrease thereafter for the Athabasca River. Moore et al. (2009) also noted increasing trends of runoff in glacierized basins in relatively colder northwest British Columbia and southwest Yukon.
10 Casassa et al. (2009) generalized that high elevation basins and/or basins with high glacierization were experiencing increasing runoff trends around the world.

The novel model falsification approach of using past glacier configuration with present climate and present glacier configuration with past climate provided a means to separate the combined and individual impacts of climate
15 change and glacier recession on the headwater hydrology and water balance of glacierized basins. With fixed glacier configurations (C3 and C4), the impact of climate on glacier hydrology could be analyzed and using the same climate to force past and present glacier configuration models (C2 and C5) provided an explicit separation of the impacts of changes in glacier configuration on glacier hydrology.

20 The warmer temperatures and increased rainfall ratio in the present climate led to increased glacier runoff from both basins. However, the reduced glacier extent in the present glacier configuration resulted in decreased runoff from both basins, counteracting the direct impacts of climate change on the basins. In summary, the outputs show that changes in both climate and basin configurations were causing changes in the melt rate and runoff. Compared to the past climate and past glacier configuration, present climate and present glacier configuration provided more
25 runoff in both basins, although there were significant losses of glacier area over the last five decades.

5 Conclusions

30 This study investigated the influence of snow and glaciers on headwater hydrology in two mountain basins in the Canadian Rockies, where a warming climate and glacier retreat continue to cause concern about changes in high mountain hydrology.

There was an increase in air temperature, mainly in daily maximum and winter minimum temperatures. Total
35 precipitation has not increased, but the rainfall ratio has increased with the shift in climate. Both present climate and present glacier caused an increase in rainfall ratio compared to past climate and past glacier. Decreases in winter precipitation were balanced by increased precipitation in the other seasons, some of which fell as rainfall. Both mass balance studies and analysis of satellite imagery show that the glaciers are losing mass and area, and that the exposure of ice at the glacier surfaces has increased. The fractional glacier retreat is lower at AGRB (6%)
40 than at PGRB (31%), whereas the fractional change in the exposure of ice is higher at AGRB (109%) than at

PGRB (78%). The retreat of the glaciers has led to reductions in glacierized areas and changes in elevation and slope of the glacier surfaces.

5 The study used a novel approach to apply present climate forcings to drive hydrological modelling using past glacier configurations and past climate forcings to drive modelling using present glacier configuration, so that the impacts of changes in glacier configuration and climate on glacier hydrology could be explicitly separated. The modelling results presented here show that glacier retreat and ablation are due to the joint effect of warming climate and an increase in ice exposure, which increased both seasonal melt and runoff. The sum of firmmelt and icemelt contributions to annual discharges increased from the past climate and glacier to the present climate and glacier at PGRB from 27 to 43% and decreased slightly at AGRB from 61% to 59%. Increased streamflow
10 discharges (3-19%) were due to climate warming and were somewhat limited by glacier retreat and areal decreases. The Model results indicated that streamflow from the glaciers increased in the present climate (2008-2018) from the past climate (1965-1975) despite reductions in glacier area and volume. Such a modelling approach is important for diagnosing the hydrological responses from a glacierized basin in the context of climate change
15 and variability. The results suggest that increased exposure of glacier ice and lower surface elevation due to glacier thinning were less influential than climate warming in increasing streamflow.

Author contribution

20 DP and JWP conceptualized the research. DP did the analysis and prepared the manuscript. JWP instrumented the research basins, guided research methods, and edited and revised the manuscript.

Competing interests

The authors declare that they have no conflict of interest.

25

Acknowledgements

The authors wish to acknowledge the decades of extremely challenging field research at Peyto and Athabasca glacier basins by dozens of scientists and students over the decades. Meriting special mention are Dr. D. Scott Munro of the University of Toronto and Michael N. Demuth of Natural Resources Canada whose legacy of
30 research on Peyto Glacier including meteorological and mass balance observations made this study possible. Special mention also goes to the prescient scientists of the International Hydrological Decade who established Peyto Glacier as a research site, including Dr. Gordon Young of Environment Canada and Wilfrid Laurier University. Additional data were provided by Environment and Climate Change Canada's Meteorological Service of Canada and Water Survey of Canada, Parks Canada and Alberta Environment and Parks. Tom Brown and Xing
35 Fang of the Centre for Hydrology supported CRHM development and operation/parameterization, respectively. Nammy Hang Kirat of The Small Earth Nepal supported satellite image interpretation and GIS maps. Funding for this study was provided by the Canada Research Chairs, Canada Foundation for Innovation, Natural Sciences and Engineering Research Council of Canada through its Discovery Grants and the Changing Cold Regions Network and the Canada First Research Excellence Fund's Global Water Futures Programme. Special thanks go to the

reviewers (anonymous reviewer#1 and Dr. Eleanor Bash) for their careful and critical review of the manuscript and their several insightful comments and constructive suggestions, which helped to improve the manuscript.

References

- Barry, R. G.: The status of research on glaciers and global glacier recession: a review, *Prog. Phys. Geogr.*, 30(3), 285–306, doi:10.1191/0309133306pp478ra, 2006.
- Bolch, T., Menounos, B. and Wheate, R.: Landsat-based inventory of glaciers in western Canada, 1985–2005, *Remote Sens. Environ.*, 114(1), 127–137, doi:10.1016/j.rse.2009.08.015, 2010.
- Casassa, G., López, P., Pouyaud, B. and Escobar, F.: Detection of changes in glacial run-off in alpine basins: Examples from North America, the Alps, central Asia and the Andes, *Hydrol. Process.*, 41, 31–41, doi:10.1002/hyp.7194, 2009.
- Castellazzi, P., Burgess, D., Rivera, A., Huang, J., Longuevergne, L. and Demuth, M. N.: Glacial Melt and Potential Impacts on Water Resources in the Canadian Rocky Mountains, *Water Resour. Res.*, 55(12), 10191–10217, doi:10.1029/2018WR024295, 2019.
- Chernos, M., MacDonald, R. J., Nemeth, M. W. and Craig, J. R.: Current and future projections of glacier contribution to streamflow in the upper Athabasca River Basin, *Can. Water Resour. J. / Rev. Can. des ressources hydriques*, 45(4), 324–344, doi:10.1080/07011784.2020.1815587, 2020.
- Clarke, G. K. C., Jarosch, A. H., Anslow, F. S., Radić, V. and Menounos, B.: Projected deglaciation of western Canada in the twenty-first century, *Nat. Geosci.*, 8(5), 372–377, doi:10.1038/ngeo2407, 2015.
- Comeau, L. E. L., Pietroniro, A. and Demuth, M. N.: Glacier contribution to the North and South Saskatchewan Rivers, in *Hydrological Processes*, vol. 23, pp. 2640–2653., 2009.
- DeBeer, C. M., Wheeler, H. S., Carey, S. K. and Chun, K. P.: Recent climatic, cryospheric, and hydrological changes over the interior of western Canada: a synthesis and review, *Hydrol. Earth Syst. Sci.*, 20, 1573–1598, doi:10.5194/hess-20-1573-2016, 2016.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kállberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J. N. and Vitart, F.: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, 137(April), 553–597, doi:10.1002/qj.828, 2011.
- Demuth, M. N. and Keller, R.: An assessment of the mass balance of Peyto glacier (1966–1995) and its relation to Recent and past-century climatic variability, in *Peyto Glacier: One Century of Science*, edited by M. N. Demuth, D. S. Munro, and G. J. Young, pp. 83–132, National Hydrology Research Institute, Saskatoon, Saskatchewan., 2006.
- Ellis, C. R., Pomeroy, J. W., Brown, T. and MacDonald, J.: Simulation of snow accumulation and melt in needleleaf forest environments, *Hydrol. Earth Syst. Sci.*, 14(6), 925–940, doi:10.5194/hess-14-925-2010, 2010.
- Fang, X. and Pomeroy, J. W.: Diagnosis of future changes in hydrology for a Canadian Rockies headwater basin, *Hydrol. Earth Syst. Sci.*, 24, 2731–2754, doi:10.5194/HESS-24-2731-2020, 2020.
- Fang, X., Pomeroy, J. W., Ellis, C. R., MacDonald, M. K., DeBeer, C. M. and Brown, T.: Multi-variable evaluation of hydrological model predictions for a headwater basin in the Canadian Rocky Mountains, *Hydrol. Earth Syst. Sci.*, 17(4), 1635–1659, doi:10.5194/hess-17-1635-2013, 2013.

- Fountain, A. G. and Tangborn, W. V.: The effect of glaciers on streamflow variations, *Water Resour. Res.*, 21(4), 579–586, doi:10.1029/WR021i004p00579, 1985.
- Gudmundsson, L.: qmap: Statistical transformations for post-processing climate model output. R package version 1.0-4., R Packag. version 1.0-4, 2016.
- 5 Harder, P. and Pomeroy, J. W.: Estimating precipitation phase using a psychrometric energy balance method, *Hydrol. Process.*, 27(May), 1901–1914, doi:10.1002/hyp.9799, 2013.
- Harder, P., Pomeroy, J. W. and Westbrook, C. J.: Hydrological resilience of a Canadian Rockies headwaters basin subject to changing climate, extreme weather, and forest management, *Hydrol. Process.*, 29(18), 3905–3924, doi:10.1002/hyp.10596, 2015.
- 10 Hopkinson, C. and Young, G. J.: The effect of glacier wastage on the flow of the Bow River at Banff, Alberta, 1951–1993, *Hydrol. Process.*, 12, 1745–1762, doi:10.1002/(SICI)1099-1085(199808/09)12:10/11<1745::AID-HYP692>3.0.CO;2-S, 1998.
- Hynčica, M. and Huth, R.: Long-term changes in precipitation phase in Europe in cold half year, *Atmos. Res.*, doi:10.1016/j.atmosres.2019.04.032, 2019.
- 15 Intsiful, A. and Ambinakudige, S.: Glacier Cover Change Assessment of the Columbia Icefield in the Canadian Rocky Mountains, Canada (1985–2018), *Geosciences*, 11(19), doi:10.3390/GEOSCIENCES11010019, 2021.
- Kehrl, L. M., Hawley, R. L., Osterberg, E. C., Winski, D. A. and Lee, A. P.: Volume loss from lower Peyto Glacier, Alberta, Canada, between 1966 and 2010, *J. Glaciol.*, 60(219), 51–56, doi:10.3189/2014JoG13J039, 2014.
- 20 Kienzle, S. W., Nemeth, M. W., Byrne, J. M. and Macdonald, R. J.: Simulating the hydrological impacts of climate change in the upper North Saskatchewan River basin, Alberta, Canada, *J. Hydrol.*, 412–413, 76–89, doi:10.1016/j.jhydrol.2011.01.058, 2012.
- Krogh, S. A. and Pomeroy, J. W.: Recent changes to the hydrological cycle of an Arctic basin at the tundra-taiga transition, *Hydrol. Earth Syst. Sci.*, 22, 3993–4014, doi:10.5194/hess-22-3993-2018, 2018.
- 25 Marks, D., Winstral, A., Flerchinger, G., Reba, M., Pomeroy, J., Link, T. and Elder, K.: Comparing Simulated and Measured Sensible and Latent Heat Fluxes over Snow under a Pine Canopy to Improve an Energy Balance Snowmelt Model, *J. Hydrometeorol.*, 9(6), 1506–1522, doi:10.1175/2008JHM874.1, 2008.
- Marshall, S. J., White, E. C., Demuth, M. N., Bolch, T., Wheate, R., Menounos, B., Beedle, M. J. and Shea, J. M.: Glacier Water Resources on the Eastern Slopes of the Canadian Rocky Mountains, *Can. Water Resour. J.*, 36(March 2010), 109–134, doi:10.4296/cwrj3602823, 2011.
- 30 Moore, R. D., Fleming, S. W., Menounos, B., Wheate, R., Fountain, A., Stahl, K., Holm, K. and Jakob, M.: Glacier change in western North America: influences on hydrology, geomorphic hazards and water quality, *Hydrol. Process.*, 23, 42–61, doi:10.1002/hyp.7162, 2009.
- Munro, D. S.: Peyto Creek hydrometeorological database (Peyto Creek Base Camp AWS), IP3 Arch. [online] Available from: www.usask.ca/ip3/data, 2011.
- 35 Neupane, R. P., Adamowski, J. F., White, J. D. and Kumar, S.: Future streamflow simulation in a snow-dominated Rocky Mountain headwater catchment, *Hydrol. Res.*, 49(4), 1172–1190, doi:10.2166/NH.2017.024, 2018.
- Pomeroy, J. W., Gray, D. M., Brown, T., Hedstrom, N. R., Quinton, W. L., Granger, R. J. and Carey, S. K.: The cold regions hydrological model: a platform for basing process representation and model structure on physical evidence, *Hydrol. Process.*, 21(19), 2650–2667, doi:10.1002/hyp.6787, 2007.
- 40

- Pradhananga, D. and Pomeroy, J. W.: Diagnosing changes in glacier hydrology from physical principles using a hydrological model with snow redistribution, sublimation, firnification and energy balance ablation algorithms, *J. Hydrol.*, 608(127545), doi:<https://doi.org/10.1016/j.jhydrol.2022.127545>, 2022.
- Pradhananga, D., Pomeroy, J., Aubry-Wake, C., Munro, D. S., Shea, J., Demuth, M., Kirat, N. H., Menounos, B. and Mukherjee, K.: Hydrometeorological, glaciological and geospatial research data from the Peyto Glacier Research Basin in the Canadian Rockies, *Earth Syst. Sci. Data*, 13, 2875–2894, doi:10.5194/essd-13-2875-2021, 2021.
- R Core Team: R: A language and environment for statistical computing, [online] Available from: <https://www.r-project.org/>, 2017.
- Reynolds, J. R. and Young, G. J.: Changes in areal extent, elevation and volume of Athabasca Glacier, Alberta, Canada, as estimated from a series of maps produced between 1919 and 1979, *Ann. Glaciol.*, 24, 60–65, 1997.
- Riedel, J. L., Wilson, S., Baccus, W., Larrabee, M., Fudge, T. J., Fountain, A. and Riedel, C. J. L.: Glacier status and contribution to streamflow in the Olympic Mountains, Washington, USA, *J. Glaciol.*, 61(225), 8–16, doi:10.3189/2015JoG14J138, 2015.
- Schiefer, E., Menounos, B. and Wheate, R.: Recent volume loss of British Columbian glaciers, Canada, *Geophys. Res. Lett.*, 34(16), 1–6, doi:10.1029/2007GL030780, 2007.
- Sedgwick, J. K. and Henoch, W. E. S.: 1966 Peyto Glacier Map, Banff National Park, Alberta. Environment Canada, IWD 1010, 1:10,000., 1975.
- Stahl, K. and Moore, R. D.: Influence of watershed glacier coverage on summer streamflow in British Columbia, Canada, *Water Resour. Res.*, 42(W06201), 1–5, doi:10.1029/2006WR005022, 2006.
- Tennant, C. and Menounos, B.: Glacier change of the Columbia Icefield, Canadian Rocky Mountains, 1919–2009, *J. Glaciol.*, 59(216), 671–686, doi:10.3189/2013JoG12J135, 2013.
- Tennant, C., Menounos, B., Wheate, R. and Clague, J. J.: Area change of glaciers in the Canadian rocky mountains, 1919 to 2006, *Cryosphere*, 6, 1541–1552, doi:10.5194/tc-6-1541-2012, 2012.
- Uppala, S. M., Kållberg, P. W., Simmons, A. J., Andrae, U., Bechtold, V. D. C., Fiorino, M., Gibson, J. K., Haseler, J., Hernandez, A., Kelly, G. A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R. P., Andersson, E., Arpe, K., Balmaseda, M. A., Beljaars, A. C. M., Berg, L. Van De, Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Hólm, E., Hoskins, B. J., Isaksen, I., Janssen, P. A. E. M., Jenne, R., McNally, A. P., Mahfouf, J.-F., Morcrette, J.-J., Rayner, N. A., Saunders, R. W., Simon, P., Sterl, A., Trenberth, K. E., Untch, A., Vasiljevic, D., Viterbo, P. and Woollen, J.: The ERA-40 re-analysis, *Q. J. R. Meteorol. Soc.*, 131(612), 2961–3012, doi:10.1256/qj.04.176, 2005.
- Wilcoxon, F.: Individual Comparisons by Ranking Methods, *Biometrics Bull.*, 1(6), 80, doi:10.2307/3001968, 1945.
- de Woul, M., Hock, R., Braun, M., Thorsteinsson, T., Jóhannesson, T. and Halldórsdóttir, S.: Firm layer impact on glacial runoff: a case study at Hofsjökull, Iceland, *Hydrol. Process.*, 20(10), 2171–2185, doi:10.1002/hyp.6201, 2006.




Figure 1: Location map of Peyto and Athabasca glacier research basins. Gray line is the border between two provinces, blue polygons are basin boundaries, and green pentagons are meteorological stations.

5





Figure 2: Monthly means of daily maximum (a, b), minimum (c, d) and mean (e, f) temperature comparison between two periods: past (1965-1975) and present (2008-2018). (a, c, and e) AGRB (b, d, f) PGRB.



5 **Figure 3: Monthly mean precipitation averaged over the two periods: past (1965-1975) and present (2008-2018). (a) AGRB (b) PGRB.**




Figure 4: Mean-monthly rainfall ratios simulated for the four model run scenarios, A-D combining past and present climate and glacier. (a) AGRB, (b) PGRB.




Figure 5: Change in landcover between the glaciers in past and present. (a) and (b) are AGRB in 1984 and 2014, respectively; (c) and (d) are PGRB in 1966 and 2014, respectively.




Figure 6: Glacier mass balance – winter, summer and annual, from the four model scenarios, A-D combining past and present climate and glacier. Error bars show the annual variability, defined as the standard error between years. (a) AGRB and (b) PGRB.

5





Figure 7: Mean annual melt, rainfall-runoff and basin runoff for the past and present glacier configuration and climate scenarios. Error bars show the annual variability, defined as the standard error between years. (a) AGRB (b) PGRB.



5

Figure 8: Monthly averaged runoff simulated using the past and present glacier and climate scenarios. (a) AGRB (b) PGRB.

Table 1: Comparison of model outputs.

Comparison #	Comparison of model scenarios		
C1	Past Climate-Past Glacier [A]	<i>VERSUS</i>	Present Climate-Present Glacier [D]
C2	Past Climate-Past Glacier [A]	<i>VERSUS</i>	Past Climate-Present Glacier [B]
C3	Past Climate-Past Glacier [A]	<i>VERSUS</i>	Present Climate-Past Glacier [C]
C4	Past Climate-Present Glacier [B]	<i>VERSUS</i>	Present Climate-Present Glacier [D]
C5	Present Climate-Past Glacier [C]	<i>VERSUS</i>	Present Climate-Present Glacier [D]

5