

Hydrology and riparian forests drive carbon and nitrogen supply and DOC:NO₃⁻ stoichiometry along a headwater Mediterranean stream

José L. J. Ledesma^{1,2}, Anna Lupon², Eugènia Martí², Susana Bernal²

5 ¹Institute of Geography and Geoecology, Karlsruhe Institute of Technology (KIT), Karlsruhe, 76131, Germany

²Centre for Advanced Studies of Blanes, Spanish National Research Council (CEAB-CSIC), Blanes, 17300, Spain

Correspondence to: José L. J. Ledesma (jose.ledesma@kit.edu)

Abstract. In forest headwater streams, metabolic processes are predominately heterotrophic and depend on both the availability of carbon (C) and nitrogen (N) and a favourable C:N stoichiometry. In this context, hydrological conditions and the presence of riparian forests adjacent to streams can play an important, yet understudied role determining dissolved organic carbon (DOC) and nitrate (NO₃⁻) concentrations and DOC:NO₃⁻ molar ratios. Here, we aimed to investigate how the interplay between hydrological conditions and riparian forest coverage drives DOC and NO₃⁻ supply and DOC:NO₃⁻ stoichiometry in an oligotrophic headwater Mediterranean stream. We analysed DOC and NO₃⁻ concentrations, and DOC:NO₃⁻ molar ratios during both base flow and storm flow conditions at three stream locations along a longitudinal gradient of increased riparian forest coverage. Further, we performed an event analysis to examine the hydroclimatic conditions that favour the transfer of DOC and NO₃⁻ from riparian soils to the stream during storms. Stream DOC and NO₃⁻ concentrations were generally low (overall average ± SD was 1.0 ± 0.6 mg C L⁻¹ and 0.20 ± 0.09 mg N L⁻¹), although significantly higher during storm flow compared to base flow conditions in all three stream sites. Optimal DOC:NO₃⁻ stoichiometry for stream heterotrophic microorganisms (corresponding to DOC:NO₃⁻ molar ratios between 4.8 and 11.7) was prevalent at the midstream and downstream sites under both flow conditions, whereas C-limited conditions were prevalent at the upstream site, which had no surrounding riparian forest. The hydroclimatic analysis of storms suggest that large and medium storm events display a distinct mechanism of DOC and NO₃⁻ mobilization. In comparison to large storms, medium storm events showed limited hydrological responses that led to significantly lower stream DOC and NO₃⁻ concentrations. During large storm events, different patterns of DOC and NO₃⁻ mobilization depending on antecedent soil moisture conditions arise: drier antecedent conditions promoted rapid elevations of riparian groundwater tables, hydrologically activating a wider and shallower soil layer, and leading to relatively higher increases in stream DOC and NO₃⁻ concentrations compared to events preceded by wet conditions. Our results suggest that (i) increased supply of limited resources during storms could sustain in-stream heterotrophic activity during high flows, especially during large storm events preceded by dry conditions, and (ii) C-limited conditions upstream were overcome downstream, likely due to higher C inputs from riparian forests present at lower elevations. The contrasting spatiotemporal patterns in DOC and NO₃⁻ availability and DOC:NO₃⁻

stoichiometry observed at the study stream suggests that groundwater inputs from riparian forests are essential for maintaining in-stream heterotrophic activity in oligotrophic, forest headwater catchments.

1 Introduction

The metabolism of lotic systems is highly dependent on environmental conditions (Bernhardt et al., 2018). In forest headwater streams, metabolism is dominated by the activity of heterotrophs because light availability is limited for primary producers and allochthonous matter inputs constitute the major energy source (Tank et al., 2018). In these environments, the supply and availability of carbon (C) and nutrients, including nitrogen (N), can constrain both the activity and composition of stream heterotrophic microorganisms (Dodds et al., 2002; Brookshire et al., 2005; Fasching et al., 2020). Consequently, biotic assimilation of both dissolved organic carbon (DOC) and nitrate (NO_3^-) have been shown to relate closely with ecosystem respiration in streams worldwide (Pastor et al., 2014; Catalán et al., 2018; Tank et al., 2018; Lupon et al., 2020a), which highlights the strong linkages between resource supply, demand, and in-stream heterotrophic activity.

Yet, even if DOC and NO_3^- are available, DOC: NO_3^- stoichiometry can further influence in-stream heterotrophic activity (Elser et al., 2000; Helton et al., 2015). Heterotrophic organisms can adapt to a wide range of resource stoichiometric ratios (Manzoni et al., 2017), and thus it is difficult to establish general reference values defining optimal conditions for in-stream heterotrophic activity. For example, global studies on stoichiometric controls for a wide range of ecosystems show discrepant reference C:N molar ratios for streams and rivers; ranging from 4.8 (Taylor and Townsend, 2010) to 11.7 (Helton et al., 2015). Modelling approaches using bacterial cultures show that heterotrophic microorganisms can perform well under C:N molar ratios ranging between 5.6 and 18.4 (Danger et al., 2008). Empirical case studies in forest headwater streams reported a C:N range of 5.9 to 13.4 for optimal microbial activity (Pastor et al., 2014). Hence, considering that ecological stoichiometry theory sets biochemical constraints for metabolic activity based on the comparison of element ratios between resources and organisms (Sterner and Elser, 2002), there is a wide range of potentially optimal stoichiometric conditions for microbial heterotrophs. Nevertheless, C:N molar ratios outside this range may lead to either C- or N-limitation for heterotrophic activity.

In this context, hydrological processes at the riparian-stream interface, and in particular the mobilization of solutes from riparian soils to streams, can play an important role in determining stream DOC and NO_3^- availability. Riparian zones are acknowledged as the major source of DOC in a wide range of forest headwater streams because their soils generally show high organic matter contents, especially in shallower layers (Dosskey & Bertsch, 1994; Bernal et al., 2005; Köhler et al., 2009; Strohmeier et al., 2013). Relatively wet conditions in the riparian zone favour the build-up of this organic matter and can also lead to denitrification, limiting the mobilization of NO_3^- (McClain et al., 1994; Cirimo and McDonnell, 1997). However, riparian zones that experience frequent dryness, such as those in Mediterranean catchments, can manifest nitrification and be net sources of NO_3^- to streams (Lupon et al., 2016a). As a result of these riparian zone characteristics, increased DOC concentrations in forest headwaters during storm events is an almost universal phenomenon (Hinton et al.,

1997; Boy et al., 2008; Yang et al., 2015; Musolff et al., 2018), whereas stream NO_3^- concentrations can increase, decrease, or remain unchanged during storms (Àvila et al., 1992; Bernal et al., 2002; Inamdar and Mitchell, 2006; Dupas et al., 2017).
65 At the same time, climate and associated soil moisture conditions preceding storm events can influence both the amount of DOC and NO_3^- stored in riparian soils and the riparian soil layers that are hydrologically connected to the stream, thus strongly influencing the supply of C and N to the aquatic compartment (Àvila et al., 1992; Fovet et al., 2015; Zimmer and McGlynn, 2018; Werner et al., 2019; Wen et al., 2020). Resulting stream DOC and NO_3^- concentrations and DOC: NO_3^- stoichiometry might therefore significantly differ between base flow and storm flow conditions, as well as among storms
70 depending on antecedent accumulated precipitation, and thereafter influence in-stream DOC and NO_3^- availability for microbial communities. Hence, studying the role of hydrology in controlling DOC and NO_3^- supply from riparian zones to adjacent streams becomes important for the subsequent understanding of in-stream heterotrophic activity.

In this study, we aimed to investigate how the interplay between hydrological processes, antecedent climate conditions, and riparian forest coverage drives DOC and NO_3^- availability and DOC: NO_3^- stoichiometry in a forest headwater Mediterranean
75 stream. In this stream, NO_3^- is the major source of N for stream heterotrophic microorganisms, accounting for more than 90% of the dissolved inorganic N (Bernal et al., 2015) and for more than 80% of the total dissolved N (unpublished data). Yet, ambient DOC and NO_3^- concentrations are relatively low compared to forest headwater streams located elsewhere (Hartmann et al., 2014; Bernal et al., 2015; Bernal et al., 2018), and we expected hydrological conditions and the presence of riparian forests to have a significant influence on the supply of these limited solutes for stream heterotrophic
80 microorganisms. In order to fulfil our aim, we analysed DOC and NO_3^- concentrations, and DOC: NO_3^- molar ratios under base flow and storm flow conditions at three stream locations along a longitudinal gradient of increasing riparian forest coverage. We define riparian zones here as near-stream areas within the catchment where riparian forest is present. In addition, we performed an event analysis to examine the hydroclimatic conditions that most favoured the transfer of DOC and NO_3^- from the riparian zone to the stream during storms, which were assumed to have a disproportionate influence on
85 solute supply. Based on this hydroclimatic event analysis, we propose a conceptual model synthesising the complex hydroclimatic and biogeochemical processes driving DOC and NO_3^- mobilization from riparian zones to streams during large storms.

2 Materials and methods

2.1 Study site characterization

90 The Font del Regàs catchment, located in the Montseny Natural Park in NE Spain (outlet at 41°50' N, 2°28' E), includes a total drainage area of 15.5 km² that ranges between 405 and 1603 m above sea level (m a.s.l) (Fig. 1). The climate is subhumid Mediterranean with mild winters, wet springs, and dry summers. Long-term (1940 to 2000) annual precipitation averaged 925 ± 151 mm year⁻¹ (average ± SD), whereas average annual air temperature during this period was 12.1 ± 2.5 °C (Lupon et al., 2016b).

95 The altitude gradient drives changes in vegetation cover in the catchment. While deciduous European beech (*Fagus sylvatica*) forests and a small proportion of heathlands (*Calluna vulgaris* and gramineae) dominate in the upper, steeper parts of the catchment, evergreen oak (*Quercus ilex*) forests dominate in the lower parts. Riparian forests, composed of a mixture of tree species including black alder (*Alnus glutinosa*), black locust (*Robinia pseudoacacia*), European ash (*Fraxinus excelsior*), and black poplar (*Populus nigra*), develop in flat, near-stream zones of the lower parts of the catchment (Lupon et al., 2016b). Both the width of the riparian forests and the total basal area of riparian trees increase downstream (Bernal et al., 2015). For its most part, Font del Regàs can be considered a closed-canopy stream (*sensu* Tank et al., 2018). For this study, we selected three stream locations along a longitudinal gradient of increased riparian forest coverage, which delineate three nested subcatchments (Fig. 1). The upstream subcatchment (746 to 1603 m a.s.l.) drains a steep area of 1.8 km² along the first 2.9 km of the Font del Regàs stream. The vegetation here is dominated by beech forests and heathlands 105 (92%), with lower proportions of oak forests (8%), and no riparian forests (i.e. no riparian zones). The midstream subcatchment (local drainage area of 6.8 km², ranging from 566 to 746 m a.s.l.) drains a cumulated area of 8.5 km² and includes 1.8 km more of the stream. Cumulated vegetation proportions here are 53% beech forests, 43% oak forests, and 4% mixed, 5 to 15 m wide riparian forests. The downstream subcatchment (local drainage area of 4.4 km², ranging from 503 to 566 m a.s.l.) drains a cumulated area of 13 km² and includes 1.4 km more of the stream, for a total of 6 km. Cumulated 110 vegetation proportions at the downstream subcatchment are 46% beech forests, 48% oak forests, and 6% riparian forests, which are well-developed and can be up to 30 m wide.

2.2 Field data collection, sampling, and laboratory analyses

The present study is based on data collected during the two-year period 9th September 2010 to 31st August 2012. All data and analyses were integrated and carried out for daily resolutions, which was the resolution of the stream chemical data.

115 Cumulated precipitation was recorded at 15 min intervals using an automatic weather station located at the valley bottom of the catchment, near the downstream site (Fig. A1). At each of the three stream locations, referred to as sites hereafter, we measured stream water levels at 15 min intervals using pressure transducers (HOBO U20-001-04) connected to autosamplers (Teledyne Isco Model 1612). Stream flow was measured every two weeks at each site by applying the salt dilution method (Gordon et al., 2004). Rating curves obtained from the relationships between stream flow and stream water level 120 measurements were used to construct daily time series of stream flow data at each site ($R^2 > 0.97$ in all cases; Lupon et al., 2016b). Stream flow was normalized by drainage area; thus, it is denoted by units of mm throughout ($N = 723$ days in all cases; Fig. A1).

Groundwater tables were recorded at 15 min intervals using a water pressure transducer (Druck PDCR 1830) within a piezometer placed 2.5 m away from the stream channel and located in a riparian zone area ca. 500 m downstream the 125 downstream site (Fig. 1), where stream flow normalized by drainage area is assumed to be equivalent. In a previous study, we showed that the dynamics of the pressure transducer dataset captured well the dynamics of manually-measured groundwater table variations in seven other piezometers located in the surrounding riparian area (Ledesma et al., 2021).

Therefore, we are confident that the recorded pattern at the pressure transducer location was representative of the groundwater table variations in the riparian zone soils in the lower parts of the catchment. The available daily groundwater table records were 664 (sporadic device failure precluded a full time series for the full study period).

Stream water samples were collected every day at noon at each of the three sites using autosamplers (Teledyne Isco Model 1612), which were installed ca. 1 m belowground to keep samples cool and to prevent biogeochemical transformations (Lupon et al., 2016b). Water samples were transported to the laboratory every 7 to 14 days, where they were filtered through pre-washed Whatman GF/F glass fibre filters (pore size = 0.7 μm) and analysed for DOC by catalytic oxidation using a Shimadzu total organic carbon (TOC) analyser, and for NO_3^- by cadmium reduction using a Technicon autoanalyzer. Sporadic autosampler failure precluded a full time series of data for the full study period. Total number of available daily DOC concentration values for upstream, midstream, and downstream sites were 537, 668, and 620, respectively. Analogously, total number of available daily NO_3^- concentration values were 552, 676, and 613, respectively. Stream DOC and NO_3^- concentrations were used to calculate the DOC: NO_3^- molar ratio for each available pair. Total number of DOC: NO_3^- molar ratio values were 537, 668, and 611 for upstream, midstream, and downstream sites, respectively (Fig. A1).

2.3 Data treatment and statistical analyses

At each site, we classified daily stream flows into base flow and storm flow. This separation can be done using a variety of approaches and time resolutions, which in most cases require certain degree of subjectivity (Hewlett and Hibbert, 1967; Buffam et al., 2001; Fovet et al., 2018; Caillon and Schelker, 2020). Here, we defined the beginning of storm flow event as the day when cumulated precipitation amounted more than 5 mm and stream flow was more than 10% higher than the previous day. We chose this precipitation threshold because hydrological responses to precipitation inputs below 5 mm day^{-1} are generally undetectable in Mediterranean catchments (Gallart et al., 2002). The end of a storm flow event was defined as the day when stream flow was less than 5% lower than the previous day. All stream flow values within event days were classified as storm flow, whereas the remaining values were classified as base flow. From the available 723 days, the upstream, midstream, and downstream sites included 128, 129, and 151 storm flow days. The small discrepancies in the storm flow correspondence among sites were likely caused by (i) differences in response times, being the downstream site slower in returning to base flow conditions; (ii) sporadic activations of intermediate intermittent tributaries; and (iii) storm origin, being eastern storms that first enter from the downstream part of the catchment more common than western storms that enter from the upper and middle parts of the catchment.

In order to investigate the influence of hydrological conditions on stream chemistry, for each site, we compared stream DOC concentrations, stream NO_3^- concentration, and stream DOC: NO_3^- molar ratios between base flow and storm flow conditions using non-parametric Wilcoxon rank-sum tests for non-normally distributed data (Zar, 2010). Moreover, in order to investigate the impact of riparian forest coverage on stream chemistry, we further performed pairwise Wilcoxon rank-sum

160 tests comparing DOC and NO_3^- concentrations, and DOC: NO_3^- molar ratios between stream sites (i.e. upstream *versus* midstream, upstream *versus* downstream, and midstream *versus* downstream), separately for base flow and storm flow conditions.

Based on previous studies assessing optimal DOC: NO_3^- molar ratios for microbial activity in lotic ecosystems, we classified stream DOC: NO_3^- molar ratios into ‘optimal’, ‘C-limited’, and ‘N-limited’ for stream heterotrophic microorganisms. We
165 used the reference DOC: NO_3^- molar ratios for streams and rivers presented in the global reviews of Taylor and Townsend (2010) and Helton et al. (2015) as the lower (i.e. 4.8) and upper (i.e. 11.7) limits delimiting the range of ‘optimal’ stoichiometric conditions. Consequently, stoichiometric conditions were considered ‘C-limited’ when stream DOC: NO_3^- molar ratios were below 4.8 and ‘N-limited’ when stream DOC: NO_3^- molar ratios were above 11.7. We explored the effect of hydrology and riparian forest coverage on DOC: NO_3^- stoichiometry by calculating the frequency of optimal, C-limited,
170 and N-limited conditions at each site and flow condition. We tested whether the frequencies of optimal, C-limited, and N-limited conditions for a given site were statistically different between base flow and storm flow by using a contingency analysis (Zar, 2010). Contingency analysis was also employed to test whether there were statistical differences in the frequency of optimal, C-limited, and N-limited conditions among sites. This analysis was done separately for base flow and storm flow conditions.

175 Finally, we computed simple linear regressions between stream DOC concentrations, stream NO_3^- concentrations, and stream DOC: NO_3^- molar ratios in order to investigate whether DOC or NO_3^- concentrations controlled changes in stoichiometry. Linear regressions were computed separately for each site and flow condition. Statistical analyses were carried out in the software JMP® Pro version 14.0.0 and the significance level for all analyses was set at $p < 0.01$.

2.4 Hydroclimatic analysis of storm events

180 We further aimed to explore the hydroclimatic characteristics that most effectively promoted the mobilization of DOC and NO_3^- into the stream via groundwater table elevation and consequent hydrological activation of upper riparian layers during storm events, which are generally associated with the largest exports of solutes from headwater catchments. For that, we identified the storm events that occurred during the study period and characterized them using nine hydroclimatic descriptors. We then related those descriptors with the observed stream DOC and NO_3^- concentrations through a partial least
185 square (PLS) regression model for the subset of storms for which clear hydrological activation of upper riparian soil layers occurred. This analysis was performed only for data from the downstream site, where the riparian zone was well-developed. Moreover, this was the only site with associated groundwater table data.

We identified storm events based on three requirements: (i) a precipitation amount during the days included in the event of at least 25 mm (events below these threshold showed marginal groundwater table responses at the temporal resolution of the
190 study), (ii) stream flow was classified as storm flow during the days of the event (not all events with precipitation amounts higher than 25 mm generated storm flow at the temporal resolution of the study), and (iii) a complete stream chemistry data

series associated with the event for the downstream site (i.e. no gaps in chemical data for the event dates were allowed as we were ultimately interested in stream chemical responses). Each event included all days classified as storm flow for the corresponding time period, plus the last day classified as base flow prior the days classified as storm flow, which was considered as the starting date of the event (Blaurock et al., 2021). This strategy ensured that the relevant magnitude and range of the hydroclimatic descriptors associated with each event were covered. Moreover, when there was a substantial decline in stream flow and riparian groundwater tables followed by a subsequent increase in both variables without reaching base flow conditions, we split this bimodal event into two events. In this case, the day immediately before the second increase in stream flow and riparian groundwater tables was considered as both the last day of event i and the first day of event $i+1$ (this was the case for events #1 and #2, presented later).

Each of the identified storm events was characterized by the following hydroclimatic descriptors: *duration* (days); precipitation amount (P , mm); accumulated precipitation seven days before the event ($P-7$, mm), as a proxy for short-range antecedent soil moisture conditions; accumulated precipitation 30 days before the event ($P-30$, mm), as a proxy for long-range antecedent soil moisture conditions; average stream flow (Q_{avg} , mm); average groundwater table (Gw_{avg} , m); and groundwater table range (ΔGw , m), calculated as the absolute difference between the highest and the lowest groundwater tables measured during the event. Moreover, the hydroclimatic characterization of each event included the slope of the linear relationship between daily riparian groundwater tables and daily stream flows (*slope*, $m\ mm^{-1}$), as a proxy for the rate at which upper riparian layers are hydrologically activated in relation to increments in stream flow (Rodhe, 1989). We anticipated that this slope would vary depending on antecedent soil moisture conditions, with higher values related to drier conditions, and that this variation would have implications for solute mobilization and stream chemistry. Finally, we also calculated the average stream flow normalized to average groundwater table (Q_{avg}/Gw_{avg} , $mm\ m^{-1}$), which served as proxy for average hydraulic conductivity in the activated layers of the riparian profile, with higher values indicating larger lateral transmissivity from the profile.

We observed large differences between two subsets of events in their groundwater table elevation (defined by their groundwater table range ΔGw), generally associated with their precipitation amount P ($R^2 = 0.67$, linear regression between P and ΔGw). We classified these two subsets of events as ‘medium storms’ ($P = 29$ to 98 mm, $\Delta Gw = 2$ to 9 cm, $N = 11$) and ‘large storms’ ($P = 67$ to 174 mm, $\Delta Gw = 21$ to 55 cm, $N = 5$), and proceeded with the PLS analyses only with the subset of large storms because, unlike medium storms, they showed clear hydrological activation of upper riparian soil layers defined by ΔGw . Pairwise Wilcoxon rank-sum tests were used to compare the hydroclimatic descriptors and the stream DOC and NO_3^- concentrations between medium and large storm events.

The nine hydroclimatic descriptors were considered as predictors in the PLS regression model, and average DOC and NO_3^- concentrations at the downstream site during each of the five large storm events were the response variables. A PLS regression model was appropriate in our case because (i) the number of predictors was large ($N = 9$) and higher than the number of observations ($N = 5$) and (ii) there was multicollinearity among predictors (not explicitly shown) (Wold et al.,

225 2001). For each response variable, a PLS regression model returns a goodness of fit (R^2Y), which is the variation of the response variable explained by the predictors, and a predictive ability (Q^2Y), which is an indication on how well the model can predict new data ($Q^2Y > 0.5$ are considered good models). In addition, the model returns a VIP (variable influence on projection) value associated with each predictor, which informs about the relative importance of the predictor in the overall model. Predictors with $VIP > 1$ are considered statistically important for the model performance (Eriksson et al., 1999). The

230 PLS regression analysis was carried out in the software SIMCA (©Umetrics AB) version 14.0.

Finally, for each of the characterized large storm events, we examined the type of hysteresis loop (clockwise, anticlockwise, or linear) of the relationship between daily riparian groundwater tables and daily stream flows. This relationship typically shows clockwise hysteresis loops, which imply that, for a given stream flow, the groundwater table is higher during the rising limb than during the falling limb, indicating that near-stream zones are the dominant water sources during the rising limb (Kendall et al., 1999; Frei et al., 2010). We compared the type of hysteresis loop among the five large storm events in order to determine potential differences in runoff generation processes and associated solute mobilization and DOC:NO_3^- stoichiometry.

3 Results

3.1 Base flow *versus* storm flow stream chemistry and changes along the stream

240 Including all sites and dates, stream DOC concentrations averaged $1.0 \pm 0.6 \text{ mg C L}^{-1}$ and ranged from 0.2 to 5.4 mg C L^{-1} . Stream DOC concentrations were significantly higher during storm flow than during base flow conditions in the three sites (Wilcoxon test, $p < 0.01$). On average, stream DOC concentrations increased similarly from base flow to storm flow in all sites: by 66% at the upstream site (from 0.8 ± 0.4 to $1.4 \pm 0.8 \text{ mg C L}^{-1}$), by 60% at the midstream site (from 1.0 ± 0.5 to $1.6 \pm 0.8 \text{ mg C L}^{-1}$), and by 55% at the downstream site (from 0.9 ± 0.4 to $1.4 \pm 0.8 \text{ mg C L}^{-1}$) (Fig. 2a-b). The variability in stream DOC concentrations was relatively similar during both flow conditions, as indicated by similar coefficient of variations (CV): 54% and 57% in the upstream site, 46% and 52% in the midstream site, and 40% and 60% in the downstream site, respectively for base flow and storm flow conditions. All pairwise comparisons between sites showed statistically significant differences in stream DOC concentrations during base flow conditions (Fig. 2a). On average, stream DOC concentrations increased by 19% from the upstream to the midstream site and then decreased by 10% from the midstream to the downstream site (Table 1). This pattern was similar during storm flow conditions, i.e. a 15% increase from the upstream to the midstream site and a 13% decrease from the midstream to the downstream site were observed, but differences in DOC concentrations were statistically significant only between the midstream and the downstream sites in this case (Fig. 2b).

250 Stream NO_3^- concentrations ranged from 0.09 to 1.1 mg N L^{-1} considering all sites and dates, and averaged $0.20 \pm 0.09 \text{ mg N L}^{-1}$. Stream NO_3^- concentrations were significantly higher during storm flow than during base flow conditions in the three

sites (Wilcoxon test, $p < 0.01$). The relative increase in stream NO_3^- concentrations from base flow to storm flow was remarkably similar among sites, i.e. 41% at the upstream site (from 0.26 ± 0.05 to 0.36 ± 0.17 mg N L^{-1}), 40% at the midstream site (from 0.15 ± 0.03 to 0.22 ± 0.11 mg N L^{-1}), and 42% at the downstream site (from 0.17 ± 0.03 to 0.24 ± 0.10 mg N L^{-1}) (Fig. 2c-d). Moreover, stream NO_3^- concentrations were consistently more variable during storm flow than during base flow, as indicated by increases in the CV: from 20% to 47%, from 21% to 52%, and from 18% to 42% in upstream, midstream, and downstream sites, respectively. All pairwise comparisons between sites showed statistically significant differences in stream NO_3^- concentrations (Fig. 2c-d). On average, stream NO_3^- concentrations decreased 40% from the upstream to the midstream site and increased nearly 10% from the midstream to the downstream site (Table 1). This pattern held during both base flow and storm flow conditions.

265 Considering the three sites pooled together, stream DOC: NO_3^- molar ratios varied from 1.0 to 28, with an average of 6.1 ± 3.3 . On average, stream DOC: NO_3^- molar ratios were 20% higher during storm flow compared to base flow conditions for the upstream (increasing from 3.8 ± 2.1 to 4.6 ± 2.0) and midstream (increasing from 7.5 ± 3.4 to 8.9 ± 3.6) sites (Wilcoxon test, $p < 0.01$) (Fig. 2e-f). There was no statistical difference in stream DOC: NO_3^- molar ratios between base flow and storm flow conditions at the downstream site (Wilcoxon test, $p = 0.08$), where the overall average was 6.3 ± 2.7 . All pairwise comparisons between sites showed statistically significant differences in stream DOC: NO_3^- molar ratios (Fig. 2e-f). These differences were consistent between base flow and storm flow conditions: stream DOC: NO_3^- molar ratios almost doubled between upstream and midstream sites and then decreased ca. 20% from the midstream to the downstream site (Table 1).

275 Stream DOC: NO_3^- molar ratios considered optimal for heterotrophic microorganisms (i.e. from 4.8 to 11.7) were not prevalent at the upstream site during either base flow or storm flow conditions (Fig. 3). Nevertheless, at this site, the frequency of optimal DOC: NO_3^- molar ratios was statistically higher during storm flow (33%) than during base flow (18%), as indicated by the contingency analysis ($p < 0.01$). The rest of the time, stream DOC: NO_3^- molar ratios were lower than 4.8, indicating that C-limited conditions for heterotrophic activity dominated in the upstream site. By contrast, optimal DOC: NO_3^- molar ratios were largely prevalent (ranging from 69% to 74% of the time) at the midstream and downstream sites during both base flow and storm flow conditions (Fig. 3). According to the contingency analysis, there was a statistical difference in the frequency of DOC: NO_3^- stoichiometric conditions between base flow and storm flow for the midstream site ($p < 0.01$). This difference was driven by an increase in N-limited conditions from 11% during base flow to 24% during storm flow, which was accompanied by a decrease in C-limited conditions from 15% during base flow to 3% during storm flow. At the downstream site, the frequency of DOC: NO_3^- stoichiometric conditions was similar between base flow and storm flow conditions (contingency analysis, $p = 0.31$), and C-limited conditions were more frequent than N-limited conditions (25% versus 6% for the combined base flow and storm flow periods). As a result of these patterns, there were differences in the frequency of DOC: NO_3^- stoichiometric conditions among sites during both base flow and storm flow (contingency analyses, $p < 0.01$).

285

During base flow, no relationship between stream DOC and NO_3^- concentrations was found, whereas they were positively related during storm flow conditions at the three sites ($R^2 > 0.39$, $p < 0.01$) (Fig. 4a-b). Strong positive linear relationships were observed between stream DOC concentrations and stream DOC: NO_3^- molar ratios during base flow conditions at the three sites ($R^2 > 0.80$, $p < 0.01$) (Fig. 4c). During storm flow conditions, these relationships weakened because data became more scattered as DOC concentrations increased, but were still statistically significant in all sites ($R^2 > 0.22$, $p < 0.01$) (Fig. 4d). By contrast, the three sites showed no relationship between stream NO_3^- concentrations and stream DOC: NO_3^- molar ratios (Fig. 4e-f).

295 3.2 Hydroclimatic analysis of storm events

A total of 16 storm events satisfied our selection requirements for further analysis (Table A1). Hydrological responses were considerably different between medium and large storm events. Compared to large storms, medium storms were characterized by (i) lower accumulated precipitation amounts, (ii) lower average stream flow, (iii) deeper groundwater tables, and (iv) smaller groundwater table ranges, i.e. smaller thickness of the riparian layer that becomes hydrologically activated during the event (Wilcoxon test, $p < 0.01$ in all cases; Fig. 5a-d). Additionally, both stream DOC and stream NO_3^- concentrations were significantly lower during the medium storm events than during the large storm events (Wilcoxon test, $p < 0.01$ in both cases; Fig. 5e-f).

In their turn, the nine hydroclimatic descriptors varied widely among the five large storm events (Table 2). Total *duration* varied between 4 and 11 days and precipitation amount (P) between 67 and 174 mm. Accumulated precipitation seven ($P-7$) and, especially, 30 ($P-30$) days before the event (as proxies for short- and long-range antecedent soil moisture conditions, respectively) showed wide ranges, i.e. $P-7$ varied between 11 and 182 mm and $P-30$ varied between 39 and 426 mm. Average stream flow (Q_{avg}) ranged from 0.8 to 4.2 mm, whereas average groundwater table (Gw_{avg}) varied more moderately, between 0.64 and 0.87 m below the soil surface. Groundwater table range (ΔGw) varied between 0.21 and 0.55 m. The slope of the linear relationship between daily riparian groundwater tables and daily stream flows (*slope*), as a proxy for the rate at which upper riparian layers are hydrologically activated in relation to increments in stream flow, ranged from 0.06 to 0.18 m mm^{-1} and was negatively related with both $P-30$ ($R^2 = 0.74$) and the average stream flow normalized to average groundwater table (Q_{avg}/Gw_{avg}) ($R^2 = 0.68$). Q_{avg}/Gw_{avg} was a proxy for average hydraulic conductivity in the activated layers of the riparian profile and ranged from 0.9 to 6.1 mm m^{-1} . During all five large storm events, stream DOC and NO_3^- concentrations were 55% to 178% higher than during base flow conditions (Table 2). Additionally, stream DOC concentrations varied more (CV = 21%) than stream NO_3^- concentrations (CV = 13%) among events.

A two-component PLS regression model applied to the five large storm events dataset explained 96% (i.e. $R^2Y = 0.96$) and 94% (i.e. $R^2Y = 0.94$) of the variation in average stream DOC and NO_3^- concentrations, respectively. The ability of the model to predict new data was high, with Q^2Y values of 0.88 and 0.82 for stream DOC and NO_3^- concentrations, respectively. Four out of the nine predictors used in the PLS regression model reached $\text{VIP} > 1$ (Fig. A2), indicating that they were important for

320 explaining the variability in the response variables. Specifically, lower $P-30$, lower Q_{avg} , higher $slope$, and lower Q_{avg}/Gw_{avg} supported higher concentrations of both DOC and NO_3^- (Fig. 6). These four predictors showed the highest loadings in the first component of the model, which explained most of the variation in stream DOC ($R^2Y = 0.68$) and NO_3^- ($R^2Y = 0.86$) concentrations. From the four important predictors, $P-30$ showed the largest loading in the second component and provided a distinct and extra statistical explanation for the DOC model compared to the NO_3^- model. Hence, this component explained
325 28% of the variation in average stream DOC concentrations, but only 8% of the variation in average stream NO_3^- concentrations.

Four of the five large storm events showed clockwise hysteresis loops in their riparian groundwater table – stream flow relationship, whereas event #5, with the largest $P-30$, displayed a slight anticlockwise loop (Fig. 7a). Stream DOC: NO_3^- molar ratios at the downstream site during the five events generally fell within the suggested optimal stoichiometric range
330 (Fig. 7b). Nevertheless, event #5 deviated again from the general pattern and showed lower stream DOC: NO_3^- molar ratios than the other events, partially falling within the range defined as C-limited conditions for heterotrophic activity.

4 Discussion

4.1 The role of the riparian zone on determining DOC, NO_3^- , and DOC: NO_3^- along the stream

Stream DOC concentrations at the Font del Regàs catchment (1.0 ± 0.6 mg C L⁻¹) were remarkably low compared with other
335 forest headwaters in boreal (de Wit et al., 2016), temperate (Musolff et al., 2018), and tropical (Boy et al., 2008) regions, but were comparable to other Mediterranean sites (Casas-Ruiz et al., 2017; Catalán et al., 2018). On the other hand, stream NO_3^- concentrations (0.20 ± 0.09 mg N L⁻¹) were somewhat higher than those typically measured in N-limited boreal streams (Blackburn et al., 2017), comparable to the low concentrations measured in tropical and Mediterranean forest headwaters (McDowell et al., 1992; Catalán et al., 2018), and notably lower than those measured in temperate sites (Musolff et al.,
340 2017). Therefore, Font del Regàs can be considered an oligotrophic stream with relatively low N and, especially, C availability. These conditions could constrain stream productivity in general and the activity of stream heterotrophs in particular, given that this is a predominantly heterotrophic system where ecosystem respiration rates are between 10- and 100-fold higher than gross primary production rates (Lupon et al., 2016c).

Clear and consistent patterns were observed for DOC and NO_3^- concentrations and DOC: NO_3^- molar ratios along the Font
345 del Regàs stream (Table 1, Fig. 2). DOC concentrations were especially low at the upstream site and increased moderately (ca. 20%) until the midstream site. This increase could be attributed to changes in catchment topography from upstream to midstream and the associated development of riparian forests in near-stream areas (Jencso et al., 2009; Musolff et al., 2018). The upstream site drains a steep area where soils are well-drained, and no riparian forest can develop. As geomorphology becomes more favourable for the development of wetter soils in the near-stream zone, riparian forests can establish and
350 support relatively larger mobilization of DOC from the riparian zone to the stream (Inamdar and Mitchell, 2006; Bernal et

al., 2018). However, DOC concentrations decreased ca. 10% from the midstream to the downstream site despite that riparian forests are well-developed along this section. This result suggests that DOC was not transported conservatively along the stream and that in-stream DOC demand was likely higher than supply between the midstream and downstream sites, as suggested also by Lupon et al. (2020b). This indication can be further supported by a more consistent optimal DOC:NO₃⁻ stoichiometry in this section compared to that observed between the upstream and midstream sites, where stoichiometric conditions evolved from C-limited to optimal.

Concentrations of NO₃⁻ were highest at the upstream site, and then notably decreased (ca. 40%) in the stretch to the midstream site. This result is in line with the idea that headwater streams can remove substantial amounts of NO₃⁻ within relatively short distances (Peterson et al., 2001). Nevertheless, previous studies in this stream suggest that in-stream processing alone might not have been enough to explain the observed decrease in NO₃⁻ concentration along the upstream-midstream section (Bernal et al., 2015). Denitrification in riparian zones developed along this section could provide groundwater inputs with low NO₃⁻ concentrations and contribute to the reduction in stream NO₃⁻, a process widely observed in temperate forest catchments (Cirimo and McDonnell, 1997). However, riparian soils at Font del Regàs can act as sources of NO₃⁻ to the stream because they are well-oxygenated and can sustain large nitrification rates (Lupon et al., 2016a). Thus, it is unclear why the upstream site showed such consistently higher NO₃⁻ concentrations with respect to the midstream site. A combination of steeper slopes (which also favour oxygenation and potential nitrification in soil water) and a vegetation dominated by beech forest (which might uptake lower amounts of NO₃⁻ than the oak and riparian forests at the midstream section) might partially explain the higher NO₃⁻ concentrations upstream and the upstream-midstream pattern (Schiff et al., 2002; Poblador et al., 2019; Simon et al., 2021). The small increase (ca. 10%) in NO₃⁻ concentrations between the midstream and downstream sites could in this case be explained by the increasing influence of NO₃⁻-rich riparian groundwater inputs in the downstream areas, though in-stream mineralization of leaf litter inputs could also contribute (Bernal et al., 2015; Lupon et al., 2015).

As a result of the observed DOC and NO₃⁻ patterns among the stream sites, DOC:NO₃⁻ molar ratios doubled from the upstream to the midstream site. This change could have important implications for stream heterotrophic activity because stoichiometric conditions shifted from predominately C-limited at the upstream site to predominately optimal at the midstream site (Fig. 3). C-limited conditions at the upstream site were clearly driven by low DOC availability, which is in accordance with previous studies showing strong C-limitation in Mediterranean and semi-arid streams (Catalán et al., 2018). Supporting this idea, we found that stream DOC concentrations were responsible for driving stream DOC:NO₃⁻ molar ratios at the three sites under almost all circumstances (Fig. 4). As the influence of the riparian forest in the catchment increased from upstream to midstream and downstream locations, potential C-limitation based on DOC:NO₃⁻ molar ratios was generally overcome, and stoichiometric optimal conditions for stream heterotrophic activity became prevalent. This result suggests that riparian zones are essential ecosystem compartments for ensuring the supply of DOC to the stream and, thus, for fuelling in-stream heterotrophic activity (Lupon et al., 2020b).

It is worth mentioning that defining an optimal range of DOC:NO₃⁻ molar ratios for in-stream heterotrophic activity is
385 difficult due to the lack of consistent reference values, and the fact that heterotrophic microorganisms can adapt to a wide
range of resource stoichiometric ratios (Manzoni et al., 2017). We acknowledged this uncertainty by considering a wide
range of DOC:NO₃⁻ values (from 4.8 to 11.7) defining potential optimal conditions for in-stream heterotrophic activity based
on published data compilations (Taylor and Townsend, 2010; Helton et al., 2015). However, this range could be even wider
if modelling and site-specific studies are considered (Danger et al., 2008; Pastor et al., 2014; Bastias et al., 2020).
390 Noteworthy, we obtained very similar frequencies of optimal, C-limited, and N-limited conditions when using a less
restrictive DOC:NO₃⁻ molar ratio range for optimal conditions (from 4.5 to 16) (Fig. A3). Therefore, we are confident that
the approach used in the present study is helpful for illustrating different stoichiometric conditions and how they relate to C
and N availability for stream heterotrophic microorganisms at the study site and other similar sites elsewhere.

395 **4.2 Changes in DOC, NO₃⁻, and DOC:NO₃⁻ associated with hydrological conditions: potential implications for in-stream heterotrophic activity**

Stream DOC and NO₃⁻ concentrations substantially increased during storm flow conditions at the three sites, a pattern
consistent with previous observations in other Mediterranean catchments and that can be explained by the hydrological
activation of organic-rich layers in near-stream zones (Àvila et al., 1992; Bernal et al., 2002). At the upstream site, the
magnitude of change between flow conditions was different between DOC (which on average increased by 66%) and NO₃⁻
400 (which on average increased by 41%), leading to an increase in the frequency of optimal DOC:NO₃⁻ stoichiometric
conditions for heterotrophic activity during storm flow. By contrast, at the midstream and downstream sites, the frequency of
optimal DOC:NO₃⁻ stoichiometry was high regardless of flow conditions; although, at the midstream site, N-limited and C-
limited conditions became more and less frequent during storm flow, respectively. Overall, these results suggest that (i)
hydrological inputs during storms increase stream DOC and NO₃⁻ availability by supplying allochthonous dissolved organic
405 matter (DOM) and nutrients from near-stream zones, potentially alleviating resource limitation, and (ii) in-stream
heterotrophic activity might be further favoured during high flows in sites with predominant C-limitation due to an increase
in the frequency of optimal stoichiometry.

Nevertheless, higher water velocities and lower water residence times during storm flows can limit the capacity of
heterotrophic microorganisms to take up essential elements from the water column, especially in headwater streams with
410 steeper slopes. Thus, the biogeochemical opportunity for in-stream processing (*sensu* Marcé et al., 2018) might be restricted
to locations downstream, as proposed by the pulse-shunt concept (Raymond et al., 2016). However, there are multiple lines
of evidence resulting from studies carried out in our site and in other forest catchments that suggest that high stream flows
might indeed offer *windows of opportunity* for in-stream heterotrophic activity in headwater locations. For example, recent
studies have reported that heterotrophic microorganisms can process some of the C and N entering the stream during high
415 flow conditions in forest headwaters across ecoregions (Seybold and McGlynn, 2018; Wollheim et al., 2018). Bernal et al.

(2019) showed that in-stream DOC and NO_3^- uptake at Font del Regàs could be as high or even higher during storm flow than during base flow. Further, Bernal et al. (2018) showed that DOM at Font del Regàs has a prominent protein-like character in both riparian groundwater and stream water during base flow conditions, which could favour rapid assimilation even during periods of short water residence times assuming DOM molecular composition maintains part or most of its labile character across flow conditions. Sporadic inputs of DOM rich in aliphatic molecules have been observed in other forest headwaters during storm flow conditions, which could further increase bioavailability and assimilation (Wilson et al., 2013; Wagner et al., 2019). Part of the in-stream biogeochemical uptake during high flows could also be triggered by extra inflows of heterotrophic bacteria mobilized from catchment soils, as suggested by Caillon and Schelker (2020). Mechanistically, high flows can also enhance the hydrological interaction with streambed sediments, where microbial assemblages develop (Li et al. 2021). Lastly, and more importantly for the present study, changes in DOC, NO_3^- , and DOC: NO_3^- along the Font del Regàs stream were remarkably similar during both base flow and storm flow conditions. This observation suggests that the pattern of resource consumption was maintained independently of stream flow conditions. All these direct and indirect pieces of evidence support that in-stream heterotrophic activity can be partially sustained during storm flows, especially in sites with limited C and N availability such as Font del Regàs.

4.3 Antecedent soil moisture conditions shape the mobilization of DOC and NO_3^- during large storm events

Understanding which hydrological and climatic conditions result in more effective resource mobilization from riparian soils to the stream is relevant in the context of the present study and for other oligotrophic streams. Here, we focused this question on storm events, which are generally associated with the largest exports of solutes from headwater catchments and have the capacity to hydrologically connect shallow, organic-rich riparian layers to the fluvial network (Godsey et al., 2009; Raymond et al., 2016; Zimmer and McGlynn, 2018).

Based on their hydrological response to precipitation, we were able to clearly identify two subsets of events. Large storms, characterized by higher precipitation amounts, produced considerable stream flow and groundwater table responses, whereas medium storms produced only moderate responses in stream flow and limited groundwater table elevations (Fig. 5). These differences in climatic and, especially, hydrological characteristics between large and medium storm events led to marked differences in the resulting stream chemistry, with both DOC and NO_3^- concentrations being significantly higher during the large storm events. This observation is likely caused by the hydrological activation of a thicker and shallower riparian layer during large storm events that leads to the mobilization of relatively larger amounts of DOC and NO_3^- stored in the riparian soil compared to the amounts mobilized from the deeper and narrower layers that are hydrologically activated during medium storm events. These results suggest that large and medium storm events display distinct mechanisms of DOC and NO_3^- mobilization from the riparian zone and that large storms are responsible for a disproportionately larger supply of these solutes.

The PLS regression analysis focussed on understanding which hydrological and climatic conditions result in more effective resource mobilization from riparian soils during large storm events. While the sample size of large storms was low, the nature of these events covered a wide range of hydroclimatic characteristic and provided useful information on the climatic conditions and hydrological processes that control DOC and NO_3^- mobilization from riparian zones to streams in forest headwater catchments. We found that long-range antecedent soil moisture conditions and the relationship between riparian groundwater tables and stream flows are essential factors for understanding changes in stream DOC and NO_3^- concentrations during large storms. Particularly, antecedent drier conditions (lower $P-30$) led to relatively higher stream DOC and NO_3^- concentrations and constituted the most effective situations for the mobilization and supply of these solutes to the Font del Regàs stream. Therefore, large storms preceded by dry conditions might be essential for heterotrophic activity in this nutrient-limited stream. The large mobilization of solutes during these circumstances can be partially explained by abrupt increases in riparian groundwater tables, i.e. rapid hydrological activation of upper riparian layers in relation to increments in stream flow (higher *slope*). Hence, according to the PLS analysis, drier antecedent conditions led to steeper slopes of the relationship between groundwater table and stream flow, which were related to lower average hydraulic conductivity in the activated layers of the riparian profile (lower Q_{avg}/Gw_{avg}), and to lower stream flows (lower Q_{avg}) (Fig. 6). A faster increment in riparian groundwater table position compared to the increment in stream flow can be explained by a lower effective hydraulic conductivity of each riparian soil layer (estimated here as Q_{avg}/Gw_{avg}) induced by the pre-existing dry conditions and an increased hydrological disconnection between the riparian zone and the rest of the catchment. This preconditioning will make precipitation more efficient at filling up the riparian profile vertically relative to the otherwise dominant lateral transmissivity dimension (Frei et al., 2012; Fovet et al., 2015; Tunaley et al., 2016). Consequently, more riparian layers will be hydrologically connected to the stream, increasing the thickness of the ‘Dominant Source Layer’ (i.e. the riparian zone depth stratum that contributes the most to water and solute fluxes to the stream, Ledesma et al., 2018) and, thus, the chances for relatively more DOC and NO_3^- mobilization, which in turn generally show higher concentrations closer to the soil surface (Ranalli and Macalady, 2010; Camino-Serrano et al., 2016). In addition, drier conditions can restrict decomposition and promote primary production and nitrification in the riparian soil and result in the built up of soluble DOC and NO_3^- pools (Lupon et al., 2016a; Arce et al., 2019; Wen et al., 2020), which will further contribute to a higher mobilization of these solutes during subsequent large storm events, eventually leading to large increases in stream DOC and NO_3^- concentrations (Fig. 8a). Conversely, the PLS analysis indicated that wetter antecedent soil moisture conditions and less steep slopes of the relationship between groundwater table and stream flow resulted in relatively lower stream DOC and NO_3^- concentrations during large storms. Under these circumstances, soil moisture will be high and lateral transmissivity will be relatively larger due to increased hydrological connectivity between the riparian zone and the rest of the catchment. In addition, during large storms preceded by wet conditions, soluble DOC and NO_3^- pools in the riparian soils can be depleted from precedent storm events (Butturini et al., 2003; Zimmer and McGlynn, 2018; Werner et al., 2019), eventually leading to relatively lower increments in stream DOC and NO_3^- concentrations (Fig. 8b). Moreover, the relatively higher stream flows during events

480 preceded by wet conditions suggest larger spatial extension of water connectivity throughout the catchment (Tunaley et al., 2016), which may further contribute to dilute solutes, especially in oligotrophic streams such as Font del Regàs. Our conceptual model of DOC and NO_3^- mobilization during storm events based on data from a sub-humid Mediterranean catchment is similar to previous conceptualizations presented for temperate sites (e.g. Werner et al., 2019; Beiter et al., 2020), suggesting that the mechanisms proposed here can be representative of other forest headwaters located in similar
485 climatic settings.

Importantly, our results additionally highlight that the relationship between riparian groundwater tables and stream flows is not static but variable depending on storm event characteristics. This relationship typically shows clockwise hysteresis loops and synchronous behaviour patterns between riparian groundwater tables and stream flows (Jung et al., 2004; Frei et al., 2010; Fovet et al., 2015). However, here we showed that clockwise hysteresis loops during large storms were not
490 consistently similar between events and that under very wet antecedent conditions the clockwise pattern can turn into anticlockwise, as seen for event #5 (Fig. 7a). Moreover, event #5 was the only large storm event for which stream $\text{DOC}:\text{NO}_3^-$ molar ratios fell outside the optimal stoichiometric range for in-stream heterotrophic activity (Fig. 7b), which was explained by a relatively lower increase in stream DOC concentrations compared to that in stream NO_3^- concentrations (Table 2). This result suggests higher depletion and dilution of riparian DOC compared to riparian NO_3^- during large storms preceded by
495 very wet conditions (Fig. 8b), implying that C rather than N limitation might unfold in Mediterranean catchments such as Font del Regàs under those circumstances (Doblas-Miranda et al., 2013). Overall, these results underscore the relevance of climatic conditions preceding storm events in controlling the relative availability of DOC and NO_3^- in streams and, ultimately, the potential activity of heterotrophic microbial assemblages.

5 Conclusions

500 Spatiotemporal patterns of stream DOC and NO_3^- concentrations and $\text{DOC}:\text{NO}_3^-$ molar ratios, and thus where and when C and N supply from terrestrial ecosystems can sustain or constrain in-stream heterotrophic activity, depend on catchment hydrological and biogeochemical processes, especially those occurring at the riparian zone. Supporting this idea, we found clear and consistent changes in DOC and NO_3^- concentrations and $\text{DOC}:\text{NO}_3^-$ molar ratios during both base flow and storm
505 flow conditions along a longitudinal gradient of increased riparian forest coverage at the Mediterranean Font del Regàs stream. This outcome indicates that catchment geomorphic and topographic features, which give rise to specific vegetation and hydromorphological features, determine stream DOC and NO_3^- dynamics by modulating riparian forest formation (Inamdar and Mitchell, 2006; Ledesma et al., 2018).

Increases in both DOC and NO_3^- concentrations during storm flow compared to base flow conditions is a common observation in forest headwaters and the Font del Regàs stream was not an exception. Nevertheless, in this study we went a
510 step further by considering how changes in $\text{DOC}:\text{NO}_3^-$ stoichiometry along the stream and between flow conditions might

515 impact in-stream heterotrophic activity. Optimal DOC:NO₃⁻ stoichiometry for heterotrophic microorganisms was prevalent at midstream and downstream locations regardless of hydrological conditions, whereas they were more frequent during storm flow than during base flow conditions upstream. We conclude that hydrology plays a critical role in controlling C and N supply as well as DOC:NO₃⁻ stoichiometric requirements in close-canopy, oligotrophic Mediterranean streams because solute mobilization during high flows can alleviate resource limitation and burst in-stream heterotrophic activity during or right after storm events.

Further, our results suggest that increases in stream DOC and NO₃⁻ concentrations during large storm events are controlled by complex hydrological processes involving antecedent soil moisture conditions. Particularly, large storms preceded by drier conditions lead to more rapid elevations of riparian groundwater tables and to larger increases in stream DOC and NO₃⁻ concentrations. As the world climate continues changing, including an increased variability in precipitation patterns in the Mediterranean ecoregion (i.e. larger precipitation events and longer dry spells), it will be critical to understand hydrological and biogeochemical processes at the stream-riparian interface, and how the aftermath hydrology plays out for in-stream heterotrophic activity and consequent ecosystem functioning in Mediterranean forest headwaters.

Data availability

525 Precipitation, stream flow, groundwater table, and stream chemistry data used in the present study are available in the research data repository HydroShare at <https://www.hydroshare.org/resource/3366012c5e254937aa661ce2a93c3140/> (Ledesma, 2021) and <https://www.hydroshare.org/resource/0b4eb61ad7544ffb970c1600927a819f/>.

Author contribution

530 JLJL: Conceptualization, Formal analysis, Funding acquisition, Methodology, Visualization, Writing – original draft preparation. AL: Conceptualization, Investigation, Resources, Writing – review & editing. EM: Conceptualization, Supervision, Writing – review & editing. SB: Conceptualization, Investigation, Resources, Supervision, Writing – review & editing.

Competing interests

The authors declare that they have no conflict of interest.

535 Acknowledgements

JLJL was funded by the project RIPARIONS granted by the European Commission through a Marie Skłodowska Curie Individual Fellowship (H2020-MSCA-IF-2018-834363) and by the Spanish Government through a Juan de la Cierva grant

(FJCI-2017-32111). AL was supported by the Catalan Government and the European Commission through a MSCA-Beatriz de Pinós grant (BP-2018-00082). SB work was supported by the CANTERA project (RTI2018-094521-B-100) and a Ramón y Cajal fellowship (RYC-2017-22643) funded by MCIN/AEI/10.13039/501100011033/FEDER Una manera de hacer Europa.

References

- Arce, M. I., Mendoza-Lera, C., Almagro, M., Catalán, N., Romaní, A. M., Martí, E., Gómez, R., Bernal, S., Foulquier, A., Mutz, M., Marcé, R., Zoppini, A., Gionchetta, G., Weigelhofer, G., del Campo, R., Robinson, C. T., Gilmer, A., Rulik, M., Obrador, B., Shumilova, O., Zlatanović, S., Arnon, S., Baldrian, P., Singer, G., Datry, T., Skoulikidis, N., Tietjen, B., and von Schiller, D.: A conceptual framework for understanding the biogeochemistry of dry riverbeds through the lens of soil science, *Earth-Sci. Rev.*, 188, 441-453, [10.1016/j.earscirev.2018.12.001](https://doi.org/10.1016/j.earscirev.2018.12.001), 2019.
- Àvila, A., Piñol, J., Rodà, F., and Neal, C.: Storm solute behavior in a montane Mediterranean forested catchment, *J. Hydrol.*, 140, 143-161, [https://doi.org/10.1016/0022-1694\(92\)90238-q](https://doi.org/10.1016/0022-1694(92)90238-q), 1992.
- 545 Bastias, E., Ribot, M., Bernal, S., Sabater, F., and Martí, E.: Microbial uptake of nitrogen and carbon from the water column by litter-associated microbes differs among litter species, *Limnol. Oceanogr.*, 65, 1891-1902, <https://doi.org/10.1002/lno.11425>, 2020.
- Beiter, D., Weiler, M., and Blume, T.: Characterising hillslope-stream connectivity with a joint event analysis of stream and groundwater levels, *Hydrol. Earth Syst. Sci.*, 24, 5713-5744, <https://doi.org/10.5194/hess-24-5713-2020>, 2020.
- 555 Bernal, S., Butturini, A., and Sabater, F.: Variability of DOC and nitrate responses to storms in a small Mediterranean forested catchment, *Hydrol. Earth Syst. Sci.*, 6, 1031-1041, <https://doi.org/10.5194/hess-6-1031-2002>, 2002.
- Bernal, S., Butturini, A., and Sabater, F.: Seasonal variations of dissolved nitrogen and DOC : DON ratios in an intermittent Mediterranean stream, *Biogeochemistry*, 75, 351-372, <https://doi.org/10.1007/s10533-005-1246-7>, 2005.
- Bernal, S., Lupon, A., Ribot, M., Sabater, F., and Martí, E.: Riparian and in-stream controls on nutrient concentrations and fluxes in a headwater forested stream, *Biogeosciences*, 12, 1941-1954, <https://doi.org/10.5194/bg-12-1941-2015>, 2015.
- 560 Bernal, S., Lupon, A., Catalán, N., Castelar, S., and Martí, E.: Decoupling of dissolved organic matter patterns between stream and riparian groundwater in a headwater forested catchment, *Hydrol. Earth Syst. Sci.*, 22, 1897-1910, <https://doi.org/10.5194/hess-22-1897-2018>, 2018.
- Bernal, S., Lupon, A., Wollheim, W. M., Sabater, F., Poblador, S., and Martí, E.: Supply, Demand, and In-Stream Retention of Dissolved Organic Carbon and Nitrate During Storms in Mediterranean Forested Headwater Streams, *Front. Environ. Sci.*, 7, 14, <https://doi.org/10.3389/fenvs.2019.00060>, 2019.
- Bernhardt, E. S., Heffernan, J. B., Grimm, N. B., Stanley, E. H., Harvey, J. W., Arroita, M., Appling, A. P., Cohen, M. J., McDowell, W. H., Hall, R. O., Read, J. S., Roberts, B. J., Stets, E. G., and Yackulic, C. B.: The metabolic regimes of flowing waters, *Limnol. Oceanogr.*, 63, S99-S118, <https://doi.org/10.1002/lno.10726>, 2018.

- 570 Blackburn, M., Ledesma, J. L. J., Näsholm, T., Laudon, H., and Sponseller, R. A.: Evaluating hillslope and riparian contributions to dissolved nitrogen (N) export from a boreal forest catchment, *J. Geophys. Res.-Biogeosci.*, 122, 324-339, <https://doi.org/10.1002/2016jg003535>, 2017.
- Blaurock, K., Beudert, B., Gilfedder, B. S., Fleckenstein, J. H., Peiffer, S., and Hopp, L.: Missing connectivity during summer drought controls DOC mobilization and export in a small, forested catchment, *Hydrol. Earth Syst. Sci. Discuss.* [preprint], <https://doi.org/10.5194/hess-2021-89>, in review, 2021.
- 575 Boy, J., Valarezo, C., and Wilcke, W.: Water flow paths in soil control element exports in an Andean tropical montane forest, *Eur. J. Soil Sci.*, 59, 1209-1227, <https://doi.org/10.1111/j.1365-2389.2008.01063.x>, 2008.
- Brookshire, E. N. J., Valett, H. M., Thomas, S. A., and Webster, J. R.: Coupled cycling of dissolved organic nitrogen and carbon in a forest stream, *Ecology*, 86, 2487-2496, <https://doi.org/10.1890/04-1184>, 2005.
- 580 Buffam, I., Galloway, J. N., Blum, L. K., and McGlathery, K. J.: A stormflow/baseflow comparison of dissolved organic matter concentrations and bioavailability in an Appalachian stream, *Biogeochemistry*, 53, 269-306, <https://doi.org/10.1023/a:1010643432253>, 2001.
- Butturini, A., Bernal, S., Nin, E., Hellin, C., Rivero, L., Sabater, S., and Sabater, F.: Influences of the stream groundwater hydrology on nitrate concentration in unsaturated riparian area bounded by an intermittent Mediterranean stream, *Water Resour. Res.*, 39, 13, <https://doi.org/10.1029/2001wr001260>, 2003.
- 585 Caillon, F., and Schelker, J.: Dynamic transfer of soil bacteria and dissolved organic carbon into small streams during hydrological events, *Aquat. Sci.*, 82, 11, <https://doi.org/10.1007/s00027-020-0714-4>, 2020.
- Camino-Serrano, M., Pannatier, E. G., Vicca, S., Luysaert, S., Jonard, M., Ciais, P., Guenet, B., Gielen, B., Peñuelas, J., Sardans, J., Waldner, P., Etzold, S., Cecchini, G., Clarke, N., Galić, Z., Gandois, L., Hansen, K., Johnson, J., Klinck, U.,
- 590 Lachmanová, Z., Lindroos, A. J., Meesenburg, H., Nieminen, T. M., Sanders, T. G. M., Sawicka, K., Seidling, W., Thimonier, A., Vanguelova, E., Verstraeten, A., Vesterdal, L., and Janssens, I. A.: Trends in soil solution dissolved organic carbon (DOC) concentrations across European forests, *Biogeosciences*, 13, 5567-5585, <https://doi.org/10.5194/bg-13-5567-2016>, 2016.
- Casas-Ruiz, J. P., Catalán, N., Gómez-Gener, L., von Schiller, D., Obrador, B., Kothawala, D. N., López, P., Sabater, S., and
- 595 Marcé, R.: A tale of pipes and reactors: Controls on the in-stream dynamics of dissolved organic matter in rivers, *Limnol. Oceanogr.*, 62, S85-S94, <https://doi.org/10.1002/lno.10471>, 2017.
- Catalán, N., Casas-Ruiz, J. P., Arce, M. I., Abril, M., Bravo, A. G., del Campo, R., Estévez, E., Freixa, A., Giménez-Grau, P., González-Ferreras, A. M., Gómez-Gener, L., Lupon, A., Martínez, A., Palacin-Lizarbe, C., Poblador, S., Rasines-Ladero, R., Reyes, M., Rodríguez-Castillo, T., Rodríguez-Lozano, P., Sanpera-Calbet, I., Tornero, I., and Pastor, A.:
- 600 Behind the Scenes: Mechanisms Regulating Climatic Patterns of Dissolved Organic Carbon Uptake in Headwater Streams, *Global Biogeochem. Cy.*, 32, 1528-1541, <https://doi.org/10.1029/2018gb005919>, 2018.

- Cirno, C. P., and McDonnell, J. J.: Linking the hydrologic and biogeochemical controls of nitrogen transport in near-stream zones of temperate-forested catchments: a review, *J. Hydrol.*, 199, 88-120, [https://doi.org/10.1016/s0022-1694\(96\)03286-6](https://doi.org/10.1016/s0022-1694(96)03286-6), 1997.
- 605 Danger, M., Daufresne, T., Lucas, F., Pissard, S., and Lacroix, G.: Does Liebig's law of the minimum scale up from species to communities?, *Oikos*, 117, 1741-1751, <https://doi.org/10.1111/j.1600-0706.2008.16793.x>, 2008.
- de Wit, H. A., Valinia, S., Weyhenmeyer, G. A., Futter, M. N., Kortelainen, P., Austnes, K., Hessen, D. O., Raike, A., Laudon, H., and Vuorenmaa, J.: Current Browning of Surface Waters Will Be Further Promoted by Wetter Climate, *Environ. Sci. Tech. Lett.*, 3, 430-435, <https://doi.org/10.1021/acs.estlett.6b00396>, 2016.
- 610 Doblas-Miranda, E., Rovira, P., Brotons, L., Martnez-Vilalta, J., Retana, J., Pla, M., and Vayreda, J.: Soil carbon stocks and their variability across the forests, shrublands and grasslands of peninsular Spain, *Biogeosciences*, 10, 8353-8361, <https://doi.org/10.5194/bg-10-8353-2013>, 2013.
- Dodds, W. K., Lopez, A. J., Bowden, W. B., Gregory, S., Grimm, N. B., Hamilton, S. K., Hershey, A. E., Mart, E., McDowell, W. H., Meyer, J. L., Morrall, D., Mulholland, P. J., Peterson, B. J., Tank, J. L., Valett, H. M., Webster, J. R., and Wollheim, W.: N uptake as a function of concentration in streams, *J. N. Am. Benthol. Soc.*, 21, 206-220, <https://doi.org/10.2307/1468410>, 2002.
- 615 Dupas, R., Musolff, A., Jawitz, J., Rao, P. S. C., Jager, C. G., Fleckenstein, J. H., Rode, M., and Borchardt, D.: Carbon and nutrient export regimes from headwater catchments to downstream reaches, *Biogeosciences*, 14, 4391-4407, <https://doi.org/10.5194/bg-14-4391-2017>, 2017.
- 620 Eriksson, L., Johansson, E., and Wold, S.: Introduction to Multi- and Megavariate Data Analysis Using Projection Methods (PCA & PLS), Umetrics, 1999.
- Fasching, C., Akotoye, C., Bizic, M., Fonvielle, J., Ionescu, D., Mathavarajah, S., Zoccarato, L., Walsh, D. A., Grossart, H. P., and Xenopoulos, M. A.: Linking stream microbial community functional genes to dissolved organic matter and inorganic nutrients, *Limnol. Oceanogr.*, 65, S71-S87, <https://doi.org/10.1002/lno.11356>, 2020.
- 625 Fovet, O., Ruiz, L., Hrachowitz, M., Faucheux, M., and Gascuel-Oudou, C.: Hydrological hysteresis and its value for assessing process consistency in catchment conceptual models, *Hydrol. Earth Syst. Sci.*, 19, 105-123, <https://doi.org/10.5194/hess-19-105-2015>, 2015.
- Fovet, O., Humbert, G., Dupas, R., Gascuel-Oudou, C., Gruau, G., Jaffrezic, A., Thelusma, G., Faucheux, M., Gilliet, N., Hamon, Y., and Grimaldi, C.: Seasonal variability of stream water quality response to storm events captured using high-
- 630 frequency and multi-parameter data, *J. Hydrol.*, 559, 282-293, <https://doi.org/10.1016/j.jhydrol.2018.02.040>, 2018.
- Frei, S., Lischeid, G., and Fleckenstein, J. H.: Effects of micro-topography on surface-subsurface exchange and runoff generation in a virtual riparian wetland - A modeling study, *Adv. Water Resour.*, 33, 1388-1401, <https://doi.org/10.1016/j.advwatres.2010.07.006>, 2010.

- Frei, S., Knorr, K. H., Peiffer, S., and Fleckenstein, J. H.: Surface micro-topography causes hot spots of biogeochemical activity in wetland systems: A virtual modeling experiment, *J. Geophys. Res.-Biogeosci.*, 117, 18, <https://doi.org/10.1029/2012jg002012>, 2012.
- Gallart, F., Llorens, P., Latron, J., and Regüés, D.: Hydrological processes and their seasonal controls in a small Mediterranean mountain catchment in the Pyrenees, *Hydrol. Earth Syst. Sci.*, 6, 527-537, <https://doi.org/10.5194/hess-6-527-2002>, 2002.
- Godsey, S. E., Kirchner, J. W., and Clow, D. W.: Concentration-discharge relationships reflect chemostatic characteristics of US catchments, *Hydrol. Process.*, 23, 1844-1864, <https://doi.org/10.1002/hyp.7315>, 2009.
- Gordon, N. D., McMahon, T. A., Finlayson, B. L., Gippel, C. J., and Nathan, R. J.: *Stream Hydrology: An Introduction for Ecologists*, Wiley, 2004.
- Hartmann, J., Lauerwald, R., and Moosdorf, N.: A brief overview of the GLObal RIver CHEmistry Database, GLORICH, *Proced. Earth Plan. Sci.*, 10, 23-27, <https://doi.org/10.1016/j.proeps.2014.08.005>, 2014.
- Helton, A. M., Ardón, M., and Bernhardt, E. S.: Thermodynamic constraints on the utility of ecological stoichiometry for explaining global biogeochemical patterns, *Ecol. Lett.*, 18, 1049-1056, <https://doi.org/10.1111/ele.12487>, 2015.
- Hewlett, J., and Hibbert, A.: Factors affecting the response of small watersheds to precipitation in humid areas, in: *Forest Hydrology*, edited by: Sopper, W., and Lull, H., Pergamon Press, New York, 1967.
- Hinton, M. J., Schiff, S. L., and English, M. C.: The significance of storms for the concentration and export of dissolved organic carbon from two Precambrian Shield catchments, *Biogeochemistry*, 36, 67-88, <https://doi.org/10.1023/a:1005779711821>, 1997.
- Inamdar, S. P., and Mitchell, M. J.: Hydrologic and topographic controls on storm-event exports of dissolved organic carbon (DOC) and nitrate across catchment scales, *Water Resour. Res.*, 42, <https://doi.org/10.1029/2005wr004212>, 2006.
- Jencso, K. G., McGlynn, B. L., Gooseff, M. N., Wondzell, S. M., Bencala, K. E., and Marshall, L. A.: Hydrologic connectivity between landscapes and streams: Transferring reach-and plot-scale understanding to the catchment scale, *Water Resour. Res.*, 45, <https://doi.org/10.1029/2008wr007225>, 2009.
- Jung, M., Burt, T. P., and Bates, P. D.: Toward a conceptual model of floodplain water table response, *Water Resour. Res.*, 40, 13, <https://doi.org/10.1029/2003wr002619>, 2004.
- Kendall, K. A., Shanley, J. B., and McDonnell, J. J.: A hydrometric and geochemical approach to test the transmissivity feedback hypothesis during snowmelt, *J. Hydrol.*, 219, 188-205, [https://doi.org/10.1016/s0022-1694\(99\)00059-1](https://doi.org/10.1016/s0022-1694(99)00059-1), 1999.
- Köhler, S. J., Buffam, I., Seibert, J., Bishop, K. H., and Laudon, H.: Dynamics of stream water TOC concentrations in a boreal headwater catchment: Controlling factors and implications for climate scenarios, *J. Hydrol.*, 373, 44-56, <https://doi.org/10.1016/j.jhydrol.2009.04.012>, 2009.
- Ledesma, J. L. J.: Daily time series (2010 - 2012) of meteorological and hydrological variables at the Font del Regàs catchment, Spain, *HydroShare*, <https://doi.org/10.4211/hs.3366012c5e254937aa661ce2a93c3140>, 2021.

- Ledesma, J. L. J., Futter, M. N., Blackburn, M., Lidman, F., Grabs, T., Sponseller, R. A., Laudon, H., Bishop, K. H., and Köhler, S. J.: Towards an Improved Conceptualization of Riparian Zones in Boreal Forest Headwaters, *Ecosystems*, 21, 297-315, <https://doi.org/10.1007/s10021-017-0149-5>, 2018.
- 670 Ledesma, J. L. J., Ruiz-Pérez, G., Lupon, A., Poblador, S., Futter, M. N., Sabater, F., and Bernal, S.: Future changes in the Dominant Source Layer of riparian lateral water fluxes in a subhumid Mediterranean catchment. *J. Hydrol.*, 595, 126014, <https://doi.org/10.1016/j.jhydrol.2021.126014>, 2021.
- Li, A. G., Bernal, S., Kohler, B., Thomas, S. A., Martí, E., and Packman, A. I.: Residence Time in Hyporheic Bioactive Layers Explains Nitrate Uptake in Streams, *Water Resour. Res.*, 57, 10.1029/2020wr027646, 2021.
- 675 Lupon, A., Gerber, S., Sabater, F., and Bernal, S.: Climate response of the soil nitrogen cycle in three forest types of a headwater Mediterranean catchment, *J. Geophys. Res.-Biogeosci.*, 120, 859-875, <https://doi.org/10.1002/2014jg002791>, 2015.
- Lupon, A., Sabater, F., Miñarro, A., and Bernal, S.: Contribution of pulses of soil nitrogen mineralization and nitrification to soil nitrogen availability in three Mediterranean forests, *Eur. J. Soil Sci.*, 67, 303-313, <https://doi.org/10.1111/ejss.12344>,
680 2016a.
- Lupon, A., Bernal, S., Poblador, S., Martí, E., and Sabater, F.: The influence of riparian evapotranspiration on stream hydrology and nitrogen retention in a subhumid Mediterranean catchment, *Hydrol. Earth Syst. Sci.*, 20, 3831-3842, <https://doi.org/10.5194/hess-20-3831-2016>, 2016b.
- Lupon, A., Martí, E., Sabater, F., and Bernal, S.: Green light: gross primary production influences seasonal stream N export
685 by controlling fine-scale N dynamics, *Ecology*, 97, 133-144, <https://doi.org/10.1890/14-2296.1>, 2016c.
- Lupon, A., Denfeld, B. A., Laudon, H., Leach, J., and Sponseller, R. A.: Discrete groundwater inflows influence patterns of nitrogen uptake in a boreal headwater stream, *Freshw. Sci.*, 39, 228-240, <https://doi.org/10.1086/708521>, 2020a.
- Lupon, A., Catalán, N., Martí, E., and Bernal, S.: Influence of Dissolved Organic Matter Sources on In-Stream Net Dissolved Organic Carbon Uptake in a Mediterranean Stream, *Water*, 12, 15, <https://doi.org/10.3390/w12061722>, 2020b.
- 690 Manzoni, S., Čapek, P., Mooshammer, M., Lindahl, B. D., Richter, A., and Šantrůčková, H.: Optimal metabolic regulation along resource stoichiometry gradients, *Ecol. Lett.*, 20, 1182-1191, <https://doi.org/10.1111/ele.12815>, 2017.
- Marcé, R., von Schiller, D., Aguilera, R., Martí, E., and Bernal, S.: Contribution of Hydrologic Opportunity and Biogeochemical Reactivity to the Variability of Nutrient Retention in River Networks, *Global Biogeochem. Cy.*, 32, 376-388, 10.1002/2017gb005677, 2018.
- 695 McClain, M. E., Richey, J. E., and Pimentel, T. P.: Groundwater nitrogen dynamics at the terrestrial-lotic interface of a small catchment in the Central Amazon Basin, *Biogeochemistry*, 27, 113-127, 1994.
- McDowell, W. H., Bowden, W. B., and Asbury, C. E.: Riparian nitrogen dynamics in two geomorphologically distinct tropical rain-forest watersheds: subsurface solute patterns, *Biogeochemistry*, 18, 53-75, <https://doi.org/10.1007/bf00002703>, 1992.

- 700 Musolff, A., Selle, B., Büttner, O., Opitz, M., and Tittel, J.: Unexpected release of phosphate and organic carbon to streams linked to declining nitrogen depositions, *Glob. Change Biol.*, 23, 1891-1901, <https://doi.org/10.1111/gcb.13498>, 2017.
- Musolff, A., Fleckenstein, J. H., Opitz, M., Büttner, O., Kumar, R., and Tittel, J.: Spatio-temporal controls of dissolved organic carbon stream water concentrations, *J. Hydrol.*, 566, 205-215, <https://doi.org/10.1016/j.jhydrol.2018.09.011>, 2018.
- 705 Pastor, A., Compson, Z. G., Dijkstra, P., Riera, J. L., Martí, E., Sabater, F., Hungate, B. A., and Marks, J. C.: Stream carbon and nitrogen supplements during leaf litter decomposition: contrasting patterns for two foundation species, *Oecologia*, 176, 1111-1121, <https://doi.org/10.1007/s00442-014-3063-y>, 2014.
- Peterson, B. J., Wollheim, W. M., Mulholland, P. J., Webster, J. R., Meyer, J. L., Tank, J. L., Marti, E., Bowden, W. B., Valett, H. M., Hershey, A. E., McDowell, W. H., Dodds, W. K., Hamilton, S. K., Gregory, S., and Morrall, D. D.: Control of nitrogen export from watersheds by headwater streams, *Science*, 292, 86-90, <https://doi.org/10.1126/science.1056874>, 2001.
- 710 Poblador, S., Thomas, Z., Rousseau-Gueutin, P., Sabaté, S., and Sabater, F.: Riparian forest transpiration under the current and projected Mediterranean climate: Effects on soil water and nitrate uptake, *Ecohydrology*, 12, 16, 10.1002/eco.2043, 2019.
- 715 Ranalli, A. J., and Macalady, D. L.: The importance of the riparian zone and in-stream processes in nitrate attenuation in undisturbed and agricultural watersheds - A review of the scientific literature, *J. Hydrol.*, 389, 406-415, <https://doi.org/10.1016/j.jhydrol.2010.05.045>, 2010.
- Raymond, P. A., Saiers, J. E., and Sobczak, W. V.: Hydrological and biogeochemical controls on watershed dissolved organic matter transport: pulse-shunt concept, *Ecology*, 97, 5-16, <https://doi.org/10.1890/14-1684.1>, 2016.
- 720 Rodhe, A.: On the generation of stream runoff in till soils, *Nord. Hydrol.*, 20, 1-8, 1989.
- Schiff, S. L., Devito, K. J., Elgood, R. J., McCrindle, P. M., Spoelstra, J., and Dillon, P.: Two adjacent forested catchments: Dramatically different NO₃⁻ export, *Water Resour. Res.*, 38, 13, <https://doi.org/10.1029/2000wr000170>, 2002.
- Seybold, E., and McGlynn, B.: Hydrologic and biogeochemical drivers of dissolved organic carbon and nitrate uptake in a headwater stream network, *Biogeochemistry*, 138, 23-48, <https://doi.org/10.1007/s10533-018-0426-1>, 2018.
- 725 Simon, J., Bilela, S., and Rennenberg, H.: Nitrogen uptake capacity of European beech (*Fagus sylvatica* L.) only partially depends on tree age, *Trees-Struct. Funct.*, 35, 1739-1745, 10.1007/s00468-021-02190-z, 2021.
- Sterner, R. W., and Elser, J. J.: *Ecological Stoichiometry: The Biology of Elements from Molecules to the Biosphere*, Princeton University Press, 2002.
- Tank, J. L., Martí, E., Riis, T., von Schiller, D., Reisinger, A. J., Dodds, W. K., Whiles, M. R., Ashkenas, L. R., Bowden, W. B., Collins, S. M., Crenshaw, C. L., Crowl, T. A., Griffiths, N. A., Grimm, N. B., Hamilton, S. K., Johnson, S. L., McDowell, W. H., Norman, B. M., Rosi, E. J., Simon, K. S., Thomas, S. A., and Webster, J. R.: Partitioning assimilatory nitrogen uptake in streams: an analysis of stable isotope tracer additions across continents, *Ecol. Monogr.*, 88, 120-138, <https://doi.org/10.1002/ecm.1280>, 2018.
- 730

- Taylor, P. G., and Townsend, A. R.: Stoichiometric control of organic carbon-nitrate relationships from soils to the sea, *Nature*, 464, 1178-1181, <https://doi.org/10.1038/nature08985>, 2010.
- 735 Tunaley, C., Tetzlaff, D., Lessels, J., and Soulsby, C.: Linking high-frequency DOC dynamics to the age of connected water sources, *Water Resour. Res.*, 52, 5232-5247, <https://doi.org/10.1002/2015wr018419>, 2016.
- Wagner, S., Fair, J. H., Matt, S., Hosen, J. D., Raymond, P., Saiers, J., Shanley, J. B., Dittmar, T., and Stubbins, A.: Molecular Hysteresis: Hydrologically Driven Changes in Riverine Dissolved Organic Matter Chemistry During a Storm
- 740 Event, *J. Geophys. Res.-Biogeosci.*, 124, 759-774, <https://doi.org/10.1029/2018jg004817>, 2019.
- Wen, H., Perdrial, J., Abbott, B. W., Bernal, S., Dupas, R., Godsey, S. E., Harpold, A., Rizzo, D., Underwood, K., Adler, T., Sterle, G., and Li, L.: Temperature controls production but hydrology regulates export of dissolved organic carbon at the catchment scale, *Hydrol. Earth Syst. Sci.*, 24, 945-966, <https://doi.org/10.5194/hess-24-945-2020>, 2020.
- Werner, B. J., Musolff, A., Lechtenfeld, O. J., de Rooij, G. H., Oosterwoud, M. R., and Fleckenstein, J. H.: High-frequency
- 745 measurements explain quantity and quality of dissolved organic carbon mobilization in a headwater catchment, *Biogeosciences*, 16, 4497-4516, <https://doi.org/10.5194/bg-16-4497-2019>, 2019.
- Wilson, H. F., Saiers, J. E., Raymond, P. A., and Sobczak, W. V.: Hydrologic Drivers and Seasonality of Dissolved Organic Carbon Concentration, Nitrogen Content, Bioavailability, and Export in a Forested New England Stream, *Ecosystems*, 16, 604-616, <https://doi.org/10.1007/s10021-013-9635-6>, 2013.
- 750 Wold, S., Sjöström, M., and Eriksson, L.: PLS-regression: a basic tool of chemometrics, *Chemometrics Intell. Lab. Syst.*, 58, 109-130, [https://doi.org/10.1016/s0169-7439\(01\)00155-1](https://doi.org/10.1016/s0169-7439(01)00155-1), 2001.
- Yang, L., Hur, J., Lee, S., Chang, S. W., and Shin, H. S.: Dynamics of dissolved organic matter during four storm events in two forest streams: source, export, and implications for harmful disinfection byproduct formation, *Environ. Sci. Pollut. Res.*, 22, 9173-9183, 2015.
- 755 Zar, J. H.: *Biostatistical Analysis*, Prentice Hall, Upper Saddle River, NJ, 2010.
- Zimmer, M. A., and McGlynn, B. L.: Lateral, Vertical, and Longitudinal Source Area Connectivity Drive Runoff and Carbon Export Across Watershed Scales, *Water Resour. Res.*, 54, 1576-1598, <https://doi.org/10.1002/2017wr021718>, 2018.

Table 1. Relative differences in average dissolved organic carbon (DOC) and nitrate (NO_3^-) concentrations and DOC: NO_3^- molar ratios between upstream and midstream, and between midstream and downstream sites at the Font del Regàs stream for the study period (September 2010 to August 2012). Positive values indicate that either solute concentrations or stoichiometric ratios increased in the downstream direction, while negative values indicate a decrease. Differences are shown separately for base flow and storm flow conditions.

Section	Flow conditions	DOC difference	NO_3^- difference	DOC: NO_3^- difference
Upstream to midstream	Base flow	+19%	-40%	+98%
Upstream to midstream	Storm flow	+15%	-40%	+95%
Midstream to downstream	Base flow	-10%	+8%	-18%
Midstream to downstream	Storm flow	-13%	+9%	-23%

765 **Table 2.** Hydroclimatic descriptors of the five large storm events identified during the study period (September 2010 to August 2012) at
the downstream site of the Font del Regàs catchment, including duration, precipitation amount (P), accumulated precipitation seven (P-7)
and 30 (P-30) days before the event, average stream flow (Q_{avg}), average groundwater table (Gw_{avg}), groundwater table range (ΔGw), slope
of the linear relationship between riparian groundwater table and stream flow (slope), and average stream flow normalized to average
770 groundwater table (Q_{avg}/Gw_{avg}). These descriptors were used as predictors in a partial least square regression analysis, in which average
dissolved organic carbon (DOC) concentrations and average nitrate (NO_3^-) concentrations during each event at the downstream site were
the response variables (also shown). The relative increase in DOC and NO_3^- concentrations with respect to average concentrations
observed during base flow conditions at the downstream site are shown within brackets.

Event	Period	Duration (days)	P (mm)	P-7 (mm)	P-30 (mm)	Q_{avg} (mm)	Gw_{avg} (m)	ΔGw (m)	Slope (m mm ⁻¹)	Q_{avg}/Gw_{avg} (mm m ⁻¹)	DOC (mg C L ⁻¹)	NO_3^- (mg N L ⁻¹)
#1	11-14/03/2011	4	95	11	39	1.7	0.74	0.36	0.16	2.3	2.35 (167%)	0.40 (140%)
#2	14-23/03/2011	10	116	102	133	3.3	0.64	0.55	0.14	5.1	2.16 (146%)	0.33 (97%)
#3	02-12/11/2011	11	134	65	116	0.8	0.87	0.21	0.18	0.9	2.45 (178%)	0.40 (138%)
#4	14-19/11/2011	6	174	14	261	4.2	0.69	0.52	0.06	6.1	1.91 (117%)	0.29 (74%)
#5	21-30/11/2011	10	67	182	426	3.4	0.74	0.21	0.07	4.5	1.36 (55%)	0.34 (102%)





Figure 1. Map of the study catchment, Font del Regàs, including weather station, sampling sites, and main hydromorphological properties. The location of the Font del Regàs catchment within Spain is shown in the inset.



780 **Figure 2.** Box plots of (a, b) dissolved organic carbon (DOC) concentrations, (c, d) nitrate (NO₃⁻) concentrations, and (e, f) DOC:NO₃⁻
 785 molar ratios for three sampling sites along the Font del Regàs stream for the study period (September 2010 to August 2012). Data are
 shown separately for base flow (left panels) and storm flow (right panels) conditions. On each box, the central mark indicates the median,
 and the bottom and top edges of the box indicate the 25th and 75th percentiles, respectively. The whiskers extend to the most extreme data
 points not considered outliers. Arrows indicate statistical increase (upward arrow) or no statistical difference (straight arrow) between base
 flow and storm flow conditions for each colour-coded site using non-parametric Wilcoxon rank-sum tests. Results from pairwise Wilcoxon
 rank-sum tests are also shown, where * indicates statistical difference (p < 0.01) and ns indicates non-significant difference.




Figure 3. Frequency of dissolved organic carbon to nitrate (DOC:NO₃⁻) molar ratios considered optimal for heterotrophic microorganisms at three sampling sites along the Font del Regàs stream for the study period (September 2010 to August 2012). The corresponding frequencies for which there was either carbon (C) or nitrogen (N) limitation are also displayed in each case. Data are shown separately for base flow and storm flow conditions. The considered range for optimal conditions (4.8 to 11.7) was based on the global reviews of Taylor and Townsend (2010) and Helton et al. (2015).

790




Figure 4. Relationships between dissolved organic carbon (DOC) concentrations, nitrate (NO_3^-) concentrations, and DOC: NO_3^- molar ratios for three sampling sites along the Font del Regàs stream for the study period (September 2010 to August 2012). Relationships are shown separately for base flow and storm flow conditions. Coefficients of determination (R^2) and slopes are shown for significant linear correlations ($p < 0.01$).






Figure 5. Box plots of (a) precipitation amount (P), (b) average stream flow (Q_{avg}), (c) average groundwater table (Gw_{avg} , with 0 value indicating the soil surface and negative values indicating cm below the soil surface), (d) groundwater table range (ΔGw), (e) average dissolved organic carbon (DOC) concentrations at the downstream site, and (f) average nitrate (NO_3^-) concentrations at the downstream site for 'large' (N = 5) and 'medium' (N = 11) storm events identified during the study period (September 2010 to August 2012). On each box, the central mark indicates the median, and the bottom and top edges of the box indicate the 25th and 75th percentiles, respectively. The whiskers extend to the most extreme data points not considered outliers. Results from pairwise Wilcoxon rank-sum tests are also shown, where * indicates statistical difference ($p < 0.01$).




805

Figure 6. Biplot of the two-component partial least square (PLS) regression model with score and loading vectors re-scaled into the -1 to +1 numerical range in order to display the relative relationships between observations (i.e. five large storm events), predictors (i.e. duration, accumulated precipitation amount (P), accumulated precipitation seven (P-7) and 30 (P-30) days before the event, average stream flow (Q_{avg}), average groundwater table (Gw_{avg}), groundwater table range (ΔGw), slope of the linear relationship between riparian groundwater table and stream flow (slope), and average stream flow normalized to average groundwater table (Q_{avg}/Gw_{avg})), and response variables (average dissolved organic carbon (DOC) concentration and average nitrate (NO₃⁻) concentration during each event at the downstream site of the Font del Regàs catchment).

810



815 **Figure 7. (a)** Relationships between daily riparian groundwater tables and daily stream flow at the downstream site of the Font del Regàs
catchment for the five large storm events identified in the study period (September 2010 to August 2012). All events displayed clockwise
hysteresis loops for this relationship (denoted by arrows), except event #5, for which the hysteresis loop was anticlockwise. In all cases,
linear regressions showed $R^2 > 0.8$ ($p < 0.01$). **(b)** Corresponding distribution of dissolved organic carbon to nitrate (DOC:NO₃⁻) molar ratios
in each of the events. On each box, the central mark indicates the median, and the bottom and top edges of the box indicate the 25th and
75th percentiles, respectively. The whiskers extend to the most extreme data points not considered outliers. Dotted lines delimitate the
820 lower (4.8) and upper (11.7) limits of stoichiometric conditions considered optimal for stream heterotrophic microorganisms. In both
panels, the darker the blue the wetter the antecedent conditions preceding the corresponding event.



825 **Figure 8.** Diagram conceptualizing the mobilization of dissolved organic carbon (DOC) and nitrate (NO_3^-) from the riparian zone to the
 830 stream at the Font del Regàs catchment during large storm events preceded by (a) dry and (b) wet conditions. Under dry antecedent
 835 conditions, the riparian groundwater table rises more rapidly, as indicated by the steeper slope of the relationship between groundwater
 table and stream flow. This pattern suggests relatively lower water lateral transmissivity (represented by blue horizontal arrows) of each
 riparian soil layer compared to large storm events preceded by wet conditions. As a result, the ‘Dominant Source Layer’, i.e. the riparian
 zone depth stratum that contributes the most to water and solute fluxes to the stream (delimited by dashed horizontal lines), will be
 thicker during large storms preceded by dry conditions because more layers will be hydrologically activated, increasing the chances for
 relatively more DOC and NO_3^- mobilization. In addition, soluble DOC and NO_3^- pools (represented by green and purple hypothetical lines,
 respectively) will accumulate in upper riparian soil layers during dry periods, and will leach and deplete during wet periods (especially that
 of DOC as C is more limited than N in this catchment). As a result, during large storms preceded by dry conditions, groundwater will flow
 through shallower riparian soil layers with larger DOC and NO_3^- soluble pools, which will lead to large increases in stream DOC and NO_3^-
 concentrations. During large storms preceded by wet conditions, groundwater will not reach the most superficial soil layers and DOC and
 NO_3^- pools will be smaller, which will result in minor changes in stream DOC concentrations and only moderate increases in stream NO_3^-
 concentrations.

Appendix A

840 **Table A1.** Hydroclimatic descriptors of the storm events identified during the study period (September 2010 to August 2012) at the downstream site of the Font del Regàs catchment, including duration, precipitation amount (P), accumulated precipitation seven (P-7) and 30 (P-30) days before the event, average stream flow (Q_{avg}), average groundwater table (Gw_{avg}), groundwater table range (ΔGw), slope of the linear relationship between riparian groundwater table and stream flow (slope), and average stream flow normalized to average groundwater table (Q_{avg}/Gw_{avg}). Average dissolved organic carbon (DOC) concentrations and average nitrate (NO_3^-) concentrations during each event at the downstream site and the relative increase in DOC and NO_3^- concentrations with respect to average concentrations observed during base flow conditions (within brackets) are also shown. Events are subdivided into medium (*) and large (#) storms.

Event	Period	Duration (days)	P (mm)	P-7 (mm)	P-30 (mm)	Q_{avg} (mm)	Gw_{avg} (m)	ΔGw (m)	Slope (m mm ⁻¹)	Q_{avg}/Gw_{avg} (mm m ⁻¹)	DOC (mg C L ⁻¹)	NO_3^- (mg N L ⁻¹)
*1	16-19/09/2010	4	52	9	36	0.4	0.97	0.04	0.10	0.4	1.01 (15%)	0.23 (36%)
*2	14-17/05/2011	4	30	22	122	0.7	0.90	0.02	0.15	0.8	0.99 (13%)	0.17 (5%)
*3	30/05-02/06/2011	4	36	0	69	0.6	0.92	0.02	0.15	0.6	1.19 (36%)	0.18 (7%)
*4	03-06/06/2011	4	50	36	96	0.6	0.92	0.03	0.14	0.7	0.83 (-5%)	0.17 (-1%)
*5	09-15/06/2011	7	34	66	148	0.6	0.91	0.03	0.15	0.7	1.10 (26%)	0.18 (7%)
*6	25/07-03/08/2011	10	74	4	48	0.5	0.94	0.04	0.16	0.5	0.82 (-7%)	0.18 (5%)
*7	23-30/10/2011	8	98	17	39	0.3	0.95	0.09	0.17	0.3	1.53 (75%)	0.23 (41%)
*8	21-23/03/2012	3	57	6	8	0.8	0.88	0.06	0.22	0.9	0.95 (8%)	0.36 (118%)
*9	03-05/04/2012	3	29	0	63	0.7	0.95	0.06	0.13	0.7	1.23 (40%)	0.16 (-2%)
*10	18-24/05/2012	7	49	10	62	0.5	0.97	0.04	0.02	0.5	0.82 (-7%)	0.19 (12%)
*11	04-09/08/2012	6	33	0	4	0.2	1.05	0.03	0.23	0.2	1.08 (23%)	0.18 (5%)
#1	11-14/03/2011	4	95	11	39	1.7	0.74	0.36	0.16	2.3	2.35 (167%)	0.40 (140%)
#2	14-23/03/2011	10	116	102	133	3.3	0.64	0.55	0.14	5.1	2.16 (146%)	0.33 (97%)
#3	02-12/11/2011	11	134	65	116	0.8	0.87	0.21	0.18	0.9	2.45 (178%)	0.40 (138%)
#4	14-19/11/2011	6	174	14	261	4.2	0.69	0.52	0.06	6.1	1.91 (117%)	0.29 (74%)
#5	21-30/11/2011	10	67	182	426	3.4	0.74	0.21	0.07	4.5	1.36 (55%)	0.34 (102%)

845






Figure A1. Time series of precipitation at Font del Regàs catchment, as well as stream flow, dissolved organic carbon (DOC) concentrations, nitrate (NO_3^-) concentrations, and DOC: NO_3^- molar ratios at three sampling sites (upstream, midstream, and downstream) along the Font Regàs stream for the period September 2010 to August 2012.



850

Figure A2. VIP (variable influence on projection) values computed for the nine predictors (duration, precipitation amount (P), accumulated precipitation seven (P-7) and 30 (P-30) days before the large storm event, average stream flow (Q_{avg}), average groundwater table (Gw_{avg}), groundwater table range (ΔGw), slope of the linear relationship between riparian groundwater table and stream flow (slope), and average stream flow normalized to average groundwater table (Q_{avg}/Gw_{avg})) included in the partial least square regression model carried out in this study. Predictors are considered important in the overall model when $VIP > 1$. Important predictors are highlighted with darker colour.

855



860

Figure A3. Frequency of dissolved organic carbon to nitrate (DOC:NO₃⁻) molar ratios considered optimal for heterotrophic microorganisms at three sampling sites along the Font del Regàs stream for the study period (September 2010 to August 2012) for two considered cases: (a) optimal range is 4.8 to 11.7 (presented in the main body of the article and based on the global reviews of Taylor and Townsend (2010) and Helton et al. (2015)) and (b) optimal range is 4.5 to 16 (a less restricted range based on values from other modelling and site-specific studies). The corresponding frequencies for which there was either carbon (C) or nitrogen (N) limitation are also displayed. Data are shown separately for base flow and storm flow conditions.