



Hydrology and riparian forests drive carbon and nitrogen supply and DOC:NO_3^- stoichiometry along a headwater Mediterranean stream

José L. J. Ledesma^{1,2}, Anna Lupon², Eugènia Martí², Susana Bernal²

5 ¹Institute of Geography and Geoecology, Karlsruhe Institute of Technology (KIT), Karlsruhe, 76131, Germany

²Centre for Advanced Studies of Blanes, Spanish National Research Council (CEAB-CSIC), Blanes, 17300, Spain

Correspondence to: José L. J. Ledesma (jose.ledesma@kit.edu)

Abstract. In forest headwater streams, metabolic processes are predominately heterotrophic and depend on both the availability of carbon (C) and nitrogen (N) and a favourable C:N stoichiometry. In this context, hydrological conditions and the presence of riparian forests adjacent to streams can play an important, yet understudied role determining dissolved organic carbon (DOC) and nitrate (NO_3^-) concentrations and DOC:NO_3^- molar ratios. Here, we aimed to investigate how the interplay between hydrological conditions and riparian forest coverage drives DOC and NO_3^- supply and DOC:NO_3^- stoichiometry in an oligotrophic headwater Mediterranean stream. We analysed DOC and NO_3^- concentrations, and DOC:NO_3^- molar ratios during both base flow and storm flow conditions at three stream locations along a longitudinal gradient of increased riparian forest coverage. Further, we performed an event analysis to examine the hydroclimatic conditions that favour the transfer of DOC and NO_3^- from riparian soils to the stream during large storms. Stream DOC and NO_3^- concentrations were generally low (overall average \pm SD was $1.0 \pm 0.6 \text{ mg C L}^{-1}$ and $0.20 \pm 0.09 \text{ mg N L}^{-1}$), although significantly higher during storm flow compared to base flow conditions in all three stream sites. Optimal DOC:NO_3^- stoichiometry for stream heterotrophic microorganisms (corresponding to DOC:NO_3^- molar ratios between 4.8 and 11.7) was prevalent at the midstream and downstream sites under both flow conditions, whereas C-limited conditions were prevalent at the upstream site, which had no surrounding riparian forest. The hydroclimatic analysis of large storm events highlighted different patterns of DOC and NO_3^- mobilization depending on antecedent soil moisture conditions: drier antecedent conditions promoted rapid elevations of riparian groundwater tables, hydrologically activating a wider and shallower soil layer, and leading to relatively higher increases in stream DOC and NO_3^- concentrations compared to events preceded by wet conditions. These results suggest that (i) increased supply of limited resources during storms can promote in-stream heterotrophic activity during high flows, especially during large storm events preceded by dry conditions, and (ii) C-limited conditions upstream were gradually overcome downstream, likely due to higher C inputs from riparian forests present at lower elevations. The contrasting spatiotemporal patterns in DOC and NO_3^- availability and DOC:NO_3^- stoichiometry observed at the study stream suggests that groundwater inputs from riparian forests are essential for maintaining in-stream heterotrophic activity in oligotrophic, forest headwater catchments.



1 Introduction

The metabolism of lotic systems is highly dependent on environmental conditions (Bernhardt et al., 2018). In forest headwater streams, metabolism is dominated by the activity of heterotrophs because light availability is limited for primary producers and allochthonous matter inputs constitute the major energy source (Tank et al., 2018). In these environments, the supply and availability of carbon (C) and nutrients, including nitrogen (N), can therefore constrain both the activity and composition of stream heterotrophic microorganisms (Dodds et al., 2002; Brookshire et al., 2005; Fasching et al., 2020). Consequently, biotic assimilation of both dissolved organic carbon (DOC) and nitrate (NO_3^-) have been shown to relate closely with ecosystem respiration in streams worldwide (Pastor et al., 2014; Catalán et al., 2018; Tank et al., 2018; Lupon et al., 2020a), which highlights the strong linkages between resource supply, demand, and in-stream heterotrophic activity.

Yet, even if DOC and NO_3^- are available, DOC: NO_3^- stoichiometry can further influence in-stream heterotrophic activity (Elser et al., 2000; Helton et al., 2015). Heterotrophic organisms can adapt to a wide range of resource stoichiometric ratios (Manzoni et al., 2017), and thus it is difficult to establish general reference values defining optimal conditions for in-stream heterotrophic activity. For example, global studies on stoichiometric controls for a wide range of ecosystems show discrepant reference C:N molar ratios for streams and rivers; ranging from 4.8 (Taylor and Townsend, 2010) to 11.7 (Helton et al., 2015). Modelling approaches using bacterial cultures show that heterotrophic microorganisms can perform well under C:N molar ratios ranging between 5.6 and 18.4 (Danger et al., 2008). Empirical case studies in forest headwater streams reported a C:N range of 5.9 to 13.4 for optimal microbial activity (Pastor et al., 2014). Hence, considering that ecological stoichiometry theory sets biochemical constraints for metabolic activity based on the comparison of element ratios between resources and organisms (Sternner and Elser, 2002), there is a wide range of potentially optimal stoichiometric conditions for microbial heterotrophs.

In this context, hydrological processes at the riparian-stream interface, and in particular the mobilization of solutes from riparian soils to streams, can play an important role in determining stream DOC and NO_3^- availability. Riparian zones are acknowledged as the major source of DOC in a wide range of forest headwater streams because their soils generally show high organic matter contents, especially in shallower layers (Dosskey & Bertsch, 1994; Bernal et al., 2005; Köhler et al., 2009; Strohmeier et al., 2013). Relatively wet conditions in the riparian zone favour the build-up of this organic matter and can also lead to denitrification, limiting the mobilization of NO_3^- (McClain et al., 1994; Cirimo and McDonnell, 1997). However, riparian zones that experience frequent dryness, such as those in Mediterranean catchments, can manifest nitrification and be net sources of NO_3^- to streams (Lupon et al., 2016a). As a result of these riparian zone characteristics, increased DOC concentrations in forest headwaters during storm events is an almost universal phenomenon (Hinton et al., 1997; Boy et al., 2008; Yang et al., 2015; Musolff et al., 2018), whereas stream NO_3^- concentrations can increase, decrease, or remain unchanged during storms (Àvila et al., 1992; Bernal et al., 2002; Inamdar and Mitchell, 2006; Dupas et al., 2017). At the same time, climate and associated soil moisture conditions preceding storm events can influence both the amount of DOC and NO_3^- stored in riparian soils and the riparian soil layers that are hydrologically connected to the stream, thus



strongly influencing the supply of C and N to streams (Àvila et al., 1992; Fovet et al., 2015; Zimmer and McGlynn, 2018; Werner et al., 2019; Wen et al., 2020). Resulting stream DOC and NO_3^- concentrations and DOC: NO_3^- stoichiometry might thus significantly differ between base flow and storm flow conditions, as well as among storms depending on antecedent accumulated precipitation, and thereafter influence in-stream DOC and NO_3^- availability for microbial communities. Therefore, studying the role of hydrology in controlling DOC and NO_3^- supply from riparian zones to adjacent streams becomes important for the subsequent understanding of in-stream heterotrophic activity.

In this study, we aimed to investigate how the interplay between hydrological processes, antecedent climate conditions, and riparian forest coverage drives DOC and NO_3^- availability and DOC: NO_3^- stoichiometry in a forest headwater Mediterranean stream. Ambient DOC and NO_3^- concentrations in this site are relatively low compared to forest headwater streams located elsewhere (Hartmann et al., 2014; Bernal et al., 2015; Bernal et al., 2018), and thus we expected hydrological conditions and the presence of riparian forests to have a significant influence on the supply of these limited solutes for stream heterotrophic microorganisms. In order to fulfil our aim, we analysed DOC and NO_3^- concentrations, and DOC: NO_3^- molar ratios under base flow and storm flow conditions at three stream locations along a longitudinal gradient of increasing riparian forest coverage. In this study, riparian zones are defined as near-stream areas within the catchment where riparian forest is present. In addition, we performed an event analysis to examine the hydroclimatic conditions that most favoured the transfer of DOC and NO_3^- from the riparian zone to the stream during large storms, which were assumed to have a disproportionate influence on solute supply. Based on this hydroclimatic event analysis, we propose a conceptual model synthesising the complex hydrological and biogeochemical processes driving DOC and NO_3^- mobilization from riparian zones to streams during large storms.

2 Materials and methods

2.1 Study site characterization

The Font del Regàs catchment, located in the Montseny Natural Park in NE Spain (outlet at $41^\circ 50'$ N, $2^\circ 28'$ E), includes a total drainage area of 15.5 km^2 that ranges between 405 and 1603 m above sea level (m a.s.l) (Fig. 1). The climate is subhumid Mediterranean with mild winters, wet springs, and dry summers. Long-term (1940 to 2000) annual precipitation averaged $925 \pm 151 \text{ mm year}^{-1}$ (average \pm SD), whereas average annual air temperature during this period was $12.1 \pm 2.5^\circ \text{C}$ (Lupon et al., 2016b).

The altitude gradient drives changes in vegetation cover in the catchment. While deciduous European beech (*Fagus sylvatica*) forests and a small proportion of heathlands (*Calluna vulgaris* and gramineae) dominate in the upper, steeper parts of the catchment, evergreen oak (*Quercus ilex*) forests dominate in the lower parts. Riparian forests, composed of a mixture of tree species including black alder (*Alnus glutinosa*), black locust (*Robinia pseudoacacia*), European ash (*Fraxinus excelsior*), and black poplar (*Populus nigra*), develop in flat, near-stream zones of the lower parts of the catchment (Lupon et



95 al., 2016b). Both the width of the riparian forests and the total basal area of riparian trees increase downstream (Bernal et al., 2015). For its most part, Font del Regàs can be considered a closed-canopy stream (*sensu* Tank et al., 2018).

For this study, we selected three stream locations along a longitudinal gradient of increased riparian forest coverage, which delineate three nested subcatchments (Fig. 1). The upstream subcatchment (746 to 1603 m a.s.l.) drains a steep area of 1.8 km² along the first 2.9 km of the Font del Regàs stream. The vegetation here is dominated by beech forests and heathlands
 100 (92%), with lower proportions of oak forests (8%), and no riparian forests (i.e. no riparian zones). The midstream subcatchment (local drainage area of 6.8 km², ranging from 566 to 746 m a.s.l.) drains a cumulated area of 8.5 km² and includes 1.8 km more of the stream. Cumulated vegetation proportions here are 53% beech forests, 43% oak forests, and 4% mixed, 5 to 15 m wide riparian forests. The downstream subcatchment (local drainage area of 4.4 km², ranging from 503 to 566 m a.s.l.) drains a cumulated area of 13 km² and includes 1.4 km more of the stream, for a total of 6 km. Cumulated
 105 vegetation proportions at the downstream subcatchment are 46% beech forests, 48% oak forests, and 6% riparian forests, which are well-developed and can be up to 30 m wide.

2.2 Field data collection, sampling, and laboratory analyses

The present study is based on data collected during the two-year period 9th September 2010 to 31st August 2012. All data and analyses were integrated and carried out for daily resolutions, which were determined by the availability of the stream
 110 chemical data.

Cumulated precipitation was recorded at 15 min intervals using an automatic meteorological station located at the valley bottom of the catchment, near the downstream site (Fig. A1). At each of the three stream locations, referred to as sites hereafter, we measured stream water levels at 15 min intervals using pressure transducers (HOBO U20-001-04) connected to autosamplers (Teledyne Isco Model 1612). Stream flow was measured fortnightly at each site by applying the salt dilution
 115 method (Gordon et al., 2004). Rating curves obtained from the relationships between stream flow and stream water level measurements were used to construct daily time series of stream flow data at each site ($R^2 > 0.97$ in all cases; Lupon et al., 2016b). Stream flow was normalized by drainage area; thus, it is denoted by units of mm throughout ($N = 723$ days in all cases; Fig. A1).

Groundwater tables were recorded at 15 min intervals using a water pressure transducer (Druck PDCR 1830) within a
 120 piezometer placed 2.5 m away from the stream channel and located in a riparian zone area ca. 500 m downstream the downstream site (Fig. 1). Stream flow normalized by drainage area is assumed equivalent at these two locations (Ledesma et al., 2021). Furthermore, the dynamics of this dataset capture well the dynamics of the groundwater table variation in the surrounding riparian area, and therefore we are confident that the recorded pattern at the monitoring location was representative of the groundwater table variations in the riparian zone soils in the lower parts of the catchment (Ledesma et al., 2021). The available daily groundwater table records were 664 (sporadic device failure precluded a full time series for
 125 the full study period).



Stream water samples were collected every day at noon at each of the three sites using autosamplers (Teledyne Isco Model 1612), which were installed ca. 1 m belowground to keep samples cool and to prevent biogeochemical transformations (Lupon et al., 2016b). Water samples were transported to the lab every 7 to 14 days, where they were filtered through pre-washed Whatman GF/F glass fibre filters (pore size = 0.7 μm) and analysed for DOC by catalytic oxidation using a Shimadzu total organic carbon (TOC) analyser, and for NO_3^- by cadmium reduction using a Technicon autoanalyzer. Sporadic autosampler failure precluded a full time series of data for the full study period. Total number of available daily DOC concentration values for upstream, midstream, and downstream sites were 537, 668, and 620, respectively. Analogously, total number of available daily NO_3^- concentration values were 552, 676, and 613, respectively. Stream DOC and NO_3^- concentrations were used to calculate the $\text{DOC}:\text{NO}_3^-$ molar ratio for each available pair. Total number of $\text{DOC}:\text{NO}_3^-$ molar ratio values were 537, 668, and 611 for upstream, midstream, and downstream sites, respectively (Fig. A1).

2.3 Data treatment and statistical analyses

At each site, we classified daily stream flows into base flow and storm flow. This separation can be done using a variety of approaches and time resolutions, which in most cases require certain degree of subjectivity (Hewlett and Hibbert, 1967; Buffam et al., 2001; Fovet et al., 2018; Caillon and Schelker, 2020). Here, we defined the beginning of a storm flow event as the day when cumulated precipitation amounted more than 5 mm and stream flow was more than 10% higher than the previous day. We chose this precipitation threshold because hydrological responses to precipitation inputs below 5 mm day^{-1} are generally undetectable in Mediterranean catchments (Gallart et al., 2002). The end of a storm flow event was defined as the day when stream flow was less than 5% lower than the previous day. All stream flow values within event days were classified as storm flow, whereas the remaining values were classified as base flow. From the available 723 days, the upstream, midstream, and downstream sites included 128, 129, and 151 storm flow days. The small discrepancies in the storm flow correspondence among sites were likely caused by (i) differences in response times, being the downstream site slower in returning to base flow conditions; (ii) sporadic activations of intermediate intermittent tributaries; and (iii) storm origin, being eastern storms that first enter from the downstream part of the catchment more common than western storms that enter from the upper and middle parts of the catchment.

In order to investigate the influence of hydrological conditions on stream chemistry, for each site, we compared stream DOC concentrations, stream NO_3^- concentration, and stream $\text{DOC}:\text{NO}_3^-$ molar ratios between base flow and storm flow conditions using non-parametric Wilcoxon signed-rank tests for non-normally distributed data (Zar, 2010). Moreover, in order to investigate the impact of riparian forest coverage on stream chemistry, we further performed pairwise Wilcoxon tests comparing DOC and NO_3^- concentrations, and $\text{DOC}:\text{NO}_3^-$ molar ratios between stream sites (i.e. upstream *versus* midstream, upstream *versus* downstream, and midstream *versus* downstream), separately for base flow and storm flow conditions.



Based on previous studies assessing optimal DOC:NO_3^- molar ratios for microbial activity in lotic ecosystems, we classified stream DOC:NO_3^- molar ratios into ‘optimal’, ‘C-limited’, and ‘N-limited’ for stream heterotrophic microorganisms. We used the reference DOC:NO_3^- molar ratios for streams and rivers presented in the global reviews of Taylor and Townsend (2010) and Helton et al. (2015) as the lower (i.e. 4.8) and upper (i.e. 11.7) limits delimiting the range of ‘optimal’ stoichiometric conditions. Consequently, stoichiometric conditions were considered ‘C-limited’ when stream DOC:NO_3^- molar ratios were below 4.8 and ‘N-limited’ when stream DOC:NO_3^- molar ratios were above 11.7. We explored the effect of hydrology and riparian forest coverage on DOC:NO_3^- stoichiometry by calculating the frequency of optimal, C-limited, and N-limited conditions at each site and flow condition. We tested whether the frequencies of optimal, C-limited, and N-limited conditions for a given site were statistically different between base flow and storm flow by using a contingency analysis (Zar, 2010). Contingency analysis was also employed to test whether there were statistical differences in the frequency of optimal, C-limited, and N-limited conditions among sites. This analysis was done separately for base flow and storm flow conditions.

Finally, we computed simple linear regressions between stream DOC concentrations, stream NO_3^- concentrations, and stream DOC:NO_3^- molar ratios in order to investigate whether DOC or NO_3^- concentrations controlled changes in stoichiometry. Linear regressions were computed separately for each site and flow condition. Statistical analyses were carried out in the software JMP® Pro version 14.0.0 and the significance level for all analyses was set at $p < 0.01$.

2.4 Hydroclimatic analysis of large storm events

We further aimed to explore the hydroclimatic characteristics that most effectively promoted the mobilization of DOC and NO_3^- from the riparian zone to the stream during large storm events, which are generally associated with the largest exports of solutes from headwater catchments. For that, we identified the largest storm events that occurred during the study period, characterized them using nine hydroclimatic descriptors, and related those descriptors with the observed stream DOC and NO_3^- concentrations through a partial least square (PLS) regression model. This analysis was performed only for data from the downstream site, for which the riparian zone was well-developed. Moreover, this was the only site for which we could compare stream flow and associated groundwater tables, which were used in some of the hydroclimatic descriptors described below.

We identified large storm events based on three requirements: (i) a precipitation amount during the days included in the event of at least 50 mm, (ii) a riparian groundwater table elevation during the event of at least 15 cm with respect to pre-event conditions, and (iii) a complete stream chemistry data series associated with the event for the downstream site (i.e. no gaps in chemical data for the event dates). There were five large storm events that fulfilled these requirements during the study period. Each event included all days classified as storm flow for the corresponding time period, plus the day classified as base flow prior the days classified as storm flow, which was considered as the starting date of the event. This strategy followed Blaurock et al. (2021) and ensured that the relevant magnitude and range of the associated hydroclimatic event



descriptors were covered. Moreover, when there was a substantial decline in stream flow and riparian groundwater tables followed by a subsequent increase without reaching base flow conditions, we split this bimodal event into two events. In this case, the day immediately before the second increase in stream flow and riparian groundwater tables was considered as both the last day of event i and the first day of event $i+1$ (this was the case for events #1 and #2, presented later).

Each of the identified large storm events was characterized by the following hydroclimatic descriptors: *duration* (days); precipitation amount (P , mm); accumulated precipitation seven days before the event ($P-7$, mm), as a proxy for short-range antecedent soil moisture conditions; accumulated precipitation 30 days before the event ($P-30$, mm), as a proxy for long-range antecedent soil moisture conditions; average stream flow (Q_{avg} , mm); average groundwater table (Gw_{avg} , m); and groundwater table range (ΔGw , m), calculated as the absolute difference between the highest and the lowest groundwater tables measured during the event. Moreover, the hydroclimatic characterization of each event included the slope of the linear relationship between daily riparian groundwater tables and daily stream flows (*slope*, m mm⁻¹), as a proxy for the rate at which upper riparian layers are hydrologically activated in relation to increments in stream flow (Rodhe, 1989). We anticipated that this slope would vary depending on antecedent soil moisture conditions, with higher values related to drier conditions, and that this variation would have implications for solute mobilization. Finally, we also calculated the average stream flow normalized to average groundwater table (Q_{avg}/Gw_{avg} , mm m⁻¹), which served as proxy for average hydraulic conductivity in the activated layers of the riparian profile, with higher values indicating larger lateral transmissivity from the profile.

These nine hydroclimatic descriptors were considered as predictors in a PLS regression model, in which average DOC and NO₃⁻ concentrations at the downstream site during each of the five large storm events were the response variables. A PLS regression model was appropriate in our case because (i) the number of predictors was large ($N = 9$) and higher than the number of observations ($N = 5$) and (ii) there was multicollinearity among predictors (not explicitly shown) (Wold et al., 2001). For each response variable, a PLS regression model returns a goodness of fit (R^2Y), which is the variation of the response variable explained by the predictors, and a predictive ability (Q^2Y), which is an indication on how well the model can predict new data ($Q^2Y > 0.5$ are considered good models). In addition, the model returns a VIP (variable influence on projection) value associated with each predictor, which informs about the relative importance of the predictor in the overall model. Predictors with $VIP > 1$ are considered statistically important for the model performance (Eriksson et al., 1999). The PLS regression analysis was carried out in the software SIMCA (©Umetrics AB) version 14.0.

Finally, for each of the characterized large storm events, we examined the type of hysteresis loop (clockwise, anticlockwise, or linear) of the relationship between daily riparian groundwater tables and daily stream flows. This relationship typically shows clockwise hysteresis loops, which imply that, for a given stream flow, the groundwater table is higher during the rising limb than during the falling limb, indicating that near-stream zones are the dominant water sources during the rising limb (Kendall et al., 1999; Frei et al., 2010). We compared the type of hysteresis loop among the five large storm events in



order to determine potential differences in runoff generation processes and associated solute mobilization and $\text{DOC}:\text{NO}_3^-$ stoichiometry.

3 Results

225 3.1 Base flow *versus* storm flow stream chemistry and changes along the stream

Including all sites and dates, stream DOC concentrations averaged $1.0 \pm 0.6 \text{ mg C L}^{-1}$ and ranged from 0.2 to 5.4 mg C L^{-1} . Stream DOC concentrations were significantly higher during storm flow than during base flow conditions in the three sites (Wilcoxon test, $p < 0.01$). On average, stream DOC concentrations increased similarly from base flow to storm flow in all sites: by 66% at the upstream site (from 0.8 ± 0.4 to $1.4 \pm 0.8 \text{ mg C L}^{-1}$), by 60% at the midstream site (from 1.0 ± 0.5 to $1.6 \pm 0.8 \text{ mg C L}^{-1}$), and by 55% at the downstream site (from 0.9 ± 0.4 to $1.4 \pm 0.8 \text{ mg C L}^{-1}$) (Fig. 2a,b). The variability in stream DOC concentrations was relatively similar during both flow conditions, as indicated by similar coefficient of variations (CV): 54% and 57% in the upstream site, 46% and 52% in the midstream site, and 40% and 60% in the downstream site, respectively for base flow and storm flow conditions. All pairwise comparisons between sites showed statistically significant differences in stream DOC concentrations during base flow conditions (Fig. 2a). On average, stream DOC concentrations increased by 19% from the upstream to the midstream site and then decreased by 10% from the midstream to the downstream site (Table 1). This pattern was similar during storm flow conditions, i.e. a 15% increase from the upstream to the midstream site and a 13% decrease from the midstream to the downstream site were observed, but differences in DOC concentrations were statistically significant only between the midstream and the downstream sites in this case (Fig. 2b).

240 Stream NO_3^- concentrations ranged from 0.09 to 1.1 mg N L^{-1} considering all sites and dates, and averaged $0.20 \pm 0.09 \text{ mg N L}^{-1}$. Stream NO_3^- concentrations were significantly higher during storm flow than during base flow conditions in the three sites (Wilcoxon test, $p < 0.01$). The relative increase in stream NO_3^- concentrations from base flow to storm flow was remarkably similar among sites, i.e. 41% at the upstream site (from 0.26 ± 0.05 to $0.36 \pm 0.17 \text{ mg N L}^{-1}$), 40% at the midstream site (from 0.15 ± 0.03 to $0.22 \pm 0.11 \text{ mg N L}^{-1}$), and 42% at the downstream site (from 0.17 ± 0.03 to $0.24 \pm 0.10 \text{ mg N L}^{-1}$) (Fig. 2c,d). Moreover, stream NO_3^- concentrations were consistently more variable during storm flow than during base flow, as indicated by increases in the CV: from 20% to 47%, from 21% to 52%, and from 18% to 42% in upstream, midstream, and downstream sites, respectively. All pairwise comparisons between sites showed statistically significant differences in stream NO_3^- concentrations (Fig. 2c,d). On average, stream NO_3^- concentrations decreased 40% from the upstream to the midstream site and increased nearly 10% from the midstream to the downstream site (Table 1). This pattern held during both base flow and storm flow conditions.

Considering the three sites pooled together, stream $\text{DOC}:\text{NO}_3^-$ molar ratios varied from 1.0 to 28 with an average of 6.1 ± 3.3 . On average, stream $\text{DOC}:\text{NO}_3^-$ molar ratios were 20% higher during storm flow compared to base flow conditions for



the upstream (increasing from 3.8 ± 2.1 to 4.6 ± 2.0) and midstream (increasing from 7.5 ± 3.4 to 8.9 ± 3.6) sites (Wilcoxon test, $p < 0.01$) (Fig. 2e,f). There was no statistical difference in stream $\text{DOC}:\text{NO}_3^-$ molar ratios between base flow and storm flow conditions at the downstream site (Wilcoxon test, $p = 0.08$), where the overall average was 6.3 ± 2.7 . All pairwise comparisons between sites showed statistically significant differences in stream $\text{DOC}:\text{NO}_3^-$ molar ratios (Fig. 2e,f). These differences were consistent between base flow and storm flow conditions: stream $\text{DOC}:\text{NO}_3^-$ molar ratios almost doubled between upstream and midstream sites and then decreased ca. 20% from the midstream to the downstream site (Table 1).

Stream $\text{DOC}:\text{NO}_3^-$ molar ratios considered optimal for heterotrophic microorganisms (i.e. from 4.8 to 11.7) were not prevalent at the upstream site during either base flow or storm flow conditions (Fig. 3). Nevertheless, at this site, the frequency of optimal $\text{DOC}:\text{NO}_3^-$ molar ratios was statistically higher during storm flow (33%) than during base flow (18%), as indicated by the contingency analysis ($p < 0.01$). The rest of the time, stream $\text{DOC}:\text{NO}_3^-$ molar ratios were lower than 4.8, indicating that C-limited conditions for heterotrophic activity dominated in the upstream site. By contrast, optimal $\text{DOC}:\text{NO}_3^-$ molar ratios were largely prevalent (ranging from 69% to 74% of the time) at the midstream and downstream sites during both base flow and storm flow conditions (Fig. 3). According to the contingency analysis, there was a statistical difference in the frequency of $\text{DOC}:\text{NO}_3^-$ stoichiometric conditions between base flow and storm flow for the midstream site ($p < 0.01$). This difference was driven by an increase in N-limited conditions from 11% during base flow to 24% during storm flow, which was accompanied by a decrease in C-limited conditions from 15% during base flow to 3% during storm flow. At the downstream site, the frequency of $\text{DOC}:\text{NO}_3^-$ stoichiometric conditions was similar between base flow and storm flow conditions (contingency analysis, $p = 0.31$), and C-limited conditions were more frequent than N-limited conditions (25% versus 6% for the combined base flow and storm flow conditions). As a result of these patterns, there were differences in the frequency of $\text{DOC}:\text{NO}_3^-$ stoichiometric conditions among sites during both base flow and storm flow (contingency analyses, $p < 0.01$).

During base flow, no relationship between stream DOC and NO_3^- concentrations was found, whereas they were positively related during storm flow conditions at the three sites ($R^2 > 0.39$, $p < 0.01$) (Fig. 4a,b). Strong positive linear relationships were observed between stream DOC concentrations and stream $\text{DOC}:\text{NO}_3^-$ molar ratios during base flow conditions at the three sites ($R^2 > 0.80$, $p < 0.01$) (Fig. 4c). During storm flow conditions, these relationships weakened because data became more scattered as DOC concentrations increased, but were still statistically significant in all sites ($R^2 > 0.22$, $p < 0.01$) (Fig. 4d). By contrast, the three sites showed no relationship between stream NO_3^- concentrations and stream $\text{DOC}:\text{NO}_3^-$ molar ratios (Fig. 4e,f). Notably, all relationships presented in Fig. 4 were similar among sites.

3.2 Hydroclimatic analysis of large storm events

The five large storm events identified during the study period varied widely considering the nine hydroclimatic descriptors associated with them (Table 2). Total *duration* varied between 4 and 11 days and precipitation amount (P) between 67 and 174 mm. Accumulated precipitation seven ($P-7$) and, especially, 30 ($P-30$) days before the event (as proxies for short- and



285 long-range antecedent soil moisture conditions, respectively) showed wide ranges, i.e. $P-7$ varied between 11 and 182 mm and $P-30$ varied between 39 and 426 mm. Average stream flow (Q_{avg}) ranged from 0.8 to 4.2 mm, whereas average groundwater table (Gw_{avg}) varied more moderately, between 0.64 and 0.87 m below the soil surface. Groundwater table range (ΔGw) varied between 0.21 and 0.55 m. The slope of the linear relationship between daily riparian groundwater tables and daily stream flows ($slope$), as a proxy for the rate at which upper riparian layers are hydrologically activated in relation to increments in stream flow, ranged from 0.06 to 0.18 m mm⁻¹ and was negatively related with both $P-30$ ($R^2 = 0.74$, $p < 0.1$) and the average stream flow normalized to average groundwater table (Q_{avg}/Gw_{avg}) ($R^2 = 0.68$, $p < 0.1$), which was a proxy for average hydraulic conductivity in the activated layers of the riparian profile and ranged from 0.9 to 6.1 mm m⁻¹ (Table 2). During all five large storm events, stream DOC and NO₃⁻ concentrations were notably higher than during base flow conditions (Table 2). Additionally, stream DOC concentrations varied more (CV = 21%) than stream NO₃⁻ concentrations (CV = 13%) among events.

A two-component PLS regression model applied to the five large storm events dataset explained 96% (i.e. $R^2Y = 0.96$) and 94% (i.e. $R^2Y = 0.94$) of the variation in average stream DOC and NO₃⁻ concentrations, respectively. The ability of the model to predict new data was notably high, with Q^2Y values of 0.88 and 0.82 for stream DOC and NO₃⁻ concentrations, respectively. Four out of the nine predictors used in the PLS regression model reached VIP > 1 (Fig. A2), indicating that they were important for explaining the variability in the response variables. Specifically, lower $P-30$, lower Q_{avg} , higher $slope$, and lower Q_{avg}/Gw_{avg} supported higher concentrations of both DOC and NO₃⁻ (Fig. 5). These four predictors showed the highest loadings in the first component of the model, which explained most of the variation in stream DOC ($R^2Y = 0.68$) and NO₃⁻ ($R^2Y = 0.86$) concentrations. From the four important predictors, $P-30$ showed the largest loading in the second component and provided a distinct and extra statistical explanation for the DOC model compared to the NO₃⁻ model. Hence, this component explained 28% of the variation in average stream DOC concentrations, but only 8% of the variation in average stream NO₃⁻ concentrations.

Four of the five large storm events showed clockwise hysteresis loops in their riparian groundwater table – stream flow relationship, whereas event #5, with the largest $P-30$, displayed a slight anticlockwise loop (Fig. 6a). Stream DOC:NO₃⁻ molar ratios at the downstream site during the five events generally fell within the suggested optimal stoichiometric range (Fig. 6b). Nevertheless, event #5 deviated again from the general pattern and showed lower stream DOC:NO₃⁻ molar ratios than the other events, partially falling within the range defined as C-limited conditions for heterotrophic activity.

4 Discussion

4.1 The role of the riparian zone on determining DOC, NO₃⁻, and DOC:NO₃⁻ along the stream

Stream DOC concentrations at the Font del Regàs catchment (1.0 ± 0.6 mg C L⁻¹) were remarkably low compared with other forest headwaters in boreal (de Wit et al., 2016), temperate (Musolff et al., 2018), and tropical (Boy et al., 2008) regions, but



were comparable to other Mediterranean sites (Casas-Ruiz et al., 2017; Catalán et al., 2018). On the other hand, stream NO_3^- concentrations ($0.20 \pm 0.09 \text{ mg N L}^{-1}$) were somewhat higher than those typically measured in N-limited boreal streams (Blackburn et al., 2017), comparable to the low concentrations measured in tropical and Mediterranean forest headwaters (McDowell et al., 1992; Catalán et al., 2018), and notably lower than those measured in temperate sites (Musolff et al., 2017). Therefore, Font del Regàs can be considered an oligotrophic stream with relatively low N and, especially, C availability. These conditions could constrain stream productivity in general and the activity of stream heterotrophs in particular, given that this is a predominantly heterotrophic system (Lupon et al., 2016c).

Clear and consistent patterns were observed for DOC and NO_3^- concentrations and DOC: NO_3^- molar ratios along the Font del Regàs stream (Table 1, Fig. 2). DOC concentrations were especially low at the upstream site and increased moderately (ca. 20%) until the midstream site. This increase could be attributed to changes in catchment topography from upstream to midstream and the associated development of riparian forests in near-stream areas (Jencso et al., 2009; Musolff et al., 2018). The upstream site drains a steep area where soils are well-drained, and no riparian forest can develop. As geomorphology becomes more favourable for the development of wetter soils in the near-stream zone, riparian forests can establish and support relatively larger mobilization of DOC from the riparian zone to the stream (Inamdar and Mitchell, 2006; Bernal et al., 2018). However, DOC concentrations decreased ca. 10% from the midstream to the downstream site despite that riparian forests are well-developed along this section. This result suggests that DOC was not transported conservatively along the stream and that in-stream DOC demand was likely higher than supply between the midstream and downstream sites, as suggested also by Lupon et al. (2020b). This indication can be further supported by a more consistent optimal DOC: NO_3^- stoichiometry in this section compared to that observed between the upstream and midstream sites, where stoichiometric conditions evolved from C-limited to optimal.

Concentrations of NO_3^- were highest at the upstream site, and then notably decreased (ca. 40%) in the stretch to the midstream site. This result is in line with the idea that headwater streams can remove substantial amounts of NO_3^- within relatively short distances (Peterson et al., 2001). Yet, results from previous studies suggest that in-stream processing alone might had not been enough for the observed large NO_3^- removal along the upstream-midstream section (Bernal et al., 2015). Riparian zones developed along this section could contribute to the reduction in stream NO_3^- by providing groundwater inputs with low NO_3^- concentrations driven by denitrification, as observed in temperate forest catchments (Cirimo and McDonnell, 1997). However, riparian soils at Font del Regàs usually act as sources of NO_3^- to the stream because they are well-oxygenated and can sustain large nitrification rates (Lupon et al., 2016a). Alternatively, we propose that the steep topography of the upstream subcatchment led to rapid drainage of aerated soils and relatively larger NO_3^- mobilization compared to the flatter near-stream zones of the midstream subcatchment, thus partially explaining the observed upstream-midstream pattern (Schiff et al., 2002). The small increase (ca. 10%) in NO_3^- concentrations between the midstream and downstream sites could in this case be explained by the increasing influence of NO_3^- -rich riparian groundwater inputs in the



downstream areas, though in-stream mineralization of leaf litter inputs could also contribute to the observed midstream-downstream pattern (Bernal et al., 2015; Lupon et al., 2015).

350 As a result of the observed DOC and NO_3^- patterns among the stream sites, DOC: NO_3^- molar ratios doubled from the upstream to the midstream site. This change could have important implications for stream heterotrophic activity because stoichiometric conditions shifted from predominately C-limited at the upstream site to predominately optimal at the midstream site (Fig. 3). C-limited conditions at the upstream site were clearly driven by low DOC availability, which is in accordance with previous studies showing strong C-limitation in Mediterranean and semi-arid streams (Catalán et al., 2018).

355 Supporting this idea, we found that stream DOC concentrations were responsible for driving stream DOC: NO_3^- molar ratios at the three sites under almost all circumstances (Fig. 4). As the influence of the riparian forest in the catchment increased from upstream to midstream and downstream locations, potential C-limitation based on DOC: NO_3^- molar ratios was generally overcome, and stoichiometric optimal conditions for stream heterotrophic activity became prevalent. This result suggests that riparian zones are essential ecosystem compartments for ensuring the supply of dissolved organic matter

360 (DOM) to the stream and, thus, for fuelling in-stream heterotrophic activity (Lupon et al., 2020b).

It is worth mentioning that defining an optimal range of DOC: NO_3^- molar ratios for in-stream heterotrophic activity is difficult due to the lack of consistent reference values, and the fact that heterotrophic microorganisms can adapt to a wide range of resource stoichiometric ratios (Manzoni et al., 2017). We acknowledge this fact by considering a range of DOC: NO_3^- values published in the literature for defining potential optimal conditions; however, this range could be more

365 restricted than in reality. Hence, site specific and modelling studies indicate that the optimal stoichiometric C:N molar ratio can include values that expand from the lower and upper limits considered in this study (Danger et al., 2008; Pastor et al., 2014; Bastias et al., 2020). Nevertheless, we obtained very similar frequencies of optimal, C-limited, and N-limited conditions when using a less restrictive DOC: NO_3^- molar ratio range for optimal conditions (from 4.5 to 16) (Fig. A3). Therefore, we are confident that the approach used in the present study is helpful for illustrating different stoichiometric

370 conditions and how they relate to C and N availability for stream heterotrophic microorganisms at the study site and other similar sites elsewhere.

4.2 Changes in DOC, NO_3^- , and DOC: NO_3^- associated with hydrological conditions: potential implications for in-stream heterotrophic activity

Stream DOC and NO_3^- concentrations substantially increased during storm flow conditions at the three sites, a pattern

375 consistent with previous observations in other Mediterranean catchments and that can be explained by the hydrological activation of organic-rich layers in near-stream zones (Àvila et al., 1992; Bernal et al., 2002). The magnitude of change between flow conditions was different for DOC and NO_3^- at the upstream site, which led to an increase in the frequency of optimal DOC: NO_3^- stoichiometric conditions for heterotrophic activity during storm flow. By contrast, at the midstream and downstream sites, the frequency of optimal DOC: NO_3^- stoichiometric conditions was high regardless of flow conditions;



380 although, at the midstream site, N-limited and C-limited conditions became more and less frequent during storm flow, respectively. Overall, these results suggest that (i) hydrological inputs during storms increase stream DOC and NO_3^- availability by supplying allochthonous DOM and nutrients from near-stream zones, which could alleviate resource limitation and (ii) in-stream heterotrophic activity might be further favoured during high flows in sites with predominant C-limitation due to increased frequency of optimal stoichiometry.

385 Nevertheless, higher water velocities and lower water residence times during storm flows can limit the capacity of heterotrophic microorganisms to take up essential elements from the water column, especially in headwater streams with steeper slopes. Thus, the biogeochemical opportunity for in-stream processing (*sensu* Marcé et al., 2018) might be restricted to locations downstream, as proposed by the pulse-shunt concept (Raymond et al., 2016). However, there are multiple lines of evidence resulting from studies carried out in our site and in other forest catchments suggesting that high stream flows

390 might indeed offer *windows of opportunity* for in-stream heterotrophic activity in headwater locations. For example, recent studies have reported that heterotrophic microorganisms can process some of the C and N entering the stream during high flow conditions in forest headwaters across ecoregions (Seybold and McGlynn, 2018; Wollheim et al., 2018). Bernal et al. (2019) showed that in-stream DOC and NO_3^- uptake at Font del Regàs could be as high or even higher during storm flow than during base flow. Further, another study from Font del Regàs showed that DOM has a prominent protein-like character

395 in both riparian groundwater and stream water (Bernal et al., 2018), which could lead to rapid assimilation even during periods of short water residence times associated with storm flow conditions. Sporadic inputs of DOM rich in aliphatic molecules have been observed in other forest headwaters during storm flow conditions, which could further increase bioavailability and assimilation (Wilson et al., 2013; Wagner et al., 2019). Part of the in-stream biogeochemical uptake during high flows could also be triggered by extra inflows of heterotrophic bacteria mobilized from catchment soils, as

400 suggested by Caillon and Schelker (2020). Mechanistically, high flows can also enhance the hydrological interaction with streambed sediments, where microbial assemblages develop (Li et al. 2021). Lastly, and more importantly for the present study, changes in DOC, NO_3^- , and $\text{DOC}:\text{NO}_3^-$ along the Font del Regàs stream were remarkably similar during both base flow and storm flow conditions. This observation suggests that the pattern of resource consumption was maintained independently of stream flow conditions. All these pieces of evidence support that in-stream heterotrophic activity can be

405 partially sustained during storm flow conditions, especially in sites with limited C and N availability such as Font del Regàs.

4.3 Antecedent soil moisture conditions shape the mobilization of DOC and NO_3^- during large storm events

Understanding which hydrological and climatic conditions result in more effective resource mobilization from riparian soils to the stream is relevant in the context of the present study and for other oligotrophic streams. Here, we focused this question on the largest storm events, which are generally associated with the largest exports of solutes from headwater catchments

410 and have the capacity to hydrologically connect shallow, organic-rich riparian layers to the fluvial network (Godsey et al., 2009; Raymond et al., 2016; Zimmer and McGlynn, 2018). For the study period, we identified and analysed five large



storms. While the sample size was low, the nature of these events covered a wide range of hydroclimatic characteristic. Thus, we believe that the hydroclimatic analysis carried out on these few study cases is helpful for shedding light on the climatic conditions and hydrological processes that control DOC and NO_3^- mobilization from riparian zones to streams and stoichiometric dynamics during large storms in forest headwater catchments.

The PLS regression analysis showed that long-range antecedent soil moisture conditions and the relationship between riparian groundwater tables and stream flows are essential factors for understanding changes in stream DOC and NO_3^- concentrations during large storms. Particularly, antecedent drier conditions (lower $P-30$) led to relatively higher stream DOC and NO_3^- concentrations and constituted the most effective situations for the mobilization and supply of these solutes to the Font del Regàs stream. Therefore, large storms preceded by dry conditions might be essential for heterotrophic activity in this nutrient-limited stream. The large mobilization of solutes during these circumstances can be partially explained by abrupt increases in riparian groundwater tables, i.e. rapid hydrological activation of upper riparian layers in relation to increments in stream flow (higher *slope*). Hence, according to the PLS analysis, drier antecedent conditions led to steeper slopes of the relationship between groundwater table and stream flow, which were related to lower average hydraulic conductivity in the activated layers of the riparian profile (lower Q_{avg}/Gw_{avg}), and to lower stream flows (lower Q_{avg}) (Fig. 5). A faster increment in riparian groundwater table position compared to the increment in stream flow can be explained by a lower effective hydraulic conductivity of each riparian soil layer (estimated here as Q_{avg}/Gw_{avg}) induced by the pre-existing dry conditions and increased hydrological disconnection between the riparian zone and the rest of the catchment. This preconditioning will make precipitation more efficient at filling up the riparian profile vertically relative to the otherwise dominant lateral transmissivity dimension (Frei et al., 2012; Fovet et al., 2015; Tunaley et al., 2016). Consequently, more riparian layers will be hydrologically connected to the stream, increasing the thickness of the ‘Dominant Source Layer’ (i.e. the riparian zone depth stratum that contributes the most to water and solute fluxes to the stream, Ledesma et al., 2018) and, thus, the chances for relatively more DOC and NO_3^- mobilization, which in turn generally show higher concentrations closer to the soil surface (Ranalli and Macalady, 2010; Camino-Serrano et al., 2016). In addition, drier conditions can restrict decomposition and promote primary production and nitrification in the riparian soil, and thus result in the built up of soluble DOC and NO_3^- pools (Lupon et al., 2016a; Arce et al., 2019; Wen et al., 2020), which will further contribute to the mobilization of these solutes during subsequent large storm events, eventually leading to large increases in stream DOC and NO_3^- concentrations (Fig. 7a). Conversely, the PLS analysis indicated that wetter antecedent soil moisture conditions and less steep slopes of the relationship between groundwater table and stream flow resulted in relatively lower stream DOC and NO_3^- concentrations during large storms. Under these circumstances, soil moisture will be high and lateral transmissivity will be relatively larger due to increased hydrological connectivity between the riparian zone and the rest of the catchment. In addition, during large storms preceded by wet conditions, soluble DOC and NO_3^- pools in the riparian soils can be depleted from precedent storm events (Butturini et al., 2003; Zimmer and McGlynn, 2018; Werner et al., 2019), eventually leading to relatively lower increments in stream DOC and NO_3^- concentrations (Fig. 7b). Moreover, the relatively higher stream flows



during events preceded by wet conditions suggest larger spatial extension of water connectivity throughout the catchment (Tunaley et al., 2016), which may further contribute to dilute solutes, especially in oligotrophic streams such as Font del Regàs.

Importantly, our results additionally highlight that the relationship between riparian groundwater tables and stream flows is not static but variable depending on storm event characteristics. This relationship typically shows clockwise hysteresis loops and synchronous behaviour patterns between riparian groundwater tables and stream flows (Jung et al., 2004; Frei et al., 2010; Fovet et al., 2015), but here we showed that clockwise hysteresis loops during large storms were not consistent in our subhumid Mediterranean catchment and under very wet antecedent conditions the clockwise pattern can turn into anticlockwise, as seen for event #5 (Fig. 6a). Moreover, event #5 was the only large storm event for which stream $\text{DOC}:\text{NO}_3^-$ molar ratios fell outside the optimal stoichiometric range for in-stream heterotrophic activity (Fig. 6b), which was explained by a relatively lower increase in stream DOC concentrations compared to that in stream NO_3^- concentrations (Table 2). This result suggests higher depletion and dilution of riparian DOC compared to riparian NO_3^- during large storms preceded by very wet conditions (Fig. 7b), implying that C rather than N limitation might unfold in Mediterranean catchments such as Font del Regàs under such circumstances (Doblas-Miranda et al., 2013). These results highlight the relevance of climatic conditions preceding storm events in controlling the relative availability of DOC and NO_3^- in streams and, ultimately, the potential activity of heterotrophic microbial assemblages.

5 Conclusions

Spatiotemporal patterns of stream DOC and NO_3^- concentrations and $\text{DOC}:\text{NO}_3^-$ molar ratios, and thus where and when C and N supply from terrestrial ecosystems can sustain or constrain in-stream heterotrophic activity, depend on catchment hydrological and biogeochemical processes, especially those occurring at the riparian zone. Supporting this idea, we found clear and consistent changes in DOC and NO_3^- concentrations and $\text{DOC}:\text{NO}_3^-$ molar ratios along a longitudinal gradient of increased riparian forest coverage of the Font del Regàs stream, during both base flow and storm flow conditions. This outcome indicates that catchment geomorphic and topographic features, which give rise to specific vegetation and hydromorphological features, determine stream DOC and NO_3^- dynamics by modulating riparian forest formation (Inamdar and Mitchell, 2006; Ledesma et al., 2018).

Both DOC and NO_3^- concentrations were relatively low along the Font del Regàs stream, but they significantly increase during storm flow compared to base flow conditions. Optimal $\text{DOC}:\text{NO}_3^-$ stoichiometry for heterotrophic microorganisms was prevalent at midstream and downstream locations regardless of hydrological conditions, whereas they were more frequent during storm flow than during base flow conditions upstream. We conclude that hydrology plays a critical role in controlling C and N supply as well as $\text{DOC}:\text{NO}_3^-$ stoichiometric requirements in close-canopy, oligotrophic Mediterranean streams because solute mobilization during high flows can alleviate resource limitation and burst in-stream heterotrophic



activity during or right after storm events. Further, our results suggest that increases in stream DOC and NO_3^- concentrations during large storm events are controlled by complex hydrological processes involving antecedent soil moisture conditions. Particularly, large storms preceded by drier conditions lead to more rapid elevations of riparian groundwater tables and to larger increases in stream DOC and NO_3^- concentrations. As the world climate continues changing, including an increased
480 variability in precipitation patterns in the Mediterranean ecoregion (i.e. larger precipitation events and longer dry spells), it will be critical to understand hydrological and biogeochemical processes at the stream-riparian interface, and how the aftermath hydrology plays out for in-stream heterotrophic activity and consequent ecosystem functioning in Mediterranean forest headwaters.

Data availability

485 Precipitation, stream flow, groundwater table, and stream chemistry data used in the present study are available in the research data repository HydroShare at <https://www.hydroshare.org/resource/3366012c5e254937aa661ce2a93c3140/> (Ledesma, 2021) and <https://www.hydroshare.org/resource/0b4eb61ad7544ffb970c1600927a819f/>.

Author contribution

JLJL: Conceptualization, Formal analysis, Funding acquisition, Methodology, Visualization, Writing – original draft
490 preparation. AA: Conceptualization, Investigation, Resources, Writing – review & editing. EM: Conceptualization, Supervision, Writing – review & editing. SB: Conceptualization, Investigation, Resources, Supervision, Writing – review & editing.

Competing interests

The authors declare that they have no conflict of interest.

500 Acknowledgements

JLJL was funded by the project RIPARIONS granted by the European Commission through a Marie Skłodowska Curie Individual Fellowship (H2020-MSCA-IF-2018-834363) and by the Spanish Government through a Juan de la Cierva grant (FJCI-2017-32111). AL was supported by the Catalan Government and the European Commission through a MSCA-Beatriu de Pinós grant (BP-2018-00082). SB work was funded by the CANTERA project (RTI2018-094521-B-I00) and a Ramon y Cajal fellowship (RYC-2017-22643) from the Spanish Ministry of Science, Innovation, and Universities and AEI/FEDER-UE.



References

- Arce, M. I., Mendoza-Lera, C., Almagro, M., Catalán, N., Romaní, A. M., Martí, E., Gómez, R., Bernal, S., Foulquier, A., Mutz, M., Marcé, R., Zoppini, A., Gionchetta, G., Weigelhofer, G., del Campo, R., Robinson, C. T., Gilmer, A., Rulik, M., Obrador, B., Shumilova, O., Zlatanović, S., Arnon, S., Baldrian, P., Singer, G., Datry, T., Skoulikidis, N., Tietjen, B., and von Schiller, D.: A conceptual framework for understanding the biogeochemistry of dry riverbeds through the lens of soil science, *Earth-Sci. Rev.*, 188, 441-453, 10.1016/j.earscirev.2018.12.001, 2019.
- Àvila, A., Piñol, J., Rodà, F., and Neal, C.: Storm solute behavior in a montane Mediterranean forested catchment, *J. Hydrol.*, 140, 143-161, [https://doi.org/10.1016/0022-1694\(92\)90238-q](https://doi.org/10.1016/0022-1694(92)90238-q), 1992.
- Bastias, E., Ribot, M., Bernal, S., Sabater, F., and Martí, E.: Microbial uptake of nitrogen and carbon from the water column by litter-associated microbes differs among litter species, *Limnol. Oceanogr.*, 65, 1891-1902, <https://doi.org/10.1002/lno.11425>, 2020.
- Bernal, S., Butturini, A., and Sabater, F.: Variability of DOC and nitrate responses to storms in a small Mediterranean forested catchment, *Hydrol. Earth Syst. Sci.*, 6, 1031-1041, <https://doi.org/10.5194/hess-6-1031-2002>, 2002.
- Bernal, S., Butturini, A., and Sabater, F.: Seasonal variations of dissolved nitrogen and DOC : DON ratios in an intermittent Mediterranean stream, *Biogeochemistry*, 75, 351-372, <https://doi.org/10.1007/s10533-005-1246-7>, 2005.
- Bernal, S., Lupon, A., Ribot, M., Sabater, F., and Martí, E.: Riparian and in-stream controls on nutrient concentrations and fluxes in a headwater forested stream, *Biogeosciences*, 12, 1941-1954, <https://doi.org/10.5194/bg-12-1941-2015>, 2015.
- Bernal, S., Lupon, A., Catalán, N., Castelar, S., and Martí, E.: Decoupling of dissolved organic matter patterns between stream and riparian groundwater in a headwater forested catchment, *Hydrol. Earth Syst. Sci.*, 22, 1897-1910, <https://doi.org/10.5194/hess-22-1897-2018>, 2018.
- Bernal, S., Lupon, A., Wollheim, W. M., Sabater, F., Poblador, S., and Martí, E.: Supply, Demand, and In-Stream Retention of Dissolved Organic Carbon and Nitrate During Storms in Mediterranean Forested Headwater Streams, *Front. Environ. Sci.*, 7, 14, <https://doi.org/10.3389/fenvs.2019.00060>, 2019.
- Bernhardt, E. S., Heffernan, J. B., Grimm, N. B., Stanley, E. H., Harvey, J. W., Arroita, M., Appling, A. P., Cohen, M. J., McDowell, W. H., Hall, R. O., Read, J. S., Roberts, B. J., Stets, E. G., and Yackulic, C. B.: The metabolic regimes of flowing waters, *Limnol. Oceanogr.*, 63, S99-S118, <https://doi.org/10.1002/lno.10726>, 2018.
- Blackburn, M., Ledesma, J. L. J., Näsholm, T., Laudon, H., and Sponseller, R. A.: Evaluating hillslope and riparian contributions to dissolved nitrogen (N) export from a boreal forest catchment, *J. Geophys. Res.-Biogeosci.*, 122, 324-339, <https://doi.org/10.1002/2016jg003535>, 2017.
- Blaurock, K., Beudert, B., Gilfedder, B. S., Fleckenstein, J. H., Peiffer, S., and Hopp, L.: Missing connectivity during summer drought controls DOC mobilization and export in a small, forested catchment, *Hydrol. Earth Syst. Sci. Discuss.* [preprint], <https://doi.org/10.5194/hess-2021-89>, in review, 2021.



- Boy, J., Valarezo, C., and Wilcke, W.: Water flow paths in soil control element exports in an Andean tropical montane
 535 forest, *Eur. J. Soil Sci.*, 59, 1209-1227, <https://doi.org/10.1111/j.1365-2389.2008.01063.x>, 2008.
- Brookshire, E. N. J., Valett, H. M., Thomas, S. A., and Webster, J. R.: Coupled cycling of dissolved organic nitrogen and
 carbon in a forest stream, *Ecology*, 86, 2487-2496, <https://doi.org/10.1890/04-1184>, 2005.
- Buffam, I., Galloway, J. N., Blum, L. K., and McGlathery, K. J.: A stormflow/baseflow comparison of dissolved organic
 matter concentrations and bioavailability in an Appalachian stream, *Biogeochemistry*, 53, 269-306,
 540 <https://doi.org/10.1023/a:1010643432253>, 2001.
- Butturini, A., Bernal, S., Nin, E., Hellin, C., Rivero, L., Sabater, S., and Sabater, F.: Influences of the stream groundwater
 hydrology on nitrate concentration in unsaturated riparian area bounded by an intermittent Mediterranean stream, *Water
 Resour. Res.*, 39, 13, <https://doi.org/10.1029/2001wr001260>, 2003.
- Caillon, F., and Schelker, J.: Dynamic transfer of soil bacteria and dissolved organic carbon into small streams during
 545 hydrological events, *Aquat. Sci.*, 82, 11, <https://doi.org/10.1007/s00027-020-0714-4>, 2020.
- Camino-Serrano, M., Pannatier, E. G., Vicca, S., Luyssaert, S., Jonard, M., Ciais, P., Guenet, B., Gielen, B., Peñuelas, J.,
 Sardans, J., Waldner, P., Etzold, S., Cecchini, G., Clarke, N., Galić, Z., Gandois, L., Hansen, K., Johnson, J., Klinck, U.,
 Lachmanová, Z., Lindroos, A. J., Meessenburg, H., Nieminen, T. M., Sanders, T. G. M., Sawicka, K., Seidling, W.,
 Thimonier, A., Vanguelova, E., Verstraeten, A., Vesterdal, L., and Janssens, I. A.: Trends in soil solution dissolved
 550 organic carbon (DOC) concentrations across European forests, *Biogeosciences*, 13, 5567-5585,
<https://doi.org/10.5194/bg-13-5567-2016>, 2016.
- Casas-Ruiz, J. P., Catalán, N., Gómez-Gener, L., von Schiller, D., Obrador, B., Kothawala, D. N., López, P., Sabater, S., and
 Marcé, R.: A tale of pipes and reactors: Controls on the in-stream dynamics of dissolved organic matter in rivers, *Limnol.
 Oceanogr.*, 62, S85-S94, <https://doi.org/10.1002/lno.10471>, 2017.
- 555 Catalán, N., Casas-Ruiz, J. P., Arce, M. I., Abril, M., Bravo, A. G., del Campo, R., Estévez, E., Freixa, A., Giménez-Grau,
 P., González-Ferreras, A. M., Gómez-Gener, L., Lupon, A., Martínez, A., Palacin-Lizarbe, C., Poblador, S., Rasines-
 Ladero, R., Reyes, M., Rodríguez-Castillo, T., Rodríguez-Lozano, P., Sanpera-Calbet, I., Tornero, I., and Pastor, A.:
 Behind the Scenes: Mechanisms Regulating Climatic Patterns of Dissolved Organic Carbon Uptake in Headwater
 Streams, *Global Biogeochem. Cy.*, 32, 1528-1541, <https://doi.org/10.1029/2018gb005919>, 2018.
- 560 Cirno, C. P., and McDonnell, J. J.: Linking the hydrologic and biogeochemical controls of nitrogen transport in near-stream
 zones of temperate-forested catchments: a review, *J. Hydrol.*, 199, 88-120, [https://doi.org/10.1016/s0022-1694\(96\)03286-6](https://doi.org/10.1016/s0022-1694(96)03286-6), 1997.
- Danger, M., Daufresne, T., Lucas, F., Pissard, S., and Lacroix, G.: Does Liebig's law of the minimum scale up from species
 to communities?, *Oikos*, 117, 1741-1751, <https://doi.org/10.1111/j.1600-0706.2008.16793.x>, 2008.
- 565 de Wit, H. A., Valinia, S., Weyhenmeyer, G. A., Futter, M. N., Kortelainen, P., Austnes, K., Hessen, D. O., Räike, A.,
 Laudon, H., and Vuorenmaa, J.: Current Browning of Surface Waters Will Be Further Promoted by Wetter Climate,
Environ. Sci. Tech. Lett., 3, 430-435, <https://doi.org/10.1021/acs.estlett.6b00396>, 2016.



- Doblas-Miranda, E., Rovira, P., Brotons, L., Martínez-Vilalta, J., Retana, J., Pla, M., and Vayreda, J.: Soil carbon stocks and their variability across the forests, shrublands and grasslands of peninsular Spain, *Biogeosciences*, 10, 8353-8361, <https://doi.org/10.5194/bg-10-8353-2013>, 2013.
- Dodds, W. K., López, A. J., Bowden, W. B., Gregory, S., Grimm, N. B., Hamilton, S. K., Hershey, A. E., Martí, E., McDowell, W. H., Meyer, J. L., Morrall, D., Mulholland, P. J., Peterson, B. J., Tank, J. L., Valett, H. M., Webster, J. R., and Wollheim, W.: N uptake as a function of concentration in streams, *J. N. Am. Benthol. Soc.*, 21, 206-220, <https://doi.org/10.2307/1468410>, 2002.
- Dupas, R., Musolff, A., Jawitz, J., Rao, P. S. C., Jager, C. G., Fleckenstein, J. H., Rode, M., and Borchardt, D.: Carbon and nutrient export regimes from headwater catchments to downstream reaches, *Biogeosciences*, 14, 4391-4407, <https://doi.org/10.5194/bg-14-4391-2017>, 2017.
- Eriksson, L., Johansson, E., and Wold, S.: Introduction to Multi- and Megavariate Data Analysis Using Projection Methods (PCA & PLS), Umetrics, 1999.
- Fasching, C., Akotoye, C., Bižić, M., Fonvielle, J., Ionescu, D., Mathavarajah, S., Zoccarato, L., Walsh, D. A., Grossart, H. P., and Xenopoulos, M. A.: Linking stream microbial community functional genes to dissolved organic matter and inorganic nutrients, *Limnol. Oceanogr.*, 65, S71-S87, <https://doi.org/10.1002/lno.11356>, 2020.
- Fovet, O., Ruiz, L., Hrachowitz, M., Fauchaux, M., and Gascuel-Oudou, C.: Hydrological hysteresis and its value for assessing process consistency in catchment conceptual models, *Hydrol. Earth Syst. Sci.*, 19, 105-123, <https://doi.org/10.5194/hess-19-105-2015>, 2015.
- Fovet, O., Humbert, G., Dupas, R., Gascuel-Oudou, C., Gruau, G., Jaffrezic, A., Thelusma, G., Fauchaux, M., Gilliet, N., Hamon, Y., and Grimaldi, C.: Seasonal variability of stream water quality response to storm events captured using high-frequency and multi-parameter data, *J. Hydrol.*, 559, 282-293, <https://doi.org/10.1016/j.jhydrol.2018.02.040>, 2018.
- Frei, S., Lischeid, G., and Fleckenstein, J. H.: Effects of micro-topography on surface-subsurface exchange and runoff generation in a virtual riparian wetland - A modeling study, *Adv. Water Resour.*, 33, 1388-1401, <https://doi.org/10.1016/j.advwatres.2010.07.006>, 2010.
- Frei, S., Knorr, K. H., Peiffer, S., and Fleckenstein, J. H.: Surface micro-topography causes hot spots of biogeochemical activity in wetland systems: A virtual modeling experiment, *J. Geophys. Res.-Biogeosci.*, 117, 18, <https://doi.org/10.1029/2012jg002012>, 2012.
- Gallart, F., Llorens, P., Latron, J., and Regüés, D.: Hydrological processes and their seasonal controls in a small Mediterranean mountain catchment in the Pyrenees, *Hydrol. Earth Syst. Sci.*, 6, 527-537, <https://doi.org/10.5194/hess-6-527-2002>, 2002.
- Godsey, S. E., Kirchner, J. W., and Clow, D. W.: Concentration-discharge relationships reflect chemostatic characteristics of US catchments, *Hydrol. Process.*, 23, 1844-1864, <https://doi.org/10.1002/hyp.7315>, 2009.
- Gordon, N. D., McMahon, T. A., Finlayson, B. L., Gippel, C. J., and Nathan, R. J.: Stream Hydrology: An Introduction for Ecologists, Wiley, 2004.



- Hartmann, J., Lauerwald, R., and Moosdorf, N.: A brief overview of the GLObal RIver CHEmistry Database, GLORICH, *Proced. Earth Plan. Sci.*, 10, 23-27, <https://doi.org/10.1016/j.proeps.2014.08.005>, 2014.
- Helton, A. M., Ardón, M., and Bernhardt, E. S.: Thermodynamic constraints on the utility of ecological stoichiometry for explaining global biogeochemical patterns, *Ecol. Lett.*, 18, 1049-1056, <https://doi.org/10.1111/ele.12487>, 2015.
- Hewlett, J., and Hibbert, A.: Factors affecting the response of small watersheds to precipitation in humid areas, in: *Forest Hydrology*, edited by: Sopper, W., and Lull, H., Pergamon Press, New York, 1967.
- Hinton, M. J., Schiff, S. L., and English, M. C.: The significance of storms for the concentration and export of dissolved organic carbon from two Precambrian Shield catchments, *Biogeochemistry*, 36, 67-88, <https://doi.org/10.1023/a:1005779711821>, 1997.
- Inamdar, S. P., and Mitchell, M. J.: Hydrologic and topographic controls on storm-event exports of dissolved organic carbon (DOC) and nitrate across catchment scales, *Water Resour. Res.*, 42, <https://doi.org/10.1029/2005wr004212>, 2006.
- Jencso, K. G., McGlynn, B. L., Gooseff, M. N., Wondzell, S. M., Bencala, K. E., and Marshall, L. A.: Hydrologic connectivity between landscapes and streams: Transferring reach-and plot-scale understanding to the catchment scale, *Water Resour. Res.*, 45, <https://doi.org/10.1029/2008wr007225>, 2009.
- Jung, M., Burt, T. P., and Bates, P. D.: Toward a conceptual model of floodplain water table response, *Water Resour. Res.*, 40, 13, <https://doi.org/10.1029/2003wr002619>, 2004.
- Kendall, K. A., Shanley, J. B., and McDonnell, J. J.: A hydrometric and geochemical approach to test the transmissivity feedback hypothesis during snowmelt, *J. Hydrol.*, 219, 188-205, [https://doi.org/10.1016/s0022-1694\(99\)00059-1](https://doi.org/10.1016/s0022-1694(99)00059-1), 1999.
- Köhler, S. J., Buffam, I., Seibert, J., Bishop, K. H., and Laudon, H.: Dynamics of stream water TOC concentrations in a boreal headwater catchment: Controlling factors and implications for climate scenarios, *J. Hydrol.*, 373, 44-56, <https://doi.org/10.1016/j.jhydrol.2009.04.012>, 2009.
- Ledesma, J. L. J.: Daily time series (2010 - 2012) of meteorological and hydrological variables at the Font del Regàs catchment, Spain, *HydroShare*, <https://doi.org/10.4211/hs.3366012c5e254937aa661ce2a93c3140>, 2021.
- Ledesma, J. L. J., Futter, M. N., Blackburn, M., Lidman, F., Grabs, T., Sponseller, R. A., Laudon, H., Bishop, K. H., and Köhler, S. J.: Towards an Improved Conceptualization of Riparian Zones in Boreal Forest Headwaters, *Ecosystems*, 21, 297-315, <https://doi.org/10.1007/s10021-017-0149-5>, 2018.
- Ledesma, J. L. J., Ruiz-Pérez, G., Lupon, A., Poblador, S., Futter, M. N., Sabater, F., and Bernal, S.: Future changes in the Dominant Source Layer of riparian lateral water fluxes in a subhumid Mediterranean catchment. *J. Hydrol.*, 595, 126014, <https://doi.org/10.1016/j.jhydrol.2021.126014>, 2021.
- Li, A. G., Bernal, S., Kohler, B., Thomas, S. A., Martí, E., and Packman, A. I.: Residence Time in Hyporheic Bioactive Layers Explains Nitrate Uptake in Streams, *Water Resour. Res.*, 57, 10.1029/2020wr027646, 2021.
- Lupon, A., Gerber, S., Sabater, F., and Bernal, S.: Climate response of the soil nitrogen cycle in three forest types of a headwater Mediterranean catchment, *J. Geophys. Res.-Biogeosci.*, 120, 859-875, <https://doi.org/10.1002/2014jg002791>, 2015.



- Lupon, A., Sabater, F., Miñarro, A., and Bernal, S.: Contribution of pulses of soil nitrogen mineralization and nitrification to soil nitrogen availability in three Mediterranean forests, *Eur. J. Soil Sci.*, 67, 303-313, <https://doi.org/10.1111/ejss.12344>, 2016a.
- Lupon, A., Bernal, S., Poblador, S., Martí, E., and Sabater, F.: The influence of riparian evapotranspiration on stream hydrology and nitrogen retention in a subhumid Mediterranean catchment, *Hydrol. Earth Syst. Sci.*, 20, 3831-3842, <https://doi.org/10.5194/hess-20-3831-2016>, 2016b.
- Lupon, A., Martí, E., Sabater, F., and Bernal, S.: Green light: gross primary production influences seasonal stream N export by controlling fine-scale N dynamics, *Ecology*, 97, 133-144, <https://doi.org/10.1890/14-2296.1>, 2016c.
- Lupon, A., Denfeld, B. A., Laudon, H., Leach, J., and Sponseller, R. A.: Discrete groundwater inflows influence patterns of nitrogen uptake in a boreal headwater stream, *Freshw. Sci.*, 39, 228-240, <https://doi.org/10.1086/708521>, 2020a.
- Lupon, A., Catalán, N., Martí, E., and Bernal, S.: Influence of Dissolved Organic Matter Sources on In-Stream Net Dissolved Organic Carbon Uptake in a Mediterranean Stream, *Water*, 12, 15, <https://doi.org/10.3390/w12061722>, 2020b.
- Manzoni, S., Čapek, P., Mooshammer, M., Lindahl, B. D., Richter, A., and Šantrůčková, H.: Optimal metabolic regulation along resource stoichiometry gradients, *Ecol. Lett.*, 20, 1182-1191, <https://doi.org/10.1111/ele.12815>, 2017.
- Marcé, R., von Schiller, D., Aguilera, R., Martí, E., and Bernal, S.: Contribution of Hydrologic Opportunity and Biogeochemical Reactivity to the Variability of Nutrient Retention in River Networks, *Global Biogeochem. Cy.*, 32, 376-388, [10.1002/2017gb005677](https://doi.org/10.1002/2017gb005677), 2018.
- McClain, M. E., Richey, J. E., and Pimentel, T. P.: Groundwater nitrogen dynamics at the terrestrial-lotic interface of a small catchment in the Central Amazon Basin, *Biogeochemistry*, 27, 113-127, 1994.
- McDowell, W. H., Bowden, W. B., and Asbury, C. E.: Riparian nitrogen dynamics in two geomorphologically distinct tropical rain-forest watersheds: subsurface solute patterns, *Biogeochemistry*, 18, 53-75, <https://doi.org/10.1007/bf00002703>, 1992.
- Musolff, A., Selle, B., Büttner, O., Opitz, M., and Tittel, J.: Unexpected release of phosphate and organic carbon to streams linked to declining nitrogen depositions, *Glob. Change Biol.*, 23, 1891-1901, <https://doi.org/10.1111/gcb.13498>, 2017.
- Musolff, A., Fleckenstein, J. H., Opitz, M., Büttner, O., Kumar, R., and Tittel, J.: Spatio-temporal controls of dissolved organic carbon stream water concentrations, *J. Hydrol.*, 566, 205-215, <https://doi.org/10.1016/j.jhydrol.2018.09.011>, 2018.
- Pastor, A., Compson, Z. G., Dijkstra, P., Riera, J. L., Martí, E., Sabater, F., Hungate, B. A., and Marks, J. C.: Stream carbon and nitrogen supplements during leaf litter decomposition: contrasting patterns for two foundation species, *Oecologia*, 176, 1111-1121, <https://doi.org/10.1007/s00442-014-3063-y>, 2014.
- Peterson, B. J., Wollheim, W. M., Mulholland, P. J., Webster, J. R., Meyer, J. L., Tank, J. L., Marti, E., Bowden, W. B., Valett, H. M., Hershey, A. E., McDowell, W. H., Dodds, W. K., Hamilton, S. K., Gregory, S., and Morrall, D. D.: Control of nitrogen export from watersheds by headwater streams, *Science*, 292, 86-90, <https://doi.org/10.1126/science.1056874>, 2001.



- 670 Ranalli, A. J., and Macalady, D. L.: The importance of the riparian zone and in-stream processes in nitrate attenuation in undisturbed and agricultural watersheds - A review of the scientific literature, *J. Hydrol.*, 389, 406-415, <https://doi.org/10.1016/j.jhydrol.2010.05.045>, 2010.
- Raymond, P. A., Saiers, J. E., and Sobczak, W. V.: Hydrological and biogeochemical controls on watershed dissolved organic matter transport: pulse-shunt concept, *Ecology*, 97, 5-16, <https://doi.org/10.1890/14-1684.1>, 2016.
- 675 Rodhe, A.: On the generation of stream runoff in till soils, *Nord. Hydrol.*, 20, 1-8, 1989.
- Schiff, S. L., Devito, K. J., Elgood, R. J., McCrindle, P. M., Spoelstra, J., and Dillon, P.: Two adjacent forested catchments: Dramatically different NO_3^- export, *Water Resour. Res.*, 38, 13, <https://doi.org/10.1029/2000wr000170>, 2002.
- Seybold, E., and McGlynn, B.: Hydrologic and biogeochemical drivers of dissolved organic carbon and nitrate uptake in a headwater stream network, *Biogeochemistry*, 138, 23-48, <https://doi.org/10.1007/s10533-018-0426-1>, 2018.
- 680 Sterner, R. W., and Elser, J. J.: *Ecological Stoichiometry: The Biology of Elements from Molecules to the Biosphere*, Princeton University Press, 2002.
- Tank, J. L., Martí, E., Riis, T., von Schiller, D., Reisinger, A. J., Dodds, W. K., Whiles, M. R., Ashkenas, L. R., Bowden, W. B., Collins, S. M., Crenshaw, C. L., Crowl, T. A., Griffiths, N. A., Grimm, N. B., Hamilton, S. K., Johnson, S. L., McDowell, W. H., Norman, B. M., Rosi, E. J., Simon, K. S., Thomas, S. A., and Webster, J. R.: Partitioning assimilatory
- 685 nitrogen uptake in streams: an analysis of stable isotope tracer additions across continents, *Ecol. Monogr.*, 88, 120-138, <https://doi.org/10.1002/ecm.1280>, 2018.
- Taylor, P. G., and Townsend, A. R.: Stoichiometric control of organic carbon-nitrate relationships from soils to the sea, *Nature*, 464, 1178-1181, <https://doi.org/10.1038/nature08985>, 2010.
- Tunaley, C., Tetzlaff, D., Lessels, J., and Soulsby, C.: Linking high-frequency DOC dynamics to the age of connected water
- 690 sources, *Water Resour. Res.*, 52, 5232-5247, <https://doi.org/10.1002/2015wr018419>, 2016.
- Wagner, S., Fair, J. H., Matt, S., Hosen, J. D., Raymond, P., Saiers, J., Shanley, J. B., Dittmar, T., and Stubbins, A.: Molecular Hysteresis: Hydrologically Driven Changes in Riverine Dissolved Organic Matter Chemistry During a Storm Event, *J. Geophys. Res.-Biogeosci.*, 124, 759-774, <https://doi.org/10.1029/2018jg004817>, 2019.
- Wen, H., Perdrial, J., Abbott, B. W., Bernal, S., Dupas, R., Godsey, S. E., Harpold, A., Rizzo, D., Underwood, K., Adler, T.,
- 695 Sterle, G., and Li, L.: Temperature controls production but hydrology regulates export of dissolved organic carbon at the catchment scale, *Hydrol. Earth Syst. Sci.*, 24, 945-966, <https://doi.org/10.5194/hess-24-945-2020>, 2020.
- Werner, B. J., Musolff, A., Lechtenfeld, O. J., de Rooij, G. H., Oosterwoud, M. R., and Fleckenstein, J. H.: High-frequency measurements explain quantity and quality of dissolved organic carbon mobilization in a headwater catchment, *Biogeosciences*, 16, 4497-4516, <https://doi.org/10.5194/bg-16-4497-2019>, 2019.
- 700 Wilson, H. F., Saiers, J. E., Raymond, P. A., and Sobczak, W. V.: Hydrologic Drivers and Seasonality of Dissolved Organic Carbon Concentration, Nitrogen Content, Bioavailability, and Export in a Forested New England Stream, *Ecosystems*, 16, 604-616, <https://doi.org/10.1007/s10021-013-9635-6>, 2013.



- Wold, S., Sjöström, M., and Eriksson, L.: PLS-regression: a basic tool of chemometrics, *Chemometrics Intell. Lab. Syst.*, 58, 109-130, [https://doi.org/10.1016/s0169-7439\(01\)00155-1](https://doi.org/10.1016/s0169-7439(01)00155-1), 2001.
- 705 Yang, L., Hur, J., Lee, S., Chang, S. W., and Shin, H. S.: Dynamics of dissolved organic matter during four storm events in two forest streams: source, export, and implications for harmful disinfection byproduct formation, *Environ. Sci. Pollut. Res.*, 22, 9173-9183, 2015.
- Zar, J. H.: *Biostatistical Analysis*, Prentice Hall, Upper Saddle River, NJ, 2010.
- Zimmer, M. A., and McGlynn, B. L.: Lateral, Vertical, and Longitudinal Source Area Connectivity Drive Runoff and
710 Carbon Export Across Watershed Scales, *Water Resour. Res.*, 54, 1576-1598, <https://doi.org/10.1002/2017wr021718>, 2018.



Table 1. Relative differences in average dissolved organic carbon (DOC) and nitrate (NO_3^-) concentrations and $\text{DOC}:\text{NO}_3^-$ molar ratios between upstream and midstream, and between midstream and downstream sites at the Font del Regàs stream for the study period (September 2010 to August 2012). Positive values indicate that either solute concentrations or stoichiometric ratios increased in the downstream direction, while negative values indicate a decrease. Differences are shown separately for base flow and storm flow conditions.

Section	Flow conditions	DOC difference	NO_3^- difference	$\text{DOC}:\text{NO}_3^-$ difference
Upstream to midstream	Base flow	+19%	-40%	+98%
Upstream to midstream	Storm flow	+15%	-40%	+95%
Midstream to downstream	Base flow	-10%	+8%	-18%
Midstream to downstream	Storm flow	-13%	+9%	-23%



Table 2. Hydroclimatic descriptors of the five large storm events identified during the study period (September 2010 to August 2012) at the downstream site of the Font del Regàs catchment, including duration, precipitation amount (P), accumulated precipitation seven (P-7) and 30 (P-30) days before the event, average stream flow (Q_{avg}), average groundwater table (Gw_{avg}), groundwater table range (ΔGw), slope of the linear relationship between riparian groundwater table and stream flow (slope), and average stream flow normalized to average groundwater table (Q_{avg}/Gw_{avg}). All these nine descriptors were used as predictors in a partial least square regression analysis, in which average dissolved organic carbon (DOC) concentrations and average nitrate (NO_3^-) concentrations during each event at the downstream site were the response variables (also shown). The relative increase in DOC and NO_3^- concentrations with respect to average concentrations observed during base flow conditions at the downstream site are shown within brackets.

Event	Period	Duration (days)	P (mm)	P-7 (mm)	P-30 (mm)	Q_{avg} (mm)	Gw_{avg} (m)	ΔGw (m)	Slope (m mm ⁻¹)	Q_{avg}/Gw_{avg} (mm m ⁻¹)	DOC (mg C L ⁻¹)	NO_3^- (mg N L ⁻¹)
#1	11-14/03/2011	4	95	11	39	1.7	0.74	0.36	0.16	2.3	2.35 (167%)	0.40 (140%)
#2	14-23/03/2011	10	116	102	133	3.3	0.64	0.55	0.14	5.1	2.16 (146%)	0.33 (97%)
#3	02-12/11/2011	11	134	65	116	0.8	0.87	0.21	0.18	0.9	2.45 (178%)	0.40 (138%)
#4	14-19/11/2011	6	174	14	261	4.2	0.69	0.52	0.06	6.1	1.91 (117%)	0.29 (74%)
#5	21-30/11/2011	10	67	182	426	3.4	0.74	0.21	0.07	4.5	1.36 (55%)	0.34 (102%)




Figure 1. Map of the study catchment, Font del Regàs, including sampling sites and main hydromorphological properties. The location of the Font del Regàs catchment within Spain is shown in the inset.





Figure 2. Distributions of (a, b) dissolved organic carbon (DOC) concentrations, (c, d) nitrate (NO₃⁻) concentrations, and (e, f) DOC:NO₃⁻ molar ratios for three sampling sites along the Font del Regàs stream for the study period (September 2010 to August 2012). Data are shown separately for base flow (left panels) and storm flow (right panels) conditions. On each box, the central mark indicates the median, and the bottom and top edges of the box indicate the 25th and 75th percentiles, respectively. The whiskers extend to the most extreme data points not considered outliers. Arrows indicate statistical increase (upward arrow) or no statistical difference (straight arrow) between base flow and storm flow conditions for each colour-coded site using non-parametric Wilcoxon signed-rank tests. Results from pairwise Wilcoxon tests are also shown, where * indicates statistical difference (p<0.01) and ns indicates non-significant difference.



740 **Figure 3.** Frequency of dissolved organic carbon to nitrate (DOC:NO₃⁻) molar ratios considered optimal for heterotrophic microorganisms at the three sampling sites along the Font del Regàs stream for the study period (September 2010 to August 2012). The corresponding frequencies for which there was either carbon (C) or nitrogen (N) limitation are also displayed in each case. Data are shown separately for base flow and storm flow conditions. The considered range for optimal conditions (4.8 to 11.7) was based on the global reviews of Taylor and Townsend (2010) and Helton et al. (2015).




Figure 4. Relationships between dissolved organic carbon (DOC) concentrations, nitrate (NO_3^-) concentrations, and DOC: NO_3^- molar ratios for three sampling sites along the Font del Regàs stream for the study period (September 2010 to August 2012). Relationships are shown separately for base flow and storm flow conditions. Coefficients of determination (R^2) and slopes are shown for significant linear correlations ($p < 0.01$).




Figure 5. Biplot of the two-component partial least square (PLS) regression model with score and loading vectors re-scaled into the -1 to +1 numerical range in order to display the relative relationships between observations (i.e. five large storm events), predictors (i.e. duration, accumulated precipitation amount (P), accumulated precipitation seven (P-7) and 30 (P-30) days before the event, average stream flow (Q_{avg}), average groundwater table (Gw_{avg}), groundwater table range (ΔGw), slope of the linear relationship between riparian groundwater table and stream flow (slope), and average stream flow normalized to average groundwater table (Q_{avg}/Gw_{avg})), and response variables (average dissolved organic carbon (DOC) concentration and average nitrate (NO_3^-) concentration during each event at the downstream site of the Font del Regàs catchment).




Figure 6. (a) Relationships between daily riparian groundwater tables and daily stream flow at the downstream site of the Font del Regàs catchment for the five large storm events identified in the study period (September 2010 to August 2012). For this relationship, all events displayed clockwise hysteresis loops (denoted by arrows), except event #5, for which the hysteresis loop was anticlockwise. In all cases, linear regressions showed $R^2 > 0.8$ ($p < 0.01$). (b) Corresponding distribution of dissolved organic carbon to nitrate (DOC:NO₃⁻) molar ratios for each of the events. On each box, the central mark indicates the median, and the bottom and top edges of the box indicate the 25th and 75th percentiles, respectively. The whiskers extend to the most extreme data points not considered outliers. Dotted lines delimitate the lower (4.7) and upper (11.7) limits of stoichiometric conditions considered optimal for stream heterotrophic microorganisms.




Figure 7. Diagram conceptualizing the mobilization of dissolved organic carbon (DOC) and nitrate (NO_3^-) from the riparian zone to the stream at the Font del Regàs catchment during large storm events preceded by (a) dry and (b) wet conditions. Under dry antecedent conditions, the riparian groundwater table rises more rapidly, as indicated by the steeper slope of the relationship between groundwater table and stream flow. This pattern suggests relatively lower water lateral transmissivity (represented by blue horizontal arrows) of each riparian soil layer compared to large storm events preceded by wet conditions. As a result, the 'Dominant Source Layer', i.e. the riparian zone depth stratum that contributes the most to water and solute fluxes to the stream (delimited by dashed horizontal lines), will be thicker during large storms preceded by dry conditions because more layers will be hydrologically activated, increasing the chances for relatively more DOC and NO_3^- mobilization. In addition, soluble DOC and NO_3^- pools (represented by green and purple hypothetical lines, respectively) will accumulate in upper riparian soil layers during dry periods, and will leach and deplete during wet periods (especially that of DOC as C is more limited than N in this type of catchments). As a result, during large storms preceded by dry conditions, groundwater will flow through shallower riparian soil layers with larger DOC and NO_3^- soluble pools, which will lead to large increases in stream DOC and NO_3^- concentrations. During large storms preceded by wet conditions, groundwater will not reach the most superficial soil layers and DOC and NO_3^- pools will be smaller, which will result in minor changes in stream DOC concentrations and only moderate increases in stream NO_3^- concentrations.

Appendix A




Figure A1. Time series of precipitation at Font del Regàs catchment, as well as stream flow, dissolved organic carbon (DOC) concentrations, nitrate (NO_3^-) concentrations, and $\text{DOC}:\text{NO}_3^-$ molar ratios at three sites (upstream, midstream, and downstream) along the Font Regàs stream for the period September 2010 to August 2012.




Figure A2. VIP (variable influence on projection) values computed for the nine predictors (duration, precipitation amount (P), accumulated precipitation seven (P-7) and 30 (P-30) days before the large storm event, average stream flow (Q_{avg}), average groundwater table (Gw_{avg}), groundwater table range (ΔGw), slope of the linear relationship between riparian groundwater table and stream flow (slope), and average stream flow normalized to average groundwater table (Q_{avg}/Gw_{avg})) included in the partial least square regression model carried out in this study. Predictors are considered important in the overall model when $VIP > 1$. Important predictors are highlighted with darker colour.




Figure A3. Frequency of dissolved organic carbon to nitrate ($DOC:NO_3^-$) molar ratios considered optimal for heterotrophic microorganisms at the three sampling sites along the Font del Regàs stream for the study period (September 2010 to August 2012) for two considered scenarios: (a) optimal range is 4.8 to 11.7 (presented in the paper and based on the global reviews of Taylor and Townsend (2010) and Helton et al. (2015)) and (b) optimal range is 4.5 to 16 (a less restricted range based on values from other site specific and modelling studies). The corresponding frequencies for which there was either carbon (C) or nitrogen (N) limitation are also displayed in each case. Data are shown separately for base flow and storm flow conditions.