

# High-resolution drought simulations and comparison to soil moisture observations in Germany

Friedrich Boeing<sup>1</sup>, Oldrich Rakovec<sup>1,2</sup>, Rohini Kumar<sup>1</sup>, Luis Samaniego<sup>1</sup>, Martin Schrön<sup>3</sup>, Anke Hildebrandt<sup>1</sup>, Corinna Rebmann<sup>1</sup>, Stephan Thober<sup>1</sup>, Sebastian Müller<sup>1</sup>, Steffen Zacharias<sup>3</sup>, Heye Bogena<sup>4</sup>, Katrin Schneider<sup>5</sup>, Ralf Kiese<sup>5</sup>, Sabine Attinger<sup>1</sup>, and Andreas Marx<sup>1</sup>

<sup>1</sup>Helmholtz Centre for Environmental Research – UFZ, Department Computational Hydrosystems, Permoserstraße 15, 04318 Leipzig, Germany

<sup>2</sup>Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Praha-Suchbát 16500, Czech Republic

<sup>3</sup>Helmholtz Centre for Environmental Research – UFZ, Department Monitoring and Exploration Technologies, Permoserstraße 15, 04318 Leipzig, Germany

<sup>4</sup>Forschungszentrum Jülich GmbH, Agrosphere Institute (IBG-3), Germany

<sup>5</sup>Karlsruhe Institute of Technology, IMK-IFU, Ecosystem Matter Fluxes, Kreuzeckbahnstr. 19, 82467 Garmisch-Partenkirchen, Germany

**Correspondence:** Friedrich Boeing (friedrich.boeing@ufz.de)

**Abstract.** Germany's 2018-2020 consecutive drought events resulted in impacts ~~related-with-several-sectors-such-as~~ in multiple sectors including agriculture, forestry, water management, energy production, and transport. ~~The~~ High-resolution information systems are key to preparedness for such extreme drought events ~~are high-resolution information systems~~. This study evaluates the new setup of the one-kilometre German drought monitor (GDM), which is based on ~~the~~ daily soil moisture (SM) simulations from the mesoscale hydrological model (mHM). The simulated SM is compared against a set of diverse observations from single profile measurements, spatially distributed sensor networks, cosmic-ray neutron stations and lysimeters at 40 sites in Germany. ~~Results~~ Our results show that the agreement of simulated and observed SM dynamics in the upper soil (0-25 cm) is especially high in the vegetative active period (0.84 median correlation R) and lower in winter (0.59 median R). ~~Lower~~ The lower agreement in winter results from methodological uncertainties in ~~simulations-as-well-as-in~~ both simulations and observations. Moderate but significant improvements between the ~~older~~ (coarser 4km resolution) ~~)-setup and the present~~ ( $\approx 1.2$ km resolution) ~~)-GDM-is-found~~ GDM in the agreement to observed SM dynamics is observed in autumn (+0.07 median R) and winter (+0.12 median R). The higher resolution of the second GDM version allows for a ~~much~~ more detailed representation of the spatial variability of SM, which is particularly beneficial for local risk assessments. Furthermore, the results underline that nationwide drought information systems depend both on appropriate simulations of the water cycle and a broad, high-quality observational soil moisture database.

## 1 Introduction

The extreme drought events since 2018 in Germany ~~lead-led~~ to multi-sectoral impacts (~~Madruga-de-Brito-et-al., 2020~~) (Madruga de Brito et al., 2020) and increased stakeholder awareness. Moreover, recent studies emphasized that extreme ~~soil-moisture~~ SM drought events will

be more likely and more severe in Central Europe under future warming scenarios (Samaniego et al., 2018; Grillakis, 2019).

20 The singularity of the 2018/19 drought within observational records in terms of consecutive multiyear water deficits ~~was~~ has been confirmed for Germany and Central Europe (Boergens et al., 2020; Hari et al., 2020; Rakovec et al., 2022). With these prospects comes an increased need for state-of-the-art information on droughts as a basis for ~~precisely-assessing-precise~~ assessment of the uniqueness and potential impacts of drought events from local to continental scales.

In recent years, several national and ~~supranational-international~~ drought monitoring systems ~~were-have been~~ developed. The

25 German Drought Monitor (GDM) was first introduced in 2014 as an information platform for agricultural droughts in Germany under [www.ufz.de/droughtmonitor](http://www.ufz.de/droughtmonitor) and is operationally updated daily (Zink et al., 2016). Core ~~of-to~~ the GDM is simulated ~~soil-moisture-SM~~ using the open-source mesoscale hydrological model (~~mHM, (Samaniego et al., 2010; Kumar et al., 2013)~~ mHM; Samaniego et al., 2010; Kumar et al., 2013). The GDM provides a near real-time status of ~~soil-moisture-SM~~ and drought in Germany with a time lag of one day due to the meteorological data availability. Information on the drought status is

30 provided for the uppermost soil layer (25 cm) and total soil column (~~depth-variable, depending on-varying depth depending on~~ the soil map) by calculating the Soil Moisture Index (~~(SMI) (Samaniego et al., 2013)~~ (SMI; Samaniego et al., 2013) and Plant Available Water (PAW). With ~~its-use-in-national-and-federal-state-agencies,~~ around 2200 media contributions in the year 2020 and more than four ~~Million-million~~ website views since 2018 alongside its use in national and federal state agencies, it proved its important role as a drought information tool in Germany. ~~Feedback-and-requests-showed~~ The feedback and requests received

35 show that the GDM is ~~both~~ used by interested public and practitioners as well as media and politics to obtain up-to-date drought information. A crucial aspect for ~~the~~ optimal use of scientific ~~,-environmental data~~ environmental data, from a practitioners point of view ~~is-the-, is~~ applicability to local purposes. ~~Targeted-Data from targeted~~ stakeholder interviews within the EDgE project (<http://edge.climate.copernicus.eu>) and in Climalert (<http://climalert.eu/>) with a core stakeholder group of 15 farmers in Central Germany supported this need. So far, hydrological models ~~that-were~~ applied at national or ~~supranational-international~~ level in

40 operational drought services were mostly run on ~~"relative"~~ relatively low spatial resolutions, e.g. with grid cell size  $5 \times 5 \text{ km}^2$  in the European drought observatory (EDO) (Sepulcre-Canto et al., 2012) or  $4 \times 4 \text{ km}^2$  in the GDM (Zink et al., 2017). The spatial resolution ~~was-is~~ mainly restricted due to input data availability, such as the soil map BUEK1000 (spatial resolution 1:1,000,000) for Germany.

Recently, an updated version of the nationwide German soil database (BGR, 2020) was published ~~at-with~~ 25 times higher res-

45 olution ~~that-enabled-to-take-a-step-forward-to-enabling~~ hydrological modelling at much higher spatial resolution ( $\approx 1.2 \times 1.2 \text{ km}^2$ , an  $\approx 11$  fold increase to the prior GDM version). Nevertheless, it was not clear how the quality of the ~~soil-moisture-SM~~ simulations would change at higher spatial resolution. In contrast to other environmental variables ~~,-for-instance,-air-temperature,-~~ it-is-much-more-it-is-very challenging to aggregate ~~soil-moisture-SM~~ to a larger scale due to its highly heterogeneous nature and measurement uncertainties (Western et al., 2004; Bogen et al., 2010; Rosenbaum et al., 2012). Simulated ~~soil-moisture~~

50 SM derived from hydrological models is the prime alternative to observed ~~soil-moisture-and-widely-employed-to-estimate~~ soil-moisture-SM and is widely employed for SM estimation on regional to global scales (Keyantash and Dracup, 2002). Nevertheless, simulations also face methodological uncertainties ~~as-well-(Marx-et-al.,-2018,-e.g.-),~~ especially under transient conditions such as those caused by climate change (Marx et al., 2018; O et al., 2020). Cammalleri et al. (2015) investigated the

use of ~~multiple~~ hydrological models for drought monitoring in Europe using SM anomalies and drought classification metrics and found that including ~~several~~ multiple hydrological models improved ~~the~~ overall performance. Furthermore, hydrological models are typically calibrated based on streamflow ~~since it~~, which represents the integral hydrological catchment response. ~~Yet, besides~~ Besides validating the modelled streamflow ~~itself~~, there is a clear need to thoroughly evaluate other water cycle components that are not used for constraining the model parameters emerges. Ideally, such validations require observations of the variable of interest that ~~a)~~ span (a) cover the same spatial scales as the model and extend ~~across~~ over different climate regimes within the study area and ~~b)~~ span (b) extend over long temporal scales, which would allow them to be termed “representative”. Although large-scale meteorology-driven SM variations can display seasonal varying length scales up to 500 km (Koster et al., 2019), small-scale SM variability largely depends on local site characteristics such as soil properties, topography and land use. Therefore, optimal drought monitoring systems over large areas ~~can only be built up using~~ should make use of the best available observation data in combination with a smart simulation system.

Enormous efforts have been and are being made to build up construct environmental observation networks from regional to global scales. Within global environmental monitoring networks such as FLUXNET ~~that~~ which focuses on measuring ecosystem carbon fluxes (Baldocchi et al., 2001), ~~soil moisture is usually~~ SM is sometimes measured in multiple depths at single profiles. However, extensive validations of simulated ~~soil moisture~~ SM from hydrological models are hampered by the limited spatial representativeness of point-scale sensors and hence require novel measurement approaches to bridge the scale gap between local observations and model resolutions. Measurements that capture the spatial structure of ~~soil moisture~~ SM at larger scales are expensive and time-consuming. ~~For this reason they~~, and for this reason are rare and only applicable in comparatively small catchments of a few tens of hectares (Bogena et al., 2010). In Germany, the infrastructure of Terrestrial Environmental Observatories (TERENO) has been established since 2008 to build up a nationwide long-term monitoring network ~~where~~ in which one of the focuses is on hydrological variables (Zacharias et al., 2011; Bogena, 2016). Many of those sites were equipped with spatially distributed measurements of ~~soil moisture~~ SM networks (SDM, Bogena et al., 2010) and Cosmic-Ray Neutron Sensors (CRNS, Zreda et al., 2012; Andreasen et al., 2017; Schrön et al., 2018)). CRNS detectors count neutrons of the natural cosmic-ray background radiation as a proxy for soil water content (Desilets et al., 2010; Köhli et al., 2021). The integral measurement footprint covers areas of 300–600 m diameter and depths of 15–70 cm, both increasing for dryer conditions (Köhli et al., 2015; Schrön et al., 2017). The CRNS method has emerged as a reliable technique to continuously monitor root-zone ~~soil moisture~~ SM at the field scale (Bogena et al., 2015; Andreasen et al., 2017) and has been used recently for the validation of land surface and hydrological models (Han et al., 2016; Iwema et al., 2017; Dimitrova-Petrova et al., 2020).

~~On the other hand, satellite-based soil moisture~~ Satellite-based SM data benefits from spatial coverage at the ~~km scale~~. ~~Still, kilometre scale, but~~ the shallow penetration of the signal in the upper few centimeters of the soil is a significant constraint. While those signals also depend on the surface condition, vegetation density and microwave frequencies, these products themselves require ground-based ~~soil moisture~~ SM observations for validation (Peng et al., 2021). ~~To evaluate the performance of the drought monitor model, the~~ The time series of SDM and CRNS observations at the TERENO sites appear to be better suited for evaluation of the drought monitor model in terms of long-term continuity and root-zone representation. In particular, the data covers recent wet (e.g. 2017) and dry (e.g. 2015, ~~2018–2020~~ 2018–2020) years, including extreme drought conditions.

Here, we evaluate ~~soil moisture simulations~~ SM simulations from mHM at the one and four ~~kilometer-scale simulated from~~ mHM-kilometre scale simulated against an unprecedented compilation of ~~soil moisture~~ SM observations from 40 locations across Germany. A wide range of climatic conditions and vegetation types is covered. Specifically, the study aims ~~at two objectives to answer two questions~~. Firstly, how well do the high-resolution German-wide ~~soil moisture~~ SM simulations capture the ~~SM-dynamics in observed soil moisture that constitute the basis for the near real-time soil moisture drought monitoring system~~ SM? Emphasis is given ~~to distinguish between different soil moisture~~ on the comparison of different SM measurement techniques due to their relevance for interpreting the evaluation results. Secondly, can ~~the simulations with higher spatial resolved~~ SM simulations at higher spatial modelling resolution including refined spatial-resolution soil input data be provided with a consistent quality ~~of soil moisture estimates~~? Higher resolution does not ~~automatically increase~~ necessarily improve the model performance ~~but, and~~ may even worsen the quality of the simulation results. To assess this ~~effect~~, the low-resolution model setup GDM-v1-2016 as well as the one kilometer setup GDM-v2-2021 are compared against multi-method ~~soil moisture~~ SM observations. Furthermore, drought characteristics estimated with both model setups are compared using annual drought intensities over the last 69 years (~~1952-2020~~ 1952-2020).

## 2 Methods and Datasets

### 2.1 The mesoscale ~~hydrological model~~ Hydrological Model (mHM)

The mesoscale ~~hydrological model~~ Hydrological Model is a grid-based spatially distributed ~~hydrologic model forced with~~ hydrological model driven by daily precipitation, temperature and potential evapotranspiration PET. It accounts for major ~~hydrologic hydrological~~ processes such as snow generation and snowmelt, canopy interception, soil infiltration, ET, deep percolation, baseflow generation, and surface runoff routing. The open-source model code ~~is available in git repository~~ is available and is under active development and maintenance (<https://git.ufz.de/mhm/mhm>). The model uses three distinct levels to organize the modelling procedures: level 0 (L0) for input data of the sub-grid physical basin characteristics, level 1 (L1) for the realization of the integrated hydrological processes and level 2 (L2) for specification of meteorological forcing inputs. An unique component of mHM is the Multiscale Parameter Regionalization (MPR) technique (Samaniego et al., 2010) that allows inferring spatial variability of the required model parameters seamlessly on various modelling scales. One of the distinguishing aspects of the MPR approach compared to other regionalization techniques is to deliver a quasi scale-invariant model performance across modelling scales and improve the transferability of model parameters to ungauged basins (Kumar et al., 2013; Rakovec et al., 2016; Samaniego et al., 2017). The ~~mHM-model~~ was applied and evaluated in ~~different climatological regions e.g. multiple climatological regions including~~ Europe (Thober et al., 2015; Rakovec et al., 2016), West Africa (Dembélé et al., 2020), India (Saha et al., 2021) and ~~US the conterminous United States~~ (Livneh et al., 2015; Rakovec et al., 2019). Within the MPR technique, the subgrid physical basin characteristics at L0 are linked to model parameters through transfer functions and a set of global parameters and subsequently upscaled to generate effective parameters at L1. The aggregation ~~has based~~ is based on a set of upscaling rules (e.g. arithmetic or harmonic mean) following ~~e.g.~~ flux conservation schemes (Samaniego et al., 2010).



A general overview on the model processes and parameterization can be obtained from Samaniego et al. (2010) and Kumar et al. (2013). ~~In the following only the soil moisture~~ Only the SM component of mHM is described here, due to its relevance for this study. The incoming precipitation and snowmelt are partitioned into root-zone ~~soil moisture~~ SM and runoff components, depending on the degree of soil saturation, using a power function similar to the well-known HBV model (Samaniego et al., 2010). The degree of non-linearity depends on the underlying vegetation and soil characteristics following the MPR framework (Samaniego et al., 2010; Kumar et al., 2013). The evapotranspiration from soil layers is estimated as a fraction of the potential evapotranspiration depending on the ~~soil moisture~~ SM stress and the fraction of vegetation roots present in each layer (Samaniego et al., 2010). The moisture stress function depends on the specification of soil-water content at a permanent wilting point, critical and saturation levels, which are determined using a set of pedo-transfer functions estimated within the MPR framework (Livneh et al., 2015; Zacharias and Wessolek, 2007).

## 2.2 Model setups at $4 \times 4 \text{ km}^2$ and $1.2 \times 1.2 \text{ km}^2$ spatial resolution

The new setup GDM-v2-2021, as used in the GDM version 2, includes several changes ~~compared to the older to the previous~~ model setup GDM-v1-2016. The main features of the two mHM ~~model~~ setups that are used in the analysis are described in Table 1. While the GDM-v1-2016 uses ~~the mHM model~~ mHM version 5.6, in GDM-v2-2021 ~~the mHM model~~ mHM was updated to version 5.10 (see <https://github.com/mhm-ufz>). The implemented changes in ~~the mHM model~~ mHM did not change the hydrological process representations related to ~~soil moisture~~ SM that were used in the simulations here. Between the setups GDM-v1-2016 and GDM-v2-2021, the projection system was changed from the projected coordinate system Gauss-Krueger 4 (EPSG:31468) to the World Geodetic coordinate ~~System~~ system (EPSG:4326). While the ~~lengths~~ size of the grid cells in the GDM-v1-2016 setup ~~were fixed on~~ was fixed at  $4 \times 4 \text{ km}^2$  (L1 level), the grid cell ~~sizes~~ size in the GDM-v2-2021 setup ~~are~~ is measured in degree, ~~where~~, As such, the grid cell ~~length~~ size varies with latitude ~~with a decreasing grid cell length~~, with grid cell width in east/west direction decreasing from 1.23 km at  $45.47.25^\circ$  N latitude (south of Germany) ~~decreasing~~ to 0.98 km at  $55.5^\circ$  N latitude (north of Germany) and a constant grid cell length of 1.7 km in north/south direction.


**Table 1.** Main features of the model setups GDM-v1-2016 and GDM-v2-2021. Core ~~of~~to the setups is the mesoscale hydrological model mHM. Vertical discretization of soil layers in the hydrological model mHM and projection system are denoted. In the spatial model resolution the Level 0 (L0) describes the subgrid variability of relevant basins characteristics. Level 1 (L1) and Level 2 (L2) describe the dominant hydrological processes and meteorological forcings, respectively. ~~Underlying datasets~~Datasets used as model inputs for soil as well as land use and geology ~~used as model inputs for mHM~~ on L0 model resolution are stated.

Setup	spatial model resolution	soil dataset	vertical soil discretization	projection	<del>land use</del> <u>land use</u> dataset	<u>hydro geology</u> dataset
GDM-v2-2021	L1 and L2: 0.01562° × 0.01562° eq. ~1.2 × 1.2 km <sup>2</sup> L0: 0.001953125° × 0.001953125°	BUEK200	4 layers: 0–5 cm 5–25 cm 25–60 cm <del>60–variable</del> <u>60–variable</u> cm	Latlon (EPSG:4326)	GLOBCOVER	GLIM
GDM-v1-2016	L1 and L2: 4 × 4 km <sup>2</sup> L0: 100 × 100 m <sup>2</sup>	BUEK1000	3 layers: 0–5 cm 5–25 cm <del>25–variable</del> <u>25–variable</u> cm	Gauss Krüger-4 (EPSG:31468)	CORINE	HUEK2

Soil ~~physical properties on~~ texture (sand and clay fraction) and mineral bulk density are derived from national digital soil maps provided by the BGR (Federal Institute for Geosciences and Natural Resources). The BUEK200 dataset (BGR, 2020) used in the GDM-v2-2021 setup substantially increased the mapping resolution compared to the BUEK1000 dataset (BGR, 1998) used in the GDM-v1-2016 setup (scale 1:1 000 000 to 1:200 000). At the time of the creation of this study, the database version of BUEK200 was ~~V0~~v0.5. Figure 1 shows the depth averaged clay contents ~~exemplary for a~~for an exemplary region in Central Germany where ~~soil moisture SM~~ observations that were used in the analysis are located. The soil ~~maps aim to~~ represent the local soil composition by identifying patterns of similar soils ~~map used for the study (BUEK), is the standardized basic soil mapping for Germany. It shows the distribution and association of soils and their properties in Germany. The map content is classified according to soil regions and soil landscapes. For each map unit a soil series is given composed of an index soil (dominant soil) and accompanying soils. The base mapping unit of the soil maps are soil series, which can include multiple soil profiles that associate soils with similar compositions. For the derivation of the BUEK200 dataset the index soils, meaning the dominant soil in a soil series, were considered. For modeling, the soil properties of the index soil within the spatial mapping unit were used to derive the model parameters.~~

The soil depths in mHM are discretized into an upper soil layer ~~for at depth~~at depth 0–25 cm, including a top layer at depth 0–5 cm, and the remaining depth of the soil profile. In the GDM-v2-2021 setup an additional layer ~~for at~~at 25–60 cm was added due to stakeholder feedback, mainly from the agricultural sector. The ~~rooting tillage~~ depth is set to 30 cm in both model setups.

~~Landuse~~The land use datasets used in the model setups GDM-v1-2016 and GDM-v2-2021 ~~are were~~are CORINE (EEA, 2009) and GLOBCOVER (ESA, 2009), respectively. Hydrogeological input data that define the aquifer properties and govern the baseflow recession rates, was derived from the HUEK200 (~~BGR, 2009~~) and GLIM (~~Hartmann, Jörg and Moosdorf, Nils, 2012~~) databases, ~~respectively~~database for GDM-v1-2016 (BGR, 2009) and the GLIM database for GDM-v2-2021 (Hartmann, Jörg and Moosdorf, Nils, 2012). Digital elevation models were derived from BKG (2010) and USGS (2017) ~~, respectively. The changes in landuse and geology datasets can influence the simulations, yet play a minor role for the soil moisture simulations compared to the change in the soil dataset, because changes of landuse data are in subgrid scale (resolution GLOBCOVER 300m, CORINE <100m) and no direct feedback of from saturated “groundwater” storage to soil moisture storage is implemented in mHM. respectively.~~



**Figure 1.** Average ~~over soil column for the~~ derived ~~soil physical properties~~ clay [%] ~~of over the soil column from the~~ BUEK1000 soil dataset used in the GDM- v1-2016 model setup (left panel) versus ~~the~~ BUEK200 soil dataset used in the GDM-v2-2021 model setup (right panel). The grid shows the respective modelling resolution L1 at which the hydrological processes are simulated (see table 1). Both setups are projected in WGS 84 (EPSG:4326).

### 2.2.1 Meteorological input data

Precipitation, as well as minimum, maximum and average air temperature, are interpolated on a daily timescale based on meteorological station data from the German Weather Service ~~DWD~~ (DWD) using external drift kriging (EDK) with elevation as the drift variable. The meteorological station data is subject to extensive quality controls (Kaspar et al., 2013). Additionally, quality controls ~~such as checking the plausible variable range~~ are implemented in the preprocessing steps of the interpolation routine ~~e.g. checking plausible variable range~~. Theoretical variograms are estimated based on all available station data to derive seamless fields of hydro-meteorological fluxes and states for entire Germany (Zink et al., 2017). An exponential model is used for precipitation and spherical models for the temperature variables. The interpolation method and variogram parameter estimation for Germany are described and evaluated in detail in Zink et al. (2017) including cross-validation metrics and comparison to the comparable REGNIE gridded dataset by the German Weather Service (Rauthe et al., 2013). ~~Potential Evaporation (PET)~~ ~~→ PET~~ is calculated using the Hargreaves-Samani Method (Hargreaves and Samani, 1985) that is based on the interpolated temperature fields (average, minimum, and maximum) ~~, where and~~ (potential) extraterrestrial radiation, ~~which~~ is computed depending on the latitude of the location and day of the year.


### 2.2.2 Multibasin ~~Model-Calibrations~~model calibrations

The unknown parameters of the mHM ~~model~~-setup GDM-v2-2021 were calibrated against observed discharge using the Kling-Gupta efficiency (KGE; Gupta et al., 2009) as the objective function. The parameter optimization was conducted using the Dynamically Dimensioned Search (DDS; Tolson and Shoemaker, 2007) algorithm (~~with~~ 1 000 iterations), which underwent  
185 detailed scrutiny, as follows. In a first step, 200 parameter sets were obtained using a multi-basin/domain-wide joint basin calibration strategy, in which a subset of ~~6-six~~ basins was randomly selected (out of 201 total ~~number-of~~ basins) and then jointly calibrated during a common period of ~~1990-2005~~ (~~See~~ 1990-2005 (see Table S1 in the Supplements)). ~~In the second step~~Subsequently, all 200 parameter sets were evaluated against the full ensemble of ~~the~~ 201 basins during an extended period of 1986–2005 (with a warming period of 5 years). The ~~single-unique~~ parameter set with the best performance in terms of the  
190 median daily KGE over 201 basins was selected and used for the consequent analysis (See Table S2 in the Supplements). This updated approach is based on the earlier calibrations of the GDM-v1-2016 setup (Zink et al., 2017) in which the Nash–Sutcliffe efficiency instead of the KGE was applied and individual single-basin instead of the multi-basin calibrations were carried out as input to the model cross-evaluation at locations that were not used for model calibration. Previous works also focused on multibasin calibrations of mHM in other regions, such as Mizukami et al. (2017); Rakovec et al. (2019). The model performance  
195 of the best cross-evaluated parameters of the GDM-v2-2021 based on daily streamflow from 201 catchments in Germany yielded a median performance of 0.761 KGE (see Fig. A1).

### 2.3 Soil ~~Moisture-Observations~~moisture observations

The ~~soil-moisture~~ SM observations used to conduct the model evaluations were gathered from the environmental observation networks TERENO (Zacharias et al., 2011) and FLUXNET (~~FLUXNET2015 Dataset (Pastorello et al., 2020))~~ (FLUXNET2015 Dataset; P  
200 as well as from the Cunnernsdorf site operated by the DWD(~~German Weather Service~~). In total, ~~soil-moisture~~ SM data from 40 locations were compiled and processed for the analysis (see Fig. 1). ~~Even though~~ Although it is not feasible to establish an evenly distributed grid of SM measurements on a national level (Vereecken et al., 2008), the available locations cover a wide range of climatic and vegetation conditions in Germany.

In total, we analysed 46 measurements from 24 grassland sites, nine crop sites, six forest sites and one site containing a forest clearing. Four of the sites have multiple measurement methods available(~~total number (n)=46~~), ~~which allows~~, which allowed  
205 comparing the evaluations between the measurement methods at single sites. ~~In total, we analyzed 24 grassland, 9 crop, 6 forest and one site with a forest clearing.~~ The elevation ranges from 4 to ~~1252-1 252~~ meters a.m.s.l., and the long-term yearly precipitation sums range from below 500 mm to more than ~~1500-1 500~~ mm. Time series lengths of the observations are between 2.8 and 17.8 years with a median (mean) of 6.5 (6.7) years. A detailed overview of the location characteristics is shown in Table  
210 3. ~~The data comprises four different soil moisture measurement methods explained in more detail in the following.~~



**Figure 2.** Soil-moisture-SM observations of 40 locations distributed over Germany were used in the soil-moisture-SM evaluations of the GDM-v2-2021 and GDM-v1-2016 model setups. The subplots zoom-to-display the different experimental sites in Germany greater detail, representing different climate gradients in Germany. The maps show the digital elevation model on the hydrological subgrid variability resolution L0 of the mHM model in the GDM-v2-2021 setup ( $0.001953125^\circ \times 0.001953125^\circ$ ). The grid corresponds to the modelling resolution L1 in the GDM-v2-2021 setup ( $0.01562^\circ \times 0.01562^\circ$  eq., which equals  $\approx 1.2 \text{ km} \times 1.2 \text{ km}^2$ ) at which the hydrological processes are simulated. Lower-The lower panel shows the distribution of different soil-moisture-SM observations depending on land use land use type, elevation and average yearly precipitation. Note that some of the 40 locations have multiple soil-moisture-SM data sources ( $n_{\text{total}} = 46$ ).



~~Soil-moisture~~The data is comprised of four different SM measurement methods. SM observations from seven FLUXNET and 16 TERENO sites in Germany based on several vertically distributed sensors within one soil profile were used (in the following abbreviated SPM). The sensor depths are described in Table 3. SPM sites used from TERENO-Northeast Observatory are further described in Itzerott et al. (2018a) and Itzerott et al. (2018b). ~~Soil-Moisture~~SM data from lysimeters are available for four sites from the ~~lysimeter-network-TERENO-SOILCan~~TERENO-SOILCan lysimeter network (Pütz et al., 2016) at the Bad Lauchstädt experimental site and the TERENO ~~Pre-alpine~~Pre-Alpine Observatory (PAO) (Kiese et al., 2018). Lysimeters are large vessels containing an undisturbed soil column to allow gravimetric measurements. Since the lysimeter vessels are closed at the bottom, water tension is adjusted to reference measurements at the same depth in the undisturbed soil close to the lysimeter (Pütz et al., 2016; Kiese et al., 2018). In the lysimeters, ~~soil-moisture~~SM is measured by single sensors in multiple depths. At each SOILCan-site multiple lysimeters are organized in hexagons (Pütz et al., 2016). The number of lysimeters per site are described in Table 3.

For three of the 40 sites (Am Grossen Bruch, Hohes Holz and Wüstebach), spatially distributed measurements (SPM) of ~~soil-moisture~~SM are available. Multiple sensors are installed in a spatial grid ~~and-at~~ different depths covering an area of some hundreds of ~~m<sup>2</sup>square metres~~. For the locations Hohes Holz ~~39-profiles~~ and Am Grossen Bruch ~~39 and 20 profiles-with-sensors in-profile with sensors at~~ multiple depths were used ~~respectively~~ (depths varied slightly between profiles depending on soil property changes). Therefore, they are not denoted explicitly). For the Wüstebach site, 51 profiles with two sensors each at 5 ~~cm~~, 20 ~~cm~~, and 50 cm depth were used (Wiekenkamp et al., 2019). The Wüstebach ~~soil-moisture~~SM measurement network is described in detail in Bogena et al. (2018).

~~Soil-moisture~~SM observations derived from Cosmic Ray Neutron Sensing (CRNS) stations were used from 17 sites (see Table 3) of the TERENO observatories (Bogena et al., 2022). The soil albedo component of cosmic-ray neutrons is particularly prone to changes of ~~soil-moisture~~SM (Desilets et al., 2010; Köhli et al., 2021). However, since neutrons are sensitive to all pools of hydrogen, the measured neutron signal is also affected by biomass (Baatz et al., 2015), intercepted water (Bogena et al., 2013; Schrön et al., 2017), and snow (Schattan et al., 2017) and therefore requires a correction of the measured signal in this respect. In this study, periods of snow cover have been excluded from the CRNS data. ~~Soil-moisture~~SM from CRNS data has been calculated by standard methods (Desilets et al., 2010; Zreda et al., 2012) and aggregated to daily time steps. This leads to typical statistical uncertainties of less than 3 vol.% (Schrön et al., 2018).

All SM data was checked according to their flagging conventions for doubtful or low-quality values. In some cases, doubtful data was removed manually after personal communication from site maintainers (e.g. ~~for some sites from~~ the TERENO PAO lysimeter sites ~~where some sites experienced~~showed doubtful data after frost in early 2017). The available SM data in the respective depths, as noted in Table 3, was aggregated to weighted vertical averages according to the soil discretization depths in mHM (0–25 cm, 25–60 cm and 0–60 cm). Highest weights were allocated when the sensor depth is located in the center of the soil depth range and weights linearly decrease towards the edges of the soil depth range. The spatial mean values were calculated for the SDM measurements based on the available sensors.

## 245 2.4 Soil moisture data preparation and evaluation metrics

Since the computation of ~~soil-moisture-drought-indice~~~~se~~~~-g~~~~-the~~ SMI (Samaniego et al., 2013) SM drought indices, including the estimation of ~~soil-moisture~~ SM probability distributions by kernel density estimates, is hampered for the available observed data due to the limited length of observed ~~soil-moisture~~ SM data (<10 years for most locations), the analysis here is based on ~~the~~ a comparison of observed to simulated ~~soil-moisture~~ SM (e.g., Samaniego et al., 2013). It is widely known that absolute ~~soil~~ moisture SM values cannot be adequately determined by a regional model (~~e.g.~~ partly due to the spatial heterogeneity), yet the hydrological model typically captures the temporal dynamics well (Koster et al., 2009). As drought is defined by the deviation from normal conditions, ~~soil-moisture-anomalies-are~~ SM anomalies were calculated. To preserve the units of volumetric ~~soil~~ moisture SM [mm/mm] and the original range of ~~soil-moisture~~ SM dynamics, standardization by dividing standard deviation ~~is~~ was not undertaken in this study. The anomalies are calculated in two ways. ~~Firstly~~ First, the mean of all values in each SM  
255 time series is subtracted.

$$\theta(anom)_{i,k} = \theta_{i,k} - \bar{\theta} \quad (1)$$

Secondly, the multi-year mean for each day of the year is subtracted to deseasonalize the anomalies. The removal of the annual average cycle of SM is necessary for the subsequent drought classification based on percentile thresholds as described in the next section.

$$260 \quad \theta(deseas - anom)_{i,k} = \theta_{i,k} - \bar{\theta}_i \quad (2)$$

~~Where~~ where  $i$  is the calendar day of the year (DOY 1, ..., 365) and  $k$  ~~each-year~~available in a data record~~is the year~~. To reduce uncertainty in the ~~calculation-of-mean-values~~ mean resulting from heterogeneous and small sample sizes, for each ~~DOY~~ a ~~a~~ i a moving window with 15 days on each side of the day was used to increase ~~sample-size~~ b ~~the sample size and~~ the multiyear mean of each ~~DOY~~ i was calculated based on the mean of randomly drawing 500 bootstrap samples. ~~The simulated~~ data were ~~Since there are some data gaps in the observed data (see Table 3 for data availability), the simulated data was~~ masked  
265 to the available observed data to allow a comparable calculation of SM seasonality. Leap days were removed before calculating the deseasonalized anomalies.

The evaluation of observed against simulated SM is based on the Spearman rank correlation coefficient (R). The Spearman rank correlation coefficient is a non-parametric measure to quantify the strength of ~~a~~ the monotonic relationship between two  
270 variables. The correlations are calculated on whole data records as well as on sub-periods (months, seasons and vegetative active period) to investigate the ~~intra-annual~~ seasonal variability in the performance metrics. Paired Wilcoxon signed-rank tests were conducted to identify significant changes between the model setups.

### 2.4.1 ~~Soil Moisture Index Computation~~ moisture index computation and ~~Analysis~~ analysis

Simulated ~~soil-moisture~~ SM by the two model setups is used to compute a Soil Moisture Index (SMI) ~~on the years 1952–2020~~  
275 following Samaniego et al. (2013) and Zink et al. (2016) ~~that enables~~ enabling a SM drought analysis based on ~~the~~ long term

SM ~~dataset~~data. The SMI for a given cell and day is estimated as

$$SMI_t = \hat{F}_T(x_t) \quad (3)$$

and it represents the quantile at the ~~soil moisture~~ SM fraction value  $x$  (normalized against the respective saturated soil water content).

280  $x_t$  denotes the simulated monthly ~~soil moisture~~ SM fraction at a time  $t$  and  $\hat{F}_T$  is the empirical distribution function estimated using non-parametric kernel density estimates. The optimal bandwidths are estimated by minimizing a cross-validation error estimate. Details regarding the computation of the SMI can be found in Samaniego et al. (2013).

The SMI drought threshold concept used in the German Drought Monitor is based on the D0-D4 classification system for droughts from the US-Drought monitor (Svoboda et al., 2002) that related drought categories to potential impact types. The drought thresholds reflect the occurrence of similar SM conditions in the past and hence indicate the potential impacts of these conditions (Zink et al., 2016). A cell at time  $t$  is under drought when  ~~$SMI_t < \tau$~~   $SMI_t < \tau$ . Here,  $\tau$  denotes that the soil water content in ~~this a~~ a cell is less than the values occurring  $\tau \times 100\%$  of the time. ~~In this study,  $\tau$  is set to 0.2 indicating a soil moisture state occurring less than 20% of the time. This threshold follows common values for drought classifications that align with typical occurrences of drought impacts, (Svoboda et al., 2002). Drought~~ The 20th percentile used as  $\tau$  in this study is defined as moderate drought conditions, that indicate conditions of “possible damages to crops and pastures”. Extreme drought conditions are defined as the 5th percentile indicating “high probability of major losses in crops and pastures”. The resulting impact of SM drought conditions need to be identified for each specific impact type based on the timing within the year and duration of drought conditions. For example, Peichl et al. (2018, 2021) identified specific monthly damage functions between the SMI and different crops using varying statistical methods. The work showed that dry SM anomalies in some months can

290 reduce yield (e.g., August, September for maize), while in other months it may increase crop yield (e.g., May for maize). Impacts of SM droughts can affect a broad range of sectors besides agriculture. Especially, the considered soil depth of the SMI is relevant for different sectors. While the drought conditions in the upper soil (0-25 cm and 0-60 cm) are more relevant to agriculture, drought in the total soil column (up to 2 meters) indicate potential impacts on water resources and the forestry sector.

300 SMI based drought statistics are calculated for the years 1952–2020 on fixed temporal (annual and vegetative active period from April to October) and spatial scales (per grid cell and aggregated for Germany).

When calculating the cumulative density functions of SM, a common statistical basis of 1951-2015 was used for both model setups. The drought intensities (DI) per year are calculated by

$$DI = \frac{1}{dA} \frac{1}{d * A} \sum_{t_0}^{t_1} \int_A [\tau - SMI_i(t)]_+ \quad (4)$$

305 with the area of interest  $A$  (here Germany) and duration  $d$  ( $t_1 - t_0$ ) in days (annual  $t_0$  Jan 1st to  $t_1$  Dec 31st and vegetative active period  $t_0$  Apr 1st to  $t_1$  Oct 31st ). The drought intensities take into account the degree of negative departure from drought conditions (hence, the extremer the drought conditions, the higher the intensities) as well the temporal aggregation length and

~~the spatial aggregation area. The~~ area under drought is calculated as the percentage of grid cells ~~with where~~ SMI < 0.2 ~~of the total German surface area (total number of grid cells)~~ averaged over the respective temporal periods. ~~For calculating the~~  
310 ~~cumulative density functions of SM, a common statistical basis of 1951–2015 was used.~~


### 3 Results and Discussion

In the following sections, the comparisons of the multi-method ~~soil moisture~~ SM observations with two hydrological model simulations are presented and discussed to investigate the proposed research objectives. In section 3.1, a comparison of ~~soil moisture observations from different measurement methods to the~~ SM observations to the simulations from the high-resolution  
315 operational model setup GDM-v2-2021 is shown. The setup allows a comparison of observations to the 0–25 cm layer as well as to the additional, deeper soil layer of 25–60 cm. In section 3.2, the differences between the two simulation setups are shown for annual drought intensities during 1952–2020 and compared to ~~soil moisture~~ SM observations. The two mHM simulations are used in their operational setups, meaning that only data and information available for ~~whole Germany have been the whole~~  
320 ~~of Germany was~~ used. Additional available information on soils or meteorological measurements at the observation sites ~~has not been was not~~ incorporated in the simulations.

#### 3.1 Comparison of high resolution simulations against observed SM dynamics


Here,  $1.2 \times 1.2 \text{ km}^2$  simulations in two soil layers (GDM-v2-2021) are compared to SM observations using four different measurement methods: Cosmic Ray Neutron Sensing (CRNS), spatially distributed measurements (SDM), single profile measurements (SPM) and lysimeter (LYSI). ~~In this study,~~ SM anomalies as well as deseasonalized SM anomalies are used.  
325 Figure 3 shows the results for three selected locations that contain both CRNS and SDM measurements for the six-year period 2014–2019. In general, the SM anomalies and deseasonalized data agree well, with ~~small deterioration of correlations in a small reduction of correlations for the~~ deseasonalized data. Furthermore, observations and simulations ~~compare well agree well both~~ in the uppermost ~~25 soil layer (0–25 cm soil layer cm~~ and in the deeper layer ~~down to 60 (25–60 cm depth)~~. The correlation strength between simulations and observations from different measurement techniques is similar for the sites Am  
330 Grossen Bruch and Hohes Holz, but ~~deviate more from each other deviates more~~ for the Wüstebach site. It is worth noting that different spatial scales are mapped by those measurements. While the SPM (not included here) represents point information, the SDM and CRNS cover an area less than  $0.1 \text{ km}^2$  and the mHM simulations cover an area of  $\approx 1.44 \text{ km}^2$ . In general, the day-to-day variability is ~~smaller lower~~ in simulations than in observations. At the forest sites Wüstebach and Hohes Holz, the day-to-day variability in the CRNS data is higher than in SDM. Several environmental factors other than SM ~~can~~ influence  
the CRNS signals (see ~~method section~~ Methods). While the changing biomass might have a low impact on the signal, it can introduce a (constant) systematic bias. Since only anomalies are analysed here, the impact of such bias on this (comparative) anomaly analysis should be minimal. Intercepted water on leaves and in the litter layer can be particularly challenging to quantify, especially in forested stations such as Hohes Holz or Wüstebach (Bogena et al., 2013; Schrön et al., 2017). It might lead to stronger dynamics in the CRNS signal during and shortly after rain events, ~~compared in comparison~~ to the model

340 output or other observation methods. Additionally, ~~at the Wüstebach site~~ partial deforestation in 2013 at the Wüstebach site  
modified SM flows ~~that resulted~~ resulting in stronger response to rainfall (Wickenkamp et al., 2019). Nevertheless, there is no  
general tendency for lower correlations at forest sites ~~compared to than~~ crop and grassland sites (see Figure 4). Crop sites show  
slightly lower correlations than grassland sites, which is expected since anthropogenic activities (e.g., crop rotation) are not ~~rep-~~  
~~resented~~ represented in mHM. Correlations display no clear tendency across the range of elevation and precipitation regimes.  
345 In general, Figure 4 reveals that the model ~~performances~~ performance does not systematically depend on site conditions. ~~As~~  
~~well, no trend of deteriorating correlations with length of~~ Moreover, no systematic relationship between correlations and the  
length of the time series can be ~~detected~~ found (see Figure 4 d)).



**Figure 3.** Soil moisture (SM) time series for the years 2014–2019 for the selected locations Am Grossen Bruch, Hohes Holz and Wüstebach, showing SDM and CRNS data against simulated data from mHM in 0–25 cm and 25–60 cm depth in the GDM-v2-2021 setup. The Hordorf site also contains both CRNS and SDM measurements, but with much shorter time series length. For visualization the stations with longer time series were selected for visualization. Correlation Coefficients Spearman rank correlation coefficients are denoted at the left side of each time series. Upper-panel Panel (a) shows SM anomalies, including seasonality (subtract mean of the whole time series) and lower panels show deseasonalized anomalies-panel (subtracting multiyear mean for each calendar day) analyzed in this study shows deseasonalized SM anomalies.






**Figure 4.** Spearman correlation coefficients of the simulated versus observed de-seasonalized soil moisture anomalies in dependence of its against site characteristics: (a) landuse/land use, (b) elevation and (c) average yearly precipitation and (d) length of the time series. See Table 3 for detailed overview per location. Colors denote the soil moisture-SM data method (Cosmic Ray Neutron Sensing (CRNS), spatially distributed measurements (SDM), single profile measurements (SPM) and lysimeter (LYSI)) and shapes the landuse/land use types reported at the locations (abbreviated as following: grass = grassland, clear= forest clearing, LYSI=Lysimeter).

Monthly Spearman correlation coefficients for all locations and measurement methods ( $n_{\text{total}}=46$ ) in at 0–25 cm depth are presented-shown in Fig. 5. The correlation coefficients show an apparent clear intra-annual-seasonal variation, with the highest values in summer/autumn months, while-and the lowest values are found in-wintermonths in winter. The highest median correlation is detected in August (0.87), while the lowest correlations-are median correlation is found in January (0.37). The spread of the correlation coefficients within the different locations is largest in winter months, with some locations having-very high-have correlations close to 1.0. Yet, While in February and March, some locations show-correlations-below-zero-for-the measurement-methods-with CRNS, SPM and LYSI measurements show correlations below zero. The intra-annual variation of performance metrics was similar to the findings of Xia et al. (2014), that extensively evaluated simulated SM from four


different hydrological models (Noah, Mosaic, SAC, VIC) in the North American Land Data Assimilation System phase 2 (NLDAS-2) dataset, which is used for drought monitoring in the United States, ~~with~~ and similarly observed generally higher correlations in summer and lower correlations in winter. ~~Lower correlations in winter can~~ The lower correlations observed in winter could be related to higher uncertainties in simulations ~~as well as~~ and observations with respect to frozen soils and snow cover. The sensor quality of SDM, SPM and LYSI in winter can be reduced during frost days. Especially, SPM and LYSI measurements can be affected by sensor failures as they rely only on few sensors compared to the spatially distributed measurements (SDM) with a larger number of sensors. Annex Figure A3 shows correlations between simulated SM, CRNS and SDM. Low correlations between simulation and observations are accompanied by low correlations between the measurement methods, especially in winter. In a climate impact study investigating low flows over Europe, it could be shown that ~~the overall~~ uncertainty due to the selection of the hydrological model dominates the overall uncertainty including the meteorological drivers in snow dominated areas (Marx et al., 2018). Furthermore, the mHM ~~model~~ does not contain a full energy balance model, which limits the description of soil frost depths.



**Figure 5.** Comparison of the simulated mHM soil moisture SM (0–25 cm) in the GDM-v2-2021 setup to observed SM anomalies without (a) and with (b) removing subtraction of the mean seasonal SM cycle for each month depicted with boxplots. Values (x shapes) of SM anomalies are plotted for each location colored by and coloured according to the SM measurement method (CRNS: Cosmic Ray Neutron Sensing; SDM: spatially distributed measurements; SPM: single profile measurements; LYSI: Lysimeter) are shown method used. Sample Note that sample sizes between measurement methods differ (CRNS: n=1617, SDM: n=3, SPM: n=23; LYSI: n=4). Data points that are not fulfilling conditions of significant correlations both significantly (p-value < 0.05) and positive correlations positively correlated are marked with x. See Figure A2 for additional plots separating the locations in subsets for detailed comparisons.

Observations using different SM measurement methods display considerably different correlations. The SPM generally scatter vary more, with large variation in winter and outliers with low performance the presence of low-performance outliers in summer months. CRNS measurements show a consistently high performance in summer months but notably low correlations in winter (especially January). As snow days were removed from the time series in the CRNS measurements, the anomaly calculation from the remaining data was impacted by a smaller and incomplete sample size. Another reason for the observed lower correlations might be related lower correlations observed might be due to the variable penetration depth of CRNS ranging, which ranges between 15 to 70 cm, depending on soil moisture depending on SM (Köhli et al., 2015; Schrön et al., 2017). This could introduce systematic and temporally variable errors and deteriorate affect the correlation between observed and simulated soil water content (Baroni et al., 2018). The comparison to Comparison to the mHM top soil (0–25 cm)

layer is ~~still-assumed-assumed to remain~~ a good compromise, since the soil water distribution is rather homogeneous between 0 and 25 cm under wet conditions, ~~while under~~. Under dry conditions the footprint is deeper and more heterogenous, but the highest sensitivity is ~~still~~ in the upper soil layers (exponential sensitivity). The SDM measurements show the most consistent performance across all months ~~,-except for May. This is with the exception of May, as~~ illustrated in Figure A2 a), ~~which displays only the locations Am Grossen Bruch and Hohes Holz, where both SDM and CRNS measurements are available.~~ All measurement methods at these sites show ~~in May and June~~ a drop in correlations to the SM simulations in May and June, while the observations have higher correlations between each other (see also Figure A3). ~~That~~ This points to deficiencies in the model, which may be related to the static vegetation module in mHM which does not include processes such as possible early onset of the growing season and ~~related-consequent~~ earlier depletion of the soil water storage. Moreover, lower correlations of deseasonalized anomalies in May are detected especially at forest locations (median of 0.63 over all forest locations). The timing of leaf unfolding in trees, usually between late April to May (Chen et al., 2018), is subject to annual fluctuations and affects evaporation from the soil and therefore SM dynamics. Figure A2 b) depicts the SPM data from FLUXNET and TERENO separately, which cover ~~different time periods~~, the time periods 1997–2014 and 2011–2019, respectively. The performance at TERENO sites is generally higher compared to the FLUXNET sites. ~~The intra-annual seasonal~~ variation of the correlations is in good agreement among both monitoring networks and time periods. ~~Higher correlations to SM simulations at the TERENO SPM sites may appear~~ The performance at TERENO sites is generally higher than at the FLUXNET sites, possibly due to a larger number of sensors installed along the soil depth in the TERENO sites which may improve the vertical averaging of SM (see Table 3).



**Figure 6.** Spearman correlation coefficients of the simulated mHM ~~soil moisture-SM~~ in the GDM-v2-2021 against observed deseasonalized ~~soil moisture-SM~~ anomalies depicted with boxplots for three depths (0–25 cm, 0–60 cm, 25–60 cm). Values of each location are ~~shown to enhance comparability between measurement methods as their sample sizes differ.~~ Colours ~~plotted and colours~~ denote the ~~sourcee~~ measurement method of ~~soil moisture-the SM~~ data (Methods: SDM: spatially distributed measurements; SPM: single profile measurements; LYSI: Lysimeter). Here only locations (~~n=26~~) ~~are taken into account~~ with measurements ~~in-at~~ 25–60 cm depth ~~are taken into account~~ (SDM: n=3, SPM: n=19; LYSI: n=4). Data points ~~that are~~ not ~~fulfilling conditions of significant correlations both significantly~~ (p-value < 0.05) and ~~positive correlations positively correlated~~ are marked with x.

395 The Spearman correlation coefficients for each season and soil depth (~~0–25 cm, 25–60 cm and average over 0–60 cm~~) is depicted in Figure 6. Note ~~that here only locations were considered that have SM data in all depths were considered~~ and CRNS data was excluded as its varying penetration depth does not allow a consistent depth-wise evaluation, ~~which~~. This leads to a smaller sample sizes of locations (n=26). Figure 6 shows that the median correlation is lower for the deeper ~~soil moisture simulations when comparing the 0–25 cm to 25–60 cm depth SM simulations~~ for all seasons, except for winter ~~during which the median R is higher in the lower 25–60 cm depth ( $\Delta + 0.06$ )~~. In spring, the lower depth 25–60 cm ~~also shows the~~ strongest negative difference to the upper depth ~~0–25 cm~~ in comparison to summer and autumn (Spring  $\Delta - 0.19$ , Summer  $\Delta - 0.12$ , Fall  $\Delta - 0.02$ ). The correlations vary more in the 25–60 cm depth between locations in all seasons, with more outliers ~~with of~~ very low correlations ~~observed~~. Since the mHM was conceptualized for dominant processes at the large scale (mesoscale), not all processes that are important at the local scale are currently accounted for (e.g. ~~variable rooting~~ ~~depths species specific root water uptake~~, lateral flow or groundwater soil water interaction). For instance, Rosenbaum et al. (2012) showed for the distributed SM measurements at the Wüstebach catchment that SM dynamics in the topsoil (5–50 cm depth) are influenced by groundwater. Processes of capillary rise are not modelled in mHM, hence it is expected that agreement to simulated SM by mHM at sites with groundwater-influence is lower compared to groundwater-distant sites. This effect should increase with depth due to increasing groundwater influence that could explain ~~the~~ lower correlations in the 25–60 cm

410 depth. ~~Also, SPM sites might be affected more than SDM since these effects could~~ more affected by this than SDM in which  
these effects can be averaged out ~~for SDM~~. Identification of groundwater characteristics at each measurement site was, however,  
out of scope for this study. ~~For the average across 0–60 cm the correlations tend to be similar to 0–25 cm. Only in spring,~~  
~~correlations over the 0–60 cm soil column are lower compared to 0–25 cm due to the low correlations observed in 25–60 cm~~  
~~during this period. In Winter the average over 0–60 cm depth shows higher correlation than both 0–25 cm ( $\Delta + 0.11$ ) and~~  
415 ~~25–60 cm ( $\Delta + 0.02$ ) alone.~~ The SDM overperform SPM and LYSI, with higher than average correlation values especially for  
the 25–60 cm depth, which underlines the assumption that local characteristics of single sensors e.g. groundwater influence and  
the resulting spatial variability of ~~soil moisture~~ SM (Famiglietti et al., 2008) are averaged out over a larger area and generally  
supports the closer scale match of SDM measurements and the  $1.2 \times 1.2$  km<sup>2</sup> simulation grid cells. It has to be noted that the  
SPM at the same sites as the SDM also show comparable high correlation values (for an overview of the locations, see Table  
420 3).

### 3.2 Comparison of different mHM model setups

In the following, the comparison between observations and the two model setups GDM-v1-2016 and GDM-v2-2021 (i.e.,  
GDM version 1 and 2), as well as drought metrics between the two simulation setups are shown and discussed. Table 2  
shows the median values of the Spearman correlation coefficients for ~~different sub-periods within the year (season~~ selected  
425 sub-periods (seasons, vegetative active period April–October) and for the full year. Considering ~~all available~~ the observed SM  
data from all locations and measurement methods, the median correlations between the two simulation setups slightly increase  
by +0.05 in GDM-v2-2021. The results on an ~~intra-annual~~ seasonal scale show a small decrease in the correlations in spring  
( $\Delta - 0.03$ ) and summer ( $\Delta - 0.01$ ), but a significant increase of correlations in fall ( $\Delta + 0.07$ ) and winter ( $\Delta + 0.12$ ) in the  
new model setup. Several of the changes in the model setup may provide explanations for the improved model agreement to  
430 observed SM dynamics in fall and winter. The higher modelling resolution of the 1 km runs may better resolve the sub-grid  
variability of cold season related processes such as snow accumulation that improves the simulated SM dynamics. As well,  
the finer spatial soil texture representation possibly contribute to an improved model representation of soil wetting/drying e.g.  
especially during saturated conditions in the cold season. When analysing the metric over the vegetative and non-vegetative  
active period (defined from April–October and November–March, respectively), the increase in median correlations is +0.03  
435 and +0.10 ~~(significant), respectively. Correlations using CRNS (n=16) and SPM (n=23) respectively.~~ Median correlations using  
CRNS and SPM measurements support the overall findings. In general, the results show that the CRNS yield higher median  
correlation than the SPM measurements for both model setups, except for spring. While the median correlation in winter  
increased by +0.17 between GDM-v1-2016 to GDM-v2-2021 for CRNS, there is only a small increase in correlation of +0.03  
for SPM. Similar results showing an overall increase in simulation performance were found by Albergel et al. (2012). In their  
440 study, the EMCWF operational and re-analysis SM product using the hydrological model H-TESSEL was improved due to  
changes in the soil hydrology in the model and an increase of model resolution. They concluded that a better representation  
of soil texture might obtain further improvements. Furthermore, De Lannoy et al. (2014) found moderate improvements in the  
agreement of SM simulations compared to observations through implementing updated soil texture information.




**Table 2.** Median ~~spearman~~Spearman rank correlation coefficients  $R$  of ~~both~~ simulated (GDM-v1-2016, GDM-v2-2021) versus observed deseasonalized ~~soil-moisture~~SM anomalies ~~in-at~~ depth 0–25 cm. ~~Spearman~~The correlation coefficients  $R$  are calculated ~~annual~~annually, ~~seasonal~~seasonally (spring=~~mar~~March, April, May; summer=~~jja~~June, July, August; fall=~~son~~September, October, November; winter=~~djf~~December, January, February) and ~~over~~ the vegetative ~~and non-vegetative~~ active period (defined from April–October and non-veg November–March ~~respectively~~). Note that some of the 40 locations have multiple ~~soil-moisture~~SM data sources (~~n-available, resulting in n = 46~~); see Table 3. Stars denote significant differences (p-value < 0.05) in correlations between the model setups according to paired Wilcoxon signed-rank test.

metric	method	setup	annual	spring	summer	fall	winter	non-veg	veg
R [-]	ALL (n=46)	GDM-v2-2021	0.78	0.65	0.85	0.84	0.49	0.59	0.84
		GDM-v1-2016	0.73	0.68	0.86	0.77	0.37	0.49	0.81
		$\Delta$	+0.05	-0.03	-0.01	+0.07 (*)	+0.12 (*)	+0.10 (*)	+0.03
	CRNS (n= <del>16</del> <u>17</u> )	GDM-v2-2021	0.81	0.63	0.88	0.86	0.60	0.65	0.86
		GDM-v1-2016	0.79	0.67	0.88	0.80	0.46	0.48	0.84
		$\Delta$	+0.02	-0.04	0.0	+0.06 (*)	+0.14 (*)	+0.17 (*)	+0.02
	SPM (n=23)	GDM-v2-2021	0.72	0.67	0.79	0.78	0.39	0.45	0.82
		GDM-v1-2016	0.71	0.69	0.80	0.76	0.34	0.42	0.77
		$\Delta$	+0.01	-0.02	-0.01	+0.02	+0.05	+0.03	+0.05


Spearman rank correlations between simulated and observed deseasonalized SM anomalies that fall below the 20th percentile in the observed SM time series are shown in Figure 7 to specifically analyse the dry anomaly spectrum. It is important to emphasize that we do not aim to estimate drought periods here, as its solid calculation requires a much longer time series. The drought estimation is performed using histograms for every grid cell and day of the year (see method section 2.4.1). Consequently, estimating robust percentiles requires time series lengths of minimum 30 years – this means that the time series length of the observational data is considered insufficient. Figure 7 a) shows a median correlation of 0.61 over all observations in the GDM-v2-2021 setup. The performance in the two model setups remains similar. However, the comparison separated between the measurements with larger spatial footprint (SDM, CRNS) and point scale measurements (SPM, LYSI) shows that the agreement between simulations and the larger footprint observations increased towards the high resolution setup, but the median agreement to the point scale SM measurement decreased. In general, the measurements with larger spatial footprint display higher agreement to the simulations. Due to the varying day-to-day variability of SM between the SM observation types and the simulations, in Figure 7 b) additionally a statistical smoothing was applied by calculating a running 30 day mean on the daily SM time series before subtraction of the seasonal SM cycle. This approach is similar to the SM preprocessing for the SMI as proposed in Zink et al. (2016). Figure 7 b) shows when smoothing is applied, the agreement between observations and simulations during dry periods can be substantially improved to a median correlation of 0.7 over all observations in the

GDM-v2-2021 setup ( $\approx 1$ -km resolution). Especially the agreement between the point scale measurements and simulations is increased to a median correlation of 0.63 in both model setups.



**Figure 7.** Correlations of deseasonalized daily SM below the 20th percentile (based on the observed SM time series) between simulations and observations (a). In (b) additionally a statistical smoothing was applied by calculating a running 30 day mean on the daily SM time series before subtraction of the seasonal cycle. The correlations are shown for all observations (n=46) and separated between observations with larger spatial footprint (n=20) including Cosmic Ray Neutron Sensing (CRNS) and spatially distributed measurements (SDM) as well as point measurements (n=26) including single profile measurements (SPM) and lysimeters (LYSI).

Next, we contrast the drought characteristics based on the two model setups to assess the differences in drought ranking and the spatial structure of drought events. Annual drought intensities aggregated over Germany based on the daily SMI using simulated SM from 1952–2020 are presented in Fig. 8 and grid-based for the last decade in Fig. 9 to compare the model setups in terms of drought classification. Fig. 8 shows only marginal differences between the model setups. They, which are slightly more prominent in the top soil compared to the total soil column. The model setups largely agree on the three years with the most intensive droughts. The ranking in the top soil during the vegetative active period shows an exception: it differs slightly due to the similar drought intensities in the years 1959, 1976 and 2003 in the top soil. 2003. The drought years are more pronounced with respect to drought intensities in the GDM-v2-2021 setup in the top soil, but in contrast, the average drought area is estimated larger in the GDM-v1-2016 setup in those years. Generally, the classification of drought years aggregated over Germany results in similar estimates using the different operational drought monitor setups.




**Figure 8.** Soil moisture SM drought intensities spatially aggregated for over Germany in during the vegetative active period (April - October) in the top soil (5–25 cm) and total soil column (up to 2 m). The size of the circles shows represents the average area under drought. The three largest drought events are denoted with numbers numbered in each panel. Colours represent the two model setup GDM-v1-2016 and GDM-v2-2021. Colours are mixed when circles from both model setups overlay.


To assess regional differences in drought characteristics between the model setups, Fig-Figure 9 shows the drought intensity maps in the vegetative active period for 2011–2020. Drought intensities are more spatially diverse in the GDM-v2-2021 setup that stem stemming from the higher granularity of the GDM-v2-2021 setup with higher resolved including higher-resolution soil information and less smoothed-smooth patterns than the GDM-v1-2016. Nevertheless, the general patterns are similar between the two setups, but regionally. Regionally large differences can be seen (e.g., the drought intensities in the Swabian and Franconian Jura region are more pronounced concerning the neighbouring areas in the GDM-v1-2016 setup – see years 2017 and 2019 for the total soil). Also Additionally, the differences in drought intensities are more pronounced in the total soil column in the last decade, which can be explained by multi-annual, cumulative effects. The current total soil drought lasts in many regions for at least three years. In Figure 10 the variance between grid cells for drought intensities during vegetation the vegetative active period are shown as semi-variograms. In general, the spatial variance is larger in the total soil than top soil. The GDM-v2-2021 setup shows a general larger spatial variance between grid cells in the top soil and larger increase with distance (see Figure 10 (a)). The spatial variance in the total soil is lower at smaller distances in the GDM-v1-2016 setup, but slightly higher at larger distances. Figure 10 (b) showing semi-variance normalized by distance (and log-sealed x-axis to

~~to improve visibility of smaller distances)~~ demonstrates that in the GDM-v2-2021 setup the distance-normalized variance of  
485 drought intensities is increased especially at small spatial scale in both the top and total soil, indicating larger local differences  
in response to drought intensities. These findings are in line with Livneh et al. (2015) who investigated the influence of different  
soil databases on resulting hydrologic fluxes. They reported that the higher variability of soil properties in the finer soil database  
generally resulted in simulations with more ~~considerable variability of variability in~~ (extreme) hydrologic responses.

We would like to highlight that in our study several changes between the operational model setups were implemented  
490 besides changing the underlying soil dataset as described in section 2.2. The changes such as the land use and geology datasets  
influence the hydrological simulations, yet they play a minor role for the SM simulations compared to the change in the soil  
dataset. The SM simulations are not influenced by the geological dataset, because no direct feedback from the saturated aquifer  
to the SM reservoir is implemented in mHM. To demonstrate the different role of the change in SM dynamics related to the  
specific soil and land use datasets temporal correlations between SM from separated model runs fixing all model settings  
495 (L1  $\approx 1.2 \times 1.2 \text{ km}^2$  resolution, default mHM parameters) only changing the soil dataset (BUEK200 – BUEK1000) and in a  
separate step only changing the land use dataset (CORINE – GLOBCOVER) are shown in Figure A4. The change of the soil  
dataset has a much larger impact on the SM simulations compared to the change of the land use dataset in these specific model  
setups. The CORINE and GLOBCOVER land use datasets both have already high horizontal resolutions ( $\approx 100 \text{ m}$  and  $300 \text{ m}$ ,  
respectively). The differences between the land use datasets mostly lie in the subgrid scale of the mHM hydrological modelling  
500 resolution and have a minor effect on the upscaled hydrological response at the L1 level (here  $\approx 1.2 \times 1.2 \text{ km}^2$ ).



**Figure 9.** Soil moisture-SM drought intensities (DI) per grid cell for (a) upper soil (5–25 cm) and (b) total soil column (up to 2 m) during the vegetative active period (April -October) in the last decade (2011–2020) for the model setups GDM-v1-2016 and GDM-v2-2021 and for the absolute differences between the setups (GDM-v1-2016 - GDM-v2-2021). The GDM-v1-2016 data was remapped to the GDM-v2-2021 grid to calculate for the differences difference calculation. Graphs including all of years from 1952 on can be found at <https://www.ufz.de/index.php?de=47252>.



**Figure 10.** Empirical semi-variograms for drought intensities during ~~vegetation-the vegetative~~ active period in upper ~~Soil-soil~~ for ~~the~~ GDM-v1-2016 and ~~the~~ GDM-v2-2021 setup. The bin size was set to 5 km (~~that corresponds to the~~ nearest larger even ~~km-kilometre~~ bin size relative to the GDM-v2-2016 modelling resolution). The length scale and nugget of the fitted exponential theoretical semi-variograms are noted in the legend. ~~In-~~Subplot (b) ~~shows~~ the ~~y-semivariance normalized by distance~~ and ~~the~~ x axis ~~are-is~~ log scaled.



**Table 3.** Overview of ~~soil-moisture-SM~~ measurement sites. Method denotes the different data sources: Cosmic Ray Neutron Sensing (CRNS), spatially distributed measurements (SDM), single profile measurements (SPM) and lysimeter (LYSI)(~~see method section for detailed description~~). The group ~~Network~~ denotes the ~~experimental-site-environmental observation~~ network ~~name~~ (for TERENO: GC =Central Germany; Rur/E = Rur/Eifel; NE = Northeast, PAO = Pre Alpine Observatory) and land use ~~shows-describes the~~ site characteristics (grass= grass-land, crop = cropland, DBF = Deciduous Broadleaved Forest, ENF = Evergreen Needled Forest, clear=clearing). For ~~Fluxnet~~FLUXNET, the original site name is ~~denoted-included in bracketsparentheses~~. Sensor depths and numbers are denoted. R Spearman correlation coefficients of ~~comparing-simulated versus observed~~ ~~de-seasonalized-soil-moisture~~ ~~deseasonalized SM~~ anomalies in the GDM-v2-2021 setup shown are based on the whole period ~~for-at~~ 0–25 cm and 25–60 cm depth.

network	site	method	land use	begin	end	availability	data $n$	elevation	precipitation	sensor $n$	sensor depth		R	R
											0–25 cm	25–60 cm	0–25 cm	25–60 cm
TERENO CG	BadLauchstädt	LYSI	crop	2016-01-01	2018-12-31	100 %	1091	118 m	498 mm	3	10	30, 50	0.73	0.71
	Ermsleben	SPM	grass	2012-01-25	2019-12-31	81 %	2343	167 m	541 mm	1	10, 20	30, 40, 50, 60	0.80	0.68
	Am Grossen Bruch	CRNS	grass	2014-06-24	2019-11-28	95 %	1892	81 m	545 mm				0.79	-
		SDM		2014-07-30	2019-11-18	84 %	1618			20	var.	var.	0.82	0.84
		SPM		2014-02-07	2019-12-31	98 %	2114			1	10, 20	30, 40, 50	0.86	0.85
	Hecklingen	SPM	grass	2013-07-05	2019-12-31	94 %	2223	93 m	525 mm	1	10, 20	30, 40, 50	0.72	0.71
	HohesHolz	CRNS	DBF	2014-08-27	2019-11-28	91 %	1745	203 m	645 mm				0.80	-
		SDM		2012-07-20	2019-12-30	96 %	2613			39	var.	var.	0.88	0.86
		SPM		2013-04-25	2019-12-31	97 %	2358			2	10, 20	30, 40, 50	0.87	0.88
	Hordorf	CRNS	crop	2016-09-29	2019-11-28	88 %	1022	80 m	554 mm				0.83	-
		SPM		2015-11-06	2019-12-31	98 %	1493			1	10, 20	30, 40, 50	0.82	0.74
TERENO Rur/E	Aachen	CRNS	crop	2012-01-13	2019-05-01	91 %	2437	216 m	875 mm				0.67	-
	Gevenich	CRNS	crop	2011-07-06	2019-01-04	91 %	2496	104 m	766 mm				0.84	-
	Heinsberg	CRNS	grass	2011-09-08	2019-05-01	94 %	2628	61 m	712 mm				0.83	-
	Kall	CRNS	grass	2011-09-14	2019-05-01	78 %	2185	492 m	861 mm				0.82	-
	Kleinau	CRNS	grass	2015-08-25	2019-04-26	87 %	1169	355 m	937 mm				0.88	-
	Merzenhausen	CRNS	crop	2011-05-18	2019-04-03	90 %	2597	91 m	767 mm				0.85	-
	Rollesbr1	CRNS	grass	2011-05-18	2018-12-31	87 %	2409	516 m	1183 mm				0.77	-
	Rollesbr2	CRNS	grass	2012-06-30	2018-12-25	86 %	2045	516 m	1183 mm				0.82	-
	Ruraue	CRNS	grass	2011-11-08	2019-01-01	90 %	2340	102 m	734 mm				0.77	-
	Schoeneseiffen	CRNS	grass	2015-08-13	2019-04-25	83 %	1120	567 m	1119 mm				0.82	-
	Selhausen	CRNS	crop	2015-03-06	2019-04-26	95 %	1441	102 m	726 mm				0.77	-
	Wildenrath	CRNS	clear	2012-05-11	2019-03-23	91 %	2273	79 m	776 mm				0.80	-
	Wüstenbach	CRNS	ENF	2011-03-12	2018-10-05	79 %	2173	614 m	1165 mm				0.44	-
		SDM		2009-01-27	2019-12-31	100 %	3989			150	10, 20 (2x)	50	0.75	0.74
TERENO NE	AltTellin	SPM	grass	2014-05-10	2019-12-30	90 %	1859	9 m	551 mm	1	10, 20	30, 40, 50	0.87	0.26
	Bentzin	SPM	grass	2013-08-19	2019-12-30	98 %	2281	5 m	568 mm	1	10, 20	30, 40, 50, 60	0.53	0.77
	Droennewitz	SPM	grass	2014-04-12	2019-12-30	73 %	1519	33 m	598 mm	1	10, 20	30, 40, 50, 60	0.59	0.66
	Goermin	SPM	grass	2013-08-19	2019-12-30	95 %	2207	7 m	569 mm	1	10, 20	30, 40, 50, 60	0.73	0.47
	Leppin	SPM	grass	2013-01-28	2019-12-30	96 %	2420	6 m	563 mm	1	10, 20	30, 40, 50, 60	0.71	0.53
	Medrow	SPM	grass	2015-07-13	2019-12-30	100 %	1627	5 m	595 mm	1	10, 20	30, 40, 50, 60	0.78	0.76
	Muehlenkamp	SPM	grass	2012-01-01	2019-12-30	82 %	2388	8 m	574 mm	1	10, 20	30, 40, 50, 60	0.71	0.37
	Ueckeritz	SPM	grass	2013-01-28	2019-12-30	93 %	2349	4 m	562 mm	1	10, 20	30, 40, 50, 60	0.63	0.61
	Wotenick	SPM	grass	2014-04-30	2019-12-30	95 %	1964	11 m	588 mm	1	10, 20	30, 40, 50, 60	0.63	0.51
	Zarnckla	SPM	grass	2013-01-23	2019-12-30	95 %	2404	6 m	590 mm	1	10, 20	30, 40, 50, 60	0.83	0.77
TERENO PAO	Fendt	LYSI	grass	2017-01-01	2019-12-31	100 %	1090	634 m	1059 mm	18	10	30, 50	0.80	0.7
	Graswang	LYSI	grass	2017-03-17	2019-12-31	93 %	948	916 m	1570 mm	6	10	30, 50	0.77	0.66
	Rottenbuch	LYSI	grass	2017-03-17	2019-12-31	93 %	948	765 m	1265 mm	12	10	30, 50	0.54	0.53
FLUXNET	Gebesee (DE-Geb)	SPM	crop	2001-01-16	2014-12-31	91 %	4657	156 m	522 mm	1	8, 16	32	0.38	0.33
	Grillenburg (DE-Gri)	SPM	grass	2006-11-21	2014-12-31	98 %	2891	394 m	856 mm	1	10	-	0.68	-
	Hainich (DE-Hai)	SPM	DBF	2002-12-27	2012-12-31	98 %	3570	420 m	774 mm	1	8, 16	32	0.72	0.57
	Klingenberg (DE-Kli)	SPM	crop	2004-11-27	2014-12-31	87 %	3208	478 m	860 mm	1	10	-	0.57	-
	Lackenberg (DE-Lkb)	SPM	ENF	2009-05-01	2013-12-31	90 %	1533	1252 m	1573 mm	1	4	-	0.44	-
	Leinefelde (DE-Lnf)	SPM	DBF	2002-05-01	2012-12-31	72 %	2796	453 m	784 mm	1	8, 16	32	0.83	0.77
	Tharandt (DE-Tha)	SPM	ENF	1997-03-06	2014-12-31	95 %	6189	369 m	791 mm	1	10	-	0.57	-
DWD	Cunnersdorf	CRNS	crop	2016-06-23	2019-12-31	95 %	1217	131 m	634 mm				0.83	0.68

#### 4 Summary and ~~Conclusions~~conclusions

This study evaluates soil moisture (SM) dynamics from two mHM ~~model simulations from operational GDM setups~~ simulations used as operational model setups in the German Drought Monitor (GDM). The increase of hydrological modelling resolution from  $4 \times 4 \text{ km}^2$  in the GDM-v1-2016 setup to  $\approx 1.2 \times 1.2 \text{ km}^2$  in the GDM-v2-2021 setup was possible due to the implementation of a higher-resolution input soil data (BUEK1000 to BUEK200). The comparisons between observed and simulated SM are conducted using various ground-based SM observations (~~n-total = 46~~) with multiple measurement methods and different climate gradients. The agreement between simulated and observed SM dynamics ~~agreement~~ is especially high in the vegetative active period (median R 0.84 in GDM-v2-2021) and lower in winter (median R 0.59 in GDM-v2-2021). ~~The improvements in capturing observed SM dynamics between the model setups GDM-v1-2016 to GDM-v2-2021 showed that an increased input soil data resolution (BUEK1000 to BUEK200) in combination with increased modelling resolution ( $4 \times 4 \text{ km}^2$  to  $\approx 1.2 \times 1.2 \text{ km}^2$ ) km resolution GDM~~ not only produces simulated SM in similar quality as the lower resolution model setup ~~but~~, but partly enhances the model ~~accuracy to simulate~~ ability to simulate observed SM dynamics. We identified significant improvements between the first and second GDM versions in terms of agreement to observed SM, with enhanced correlations during fall (+0.07 median) and winter (+0.12 median). However, the overall improvements were relatively small, partly because the lower resolution model setup ( $4 \times 4 \text{ km}$  grid cells) was already capturing the observed SM dynamics well. Annual drought statistics and ranking based on drought intensities and average area under drought computed on the time frame 1952–2020 were robust between the model setups, with only minor differences on the scale of Germany. The spatial structures in the ~~higher-resolved~~ higher-resolution GDM-v2-2021 setup, including an updated soil map, display larger granularity and spatially more diverse responses to drought, allowing a more refined representation of spatial SM heterogeneity. The higher spatial resolution ~~that is~~ achieved is of great relevance, especially concerning local risk assessments.

~~We focused the model comparisons on analyzing SM dynamics for their relevance for SM droughts, which are defined as a negative deviation from normal SM conditions. A way forward to better absolute estimations of simulated SM (that was not investigated here), which could improve the model internal flux partitioning, can be the integration of observed SM data in the model calibration itself. This kind of approach has been successfully demonstrated using, e.g. CRNS data in the Rur catchment in Germany (Baatz et al., 2017) or remotely sensed SM in the Danube catchment (Wanders et al., 2014). An extended model validation of the SM component of mHM forced with onsite precipitation, local soil maps with soil physical property information on even higher resolution (e.g. BUEK25 or BUEK50) would help to further understand current limitations of mHM to model SM dynamics and separate the analyses from the limited data availability at the scale of Germany.~~


~~The results prominently~~ The results underline the importance of long-term measurement series for developing and optimising ~~such data products~~ data products such as the GDM. Good coverage of relevant environmental gradients with suitable measurement networks is essential ~~, especially because of~~ due to rapidly changing environmental conditions. The direct comparison of the different measurement methods for recording ~~soil moisture~~ SM showed the importance of measurement methods such as CRNS or SDM, which allow better estimates of mean ~~soil moisture conditions to be determined for~~ SM conditions across larger areas. ~~Also~~ However, the temporal ~~availability concerning record lengths~~ and spatial availability ~~, e.g. national coverage~~

535 ~~of spatially representative SDM and CRNS measurements~~ still limits the studies, such as the one presented here, in terms of statistical robustness. Furthermore, we did not ~~analyze~~ analyse deeper soil depths (> 60 cm), ~~because as~~ most measurement sites do not have SM data ~~in these at those~~ depths. Continuous improvements of the SM observational database will be beneficial for future hydrological model evaluations. For future studies, a solution to the variable penetration depth of CRNS could be to compare observed and simulated neutron counts directly by using the COSMIC forward model (Shuttleworth et al., 2013).  
540 ~~It, which~~ is designed to account for irregular ~~soil moisture~~ SM profiles in all modelled depth layers. While COSMIC has been already implemented in mHM, its proper parameterization would require dedicated research and is ~~out outside~~ of the scope of this study, ~~where we focus on first-order comparisons to conventional soil moisture products~~. Regarding the SDM measurements, a source of uncertainty ~~lies in calculating the spatially averaged value~~ remains in the calculation the spatial average. The mean calculation is ~~critical due to a~~ challenging due to the varying number of available sensors in the measurement grids over  
545 time. A robust mean calculation with advanced sensor weighting is currently ~~the subject of a~~ subject of active research.


~~Several questions arise~~ We compared the model simulations in terms of SM dynamics for their relevance for SM droughts, which are defined as a negative deviation from normal SM conditions. The integration of observed SM data in the model calibration itself could improve the absolute estimations of simulated SM and the model internal flux partitioning. This approach has been successfully demonstrated using CRNS data in the Rur catchment in Germany (Batz et al., 2017) and  
550 remotely sensed SM in the Danube catchment (Wanders et al., 2014). An extended model validation of the SM component of mHM forced with onsite precipitation and local soil maps with soil physical property information on even higher resolution (e.g. BUEK25 or BUEK50) would help to further understand current limitations of mHM in modelling SM dynamics and separate the analyses from the limited data availability at the scale of Germany.

Several other aspects are relevant to further improve SM states' simulations with mHM on a national scale in Germany  
555 (or larger towards a continental scale). ~~What are the limitations of the operational German-wide model setup concerning model input data? What are the current limitations of process representation? Which hydrological processes become relevant at smaller scales?~~ A decisive input that influences hydrological model performance is precipitation (Mo et al., 2012). Model performance of mHM was related to rain gauge density on a European scale by Rakovec et al. (2016). While Germany has a very dense meteorological station network, local precipitation can still differ significantly from the interpolated products. ~~Even~~  
560 ~~though~~ Although, Samaniego et al. (2013) showed that the interpolation results on daily precipitation data here compared to the high resolution German Weather Service reanalysis product REGNIE (Rauthe et al., 2013) only differ marginally, the difference of local precipitation from interpolated values is expected to have a large influence on SM dynamics. Thus, improvements in the interpolated precipitation may result in increased model performance. ~~Besides, we may achieve~~ Additionally, a more precise estimation of potential evapotranspiration may be achieved by implementing the Penman-Monteith methods.


565 Finally, we conclude that the resolution of  $\approx 1.2 \times 1.2 \text{ km}^2$  is currently the best compromise between the need for increased model resolution (user perspective) and the current data availability and process representation in mHM (scientific perspective). We emphasise the need for continuous dialogues between stakeholders and the scientific community to improve the underlying model system ~~as well as~~ alongside the provision of user-tailored drought information.




**Figure A1.** Results of mHM multibasin model calibration based on streamflow data from 201 catchments. Left: spatial map of KGE for each basin. Right: KGE Cumulative density function of setup 200 parameter sets, generated by random sampling of the basins. Bold red marks the selected parameter set.



**Figure A2.** Spearman correlation coefficients of simulated soil moisture by mHM in the GDM-v2-2021 setup versus observed ~~deseasonalized~~ ~~de-seasonalized~~ soil moisture anomalies for each month as a supplement to ~~fig~~Fig. 5 by (a) comparing ~~the~~ locations Hohes Holz and Am Grossen Bruch equipped with CRNS, ~~spatially-distributed-SDM~~ and ~~single-profile-measurements-of-SPM~~ soil moisture ~~measurements~~ and (b) comparing FLUXNET (n=7) and TEREÑO (n=20) SPM data.



**Figure A3.** Monthly ~~spearman~~ Spearman rank correlation coefficients for ~~years~~ 2014 - 2019 ~~on~~ between deseasonalized SM anomalies simulated by mHM ~~against~~ and SM observations from CRNS and SDM measurements and between the ~~SM~~ CRNS and SDM observations for the three locations Am Grossen Bruch, Hohes Holz and Wüstebach ~~that have both CRNS and SDM measurements available~~.



**Figure A4.** Correlations between simulated daily SM in the upper soil (5–25 cm) and the total soil column (up to 2 m) using model runs for the time period 1991–2019 keeping the model settings identical (L1 1.2km<sup>2</sup> resolution, default mHM parameters) except changing soil dataset BUEK200 versus BUEK1000 (left) and secondly changing landcover dataset CORINE versus GLOBCOVER (right).



*Code and data availability.* Observational and simulated SM data (<https://www.doi.org/10.48758/ufz.12541>) as well as simulated SMI drought characteristics (DOI: <https://www.doi.org/10.48758/ufz.12534>) that were used in the study are available in UFZ Data Investigation Portal. Open-source mHM code is available at <https://github.com/mhm-ufz> and SMI code <https://doi.org/10.5281/zenodo.5842486>. TERENO and FLUXNET soil moisture data can be obtained at <https://ddp.tereno.net/ddp/> and <https://fluxnet.org/>, respectively.

575 *Author contributions.* F.B., L.S., A.H., C.R., M.S., A.M., R.Kumar developed the concept for the manuscript; F.B. conducted simulations, SM data compilation, data analyses, first manuscript drafts; O.R., F.B. L.S., R.Kumar, A.M. contributed to develop the GDM-v2-2021 setup; O.R. calibrated the GDM-v2-2021 setup; R.Kumar., M.S., O.R., A.M., A.H., L.S., S.T. helped to improve the analyses and manuscript; S.M. and S.T. supported mHM model development and technical maintenance of the GDM; M.S., C.R., R.Kiese, K.S., H.B. provided SM observation data and helped with interpreting the data/improve the manuscript; S.Z. assisted answering questions related to soil processes and  
580 improve the manuscript.

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* This work is partly funded by Helmholtz-Climate-Initiative (HI-CAM) in the Helmholtz Associations Initiative and Networking Fund. The authors are responsible for the content of this publication. We acknowledge the TERENO community and FLUXNET community for providing soil moisture observational data. Special acknowledgment goes to Hans-Jörg Vogel (UFZ), René Zahl (UFZ), Christian Hohmann (GFZ), Ingo Heinrich (GFZ) and Falk Böttcher (DWD) for providing soil moisture observations. We kindly acknowledge the  
585 German Weather Service (DWD), the European Environmental Agency (EEA), the Federal Institute for Geosciences and Natural Resources (BGR), the Federal Agency for Cartography and Geodesy (BKG), the European Space Agency (ESA), the U.S. Geological Survey (USGS), the Global Runoff Data Centre (GRDC) as data providers. The scientific results have (in part) been computed at the High-Performance Computing (HPC) Cluster EVE, a joint effort of both the Helmholtz Centre for Environmental Research - UFZ (<http://www.ufz.de/>) and the  
590 German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig (<http://www.idiv-biodiversity.de/>). We would like to thank the administration and support staff of EVE who keep the system running and support us with our scientific computing needs: Thomas Schnicke, Ben Langenberg, Guido Schramm, Toni Harzendorf, Tom Stempel and Lisa Schurack from the UFZ, and Christian Krause from iDiv. [We thank Lily-belle Sweet for her help to improve the language of the manuscript.](#) Finally, we want to thank two anonymous reviewers, [the reviewer René Orth](#) and the Editor for their constructive comments, which improved the quality of this study.

- Albergel, C., De Rosnay, P., Balsamo, G., Isaksen, L., and Muñoz-Sabater, J.: Soil moisture analyses at ECMWF: Evaluation using global ground-based in situ observations, *Journal of Hydrometeorology*, 13, 1442–1460, <https://doi.org/10.1175/JHM-D-11-0107.1>, 2012.
- Andreasen, M., Jensen, K. H., Desilets, D., Franz, T. E., Zreda, M., Bogen, H. R., and Looms, M. C.: Status and Perspectives on the Cosmic-Ray Neutron Method for Soil Moisture Estimation and Other Environmental Science Applications, *Vadose Zone Journal*, 16, vzj2017.04.0086, <https://doi.org/https://doi.org/10.2136/vzj2017.04.0086>, <https://acsess.onlinelibrary.wiley.com/doi/abs/10.2136/vzj2017.04.0086>, 2017.
- Baatz, R., Bogen, H. R., Hendricks Franssen, H., Huisman, J. A., Montzka, C., and Vereecken, H.: An empirical vegetation correction for soil water content quantification using cosmic ray probes, *Water Resources Research*, 51, 2030–2046, <https://doi.org/10.1002/2014WR016443>, <https://onlinelibrary.wiley.com/doi/abs/10.1002/2014WR016443>, 2015.
- Baatz, R., Hendricks Franssen, H.-J., Han, X., Hoar, T., Bogen, H. R., and Vereecken, H.: Evaluation of a cosmic-ray neutron sensor network for improved land surface model prediction, *Hydrology and Earth System Sciences*, 21, 2509–2530, <https://doi.org/https://doi.org/10.5194/hess-21-2509-2017>, <https://hess.copernicus.org/articles/21/2509/2017/>, 2017.
- Baldocchi, D., Falge, E., Gu, L., Olson, R., Hollinger, D., Running, S., Anthoni, P., Bernhofer, C., Davis, K., Evans, R., Fuentes, J., Goldstein, A., Katul, G., Law, B., Lee, X., Malhi, Y., Meyers, T., Munger, W., Oechel, W., U. K. T. P., Pilegaard, K., Schmid, H. P., Valentini, R., Verma, S., Vesala, T., Wilson, K., and Wofsy, S.: FLUXNET: A New Tool to Study the Temporal and Spatial Variability of Ecosystem-Scale Carbon Dioxide, Water Vapor, and Energy Flux Densities, *Bulletin of the American Meteorological Society*, 82, 2415–2434, [https://doi.org/10.1175/1520-0477\(2001\)082<2415:FANTTS>2.3.CO;2](https://doi.org/10.1175/1520-0477(2001)082<2415:FANTTS>2.3.CO;2), [https://journals.ametsoc.org/view/journals/bams/82/11/1520-0477\\_2001\\_082\\_2415\\_fantts\\_2\\_3\\_co\\_2.xml](https://journals.ametsoc.org/view/journals/bams/82/11/1520-0477_2001_082_2415_fantts_2_3_co_2.xml), 2001.
- Baroni, G., Scheffele, L., Schrön, M., Ingwersen, J., and Oswald, S.: Uncertainty, sensitivity and improvements in soil moisture estimation with cosmic-ray neutron sensing, *Journal of Hydrology*, 564, 873–887, <https://doi.org/10.1016/j.jhydrol.2018.07.053>, <https://linkinghub.elsevier.com/retrieve/pii/S0022169418305675>, 2018.
- BGR: Digital soil map of Germany 1 : 1,000,000 (BUEK 1000), 1998.
- BGR: Hydrogeological map of Germany: 200,000 (HUEK 200), 2009.
- BGR: Digital soil map of Germany 1 : 200,000 (BUEK 200) v0.5, 2020.
- BKG: Digital Elevation Model (DEM), 2010.
- Boergens, E., Güntner, A., Dobslaw, H., and Dahle, C.: Quantifying the Central European Droughts in 2018 and 2019 With GRACE Follow-On, *Geophysical Research Letters*, 47, <https://doi.org/10.1029/2020GL087285>, 2020.
- Bogen, H., Herbst, M., Huisman, J., Rosenbaum, U., Weuthen, A., and Vereecken, H.: Potential of Wireless Sensor Networks for Measuring Soil Water Content Variability, *Vadose Zone Journal*, 9, 1002–1013, <https://doi.org/10.2136/vzj2009.0173>, <http://doi.wiley.com/10.2136/vzj2009.0173>, 2010.
- Bogen, H., Montzka, C., Huisman, J., Graf, A., Schmidt, M., Stockinger, M., von Hebel, C., Hendricks-Franssen, H., van der Kruk, J., Tappe, W., Lücke, A., Baatz, R., Bol, R., Groh, J., Pütz, T., Jakobi, J., Kunkel, R., Sorg, J., and Vereecken, H.: The TERENO-Rur Hydrological Observatory: A Multiscale Multi-Compartment Research Platform for the Advancement of Hydrological Science, *Vadose Zone Journal*, 17, 180055, <https://doi.org/10.2136/vzj2018.03.0055>, <http://doi.wiley.com/10.2136/vzj2018.03.0055>, 2018.
- Bogen, H. R.: TERENO: German network of terrestrial environmental observatories, *Journal of large-scale research facilities JLSRF*, 2, 52, <https://doi.org/10.17815/jlsrf-2-98>, <http://jlsrf.org/index.php/lrf/article/view/98>, 2016.

- Bogena, H. R., Huisman, J. A., Baatz, R., Hendricks Franssen, H.-J., and Vereecken, H.: Accuracy of the cosmic-ray soil water content probe in humid forest ecosystems: The worst case scenario: Cosmic-Ray Probe in Humid Forested Ecosystems, *Water Resources Research*, 49, 5778–5791, <https://doi.org/10.1002/wrcr.20463>, <http://doi.wiley.com/10.1002/wrcr.20463>, 2013.
- 635 Bogena, H. R., Huisman, J. A., Güntner, A., Hübner, C., Kusche, J., Jonard, F., Vey, S., and Vereecken, H.: Emerging methods for noninvasive sensing of soil moisture dynamics from field to catchment scale: a review, *WIREs Water*, 2, 635–647, <https://doi.org/10.1002/wat2.1097>, <https://onlinelibrary.wiley.com/doi/10.1002/wat2.1097>, 2015.
- Bogena, H. R., Schrön, M., Jakobi, J., Ney, P., Zacharias, S., Andreasen, M., Baatz, R., Boorman, D., Duygu, M. B., Eguibar-Galán, M. A., Fersch, B., Franke, T., Geris, J., González Sanchis, M., Kerr, Y., Korf, T., Mengistu, Z., Mialon, A., Nasta, P., Nitychoruk, J., Pisinaras, V., Rasche, D., Rosolem, R., Said, H., Schattan, P., Zreda, M., Achleitner, S., Albentosa-Hernández, E., Akyürek, Z., Blume, T., del Campo, A., Canone, D., Dimitrova-Petrova, K., Evans, J. G., Ferraris, S., Frances, F., Gisolo, D., Güntner, A., Herrmann, F., Iwema, J., Jensen, K. H., Kunstmann, H., Lidón, A., Looms, M. C., Oswald, S., Panagopoulos, A., Patil, A., Power, D., Rebmann, C., Romano, N., Scheffele, L., Seneviratne, S., Weltin, G., and Vereecken, H.: COSMOS-Europe: a European network of cosmic-ray neutron soil moisture sensors, *Earth System Science Data*, 14, 1125–1151, <https://doi.org/10.5194/essd-14-1125-2022>, [https://essd.copernicus.org/articles/14/](https://essd.copernicus.org/articles/14/1125/2022/)  
645 1125/2022/, 2022.
- Cammalleri, C., Micale, F., and Vogt, J.: On the value of combining different modelled soil moisture products for European drought monitoring, *Journal of Hydrology*, 525, 547–558, <https://doi.org/10.1016/j.jhydrol.2015.04.021>, <http://dx.doi.org/10.1016/j.jhydrol.2015.04.021>, publisher: Elsevier B.V., 2015.
- Chen, L., Huang, J.-G., Ma, Q., Hänninen, H., Rossi, S., Piao, S., and Bergeron, Y.: Spring phenology at different altitudes is becoming more uniform under global warming in Europe, *Global Change Biology*, 24, 3969–3975, <https://doi.org/10.1111/gcb.14288>, <https://onlinelibrary.wiley.com/doi/10.1111/gcb.14288>, 2018.
- De Lannoy, G. J. M., Koster, R. D., Reichle, R. H., Mahanama, S. P. P., and Liu, Q.: An updated treatment of soil texture and associated hydraulic properties in a global land modeling system, *Journal of Advances in Modeling Earth Systems*, 6, 957–979, <https://doi.org/10.1002/2014MS000330>, <http://doi.wiley.com/10.1002/2014MS000330>, 2014.
- 655 Dembélé, M., Ceperley, N., Zwart, S. J., Salvatore, E., Mariethoz, G., and Schaeffli, B.: Potential of satellite and reanalysis evaporation datasets for hydrological modelling under various model calibration strategies, *Advances in Water Resources*, 143, 103 667, <https://doi.org/10.1016/j.advwatres.2020.103667>, <https://linkinghub.elsevier.com/retrieve/pii/S030917082030230X>, 2020.
- Desilets, D., Zreda, M., and Ferré, T. P. A.: Nature’s neutron probe: Land surface hydrology at an elusive scale with cosmic rays: NATURE’S NEUTRON PROBE, *Water Resources Research*, 46, <https://doi.org/10.1029/2009WR008726>, [http://doi.wiley.com/10.1029/](http://doi.wiley.com/10.1029/2009WR008726)  
660 2009WR008726, 2010.
- Dimitrova-Petrova, K., Geris, J., Wilkinson, M. E., Rosolem, R., Verrot, L., Lilly, A., and Soulsby, C.: Opportunities and challenges in using catchment-scale storage estimates from cosmic ray neutron sensors for rainfall-runoff modelling, *Journal of Hydrology*, 586, 124 878, <https://doi.org/10.1016/j.jhydrol.2020.124878>, <https://linkinghub.elsevier.com/retrieve/pii/S0022169420303383>, 2020.
- EEA: CORINE Land Cover 1990, 2000 and 2006, 2009.
- 665 ESA: Global Land Cover Map for 2009, [http://due.esrin.esa.int/files/Globcover2009\\_V2.3\\_Global\\_.zip](http://due.esrin.esa.int/files/Globcover2009_V2.3_Global_.zip), 2009.
- Famiglietti, J. S., Ryu, D., Berg, A. A., Rodell, M., and Jackson, T. J.: Field observations of soil moisture variability across scales: SOIL MOISTURE VARIABILITY ACROSS SCALES, *Water Resources Research*, 44, <https://doi.org/10.1029/2006WR005804>, <https://onlinelibrary.wiley.com/doi/10.1029/2006WR005804>, 2008.

- Grillakis, M. G.: Increase in severe and extreme soil moisture droughts for Europe under climate change, *Science of The Total Environment*, 660, 1245–1255, <https://doi.org/10.1016/j.scitotenv.2019.01.001>, <https://linkinghub.elsevier.com/retrieve/pii/S0048969719300014>, 2019.
- Gupta, H. V., Kling, H., Yilmaz, K. K., and Martinez, G. F.: Decomposition of the mean squared error and NSE performance criteria: Implications for improving hydrological modelling, *Journal of Hydrology*, 377, 80–91, <https://doi.org/10.1016/j.jhydrol.2009.08.003>, <https://www.sciencedirect.com/science/article/pii/S0022169409004843>, 2009.
- Han, X., Hendricks Franssen, H.-J., Jiménez Bello, M. \., Rosolem, R., Bogen, H., Alzamora, F. M., Chanzy, A., and Vereecken, H.: Simultaneous soil moisture and properties estimation for a drip irrigated field by assimilating cosmic-ray neutron intensity, *Journal of Hydrology*, 539, 611–624, <https://doi.org/10.1016/j.jhydrol.2016.05.050>, <https://linkinghub.elsevier.com/retrieve/pii/S0022169416303171>, 2016.
- Hargreaves, G. H. and Samani, Z. A.: Reference Crop Evapotranspiration from Temperature, *Applied Engineering in Agriculture*, 1, 96–99, <https://doi.org/10.13031/2013.26773>, [http://ce.nmsu.edu/\\$sim\\$zsamani/papers/Hargreaves\\_Samani\\_85.pdf%5Cnhttp://elibrary.asabe.org/abstract.asp?JID=3&AID=26773&CID=aeaj1985&v=1&i=2&T=1](http://ce.nmsu.edu/$sim$zsamani/papers/Hargreaves_Samani_85.pdf%5Cnhttp://elibrary.asabe.org/abstract.asp?JID=3&AID=26773&CID=aeaj1985&v=1&i=2&T=1), 1985.
- Hari, V., Rakovec, O., Markonis, Y., Hanel, M., and Kumar, R.: Increased future occurrences of the exceptional 2018–2019 Central European drought under global warming, *Scientific Reports*, 10, <https://doi.org/10.1038/s41598-020-68872-9>, <https://doi.org/10.1038/s41598-020-68872-9>, ISBN: 4159802068872 Publisher: Nature Publishing Group UK, 2020.
- Hartmann, Jörg and Moosdorf, Nils: Global Lithological Map Database v1.0 (gridded to 0.5° spatial resolution), supplement to: Hartmann, Jens; Moosdorf, Nils (2012): The new global lithological map database GLiM: A representation of rock properties at the Earth surface. *Geochemistry, Geophysics, Geosystems*, 13, Q12004, <https://doi.org/10.1594/PANGAEA.788537>, <https://doi.pangaea.de/10.1594/PANGAEA.788537>, type: dataset, 2012.
- Itzerott, S., Hohmann, C., Stender, V., Maass, H., Borg, E., Renke, F., Jahncke, D., Berg, M., Conrad, C., and Spengler, D.: TERENO (Northeast), Climate stations of the GFZ German Research Centre for Geosciences (GFZ), <https://doi.org/10.5880/TERENO.GFZ.CL.2018.ALL>, <http://dataservices.gfz-potsdam.de/tereno-new/showshort.php?id=escidoc:3508888>, type: dataset, 2018a.
- Itzerott, S., Hohmann, C., Stender, V., Maass, H., Borg, E., Renke, F., Jahncke, D., Berg, M., Conrad, C., and Spengler, D.: TERENO (Northeast), Soil moisture stations of the GFZ German Research Centre for Geosciences (GFZ), <https://doi.org/10.5880/TERENO.GFZ.SM.2018.ALL>, <https://dataservices.gfz-potsdam.de/tereno-new/showshort.php?id=escidoc:3547898>, type: dataset, 2018b.
- Iwema, J., Rosolem, R., Rahman, M., Blyth, E., and Wagener, T.: Land surface model performance using cosmic-ray and point-scale soil moisture measurements for calibration, *Hydrology and Earth System Sciences*, 21, 2843–2861, <https://doi.org/10.5194/hess-21-2843-2017>, <https://hess.copernicus.org/articles/21/2843/2017/>, 2017.
- Kaspar, F., Müller-Westermeier, G., Penda, E., Mächel, H., Zimmermann, K., Kaiser-Weiss, A., and Deutschländer, T.: Monitoring of climate change in Germany – data, products and services of Germany’s National Climate Data Centre, *Advances in Science and Research*, 10, 99–106, <https://doi.org/10.5194/asr-10-99-2013>, <https://asr.copernicus.org/articles/10/99/2013/>, 2013.
- Keyantash, J. and Dracup, J.: *The Quantification of Drought: An Evaluation of Drought Indices*, American Meteorological Society, 2002.
- Kiese, R., Fersch, B., Baessler, C., Brosy, C., Butterbach-Bahl, K., Chwala, C., Dannenmann, M., Fu, J., Gasche, R., Grote, R., Jahn, C., Klatt, J., Kunstmann, H., Mauder, M., Rödiger, T., Smiatek, G., Soltani, M., Steinbrecher, R., Völksch, I., Werhahn, J., Wolf, B., Zeeman, M., and Schmid, H.: The TERENO Pre-Alpine Observatory: Integrating Meteorological, Hydrological, and Biogeochemical Measurements and Modeling, *Vadose Zone Journal*, 17, 180060, <https://doi.org/10.2136/vzj2018.03.0060>, <http://doi.wiley.com/10.2136/vzj2018.03.0060>, 2018.

- Koster, R. D., Guo, Z., Yang, R., Dirmeyer, P. A., Mitchell, K., and Puma, M. J.: On the Nature of Soil Moisture in Land Surface Models, *Journal of Climate*, 22, 4322–4335, <https://doi.org/10.1175/2009JCLI2832.1>, <http://journals.ametsoc.org/doi/10.1175/2009JCLI2832.1>, 2009.
- Koster, R. D., Reichle, R. H., Schubert, S. D., and Mahanama, S. P.: Length Scales of Hydrological Variability as Inferred from SMAP Soil  
710 Moisture Retrievals, *Journal of Hydrometeorology*, 20, 2129–2146, <https://doi.org/10.1175/JHM-D-19-0070.1>, <http://journals.ametsoc.org/doi/10.1175/JHM-D-19-0070.1>, 2019.
- Kumar, R., Samaniego, L., and Attinger, S.: Implications of distributed hydrologic model parameterization on water fluxes at multiple scales and locations: DISTRIBUTED HYDROLOGIC MODEL PARAMETERIZATIONS, *Water Resources Research*, 49, 360–379, <https://doi.org/10.1029/2012WR012195>, <http://doi.wiley.com/10.1029/2012WR012195>, 2013.
- 715 Köhli, M., Schrön, M., Zreda, M., Schmidt, U., Dietrich, P., and Zacharias, S.: Footprint characteristics revised for field-scale soil moisture monitoring with cosmic-ray neutrons, *Water Resources Research*, 51, 5772–5790, <https://doi.org/https://doi.org/10.1002/2015WR017169>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015WR017169>, 2015.
- Köhli, M., Weimar, J., Schrön, M., Baatz, R., and Schmidt, U.: Soil Moisture and Air Humidity Dependence of the Above-Ground Cosmic-Ray Neutron Intensity, *Frontiers in Water*, 2, <https://doi.org/10.3389/frwa.2020.544847>, [https://www.frontiersin.org/articles/10.3389/frwa.](https://www.frontiersin.org/articles/10.3389/frwa.2020.544847/full)  
720 2020.544847/full, 2021.
- Livneh, B., Kumar, R., and Samaniego, L.: Influence of soil textural properties on hydrologic fluxes in the Mississippi river basin: Influence of Soil Textural Properties on Hydrologic Fluxes, *Hydrological Processes*, 29, 4638–4655, <https://doi.org/10.1002/hyp.10601>, <http://doi.wiley.com/10.1002/hyp.10601>, 2015.
- Madruza de Brito, M., Kuhlicke, C., and Marx, A.: Near-real-time drought impact assessment: a text mining approach on the 2018/19  
725 drought in Germany, *Environmental Research Letters*, 15, 1040a9, <https://doi.org/10.1088/1748-9326/aba4ca>, <https://iopscience.iop.org/article/10.1088/1748-9326/aba4ca>, 2020.
- Marx, A., Kumar, R., Thober, S., Rakovec, O., Wanders, N., Zink, M., Wood, E. F., Pan, M., Sheffield, J., and Samaniego, L.: Climate change alters low flows in Europe under global warming of 1.5, 2, and 3 °C, *Hydrology and Earth System Sciences*, 22, 1017–1032, <https://doi.org/10.5194/hess-22-1017-2018>, <https://hess.copernicus.org/articles/22/1017/2018/>, 2018.
- 730 Mizukami, N., Clark, M. P., Newman, A. J., Wood, A. W., Gutmann, E. D., Nijssen, B., Rakovec, O., and Samaniego, L.: Towards seamless large-domain parameter estimation for hydrologic models: LARGE-DOMAIN MODEL PARAMETERS, *Water Resources Research*, 53, 8020–8040, <https://doi.org/10.1002/2017WR020401>, <http://doi.wiley.com/10.1002/2017WR020401>, 2017.
- Mo, K. C., Chen, L.-C., Shukla, S., Bohn, T. J., and Lettenmaier, D. P.: Uncertainties in North American Land Data Assimilation Systems over the Contiguous United States, *Journal of Hydrometeorology*, 13, 996–1009, <https://doi.org/10.1175/JHM-D-11-0132.1>, [http://journals.](http://journals.ametsoc.org/doi/10.1175/JHM-D-11-0132.1)  
735 [ametsoc.org/doi/10.1175/JHM-D-11-0132.1](http://journals.ametsoc.org/doi/10.1175/JHM-D-11-0132.1), 2012.
- O, S., Dutra, E., and Orth, R.: Robustness of Process-Based versus Data-Driven Modeling in Changing Climatic Conditions, *Journal of Hydrometeorology*, 21, 1929–1944, <https://doi.org/10.1175/JHM-D-20-0072.1>, <https://journals.ametsoc.org/view/journals/hydr/21/9/jhmd200072.xml>, 2020.
- Orth, R., O, S., Zscheischler, J., Mahecha, M. D., and Reichstein, M.: Contrasting biophysical and societal impacts of hydro-meteorological extremes, *Environmental Research Letters*, 17, 014 044, <https://doi.org/10.1088/1748-9326/ac4139>, [https://iopscience.iop.org/article/10.](https://iopscience.iop.org/article/10.1088/1748-9326/ac4139)  
740 1088/1748-9326/ac4139, 2022.
- Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y.-W., Poindexter, C., Chen, J., Elbashandy, A., Humphrey, M., Isaac, P., Polidori, D., Reichstein, M., Ribeca, A., van Ingen, C., Vuichard, N., Zhang, L., Amiro, B., Ammann, C., Arain, M. A., Ardö, J.,

- Arkebauer, T., Arndt, S. K., Arriga, N., Aubinet, M., Aurela, M., Baldocchi, D., Barr, A., Beamesderfer, E., Marchesini, L. B., Bergeron, O., Beringer, J., Bernhofer, C., Berveiller, D., Billesbach, D., Black, T. A., Blanken, P. D., Bohrer, G., Boike, J., Bolstad, P. V., Bonal, D., Bonnefond, J.-M., Bowling, D. R., Bracho, R., Brodeur, J., Brümmer, C., Buchmann, N., Burban, B., Burns, S. P., Buysse, P., Cale, P., Cavagna, M., Cellier, P., Chen, S., Chini, I., Christensen, T. R., Cleverly, J., Collalti, A., Consalvo, C., Cook, B. D., Cook, D., Coursolle, C., Cremonese, E., Curtis, P. S., D'Andrea, E., da Rocha, H., Dai, X., Davis, K. J., Cinti, B. D., Grandcourt, A. d., Ligne, A. D., De Oliveira, R. C., Delpierre, N., Desai, A. R., Di Bella, C. M., Tommasi, P. d., Dolman, H., Domingo, F., Dong, G., Dore, S., Duce, P., Dufrêne, E., Dunn, A., Dušek, J., Eamus, D., Eichmann, U., ElKhidir, H. A. M., Eugster, W., Ewenz, C. M., Ewers, B., Famulari, D., Fares, S., Feigenwinter, I., Feitz, A., Fensholt, R., Filippa, G., Fischer, M., Frank, J., Galvagno, M., Gharun, M., Gianelle, D., Gielen, B., Gioli, B., Gitelson, A., Goded, I., Goeckede, M., Goldstein, A. H., Gough, C. M., Goulden, M. L., Graf, A., Griebel, A., Gruening, C., Grünwald, T., Hammerle, A., Han, S., Han, X., Hansen, B. U., Hanson, C., Hatakka, J., He, Y., Hehn, M., Heinesch, B., Hinko-Najera, N., Hörtnagl, L., Hutley, L., Ibrom, A., Ikawa, H., Jackowicz-Korczynski, M., Janouš, D., Jans, W., Jassal, R., Jiang, S., Kato, T., Khomik, M., Klatt, J., Knohl, A., Knox, S., Kobayashi, H., Koerber, G., Kolle, O., Kosugi, Y., Kotani, A., Kowalski, A., Kruijt, B., Kurbatova, J., Kutsch, W. L., Kwon, H., Launiainen, S., Laurila, T., Law, B., Leuning, R., Li, Y., Liddell, M., Limousin, J.-M., Lion, M., Liska, A. J., Lohila, A., López-Ballesteros, A., López-Blanco, E., Loubet, B., Loustau, D., Lucas-Moffat, A., Lüers, J., Ma, S., Macfarlane, C., Magliulo, V., Maier, R., Mammarella, I., Manca, G., Marcolla, B., Margolis, H. A., Marras, S., Massman, W., Mastepanov, M., Matamala, R., Matthes, J. H., Mazzenga, F., McCaughey, H., McHugh, I., McMillan, A. M. S., Merbold, L., Meyer, W., Meyers, T., Miller, S. D., Minerbi, S., Moderow, U., Monson, R. K., Montagnani, L., Moore, C. E., Moors, E., Moreaux, V., Moureaux, C., Munger, J. W., Nakai, T., Neiryneck, J., Nesic, Z., Nicolini, G., Noormets, A., Northwood, M., Nosetto, M., Nouvellon, Y., Novick, K., Oechel, W., Olesen, J. E., Ourcival, J.-M., Papuga, S. A., Parmentier, F.-J., Paul-Limoges, E., Pavelka, M., Peichl, M., Pendall, E., Phillips, R. P., Pilegaard, K., Pirk, N., Posse, G., Powell, T., Prasse, H., Prober, S. M., Rambal, S., Rannik, A., Raz-Yaseef, N., Rebmann, C., Reed, D., Dios, V. R. d., Restrepo-Coupe, N., Reverter, B. R., Roland, M., Sabbatini, S., Sachs, T., Saleska, S. R., Sánchez-Cañete, E. P., Sanchez-Mejia, Z. M., Schmid, H. P., Schmidt, M., Schneider, K., Schrader, F., Schroder, I., Scott, R. L., Sedláč, P., Serrano-Ortíz, P., Shao, C., Shi, P., Shironya, I., Siebicke, L., Šigut, L., Silberstein, R., Sirca, C., Spano, D., Steinbrecher, R., Stevens, R. M., Sturtevant, C., Suyker, A., Tagesson, T., Takanashi, S., Tang, Y., Tapper, N., Thom, J., Tomassucci, M., Tuovinen, J.-P., Urbanski, S., Valentini, R., van der Molen, M., van Gorsel, E., van Huissteden, K., Varlagin, A., Verfaillie, J., Vesala, T., Vincke, C., Vitale, D., Vygodskaya, N., Walker, J. P., Walter-Shea, E., Wang, H., Weber, R., Westermann, S., Wille, C., Wofsy, S., Wohlfahrt, G., Wolf, S., Woodgate, W., Li, Y., Zampedri, R., Zhang, J., Zhou, G., Zona, D., Agarwal, D., Biraud, S., Torn, M., and Papale, D.: The FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy covariance data, *Scientific Data*, 7, 225, <https://doi.org/10.1038/s41597-020-0534-3>, <https://www.nature.com/articles/s41597-020-0534-3>, 2020.
- Peichl, M., Thober, S., Meyer, V., and Samaniego, L.: The effect of soil moisture anomalies on maize yield in Germany, *Natural Hazards and Earth System Sciences*, 18, 889–906, <https://doi.org/10.5194/nhess-18-889-2018>, <https://nhess.copernicus.org/articles/18/889/2018/>, 2018.
- Peichl, M., Thober, S., Samaniego, L., Hansjürgens, B., and Marx, A.: Machine-learning methods to assess the effects of a non-linear damage spectrum taking into account soil moisture on winter wheat yields in Germany, *Hydrology and Earth System Sciences*, 25, 6523–6545, <https://doi.org/10.5194/hess-25-6523-2021>, <https://hess.copernicus.org/articles/25/6523/2021/>, 2021.
- Peng, J., Albergel, C., Balenzano, A., Brocca, L., Cartus, O., Cosh, M. H., Crow, W. T., Dabrowska-Zielinska, K., Dadson, S., Davidson, M. W., de Rosnay, P., Dorigo, W., Gruber, A., Hagemann, S., Hirschi, M., Kerr, Y. H., Lovergine, F., Mahecha, M. D., Marzahn, P., Mattia, F., Musial, J. P., Preusmann, S., Reichle, R. H., Satalino, G., Silgram, M., van Bodegom, P. M., Verhoest, N. E., Wagner, W., Walker, J. P., Wegmüller, U., and Loew, A.: A roadmap for high-resolution satellite soil moisture applications – confronting product

- characteristics with user requirements, *Remote Sensing of Environment*, 252, 112–162, <https://doi.org/10.1016/j.rse.2020.112162>, <https://linkinghub.elsevier.com/retrieve/pii/S0034425720305356>, 2021.
- 785 Pütz, T., Kiese, R., Wollschläger, U., Groh, J., Rupp, H., Zacharias, S., Priesack, E., Gerke, H. H., Gasche, R., Bens, O., Borg, E., Baessler, C., Kaiser, K., Herbrich, M., Munch, J.-C., Sommer, M., Vogel, H.-J., Vanderborght, J., and Vereecken, H.: TERENO-SOILCan: a lysimeter-network in Germany observing soil processes and plant diversity influenced by climate change, *Environmental Earth Sciences*, 75, 1242, <https://doi.org/10.1007/s12665-016-6031-5>, <https://doi.org/10.1007/s12665-016-6031-5>, 2016.
- Rakovec, O., Kumar, R., Mai, J., Cuntz, M., Thober, S., Zink, M., Attinger, S., Schäfer, D., Schrön, M., and Samaniego, L.: Multiscale and Multivariate Evaluation of Water Fluxes and States over European River Basins, *Journal of Hydrometeorology*, <https://doi.org/10.1175/JHM-D-15-0054.1>, 2016.
- 790 Rakovec, O., Mizukami, N., Kumar, R., Newman, A. J., Thober, S., Wood, A. W., Clark, M. P., and Samaniego, L.: Diagnostic Evaluation of Large-Domain Hydrologic Models Calibrated Across the Contiguous United States, *Journal of Geophysical Research: Atmospheres*, 124, 13 991–14 007, <https://doi.org/10.1029/2019JD030767>, <https://onlinelibrary.wiley.com/doi/10.1029/2019JD030767>, 2019.
- Rakovec, O., Samaniego, L., Hari, V., Markonis, Y., Moravec, V., Thober, S., Hanel, M., and Kumar, R.: The 2018–2020 Multi-Year Drought Sets a New Benchmark in Europe, *Earth’s Future*, 10, <https://doi.org/10.1029/2021EF002394>, <https://onlinelibrary.wiley.com/doi/10.1029/2021EF002394>, 2022.
- 795 Rauthe, M., Steiner, H., Riediger, U., Mazurkiewicz, A., and Gratzki, A.: A Central European precipitation climatology – Part I: Generation and validation of a high-resolution gridded daily data set (HYRAS), *Meteorologische Zeitschrift*, pp. 235–256, <https://doi.org/10.1127/0941-2948/2013/0436>, [https://www.schweizerbart.de/papers/metz/detail/22/81060/A\\_Central\\_European\\_precipitation\\_climatology\\_Part\\_?af=crossref](https://www.schweizerbart.de/papers/metz/detail/22/81060/A_Central_European_precipitation_climatology_Part_?af=crossref), 2013.
- 800 Rosenbaum, U., Bogen, H. R., Herbst, M., Huisman, J. A., Peterson, T. J., Weuthen, A., Western, A. W., and Vereecken, H.: Seasonal and event dynamics of spatial soil moisture patterns at the small catchment scale: DYNAMICS OF CATCHMENT-SCALE SOIL MOISTURE PATTERNS, *Water Resources Research*, 48, <https://doi.org/10.1029/2011WR011518>, <http://doi.wiley.com/10.1029/2011WR011518>, 2012.
- 805 Saha, T. R., Shrestha, P. K., Rakovec, O., Thober, S., and Samaniego, L.: A drought monitoring tool for South Asia, *Environmental Research Letters*, 16, 054 014, <https://doi.org/10.1088/1748-9326/abf525>, <https://iopscience.iop.org/article/10.1088/1748-9326/abf525>, 2021.
- Samaniego, L., Kumar, R., and Attinger, S.: Multiscale parameter regionalization of a grid-based hydrologic model at the mesoscale: MULTISCALE PARAMETER REGIONALIZATION, *Water Resources Research*, 46, <https://doi.org/10.1029/2008WR007327>, <http://doi.wiley.com/10.1029/2008WR007327>, 2010.
- 810 Samaniego, L., Kumar, R., and Zink, M.: Implications of parameter uncertainty on soil moisture drought analysis in Germany, *Journal of Hydrometeorology*, 14, 47–68, <https://doi.org/10.1175/JHM-D-12-075.1>, ISBN: 1525-755X, 2013.
- Samaniego, L., Kumar, R., Thober, S., Rakovec, O., Zink, M., Wanders, N., Eisner, S., Müller Schmied, H., Sutanudjaja, E. H., Warrach-Sagi, K., and Attinger, S.: Toward seamless hydrologic predictions across spatial scales, *Hydrology and Earth System Sciences*, 21, 4323–4346, <https://doi.org/https://doi.org/10.5194/hess-21-4323-2017>, <https://hess.copernicus.org/articles/21/4323/2017/>, 2017.
- 815 Samaniego, L., Thober, S., Kumar, R., Wanders, N., Rakovec, O., Pan, M., Zink, M., Sheffield, J., Wood, E. F., and Marx, A.: Anthropogenic warming exacerbates European soil moisture droughts, *Nature Climate Change*, 8, 421–426, <https://doi.org/10.1038/s41558-018-0138-5>, <https://www.nature.com/articles/s41558-018-0138-5>, 2018.



- Schattan, P., Baroni, G., Oswald, S. E., Schöber, J., Fey, C., Kormann, C., Huttenlau, M., and Achleitner, S.: Continuous monitoring of snow-pack dynamics in alpine terrain by aboveground neutron sensing: ALPINE SNOWPACK MONITORING BY CRNS, *Water Resources Research*, 53, 3615–3634, <https://doi.org/10.1002/2016WR020234>, <http://doi.wiley.com/10.1002/2016WR020234>, 2017.
- Schrön, M., Köhli, M., Scheffele, L., Iwema, J., Bogen, H. R., Lv, L., Martini, E., Baroni, G., Rosolem, R., Weimar, J., Mai, J., Cuntz, M., Rebmann, C., Oswald, S. E., Dietrich, P., Schmidt, U., and Zacharias, S.: Improving calibration and validation of cosmic-ray neutron sensors in the light of spatial sensitivity, *Hydrology and Earth System Sciences*, 21, 5009–5030, <https://doi.org/10.5194/hess-21-5009-2017>, <https://hess.copernicus.org/articles/21/5009/2017/>, 2017.
- Schrön, M., Zacharias, S., Womack, G., Köhli, M., Desilets, D., Oswald, S. E., Bumberger, J., Mollenhauer, H., Kögler, S., Remmler, P., Kasner, M., Denk, A., and Dietrich, P.: Intercomparison of cosmic-ray neutron sensors and water balance monitoring in an urban environment, *Geoscientific Instrumentation, Methods and Data Systems*, 7, 83–99, <https://doi.org/10.5194/gi-7-83-2018>, <https://gi.copernicus.org/articles/7/83/2018/>, 2018.
- Sepulcre-Canto, G., Horion, S., Singleton, A., Carrao, H., and Vogt, J.: Development of a Combined Drought Indicator to detect agricultural drought in Europe, *Natural Hazards and Earth System Sciences*, 12, 3519–3531, <https://doi.org/10.5194/nhess-12-3519-2012>, <https://nhess.copernicus.org/articles/12/3519/2012/>, 2012.
- Shuttleworth, J., Rosolem, R., Zred, M., and Franz, T.: The COsmic-ray Soil Moisture Interaction Code (COSMIC) for use in data assimilation, *Hydrology and Earth System Sciences*, 17, 3205–3217, <https://doi.org/10.5194/hess-17-3205-2013>, <https://hess.copernicus.org/articles/17/3205/2013/>, 2013.
- Svoboda, M., LeComte, D., Hayes, M., Heim, R., Gleason, K., Angel, J., Rippey, B., Tinker, R., Palecki, M., Stooksbury, D., Miskus, D., and Stephens, S.: THE DROUGHT MONITOR, *Bulletin of the American Meteorological Society*, 83, 1181–1190, <https://doi.org/10.1175/1520-0477-83.8.1181>, [https://journals.ametsoc.org/view/journals/bams/83/8/1520-0477-83\\_8\\_1181.xml](https://journals.ametsoc.org/view/journals/bams/83/8/1520-0477-83_8_1181.xml), 2002.
- Thober, S., Kumar, R., Sheffield, J., Mai, J., Schäfer, D., and Samaniego, L.: Seasonal Soil Moisture Drought Prediction over Europe Using the North American Multi-Model Ensemble (NMME), *Journal of Hydrometeorology*, 16, 2329–2344, <https://doi.org/10.1175/JHM-D-15-0053.1>, <http://journals.ametsoc.org/doi/10.1175/JHM-D-15-0053.1>, 2015.
- Tolson, B. A. and Shoemaker, C. A.: Dynamically dimensioned search algorithm for computationally efficient watershed model calibration, *Water Resources Research*, 43, <https://doi.org/10.1029/2005WR004723>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2005WR004723>, 2007.
- USGS: Global Multi-resolution Terrain Elevation Data 2010 (GMTED2010), <https://doi.org/10.5066/F7J38R2N>, <https://www.usgs.gov/centers/eros/science/usgs-eros-archive-digital-elevation-global-multi-resolution-terrain-elevation>, type: dataset, 2017.
- Vereecken, H., Huisman, J. A., Bogen, H., Vanderborght, J., Vrugt, J. A., and Hopmans, J. W.: On the value of soil moisture measurements in vadose zone hydrology: A review: SOIL MOISTURE AND HYDROLOGY, *Water Resources Research*, 44, <https://doi.org/10.1029/2008WR006829>, <http://doi.wiley.com/10.1029/2008WR006829>, 2008.
- Wanders, N., Bierkens, M. F. P., de Jong, S. M., de Roo, A., and Karssen, D.: The benefits of using remotely sensed soil moisture in parameter identification of large-scale hydrological models, *Water Resources Research*, 50, 6874–6891, <https://doi.org/10.1002/2013WR014639>, <http://doi.wiley.com/10.1002/2013WR014639>, 2014.
- Western, A. W., Zhou, S.-L., Grayson, R. B., McMahon, T. A., Blöschl, G., and Wilson, D. J.: Spatial correlation of soil moisture in small catchments and its relationship to dominant spatial hydrological processes, *Journal of Hydrology*, 286, 113–134, <https://doi.org/10.1016/j.jhydrol.2003.09.014>, <https://linkinghub.elsevier.com/retrieve/pii/S0022169403003809>, 2004.

- 855 Wickenkamp, I., Huisman, J. A., Bogen, H. R., and Vereecken, H.: Effects of Deforestation on Water Flow in the Vadose Zone, *Water*, 12, 35, <https://doi.org/10.3390/w12010035>, <https://www.mdpi.com/2073-4441/12/1/35>, 2019.
- Xia, Y., Sheffield, J., Ek, M. B., Dong, J., Chaney, N., Wei, H., Meng, J., and Wood, E. F.: Evaluation of multi-model simulated soil moisture in NLDAS-2, *Journal of Hydrology*, 512, 107–125, <https://doi.org/10.1016/j.jhydrol.2014.02.027>, <http://dx.doi.org/10.1016/j.jhydrol.2014.02.027>, publisher: Elsevier B.V., 2014.
- 860 Zacharias, S. and Wessolek, G.: Excluding Organic Matter Content from Pedotransfer Predictors of Soil Water Retention, *Soil Science Society of America Journal*, 71, 43–50, <https://doi.org/10.2136/sssaj2006.0098>, <http://doi.wiley.com/10.2136/sssaj2006.0098>, 2007.
- Zacharias, S., Bogen, H., Samaniego, L., Mauder, M., Fuß, R., Pütz, T., Frenzel, M., Schwank, M., Baessler, C., Butterbach-Bahl, K., Bens, O., Borg, E., Brauer, A., Dietrich, P., Hajnsek, I., Helle, G., Kiese, R., Kunstmann, H., Klotz, S., Munch, J. C., Papen, H., Priesack, E., Schmid, H. P., Steinbrecher, R., Rosenbaum, U., Teutsch, G., and Vereecken, H.: A Network of Terrestrial Environmental Observatories
- 865 in Germany, *Vadose Zone Journal*, 10, 955–973, <https://doi.org/10.2136/vzj2010.0139>, <http://doi.wiley.com/10.2136/vzj2010.0139>, 2011.
- Zink, M., Samaniego, L., Kumar, R., Thober, S., Mai, J., Schäfer, D., and Marx, A.: The German drought monitor, *Environmental Research Letters*, 11, 074002, <https://doi.org/10.1088/1748-9326/11/7/074002>, <https://iopscience.iop.org/article/10.1088/1748-9326/11/7/074002>, 2016.
- Zink, M., Kumar, R., Cuntz, M., and Samaniego, L.: A high-resolution dataset of water fluxes and states for Germany accounting for
- 870 parametric uncertainty, *Hydrology and Earth System Sciences*, 21, 1769–1790, <https://doi.org/10.5194/hess-21-1769-2017>, iSBN: 1607-7938, 2017.
- Zreda, M., Shuttleworth, W. J., Zeng, X., Zweck, C., Desilets, D., Franz, T., and Rosolem, R.: COSMOS: the COsmic-ray Soil Moisture Observing System, *Hydrology and Earth System Sciences*, 16, 4079–4099, <https://doi.org/https://doi.org/10.5194/hess-16-4079-2012>, <https://hess.copernicus.org/articles/16/4079/2012/>, 2012.