

The influence of vegetation water dynamics on the ASCAT backscatter-incidence angle relationship in the Amazon

Ashwini Petchiappan¹, Susan C. Steele-Dunne², Mariette Vreugdenhil³, Sebastian Hahn³, Wolfgang Wagner³, and Rafael Oliveira⁴

¹Department of Water Management, Delft University of Technology, Stevinweg 1, Delft 2600 GA, The Netherlands

²Department of Geoscience and Remote Sensing, Delft University of Technology, Stevinweg 1, Delft 2600 GA, The Netherlands

³Department of Geodesy and Geo-Information, TU Wien, Vienna 1040, Austria

⁴Department of Plant Biology, Institute of Biology P.O.Box: 6109, University of Campinas – UNICAMP 13083-970, Campinas, SP, Brazil

Correspondence: Susan Steele-Dunne (s.c.steele-dunne@tudelft.nl)

Abstract. Microwave observations are sensitive to plant water content and could therefore provide essential information on biomass and plant water status in ecological and agricultural applications. The combined data record of the C-band scatterometers on ERS 1/2, the Metop series and the planned Metop Second Generation satellites will span over 40 years, which would provide a long-term perspective on the role of vegetation in the climate system. Recent research has indicated that the unique viewing geometry of ASCAT could be exploited to observe vegetation water dynamics. The incidence angle dependence of backscatter can be described with a second order polynomial, the slope and curvature of which are related to vegetation. In a study limited to grasslands, seasonal cycles, spatial patterns and interannual variability in the slope and curvature were found to vary among grassland types and were attributed to differences in moisture availability, growing season length and phenological changes. To exploit ASCAT slope and curvature for global vegetation monitoring, their dynamics over a wider range of vegetation types needs to be quantified and explained in terms of vegetation water dynamics. Here, we compare ASCAT data with meteorological data and GRACE Equivalent Water Thickness (EWT) to explain the dynamics of ASCAT backscatter, slope and curvature in terms of moisture availability and demand. We consider differences in the seasonal cycle, diurnal differences, and the response to the 2010 and 2015 droughts across ecoregions in the Amazon basin and surroundings. Results show that spatial and temporal patterns in backscatter reflect moisture availability indicated by GRACE EWT. Slope and curvature dynamics vary considerably among the ecoregions. The evergreen forests, often used as a calibration target, exhibit very stable behaviour even under drought conditions. The limited seasonal variation follows changes in the radiation cycle, and may indicate phenological changes such as litterfall. In contrast, the diversity of land cover types within the Cerrado region results in considerable heterogeneity in terms of the seasonal cycle and the influence of drought on both slope and curvature. Seasonal flooding in forest and savanna areas also produced a distinctive signature in terms of the backscatter as a function of incidence angle. This improved understanding of the incidence angle behaviour of backscatter increases our ability to interpret and make optimal use of the ASCAT data record and Vegetation Optical Depth products for vegetation monitoring.

23 1 Introduction

24 Microwave remote sensing observations are sensitive to plant water content, which depends on above ground biomass and plant
25 water status (Konings et al., 2019; Owe et al., 2001; Jackson et al., 1982). Data from active and passive microwave sensors can
26 provide valuable information about vegetation in a range of applications in ecological and agricultural monitoring (Konings
27 et al., 2019; Chaparro et al., 2019; Rao et al., 2019; Steele-Dunne et al., 2017; Tian et al., 2016; Andela et al., 2013; Saatchi
28 et al., 2013; Liu et al., 2013; McNairn et al., 2000; Wagner et al., 1999). In particular, Vegetation Optical Depth (VOD) products
29 derived from various passive and active microwave sensors are increasingly used for biomass monitoring (Liu et al., 2015),
30 drought monitoring (Liu et al., 2018), wildfire risk assessment (Forkel et al., 2019) and have been related to Gross Primary
31 Production (Teubner et al., 2018, 2019), carbon stocks (Chaparro et al., 2019) and drought-driven tree mortality (Rao et al.,
32 2019). Currently VOD datasets are available from single sensor passive microwave observations, such as SMAP (Konings
33 et al., 2016), SMOS (Fernandez-Moran et al., 2017) and AMSR2 (Owe et al., 2001; De Jeu, 2003), and active microwave
34 observations such as ASCAT (Vreugdenhil et al., 2016). Furthermore, long-term data records are available that combine VOD
35 from different sensors (Moesinger et al., 2020; Liu et al., 2011).

36 The current study is motivated by the availability of consistent C-band data from 1991 to at least 2030, and its potential
37 value as a long-term data record for vegetation monitoring. The Advanced Scatterometer (ASCAT) is a real aperture radar
38 operating at 5.255 GHz with VV polarization. There are currently three ASCAT instruments in orbit on Metop-A, Metop-B
39 and Metop-C, launched in October 2006, September 2012 and November 2018 respectively. ASCAT builds on the success
40 of the European Scatterometer (ESCAT) which flew on the ERS-1/2 satellites from 1991-2011 (Attema, 1991; Figa-Saldaña
41 et al., 2002; Wagner et al., 2013)). Continuation of the ESCAT/ASCAT record is ensured by the plans to launch SCA on
42 Metop-SG in 2024 (Stoffelen et al., 2017). Using data from a single series of satellites with identical and inter-calibrated
43 instruments circumvents many of the challenges of reconciling data using different frequencies, viewing geometries and orbit
44 characteristics. The continuity from ERS to Metop and Metop-SG ensures an internally consistent data product for at least 40
45 years, rendering it ideal to study the role of vegetation in the climate system.

46 Many early studies demonstrated the sensitivity of ESCAT and ASCAT backscatter to vegetation, and explored the potential
47 value of these data for vegetation monitoring (Wismann et al., 1995; Frison et al., 1998; Woodhouse et al., 1999; Jarlan et al.,
48 2002; Steele-Dunne et al., 2012; Schroeder et al., 2016). These studies focused on spatial and temporal variations in backscatter
49 normalized to some reference angle. Here, the focus is on the potential information content of the incidence angle behaviour
50 of backscatter, and particularly the so-called "Dynamic Vegetation Parameters" describing the incidence angle behaviour of
51 backscatter as calculated in the TU Wien Soil Moisture Retrieval (TUW SMR) algorithm (Hahn et al., 2017).

52 The ASCAT Dynamic Vegetation Parameters refer to the parameters of the second order Taylor polynomial used to describe
53 the incidence angle (θ) dependence of backscatter σ° . This is described as follows:

$$54 \quad \sigma^\circ(\theta) = \sigma^\circ(\theta_r) + \sigma'(\theta) \cdot (\theta - \theta_r) + \frac{1}{2} \cdot \sigma''(\theta_r) \cdot (\theta - \theta_r)^2 \quad [dB] \quad (1)$$

55 where $\sigma^\circ(\theta_r)$, $\sigma'(\theta_r)$ and $\sigma''(\theta_r)$ are the normalized backscatter, slope and curvature at some reference angle θ_r . In the TUW
56 SMR algorithm, this expression is used to normalize backscatter values from different incidence angles to a reference angle
57 θ_r . It is also used to account for the influence of vegetation on backscatter as the incidence angle behaviour of σ° depends on
58 whether total backscatter is dominated by surface scattering from the soil, volume scattering from the vegetation, or multiple
59 scattering (Wagner et al., 1999; Naeimi et al., 2009; Hahn et al., 2017). In other words, slope and curvature are calculated
60 and used to account for the influence of vegetation in the soil moisture retrieval. An increase in soil moisture results in an
61 increase in backscatter at all incidence angles, while a change in the vegetation (due to growth cycle or water status) changes
62 the sensitivity of backscatter to incidence angle, i.e. it results in a change in slope and curvature. So, the slope and curvature
63 provide complementary information to the normalized backscatter.

64 Results from Steele-Dunne et al. (2019) suggest that considering the slope ($\sigma'(\theta)$) and curvature ($\sigma''(\theta)$) dynamics in com-
65 bination with the backscatter could yield valuable insights into vegetation water dynamics. Seasonal cycles, spatial patterns
66 and interannual variability in the slope varied between grassland cover type reflecting variations in soil moisture availability
67 and growing season length. Slope is considered an indication of vegetation density, or above ground fresh biomass, which is
68 a combination of dry biomass and vegetation water content. Results also suggested that curvature variations were influenced
69 by the total water content, but also its vertical distribution within the vegetation and the geometry of constituents. Contiguous
70 anomalies were observed in both slope and curvature during drought periods, suggesting that the slope and curvature provide
71 insight into when the severity of a soil moisture anomaly is enough to impact vegetation. Diurnal variations were also observed
72 and attributed to sub-daily variations in the dominant scattering mechanism due to changes in the vertical moisture distribution
73 of the grasses. More recently, Pfeil et al. (2020) observed a “spring peak” in slope values around April in broadleaf deciduous
74 forest in Europe. Using LAI and data from the Pan European Phenological database (PEP725) (Templ et al., 2018) they argued
75 that this spring peak in ASCAT slope coincides with spring activation, particularly the increase in water content of bare twigs
76 and branches prior to leaf out in broadleaf deciduous forests. ASCAT slope and curvature therefore seem to be sensitive to
77 changes in vegetation water content and structure of vegetation.

78 The goal of this study is to improve our understanding of the ASCAT backscatter-incidence angle relationship and how
79 they might be used to monitor vegetation water dynamics. The Amazon basin and its surroundings has been chosen as a study
80 area as it provides a wide range in terms of expected variability in ASCAT backscatter, slope and curvature. Backscatter in
81 the evergreen forest was considered so stable that this region has been used for satellite radar calibration (Birrer et al., 1982).
82 In contrast, seasonal changes in the Cerrado are expected to yield strong annual cycles in backscatter, slope and curvature.
83 Seasonal cycles and diurnal differences in ASCAT backscatter, slope and curvature will be determined for several ecoregions
84 of interest. These will be compared to meteorological data and GRACE terrestrial water storage variations to relate the ASCAT
85 backscatter, slope and curvature to moisture availability and demand. Finally, we will investigate whether there are anomalies
86 in the ASCAT backscatter, slope and curvature as a result of the 2010 and 2015 droughts.




Figure 1. Study Area. The map is colored by biome, and nine ecoregions of interest are highlighted based on the dataset of Olson et al. (2001). The six forest ecoregions of interest are Napo moist forest (fNW), Guianan moist forests (fNE), Southwest Amazon moist forests (fSW), Madeira-Tapajos moist forest (fSE), Jurua-Perez moist forests (fC) and the Marajo Varzea flooded forests (ff). The three savanna ecoregions of interest are the Cerrado (sC), Guianan Savanna (sG) and Beni Savanna (sB).

2 Data and Methods

2.1 Study Area

Figure 1 shows the extent of the study domain, highlights the biomes (by color) and outlines the ecoregions of interest identified in the WWF Terrestrial Ecoregions dataset (WWF, 2019) and described by Olson et al. (2001). The study domain extends from 9°N to 19°S, and 44°W to 80°W. Most of the study region is covered by the Amazon rain forest, which extends over 5.3 million km² (Soares et al., 2006). Six forest ecoregions are investigated here:

1. The Napo moist forests (fNW), located in northwest Amazonia, receive some of the highest amounts of annual precipitation in the biome, reaching up to 4000 mm in some parts. This highly biodiverse region has canopies reaching 40 m.
2. The Guianan moist forests (fNE) are one of the largest continuous stretches of relatively pristine lowland tropical rainforest in the world. There are two distinct wet seasons: from December to January and from May to August. The floral diversity is rich, with multi-tiered vegetation of 40 m tall trees with herbaceous plants below. The dry season (September-November) can see a substantial reduction in leaves, although the forest is evergreen.

- 100 3. The Southwest Amazon moist forests (fSW) have significant variations in topography and soil characteristics, leading
101 to extremely high biodiversity. The size and orientation of the ecoregion means that climatic conditions vary markedly
102 – the north being wetter and having less seasonal variability compared to the south. The inaccessibility of the region has
103 aided in its conservation.
- 104 4. The Madeira-Tapajós moist forests (fSE) are transected by the Transamazon Highway, and have high levels of urbaniza-
105 tion and deforestation. There are characteristic liana (woody vine) forests with a lower (< 25 m) and more open canopy
106 than the typical humid terra firme forests.
- 107 5. The Juruá-Perez moist forests (fC) are largely intact forests in the low Amazon Basin. The canopy can reach up to 30 m,
108 with some patches of open canopy.
- 109 6. The seasonally flooded forest, Marajó várzea (ff), is located at the mouth of the Amazon River. The vegetation is dom-
110 inated by palms, and shorter than surrounding forests. It has areas with tidal flows from the Atlantic Ocean, as well as
111 seasonally and permanently inundated forests. The annual seasonal flooding occurs during the peak precipitation period
112 between January-May (Camarão et al., 2002).

113 Three savanna ecoregions are also considered in this study:

- 114 1. The Cerrado (sC) borders the Amazon biome to the southeast. It occupies an area of 2 million km² in the Brazilian
115 Central Plateau and is the second most extensive biome in South America (Oliveira et al., 2005). The vegetation cover
116 varies from closed tree canopy to grasslands with low shrubs only (Eiten, 1972).
- 117 2. The Guianan savanna (sG) consists of forest patches encircled by extensive grasslands and shrub formations. The area
118 is more susceptible to vegetation fires than typical humid moist forest environments and the dry season lasts from
119 December-March.
- 120 3. The Beni savanna (sB) is a wetland region with riverine gallery forests and small forest islands. The landscape is domi-
121 nated by the palm species *Attalea princeps* (Hordijk et al., 2019). Seasonal flooding occurs in up to half the region for 4
122 to 9 months, peaking in March-April (Hamilton et al., 2004).

123 Three Köppen-Geiger Climate Classes (KGCC) cover most of the study region (Fig. 2). The evergreen forest regions are
124 classified as Af (tropical fully humid) or Am (tropical monsoonal), and the savanna regions have Aw (tropical winter dry)
125 climate (Bradley et al., 2011). The annual precipitation in the forests can exceed 2000-3000 mm, with less than 100 mm
126 rainfall for up to three months in the year. The savannas have a wet season extending for 5-8 months, with an annual total of
127 1000-2000 mm (Bradley et al., 2011). Net radiation peaks in the winter months, due to the absence of cloud cover in the dry
128 season (Liu et al., 2018). Two major droughts occurred in the region during the study period, in 2010 and 2015 (Jiménez-Muñoz
129 et al., 2016; Marengo et al., 2011) and are of particular interest in this study.




Figure 2. Köppen-Geiger climate zones in the study area (Source: (Peel et al., 2007)).

2.2 ASCAT data

The Advanced Scatterometer (ASCAT) data were processed using the same procedure as Steele-Dunne et al. (2019). Metop-A ASCAT SZR Level 1b Fundamental Climate Data Record, resampled at a 12.5 km swath grid, were obtained from the EU-METSAT Data Centre for the period 2007 to 2016. Following the procedure described by Naeimi et al. (2009), the backscatter observations were resampled to a fixed Earth grid using a Hamming window function. The slope and curvature were calculated from the ASCAT backscatter observations using the method introduced by Metzler (2013) and described by Hahn et al. (2017). The ASCAT instrument on-board the Metop satellites has three antennas on either side, oriented at 45° (fore), 90° (mid) and 135° (aft) to the satellite track. As a result, three independent measurements of each location on the surface are obtained almost instantaneously. These so-called “backscatter triplets” (Hahn et al., 2017) are used to calculate an instantaneous backscatter slope, also known as the “local slope”:

$$\sigma'(\frac{\theta_{mid} - \theta_{a/f}}{2}) = \frac{\sigma_{mid}^{\circ}(\theta_{mid}) - \sigma_{a/f}^{\circ}(\theta_{a/f})}{\theta_{mid} - (\theta_{a/f})} \quad [dB/deg] \quad (2)$$

where *mid*, *a* and *f* indicate the backscatter measurements from the mid-, aft- and fore-beams respectively. Following the approach of Metzler (2013), an Epanechnikov kernel (with width $\lambda=21$) is used to weight the local slope values by their temporal distance from a given day of interest. This yields an estimate of slope and curvature for a given day, based on all local slope values within a 42-day window. For a more detailed description of their derivation, the reader is referred to Steele-Dunne et al. (2019).

(Anderson et al., 2011) showed a calibration accuracy of Metop ASCAT backscatter of 0.15 - 0.25 dB. However, radiometric accuracy is expected to be better (i.e. less noisy) over stable, homogeneous targets (e.g. evergreen rainforest). To reduce noise,

the backscatter data is averaged in space (over the ecoregions of interest) and/or time (to monthly or dekadal intervals). [The number of grid points averaged is provided in Tables S1 and S2. Data are available every 1-2 days \(Wagner et al., 2013\).](#) Observations from the descending and ascending overpasses are unlikely to occur on the same day. Hence, the σ_{40}° data were aggregated into 10-day intervals (dekads). Unless otherwise indicated, the analysis uses data from the descending pass only (~ 10 am). Diurnal differences ~~refers~~[refer](#) to the values from the descending overpass (~ 10 am) minus the values from the ascending overpass (~ 10 pm).

2.3 Water Dynamics data

Downwelling shortwave radiation at the surface and specific humidity were obtained from the Princeton meteorological dataset (Sheffield et al., 2006). These data have a $0.5^{\circ} \times 0.5^{\circ}$ daily resolution. Precipitation data were obtained from the Global Precipitation Climatology Product (GPCP) Precipitation Level 3 Monthly 0.5-Degree V3.0 beta dataset (Huffman et al., 2009). Precipitation, radiation and humidity are hypothesized to be the main atmospheric forcing for vegetation activity in the Amazon ((Nemani et al., 2003)). Therefore, these three forcings are compared to slope and curvature. As they are on similar temporal and spatial scales, quantitative comparisons are performed. Data from the Gravity Recovery and Climate Experiment (GRACE) mission were used to provide insight into terrestrial water storage variations (Landerer and Swenson, 2012; Swenson and Wahr, 2006). Here, we used the equivalent water thickness (EWT) from the GRACE Tellus dataset which is available at $1^{\circ} \times 1^{\circ}$, monthly resolution from the NASA JPL Physical Oceanography Distributed Active Archive Center (PO DAAC). These data give the relative change in EWT with respect to a baseline, the method of calculation for which is explained by Wahr et al. (1998). These data provide information on fluctuations in EWT on monthly to inter-annual timescales. [Note that EWT includes variations in all terrestrial water storage terms including groundwater ~~soil moisture, vegetation~~ and surface water, in addition to the variables of interest in this paper, namely soil moisture and vegetation. Furthermore, EWT is based on monthly data with a spatial resolution of hundreds of kilometers. Statistical comparisons between the EWT and ASCAT would be strongly influenced by the sensitivity of EWT to ground- and surface water and by artefacts of the difference in spatial and temporal scale between the two products.](#) Therefore, EWT is only qualitatively compared to backscatter, which is affected by soil moisture and vegetation.

Seasonal cycles were determined for precipitation, radiation, humidity, and EWT by averaging data from the entire study duration. Anomalies in precipitation during the drought years were also calculated (as drought year values minus climatology) to provide an indicator of the water stress against which to compare the backscatter, slope and curvature anomalies.

3 Results and Discussion

3.1 Seasonal Climatology

Figure 3 shows the mean and range of normalized backscatter (σ_{40}°), slope and curvature for the study period (2007-16). In general, the spatial patterns in the mean and range of all three quantities reflect the spatial patterns in land cover expected

179 from Fig. 1. It is striking that even the influence of the riverine network on the vegetation cover is discernible in the maps,
180 particularly that of the mean backscatter (Fig. 3(a)). Striping effects are visible in several of the maps, particularly that of the
181 range in curvature (Fig. 3(f)). This is due to the backscatter observations at the swath edges being available only at very high
182 or very low incidence angles, which skews the calculation of the slope and curvature. This effect is particularly noticeable in
183 forest regions where the natural dynamic range in both quantities is limited.

184 Mean backscatter is highest, with the least variability, in the evergreen forest regions (Fig. ??S1). Mean backscatter is 2-
185 2.5 dB lower in the savanna areas, but the range is up to 3 dB, compared to just 0.5 to 1 dB in the forest. The stability of the
186 forest is also apparent in the maps of slope and curvature. Though there is some variability among the forest ecoregions, the
187 most striking differences in slope and curvature are between the forest and savanna areas. Limited structural and water content
188 changes in the forest canopy result in a limited range of slope and curvature values in the forest ecoregions. The range of
189 both slope and curvature are highest in the Cerrado areas (Fig. ??S1). One interesting feature is the difference in mean slope
190 between the Guianan savanna (sG) in the north and the Cerrado (sC) region in the south. The Guianan savanna, with sparse
191 vegetation, has low mean slope values. The Cerrado, on the other hand, shows mean values higher than the evergreen forests.
192 This is unexpected since slope is generally considered a measure of “vegetation density”, and the evergreen forests are much
193 denser than savannas. This will be discussed in detail in Sect. 3.1.1. Seasonal flooding of the Marajó várzea (the seasonally
194 flooded forest) and Beni savannas ensure that both ecoregions have strong seasonal cycles in all three quantities. These will be
195 discussed separately in Sect. 3.1.2.

196 The mean seasonal cycles in backscatter for all ecoregions of interest are compared in Fig. 4 (a). This highlights the contrast
197 between the very stable evergreen forest regions and the flooded forest and savanna areas. The mean backscatter value is
198 high, with limited seasonal amplitude in the evergreen forest regions. Backscatter variations are so limited in these areas that
199 they have long been used as calibration targets for spaceborne radar (Birrer et al., 1982; Kennett and Li, 1989; Frison and
200 Mougin, 1996; Hawkins et al., 2000). In contrast, backscatter is generally low, but also exhibits strong seasonal variations in
201 the flooded forest and savanna areas. Figures 4 (b-f) show the seasonal variation in backscatter split out for five ecoregions
202 of interest, against the corresponding climatologies of precipitation and EWT. As the evergreen forest ecoregions showed
203 very similar climatologies, only the Jurua-Purus moist forest is shown as a separate plot. In all of the ecoregions, the maximum
204 backscatter occurs during the wet season, and a decrease in backscatter is observed during the dry season, though the amplitude
205 of the variations is obviously much smaller in the forest ecoregions. In each ecoregion, there is clear agreement between
206 the seasonality of EWT and backscatter. This indicates that backscatter is influenced by moisture availability in terms of
207 total terrestrial water storage, which includes groundwater storage. It is noteworthy that this temporal consistency between
208 backscatter and EWT is apparent for both forest (fC in Fig. 4 (b)) and the Guianan Savanna (sG in 4 (e)) despite the contrast
209 between almost zero (0.25 dB) variability in backscatter in fC and the 2.5 dB seasonal cycle in sG. Figure ??S2 shows
210 temporal correlation between backscatter and precipitation is low for all ecoregions. A strong negative correlation and strong
211 positive correlation are found with radiation and humidity for lags between -2 and 2 months, indicating that backscatter is
212 lowest during drier periods with higher radiation and lower specific humidity.




Figure 3. Mean and range of ASCAT normalized backscatter, slope and curvature in the study period (2007-16). Note that there are no data gaps, so white indicates that the quantity has a value equal to or less than the minimum value indicated on the colorbar.




Figure 4. Climatologies of backscatter for all ecoregions; five evergreen forest (dark green), flooded forest (cyan) and three savanna (light green) (a). Plot (b) to (f) show climatology of backscatter (green line) with precipitation (bars) and EWT (blue line) per ecoregion. Note the different y-axes and that only the Jurua-Purus moist forest (fC) is shown as it is similar to the other evergreen forests.

213 Figure 5 (a) summarizes the mean seasonal cycle in the slope for the ecoregions of interest. The difference between ecore-
214 gions is more pronounced than for backscatter. The seasonal cycle for the evergreen forest ecoregions are similar in magnitude
215 but there are minor differences in the timing of the peak. The differences between the savanna regions are more pronounced
216 than for backscatter. Significant differences can be seen in the mean slope value, as well as the amplitude and timing of the
217 seasonal cycle of slope values for each ecoregion of interest.

218 In Fig. 5(b-f), the seasonal cycle of slope in each ecoregion is compared to the corresponding cycles of radiation, specific
219 humidity and precipitation which drive photosynthetic activity in the region. Note again that only the Jurua-Purus moist forest is
220 shown as a separate plot. Furthermore, Fig. ??S2 illustrates the temporal correlation between slope and precipitation, radiation
221 and specific humidity. In the Jurua-Purus moist forests (Fig. 5(b)), the change in slope is one-tenth that observed in the other
222 ecoregions. The variations in radiation and specific humidity are also very limited. Nonetheless, the seasonal cycle of the slope
223 follows that of the radiation with a lag of about 30 days (Fig. ??S2, $R=0.75$ at lag -1). This can be explained by the fact
224 that the vegetation phenology in this tropical evergreen forest is driven by radiation (Romatschke and Houze Jr, 2013). The
225 photosynthetic capacity depends on the available solar energy (Borchert et al., 2015). Energy availability drives transpiration
226 and the accumulation of leafy biomass. This increases volume scattering from the canopy and therefore leads to an increase
227 in the slope. Similar results were observed for the other forest ecoregions. In the Marajo varzea flooded forest (Fig. 5(c)), the
228 variation in slope is much larger, and the seasonal cycle is clearly out of phase with that of the radiation. The seasonal variations
229 in slope in this ecoregion are dominated by the influence of surface flooding rather than vegetation water content variations
230 (Sect. 3.1.2).

231 In the Cerrado (Fig. 5(d)), there is a significant variation in specific humidity, and radiation as well as a strong seasonal cycle
232 in precipitation. The peak in slope occurs during the driest time of year, when radiation is at a maximum and specific humidity
233 and precipitation are at a minimum. Recall from Fig. 4, that this is also during the minimum EWT and backscatter period. This
234 is also illustrated in Fig. ??S2 where strong negative correlations are found between slope and humidity. Correlations between
235 slope and radiation are lower, and the highest correlation occurs at a lag of two months, i.e. slope leads radiation. Section
236 3.1.1 provides a detailed analysis of the vegetation types within the Cerrado ecoregion to better understand these variations.
237 The slope values in the Guianan Savanna (Fig. 5(e)) are the lowest observed in all ecoregions, and also have the smallest
238 variations among the non-forest cover types which are not strongly related to precipitation, radiation or specific humidity. This
239 is consistent with the relatively low, but stable vegetation density associated with grasslands (Steele-Dunne et al., 2019). In
240 the Beni Savanna (Fig. 5(f)), on the other hand, slope varies as much as in the Cerrado, and there is a very clear relationship
241 between the slope and the atmospheric forcing data (Fig. 5 (f)). The maximum slope occurs at the peak of precipitation, EWT
242 (from Fig. 4) and humidity. The minimum slope occurs during the dry season at the minimum in precipitation, humidity and
243 EWT. This is consistent with the interpretation of slope as an indicator of vegetation density as the vegetation cover in this
244 savanna changes dramatically in response to atmospheric forcing. This is also illustrated in Fig. ??S2, where high correlations
245 are observed between slope and humidity with small lags. The contrast in the seasonal cycles in slope in Fig. 5 reflect the
246 diversity of the vegetation cover types in the ecoregions and their varied response to moisture supply and demand.




Figure 5. Climatologies of slope for all ecoregions; five evergreen forest (dark green), flooded forest (cyan) and three savanna (light green) (a). Plot (b) to (f) show climatology of slope (green line) with precipitation (bars) and specific humidity (blue line) and radiation (red line) per ecoregion. Note the different y-axes and that only the Jurua-Purus moist forest (fC) is shown as it is similar to the other evergreen forests.

Figure 6 (a) shows the mean seasonal cycles in curvature for the regions of interest. The differences in the amplitudes of the seasonal cycles vary considerably among the regions. While the evergreen forests vary less than 0.0005 dB/deg^2 , variations in the wetland regions (Beni savanna and Marajó várzea) are an order of magnitude larger. Aside from the Guianan savanna, the timing of the seasonal cycle is similar across all ecoregions. Previous research has suggested that curvature is related to vegetation phenology and structure (Steele-Dunne et al., 2019). Since the vegetation phenology in much of the forested region is radiation-driven, we hypothesize that the curvature seasonality is related to the radiation and evaporative demand. In the Amazon rainforest, Borchert et al. (2015) observed that leaf flushing and flowering in adult trees of numerous species coincided with the rise and decline of insolation. Wagner et al. (2016) made a similar observation about leaf flushing and rising insolation in July, and also noted that the litterfall peak occurs when evaporative demand is highest and can persist through the dry season. Figure 6 (b) shows that although the changes in curvature are very small in the rainforest, the peak occurs in July on the rising limb of the radiation data, and when the specific humidity is near its minimum. Figures 6 (b-f) and Fig. [S2](#) show the strong correspondence between curvature and radiation (positive correlation at a lag of 2 months) and specific humidity (negative correlation with a lag of -1 month) and that the highest values of curvature generally correspond to lower humidity, higher solar radiation and lower precipitation. This suggests that higher values of curvature may be related to litterfall during periods of high evaporative demand. It is also noteworthy that the curvature values in the Guianan savanna (Fig. 6 (e)) are positive for much of the year, consistent with the dominance of grass cover in this region.

3.1.1 Cerrado

As described in Section 3, the Cerrado shows a peak in slope, which indicates increased volume scattering, at a time of low precipitation and humidity, maximum radiation and low backscatter. To better understand these variations backscatter, slope and curvature are analyzed [for the entire Cerrado region per land cover class. An overview of the number of used grid points per land cover can be found in \(Supplementary\) Table S2.](#) Figure 7 provides a detailed map of the Copernicus Global Land Service Land Cover within the Cerrado region Buchhorn et al. (2020). The dominant cover types are herbaceous cover and shrubland, with patches of cropland and forest. Figures 8 and 9 show the spatial patterns and boxplot per land cover type of mean, maximum and the DOY of the maximum for backscatter, slope and curvature. The mean backscatter varies between -13 and -7 dB and is highest for forest regions and lowest for croplands. The DOY for the maximum backscatter varies with latitude, from December to January in the southern region to April in the northern region. As expected, the highest backscatter corresponds with the months of highest precipitation and EWT, the minimum in backscatter corresponds with the months of lowest moisture availability (Fig. 4). The seasonal dynamics in backscatter are strongest in cropland. This may be related to the higher sensitivity to surface soil moisture in croplands and low backscatter may be related to dry surface soil conditions. The slope mean and maximum values show a decrease from shrubs to herbaceous to cropland, decreasing with vegetation density as expected. Forests are characterised by high mean and maximum slope values. The seasonal dynamics and DOY of the maximum slope vary strongly with land cover type. In croplands, the maximum slope, i.e. where volume scattering is highest, occurs between DOY 340-150. This corresponds to the highest precipitation and EWT, indicating increased vegetation density. In natural vegetation, such as herbaceous cover, shrubs and forests, the highest slope occurs between day 200 and 300 and




Figure 6. Climatologies of curvature for all ecoregions; five evergreen forest (dark green), flooded forest (cyan) and three savanna (light green) (a). Plot (b) to (f) show climatology of curvature (green line) with precipitation (bars) and specific humidity (blue line) and radiation (red line) per ecoregion. Note the different y-axes and that only the Jurua-Purus moist forest (fC) is shown as it is similar to the other evergreen forests.

281 coincides with the minimum in precipitation and EWT but with maximum radiation (Fig. 5). This is illustrated in Figure 10,
 282 where slope and radiation dynamics for different land cover classes are depicted. To exclude confounding effects due to
 283 heterogeneous land cover within ASCAT pixels, we used only pixels with a dominant land cover fraction of $> 80\%$. The slope
 284 dynamics in cropland are following the precipitation dynamics and have their peak during the wet season. Herbaceous cover
 285 shows two peaks in slope, one coinciding with the wet season at the beginning of the year, and a higher peak coinciding with
 286 the dry season and maximum in radiation. The increase in slope coincides with the onset of the increase in radiation. In shrubs
 287 and forests, slope starts to increase after the wet season, but before the increase in radiation (Fig. 10). This counter-intuitive
 288 behavior of the slope over natural vegetation can be explained by the variability in limiting factors to vegetation activity. Within
 289 the Cerrado region, vegetation can be moisture limited or energy limited (Nemani et al., 2003), depending on location and land
 290 cover type. Contrary to crops, natural vegetation types such as herbaceous vegetation, shrublands and forests have deeper root
 291 systems and they can tap into deeper water reservoirs. This enables them to increase photosynthesis and leaf development
 292 slightly before or at the onset of increasing radiation even though precipitation is at its minimum. The increase in vegetation
 293 activity will lead to increased volume scattering and a flatter backscatter over all incidence angle and subsequent higher slope.
 294 Chave et al. (2010) found that, among the tropical forest types in South America, the highest seasonality in litterfall was
 295 observed in "low" stature forests, such as those found in the Cerrado. They also cite Wright and Van Schaik (1994) to argue that
 296 seasonality of solar radiation rather than precipitation may be the most important trigger for leaf flushing and leaf abscission.
 297 Croplands and herbaceous vegetation show positive curvatures, whereas forests are characterised by negative curvatures with
 298 the maximum values occurring between DOY 200 and 300 across the Cerrado. The positive curvature for crops and herbaceous
 299 vegetation can be explained by the vertical structure of the vegetation.

300 3.1.2 Seasonal Flooding

301 Fig. 11 shows the striking effect of seasonal flooding on the incidence angle dependence of backscatter. This relationship was
 302 obtained using Equation 1 for a reference angle of 40° with the climatological mean values of σ_{40}° , slope and curvature for
 303 several days during the year. The flooded period is indicated in shades of blue. First, note that σ_{40}° is around 2 dB higher during
 304 the seasonal flooding. Under forest/woody vegetation, this is due to a combination of double bounce scattering between the
 305 surface and trunks, and multi-path scattering between the surface and the vegetation (Townsend, 2002).

306 Recall from Fig. 5 that the slope is slightly higher during this period as this multiple scattering is apparently slightly less
 307 sensitive to incidence angle than scattering from the vegetation during non-flooded period. However, the most noteworthy
 308 difference is in the curvature. In both ecoregions, the curvature changes considerably and even changes sign during the flooded
 309 period. This illustrates that the curvature includes useful information on changes in the scattering mechanisms, which are
 310 related to physical changes at the land surface.

311 3.2 Diurnal Differences

312 Figure 12 shows the mean diurnal differences for backscatter and EWT in the study area for alternate months in the year, where
 313 positive values indicate that values are higher during the descending (10 am) overpass than those from the ascending (10 pm)




Figure 7. Dominant land cover type (left) and fraction (right) derived from the Copernicus Global Land Service Land Cover (2015) for the Cerrado region Buchhorn et al. (2020)

overpass. The diurnal differences in backscatter are generally very small, with maximum values less than 0.15 dB. Although this is unquestionably close to the limits of the ASCAT sensor in terms of radiometric accuracy, these results are based on monthly averages, which is expected to reduce noise. Furthermore, there is a clear seasonal variation, broadly following that of EWT, demonstrating that patterns are likely not a result of noise.

For most of the domain, especially the evergreen forests, high values in EWT coincide with negative diurnal differences in backscatter and vice versa. During periods of maximum EWT, the backscatter is higher in the evening than in the morning. This is consistent with the finding that precipitation in tropical South America (since it is generally produced by convective systems) predominantly occurs in the late afternoons and evenings (Romatschke and Houze Jr, 2013). Hence, these higher backscatter values are due to the canopy being wetter in the evening.

During the drier periods (e.g. September (h, k) and November (i,l) in the south of the study area), backscatter is higher at 10 am than at 10 pm, consistent with the loss of moisture through transpiration during the day. In a light-limited evergreen forest such as the Amazon (rather than a water-limited forest), the canopy photosynthetic capacity seasonality is driven by radiation (Wagner et al., 2016). When the plants are phenologically active, they lose water during the daytime through transpiration resulting in lower evening backscatter values. The results in Fig. 12 are consistent with the findings of previous studies by Frolking et al. (2011) and Friesen et al. (2012) who also found the morning backscatter over Amazonia to be higher (on average) than the evening values due to higher water content in the vegetation. In the areas surrounding the evergreen forests, the patterns can be less straight-forward. Note, for example, that the diurnal difference in σ_{40}^0 in the Guianan savanna (sG) consistently has the opposite sign to that of the surrounding forest.




Figure 8. Mean, maximum and day of year of maximum for backscatter, slope and curvature over the Cerrado.




Figure 9. Time series averaged per land cover class and boxplots of mean, maximum and day of year of maximum for backscatter, slope and curvature over the Cerrado.




Figure 10. Seasonal cycle of slope and radiation per land cover class in the Cerrado region. Only ASCAT pixels in which the fraction of the dominant land cover type exceeds 80% are included.




Figure 11. Averaged backscatter as a function of incidence angle for several dates in the Marajo varzea (a) and Beni Savanna (b) ecoregions.




Figure 12. Maps of monthly mean diurnal differences in σ_{40}^0 (morning minus evening passes) and monthly mean Equivalent Water Thickness (EWT) from GRACE for different months in the year.

332 In Fig. 13, the seasonal cycle of the diurnal difference in σ_{40}^0 is compared to those of the radiation, precipitation and EWT
 333 for each of the ecoregions of interest. Figure 13(a) is indicative of the seasonal variations observed across the evergreen forest
 334 ecoregions. Note that the diurnal differences are very small (< 0.06 dB). Recall from Fig. 4, 5 and 6 that the backscatter, slope
 335 and curvature in these evergreen forests was essentially stable throughout the year, so even this small diurnal difference is
 336 noteworthy given the limited seasonal variation. As mentioned in the discussion of Fig. 12, evening values are higher than
 337 morning values during the EWT maximum and vice versa. Diurnal differences are larger in the Marajo varzea (Fig. 13(b)), but
 338 interpreting their seasonal variation is complicated by the seasonal inundation. In the Cerrado and Beni Savanna ecoregions
 339 (Fig. 13(c) and (e)), the diurnal differences in backscatter are almost twice as large as those observed in the evergreen forest
 340 regions. Morning values are up to 0.1 dB higher than evening values during the dry season due to loss of plant moisture during
 341 the day. Similar to the forest regions, evening backscatter values are higher during the rainy season. The Guianan savanna
 342 (Fig. 13(d)) is quite distinct in that morning backscatter is up to 0.15 dB higher than evening backscatter during the EWT
 343 and backscatter peak. One possible explanation for this unusual seasonal cycle is that it is related to a change in the relative
 344 dominance of the forests and grasslands in the backscatter signal. The transition from positive to negative curvature values
 345 during the EWT peak indicate an increased contribution from tree patches and shrubs during the wetter period. The higher
 346 backscatter in the morning may be due to water uptake in the trees during the night.




Figure 13. Seasonal cycle of diurnal difference in backscatter (black line), radiation (red line), EWT (blue line), and precipitation (bars) for different cover types. Green (yellow) fill indicates days in which backscatter is higher (lower) in the morning than in the evening.

3.3 The 2010 and 2015 droughts

During the study period (2007-16), two major droughts occurred in Amazonia, in 2010 and 2015. Figure 14 shows the spatial distribution of anomalies in σ_{40}^o , slope and curvature during the peak of the droughts from June to September 2010 and October 2010 to December 2015. Two regions of interest are indicated in the maps, the savanna Cerrado (sC) ecoregion and Southwest Amazon moist forests (fsW). The 2010 drought was most severe over southern and western Amazonia (Panisset et al., 2018). The 2015 drought was considered a “record-breaking” event with stronger warming than that seen in previous events (Jiménez-Muñoz et al., 2016). According to Panisset et al. (2018), there was a “pronounced lack of rainfall availability during late spring and early summer”. The 2015 drought was more widespread than the event in 2010, and strongest in eastern Amazonia.

355 Negative anomalies are observed in σ_{40}^o , especially in the southern regions and in the Cerrado in 2010 and in eastern regions
356 in 2015. Although anomalies are small over forests they are robust. Recall from section 2 that the backscatter noise level is
357 low over evergreen forests and noise is further reduced through temporal aggregation. Note that the most eastern part of the
358 Cerrado shows positive anomalies in 2010. The forests in fsW show minor negative anomalies (<0.1 dB) in σ_{40}^o in 2010 and
359 slightly stronger negative anomalies in 2015. Negative anomalies in backscatter from QSCAT were also observed during the
360 2005 drought (Saatchi et al. (2013); Frohling et al. (2017)). No significant spatial or temporal anomalies were observed in
361 the diurnal differences in backscatter during the drought years. The slope and curvature do not show clear spatial patterns in
362 anomalies during the 2010 drought, although the southern region shows slightly more positive anomalies. A clear positive
363 anomaly can be observed in the slope in eastern Amazonia in the 2015 drought. The curvature shows less clear patterns,
364 although a striping pattern can be seen, likely related to swaths.

365 Figure 15 shows the time series of anomalies in backscatter, slope and curvature for the moist forests in fsW and the Cerrado
366 region for the 2010 and 2015 drought. The backscatter, slope and curvature over the closed evergreen forest in fsW shows very
367 little variation (both in time and space) during both droughts. A slight increase up to 0.002 dB/deg in slope can be observed
368 during the peak of the 2015 drought. This demonstrates that the fsW forests are stable for satellite calibration. The Cerrado
369 shows varying responses depending on land cover type and are therefore investigated further. Negative anomalies in σ_{40}^o in
370 cropland and herbaceous land cover can be seen during both droughts. Especially during the 2010 drought the croplands in the
371 Cerrado are strongly affected, with a negative anomaly of >-1 dB for some pixels. During the more extensive drought in 2015,
372 σ_{40}^o in forest is also affected and negative anomalies up to -1.5 dB are observed. The slope shows minor positive anomalies
373 during the peak of the drought in 2010. In an analysis of drought impact on VOD over the forests in southern Amazonia,
374 Liu et al. observed similar positive anomalies in VOD from May to August during the 2010 drought. Negative anomalies in
375 VOD were only observed during later stages of the drought, from August to October. In 2010, negative slope anomalies in
376 the Cerrado are observed from October on. During the 2015 drought strong positive anomalies in slope and curvature are
377 present over the Cerrado especially in forests. Contrary to the drought of 2010, the peak of the 2015 drought occurs during
378 the precipitation season in the Cerrado. Normally the precipitation season is characterised by lower radiation, and the positive
379 anomalies in radiation during the drought might enhance vegetation growth. Note that the very small fluctuations in backscatter
380 observed in Figures 14 and 15 may only be scientifically evaluated in rainforest regions, where the spatio-temporal backscatter
381 dynamics (radiometric variations) are among the most stable in the world.

382 4 Conclusions

383 In this study, ASCAT backscatter, slope and curvature were analyzed in conjunction with meteorological data and terrestrial
384 water storage from GRACE in the Amazon region. Previous results, limited to grasslands, had suggested that the slope and
385 curvature contained useful information for monitoring vegetation water dynamics. However, the current study is the first to
386 attempt to explain the spatial and temporal variations in slope and curvature in terms of seasonal variations in moisture avail-




Figure 14. Spatial patterns in anomalies in backscatter, slope and curvature in response to the 2010 and 2015 droughts.




Figure 15. Time series of anomalies in backscatter, slope and curvature for moist forest (fsW) and Cerrado region. The shaded areas indicate the 5th and 95th percentile. The peak drought intervals (June-September 2010 and September-December 2015) are shown within dashed-dotted blue lines.

ability and demand. Furthermore, it confirms that the conclusions of Steele-Dunne et al. (2019) can be extended to a wide range of cover types.

Results show that the unique viewing geometry of ASCAT provides valuable insight into vegetation water dynamics across a diverse range of ecoregions. The timing of the seasonal cycle of normalized backscatter was consistent with that of GRACE EWT, with the maximum (minimum) normalized backscatter coinciding with the maximum (minimum) EWT in all ecoregions. Spatial patterns in mean and range of slope reflect the ecoregions within the study area. The seasonal cycle in slope was found to follow the moisture availability and demand indicated by meteorological data and their influence on phenology. A detailed analysis per land cover type over the Cerrado demonstrated this. Slope dynamics were concurrent with precipitation in croplands and herbaceous cover, although herbaceous cover showed a second peak coinciding with the maximum in radiation. Slope dynamics in shrubs and forest corresponded with radiation, although the onset in increasing slope preceded the onset of increasing radiation. This may be due to leaf flushing, but it is difficult to draw a firmer conclusion given the limited availability of ground data (Chave et al., 2010). While the mechanism driving these variations in slope may not be immediately clear, it is important to note that there are open questions around the process of litterfall and its relation to precipitation and radiation in general. A recent study from Hashimoto et al. (2021) demonstrated that the temporal density of optical data from the Advanced Baseline Imager (ABI) onboard the Geostationary Operational Environmental Satellite 16 (GOES-16) yields unprecedented detail on the seasonality of NDVI and LAI in the evergreen Amazon forests. A comparison of ASCAT slope and curvature and ABI data may yield additional insight into the connection between slope, curvature and litterfall in the various ecoregions of our study area. Consistent with the findings of Steele-Dunne et al. (2019) in a study limited to grasslands, variations in curvature seem to be related to phenological change. The highest values of curvature coincide with periods of high evaporative demand (e.g. high radiation, lower humidity and lower precipitation). This suggests a link between curvature and phenological changes such as leaf flushing and litterfall. For example, the curvature peak in July in the rainforest occurs during rising insolation, and coincides with leaf flushing. Areas affected by seasonal flooding exhibited dramatic changes in both backscatter and curvature due to a suspected increase in multiple scattering between water on the surface and the vegetation.

Diurnal variations (i.e. the difference between morning and evening overpasses) were generally small, particularly in the evergreen forests. Nonetheless, their relation to the timing of precipitation highlights the importance of overpass time in using microwave observations for vegetation monitoring. Diurnal differences in backscatter during the dry season are dominated by transpiration losses. Long-term monitoring of these diurnal differences could provide insight into moisture availability and its influence on transpiration and vegetation functioning (Konings et al., 2021). Consistent with previous studies on the effect of drought on the backscatter signal over the Amazon forests (Frolking et al., 2011; Saatchi et al., 2013), a negative anomaly in backscatter was observed during the 2010 and 2015 drought, although this was minor for the moist forests, strong anomalies were observed in the Cerrado. The slope showed positive anomalies during the drought events in the Cerrado, similar to positive anomalies in VOD over forests observed by Liu et al. who attributed this to enhanced canopy growth due to increased radiation. Persistent positive anomalies in radiation were observed over the Cerrado, especially in 2015. The analysis confirms the confounding effects of mechanisms driving variation in slope in these regions.

421 For regions with non-closed-canopy conditions and significant soil contribution, the water sensitivity of the slope and
422 curvature may be influenced, or even dominated by soil moisture dynamics (Greimeister-Pfeil et al., 2022). Furthermore,
423 the various storage compartments (soil, vegetation) are linked by the soil-plant-atmosphere system. An improved physical
424 understanding of the influence of both soil and vegetation on slope and curvature is essential. Future research should also
425 include forward electro-magnetic modelling of multi-angular backscatter (i.e slope and curvature) to improve our understanding
426 of how they relate to vegetation water and biomass variations as well as soil moisture.

427 The improved understanding of slope and curvature gained in this study is valuable in terms of our ability to use ASCAT
428 for vegetation monitoring, and specifically for vegetation water dynamics. ~~The slope and curvature are used to produce the~~
429 ~~VOD from ASCAT (Vreugdenhil et al., 2016)~~Slope and curvature may be influenced by the number and distribution of the
430 scatterers, and their dielectric properties, all of which influence the optical depth i.e. the attenuation of the signal by the
431 vegetation.Our improved understanding of the slope and curvature and how they are affected by vegetation structure and
432 water content, and interactions between the soil and vegetation is essential to improve our ability to interpret and optimally
433 use VOD derived from ASCAT. Therefore, this research contributes directly to the continued development of the ASCAT
434 VOD products. For example, it provides further insights in the VOD calculated from ASCAT by Vreugdenhil et al. (2016)
435 , where the main temporal dynamics stem from the slope and curvature. Furthermore, the fact that the slope and curvature
436 themselves reveal different aspects of the vegetation response to the balance between moisture availability and demand means
437 they are potentially useful low-level observables, i.e. they are obtained with minimal processing, and avoid the assumptions and
438 simplifications required to retrieve geophysical variables. The results of this study suggest that their information content can
439 be directly exploited to monitor vegetation water dynamics. The current study was performed over different land cover types,
440 demonstrating the potential to study vegetation water dynamics with these observables over different regions. However this
441 research also confirms the need for further research to overcome the limited understanding of the spatio-temporal dynamics
442 of slope compared to environmental drivers and effects in structure of vegetation. A lot of our understanding of the incidence
443 angle dependence of backscatter is based on experiments with tower-based or airborne radar systems conducted in the 1970s
444 to 1990s (e.g. Ulaby (1975); Ferrazzoli et al. (1992)) to optimize the design of spaceborne radar systems. However, these
445 experiments were generally focused on classification, soil moisture or LAI/biomass retrieval. Radar data were limited in space
446 and/or time, and water dynamics (beyond soil moisture) were often not considered. Recent studies have focused on the relation
447 between water dynamics in vegetation and tower-based radar backscatter response(e.g. (Vermunt et al., 2020; Khabbazan et al.,
448 2022)), but not at the slope of the backscatter incidence angle relationship. So, in any first-order ground validation, we advocate
449 the inclusion of incidence angle dependence. Nonetheless, field-based experimental campaigns have the disadvantage that they
450 are very localized. Thus, studies like the one presented here, to explore ASCAT dynamic vegetation parameters and explain
451 the variations in terms of modeled or observed geophysical variables are equally valuable and needed, because they allow us
452 to study a wide range of cover and climate types and the impact of events such as drought. Based on this, and considering
453 the planned SCA instrument on Metop-SG, incidence angle variations should be studied in more detail and be considered as
454 a potentially valuable source of useful information. Ongoing research is focused on using data-driven and radiative transfer
455 modeling approaches to investigate the sensitivity of slope and curvature to physical changes at the land surface including also

456 different regions and cover types. ~~Mean (top row) and range (bottom row) for backscatter, slope and curvature for different~~
457 ~~ecoregions. Correlation between backscatter (top row), slope (middle row) and curvature (bottom row) with precipitation,~~
458 ~~radiation and humidity for the different ecoregions for different lag times.~~

459 *Author contributions.* AP, SSD and MV were responsible for the conceptualization, methodology, formal analysis, investigation, visual-
460 ization and writing (original draft preparation). SH provided resources (ASCAT data). RO contributed to the investigation. SSD and MV
461 provided supervision. All authors contributed to writing (review and editing).

462 *Competing interests.* The authors declare that no competing interests are present.

463 *Acknowledgements.* Susan Steele-Dunne was supported by The Netherlands Organization for Scientific Research (NWO) User Support
464 Programme Space Research (Project ALWGO.2018.036 - 'A new perspective on global vegetation water dynamics from radar satellite
465 data') and NWO Vidi Grant 14126. Mariette Vreugdenhil was supported by ESA's Living Planet Fellowship SHRED (contract number
466 4000125441/18/I-NS).

- 468 Andela, N., Liu, Y. Y., van Dijk, A., de Jeu, R. A. M., and McVicar, T. R.: Global changes in dryland vegetation dynamics (1988-2008)
 469 assessed by satellite remote sensing: comparing a new passive microwave vegetation density record with reflective greenness data, *Bio-*
 470 *geosciences*, 10, 6657, <http://search.proquest.com/openview/75d8704b5c57105c1e46b3b2d6ee313f/1?pq-origsite=gscholar>, 2013.
- 471 Anderson, C., Figa, J., Bonekamp, H., Wilson, J. J. W., Verspeek, J., Stoffelen, A., and Portabella, M.: Validation of Backscat-
 472 ter Measurements from the Advanced Scatterometer on MetOp-A, *Journal of Atmospheric and Oceanic Technology*, 29, 77 – 88,
 473 <https://doi.org/10.1175/JTECH-D-11-00020.1>, 2011.
- 474 Attema, E. P.: The active microwave instrument on-board the ERS-1 satellite, *Proceedings of the IEEE*, 79, 791–799, publisher: IEEE, 1991.
- 475 Birrer, I., Bracalente, E., Dome, G., Sweet, J., and Berthold, G.: σ signature of the Amazon rain forest obtained from the SeaSat scatterometer,
 476 *IEEE Transactions on Geoscience and Remote Sensing*, GE-20, 11–17, 1982.
- 477 Borchert, R., Calle, Z., Strahler, A. H., Baertschi, A., Magill, R. E., Broadhead, J. S., Kamau, J., Njoroge, J., and Muthuri, C.: Insolation and
 478 photoperiodic control of tree development near the equator, *New Phytologist*, 205, 7–13, 2015.
- 479 Bradley, A. V., Gerard, F. F., Barbier, N., Weedon, G. P., Anderson, L. O., Huntingford, C., Aragão, L. E., Zelazowski, P., and Arai, E.:
 480 Relationships between phenology, radiation and precipitation in the Amazon region, *Global Change Biology*, 17, 2245–2260, 2011.
- 481 Buchhorn, M., Smets, B., Bertels, L., Roo, B. D., Lesiv, M., Tsendbazar, N.-E., Herold, M., and Fritz, S.: Copernicus Global Land Service:
 482 Land Cover 100m: collection 3: epoch 2015: Globe, <https://doi.org/10.5281/zenodo.3939038>, type: dataset, 2020.
- 483 Camarão, A., Júnior, L., Dutra, S., et al.: Flooded pasture production for grazing buffalo in the brazilian Amazon region., in: Embrapa
 484 Amazônia Oriental-Artigo em anais de congresso (ALICE), pp. 68–82, In: Buffalo Symposium Of Americas, 1., 2002, Belém. Proceedings
 485 of the ..., 2002.
- 486 Chaparro, D., Duveiller, G., Piles, M., Cescatti, A., Vall-llossera, M., Camps, A., and Entekhabi, D.: Sensitivity of L-band vegetation optical
 487 depth to carbon stocks in tropical forests: a comparison to higher frequencies and optical indices, *Remote Sensing of Environment*, 232,
 488 111 303, <https://doi.org/10.1016/j.rse.2019.111303>, 2019.
- 489 Chave, J., Navarrete, D., Almeida, S., Álvarez, E., Aragão, L. E. O. C., Bonal, D., Châtelet, P., Silva-Espejo, J. E., Goret, J.-Y., von Hilde-
 490 brand, P., Jiménez, E., Patiño, S., Peñuela, M. C., Phillips, O. L., Stevenson, P., and Malhi, Y.: Regional and seasonal patterns of litterfall
 491 in tropical South America, *Biogeosciences*, 7, 43–55, <https://doi.org/10.5194/bg-7-43-2010>, publisher: Copernicus GmbH, 2010.
- 492 De Jeu, R. A.: Retrieval of land surface parameters using passive microwave remote sensing, PhD diss., Vrije Universiteit Amsterdam, 2003.
- 493 Eiten, G.: The cerrado vegetation of Brazil, *The Botanical Review*, 38, 201–341, 1972.
- 494 Fernandez-Moran, R., Al-Yaari, A., Mialon, A., Mahmoodi, A., Al Bitar, A., De Lannoy, G., Rodriguez-Fernandez, N., Lopez-Baeza, E.,
 495 Kerr, Y., and Wigneron, J.-P.: SMOS-IC: An Alternative SMOS Soil Moisture and Vegetation Optical Depth Product, *Remote Sensing*, 9,
 496 457, <https://doi.org/10.3390/rs9050457>, 2017.
- 497 Ferrazzoli, P., Paloscia, S., Pampaloni, P., Schiavon, G., Solimini, D., and Coppo, P.: Sensitivity of microwave measurements to vegetation
 498 biomass and soil moisture content: a case study, *IEEE Transactions on Geoscience and Remote Sensing*, 30, 750–756, [http://ieeexplore.](http://ieeexplore.ieee.org/xpls/abs_all.jsp?arnumber=158869)
 499 [ieee.org/xpls/abs_all.jsp?arnumber=158869](http://ieeexplore.ieee.org/xpls/abs_all.jsp?arnumber=158869), 1992.
- 500 Figa-Saldaña, J., Wilson, J. J., Attema, E., Gelsthorpe, R., Drinkwater, M. R., and Stoffelen, A.: The advanced scatterometer (ASCAT) on
 501 the meteorological operational (MetOp) platform: A follow on for European wind scatterometers, *Canadian Journal of Remote Sensing*,
 502 28, 404–412, publisher: Taylor & Francis, 2002.

503 Forkel, M., Andela, N., Harrison, S. P., Lasslop, G., van Marle, M., Chuvieco, E., Dorigo, W., Forrest, M., Hantson, S., Heil, A., Li, F.,
504 Melton, J., Sitch, S., Yue, C., and Arneeth, A.: Emergent relationships with respect to burned area in global satellite observations and
505 fire-enabled vegetation models, *Biogeosciences*, 16, 57–76, <https://doi.org/10.5194/bg-16-57-2019>, publisher: Copernicus GmbH, 2019.

506 Friesen, J., Steele-Dunne, S. C., and van de Giesen, N.: Diurnal differences in global ERS scatterometer backscatter observations of the land
507 surface, *IEEE Transactions on Geoscience and Remote Sensing*, 50, 2595–2602, 2012.

508 Frison, P.-L. and Mougin, E.: Use of ERS-1 wind scatterometer data over land surfaces, *IEEE Transactions on Geoscience and Remote*
509 *Sensing*, 34, 550–560, 1996.

510 Frison, P. L., Mougin, E., and Hiernaux, P.: Observations and interpretation of seasonal ERS-1 wind scatterometer data over northern Sahel
511 (Mali), *Remote Sensing of Environment*, 63, 233–242, publisher: Elsevier, 1998.

512 Frolking, S., Milliman, T., Palace, M., Wisser, D., Lammers, R., and Fahnestock, M.: Tropical forest backscatter anomaly evident in SeaWinds
513 scatterometer morning overpass data during 2005 drought in Amazonia, *Remote Sensing of Environment*, 115, 897–907, 2011.

514 Frolking, S., Hagen, S., Braswell, B., Milliman, T., Herrick, C., Peterson, S., Roberts, D., Keller, M., and Palace, M.: Evaluating multiple
515 causes of persistent low microwave backscatter from Amazon forests after the 2005 drought, *PloS one*, 12, e0183 308, 2017.

516 Greimeister-Pfeil, I., Wagner, W., Quast, R., Hahn, S., Steele-Dunne, S., and Vreugdenhil, M.: Analysis of short-term soil moisture effects
517 on the ASCAT backscatter-incidence angle dependence, *Science of Remote Sensing*, P, pp (Accepted), 2022.

518 Hahn, S., Reimer, C., Vreugdenhil, M., Melzer, T., and Wagner, W.: Dynamic characterization of the incidence angle dependence of backscat-
519 ter using metop ASCAT, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 10, 2348–2359, 2017.

520 Hamilton, S. K., Sippel, S. J., and Melack, J. M.: Seasonal inundation patterns in two large savanna floodplains of South America: the Llanos
521 de Moxos (Bolivia) and the Llanos del Orinoco (Venezuela and Colombia), *Hydrological Processes*, 18, 2103–2116, 2004.

522 Hashimoto, H., Wang, W., Dungan, J. L., Li, S., Michaelis, A. R., Takenaka, H., Higuchi, A., Myneni, R. B., and Nemani, R. R.: New
523 generation geostationary satellite observations support seasonality in greenness of the Amazon evergreen forests, *Nature Communications*,
524 12, 684, <https://doi.org/10.1038/s41467-021-20994-y>, number: 1 Publisher: Nature Publishing Group, 2021.

525 Hawkins, R., Attema, E., Crapolicchio, R., Lecomte, P., Closa, J., Meadows, P., and Srivastava, S.: Stability of Amazon Backscatter at C-
526 Band: Spaceborne Results from ERS-1/2 and RADARSAT-1., in: *SAR workshop: CEOS Committee on Earth Observation Satellites*, vol.
527 450, p. 99, 2000.

528 Hordijk, I., Meijer, F., Nissen, E., Boorsma, T., and Poorter, L.: Cattle affect regeneration of the palm species *Attalea princeps* in a Bolivian
529 forest–savanna mosaic, *Biotropica*, 51, 28–38, 2019.

530 Huffman, G. J., Adler, R. F., Bolvin, D. T., and Gu, G.: Improving the global precipitation record: GPCP Version 2.1, *Geophysical Research*
531 *Letters*, 36, <https://doi.org/https://doi.org/10.1029/2009GL040000>, 2009.

532 Jackson, T. J., Schmugge, T. J., and Wang, J. R.: Passive microwave sensing of soil moisture under vegetation canopies, *Water Resources*
533 *Research*, 18, 1137–1142, <https://doi.org/10.1029/WR018i004p01137>, 1982.

534 Jarlan, L., Mougin, E., Frison, P. L., Mazzega, P., and Hiernaux, P.: Analysis of ERS wind scatterometer time series over Sahel (Mali),
535 *Remote Sensing of Environment*, 81, 404–415, [https://doi.org/10.1016/S0034-4257\(02\)00015-9](https://doi.org/10.1016/S0034-4257(02)00015-9), 2002.

536 Jiménez-Muñoz, J. C., Mattar, C., Barichivich, J., Santamaría-Artigas, A., Takahashi, K., Malhi, Y., Sobrino, J. A., and Van Der Schrier, G.:
537 Record-breaking warming and extreme drought in the Amazon rainforest during the course of El Niño 2015–2016, *Scientific reports*, 6,
538 33 130, 2016.

539 Kennett, R. G. and Li, F. K.: Seasat over-land scatterometer data. II. Selection of extended area and land-target sites for the calibration of
540 spaceborne scatterometers, *IEEE Transactions on Geoscience and Remote Sensing*, 27, 779–788, 1989.

541 Khabbazan, S., Steele-Dunne, S. C., Vermunt, P., Judge, J., Vreugdenhil, M., and Gao, G.: The influence of surface canopy water on the
542 relationship between L-band backscatter and biophysical variables in agricultural monitoring, *Remote Sensing of Environment*, 268,
543 112 789, <https://doi.org/10.1016/j.rse.2021.112789>, 2022.

544 Konings, A. G., Piles, M., Rötzer, K., McColl, K. A., Chan, S. K., and Entekhabi, D.: Vegetation optical depth and scattering
545 albedo retrieval using time series of dual-polarized L-band radiometer observations, *Remote Sensing of Environment*, 172, 178–189,
546 <https://doi.org/10.1016/j.rse.2015.11.009>, 2016.

547 Konings, A. G., Rao, K., and Steele-Dunne, S. C.: Macro to micro: microwave remote sensing of plant water con-
548 tent for physiology and ecology, *New Phytologist*, 223, 1166–1172, <https://doi.org/https://doi.org/10.1111/nph.15808>, _eprint:
549 <https://nph.onlinelibrary.wiley.com/doi/pdf/10.1111/nph.15808>, 2019.

550 Konings, A. G., Saatchi, S. S., Frankenberg, C., Keller, M., Leshyk, V., Anderegg, W. R., Humphrey, V., Matheny, A. M., Trugman, A.,
551 Sack, L., et al.: Detecting forest response to droughts with global observations of vegetation water content, *Global change biology*, 27,
552 6005–6024, 2021.

553 Landerer, F. W. and Swenson, S.: Accuracy of scaled GRACE terrestrial water storage estimates, *Water resources research*, 48, 2012.

554 Liu, Y. Y., de Jeu, R. A., McCabe, M. F., Evans, J. P., and van Dijk, A. I.: Global long-term passive microwave satellite-based retrievals of
555 vegetation optical depth, *Geophysical Research Letters*, 38, <http://onlinelibrary.wiley.com/doi/10.1029/2011GL048684/full>, 2011.

556 Liu, Y. Y., Dijk, A. I., McCabe, M. F., Evans, J. P., and Jeu, R. A.: Global vegetation biomass change (1988–2008) and attribution to
557 environmental and human drivers, *Global ecology and biogeography*, 22, 692–705, [http://onlinelibrary.wiley.com/doi/10.1111/geb.12024/](http://onlinelibrary.wiley.com/doi/10.1111/geb.12024/full)
558 full, 2013.

559 Liu, Y. Y., Van Dijk, A. I., De Jeu, R. A., Canadell, J. G., McCabe, M. F., Evans, J. P., and Wang, G.: Recent reversal in loss of global
560 terrestrial biomass, *Nature Climate Change*, 5, 470–474, <http://www.nature.com/nclimate/journal/v5/n5/abs/nclimate2581.html>, 2015.

561 Liu, Y. Y., van Dijk, A. I., Miralles, D. G., McCabe, M. F., Evans, J. P., de Jeu, R. A., Gentine, P., Huete, A., Parinussa, R. M., Wang, L.,
562 et al.: Enhanced canopy growth precedes senescence in 2005 and 2010 Amazonian droughts, *Remote sensing of environment*, 211, 26–37,
563 2018.

564 Marengo, J. A., Tomasella, J., Alves, L. M., Soares, W. R., and Rodriguez, D. A.: The drought of 2010 in the context of historical droughts
565 in the Amazon region, *Geophysical Research Letters*, 38, 2011.

566 McNairn, H., Van der Sanden, J. J., Brown, R. J., and Ellis, J.: The potential of RADARSAT-2 for crop mapping and assessing crop condition,
567 in: *Second International Conference on Geospatial Information in Agriculture and Forestry*, Lake Buena Vista, FL, 2000.

568 Metzler, T.: Vegetation Modelling in WARP 6.0, in: *Proc. EUMETSAT Meteorological Satellite Conf.*, pp. 1–7, Vienna, Austria, 2013.

569 Moesinger, L., Dorigo, W., de Jeu, R., van der Schalie, R., Scanlon, T., Teubner, I., and Forkel, M.: The global long-term microwave
570 Vegetation Optical Depth Climate Archive (VODCA), *Earth System Science Data*, 12, 177–177, 2020.

571 Naeimi, V., Scipal, K., Bartalis, Z., Hasenauer, S., and Wagner, W.: An Improved Soil Moisture Retrieval Algorithm for
572 ERS and METOP Scatterometer Observations, *IEEE Transactions on Geoscience and Remote Sensing*, 47, 1999–2013,
573 <https://doi.org/10.1109/TGRS.2008.2011617>, 2009.

574 Nemani, R. R., Keeling, C. D., Hashimoto, H., Jolly, W. M., Piper, S. C., Tucker, C. J., Myneni, R. B., and Running, S. W.: Climate-driven
575 increases in global terrestrial net primary production from 1982 to 1999, *Science*, 300, 1560–1563, 2003.

576 Oliveira, R., Bezerra, L., Davidson, E., Pinto, F., Klink, C., Nepstad, D., and Moreira, A.: Deep root function in soil water dynamics in
577 cerrado savannas of central Brazil, *Functional Ecology*, 19, 574–581, 2005.

578 Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V., Underwood, E. C., D'amico, J. A., Itoua, I., Strand,
579 H. E., Morrison, J. C., et al.: Terrestrial Ecoregions of the World: A New Map of Life on Earth A new global map of terrestrial ecoregions
580 provides an innovative tool for conserving biodiversity, *BioScience*, 51, 933–938, 2001.

581 Owe, M., de Jeu, R., and Walker, J.: A methodology for surface soil moisture and vegetation optical depth retrieval using the microwave
582 polarization difference index, *IEEE Transactions on Geoscience and Remote Sensing*, 39, 1643–1654, [http://ieeexplore.ieee.org/xpls/abs_](http://ieeexplore.ieee.org/xpls/abs_all.jsp?arnumber=942542)
583 [all.jsp?arnumber=942542](http://ieeexplore.ieee.org/xpls/abs_all.jsp?arnumber=942542), 2001.

584 Panisset, J. S., Libonati, R., Gouveia, C. M. P., Machado-Silva, F., França, D. A., França, J. R. A., and Peres, L. F.: Contrasting patterns of
585 the extreme drought episodes of 2005, 2010 and 2015 in the Amazon Basin, *International Journal of Climatology*, 38, 1096–1104, 2018.

586 Peel, M. C., Finlayson, B. L., and McMahon, T. A.: Updated world map of the Köppen-Geiger climate classification, *Hydrology and Earth*
587 *System Sciences Discussions*, 4, 439–473, 2007.

588 Pfeil, I., Wagner, W., Forkel, M., Dorigo, W., and Vreugdenhil, M.: Does ASCAT observe the spring reactivation in temperate deciduous
589 broadleaf forests?, *Remote Sensing of Environment*, 250, 112 042, 2020.

590 Rao, K., Anderegg, W. R. L., Sala, A., Martínez-Vilalta, J., and Konings, A. G.: Satellite-based vegetation optical depth as an indicator of
591 drought-driven tree mortality, *Remote Sensing of Environment*, 227, 125–136, <https://doi.org/10.1016/j.rse.2019.03.026>, 2019.

592 Romatschke, U. and Houze Jr, R. A.: Characteristics of precipitating convective systems accounting for the summer rainfall of tropical and
593 subtropical South America, *Journal of Hydrometeorology*, 14, 25–46, 2013.

594 Saatchi, S., Asefi-Najafabady, S., Malhi, Y., Aragão, L. E., Anderson, L. O., Myneni, R. B., and Nemani, R.: Persistent effects of a severe
595 drought on Amazonian forest canopy, *Proceedings of the National Academy of Sciences*, 110, 565–570, 2013.

596 Schroeder, R., McDonald, K. C., Azarderakhsh, M., and Zimmermann, R.: ASCAT MetOp-A diurnal backscatter observations of recent
597 vegetation drought patterns over the contiguous US: An assessment of spatial extent and relationship with precipitation and crop yield,
598 *Remote sensing of environment*, 177, 153–159, publisher: Elsevier, 2016.

599 Sheffield, J., Goteti, G., and Wood, E. F.: Development of a 50-year high-resolution global dataset of meteorological forcings for land surface
600 modeling, *Journal of climate*, 19, 3088–3111, 2006.

601 Soares, B. S., Nepstad, D. C., Curran, L. M., Cerqueira, G. C., Garcia, R. A., Ramos, C. A., Voll, E., McDonald, A., Lefebvre, P., and
602 Schlesinger, P.: Modelling conservation in the Amazon basin, *Nature*, 440, 520–523, 2006.

603 Steele-Dunne, S. C., Friesen, J., and van de Giesen, N.: Using diurnal variation in backscatter to detect vegetation water stress, *IEEE*
604 *Transactions on Geoscience and Remote Sensing*, 50, 2618–2629, 2012.

605 Steele-Dunne, S. C., McNairn, H., Monsivais-Huertero, A., Judge, J., Liu, P.-W., and Papathanassiou, K.: Radar remote sensing of agricultural
606 canopies: A review, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 10, 2249–2273, 2017.

607 Steele-Dunne, S. C., Hahn, S., Wagner, W., and Vreugdenhil, M.: Investigating vegetation water dynamics and drought using Metop ASCAT
608 over the North American Grasslands, *Remote Sensing of Environment*, 224, 219–235, 2019.

609 Stoffelen, A., Aaboe, S., Calvet, J.-C., Cotton, J., De Chiara, G., Saldana, J. F., Mouche, A. A., Portabella, M., Scipal, K., and Wagner,
610 W.: Scientific developments and the EPS-SG scatterometer, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote*
611 *Sensing*, 10, 2086–2097, 2017.

612 Swenson, S. and Wahr, J.: Post-processing removal of correlated errors in GRACE data, *Geophysical Research Letters*, 33, 2006.

613 Templ, B., Koch, E., Bolmgren, K., Ungersböck, M., Paul, A., Scheifinger, H., Busto, M., Chmielewski, F.-M., Hájková, L., Hodzić, S., et al.:
614 Pan European Phenological database (PEP725): a single point of access for European data, *International journal of biometeorology*, 62,
615 1109–1113, 2018.

616 Teubner, I., Forkel, M., Jung, M., Liu, Y., Miralles, D., Parinussa, R., van der Schalie, R., Vreugdenhil, M., Schwalm, C., Tramontana, G.,
617 Camps-Valls, G., and Dorigo, W.: Assessing the relationship between microwave vegetation optical depth and gross primary production,
618 *International Journal of Applied Earth Observation and Geoinformation*, 65, 79–91, <https://doi.org/10.1016/j.jag.2017.10.006>, 2018.

619 Teubner, I., Forkel, M., Camps-Valls, G., Jung, M., Miralles, D., Tramontana, G., van der Schalie, R., Vreugdenhil, M., Möisinger, L., and
620 Dorigo, W.: A carbon sink-driven approach to estimate gross primary production from microwave satellite observations, *Remote Sensing*
621 *of Environment*, 229, 100–113, <https://doi.org/10.1016/j.rse.2019.04.022>, 2019.

622 Tian, F., Brandt, M., Liu, Y. Y., Verger, A., Tagesson, T., Diouf, A. A., Rasmussen, K., Mbow, C., Wang, Y., and Fensholt, R.: Remote sensing
623 of vegetation dynamics in drylands: Evaluating vegetation optical depth (VOD) using AVHRR NDVI and in situ green biomass data over
624 West African Sahel, *Remote Sensing of Environment*, 177, 265–276, <https://doi.org/10.1016/j.rse.2016.02.056>, 2016.

625 Townsend, P.: Relationships between forest structure and the detection of flood inundation in forested wetlands using C-band SAR, *Interna-*
626 *tional Journal of Remote Sensing*, 23, 443–460, 2002.

627 Ulaby, F.: Radar response to vegetation, *IEEE Transactions on Antennas and Propagation*, 23, 36–45,
628 <https://doi.org/10.1109/TAP.1975.1140999>, 1975.

629 Vermunt, P. C., Khabbazan, S., Steele-Dunne, S. C., Judge, J., Monsivais-Huertero, A., Guerriero, L., and Liu, P.-W.: Response of Subdaily
630 L-Band Backscatter to Internal and Surface Canopy Water Dynamics, *IEEE Transactions on Geoscience and Remote Sensing*, pp. 1–16,
631 <https://doi.org/10.1109/TGRS.2020.3035881>, conference Name: IEEE Transactions on Geoscience and Remote Sensing, 2020.

632 Vreugdenhil, M., Dorigo, W. A., Wagner, W., De Jeu, R. A., Hahn, S., and Van Marle, M. J.: Analyzing the vegetation parameterization in
633 the TU-Wien ASCAT soil moisture retrieval, *IEEE Transactions on Geoscience and Remote Sensing*, 54, 3513–3531, 2016.

634 Wagner, F. H., Hérault, B., Bonal, D., Stahl, C., Anderson, L. O., Baker, T. R., Sebastian Becker, G., Beeckman, H., D Souza, B., Cesar Boto-
635 sso, P., et al.: Climate seasonality limits leaf carbon assimilation and wood productivity in tropical forests, *Biogeosciences*, 13, 2537–2562,
636 2016.

637 Wagner, W., Lemoine, G., Borgeaud, M., and Rott, H.: A study of vegetation cover effects on ERS scatterometer data, *IEEE Transactions on*
638 *Geoscience and Remote Sensing*, 37, 938–948, 1999.

639 Wagner, W., Hahn, S., Kidd, R., Melzer, T., Bartalis, Z., Hasenauer, S., Figa-Saldaña, J., de Rosnay, P., Jann, A., Schneider, S., Komma,
640 J., Kubu, G., Brugger, K., Aubrecht, C., Züger, J., Gangkofner, U., Kienberger, S., Brocca, L., Wang, Y., Blöschl, G., Eitzinger, J.,
641 Steinnocher, K., Zeil, P., and Rubel, F.: The ASCAT Soil Moisture Product: A Review of Its Specifications, Validation Results, and
642 Emerging Applications, *Meteorologische Zeitschrift*, 22, 5–33, <https://doi.org/10.1127/0941-2948/2013/0399>, 2013.

643 Wahr, J., Molenaar, M., and Bryan, F.: Time variability of the Earth’s gravity field: Hydrological and oceanic effects and their possible
644 detection using GRACE, *Journal of Geophysical Research: Solid Earth*, 103, 30 205–30 229, 1998.

645 Wismann, V. R., Boehnke, K., and Schmulilius, C.: Monitoring ecological dynamics in Africa with the ERS-1 scatterometer, in: 1995 Inter-
646 national Geoscience and Remote Sensing Symposium, IGARSS’95. Quantitative Remote Sensing for Science and Applications, vol. 2,
647 pp. 1523–1525, IEEE, 1995.

648 Woodhouse, I. H., van der Sanden, J. J., and Hoekman, D. H.: Scatterometer observations of seasonal backscatter variation over tropical rain
649 forest, *IEEE transactions on geoscience and remote sensing*, 37, 859–861, publisher: IEEE, 1999.

650 Wright, S. J. and Van Schaik, C. P.: Light and the phenology of tropical trees, *The American Naturalist*, 143, 192–199, 1994.

651 WWF: Terrestrial Ecoregions | Biome Categories | WWF, <https://www.worldwildlife.org/biome-categories/terrestrial-ecoregions>, 2019.