

## A multivariate-driven approach disentangling the abrupt reduction of near-natural Iberian streamflow post-1980

Amar Halifa-Marín<sup>1</sup>, Miguel A. Torres-Vázquez<sup>1</sup>, Enrique Pravia-Sarabia<sup>1</sup>, Marc Lemus-Cánovas<sup>2</sup>, Pedro Jiménez-Guerrero<sup>1</sup>, Juan Pedro Montávez<sup>1</sup>

<sup>1</sup>Regional Atmospheric Modelling (MAR) Group, Regional Campus of International Excellence Campus Mare Nostrum (CEIR), University of Murcia, 30100 Murcia, Spain.

<sup>2</sup>Climatology Group, Department of Geography, University of Barcelona, 08001 Barcelona, Spain.

10 Correspondence to: Pedro Jiménez-Guerrero (pedro.jimenezguerrero@um.es)

### Abstract

15 Whereas the literature still debates how several human/natural factors contributed to the recent streamflow decline in the Iberian Peninsula, a significant abrupt decrease of Winter Precipitation (WP) has been noticed in this area since 1980s related to large-scale atmospheric drivers. This contribution assesses its potential propagation into long-term streamflow series. For this purpose, the novel dataset of Near Natural Water Inflows to Reservoirs of Spain (NENWIRES) was created. The results highlight that those higher decreases of Winter Water Inflows (WWI) are always found related to WP reductions. Whereas WP declining was strongly provoked by the enhancement of NAOi, the WWI reductions could not be essentially linked to its behaviour in several NENWIRES catchments. Instead the intensification of permanent drought and forest extension promoted WWI reductions over the target area. In fact, these mechanisms allowed to understand why WWI reductions were higher than WP weakening. Summarizing, most humid catchments registered a WWI decline, mainly promoted by the NAOi enhancement, while its extension of forest and evapotranspiration rises seem to explain the WWI losses in the semiarid environments. This contribution sheds light on the recent debate about magnitude/drivers of streamflow decline over southern European regions. Furthermore, it might help water planning with the goal of mitigating the climate change impacts affecting the water cycle.

**Eliminado:** Disentangling the scarcity...brupt reduction of near-natural Iberian hydrological resources ...streamflow post-since ...980s: a multivariate-driven approach (... [1])

**Con formato:** Fuente: 12 pto

**Eliminado:** A

**Eliminado:** declining...ecline in the Iberian Peninsula, a significant abrupt decrease of Winter Precipitation (WP) has been noticed (... [2])

**Eliminado:** in the Iberian Peninsula

**Eliminado:** ...cale atmospheric drivers. This contribution p (... [3])

**Eliminado:** the ...ong-term variability of water resources...streamflow seriesbased on a multivariate-driven approach... For this purpose, the novel dataset of Near Natural Water Inflows to Reservoirs of Spain (NENWIRES) was created. Results confirm that Winter Water Inflows (WWI) have been modulated by the sudden decline in WP (... [4])

**Eliminado:** Our...he results highlight main finding is (... [5])

**Eliminado:** .

**Eliminado:** always ...re always found in relation (... [6])

**Eliminado:** Winter Water Inflows (...WI) (... [7])

**Eliminado:** These drastic reductions of WP/WWI were mainly controlled by the enhancement of the positive phase of the North Atlantic Oscillation (NAOi+). Nonetheless, our results also highlight the anthropogenic/physical causes contributing to the 1980s shift in the hydroclimate series. ...nstead tT...e rise ...ntensification of temperature (... [8])

**Eliminado:** (SPEI12)

**Eliminado:** .

**Eliminado:** .

**Eliminado:** the cropland abandonment and ...orest extension provoked ...romoted evapotranspiration gains and (... [9])

**Eliminado:** the

**Eliminado:** run-off ...WI weakening (... [10])

**Eliminado:** there

**Eliminado:** ...ummarizing, NENWIRES ...ost humid catchments registered the decrease of (... [11])

**Eliminado:** ing

**Eliminado:** + persistence/frequency... while the extensionland greening-up... (... [12])

**Eliminado:** ET ...vapotranspiration rises, it seems, (... [13])

**Eliminado:** Iberian ...emiarid environments. This contribution sheds some ... (... [14])

**Eliminado:** declining

**Eliminado:** Therefore

**Eliminado:** to

## 125 1 Introduction

The Mediterranean region shows the strongest and the most consistent pattern of significant streamflow decline worldwide as a result of climate change (Gudmundsson et al., 2021). Water planning in this area thus faces the challenge of securing the sustainability of natural and human systems. Freshwater scarcity poses an incipient risk (Tramblay et al., 2020) given that intensified droughts (climate drivers) coexist with the increase of human-induced water requirements (Polade et al., 2017; Vicente-Serrano et al., 2019). So, current studies need to shed light on the water resource variability/modelling based on multivariate approaches (e.g. Teuling et al., 2019, Massei et al., 2020).

Focusing on the Iberian Peninsula (IP), the scientific literature generally reports a streamflow decline over the last decades (e.g. Lorenzo-Lacruz et al., 2012). Nevertheless, the role of physical/anthropogenic factors driving these reductions are still under debate.

On the one hand, the decrease of recent Winter Precipitation (WP) has been robustly reported (e.g. de Luis et al., 2010; Lorenzo-Lacruz et al., 2013). That decrease was reasonably associated to the variability of the North Atlantic Oscillation index (NAOi) (e.g. Trigo et al., 2004). Nonetheless, Guerreiro et al. (2014) also pointed out abrupt WP decreases within the Tagus Basin since the late(early) 1970s(1980s). Other works also quantified the same drastic reduction of WP in the adjoining watersheds, Jucar and Guadalquivir (Gómez-Martínez et al., 2018, Halifa-Marín et al., 2021). They discussed 1) whether WP losses were promoted by a gradual decline or an abrupt shift; 2) the potential propagation of the WP abrupt shift have occurred into the streamflow series; and 3) its relationship with the NAOi enhancement, phenomenon which has been widely described in the scientific literature (e.g. Luo & Gong, 2006, Wang et al., 2014). The lack of knowledge within this topic does not only concern the IP region. An abrupt change of streamflow was also reported in central/northern Europe (Hannaford et al., 2013), with a positive trend at annual/wintertime scales (e.g. Stahl et al., 2010; Vicente-Serrano et al., 2019). However, while those authors generally accepted the potential links between NAOi and the changes in wintertime streamflow over Europe, they highlighted the importance of carrying out long record analyses in southern Europe, where data are sparse. They suggest that more long-term series are needed to determine whether

Con formato: Justificado, Interlineado: 1,5 líneas

Eliminado: declining

Eliminado: , where

Eliminado: ying

Eliminado: s

Con formato: Color de fuente: Texto 1, Inglés (americano)

Eliminado: detects

Eliminado: declining

Eliminado: through

Eliminado: robustness

Eliminado: decreases

Eliminado: reported

Eliminado:

Eliminado: , which were

Eliminado: ,

Eliminado: ,

Eliminado: literature

Eliminado: to

Eliminado: where it was registered its

Eliminado: such

Eliminado: changes in

Eliminado:

recent tendencies towards a decreased runoff in that region are found in longer records and whether this fact is related to the atmospheric circulation.

Eliminado: is

Eliminado: seen

On the other hand, several contributions have concluded that the streamflow decline was exacerbated by the temperature/evapotranspiration rise in the IP (e.g. Vicente-Serrano et al., 2014). Similar conclusions were obtained for the Mediterranean basin (García-Ruiz et al., 2011) and Europe (Teuling et al., 2019). All these works highlighted the role of a warmer climate and reforestation/afforestation processes into the evapotranspiration rise since potential evapotranspiration (ETP) also increases in response to global warming. Peña-Angulo et al. (2021) reported that the human-induced land greening-up processes through the 20<sup>th</sup> century contributed to the Iberian streamflow decline (intensifying the hydrological droughts). In this line, according to Vicente-Serrano et al. (2019), human-induced land cover changes (e.g. afforestation/irrigation) mainly explain the streamflow decline in the IP. Also, the role of other impacts was assessed (e.g. the construction of dams, Lorenzo-Lacruz et al., 2012) as well as the time-lag in the hydrological response caused by the permeability of soils (Lorenzo-Lacruz et al., 2013).

Eliminado: investigations

Eliminado: ing

Eliminado: C

Eliminado: decline

Eliminado: ing

Eliminado: ,

Eliminado: (

Eliminado: we found

Eliminado: that modulate

Eliminado: ing

Eliminado: more

Eliminado: an

Eliminado: to

Eliminado: Whereas

Eliminado: , Meanwhile

Eliminado: ,

Con formato: Fuente: Cursiva

Con formato: Derecha

Eliminado:

Eliminado:

Eliminado:

Eliminado:

Eliminado:

Eliminado:

Eliminado: →

Eliminado: ,

Con formato: Fuente: Cursiva

Con formato: Fuente: Cursiva

Con formato: Fuente: Cursiva

Therefore, noticeable divergences are found in the state-of-the-art regarding the importance of anthropogenic/physical drivers modulating the Iberian streamflow, which inspire a strong need for further scientific knowledge to ensure efficient water management over this target region. Also, uncertainties persist about how the wintertime NAOi enhancement has affected the Iberian streamflow variability. As far as the authors are concerned, an abrupt change of streamflows was not fully assessed in the region. At the same time as it appears likely that NAOi enhancement provoked a sudden decrease of WP, several authors have concluded that human perturbations determined the recent streamflow behaviour instead of climate drivers.

Generally, it is well-known that hydrological response can be approximated by a simple equation (Eq. 1):

$$RR = P - E - I, \quad (\text{Eq. 1})$$

where run-off (RR) depends on the subtraction between precipitation gains (P) and the accumulation of Evaporation (E) and Infiltration (I) losses.

RR refers to surface/underground run-off that flows outside of catchment boundaries. The estimation of evaporation is conducted by several factors (Eq. 2):

$$E = Ed + Ec + Et, \quad (\text{Eq. 2})$$

where  $E$  depends on the accumulation of losses due to evaporation processes from soils ( $Ed$ ), evaporation processes from vegetation ( $Ec$ ) and evapotranspiration ( $Et$ ).

So, it is assumed that  $E$  depends on vegetation, being  $Ed$  higher under lower vegetation density and  $Ec$  increases in response to the vegetation extension. Clearly,  $Et$  also rises due to the extension of vegetation, but its magnitude depends on soil-moisture budgets. However, at seasonal/monthly scales, the importance of evapotranspiration prevails in the estimations of evaporation. Henceforth, vegetation changes/cover have to be established to estimate variations in evaporation. Furthermore, it is well-known that other factors, as temperature, can impact the estimations of evaporation (i.e. through changes in evapotranspiration,  $Et$ ). Finally, the estimation of infiltration depends on terrain characteristics such as potential water storage/recharge, permeability/porosity of soils and the moisture content, which depends on the prevailing climate conditions (e.g. precipitation records). So, as stated in Eq. 1, natural variability of streamflow is conducted by changes in the parameters  $P$ ,  $E$ , and  $I$ .

Under this umbrella, this work assesses the variability of wintertime Iberian streamflow in a long-term analysis (1952/2018), identifying reservoirs where their water inflows have limited human influence (near-natural environments). The relevance of wintertime changes of precipitation/streamflow resides in its contribution to water resources, given that winter is the wet/recharge season in most areas of the IP. In addition, to the extent water planning is concerned here, the occurrence of abrupt decreases in streamflow records might severely affect the sustainability of natural/human systems (more than gradual changes).

Likewise, the assessment of water resources needs to focus on near-natural catchments (e.g. Stahl et al., 2010, Hannaford et al., 2013, Vicente-Serrano et al., 2014), in order to reduce the uncertainties added by human-induced perturbations. Because the streamflow depends on several factors with opposing effects, changes in the streamflow should be analyzed at small scales where individual factors can be understood, rather than at larger river-basin scales (e.g. Teuling et al., 2019). This approach allows to fully assess the

**Eliminado:**  $E$   
**Con formato:** Fuente: Cursiva

**Eliminado:** .  
**Eliminado:** .

**Con formato:** Fuente: Cursiva  
**Con formato:** Derecha

**Con formato:** ... [15]

**Con formato:** Fuente: Cursiva  
**Eliminado:** . Whereas

**Eliminado:** is ... higher under lower vegetation density and, ... [16]

**Con formato:** Fuente: Cursiva  
**Con formato:** Inglés (americano)

**Con formato:** Fuente: Cursiva, Inglés (americano)  
**Con formato:** ... [17]

**Con formato:** Fuente: Cursiva  
**Eliminado:** Et...vapotranspiration prevails in the E...stimations of evaporation. Henceforth, consequently it is ...established th... [18]

**Eliminado:** E...variations in evaporation. Furthermore, it is well-known that other factors, as temperature, can impacts...on th... [19]

**Eliminado:** promoting  
**Eliminado:** T...e I...estimation of infiltration depends on terrain characteristics such as potential water storage/recharge, permeability/porous...osity of soils, ... [20]

**Con formato:** Fuente: Cursiva  
**Con formato:** ... [21]

**Eliminado:** the ...ost areas of the IP. In addition, to the extent that water planning is concerned here, more than gradual changes, ...he occurrence of abrupt decreases in streamflow records might severely impact...ffect to ... [22]

**Eliminado:** analysed...alyzed at small scales where individual factors can be understood, rather than at larger river-... [23]

**Eliminado:** It

hydrological response to global change forcings and pressures, such as changes in atmospheric circulation patterns and/or human perturbations. In addition, the Iberian flow regimes are controlled by quite different climate conditions, which helps to assess these perturbations to a wide variety of hydroclimate areas. Therefore, this contribution aims to 1) characterize/verify post-1980 changes into near-natural water inflows series, and 2) disentangle how climate/human drivers have contributed to the magnitude of post-1980 change.

The following list of tasks has been addressed to achieve both objectives: 1) quantify the trend and the abrupt changes of water inflows series; 2) analyze whether the NAOi plays a leading role in those changes of water inflows detected; 3) identify how permanent meteorological droughts modulate water inflows changes; 4) estimate the contribution of forest extension (as a consequence of agricultural abandonment, human-induced) to the evolution of Winter Water Inflows (WWI); and 5) examine the results to provide an identification of the main precursor for WWI changes in the target catchments. For that, the initial working hypothesis is that the climate (NAOi/WP behaviour) essentially controls the reduction of water inputs, while the impact of human perturbations is weaker. The magnitude of change in any parameter of the RR equation is assumed to be important to understand the variations of WWI.

## 2 Data and Methods

### 2.1 Data Sources

#### 2.1.1 Spanish Near-Natural Water Inflows to Reservoirs

In this study monthly water inflows recharging the Spanish reservoirs network (376 series) have been analysed (CEDEX, 2021). Water inflows are estimated by water management agencies from the reservoir outflow while accounting for reservoir storage changes at daily scale. For the Spanish Ministry of Environment (currently called *Ministerio para la Transición Ecológica y Reto Demográfico*) these series thus represent the official dataset of drained streamflow to the country reservoirs. Given that this analysis focuses on the variability of near-natural water inflows, data were collected from the repository using 36 series (9.6%), composing the NEar-Natural Water Inflows to REservoirs of Spain (NENWIRES) dataset.

Eliminado: the

Eliminado: /

Eliminado: at

Eliminado: s

Eliminado: ing

Eliminado: the

Eliminado: ing

Eliminado: For that purpose, at least, the

Eliminado: into

Eliminado: analyse

Eliminado: if

Eliminado: er-

Eliminado: evolution

Eliminado: principal

Eliminado: our

Eliminado: led

Eliminado: While we assume that

Eliminado: t

Eliminado: variations

Movido (inserción)[1]

Movido hacia arriba[1]: Their results might suggest that the sudden fall of water resources could have occurred in other

Eliminado: ¶ ... [24]

Eliminado: Droughts and floods keep increasing their imp... [25]

Eliminado: Data Sources

Eliminado: Spanish

Eliminado: T

Eliminado: e identification of the number of reservoirs wh... [26]

Eliminado: that recharging

Eliminado: . These hydrological series can be downloaded... [27]

Con formato ... [28]

Con formato ... [29]

Eliminado: our

Eliminado: i

Eliminado: wWe

Eliminado: 36 water inflow series (9.6%)

Eliminado: total

Eliminado: (375 series)

Eliminado: which is called. Then, the

Eliminado: t was created

475 Further details of NENWIRES reservoirs (Appendix A, Table A1) and methods used to estimate the water inflows series are included in Appendix A.

480 In addition, the procedures to select the NENWIRES catchments follow commonly used methods (e.g. Hannaford et al., 2013) in this field of knowledge: 1) verification that water inflows are not affected by large perturbations, excluding those series impacted by water regulation (e.g. damming) and urban/irrigation extractions; 2) long-term series are prioritized, and water inflows records must cover at least 46 years (70%) through the study period (1952-2018); 3) series must provide continuously records for at least 33 years (50%); and 4) records must not be reconstructed, and homogeneous quality controls have been applied to series.

485 The dimension of NENWIRES catchments is generally small with an average area of 929 km<sup>2</sup>. The compiled data thus provides near-natural water inflows records spanning the continental Spain (Fig. 1). Likewise, the NENWIRES basins also mention the headwaters of the transboundary basins (Douro, Tagus, Guadiana), so the findings properly considered the streamflow evolution of IP. The boundaries of their drainage basins were provided by the IDE (2021).

Eliminado:

Eliminado:

Eliminado: our

Eliminado: the

Eliminado: we verify

Eliminado: that

Eliminado: We exclude

Eliminado: that are

Eliminado: ,

Eliminado: prioritised

Eliminado:

Eliminado: ; and 5)

Eliminado: our

Eliminado: analysed

Eliminado: Also, the

Eliminado:

Eliminado: catchments draining


Eliminado: to the NENWIRES reservoirs can be downloaded

Eliminado: from

Eliminado: , which provides the area of Spanish principal/secondary basins... The NENWIRES dataset provides near-natural water inflows series spread throughout the peninsular territory of Spain (Fig. 1). Only the smaller territories of Canary/Balearic Islands, Galician, Catalonian, and Andalusian Basins were not studied. Given that the headwaters of several transboundary IHB (Douro, Tagus, Guadiana) are in the Spanish territory, our findings represent properly the variability of Iberian Water Resources. ¶

Eliminado: The NENWIRES dataset provides near-natural water spread throughout the peninsular territory of Spain (Fig. 1). Only the smaller territories of Canary/Balearic Islands, Galician, Catalonian, and Andalusian Basins were not studied. Given that the headwaters of several transboundary IHB (Douro, Tagus, Guadiana) are in the Spanish territory, our findings represent properly the variability of Iberian Water Resources. ¶

Con formato: Color de fuente: Automático



530 **Figure 1.** Panels show (a) the orography in the Iberian Peninsula (IP), and (b) the NENWIRES reservoirs grouped by Iberian Hydrological Basins (IHB). In the left panel, light blue contours represent the IHB boundaries in the continental Spain, and black contours mention the altitude above sea level (each 500 meters).

### 2.1.2 Climate Data

535 The Spanish Precipitation Gridded Dataset has been recently published by the Spanish Meteorological Agency (AEMET). The interpolation procedures manage 3,236 rain gauges. This dataset currently covers the period from January 1951 to December 2020 at daily scale. Its spatial resolution reaches  $\sim 0.05 \times 0.05^\circ$ , covering the Spanish territory except the Canary Islands. The number of records managed and its spatial resolution motivates the usage of this dataset. In the case of the Temperature Gridded Dataset of Spain, 540 the interpolation manages 1,800 gauges, providing the daily maximum and minimum temperature. Both datasets can be downloaded in netCDF format through the AEMET Website (AEMET, 2021). Finally, the monthly NAO index (NAOi) was collected from the National Oceanic and Atmospheric Administration (NOAA), which covers the period from January 1950 to nearly real-time (NOAA, 2021). This index is a simplification of a large-scale atmospheric circulation pattern over the North Atlantic, 545 which usefully helps to understand the principal moisture sources reaching the IP (e.g. Trigo et al., 2004).

**Eliminado:**

**Eliminado:** T

**Eliminado:** );

**Eliminado:** the Iberian Hydrological Basins (IHB). Purple regions represent the IHB boundaries within the Spanish territory

**Eliminado:** Soft

**Eliminado:** represent

**Eliminado:** each 500 meters

**Eliminado:** †

† The procedures to identify the NENWIRES catchments are summarized as: 1) we verify that water inflows are not affected by human-induced disturbances. We thus checked that upstream water inflows are not impacted by dams, channels or other water works. However, some dams related to flood risk control or hydropower production were included because they do not disturb the monthly/seasonal variability; 2) Long-term series are needed to conclude significantly statistical findings, so water inflows records must cover at least 45 years through the period 1952/2018; and 3) In addition, water inflow series must provide continuously records for at least 30 years. Thus, missing values should be unusual. †

**Eliminado:** created

**Eliminado:**  $\sim 5\text{km}^2$

**Eliminado:** The robustness of the analysis improves due to

**Eliminado:** t

**Eliminado:** observational series interpolated

**Eliminado:** ,

**Eliminado:** the

**Eliminado:** high-

**Eliminado:** to use

**Eliminado:** The Temperature Gridded Dataset of Spain has technical characteristics similar to those of the precipitation dataset aforementioned.

**Eliminado:** this case

**Eliminado:** . The temperature dataset

**Eliminado:** provided

**Eliminado:** (TX)

**Eliminado:** (TN) at daily scale.

**Eliminado:** accessible

**Eliminado:** One the other hand, the NCEP/NCAR Reanalysis I has been used in many meteorological studies, providing  $1 \times 1^\circ$  global climate data framed between  $90^\circ\text{N}/\text{S}$  latitude. The high quality of this reanalysis dataset in the northern hemisphere is due to the high density of the meteorological observations worldwide. The t(... [30]

**Eliminado:** montly

**Eliminado:** taken

**Eliminado:** .



The negative phases of the NAO index (NAOi-) are related to low systems affecting the IP, promoting moisture fluxes from the SW, whereas NAOi+ represents the opposite synoptic pattern.

Eliminado: . W

### 605 2.1.3 Soils Permeability

Hydrological data collected from aquifers is not considered in this contribution. Instead, the groundwater modulation to WWI is estimated assessing the permeability of soils. It is assumed that this dataset helps to estimate the behaviour of  $f$  parameter into Eq. 1. The Permeability of Soils Dataset, provided by the Spanish Geological Survey (IGME), classifies soils into 9 (4) types (groups): 1) A1 and A2 represent alluvial deposits and colluvial soils formed by very permeable porous banks; 2) B1 and B2 represent carbonate bedrock, which are very permeable due to cracking or karstification processes; 3) C1, C2 and C3 represent well-drained volcanic soils, which are not common; and 4) D1 and D2 represent low permeability and impermeable conditions, respectively. This dataset quantifies the percentage of each type of soil in relation to the basin dimension. In this study, these types of soils are classified in 4 groups: A/B1 soils are Very-High Permeable Soils (VHPS); A/B2 soils are High Permeable Soils (HPS); D1 soils are Low Permeable Soils (LPS); and D2 soils are No Permeable Soils (NPS). The dataset is online accessible through the IGME Website (IGME, 2021).

Eliminado: We do not manage

Eliminado: h

Eliminado: we did an estimation of

Eliminado: RR e

Eliminado: uation

Con formato: Fuente: Cursiva

Eliminado: was used in this contribution. It

Eliminado: respectively

Eliminado: we classify

Eliminado: .

Eliminado: .

Eliminado: In this study, Permeable Soils include A1/2 and B1/2 types. Oppositely, Impermeable Soils are only represented by D2 soils. ...

Eliminado: is

### 2.1.4 Land Cover Changes

As recommended by Teuling et al. (2019) for studies covering this topic, the Historic Land Dynamics Assessment (HILDA, v2.0) model reconstruction of historic land cover/use change (Fuchs et al., 2015) has been used. This dataset is based on multiple harmonized and consistent data streams including remote sensing, national inventories, aerial photographs, statistics, old encyclopedias, and historic maps to reconstruct historic land cover. The spatial resolution reaches 1x1 km, whereas the time coverage ranges from 1900 to 2010 in decadal time steps (HILDA, 2021). The reconstruction provides information for six different land cover/use categories: forest, grassland, cropland, settlements/urban, water bodies and other. Only the changes of forest cover in the NENWIRES catchments have been quantified. The gross land changes were studied, computing the sum of all area gains and losses occurring within an area and period. Changes in vegetation are used to estimate the  $E$  parameter into Eq. 1.

Eliminado: in

Eliminado: in

Eliminado: such as

Eliminado: <sup>2</sup>

Eliminado: We only quantify analyse

Eliminado: which compute

Eliminado: RR equation

Con formato: Fuente: Cursiva



## 2.2 Analysis Procedures

### 2.2.1 Study Period and Preprocessing

The study period covers from October 1951 to September 2018. Only the extended wintertime season (December to March, DJFM) is considered here, given that 1) the extended winter allows to study the contribution of snowfalls into streamflow series (early snowmelt); 2) the relation between WP and annual streamflow is intensive because the recharge of reservoirs/aquifers mainly occur at wintertime (Lorenzo-Lacruz et al., 2012); 3) WP shows the strongest and the most consistent pattern of significant decline in the IP (De Luis et al., 2010); and 3) the NAOi enhancement has been reported on wintertime. WWI generally explain more than 50% of annual records in NENWIRES catchments (Fig. 2). However, a higher contribution to annual water inputs was not confirmed within the extended winter for several NENWIRES catchments (R3, 9, 20, 21, 25, 33 and 36; 19%), where the peak of the hydrograph is driven by Mediterranean heavy rainfalls (spring/autumn) or snow accumulation/melt processes (late spring and early summer).

On the other hand, processing of NetCDF files for climate data was conducted with CDO software (Schulzweida, 2019). For converting from daily to seasonal scale (large winter, DJFM), *seassum* and *seasmean* CDO functions were used. After calculating the large winter accumulation of WP and average of maximum and minimum temperatures (WTX and WTN, respectively), its spatial average within the target watersheds is computed. This procedure was implemented in RCRAN language, loading the NetCDF files with the *brick* function (Hijmans, 2021). In addition, the catchments polygons (ESRI shapefile) were read with the *readOGR* function (Bivand, 2021). Then the gridded climate data is cropped within the boundaries of catchments using the *mask* function, and consecutively the spatial average of timesteps is estimated with the *cellStats* function (Hijmans, 2021). The extended winter accumulation/average of record series (WWI and NAOi) was also developed in RCRAN.

Eliminado: Define the

Eliminado: But we o

Eliminado: study

Eliminado: large

Eliminado: Winter Water Inflows (

Eliminado: )

Eliminado: large

Eliminado: , the

Eliminado: the

Eliminado: /

Eliminado: having calculated

Eliminado: /

Eliminado: we compute

Eliminado: we read

Con formato: Fuente: Cursiva

Eliminado: we cropped

Con formato: Fuente: Cursiva

Eliminado: we estimated


Con formato: Fuente: Cursiva

Eliminado: On the other hand, large

Con formato: Fuente: Cursiva

Eliminado: the

Mean Annual Hydrograph of NENWIRES Series



**Eliminado: 2.2.1 Steps to Define the Study Period**  
 To establish links between these multidisciplinary data sources, their time coverage should be consistent. Several datasets start only from 1951 (NAOI, Precipitation, TX/TN). Henceforth, the period covers from October 1951 to September 2018. Also, in order to simplify the analysis, the large wintertime scale (December to March, DJFM) was considered. Consequently, we have considered that 1) the relation between WP and summer droughts are intensive, because recharge of reservoirs/aquifers mainly occur at wintertime (Lorenzo-Lacruz et al., 2012); and 2) WP shows the strongest and the most consistent pattern of significant decline in the IP (De Luis et al., 2010, Halifa-Marín et al., 2021). In fact, we confirmed that Winter Water Inflows (WWI) explain more than 50% of annual water inflows (Fig. 2).

**Eliminado: A n**  
**Eliminado: has been created**  
**Eliminado: large**

**Eliminado: Figure 2.** Hydrograph of NENWIRES dataset. It represents the monthly water inflows (MWI) to the annual water inflows (AWI) from 1952 to 2018. The total average of MWI is showed in the right axis.

**2.2.2 Pre-processing data**  
 Processing of NetCDF files for climate data was conducted with CDO software (Schulzweida, 2019). For converting from daily to seasonal scale (DJFM), the seasum/season CDO functions were used. After having calculated the WP and winter maximum and minimum temperatures (WTX/WTN), we compute its spatial average within the target watersheds. This procedure was implemented in RCRAN language, loading the NetCDF files with the brick function (Hijmans, 2021). In addition, we read the catchments polygo...

**Eliminado: 3**

**Eliminado: of hydroclimate variables**

**Eliminado: Our**

**Eliminado: S**

**Eliminado: To**

**Eliminado: our**

**Eliminado: we quantified**

**Eliminado: First,**

**Eliminado: f**

**Eliminado: (shown in Section 3, Fig. 3)**

**Eliminado: was obtained**

**Con formato: Fuente: Cursiva**

**Eliminado: 1**

**Eliminado: 1**

**Con formato: Derecha**

**Con formato: Fuente: Cursiva**

**Con formato: Fuente: Cursiva**

**Eliminado: Their statistical significance at  $\alpha = .05$  has been tested. Henceforth, the same statistical significance was always test...**

**Figure 2.** Novel hydrograph version for NENWIRES series. Each cell in the matrix shows the relative monthly contribution (rows) to the annual accumulation (AWI) in each NENWIRES catchment (columns). The monthly contribution is based on records along the study period. The monthly average of the dataset is highlighted in the right axis. The horizontal black lines highlight the extended wintertime season (DJFM).

**2.2.2 Statistical analysis**

The statistical procedures were fully conducted in RCRAN language. In order to address task 1, the trend of magnitudes for WWI throughout the study period was obtained. This trend analysis was conducted using the Sen's slope test with *sens.slope* function (Pohlert et al., 2018). To allow a proportional discussion of trend analysis, slope estimates were standardized (ZSS) as follows (Eq. 3):

$$ZSS_{WWI} = \frac{SS_{WWI}}{x_{WWI}}, \quad \text{(Eq. 3)}$$

where standardized Sen's slope (ZSS) is the coefficient between Sen's Slope estimate (SS) and the mean WWI for study catchments.

Significant trend estimates are considered with p-value  $\geq 0.95$ , which is required in all statistical procedures. Once the ZSS magnitudes were quantified, the potential propagation of WP abrupt change to WWI since the 1980s was assessed (addressing the aim established in task 1 of this contribution). For that purpose, the most probable change was identified through the Pettitt's Homogeneity method, according to *pettitt.test* function (Pohlert, 2018). Likewise, its significance was externally evaluated with the non-parametric Mann-Whitney U test through the *wilcox.test* function (Hothorn et al., 2019). First, the breakpoint (BP) was computed for standardized series of average WWI/WP in the dataset, which were normalized through the *scale* function (R Core Team, 2021). The average WWI/WP was quantified through the *mean* function (R Core Team, 2021). A similar method was applied to both variables at catchment analysis. Then, the Relative Change (RC) of WWI/WP was quantified after the more frequent significant BP registered in the series, 1979/1980. RC was calculated as follows (Eq. 4):

$$RC = \frac{MS_{P2} - MS_{P1}}{MS_{P1}} \quad (\text{Eq. 4})$$

where RC is estimated by subtracting the average (horizontal bars) of WWI/WP during the first period ( $MS_{P1}$ ) to its average during the last period ( $MS_{P2}$ ), standardizing with the early mean ( $MS_{P1}$ ). The average was quantified with *mean* function (R Core Team, 2021).

In order to address the task 2, the break point (BP) in the NAOi series was checked using the aforementioned methods. In addition, the correlation between WP/WWI and NAOi was also calculated. To this end, the *cor.test* function was used under Pearson's method (R Core Team, 2021), which provides its p-value. Previously, the series were detrended using the *detrend* function (Borchers, 2019). These correlation coefficients were related to the RC of WWI/WP (estimations based on Eq. 4) through the *lm* function (R Core Team, 2021). This function was used to fit linear models (regression) between variables. The function *lm* will be used afterwards in the analysis presented in Section 3. In addition, the correlation between WWI and WP was also quantified.

The abovementioned methods allow to identify those basins where WWI changes could not be explained only with the NAOi/WP variability. So, the role of several factors was analyzed (tasks 3/4). Preliminary, the modulation of WWI by the magnitude of persistent drought conditions was evaluated. The magnitude

**Eliminado:** above

**Con formato:** Fuente: Cursiva

**Eliminado:** /...ask 1 of this...contribution). For that purpose, we identified ... [33]

**Con formato** ... [34]

**Eliminado:** we compute ...he breakpoint (BP) was computed for standardized series of average WWI/WP in the dataset (Fig. ... [35]

**Con formato** ... [36]

**Eliminado:** S...imilar method was applied to both variables at catchment analysis (Fig. 5)... Then, the Relative Change (RC) of WWI/WP was quantified after the more frequent significant BP registered in the series, 1979/1980 (Fig. 6)... The ...C was calculated as follows (Eq. 4) ... [37]

**Eliminado:** ... (Eq. 4) ... [38]

**Con formato:** Derecha

**Eliminado:** the

**Con formato** ... [39]

**Eliminado:** keeping similar methods abovementioned, we checked the break point (BP) in the NAOi series was checked...using the aforementioned methods. In addition, the correlation between WP/WWI and NAOi was also calculated. (Fig. 7). ... [40]

**Con formato** ... [41]

**Eliminado:** 2

**Con formato** ... [42]

**Eliminado:** in...fterwards in the following ...analysis of...resented in S... [43]

**Eliminado:** (Fig. 9)

**Con formato:** Sin Resaltar

**Eliminado:** we analysed ...he role of several factors was analyzed (response to our ...asks 3/4). We p...reliminary, evaluated ... [44]

of drought conditions is used as an estimation of moisture in soils, which impacts the hydrological response (run-off/infiltration processes) in the watersheds. To this end, the mean wintertime drought intensity based on the previous 6/12 months was analyzed. It required the quantification Standardised Precipitation-Evapotranspiration Index (SPEI) (Berguería & Vicente-Serrano, 2017), which needs the estimation of potential evapotranspiration (ETP) using the Hargreaves method. SPEI was computed with the *spei/hargreaves* functions. These estimations allow to assess the changes of ETP in relation to the temperature evolution. The average of SPEI6/12 (through the mean function) are presented in Section 3. Likewise, the most probable BP was also detected for those SPEI series (through the *pettitt.test* function). Finally, the absolute change of SPEI6/12 post-1980 is also shown in Section 3. The SPEI change quantified the difference between its average along 1952/1979 and 1980/2018.

In order to understand the hydrological response during the most humid winters, the potential time-lag that permeability of soils motivates on the water generation (WWI) was evaluated in the NENWIRES catchments. This quantification allows us to understand how persistent/heavy rainfall events are linked to the recharge of aquifers, and the relationship between these modes of recharge and WWI series. Both estimations are crucial to understand the contribution of the infiltration parameter into WWI changes (parameter  $I$  in Eq. 1). This analysis is of particular interest to the limestone environments where the permeability of soils (its porosity) guarantees high infiltration rates. So, the deviation between extreme percentiles of WWI/WP in NENWIRES basins is quantified. The QQ-Deviation test (QQD) was calculated as shown in Eq. 5:

$$QQD = P_{90}(Z_{WWI}) - P_{90}(Z_{WP}) \quad \text{(Eq. 5)}$$

where Quantile-Quantile Deviation (QQD) is the difference between standardized WWI/WP anomalies averaged (horizontal bars) to the points over 90<sup>th</sup> percentile ( $Z_{WWI}/Z_{WP}$ ). Higher values of QQD thus show a greater water generation whether persistent/heavy rainfalls events occur (most humid winters), whereas lower (below 0) values show the opposite relationship.

In addition, in order to assess the change of forest cover, the HILDA dataset was used. Given that this study focuses on near-natural catchments (headwaters, depopulated areas), the changes of forest cover

**Eliminado:** on ...he hydrological response (run-off/infiltration processes) in the watersheds. To this end, we analysed ...he mean wintertime drought intensity based on the previous 6/12 months was analyzed. It required the SPEI ...uantification Standardised Precipitation-Evapotranspiration Index (SPEI) (Berguería & Vicente-Serrano, 2017), which needs the estimation of ETP...otential evapotranspiration (ETP) estimation under ... [45]

**Eliminado:** (Berguería & Vicente-Serrano, 2017)... As well as t...ese estimations allow us ...o assess the changes of ETP in relation to the temperature evolution. The average of SPEI6/12 (through the mean function) are presented in the ...s...ction 3. (Fig. 8). ... [46]

**Con formato:** Fuente: Cursiva

**Con formato:** Fuente: Cursiva

**Eliminado:** the s...ction 3. (Fig. 9). ... [47]

**Eliminado:** We also aimed to

**Eliminado:** ...nderstand the hydrological response during the most humid winters. For that... the potential time-lag that permeability of soils motivates on the water generation (WWI) was evaluated in the NENWIRES catchments. This quantification allows us to understand how 1)...persistent/heavy rainfall events are linked to the recharge of aquifers, and 2) ...he relationship between these modes of recharge and WWI series. Both estimations are crucial to understand the contribution of I ... [48]

**Eliminado:** in terms of RR equation...q. 1). This analysis is of particular interest to the limestone environments where the permeability of soils (its porosity) guarantees high infiltration rates. So, we quantified ...he deviation between extreme percentiles of WWI/WP in NENWIRES basins (Fig. 9)...s quantified. The QQ-Deviation test (QQD) was calculated as shown in (...q. 53) ... [49]

**Con formato:** Fuente: Cursiva

**Eliminado:** ...Eq. 53 ... [50]

**Con formato:** Derecha

**Con formato** ... [51]

**Eliminado:** Once the trend estimates were computed, the potential propagation of WP abrupt change to WWI since the 1980s was quantified. For that purpose, Pettitt's Homogeneity test was used according to pettitt.test function (Pohler, 2018). ... [52]

**Eliminado:** ... Then, the Relative Change (RC) after the change point detected on WWI/WP was also quantified. The RC was calculated as follows (Eq. 2):  

$$RC = \frac{MS_{P2} - MS_{P1}}{MS_{P1}}$$
(2) ... [52]

**Eliminado:** The significant change point detected was evaluated with the non-parametric Mann-Whitney U test through the ... [53]

**Eliminado:** On the other hand...n addition, t ... [54]

**Eliminado:** T... assess...s the change of forest cover, the HILDA dataset was used., ... [55]

**Eliminado:** our...his study focus...s on near-natural catchments (headwaters, depopulated areas), we considered ... [56]

were considered, as well as the agricultural abandonment and other human-induced land cover changes. HILDA provides interdecadal estimates of forest area. For our study period, a estimation per decade is available from 1950 to 2010 (7 time-steps). We thus computed the average of their interdecadal relative changes (DRC) as indicated in Eq. 6;

$$RC = \frac{\sum_{i=1}^{n-1} DRC_i}{n-1}; DRC_i = \frac{DC_{i+1} - DC_i}{DC_i}, \quad (\text{Eq. 6})$$

where the decadal cover (DC) represents the forest area in each time-step, and the mean RC (RC) is estimated by the average of interdecadal changes (DRC). Therefore, this method allows us to evaluate the changes of the forest area along the entire study period (shown in Fig. 5A, Appendix A).

All those methods allow to disentangle how climate/human drivers have contributed to the magnitude of post-1980 change, and to assess the contribution of abovementioned variables to estimate the principal precursor that promoted WWI changes in target catchments. Basically, a clustering methodology was applied through the K-Means algorithm (kmeans function, R Core Team, 2021), keeping the default algorithm settings. Once basins were classified, indicators of studied variables were computed for each cluster. These indicators refer to the average of variables within each cluster.

### 3 Results and Discussion

#### 3.1 Recent Evolution of Wintertime Iberian Near-Natural Water Inflows

The mean WWI ranged from 5 to 824 hm<sup>3</sup> in the NENWIRES basins (Fig. 3a). Higher records were observed over the western/central sector (400-800 hm<sup>3</sup>), and northern/southern areas (250-400 hm<sup>3</sup>). The lower WWI were registered in the eastern/southern coast (<= 250 hm<sup>3</sup>). Meanwhile, the trends ranged from -1.8 to 0.1%/yr (Fig. 3b). 97% of WWI series have decreased in the study period. WWI only increased in one basin, but not significantly. These WWI reductions were significant in most of the catchments (55%), mainly over central and eastern sectors. Also, significant reductions frequently occurred in the most humid basins. The results thus depict significant reductions in the headwater of the Tagus, Ebro, Douro, Segura, Jucar, and Mino basins (as defined in Fig. 1). These results agree with similar quantifications in the literature (e.g. Lorenzo-Lacruz et al., 2012), whereas this contribution is of interest to confirm the

- Eliminado: also as estimation of the
- Eliminado: Then, one ... or our study period, a value ... [57]
- Eliminado: RC
- Eliminado: (
- Eliminado: 3
- Eliminado: 4... [58]
- Con formato: Fuente: Cursiva
- Eliminado: ... (Eq. 64) ... [59]
- Eliminado: 3
- Con formato: Derecha
- Con formato: Fuente: Cursiva ... [60]
- Con formato: Fuente: Cursiva
- Eliminado: full
- Eliminado: ¶
- Eliminado: response the aim 2 (task 5)... as well as we assessed... to assess the contribution of abovementioned variables to estimate the principal precursor that promoted WWI changes in target catchments. Basically, we apply a ... clustering methodology was applied through the K-Means algorithm. To this end, the ... [61]
- Eliminado:
- Eliminado:
- Eliminado: n was used (
- Con formato: Fuente: Cursiva
- Eliminado: ¶
- In addition, the correlation between hydroclimate variables was estimated. To this end, the cor.test function was used under Pearson's method (R Core Team, 2021). This function automatically provides the p-value associated to the correlation coefficients. Previously, the series were detrended using the detrend function (Borchers, 2019). ¶ [62]
- Eliminado: In the end, the analysis of meteorological drou... [63]
- Eliminado: Variability ... Recent Evolution of Wintertime... [64]
- Eliminado: The results shown in Fig. 3a indicate that the
- Eliminado: In the NENWIRES basins the
- Eliminado: ...ranged from 5 to 824 hm<sup>3</sup> ... in the NENWIRES... [65]
- Eliminado: Larger
- Eliminado: /yr
- Eliminado:
- Eliminado: /yr
- Eliminado: less than... 250 hm<sup>3</sup>). Meanwhile, the trends ... [66]
- Eliminado: %/yr... (Fig. 3b). 97% of WWI series have dec... [67]
- Eliminado: shown
- Eliminado: our

reported changes of WWI within non-regulated catchments (Vicente-Serrano et al., 2014). Likewise, Vicente-Serrano et al. (2019) described negative trends in annual streamflows in continental Spain, which, according to these results, could be provoked by WWI changes.




Figure 3. (a) Mean WWI, and (b) its Standardized Sen's slope trend estimates computed after Eq. 3 (Section 2.2.2). For the trends, symbols represent positive trend (filled triangle/point-up) and negative trend (filled triangle/point-down). In the left panel, significant estimates are indicated with a black outline.

Contours of orography (black) and IHB of continental Spain (purple) were also added (see Fig. 1).

### 3.2 Abrupt changes or trend of Wintertime Water Inflows?

This section intends to shed some light to the question raised by Guerreiro et al. (2014) about whether WP gradually or suddenly changed in the IP since early 1980s. The overall negative trends of WWI could obscure an abrupt change post-1980, which has been recently noticed for the WP in the IP (e.g. Halifa-Marín et al., 2021). Thus, the most probable break point (BP) in average series of WP/WWI series (see Section 2.2.2) is identified. The average series probably are not representative for all NENWIRES catchments but give a general insight into the WWI variability.

Eliminado: the

Eliminado: our

Eliminado: eh

Eliminado: These WWI reductions were significant in most of catchments (55%), mainly over central and eastern sectors. Also, significant reductions frequently occurred in the most humid basins. WWI only increased in one basin, but not significantly. The results thus depict significant reductions in the headwater of the Tagus, Ebro, Douro, Segura, Júcar, and Mino basins.

Eliminado: Panels show

Eliminado: the

Eliminado: Wintertime Water Inflows (

Eliminado: )

Eliminado: l

Eliminado: S

Eliminado: ¶

Eliminado: ¶

Eliminado: Evidence of Climate-Driven Changes on Wintertime Water Inflows

Eliminado: Echoing the question made by

Eliminado: if

Eliminado: , our first main is objective objective has been to answer that question

Eliminado: ,

Eliminado: response to the heading of this chapter.

Eliminado: A lead role of WP to modulate the streamflow was already described in Trigo et al. (2004) and Lorenzo-Lacruz et al. (2012). However, recent studies have noticed a sudden decline of WP in the Tagus, Júcar, and Guadalquivir basins since the early 1980s (Guerreiro et al., 2014, Gómez-Martínez et al., 2018, Halifa-Marín et al., 2021).

Eliminado: We t

Eliminado: identify

Eliminado: To analyse

Eliminado: potential propagation of


Eliminado: this

Eliminado: abrupt decline into the W

Eliminado: variability

Eliminado: ,

230 Fig. 4 shows that WWI had a significant abrupt BP since 1979 in the NENWIRES dataset. The average  
of WWI was 175.6 hm<sup>3</sup> until 1979, while it shrinks the 30% since 1980. Gómez-Martínez et al. (2018)  
also identified a significant BP post-1980 on annual streamflow records in the Jucar/Turia headwaters.  
The results presented here suggest that it was provoked by the abrupt decrease registered in WWI series  
since 1979. Likewise, the detection of the most probable BP for WWI was also performed at catchment  
scale. This analysis shows that WWI shift was detected in 1979 (41%), as well as 1978 (17%) and 1980  
235 (14%).



240 **Figure 4.** Standardized series from average Wintertime Precipitation (WP, top) and Wintertime Water Inflows (WWI, bottom) in the NENWIRES dataset, during the period 1952/2018. Bars represent negative (red) and positive (blue) values. Vertical blue/black lines represent the most probable BP detected through the Pettitt's Homogeneity Test (Section 2.2.2).


Hence, a BP was found in 72% of WWI series between 1978/1980, although only 55% of these detections were significant (Fig. 5a). We found those significant BP in the headwater of Jucar, Tajo, Segura, Guadalquivir and Douro basins. Given that the uncertainties of the most probable change methods are wide, the results for those basins where WWI change between 1978 and 1980 agree with the post-1980

- Eliminado: In
- Eliminado: it is shown
- Eliminado: the significant break point in the total average of WP/WWI was calculated (Fig. 4). Both variables
- Eliminado: shift
- Con formato: Sin Resaltar
- Eliminado: , which our results
- Eliminado: to
- Eliminado: also

- Eliminado: we consider that



255 changes in streamflows reported by Gómez-Martínez et al. (2018). Furthermore, a source of error in the BP estimation could be caused by missing values in WWI series. However, we have identified a significant BP from 1978/1980 in case of full series and series with missing values.



260 **Figure 5.** (a) Break point (BP) detected for WP series, and (b) Id. for WWI series through methods used in Fig. 4. Grey squares represent a BP not detected between 1978 and 1980. Larger squares (framed into black borders) represent the significantly BP detected in both panels. Contours of orography (black) and IHB of continental Spain (purple) were also added (see Fig. 1).

### 265 **3.3 Does Precipitation Control the Abrupt Change of Wintertime Water Inflows?**

The abovementioned references recently reported a significant abrupt decrease of WP since the 1980s across several Iberian basins, which motivates essentially this study. Once the BP has been detected in WWI series, a similar analysis was conducted for WP records. The BP detected for average WP was found in 1979 (Fig. 4), matching the estimation for average WWI. Meanwhile, the average WP decreased by -21% since 1979, a lower reduction than the average WWI change. Furthermore, a high correlation was found between both average series (0.87). For WP, the analysis at basin scale shows that 75% of

- Eliminado: our
- Eliminado: promoted
- Eliminado: the
- Eliminado: into
- Eliminado: ,
- Eliminado: although
- Eliminado: estimate
- Con formato: Sin Resaltar
- Con formato: Sin Resaltar
- Con formato: Sin Resaltar
- Eliminado: for
- Eliminado: the BP in case of full
- Eliminado: , as well as in series where
- Eliminado: appear
- Con formato: Sin Resaltar
- Con formato: Sin Resaltar
- Con formato: Sin Resaltar

- Eliminado: performed
- Eliminado: to
- Eliminado: reduced
- Eliminado: the
- Eliminado: which means
- Eliminado: we found
- Eliminado: In case of

290 precipitation series have a BP in 1979, which is significantly detected for 58% of the series, generally in  
the southern/western IP (Fig. 5b). However, the BP detection differed for eastern/northern IP. Therefore,  
WP generally changed in 1979, and WWI also varied between 1978 and 1980. These results agree with  
those of Gómez-Martínez et al. (2018) and Halifa-Marín et al. (2021), who noticed the concomitant  
WWI/WP abrupt changes in the headwater of southern/eastern IP since 1980, while Guerreiro et al. (2014)  
295 do not always find the BP of WP in 1979 for western IP (e.g. Tagus basin). Different time slices were  
analysed in all these investigations, which could explain the divergences identifying the BP in WP/WWI  
series. In conclusion, it seems very likely that the reduction of precipitation provoked the WWI decrease.  
However, the average WP decrease was more important than the corresponding decrease of average  
WWI.

300 After characterizing the BP of WWI/WP, the RC was quantified at basin scale for both variables since  
1979/1980 (Fig. 6). So, the RC of WP ranged from -40% to 10% (Fig. 6a). 61% of catchments registered  
a significant RC of WP. A significant RC was observed in the central areas and western/southern IP.  
Those negative RCs also occurred where a significant post-1980 BP was not detected (i.e., northern IP).  
305 Conversely, a positive RC of WP was estimated over the eastern IP. For WWI, all catchments registered  
losses (Fig. 6b). The RC of WWI ranged from -60% to -3%. Major WWI decreases were observed in the  
headwater of Tagus, Jucar, and Segura basins (central-eastern IP, see Fig. 1). The magnitude of WWI  
losses is generally higher than the reductions of WP. That is, WWI also decreased where WP increased  
post-1980. This converse pattern suggests a poor or moderate relationship between WP and WWI in  
310 several NENWIRES basins. Vicente-Serrano et al. (2019) have already evidenced the poor relationship  
between climate and streamflow in basins of southern Spain. The results presented here match these  
conclusions, given that increases(decreases) of RC were found for WP(WWI) over southern IP. However,  
larger areas of Spain/IP registered WWI reductions caused by the abrupt decrease of WP (Fig. 4).  
Questions arise, however, about why an abrupt decrease of WP has been registered. Hence, once could  
315 wonder about the role that NAOi plays into WP changes. This question is analysed in the following  
subsection.

Eliminado: the

Eliminado: Our

Eliminado: . W

Eliminado: always

Eliminado: covers

Eliminado: higher than

Eliminado: (

Eliminado: ¶  
Figure 4. Panels show the sStandardized series from Mean average Wintertime Precipitation (WP, upper) and Mean Wintertime Water Inflows (WWI, lower) in the NENWIRES dataset, during the period 1952/2018. Bars represent negative (red) and positive (blue) values. Vertical blue/black lines represents the most probable change BP detected through the Pettitt's Homogeneity Test (Section 2.2.2). (break point, BP).¶

Eliminado: ere

Eliminado: The

Eliminado: .

Eliminado: in

Eliminado: generally

Eliminado: opposite

Eliminado: /

Eliminado: Our

Eliminado: matched with their

Eliminado: we found

Eliminado:

Eliminado:

Eliminado: there

Eliminado: we notice that


Eliminado: see

Eliminado: of

Eliminado: We thus

Eliminado: ed what role

Eliminado: , which



355 **Figure 6.** (a) Relative Change (RC) of WWI since 1979/1980; and (b) Id. for WP. The RC is computed  
 as Eq. 4 (Section 2.2.2). Symbols represent positive RC (filled triangle point-up) and negative RC (filled  
 triangle point down), being marked with a black outline the significant estimates. Contours of orography  
 (black) and IHB of continental Spain (purple) were also added (see Fig. 1).

360 **3.4 NAOi Enhancement Causing the Post-1980 Abrupt Decrease of Precipitation**

The previous results have shown that WP/WWI suddenly decreased since early 1980s. The scientific literature has already warned about the WP abrupt decreases in several Iberian basins, which were preliminary associated to the NAOi enhancement (Guerreiro et al., 2014, Gómez-Martínez et al., 2018, Halifa-Marín et al., 2021). So, a significant BP post-1980 in NAOi series was detected in this contribution (Appendix A, Fig. A1a). Average NAOi changed from -0.35 (1952/1979) to 0.38 (1980/2018). Almost identical composites of SLP/Z500/U/V-W between NAOi-/NAOi+ phases and before/after 1980 were obtained (Appendix A, Fig. A1b-c), which confirms that winters post-1980 generally presented NAOi+ phases. A higher frequency of NAOi+ should explain the WP declining over the IP, and its propagation in WWI records, according to the literature (Trigo et al., 2004). In fact, WP/WWI (Fig. 4) and NAOi

Eliminado: The

Eliminado: aforementioned

Eliminado: demonstrated


Eliminado: we also detected

Eliminado: We obtained a

Eliminado: registered

Eliminado: in

(Appendix A, Fig. A1a) have shown a significant BP since 1979/1980. Changes post-1980 in these variables are physically coherent with the behaviour of NAO. Also, WP/WWI highly correlated with the NAOi in the NENWIRES catchments (Fig. 7). The correlation WP/NAOi ranged from -0.75 to -0.1. Generally, these correlations were significant except in the case of eastern/northern IP (see Appendix A, Fig. A2). Higher correlations were found in those basins where BPs of WP were detected in 1979 (Fig. 7b).



**Figure 7.** Correlation between WWI (a) and WP (b) with NAOi (X axis) in relation to the RC of each variable (Y axis). Red squares show the significant correlation values. Black circles show the significant RC of each variable. Painted spheres show the BP detected through the same rules (colour palette) used in Fig. 5. Purple line/text shows the linear regression coefficients.

Likewise, a strong relationship between the NAOi/WP correlation and RC of WP was also found. A significant adjusted  $R^2 = 0.82$  was quantified through linear regression between both variables, which allows to attribute those larger decreases of WP to the NAOi post-1980 enhancement. Meanwhile, focusing on WWI/NAOi links, a similar spatial pattern (less intense) although non-significant positive correlations were observed over the eastern IP (Appendix A, Fig. A2). In this case, correlation coefficients

**Eliminado:** One the one hand, t

**Eliminado:** ¶

**Eliminado:** ¶

**Eliminado:** After characterizing the precipitation-driven shift of WWI, changes of WP/WWI were assessed (Fig. 6). The relative change of WP ranged from -40% to 10%. 61% of catchments registered a significant relative change of WP. WP losses generally appeared from 1980. The significant relative change was observed in the central areas and western/southern IP. Those negative relative changes also occurred where the breaking point was not detected (i.e., northern IP). Conversely, a positive relative change of WP was estimated in the eastern IP. For WWI, all catchments registered losses. The relative change of WWI ranged from -60% to -3%. Major WWI decreases were observed in the headwater of Tagus, Jucar, and Segura basins. These territories store an important part of water resources in the IP, but these WWI losses generally are higher than reductions of WP. That is, WWI decreased where the relative changes of WP increased. This incoherent pattern suggests a poor relationship between WP and WWI in several NENWIRES basins. ¶

**Eliminado:** ¶

**Figure 6.** Panels show (a) The Relative Change (RC) of WWIP since 1979/1980; and (b) Similarly for WPWI. The RC is computed as Eq. 2 (Section 2.2.2). Symbols represent positive RC (filled triangle point-up) and negative RC (filled triangle point-down), being marked with a black outline the significant estimates. Contours of orography (black) and IHB of continental Spain (purple) were also added (see Fig. 1). ¶

**Eliminado:** ¶

At this point, results are not conclusive about the incoherent WP/WWI changes through the study period: 1) an abrupt shift of WP was observed, causing the decline of WWI, but climate dynamics promoting the WP change were not clarified; 2) at the same time, an increase of WP was found, but WWI decreased in several catchments; and 3) larger losses were found in WWI than in WP. ¶

**Eliminado:** ¶

**Eliminado:** 3.3 Wintertime Iberian Meteorological Droughts caused by NAO+¶  
The aforementioned results have demonstrated that WP suddenly decreased since early 1980s. The scientific literature had already warned about the WP abrupt decreases in some IHB, which were linked to the large-scale influence (Guerreiro et al., 2014, Gómez-Martínez et al., 2018, Halifa-Marin et al., 2021). They noticed an abrupt shift of wintertime NAOi since late 1970s/early 1980s. In this sense, a significant break point for wintertime NAOi since 1979/1980. ¶ [68]

**Eliminado:** ¶

... [69]

**Eliminado:** 8

**Eliminado:** Panels show the c

**Eliminado:** in the

**Eliminado:**

**Eliminado:** The wintertime correlation between NAOi and (a) precipitation; and (b) water inflows. Symbols represent positive correlation (filled triangle point-up) and negative correlation (... [70]

**Con formato:** Superíndice

525 ranged from -0.6 to 0.2 (Fig. 7a). However, the relationship between NAOi/WP was clearly more intense  
than NAOi/WWI. Linear regression between WWI/NAOi correlation and RC of WWI shows a poor  
relationship. Therefore, the Iberian WWI abrupt reductions, understandably, depends on NAOi  
enhancement post-1980, whereas their magnitudes might not be essentially provoked by the WP  
declining.

530 Likewise, NAOi shift was the principal precursor of WP decreases, especially in areas severely affected  
by Atlantic fronts (e.g. precipitation events coming from Atlantic Ocean, where NAOi influence is crucial  
to the precipitation regime). Meanwhile, uncertainties about the drivers which have promoted the NAOi  
shift persist (Luo & Gong, 2006, Wang et al., 2014). If NAOi enhancement persists due to anthropogenic  
535 climate change (one of the key issues currently under discussion, e.g. Wang et al., 2014), the water  
resource could strongly be affected in those areas where NAOi+ increases the meteorological drought  
events/severity (e.g. IP, Southern Europe).

At this point, the results inspired the aim to fully understanding the causes of the post-1980 changes into  
540 WWI records, given that it has been depicted that several series were not driven by NAOi/WP changes  
(WWI declining shows inverse/higher magnitude than WP reductions). Also, the results of this  
contribution found similar divergences as noticed by state of the art literature. Whereas Gudmundsson et  
al. (2021) attributed the negative streamflow trends in the Mediterranean to warmer climate consequences  
(e.g. decreasing precipitation, increase of temperature), Vicente-Serrano et al. (2019) inferred that human-  
545 induced factors are sometimes more important than climate for understanding streamflow trends in Spain  
(e.g. irrigation). Moreover, Peña-Angulo et al. (2021) conclude that changes in vegetation have a strong  
impact on the relationship between climatic and hydrological drought over time. Henceforth, their  
conclusions advice to assess the contribution of other factors to the WWI changes in the following section.

550

### 3.4. Propagation of Persistent Meteorological Droughts into Water Inflows Series

The SPEI6/12 index was quantified in NENWIRES basins, in order to analyse how the drought conditions might impact on WWI records (Section 2.2.2). The average of wintertime SPEI6/12 is shown in Fig. 8.




Figure 8. (a) Mean wintertime SPEI12 registered in the NENWIRES catchments (top) and (b) SPEI6 (bottom). Blue(red) bars represent positive(negative) estimations. The most likely change point (BP), computed through the methods used in Fig. 4, is shown for both series (light blue and dark lines).

The BP was detected in 1980 (SPEI12) and 1970 (SPEI6). Clearly negative estimations turn more frequent in both series since its BP was detected. For SPEI12, the BP detection agreed with the BP of WWI/WP series. The average SPEI12 changed from 0.37 to -0.27 since 1980. Also, persistent drought conditions prevailed in 70% (25) of post-1980 winters. These results help understanding a higher WWI decline. In those basins where SPEI12 decreases (increase of meteorological droughts conditions), WP events probably need to supply the water stress of vegetation and lower water reserves in aquifers. A simultaneous WP decline was also reported from 1980 here. In addition, SPEI12 changes also refer to the

Eliminado: The Frequency of Meteorological Droughts

Eliminado: ere

Eliminado: the

Eliminado: In case of

Eliminado: coincided

Eliminado: the

Eliminado: winters

Eliminado: to understand

Eliminado: magnitude of

Eliminado: ing

Eliminado: ly

Eliminado: ing

Eliminado: in our results

580 increase of ETP promoted by the temperature rise. The decrease of SPEI12 estimations in headwaters of Spain already has been mentioned by the literature (Vicente-Serrano et al., 2014; Peña-Angulo et al., 2021). The mechanisms that could promote the droughts intensification are the decrease of cloud cover, insolation and maximum temperature rise due to NAOi enhancement. However, further work is needed to verify this assertion. In fact, an increase of maximum temperature (~0.9°C) from 1980 is generally reported in the NENWIRES catchments (Appendix A, Fig. A3). In addition, the potential impacts on the recharge/reserve in groundwaters due to the increase of permanent meteorological droughts (SPEI12) has been addressed.

Eliminado: M

Con formato: Fuente: 12 pto, Inglés (americano)

Con formato: Fuente: 12 pto, Inglés (americano)

Eliminado: we generally reported

590 To this end, the permeability of soils was evaluated. Continental Spain shows a wide variety of soils, with permeable(impermeable) soils prevailing over central/eastern(western) IP (Appendix A, Fig. A4). The permeable soils exceeded 80% in southeastern catchments, where limestone soils are abundant. Conversely, impermeable soils reached 100% over the western IP. So, the relationship between permeability of soils and WP/WWI correlation was assessed (Fig. 9a). These correlation coefficients ranged from 0.3 to 0.9. Generally, WP/WWI are highly correlated (0.8-0.9) where impermeable soils prevail. It is well-known that correlation between WP and WWI is more intense within impermeable basins because the run-off response is instantaneous (Lorenzo-Lacruz et al., 2013). Conversely, some permeable watersheds have registered poor correlation between WP and WWI.

Eliminado: The c

Eliminado: where

Eliminado: in the

Eliminado: In contrast

Eliminado: in

Eliminado: we assessed

Eliminado:

Eliminado: Is

Con formato: Sin Resaltar

500 After WP/WWI correlation was estimated, the QQ-Deviation (QQD) (Eq. 5, Fig. 9) was quantified (the reader is referred to Section 2.2.2 for further information about QOD quantification). A significant adjusted  $R^2$  above 0.35 is found in the linear regression analysis between QQD and WWI/WP correlation. So, the role of permeable soils to generate higher WWI was mostly proved during humid winters (Fig. 9a). Whereas QQD showed negative estimates over the northern/western IP (impermeable soils), positive estimates have been found in the high-permeable catchments (eastern/southern IP). The spatial pattern of QQD and WWI/WP correlation are shown in Fig. A5 of the Appendix A. These results can be summarized as follows: 1) porous watershed can infiltrate a larger volume of water; 2) Iberian extreme events of WP

Eliminado: 4

Eliminado: 6

Con formato: Normal;Texto

Eliminado: .

Eliminado: should be revised to understand the QQD quantification

Eliminado: We find a

Eliminado: the



are characterized by the persistence of rainy days, even for several weeks; 3) which allows the water accumulation into aquifers; 4) generating an underground baseline flow joint to the surface run-off,

**Con formato:** Fuente: Negrita, Color de fuente: Automático

**Eliminado:** To estimate the previous conditions of soil moisture and the water reserve of aquifers each winter, the changes of SPEI12 (Fig. 9) have been quantified. The results depict an increase of Persistent Moderate Droughts (PMD) conditions preceding the winter season from 1980. All catchments have recorded the increase of PMD events. The increase of PMD frequency ranged from 0.5 to 7.5 events/decade. Likewise, the rise of PMD was higher over central/southern IP. Also, the results prove the increases of Persistent Severe Droughts (PSD) events, ranging from -0.5 to 2.1 events/decade. The higher increase of PDS appeared on northern/western IP. These estimates highlight the positive variations of meteorological droughts conditions, which can affect to the hydrological response in the target catchments. Large increases of MMD events often occur where WP is limited to explain the reductions of WWI. The links between rising temperatures with increasing meteorological droughts has been unrevealed for the IP (Vicente-Serrano et al., 2014). The results here suggest that the increased temperature and ETP have been encouraged by the NAO shift, given that winter maximum temperatures significantly correlated with WP (-0.45).

**Eliminado:** the response between WP and WWI anomalies is higher in relation to common/dry winters. This time-lag effect probably depends on the hydrological response, water yield and the capacity of aquifers to store water. Likewise, we find a significant adjusted  $R^2$  above 0.2 between QQD and RC of WWI (Fig. 9b). In the same panel, we show that, precisely, the intensification post-1980 of persistent drought conditions (SPEI12) was higher in those basins which registered larger RC of WWI decreases and QQD is higher. These results suggest that those basins, where the groundwater reserve highly contributes to WWI, have been affected by the increase of droughts conditions. There, higher reductions post-1980 of WWI than WP are physically consistent because the implications of drought intensification for the hydrological response, especially under permeable conditions, where WWI also depends on groundwater reserve/flow. Future works will allow to confirm this assertion. In brief, our results suggest that changes of I parameter have influenced the WWI reductions, it seems, given that prevailing moisture content has decreased as result of precipitation losses at annual scale.

**Eliminado: 3.5. Permeable Soils Holding the Time-lag of Hydrological Response**

Two types of target catchments were found depending on its permeability characteristics: 1) those where permeable soils prevailing, which were mostly found over the central/eastern IP; and 2) those where impermeable soils prevailed (western IP). The permeable soils exceeded 80% in southeastern catchments (Fig. 10). In contrast, impermeable soils reached 100% in the western (... [71])

**Eliminado: 11**

**Eliminado:** The wintertime correlation between water inflows and precipitation; and (b) the QQ-deviation in the most humid monthly records (computed after Eq. 4). All correlation coefficients were significant. For QQ-deviation, symbols represent positive dev (... [72])


**Eliminado:** In case of

**Eliminado:** s

**Con formato:** Superíndice

**Eliminado:** ¶

625



**Figure 9.** (a) Symbols represent the relationship between Q-Q Deviation (X-axis) and Correlation estimated for WWI and WP (Y-axis), whereas are painted through the prevailing type of soils in the catchments (permeability characteristics). Types of soils depending on its permeability conditions are defined in Methods (Section 2.1.3). (b) Symbols represent links between Q-Q Deviation (X-axis) and RC of WWI (Y-axis), which are painted through the absolute change of Mean Wintertime SPEI12 before/after 1980. The purple line shows the linear regression between X-Y variables, which Adj.  $R^2$  and p-value are also mentioned.

630

The response between WP and WWI anomalies is higher in relation to common/dry winters. This time-lag effect probably depends on the hydrological response, water yield and the capacity of aquifers to store water. Likewise, a significant adjusted  $R^2$  above 0.2 between QQD and RC of WWI is found (Fig. 9b). The same panel shows that the intensification post-1980 of persistent drought conditions (SPEI12) was

635

higher in those basins which registered higher RCs of WWI decreases and a higher QQD. These results suggest that those basins, where the groundwater reserve highly contributes to WWI, have been affected by the increase of droughts conditions. There, higher reductions post-1980 of WWI than WP are physically consistent because the implications of drought intensification for the hydrological response, especially under permeable conditions (where WWI also depends on groundwater reserve/flow). Future works should be devoted to further research on this assertion. In brief, the results presented here suggest that changes of infiltration ( $I$  parameter in Eq. 1) have influenced the WWI reductions, given that prevailing moisture content has decreased as a result of precipitation losses at annual scale.

Con formato: Fuente: Cursiva

### 3.5 Does Land Greening-up Amplify the Water Inflows Decline?

The scientific literature has widely described that the forest extensions grew thanks to the cropland abandonment and afforestation works in Continental Spain (e.g. Peña-Angulo et al., 2021). Those authors also linked the revegetation of headwaters with an intensification of hydrological droughts. Therefore, in order to understand the higher magnitude of WWI losses than WP declining, the role of forest cover changes into the WWI variability (e.g. water generation) was further explored in the NENWIRES basins. There, the extension of forest cover was generally registered from 1950 to 2010 (Appendix A, Fig. A5b). The RC of forest cover ranged from -12% to 15%. Forest areas extended in 67% of the catchments through the study period (greening-up), whereas its cover mainly did not extent over the northern/southeastern IP. Also, the greening-up was limited where forest already exceeded 80% of watershed dimension in 1950 (Appendix A, Fig. A5a).

Eliminado: ¶

Con formato: Color de fuente: Automático

Eliminado: 6... ..Does LL

[73]

Eliminado: ing ...he Water Inflows Declining

[74]

Eliminado: Promoting the...Water Stress

[75]

Eliminado: L...terature has widely described that the forest extensions grew thanks to the cropland abandonment and afforestation works in the

[76]

Eliminado: Such

Eliminado: After having described most of the physical drivers... therefore, in order to understand the higher magnitude of WWI losses than WP declining, the role of land cover...rest cover changes into the NENWIRES hydrological response

[77]

Eliminado: The results widely registered ...t...ere, the extension of forest cover was generally registered from 1950 to 2010 (Appendix A, Fig. 1...5b2.... The relative changes...C of forest cover ranged from -60...2% to 90...5% until 2010... Those forest extensions grew thanks to the cropland abandonment and afforestation works. ...rom 1950 to 2010, f...rest areas extended in 67% of the catchments through the study period (greening-up), whereas, Forest areas

[78]

Eliminado: in the

Eliminado: area ...imension in 1950 (Appendix A, Fig. A5a)

[79]

Gains of forest cover mostly occurred in the semiarid basins (Fig. 10a,b) where lower mean WP is recorded. Meanwhile the most humid basins registered a reduction of forest cover. This agrees to the fact that higher dimension of forest cover impacts on ETR and run-off, as shown by previous works (e.g. García-Ruiz et al., 2011, Teuling et al., 2019). So, it is reasonable to assume that the greening-up has contributed to increase the reduction of WWI in NENWIRES catchments, especially in semiarid environments. This assertion agrees with the results of Peña-Angulo et al. (2021), who confirmed the

Eliminado: Our results show that g

Eliminado: ...where lower mean WP is recorded. Meanwhile the most humid basins registered the... reduction of forest cover. Literature...his agrees to conclude...he fact that higher dimension of forest cover impacts on ETR and run-off, as shown by previous works ...

[80]

Eliminado: , so ...t is not unreasonable...easonable to assume that the greening-up has contributed to increase the reduction of WWI in NENWIRES catchments, especially in semiarid environments. This assertion agrees to

[81]


Con formato: Sin Resaltar

implications of forest extension in the occurrence of hydrological droughts more intense than simultaneous meteorological droughts. Those authors do not discriminate between basins based on their precipitation regime. Therefore, the results shown here warn of the potential  $EI$  gains (see Eq. 1) ( $E$  parameter) as a consequence of the greening-up (e.g. extension of forest) that has been observed into NENWIRES catchments, which coexists with the temperature rise as a result of global warming.

In brief, one of the main results of this contribution is that a higher magnitude of WWI reductions is found in relation to WP losses, leading to the assumption that WP was not the only cause for the WWI decline.

So, basically, this study evaluated how several causes ( $E$  and  $I$  parameters in Eq. 1) have contributed to post-1980 sudden losses of Iberian WWI. To understand how each of them modulates the WWI changes, a K-Means clustering was conducted. The clustering shows catchments where WWI changes are explained by similar mechanisms/factors (Fig. 10f). 4 clusters that differed in contribution of each precursor are identified (Appendix A, Table A2). Cluster 1 (C1) consist of 11.1% of dataset, which are well differentiated from other NENWIRES basins. These basins 1) registered the lower mean WP (164.7 mm/yr), 2) had the most important extension of forest cover, 3) registered a very poor correlation of WWI with NAOi, 4) presented a lower intensification of droughts (SPEI12 decrease), 5) had high permeable soils and 6) showed a higher magnitude of WWI reductions in relation to WP losses (-32.4%). It seems that the WP losses and forest extension drove the higher magnitude of WWI reductions in those basins.

- Con formato: Sin Resaltar
- Con formato: Sin Resaltar
- Eliminado: Whereas they
- Eliminado: distinct
- Con formato: Sin Resaltar
- Eliminado: our
- Eliminado: ed
- Con formato: Fuente: Cursiva
- Con formato: Fuente: Cursiva
- Eliminado: ¶
- Eliminado: we have found
- Eliminado: which allow
- Eliminado: did not only provoke the
- Eliminado: ing
- Eliminado: our
- Eliminado: consisted of evaluating
- Con formato: Fuente: Cursiva
- Con formato: Fuente: Cursiva
- Eliminado: into the RR equation
- Con formato: Inglés (americano)
- Con formato: Inglés (americano)
- Eliminado: In order to
- Con formato: Inglés (americano)
- Eliminado: performed
- Con formato: Sin Resaltar
- Con formato: Sin Resaltar
- Con formato: Sin Resaltar
- Eliminado: We identify
- Eliminado: The
- Eliminado: another
- Eliminado: registered
- Eliminado: ,
- Eliminado: ,
- Eliminado: WWI has
- Eliminado: ,
- Eliminado: conducted



**Figure 10.** Scatter plot of relative changes of WP (X axis) and WWI (Y axes). Symbols are filled by the magnitude of several variables: (a) mean WP; (b) RC of forest cover computed after Eq. 6; (c) absolute change of SPEI12; (d) correlation between WWI and NAOi; (e) percentage of permeable soils; and (f) K-Means clustering depending on abovementioned variables.

**Eliminado:** Finally, the results indicate that the lower mean WP, the larger forest extension through the study period. In this sense, the significant role of land greening-up to decrease the WWI in semiarid catchments was proved. On the other hand, the pattern of WWI changes differed. The relative change of WWI ranged from -65% to 17% under the semiarid conditions; from -45% to -2% under transitional conditions; and from -28% to -19% under humid conditions (Fig. 13). The relative changes of WI were higher than the decline of WP over semiarid catchments. A high correlation between the total relative changes of forest area from 1950 to 2010 and mean WP was found (-0.68). The total relative changes of forest area were higher over semiarid basins. Also, a larger decrease of relative changes of WWI was quantified related to those of WP, precisely over semiarid catchments. The results clearly depict how the vegetation control of WWI changes. The magnitude of the relative changes for WP/WWI is similar in the transitional and humid catchments. However, semiarid environment, where forest widely extended, registered a higher relative change of WWI. Furthermore, the NAOi cannot be related to the hydroclimate variability in these watersheds. Therefore, the links with NAO are not robust in the semiarid catchments, whereas a modulation of WWI reductions due to NAOi was observed in most of target basins. ¶

**Eliminado:** Panels show the s

**Eliminado:** with

**Eliminado:** RC

**Eliminado:** x

**Eliminado:** y

**Con formato:** Fuente: Negrita

**Eliminado:** (a),

**Eliminado:** b

**Eliminado:** .

**Eliminado:** (c)

**Eliminado:**

**Eliminado:** (d),

**Eliminado:**

**Eliminado:** (e),

**Eliminado:** (f)

**Eliminado:** ¶

The ETP is assumed to increase due to forest extension, while SPEI12 quantifications did not intensify since 1980. So, although the WP does not change significantly, WWI reduced by -29.6% because the rise of outputs affecting the water yield. However, hydrological modelling is needed to confirm this assertion.

Cluster 2 (C2) covers the 16.7% of basins where 1) mean WP is 355.2 mm/yr; 2) permeable soils are abundant (70.1%); 3) WWI highly correlated with NAOi; 4) SPEI12 severely decreased; 5) forest cover extended 2.5%; and 6) WWI reduced more than WP (-10.8%). WWI changes are seemingly provoked by NAOi enhancement, whereas the magnitude of reduction was intensified by the amplification of permanent meteorological droughts since 1980, which affected to the groundwater reserves, given that C2 consist of, generally, permeable soils. Clearly, the extension of forest also contributes to reduce WWI, as suggested for C1 basins, but in the case of C2, forest extension was less important.

Likewise, Cluster 3 (C3) includes 27.7% of catchments where 1) mean WP is 232.4 mm/yr; 2) permeable soils are less frequent (40.5%); 3) WWI correlated with NAOi (-0.4); 4) SPEI12 decreased -0.9 since 1980; 5) the extension of forest cover was important (4.7%); and 6) WWI reduced more than WP (-17.5%). In this group of basins, the higher reduction of WWI is driven by the same processes as for C2, but C3 registered higher intensification of SPEI12 (duration of meteorological droughts) and extension of forest. Oppositely, the influence of NAOi is less important for C3. Also, the RC of WP was weaker.

Finally, Cluster 4 (C4) covers 44.4% of basins. Most humid basins are grouped into C4, where 1) mean WP is 445.2 mm/yr; 2) soils are mainly not permeable; 3) WWI highly correlated with NAOi (-0.5); 4) SPEI12 decreased -0.5; 5) forest cover does not change (-0.1); and 6) WWI changed more than WP (-7.1), however the magnitude of change was almost similar for both variables. Clearly, in the C4 basins, WWI changes were driven by NAOi enhancement and amplification of meteorological droughts.

Likewise, given that C4 basin are generally impermeable, the groundwater reserve cannot supply the decrease of surface run-off during dry events.

Eliminado: We assume that

Eliminado: ETR increased

Eliminado: affected to the

Eliminado: works

Eliminado: The

Eliminado: joint

Eliminado: ,

Eliminado: ,

Eliminado: ,

Eliminado: i,

Eliminado: ,

Eliminado: ,

Eliminado: It seems that

Eliminado: like it was

Eliminado: the

Eliminado: consist

Eliminado: of

Eliminado: ,

Eliminado: ,

Eliminado: ,

Eliminado: ,

Eliminado: it seems that

Eliminado: conducted

Eliminado: than

Eliminado: influence

Eliminado: , as well as

Eliminado: the

Eliminado: consist

Eliminado: of

Eliminado: ,

Eliminado: ,

Eliminado: higly

Eliminado: ,

Eliminado: ,

Eliminado: ,

Eliminado: conducted

Eliminado: ¶  
Figure 13. The RC of WWI (bottom right), the RC of WWI minus RC of WP (top left), the RC of forest area from 1950 to 1980 (top right), and NAOi (bottom left) were related to the MWP of ... [82]

Con formato: Color de fuente: Automático

#### 4. Final Remarks

This contribution focused mainly on disentangling the abrupt WWI reduction post-1980, echoing the call made in Halifa-Marín et al. (2021), for assessing the potential propagation of abrupt changes in WP records into streamflow records. Therefore, the main findings of this contribution can be summarized as:

- 1) The NENWIRES dataset was created to analyze the recent evolution of near-natural Iberian streamflow draining to reservoirs in headwater catchments (NENWIRES). We identify a significant reduction of the WWI, that is related to an abrupt change after 1980 for most of basins. This change agrees with the abrupt change reported for WP by other authors and confirmed in this contribution. This allows to analyze the changes reported here as the differences between two periods (1952/1979-1980/2018).
- 2) The decrease of WP is the main driver of the decline of WWI. However, a higher magnitude of WWI losses rather than WP reductions has been generally quantified. These extra water losses depend on the extension of vegetation ( $E$  parameter in Eq. 1), and moisture content of soils (e.g. its permeability characteristics,  $I$  parameter in Eq. 1), which, in turn, also differed between the variety of climate conditions (e.g. precipitation regime).
- 3) A strong reduction of WWI has been observed in a set of semiarid catchments, where WP slightly increased, and where the NAOi does not exert an important influence. Those basins (grouped in the C1 cluster) are characterized by permeable soils, which are affected by the most important extension of vegetation. So, it is assumed that this greening-up in semiarid environments, where water storage of soils is abundant, has provoked significant evapotranspiration rises. Consequently, those water losses have decreased the WWI records. Similar assertions are concluded for catchments classified in the C3 cluster, where, however, WP is promoted by the NAOi influence, although they also are semiarid environments. C3 catchments registered the higher decreases of WWI.

**Eliminado:** Whereas Gudmundsson et al. (2021) linked the strongest declining of water resources in the Mediterranean to anthropogenic global warming, Vicente-Serrano et al. (2019) attributed the lead role of human-induced impacts to the decrease of streamflow in southern Europe. According to these latter authors, the extension of irrigated areas, agricultural intensification, and natural revegetation of marginal lands of mountain areas were the main drivers of decreases in streamflow, specially over the southern IP. Peña-Angulo et al. (2021) have also demonstrated the intensification of meteorological droughts due to the vegetation extension in the IP. A similar conclusion was presented for the Mediterranean basin (García-Ruiz et al., 2011) and Europe (Teuling et al., 2019). The NENWIRES dataset were created to shed light on the observed variability of near-natural Iberian hydrological resources dur... [83]

**Eliminado:** 5... Conclusions ... [84]

**Eliminado:** We...his contribution particularly ... [85]

**Eliminado:** This work has assessed the long-term variability ... [86]

**Eliminado:** summarizing, ...he main findings of this contri... [87]

**Eliminado:** analyse ... [88]

**Con formato:** ... [89]

**Eliminado:** is coincident ... [90]

**Eliminado:** for ... [91]

**Eliminado:** also ...n this work ... [92]

**Con formato:** ... [93]

**Con formato:** ... [94]

**Eliminado:** permits us ... [95]

**Con formato:** ... [96]

**Con formato:** ... [97]

**Con formato:** Color de fuente: Texto 1 ... [98]

**Con formato:** Párrafo de lista ... [99]

**Eliminado:** declining ... [100]

**Eliminado:** we generally have quantified ... [101]

**Con formato:** ... [102]

**Con formato:** Inglés (americano) ... [103]

**Con formato:** ... [104]

**Eliminado:** , it seems,... [105]

**Eliminado:** into the RR estimation...n Eq. 1), and moist... [106]

**Con formato:** Fuente: Cursiva ... [107]

**Con formato:** Color de fuente: Texto 1 ... [108]

**Con formato:** Normal;Texto, Sin viñetas ni numeración ... [109]

**Eliminado:** S ... [110]

**Eliminado:** , ... [111]

**Eliminado:** registerted ... [112]

4) Another set of most humid catchments (C4 cluster) is characterized by impermeable soils, and unchanged vegetation cover. There, the magnitude of WWI and WP reductions is almost similar. Meanwhile, the WP of catchments grouped into C2 cluster is characterized by a strong influence of the NAOi, and the extension of vegetation. Then, higher reductions of WWI are observed in relation to C4 catchments.

These assertions allow to conclude that a higher mean precipitation induces a minor role of evapotranspiration/infiltration losses into streamflow changes. Nonetheless, water generation in semiarid catchments is widely affected by those losses, which can play the main role, as important as the decrease of precipitation. In terms of Eq. 1, in case of lower values of precipitation ( $P$ ), the importance of  $E_t$  (vegetation cover) and  $I$  (water storage) is large. When  $P$  is higher,  $RR$  is almost similar as  $P$ , as well as when  $E$  and  $I$  are weaker.

Last, the conclusions to this contribution confirm that the initial hypothesis was not accurate, since WWI changes essentially do not depend on climate in the set of NENWIRES catchments. Abrupt changes do not impact on water planning in the same way as gradual changes, so that policies adapted to this mode of climate variability could be necessary. Future works will have to deal with the improvement of the scientific knowledge about shifts of the NAOi and their implications to hydrological resources in southern Europe. The findings presented here thus encourage the need to develop a deeper knowledge about NAOi variability under a warmer climate and conduct high-resolution modelling considering the water losses because of vegetation extension and moisture-soil content, especially in semiarid environments where the availability of freshwater is crucial. Henceforth, this work can contribute to the mitigation of warmer climate impacts on water cycle/planning in the Mediterranean area, the JP or other global semiarid environments.

Data availability. All data sets used in the current study are publicly available from the indicated references or sources (See assets in the doi of manuscript). Also, we will be pleased to send the NENWIRES dataset under request.

- Eliminado: Other
- Eliminado: characterised
- Con formato: Color de fuente: Texto 1, Inglés (americano)
- Con formato: Color de fuente: Texto 1, Inglés (americano)
- Con formato: Inglés (americano)
- Con formato: Sangría: Izquierda: 1,27 cm, Sin viñetas ni numeración
- Eliminado: greater
- Eliminado: principal
- Eliminado: decreases
- Eliminado: the RR equation
- Eliminado: , which was defined in the introduction
- Eliminado: P
- Con formato: Fuente: Cursiva
- Con formato: Fuente: Cursiva
- Eliminado: than
- Con formato: Fuente: Cursiva
- Con formato: Fuente: Cursiva
- Con formato: Fuente: Cursiva
- Con formato: Fuente: Cursiva
- Eliminado: †
- Con formato: Fuente: Cursiva
- Con formato: Fuente: Cursiva
- Eliminado: In the end
- Eliminado: conclusions
- Eliminado: our
- Eliminado: wrong
- Eliminado: because
- Eliminado: Whereas a
- Eliminado: improve
- Eliminado: shifts
- Eliminado: Our
- Eliminado: better
- Eliminado: in
- Eliminado: the
- Eliminado: we hope that our work
- Eliminado: /
- Con formato: Color de fuente: Texto 1
- Eliminado: <#>WP decline has shown a lower magnitu (... [99])
- Eliminado: full



*Author contributions.* A.H-M conceived the original idea and designed the overall study. A.H-M, E.P-S and M.T-V developed the NENWIRES dataset. A.H-M, and JP.M performed the analysis. All co-authors contributed to the interpretation of the results. A.H-M led the writing of the paper, with contributions of

310 P.J-G and JP.M.

Eliminado: ¶

*Competing interests.* The authors declare that they have no conflict of interest.

315 *Acknowledgements.* The authors thank the reviewers and the editor of the manuscript for their valuable contributions and fruitful discussions.

**Eliminado:** The authors acknowledge the ACEX project (CGL2017-87921-R) of the *Ministerio de Economía y Competitividad/Agencia Estatal de Investigación* of Spain, the ECCE project (PID2020-115693RB-I00) of *Ministerio de Ciencia e Innovación/Agencia Estatal de Investigación* (MCIN/AEI/10.13039/501100011033/) and the European Regional Development Fund (ERDF/ FEDER *Una manera de hacer Europa*). Also, t

320 *Funding.* The authors acknowledge the ECCE project (PID2020-115693RB-I00) of *Ministerio de Ciencia e Innovación/Agencia Estatal de Investigación* (MCIN/AEI/10.13039/501100011033/). A.H-M thanks his predoctoral contract FPU18/00824 to the *Ministerio de Ciencia, Innovación y Universidades* of Spain. E.P-S thanks for his predoctoral contract to the ACEX project. M.L-C thanks his predoctoral contract FPU17/02166 to the *Ministerio de Ciencia, Innovación y Universidades* of Spain.

**Eliminado:** A.H-M thanks his predoctoral contract FPU18/00824 to the *Ministerio de Ciencia, Innovación y Universidades* of Spain. E.P-S thanks for his predoctoral contract to the ACEX project. M.L-C thanks his predoctoral contract FPU17/02166 to the *Ministerio de Ciencia, Innovación y Universidades* of Spain.

**Con formato:** Fuente: Cursiva

325 *Review statement.* This paper was edited by Erwin Zehe and reviewed by A.J. Teuling and one anonymous referee.

Eliminado: ¶

**Con formato:** Fuente: Times New Roman, 12 pto, Color de fuente: Texto 1, Inglés (americano)

**Con formato:** Color de fuente: Texto 1, Inglés (americano)

**Con formato:** Fuente: Times New Roman, 12 pto, Color de fuente: Texto 1, Inglés (americano)

**Con formato:** Fuente: Times New Roman, 12 pto, Color de fuente: Texto 1, Inglés (americano)

**Con formato:** Fuente: Times New Roman, 12 pto, Color de fuente: Texto 1, Inglés (americano)

## 325 References

AEMET, Precipitation/Temperature Gridded Dataset, available at: [http://www.aemet.es/es/serviciosclimaticos/cambio\\_climat/datos\\_diarios?w=2](http://www.aemet.es/es/serviciosclimaticos/cambio_climat/datos_diarios?w=2), last access: 20 October 2021.

330 Berguería, S., Vicente-Serrano, S.M.: SPEI: Calculation of the Standardised Precipitation-Evapotranspiration Index, R package version 1.7, available at: <https://cran.r-project.org/web/packages/SPEI/index.html>, last access: 20 October 2021.

- Bivand, R.: Rgdal: Bindings for the 'Geospatial' Data Abstraction Library, R package version 3.4.13, available at: <https://cran.r-project.org/web/packages/rgdal/index.html>, last access: 20 October 2021.
- Borchers, H.W.: Pracma: Practical Numerical Math Functions, R package version 2.2.9, available at:  
350 <https://CRAN.R-project.org/package=pracma>, last access: 20 October 2021.
- CEDEX, Spanish Hydrological Repository, available at:  
<https://ceh.cedex.es/anuarioaforos/demarcaciones.asp>, last access: 20 October 2021.
- de Luis, M., Brunetti, M., González-Hidalgo, J. C., Longares, L. A., & Martín-Vide, J.: Changes in  
seasonal precipitation in the Iberian Peninsula during 1946-2005, *Global and Planetary Change*,  
355 74, 27-33, doi:10.1016/j.gloplacha.2010.06.006, 2010.
- Fuchs, R., Herold, M., Verburg, P. H., Clevers, J. G. P. W., and Eberle, J.: Gross Changes in  
Reconstructions of Historic Land Cover/Use for Europe between 1900 and 2010, *Glob. Change  
Biol.*, 21, 299-313, <https://doi.org/10.1111/gcb.12714>, 2015.
- García-Ruiz, J. M., López-Moreno, J. I., Vicente-Serrano, S. M., Lasanta-Martínez, T., & Beguería, S.:  
360 Mediterranean water resources in a global change scenario, *Earth-Science Reviews*, 105(3-4), 121-  
139, <https://doi.org/10.1016/j.earscirev.2011.01.006>, 2011.
- Gómez-Martínez, G., Pérez-Martín, M. A., Estrela-Monreal, T., & del-Amo, P.: North Atlantic  
Oscillation as a Cause of the Hydrological Changes in the Mediterranean (Júcar River, Spain),  
*Water Resources Management*, 32(8), 2717-2734, doi:10.1007/s11269-018-1954-0, 2018.
- 365 Gudmundsson, L., Boulange, J., Do, H. X., Gosling, S. N., Grillakis, M. G., Koutroulis, A. G., ... &  
Zhao, F.: Globally observed trends in mean and extreme river flow attributed to climate change,  
*Science*, 371(6534), 1159-1162, 2021.
- Guerreiro, S. B., Kilsby, C. G., & Serinaldi, F.: Analysis of time variation of rainfall in transnational  
basins in Iberia: Abrupt changes or trends?, *International Journal of Climatology*, 34, 114-133,  
370 doi:10.1002/joc.3669, 2014.
- Halifa-Marín, A., Lorente-Plazas, R., Pravia-Sarabia, E., Montávez, J. P., & Jiménez-Guerrero, P.  
(2021). Atlantic and Mediterranean influence promoting an abrupt change in winter precipitation  
over the southern Iberian Peninsula, *Atmospheric Research*, 253, 105485,  
<https://doi.org/10.1016/j.atmosres.2021.105485>, 2018.

Código de campo cambiado

Código de campo cambiado

Código de campo cambiado

- 375 Hannaford, J., Buys, G., Stahl, K., and Tallaksen, L. M.: The influence of decadal-scale variability on trends in long European streamflow records, *Hydrol. Earth Syst. Sci.*, 17, 2717–2733, <https://doi.org/10.5194/hess-17-2717-2013>, 2013.
- Hijmans, R.J.: Raster: Geographic Data Analysis and Modeling, R package version 3.4.13, available at: <https://cran.r-project.org/web/packages/raster/index.html>, last access: 20 October 2021.
- 380 HILDA, HIstoric Land Dynamics Assessment Dataset, available at: <https://www.wur.nl/en/Research-Results/Chair-groups/Environmental-Sciences/Laboratory-of-Geo-information-Science-and-Remote-Sensing/Models/Hilda.htm>, last access: 20 October 2021.
- Hothorn, T., Winell, H., Hornik, K., van de Wiel, M.A., Zeileis, A.: Coin: Conditional Inference Procedures in a Permutation Test Framework, R package version 1.3-1, available at: <https://CRAN.R-project.org/package=coin>, last access: 20 October 2021.
- 385 IDE, Spanish Catchment Boundaries Dataset, available at: <https://www.miteco.gob.es/es/cartografia-y-sig/ide/descargas/agua/cuencas-y-subcuencas.aspx>, last access: 20 October 2021.
- IGME, Mid-Resolution Permeability of Soils Dataset, available at: [http://mapas.igme.es/Servicios/default.aspx#IGME Permeabilidad 1M](http://mapas.igme.es/Servicios/default.aspx#IGME%20Permeabilidad%201M), last access: 20 October 2021.
- 390 Lorenzo-Lacruz, J., Vicente-Serrano, S. M., López-Moreno, J. I., Morán-Tejeda, E., and Zabalza, J.: Recent Trends in Iberian Streamflows (1945–2005), *J. Hydrol.*, 414–415, 463–475, <https://doi.org/10.1016/j.jhydrol.2011.11.023>, 2012.
- Lorenzo-Lacruz J, Vicente-Serrano SM, González-Hidalgo JC, López-Moreno JI, Cortesi N.: Hydrological drought response to meteorological drought in the Iberian Peninsula, *Clim Res* 58:117-131, <https://doi.org/10.3354/cr01177>, 2013.
- 395 Luo, D., & Gong, T.: A possible mechanism for the eastward shift of interannual NAO action centers in last three decades, *Geophysical research letters*, 33(24). <https://doi.org/10.1029/2006GL027860>, 2006.
- 400 Massei, N., Kingston, D. G., Hannah, D. M., Vidal, J. P., Dieppois, B., Fossa, M., ... & Laignel, B.: Understanding and predicting large-scale hydrological variability in a changing environment,

Código de campo cambiado

Código de campo cambiado

Código de campo cambiado

Código de campo cambiado

Código de campo cambiado

Código de campo cambiado

Proceedings of the International Association of Hydrological Sciences, 383, 141-149.  
<https://doi.org/10.5194/piahs-383-141-2020>, 2020.

NOAA, The North Atlantic Oscillation Index, available at:  
 405 <https://www.cpc.ncep.noaa.gov/products/precip/CWlink/pna/nao.shtml>, last access: 20 October 2021.

410 [Peña-Angulo, D., Vicente-Serrano, S. M., Domínguez-Castro, F., Noguera, I., Tomas-Burguera, M., López-Moreno, J. I., ... & El Kenawy, A.; Unravelling the role of vegetation on the different trends between climatic and hydrologic drought in headwater catchments of Spain. \*Anthropocene\*, 36, 100309, <https://doi.org/10.1016/j.ancene.2021.100309>, 2021](#)

Pohlert, T.: Trend: Non-parametric Trend Tests and Change-Point Detection, R package version 1.1.1, available at: <https://CRAN.R-project.org/package=trend>, last access: 20 October 2021

Polade, S.D., Gershunov, A., Cayan, D.R. et al.: Precipitation in a warming world: Assessing projected hydro-climate changes in California and other Mediterranean climate regions, *Sci Rep* 7, 10783,  
 415 <https://doi.org/10.1038/s41598-017-11285-y>, 2017.

R Core Team: A language and environment for statistical computing, R Foundation for Statistical Computing, at available at: <https://rdrr.io/r/base/base-package.html>, last access: 20 October 2021.

Schulzweida, U.: CDO user guide, Climate Data Operator, available at:  
<https://code.mpimet.mpg.de/projects/cdo/embedded/index.html>, last access: 20 October 2021.

420 Stahl, K., Hisdal, H., Hannaford, J., Tallaksen, L. M., van Lanen, H. A. J., Sauquet, E., Demuth, S., Fendekova, M., and Jódar, J.: Streamflow trends in Europe: evidence from a dataset of near-natural catchments, *Hydrol. Earth Syst. Sci.*, 14, 2367–2382, <https://doi.org/10.5194/hess-14-2367-2010>, 2010.

Teuling, A. J., de Badts, E. A. G., Jansen, F. A., Fuchs, R., Buitink, J., Hoek van Dijke, A. J., and  
 425 Sterling, S. M.: Climate change, reforestation/afforestation, and urbanization impacts on evapotranspiration and streamflow in Europe, *Hydrol. Earth Syst. Sci.*, 23, 3631–3652, <https://doi.org/10.5194/hess-23-3631-2019>, 2019.

Tramblay, Y., Llasat, M.C., Randin, C. et al.: Climate change impacts on water resources in the Mediterranean, *Reg Environ Change*, 20, 83, <https://doi.org/10.1007/s10113-020-01665-y>, 2020.

Con formato: Color de fuente: Texto 1

Con formato: Color de fuente: Texto 1, Inglés (americano)

Con formato: Color de fuente: Texto 1, Inglés (americano)

Con formato: Fuente: Sin Cursiva, Color de fuente: Texto 1, Inglés (americano)

Con formato: Color de fuente: Texto 1, Inglés (americano)

Con formato: Fuente de párrafo predeter., Color de fuente: Texto 1, Inglés (americano)

Código de campo cambiado

Eliminado: PSL, NCEP/NCAR Reanalysis 1 Repository, available at: <https://psl.noaa.gov/data/gridded/data.ncep.reanalysis.html>. last access: 20 October 2021.

Código de campo cambiado

Código de campo cambiado

- Trigo, R. M., Pozo-Vázquez, D., Osborn, T. J., Castro-Díez, Y., Gámiz-Fortis, S., & Esteban-Parra, M.  
435 J.: North Atlantic oscillation influence on precipitation, river flow and water resources in the Iberian  
Peninsula, *International Journal of Climatology*, 24(8), 925–944, doi:10.1002/joc.1048, 2004.
- Vicente-Serrano, S. M., Lopez-Moreno, J. I., Beguería, S., Lorenzo-Lacruz, J., Sanchez-Lorenzo, A.,  
García-Ruiz, J. M., ... & Espejo, F.: Evidence of increasing drought severity caused by temperature  
rise in southern Europe, *Environmental Research Letters*, 9(4), 044001,  
440 <https://doi.org/10.1088/1748-9326/9/4/044001>, 2014.
- Vicente-Serrano, S. M., Peña-Gallardo, M., Hannaford, J., Murphy, C., Lorenzo-Lacruz, J., Dominguez-  
Castro, F., ... & Vidal, J. P.: Climate, irrigation, and land cover change explain streamflow trends  
in countries bordering the northeast Atlantic, *Geophysical Research Letters*, 46(19), 10821-10833,  
<https://doi.org/10.1029/2019GL084084>, 2019.
- 445 Wang, Y. H., Magnusdottir, G., Stern, H., Tian, X., & Yu, Y.: Uncertainty estimates of the EOF-derived  
North Atlantic Oscillation, *Journal of climate*, 27(3), 1290-1301. <https://doi.org/10.1175/JCLI-D-13-00230.1>, 2014.

Código de campo cambiado

Código de campo cambiado

Eliminado: ¶