

# 1 Spatiotemporal responses in crop water footprint and benchmark 2 under different irrigation techniques to climate change scenarios in 3 China

4 Zhiwei Yue<sup>1,3,\*</sup>, Xiangxiang Ji<sup>1,3,\*</sup>, La Zhuo<sup>2,3,4,5</sup>, Wei Wang<sup>4,5</sup>, Zhibin Li<sup>4,5</sup>, Pute Wu<sup>2,3,4,5</sup>

5 <sup>1</sup>College of Water Resources and Architectural Engineering, Northwest A&F University, Yangling 712100, China

6 <sup>2</sup>Institute of Soil and Water Conservation, Northwest A&F University, Yangling 712100, China

7 <sup>3</sup>Institute of Water-saving Agriculture in Arid Regions of China, Northwest A&F University, Yangling 712100, China

8 <sup>4</sup>Institute of Soil and Water Conservation, Chinese Academy of Sciences and Ministry of Water Resources, Yangling 712100,  
9 China

10 <sup>5</sup> University of Chinese Academy of Sciences, Beijing 100049, China.

11

12 \*The authors contribute equally.

13 Correspondence to: La Zhuo ([zhuola@nwafu.edu.cn](mailto:zhuola@nwafu.edu.cn); [lzhuo@ms.iswc.ac.cn](mailto:lzhuo@ms.iswc.ac.cn)) and Pute Wu ([gjzwpt@hotmail.com](mailto:gjzwpt@hotmail.com))

14

15 **Abstract.** Adaptation to future climate change with limited water resources is a major global challenge to sustainable and  
16 sufficient crop production. However, the large-scale responses of crop water footprint and its associated benchmarks under  
17 various irrigation techniques to future climate change scenarios remain unclear. The present study quantified the responses of  
18 maize and wheat water footprint per unit yield (WF,  $\text{m}^3 \text{t}^{-1}$ ) and corresponding WF benchmarks under two representative  
19 concentration pathways (RCPs) in the 2030s, 2050s, and 2080s at a 5-arc minute grid level in China. The AquaCrop model  
20 with the outputs of six global climate models in Coupled Model Intercomparison Project Phase 5 (CMIP5) as its input data  
21 was used to simulate the WF of maize and wheat. The differences among rain-fed and furrow-, micro-, and sprinkler-irrigated  
22 wheat and maize were identified. Compared with the baseline year (2013), maize WF will increase under both RCP2.6 and  
23 RCP8.5, by 17 % and 13 %, respectively, until the 2080s. Wheat WF will increase under RCP2.6 (by 12 % until the 2080s)  
24 and decrease by 12 % under RCP8.5 until the 2080s, with a higher increase in wheat yield and decrease in wheat WF due to  
25 the higher  $\text{CO}_2$  concentration in 2080s under RCP8.5. WF will increase the most for rain-fed crops. Relative to rain-fed crops,  
26 micro irrigation and sprinkler irrigation result in the smallest increases in WF for maize and wheat, respectively. These water-  
27 saving managements will mitigate the negative impact of climate change more effectively. The WF benchmarks of maize and  
28 wheat in the humid zone (~overall average at  $680 \text{ m}^3 \text{ t}^{-1}$  for maize and  $873 \text{ m}^3 \text{ t}^{-1}$  for wheat at 20<sup>th</sup> percentile) are 13–32 %  
29 higher than those in the arid zone (~ overall average at  $601 \text{ m}^3 \text{ t}^{-1}$  for maize and  $753 \text{ m}^3 \text{ t}^{-1}$  for wheat). The differences in WF  
30 benchmarks among various irrigation techniques are more significant in the arid zone, which can be as high as 57%, for 20<sup>th</sup>  
31 percentile WF benchmarks of  $1020 \text{ m}^3 \text{ t}^{-1}$  for sprinkler-irrigated wheat and  $648 \text{ m}^3 \text{ t}^{-1}$  for micro-irrigated wheat. Nevertheless,  
32 WF benchmarks will not respond to climate changes as dramatically as the WF in the same area, especially in the area with  
33 limited agricultural development. The present study demonstrated that the visible different responses to climate change in  
34 terms of crop water consumption, water use efficiency, and WF benchmarks under different irrigation techniques cannot be

35 ignored. It also lays the foundation for future investigations into the influences of irrigation methods, RCPs, and crop types on  
36 WF and its benchmarks in response to climate change in all agricultural regions worldwide.

## 37 **1 Introduction**

38 The progressive decline in water resource availability is a major impediment to global food production security (Pastor  
39 et al., 2019; Trnka et al., 2019; Konapala et al., 2020). Food crops are the main source of human nutrition (Myers et al., 2017;  
40 Lobell and Gourdji, 2012). Humans depend on food crops for ~47 % of their daily protein intake (FAO, 2021). However, as a  
41 result of human activity, the climate system is changing and global warming is a significant characteristic of this process (IPCC,  
42 2021). Since the 1980s, each successive decade has been warmer than any preceding one after 1850 (Kappelle, 2020). Climate  
43 change affects water consumption and crop yield by altering precipitation, temperature, carbon dioxide (CO<sub>2</sub>) concentration,  
44 and other factors during crop growth (Hatfield and Dold, 2019). Crop adaptation to future climate change with limited water  
45 resources has become a major challenge in sustainable crop production and supply worldwide.

46 The water footprint per unit crop (WF, m<sup>3</sup> t<sup>-1</sup>) (Hoekstra, 2003) is the amount of water consumed by the crop per unit  
47 yield during crop growth within a certain region. It includes blue WF (surface and groundwater), green WF (precipitation that  
48 will not become runoff), and grey WF (freshwater that assimilates pollutants from human activities) (Hoekstra et al., 2011).  
49 Blue and green WF are collectively known as consumptive WF, and grey WF is also called degradative WF (Hoekstra, 2013).  
50 Unlike traditional crop water productivity and other agricultural water metrics, WF covers water consumption, sources, and  
51 spatiotemporal dimensions during the crop growth period. Therefore, water consumption intensity and efficiency for irrigated  
52 and rain-fed **growing modes** may be compared. WF is an effective indicator of the sustainability of regional water use and  
53 optimal water resource allocation (Xu et al., 2019; Mali et al., 2021). The present study focuses exclusively on consumptive  
54 WF, which depends on crop yield and the intensity of water consumption per unit planted area.

55 Several studies have been conducted on the responses of WF to future climate change. Nevertheless, no consensus has  
56 been reached. Certain scholars believe that future climate change will weaken food crop production security. Ahmadi et al.  
57 (2021) reported that maize WF in the Qazvin Plain of India will increase by 42 % and 147 % under representative concentration  
58 pathways (RCP) 4.5 and RCP8.5, respectively, by 2061–2080. Zheng et al. (2020) found that rice yield in Henan and Jiangsu  
59 Provinces (China) will decrease, while WF will increase under four RCPs at various stages of the 21<sup>st</sup> century. Other scholars  
60 believe that crop yield may actually benefit from future increases in precipitation and atmospheric CO<sub>2</sub> concentration. Jans et  
61 al. (2021) considered the combined effects of changes in climatic factors, such as temperature, precipitation, and rising  
62 atmospheric CO<sub>2</sub> concentration, and predicted that between 2011 and 2099, global cotton yield will increase by > 50 % and  
63 WF will decrease by 30 % under RCP8.5. Arunrat et al. (2020) found that in the present century, the yield of individual and  
64 large-scale rice farms in Thailand will increase by 1–30 % and 2–31 %, respectively, while WF will decrease by 10–43 % and  
65 1–67 %, respectively, under RCP4.5. Significant spatiotemporal differences in WF under various irrigation techniques have  
66 been confirmed at the site (Chukalla et al., 2015) and regional (Wang et al., 2019) scales. However, current large-scale studies

67 on the responses of WF to environmental change are usually based on simulations assuming adequate furrow irrigation. These  
68 studies exclude comparisons between various irrigation techniques and the differences in their influences on crop WFs.  
69 Although Dai et al. (2020) optimised maize and wheat cropping patterns under RCP4.5 and RCP8.5 with consideration of  
70 various irrigation modes in the Huaihe River Basin in China by 2050, they only considered blue water.

71 Magnitudes and constitution of crop WF vary widely among regions and areas (Mekonnen and Hoekstra, 2011). To  
72 encourage water users to reduce WF to a reasonable level, Hoekstra (2013, 2014) recommended establishing WF benchmarks  
73 for different products as they facilitate prudent water allocation and fair water resource sharing among sectors and users  
74 (Hoekstra, 2013). On the large-scale, specific WF benchmarks can be set for crops grown on different farms within the same  
75 region (Mekonnen and Hoekstra, 2014). A previous study demonstrated the sensitivity of WF benchmarks to climate zones  
76 (Zhuo et al., 2016a). WF benchmarks significantly differ among irrigation techniques, especially in arid zones (Wang et al.,  
77 2019). However, little is known about the responses of WF benchmarks under different irrigation techniques to future climate  
78 change.

79 To investigate the influence of future climate change on large-scale WF and benchmarks under diverse irrigation  
80 techniques, maize and wheat grown in mainland China were the subjects of this study. We used the outputs of six global  
81 climate models (GCMs) (Table 1), including three models each for relatively wet and dry climate outputs, in Coupled Model  
82 Intercomparison Project Phase 5 (CMIP5). We then used the AquaCrop model to simulate the spatiotemporal responses of  
83 blue and green WF and corresponding WF benchmarks for wheat and maize in the 2030s (2020–2049), 2050s (2040–2069),  
84 and 2080s (2070–2099) under RCP2.6 and RCP8.5 at a 5-arc minute grid resolution. We distinguished between rain-fed and  
85 irrigated **growing modes** and among furrow, micro, and sprinkler irrigation.

86 As of 2019, China was the world's second largest maize and largest wheat producer, accounting for 23 % and 17 % of  
87 total global production, respectively (FAO, 2021). China's cereal production has helped stabilise global food production and  
88 supply. In 2019, the planted areas of maize and wheat in China were 41 million ha and 24 million ha, respectively, and  
89 accounted for 25 % and 14 % of the national total croplands, respectively (NBSC, 2021). Cereal production consumes  
90 substantial volumes of water in China, and these quantities change over time. Zhuo et al. (2019) reported that maize water  
91 consumption increased by 49 % between 2000 and 2013 as planted areas and feed demand increased. Conversely, Wang et al.  
92 (2019) reported that wheat planted and irrigated areas decreased and water consumption slightly declined (4.4 %) from 2000  
93 to 2014. Other studies reported that maize and wheat consume relatively more water in the North than the South of China (Tian  
94 et al., 2019; Wang et al., 2019). Developing water-saving irrigation has become an important way to alleviate the prominent  
95 contradiction between water resources utilization and grain production in China. According to NBSC (2021), the area of water-  
96 saving irrigation projects in China in 2019 was 37 million ha, including 7 million ha for micro irrigation. Therefore, micro  
97 irrigation does apply to food crops in China despite the limited irrigated area. For instance, in Xinjiang province, the area of  
98 micro irrigated maize and wheat was 0.033 million ha in 2009 (CIDDC, 2022), of which the wheat area dominated at up to  
99 0.031 million ha (Wang et al., 2011). Meanwhile, some scholars are conducting research on micro irrigated maize (Bai and  
100 Gao, 2021; Guo et al., 2021) and wheat (Li et al., 2021; Zain et al., 2021) in China, especially in the North. Therefore, the

101 water consumption rates of these staple crops under future climate change scenarios with different irrigation techniques should  
 102 be closely monitored to ensure water supply and food crop production security in China and worldwide. Compared to existing  
 103 literatures **on evaluation of WFs of crop production under climate change scenarios (e.g., Karandish et al., 2022)**, the  
 104 innovations of the current research are embodied in two points. The present study clarifies large-scale spatiotemporal responses  
 105 of WF to future climate change scenarios under different irrigation techniques for the first time. **This analysis is also the first**  
 106 **to explore the large-scale future changes in WF benchmarks under different irrigation techniques.**

107

108 **Table 1.** Inventory of global climate models (GCMs) used in the current study.

GCM	Institute	Reference	Type
CCCMA-CanESM2	Canadian Centre for Climate Modelling and Analysis	Arora et al. (2011); von Salzen et al. (2013)	Wet
CESM1-CAM5	National Science Foundation, Department of Energy, National Center for Atmospheric Research	Hurrell et al. (2013)	
GFDL-CM3	NOAA Geophysical Fluid Dynamics Laboratory	Delworth et al. (2006); Donner et al. (2011)	
FIO-ESM	The First Institute of Oceanography, State Oceanic Administration, China	Qiao et al. (2013)	Dry
GISS-E2R	NASA Goddard Institute for Space Studies USA	Schmidt et al. (2006); Schmidt et al. (2014)	
IPSL-CM5A-MR	Institute Pierre Simon Laplace	Dufresne et al. (2013)	

109


## 110 **2 Method and data**

### 111 **2.1 Research set-up**

112 We studied the spatiotemporal responses of blue and green WF and corresponding WF benchmarks for two crops (maize  
 113 and wheat) to future climate change under two climate change scenarios (RCP2.6 and RCP8.5) using four different **growing**  
 114 **modes** (rain-fed and furrow-, micro-, and sprinkler-irrigated). First, we determined the baseline year. Second, we considered  
 115 different **growing modes** to quantify WF and corresponding WF benchmarks of two crops in the baseline year and future year  
 116 levels under two climate change scenarios. Finally, the spatiotemporal responses of crop WF and corresponding WF  
 117 benchmarks to future climate change were analysed (Fig. 1).

118

119



**Figure 1.** Flow chart for the study.

120  
121  
122

## 123 2.2 Determining the baseline year

124 **Determining the baseline year is needed for a comparison between future and current conditions.** Climate determines the  
125 annual variability of WF (Zhuo et al., 2014), and the baseline year should be determined when there is a relative balance  
126 between aridity and moisture. Hence, the aridity index (AI) was used here. Annual reference evapotranspiration ( $ET_0$ , mm)  
127 and precipitation (PR, mm) in China were calculated (Harris et al., 2014). Then, the AI was calculated, and climate change  
128 trends from 2000 to 2014 were analysed. The year 2013 was designated the baseline as its drought level was nearest the 15-  
129 year national average. The AI was calculated according to the method of Middleton and Thomas (1997):

$$AI = \frac{PR}{ET_0}, \quad (1)$$

## 130 2.3 Water footprint per unit crop calculation

131 WF ( $m^3 t^{-1}$ ) comprises blue WF ( $WF_b$ ,  $m^3 t^{-1}$ ) and green WF ( $WF_g$ ,  $m^3 t^{-1}$ ):

$$WF = WF_b + WF_g, \quad (2)$$

132 where  $WF_b$  and  $WF_g$  were calculated as the quotient of the blue ( $CWU_b$ ,  $m^3 ha^{-1}$ ) and green ( $CWU_g$ ,  $m^3 ha^{-1}$ ) components of  
 133 crop water use ( $CWU$ ,  $m^3 ha^{-1}$ ) and crop yield ( $Y$ ,  $t ha^{-1}$ ), respectively.  $CWU_b$  and  $CWU_g$  were equivalent to the cumulation  
 134 of daily evapotranspiration ( $ET$ ,  $mm d^{-1}$ ) throughout the whole crop growth period (Hoekstra et al., 2011):

$$WF_b = \frac{CWU_b}{Y} = \frac{10 \times \sum_{d=1}^{lgp} ET_b}{Y}, \quad (3)$$

$$WF_g = \frac{CWU_g}{Y} = \frac{10 \times \sum_{d=1}^{lgp} ET_g}{Y}, \quad (4)$$

135 where  $ET_b$  and  $ET_g$  (mm) refer to the blue and green water evapotranspiration, respectively, and  $lgp$  refers to the number of  
 136 days of the crop growth period. The coefficient, 10, is a unit conversion factor, transforming the water depth of  $ET$  (mm) into  
 137 the water amount per unit land area of  $CWU$  ( $m^3 ha^{-1}$ ).

138 The  $ET$  and  $Y$  per grid for each crop were simulated by the AquaCrop model based on the dynamic daily soil water  
 139 balance (Mekonnen and Hoekstra, 2010):

$$S_{[t]} = S_{[t-1]} + PR_{[t]} + IRR_{[t]} + CR_{[t]} - ET_{[t]} - RO_{[t]} - DP_{[t]}, \quad (5)$$

140 where  $S_{[t]}$  and  $S_{[t-1]}$  (mm) refer to the water content in soil when the day,  $t$ , ends and begins, respectively;  $PR_{[t]}$  (mm) is the  
 141 amount of precipitation on day,  $t$ ;  $IRR_{[t]}$  (mm) is the amount of water used for irrigation;  $CR_{[t]}$  (mm) is the capillary rise to the  
 142 crop root zone from the shallow groundwater;  $RO_{[t]}$  (mm) is the water lost by surface runoff due to precipitation; and  $DP_{[t]}$   
 143 (mm) is the water lost by deep percolation caused by excessive precipitation or irrigation. It was assumed that  $CR_{[t]} = 0$  as the  
 144 ground water depth was  $> 1$  m (Allen et al., 1998).  $RO_{[t]}$  was calculated using the Soil Conservation Service curve-number  
 145 (CN) equation (USDA, 1964; Rallison, 1980):

$$RO_{[t]} = \frac{(PR_{[t]} - I_a)^2}{PR_{[t]} + S - I_a}, \quad (6)$$

$$S = 254 \left( \frac{100}{CN} - 1 \right), \quad (7)$$

146 where  $S$  (mm) is the potential maximum water storage, and  $I_a$  (mm) is the initial amount of water loss before the runoff  
 147 formation.

148 By tracking the daily flow of water in and out of the crop root zone, we separated the daily blue and green soil water  
 149 balances (Zhuo et al., 2016b):

$$S_{b[t]} = S_{b[t-1]} + (PR_{[t]} + IRR_{[t]} - RO_{[t]}) \times \frac{IRR_{[t]}}{PR_{[t]} + IRR_{[t]}} - (DP_{[t]} + ET_{[t]}) \times \frac{S_{b[t-1]}}{S_{[t-1]}}, \quad (8)$$

$$S_{g[t]} = S_{g[t-1]} + (PR_{[t]} + IRR_{[t]} - RO_{[t]}) \times \frac{PR_{[t]}}{PR_{[t]} + IRR_{[t]}} - (DP_{[t]} + ET_{[t]}) \times \frac{S_{g[t-1]}}{S_{[t-1]}}, \quad (9)$$

150 where  $S_{b[t]}$  and  $S_{b[t-1]}$  (mm) are the blue water content in soil when the day,  $t$ , ends and begins, respectively; and  $S_{g[t]}$  and  $S_{g[t-1]}$   
 151 (mm) are the green water content in soil when the day,  $t$ , ends and begins, respectively. It is assumed that the initial soil water  
 152 content before the crop growth period is green water.

153 In AquaCrop, the daily transpiration ( $Tr_{[t]}$ , mm) calculates the daily shoot biomass production ( $B$ , kg) using the normalised  
 154 crop biomass water productivity ( $WP^*$ ,  $kg m^{-2}$ ) (Raes et al., 2017):

$$B = WP^* \times \sum \frac{Tr_{[t]}}{ET_{0[t]}} \quad (10)$$

155 where  $WP^*$  is normalised to consider  $CO_2$  concentration, reference evapotranspiration ( $ET_0$ ), and crop classes (C3 or C4) so  
 156 that it is applicable to various locations and seasons. Water productivity remains constant for specific crops.  $Y$ , as the  
 157 harvestable portion of final  $B$ , is calculated by multiplying  $B$  with the adjusted reference Harvest Index ( $HI_0$ , %):

$$Y = f_{HI} \times HI_0 \times B \quad (11)$$

158 where  $f_{HI}$  is a correction factor for  $HI_0$ . It considers the water and temperature stresses during the crop growth period. Being  
 159 consistent with the existing widely used **scaling** method (Mekonnen and Hoekstra, 2011; Zhuo et al., 2016b, 2016c, 2019;  
 160 Wang et al., 2019; Mialyk et al., 2022), the simulated  $Y$  per grid for each crop in 2013 was validated via scaling model  
 161 simulation outputs to correspond with the crop yield statistics data at the provincial level (NBSC, 2021). With the consistent  
 162 **scaling factors for the  $Y$  simulation and crop parameters including the crop calendar,  $WP^*$ ,  $HI_0$ , and the maximum root depth**  
 163 which represent the existing agricultural production level, climate was the only variable for future scenario simulations.

164 In the simulation, different **growing modes**, namely rain-fed and three different irrigation techniques (furrow, micro, and  
 165 sprinkler irrigation), were considered. The irrigation schedule of three irrigation techniques in the model was the Generation  
 166 of Irrigation Schedule, namely the generation of an irrigation schedule by specifying a time and depth criterion for planning  
 167 or evaluating a potential irrigation strategy. **The time criterion we used was Allowable depletion (%), namely the percentage**  
 168 **of the Readily Available soil Water (RAW) that can be depleted before irrigation water has to be applied. The depth criterion**  
 169 **we used was the Back to field capacity as the extra water on top of the amount of irrigation water required to bring the root**  
 170 **zone back to field capacity. The water quality was expressed by the Electrical conductivity ( $dS\ m^{-1}$ ) of the irrigation water.**  
 171 **The soil surface wetted (%), an indicative value for the fraction of soil surface wetted, was used to select irrigation techniques.**  
 172 **Table 2** shows the parameters of three irrigation techniques (Raes et al., 2017). We can adjust the simulated  $ET$  and  $Y$  according  
 173 to the performance of the irrigation schedule.

174

175 **Table 2.** Parameters of three irrigation techniques.

Irrigation technique	From day	Time criterion	Depth criterion	Water quality	Soil surface wetted
		Allowable depletion (%)	Back to field capacity (+/- mm)	Electrical conductivity ( $dS\ m^{-1}$ )	
Furrow	1	50	10	1.5	80
Micro	1	20	10	0	40
Sprinkler	1	50	10	1.5	100

176

## 177 2.4 Benchmarking consumptive WF in crop production

178 Based on the work of Mekonnen and Hoekstra (2014), we ranked grid-level WF for each crop in ascending order of size  
 179 against the corresponding cumulative percentages of the total crop production. The annual WF of 20 % or 25 % of the producers

180 with the highest water productivity in China was set as the annual WF benchmark. The climate zones should be divided when  
 181 WF benchmarks are established (Zhuo et al., 2016a). To this end, the AI partitioned China into arid (< 0.5) and humid (> 0.5)  
 182 zones based on the annual  $ET_0$  and PR from 2000 to 2014 at a 30-arc minute grid resolution (Harris et al., 2014) (Fig. 2).  
 183



184  
 185 **Figure 2.** Regions and climate zones of mainland China.  
 186

## 187 2.5 Data sources

188 Monthly climate data, such as maximum ( $T_x$ ), minimum air temperature ( $T_n$ ), precipitation (PR), and reference  
 189 evapotranspiration ( $ET_0$ ), from 2000 to 2014 at a resolution of 30-arc minute were derived from the CRU-TS 3.24 dataset  
 190 (Harris et al., 2014; CEDA, 2018). The mean annual atmospheric  $CO_2$  concentration (ppm) from 2000 to 2014 was obtained  
 191 from the Mauna Loa Observatory, Hawaii, USA (NOAA, 2018). The downscaled outputs of six GCMs at a 5-arc minute grid  
 192 resolution in the 2030s, 2050s, and 2080s were obtained from the Climate Change, Agriculture and Food Security (CCAFS)  
 193 database (Navarro-Racines et al., 2020; CCAFS, 2015). As the CCAFS database has no  $ET_0$  data, we calculated  $ET_0$  for each  
 194 climate scenario using temperature inputs via the FAO Penman-Monteith method with missing data as described by Allen et  
 195 al. (1998). The projected  $CO_2$  concentrations under RCP2.6 and RCP8.5 were obtained from van Vuuren et al. (2007) and  
 196 Riahi et al. (2007), respectively. To make the model simulation more in line with the actual situation in China, we reset the  
 197 maximum root depth ( $Z_x$ ) according to the FAO-56 recommendation (Allan et al., 1998). **The FAO-56 recommended values**  
 198 **provide clear range of the  $Z_x$  for each type of crops for typical climatic zones.** In addition, we further combined the literature  
 199 research on maize and wheat in China to reset the  $HI_0$  (Zhuo et al., 2016c). The other parameters used in AquaCrop were


200 derived from Raes et al. (2017). Soil texture data and soil water capacity data at a 5-arc minute grid resolution were acquired  
201 from the ISRIC Soil and Terrain database (Dijkshoorn et al., 2008) and ISRIC-WISE dataset (Batjes, 2012), respectively. The  
202 planted areas for each irrigated or rain-fed crop at a 5-arc minute grid resolution were acquired from the MIRCA2000 dataset  
203 (Portmann et al., 2010). We divided these planted areas into different parts subjected to various irrigation techniques using  
204 statistical yearbook data (NBSC, 2021). Provincial-level crop yield statistics data were procured from the National Bureau of  
205 Statistics of China (NBSC, 2021).

## 206 **3 Results**

### 207 **3.1 Future climate change trends in maize and wheat planted areas**

208 In the baseline year 2013, the average annual reference evapotranspiration ( $ET_0$ ) and precipitation (PR) in the planted  
209 areas of two crops were 941 mm and 727 mm, respectively. Compared with the baseline level of 2013, the average annual  $ET_0$   
210 and PR in the planted areas of two crops will both increase under two RCPs, and the increase in  $ET_0$  exceeded that of PR.  $ET_0$   
211 will increase by 17 % and 29 % under RCP2.6 and RCP8.5, respectively, until the 2080s. However, PR will increase by 8 %  
212 and 14 %, respectively. The increases under RCP8.5 (18–29 % and 3–14 % for  $ET_0$  and PR, respectively) were much higher  
213 than those under RCP2.6 (16–17 % and 4–8 % for  $ET_0$  and PR, respectively). Climate change will be relatively more intense  
214 under RCP8.5. The increases in  $ET_0$  were concentrated from April to August (14–39 mm). The increases in PR were  
215 concentrated between June and August (8–20 mm and 12–28 mm, respectively). However, PR will decline in May, July,  
216 November, and December, and it will decline more in May ( $\leq 9$  mm until the 2030s) (Fig. 3a, b). Water and heat resources  
217 were unevenly distributed in the planted areas of the two crops in 2013.  $ET_0$  was relatively higher in East Coast and North  
218 China. PR distribution was comparatively higher in the South and lower in the North (Fig. S4). Compared with 2013,  $ET_0$  and  
219 PR for the most heavily planted areas will increase under both scenarios until the 2080s. The areas with a relatively greater  
220 increase in  $ET_0$  were distributed mainly in Southwest and Northeast (Fig. 3c, e), and PR increased relatively faster in Northwest  
221 and Jing-Jin (Fig. 3d, f).  $ET_0$  decreased mainly in Xinjiang and Inner Mongolia (Fig. 3c, e), and PR decreased mainly in  
222 Xinjiang, Tibet, Northeast, and South Coast (Fig. 3d, f). However, the areas where  $ET_0$  decreased were 86–94 % smaller than  
223 those where PR decreased.

224



**Figure 3.** Future climate projections for the maize and wheat planted zones in China.


225  
226  
227

### 228 3.2 WF distribution in the baseline year 2013

229 The national average WF for wheat ( $1,008 \text{ m}^3 \text{ t}^{-1}$ ) was higher than that for maize ( $813 \text{ m}^3 \text{ t}^{-1}$ ) in the baseline year 2013.  
230 The corresponding blue WF proportions were 37 % and 20 %, respectively. The reason for this discrepancy is that maize is a  
231 C4 crop while wheat is a C3 crop. C4 crops have a relatively higher  $\text{CO}_2$  fixation efficiency and faster photosynthetic rate than

232 C3 crops. Hence, maize can accumulate comparatively more yield than wheat under the same water consumption condition  
 233 (Wang et al., 2012). Figure 4 shows that the high  $WF_g$  value was mainly distributed in areas with relatively greater precipitation  
 234 during crop growth, i.e., abundant green water resources. The main component of WF is  $WF_g$ ; therefore, the high maize WF  
 235 was mainly distributed in Northwest (Fig. 4a), while the high wheat WF was mainly distributed in Southwest and South Coast  
 236 (Fig. 4b). Elevated  $ET_0$  and insufficient precipitation can increase blue water consumption in food production. Thus, the high  
 237  $WF_b$  value was mainly distributed in areas with uneven water and heat resource distributions during crop growth. The high  
 238 maize  $WF_b$  was mainly distributed in Northwest and East Coast (Fig. 4c), while that of wheat was distributed mainly in North  
 239 China (Fig. 4d). In all grids, the proportions of  $WF_b$  and  $WF_g$  were up to 68 % (wheat in Xinjiang) (Table S2) and 98 % (maize  
 240 in Hainan) (Table S1), respectively.

241



242

243

244


**Figure 4.** WF of maize and wheat in China in 2013.

245 A comparison of rain-fed and irrigation techniques demonstrated that the WF of maize and wheat under furrow and  
246 sprinkler irrigation was higher than that under rain-fed in 2013. The WF of micro-irrigated crops was lower than that of rain-  
247 fed crops. The WF of maize ( $850 \text{ m}^3 \text{ t}^{-1}$ ) and wheat ( $1,170 \text{ m}^3 \text{ t}^{-1}$ ) was highest under furrow and sprinkler irrigation, respectively.  
248 For wheat under all three irrigation techniques,  $\text{WF}_b$  was dominant (54–65 %). However,  $\text{WF}_b$  for maize was only dominant  
249 under micro irrigation (61 %). Micro-irrigated ( $9.55 \text{ t ha}^{-1}$  for maize and  $5.46 \text{ t ha}^{-1}$  for wheat) and rain-fed ( $5.76 \text{ t ha}^{-1}$  for  
250 maize and  $4.51 \text{ t ha}^{-1}$  for wheat) crops had the highest and lowest yield, respectively, in 2013. The responses of maize yield to  
251 rain-fed and various irrigation techniques were stronger than those of wheat yield (Fig. 4e, f).

### 252 3.3 Spatiotemporal responses of WF to future climate change

253 Compared with the baseline year 2013 and at the national average level, maize WF will increase under both RCP2.6 and  
254 RCP8.5, by 17 % and 13 %, respectively, until the 2080s. The WF of wheat will increase under RCP2.6 (by 12 % until the  
255 2080s) but decrease by 12 % under RCP8.5 until the 2080s (Fig. 5a). The increases in  $\text{CO}_2$  concentration and, by extension,  
256 yield gain, will be lower under RCP2.6 than RCP8.5. During the same period, the increases in WF under RCP2.6 will be 1–  
257 3 % higher for maize and 2–10 % higher for wheat than those under RCP8.5. There will be relatively smaller differences in  
258  $\text{CO}_2$  concentration between climate scenarios of the 2030s (431 ppm under RCP2.6 and 449 ppm under RCP8.5). Thus, the  
259 differences in WF between RCPs will be smaller before the 2030s and larger after the 2050s. The WF of irrigated wheat under  
260 RCP8.5 will decline by 3 % until the 2050s and by 15 % until the 2080s. The increase in WF will be highest under rain-fed,  
261 and the WF of rain-fed maize and wheat under RCP2.6 will increase by 19 % and 24 %, respectively, until the 2080s. By  
262 contrast, the WF of irrigated maize and wheat under RCP2.6 will only increase by 13 % and 7 %, respectively, until the 2080s  
263 (Fig. 5a). A comparison of the various irrigation techniques demonstrated that the WFs of wheat and maize respond differently  
264 under the same scenario. The increase in WF amplitude for maize will be highest under furrow irrigation (14 % and 11 %  
265 under RCP2.6 and RCP8.5 until the 2080s, respectively) and lowest under micro irrigation (5 % and 2 % under RCP2.6 and  
266 RCP8.5 until the 2080s, respectively). The WF of sprinkler-irrigated wheat under RCP8.5 will decline by 1 % until the 2030s.  
267 The WF of wheat under micro irrigation had the highest increase (9 % until the 2080s under RCP2.6) and the lowest decrease  
268 (14 % until the 2080s under RCP8.5). The WF of wheat under sprinkler irrigation had the lowest increase (only 2 % until the  
269 2080s under RCP2.6) and the highest decrease (19 % until the 2080s under RCP8.5) (Fig. 5b).

270



**Figure 5.** WF of maize and wheat in 2013 and future year levels under various climate change scenarios in China.

271

272

273

274

275

276

277

278

279

280

281

282

283


284

285

286

The spatial distribution of the relative changes in maize and wheat WF from 2013 to the 2080s showed regional differences. The WF will increase for 90–93 % of all areas planted with maize (Fig. 6a, b), and it will increase for 78 % of all areas planted with wheat under RCP2.6 (Fig. 6c) and decrease for 81 % of all areas planted with wheat under RCP8.5 (Fig. 6d). Increases in  $ET_0$  lead to increases in WF, while decreases in PR lead to increases in  $WF_b$  (Fig. S6). Hence, the regions with relatively greater increases in WF were mainly distributed where  $ET_0$  strongly increased and PR slightly increased or even decreased. In Yunnan, maize WF increased by 44 % and 38 % under RCP2.6 and RCP8.5, respectively. In Guangxi, wheat WF increased by 50 % and 16 % under RCP2.6 and RCP8.5, respectively (Table S5). Comparison of rain-fed and various irrigation techniques revealed that the WF of each crop responded uniquely to latitudinal and longitudinal climate change under the same scenario. The responses of maize WF to climate change with latitude were relatively consistent. It increased by 27–43 % at 19–26 °N and ~51 °N latitude and decreased at ~44 °N latitude. By contrast, the responses of WF for rain-fed maize were more sensitive at ~40 °N and ~52 °N latitude. The responses of maize WF vary widely within 74–100 °E longitude. The WF of maize under rain-fed and furrow and sprinkler irrigation declined at 74–90 °E longitude. The increase in WF for maize under rain-fed at 93–98 °E longitude was 3–51 % higher than the increase in WF for maize under furrow and

287 sprinkler irrigation. The WF of micro-irrigated maize decreased at 74–95 °E longitude (Fig. 6a, b). The responses of wheat  
 288 WF to climate change with latitude and longitude were relatively consistent. However, in certain areas, there were large  
 289 differences in wheat WF between rain-fed and the three irrigation techniques. The WF of wheat under rain-fed decreased at  
 290 74–80 °E longitude and by more than the WF of wheat under the three irrigation techniques at the same longitude range. The  
 291 increases in the WF of wheat under rain-fed at ~93 °E and ~122 °E longitude and ~22 °N latitude were significantly higher  
 292 than the increases in WF of wheat under the three irrigation techniques (Fig. 6c, d).  
 293




294  
 295 **Figure 6.** Spatial distributions in relative changes  $\Delta$  (%) in WF (bottom left panel) with longitudinal (top panel) and latitudinal (right  
 296 panel) changes under different irrigation techniques applied to both crops under two scenarios from 2013 to the 2080s.

297

298        WF is determined by both crop yield (Y) and crop water use (CWU). We compared the relationships between the relative  
299 changes in WF ( $\Delta WF$ ) and corresponding Y ( $\Delta Y$ ) and CWU ( $\Delta CWU$ ) (Fig. 7). The  $\Delta WF$  of maize and wheat under future  
300 climate change scenarios was inversely proportional to  $\Delta Y$  and directly proportional to  $\Delta CWU$ . Nevertheless,  $\Delta WF$  was  
301 relatively more sensitive to  $\Delta Y$ . When  $\Delta Y$  was 25 %,  $\Delta WF$  of wheat under RCP2.6 and maize was approximately -25 %, while  
302  $\Delta WF$  of wheat under RCP8.5 was approximately -10 %. When  $\Delta CWU$  was 25 %,  $\Delta WF$  of wheat under RCP2.6 and maize  
303 was ~20 %, while  $\Delta WF$  of wheat under RCP8.5 was approximately -8 % (Fig. 7a, b). The responses of  $\Delta WF$  of maize were  
304 more sensitive to  $\Delta Y$  and  $\Delta CWU$  than those of wheat. The responses of  $\Delta WF$  of maize and wheat under RCP2.6 were more  
305 sensitive to  $\Delta Y$  and  $\Delta CWU$  than those under RCP8.5. Comparison of rain-fed and various irrigation techniques revealed that  
306 the correlation between  $\Delta WF$  and  $\Delta Y$  was stronger for rain-fed crops. For rain-fed maize,  $R^2$  can reach 0.55 (Fig. 7a).  $\Delta WF$   
307 and  $\Delta CWU$  were strongly correlated for irrigated crops, and  $\Delta WF$  and  $\Delta CWU$  were especially strongly correlated for crops  
308 under micro irrigation ( $R^2$  can reach 0.98 for wheat) (Fig. 7b). We also determined the relationship between  $\Delta WF_b$  and  $\Delta CWU_b$ ,  
309 was similar but more significant than that between  $\Delta WF$  and  $\Delta CWU$  (Fig. 7c).

310



311

312 **Figure 7.** Relationships between relative changes  $\Delta$  (%) in (a) Y and corresponding WF, (b) CWU and corresponding WF, and (c) CWU<sub>b</sub>

313

and corresponding WF<sub>b</sub> of two crops under RCP2.6 and RCP8.5 from 2013 to the 2080s.

314

### 315 3.4 Spatiotemporal WF benchmarks responses to climate change

316 Table 3 shows the WF benchmarks of maize and wheat among various irrigation techniques and climate zones in 2013  
 317 and future year levels. The WF benchmarks of maize and wheat in the humid zone were 13–32 % higher than those in the arid  
 318 zone, which is similar to results obtained by Wang et al. (2019). In the same climate zone, WF benchmarks of wheat were  
 319 generally 2–35 % higher than those of maize. However, in the humid zone, the WF benchmark for the 25th production  
 320 percentile of maize was 3 % higher than that of wheat under RCP8.5 in the 2080s. In the arid zone, WF benchmarks of rain-  
 321 fed maize were 13–34 % higher than those of irrigated maize. In the humid zone of the future, WF benchmarks of rain-fed  
 322 wheat were 2–7 % higher than those of irrigated wheat. In general, WF benchmarks of sprinkler-irrigated crops were higher,  
 323 while those of micro-irrigated crops were lower. The differences in WF benchmarks among various irrigation techniques were  
 324 more significant in the arid zone. WF benchmarks of the crops under micro irrigation were 30–38 % lower than those under  
 325 sprinkler irrigation in the arid zone. The difference in the humid zone was only 8–14 %, which is also consistent with the study  
 326 by Wang et al. (2019). In the humid zone, however, WF benchmarks of maize under furrow irrigation were 7–21 % higher  
 327 than those under sprinkler irrigation.

328

329 **Table 3.** WF benchmarks ( $\text{m}^3 \text{t}^{-1}$ ) of maize and wheat for different climate zones in 2013 and future year levels under two climate change  
 330 scenarios in China.

Climate zones	Crop	Type	WF ( $\text{m}^3 \text{t}^{-1}$ ) at different production percentile*					
			20th			25th		
			2013	RCP2.6	RCP8.5	2013	RCP2.6	RCP8.5
Arid	Maize	Total	601	(577, 576, 580)	(589, 584, 566)	623	(661, 658, 655)	(655, 652, 634)
		Irrigated	522	(505, 504, 506)	(503, 503, 496)	548	(508, 507, 511)	(507, 509, 501)
		Furrow	618	(658, 658, 658)	(654, 654, 642)	654	(693, 693, 691)	(689, 687, 674)
		Micro	466	(455, 454, 456)	(456, 454, 440)	477	(459, 458, 460)	(458, 460, 446)
		Sprinkler	700	(727, 725, 723)	(722, 719, 708)	706	(729, 729, 726)	(724, 721, 710)
		Rain-fed	599	(661, 661, 662)	(652, 649, 630)	618	(682, 679, 671)	(672, 667, 652)
	Wheat	Total	753	(776, 764, 781)	(765, 707, 620)	768	(829, 816, 828)	(809, 756, 666)
		Irrigated	754	(776, 764, 781)	(765, 707, 620)	768	(830, 816, 829)	(810, 757, 666)
		Furrow	830	(850, 840, 850)	(830, 774, 680)	940	(885, 875, 887)	(868, 809, 712)
		Micro	648	(701, 690, 705)	(694, 643, 562)	670	(717, 705, 721)	(707, 654, 572)
		Sprinkler	1020	(1003, 998, 1007)	(989, 920, 811)	1032	(1034, 1028, 1038)	(1019, 948, 837)
		Rain-fed	692	(743, 734, 753)	(729, 692, 618)	692	(790, 772, 791)	(769, 737, 653)
Humid	Maize	Total	680	(761, 754, 752)	(756, 752, 739)	718	(813, 807, 807)	(809, 806, 785)
		Irrigated	743	(905, 905, 908)	(902, 900, 881)	782	(939, 939, 944)	(937, 936, 916)
		Furrow	762	(925, 926, 930)	(921, 921, 901)	801	(943, 942, 948)	(940, 939, 919)
		Micro	649	(709, 704, 707)	(694, 696, 683)	660	(734, 726, 732)	(721, 726, 708)
		Sprinkler	713	(770, 771, 768)	(764, 762, 750)	737	(813, 814, 812)	(808, 806, 793)
		Rain-fed	631	(712, 703, 707)	(710, 702, 678)	656	(744, 737, 737)	(740, 736, 716)
	Wheat	Total	873	(933, 932, 946)	(921, 851, 752)	887	(944, 942, 957)	(931, 860, 760)

Irrigated	887	(914, 914, 924)	(900, 841, 744)	897	(925, 926, 937)	(912, 849, 752)
Furrow	887	(914, 914, 925)	(901, 841, 744)	896	(925, 927, 937)	(913, 849, 752)
Micro	820	(821, 826, 838)	(804, 753, 665)	833	(830, 839, 849)	(812, 759, 671)
Sprinkler	933	(949, 944, 955)	(936, 872, 770)	946	(958, 953, 964)	(944, 880, 777)
Rain-fed	812	(973, 958, 984)	(950, 863, 757)	831	(989, 973, 998)	(964, 877, 763)


\*The three numbers in brackets are the values of 2030s, 2050s and 2080s.

331

332 Compared with the baseline year, 2013, the changes in maize and wheat WF benchmarks under future climate change  
333 scenarios are similar to the changes in WF. However, the WF benchmark for the 20th production percentile of maize will  
334 decline by 2–6 % in the arid zone. WF benchmarks of wheat under RCP8.5 will decrease by 2–6 % and 13–18 % until the  
335 2050s and the 2080s, respectively. The increasing range of the WF benchmark for the 25th production percentile of maize was  
336 7–8 % higher in the humid zone than that in the arid zone. The increasing range of the WF benchmark for the 20th production  
337 percentile of wheat was 4–5 % higher in the humid zone than that in the arid zone. WF benchmarks of maize and wheat  
338 increased to a greater extent under RCP2.6 but decreased to a greater extent under RCP8.5. WF benchmarks of rain-fed crops  
339 increased more than those of irrigated crops in the same climate zone. Nevertheless, the increase in WF benchmarks was 7–  
340 11 % lower for rain-fed than irrigated maize in the humid zone. WF benchmarks of maize and wheat generally increased  
341 relatively more under furrow irrigation and comparatively less under sprinkler irrigation. However, under RCP2.6, the growth  
342 rate of the WF benchmark for the 20th production percentile of wheat was 5–6 % higher under micro irrigation than that under  
343 furrow irrigation in the arid zone. The increase in the WF benchmark for the 20th production percentile of wheat was 0.19–  
344 2 % higher under sprinkler irrigation than that under micro irrigation in the humid zone (Table 3).

345 Figure 8 shows the spatial distribution of the relative changes in the WF of maize and wheat compared with the benchmark  
346 for the 25th production percentile in 2013 and the 2080s. In 2013, the WF for 81 % and 79 % of the maize and wheat planted  
347 areas, respectively, was higher than its benchmark. The maize planted areas with WF below the benchmark were distributed  
348 mainly in Xinjiang in the arid zone and northeast Inner Mongolia in the humid zone (Fig. 8a). The wheat planted areas with  
349 WF below the benchmark were distributed mainly in Xinjiang in the arid zone and Qinghai (Fig. 8d). Under future climate  
350 change scenarios, the maize and wheat planted areas with the WF below the benchmark will slightly decrease in the 2080s.  
351 These areas are mainly distributed in Heilongjiang, Tibet, southern Gansu, and Sichuan in the humid zone for maize; and  
352 Henan and Tibet in the humid zone and Qinghai for wheat. This is because that the annual  $ET_0$  will increase relatively faster  
353 in Heilongjiang and Tibet, which will lead to a greater increase in  $WF_b$ . The annual PR in other regions will significantly  
354 increase, which will result in a greater increase in  $WF_g$ . Maize and wheat planted areas under RCP8.5 with WF below the  
355 benchmark will decrease by 5 % and 4 %, respectively, until the 2080s.

356



357

358 **Figure 8.** Relative changes  $\Delta$  (%) in the WF of maize and wheat compared with the benchmark for the 25th production percentile in 2013  
 359 and the 2080s under RCP2.6 and RCP8.5 in different climate zones of China.

360

### 361 3.5 Discussion

362 This study analysed and compared the WF and WF benchmarks responses of wheat and maize under rain-fed and various  
 363 irrigation conditions and forecasted their responses to future climate change scenarios in China. Under the background that the  
 364 annual  $ET_0$  and PR will both increase but  $ET_0$  will increase faster, maize WF will increase under both RCP2.6 and RCP8.5.  
 365 Wheat WF will increase under RCP2.6 but decrease under RCP8.5 until the 2080s. Rain-fed crops had higher ranges of  
 366 increasing WF, which is consistent with Rosa et al. (2020). The increasing ranges of maize and wheat WF were lowest under  
 367 micro irrigation and sprinkler irrigation, respectively. Therefore, the implementation of water-saving irrigation techniques  
 368 (micro and sprinkler irrigation) may help mitigate the adverse effects of future climate change on agriculture, which is in line  
 369 with Dai et al. (2020). Under future climate change, WF benchmarks will be modified in a manner resembling that for WF.  
 370 However, the former changes will not be as significant as the latter in the same area.

371 In 2013, the WF of maize was lower than that of wheat. Nevertheless, maize WF is expected to increase more rapidly  
372 than wheat WF under future climate change scenarios. C4 crops such as maize have higher photosynthetic rates than C3 crops  
373 such as wheat. However, C4 crops are less sensitive to elevated atmospheric CO<sub>2</sub> than C3 crops (Bowes, 1993). Hence, while  
374 maize yield is higher than wheat yield, the former increases less than the latter. We compared current results against those of  
375 previous studies in Table 4. The differences we determined for the relative changes in maize and wheat WF between years and  
376 RCPs resembled those reported by Zhuo et al. (2016d). However, these authors also considered other factors, such as harvested  
377 crop area, technology, diet, and population, that could partially offset the adverse effects of future climate change. Therefore,  
378 maize and wheat WF will decline in the future according to Zhuo et al. (2016d). Fader et al. (2010) studied relative global-  
379 scale changes in maize WF for 2050. Their analysis was conducted in the opposite direction of that of the present study on  
380 China. Moreover, the two studies differed in terms of climate scenario, research area, and crop model. Winter wheat WF in  
381 Germany and Italy will decline by 2050 according to Garofalo et al. (2019). Nevertheless, our research showed that winter  
382 wheat WF will increase in China by 2050. The crop water use in Germany and Italy changes more smaller than that in China.  
383 However, our observed differences in the relative changes in WF between RCPs were consistent with those of Garofalo et al.  
384 (2019); namely, under RCP8.5, WF will either decrease more or increase less.

385

386 **Table 4.** Comparison of the results between current and previous studies.

Reference	Year	Study case	Scenario	Relative changes in WF (%)
Zhuo et al. (2016d)	2030	China Maize	RCP2.6 / RCP8.5	-38–32 / -10–0
		China Wheat		-25–17 / -20–11
	2050	China Maize		-51–43 / -22–8
	China Wheat	-36–27 / -38–27		
Current study	2030s (2020–2049)	China Maize	RCP2.6 / RCP8.5	17 / 16
		China Wheat		11 / 9
	2050s (2040–2069)	China Maize		16 / 15
	China Wheat	10 / 0.20		
Fader et al. (2010)	2041–2070	Global Maize	SRES A2	-0.44–0.35
Current study	2050s (2040–2069)	China Maize	RCP2.6 / RCP8.5	16 / 15
Garofalo et al. (2019)	2050	Germany Winter wheat	RCP4.5 / RCP8.5	-24 / -26
		Italy Winter wheat		-5 / -6
Current study	2050s (2040–2069)	China Winter wheat	RCP2.6 / RCP8.5	10 / 0.60

387

388 In the future, the spatial distributions of maize and wheat WF will change considerably. By contrast, the spatial  
389 distributions of WF benchmarks will negligibly change. This phenomenon is comparatively more pronounced in the area with  
390 limited agricultural development. In 2013, Guizhou and Guangxi had the highest maize and wheat WF (1,317 m<sup>3</sup> t<sup>-1</sup> and 3,720  
391 m<sup>3</sup> t<sup>-1</sup>, respectively) (Table S1, S2). In the humid zone, maize WF in Guizhou and wheat WF in Guangxi will increase by 37 %  
392 and 50 %, respectively, under RCP2.6 and by 33 % and 16 %, respectively, under RCP8.5 until the 2080s (Table S5).  
393 Nevertheless, the WF benchmarks for the 25th production percentile of maize and wheat in the humid zone will only increase  
394 by 12 % and 8 %, respectively, under RCP2.6 and increase by 9 % and decrease by 14 %, respectively, under RCP8.5. These

395 areas will nonetheless have great potential for agricultural water conservation in the future. If maize and wheat WF in various  
396 regions of China can be reduced to the benchmark for the 25th production percentile, the total CWU can be reduced by 45–66  
397 billion m<sup>3</sup> (~14–17 %). Rain-fed agriculture can save 27–40 billion m<sup>3</sup> (~18–22 %), water which is more than that conserved  
398 by irrigation. In irrigated agriculture, furrow irrigation has a comparatively high water-saving potential (17–22 billion m<sup>3</sup>;  
399 ~11–12 %). To optimise the agricultural water-saving potential in China, we must either reduce WF or prevent it from  
400 increasing, either by enhancing crop yield or decreasing CWU. However, this goal can only be realised with the support of  
401 relevant policies and management practices. The annual PR is relatively low, and the ET<sub>0</sub> is relatively high in North China.  
402 Shortage of water for agriculture is a major bottleneck in the development of local agriculture there. However, furrow irrigation  
403 is mainly applied in these areas (Fig. S3). Hence, irrigation water use efficiency is low and WF<sub>b</sub> is high. High-efficiency,  
404 water-saving micro irrigation, and sprinkler irrigation could replace furrow irrigation in these areas so that CWU and WF  
405 decrease. The planted areas in the South have abundant precipitation but limited distribution (Fig. S2) and high WF (Fig. 4a,  
406 b). WF can be mitigated by implementing ground cover techniques (ex. straw return, mulch) to reduce soil evaporation and by  
407 improving farmer skills. WF can also be reduced by optimizing the structure of crop planting. Crops and varieties best adapted  
408 to local climate conditions and climate change can lower irrigation requirements and reduce WF.

409 To make climate models comparable and promote their development, The World Climate Research Program (WCRP)  
410 has developed and promoted the CMIP since 1995 (Meehl et al., 1997, 2000). Its current iteration is CMIP Phase 6 (CMIP6),  
411 which will be used in the forthcoming Intergovernmental Panel on Climate Change's Sixth Assessment Report (IPCC AR6).  
412 GCMs and their associated research results based on CMIP5 provided vital support for IPCC's Fifth Assessment Report (IPCC  
413 AR5). CMIP5 proposed four RCP scenarios (RCP2.6, RCP4.5, RCP6.0, and RCP8.5) by considering greenhouse gas (GHG)  
414 emissions and concentrations, atmospheric pollutant concentrations, and land use in the 21<sup>st</sup> century (Moss et al., 2008).  
415 However, no specific socio-economic assumptions were made. The Scenario Model Intercomparison Project (ScenarioMIP),  
416 as the primary activity within CMIP6, will provide a series of new climate scenarios that consider social factors related to  
417 climate change adaptation and impacts. They will be based on the combined application of shared socioeconomic pathways  
418 (SSPs) and RCPs and will compensate for the limitations of the RCPs in CMIP5 (O'Neill et al., 2016). The climate models in  
419 CMIP5 and CMIP6 can both effectively simulate changes in potential evapotranspiration (Liu et al., 2020) and precipitation  
420 (Müller et al., 2021) in most parts of the world. Müller et al. (2021) reported that CMIP5 and CMIP6 simulate increasing trends  
421 in temperature in a similar fashion. Nevertheless, the simulation generated by CMIP6 is higher than that by CMIP5.  
422 Notwithstanding, CMIP5 and CMIP6 are reasonably consistent and similar in terms of their abilities to predict future climate  
423 changes. This study focused on the responses of crop production to future climate change. It mainly considered the influences  
424 of GHG emission- and concentration-driven climate change and excluded the influences of alterations in socioeconomic  
425 development. Therefore, we implemented CMIP5 in our current research.

426 Three are two methods of establishing WF benchmarks (Hoekstra, 2013). Method 1 is based on yield accumulation  
427 statistical analysis. Due to the variability of WFs found across regions and among producers within a region, for each crop, we  
428 can select the WF of 20 % or 25 % of the producers with the highest water productivity as the WF benchmark (Mekonnen and

429 Hoekstra, 2014). Method 2 is based on the available optimal technique analysis. We can compare the WFs at each location  
430 under different agricultural management practices and take the WF associated with optimal practice, which results in the  
431 smallest WF, as the WF benchmark (Chukalla et al., 2015). Both methods establish WF benchmarks based on the maximum  
432 reasonable water consumption in each step of the product's supply chain (Hoekstra, 2014). Method 1 is suitable for large-scale  
433 application. The differences in environmental conditions (such as climate) and development conditions should be considered  
434 comprehensively (Mekonnen and Hoekstra, 2014; Zhuo et al., 2016a). The drawback of Method 1 is that no matter what spatial  
435 scope one takes in grouping producers, within that scope there will still be variability from place to place even if the differences  
436 in regional environmental and development conditions are taken into account (Schyns et al., 2022). Method 2 is suitable for  
437 smaller scale and overcomes this drawback of Method 1 to some extent. The Method 2's drawback is that it has the higher  
438 requirements on the setting and simulation of different agricultural management practices. We mainly want to explore the  
439 response of large-scale WF to future climate change under specific irrigation technique, that is, each irrigation technique has  
440 its corresponding WF benchmarks. And only one agricultural management practice, that is irrigation, is considered here.  
441 Therefore, we choose Method 1. A combination of methods should be established. If conditions permit, we strongly  
442 recommend that Method 1 and Method 2 are combined to establish small-scale WF benchmarks. Different agricultural  
443 management practices, such as irrigation, mulching techniques and so on, can be combined to further determine WF  
444 benchmarks.

445 The sources of uncertainty in research on the responses of crop production to climate change include GCMs, climate  
446 scenarios, crop models, and their interactions (Wang et al., 2020). Semenov and Stratonovitch (2010) proposed that the use of  
447 multiple GCMs can reduce the uncertainty associated with them. We selected three GCMs each for wet and dry climate outputs  
448 to encompass a broad climate prediction scenario. To objectively and comprehensively project the future climate change trends  
449 of China, we selected two extreme RCPs, namely, RCP2.6 and RCP8.5. Wang et al. (2020) suggested that crop models are the  
450 main source of uncertainty in predicting wheat yield in China under future climate change. The application of various crop  
451 models and parameter settings inevitably lead to different yield forecasts (Asseng et al., 2013). Hence, the use of AquaCrop  
452 alone may introduce uncertainty into WF forecasting.

453 The present study had certain limitations in terms of the assumptions it made for the simulation. First, we assumed that  
454 the crop parameters (such as planting calendar,  $HI_0$ , and  $Z_x$ ) for each crop under the identical **growing mode** (irrigated or rain-  
455 fed) were constant on a spatiotemporal scale. Yoon and Choi (2020) proposed that future increases in temperature and  
456 precipitation might shorten the crop growth period. Xiao et al. (2020) indicated that the winter wheat and summer maize  
457 growing periods will be lengthened and shortened, respectively, under future climate change. However, we did not consider  
458 future changes in the crop growth period. Second, we assumed a constant soil surface moisture rate for each grid under the  
459 various irrigation techniques. Third, it was assumed that the observed changes in the planted areas in 2013 were based on the  
460 2000 raster database, and we ignored the migration of planted areas. Finally, we assumed that the maize and wheat planted  
461 areas will not change in the future and would remain consistent with baseline year 2013. Thus, we did not consider future  
462 development of cultivated lands.

463 The core content of this study was to quantify the responses of maize and wheat WF and WF benchmarks to future climate  
464 change under various irrigation techniques. Future research must improve the accuracy of the crop model simulation and  
465 reduce the uncertainty of climate prediction associated with using different GCMs. Moreover, this study only considered future  
466 climate change scenarios. Future investigations should also consider the influence of changes in technological development,  
467 land use, **growing modes**, and so on.

#### 468 **4 Conclusions**

469 This study explored the responses of maize and wheat WF accounting and benchmarking to future climate change in  
470 China. The crops were subjected to various irrigation techniques. The year 2013 was the baseline, and WF and its benchmarks  
471 were quantified for each crop under rain-fed and irrigation (furrow, micro, and sprinkler) management techniques in the 2030s,  
472 2050s, and 2080s under RCP2.6 and RCP8.5 at a 5-arc grid scale. The AquaCrop model with the outputs of six GCMs in  
473 CMIP5 as its input data was used to simulate the WF of maize and wheat. The results show that: (1) Compared with 2013, the  
474 annual  $ET_0$  and PR in the maize and wheat planted areas of China will both increase; however, the former will increase faster  
475 than the latter. (2) Maize WF will increase under both RCP2.6 and RCP8.5 by 17 % and 13 %, respectively, until the 2080s.  
476 Wheat WF will increase under RCP2.6 (by 12 % until the 2080s) but decrease by 12 % under RCP8.5 until the 2080s. Rain-  
477 fed crops were more vulnerable to the adverse impacts of future climate change, and their WF increased to a greater extent  
478 than that of irrigated crops. Micro irrigation and sprinkler irrigation resulted in the lowest increases in WF for maize and wheat,  
479 respectively. Hence, these water-saving irrigation practices effectively mitigated the negative impact of climate change. (3)  
480 Within different climate zones and under various irrigation techniques, there will be significant differences in the responses of  
481 WF benchmarks to future climate change. The changes in WF and its benchmarks will be similar in response to future climate  
482 change. The rate of increase in WF benchmarks for sprinkler-irrigated crops will generally be lower than those for rain-fed,  
483 micro-irrigated, and furrow-irrigated crops within the same climate zone. However, the change in the spatial distribution of  
484 WF benchmarks will not be as significant as that of WF itself. Moreover, this difference will be more pronounced in the region  
485 with low agricultural development. Additionally, this study also demonstrated that the agricultural water in China still has  
486 substantial water-saving potential and can be effectively conserved.

487

488 **Data availability.** Data sources are listed in Sect. 2.5. Data generated in this paper are available by contacting La Zhuo.

489 **Competing interests.** The authors declare that they have no conflict of interest.

490 **Author contribution**

491 La Zhuo and Pute Wu designed the study. Zhiwei Yue and Xiangxiang Ji carried it out, and prepared the manuscript with  
492 contributions from all co-authors.

493 **Acknowledgements**

494 The study is financially supported by the Program for Cultivating Outstanding Talents on Agriculture, Ministry of Agriculture  
495 and Rural Affairs, People's Republic of China [13210321], and the National Natural Science Foundation of China Grants  
496 [51809215].

497

498 **References**

- 499 Ahmadi, M., Etedali, H. R., and Elbeltagi, A.: Evaluation of the effect of climate change on maize water footprint under RCPs  
500 scenarios in Qazvin plain, Iran, *Agr. Water Manage.*, 254, 106969, <https://doi.org/10.1016/j.agwat.2021.106969>, 2021.
- 501 Allen, R. G., Pereira, L. S., Raes, D., and Smith, M.: Crop evapotranspiration-Guidelines for computing crop water  
502 requirements-FAO Irrigation and drainage paper 56, 300, FAO, Rome, Italy, 1998.
- 503 Arora, V. K., Scinocca, J. F., Boer, G. J., Christian, J. R., Denman, K. L., Flato, G. M., Kharin, V. V., Lee, W. G., and  
504 Merryfield, W. J.: Carbon emission limits required to satisfy future representative concentration pathways of greenhouse  
505 gases, *Geophys. Res. Lett.*, 38, 387–404, <https://doi.org/10.1029/2010GL046270>, 2011.
- 506 Arunrat, N., Pumijumnong, N., Sreenonchai, S., Chareonwong, U., and Wang, C.: Assessment of climate change impact on  
507 rice yield and water footprint of large-scale and individual farming in Thailand, *Sci. Total Environ.*, 726, 137864,  
508 <https://doi.org/10.1016/j.scitotenv.2020.137864>, 2020.
- 509 Asseng, S., Ewert, F., Rosenzweig, C., Jones, J. W., Hatfield, J. L., Ruane, A. C., Boote, K. J., Thorburn, P. J., Rötter, R. P.,  
510 Cammarano, D., Brisson, N., Basso, B., Martre, P., Aggarwal, P. K., Angulo, C., Bertuzzi, P., Biernath, C., Challinor, A.  
511 J., Doltra, J., Gayler, S., Goldberg, R., Grant, R., Heng, L., Hooker, J., Hunt, L. A., Ingwersen, J., Izaurralde, R. C.,  
512 Kersebaum, K. C., Müller, C., Naresh Kumar, C., Nendel, C., O'Leary, G., Olesen, J. E., Osborne, T. M., Palosuo, T.,  
513 Priesack, E., Ripoche, D., Semenov, M. A., Shcherbak, I., Steduto, P., Stöckle, C., Stratonovitch, P., Streck, T., Supit, I.,  
514 Tao, F., Travasso, M., Waha, K., Wallach, D., White, J. W., Williams, J. R., and Wolf, J.: Uncertainty in simulating wheat  
515 yields under climate change, *Nat. Clim. Chang.*, 3, 827–832, <https://doi.org/10.1038/nclimate1916>, 2013.
- 516 Bai, T. and Gao, J.: Optimization of the nitrogen fertilizer schedule of maize under drip irrigation in Jilin, China, based on  
517 DSSAT and GA, *Agric Water Manag.*, 244, 106555, <https://doi.org/10.1016/j.agwat.2020.106555>, 2021.
- 518 Batjes, N.: ISRIC-WISE Derived Soil Properties on a 5 by 5 Arc-Minutes Global Grid (Ver. 1.2), ISRIC, Wageningen, The  
519 Netherlands, available at: <https://www.isric.org>, 2012.

520 Bowes, G.: Facing the Inevitable: Plants and Increasing Atmospheric CO<sub>2</sub>, *Annu. Rev. Plant Phys. Plant Mol. Biol.*, 44, 309–  
521 332, <https://doi.org/10.1146/annurev.pp.44.060193.001521>, 1993.

522 CCAFS: CCAFS-Climate Statistically Downscaled Delta Method data, Climate Change, Agriculture and Food Security,  
523 available at: [www.ccafs-climate.org](http://www.ccafs-climate.org), 2015.

524 CIDDC: China Irrigation and Drainage Development Center, China, available at:  
525 <http://www.jsdg.com.cn/temp/Index/Display.asp?NewsID=12313>, last access: 14 April 2022.

526 CEDA: Climatic Research Unit (CRU) time-series datasets of variations in climate with variations in other phenomena, NCAS  
527 British Atmospheric Data Centre, date of citation, available at: <http://catalogue.ceda.ac.uk/uuid>, 2018.

528 Chukalla, A. D., Krol, M. S., and Hoekstra, A. Y.: Green and blue water footprint reduction in irrigated agriculture: Effect of  
529 irrigation techniques irrigation strategies and mulching, *Hydrol. Earth Syst. Sci.*, 19, 4877–4891,  
530 <https://doi.org/10.5194/hess-19-4877-2015>, 2015.

531 Dai, C., Qin, X. S., Lu, W. T., and Huang, Y.: Assessing adaptation measures on agricultural water productivity under climate  
532 change: A case study of Huai River Basin, China, *Sci. Total Environ.*, 721, 137777,  
533 <https://doi.org/10.1016/j.scitotenv.2020.137777>, 2020.

534 Delworth, T. L., Broccoli, A. J., Rosati, A., Stouffer, R. J., Balaji, V., Beesley, J. A., Cooke, W. F., Dixon, K. W., Dunne, J.,  
535 Dunne, K. A., Durachta, J. W., Findell, K. L., Ginoux, P., Gnanadesikan, A., Gordon, C. T., Griffies, S. M., Gudgel, R.,  
536 Harrison, M. J., Held, I. M., Hemler, R. S., Horowitz, L. W., Klein, S. A., Knutson, T. R., Kushner, P. J., Langenhorst,  
537 A. R., Lee, H. C., Lin, S. J., Lu, J., Malyshev, S. L., Milly, P. C. D., Ramaswamy, V., Russell, J., Schwarzkopf, M. D.,  
538 Shevliakova, E., Sirutis, J. J., Spelman, M. J., Stern, W. F., Winton, M., Wittenberg, A. T., Wyman, B., Zeng, F., and  
539 Zhang, R.: GFDL’s CM2 Global Coupled Climate Models. Part I: Formulation and Simulation Characteristics, *J. Clim.*,  
540 19, 643–674, <https://doi.org/10.1175/JCLI3629.1>, 2006.

541 Dijkshoorn, J. A., Engelen, V. W. P. V., and Huting, J. R. M.: Soil and landform properties for LADA partner countries  
542 (Argentina, China, Cuba, Senegal, South Africa and Tunisia), ISRIC–World Soil Information and FAO, Wageningen, the  
543 Netherlands, available at: <https://www.isric.org/>, <https://doi.org/10.13031/2013.42676>, 2008.

544 Donner, L. J., Wyman, B. L., Hemler, R. S., Horowitz, L. W., Ming, Y., Zhao, M., Golaz, J. C., Ginoux, P., Lin, S. J.,  
545 Schwarzkopf, M. D., Austin, J., Alaka, G., Cooke, W. F., Delworth, T. L., Freidenreich, S. M., Gordon, C. T., Griffies,  
546 S. M., Held, I. M., Hurlin, W. J., Klein, S. A., Knutson, T. R., Langenhorst, A. R., Lee, H. C., Lin, Y., Magi, B. I.,  
547 Malyshev, S. L., Milly, P. C. D., Naik, V., Nath, M. J., Pincus, R., Ploshay, J. J., Ramaswamy, V., Seman, C. J.,  
548 Shevliakova, E., Sirutis, J. J., Stern, W. F., Stouffer, R. J., Wilson, R. J., Winton, M., Wittenberg, A. T., and Zeng, F.:  
549 The Dynamical Core, Physical Parameterizations, and Basic Simulation Characteristics of the Atmospheric Component  
550 AM3 of the GFDL Global Coupled Model CM3, *J. Clim.*, 24, 3484–3519, <https://doi.org/10.1175/2011JCLI3955.1>, 2011.

551 Dufresne, J. L., Foujols, M. A., Denvil, S., Caubel, A., Marti, O., Aumont, O., Balkanski, Y., Bekki, S., Bellenger, H., Benschila,  
552 R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., Noblet,  
553 N. D., Duvel, J. P., Ethé, C., Fairhead, L., Fichefet, T., Flavoni, S., Friedlingstein, P., Grandpeix, J. Y., Guez, L., Guilyardi,

554 E., Hauglustaine, D., Hourdin, F., Idelkadi, A., Ghattas, J., Joussaume, S., Kageyama, M., Krinner, G., Labetoulle, S.,  
555 Lahellec, A., Lefebvre, M. P., Lefevre, F., Levy, C., Li, Z. X., Lloyd, J., Lott, F., Madec, G., Mancip, M., Marchand, M.,  
556 Masson, S., Meurdesoif, Y., Mignot, J., Musat, I., Parouty, S., Polcher, J., Rio, C., Schulz, M., Swingedouw, D., Szopa,  
557 S., Talandier, C., Terray, P., Viovy, N., and Vuichard, N.: Climate change projections using the IPSL-CM5 Earth System  
558 Model: from CMIP3 to CMIP5, *Clim. Dyn.*, 40, 2123–2165, <https://doi.org/10.1007/s00382-012-1636-1>, 2013.

559 Fader, M., Rost, S., Müller, C., Bondeau, A., and Gerten, D.: Virtual water content of temperate cereals and maize: Present  
560 and potential future patterns, *J. Hydrol.*, 384, 218–231, <https://doi.org/10.1016/j.jhydrol.2009.12.011>, 2010.

561 FAO: FAOSTAT on-line database, Food and Agriculture Organization of the United Nation, Rome, Italy, available at:  
562 <http://www.fao.org/faostat/en/#data/QC>, 2021.

563 Garofalo, P., Ventrella, D., Kersebaum, K. C., Gobin, A., Trnka, M., Giglio, L., Dubrovský, M., and Castellini, M.: Water  
564 footprint of winter wheat under climate change: Trends and uncertainties associated to the ensemble of crop models, *Sci.*  
565 *Total Environ.*, 658, 1186–1208, <https://doi.org/10.1016/j.scitotenv.2018.12.279>, 2019.

566 Guo, H., Li, S., Wong, F. L., Qin, S., Wang, Y., Yang, D., and Lam, H. M.: Drivers of carbon flux in drip irrigation maize  
567 fields in northwest China, *Carbon Balance Manag.*, 16, 12, <https://doi.org/10.1186/s13021-021-00176-5>, 2021.

568 Harris, I., Jones, P. D., Osborn, T. J., and Lister, D. H.: Updated high-resolution grids of monthly climatic observations - the  
569 CRU TS3.10 Dataset, *Int. J. Climatol.*, 34, 623–642, <https://doi.org/10.1002/joc.3711>, 2014.

570 Hatfield, J. L. and Dold, C.: Water-Use Efficiency: Advances and Challenges in a Changing Climate, *Front. Plant Sci.*, 10,  
571 103, <https://doi.org/10.3389/fpls.2019.00103>, 2019.

572 Hoekstra, A. Y. (ed.): Virtual water trade: Proceedings of the International Expert Meeting on Virtual Water Trade, Delft, the  
573 Netherlands, 12–13 December 2002, Value of Water Research Report Series No. 12, UNESCO-IHE, Delft, The  
574 Netherlands, 2003.

575 Hoekstra, A. Y.: The water footprint of modern consumer society, Routledge, London, UK, 208 pp, 2013.

576 Hoekstra, A. Y.: Sustainable, efficient, and equitable water use: the three pillars under wise freshwater allocation, *WIREs*  
577 *Water*, 1, 31–40, <https://doi.org/10.1002/wat2.1000>, 2014.

578 Hoekstra, A. Y., Chapagain, A. K., Aldaya, M. M., and Mekonnen, M. M.: The Water Footprint Assessment Manual: Setting  
579 the Global Standard, Earthscan, London, UK, 2011.

580 Hurrell, J. W., Holland, M. M., Gent, P. R., Ghan, S., Kay, J. E., Kushner, P. J., Lamarque, J. F., Large, W. G., Lawrence, D.,  
581 Lindsay, K., Lipscomb, W. H., Long, M. C., Mahowald, M., Marsh, D. R., Neale, R. B., Rasch, P., Vavrus, S., Vertenstein,  
582 M., Bader, D., Collins, W. D., Hack, J. J., Kiehl, J., and Marshall, S.: The Community Earth System Model: A Framework  
583 for Collaborative Research, *Bull. Am. Meteorol. Soc.*, 94, 1339–1360, <https://doi.org/10.1175/BAMS-D-12-00121.1>,  
584 2013.

585 IPCC: Summary for Policymakers. In: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I  
586 to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Masson-Delmotte, V.,  
587 Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. L., Huang, M.,

588 Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., Cambridge  
589 University Press, In Press, 2021.

590 Jans, Y., Bloh, W. V., Schaphoff, S., and Müller, C.: Global cotton production under climate change – Implications for yield  
591 and water consumption, *Hydrol. Earth Syst. Sci.*, 25, 2027–2044, <https://doi.org/10.5194/hess-25-2027-2021>, 2021.

592 Kappelle, M.: WMO Statement on the State of the Global Climate in 2019, World Meteorological Organization, Geneva,  
593 Switzerland, 2020.

594 Karandish, F., Nouri, H., and Schyns, J. F.: Agricultural adaptation to reconcile food security and water sustainability under  
595 climate change: the case of cereals in Iran, *Earths Future*, 10, e2021EF002095, <https://doi.org/10.1029/2021EF002095>,  
596 2022.

597 Konapala, G., Mishra, A. K., Wada, Y., and Mann, M. E.: Climate change will affect global water availability through  
598 compounding changes in seasonal precipitation and evaporation, *Nat. Commun.*, 11, 3044,  
599 <https://doi.org/10.1038/s41467-020-16757-w>, 2020.

600 Li, H., Mei, X., Nangia, V., Guo, R., Liu, Y., Hao, W., and Wang, J.: Effects of different nitrogen fertilizers on the yield,  
601 water- and nitrogen-use efficiencies of drip-fertigated wheat and maize in the North China Plain, *Agric Water Manag.*,  
602 243, 106474, <https://doi.org/10.1016/j.agwat.2020.106474>, 2021.

603 Liu, X., Li, C., Zhao, T., and Han, L.: Future changes of global potential evapotranspiration simulated from CMIP5 to CMIP6  
604 models, *Atmos. Oceanic Sci Lett.*, 13, 568–575, <https://doi.org/10.1080/16742834.2020.1824983>, 2020.

605 Lobell, D. B. and Gourdji, S. M.: The influence of climate change on global crop productivity, *Plant Physiol.*, 160, 1686–1697,  
606 <https://doi.org/10.1104/pp.112.208298>, 2012.

607 Mali, S. S., Shirsath, P. B., and Islam, A.: A high-resolution assessment of climate change impact on water footprints of cereal  
608 production in India, *Sci. Rep.*, 11, 8715, <https://doi.org/10.1038/s41598-021-88223-6>, 2021.

609 Meehl, G. A., Boer, G. J., Covey, C., Latif, M., and Stouffer, R. J.: Intercomparison makes for a better climate model, *Eos.*  
610 *Trans. Amer. Geophys. Union*, 78, 445–451, <https://doi.org/10.1029/97EO00276>, 1997.

611 Meehl, G. A., Boer, G. J., Covey, C., Latif, M., and Stouffer, R. J.: The Coupled Model Intercomparison Project (CMIP), *Bull.*  
612 *Am. Meteorol. Soc.*, 81, 313–318, [https://doi.org/10.1175/1520-0477\(2000\)081<0313:TCMIPC>2.3.CO;2](https://doi.org/10.1175/1520-0477(2000)081<0313:TCMIPC>2.3.CO;2), 2000.

613 Mekonnen, M. M. and Hoekstra, A. Y.: A global and high-resolution assessment of the green, blue and grey water footprint  
614 of wheat, *Hydrol. Earth Syst. Sci.*, 14, 1259–1276, <https://doi.org/10.5194/hess-14-1259-2010>, 2010.

615 Mekonnen, M. M. and Hoekstra, A. Y.: The green, blue and grey water footprint of crops and derived crop products, *Hydrol.*  
616 *Earth Syst. Sci.*, 15, 1577–1600, <https://doi.org/10.5194/hess-15-1577-2011>, 2011.

617 Mekonnen, M. M. and Hoekstra, A. Y.: Water footprint benchmarks for crop production: A first global assessment, *Ecol.*  
618 *Indic.*, 46, 214–223, <https://doi.org/10.1016/j.ecolind.2014.06.013>, 2014.

619 Mialyk, O., Schyns, J. F., Booij, M. J., and Hogeboom, R. J.: Historical simulation of maize water footprints with a new global  
620 gridded crop model ACEA, *Hydrol. Earth Syst. Sci.*, 26, 923–940, <https://doi.org/10.5194/hess-26-923-2022>, 2022.

621 Middleton, N. and Thomas, D. S. G.: World atlas of desertification, Arnold, London, UK, 1997.

622 Moss, R., Babiker, M., Brinkman, S., Calvo, E., Carter, T., Edmonds, J., Elgizouli, I., Emori, S., Erda, L., Hibbard, K., Jones,  
623 R., Kainuma, M., Kelleher, J., Lamarque, J. F., Manning, M., Matthews, B., Meehl, J., Meyer, L., Mitchell, J.,  
624 Nakicenovic, N., O'Neill, B., Pichs, R., Riahi, K., Rose, S., Runci, P., Stouffer, R., van Vuuren, D., Weyant, J., Wilbanks,  
625 T., van Ypersele, J. P., and Zurek, M.: Towards New Scenarios for Analysis of Emissions, Climate Change, Impacts, and  
626 Response Strategies, IPCC Expert Meeting Report, 19–21 September, 2007, Noordwijkerhout, Netherlands,  
627 Intergovernmental Panel on Climate Change (IPCC), Geneva, Switzerland, 132 pp, 2008.

628 Müller, C., Franke, J., Jägermeyr, J., Ruane, A. C., Elliott, J., Moyer, E., Heinke, J., Falloon, P. D., Folberth, C., Francois, L.,  
629 Hank, T., César Izaurralde, R., Jacquemin, I., Liu, W., Olin, S., Pugh, T. A. M., Williams, K., and Zabel, F.: Exploring  
630 uncertainties in global crop yield projections in a large ensemble of crop models and CMIP5 and CMIP6 climate scenarios,  
631 *Environ. Res. Lett.*, 16, 034040, <https://doi.org/10.1088/1748-9326/abd8fc>, 2021.

632 Myers, S. S., Smith, M. R., Guth, S., Golden, C. D., Vaitla, B., Mueller, N. D., Dangour, A. D., and Huybers, P.: Climate  
633 Change and Global Food Systems: Potential Impacts on Food Security and Undernutrition, *Annu. Rev. Public Health*, 38,  
634 259–277, <https://doi.org/10.1146/annurev-publhealth-031816-044356>, 2017.

635 Navarro-Racines, C., Tarapues, J., Thornton, P., Jarvis, A., and Ramirez-Villegas, J.: High-resolution and bias-corrected  
636 CMIP5 projections for climate change impact assessments, *Sci. Data*, 7, 7, <https://doi.org/10.1038/s41597-019-0343-8>,  
637 2020.

638 NBSC: National Data, China, National Bureau of Statistics, Beijing, China, available at: <https://data.stats.gov.cn/>, 2021.

639 NOAA: National Oceanic and Atmospheric Administration, U.S, available at: <https://www.esrl.noaa.gov>, 2018.

640 O'Neill, B. C., Tebaldi, C., van Vuuren, D. P., Eyring, V., Friedlingstein, P., Hurtt, G., Hurtt, R., Kriegler, E., Lamarque, J.  
641 F., Lowe, J., Meehl, G. A., Moss, R., Riahi, K., and Sanderson, B. M.: The Scenario Model Intercomparison Project  
642 (ScenarioMIP) for CMIP6, *Geosci. Model Dev.*, 9, 3461–3482, <https://doi.org/10.5194/gmd-9-3461-2016>, 2016.

643 Pastor, A. V., Palazzo, A., Havlik, P., Biemans, H., Wada, Y., Obersteiner, M., Kabat, P., and Ludwig, F.: The global nexus  
644 of food–trade–water sustaining environmental flows by 2050, *Nat. Sustain.*, 2, 499–507, [https://doi.org/10.1038/s41893-](https://doi.org/10.1038/s41893-019-0287-1)  
645 [019-0287-1](https://doi.org/10.1038/s41893-019-0287-1), 2019.

646 Portmann, F. T., Siebert, S., and Döll, P.: MIRCA2000-Global monthly irrigated and rainfed crop areas around the year 2000:  
647 A new high-resolution data set for agricultural and hydrological modelling, *Global Biogeochem. Cy.*, 24,  
648 <https://doi.org/10.1029/2008gb003435>, 2010.

649 Qiao, F., Song, Z., Bao, Y., Song, Y., Shu, Q., Huang, C., and Zhao, W.: Development and evaluation of an Earth System  
650 Model with surface gravity waves, *J. Geophys. Res. Ocean.*, 118, 4514–4524, <https://doi.org/10.1002/jgrc.20327>, 2013.

651 Raes, D., Steduto, P., Hsiao, T. C., and Fereres, E.: Reference manual, Chapter 2, AquaCrop model, Version 6.0, Food and  
652 Agriculture Organization of the United Nations, Rome, Italy, 2017.

653 Rallison, R. E.: Origin and evolution of the SCS runoff equation, in: Symposium on Watershed Management, Boise, Idaho,  
654 United States, 21–23 July, 912–924, 1980.

655 Riahi, K., Gruebler, A., and Nakicenovic, N.: Scenarios of long-term socio-economic and environmental development under  
656 climate stabilization, *Technol. Forecast. Soc. Chang.*, 74, 887–935, <https://doi.org/10.1016/j.techfore.2006.05.026>, 2007.

657 Rosa, L., Chiarelli, D. D., Sangiorgio, M., Beltran-Peña, A. A., Rulli, M. C., D’Odorico, P., and Fung, I.: Potential for  
658 sustainable irrigation expansion in a 3° C warmer climate, *Proc. Natl. Acad. Sci. U. S. A.*, 117, 29526–29534,  
659 <https://doi.org/10.1073/pnas.2017796117>, 2020.

660 Schmidt, G. A., Kelley, M., Nazarenko, L., Ruedy, R., Russell, G. L., Aleinov, I., Bauer, M., Bauer, S. E., Bhat, M. K., Bleck,  
661 R., Canuto, V., Chen, Y. H., Cheng, Y., Clune, T. L., Genio, A. D., Fainchtein, R. D., Faluvegi, G., Hansen, J. E., Healy,  
662 R. J., Kiang, N. Y., Koch, D., Lacis, A. A., LeGrande, A. N., Lerner, J., Lo, K. K., Matthews, E. E., Menon, S., Miller,  
663 R. L., Oinas, V., Oloso, A. O., Perlwitz, J. P., Puma, M. J., Putman, W. M., Rind, D., Romanou, A., Sato, M., Shindell,  
664 D. T., Sun, S., Syed, R. A., Tausnev, N., Tsigaridis, K., Unger, N., Voulgarakis, A., Yao, M. S., and Zhang, J.:  
665 Configuration and assessment of the GISS ModelE2 contributions to the CMIP5 archive, *J. Adv. Model. Earth Syst.*, 6,  
666 141–184, <https://doi.org/10.1002/2013MS000265>, 2014.

667 Schmidt, G. A., Ruedy, R., Hansen, J. E., Aleinov, I., Bell, N., Bauer, M., Bauer, S., Cairns, B., Canuto, V., Cheng, Y.,  
668 Delgenio, A. D., Faluvegi, G., Friend, A. D., Hall, T. M., Hu, Y., Kelley, M., Kiang, N. Y., Koch, D., Lacis, A. A., Lerner,  
669 J., Lo, K. K., Miller, R. L., Nazarenko, L., Oinas, V., Perlwitz, J., Perlwitz, J., Rind, D., Romanou, A., Russell, G. L.,  
670 Sato, M., Shindell, D. T., Stone, P. H., Sun, S., Tausnev, N., Thresher, D., and Yao, M. S.: Present-Day Atmospheric  
671 Simulations Using GISS ModelE: Comparison to In Situ, Satellite, and Reanalysis Data, *J. Clim.*, 19, 153–192,  
672 <https://doi.org/10.1175/JCLI3612.1>, 2006.

673 Schyns, J. F., Hogeboom, R. J., and Krol, M. S.: 4 - Water Footprint Assessment: towards water-wise food systems, in: *Food*  
674 *Systems Modelling*, edited by: Peters, C., and Thilmany, D., Academic Press, Salt Lake City, USA, 63–88,  
675 <https://doi.org/10.1016/B978-0-12-822112-9.00006-0>, 2022.

676 Semenov, M. A. and Stratonovitch, P.: Use of multi-model ensembles from global climate models for assessment of climate  
677 change impacts, *Clim. Res.*, 41, 1–14, <https://doi.org/10.3354/cr00836>, 2010.

678 Tian, Y., Ruth, M., Zhu, D., Ding, J., and Morris, N.: A Sustainability Assessment of Five Major Food Crops' Water Footprints  
679 in China from 1978 to 2010, *Sustainability*, 11, 1–20, <https://doi.org/10.3390/su11216179>, 2019.

680 Trnka, M., Feng, S., Semenov, M. A., Olesen, J. E., Kersebaum, K. C., Rötter, R. P., Semerádová, D., Klem, K., Huang, W.,  
681 Ruiz-Ramos, M., Hlavinka, P., Meitner, J., Balek, J., Havlík, P., and Büntgen, U.: Mitigation efforts will not fully alleviate  
682 the increase in water scarcity occurrence probability in wheat-producing areas, *Sci. Adv.*, 5, eaau2406,  
683 <https://doi.org/10.1126/sciadv.aau2406>, 2019.

684 USDA: Estimation of direct runoff from storm rainfall, Section 4 Hydrology, Chapter 4, *National Engineering Handbook*,  
685 Washington DC, USA, 1–24, 1964.

686 van Vuuren, D. P., den Elzen, M. G. J., Lucas, P. L., Eickhout, B., Strengers, B. J., van Ruijven, B., Wonink, S., and van Houdt,  
687 R.: Stabilizing greenhouse gas concentrations at low levels: an assessment of reduction strategies and costs, *Clim. Change*,  
688 81, 119–159, <https://doi.org/10.1007/s10584-006-9172-9>, 2007.

689 von Salzen, K., Scinocca, J. F., McFarlane, N. A., Li, J., Cole, J. N. S., Plummer, D., Verseghy, D., Reader, M. C., Ma, X.,  
690 Lazare, M., and Solheim, L.: The Canadian Fourth Generation Atmospheric Global Climate Model (CanAM4). Part I:  
691 Representation of Physical Processes, *Atmosphere-Ocean*, 51, 104–125, <https://doi.org/10.1080/07055900.2012.755610>,  
692 2013.

693 Wang, B., Feng, P., Liu, D. L., O’Leary, G. J., Macadam, I., Waters, C., Asseng, S., Cowie, A., Jiang, T., Xiao, D., Ruan, H.,  
694 He, J., and Yu, Q.: Sources of uncertainty for wheat yield projections under future climate are site-specific, *Nat. Food.*,  
695 1, 720–728, <https://doi.org/10.1038/s43016-020-00181-w>, 2020.

696 Wang, C., Guo, L., Li, Y., and Wang, Z.: Systematic Comparison of C3 and C4 Plants Based on Metabolic Network Analysis,  
697 *BMC Syst Biol.*, 6, S9, <https://doi.org/10.1186/1752-0509-6-S2-S9>, 2012.

698 Wang, J., Gao, S., Xu, Y., and Wang, H.: Application and Existing Problems of Drip Irrigation for Wheat in Xinjiang, in: 2011  
699 International Conference on Agricultural and Natural Resources Engineering (ANRE 2011), Intelligent Information  
700 Technology Application Association, Singapore, 3 July 2011, 25–29, 2011.

701 Wang, W., Zhuo, L., Li, M., Liu, Y., and Wu, P.: The Effect of Development in Water-Saving Irrigation Techniques on Spatial-  
702 Temporal Variations in Crop Water Footprint and Benchmarking, *J. Hydrol.*, 577, 123916,  
703 <https://doi.org/10.1016/j.jhydrol.2019.123916>, 2019.

704 Xiao, D., Liu, D. L., Wang, B., Feng, P., Bai, H., and Tang, J.: Climate change impact on yields and water use of wheat and  
705 maize in the North China Plain under future climate change scenarios, *Agr. Water Manage.*, 238, 106238, 2020.

706 Xu, Z., Chen, X., Wu, S. R., Gong, M., Du, Y., Wang, J., Li, Y., and Liu, J.: Spatial-temporal assessment of water footprint,  
707 water scarcity and crop water productivity in a major crop production region, *J. Clean. Prod.*, 224, 375–383,  
708 <https://doi.org/10.1016/j.jclepro.2019.03.108>, 2019.

709 Yoon, P. R. and Choi, J. Y.: Effects of shift in growing season due to climate change on rice yield and crop water requirements,  
710 *Paddy Water Environ.*, 18, 291–307, <https://doi.org/10.1007/s10333-019-00782-7>, 2020.

711 Zain, M., Si, Z., Chen, J., Mehmood, F., Rahman, S. U., Shah, A. N., Li, S., Gao, Y., and Duan, A.: Suitable nitrogen  
712 application mode and lateral spacing for drip-irrigated winter wheat in North China Plain, *PLoS One*,  
713 <https://doi.org/10.1371/journal.pone.0260008>, 2021.

714 Zheng, J., Wang, W., Ding, Y., Liu, G., Xing, W., Cao, X., and Chen, D.: Assessment of climate change impact on the water  
715 footprint in rice production: Historical simulation and future projections at two representative rice cropping sites of China,  
716 *Sci. Total Environ.*, 709, 136190, <https://doi.org/10.1016/j.scitotenv.2019.136190>, 2020.

717 Zhuo, L., Liu, Y., Yang, H., Hoekstra, A. Y., Liu, W., Cao, X., Wang, M., and Wu, P.: Water for maize for pigs for pork: An  
718 analysis of inter-provincial trade in China, *Water Res.*, 166, 115074, <https://doi.org/10.1016/j.watres.2019.115074>, 2019.

719 Zhuo, L., Mekonnen, M. M., and Hoekstra, A. Y.: Sensitivity and uncertainty in crop water footprint accounting: a case study  
720 for the Yellow River basin, *Hydrol. Earth Syst. Sci.*, 18, 2219–2234, <https://doi.org/10.5194/hess-18-2219-2014>, 2014.

721 Zhuo, L., Mekonnen, M. M., and Hoekstra, A. Y.: Benchmark levels for the consumptive water footprint of crop production  
722 for different environmental conditions: a case study for winter wheat in China, *Hydrol. Earth Syst. Sci.*, 20, 4547–4559,  
723 <https://doi.org/10.5194/hess-20-4547-2016>, 2016a.

724 Zhuo, L., Mekonnen, M. M., Hoekstra, A. Y., and Wada, Y.: Inter- and intra-annual variation of water footprint of crops and  
725 blue water scarcity in the Yellow River basin (1961–2009), *Adv. Water Resour.*, 87, 29–41,  
726 <https://doi.org/10.1016/j.advwatres.2015.11.002>, 2016b.

727 Zhuo, L., Mekonnen, M. M., and Hoekstra, A. Y.: The effect of inter-annual variability of consumption, production, trade and  
728 climate on crop-related green and blue water footprints and inter-regional virtual water trade: A study for China (1978-  
729 2008), *Water Res.*, 94, 73–85, <https://doi.org/10.1016/j.watres.2016.02.037>, 2016c.

730 Zhuo, L., Mekonnen, M. M., and Hoekstra, A. Y.: Consumptive water footprint and virtual water trade scenarios for China —  
731 With a focus on crop production, consumption and trade, *Environ. Int.*, 94, 211–223,  
732 <https://doi.org/10.1016/j.envint.2016.05.019>, 2016d.